

# TABLES

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites—School Health Education Profiles, Principals' and Teachers' Surveys, 2000

**STATE SURVEYS**

Site	Principals' surveys		Teachers' surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
<b>Weighted Data</b>				
Alabama	304	89	271	79
Alaska	265	82	236	73
Arkansas	328	78	314	74
California	369	73	NA*	NA
Delaware	56	85	47	71
Hawaii	65	77	62	74
Idaho	215	89	202	84
Illinois**	391	85	356	77
Iowa	322	93	294	84
Maine	215	98	200	91
Massachusetts	573	82	563	81
Michigan	331	84	298	76
Minnesota	312	84	306	82
Missouri	334	80	335	80
Montana	255	74	248	72
Nebraska	333	82	318	78
New Hampshire	167	80	154	73
New Jersey	316	75	307	73
North Dakota	163	76	155	72
Ohio	368	79	344	74
Oklahoma	332	78	326	77
Pennsylvania**	305	71	NA	NA
Tennessee	254	73	254	73
Utah	192	72	188	70
Virginia	275	75	261	71
West Virginia	259	89	235	80
<b>Unweighted Data</b>				
California	NA	NA	331	66
Georgia	231	63	217	59
Indiana	262	67	256	65
Kentucky	282	65	229	53
Louisiana**	214	62	207	60
Maryland**	166	53	155	50
Pennsylvania**	NA	NA	285	66
South Carolina	229	53	228	53
South Dakota	148	60	NA	NA
Texas**	340	67	321	64

(continued)

TABLE 1. Sample Sizes and Response Rates, Selected U.S. Sites—School Health Education Profiles, Principals' and Teachers' Surveys, 2000 (*continued*)**LOCAL SURVEYS**

Site	Principals' surveys		Teachers' surveys	
	Sample size	Response rate (%)	Sample size	Response rate (%)
<b>Weighted Data</b>				
Chicago	242	79	235	77
Dallas	43	86	43	86
District of Columbia	40	89	39	87
Ft. Lauderdale	57	78	57	78
Los Angeles	100	79	98	77
Miami	87	82	91	86
New Orleans	24	100	24	100
Orange County	37	82	39	87
Palm Beach	38	88	36	84
Philadelphia	98	78	92	74
San Diego	43	93	45	98
San Francisco	33	75	31	70
<b>Unweighted Data</b>				
Houston	35	58	36	60

\* NA = not available.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 2. Percentage of Schools That Required Health Education in Grades 6–12 and, Among Those Schools, Percentage That Taught ≥1 Separate Health Education Course, Selected U.S. Sites—School Health Education Profiles, Principals’ Surveys, 2000

**STATE SURVEYS**

Site	Required health education	Taught ≥1 separate health education course
<b>Weighted Data</b>		
Alabama	87.3	84.8
Alaska	86.8	97.7
Arkansas	96.0	99.4
California	83.3	78.1
Delaware	98.1	98.1
Hawaii	100.0	100.0
Idaho	97.2	98.0
Illinois*	93.9	93.6
Iowa	75.7	93.0
Maine	90.1	96.3
Massachusetts	92.3	95.8
Michigan	82.4	94.0
Minnesota	98.4	99.6
Missouri	86.9	91.6
Montana	93.2	89.2
Nebraska	89.9	89.0
New Hampshire	90.1	95.0
New Jersey	96.7	97.9
North Dakota	94.2	96.8
Ohio	97.9	99.1
Oklahoma	31.4	80.1
Pennsylvania*	96.8	99.6
Tennessee	84.3	89.8
Utah	97.8	99.4
Virginia	90.6	89.9
West Virginia	98.4	96.8
<b>Unweighted Data</b>		
Georgia	99.1	97.3
Indiana	98.1	97.2
Kentucky	91.0	97.6
Louisiana*	91.5	87.3
Maryland*	91.9	92.3
South Carolina	85.8	82.7
South Dakota	79.6	91.0
Texas*	84.5	91.2
<b>State Median</b>	<b>91.7</b>	<b>95.4</b>

(continued)

TABLE 2. Percentage of Schools That Required Health Education in Grades 6–12 and, Among Those Schools, Percentage That Taught  $\geq 1$  Separate Health Education Course, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	Required health education	Taught $\geq 1$ separate health education course
<b>Weighted Data</b>		
Chicago	81.8	73.7
Dallas	64.8	100.0
District of Columbia	88.0	97.0
Ft. Lauderdale	88.4	91.6
Los Angeles	98.9	98.9
Miami	67.9	78.7
New Orleans	100.0	100.0
Orange County	86.5	69.0
Palm Beach	80.6	88.1
Philadelphia	93.4	94.8
San Diego	0.0	NA**
San Francisco	93.8	73.3
<b>Unweighted Data</b>		
Houston	91.4	100.0
<b>Local Median</b>	<b>88.0</b>	<b>93.2</b>

\* Survey did not include students from one of the state's large school districts.

\*\* NA = not available.

TABLE 3. Percentage of Schools With a Required Health Education Course That Required Teachers To Use Standards, a Specific Curriculum, Guidelines, Framework, or Other Selected Materials, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

**STATE SURVEYS**

Site	National Health Education Standards	State, district, or school curriculum, guidelines, or framework	Materials from health organizations	Commercial teacher's guide
<b>Weighted Data</b>				
Alabama	46.3	99.5	55.2	72.1
Alaska	31.4	95.7	25.5	45.5
Arkansas	37.7	97.1	36.2	68.4
Delaware	24.5	85.2	10.5	34.0
Hawaii	60.8	90.5	16.3	21.6
Idaho	16.5	87.7	24.6	55.9
Illinois*	32.2	95.4	36.6	58.9
Iowa	32.4	91.9	33.1	41.4
Maine	26.9	92.3	19.1	20.5
Massachusetts	37.1	96.9	42.3	37.7
Michigan	18.5	93.0	38.2	39.9
Minnesota	29.0	92.1	33.5	35.8
Missouri	35.6	96.4	31.2	51.1
Montana	25.8	87.3	31.8	46.1
Nebraska	24.7	90.0	34.4	52.1
New Hampshire	24.1	81.6	26.6	34.6
New Jersey	37.3	99.6	48.4	56.1
North Dakota	20.4	81.0	26.7	44.2
Ohio	33.3	98.6	27.5	54.0
Oklahoma	30.2	96.6	59.0	55.0
Tennessee	32.0	98.2	33.1	61.5
Utah	22.3	98.9	26.5	39.8
Virginia	26.0	95.7	41.1	68.0
West Virginia	40.5	99.5	40.6	80.1
<b>Unweighted Data</b>				
California	35.8	97.6	48.0	51.7
Georgia	34.2	100.0	41.7	76.7
Indiana	46.8	97.0	42.1	72.1
Kentucky	39.5	98.0	29.9	53.9
Louisiana*	24.1	94.1	44.9	69.3
Maryland*	44.5	98.5	43.1	50.4
Pennsylvania*	41.2	95.9	44.7	51.3
South Carolina	40.8	94.3	38.8	69.7
Texas*	29.6	97.5	42.5	68.5
<b>State Median</b>	<b>32.2</b>	<b>95.9</b>	<b>36.2</b>	<b>52.1</b>

(continued)

TABLE 3. Percentage of Schools With a Required Health Education Course That Required Teachers To Use Standards, a Specific Curriculum, Guidelines, Framework, or Other Selected Materials, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

### LOCAL SURVEYS

Site	National Health Education Standards	State, district, or school curriculum, guidelines, or framework	Materials from health organizations	Commercial teacher's guide
<b>Weighted Data</b>				
Chicago	47.0	97.4	60.0	54.7
Dallas	59.4	100.0	78.7	78.2
District of Columbia	30.6	93.9	45.2	66.6
Ft. Lauderdale	43.1	100.0	73.9	72.1
Los Angeles	37.1	100.0	34.8	41.9
Miami	40.6	98.3	70.8	62.0
New Orleans	73.7	100.0	75.0	66.7
Orange County	52.4	100.0	45.5	52.4
Palm Beach	41.6	100.0	64.9	65.0
Philadelphia	27.1	96.9	38.3	32.7
San Diego	NA**	NA	NA	NA
San Francisco	56.3	95.2	75.0	66.7
<b>Unweighted Data</b>				
Houston	50.0	100.0	77.4	58.1
<b>Local Median</b>	<b>45.1</b>	<b>100.0</b>	<b>67.9</b>	<b>63.5</b>

\* Survey did not include students from one of the state's large school districts.

\*\* NA = not available.

TABLE 4. Percentage of Schools That Tried To Increase Student Knowledge in Specific Topics,\* Selected U.S. Sites—  
School Health Education Profiles, Teachers’ Surveys, 2000

**STATE SURVEYS**

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV** prevention	Physical activity and fitness	Pregnancy prevention	STD <sup>§</sup> prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Alabama	100.0	94.9	98.0	98.1	88.8	94.0	82.4	100.0	90.0
Alaska	95.4	88.6	86.5	92.4	71.5	75.2	73.4	92.9	79.3
Arkansas	99.3	92.1	96.3	92.7	80.5	87.5	82.9	99.7	82.1
Delaware	100.0	86.5	100.0	95.7	97.5	97.5	77.0	100.0	88.9
Hawaii	100.0	98.6	100.0	97.2	94.4	100.0	69.2	100.0	94.4
Idaho	98.9	96.8	94.8	95.6	70.6	89.1	77.3	98.8	78.3
Illinois <sup>§§</sup>	99.7	91.0	98.6	92.5	82.3	95.3	73.0	98.9	77.2
Iowa	98.5	94.5	98.1	93.9	88.8	94.6	73.7	99.1	72.4
Maine	97.3	93.0	97.7	88.1	84.7	93.2	70.1	98.2	84.2
Massachusetts	98.8	95.2	96.9	94.0	84.1	90.6	74.3	99.2	94.9
Michigan	99.2	90.7	96.6	93.2	79.1	92.1	57.6	97.9	82.5
Minnesota	98.9	91.4	98.6	92.2	90.6	95.1	81.5	99.3	82.0
Missouri	99.6	95.3	96.2	94.6	82.0	90.2	73.2	99.2	81.8
Montana	98.0	90.5	94.8	97.0	76.5	84.0	60.7	98.8	81.9
Nebraska	97.8	90.2	95.1	91.6	76.0	85.1	67.1	98.1	74.0
New Hampshire	98.0	94.1	96.2	92.2	83.5	90.5	73.1	97.1	80.1
New Jersey	100.0	93.2	99.6	95.3	89.1	97.1	76.6	100.0	87.0
North Dakota	100.0	94.3	96.9	94.3	74.2	89.6	77.9	100.0	83.1
Ohio	99.0	94.9	NA <sup>+</sup>	92.8	89.5	NA	77.7	100.0	78.3
Oklahoma	94.6	86.9	94.3	94.6	77.9	77.2	56.6	94.4	80.4
Tennessee	98.8	96.5	98.1	98.1	85.9	91.6	74.6	100.0	82.8
Utah	99.4	96.3	99.4	95.8	76.7	91.0	90.4	99.4	91.9
Virginia	98.8	92.8	97.4	97.3	86.0	94.2	70.1	97.4	78.2
West Virginia	100.0	96.2	98.1	97.3	87.7	93.0	79.9	100.0	93.0
<b>Unweighted Data</b>									
California	99.5	93.4	99.0	93.4	91.0	96.2	67.5	99.0	82.9
Georgia	100.0	93.6	98.0	95.1	89.2	93.1	83.8	100.0	89.7
Indiana	100.0	94.9	99.6	96.2	92.6	98.3	84.1	99.6	85.5
Kentucky	99.5	97.0	98.0	96.5	87.4	94.5	75.0	99.5	87.4
Louisiana <sup>§§</sup>	97.7	85.9	74.7	93.6	45.0	62.4	58.0	98.2	87.2
Maryland <sup>§§</sup>	100.0	92.3	97.7	90.8	91.3	95.3	77.3	100.0	84.7
Pennsylvania <sup>§§</sup>	100.0	91.0	99.6	96.6	87.6	94.8	68.8	100.0	75.5
South Carolina	97.5	93.6	98.7	92.3	87.9	93.8	56.7	98.7	78.3
Texas <sup>§§</sup>	99.2	93.9	96.3	93.9	80.7	89.8	76.5	98.8	85.0
<b>State Median</b>	<b>99.2</b>	<b>93.6</b>	<b>97.8</b>	<b>94.3</b>	<b>85.9</b>	<b>93.1</b>	<b>74.3</b>	<b>99.2</b>	<b>82.8</b>

(continued)



TABLE 4. Percentage of Schools That Tried To Increase Student Knowledge in Specific Topics,\* Selected U.S. Sites—  
School Health Education Profiles, Teachers' Surveys, 2000 (continued)

#### LOCAL SURVEYS

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV** prevention	Physical activity and fitness	Pregnancy prevention	STD§ prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Chicago	98.1	92.9	95.8	96.1	79.5	88.2	50.6	98.0	90.8
Dallas	100.0	100.0	100.0	100.0	91.0	100.0	95.2	100.0	100.0
District of Columbia	100.0	93.5	100.0	93.5	93.9	100.0	50.2	97.0	87.4
Ft. Lauderdale	100.0	98.1	98.2	92.0	91.0	98.1	77.9	100.0	86.1
Los Angeles	100.0	100.0	100.0	95.7	94.4	98.9	81.5	100.0	87.2
Miami	100.0	95.0	100.0	84.9	91.6	98.4	77.4	98.3	89.9
New Orleans	100.0	100.0	100.0	95.2	95.2	100.0	66.7	95.0	90.5
Orange County	100.0	95.8	100.0	95.5	100.0	100.0	81.8	100.0	87.0
Palm Beach	100.0	100.0	100.0	92.7	96.3	96.2	85.0	100.0	92.5
Philadelphia	100.0	90.1	100.0	97.4	86.1	94.6	63.8	98.4	85.6
San Diego	NA	NA	NA	NA	NA	NA	NA	NA	NA
San Francisco	100.0	95.0	100.0	100.0	90.0	95.0	94.4	100.0	100.0
<b>Unweighted Data</b>									
Houston	100.0	96.9	100.0	96.9	96.9	100.0	81.3	100.0	97.0
<b>Local Median</b>	<b>100.0</b>	<b>96.4</b>	<b>100.0</b>	<b>95.6</b>	<b>92.8</b>	<b>98.6</b>	<b>79.6</b>	<b>100.0</b>	<b>90.2</b>

\* In a required health education course.

\*\* HIV = human immunodeficiency virus.

§ STD = sexually transmitted disease.

§§ Survey did not include students from one of the state's large school districts.

+ NA = not available.

TABLE 5. Percentage of Schools That Tried To Improve Specific Student Skills,\* Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

**STATE SURVEYS**

Site	Analysis of media messages	Communication	Decision making	Goal setting	Conflict resolution	Resisting peer pressure	Stress management
<b>Weighted Data</b>							
Alabama	79.0	89.5	97.4	92.8	91.8	97.5	89.2
Alaska	65.3	87.2	93.8	87.2	82.7	88.9	78.5
Arkansas	68.5	90.5	97.0	95.0	88.7	97.5	90.4
Delaware	88.2	95.7	98.1	93.8	100.0	98.1	96.2
Hawaii	92.4	97.2	98.6	95.8	81.5	97.2	91.3
Idaho	78.0	91.2	97.8	94.1	85.0	94.6	91.8
Illinois**	80.0	90.2	97.3	92.3	82.8	96.5	88.6
Iowa	81.1	89.7	96.1	92.0	78.7	94.3	91.6
Maine	90.2	95.0	96.7	96.1	81.2	95.6	89.2
Massachusetts	93.6	94.6	99.4	93.3	92.6	98.2	89.5
Michigan	79.6	91.3	98.1	88.9	83.9	97.7	82.0
Minnesota	85.3	90.6	98.6	91.6	80.6	96.5	90.1
Missouri	76.3	93.1	98.6	95.0	88.6	98.3	94.7
Montana	78.8	86.5	91.2	84.8	81.6	94.8	82.5
Nebraska	73.4	87.0	95.1	89.3	80.5	94.9	86.3
New Hampshire	88.0	93.8	99.1	89.7	86.4	95.9	87.1
New Jersey	91.5	94.6	99.3	95.2	92.4	98.9	89.1
North Dakota	81.0	91.8	97.7	89.8	84.7	96.1	93.2
Ohio	85.8	90.8	97.3	92.3	85.4	98.4	89.6
Oklahoma	62.3	85.4	98.2	87.4	92.7	96.3	78.2
Tennessee	70.0	89.4	95.8	91.4	85.6	96.3	84.9
Utah	91.7	97.7	97.6	97.6	93.2	98.0	98.8
Virginia	86.3	92.8	99.1	94.9	86.8	96.6	86.1
West Virginia	85.6	96.1	98.6	98.6	95.8	99.0	93.8
<b>Unweighted Data</b>							
California	80.6	87.7	95.7	89.1	79.1	96.7	75.8
Georgia	85.4	94.1	98.5	93.7	93.2	98.5	93.2
Indiana	90.6	95.7	99.6	94.0	89.7	99.1	96.6
Kentucky	80.2	90.4	98.5	92.5	88.4	97.5	92.3
Louisiana**	79.8	88.8	94.7	94.7	89.5	97.1	86.5
Maryland**	91.4	92.2	97.7	93.8	90.8	98.5	92.2
Pennsylvania**	86.5	92.9	97.0	90.3	82.0	98.5	89.6
South Carolina	75.0	89.8	96.9	95.0	85.5	95.6	81.9
Texas**	73.0	93.9	95.5	93.1	89.1	99.2	91.9
<b>State Median</b>	<b>81.0</b>	<b>91.3</b>	<b>97.7</b>	<b>93.1</b>	<b>86.4</b>	<b>97.2</b>	<b>89.6</b>

(continued)

TABLE 5. Percentage of Schools That Tried To Improve Specific Student Skills,\* Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)**LOCAL SURVEYS**

Site	Analysis of media messages	Communication	Decision making	Goal setting	Conflict resolution	Resisting peer pressure	Stress management
<b>Weighted Data</b>							
Chicago	59.5	87.9	94.0	93.3	93.3	94.7	77.5
Dallas	78.2	95.7	100.0	100.0	95.7	95.2	95.2
District of Columbia	75.1	90.3	93.9	84.6	84.6	93.9	72.1
Ft. Lauderdale	81.2	92.8	95.6	95.6	91.1	98.2	88.2
Los Angeles	90.1	96.7	97.8	95.6	89.1	99.0	80.3
Miami	76.3	93.1	96.5	96.5	96.7	98.2	84.4
New Orleans	57.1	90.5	100.0	90.5	90.5	95.2	81.0
Orange County	83.3	95.8	95.8	95.8	100.0	100.0	95.8
Palm Beach	77.6	100.0	100.0	100.0	96.2	96.3	100.0
Philadelphia	76.2	90.1	98.8	94.4	89.5	97.8	73.2
San Diego	NA <sup>§</sup>	NA	NA	NA	NA	NA	NA
San Francisco	84.2	100.0	100.0	95.0	90.0	100.0	94.7
<b>Unweighted Data</b>							
Houston	81.8	100.0	100.0	100.0	97.0	100.0	96.9
<b>Local Median</b>	<b>77.9</b>	<b>94.4</b>	<b>98.3</b>	<b>95.6</b>	<b>92.2</b>	<b>98.0</b>	<b>86.3</b>

\* In a required health education course.

\*\* Survey did not include students from one of the state's large school districts.

§ NA = not available.

TABLE 6. Percentage of Schools That Taught Specific Topics Related to HIV\* Infection/AIDS\*\* Prevention,<sup>§</sup> Selected U.S. Sites—School Health Education Profiles, Teachers’ Surveys, 2000

**STATE SURVEYS**

Site	Abstinence to avoid HIV	How HIV is transmitted	How to correctly use a condom	Condom efficacy	Number of young people who get HIV	How to find information on HIV
<b>Weighted Data</b>						
Alabama	96.5	97.0	31.6	71.3	92.1	90.4
Alaska	78.8	82.7	37.4	64.2	70.9	64.2
Arkansas	92.0	93.4	25.6	62.6	85.2	78.0
Delaware	100.0	95.1	56.5	76.6	90.8	88.3
Hawaii	98.6	98.6	68.5	84.3	92.0	91.5
Idaho	93.3	93.2	23.1	57.8	87.6	76.3
Illinois <sup>SS</sup>	97.5	98.6	34.1	73.0	88.0	78.4
Iowa	96.1	96.6	48.7	83.4	89.6	82.0
Maine	96.5	97.2	61.7	84.1	85.6	86.6
Massachusetts	95.0	95.4	52.3	78.0	87.6	82.1
Michigan	95.0	94.8	31.3	67.4	86.2	79.9
Minnesota	97.9	98.0	46.6	83.5	91.8	83.8
Missouri	94.0	94.7	27.6	65.7	85.5	79.4
Montana	92.5	91.3	34.2	68.5	88.1	81.3
Nebraska	89.4	91.6	27.4	63.0	81.9	70.2
New Hampshire	94.1	94.4	55.4	77.4	86.5	82.1
New Jersey	97.9	99.3	58.0	81.7	91.5	87.5
North Dakota	92.4	93.8	25.9	60.8	85.7	72.9
Ohio	97.3	NA <sup>+</sup>	NA	NA	90.4	NA
Oklahoma	90.4	88.7	29.8	63.8	82.9	75.5
Tennessee	95.1	94.6	34.0	64.0	85.4	83.6
Utah	98.8	98.8	9.5	43.2	95.4	81.5
Virginia	92.9	93.8	NA	NA	86.5	83.5
West Virginia	94.3	95.3	48.2	76.1	90.2	86.4
<b>Unweighted Data</b>						
California	96.7	97.2	53.8	82.4	88.6	84.3
Georgia	98.0	97.5	27.4	76.5	92.4	84.3
Indiana	98.7	98.7	NA	NA	95.6	88.2
Kentucky	96.4	97.4	41.2	68.9	87.8	77.4
Louisiana <sup>SS</sup>	70.3	69.0	14.3	40.6	67.7	61.7
Maryland <sup>SS</sup>	94.6	95.4	46.5	80.0	90.0	83.7
Pennsylvania <sup>SS</sup>	99.2	99.2	50.8	82.2	95.1	86.4
South Carolina	98.1	97.4	53.2	71.2	86.4	69.5
Texas <sup>SS</sup>	95.0	93.8	27.4	63.8	85.1	76.3
<b>State Median</b>	<b>95.1</b>	<b>95.3</b>	<b>35.8</b>	<b>71.2</b>	<b>87.8</b>	<b>82.1</b>

(continued)

TABLE 6. Percentage of Schools That Taught Specific Topics Related to HIV\* Infection/AIDS\*\* Prevention,<sup>§</sup> Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	Abstinence to avoid HIV	How HIV is transmitted	How to correctly use a condom	Condom efficacy	Number of young people who get HIV	How to find information on HIV
<b>Weighted Data</b>						
Chicago	92.3	93.1	44.5	67.2	86.3	83.5
Dallas	100.0	100.0	29.8	73.9	100.0	95.7
District of Columbia	97.1	97.0	90.9	87.9	93.9	97.0
Ft. Lauderdale	98.1	98.1	75.6	84.0	98.1	95.5
Los Angeles	98.9	100.0	69.7	91.0	93.3	85.6
Miami	100.0	100.0	80.9	91.2	100.0	98.4
New Orleans	100.0	100.0	85.7	100.0	100.0	100.0
Orange County	100.0	100.0	54.2	95.8	95.8	91.3
Palm Beach	100.0	100.0	62.9	96.1	96.2	92.5
Philadelphia	97.3	98.4	61.5	81.0	92.8	92.1
San Diego	NA	NA	NA	NA	NA	NA
San Francisco	100.0	100.0	90.5	95.2	95.2	95.2
<b>Unweighted Data</b>						
Houston	100.0	100.0	51.6	90.6	100.0	100.0
<b>Local Median</b>	<b>100.0</b>	<b>100.0</b>	<b>66.3</b>	<b>90.8</b>	<b>96.0</b>	<b>95.4</b>

\* HIV = human immunodeficiency virus.

\*\* AIDS = acquired immunodeficiency syndrome.

§ In a required health education course.

§§ Survey did not include students from one of the state's large school districts.

+ NA = not available.

TABLE 7. Percentage of Schools in Which a Specific Person Was Responsible for Coordinating Health Education, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000

**STATE SURVEYS**

Site	School district administrator*	School administrator**	Health education teacher	School nurse	No coordinator
<b>Weighted Data<sup>§</sup></b>					
Alabama	16.2	32.6	39.4	6.7	5.2
Alaska	24.3	31.7	28.3	0.9	14.9
Arkansas	19.6	26.4	47.8	1.9	4.2
California	22.1	22.3	39.8	5.7	10.2
Delaware	15.8	13.7	68.5	0.0	2.0
Hawaii	4.7	14.7	75.6	0.0	5.0
Idaho	21.7	18.3	54.4	2.1	3.6
Illinois <sup>§§</sup>	23.4	24.2	46.7	0.3	5.4
Iowa	31.5	15.1	44.1	3.1	6.2
Maine	14.9	10.5	59.1	8.0	7.5
Massachusetts	60.1	9.3	25.2	2.3	3.1
Michigan	35.9	18.3	40.3	1.5	4.0
Minnesota	23.0	14.0	58.6	0.8	3.6
Missouri	28.8	22.2	42.7	4.3	2.0
Montana	19.8	13.0	60.8	2.0	4.3
Nebraska	23.8	21.7	41.8	3.0	9.7
New Hampshire	9.9	19.3	53.5	9.4	7.8
New Jersey	52.0	19.9	19.7	5.8	2.5
North Dakota	19.5	25.5	49.2	0.0	5.7
Ohio	30.1	18.3	47.7	0.3	3.6
Oklahoma	19.8	35.3	16.6	13.8	14.5
Pennsylvania <sup>§§</sup>	40.3	20.8	36.6	0.4	1.9
Tennessee	21.6	30.8	32.3	2.7	12.5
Utah	19.2	20.7	57.8	1.2	1.2
Virginia	25.2	17.4	51.3	1.7	4.4
West Virginia	14.8	22.5	60.2	0.0	2.5
<b>Unweighted Data<sup>§</sup></b>					
Georgia	29.0	27.6	41.9	0.9	0.5
Indiana	21.0	20.2	56.0	0.0	2.9
Kentucky	4.4	28.4	62.8	0.8	3.6
Louisiana <sup>§§</sup>	21.1	17.4	55.3	1.1	5.3
Maryland <sup>§§</sup>	44.7	5.9	44.7	0.7	3.9
South Carolina	24.5	23.6	42.8	2.9	6.3
South Dakota	23.2	27.5	32.6	1.4	15.2
Texas <sup>§§</sup>	26.5	26.2	37.5	2.5	7.3
<b>State Median</b>	<b>22.5</b>	<b>20.7</b>	<b>45.7</b>	<b>1.6</b>	<b>4.4</b>

(continued)

TABLE 7. Percentage of Schools in Which a Specific Person Was Responsible for Coordinating Health Education, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	School district administrator*	School administrator**	Health education teacher	School nurse	No coordinator
<b>Weighted Data<sup>§</sup></b>					
Chicago	3.9	23.1	44.1	11.5	17.4
Dallas	27.2	14.9	37.0	6.1	14.9
District of Columbia	2.6	0.0	86.7	5.5	5.2
Ft. Lauderdale	19.8	25.1	50.8	0.0	4.3
Los Angeles	12.2	25.5	57.4	1.2	3.7
Miami	26.3	42.9	23.9	1.5	5.4
New Orleans	34.8	8.7	56.5	0.0	0.0
Orange County	26.7	30.0	40.0	0.0	3.3
Palm Beach	27.9	35.6	33.7	0.0	2.9
Philadelphia	21.5	12.6	60.1	2.2	3.6
San Diego	55.1	22.4	0.0	15.0	7.4
San Francisco	12.5	16.7	70.8	0.0	0.0
<b>Unweighted Data<sup>§</sup></b>					
Houston	21.9	15.6	62.5	0.0	0.0
<b>Local Median</b>	<b>21.9</b>	<b>22.4</b>	<b>50.8</b>	<b>1.2</b>	<b>3.7</b>

\* District superintendent or district health education or curriculum coordinator.

\*\* Principal or school curriculum coordinator.

§ Percentages for each row might not add up to 100.0 due to rounding.

§§ Survey did not include students from one of the state's large school districts.

TABLE 8. Percentage of Schools in Which Health Education Teachers Planned or Coordinated Health-Related Projects or Activities With Other Groups, Selected U.S. Sites—School Health Education Profiles, Teachers' Survey, 2000

**STATE SURVEYS**

Site	Physical education staff	School health services staff	School mental health staff	Food service staff	Community members
<b>Weighted Data</b>					
Alabama	69.1	68.2	55.5	26.1	45.7
Alaska	47.6	45.3	52.9	17.3	56.1
Arkansas	64.8	61.5	39.5	14.9	39.8
Delaware	90.1	82.7	56.9	16.4	51.0
Hawaii	68.6	45.4	50.7	14.2	68.9
Idaho	62.7	57.0	52.9	12.9	49.7
Illinois*	65.0	60.0	50.4	13.4	40.6
Iowa	56.1	73.6	47.7	16.6	53.9
Maine	67.8	75.9	56.0	19.1	57.0
Massachusetts	76.4	85.5	78.9	28.6	74.6
Michigan	57.1	37.7	50.8	15.1	50.1
Minnesota	68.9	74.5	62.0	22.4	59.1
Missouri	80.7	80.5	57.5	24.9	51.9
Montana	82.8	49.2	51.3	18.2	47.5
Nebraska	61.6	61.1	36.0	13.9	42.3
New Hampshire	53.9	79.0	63.8	23.6	56.4
New Jersey	81.6	80.9	61.8	11.8	48.3
North Dakota	61.6	30.8	49.2	23.9	45.7
Ohio	67.7	68.8	51.7	11.4	51.8
Oklahoma	48.6	47.0	51.6	29.1	30.7
Tennessee	65.0	56.7	54.0	23.4	48.9
Utah	60.2	46.2	55.9	8.4	63.2
Virginia	83.5	74.0	50.3	15.8	50.0
West Virginia	73.6	74.7	58.2	25.3	54.2
<b>Unweighted Data</b>					
California	48.1	53.3	48.1	11.9	46.2
Georgia	82.8	43.7	50.5	25.0	54.0
Indiana	74.5	79.4	54.0	14.4	60.4
Kentucky	76.8	68.6	54.8	18.1	60.6
Louisiana*	70.9	60.8	46.9	20.7	34.2
Maryland*	68.2	67.8	53.8	9.7	50.3
Pennsylvania*	87.9	78.0	57.1	17.9	59.1
South Carolina	67.9	68.8	42.3	24.8	44.6
Texas*	60.5	62.1	42.5	11.4	41.6
<b>State Median</b>	<b>67.9</b>	<b>67.8</b>	<b>52.9</b>	<b>17.3</b>	<b>50.3</b>

(continued)



TABLE 8. Percentage of Schools in Which Health Education Teachers Planned or Coordinated Health-Related Projects or Activities With Other Groups, Selected U.S. Sites—School Health Education Profiles, Teachers' Survey, 2000 (*continued*)

### LOCAL SURVEYS

Site	Physical education staff	School health services staff	School mental health staff	Food service staff	Community members
<b>Weighted Data</b>					
Chicago	62.1	66.4	59.3	29.9	38.2
Dallas	39.4	67.6	38.1	14.1	43.3
District of Columbia	100.0	95.0	71.3	36.9	73.8
Ft. Lauderdale	63.2	37.6	51.9	30.9	44.8
Los Angeles	35.8	63.5	49.5	11.5	53.2
Miami	52.9	36.6	64.4	16.8	45.7
New Orleans	95.8	91.7	62.5	56.5	66.7
Orange County	44.7	48.7	38.5	10.5	48.7
Palm Beach	45.7	74.9	56.9	14.2	49.7
Philadelphia	88.0	85.1	60.2	24.7	45.9
San Diego	49.1	86.7	64.5	11.2	71.1
San Francisco	67.9	82.1	81.5	11.1	74.1
<b>Unweighted Data</b>					
Houston	97.1	88.6	64.7	17.1	73.5
<b>Local Median</b>	<b>62.1</b>	<b>74.9</b>	<b>60.2</b>	<b>16.8</b>	<b>49.7</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 9. Percentage of Schools in Which the Lead Health Education Teacher Had Professional Preparation in a Specific Area, Selected U.S. Sites—School Health Education Profiles, Teachers’ Surveys, 2000

**STATE SURVEYS**

Site	Health and physical education	Health education only	Physical education only	Science or other education degree	Nursing or counseling	Other*
<b>Weighted Data**</b>						
Alabama	53.1	5.6	17.4	18.0	3.8	2.2
Alaska	12.4	1.9	8.1	52.5	11.7	13.4
Arkansas	74.2	2.2	15.5	4.5	0.8	2.8
Delaware	70.5	4.3	18.0	5.0	2.2	0.0
Hawaii	51.6	7.8	19.4	13.9	3.6	3.6
Idaho	57.5	4.1	17.6	16.6	1.1	3.0
Illinois <sup>§</sup>	41.8	12.6	21.7	18.7	2.3	2.8
Iowa	36.1	7.3	15.3	29.0	5.6	6.8
Maine	40.7	19.5	10.7	18.3	8.3	2.5
Massachusetts	46.6	22.7	5.0	10.1	10.6	5.0
Michigan	31.9	12.3	16.1	30.4	3.4	6.0
Minnesota	67.4	9.1	8.5	9.1	2.3	3.4
Missouri	50.4	3.1	17.9	22.2	3.1	3.3
Montana	58.2	2.7	14.9	18.6	0.8	4.7
Nebraska	37.7	3.2	20.1	28.8	4.9	5.3
New Hampshire	23.3	18.3	15.6	20.9	18.4	3.5
New Jersey	64.3	8.8	3.7	6.1	13.9	3.2
North Dakota	36.7	3.2	17.8	36.4	2.5	3.3
Ohio	67.9	10.5	5.4	11.6	1.0	3.6
Oklahoma	35.0	2.1	7.9	27.5	17.3	10.2
Tennessee	54.4	4.5	9.0	19.1	9.1	4.0
Utah	43.6	14.3	15.4	19.5	1.5	5.7
Virginia	82.3	1.2	7.8	4.3	1.4	3.0
West Virginia	70.6	6.0	10.3	9.9	0.0	3.1
<b>Unweighted Data**</b>						
California	20.2	9.2	10.3	42.9	6.4	11.0
Georgia	68.3	7.9	5.0	11.9	2.5	4.5
Indiana	72.6	6.6	12.8	5.8	0.0	2.2
Kentucky	56.1	13.2	13.2	9.3	3.9	4.4
Louisiana <sup>§</sup>	74.1	1.7	10.3	9.2	1.7	2.9
Maryland <sup>§</sup>	48.9	20.7	16.3	11.1	3.0	0.0
Pennsylvania <sup>§</sup>	90.2	4.1	3.4	1.5	0.4	0.4
South Carolina	43.8	3.8	27.6	16.2	4.8	3.8
Texas <sup>§</sup>	45.0	9.2	13.1	16.0	6.0	10.6
<b>State Median</b>	<b>51.6</b>	<b>6.6</b>	<b>13.2</b>	<b>16.2</b>	<b>3.1</b>	<b>3.5</b>

(continued)

TABLE 9. Percentage of Schools in Which the Lead Health Education Teacher Had Professional Preparation in a Specific Area, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)

## LOCAL SURVEYS

Site	Health and physical education	Health education only	Physical education only	Science or other education degree	Nursing or counseling	Other
<b>Weighted Data**</b>						
Chicago	37.3	3.0	12.9	28.9	9.9	8.0
Dallas	45.5	12.4	0.0	32.3	0.0	9.8
District of Columbia	84.0	5.3	5.3	0.0	0.0	5.4
Ft. Lauderdale	32.7	8.4	10.9	36.3	6.7	5.0
Los Angeles	14.3	27.1	4.7	45.5	0.0	8.4
Miami	20.2	16.4	0.0	49.5	3.7	10.2
New Orleans	95.0	0.0	5.0	0.0	0.0	0.0
Orange County	22.2	27.8	2.8	38.9	2.8	5.6
Palm Beach	49.8	20.7	5.9	20.7	0.0	2.9
Philadelphia	86.9	1.1	3.5	5.6	1.2	1.6
San Diego	0.0	20.0	0.0	29.0	37.7	13.4
San Francisco	15.4	7.7	0.0	65.4	0.0	11.5
<b>Unweighted Data**</b>						
Houston	87.1	3.2	9.7	0.0	0.0	0.0
<b>Local Median</b>	<b>37.3</b>	<b>8.4</b>	<b>4.7</b>	<b>29.0</b>	<b>0.0</b>	<b>5.6</b>

\* Includes kinesiology, exercise science, public health and other.

\*\* Percentages for each row might not add up to 100.0 because of rounding.

§ Survey did not include students from one of the state's large school districts.

TABLE 10. Percentage of Schools in Which the Lead Health Education Teacher Had Received  $\geq 4$  Hours of Staff Development During the Preceding 2 Years in Specific Health Education Topics, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

**STATE SURVEYS**

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV* prevention	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Alabama	56.3	29.8	50.2	51.8	29.7	36.3	27.6	37.7	56.5
Alaska	40.6	18.1	30.9	22.9	14.0	17.3	30.7	25.8	41.5
Arkansas	46.8	24.3	43.8	41.2	27.3	33.9	24.8	32.1	44.6
Delaware	47.5	26.9	35.3	47.7	28.8	28.8	17.3	15.4	49.0
Hawaii	41.5	27.9	48.4	26.3	24.0	31.0	14.1	30.9	41.1
Idaho	52.7	32.9	71.1	44.1	23.0	49.8	26.0	41.0	51.7
Illinois <sup>§</sup>	43.9	16.8	41.4	36.6	19.6	28.9	18.8	23.1	45.8
Iowa	43.8	27.9	40.7	37.1	19.0	27.2	16.8	22.9	38.6
Maine	43.5	29.6	55.3	36.7	28.2	40.5	30.1	33.1	53.4
Massachusetts	65.3	44.8	50.0	46.0	30.7	39.6	35.1	51.1	73.3
Michigan	47.9	37.1	51.4	38.9	28.8	46.6	21.7	41.8	52.9
Minnesota	48.4	30.6	54.3	43.1	37.4	44.8	26.1	41.5	47.9
Missouri	60.6	31.1	38.4	45.0	24.2	32.2	27.1	36.2	55.0
Montana	44.0	30.8	51.6	48.4	25.1	39.4	19.9	36.8	50.5
Nebraska	43.1	27.3	30.3	33.7	18.5	22.4	20.7	27.2	36.1
New Hampshire	56.2	41.2	44.7	49.5	30.8	45.9	31.3	38.6	54.5
New Jersey	54.1	23.5	48.5	42.3	33.9	40.8	21.1	33.8	54.6
North Dakota	51.3	29.9	43.2	41.3	17.7	30.5	26.8	35.5	48.0
Ohio	36.5	20.3	NA <sup>§§</sup>	27.8	NA	NA	16.7	20.8	35.3
Oklahoma	59.8	22.3	74.3	24.9	28.3	49.6	27.8	33.0	57.4
Tennessee	51.9	28.1	57.3	45.1	29.9	40.4	23.0	37.6	63.8
Utah	79.6	70.8	88.0	48.3	63.4	80.7	73.0	78.5	73.1
Virginia	45.3	29.4	45.2	61.9	21.5	31.3	21.2	32.8	43.9
West Virginia	55.0	34.8	40.2	44.5	21.0	29.6	19.8	58.6	58.2
<b>Unweighted Data</b>									
California	50.9	27.0	53.6	37.7	31.3	43.6	23.9	52.1	52.1
Georgia	44.2	25.2	56.7	46.7	31.8	50.2	18.3	31.6	44.1
Indiana	39.4	18.8	49.0	34.3	20.4	32.4	13.6	25.9	42.7
Kentucky	48.6	22.8	40.7	42.1	23.3	29.5	18.8	34.2	52.9
Louisiana <sup>§</sup>	56.7	25.4	33.2	47.7	18.6	28.8	23.6	41.5	53.0
Maryland <sup>§</sup>	69.8	33.6	74.2	50.3	41.3	60.3	31.8	56.0	46.6
Pennsylvania <sup>§</sup>	52.1	23.5	48.0	47.1	24.1	36.2	28.1	32.1	52.3
South Carolina	36.8	22.3	52.9	40.3	31.4	41.6	14.1	30.8	32.7
Texas <sup>§</sup>	37.1	25.5	30.6	46.9	16.7	24.0	19.9	24.8	48.1
<b>State Median</b>	<b>48.4</b>	<b>27.9</b>	<b>48.4</b>	<b>43.1</b>	<b>26.2</b>	<b>36.3</b>	<b>23.0</b>	<b>33.8</b>	<b>50.5</b>

(continued)

TABLE 10. Percentage of Schools in Which the Lead Health Education Teacher Had Received  $\geq 4$  Hours of Staff Development During the Preceding 2 Years in Specific Health Education Topics, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV* prevention	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Chicago	49.5	33.3	54.2	40.4	34.4	48.3	23.5	34.7	58.2
Dallas	48.4	31.0	55.1	36.0	45.0	64.8	52.5	43.8	48.2
District of Columbia	60.5	51.6	84.7	70.9	63.4	74.0	18.6	47.4	65.8
Ft. Lauderdale	56.5	24.0	80.6	38.7	33.5	64.7	31.9	48.2	48.2
Los Angeles	54.4	27.6	72.9	20.8	42.9	61.6	28.0	70.8	46.9
Miami	35.8	19.9	70.4	28.0	42.8	57.6	26.8	28.9	33.5
New Orleans	83.3	66.7	87.5	82.6	69.6	87.5	52.2	66.7	79.2
Orange County	53.8	28.2	92.3	21.1	42.1	79.5	38.5	30.8	61.5
Palm Beach	65.4	42.6	68.4	31.2	45.3	51.1	48.5	42.5	65.4
Philadelphia	44.9	29.5	58.7	51.0	38.7	52.0	13.1	28.4	46.4
San Diego	100.0	11.0	100.0	13.3	97.7	97.7	75.7	100.0	93.4
San Francisco	78.6	42.9	74.1	34.6	32.1	50.0	32.1	76.9	81.5
<b>Unweighted Data</b>									
Houston	85.7	50.0	88.6	91.4	66.7	85.7	55.9	79.4	80.0
<b>Local Median</b>	<b>56.5</b>	<b>31.0</b>	<b>74.1</b>	<b>36.0</b>	<b>42.9</b>	<b>64.7</b>	<b>32.1</b>	<b>47.4</b>	<b>61.5</b>

\* HIV = human immunodeficiency virus.

\*\* STD = sexually transmitted disease.

§ Survey did not include students from one of the state's large school districts.

§§ NA = not available.

TABLE 11. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development in Specific Health Education Topics, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

**STATE SURVEYS**

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV* prevention	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Alabama	72.4	65.8	73.3	59.7	64.4	70.3	70.7	66.7	77.3
Alaska	70.0	61.4	60.9	52.5	55.5	58.9	72.0	63.4	76.7
Arkansas	73.2	57.7	70.5	59.7	57.4	65.2	71.9	68.4	77.7
Delaware	63.4	62.9	62.1	68.6	47.3	63.3	69.0	52.0	77.3
Hawaii	85.9	79.3	85.0	75.1	79.3	84.0	84.0	87.3	91.5
Idaho	72.0	65.9	67.3	54.5	57.8	62.0	78.6	65.9	79.9
Illinois <sup>§</sup>	72.5	59.7	64.3	52.1	55.0	65.1	70.4	60.8	79.0
Iowa	57.8	46.5	56.3	49.2	51.0	57.6	60.1	50.0	68.0
Maine	60.4	55.4	50.1	46.2	44.2	50.0	66.2	50.4	77.9
Massachusetts	76.3	69.6	68.9	59.2	65.7	74.3	82.6	65.5	83.5
Michigan	64.1	54.2	57.6	52.6	50.7	61.0	65.6	57.9	72.0
Minnesota	67.7	57.9	67.9	53.4	60.9	68.7	71.5	61.8	64.3
Missouri	66.5	62.7	65.0	56.5	55.2	61.9	67.7	60.0	77.4
Montana	68.4	67.1	65.3	62.7	59.2	64.5	73.5	66.4	78.0
Nebraska	65.0	49.6	55.1	45.9	43.9	52.5	61.3	56.0	72.1
New Hampshire	81.1	75.4	75.5	64.1	66.2	76.9	83.7	71.7	85.4
New Jersey	79.6	71.1	79.5	68.4	70.7	80.3	84.9	77.0	88.2
North Dakota	54.2	53.3	52.1	53.9	45.8	49.7	65.1	51.5	71.4
Ohio	65.8	51.0	NA <sup>§§</sup>	49.3	NA	NA	65.6	60.1	74.6
Oklahoma	71.8	52.2	68.6	54.2	63.2	65.7	73.7	67.5	81.0
Tennessee	72.1	62.3	66.3	58.1	57.7	62.8	74.9	60.6	81.9
Utah	73.7	69.5	69.7	58.5	65.7	67.8	84.7	71.4	86.5
Virginia	63.1	65.9	58.3	66.4	49.1	53.4	67.8	58.4	75.1
West Virginia	67.7	69.7	70.5	63.4	64.5	69.5	76.1	68.1	78.4
<b>Unweighted Data</b>									
California	66.1	57.5	61.1	49.7	54.8	60.3	68.8	54.5	76.1
Georgia	78.0	63.7	80.4	64.5	70.8	77.1	74.3	72.4	86.0
Indiana	63.7	44.5	58.7	52.3	49.6	57.0	66.9	52.7	72.0
Kentucky	69.7	61.6	70.6	58.3	59.7	67.4	73.5	61.9	82.1
Louisiana <sup>§</sup>	75.0	68.2	72.2	68.7	55.2	66.7	76.0	69.4	81.8
Maryland <sup>§</sup>	72.9	59.7	71.5	46.2	59.7	72.9	71.7	63.2	80.3
Pennsylvania <sup>§</sup>	74.7	65.6	80.8	67.8	66.1	76.1	73.7	70.3	83.3
South Carolina	71.1	68.8	72.4	64.1	65.7	69.5	74.2	65.1	77.5
Texas <sup>§</sup>	73.6	65.9	70.4	58.3	59.3	67.0	72.7	64.5	76.8
<b>State Median</b>	<b>71.1</b>	<b>62.7</b>	<b>68.2</b>	<b>58.3</b>	<b>58.5</b>	<b>65.4</b>	<b>72.0</b>	<b>63.4</b>	<b>77.9</b>

(continued)

TABLE 11. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development in Specific Health Education Topics, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)

## LOCAL SURVEYS

Site	Alcohol or other drug-use prevention	Dietary behavior and nutrition	HIV* prevention	Physical activity and fitness	Pregnancy prevention	STD** prevention	Suicide prevention	Tobacco-use prevention	Violence prevention
<b>Weighted Data</b>									
Chicago	76.3	72.7	67.7	69.6	67.4	67.5	75.3	71.0	86.5
Dallas	79.7	46.7	69.9	41.5	58.2	75.2	56.7	54.5	61.3
District of Columbia	71.8	74.7	82.3	57.8	72.0	80.0	82.3	64.2	84.3
Ft. Lauderdale	62.3	70.8	59.3	48.0	64.0	62.6	72.2	47.7	77.7
Los Angeles	67.5	57.5	63.5	45.1	53.1	61.3	68.7	59.7	77.0
Miami	77.9	76.9	70.3	57.0	67.1	72.6	73.5	72.1	79.8
New Orleans	91.7	79.2	83.3	75.0	87.5	95.8	79.2	79.2	83.3
Orange County	81.6	66.7	74.4	47.4	74.4	84.6	71.1	66.7	79.5
Palm Beach	67.4	67.6	69.5	45.4	58.6	73.3	69.4	57.1	75.4
Philadelphia	73.6	71.8	77.3	68.7	80.7	78.2	85.1	70.2	88.3
San Diego	75.5	37.8	84.4	26.6	64.3	66.6	68.8	55.4	84.4
San Francisco	60.7	66.7	60.7	63.0	57.7	65.4	81.5	53.8	81.5
<b>Unweighted Data</b>									
Houston	94.3	71.4	85.7	82.9	79.4	82.9	91.4	91.4	97.1
<b>Local Median</b>	<b>75.5</b>	<b>70.8</b>	<b>70.3</b>	<b>57.0</b>	<b>67.1</b>	<b>73.3</b>	<b>73.5</b>	<b>64.2</b>	<b>81.5</b>

\* HIV = human immunodeficiency virus.

\*\* STD = sexually transmitted disease.

§ Survey did not include students from one of the state's large school districts.

§§ NA = not available.

TABLE 12. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development in Specific Teaching Methods, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000

**STATE SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Teaching skills for behavior change
<b>Weighted Data</b>					
Alabama	45.7	33.1	17.6	56.1	44.3
Alaska	45.9	61.2	30.3	53.1	42.0
Arkansas	41.0	39.0	21.4	48.7	40.5
Delaware	39.0	36.5	10.7	48.4	34.5
Hawaii	26.8	31.5	33.1	48.1	22.6
Idaho	33.7	21.2	9.8	41.6	38.1
Illinois*	35.1	23.3	14.5	49.9	39.0
Iowa	43.3	28.0	13.6	46.8	43.6
Maine	38.8	12.4	8.0	46.3	32.3
Massachusetts	38.2	32.7	15.5	61.4	49.9
Michigan	30.3	29.4	9.5	56.2	47.7
Minnesota	33.7	38.1	18.7	44.3	38.3
Missouri	46.7	41.9	16.5	61.8	50.3
Montana	32.8	24.1	4.0	48.8	43.8
Nebraska	32.9	40.5	15.9	45.7	42.0
New Hampshire	53.4	18.2	6.1	67.4	60.1
New Jersey	41.8	36.1	12.5	51.1	42.4
North Dakota	30.6	19.1	2.4	40.2	45.7
Ohio	36.6	21.8	7.3	42.0	35.7
Oklahoma	51.1	66.2	20.1	50.0	48.2
Tennessee	47.4	35.7	18.9	63.8	53.7
Utah	30.1	44.1	34.3	67.1	52.4
Virginia	33.3	38.3	16.6	53.8	37.7
West Virginia	30.0	20.8	9.1	58.1	47.1
<b>Unweighted Data</b>					
California	46.9	65.2	59.2	58.4	45.5
Georgia	35.8	34.3	20.8	51.2	40.5
Indiana	40.6	26.5	12.0	48.8	41.4
Kentucky	53.2	42.3	12.3	64.4	51.6
Louisiana*	33.7	27.7	8.8	60.8	46.2
Maryland*	50.3	55.7	21.6	65.8	50.0
Pennsylvania*	51.2	27.7	12.5	56.2	42.7
South Carolina	31.4	34.2	11.5	53.2	37.9
Texas*	57.1	56.7	35.8	59.3	52.4
<b>State Median</b>	<b>38.8</b>	<b>34.2</b>	<b>14.5</b>	<b>53.1</b>	<b>43.6</b>

(continued)



TABLE 12. Percentage of Schools in Which the Lead Health Education Teacher Received Staff Development in Specific Teaching Methods, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Teaching skills for behavior change
<b>Weighted Data</b>					
Chicago	57.8	44.4	32.2	60.9	57.3
Dallas	39.0	73.1	50.7	85.1	43.3
District of Columbia	46.3	66.5	38.7	54.0	66.8
Ft. Lauderdale	48.6	72.8	75.3	72.1	51.1
Los Angeles	58.6	75.8	85.6	81.7	58.7
Miami	34.3	56.2	57.5	61.8	34.1
New Orleans	37.5	41.7	25.0	66.7	70.8
Orange County	54.1	68.4	77.8	64.9	48.6
Palm Beach	43.0	79.8	73.4	76.4	47.1
Philadelphia	23.5	41.3	16.2	56.9	42.8
San Diego	11.1	64.4	20.1	60.0	46.7
San Francisco	46.2	65.4	38.5	73.1	57.7
<b>Unweighted Data</b>					
Houston	70.6	77.1	77.1	76.5	80.0
<b>Local Median</b>	<b>46.2</b>	<b>66.5</b>	<b>50.7</b>	<b>66.7</b>	<b>51.1</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 13. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development in Specific Teaching Methods, Selected U.S. Sites—School Health Education Profiles, Teachers’ Surveys, 2000

**STATE SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Teaching skills for behavior change
<b>Weighted Data</b>					
Alabama	64.5	55.2	48.3	66.3	77.3
Alaska	64.7	55.2	54.7	61.6	79.8
Arkansas	56.6	49.2	45.0	63.6	75.1
Delaware	67.3	56.4	61.6	69.2	79.5
Hawaii	84.2	70.0	77.6	83.1	87.8
Idaho	59.2	54.0	51.3	65.8	79.3
Illinois*	55.0	44.7	39.5	55.7	74.4
Iowa	48.3	38.4	31.9	54.9	70.1
Maine	51.3	45.3	32.2	58.2	74.1
Massachusetts	69.2	66.1	54.8	70.5	84.5
Michigan	60.1	46.9	35.4	65.5	79.6
Minnesota	54.0	52.0	44.5	59.8	71.1
Missouri	57.9	43.9	37.7	60.4	77.6
Montana	62.1	43.9	35.6	64.8	75.1
Nebraska	48.5	43.9	34.9	44.5	65.8
New Hampshire	71.4	58.1	43.2	79.6	88.7
New Jersey	76.7	65.6	64.0	72.5	86.1
North Dakota	47.7	33.2	19.7	49.4	69.6
Ohio	58.0	40.6	31.2	59.7	71.9
Oklahoma	61.7	57.9	49.4	57.5	79.0
Tennessee	63.5	48.2	53.0	57.3	76.5
Utah	69.6	67.5	63.3	69.6	80.7
Virginia	65.8	49.8	49.2	53.6	75.2
West Virginia	58.6	47.1	37.9	61.0	72.9
<b>Unweighted Data</b>					
California	58.0	53.8	45.6	56.4	73.5
Georgia	65.7	64.3	64.6	70.0	77.1
Indiana	51.0	39.1	36.3	59.3	69.8
Kentucky	58.7	52.8	41.7	60.2	72.1
Louisiana*	64.6	60.8	45.1	68.0	76.8
Maryland*	58.9	51.3	41.5	60.3	80.1
Pennsylvania*	68.2	51.1	43.2	67.4	81.4
South Carolina	69.9	67.6	58.3	62.5	81.0
Texas*	61.5	59.3	57.9	61.0	74.4
<b>State Median</b>	<b>61.5</b>	<b>52.0</b>	<b>45.0</b>	<b>61.0</b>	<b>76.8</b>

(continued)

TABLE 13. Percentage of Schools in Which the Lead Health Education Teacher Wanted Staff Development in Specific Teaching Methods, Selected U.S. Sites—School Health Education Profiles, Teachers' Surveys, 2000 (*continued*)**LOCAL SURVEYS**

Site	Teaching students with physical or cognitive disabilities	Teaching students of various cultural backgrounds	Teaching students with limited English proficiency	Using interactive teaching methods	Teaching skills for behavior change
<b>Weighted Data</b>					
Chicago	79.9	71.0	62.7	73.2	87.2
Dallas	73.1	70.5	70.8	73.1	77.9
District of Columbia	82.0	76.8	79.5	79.5	79.5
Ft. Lauderdale	67.9	57.2	59.0	65.5	78.7
Los Angeles	56.2	51.9	48.6	62.2	77.4
Miami	66.2	63.6	52.4	61.7	77.2
New Orleans	79.2	82.6	73.9	95.8	87.5
Orange County	63.2	54.1	38.9	68.4	81.6
Palm Beach	54.0	56.9	46.9	61.5	76.8
Philadelphia	83.4	77.3	70.2	76.2	86.9
San Diego	66.7	68.8	62.2	51.0	62.2
San Francisco	77.8	77.8	59.3	63.0	77.8
<b>Unweighted Data</b>					
Houston	88.6	77.1	77.1	77.1	88.6
<b>Local Median</b>	<b>73.1</b>	<b>70.5</b>	<b>62.2</b>	<b>68.4</b>	<b>78.7</b>

---

\* Survey did not include students from one of the state's large school districts.

TABLE 14. Percentage of Schools That Received Parental Feedback About Health Education in Their School and, Among Those Schools, Percentage That Received a Specific Type of Feedback, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000

**STATE SURVEYS**

Site	Received parental feedback	Type of parental feedback received*		
		Mainly positive	Mainly negative	Equally positive and negative
<b>Weighted Data</b>				
Alabama	49.8	90.4	0.7	8.9
Alaska	39.0	79.3	4.0	16.7
Arkansas	47.2	90.4	0.0	9.6
California	54.4	88.7	1.0	10.3
Delaware	45.2	84.4	0.0	15.6
Hawaii	46.0	79.9	0.0	20.1
Idaho	61.1	85.6	0.7	13.6
Illinois**	51.9	90.4	1.5	8.1
Iowa	47.5	84.1	1.3	14.6
Maine	52.2	90.9	0.9	8.2
Massachusetts	65.9	92.3	1.4	6.3
Michigan	58.2	88.7	1.0	10.3
Minnesota	60.2	89.4	0.6	10.0
Missouri	53.2	86.8	0.6	12.6
Montana	54.3	88.2	1.4	10.4
Nebraska	40.7	91.0	0.7	8.3
New Hampshire	57.4	83.1	1.0	15.9
New Jersey	62.5	86.1	1.1	12.8
North Dakota	43.7	87.2	0.0	12.8
Ohio	47.3	86.3	1.8	11.9
Oklahoma	37.2	86.9	0.8	12.3
Pennsylvania**	59.7	91.9	1.2	6.9
Tennessee	53.5	87.1	0.7	12.2
Utah	65.6	90.6	1.6	7.9
Virginia	58.7	87.2	1.4	11.5
West Virginia	59.9	90.3	0.7	9.0
<b>Unweighted Data</b>				
Georgia	63.9	91.2	1.4	7.5
Indiana	53.8	88.7	2.1	9.2
Kentucky	49.1	86.8	0.0	13.2
Louisiana**	41.4	93.1	0.0	6.9
Maryland**	52.8	93.0	2.3	4.7
South Carolina	47.6	89.9	2.8	7.3
South Dakota	30.4	88.9	0.0	11.1
Texas**	44.4	92.7	0.7	6.6
<b>State Median</b>	<b>52.5</b>	<b>88.7</b>	<b>1.0</b>	<b>10.3</b>

(continued)

TABLE 14. Percentage of Schools That Received Parental Feedback About Health Education in Their School and, Among Those Schools, Percentage That Received a Specific Type of Feedback, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000 (*continued*)

### LOCAL SURVEYS

Site	Received parental feedback	Type of parental feedback received*		
		Mainly positive	Mainly negative	Equally positive and negative
<b>Weighted Data</b>				
Chicago	50.6	87.8	0.8	11.4
Dallas	49.5	95.5	0.0	4.5
District of Columbia	61.5	58.2	0.0	41.8
Ft. Lauderdale	56.9	92.6	3.7	3.7
Los Angeles	64.0	92.4	0.0	7.6
Miami	53.0	89.2	0.0	10.8
New Orleans	66.7	93.8	0.0	6.3
Orange County	59.5	86.4	0.0	13.6
Palm Beach	53.6	100.0	0.0	0.0
Philadelphia	44.5	86.2	0.0	13.8
San Diego	69.9	93.3	3.2	3.4
San Francisco	60.6	90.0	0.0	10.0
<b>Unweighted Data</b>				
Houston	57.1	90.0	0.0	10.0
<b>Local Median</b>	<b>57.1</b>	<b>90.0</b>	<b>0.0</b>	<b>10.0</b>

\* Percentages for each row might not add up to 100.0 because of rounding.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 15. Percentage of Schools With a Written Policy on HIV\*-Infected Students or School Staff and, Among Those Schools, Percentage That Addressed Specific Topics, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000

STATE SURVEYS

Site	Topic addressed by a written policy						
	Had a written policy	Attendance of students**	Protection from discrimination <sup>§</sup>	Confidentiality <sup>§</sup>	Worksite safety	Confidential counseling <sup>§</sup>	Communication of policy to students, staff, and parents
<b>Weighted Data</b>							
Alabama	57.6	97.6	97.5	98.8	99.3	83.1	91.4
Alaska	43.0	93.8	96.5	98.2	95.8	75.8	88.8
Arkansas	44.1	96.3	96.1	96.0	94.5	81.9	86.4
California	46.1	96.3	95.1	97.5	97.4	74.3	86.6
Delaware	50.7	100.0	100.0	100.0	96.2	84.3	92.3
Hawaii	60.1	89.7	100.0	100.0	100.0	69.0	90.0
Idaho	64.8	95.4	95.4	96.2	94.6	67.9	74.3
Illinois <sup>§§</sup>	50.9	93.7	97.3	98.4	95.8	67.4	80.2
Iowa	48.6	94.9	94.2	98.1	98.1	63.8	82.1
Maine	70.4	96.5	97.9	99.3	97.3	74.6	83.5
Massachusetts	61.9	94.6	97.9	99.4	98.5	74.5	86.1
Michigan	50.9	96.4	97.6	97.6	97.0	78.2	84.6
Minnesota	52.3	91.6	93.9	98.0	98.7	73.6	86.3
Missouri	54.8	94.8	97.8	98.9	96.0	76.3	87.0
Montana	62.1	94.0	94.4	97.3	97.8	71.7	89.0
Nebraska	58.4	91.3	93.6	92.4	92.0	63.3	81.4
New Hampshire	75.4	96.4	95.6	98.0	98.2	71.4	88.0
New Jersey	60.7	97.8	98.4	99.5	97.3	75.9	85.6
North Dakota	56.5	91.4	94.5	98.9	97.9	73.2	86.0
Ohio	NA <sup>+</sup>	NA	NA	NA	NA	NA	NA
Oklahoma	61.5	94.0	97.0	96.5	98.0	76.3	86.3
Pennsylvania <sup>§§</sup>	62.2	95.7	98.5	98.4	96.0	69.0	79.8
Tennessee	62.8	96.1	98.7	98.7	99.2	84.6	88.9
Utah	61.1	96.1	99.0	100.0	99.1	75.8	81.0
Virginia	48.7	98.5	97.0	99.2	96.9	76.7	84.8
West Virginia	41.4	96.9	99.0	99.0	98.0	83.6	92.0
<b>Unweighted Data</b>							
Georgia	49.3	94.4	99.1	99.1	96.3	81.1	86.9
Indiana	54.5	89.7	95.6	98.5	100.0	68.7	85.2
Kentucky	26.7	84.3	89.9	95.8	97.1	80.9	76.8
Louisiana <sup>§§</sup>	34.6	94.0	95.4	98.5	100.0	82.5	85.5
Maryland <sup>§§</sup>	56.8	94.4	96.6	100.0	98.9	77.3	84.3
South Carolina	56.0	95.8	97.5	99.2	96.7	84.0	86.3
South Dakota	54.4	92.3	94.7	96.2	96.1	64.5	84.4
Texas <sup>§§</sup>	40.8	93.9	96.2	99.2	90.9	83.3	86.4
<b>State Median</b>	<b>54.8</b>	<b>94.8</b>	<b>97.0</b>	<b>98.5</b>	<b>97.3</b>	<b>75.8</b>	<b>86.1</b>

(continued)

TABLE 15. Percentage of Schools With a Written Policy on HIV\*-Infected Students or School Staff and, Among Those Schools, Percentage That Addressed Specific Topics, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000 (continued)

### LOCAL SURVEYS

Site	Topic addressed by a written policy						
	Had a written policy	Attendance of students**	Protection from discrimination <sup>§</sup>	Confidentiality <sup>§</sup>	Worksite safety	Confidential counseling <sup>§</sup>	Communication of policy to students, staff, and parents
<b>Weighted Data</b>							
Chicago	58.5	95.9	97.5	98.4	98.5	83.2	86.1
Dallas	41.5	78.3	100.0	100.0	100.0	94.1	78.3
District of Columbia	82.7	91.0	100.0	97.1	96.8	87.4	96.8
Ft. Lauderdale	73.4	85.5	97.0	97.1	97.1	78.2	87.7
Los Angeles	67.5	96.8	100.0	100.0	98.5	81.8	86.7
Miami	73.5	98.5	98.3	100.0	100.0	89.0	86.9
New Orleans	75.0	100.0	100.0	100.0	100.0	94.4	100.0
Orange County	37.8	92.9	100.0	100.0	100.0	85.7	84.6
Palm Beach	57.9	100.0	100.0	100.0	100.0	91.4	84.1
Philadelphia	55.4	98.1	98.1	100.0	96.1	87.1	89.9
San Diego	100.0	95.3	100.0	100.0	100.0	0.0	100.0
San Francisco	45.5	92.9	100.0	100.0	100.0	100.0	80.0
<b>Unweighted Data</b>							
Houston	75.8	83.3	91.7	95.8	87.5	91.3	83.3
<b>Local Median</b>	<b>67.5</b>	<b>95.3</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>87.4</b>	<b>86.7</b>

\* HIV = human immunodeficiency virus.

\*\* For HIV-infected students.

§ For HIV-infected students and staff members.

§§ Survey did not include students from one of the state's large school districts.

+ NA = not available.

TABLE 16. Percentage of Schools With a Policy Prohibiting Cigarette Smoking by Students and, Among Those Schools, Percentage That Had a Policy Prohibiting Cigarette Smoking in Specific Locations, Selected U.S. Sites—  
School Health Education Profiles, Principals' Survey, 2000

**STATE SURVEYS**

Site	Adopted a policy	Location			
		In school buildings	On school grounds	In school buses	At off-campus events
<b>Weighted Data</b>					
Alabama	100.0	99.3	99.3	99.3	94.5
Alaska	98.1	100.0	99.3	95.4	96.1
Arkansas	99.7	99.7	99.7	100.0	99.1
California	98.8	100.0	100.0	99.1	98.0
Delaware	100.0	100.0	98.2	96.4	87.4
Hawaii	100.0	100.0	100.0	100.0	100.0
Idaho	99.6	100.0	100.0	100.0	99.0
Illinois*	98.7	99.7	99.7	98.6	97.1
Iowa	99.1	99.7	99.1	99.4	97.5
Maine	99.5	100.0	100.0	99.5	94.3
Massachusetts	99.3	100.0	99.4	99.3	94.4
Michigan	97.6	100.0	100.0	99.6	99.1
Minnesota	97.4	100.0	98.6	100.0	94.5
Missouri	100.0	99.7	99.4	99.7	96.0
Montana	99.1	100.0	99.2	100.0	97.6
Nebraska	96.1	99.1	98.8	97.6	96.6
New Hampshire	99.3	100.0	100.0	98.0	91.5
New Jersey	97.4	99.7	98.7	99.0	92.2
North Dakota	98.1	100.0	100.0	99.4	98.2
Ohio	99.7	100.0	100.0	100.0	99.7
Oklahoma	99.4	99.4	99.0	99.0	95.9
Pennsylvania*	99.3	100.0	100.0	99.7	96.2
Tennessee	99.6	99.6	99.6	99.6	97.1
Utah	100.0	100.0	99.4	100.0	97.8
Virginia	99.7	100.0	100.0	99.6	94.0
West Virginia	100.0	100.0	100.0	100.0	98.0
<b>Unweighted Data</b>					
Georgia	100.0	99.6	99.6	99.6	98.3
Indiana	99.2	100.0	100.0	100.0	98.4
Kentucky	96.8	100.0	98.5	100.0	84.7
Louisiana*	99.5	99.5	99.5	99.5	94.7
Maryland*	99.4	100.0	100.0	100.0	99.4
South Carolina	99.6	99.6	99.5	99.5	95.9
South Dakota	99.3	99.3	99.3	98.6	95.1
Texas*	99.1	100.0	100.0	100.0	99.7
<b>State Median</b>	<b>99.4</b>	<b>100.0</b>	<b>99.6</b>	<b>99.6</b>	<b>96.8</b>

(continued)



TABLE 16. Percentage of Schools With a Policy Prohibiting Cigarette Smoking by Students and, Among Those Schools, Percentage That Had a Policy Prohibiting Cigarette Smoking in Specific Locations, Selected U.S. Sites—  
School Health Education Profiles, Principals' Survey, 2000 (*continued*)

### LOCAL SURVEYS

Site	Adopted a policy	Location			
		In school buildings	On school grounds	In school buses	At off-campus events
<b>Weighted Data</b>					
Chicago	92.5	99.6	98.1	98.5	97.2
Dallas	97.5	100.0	100.0	100.0	100.0
District of Columbia	100.0	100.0	97.3	100.0	97.5
Ft. Lauderdale	98.5	100.0	100.0	100.0	97.9
Los Angeles	98.0	100.0	100.0	100.0	100.0
Miami	96.7	100.0	100.0	100.0	92.4
New Orleans	100.0	100.0	100.0	100.0	91.3
Orange County	97.3	100.0	100.0	100.0	100.0
Palm Beach	100.0	100.0	100.0	100.0	100.0
Philadelphia	95.8	98.8	96.8	97.8	91.6
San Diego	100.0	97.7	97.7	97.7	97.7
San Francisco	97.0	100.0	100.0	100.0	100.0
<b>Unweighted Data</b>					
Houston	100.0	100.0	100.0	100.0	100.0
<b>Local Median</b>	<b>98.0</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>97.9</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 17. Percentage of Schools That Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites—School Health Education Profiles, Principals' Survey, 2000

**STATE SURVEYS**

Site	Referred to school counselor	Referred to school administrator	Encouraged to participate in cessation program	Required to participate in cessation program	Placed in detention	Suspended from school	Informed parents or guardians
<b>Weighted Data</b>							
Alabama	43.4	98.2	34.1	14.0	54.8	74.1	98.6
Alaska	52.3	93.7	47.9	23.0	51.0	65.1	93.3
Arkansas	42.3	99.0	28.5	9.6	48.3	73.9	97.7
California	64.1	98.9	64.0	45.3	57.3	80.5	99.4
Delaware	66.9	96.3	52.4	20.4	52.6	75.8	100.0
Hawaii	52.5	100.0	66.6	19.7	59.7	86.1	100.0
Idaho	73.9	100.0	67.4	53.4	46.7	77.3	100.0
Illinois**	54.8	97.9	43.2	17.1	51.9	81.4	98.7
Iowa	64.7	97.7	54.4	19.6	39.5	65.3	98.3
Maine	81.1	100.0	73.1	41.8	34.7	85.2	99.0
Massachusetts	67.9	98.9	69.3	38.4	61.8	73.1	98.0
Michigan	60.0	97.5	59.6	31.1	37.3	85.0	98.0
Minnesota	65.4	97.5	66.9	37.1	43.2	73.8	97.2
Missouri	42.6	98.8	29.4	12.8	59.4	68.3	98.5
Montana	69.9	98.8	56.9	38.9	60.6	73.2	99.6
Nebraska	47.3	98.8	43.4	18.0	57.5	67.9	98.9
New Hampshire	73.3	98.7	64.9	33.6	33.0	87.6	95.5
New Jersey	76.4	97.6	58.4	30.0	62.7	76.2	95.4
North Dakota	65.5	98.8	42.3	18.1	58.5	64.0	100.0
Ohio	52.6	99.2	61.2	31.2	34.2	85.4	99.3
Oklahoma	49.5	98.1	33.0	13.5	63.1	71.6	97.5
Pennsylvania**	64.3	99.7	61.6	29.6	44.5	68.4	99.0
Tennessee	47.1	99.1	39.6	25.2	39.4	72.1	99.5
Utah	52.9	97.6	75.5	56.5	36.4	70.1	98.8
Virginia	59.2	99.2	57.5	37.4	44.3	88.0	98.4
West Virginia	78.2	100.0	61.9	56.8	57.5	83.2	99.2
<b>Unweighted Data</b>							
Georgia	43.8	99.6	40.2	12.0	39.1	66.5	99.5
Indiana	51.8	97.7	55.3	25.8	39.6	90.5	98.0
Kentucky	52.5	98.5	51.9	25.7	66.3	68.8	98.1
Louisiana**	51.3	98.6	36.4	14.6	47.2	86.9	98.6
Maryland**	61.3	100.0	67.8	57.7	40.9	89.9	100.0
South Carolina	60.2	100.0	39.1	21.6	46.0	87.9	98.6
South Dakota	63.6	96.5	40.3	15.8	59.4	56.8	98.6
Texas**	54.9	99.1	35.8	23.9	61.6	43.1	98.1
<b>State Median</b>	<b>59.6</b>	<b>98.8</b>	<b>54.9</b>	<b>25.4</b>	<b>49.7</b>	<b>74.0</b>	<b>98.6</b>

(continued)

TABLE 17. Percentage of Schools That Took Specific Actions When Students Were Caught Smoking Cigarettes,\* Selected U.S. Sites—School Health Education Profiles, Principals' Survey, 2000 (continued)

### LOCAL SURVEYS

Site	Referred to school counselor	Referred to school administrator	Encouraged to participate in cessation program	Required to participate in cessation program	Placed in detention	Suspended from school	Informed parents or guardian
<b>Weighted Data</b>							
Chicago	59.7	88.2	33.5	17.2	69.8	72.7	93.7
Dallas	34.8	95.0	37.1	24.7	87.1	64.5	95.0
District of Columbia	51.1	85.0	21.3	8.6	46.8	53.6	74.3
Ft. Lauderdale	62.5	98.5	59.3	29.4	52.5	76.5	98.5
Los Angeles	82.5	86.5	64.3	86.8	63.4	81.5	100.0
Miami	85.2	96.3	66.9	36.6	60.3	60.0	97.7
New Orleans	52.2	100.0	56.5	39.1	91.3	75.0	100.0
Orange County	79.4	97.2	76.5	44.1	54.3	77.1	97.1
Palm Beach	78.4	100.0	69.4	72.0	29.5	100.0	100.0
Philadelphia	63.5	95.5	47.8	15.0	74.0	79.0	95.5
San Diego	85.2	97.6	65.1	90.5	48.6	73.1	100.0
San Francisco	100.0	92.9	69.0	53.3	56.7	50.0	100.0
<b>Unweighted Data</b>							
Houston	71.9	97.1	63.6	30.3	78.8	90.9	97.1
<b>Local Median</b>	<b>71.9</b>	<b>96.3</b>	<b>63.6</b>	<b>36.6</b>	<b>60.3</b>	<b>75.0</b>	<b>97.7</b>

\* Among schools with a policy prohibiting cigarette smoking by students.

\*\* Survey did not include students from one of the state's large school districts.

TABLE 18. Percentage of Schools That Prohibited Tobacco Advertising in Specific Places, Through Sponsorship of School Events, and on Student Apparel, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys 2000

**STATE SURVEYS**

Site	Prohibited advertising in school buildings, on school grounds, on school buses, and in school publications	Prohibited sponsorship of school events	Prohibited tobacco brand-name apparel
<b>Weighted Data</b>			
Alabama	92.5	88.1	95.3
Alaska	87.4	86.8	75.0
Arkansas	91.1	89.2	94.9
California	91.3	92.1	95.8
Delaware	92.6	85.0	63.9
Hawaii	96.9	96.9	84.7
Idaho	90.4	88.0	95.7
Illinois*	93.3	91.2	93.4
Iowa	91.6	91.5	94.9
Maine	93.5	91.1	88.4
Massachusetts	94.3	91.0	56.9
Michigan	90.1	88.5	87.7
Minnesota	94.0	90.5	87.2
Missouri	92.7	89.3	92.3
Montana	89.7	87.3	93.5
Nebraska	90.3	92.0	93.7
New Hampshire	93.3	84.8	84.8
New Jersey	93.6	89.2	59.8
North Dakota	89.2	89.0	92.7
Ohio	92.6	89.3	94.7
Oklahoma	89.7	89.9	96.9
Pennsylvania*	93.7	92.0	87.8
Tennessee	93.0	91.4	89.5
Utah	94.3	95.7	96.8
Virginia	91.5	91.3	86.4
West Virginia	94.5	93.0	89.5
<b>Unweighted Data</b>			
Georgia	91.6	90.2	92.0
Indiana	92.6	90.2	95.0
Kentucky	91.2	86.8	84.5
Louisiana*	92.0	86.2	95.3
Maryland*	93.4	91.5	89.7
South Carolina	92.5	89.1	86.0
South Dakota	87.5	91.0	93.8
Texas*	92.5	91.6	98.2
<b>State Median</b>	<b>92.5</b>	<b>90.2</b>	<b>92.1</b>

(continued)

TABLE 18. Percentage of Schools That Prohibited Tobacco Advertising in Specific Places, Through Sponsorship of School Events, and on Student Apparel, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys 2000 (*continued*)**LOCAL SURVEYS**

Site	Prohibited advertising in school buildings, on school grounds, on school buses, and in school publications	Prohibited sponsorship of school events	Prohibited tobacco brand-name apparel
<b>Weighted Data</b>			
Chicago	91.3	86.0	80.5
Dallas	90.4	86.0	85.8
District of Columbia	94.8	69.1	50.1
Ft. Lauderdale	97.0	96.4	82.9
Los Angeles	92.1	99.1	92.2
Miami	93.2	90.8	82.7
New Orleans	91.7	82.6	91.7
Orange County	81.1	83.8	86.5
Palm Beach	93.8	89.9	97.7
Philadelphia	87.3	90.7	39.8
San Diego	97.7	100.0	95.1
San Francisco	87.5	93.9	71.9
<b>Unweighted Data</b>			
Houston	97.1	94.1	97.1
<b>Local Median</b>	<b>92.1</b>	<b>90.7</b>	<b>85.8</b>

\* Survey did not include students from one of the state's large school districts.

TABLE 19. Percentage of Schools That Implemented Safety and Security Measures, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000

**STATE SURVEYS**

Site	Required visitors to report to main office	Maintained a closed campus	Used staff to monitor halls	Checked bags, desks, and lockers	Prohibited backpacks	Required school uniforms	Used metal detectors	Had uniformed police
<b>Weighted Data</b>								
Alabama	100.0	96.9	91.9	73.8	28.9	24.0	30.7	33.1
Alaska	84.6	33.7	67.9	26.5	9.5	1.6	1.4	6.5
Arkansas	100.0	93.6	91.6	63.2	10.0	2.8	23.9	32.5
California	99.7	84.9	88.2	36.4	2.6	16.8	9.8	45.9
Delaware	100.0	91.1	88.9	21.8	33.7	3.6	0.0	41.1
Hawaii	100.0	100.0	92.2	7.4	0.0	17.6	0.0	54.9
Idaho	97.1	53.2	88.6	48.5	18.0	0.0	0.5	43.8
Illinois*	99.0	79.4	79.6	46.1	36.5	4.0	5.9	24.0
Iowa	95.9	71.9	80.2	34.5	26.6	0.0	0.3	14.6
Maine	98.1	81.6	68.9	20.2	25.2	0.0	2.3	24.5
Massachusetts	98.1	89.7	83.3	24.6	26.1	5.1	2.3	30.2
Michigan	99.1	84.4	86.2	46.4	27.0	6.5	7.4	39.5
Minnesota	98.3	67.0	84.7	37.9	21.3	0.7	1.0	34.8
Missouri	100.0	95.9	87.1	61.7	23.1	1.5	7.4	35.0
Montana	98.4	38.0	83.9	46.8	14.4	0.7	1.3	12.0
Nebraska	95.4	65.1	84.4	41.5	17.3	0.8	0.9	12.3
New Hampshire	100.0	77.4	71.3	22.3	20.9	0.0	1.3	25.6
New Jersey	100.0	84.1	87.2	37.1	24.0	7.3	2.8	39.4
North Dakota	93.3	53.2	87.4	34.4	13.8	1.9	0.0	7.9
Ohio	99.6	90.0	81.3	50.5	47.8	3.1	6.9	26.2
Oklahoma	98.4	73.6	92.6	68.3	11.4	2.5	13.7	21.5
Pennsylvania*	100.0	95.6	86.3	42.2	46.0	3.3	11.4	29.5
Tennessee	100.0	97.0	91.6	71.8	18.4	5.9	24.4	44.8
Utah	98.9	55.3	86.2	44.0	10.1	6.6	1.7	51.4
Virginia	100.0	93.4	93.2	44.4	19.1	4.4	25.0	69.2
West Virginia	100.0	96.5	92.6	70.8	54.1	0.0	12.1	21.5
<b>Unweighted Data</b>								
Georgia	100.0	97.8	92.4	67.4	23.2	1.3	36.0	65.9
Indiana	99.6	93.8	82.1	47.2	52.7	0.8	9.4	28.1
Kentucky	100.0	94.6	93.5	62.8	39.1	10.8	20.0	42.4
Louisiana*	99.5	98.1	90.1	77.5	13.3	53.3	49.3	31.9
Maryland*	100.0	95.2	86.7	37.0	54.2	2.4	1.2	36.7
South Carolina	100.0	95.2	93.8	67.7	12.7	6.2	34.7	83.3
South Dakota	93.8	52.8	77.8	45.1	16.0	3.5	0.0	12.5
Texas*	99.4	81.2	92.2	64.0	13.9	10.5	12.0	52.2
<b>State Median</b>	<b>99.6</b>	<b>87.3</b>	<b>87.1</b>	<b>45.6</b>	<b>21.1</b>	<b>3.2</b>	<b>6.4</b>	<b>32.8</b>

(continued)

TABLE 19. Percentage of Schools That Implemented Safety and Security Measures, Selected U.S. Sites—School Health Education Profiles, Principals' Surveys, 2000 (*continued*)**LOCAL SURVEYS**

Site	Required visitors to report to main office	Maintained a closed campus	Used staff to monitor halls	Checked bags, desks, and lockers	Prohibited backpacks	Required school uniforms	Used metal detectors	Had uniformed police
<b>Weighted Data</b>								
Chicago	100.0	95.9	94.5	71.9	32.7	76.1	81.8	92.9
Dallas	97.8	100.0	100.0	92.9	12.0	4.9	92.7	92.6
District of Columbia	100.0	92.3	97.6	89.9	36.3	31.2	92.6	94.5
Ft. Lauderdale	100.0	87.3	92.2	19.5	2.1	20.2	10.2	100.0
Los Angeles	100.0	94.4	97.8	90.8	1.9	45.2	93.3	88.7
Miami	98.8	80.5	87.0	59.3	7.9	35.7	49.9	90.7
New Orleans	100.0	95.7	95.8	83.3	8.3	87.5	91.3	95.8
Orange County	100.0	100.0	100.0	37.8	5.4	0.0	8.1	94.6
Palm Beach	100.0	100.0	100.0	43.4	3.1	12.4	4.7	100.0
Philadelphia	100.0	97.9	94.7	46.8	12.1	2.1	35.7	81.4
San Diego	100.0	100.0	95.3	12.5	2.4	31.9	2.3	63.1
San Francisco	100.0	78.8	87.9	6.1	0.0	24.2	0.0	56.3
<b>Unweighted Data</b>								
Houston	100.0	91.2	100.0	64.7	45.5	79.4	23.5	91.2
<b>Local Median</b>	<b>100.0</b>	<b>95.7</b>	<b>95.8</b>	<b>59.3</b>	<b>7.9</b>	<b>31.2</b>	<b>35.7</b>	<b>92.6</b>

\* Survey did not include students from one of the state's large school districts.

