

## **APPENDIX C:**

### ***INTERNATIONAL CLASSIFICATION OF DISEASES, 9TH REVISION, CLINICAL MODIFICATION***

#### **ADDENDUM AND CONVERSION TABLE**

The *International Classification of Diseases, 9th Revision, Clinical Modification* has been used for coding medical data in the NHDS since 1979. However, since 1986, the classification system has undergone annual updating, which includes assignment of new diagnostic and procedure codes, fourth and/or fifth digit expansion of codes, and code deletions. These changes have been released for the years 1986 through 2007, except for 1999 when no Addendum was released because of concern about possible complications for instituting coding changes prior to the millennium crossover. These changes are published in an Addendum and become effective October 1 of the calendar year. Addenda are developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Centers for Medicare and Medicaid Services (formerly the Health Care Financing Administration).

In 2002, the ICD-9-CM Coordination and Maintenance Committee created procedure Chapter 00 – Procedures and Interventions, Not Elsewhere Classified – as a way of handling space limitations in the existing hierarchical structure and alleviating inappropriate categorization of new procedures. Since October addendum changes are not implemented in the NHDS until the following data collection year, 2003 was the first year these codes were used.

Users who wish to produce trend statistics must be aware of the dynamic nature of the ICD-9-CM coding system. Because of the Addendum changes, estimates for some codes may show discontinuity across years. Also, the meaning of the original and revised code may be different, thus causing problems with interpretation across years. Further, to address the potential problem that October coding changes would cause for calendar year estimation, all coding which was consistent with the October Addendum of the survey year was changed back to the coding previous to the Addendum changes. This “backward coding” was performed for the 1992 through 2007 data years, and helps to avoid the problem of partial year estimates being mistaken for full year estimates.

In order to assist users in data analysis, a conversion table is provided that shows for each new code, its date of introduction and the previously assigned code equivalent, which had been used for reporting the selected diagnosis or procedure prior to issuance of the new code.

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
005.81, 005.89	1995	005.8
007.4	1997	007.8
007.5	2000	007.8
008.00-008.09	1992	008.0
008.43-008.47	1992	008.49
008.61-008.69	1992	008.6
031.2	1997	031.8
038.10-038.11, 038.19	1997	038.1
040.82	2002	040.89
041.00-041.05, 041.09	1992	041.0
041.04 (Code title restated)	1997	041.04
041.10-041.19	1992	041.1
041.81-041.85, 041.89	1992	041.8
041.86	1995	041.84
042	1994	042.0-042.2, 042.9, 043.0-043.3, 043.9, 044.0, 044.9 (codes deleted)
042.0-044.9	1986	279.19
<b>052.2</b>	<b>2006</b>	<b>052.7</b>
<b>053.14</b>	<b>2006</b>	<b>053.19</b>
<b>054.74</b>	<b>2006</b>	<b>054.79</b>
066.4	2002	066.3
066.40-066.42, 066.49	2004	066.4
070.20-070.21	1991	070.2
070.22	1994	070.20
070.23	1994	070.21
070.30-070.31	1991	070.3
070.32	1994	070.30
070.33	1994	070.31
070.41-070.43	1991	070.4
070.44	1994	070.41
070.49	1991	070.4
070.51-070.53	1991	070.5
070.54	1994	070.51

**Changes in ICD-9-CM diagnosis codes, 1986-2006**

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
070.59	1991	070.5
070.70	2004	070.51
070.71	2004	070.41
077.98-077.99	1993	077.9
078.10-078.11, 078.19	1993	078.1
078.88	1993	078.89
079.4	1993	079.8
079.50-079.53, 079.59	1993	079.8
079.6	1996	079.89
079.81	1995	079.89
079.82	2003	079.89
079.88-079.89	1993	079.8
079.98-079.99	1993	079.9
082.40-082.41, 082.49	2000	082.8
088.81, 088.89	1989	088.8
088.82	1993	088.89
099.40-099.49	1992	099.4
099.50-099.59	1992	078.89
112.84-112.85	1992	112.89
114.4-114.5	1993	114.3
176.0-176.9	1991	173.0-173.9
203.00	1991	203.0
203.01	1991	V10.79
203.10	1991	203.1
203.11	1991	V10.79
203.80	1991	203.8
203.81	1991	V10.79
204.00	1991	204.0
204.01	1991	V10.61
204.10	1991	204.1
204.11	1991	V10.61
204.20	1991	204.2
204.21	1991	V10.61
204.80	1991	204.8
204.81	1991	V10.61

### Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
204.90	1991	204.9
204.91	1991	V10.61
205.00	1991	205.0
205.01	1991	V10.62
205.10	1991	205.1
205.11	1991	V10.62
205.20	1991	205.2
205.21	1991	V10.62
205.30	1991	205.3
205.31	1991	V10.62
205.80	1991	205.8
205.81	1991	V10.62
205.90	1991	205.9
205.91	1991	V10.62
206.00	1991	206.0
206.01	1991	V10.63
206.10	1991	206.1
206.11	1991	V10.63
206.20	1991	206.2
206.21	1991	V10.63
206.80	1991	206.8
206.81	1991	V10.63
206.90	1991	206.9
206.91	1991	V10.63
207.00	1991	207.0
207.01	1991	V10.69
207.10	1991	207.1
207.11	1991	V10.69
207.20	1991	207.2
207.21	1991	V10.69
207.80	1991	207.8
207.81	1991	V10.69
208.00	1991	208.0
208.01	1991	V10.60
208.10	1991	208.1
208.11	1991	V10.60
208.20	1991	208.2
208.21	1991	V10.60
208.80	1991	208.8
208.81	1991	V10.60
208.90	1991	208.9
208.91	1991	V10.60

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
237.70-237.72	1990	237.7
<b>238.71-238.76, 238.79</b>	<b>2006</b>	<b>238.7</b>
250.02	1993	250.90
250.03	1993	250.91
250.12	1993	250.10
250.13	1993	250.11
250.22	1993	250.20
250.23	1993	250.21
250.32	1993	250.30
250.33	1993	250.31
250.42	1993	250.40
250.43	1993	250.41
250.52	1993	250.50
250.53	1993	250.51
250.62	1993	250.60
250.63	1993	250.61
250.72	1993	250.70
250.73	1993	250.71
250.82	1993	250.80
250.83	1993	250.81
250.92	1993	250.90
250.93	1993	250.91
252.00-252.02, 252.08	2004	252.0
255.10-255.14	2003	255.1
256.31-256.39	2001	256.3
259.5	2005	257.8
273.4	2004	277.6
275.40-275.42, 275.49	1997	275.4
276.50-276.52	2005	276.5
277.02-277.03, 277.09	2002	277.00
<b>277.30-277.31, 277.39</b>	<b>2006</b>	<b>277.3</b>
277.7	2001	277.8
277.81-277.84, 277.89	2003	277.8

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
277.85-277.86	2004	277.89
277.87	2004	277.89, 758.89
278.00-278.01	1995	278.0
278.02	2005	278.00
282.41-282.42, 282.49	2003	282.4
282.64	2003	282.63
282.68	2003	282.69
283.10-283.11, 283.19	1993	283.1
<b>284.01, 284.09</b>	<b>2006</b>	<b>284.0</b>
<b>284.1</b>	<b>2006</b>	<b>284.8</b>
<b>284.2</b>	<b>2006</b>	<b>284.8; 285.8</b>
285.21-285.22, 285.29	2000	285.8
287.30-287.33, 287.39	2005	287.3
<b>288.00-288.04, 288.09</b>	<b>2006</b>	<b>288.0</b>
<b>288.4</b>	<b>2006</b>	<b>288.0</b>
<b>288.50-288.51</b>	<b>2006</b>	<b>288.0; 288.8</b>
<b>288.59</b>	<b>2006</b>	<b>288.0</b>
<b>288.60-288.65, 288.69</b>	<b>2006</b>	<b>288.8</b>
289.52	2003	289.59
<b>289.53</b>	<b>2006</b>	<b>288.0</b>
289.81-289.82	2003	289.8
<b>289.83</b>	<b>2006</b>	<b>289.89</b>
289.89	2003	289.8
291.81	1996	291.8
291.82	2005	291.89
291.89	1996	291.8
292.85	2005	292.89
293.84	1996	293.89
294.10-294.11	2000	294.1
300.82	1996	300.81
305.1	1994	305.10, 305.11, 305.12, 305.13 (Codes deleted)

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
312.81-312.82, 312.89	1994	312.8
315.32	1996	315.39
320.81-320.89	1992	320.8
<b>323.01-323.02</b>	<b>2006</b>	<b>320.0</b>
<b>323.41-323.42</b>	<b>2006</b>	<b>323.4</b>
<b>323.51-323.52</b>	<b>2006</b>	<b>323.5</b>
<b>323.61-323.63</b>	<b>2006</b>	<b>323.6</b>
<b>323.71-323.72</b>	<b>2006</b>	<b>323.7</b>
<b>323.81-323.82</b>	<b>2006</b>	<b>323.8</b>
327.00	2005	780.51; 780.52
327.01	2005	780.51; 780.52
327.02	2005	307.41
327.09	2005	780.51; 780.52
327.10-327.14	2005	780.53; 780.54
327.15	2005	307.43
327.19	2005	780.53; 780.54
327.20-327.27	2005	780.57
327.29	2005	780.51; 780.53; 780.57
327.30-327.37; 327.39	2005	307.45
327.40-327.44; 327.49	2005	780.59
327.51	2005	780.58
327.52	2005	729.82
327.53	2005	306.8
327.59	2005	780.58
327.8	2005	780.50
331.11, 331.19	2003	331.1
331.82	2003	331.89
<b>331.83</b>	<b>2006</b>	<b>310.1</b>
<b>333.71-333.72, 333.79</b>	<b>2006</b>	<b>333.7</b>
<b>333.85</b>	<b>2006</b>	<b>333.82</b>
333.92-333.93	1994	333.99
<b>333.94</b>	<b>2006</b>	<b>333.99</b>
337.20-337.22, 337.29	1993	337.9
337.3	1998	337.9
<b>338.0</b>	<b>2006</b>	<b>Code to Pain, by site or 348.8</b>
<b>338.11-338.12</b>	<b>2006</b>	<b>Code to Pain, by site</b>
<b>338.18-338.19</b>	<b>2006</b>	<b>Code to Pain, by site</b>
<b>338.21-338.22</b>	<b>2006</b>	<b>Code to Pain, by site</b>

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
<b>338.28-338.29</b>	<b>2006</b>	<b>Code to Pain, by site</b>
<b>338.3-338.4</b>	<b>2006</b>	<b>Code to Pain, by site</b>
<b>341.20</b>	<b>2006</b>	<b>323.9</b>
<b>341.21</b>	<b>2006</b>	<b>323.8</b>
<b>341.22</b>	<b>2006</b>	<b>323.9</b>
342.00-342.02	1994	342.0
342.10-342.12	1994	342.1
342.80-342.82	1994	342.9
342.90-342.92	1994	342.9
344.00-344.04, 344.09	1994	344.0
344.30-344.32	1994	344.3
344.40-344.42	1994	344.4
344.81, 344.89	1993	344.8
345.00-345.01	1989	345.0
345.10-345.11	1989	345.1
345.40-345.41	1989	345.4
345.50-345.51	1989	345.5
345.60-345.61	1989	345.6
345.70-345.71	1989	345.7
345.80-345.81	1989	345.8
345.90-345.91	1989	345.9
346.00-346.01	1992	346.0
346.10-346.11	1992	346.1
346.20-346.21	1992	346.2
346.80-346.81	1992	346.8
346.90-346.91	1992	346.9
347.00-347.01	2004	347
347.10-347.11	2004	347
348.30-348.31, 348.39	2003	348.3
355.71	1993	354.4
355.79	1993	355.7
357.81-357.82, 357.89	2002	357.8
358.00-358.01	2003	358.0
359.81, 359.89	2002	359.8



### Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
362.03-362.07	2005	362.02
365.83	2002	365.89
371.82	1992	371.89
372.81, 372.89	2000	372.8
374.87	1990	374.89
<b>377.43</b>	<b>2006</b>	<b>377.49</b>
<b>379.60-379.63</b>	<b>2006</b>	<b>379.99</b>
380.03	2004	733.99
<b>389.15-389.16</b>	<b>2006</b>	<b>389.18</b>
403.00-403.01	1989	403.0
403.10-403.11	1989	403.1
403.90-403.91	1989	403.9
404.00-404.03	1989	404.0
404.10-404.13	1989	404.1
404.90-404.93	1989	404.9
410.00-410.02	1989	410.0
410.10-410.12	1989	410.1
410.20-410.22	1989	410.2
410.30-410.32	1989	410.3
410.40-410.42	1989	410.4
410.50-410.52	1989	410.5
410.60-410.62	1989	410.6
410.70-410.72	1989	410.7
410.80-410.82	1989	410.8
410.90-410.92	1989	410.9
411.81	1989	410.9
411.89	1989	411.8
414.00-414.01	1994	414.0
414.02-414.03	1994	996.03
414.04-414.05	1996	414.00
414.06	2002	414.00
414.07	2003	414.06
414.12	2002	414.11

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
415.11	1995	997.3 + 415.1
415.19	1995	415.1
426.82	2005	794.31
428.20-428.23	2002	428.0
428.30-428.33	2002	428.0
428.40-428.43	2002	428.0
429.71, 429.79	1989	410.0-410.9
<b>429.83</b>	<b>2006</b>	<b>429.89</b>
433.00-433.01	1993	433.0
433.10-433.11	1993	433.1
433.20-433.21	1993	433.2
433.30-433.31	1993	433.3
433.80-433.81	1993	433.8
433.90-433.91	1993	433.9
434.00-434.01	1993	434.0
434.10-434.11	1993	434.1
434.90-434.91	1993	434.9
435.3	1995	435.0 + 435.1
437.7	1992	780.9
438.0	1997	294.9 + 438
438.10	1997	784.5 + 438
438.11	1997	784.3 + 438
438.12	1997	784.4 + 438
438.19	1997	784.5 + 438
438.20	1997	342.90 + 438
438.21	1997	342.91 + 438
438.22	1997	342.92 + 438
438.30	1997	344.40 + 438
438.31	1997	344.41 + 438
438.32	1997	344.42 + 438
438.40	1997	344.30 + 438
438.41	1997	344.31 + 438
438.42	1997	344.32 + 438
438.50-438.52	1997	344.89 + 438
438.53	1998	438.50
438.6-438.7	2002	438.89

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
438.81	1997	784.69 + 438
438.82	1997	787.2 + 438
438.83-438.85	2002	438.89
438.89	1997	438
438.9	1997	438
440.20-440.22	1992	440.2
440.23	1993	440.20 + (707.1 or 707.8 or 707.9)
440.24	1993	440.20 + 785.4
440.29	1993	440.2
440.30-440.32	1994	996.1
441.00-441.03	1994	441.0
441.6	1993	441.1 + 441.3
441.7	1993	441.2 + 441.4
443.21	2002	442.81
443.22	2002	442.2
443.23	2002	442.1
443.24, 443.29	2002	442.89
443.82	2005	443.89
445.01-445.02	2002	440.29
445.81	2002	440.1
445.89	2002	440.8
446.20-446.21, 446.29	1990	446.2
451.82-451.84	1993	451.89
453.40-453.42	2004	453.8
454.8	2002	454.9
458.2	1995	997.9 + 458.9
458.21, 458.29	2003	458.2
458.8	1997	458.9
459.10-459.13; 459.19	2002	459.1
459.30-459.33; 459.39	2002	459.89
464.00-464.01	2001	464.0
464.50-464.51	2001	464.0
466.11, 466.19	1996	466.1

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
474.0 (Code title restated)	1997	474.0
474.00-474.02	1997	474.0
477.1	2000	477.8
477.2	2004	477.8
<b>478.11, 478.19</b>	<b>2006</b>	<b>478.1</b>
480.3	2003	480.8
482.30-482.39	1992	482.3
482.40-482.41, 482.49	1998	482.4
482.81-482.83, 482.89	1992	482.8
482.84	1997	482.83
483.0	1992	483
483.1	1996	078.88 + 484.8
483.8	1992	483
491.20-491.21	1991	491.2
491.22	2004	491.21
493.02	2000	493.00
493.12	2000	493.10
493.20	1989	493.90
493.21	1989	493.91
493.22	2000	493.20
493.81	2003	519.1
493.82	2003	493.90-493.91
493.92	2000	493.90
494.0-494.1	2000	494
512.1	1994	997.3
517.3	2003	282.62
518.6	1997	518.89
<b>518.7</b>	<b>2006</b>	<b>997.3</b>
518.81	1987	799.1
518.82-518.89	1987	518.8
518.83, 518.84	1998	518.81
519.00-519.02; 519.09	1998	519.0
<b>519.11, 519.19</b>	<b>2006</b>	<b>519.1</b>

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
521.00-521.05, 521.09	2001	521.0
521.06-521.08	2004	521.09
521.10-521.15	2004	521.1
521.20-521.25	2004	521.2
521.30-521.35	2004	521.3
521.40-521.45	2004	521.4
<b>521.81</b>	<b>2006</b>	<b>873.63; 873.73</b>
<b>521.89</b>	<b>2006</b>	<b>521.8</b>
<b>523.00-523.01</b>	<b>2006</b>	<b>523.0</b>
<b>523.10-523.11</b>	<b>2006</b>	<b>523.1</b>
523.20-523.25	2004	523.2
<b>523.30-523.33</b>	<b>2006</b>	<b>523.3</b>
<b>523.40-523.42</b>	<b>2006</b>	<b>523.4</b>
524.00-524.06, 524.09	1992	524.0
524.07	2004	524.09
524.10-524.12, 524.19	1992	524.1
524.20-524.29	2004	524.2
524.30-524.37, 524.39	2004	524.3
524.50-524.57, 524.59	2004	524.5
524.60-524.63, 524.69	1991	524.6
524.64	2004	524.69
524.70-524.77, 524.79	1992	524.8
524.75-524.76	2004	524.79
524.81-524.82, 524.89	2004	524.8
525.10-525.13, 525.19	2001	525.1
525.20-525.26	2004	525.2
525.40-525.44	2005	525.10
525.50-525.54	2005	525.10
<b>525.60-525.67; 525.69</b>	<b>2006</b>	<b>525.8</b>
<b>526.61-526.63; 526.69</b>	<b>2006</b>	<b>526.8</b>
<b>528.00-528.02; 528.09</b>	<b>2006</b>	<b>528.0</b>
528.71-528.72, 528.79	2004	528.7
530.10-530.11, 530.19	1993	530.1
530.12	2001	530.10
530.20-530.21	2003	530.2
530.81	1993	530.1
530.82-530.84, 530.89	1993	530.8
530.85	2003	530.2
530.86-530.87	2004	997.4

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
535.00-535.01	1991	535.0
535.10-535.11	1991	535.1
535.20-535.21	1991	535.2
535.30-535.31	1991	535.3
535.40-535.41	1991	535.4
535.50-535.51	1991	535.5
535.60-535.61	1991	535.6
536.3	1994	536.8
536.40-536.42, 536.49	1998	997.4
537.82	1990	537.89
537.83	1991	537.82
537.84	2002	531.00
<b>538</b>	<b>2006</b>	<b>558.9</b>
556.0-556.6, 556.8-556.9	1994	556
558.3	2000	558.9
562.02	1991	562.00
562.03	1991	562.01
562.12	1991	562.10
562.13	1991	562.11
564.00-564.09	2001	564.0
564.81, 564.89	1998	564.8
567.21-567.23, 567.29	2005	567.2
567.31	2005	728.89
567.38	2005	567.2
567.39	2005	567.9
567.81-567.82, 567.89	2005	567.8
569.60-569.61, 569.69	1995	569.6
569.62	1998	569.69
569.84	1990	557.1
569.85	1991	569.84
569.86	2002	569.82
574.60	1996	574.00 + 574.30
574.61	1996	574.01 + 574.31
574.70	1996	574.10 + 574.40
574.71	1996	574.11 + 574.41

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
574.80	1996	574.00 + 574.10, 574.30 + 574.40
574.81	1996	574.01 + 574.11, 574.31 + 574.41
574.90	1996	574.20 + 574.50
574.91	1996	574.21 + 574.51
575.10-575.11	1996	575.1
575.12	1996	575.0 + 575.1
585.1-585.6, 585.9	2005	585
588.81, 588.89	2004	588.8
593.70-593.73	1994	593.7
596.51-596.53	1992	596.5
596.54	1992	344.61
596.55-596.59	1992	596.5
599.60, 599.69	2005	599.6
599.81-599.89	1992	599.8
600.0-600.3, 600.9	2000	600
600.00-600.01	2003	600.0
600.10-600.11	2003	600.1
600.20-600.21	2003	600.2
600.90-600.91	2003	600.9
602.3	2001	602.8
607.85	2003	607.89
<b>608.20-608.24</b>	<b>2006</b>	<b>608.2</b>
608.82	2001	608.83
608.87	2001	608.89
<b>616.81; 616.89</b>	<b>2006</b>	<b>616.8</b>
618.00-618.05, 618.09	2004	618.0
618.81-618.83	2004	618.8
<b>618.84</b>	<b>2006</b>	<b>618.1</b>
618.89	2004	618.8
621.30-621.33	2004	621.3
622.10-622.12	2004	622.1

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
629.20-629.23	2004	629.8
<b>629.29</b>	<b>2006</b>	<b>629.20</b>
<b>629.81</b>	<b>2006</b>	<b>629.9</b>
<b>629.89</b>	<b>2006</b>	<b>629.8</b>
633.00-633.01	2002	633.0
633.10-633.11	2002	633.1
633.20-633.21	2002	633.2
633.80-633.81	2002	633.8
633.90-633.91	2002	633.9
645.00-645.01, 645.03	1991	645.0-645.1, 645.3 (amended 10/02/2004)
645.10-645.11, 645.13	2000	645.00-645.01, 645.03
645.20-645.21, 645.23	2000	645.00-645.01, 645.03
<b>649.00-649.04</b>	<b>2006</b>	<b>648.40-648.44</b>
<b>649.10-649.14</b>	<b>2006</b>	<b>646.10-646.14</b>
<b>649.20-649.24</b>	<b>2006</b>	<b>V23.89</b>
<b>649.30</b>	<b>2006</b>	<b>641.30; 648.90;666.30</b>
<b>649.31</b>	<b>2006</b>	<b>641.31; 648.91</b>
<b>649.32</b>	<b>2006</b>	<b>648.92;666.32</b>
<b>649.33</b>	<b>2006</b>	<b>641.33; 648.93</b>
<b>649.34</b>	<b>2006</b>	<b>648.94; 666.34</b>
<b>649.40-649.44</b>	<b>2006</b>	<b>648.90-648.94</b>
<b>649.50-649.51</b>	<b>2006</b>	<b>641.90-641.91</b>
<b>649.53</b>	<b>2006</b>	<b>641.93</b>
<b>649.60-649.64</b>	<b>2006</b>	<b>646.80-646.84</b>
651.30-651.31, 651.33	1989	651.00-651.01, 651.03
651.40-651.41, 651.43	1989	651.10-651.11, 651.13
651.50-651.51, 651.53	1989	651.20-651.21, 651.23
651.60-651.61, 651.63	1989	651.80-651.81, 651.83
651.70-651.71, 651.73	2005	651.8
654.20-654.21, 654.23	1990	654.2, 654.9
654.90-654.94	1990	654.2, 654.9
655.70-655.71, 655.73	1997	655.8
657.00-657.01, 657.03	1991	657.0-657.1, 657.3 (amended 10/02/2004)
659.60-659.61, 659.63	1992	659.80-659.81, 659.83
659.70-659.71, 659.73	1998	656.30-656.31, 656.33



### Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
665.10, 665.11	1992	665.10-665.12, 665.14 <sup>1</sup>
670.00, 670.02, 670.04	1991	670.0-670.1, 670.3 (amended 10/02/2004)
672.00, 672.02, 672.04	1991	672.0-672.1, 672.3 (amended 10/02/2004)
674.50-674.54	2003	674.80, 674.82, 674.84
677	1994	None
686.00-686.01, 686.09	1997	686.0
690.10, 690.18	1995	690
690.11	1995	691.8 + 704.8
690.12	1995	691.8
690.8	1995	690
692.72-692.74	1992	692.79
692.75	2000	692.79
692.76-692.77	2001	692.71
692.82-692.83	1992	692.89
692.84	2004	692.89
702.0-702.8	1991	702
702.11, 702.19	1994	702.1
704.02	1993	704.09
705.21-705.22	2004	780.8
707.00-707.07, 707.09	2004	707.0
707.10-707.15, 707.19	2000	707.1
709.00-709.01, 709.09	1994	709.0
710.5	1992	288.3, 729.1
718.70-718.79	2001	718.80-718.89
719.7	2003	719.70, 719.75-719.79 (codes deleted)
727.83	2000	727.89

### Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
728.86	1995	729.4
728.87	2003	728.9
728.88	2003	728.89
<b>729.71-729.73;729.79</b>	<b>2006</b>	<b>729.9</b>
<b>731.3</b>	<b>2006</b>	<b>733.99</b>
733.10-733.16, 733.19	1993	733.1
733.93	2001	733.16
733.94-733.95	2001	733.19
738.10-738.19	1992	738.1
747.60-747.64, 747.69	1993	747.6
747.82	1993	747.89
747.83	2002	747.89, 747.9
752.51-752.52	1996	752.5
752.61-752.63	1996	752.6
752.64-752.65, 752.69	1996	752.8
752.81, 752.89	2003	752.8
753.10-753.17, 753.19	1990	753.1
753.20-753.23, 753.29	1996	753.2
756.70-756.71, 756.79	1997	756.7
758.31-758.33, 758.39	2004	758.3
758.81	1996	758.8
758.89	1996	758.9
759.81-759.82, 759.89	1989	759.8
759.83	1994	759.89
760.75	1991	760.79
760.76	1994	760.79
760.77-760.78	2005	760.79
763.81-763.83, 763.89	1998	763.8
763.84	2005	770.1
764.00-764.09	1988	764.0
764.10-764.19	1988	764.1
764.20-764.29	1988	764.2
764.90-764.99	1988	764.9

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
765.00-765.09	1988	765.0
765.10-765.19	1988	765.1
765.20-765.24	2002	765.00-765.09
765.25-765.29	2002	765.10-765.19
766.21-766.22	2003	766.2
767.11-767.19	2003	767.1
<b>768.7</b>	<b>2006</b>	<b>768.9</b>
770.10-770.18	2005	770.1
770.81-770.84, 770.89	2002	770.8
770.85-770.86	2005	770.1
<b>770.87</b>	<b>2006</b>	<b>770.89</b>
<b>770.88</b>	<b>2006</b>	<b>768.9</b>
771.81-771.83, 771.89	2002	771.8
772.10-772.14	2001	772.1
<b>775.81, 775.89</b>	<b>2006</b>	<b>775.8</b>
779.7	2001	772.1
779.81-779.82	2002	779.8
779.83	2003	779.89
779.84	2005	None (omit code)
<b>779.85</b>	<b>2006</b>	<b>779.89</b>
779.89	2002	779.8
780.01-780.02, 780.09	1992	780.0
780.03	1993	780.01
780.31, 780.39	1997	780.3
<b>780.32</b>	<b>2006</b>	<b>780.39</b>
780.57	1992	780.51, 780.53
780.58	2004	780.59
780.71, 780.79	1998	780.7
780.91-780.92	2002	780.9
780.93-780.94	2003	780.99
780.95	2005	780.99
<b>780.96, 780.97</b>	<b>2006</b>	<b>780.99</b>
780.99	2002	780.9
781.8	1994	781.9
781.91-781.92, 781.99	2000	781.9

### Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
781.93	2002	723.5
781.94	2003	781.99
783.21	2000	783.2
783.22	2000	783.4
783.40-783.43	2000	783.4
783.7	2000	783.4
<b>784.91</b>	<b>2006</b>	<b>473.9</b>
<b>784.99</b>	<b>2006</b>	<b>784.9</b>
785.52	2003	785.59
786.03-786.07	1998	786.09
787.01-787.03	1994	787.0
787.91	1995	558.9
787.99	1995	787.9
788.20-788.21, 788.29	1993	788.2
788.30-788.37; 788.39	1992	788.3
788.38	2004	788.39
788.41-788.43	1993	788.4
788.61-788.62	1993	788.6
788.63	2003	788.69
<b>788.64-788.65</b>	<b>2006</b>	<b>788.69</b>
788.69	1993	788.6
789.00-789.07, 789.09	1994	789.0
789.30-789.37, 789.39	1994	789.3
789.40-789.47, 789.49	1994	789.4
789.60-789.67, 789.69	1994	789.6
790.01, 790.09	2000	790.0
790.21-790.22, 790.29	2003	790.2
790.91, 790.93, 790.99	1993	790.9
790.92	1993	286.9
790.94	1997	790.99
790.95	2004	790.99
792.5	2000	792.9
793.80-793.81, 793.89	2001	793.8
<b>793.91; 793.99</b>	<b>2006</b>	<b>793.9</b>
795.00-795.02, 795.09	2002	795.0

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
795.03, 795.04	2004	622.1
795.05, 795.08	2004	795.09
<b>795.06</b>	<b>2006</b>	<b>795.04</b>
795.31-795.39	2002	795.3
795.71	1994	795.8 (Code deleted)
795.79	1994	795.7
795.8	1986	795.7
<b>795.81-795.82, 795.89</b>	<b>2006</b>	<b>796.9</b>
796.5	1997	796.9
796.6	2004	796.9
799.01-799.02	2005	799.0
799.81, 799.89	2003	799.8
813.45	2002	813.42
823.40-823.42	2002	823.80-823.82
840.7	2001	840.8
850.11-850.12	2003	850.1
864.05	1992	864.09
864.15	1992	864.19
909.5	1994	909.9
922.31-922.33	1996	922.3
925.1-925.2	1993	925
<b>958.90-958.93; 958.99</b>	<b>2006</b>	<b>958.8</b>
959.0 (Code title restated)	1997	959.0
959.01	1997	854.00
959.09	1997	959.0
959.11-959.14, 959.19	2003	959.1
965.61, 965.69	1998	965.6
989.81-989.84, 989.89	1995	989.8
<b>995.20-995.23</b>	<b>2006</b>	<b>995.2</b>
<b>995.27; 995.29</b>	<b>2006</b>	<b>995.2</b>
995.50-995.55, 995.59	1996	995.5

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
995.60-995.69	1993	995.0
995.7	2000	None
995.80, 995.82-995.85	1996	995.81
995.81 (Code title restated)	1996	995.81
995.86	1998	995.89
995.90-995.94	2002	038.0-038.9
996.04	1994	996.09
996.40-996.47, 996.49	2005	996.4
996.51-996.59	1987	996.5
996.55	1998	996.52
996.56	1998	996.59
996.57	2003	996.59
996.60-996.69	1989	996.6
996.68	1998	996.69
996.70-996.79	1989	996.7
996.80-996.84, 996.86, 996.89	1987	996.8
996.85	1990	996.89
996.87	2000	996.89
997.00-997.01, 997.09	1995	997.0
997.02	1995	997.9 + 430-434, 436
997.71	2001	997.4
997.72	2001	997.5
997.79	2001	997.2
997.91, 997.99	1995	997.9
998.11-998.12	1996	998.1
998.13	1996	998.89
998.31-998.32	2002	998.3
998.51, 998.59	1996	998.5
998.81-998.82, 998.89	1994	998.8
998.83	1996	998.89
V01.71, V01.79	2004	V01.7
V01.81, V01.89	2002	V01.8
V01.82	2003	V01.89
V01.83-V01.84	2004	V01.89
V02.51-V02.52, V02.59	1998	V02.5
V02.60-V02.62, V02.69	1997	V02.6
V03.81-V03.82, V03.89	1994	V03.8
V04.81-V04.82, V04.89	2003	V04.8

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
V05.3-V05.4	1993	V05.8
V06.5-V06.6	1994	V06.8
V07.31, V07.39 V07.4	1994 1992	V07.3 V07.8
V08	1994	044.9, 795.8 (codes deleted)
V09.0-V09.91	1993	None
V10.48 V10.53	1998 2001	V10.49 V10.59
V12.00-V12.03, V12.09 V12.40-V12.41, V12.49 V12.42 V12.50-V12.52, V12.59 V12.60-V12.61; V12.69 V12.70-V12.72, V12.79	1994 1997 2005 1995 2005 1994	V12.0 V12.4 V12.49 V12.5 V12.6 V12.7
V13.00-V13.01, V13.09 V13.02-V13.03 V13.21, V13.29 V13.61, V13.69	1994 2005 2002 1998	V13.0 V13.09 V13.2 V13.6
V15.01-V15.09 V15.41-V15.42, V15.49 V15.82 V15.84-V15.86 V15.87 V15.88	2000 1996 1994 1995 2003 2005	V15.0 V15.4 305.13 (code deleted) V15.89 V15.89 V15.49
V16.40-V16.43, V16.49 V16.51, V16.59	1997 1998	V16.4 V16.5
V17.81, V17.89	2005	V17.8
<b>V18.51; V18.59</b> V18.61, V18.69 V18.9	<b>2006</b> 1998 2005	<b>V18.5</b> V18.6 V19.8
V21.30-V21.35	2000	None
V23.41, V23.49	2002	V23.4

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
V23.7	1989	V23.8
V23.81-V23.84, V23.89	1998	V23.8
V25.03	2003	V25.01
V25.43	1992	V25.49
V25.5	1992	V25.8
V26.21-V26.22, V26.29	2000	V26.2
V26.31-V26.33	2005	V26.3
<b>V26.34-V26.35</b>	<b>2006</b>	<b>V26.31</b>
<b>V26.39</b>	<b>2006</b>	<b>V26.32</b>
V26.51, V26.52	1998	None
V28.6	1997	V28.8
V29.0-V29.1, V29.8	1992	V71.8
V29.2	1994	V29.8
V29.3	1998	V29.8
V29.9	1992	V71.9
V30.00-V30.01	1989	V30.0
V31.00-V31.01	1989	V31.0
V32.00-V32.01	1989	V32.0
V33.00-V33.01	1989	V33.0
V34.00-V34.01	1989	V34.0
V35.00-V35.01	1989	V35.0
V36.00-V36.01	1989	V36.0
V37.00-V37.01	1989	V37.0
V39.00-V39.01	1989	V39.0
V42.81-V42.83, V42.89	1997	V42.8
V42.84	2000	V42.89
V43.21-V43.22	2003	V43.2
V43.60-43.66, V43.69	1994	V43.6
V43.81-V43.82, V43.89	1995	V43.8
V43.83	1998	V43.89
V44.50-V44.52, V44.59	1998	V44.5
V45.00, V45.02, V45.09	1994	V45.89
V45.01	1994	V45.0
V45.51	1994	V45.5
V45.52, V45.59	1994	V45.89
V45.61, V45.69	1997	V45.6



## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
V45.71	1997	611.8
V45.72	1997	569.89
V45.73	1997	593.89
V45.74	2000	593.89, 596.8
V45.75	2000	V45.89
V45.76	2000	518.89
V45.77	2000	602.8, 607.89, 608.89, 620.8, 621.8, 622.8
V45.78	2000	360.89
V45.79	2000	255.8, 289.59, 388.8, 569.49, 577.8
V45.82	1994	V45.89
V45.83	1995	V45.89
V45.84	2001	None
V45.85	2003	V45.89
<b>V45.86</b>	<b>2006</b>	<b>V45.89</b>
V46.11-V46.12	2004	V46.1
V46.13-V46.14	2005	V46.11
V46.2	2002	V46.8
V49.60-V49.67	1994	V49.5
V49.70-V49.77	1994	V49.5
V49.81	2000	None
V49.82	2001	None
V49.83	2004	None
V49.84	2005	V49.89
V49.89	2000	V49.8
V50.41-V50.42, V50.49	1994	V50.8
V53.01-V53.02, V53.09	1997	V53.0
V53.31-V53.32, V53.39	1994	V53.9
V53.90-V53.91, V53.99	2003	V53.9
V54.01-V54.02, V54.09	2003	V54.0
V54.10-V54.17, V54.19	2002	V54.8
V54.20-V54.27, V54.29	2002	V54.8
V54.81, V54.89	2002	V54.8
V56.1	1995	V58.89
V56.1 (Code title restated)	1998	V56.1
V56.2	1998	V56.1
V56.31-V56.32	2000	V56.8
V57.21-V57.22	1994	V57.2

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
V58.11	2005	V58.1
V58.12	2005	140-208; 230-239
<b>V58.30-V58.32</b>	<b>2006</b>	<b>V58.3</b>
V58.41, V58.49	1994	V58.4
V58.42, V58.43	2002	V58.49
V58.44	2004	V58.71-V58.78
V58.61, V58.69	1995	V67.51
V58.62	1998	V58.69
V58.63-V58.65	2003	V58.69
V58.66-V58.67	2004	V58.69
V58.71-V58.78	2002	V58.49
V58.81, V58.89	1994	V58.8
V58.82	1995	V58.89
V58.83	2000	V58.89
V59.01-V59.02, V59.09	1995	V59.0
V59.6	1995	V59.8
V59.70-V59.74	2005	V59.8
V61.10-V61.12	1996	V61.1
V61.22	1996	V61.21
V62.83	1996	V65.49
V62.84	2005	V62.89
V64.00-V64.05	2005	V64.0
V64.06	2005	V64.2
V64.07-V64.09	2005	V64.0
V64.4	1997	None
V64.41-V64.43	2003	V64.4
V65.11, V65.19	2003	V65.1
V65.40-V65.45, V65.49	1994	V65.4
V65.46	2003	V65.1
V66.7	1996	None
V67.00-V67.01, V67.09	2000	V67.0
V69.0-V69.3	1994	None
V69.4	2004	V69.8
V69.5	2005	V69.8
V69.8-V69.9	1994	None
V71.81, V71.89	2000	V71.8
V71.82-V71.83	2002	V71.89

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
<b>V72.11,V72.19</b>	<b>2006</b>	<b>V72.1</b>
V72.31-V72.32	2004	V72.3
V72.40-V72.41	2004	V72.4
V72.42	2005	V22.0-V22.1
V72.81-V72.85	1993	V72.8
V72.86	2005	V72.83
V73.88-V73.89	1993	V73.8
V73.98-V73.99	1993	V73.9
V76.10-V76.12, V76.19	1997	V76.1
V76.44-V76.45	1998	V76.49
V76.46-V76.47	2000	V76.49
V76.50-V76.52	2000	V76.49
V76.81, V76.89	2000	V76.8
V77.91, V77.99	2000	V77.9
<b>V82.71</b>	<b>2006</b>	<b>V26.31</b>
<b>V82.79</b>	<b>2006</b>	<b>V26.32</b>
V82.81, V82.89	2000	V82.8
V83.01-V83.02	2001	None
V83.81	2002	None
V83.89	2002	V19.8
V84.01-V84.04, V84.09	2004	None
V84.8	2004	None
V85.0	2005	None
V85.1	2005	None
V85.21-V85.25	2005	None
V85.30-V85.39	2005	None
V85.4	2005	None
<b>V85.51-V85.54</b>	<b>2006</b>	<b>None</b>
<b>V86.0-V86.1</b>	<b>2006</b>	<b>None</b>
E854.8	1995	E858.8
E869.4	1994	E869.8
E880.1	1995	E884.9
E884.3-E884.4	1995	E884.2

## Changes in ICD-9-CM diagnosis codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
E884.5-E884.6	1995	E884.9
E885.0	2002	E885.9
E885.1-E885.4, E885.9	2000	E885
E888.0	2001	E920
E888.1, E888.8-E888.9	2001	E888
E906.5	1995	E906.3
E908.0-E908.4, E908.8-E908.9	1995	E908
E909.0-E909.4, E909.8-E909.9	1995	E909
E917.3, E917.4	2001	E917.9
E917.5	2001	E886.0
E917.6	2001	E917.1
E917.7, E917.8	2001	E888
E920.5	1995	E920.4
E922.4	1997	E917.9
E922.5	2002	E922.8
E924.2	1995	E924.0
E928.3	2000	E928.8
E928.4, E928.5	2003	E928.8
E955.6	1997	E955.9
E955.7	2002	E955.4
E967.2	1996	E967.0
E967.3	1996	None
E967.4-E967.8	1996	E967.1
E968.5	1995	E968.8
E968.6	1997	E968.8
E968.7	2000	E968.8
E985.6	1997	E985.4
E985.7	2002	E985.4
E979.0-E979.9	2002	E960.0-E966, E968.0-E968.9

**Changes in ICD-9-CM diagnosis codes, 1986-2006**

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
E999.0	2002	E999
E999.1	2002	E969

<sup>1</sup>The title for the subcategory, 665.1, has been changed, making the fifth-digit subclassification, 665.12 and 665.14, invalid.

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
00.01-00.03, 00.09	2002	99.99
00.10	2002	99.25
00.11	2002	99.19
00.12	2002	93.98
00.13	2002	99.29
00.14	2002	99.21
00.15	2003	99.28
00.16	2004	None
00.17	2004	99.29
00.18	2005	99.29
00.21	2004	88.71
00.22	2004	88.73
00.23	2004	88.77
00.24	2004	88.72
00.25	2004	88.75
00.28-00.29	2004	88.79
00.31-00.35, 00.39	2004	None
00.40-00.43	2005	None
<b>00.44</b>	<b>2006</b>	<b>None</b>
00.45-00.48	2005	None
00.50	2002	37.80-37.87 & 37.70-37.74, 37.76
00.51	2002	37.94
00.52	2002	None
00.53	2002	37.80-37.87
00.54	2002	37.96
00.55	2002	39.90
<b>00.56</b>	<b>2006</b>	<b>89.63</b>
<b>00.57</b>	<b>2006</b>	<b>37.79</b>
00.61-00.62	2004	39.50
00.63-00.65	2004	00.55, 39.90
00.66	2005	36.01 (deleted), 36.02 (deleted), 36.05 (deleted)
00.70-00.73	2005	81.53
00.74-00.76	2005	None
<b>00.77</b>	<b>2006</b>	<b>None</b>
00.80-00.84	2005	81.55
<b>00.85</b>	<b>2006</b>	<b>81.51</b>
<b>00.86- 00.87</b>	<b>2006</b>	<b>81.52</b>

### Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
00.91-00.93	2004	None
01.26-01.27	2005	None
<b>01.28</b>	<b>2006</b>	<b>01.26</b>
02.96	1992	89.19
03.90	1987	03.99 (Insertion of catheter)
05.25	1995	39.7
11.75	1989	11.79
11.76	1989	11.62
<b>13.90, 13.91</b>	<b>2006</b>	<b>13.9 (code deleted)</b>
20.96-20.98	1986	20.95
22.12	1988	22.11
26.12	1988	26.11
27.64	2004	27.69
29.31	1991	83.02
29.32, 29.33, 29.39	1991	29.3
31.45	1988	31.43-31.44
31.95	1989	31.75
32.01, 32.09	1989	32.0
32.22	1995	32.29, 32.9
<b>32.23- 32.26</b>	<b>2006</b>	<b>32.29</b>
32.28	1989	32.29
33.27	1987	33.22 + 33.27
33.28	1987	33.27
33.29	1987	33.28-33.29
33.50-33.52	1995	33.5
33.6	1990	33.5 + 37.5
<b>33.71, 33.78-33.79</b>	<b>2006</b>	<b>33.22, 96.05</b>
34.05	1994	34.99
<b>35.55</b>	<b>2006</b>	<b>35.53</b>
35.84	1988	35.82

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
35.96	1986	35.03
36.00-36.03	1986	36.0
36.04	1986	39.97
36.05	1986	36.01 <sup>1</sup> , 36.02
36.05	1987	36.01
36.06	1995	36.01, 36.02, 36.03, 36.05
36.07	2002	36.06
36.09	1986	36.0
36.09	1991	36.00 (Code deleted)
36.17	1996	36.19
36.31, 36.32	1998	36.3
<b>36.33, 36.34</b>	<b>2006</b>	<b>36.32</b>
36.39	1998	36.3
<b>37.20</b>	<b>2006</b>	<b>37.26</b>
37.26-37.27	1988	37.29
37.28	2001	88.72
37.34	1988	37.33
37.35	1997	37.33
37.41, 37.49	2005	37.99
37.51	2003	37.5
37.52	2003	37.62
37.53, 37.54	2003	37.63
37.65, 37.66	1995	37.62
37.67	1998	37.4
37.68	2004	37.62
37.70 (Leads only)	1987	37.70 (Leads/device)
37.71-37.72 (Leads only)	1987	37.74 (Leads/device)
37.73 (Leads only)	1987	37.73 (Leads/device)
37.74 (Leads only)	1987	37.76 (Leads/device)
37.75 (Leads only)	1987	37.89 (Leads/device)
37.76 (Leads only)	1987	37.81 (Leads/device)
37.77 (Leads only)	1987	37.83-37.84 (Leads/device)
37.78	1987	37.71-37.72
37.79	1987	86.09
37.80-37.87	1992	89.49 (Code deleted; this procedure is included in the code for pacemaker insertion/replacement)
37.80-37.83 (Device only)	1987	37.73-37.77 (Leads/device)
37.85-37.87	1987	37.85
37.89	1987	37.86 + 37.89
37.90	2004	37.99
37.94-37.98	1986	37.99
38.22	1986	38.29



## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
38.44 (Abdominal aorta only)	1986	38.44 (Entire aorta)
38.45 (Thoracic aorta added)	1989	38.44-38.45
38.95	1989	38.93
39.28	1991	39.29
39.50	1995	39.59
39.65	1988	39.61
39.66	1990	39.65
39.71, 39.79	2000	39.52
39.72	2002	39.79
39.73	2005	39.79
<b>39.74</b>	<b>2006</b>	<b>38.01, 38.02</b>
39.90	1996	39.50
41.00-41.03	1988	41.0
41.04	1994	99.79
41.05, 41.06	1997	None
41.07	2000	41.04
41.08	2000	41.05
41.09	2000	41.01
42.25	1988	42.24
42.33	1989	42.32, 42.39
42.33	1990	42.91
43.11	1989	43.1
43.19	1989	43.1, 43.2
43.41	1989	43.41, 43.49
44.21, 44.29	1986	44.2
44.22	1986	44.99
44.32	2001	44.39
44.38	2004	44.39
44.43	1989	43.49, 45.32
44.44	1989	38.86
44.49	1989	43.0
44.67	2004	44.66
44.68	2004	44.69
44.93-44.94	1986	44.99
44.95, 44.96	2004	44.69
44.97, 44.98	2004	44.99
45.16	1988	45.14 (45.15 before 1987)
45.30	1989	45.31, 45.32
45.42	1988	45.41
45.43	1989	45.49

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
45.75	1988	48.66 (Code deleted; Hartmann resection added)
45.95	1987	45.93
46.13	1992	46.12 (Code deleted)
46.32	1989	46.39
46.85	1989	46.99
46.97	2000	46.99
47.01, 47.09	1996	47.0
47.11, 47.19	1996	47.1
48.36	1995	45.42
49.31, 49.39	1989	49.3
49.75, 49.76	2002	49.79
<b>50.23- 50.26</b>	<b>2006</b>	<b>50.29</b>
51.10	1989	51.97
51.11	1989	51.11, 51.97
51.14	1989	51.12
51.15	1989	51.97
51.21	1996	51.22, 51.23
51.22	1991	51.21 (Code deleted), 51.22
51.23	1991	51.22
51.24	1996	51.22, 51.23
51.64	1989	51.69
51.84-51.88	1989	51.97
51.97	1986	52.91, 51.99 or 51.82
51.98	1986	51.99
52.13	1989	51.97, 52.91
52.14	1989	52.11
52.21-52.22	1989	52.2
52.84-52.86	1996	99.29
52.93	1989	52.93 + 52.91
52.94	1989	52.09
52.97, 52.98	1989	52.91
52.99	1989	52.93, 52.94, 52.99
54.24	1987	54.23
54.25	1993	54.98
54.51, 54.59	1996	54.5
55.03-55.04	1986	55.02

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
<b>55.32-55.35</b>	<b>2006</b>	<b>55.39</b>
56.33-56.34	1987	56.33
56.35	1987	45.12
57.17-57.18	1989	57.21
57.22	1989	57.22, 57.82
58.31, 58.39	1990	58.3
58.93	1986	57.99
59.03	1996	59.02
59.12	1996	59.11
59.72	1995	59.79
59.96	1986	59.95
60.21, 60.29	1995	60.2
60.95	1991	60.99
60.96, 60.97	2000	60.29
64.97	1986	64.95
65.01, 65.09	1996	65.0
65.13	1996	65.12
65.14	1996	65.19
65.23	1996	65.21
65.24	1996	65.22
65.25	1996	65.29
65.31, 65.39	1996	65.3
65.41, 65.49	1996	65.4
65.53	1996	65.51
65.54	1996	65.52
65.63	1996	65.61
65.64	1996	65.62
65.74	1996	65.71
65.75	1996	65.72
65.76	1996	65.73
65.81, 65.89	1996	65.8
66.01	1992	66.0
66.02	1992	66.73
67.51, 67.59	2001	67.5
68.15	1987	68.14

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
68.16	1987	68.13
68.23	1996	68.29
68.31, 68.39	2003	68.3
<b>68.41, 68.49</b>	<b>2006</b>	<b>68.4 (code deleted)</b>
68.51, 68.59	1996	68.5
<b>68.61. 68.69</b>	<b>2006</b>	<b>68.6 (code deleted)</b>
<b>68.71, 68.79</b>	<b>2006</b>	<b>68.7 (code deleted)</b>
68.9	1992	68.4
74.3	1992	69.11 (Code deleted)
75.37	1998	99.29
75.38	2001	75.34
77.56	1989	77.89, 78.49, 81.18
77.57	1989	77.89, 80.48, 81.18, 83.85
77.58	1989	77.59, 81.18
78.10	1991	78.40
78.11	1991	78.41
78.12	1991	78.42
78.13	1991	78.43
78.14	1991	78.44
78.15	1991	78.45
78.16	1991	78.46
78.17	1991	78.47
78.18	1991	78.48
78.19	1991	78.49
78.20	1991	78.10, 78.20, 78.30
78.21	1991	78.11, 78.31
78.22	1991	78.12, 78.22, 78.32
78.23	1991	78.13, 78.23, 78.33
78.24	1991	78.14, 78.34
78.25	1991	78.15, 78.25, 78.35
78.27	1991	78.17, 78.27, 78.37
78.28	1991	78.18, 78.38
78.29	1991	78.11, 78.16, 78.19, 78.29, 78.39
78.39	1991	78.31
78.90 <sup>2</sup>	1987	78.40
78.91 <sup>2</sup>	1987	78.41
78.92 <sup>2</sup>	1987	78.42
78.93 <sup>2</sup>	1987	78.43
78.94 <sup>2</sup>	1987	78.44
78.95 <sup>2</sup>	1987	78.45
78.96 <sup>2</sup>	1987	78.46

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
78.97 <sup>2</sup>	1987	78.47
78.98 <sup>2</sup>	1987	78.48
78.99 <sup>2</sup>	1987	78.49
80.50-80.59	1986	80.5
81.03	1989	81.02
81.04-81.05	1989	81.03, 81.04, 81.05
81.06-81.07	1989	81.06, 81.07
81.08	1989	81.06, 81.07, 81.08
81.09	1989	81.08
81.18	2005	81.99
81.30-81.39	2001	81.09
81.40	1989	81.69
81.51	1989	81.51, 81.59
81.52	1989	81.61-81.64
81.53	1989	81.51, 81.59, 81.61-81.64
81.54-81.55	1989	81.41
81.56	1989	81.48
81.57	1989	81.31, 81.39
81.59	1989	81.39
81.61 <sup>3</sup>	2002	81.00-81.08, 81.30-81.39
81.62-81.64	2003	None
81.65, 81.66	2004	78.49
81.72	1989	81.79
81.73-81.74	1989	81.86 (Code deleted)
81.75	1989	81.87 (Code deleted)
81.79	1989	81.79, 81.87
81.80	1989	81.81
81.97	1992	81.59
84.51, 84.52	2002	None
84.53, 84.54	2004	78.30, 78.32-78.35, 78.37-78.39
84.55, 84.59	2004	None
84.56-84.57	2005	None
84.58	2005	84.59
84.60-84.69	2004	80.51
84.71-84.73	2005	None
85.95, 85.96	1987	85.99
86.06	1987	86.09
86.07	1990	86.09
86.27	1986	86.22-86.23
86.28	1988	86.22
86.67	1998	86.65

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
86.93	1987	86.89
86.94-86.96	2004	02.93, 03.93, 04.92
86.97	2005	86.94
86.98	2005	86.95
88.90	1986	88.39
88.91	1986	89.15
88.92	1986	89.39
88.93	1986	89.15
88.94	1986	89.39
88.95	1986	89.29
88.96	2002	88.91-88.97
88.97	1989	88.99
88.98	1989	88.90
88.99	1986	89.39
89.10, 89.19	1989	89.15
89.17-89.18	1988	89.15
89.49	2004	89.59
89.50	1991	89.54
89.60	2002	89.65
92.20	2005	92.28
92.3	1995	01.59, 04.07, 07.63, 07.68
92.30-92.33, 92.39	1998	92.3
93.90	1988	93.92
94.61-94.69	1989	94.25
96.29	1998	96.39
96.6	1986	96.35
96.70-96.72	1991	93.92 (code deleted)
97.05	1989	51.97
97.44	2001	37.64
98.51-98.52, 98.59	1989	59.96 (code deleted)
99.00	1995	99.02
99.10	1998	99.29
99.15	1986	99.29
99.20	1998	99.29
99.28	1994	99.25
99.71-99.79 <sup>4</sup>	1988	99.07

## Changes in ICD-9-CM in procedure codes, 1986-2006

<u>Current code(s) assignment</u>	<u>Effective October 1</u>	<u>Previous code(s) assignment</u>
99.75	2000	99.29
99.76	2002	99.79
99.77	2002	None
99.78	2004	99.71
99.85	1987	93.35
99.86	1987	93.39
99.88	1988	99.83

<sup>1</sup>Before October 1986 contents of current code 36.05 would have been assigned to 36.0.

<sup>2</sup>Codes 78.90-78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90-78.99 were reassigned to codes 78.40-78.49.

<sup>3</sup>Code 81.61 was deleted effective 10/01/2005.

<sup>4</sup>Codes 99.71-99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71-99.79 were reclassified as "Therapeutic apheresis".