Persons with disabilities experiencing problems accessing PDF file should contact nchsed@cdc.gov, or call 301-458-4688

HEALTH STATISTICS

FROM THE U. S. NATIONAL HEALTH SURVEY

OLDER PERSONS

selected health characteristics

United States

July 1957 - June 1959

Selected statistics relating to limitation of activity and mobility, disability days, chronic and acute conditions, and physician and dental visits for persons 45 years and older. Based on data collected in household interviews during the period July 1957-June 1959.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE Arthur S. Flemming, Secretary

Public Health Service Leroy E. Burney, Surgeon General

Division of Public Health Methods William H. Stewart, M. D., Chief

U. S. NATIONAL HEALTH SURVEY

Forrest E. Linder, Ph. D., Director
Theodore D. Woolsey, Assistant Director
Alice M. Waterhouse, M. D., Chief Medical Advisor
James E. Kelly, D. D. S., Dental Advisor
Walt R. Simmons, Statistical Advisor
O. K. Sagen, Ph. D., Chief, Special Studies
Philip S. Lawrence, Sc. D., Chief, Health Interview Survey
Margery R. Cunningham, Staff Assistant
Robert T. Little, Chief, Automatic Data Processing

The U. S. National Health Survey is a continuing program under which the Public Health Service makes studies to determine the extent of illness and disability in the population of the United States and to gather related information. It is authorized by Public Law 652, 84th Congress.

CO-OPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies. For the Health Interview Survey the Bureau of the Census designed and selected the sample, conducted the household interviews, and processed the data in accordance with specifications established by the Public Health Service.

CONTENTS

	Page
Introduction	1
Summary	2
Source and Description of Data	3
Disability Among Older Persons Long-Term Disability	4
Disability Days	6
Detailed Tables	8
Illness Among Older Persons	30
Prevalence of Chronic Conditions	30
Incidence of Acute Conditions	32
Detailed Tables	34
Use of Medical and Dental Services by	
Older Persons	45
Physician Visits	45
Dental Visits	46
Detailed Tables	48
Population	58
Detailed Tables	58
Appendix 1. Technical Notes On Methods	63
Background of This Report	63
Statistical Design of the Health Interview	
Survey	63
General Qualifications	64
Reliability of Estimates	64
Appendix II. Definitions of Certain Terms	
Used in This Report	68
Demographic, Social, and Economic Terms	68
Location of Residence Terms	68
Terms Defining Morbidity Conditions	69
Terms Relating to Conditions Terms Relating to Disability	69 70
Disability Days	70
Long-Term Disability	70
Terms Relating to Persons Injured	71
Medical Care Terms	71
Dental Care Terms	72
Amondia III Oversionaciae	70

SYMBOLS AND NOTES	
Data not available (three dashes)	
Category not applicable (three dots)	• •••
Quantity is zero (1 dash)	-
Magnitude greater than zero but less than one-half of the unit used	0 or 0.0
Magnitude of the sampling error precludes showing separate estimates	(*)
NOTE: Due to rounding detailed figures within tables may not add to totals	

OLDER PERSONS

INTRODUCTION

The average life expectancy at birth of persons living in the United States had increased from 59.7 years in 1930 to 69.6 years by 1956. These years of added lifetime point up the need for health statistics pertaining to older persons in the United States, a population group particularly susceptible to the disabling and degenerative effects of illness. This report has been prepared because of the need for more comprehensive health data by persons and agencies involved with the care and support of the aged.

The title of this report, Selected Health Characteristics of Older Persons, suggests that the contents are confined to information about persons 65 years of age and older, those usually identified as the aged population. However, it will be found that many of the tables contain data for persons 45 to 64 years of age. The inclusion of health data for the younger group not only provides a baseline from which the continuing effects of aging can be measured, but offers some insight into the health characteristics of persons approaching the aged status.


While data collected in the U.S. National Health Survey include many phases of health experience, emphasis in this report has been placed on areas appropriate to older persons in the population. These areas include: chronic diseases and impairments; the limitation of activity and mobility resulting from chronic conditions; temporary disability due to disease measured in terms of restricted-activity, bed-disability, and work-loss days; and the use of medical and dental services. Likewise, related demographic factors including income, place of residence, usual activity, and family relationship were selected because of their importance to the health experience of older persons.

A general qualification of the material presented in this report is its restriction to the health experience of the civilian, noninstitutional population of the United States. Areas particularly affected by the exclusion of institutionalized persons are the prevalence estimates for certain chronic conditions, and the extent of activity and mobility limitation.


This report was prepared by Geraldine A.Gleeson, of the U. S. National Health Survey staff.

SUMMARY


From data collected by household interview in the National Health Survey during the period July 1957-June 1959, it is estimated that 77 percent of the persons in the civilian noninstitutionalized population 65 years of age or older had one or more chronic conditions. While some of these conditions were relatively minor afflictions, such as sinusitis, hay fever, or bronchitis, many were more serious conditions such as high blood pressure, heart disease, or diabetes. Among persons 45-54 years of age, 56 percent had one or more chronic conditions, and this percentage increased with age to 83 percent among persons 75 years of age or older.




There was a similar relationship with age in the number of persons with partial or major limitation of activity due to chronic conditions; about 13 percent of persons in the age group 45-54 years were unable to carry on their usual activity or had some limitation in the kind or amount of activity. This percentage of partial or major limitation of activity increased to 55 percent among persons 75 years of age or older.



Persons 65 years of age and older had on the average 14.2 days of bed disability per year because of illness due to chronic or acute conditions. A day of bed disability is defined as one on which a person was confined to bed for all or most of the day because of illness.



It is estimated that persons 65 years of age and older used physician services at a rate of 6.8 physician visits perperson per year. A physician visit is defined as consultation with a physician, in person or by telephone, for examination, diagnosis, treatment, or advice. The average number of physician visits increased from 5.0 visits per person among those 45-54 years of age to 7.3 visits per person among those 75 years and older.



The rate of dental visits for persons 65 years and older was 0.8 visits per person per year. This comparatively low rate is attributable to the high proportion of edentulous persons (approximately 60 percent) in this age segment.

SOURCE AND DESCRIPTION OF DATA

The material presented in this report is based on data collected in a continuous nationwide survey conducted by household interview. Each week a randomly selected sample of households is interviewed to obtain information relating to the health characteristics of each member of the household.

The information is based on data collected during the 2-year period, July 1957-June 1959. During this period interviews were obtained in approximately 73,000 households covering about 235,000 persons.

From these collected data certain health topics were selected for presentation in this report because of their importance in describing the health characteristics of older persons. Also taken into account in this selection of topics was the reliability and completeness of household-interview data pertaining to this segment of the population. This latter criterion for selection accounts for the exclusion of hospital data from this report. The health interview as conducted by the National Health Survey covers the hospital experience during the past year only of persons living in the household at time of interview. The hospital experience of persons who died prior to the date of interview, but who were hospitalized at any time during the previous year, is automatically excluded from the data by this procedure. Since the mortality rate as well as the rate of hospitalization is high among older persons in the population, it is known that the hospital data collected by the survey considerably understate the total hospital experience for persons 45 years and older.

The body of this report is divided into 3 sections covering the following general health areas: (1) disability among older persons measured in terms of limitation of activity and mobility, and in terms of days of disability; (2) the prevalence of selected chronic conditions and the incidence

of acute conditions among older persons, and (3) medical and dental care in terms of physician and dental visits. Each section contains text material, illustrative graphic and tabular data, followed by a number of detailed tables. Many of the detailed tables contain summary age groupings at 45 years and over, 65 years and over, and 75 years and over, as well as more detailed age groupings. Population estimates used in the computation of rates shown in the report are presented in tables 40-44.

No attempt has been made to present a comprehensive analysis of the tabulated data shown in this report. Rather, emphasis has been placed on describing the material presented, pointing out its limitations and qualifications, and defining the concepts basic to its collection and preparation. The brief discussion of the tabular material in each of the sections is restricted, for the most part, to the age group 65 years and older.

Included in Appendix l is a brief description of the survey design and methods used in estimation. Since all of the data contained in this report are estimates based on a sample of the population rather than on the entire population, they are subject to sampling errors. While the sampling errors for most of the estimates are of relatively low magnitude, where an estimated number or the numerator or denominator of a rate or percentage is small, the sampling error may be high. Tables of sampling errors and instructions for their use are also presented in Appendix l.

Definitions of the terms used in this report may be found in Appendix II. Since many of the terms have specialized meanings it is suggested that the reader familiarize himself with these definitions. A facsimile of the health-interview questionnaire used during the period July 1958–June 1959 is presented in Appendix III.

DISABILITY AMONG OLDER PERSONS

The National Health Survey measures two aspects of disability due to disease. The first, long-term disability, is subdivided into inability to carry on all or part of one's regular activities, designated as limitation of activity, and inability to move about freely or complete confinement to the house, termed limitation of mobility. The other aspect of disability refers to comparatively short periods of reduced activity described as days of restricted activity, bed disability, and loss of time from work.

Other National Health Survey publications on disability, containing data for all age groups, are: Disability Days (Series B-No. 10), and Limitation of Activity and Mobility (Series B-No. 11).

Long-Term Disability

For each person for whom a chronic condition was reported during the interview, the respondent was shown one of the Cards C through F (reproduced in Appendix III), and was asked which statement on the card best fitted the activity limitation status of that person. The cards varied in wording depending upon the usual activity of the person, but the categories of limitation listed on the cards were consistent in that each card described 4 broad limitation groups. For the population considered in this report, these groups consisted of:

- Persons completely unable to engage in the usual activities of working or keeping house.
- (2) Persons able to engage in these activities but with restrictions relating to type or amount of activity.
- (3) Persons able to participate fully in these activities but with limitations involving recreational or community activities.
- (4) Persons with no activity restrictions although they were reported to have one or more chronic conditions.

In this report, category (1) is defined as major limitation of activity. Categories (2) and (3) are combined to describe partial limitation of activity, and category (4) designates persons whose degree of activity limitation is none.

Persons with one or more chronic conditions were also classified by the extent of their limitation of mobility (Card G, Appendix III). One of the 4 degrees of mobility limitation described on the card was selected by the respondent for each person with one or more chronic conditions without regard to the major activity of the person. As in the case of activity limitation, the 4 categories have been combined into 3 limitation groups: Major limitation of mobility describes persons who were confined to the house; partial limita-

tion characterizes those who were limited or needed help in getting around outside the house, and <u>no</u> limitation of mobility describes those who had no limitation of mobility even though they were reported as having one or more chronic conditions.

Most of the terms used in this report are defined in Appendix II. However, one factor, family relationship, used in tables 13 and 14 in this section of the report, has not been defined. In the tabulation of mass data it is difficult to classify a descriptive factor such as family relationship into clear-cut categories. Only because of the general interest in the "living arrangements" of older persons has an attempt been made to devise some very broad categories. As they pertain to older persons, they are described as follows: (1) Living alone or with nonrelatives; this category includes, in general, widowed and other unmarried persons living alone or with unrelated persons or families; (2) living with relatives—married: this group includes, for the most part, married couples living in their own home or in the home of relatives; and (3) living with relatives—other status: this category includes unmarried persons, such as widowed parents living with children and other unmarried persons living with relatives.

- 1. Among persons 65 years and older about 77 percent had one or more chronic conditions (table A).
- 2. Approximately 42 percent of the population 65 years and older had some degree of chronic activity limitation, and about 18 percent had chronic mobility limitation.
- 3. About 37 percent of persons 65 years and older living inside urbanized areas had some degree of activity limitation as compared with 43 percent in other urban places, 46 percent in rural-nonfarm areas, and 51 percent in rural-farm areas (table 2). This increasingly higher percentage of activity limitation as population density decreased was noted in each of the age groups, 45-54, 55-64, 65-74, and 75+ (table B). Essentially the same pattern was apparent for chronic mobility limitation.
- 4. About 48 percent of the persons 65 years and older with family income less than \$2,000 had chronic limitation of activity. Among persons in this age group with family income of \$7,000 or more, 37 percent reported chronic limitation of activity (fig. 1). The extent to which family income and chronic limitation of activity are causally related cannot be determined from these data.
- 5. Approximately 23 percent of the persons 65 years and older who reported their major activity during the past year as "usually working" had some degree of chronic activity limitation, as compared with 36 percent of those whose major activity was keeping house, and 55 percent of those

Table A. Number and percent of persons 45 years and older with one or more chronic conditions by sex and age: United States, July 1957-June 1959

Age		er of per n thousan		Percent of population			
	Both sexes	Male	Female	Both sexes	Male	Female	
All ages-45+	32,147	14,703	17,444	65.0	62.4	67.4	
45-54 55-64 65+ 65-74 75+	11,199 9,610 11,338 7,197 4,141	5,217 4,457 5,030 3,295 1,735	5,983 5,153 6,308 3,902 2,406	56.5 64.4 77.3 74.2 83.3	53.9 62.0 75.2 72.7 80.5	58.9 66.5 79.0 75.5 85.6	

Table B. Percent of persons with partial or major limitation of activity by residence and age: United States, July 1957-June 1959

Residence		Ago	2	
	45-54	55-64	65-74	75+
Inside urbanized areas Other urban places Rural nonfarm Rural farm	11.0 13.2 13.8 16.7	18.3 22.1 24.6 26.9	31.7 35.6 39,5 43.4	49.5 56.3 58.5 65.8




Figure 1. Percent of persons 65 years and older with partial or major limitation of activity or mobility by family income.

who were retired. The high percentage of chronic activity limitation among those classified as "other" in major activity status, was probably due to persons who were unable to work during the past year and did not consider themselves as retired. Even though such persons represent only a small proportion of the total age group, the percentage in the group who were limited in kind or amount of work or were completely unable to work, would, of course, be quite large. Retired persons were, in the interview, asked to select their activity limitation status from Card C (Appendix III), which refers to ability to work; therefore, the percentage of retired persons with activity limitation represents the proportion who were limited in their ability to work, even though working was not their current major activity.

6. About 40 percent of persons 65 years and older living alone or with nonrelatives had partial or major limitation of activity. Approximately this same proportion were limited in their activity among persons in the category—married, living with relatives. However, 48 percent of those in the category—other marital status, living with relatives—had some degree of activity limitation.

Table C. Percent of persons with partial or major limitation of activity or mobility by family relationship and age: United States, July 1957-June 1959

Family relationship	Age						
· · · · · · · · · · · · · · · · · · ·	45-54	55-64	65-74	75+			
	Li	imitation	of activity				
Living alone or with nonrelatives Living with relatives Married Other status	13.9 12.6 12.1 17.0	23.2 21.1 20.7 23.3		51.3 56.6 54.2 58.9			
	Li	imitation	of mobility				
Living alone or with nonrelatives Living with relatives Married Other status	3.3 2.1 1.9 4.2	6.1 5.7 5.5 7.4	10.3 12.2 11.0 15.7	24.9 32.5 26.3 38.3			

This latter group includes, for the most part, widowed parents living with children, and other unmarried persons living with relatives. Corresponding percentages in relation to limitation of mobility were: 16 percent of those living alone or with nonrelatives, 15 percent of those married, living with relatives, and 27 percent of those in the group—other marital status, living with relatives. Comparable percentages showing more age detail are presented in table C.

Disability Days

The concept of disability due to illness or injury expressed in terms of days differs from the concept of activity and mobility limitation in that the former presents a quantitative rather than a qualitative measure of disability. Moreover, disability days may result from acute as well as chronic conditions.

A day of restricted activity is defined in the survey as a day when a person cut down on his usual activities for the whole of that day on account of illness or injury. A day lost from work or a day spent in bed because of illness is, of course, a day of restricted activity. A day of bed disability is defined as a day on which a person spent all or most of the day in bed because of an illness or an injury. A day spent in a hospital is considered to be a day of bed disability even if the person was not actually in bed at the hospital. A day is counted as lost from work if the person would have been going to work at a job or business on that day but instead lost the entire time that he would have worked on that day because of an illness or an injury.

It is possible for a day of disability to have resulted from more than one condition. In describing the characteristics of conditions, the day is ascribed in the statistics to each condition that was reported to have caused the disability. However, in tabulating disability of persons, the day is counted only once as a day of disability for the person involved. Only person-days of disability are shown in this section of the report.

1. An average annual total of 625 million restricted-activity days and approximately 209 million bed-disability days were estimated for persons 65 years and older. These figures represent an average of 42,6 restricted-activity




Figure 2. Restricted-activity days per person per year for persons 45 years and older by age and

days per person per year of which 14.2 were days of bed disability. In general, the rate of disability in terms of restricted activity and bed disability was higher for females than for males 65 years and older (fig. 2).

2. The average number of restricted-activity and bed-disability days per year per person 65 years and older showed a marked increase as the amount of family income declined. The causal relationship of incapacitating illness and low income is a complex subject beyond the scope of this report (fig. 3).

3. Persons 65 years and older who reported their major activity as "usually working" had on the average 23.3 days of restricted activity, of which 5.2 days involved bed disability. Persons 65 years and older whose major activity was keeping house had on the average 39.9 restricted-activity days with 11.0 days of bed disability. Since the working group in this age category is heavily weighted by males, and those keeping house are females, the disability data reflect a sex differential. As would be expected the disability rates were considerably higher for re-




Figure 3. Disability days per person per year for persons 65 years and older by family income.

tired persons and for those in the "other" category (tables 18 and 19), which in this older age




Figure 4. Disability days per person per year for persons 65 years and older by major activity.

group consisted mainly of persons who were unable to work but did not consider themselves as "retired" (fig. 4).

The rate of work-loss days among "usually working" persons 65 years and older showed a pattern similar to that for other disability days in relation to family income. The highest rate of work loss, 13.6 days per "usually working" person per year, was among persons with known family income less than \$2,000, and the lowest rate, 8.5 days, for persons with known family income \$7,000 or more. The total workloss rate for persons 65 years and older is shown in table 20. Table 20 differs from table 17, which shows work loss only for "usually working" persons, in that it includes work loss for persons whose major activity status was keeping house, retired, or other, but who worked at some time during the past year.

DETAILED TABLES

		Page
	LONG-TERM DISABILITY	
Table 1.	Average number of persons 45 years and older according to chronic limitation of activity by age and residence: United States, July 1957-June 1959	10
2.	Percent distribution of persons 45 years and older according to chronic limitation of activity by age and residence: United States, July 1957-June 1959	11
3.	Average number of persons 45 years and older according to chronic limitation of mobility by age and residence: United States, July 1957-June 1959	12
4.	Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and residence: United States, July 1957-June 1959	13
5.	Average number of persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959	14
6.	Percent distribution of persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959-	15
7.	Average number of persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959	16
8.	Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959-	17
9.	Average number of persons 45 years and older according to chronic limitation of activity by age and major activity: United States, July 1957-June 1959	18
10.	Percent distribution of persons 45 years and older according to chronic limitation of activity by age and major activity: United States, July 1957-June 1959	19
11.	Average number of persons 45 years and older according to chronic limitation of mobility by age and major activity: United States, July 1957-June 1959	20
12.	Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and major activity: United States, July 1957-June 1959	21
13.	Average number and percent distribution of persons 65 years and older according to chronic limitation of activity by sex and family relationship: United States, July 1957-June 1959	22
14.	Average number and percent distribution of persons 65 years and older according to chronic limitation of mobility by sex and family relationship: United States, July 1957-June 1959	23
	DISABILITY DAYS	
15.	Average annual number of restricted-activity days and number of restricted-activity days per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959	24
16.	Average annual number of bed-disability days and number of bed-disability days per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959	25

DETAILED TABLES-Continued

		Page
	DISABILITY DAYS—Continued	
Table 17	Average annual number of work-loss days for "usually working" persons and number of work-loss days per "usually working" person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959	26
18	Average annual number of restricted-activity days and number of restricted-activity days per person per year for persons 45 years and older by major activity, sex, and age: United States, July 1957-June 1959	27
19	Average annual number of bed-disability days, and bed-disability days per person per year for persons 45 years and older by major activity, sex, and age: United States, July 1957-June 1959	28
20	Average annual number of work-loss days and number of work-loss days per person per year for persons 45 years and older by major activity, sex, and age:	. 20

Table 1. Average number of persons 45 years and older according to chronic limitation of activity by age and residence: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11]

		Persons	Per		h l+ chro itions	nic
Age and residence	Total persons	with no chronic		Limita	tion of a	ctivity
	persons	conditions	Total		Partial	T .
				None	Partial	Major
•	A	verage numbe	r of per	sons in	thousands	;
All ages-45+	49,429	17,282	32,147	20,235	8,868	3,04
Jrban, total	32,311	12,062	20,249	13,132	5,191	1,92
Inside urbanized areas	23,415	9,150	14,265	9,426	3,504	1,33
Other urban places	8,895	2,911	5,984	3,706	1,686	59
Rural, total	17,118	5,221	11,898	7,102	3,678	1,11
Nonfarm	11,023	3,512	7,510	4,563	2,206	74
Farm	6,096	1,708	4,388	2,539	1,472	37
45-54	19,833	8,634	11,199	8,679	2,211	30
Urban, total	12,944	5,887	7,057	5,556	1,308	19
Inside urbanized areas	9.580	4,469	5,111	4,052	921	13
Other urban places	3,364	1,418	1,947	1,503	386	5
Rural, total	6,888	2,747	4,142	3,124	903	11
Nonfarm	4,501	1,862	2,640	2,021	534	8
Farm	2,387	885	1,502	1,102	368	3
55-6 <u>4</u>	14,930	5,320	9,610	6,419	2,556	63
Urban, total	9,910	3,811	6,098	4,184	1,498	41
Inside urbanized areas	7,303	2,927	4,376	3,037	1,032	30
Other urban places	2,607	884	1,722	1,147	466	111
Rural, total	5,020	1,509	3,511	2,235	1,058	21
Nonfarm	3,020	998	2,175	1,395	637	14
Farm	1,847	511	1,336	840	421	7
65+	14,667	3,328	11,338	5,137	4,102	2,09
Urban, total	9,457	2,363	7,093	3,393	2,385	1,31
Inside urbanized areas	6,532	1,754	4,778	2,337	1,550	89
Other urban places	2,924	609	2,315	1,057	834	42
Rural, total	5,210	965	4,245	1,743	1,717	78
Nonfarm	3,348	653	2,695	1,147	1,035	51
Farm	1,862	312	1,550	597	683	27
65-74	9,698	2,500	7,197	3,739	2,545	91
Jrban, total	6,330	1,765	4,564	2,484	1,497	58
Inside urbanized areas	4,451	1,306	3,145	1,733	998	41
Other urban places	1,878	459	1,419	751	500	16
Rural, total	3,368	735	2,633	1,255	1,048	32
Nonfarm	2,155	496	1,659	808	627	22
Farm	1,213	239	973	447	421	10
75+	4,969	828	4,141	1,398	1,557	1,18
Jrban, total	3,127	598	2,529	910	888	73
Inside urbanized areas	2,081	448	1,633	603	553	47
Other urban places	1,046	151	895	306	335	25
		230	1,612	488	669	45
Rural, total	1,842	230	,	700	1 007	
Rural, total	1,042	157	1,036	338	408	29

Table 2. Percent distribution of persons 45 years and older according to chronic limitation of activity by age and residence: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

		Persons with no	Persons with 1+ chronic conditions				
Age and residence	Total persons	chronic		Limita	tion of a	ctivity	
		conditions	Total	None	Partial	Major	
* **		Pero	ent dist	ribution	<u> </u>	<u> </u>	
		1610	ent dist	TIDUCTON	•		
All ages-45+	100.0	35.0	65.0	40.9	17.9	6.2	
Urban, total	100.0	37.3	62.7	40.6	16.1	6.0	
Inside urbanized areas	100.0	39.1	60.9	40.3	15.0	5.7	
Other urban places	100.0	32.7	67.3	41.7	19.0	6.6	
Rural, total	100.0	30.5	69.5	41.5	21.5	6.5	
Nonfarm	100.0	31.9	68.1	41.4	20.0	6.7	
Farm	100.0	28.0	72.0	41.7	24.1	6.2	
45-54	100.0	43.5	56.5	43.8	11.1	1.6	
Urban, total	100.0	45.5	54.5	42.9	10.1	1.5	
Inside urbanized areas	100.0	46.6	53.4	42.3	9.6	1.4	
Other urban places	100.0	42.2	57.9	44.7	11.5	1.7	
Rural, total	100.0	39.9	60.1	45.4	13.1	1.7	
Nonfarm	100.0	41.4	58.7	44.9	11.9	1.9	
Farm	100.0	37.1	62.9	46.2	15.4	1.3	
55-64	100.0	35.6	64.4	43.0	17.1	4.3	
Urban, total	100.0	38.5	61.5	42.2	15.1	4.2	
Inside urbanized areas	100.0	40.1	59.9	41.6	14.1	4.2	
Other urban places	100.0	33.9	66.1	44.0	17.9	4.2	
Rural, total	100.0	30.1	69.9	44.5	21.1	4.3	
Nonfarm	100.0	31.5	68.5	44.0	20.1	4.5	
Farm	100.0	27.7	72.3	45.5	22.8	4.1	
65+	100.0	22.7	77.3	35.0	28.0	14.3	
Urban, total	100.0	25.0	75.0	35.9	25.2	13.9	
Inside urbanized areas	100.0	26.9	73.1	35.8	23.7	13.6	
Other urban places	100.0	20.8	79.2	36.1	28.5	14.5	
Rural, total	100.0	18.5	81.5	33.5	33.0	15.0	
Nonfarm	100.0	19.5	80.5	34.3	30.9	15.4	
Farm	100.0	16.8	83.2	32.1	36.7	14.5	
65-74	100.0	25.8	74.2	38.6	26.2	9.4	
Urban, total	100.0	27.9	72.1	39.2	23.6	9.2	
Inside urbanized areas	100.0	29.3	70.7	38.9	22.4	9.3	
Other urban places	100.0	24.4	75.6	40.0	26.6	9.0	
Rural, total	100.0	21.8	78.2	37.3	31.1	9.8	
Nonfarm	100.0	23.0	77.0	37.5	29.1	10.4	
NonfarmFarm	100.0	19.7	80.2	36.9	34.7	8.7	
75+	100.0	16.7	83.3	28.1	31.3	23.9	
Urban, total	100.0	19.1	80.9	29.1	28.4	23.4	
Inside urbanized areas	100.0	21.5	78.5	29.0	26.6	22.9	
Other urban places	1 100 A	1. 14 41	0,7 0,1		1 3/11	J 74.3	
Other urban places	100.0	14.4 12.5	85.6 87.5	29.3 26.5	32.0 36.3	24.3 24.7	
Other urban places	100.0 100.0 100.0	14.4 12.5 13.2	87.5 86.8	26.5 28.3	36.3 34.2	24.3 24.7 24.3	

Table 3. Average number of persons 45 years and older according to chronic limitation of mobility by age and residence: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix [1]

Age and residence	Total	Persons with no	Per	cond	th 1+ chro		
and legities	persons	chronic	m-4-1	Limita	tion of m	obility	
		conditions	Total	None	Partial	Major	
	Average number of persons in thousands						
All ages-45+	49,429	17,282	32,147	28,186	3,043	918	
Urban, total	32,311	12,062	20,249	17,815	1,842	592	
Inside urbanized areas	23,415	9,150	14,265	12,601	1,251	414	
Other urban places	8,895	2,911	5,984	5,215	591	178	
Rural, total	17,118	5,221	11,898	10,370	1,201	326	
. Nonfarm	11,023	3,512	7,510	6,544	758	208	
Farm	6,096	1,708	4,388	3,826	443	119	
45-54	19,833	8,634	11,199	10,762	351	86	
Urban, total	12,944	5,887	7,057	6,787	210	60	
Inside urbanized areas	9,580	4,469	5,111	4,915	155	41	
Other urban places	3,364	1,418	1,947	1,872	55	(*)	
Rural, total	6,888	2,747	4,142	3,975	141	(*)	
Nonfarm	4,501	1,862	2,640	2,533	- 89	(*)	
Farm	2,387	885	1,502	1,442	52	(*)	
55-64	14,930	5,320	9,610	8,744	675	191	
Urban, total	9,910	3,811	6,098	5,557	420	122	
. Inside urbanized areas	7,303	2,927	4,376	3,985	300	91	
Other urban places	2,607	884	1,722	1,572	120	30	
Rural, total	5,020	1,509	3,511	3,187	255	69	
Nonfarm	3,173	998	2,175	1,968	165	42	
Farm	1,847	511	1,336	1,219	90	(*)	
65+	14,667	3,328	11,338	8,679	2,017	641	
Urban, total	9,457	2,363	7,093	5,471	1,212	410	
Inside urbanized areas	6,532	1,754	4,778	3,701	797	281	
Other urban places	2,924	609	2,315	1,770	416	129	
Rural, total	5,210	965	4,245	3,208	805	232	
Nonfarm	3,348	653	2,695	2,043	505	147	
Farm	1,862	312	1,550	1,165	300	85	
65-75	9,698	2,500	7,197	6,053	924	219	
llukan tatal	6 330		1. 561	2 0/0	5.67	1.0	
Urban, total Inside urbanized areas	6,330	1,765	4,564	3,849	567	148	
Other urban places	4,451 1,878	1,306 459	3,145	2,652	387 180	106	
Rural, total	3,368	735	1,419 2,633	1,197 2,205	357	42	
Nonfarm	2,155	496	1,659	1,382	229	71 48	
Farm	1,213	239	973	823	128	· (*)	
75+	4,969	828	4,141	2,626	1,093	422	
				,	,		
Urban, total	3,127	598	2,529	1,622	645	262	
Inside urbanized areas	2,081	448	1,633	1,049	409	175	
Other urban places	1,046	151	895	573	236	87	
Rural, total	1,842	230	1,612	1,004	448	161	
NonfarmFarm	1,193	157	1,036	661	275	99	
× 97ma=+	1 649 1	73]	576	342	172	62	

Table 4. Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and residence: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

	Tet-1	Persons	Per		h 1+ chro	onic
Age and residence	Total persons	with no chronic	·		tion of n	obility
		conditions	Total	l	T	
	ļ	<u> </u>	<u> </u>	None	Partial	Major
		Perc	ent dist	ribution	ı	
All ages-45+	100.0	35.0	65.0	57.0	6.2	1.9
Urban, total	100.0	37.3	62.7	55.1	5.7	1.8
Inside urbanized areas	100.0	39.1	60.9	53.8	5.3	1.8
Other urban placesRural, total	100.0	32.7	67.3	58.6	6.6	2.0
Nonfarm	100.0	30.5 31.9	69.5	60.6	7.0	1.9
Farm	100.0	28.0	68.1 72.0	59.4 62.8	7.3	1.9
raim	100.0	20.0	/2.0	02.0	/.3	2.0
45-54	100.0	43.5	56.5	54.3	1.8	0.4
Urban, total	100.0	45.5	54.5	52.4	1.6	0.5
Inside urbanized areas	100.0	46.6	53.4	51.3	1.6	0.4
Other urban places	100.0	42.2	57.9	55.6	1.6	(*)
Rural, total	100.0	39.9	60.1	57.7	2.0	(*)
Nonfarm	100.0	41.4	58.7	56.3	2.0	(*)
Farm	100.0	37.1	62.9	60.4	2.2	(*)
55-64	100.0	35.6	64.4	58.6	4.5	1.3
Urban, total	100.0	20.5	(1.5	56.1	, ,	١.,
Inside urbanized areas	100.0	38.5	61.5	56.1	4.2	1.2
Other urban places	100.0	40.1 33.9	59.9	54.6	4.1	1.2
Rural, total	100.0	30.1	66.1 69.9	60.3	4.6	1.2
Nonfarm	100.0	31.5	68.5	63.5 62.0	5.1	1.4 1.3
Farm	100.0	27.7	72.3	66.0	4.9	(*)
65+	100.0	22.7	77.3	59.2	13.8	4.4
Urban, total	100.0	25.0	75.0	57.9	12.8	4.3
Inside urbanized areas	100.0	26.9	73.1	56.7	12.2	4.3
Other urban places	100.0	20.8	79.2	60.5	14.2	4.4
Rural, totalNonfarm	100.0	18.5	81.5	61.6	15.5	4.5
Farm	100.0	19.5 16.8	80.5 83.2	61.0 62.6	15.1 16.1	4.4
· · · · · · · · · · · · · · · · · · ·	; 100.0	10.0	05.2	.02.0	10.1	4.6
65-74	100.0	25.8	74.2	62.4	9.5	2.3
Urban, total	100.0	27.9	72.1	60.8	9.0	2.3
Inside urbanized areas	100.0	29.3	70.7	59.6	8.7	2.4
Other urban places	100.0	24.4	75.6	63.7	9.6	2.2
Rural, total	100.0	21.8	78.2	65.5	10.6	2.1
Nonfarm	, 100.0	23.0	77.0	64.1	10.6	2.2
Farm	100.0	19.7	80.2	67.8	10.6	(*)
75+	100.0	16.7	83.3	52.8	22.0	8.5
Urban, total	100.0	19.1	80.9	51.9	20.6	8.4
Inside urbanized areas	100.0	21.5	78.5	50.4	19.7	8.4
Other urban places	100.0	14.4	85.6	54.8	22.6	8.3
Rural, total	100.0	12.5	87.5	54.5	24.3	8.7
NonfarmFarm	100.0	13.2	86.8	55.4	23.1	8.3
Farm	100.0	11.2	88.8	52.7	26.5	9.6
——————————————————————————————————————						L

Table 5. Average number of persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1]

	Persons		Per	Persons with 1+ chronic conditions			
Age and family income	Total persons	with no chronic		Limita	tion of a	ctivity	
	1	conditions	Total	None	Partial	Major	
	Ä	verage numbe	r of per	sons in	thousands		
A11 ages-45+			1 1	1 .;	1	1.	
Total	49,429	17,282	32,147	20,235	8,868	3,044	
Under \$2,000	11,394	2,726	8,668	4,174	3,170	1,324	
\$2,000-3,999	10,713	3,598	7,115	4,406	2,059	650	
\$4,000-6,999	13,218	5,179	8,039	5,689	1,833	518	
\$7,000+	9,666	3,936	5,730	4,321	1,141	268	
Unknown	4,438	1,843	2,595	1,646	666	283	
<u>45-54</u>)))			
Total	19,833	8,634	11,199	8,679	2,211	309	
Under \$2,000	2,452	882	1,569	975	484	110	
\$2,000-3,999	3,886	1,611	2,275	1,665	533	78	
\$4,000-6,999	6,662	2,977	$\frac{3}{686}$	2,993	632	60	
Unknown	5,330 1,503	2,411 753	2,919 749	2,458 588	428 134	33 28	
Utiktiowti	1,505	/33	/43	J00.	154	1 2°	
<u>55-64</u>					ĺ		
Total	14,930	5,320	9,610	6,419	2,556	635	
Under \$2,000\$2,000-3,999	3,004	784	2,221	1,180	783	257	
\$2,000-3,999	3,471	1,187	2,284	1,499	624	161	
\$4,000-6,999	4,144	1,575	2,568	1,850	599	119	
\$7,000+	2,804	1,123	1,681	1,306	335	. 40	
Unknown	1,507	651	856	583	214	59	
<u>65+</u>	1	:	}			·	
Total	14,667	3,328	11,338	5,137	4,102	2,099	
Under \$2,000	5,938	1,060	4,878	2,018	1,902	958	
\$2,000-3,999	3,356	801	2,555	1,242	902	411	
\$4,000-6,999 \$7,000+	2,412	627 402	1,785	845	601	339	
Unknown	1,532	439	1,130	557 475	378 319	195 196	
	1,120	1		""		•	
65-74	0.600	2 500	7 107	2 720	0.5/6	012	
Total	9,698	2,500	7,197	3,739	2,545	913	
Under \$2,000	3,656	764	2,893	1,375	1,101	416	
\$2,000-3,999	2,463	639	1,824	995	623	206	
\$4,000-6,999	1,643	472 300	1,171 724	636 412	230	134	
\$7,000+	912	326	586	321	190	83 74	
7 <u>5+</u>		}				' '	
751. Total	4,969	828	4,141	1,398	1,557	1,186	
Under \$2,000		 	+	1	801	T	
\$2,000-3,999	2,281	296 162	1,985 732	643	279	341 206	
\$4,000-6,999	769	155	614	209	200	205	
\$7.000+	508	102	406	145	148	113	
Unknown	517	113	404	154	128	122	
UIIAIIUWII	1 31	1 113	107	1 -2-	1-0	1	

Table 6. Percent distribution of persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	1	Persons	Per		h 1+ chro	nic	
	Total	with no	conditions				
Age and family income	persons	chronic	,	Limita	tion of a	ctivity	
	persons	conditions	Total	None	Partial	Major	
		L			1	13	
<u>All ages-45+</u>		Pero	ent distribution				
Total	100.0	35.0	65.0	40.9	17.9	6.2	
Under \$2,000\$2,000-3,999	100.0	23.9	76.1	36.6	27.8	11.6	
\$2.000-3.999	100.0	33.6	66.4	41.1	19.2	6.1	
\$4.000-6.999	100.0	39.2	60.8	43.0	13.9	3.9	
\$7,000+	100.0	40.7	59.3	44.7	11.8	2.8	
\$4,000-6,999 \$7,000+ Unknown	100.0	41.5	58.5	37.1	15.0	6.4	
<u>45-54</u>				·			
Total	100.0	43.5	56.5	43.8	11.1	1.6	
Inder \$2,000	100.0	36.0	64.0	39.8	19.7	4.5	
\$2,000-3,999	100.0	41.5	58.5	42.8	13.7	2.0	
34,000-6,999	100.0	44.7	55.3	44.9	9.5	0.9	
7,000+	100.0	45.2	54.8	46.1	8.0	0.6	
Inknown	100.0	50.1	49.8	39.1	8.9	1.9	
	100.0	50.12	'''	37.1	"		
<u>55-64</u> Total	100.0	25.6	64.4	42.0	17.] , .	
	100.0	35.6	64.4	43.0	17.1	4.:	
Inder \$2,000	100.0	26.1	73.9	39.3	26.1	8.6	
22,000-3,999	100.0	34.2	65.8	43.2	18.0	4.6	
4,000-6,999	100.0	38.0	62.0	44.6	14.5	2.9	
7,000+	100.0	40.0	60.0	46.6	11.9	1.4	
Inknown	100.0	43.2	56.8	38.7	14.2	3.9	
<u>65+</u>	•						
Total	100.0	22.7	77.3	35.0	28.0	14.3	
Inder \$2,000	100.0	17.9	82.1	34.0	32.0	16.1	
\$2,000-3,999	100.0	23.9	76.1	37.0	26.9	12.	
34,000-6,999	100.0	26.0	74.0	35.0	24.9	14.	
67.000 +	100.0	26.2	73.8	36.4	24.7	12.7	
Inknown	100.0	30.7	69.3	33.3	22.3	13.7	
<u>65-74</u>					ļ		
Total	100.0	25.8	74.2	38.6	26.2	9.4	
Under \$2,000	100.0	20.9	79.1	37.6	30.1	11.4	
2,000-3,999	100.0	25.9	74.1	40.4	25.3	8.4	
4,000-6,999	100.0	28.7	71.3	38.7	24.4	8.	
67,000+	100.0	29.3	70.7	40.2	22.5	8.	
Jnknown	100.0	35.7	64.3	35.2	20.9	8.1	
				*			
<u>75+</u>	100.0		00.5	00.5			
Total	100.0	16.7	83.3	28.1	31.3	23.9	
Jnder \$2,000	100.0	13.0	87.0	28.2	35.1	23.	
\$2,000-3,999	100.0	18.1	82.0	27.7	31.2	23.1	
\$4,000-6,999	100.0	20.2	79.8	27.2	26.0	26.7	
\$7.000+	100.0	20.1	79.9	28.5	29.1	22.2	
Unknown	100.0	21.9	78.1	29.8	24.8	23.6	
•	ì	i	l I		ł	1	

Table 7. Average number of persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11]

	Total	Persons with no	Persons with 1+ chronic conditions				
Age and family income	persons	chronic conditions	Total	Limitation of mobility			
	<u> </u>	<u> </u>	<u> </u>	None	Partial	Major	
All ages-45+	A	verage numbe	er of per	sons in	thousands	3	
Total	49,429	17,282	32,147	28,186	3,043	918	
Under \$2,000	11,394	2,726	8,668	6,949	1,346	374	
\$2,000-3,999	10,713	3,598	7,115	6,307	620	187	
\$4,000-6,999	13,218	5,179	8,039	7,336	536	167	
\$7,000+ Unknown	9,666	3,936	5,730	5,338	304	88	
	4,438	1,843	2,595	2,255	238	102	
<u>45-54</u>							
Total	19,833	8,634	11,199	10,762	351	86	
Under \$2,000	2,452	882	1,569	1,423	120	26	
\$2,000-3,999	3,886	1,611	2,275	2,178	79	(*)	
\$4,000-6,999	6,662	2,977	3,686	3,572	89	24	
\$7,000+	5,330	2,411	2,919	2,863	45 (*)	(*)	
· ·	1,503	753	749	725	(*)	(*)	
<u>55-64</u>			,				
Total	14,930	5,320	9,610	8,744	675	191	
Under \$2,000	3,004	784	2,221	1,877	266	77	
\$2,000-3,999 \$4,000-6,999	3,471	1,187	2,284	2,080	157	47	
\$7,000+	4,144 2,804	1,575 1,123	2,568 1,681	2,398 1,606	135 67	35 (*)	
Unknown	1,507	651	856	782	50	24	
<u>65+</u>							
Total	14,667	3,328	11,338	8,679	2,017	641	
Under \$2,000	5,938	1,060	4,878	3,648	959	270	
\$2,000-3,999	3,356	801	2,555	2,049	385	122	
\$4,000-6,999	2,412	627	1,785	1,366	312	107	
\$7,000+	1,532	402	1,130	869	192	69	
Unknown	1,428	439	990	748	169	73	
<u>65-75</u>			÷				
Total	9,698	2,500	7,197	6,053	924	219	
Under \$2,000	3,656	764	2,893	2,377	428	88	
\$2,000-3,999	2,463	639	1,824	1,570	203	51	
\$4,000-6,999	1,643	472	1,171	998	134	39	
\$7,000+	1,024 912	300 326	724 586	616 493	88 71	20	
	312	340	586	493	71	22	
<u>75+</u>	1			l			
Total	4,969	828	4,141	2,626	1,093	422	
Under \$2,000	2,281	296	1,985	1,271	531	183	
\$2,000-3,999	893	162	732	478	182	7 1	
\$4,000-6,999	769	155	614	367	178	69	
\$4,000-6,999 \$7,000+	508 517	102	406 404	253 255	104	49	
OHERIOWIT	517	113	404	(,,	98	51	

Table 8. Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix [1]

			£=				
	Total	Persons with no	Persons with 1+ chronic conditions				
Age and family income	persons	chronic		Limitation of mobility			
	1	conditions	Total	None	Partial	Major	
	Percent distribution						
All ages-45+	1				1	•	
Total	100.0	35.0	65.0	57.0	6.2	1.9	
Under \$2,000	100.0	23.9	76.1	61.0	11.8	3.3	
\$2,000-3,999	100.0	33.6	66.4	58.9	5.8	1.7	
\$4,000-6,999	100.0	39.2	60.8	55.5	4.1	1.3	
\$7,000+	100.0	40.7	59.3	55.2	3.1	0.9	
\$7,000+	100.0	41.5	58.5	50.8	5.4	2.3	
<u>45-54</u>	,						
Total	100.0	43.5	56.5	54.3	1.8	0.4	
Under \$2,000	100.0	36.0	64.0	58.0	4.9	1.1	
\$2,000-3,999	100.0	41.5	58.5	56.0	2.0	(*)	
\$4,000-6,999	100.0	44.7	55.3	53.6	1.3	0.4	
\$7,000+	100.0	45.2	54.8	53.7	0.8	(*)	
Unknown	100.0	50.1	49.8	48.2	(*)	(*)	
<u>55-64</u>						l	
Total	100.0	35.6	64.4	58.6	4.5	1,3	
Under \$2,000	100.0	26.1	73.9	62.5	8.9	2.6	
\$2,000-3,999	100.0	34.2	65.8	59.9	4.5	1.4	
\$4,000-6,999	100.0	38.0	62.0	57.9	3.3	0.8	
\$7,000+	100.0	40.0	60.0	57.3	2.4	(*)	
•	100.0	43.2	56.8	51.9	3.3	1.6	
<u>65+</u>	1						
Total	100.0	22.7	77.3	59,2	13.8	4.4	
Under \$2,000	100.0	17.9	82.1	61.4	16.2	4.5	
\$2,000-3,999	100.0	23.9	76.1	61.1	11.5	3.6	
\$4,000-6,999	100.0	26.0	74.0	56.6	12.9	4.4	
\$7,000+	100.0	26.2 30.7	73.8 69.3	56.7 52.4	12.5	4.5 5.1	
· · · · · · · · · · · · · · · · · · ·	, 100,.0	30.7	07.5	32.4	11.0	3.1	
<u>65-74</u>]		
Total	100.0	25.8	74.2	62.4	9.5	2,3	
Under \$2,000\$2,000-3,999	100.0	20.9	79.1	65.0	11.7	2.4	
\$2,000-3,999	100.0	25.9	74.1	63.7	8.2	2.1	
\$4,000-6,999 \$7,000+	100.0	28.7	71.3	60.7	8.2	2.4	
Unknown	100.0	29.3	70.7	60.2	8.6	2.0	
· ·	100.0	35.7	64.3	54.1	7.8	2.4	
<u>75+</u> Total]		0.5			. .	
	100.0	16.7	83.3	52.8	22.0	8.5	
Under \$2,000	100.0	13.0	87.0	55.7	23.3	8.0	
\$2,000-3,999 \$4,000-6,999	100.0	18.1	82.0	53.5	20.4	8.0	
\$7,000+	100.0	20.2	79.8 79.9	47.7 49.8	23.1 20.5	9.0 9.6	
Unknown	100.0	20.1 21.9	78.1	49.8	19.0	9.9	
			, 0.1				

Table 9. Average number of persons 45 years and older according to chronic limitation of activity by age and major activity: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	Total	Persons with no	Per		h l+ chro	onic		
Age and major activity	persons	chronic conditions	Total	Limita	tion of a	ctivity		
	ļ	1	l	None	Partial	Major		
<u>All ages-45+</u>	Average number of persons in thousands							
All persons	49,429	17,282	32,147	20,235	8,868	3,044		
Usually working Keeping house Retired Other	24,163 16,863 6,134 2,269	10,613 5,044 1,189 436	13,550 11,819 4,945 1,833	10,668 7,533 1,509 525	2,654 4,038 1,574 602	228 248 1,861 706		
45-54								
All persons	19,833	8,634	11,199	8,679	2,211	309		
Usually working	13,083 5,907 83 759	6,237 2,188 (*) 193	6,846 3,719 68 566	5,692 2,767 (*) 211	1,069 915 (*) 203	85 38 35 152		
All persons	14,930	5,320	9,610	6,419	2,556	635		
Usually working Keeping house Retired Other	8,369 5,134 678 750	3,457 1,593 119 151	4,911 3,542 558 599	3,799 2,326 122 171	1,024 1,156 160 216	88 60 276 211		
<u>65+</u>								
All persons	14,667	3,328	11,338	5,137	4,102	2,099		
Usually working Keeping house Retired Other	2,711 5,822 5,373 760	918 1,264 1,055 92	1,793 4,558 4,318 669	1,177 2,440 1,377 143	561 1,967 1,391 183	55 151 1,550 343		
<u>65-74</u>	}							
All persons	9,698	2,500	7,197	3,739	2,545	913		
Usually working Keeping house Retired Other	2,312 4,014 2,973 398	810 961 670 60	1,502 3,052 2,304 338	1,009 1,783 854 94	456 1,198 784 108	38 71 666 137		
<u>75+</u>				-				
All persons	4,969	828	4,141	1,398	1,557	1,186		
Usually working Keeping house Retired Other	399 1,808 2,400 362	109 303 385 (*)	291 1,505 2,015 330	168 657 523 49	105 769 608 76	(*) 79 884 205		

Table 10. Percent distribution of persons 45 years and older according to chronic limitation of activity by age and major activity: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

	Total	Persons with no	Per		h l+ chro itions	nic
Age and major activity	persons	chronic	,	Limita	tion of a	ctivity
		conditions	Total	None	Partial	Major
		Perc	ent dist	ribution		
All ages-45+			ent arst	LIDUCION	·	
All persons	100.0	35.0	65.0	40.9	17.9	6.2
Usually working	100.0	,43.9	56.1	44.2	11.0	0.9
Keeping house	100.0	29.9	70.1	44.7	23.9	1.5
RetiredOther	100.0	19.4 19.2	80.6 80.8	24.6 23.1	25.7	30.3 31.1
Uther	100.0	19.2		23.1	26.5	. 31.1
<u>45-54</u>			[.			
All persons	100.0	43.5	56.5	43.8	11.1	1.6
Usually working	100.0	47.7	52.3	43.5	8.2	0.6
Keeping house	100.0	37.0	63.0	46.8	15.5	0.6
Retired	100.0	(*)	81.9	(*)	(*)	42.2
Other	100.0	25.4	74.6	27.8	26.7	20.0
<u>55-64</u>						} [
All persons	100.0	35.6	64.4	43.0	17.1	4.3
Usually working	100.0	41.3	58.7	45.4	12.2	1.1
	100.0	31.0	69.0	45.3	22.5	1.2
Retired	100.0	17.6	82.3	18.0	23.6	40.7
Other	100.0	20.1	79.9	22.8	28.8	28.1
<u>65+</u>					i	
All persons	100.0	22.7	77.3	35.0	28.0	14.3
Usually working	100.0	33.9	66.1	43.4	20.7	2.0
Keeping house	100.0	21.7	78.3	41.9	33.8	2.6
Retired	100.0	19.6	80.4	25.6	25.9	28.8
Other	100.0	12.1	88.0	18.8	24.1	45.1
<u>65-74</u>		,			į	
All persons	100.0	25.8	74.2	38.6	26.2	9.4
Usually working	100.0	35.0	65.0	42.6	19.7	1
Keeping house	100.0	35.0 23.9	65.0 76.0	43.6 44.4	29.8	1.6
Retired	100.0	22.5	77.5	28.7	26.4	
Other	100.0	15.1	84.9	23.6	27.1	34.4
	1					
<u>75+</u>				00.5		
All persons	100.0	16.7	83.3	28.1	31.3	23.9
Usually working	100.0	27.3	72.9	42.1	26.3	(*)
Keeping house	100.0	16.8	83.2	36.3	42.5	1
RetiredOther	100.0	16.0	84.0	21.8	25.3	36.8 56.
Utner	100.0	(*)	91.2	13.5	21.0	30.

Table 11. Average number of persons 45 years and older according to chronic limitation of mobility by age and major activity: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11]

	<u> </u>		Persons with 1+ chronic				
Age and major activity	Total	Persons with no	conditions				
	persons	chronic	Total	Limita	obility		
		conditions	local	None	Partial	Major	
All ages-45+	Av	erage number	of pers	ons in t	housands		
All persons	49,429	17,282	32,147	28,186	3,043	918	
Usually working	24,163	10,613	13,550	13,184	310	57.	
Keeping house	16,863	5,044	11,819	10,461	1,174	184	
Retired	6,134	1,189	4,945	3,372	1,153	420	
Other	2,269	436	1,833	1,170	406	257	
<u>45-54</u>				·			
All persons	19,833	8,634	11,199	10,762	351	86	
Usually working	13,083	6,237	6,846	6,722	104	(*)	
Keeping house	5,907	2,188	3,719	3,559	133	(*)	
Retired	83	(*)	68	39	. 26	(*)	
Other	759	193	566	442	88	35	
<u>55-64</u>		·					
All persons	14,930	5,320	9,610	8,744	675	191	
Usually working	8,369	3,457	4,911	4,773	113	(*)	
Keeping house	5,134	1,593	3,542	3,212	282	47	
Retired	678	119	558	345	160	52	
Other	750	151	599	413	119	67	
<u>65+</u>	<u>.</u>						
All persons	14,667	3,328	11,338	8,679	2,017	641	
Usually working	2,711	918	1,793	1,688	92	· (*)	
Keeping house	5,822	1,264	4,558	3,689	759	ìı́o	
Retired	5,373	1,055	4,318	2,987	967	364	
Other	760	92	669	314	199	155	
<u>65-74</u>							
All persons	9,698	2,500	7,197	6,053	924	219	
Usually working	2,312	810	1,502	1,435	61	(*)	
Keeping house	4,014	961	3,052	2,624	381	48	
Retired	2,973	670	2,304	1,791	396	117	
Other	398	60	338	203	87	49	
<u>75+</u>						}	
All persons	4,969	828	4,141	2,626	1,093	422	
Usually working	399	109	291	253	31	(*)	
Keeping house	1,808	303	1,505	1,065	379	62	
RetiredOther	2,400 362	385 32	2,015 330	1,196 112	571 112	247 107	

Table 12. Percent distribution of persons 45 years and older according to chronic limitation of mobility by age and major activity: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

	Total	Persons with no	Persons with 1+ chronic conditions				
Age and major activity	persons	chronic conditions	Total	Limita	tion of m	obility	
		Conditions		None	Partial	Major	
	Percent distribution						
All ages-45+				1		1	
All persons	100.0	35.0	65.0	57.0	6.2	1.9	
Usually working	100.0	43.9	56.1	54.6	1.3	0.2	
Keeping house	100.0	29.9	70.1	62.0	7.0	1.1	
Retired	100.0	19.4	80.6	55.0	18.8	6.8	
Other	100.0	19.2	80.8	51.6	17.9	11.3	
<u>45-54</u>		•					
All persons	100.0	43.5	56.5	54.3	1.8	0.4	
Usually working	100.0	47.7	52.3	51.4	0.8	(*)	
Keeping house	100.0	37.0	63.0	60.3	2.3	(*)	
Retired	100.0	(*)	81.9	47.0	31.3	(*)	
Other	100.0	25.4	74.6	58.2	11.6	4.6	
<u>55-64</u>							
All persons	100.0	35.6	64.4	58.6	4.5	1.3	
Usually working	100.0	41.3	58.7	57.0	1.4	(*)	
Keeping house	100.0	31.0	69.0	62.6	5.5	0.9	
Retired	100.0	17.6	82.3	50.9	23.6	7.7	
Other	100.0	20.1	79.9	55.1	15.9	8.9	
<u>65+</u>					,		
All persons	100.0	22.7	77.3	59.2	13.8	4.4	
Usually working	100.0	33.9	66.1	62.3	3.4	(*)	
Keeping house	100.0	21.7	78.3	63.4	13.0	1.9	
Retired	100.0	19.6	80.4	55.6	18.0	6.8	
Other	100.0	12.1	88.0	41.3	26.2	20.4	
<u>65-74</u>							
All persons	100.0	25.8	74.2	62.4	9.5	2.3	
Usually working	100.0	35.0	65.0	62.1	2.6	(*)	
Keeping house	100.0	23.9	76.0	65.4	9.5	1.2	
Retired	100.0	22.5	77.5	60.2	13.3	3.9	
Other	100.0	15.1	84.9	51.0	21.9	12.3	
<u>75+</u>							
All persons	100.0	16.7	83.3	52.8	22.0	8.5	
Usually working	100.0	27.3	72.9	63.4	7.8	(*)	
Keeping house	100.0	16.8	83.2	58.9	21.0	3.4	
Retired	100.0	16.0	84.0	49.8	23.8	10.3	
Other	100.0	8.8	91.2	30.9	30.9	29.6	

Table 13. Average number and percent distribution of persons 65 years and older according to chronic limitation of activity by sex and family relationship: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

•	ł	.	Per		th l+ chro	nic	
	m-4-1	Persons	<u></u>	cond	ditions		
Sex and family relationship	Total	with no	1	Limita	ation of a	ion of activity	
• • • • • • • • • • • • • • • • • • •	persons	chronic conditions	Total		l		
		conditions		None	Partial	Major	
•	2.5		<u> </u>		l	L	
		verage nimbe	er of per	sons in	thousands		
	Average number of persons in thousands						
Both sexes-65+	14,666	3,328	11,338	5,137	4,102	2,099	
Living alone or with nonrelatives	3,372	617	2,755	1,412	961	382	
Living with relatives	11,294	2,711	8,583	3,725	3,141	1,717	
Married	7,494	1,866	5,628	2,611	2,079	938	
Other status	3,800	845		1,114		779	
		- 1.5	-,,,,,	-,	1,002	,,,	
Male-65+	6,687	1,657	5,030	2,096	1,636	1,298	
Living alone or with nonrelatives	1,096	220	876	366	255	256	
Living with relatives	5,591	1,437	4,154		255	256	
Married	4,641	1,193	3,448	1,731	1,381	1,042	
Other status	950	244	705	1,486 245	1,186	777	
other status		244	705	245	195	265	
Female-65+	7,980	1,671	6,308	3,040	2,466	801	
Living alone or with nonrelatives	2,277	398	1,879	1,046	706	107	
Living with relatives	5,703	1,274	4,429	1,994	706 1.760	127	
Married	2,852	673	- 1	- 1		674	
Other status	2,851	601	2,180 2,249	1,125 869	894 867	161 514	
	_,051	0011	-,,	0071	5071	214	
		Perc	ent dist	ribution	1		
Both sexes-65+	100.0	22.7	77.3	35.01	28.0	14.3	
•					20.0		
Living alone or with nonrelatives	100.0	18.3	81.7	41.9	28.5	11.3	
Living with relatives	100.0	24.0	76.0	33.0	27.8	15.2	
Married	100.0	24.9	75.1	34.8	27.7	12.5	
Other status	100.0	22.2	77.8	29.3	27.9	20.5	
·							
Male-65+	100.0	24.8	75.2	31.3	24.5	19.4	
Living alone or with nonrelatives	100.0	20.1	79.9	33.4	23.3	23.4	
Living with relatives	100.0	25.7	74.3	31.0	24.7	18.6	
Married	100.0	25.7	74.3	32.0	25.6		
Other status	100.0	25.7	74.2	25.8	20.5	16.7 27.9	
	200.0	-3.7	/	23.0	20.5	27.3	
Female-65+	100.0	20.9	79.0	38.1	30.9	10.0	
Living alone or with nonrelatives	100.0	17 -	02 5	45.0	21.0		
Living with relatives		17.5	82.5	45.9	31.0	5.6	
Married	100.0	22.3	77.7	35.0	30.9	11.8	
Other status		23.6	76.4	39.4	31.3	5.6	
Other Status	100.0	21.1	78.9	30.5	30.4	18.0	

Table 14. Average number and percent distribution of persons 65 years and older according to chronic limitation of mobility by sex and family relationship: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II]

		Persons	Persons with 1+ chronic conditions					
Sex and family relationship	Total persons	with no chronic	-	Limitation of mobility				
	conditio		Total	None	Partial	Major		
	Average number of persons in thousands							
Both sexes-65+	14,666	3,328	11,338	8,679	2,017	641		
Living alone or with nonrelatives	3,372	617	2,755	2,216	443	96		
Living with relatives	11,294	2,711	8,583	6,463	1,574	545		
Married	7,494	1,866	5,628	4,534	847	247		
Other status	3,800	845	2,955	1,929	728	298		
Male-65+	6,687	1,657	5,030	4,023	781	226		
Living alone or with nonrelatives	1,096	220	876	716	135	25		
Living with relatives	5,591	1,437	4,154	3,307	646	202		
Married	4,641	1,193	3,448	2,804	502	142		
Other status	950	244	705	503	143	59		
Female-65+	7,980	1,671	6,308	4,656	1,236	415		
Living alone or with nonrelatives	2,277	398	1,879	1,500	308	72		
Living with relatives	5,703	1,274	4,429	3,157	929	344		
Married	2,852	673	2,180	1,730	344	105		
Other status	2,851	601	2,249	1,426	585	239		
		Perc	ent dist	ribution	ı	•		
Both sexes-65+	100.0	22.7	77.3	59.2	13.8	4.4		
Living alone or with nonrelatives	100.0	18.3	81.7	65.7	13.1	2.8		
Living with relatives	100.0	24.0	76.0	57.2	13.9	4.8		
Married	100.0	24.9	75.1	60.5	11.3	3.3		
Other status	100.0	22.2	77.8	50.8	19.2	7.8		
Male-65+	100.0	24.8	75.2	60.2	11.7	3.4		
Living alone or with nonrelatives	100.0	20.1	79.9	65.3	12.3	2.3		
Living with relatives	100.0	25.7	74.3	59.1	11.6	3.6		
Married	100.0	25.7	74.3	60.4	10.8	3.1		
Other status	100.0	25.7	74.2	52.9	15.1	6.2		
Female-65+	100.0	20.9	79.0	58.3	15.5	5.2		
	100.0	17.5	82.5	65.9	13.5	3.2		
Living alone or with nonrelatives								
Living with relatives	100.0	22.31	77.7	55.4	16.31	6.0		
	100.0 100.0	22.3 23.6	77.7	55.4 60.7	16.3 12.1	6.0 3.7		

Table 15. Average annual number of restricted-activity days and number of restricted-activity days per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

		`	Family	income				
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown		
	Average number of restricted-activity days in thousands							
Both sexes								
All ages-45+	1,416,504	539,305	313,720	276,277	169,304	117,90		
45-54	377,366	81,499	88,448	102,981	74,900	29,53		
55-64	414,257	146,537	94,673	92,584	45,772	34,69		
65+	624,881	311,268	130,599	80,711	48,632	53,67		
65-74 75+	371,062 253,819	174,882 136,386	87,766 42,833	51,303 29,409	28,639 19,993	28,47 25,19		
Male Male	233,013	130,300	42,033	27,407	1,7,755	23,13		
	(07.005	005 071	1// //0	110 271	7/ 5/5	/		
A11 ages-45+	607,835	235,271	144,448	112,371	74,565	41,18		
45-54	156,593	33,212	39,821	38,928	35,679	8,95		
55-64 65+	180,149 271,093	61,032 141,027	42,699 61,928	43,625 29,819	20,612 18,274	12,18 20,04		
65-74	172,635	82,308	44,671	21,552	11,695	12,41		
75+	98,458	58,720	17,257	8,267	6,579	7,63		
<u>Female</u>								
11 ages-45+	808,669	304,033	169,272	163,905	94,739	76,71		
45-54	220,773	48,287	48,627	64,053	39,221	20,58		
55-64	234,108	85,505	51,974	48,960	25,160	22,50		
65+ 65-74	353,788	170,241	68,671	50,893	30,358	33,62		
75+	198,427 155,361	92,574 77,666	43,095 25,576	29,751 21,141	16,945	16,06 17,56		
	Number of	restricte	d-activit	y days pe	r person	per year		
Both sexes	į		I	1	f	ι		
All ages-45+	28.7	47.3	29.3	20.9	17.5	26.		
45-54	19.0	33.2	22.8	15.5	14.1	19.		
55-64	27.7	48.8	27.3	22.3	16.3	23.		
65+ 65-74	42.6	52.4	38.9	33.5	31.7	37. 31.		
75+	38.3 51.1	47.8 59.8	35.6 48.0	31.2 38.2	28.0 39.4	48.		
Male_								
 All ages-45+	25.8	49.2	28.3	16.7	14.9	21.		
45-54	16.2	33.2	21.9	11.3	13.0	13.		
55-64	25.1	51.9	26.9	20.3	13.1	17		
65+	40.5	54.1	36.3	26.7	26.6	35.		
65-74 75+	38.1	52.0	35.1	27.4	23.6	31.		
	45.6	57.5	39.7	25.3	34.3	42.		
Female All ages-45+	21.2	46.0	20.2	25.2	20.2	20		
	31.2	46.0	30.2	25.2	20.3	30		
45-54	21.7	33.2	23.5	19.9	15.1	24.		
55-64	30.2	46.8	27.6	24.6	20.4	27.		
65-74	44.3	51.1	41.6	39.2	35.9	39 30		
75+	38.4 55.2	44.7 61.7	36.2 55.7	34.8 47.7	32.1 42.3	52		
, , ,	33.2	01.7	35.7	, 7/./	72.3	1 72		

Table 16. Average annual number of bed-disability days and number of bed-disability days per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix [1]

are diven in obbendix [1]								
Carry and any			Family	income		. •		
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown		
,	Average	number of	bed disa	bility da	ys in the	usands		
Both sexes								
All ages-45+	475,263	174,992	107,058	95,548	52,317	45,349		
45-54	127,834	28,089	30,096	34,243	22,648	12,758		
55-64	138,801	48,962	32,037	32,124	13,154	12,525		
65+	208,628	97,941	44,924	29,181	16,515	20,066		
65-74	112,282	53,601	26,242	14,640	8,346	9,452		
75 +	96,346	44,340	18,682	14,541	8,169	10,615		
<u>Male</u>						İ		
All ages-45+	200,164	75,192	50,951	37,670	19,885	16,466		
45-54	51,212	10,549	13,394	13,734	9,208	4,328		
55-64	60,174	20,948	13,903	14,767	5,972	4,583		
65+	88,778	43,695	23,654	9,169	4,705	7,555		
65-74	53,434	26,090	15,280	5,097	2,159	4,809		
75+	35,344	17,606	8,374	4,072	2,547	.2,746		
<u>Female</u>								
A11 ages-45+	275,099	99,801	56,107	57,877	32,431	28,883		
45-54 <u>-</u>	76,622	17,540	16,702	20,509	13,440	8,430		
55-64	78,628	28,014	18,134	17,356	7,181	7,941		
65+	119,850	54,246	21,271	20,012	11,810	12,512		
65-74 75+	58,847	27,512	10,962	9,543	6,188	4,643		
/5+	61,003	26,734	10,308	10,469	5,622	i 7,869		
ned e	Number	r of bed-	disabilit	y days pe	r person	per year		
Both sexes	1	İ	1		!	I		
All ages-45+	9.6	15.4	10.0	7.2	5.4	10.2		
45-54	6.4	11.5	7.7	5.1	4.2	8.5		
55-64	9.3	16.3	9.2	7.8	4.7	8.3		
65+	14.2	16.5	13.4	12.1	10.8	14.1		
65-74	11.6	14.7	10.7	8.9	8.2	10.4		
75+	19.4	19.4	20.9	18.9	16.1	20.5		
<u>Male</u>								
All ages-45+	8.5	15.7	10.0	5.6	4.0	8.5		
45-54	5.3	10.6	7.4	4.0	3.4	6.5		
55-64	8.4	17.8	8.8	6.9	3.8	6.6		
65+	13.3	16.8	13.9	8.2	6.8	13.2		
65-74	11.8	16.5	12.0	6.5	4.4	12.3		
75+	16.4	17.2	19.3	12.5	13.3	15.1		
<u>Female</u>								
All ages-45+	10.6	15.1	10.0	8.9	6.9	11.5		
45-54	7.5	12.1	0 1	6.4	5.2	10 1		
55-64	10.1	15.3	8.1 9.6	8.7	5.8	10.1 9.8		
65+	15.0	16.3	12.9	15.4	14.0	14.6		
65-74	11.4	13.3	9.2	11.2	11.7	8.9		
75+	21.7	21.2	22.5	23.6	17.7	23.5		
		,						

Table 17. Average annual number of work-loss days for "usually working" persons and number of work-loss days per "usually working" person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

are given in Appendix II]									
	Family income								
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown			
	Avera	ige numbe	r of work	-loss day	s in thou	sands "			
Both sexes				•					
All ages-45+	189,359	40,385	46,268	53,708	36,086	12,913			
•									
45-54 55-64	90,428	13,213	22,355	27,321	21,233	6,305			
65+	72,400	16,771 10,401	17,582 6,330	21,441 4,946	11,324	5,283 1,325			
65-74	23,005		5,203	4,662	3,074	1,299			
75+	3,526	1,634	1,127	(*)	(*)	(*)			
<u>Male</u>					•				
All ages-45+	139,265	29,356	33,434	40,546	27,641	8,287			
45-54	64,905	9,369	16,485	19,502	15,413	4,137			
55-64 65+	51,010	11,188	11,357	16,735	8,812	2,918			
65-74	23,349 20,498	8,800 7,510	5,592 4,542	4,309 4,277	3,416 2,961	1,233 1,207			
75+	2,851	1 -	1,049	(*)	(*)	(*)			
Female			'			}			
All ages-45+	50,095	11,029	12,833	13,162	8,445	4,626			
45-54	25,523	3,844	5,870	7,819	5,820	2,168			
55-64	21,390		6,225	4,706	2,511	2,365			
65+	3,182	1,601	(*)	(*)	(*)	(*)			
65-74 75+	2,507		(*)	(*)	(*)	(*)			
/5+	(*)	(*)	(*)	(*)	(*)	(*)			
	Numbe	er of wor	k-loss da	ys per "u	isually wo	rking"			
Both sexes			person	per year					
All ages-45+	7.8	12.8	9.1	6.9	5.9	6.5			
45-54	6.9	10.7	9.1	5.9	5.6	6.8			
55-64	8.7	14.4	9.1	8.2	6.0	6.7			
65-74	9.8	13.6	9.0	8.9	8.5	4.9			
75+	10.0	14.1	8.6 11.6	9.5 (*)	8.4	5.6			
Male Male				` ` ′	` `	\ ` `			
All ages-45+	8.1	15.1	9.8	7.0	_6.1	5.7			
45-54	7.2		10.2	5.8	5.7	6.6			
55-64	8.4	16.2	8.9	8.4	5.9	4.9			
65+	11.1	16.7	10.7	9.4	9.4	5.6			
65-74	11.6	18.0	10.2	10.6	9.3	6.4			
•••	8.9	11.9	13.0	(*)	(*)	(*)			
<u>Female</u>			l		ļ <u>.</u> .				
All ages-45+	7.2	9.1	7.7	6.6	5.4	8.6			
45-54	6.3	7.6	7.0	6.2	5.2	7.2			
55-64	9.2	11.8	9.5	7.6	6.4	12.9			
65-74	5.2 4.7	6.7	(*)	(*)	(*)	(*)			
75+	(*)	6.2 (*)	(*)	(*)	(*)	(*)			
• • •	` '	l `´	l `´	l `´	\ ` ′	1			

Table 18. Average annual number of restricted-activity days and number of restricted-activity days per person per year for persons 45 years and older by major activity, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

	Major activity (45 years and older)					
Sex and age	Total	Usually working	Keeping house	Retired	Other	
	Average number of restricted-activity days in thousands					
Both sexes						
All ages-45+	1,416,504	368,228	558,428	316,738	173,111	
45-54	377,366	171,288	151.851	6,555	47,672	
55-64	414,257	133,655	174,234	47,385	58,983	
65+	624,881	63,285	232,343	262,797	66,456	
65-74	371,062 253,819	51,445	149,832 82,511	137,046 125,751	32,739 33,717	
Male	253,619	11,840	02,511	125,751	. 33,717	
			415			
A11 ages-45+	607,835	267,112	(*)	233,177	105,728	
45-54	156,593	118,802	(*)	5,308	31,734	
55-64 65 +	180,149	96,498	(*) (*)	39,431	44,137	
65-74	271,093 172,635	51,811 41,296	(*) (*)	188,437 111,390	29,858 19,263	
75+	98,458	10,515	(*)	77,048	10,594	
Female	75,125	10,010		,		
111 ages-45+	808,669	101,116	556,609	83,561	67,383	
45-54	220,773	52,486	151,101	1,248	15,938	
55-64	234,108	37,157	174,151	7,954	14,846	
65+	353,788	11,474	231,356	74,360	36,598	
65-74	198,427	10,149	149,147 82,209	25,656 48,704	13,476 23,123	
/ 	155,361	1,325	62,209	40,704	23,12.	
	Number of restricted-activity days per person per year					
Both sexes	-	,	,	, cur		
All ages-45+	28.7	15.2	33.1	51.6	76.3	
45-54	19.0	13.1	25.7	79.0	62.8	
55-64	27.7	16.0	33.9	69.9	78.6	
65+ 65-74	42.6	23.3	39.9	48.9	87.4	
75+	38.3	22.3	37.3	46.1	82.	
Male	51.1	29.7	45.6	52.4	93.1	
All ages-45+	25.8	15.5	(*)	47.9	71.9	
·						
45-54 55-64	16.2	13.1	(*)	79.2	56.5	
65+	25.1 40.5	16.0 24.7	(*) (*)	70.2 44.4	77.7	
65-74	38.1	23.3	(*)	44.4	87.6 80.3	
75+	45.6	32.8	(*)	44.5	104.9	
<u>Female</u>						
All ages-45+	31.2	14.5	33.1	66.1	84.4	
45-54	21.7	13.0	25.6	73.4	80.9	
55-64	30.2	16.0	34.0	69.2	82.0	
65+	44.3	18.6	39.8	65.6	87.	
65-74 75+	38.4	18.9	37.2	55.4	85.3	
/	55.2	16.8	45.6	72.8	88.6	

Table 19. Average annual number of bed-disability days, and bed-disability days per person per year for persons 45 years and older by major activity, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

Sex and age	Major activity (45 years and older)					
	Total	Usually working	Keeping house	Retired	Other	
	Average number of bed-disability days in thousands					
Both sexes						
All ages-45+	475,263	117,347	167,138	120,820	69,958	
45-54	127,834	58,274	50,365	2,470	16,72	
55-64	138,801	45,002	52,959	16,662	24,178	
65+	208,628	14,071	63,814	101,688	29,05	
65-74	112,282	11,354	40,345	47,961	12,622	
75+	96,346	2,717	23,469	53,728	16,432	
<u>Male</u>						
All ages-45+	200,164	80,221	(*)	79,977	39,338	
45-54	51,212	37,857	(*)	1,556	11,238	
55-64	60,174	31,095	(*)	11,842	17,17	
65+	88,778	11,270	(*)	66,579	10,929	
65-74	53,434	8,848	(*)	37,643	6,942	
75+	35,344	2,421	(*)	28,936	3,987	
· <u>Female</u>						
All ages-45+	275,099	37,126	166,511	40,843	30,620	
45-54	76,622	20,417	49,804	(*)	5,487	
55-64	78,628	13,907	52,893	4,820	7,008	
65+	119,850	2,801	63,814	35,109	18,126	
65-74	58,847	2,506	40,345	10,317	5,680	
75+	61,003	(*)	23,469	24,792	12,446	
Both sexes	Number of bed-disability days per person per ye					
All ages-45+	9.6	4.9	9.9	19.7	30.	
45-54	6.4	4.5	8.5	29.8	22.0	
55-64	9.3	5.4	10.3	24.6	32.	
65+	14.2	5.2	11.0	18.9	38.	
65-74 75+	11.6	4.9	10.1	16.1	31.	
*	19.4	6.8	13.0	22.4	45.4	
Male						
All ages-45+	8.5	4.7	(*)	16.4	26.	
45-54	5.3	4.2	(*)	23.2	20.0	
55-64	8.4	5.1	(*)	21.1	30.	
65+	13.3	5.4	. (*)	15.7	32.0	
65-74	11.8	5.0	(*)	15.0	28.	
75+	16.4	7.5	(*)	16.7	39.	
<u>Female</u>						
All ages-45+	10.6	5.3	9.9	32.3	38.4	
45-54	7.5	5.0	8.4	(*)	27.	
55-64	10.1	6.0	10.3	41.9	38.	
))-04			1	1		
65+	15.0	4.5	11.0	31.0	43.3	
65+ 65-74	15.0 11.4	4.5 4.7	11.0	31.0 22.3	43. 35.	

Table 20. Average annual number of work-loss days and number of work-loss days per person per year for persons 45 years and older by major activity, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

	Major activity (45 years and older)					
Sex and age	Total	Usually working	Keeping house	Retired	Other	
	Average number of work-loss days in thousands					
Both_sexes						
A11 ages-45+	257,744	189,359	19,977	12,912	35,497	
45-54	114,097	90,428	9,279	1,174	13,216	
55-64	97,282	72,400	5,149	5,084	14,649	
65+	46,364	26,531	5,547	6,654	7,632	
65-74	38,144	23,005	4,807	4,792	5,539	
	8,221	3,526	740	1,862	2,093	
<u>Male</u>				ļ		
All ages-45+	182,622	139,265	(*)	11,329	31,457	
45-54	76,631	64,905	(*)	1,174	10,551	
55-64	69,333	51,010	(*)	4,144	14,096	
65+	36,659	23,349	(*)	6,011	6,809	
65-74 75 +	30,591	20,498	(*)	4,430	5,174	
,3.	6,067	2,851	(*)	1,581	1,635	
<u>Female</u>		,				
All ages-45+	75,122	50,095	19,403	1,583	4,040	
45-54	37,467	25,523	9,279	(*)	2,665	
55-64	27,950	21,390	5,066	(*)	(*)	
65+	9,705	3,182	5,058	(*)	(*)	
65-74	7,552	2,507	4,318	(*)	(*)	
75+	2,153	l (*)	(*)	(*)	(*)	
Both sexes	Number of work-loss days per person per year					
All ages-45+	5 .2 ·	7.8	1.2	2.1	15.6	
					,	
45-54 55-64	5.8	6.9	1.6	14.1	17.4	
65+	6.5 3.2	8.7 9.8	1.0	7.5 1.2	19.5 10.0	
65-74	3.9	10.0	1.2	1.6	13.9	
75+	1.7	8.8	0.4	0.8	5.8	
Male						
All ages-45+	7.8	8.1	(*)	2.3	21.4	
45-54	7.0	7.0	(4)	17.5	10.0	
55-64	7.9 9.7	7.2 8.4	(*) (*)	17.5	18.8 24.8	
654	5.5	11.1	(*)	1.4	20.0	
65-74	6.8	11.6	(*)	1.8	21.6	
75+	2.8	8.9	(*)	0.9	16.2	
<u>Female</u>		j				
All ages-45+	2.9	7.2	1.2	1.3	5.1	
45-54	3.7	6.3	1.6	(*)	13.5	
55-64	3.6	9.2	1.0	(*)	(*)	
•	1.2	5.2	0.9	(*)	(*)	
65+		1				
65+ 65-74 75+	1.5	4.7	1.1 (*)	(*) (*)	(*) (*)	

ILLNESS AMONG OLDER PERSONS

Illness among older persons in this section of the report is considered in terms of illness associated with selected chronic conditions and illness from acute conditions. Greater emphasis has been given to chronic conditions because they constitute a significant health problem in the older segment of the population.

In the National Health Survey, data on chronic and acute conditions are based primarily on replies to a series of "illness-recall" questions. Most of these questions, designed to assist the respondent in reporting illnesses as accurately and completely as possible, refer to illnesses occurring during the 2-week period prior to the week of interview. Other questions were designed to obtain reports of chronic conditions and impairments which were prevalent at the time of the interview.

Whenever an injury, or the residual effects of an injury was reported, information about the circumstances of the original accident was recorded in table A (see Appendix III). Only one entry was made in table A for each accident incurred by a person, irrespective of the number of injuries which he suffered as a result of the specific accident.

Prevalence of Chronic Conditions

A condition, for purposes of the survey, is considered to be chronic if it is described by the respondent in terms of one of the conditions on the "Check List of Chronic Conditions" or in terms of one of the impairments on the "Check List of Impairments" (Cards A and B, Appendix III), or is reported as having been present for more than 3 months at time of interview.

The health-interview phase of the National Health Survey measures the presence of disease or illness in terms of cases which the respondent is aware of, remembers, and considers of sufficient importance to report. Thus, the prevalence of chronic conditions based on this kind of information may differ widely from prevalence estimates made on findings in clinical studies where conditions are detected through recognized diagnostic tests and clinically significant symptoms. In general, the reporting of conditions which have been seen recently by a physician or have caused some disability is considered to be fairly accurate and complete. An exception to this is, of course, the reporting of conditions which respondents are sometimes reluctant to mention, such as mental illness and malignant neoplasm.

Because of the many factors that influence the accuracy and completeness of reporting of chronic conditions in the household interview, it has been the policy of the National Health Survey to prepare individual reports dealing with specific conditions or condition categories. This has made it possible for each report to enlarge upon the limitations and qualifications of data pertaining to the particular diagnostic category. Also, it has permitted the consideration of prevalence estimates for each category in relation to estimates from clinical and from other survey studies.

Chronic condition groups for which data have been released by the National Health Survey in the B Series of reports are: Impairments (No. 9). Chronic Respiratory Conditions (No. 12), Heart Conditions and High Blood Pressure (No. 13). Peptic Ulcers (No. 17), and Arthritis and Rheumatism (No. 20). Several other reports are in preparation. Even though it is beyond the scope of this report to repeat all of the proper reservations for each of these condition groups, information relevant to the older population from the listed reports and those now in preparation has been abstracted and summarized for the convenience of the reader. It is suggested that users of the data refer to the B Series reports for a more complete background.

The chronic condition groups shown in this report are listed below with their equivalent International List Numbers or Supplementary Impairment Code Numbers:

Chronic Condition Group	International Classification Code Numbers, 1955 Revision
Heart conditions	410-443
High blood pressure	444-447
Diabetes	260
Peptic ulcer	540-542
Arthritis and rheumatism	720-727
Hernia	560-561
Asthma-hay fever	240-241
Chronic bronchitis	502
Impairment Group	Supplementary Impairment Code Numbers
Visual impairments	X00-X05
Hearing impairments	X06-X09
Paralysis of major extremities	
and/or trunk	X40-X49; X50-X59;
	X60-X69

- 1. Of the approximate 14,667,000 noninstitutionalized persons 65 years and older in the population, about 23 percent had no chronic conditions; 26 percent had 1 chronic condition; 20 percent, 2 conditions; and 31 percent, 3 or more chronic conditions (table D).
- 2. With the exception of diabetes, peptic ulcer, asthma-hay fever, and bronchitis, the prevalence of the chronic conditions included in this report increased consistently with age. Typical prevalence trends by age are shown in figure 5.
- 3. Prevalence rates among persons 65 years and older for all of the chronic conditions included in this report were higher in rural-nonfarm areas than in urban areas. Rates for persons 65 years and older living on farms in rural areas were not essentially different from rates for persons in rural-nonfarm areas.

Table D. Distribution of persons 45 years and older by age and chronic condition status: United States, July 1957-June 1959

Age	Total persons	Number of chronic conditions				
		. 0	1	2	3+	
All ages-45+	100.0	35.0	28.1	17.2	19.6	
45-64	100.0	40.1	29.1	15.9	14.8	
65+	100.0	22.7	25.8	20.4	31.1	




Figure 5. Prevalence of selected chronic conditions among persons 45 years and older by age.

The percentage of cases that had never been seen by a physician was negligible in the diagnostic categories-heart conditions, high blood pressure, diabetes, peptic ulcer, and paralysis of extremities and/or trunk. Less than 10 percent of the cases among those 65 years and older in the categories, bronchitis and visual impairment, were never medically attended. About one seventh of the hernia cases, and one fifth of the asthma-hay fever and arthritis and rheumatism cases were reported as having never been seen by a physician. Although persons who were reported as having these medically unattended conditions may have had symptoms which they interpreted as the condition, the accuracy of their self-diagnosis is subject to doubt. About 40 percent of the hearing impairments for this age group had never been medically attended (fig. 6).

- 5. Approximately one half of the cases of paralysis of extremities and/or trunk among persons 65 years and older caused major limitation of activity. About one fourth of the heart conditions and visual impairments resulted in major activity limitation. In all other condition categories less than 15 percent of the cases caused major limitation of activity.
- 6. The relative amount of disability associated with the condition groups included in this report is shown in figure 7. This is in terms of the average annual number of restricted-activity and bed-disability days per person 65 years and older with the condition.




Figure 6. Percent of cases medically attended under care and not under care and never medically attended for selected chronic conditions among persons 65 years and older.




Figure 7. Disability days associated with selected chronic conditions per person with the condition per year among persons 65 years and older.

Tabular information on disability associated with chronic conditions can be found in tables 26 and 27. The reader is again reminded that it is possible for a day of disability to have been the result of more than one condition. In such cases the day is ascribed in the statistics to each condition that was reported to have caused the disability.

Incidence of Acute Conditions

In the National Health Survey, an acute condition is defined as a condition which has lasted less than 3 months, and has involved either medical attention or 1 or more days of restricted activity. However, the conditions and impairments listed on Cards A and B (Appendix III) are always considered as chronic regardless of duration of the condition.

With the increased emphasis placed on chronic conditions among older persons during recent years, there has been a tendency to underestimate the importance of illness and disability due to acute conditions among older members of the population. While the rate of incidence of acute conditions among persons 45 years and older is no higher than for other adults in the population, it is evident from data presented in this report that the disability from acute conditions to which older persons are subject is by no means negligible.

To demonstrate the impact of the Asian influenza epidemic of 1957-58 on older persons, data by type of acute condition are presented separately for the 2 years covered in this report. However, data pertaining to injuries by type of injury and to persons injured by class of accident represent average annual incidence based on the 2 years of data collection.

Other National Health Survey publications in the B Series of reports pertaining to acute conditions, and containing more complete information by age group, include: Acute Conditions, July 1957-June 1958 (No. 6), Persons Injured, July 1957-June 1958 (No. 8), Types of Injuries, July 1958-June 1959 (No. 16), and Acute Conditions, July 1958-June 1959 (No. 18).

The acute condition groups listed in this report with equivalent International Classification Code numbers are listed below. It should be noted that only those conditions meeting the acute conditions definitions (see Appendix II) are included in tables in this publication.

Acute Condition Group

Infectious and parasitic Upper respiratory Other respiratory Digestive system

Fractures, dislocations, sprains, and strains Open wounds and lacerations Contusions and superficial injuries Other current injuries All other acute conditions

International Classification Code Numbers, 1955 Revision

020-138 470-475; 517 480-501; 518-527; 783 530-539; 543-553; 570-587; 784; 785

N800-N848 N870-N908

N910-N929 N850-N869; N930-N999 All other acute condition code numbers

- 1. During the period July 1957-June 1958, about 23.5 million acute conditions, a rate of 1,626 conditions per 1,000 persons, were reported for persons 65 years and older. During a comparable period, July 1958-June 1959, about 20 million conditions, a rate of 1,339 per 1,000 persons, were reported for persons in this age group (table E). The higher rate during July 1957-June 1958 was due, for the most part, to the greater number of conditions in the "other respiratory conditions" category, which includes influenza, pneumonia, and acute bronchitis.
- 2. For persons 65 years and older an annual average of 3,433,000 injuries involving a total of 14,743,000 bed-disability days were reported. Injuries, like other acute conditions, are included in these figures only if they involved medical attention or caused restriction of the person's activities. About 85 percent of the bed-disability days reported were associated with fractures, dislocations, sprains, strains, contusions, and superficial injuries.
- 3. About 3,352,000 persons 65 years and older were injured annually. This constitutes a rate of 228.5 persons per 1,000 population. About two thirds of the persons were injured in accidents occurring in the home (table F).

Table E. Incidence of acute conditions among persons 45 years and older: United States, July 1957-June 1958, July 1958-June 1959

	July	1957-June	1958	July	1958-June	1959
Acute conditions	All ages 45+	45-64	65+	All ages 45+	45-64	65+
	N	umber of	acute con	ditions in	thousand	s
Total acute conditions-	84,318	60,726	23,592	69,852	50,006	19,846
Upper respiratory conditions Other respiratory conditions Other acute conditions	28,406 27,961 27,951	19,966 21,099 19,661	•		19,489 10,124 20,393	•
	Number o	f acute c	onditions	per 1,000	persons	per year
Total acute conditions-	1,721.4	1,761.7	1,625.7	1,400.5	1,426.5	1,339.0
Upper respiratory conditions Other respiratory conditions	579.9 570.8	579.2 612.1	581.6 472.9	553.8 282.5	556.0 288.8	548.7 267.5
Other acute conditions	570.6	570.4	571.2	564.2	581.7	522

NOTE: Since the estimates shown in this table are based on single years of data collection, the standard errors are 1.25 times the standard errors shown in Appendix 1.

Table F. Persons injured by class of accident for persons 45 years and older: United States, July 1957-June 1959

			A	ge		
Class of accident	All ages 45+	45-64	65+	All ages 45+	45-64	65+
	Number of persons in thousands			Per 1,000 persons per year		
All classes	11,460	8,108	3,352	231.8	233.2	228.5
Motor vehicle While at work Home Other and unknown	1,049 2,638 5,163 2,611	865 2,381 2,870 1,993	184 258 2,293 618	21.2 53.4 104.5 52.8	24.9 68.5 82.6 57.3	12.5 17.6 156.3 42.1

DETAILED TABLES

			Page
		CHRONIC CONDITIONS	
[able	21.	Prevalence of selected chronic conditions among persons 45 years and older by age and sex: United States, July 1957-June 1959	35
	22.	Prevalence of selected chronic conditions among persons 45 years and older by age and residence: United States, July 1957-June 1959	36
	23.	Number and percent distribution of selected chronic conditions among persons 45 years and older by medical attention and age: United States, July 1957-June 1959	37
	24.	Number and percent distribution of selected chronic conditions among persons 45 years and older according to degree of chronic activity limitation caused by the condition by age: United States, July 1958-June 1959	38
	25.	Number and percent distribution of selected chronic conditions among persons 45 years and older according to degree of chronic mobility limitation caused by the condition by age: United States, July 1958-June 1959	39
	26,	Average annual number of disability days for selected chronic conditions for persons 45 years and older by sex and age: United States, July 1957-June 1959	40
-	27.	Number of disability days for selected chronic conditions per person with the condition per year for persons 45 years and older by sex and age: United States, July 1957-June 1959	41
		ACUTE CONDITIONS	
	28.	Incidence of acute conditions among persons 45 years and older by type of condition and sex: United States, July 1957-June 1958, July 1958-June 1959	42
	29.	Bed-disability days associated with acute conditions and bed-disability days per person per year for persons 45 years and older by type of condition and sex: United States, July 1957-June 1958, July 1958-June 1959	43
	30.	Average annual incidence of injuries and number of bed-disability days associated with injuries for persons 45 years and older by type of injury and age: United States, July 1957-June 1959	44

Table 21. Prevalence of selected chronic conditions among persons 45 years and older by age and sex:
United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

				-								
				, <u></u>		Age						
Sex and selected conditions	All ages 45+	45-54	55-64	Total 65+	65-74	75+	All ages 45+	45-54	55-64	Total 65+	65-74	75+
		Number	of chro	nic con	ditions							
Both sexes				usands			i	Rate p	er 1,00	0 popul	at1on	
Heart conditions	4,046	726	1,137	2,183	1,249	935	81.9	36.6	76.2	148.8	128.8	188.2
High blood pressure	4,212	1,002	1,315	1,894	1,246	649	85.2	50.5	88.1	129.1	128.5	130.6
Diabetes	1,264	246	424	593	416	177	25.6	12.4	28.4	40.4	42.9	35.6
Peptic ulcer	1,286	569	390	327	256	70	26.0	28.7	26.1	22.3	26.4	14.1
Arthritis and rheuma-												
tism	8,920	2,253	2,769	3,898	2,477	1,421	180.5	113.6	185.5	265.8	255.4	286.0
Hernia	1,658	376	481	801	480	321	33.5	19.0	32.2	54.6	49.5	64.6 47.7
Asthma-hay fever Chronic bronchitis	2,880 780	1,170 252	924 -250	786 277	549 191	237 86	58.3 15.8	59.0 12.7	61.9 16.7	53.6 18.9	56.6 19.7	17.3
GILORIC BIOREMICIS] /65	.252		-//	1,1	00	15.0	12.7	10.7	10.7	'''	1,.3
Visual impairments	2,353	376	463	1,514	687	827	47.6	19.0	31.0	103.2	70.8	166.4
Deafness and other		756	001	0 500			06.4	20.1			300 0	050 /
hearing impairments Paralysis of major ex-	4,270	756	994	2,520	1,231	1,289	86.4	38.1	66.6	171.8	126.9	259.4
tremities and/or trunk-	612	111	174	328	161	167	12.4	5.6	11.7	22.4	16.6	33.6
Male												
Heart conditions	2,041	419	622	1,001	639	362	86.7	43.3	86.6	149.7	141.1	167.8
High blood pressure	1,116	289	336	491	329	163	47.4	29.9	46.8	73.4	72.6	75.6
Diabetes	513	108	181	224	156	68	21.8	11.2	25.2	33.5	34.4	31.5
Peptic ulcer	892	392	280	221	176	. 45	37.9	40.5	39.0	33.0	38.9	20.9
Arthritis and rheuma-							1					
tism	3,149	808	998	1,343	898	444	133.7	83.5	138.9	200.8	198.2	205.8
Hernia	1,301	272 519	372 469	657 424	392 300	265 124	55.3 59.9	28.1 53.6	51.8 65.3	98.3 63.4	86.5 66.2	122.9 57.5
Asthma-hay fever Chronic bronchitis	365	105	125	135	88	48	15.5	10.9	17.4	20.2	19.4	22.3
Ontonic bronenicia	1 303	103	123	1 233		10	13.3	10.5	-,,-	-0	27,17	
Visual impairments	1,077	202	240	635	313	322	45.7	20.9	33.4	95.0	69.1	149.3
Deafness and other	1,0,,	202	240	033	313	, ,,,,	73.7		33.4	1 33.0	07.1	147.5
hearing impairments	2,397	442	617	1,338	701	636	101.8	45.7	85.9	200.1	154.7	294.9
Paralysis of major ex- tremities and/or trunk-	330	58	100	171	93	79	14.0	6.0	13.9	25.6	20.5	36.6
tremittes and/of trunk-	330	50	100	1/1	,,,	,,	14.0	0.0	13.7	25.0	20.5	30.0
Female				[[
Heart conditions	2,004	307	515	1,182	609	573	77.4	30.2	66.5	148.1	117.9	203.8
High blood pressure	3,096	713	980	1,403	917	486	119.6	70.2	126.5	175.8	177.5	172.8
Diabetes	751	139	244	369	260	109	29.0	13.7	31.5	46.2	50.3	38.8
Peptic ulcer	394	178	110	106	81	25	15.2	17.5	14.2	13.3	15.7	8.9
Arthritis and rheuma-					`		Ì					
tism	5,771	1,445	1,771	2,555	1,579	977	223.0	142.3	228.6	320.2	305.6	347.4
Hermia	357	104	109	144	88	56	13.8	10.2	14.1	18.0	17.0	19.9
Asthma-hay fever	1,469	651	456	362	250	113	56.8	64.1	58.9	45.4	48.4	40.2
Chronic bronchitis	415	148	126	142	104	38	16.0	14.6	16.3	17.8	20.1	13.5
Viewal impelments	1,276	17/	223	879	374	505	49,3	17.1	28.8	110,2	72.4	179.6
Visual impairments Deafness and other]	1	ļ	j		1			j	j		
hearing impairments Paralysis of major ex-	1,873	313	377	1,183	530	653	72.4	30.8	48.7	148.2	102.6	232.2
tremities and/or trunk-	283	52	74	156	68	88	10.9	5.1	9.6	19.5	13.2	31.3
cremities and/of trunk-	263			1,30								

Table 22. Prevalence of selected chronic conditions among persons 45 years and older by age and residence: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civillan noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

												
		·				Age	·	 -	,			
Residence and selected conditions	All ages 45+	45-54	55-64	Total 65+	65-74	75+	All ages 45+	45-54	55-64	Total 65+	65-74	75+
<u>Urban</u>		Number		onic co ousands	ndition	s	Rate per 1,000 population					
Heart conditions High blood pressure Diabetes Peptic ulcer	2,470 2,645 802 792	463 647 157 350	709 838 268 248	1,298 1,161 377 194	765 779 264 143	533 382 112 50	76.4 81.9 24.8 24.5	35.8 50.0 12.1 27.0	71.5 84.6 27.0 25.0	137.3 122.8 39.9 20.5	120.9 123.1 41.7 22.6	170.5 122.2 35.8 16.0
Arthritis and rheuma- tism	5,352 975 1,762 510	1,356 218 744 165	1,657 285 560 172	2,339 471 458 174	1,496 281 327 118	843 190 131 56	165.6 30.2 54.5 15.8	104.8 16.8 57.5 12.7	167.2 28.8 56.5 17.4	247.4 49.8 48.4 18.4	236.4 44.4 51.7 18.6	269.6 60.8 41.9 17.9
Visual impairments	1,409	220	272	917	428	489	43.6	17.0	27.4	97.0	67.6	156.4
Deafness and other hearing impairments Paralysis of major ex-	2,619	487	629	1,503	725	778	81.1	37.6	63.5	159.1	114.6	248.8
tremities and/or trunk-	376	79	110	187	100	88	11.6	6.1	11.1	19.8	15.8	28.1
Rural nonfarm												
Heart conditions High blood pressure Diabetes-1 Peptic ulcer	1,009 990 297 296	170 212 · 54 132	274 297 99 85	566 481 144 78	306 307 103 67	259 174 41 (*)	91.5 89.8 26.9 26.9	37.8 47.1 12.0 29.3	86.4 93.6 31.2 26.8	169.1 143.7 43.0 23.3	142.0 142.5 47.8 31.1	217.1 145.9 34.4 (*)
Arthritis and rheuma- tism	2,107 398 730 178	506 94 276 61	646 101 245 52	955 203 210 65	590 119 142 49	366 84 68 (*)	191.1 36.1 66.2 16.1	112.4 20.9 61.3 13.6	203.6 31.8 77.2 16.4	285.2 60.6 62.7 19.4	273.8 55.2 65.9 22.7	306.8 70.4 57.0 (*)
Visual impairments Deafness and other hearing impairments Paralysis of major ex- tremities and/or trunk-	605 1,072	95 169 (*)	117 242 47	393 662 86	175 323 40	218 339 46	54.9 97.3	21.1 37.5 (*)	36.9 76.3 14.8	117.4 197.7 25.7	81.2 149.9	182.7 284.2 38.6
Rural farm	133	()	7,		~		13.5	()	14.0		10.0	30.0
Heart conditions High blood pressure Diabetes Peptic ulcer	567 577 • 165 198	93 143 35 87	154 181 57 56	319 252 72 55	177 160 49 47	142 92 23 (*)	93.0 94.7 27.1 32.5	39.0 59.9″ 14.7 36.4	83.4 98.0 30.9 30.3	171.3 135.3 38.7 . 29.5	146.0 132.0 40.4 38.8	218.8 141.8 35.4 (*)
Arthritis and rheuma- tism	1,461 285 387 91	392 64 150 27	466 95 120 26	604 127 118 38	392 80 81 25	212 47 37 (*)	239.7 46.8 63.5 14.9	164.2 26.8 62.8 11.3	252.3 51.4 65.0 14.1	324.4 68.2 63.4 20.4	323.4 66.0 66.8 20.6	326.7 72.4 57.0 (*)
Visual impairments Deafness and other	339	61	74	204	83	121	55.6	25.6	40.1	109.6	68.5	186.4
hearing impairments Paralysis of major ex- tremities and/or trunk-	579 84	100 (*)	123 (*)	356 54	183	172 33	95.0	41.9 (*)	66.6 (*)	191.2 29.0	151.0	265.0 50.8

Table 23. Number and percent distribution of selected chronic conditions among persons 45 years and older by medical attention and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957—June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11]

	·	Medic atte	ally nded	Never medi-		Medic atte	ally nded	Never
Age and selected conditions	Total	Under care	Not under care	cally at- tended	Total	Under care	Not under care	cally at- tended
	Number	of chro	nic cond usands	itions	Pe	rcent di	stributi.	on
A11 ages-45+			1			1		
Heart conditions High blood pressure Diabetes Peptic ulcer	4,046 4,212 1,264 1,286	3,220 3,080 1,162 1,010	774 1,066 99 258	52 66 (*) (*)	100.0 100.0 100.0 100.0	79.6 73.1 91.9 78.5	19.1 25.3 7.8 20.1	1.3 1.6 (*) (*)
Arthritis and rheumatism Hernia	8,920 1,658 2,880 780	3,619 638 1,397 298	3,639 801 952 419	1,662 219 532 63	100.0 100.0 100.0 100.0	40.6 38.5 48.5 38.2	40.8 48.3 33.1 53.7	18.6 13.2 18.5 8.1
Visual impairments Deafness and other hearing	2,353	971	1,194	188	100.0	41.3	50.7	8.0
impairments	4,270 612	649 320	2,069 272	1,552 21	100.0	15.2 52.3	48.5 44.4	36.3
<u>45-64</u>								
Heart conditions High blood pressure Diabetes Peptic ulcer	1,863 2,317 671 959	1,407 1,645 615 764	433 633 55 180	(*) 40 (*) (*)	100.0 100.0 100.0	75.5 71.0 91.7 79.7	23.2 27.3 8.2 18.8	(*) 1.7 (*) (*)
Arthritis and rheumatism Hernia Asthma-hay fever	5,022 857 2,094	1,954 298 1,037	2,146 458 693	923 102 364	100.0 100.0 100.0	38.9 34.8 49.5	42.7 53.4 33.1	18.4 11.9 17.4
Chronic bronchitis	503	189	277	37	100.0	37.6	55.1	7.4
Visual impairments Deafness and other hearing impairments	839 1,750	354 294	408 954	77 502	100.0	42.2 16.8	48.6 54.5	9.2
Paralysis of major extremities and/or trunk	285	145	129	(*)	100.0	50.9	45.3	(*)
<u>65+</u>				-				
Heart conditions	2,183 1,894 593 327	1,814 1,435 547 246	341 433 45 78	29 26 (*) (*)	100.0 100.0 100.0 100.0	83.1 75.8 92.2 75.2	15.6 22.9 7.6 23.9	1.3 1.4 (*) (*)
Arthritis and rheumatism Hernia	3,898 801 786 277	1,665 340 360 109	1,493 344 258 142	740 117 168 26	100.0 100.0 100.0 100.0	42.7 42.4 45.8 39.4	38.3 42.9 32.8 51.3	19.0 14.6 21.4 9.4
Visual impairments Deafness and other hearing	1,514	617	786	111	100.0	40.8	51.9	7.3
impairmentsParalysis of major extremi-	2,520	355	1,115	1,050	100.0	14.1	44.2	41.7
ties and/or trunk	328	175	143	(*)	100.0	53.4	43.6	(*)

Table 24. Number and percent distribution of selected chronic conditions among persons 45 years and older according to degree of chronic activity limitation caused by the condition by age: United States, July 1958-June 1959

Data are based on household interviews during July 1958-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

	Total	Limitat	ion of a	ctivity	Total	Limitat	ion of a	ctivit y	
Age and selected conditions	condi- tions	None	Par- tial	Major	condi- tions	None	Par- tial	Major	
,	Number	of chro		itions	P	ercent d	istribut	Lon	
All ages-45+		1							
Heart conditions	4,046	1,473	1,750	823	100.0	36.4	43.3	20.	
High blood pressure	4,212	3,142	863	207	100.0	74.6	20.5	4.	
DiabetesPeptic ulcer	1,264 1,286	937 1,028	218 182	109 75	100.0	74.1 79.9	17.2 14.2	8.0 5.8	
Arthritis and rheumatism	8,920	6,486	1,883	551	100.0	72.7	21.1	6.	
Hernia	1,658	1,251	333	74	100.0	75.5	20.1	4.	
Asthma-hay fever	2,880	2,487	277	116	100.0	86.4	9.6	4.0	
Chronic bronchitis	780	670	70	40	100.0	85.9	9.0	5.7	
Visual impairments Deafness and other hearing	2,353	1,454	474	425	100.0	61.8	20.1	18.1	
impairments	4,270	3,964	172	134	100.0	92.8	4.0	3.1	
Paralysis of major extremities and/or trunk	612	147	155	310	100.0	24.1	25.3	50.7	
<u>45-64</u>									
Heart conditions	1,863	765	846	252	100.0	41.1	45.4	13.5	
High blood pressure	2,317	1,818	446	54	100.0	78.5	19.2	2.	
DiabetesPeptic ulcer	671 959	529 799	114 130	28 30	100.0 100.0	78.8 83.3	17.0 13.6	3.	
Arthritis and rheumatism	5,022	3,942	943	138	100.0	78.5	18.8	2.	
Hernia	857	669	169	(*)	100.0	78.1	19.7	(*)	
Asthma-hay fever	2,094	1,871	192	`31	100.0	89.4	9.2	ì.	
Chronic bronchitis	503	449	41	(*)	100.0	89.3	8.2	(*)	
Visual impairments Deafness and other hearing	839	601	147	90	100.0	71.6	17.5	10.7	
impairments	1,750	1,643	77	30	100.0	93.9	4.4	1.7	
Paralysis of major extremi- ties and/or trunk	285	69	89	126	100.0	24.2	31.2	44.2	
<u>65+</u>									
Heart conditions	2,183	709	904	570	100.0	32.5	41.4	26.	
High blood pressure	1,894	1,324	418	153	100.0	69.9	22.1	8.	
Diabetes	593	408	103	81	100.0	68.8	17.4	13.	
Peptic ulcer	327	230	52	45	100.0	70.3	15.9	13.	
Arthritis and rheumatism	3,898	2,545	941	413	100.0	65.3	24.1	10.	
Hernia	801	582	164	55	100.0	72.7	20.5	6.9	
Asthma-hay feverChronic bronchitis	786 277	616 221	85 29	85 27	100.0 100.0	78.4 79.8	10.8 10.5	10.8	
Visual impairments	1,514	853	327	335	100.0	56.3	21.6	22.	
Deafness and other hearing impairments	2,520	2,321	96	104	100.0	92.1	3.8	4.:	
Paralysis of major extremities and/or trunk	328	78	66	184	100.0	23.8	20.1	56.	

NOTE: Since the estimates shown in this table are based on single years of data collection, the standard errors are 1.25 times the standard errors shown in Appendix 1.

Table 25. Number and percent distribution of selected chronic conditions among persons 45 years and older according to degree of chronic mobility limitation caused by the condition by age: United States, July 1958-June 1959

[Data are based on household interviews during July 1958—June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 11]

	Total	Limitat	ion of n	obility	Total	Limitat	ion of n	obility
Age and selected conditions	condi- tions	None	Par- tial	Major	condi- tions	None	Par- tial	Major
A11 ann 451	Number	of chro	nic cond usands	litions	Pe	ercent di	stributi	Lon
<u>All ages-45+</u>	[1	Ì		1	ì
Heart conditions	4,046 4,212	3,091 3,906	758 241	196 65	100.0 100.0	76.4 92.7	18.7 5.7	4.8 1.5
Peptic ulcer	1,264 1,286	1,126 1,229	98 39	40 (*)	100.0 100.0	89.1 95.6	7.8 3.0	3.2 (*)
Arthritis and rheumatism	8,920 1,658	7,891 1,566	846 81	183	100.0 100.0	88.5 94.5	9.5 4.9	2.1 (*)
Astham-hay fever	2,880 780	2,778 732	92 44	(*) (*)	100.0	96.5 93.8	3.2 5.6	(*)
Visual impairments Deafness and other hearing	2,353	1,784	454	115	100.0	75.8	19.3	4.9
impairments Paralysis of major extremi-	4,270	4,127	117	26	100.0	96.7	2.7	0.6
ties and/or trunk	612	222	226	164	100.0	36.3	36.9	26.8
<u>45-64</u>								
Heart conditions	1,863 2,317	1,611 2,233	209 76	43 (*)	100.0 100.0	86.5 96.4	11.2 3.3	2.3 (*)
DiabetesPeptic ulcer	671 959	641 · 932	(*) (*)	(*) (*)	100.0 100.0	95.5 97.2	(*) (*)	(*) (*)
Arthritis and rheumatism	5,022 857	4,728 832	269 22	26 (*)	100.0 100.0	94.1 97.1	5.4 2.6	0.5 (*)
Asthma-hay feverChronic bronchitis	2,094 503	2,056 481	34 (*)	(*) (*)	100.0 100.0	98.2 95.6	1.6 (*)	(*) (*)
Visual impairments Deafness and other hearing	839	723	93	23	100.0	86.2	11.1	2.7
impairments Paralysis of major extremi-	1,750	1,717	29	(*)	100.0	98.1	1.7	(*)
ties and/or trunk	285	119	112	54	100.0	41.8	39.3	18.9
<u>65+</u>							V	
Heart conditions High blood pressure Diabetes	2,183 1,894 593	1,480 1,673 485	549 165 81	154 57 27	100.0 100.0 100.0	67.8 88.3 81.8	25.1 8.7 13.7	7.1 3.0
Peptic ulcer	327	297	22	(*)	100.0	90.8	6.7	4.6 (*)
Arthritis and rheumatism	3,898 801	3,163 734	577 58	158 (*)	100.0 100.0	81.1 91.6	14.8 7.2	4.1 (*)
Asthma-hay feverChronic bronchitis	786 277	723 251	58 25	(*) (*)	100.0 100.0	92.0 90.6	7.4 9.0	(*) (*)
Visual impairments Deafness and other hearing	1,514	1,061	361	93	100.0	70.1	23.8	6.1
impairments Paralysis of major extremi-	2,520	2,410	88	22	100.0	95.6	3.5	0.9
ties and/or trunk	328	103	114	111	100.0	31.4	34.8	33.8

NOTE: Since the estimates shown in this table are based on single years of data collection, the standard errors are 1.25 times the standard errors shown in Appendix 1.

Table 26. Average annual number of disability days for selected chronic conditions for persons 45 years and older by sex and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

		.eu-activi	ty days	Bed-disability days				Work-loss days for "usually working" persons		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	
All ages 45+ .		A	verage nu	mber of d	isability	days in	thousands			
Heart conditions	211,446	100,265	111,181	74,711	32,679	42,032	15,077	13,551	1,526	
High blood pressure	80,094	23,169	56,925	25,583	6,137	19,447	7,374	4,129	3,245	
DiabetesPeptic ulcer	27,203	10,297	16,907	12,212	4,046	8,166	1,485	(*)	(*)	
reptic dicer	30,829	22,762	8,066	8,733	6,308	2,426	7,616	7,342	(*)	
Arthritis and rheumatism	212,161	74,273	137,888	52,778	18,329	34,450	12,776	8,702	4,074	
Hernia	29,303	19,345	9,958	9,797	6,091	3,707	5,099	4,396	(*)	
Asthma-hay feverChronic bronchitis	47,598 15,690	28,204 8,521	19,393 7,169	12,881 5,157	6,499 3,125	6,382 2,032	4,565 2,818	4,127 1,698	(*) 1,120	
	13,000	0,522	,,10,	3,237	3,123	_,05_	-,010	1,000	1,120	
Visual impairments Deafness and other hearing	44,177	19,567	24,610	9,546	4,747	4,798	1,520	1,404	(*)	
impairments	7,588	3,427	4,162	2,082	1,425	(*)	(*)	(*)	(*)	
Paralysis of major extremi- ties and/or trunk	45,410	26,395	19,016	20,321	11,703	8,618	(+)	(+)	(*)	
ties and/or trunk	45,410	20,393	19,010	20,321	11,703	3,010	(*)	(*)	(*)	
<u>45-64</u>										
Heart conditions	88,770	46,514	42,256	27,767	12,870	14,897	11,562	10,259	1,303	
High blood pressure Diabetes	41,340 13,681	11,074	30,266	12,511	3,246	9,265	5,712	2,506	3,206	
Peptic ulcer	20,538	6,294 14,754	7,388 5,784	5,172 5,523	2,410 4,117	2,762 1,407	1,485 7,444	(*) 7,170	(*) (*)	
		_	-						ŀ	
Arthritis and rheumatism	96,913 13,632	33,497 8,369	63,415 5,264	24,032 3,622	8,404 2,338	15,628 1,283	9,910 4,207	5,891 3,494	4,020 (*)	
Asthma-hay fever	28,433	15,994	12,438	7,925	3,706	4,219	4,398	3,960	(*)	
Chronic bronchitis	9,651	5,237	4,414	2,804	1,479	1,325	2,670	1,551	1,120	
Visual impairments	11,188	6,409	4,780	2,908	1,981	(*)	(*)	(*)	(*)	
Deafness and other hearing impairments	3,705	1,889	1,816	1,174	(*)	(4)	(4)	(4)	/43	
Paralysis of major extremi-	3,703	1,009	1,010	1,1/4	(*)	(*)	(*)	(*)	(*)	
ties and/or trunk	17,215	11,589	5,626	8,071	4,842	3,229	(*)	(*)	(*)	
<u>65+</u>	į									
Heart conditions	122,676	53,751	68,925	46,944	19,808	27,135	3,516	3,292	(*)	
High blood pressure	38,754	12,095	26,659	13,072	2,891	10,181	1,662	1,623	(*)	
Diabetes	13,522	4,003	9,519	7,040	1,637	5,403	(*)	(*)	(*)	
Peptic ulcer	10,291	8,008	2,283	3,210	2,191	1,019	(*)	(*)	(*)	
Arthritis and rheumatism	115,248	40,776	74,472	28,747	9,925	18,822	2,866	2,812	(*)	
Hernia	15,671	10,977	4,694	6,176	3,753	2,423	(*)	(*)	(*)	
Asthma-hay fever	19,165	12,210	6,955	4,956	2,793	2,164	(*) (*)	(*)	(*)	
Chronic bronchitis	6,039	3,285	2,755	2,353	1,646	(*)	(*)	(*)	(*)	
Visual impairments	32,989	13,158	19,831	6,638	2,766	3,871	(*)	(*)	(*)	
Deafness and other hearing	3,883	1,537	2,346	(*)	(*)	(*)	(*)	(*)	(*)	
Paralysis of major extremi- ties and/or trunk	28,196	14,806	13,390	12,250	6,861	5,389	(*)	(*)	(*)	

Table 27. Number of disability days for selected chronic conditions per person with the condition per year for persons 45 years and older by sex and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

Age and selected conditions	Restrict	ed-activi	ty days	Bed-d	isability	days		-loss day ally work persons	
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All ages-45+	N	umber of	disabilit	y days pe	r person	with the	condition	per year	:
Heart conditions	52.3	49.1	55.5	18.5	16.0	21.0	13.5	14.7	8.0
High blood pressure	19.0	20.8	18.4	6.1	5.5	6.3	6,2	6.5	5.9
Diabetes	21.5	20.1	22.5	9.7	7.9	10.9	4.0	(*)	(*)
Peptic ulcer	24.0	25.5	20.5	6.8	7.1	6.2	10.3	11.6	(*)
Arthritis and rheumatism	23.8	23.6	23.9	5.9	5.8	6.0	4.4	4.7	3.8
Hernia	17.7	14.9	27.9	5.9	4.7	10.4	6.6	6.0	(*)
Asthma-hay fever	16.5	20.0	13.2	4.5	4.6	4.3	3.4	4.4	(*)
Chronic bronchitis	20.1	23.3	17.3	6.6	8.6	4.9	9.5	8.8	10.9
Visual impairments Deafness and other hearing	18.8	18.2	19.3	4.1	4.4	3.8	3.1	3.5	(*)
impairments Paralysis of major extremi-	1.8	1.4	2.2	0.5	0.6	(*)	(*)	(*)	(*)
ties and/or trunk	74.2	80.0	67.2	33.2	35.5	30.5	(*)	(*)	(*)
<u>45-64</u>									
Heart conditions	47.6	44.7	51.4	14.9	12.4	18.1	13.0	14.1	8.1
High blood pressure	17.8	17.7	17.9	5.4	5.2	5.5	5.8	4.8	6.9
Diabetes	. 20 . 4	21.8	19.3	7.7	8.3	7.2	5.2	(*)	(*)
Peptic ulcer	21.4	22.0	20.2	5.8	6.1	4.9	11.0	12.4	(*)
Arthritis and rheumatism	19.3	18.5	19.7	4.8	4.7	4.9	4.2	4.0	4.5
Hernia	15.9	13.0	24.7	4.2	3.6	6.0	7.2	6.5	(*)
Asthma-hay fever	13.6	16.2	11.2	3.8	3.8	3.8	3.6	4.7	(*)
Chronic broachitis	19.2	22.8	16.2	5.6	6.4	4.9	10.2	9.3	11.8
Visual impairments Deafness and other hearing	13.3	14.5	12.0	3.5	4.5	(*)	(*)	(*)	(*)
impairments	2.1	1.8	2.6	0.7	(*)	(*)	(*)	(*)	(*)
Paralysis of major extremities and/or trunk	60.4	73.3	44.3	28.3	30.6	25.4	(*)	(*)	(*)
<u>65+</u>									
Heart conditions	56.2	53.7	58.3	21.5	19.8	23.0	15.5	16.8	(*)
High blood pressure	20.5	24.6	19.0	6.9	5.9	7.3	8.0	13.4	(*)
Diabetes	22.8	17.9	25.8	11.9	7.3	14.6	(*)	(*)	(*)
Peptic ulcer	31.5	35.2	21.5	9.8	9.9	9.6	(*)	(*)	(*)
Arthritis and rheumatism	29.6	30.4	29.1	7.4	7.4	7.4	5.4	7.7	(*)
Hernia	19.6	16.7	32.6	7.7	5.7	16.8	(*)	(*)	(*)
Asthma-hay fever	24.4	28.8	19.2	6.3	6.6	6.0	(*)	(*)	(*)
Chronic bronchitis	21.8	24.3	19.4	8.5	12.2	(*)	(*)	(*)	(*)
Visual impairments Deafness and other hearing	21.8	20.7	22.6	4.4	4.4	4.4	(*)	(*)	(*)
impairments	1.5	1.1	2.0	(*)	(*)	(*)	(*)	(*)	(*)
Paralysis of major extremities and/or trunk	86.0	86.6	85.8	37.3	40.1	34.5	(*)	(*)	(*)

Table 28. Incidence of acute conditions among persons 45 years and older by type of condition and sex: United States, July 1957-June 1958, July 1958-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

						· ·				
	July 1	957-June	1958	July 1	.958-June	1959				
Type of condition	Both sexes	Male	Female	Both sexes	Male	Female				
Age-45+	Number of acute conditions in thousands									
Total conditions	84,318	36,644	47,674	69,852	30,084	39,768				
Infectious and parasitic diseases- Jpper respiratory conditions	3,135 28,406	1,305 12,737	1,830 15,669	4,305 27,622	1,559 11,659	2,746 15,964				
Other respiratory conditions Digestive system conditions	27,961 4,595	12,586 1,696	15,375 2,899	14,088 3,902	6,054 2,056	8,034 1,846				
Injuries All other acute conditions	12,512 7,709	5,588 2,732	6,924 4,977	11,878 8,057	5,461 3,294	6,417 4,763				
Age-45+	Number o	f acute c	onditions	per 1,000 p	ersons pe	r year				
Total conditions	1,721.4	1,567.3	1,862.1	1,400.5	1,268.8	1,519.9				
Infectious and parasitic diseases- Jpper respiratory conditions	64.0 579.9	55.8 544.8	71.5 612.0	86.3 553.8	65.8 491.7	104.9 610.1				
Other respiratory conditions Digestive system conditions	570.8 93.8	538.3 72.5	600.5 113.2	282.5 78.2	255.3 86.7	307.1 70.6				
Injuries All other acute conditions	255.4 157.4	239.0 116.9	270.4 194.4	238.2 161.5	230.3 138.9	245.3 182.0				

NOTE: Since the estimates shown in this table are based on single years of data collection, the standard errors are 1.25 times the standard errors shown in Appendix 1.

Table 29. Bed-disability days associated with acute conditions and bed-disability days per person per year for persons 45 years and older by type of condition and sex: United States, July 1957-June 1958, July 1958-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

		~ 				
Type of condition	July 1	957 - June	1958	July 19	58-June 1	959
	Both sexes	Male	Female	Both sexes	Male	Female
Ages-45+	Nu	mber of b	ed-disabi	lity days in	thousands	
Total conditions	227,177	94,173	133,004	167,937	71,120	96,81
Infectious and parasitic diseases- Upper respiratory conditions	10,494 37,329	4,421 13,925	6,073 23,404	10,834 34,501	4,158 13,108	6,67 21,39
Other respiratory conditions Digestive system conditions	117,177 9,212	54,374 1,695	62,803 7,518	49,686 8,223	20,639 3,222	29,04 5,00
InjuriesAll other acute conditions	34,334 18,631	13,984 5,775	20,350 12,856	42,160 22,533	18,712 11,281	23,44 11,25
	Number	of bed-d	lisability	days per per	son per y	ear
Ages-45+						
Total conditions	4.6	4.0	5.2	3.4	3.0	3.
Infectious and parasitic diseases- Upper respiratory conditions	0.2	0.2	0.2 0.9	0.2 0.7	0.2	0
Other respiratory conditions Digestive system conditions	2.4	2.3 0.1	2.5 0.3	1.0 0.2	0.9 0.1	1. 0.
InjuriesAll other acute conditions	0.7 0.4	0.6 0.2	0.8 0.5	0.8 0.5	0.8 0.5	0.

NOTE: Since the estimates shown in this table are based on single years of data collection, the standard errors are 1.25 times the standard errors shown in Appendix I.

Table 30. Average annual incidence of injuries and number of bed-disability days associated with injuries for persons 45 years and older by type of injury and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

Type of injury		Age	
-, · · · ·	All ages 45+	45-64	65+
	Number o	of injuries i	n thousands
Total	12,195	8,762	3,433
Fractures, dislocations, sprains, and strainsOpen wounds and lacerations	4,798 1,753	3,405 1,410	1,393 (*)
Contusions and superficial injuriesOther current injuries	2,925 2,719		1,076 (*)
		of bed-days a with injurie	
Total	38,247	23,504	14,743
Fractures, dislocations, sprains, and strains Open wounds and lacerations	25,581 2,619	15,946 1,677	9,635 (*)
Contusions and superficial injuries Other current injuries	5,854 4,193	3,028 2,853	2,826 (*)

USE OF MEDICAL AND DENTAL SERVICES BY OLDER PERSONS

The utilization of health facilities, with particular emphasis on physician services, is considered at some length in this report. The data in this section provide information pertaining to the general use of medical services by older persons, as well as some insight into variations in level of use by income groups and urban and rural residence. The rates of physician visits for persons 45-64 years of age, shown in the detailed tables, are of particular importance because they provide a baseline from which to measure the increased need for medical care by persons 65 years and older. Hence, the rates may be useful in projecting future medical care needs of persons as they grow older.

Information on the volume and rate of physician and dental visits, with more detailed data by age groups, can be found in the following National Health Survey publications: Dental Care (Series B-Nos. 14 and 15), and Physician Visits (Series B-No. 19).

Physician Visits

A physician visit, in the National Health Survey, is defined as consultation with a physician, in person or by telephone, for examination, diagnosis, treatment, or advice. The visit was considered to be a physician visit if the service was provided directly by the physician or by a nurse or technician acting under a physician's super-




Figure 8. Number of physician visits per person per year for persons 65 years and older by residence and sex.

vision. In this definition "physician" includes doctors of medicine and osteopathic physicians. (For a more complete definition of physician visit, see Appendix II.)

1. Approximately 100 million physician visits were made during an average year by persons 65 years and older. This represents a rate of 6.8 physician visits per person per year.

2. The rate of physician visits was higher for females than for males in all residence categories, with the most significant sex differential in other urban places (fig. 8).

3. Persons 65 years and older living as member of families whose known income was less than \$2,000 had on the average 6.5 physician visits per year, while persons in this age group in families with known income of \$7,000 or more consulted a physician on the average 8.7 times per year. Physician visits in each of the income intervals were significantly higher for females than for males.

Table G. Percent distribution of physician visits by place of visit for persons 65 years and older: United States, July 1957-June 1959

	Physician	visits
Place of visit	Number (in thousands)	Percent distri- bution
Total visits	99,159	100.0
Office Home Hospital clinic Telephone Other and un- known (in- cluding indus-	59,731 22,595 6,765 7,141	60.2 22.8 6.8 7.2
try health unit)	2,928	3.0

4. Of the 6.8 average annual physician visits for persons 65 years and older, 4.1 were in the physician's office, 1.5 were in the home, and the remaining 1.2 visits in hospital clinic, industry health unit, in other and unknown places, or by telephone. The percent distribution of physician visits for persons 65 years and older is shown in table G.

5. Persons 65 years and older who reported no chronic conditions consulted a physician on the average of 2,2 times per year. Persons 65 years and older with one or more chronic conditions, but with no limitation of activity, had 5.3 physician visits per year, while persons with partial limitation of activity reported 8.5 visits per year, and those with major limitation of activity 14.3 visits per year (fig. 9). Within each of the activity limitation categories, the rate of physician visits was higher as the amount of family income increased.




Figure 9. Number of physician visits per person per year for persons 65 years and older by chronic condition status and degree of activity limitation.

6. Persons 65 years and older with one or more chronic conditions, but with no limitation of mobility had 6.6 physician visits per year. Those with partial limitation and those with major limitation had 11.4 and 18.9 physician visits, respectively. The average number of physician visits for persons with major limitation of mobility decreased appreciably with age, from 35.3 visits per person 45-54 years to 17.7 visits per person 75 years and older. This decrease was less marked among those with partial limitation of mobility, and the rate was quite stable among persons with no limitation (fig. 10).




Figure 10. Number of physician visits per person per year among persons with one or more chronic conditions by degree of limitation of mobility by age.

Dental Visits

The volume of dental visits is less sensitive than the number of physician visits as a measure of the use of health facilities by the older population because of the high proportion of edentulous persons in this segment of the population.

Table H. Number and percent of edentulous persons by age: United States, July 1957-June 1959

	Edentulous persons				
Age	Number (in thousands)	Percent of pop- ulation			
All ages 45+	18,830	38.1			
45-54 55-64 65+ 65-74 75+	4,433 5,685 8,712 5,369 3,343	22.4 38.1 59.4 55.4 67.3			

Data presented in National Health Survey Publication No. 15, Series B, showing volume of dental visits by type of service indicate that approximately one third of all visits made by persons 65 years and older were for denture work.

Each visit to a dentist's office for treatment or advice was considered to be a dental visit. The visit may have involved services provided directly by the dentist or by a dental hygienist acting under a dentist's supervision.

1. Persons 65 years and older made an average of 0.8 visits to the dentist per person per

year. This low average number of visits, compared with 1.5 visits per person per year in the total population (No. 15, Series B), is due to the high proportion (approximately 60 percent) of edentulous persons in the age group (table H).

2. Similar to the pattern for physician visits, the rate of dental visits for persons 65 years and older was progressively higher as the amount of family income increased.

DETAILED TABLES

			Page
		PHYSICIAN VISITS	
Table	31.	Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by residence, sex, and age: United States, July 1957-June 1959	49
	32.	Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959	50
	33.	Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by place of visit, sex, and age: United States, July 1957-June 1959	51
	34.	Average annual number of physician visits for persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959	52
	35.	Number of physician visits per person per year for persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959	53
	36.	Average annual number of physician visits for persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959	54
	37.	Number of physician visits per person per year for persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959	55
		DENTAL VISITS	
	38.	Average annual number of dental visits and number of dental visits per person per year for persons 45 years and older by residence, sex, and age: United States, July 1957-June 1959	56
-	39.	Average annual number of dental visits and number of dental visits per person per year for persons 45 years and older by family income and age: United States July 1957-June 1959	57

Table 31. Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by residence, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

			. Re	sidence			
	A11		Urban			Rural	
Sex and age	areas		Inside	Other			
. Dex and age	urcus	Total	urbanized	urban	. Total	Nonfarm	Farm
			areas	places	,		
	A	verage nu	mber of phy	sician v	isits in	thousand	s
Both sexes			1	I	1		
11 ages-45+	286,093	192,830	141,027	51,803	93,263	64,322	28,9
45-54	100,021	68,081	51,617	16,465	31,940	23,373	8,5
55-64	86,913	59,060	44,228	14,832	27,853	18,624	9,2
65+	99,159	65,689	45,183	20,506	33,471	22,325	11,1
65-74	62,794	42,392	29,733	12,659	20,402	13,259	7,1
75+	36,365	23,297	15,450	7,847	13,068	9,065	4,0
Male							
11 ages-45+	119,082	75,828	55,970	19,858	43,254	29,454	13,8
45-54	40,536	25,881	18,879	7,002	14,655	11,084	3,5
55-64	37,879	25,145	19,138	6,007	12,734	8,183	4,5
65+	40,667	24,802	17,952	6,849	15,865	10,187	5,6
65-74	26,371	16,458	12,268	4,190	9,913	6,254	3,6
75+	14,296	8,344	5,684	2,659	5,953	3,932	2,0
Female	14,250	0,544	3,004	2,035	5,555	3,752	2,0
11 ages-45+	167 011	117 002	85.057	31,945	50,009	34,868	15 1
-	167,011	1	85,057				15,1
45-54	59,486	42,201	32,738	9,463	17,285	12,289	4,9
55-64	49,034	33,915	25,089	8,825	15,119	10,441	4,6
65+	58,492	40,887	27,230	13,657	17,605	12,138	5,4
65-74	36,424	25,934	17,465	8,469	10,490	7,005	3,4
75+	22,069	14,953	9,765	l 5,187	7,116	5,133 l	1,9
	N	lumber of	nhysician y	isits pe	r person	per year	
			physician v	_	•		
Both sexes		1 .	I	٠ - ٥		 ! !	,
11 ages-45+	5.8	6.0	6.0	5.8	5.4	5.8	4
	5.8	1 .	I	5.8		 ! !	
11 ages-45+		6.0	6.0		5.4	5.8	3
11 ages-45+	5.0	6.0 5.3	6.0 5.4	4.9	5.4 4.6	5.8	
11 ages-45+ 45-54 55-64 65+	5.0 5.8 6.8	6.0 5.3 6.0 6.9	6.0 5.4 6.1 6.9	4.9 5.7 7.0	5.4 4.6 5.5 6.4	5.8 5.2 5.9 6.7	. 5
11 ages-45+ 45-54 55-64	5.0 5.8	6.0 5.3 6.0	6.0 5.4 6.1	4.9 5.7	5.4 4.6 5.5	5.8 5.2 5.9	
11 ages-45+ 45-54 55-64 65+ 65-74 75+ Male	5.0 5.8 6.8 6.5	5.3 6.0 6.9 6.7	6.0 5.4 6.1 6.9 6.7	4.9 5.7 7.0 6.7	5.4 4.6 5.5 6.4 6.1	5.8 5.2 5.9 6.7 6.2	
11 ages-45+ 45-54 55-64 65+ 65-74 75+ Male	5.0 5.8 6.8 6.5	5.3 6.0 6.9 6.7	6.0 5.4 6.1 6.9 6.7	4.9 5.7 7.0 6.7	5.4 4.6 5.5 6.4 6.1	5.8 5.2 5.9 6.7 6.2	3 5 6 5
11 ages-45+ 45-54 55-64 65-74 75+ Male 11 ages-45+ 45-54	5.0 5.8 6.8 6.5 7.3	6.0 5.3 6.0 6.9 6.7 7.5	6.0 5.4 6.1 6.9 6.7 7.4	4.9 5.7 7.0 6.7 7.5	5.4 4.6 5.5 6.4 6.1 7.1	5.8 5.2 5.9 6.7 6.2 7.6	3 5 6 5
11 ages-45+ 45-54 55-64 65+ 65-74 75+ <u>Male</u> 11 ages-45+	5.0 5.8 6.8 6.5 7.3	6.0 5.3 6.0 6.9 6.7 7.5	6.0 5.4 6.1 6.9 6.7 7.4 5.2	4.9 5.7 7.0 6.7 7.5	5.4 4.6 5.5 6.4 6.1 7.1	5.8 5.2 5.9 6.7 6.2 7.6	3 5 6 5 6
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5	4.9 5.7 7.0 6.7 7.5 4.9	5.4 4.6 5.5 6.4 6.1 7.1 5.0	5.8 5.2 5.9 6.7 6.2 7.6 5.4	3 5 6 5 6 4 2 4
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6	5.4 4.6 5.5 6.4 6.1 7.1 5.0 4.1 5.0 6.1	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3	2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2	5.4 4.6 5.5 6.4 6.1 7.1 5.0 4.1 5.0 6.1 5.8	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9	2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6	5.4 4.6 5.5 6.4 6.1 7.1 5.0 4.1 5.0 6.1	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3	2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 6.1 5.8 6.6	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0	
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 4.1 5.8 6.6	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0	
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 6.1 5.8 6.6 5.9	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 6.1 5.8 6.6 5.9	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 6.1 5.8 6.6 5.9	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
11 ages-45+	5.0 5.8 6.8 6.5 7.3 5.1 4.2 5.3 6.1 5.8 6.6	6.0 5.3 6.0 6.9 6.7 7.5 5.1 4.2 5.4 6.1 5.9 6.6	6.0 5.4 6.1 6.9 6.7 7.4 5.2 4.2 5.5 6.3 6.1 6.8 6.8	4.9 5.7 7.0 6.7 7.5 4.9 4.4 5.0 5.6 5.2 6.3	5.4 4.6 5.5 6.4 6.1 7.1 5.0 6.1 5.8 6.6 5.9	5.8 5.2 5.9 6.7 6.2 7.6 5.4 4.8 5.3 6.3 5.9 7.0 6.3	4 3 5 6 5 6 4 2 4 5 5 6 6 5 6 6 7 6 7 6 7 6 7 6 7 7 7 8 7 8 7 8 7 8

Table 32. Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by family income, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

Sou and and			Family	income		
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown
Both sexes	Average number of physician visits in thousa				ands	
All ages-45+	286,093	66,374	62,108	74,467	59,031	24,114
45-54	100,021	11,446	19,618	33,808	27,844	7,304
55-64	86,913	16,280	20,353	24,051	17,817	8,410
65+	99,159	38,648	22,136	16,607	13,369	8,400
65-74 75 +	62,794	23,007	15,854	10,972	8,042	4,919
• • •	36,365	15,641	6,282	5,635	5,327	3,480
Male				ļ		
All ages-45+	119,082	24,839	25,439	32,904	27,265	8,634
45-54	40,536	3,861	8,172	13,648	12,315	2,539
55-64	37,879	5,691	7,671	11,917	9,671	2,929
65-74	40,667	15,287	9,596	7,338	5,279	3,166
75+	26,371 14,296	8,333	7,125	5,484	3,541	1,887
/)	14,290	6,954	. 2,471	1,854	1,738	1,280
<u>Female</u>						
All ages-45+	167,011	41,535	36,669	41,563	31,765	15,480
45-54	59,486	7,585	11,446	20,160	15,529	4,765
55-64	49,034	10,589	12,683	12,134	8,146	5,482
65+	58,492	23,361	12,540 [.]	9,269	8,090	5,233
65-74	36,424	14,674	8,729	5,488	4,501	3,033
75 +	22,069	8,687	3,811	3,781	3,589	2,201
Both sexes	Nu	mber of ph	ysician vi	sits per p	erson per	year
All ages-45+	5.8	5.8	5.8	5.6	6.1	5.4
45-54	5.0	4.7		E 1	5 2	, 0
55-64	5.8	5.4	5.0 5.9	5.1 5.8	5.2 6.4	4.9 5.6
65+	6.8	6.5	6.6	6.9	8.7	5.9
65-74	6.5	6.3	6.4	6.7	7.9	5.4
75 +	7.3	6.9	7.0	7.3	10.5	6.7
	·					
<u>Male</u>						
All ages-45+	5.1	.5.2	5.0	4.9	5.5	4.5
45-54	4.2	3.9	4.5	4.0	4.5	3.8
55-64	5.3	4.8	4.8	5.5	6.1	4.2
65+	6.1	5.9	5.6	6.6	7.7	5.5
65-74 75+	5.8	5.3	5.6	7.0	7.1	4.8
, , ,	6.6	6.8	5.7	5.7	9.1	7.0
<u>Female</u>						
All ages-45+	6.5	6,3	6.5	6.4	6.8	6.2
45-54	5.9	5.2	5.5	6.3	6.0	5.7
55-64	6.3	5.8	6.7	6.1	6.6	6.7
65+	7.3	7.0	7.6	7.1	9.6	6.1
65-74	7.0	7.1	7.3	6.4	8.5	5.8
/	7.8	6.9	8.3	8.5	11.3	6.6

Table 33. Average annual number of physician visits and number of physician visits per person per year for persons 45 years and older by place of visit, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

			P	lace of vi	sit.		
Sex and age	Total	Office	Home	Hospital clinic	Company or industry health unit	Telephone	Other and unknown
Both sexes		Average	number o	f physicia	n visits i	n thousands	•
All ages-45+	286,093	192,538	39,486	24,626	3,253	17,445	8,745
45-54 55-64	100,021 86,913	71,684 61,122	7,982 8,909	9,678 8,184	1,826 1,188	5,385 4,919	3,466 2,590
65+ 65-74 75+	99,159 62,794 36,365	59,731 41,613 18,118	22,595 10,427 12,168	6,765 4,378 2,387	(*) (*) (*)	7,141 4,612 2,529	2,690 1,526 1,164
<u>Male</u>		·					-
All ages-45+	119,082	80,417	14,380	11,214	2,500	6,719	3,852
45-54 55-64 65+ 65-74 75+	40,536 37,879 40,667 26,371 14,296	28,868 26,260 25,288 18,005 7,283	3,087 3,369 7,924 3,698 4,226	4,153 3,650 3,411 1,999 1,411	1,360 1,012 (*) (*) (*)	1,855 2,108 2,756 1,758 998	1,212 1,480 1,160 (*) (*)
<u>Female</u> All ages-45+	167,011	112 120	25,107	13,413	(*)	10,726	4,894
45-54 55-64	59,486	42,816 34,862	4,896 5,541	5,525 4,534	(*)	3,530 2,811	2,253 1,110
65+ 65-74	49,034 58,492 36,424	34,442 23,608	14,670 6,728	3,354 2,379	(*) (*)	4,385 2,854	1,530 (*)
75+	22,069	_	-			n per year	l (*)
<u>Both sexes</u> All ages-45+	5.8	3.9	0.8	0.5	0.1	0.4	0.2
45-54 55-64 65+ 65-74	5.0 5.8 6.8 6.5	3.6 4.1 4.1 4.3	0.4 0.6 1.5	0.5 0.5 0.5 0.5	0.1 0.1 (*) (*)	0.3 0.3 0.5 0.5	0.2 0.2 0.2 0.2
75+ Male	7.3	3.6	2.4	0.5	(*)	0.5	0.2
All ages-45+	5.1	3.4	0.6	0.5	0.1	0.3	0.2
45-54 55-64 65+ 65-74 75+	4.2 5.3 6.1 5.8 6.6	3.0 3.7 3.8 4.0 3.4	0.3 0.5 1.2 0.8 2.0	0.4 0.5 0.5 0.4 0.7	0.1 0.1 (*) (*) (*)	0.2 0.3 0.4 0.4 0.5	0.1 0.2 0.2 (*) (*)
Female ,							
All ages-45+	6.5	4.3	1.0	0.5	(*)	0.4	0.2
45-54 55-64 65+ 65-74	5.9 6.3 7.3 7.0	4.2 4.5 4.3 4.6	0.5 0.7 1.8 1.3	0.5 0.6 0.4 0.5	(*) (*) (*) (*)	0.3 0.4 0.5 0.6	0.2 0.1 0.2 (*)
75+	7.8	3.9	2.8	(*)	(*)	0.5	(*)

Table 34. Average annual number of physician visits for persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

	m-1-1	Physician visits	Physicia		or persons	with 1+		
Age and family income	Total physician visits	for persons with no			chronic conditions Limitation of activity			
	V13123	chronic condition	Total	None	Partial	Major		
	Ave	rage number	of physic	ion vieite	in thouse	nde		
All ages-45+	1		or physic	10. 120100	III CHOUSE			
		ı	1	1	!			
Total	286,093	38,211	247,882	112,848	85,845	49,189		
Under \$2,000	66,374	4,822	61,552	18,769	25,118	17,666		
\$2,000-3,999	62,108	8,032	54,075	23,751	19,264	11,060		
\$4,000-6,999	74,467	12,163	62,304	33,131	19,241	9,932		
\$7,000+ Unknown	59,031	9,858	49,172	27,547	15,399	6,226		
	24,114	3,336	20,778	9,650	6,823	4,304		
<u>45-54</u>								
Total	100,021	20,141	79,880	49,158	23,828	6,894		
Under \$2,000	11,446	1,531	9,916	3,562	4,757	1,597		
\$2,000-3,999	19,618	4,180	15,438	8,442	5,322	1,674		
\$4,000-6,999	33,808	7,491	26,318	17,862	6,496	1,960		
\$7,000+	27,844	5,608	22,237	15,400	5,718	1,119		
Unknown	7,304	1,332	5,972	3,892	1,535	(*)		
<u>55-64</u>								
Total	86,913	10,916	75,997	36,548	27,149	12,301		
Under \$2,000	16,280	1,182	15,098	5,241	6,121	3,736		
\$2,000-3,999	20,353	2,245	18,109	8,753	6,099	3,257		
\$4,000-6,999	24,051.	3,120	20,932	11,218	6,827	2,887		
\$7,000+	17,817	3,261	14,556	8,305	5,366	(*)		
Unknown	8,410	1,108	7,302	3,031	2,737	1,534		
<u>65+</u>								
Total	99,159	7,155	92,004	27,143	34,868	29,994		
Under \$2,000	38,648	2,110	36,538	9,966	14,239	12,333		
\$2,000-3,999	22,136	1,608	20,529	6,556	7,844	6,129		
\$4,000-6,999 \$7,000+	16,607	1,552	15,054	4,052	5,918	5,085		
\$7,000+ Unknown	13,369	(*)	12,379	3,842	4,316	4,221		
	8,400	(*)	7,504	2,727	2,551	2,225		
<u>65-74</u>]				}			
Total	62,794	5,235	57,559	19,609	23,582	14,368		
Under \$2,000	23,007	1,474	21,533	6,807	9,029	5,698		
\$2,000-3,999	15,854	1,309	14,546	5,221	6,102	3,223		
\$4,000-6,999	10,972	1,106	9,866	3,197	3,940	2,729		
\$7,000+	8,042	(*)	7,324	2,676	2,821	1,827		
Unknown	4,919	(*)	4,291	1,708	1,691	(*)		
<u>75+</u>	l							
Total	36,365	1,920	34,445	7,534	11,286	15,625		
Under \$2,000	15,641	(*)	15,005	3,159	5,211	6,635		
\$2,000-3,999	6,282	(*)	5,983	1,335	1,742	2,906		
\$4,000-6,999	5,635	(*)	5,189	855	1,978	2,356		
\$7,000+	5,327	(*)	5,056	1,166	1,495	2,395		
Unknown	3,480	(*)	3,213	1,019	(*)	1,334		
					L			

Table 35. Number of physician visits per person per year for persons 45 years and older according to chronic limitation of activity by age and family income: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

estimates are given in Appendix I. Det	initions of te	erms are giver	ın Appendi	× 11]			
(Total	Physician visits for	Physician visits for persons with 1+ chronic conditions				
Age and family income	physician visits	persons with no chronic	Total	Limitation of activity			
		condition		None	Partial	Major	
	Num	ber of phys	ician vis	its per per	son per ye	ar	
A11 ages-45+		I	I	!		ı	
Total	5.8	2.2	7.7	5.6	9.7	16.2	
Under \$2,000	5.8	1.8	7.1	4.5	7.9	13.3	
\$2,000-3,999	5.8	2.2	7.6	5.4	9.4	17.0	
\$4,000-6,999	5.6	2.3	7.8	5.8	10.5	19.2	
\$7,000+	6.1	2.5	8.6		13.5	23.2	
Unknown	5.4	1.8	8.0	5.9	10.2	15.2	
<u>45-54</u>						·	
Total	5.0	2.3	7.1	5.7	10.8	22.3	
Under \$2,000	4.7	1.7	6.3	3.7	9.8	14.5	
\$2,000-3,999	5.0	2.6	6.8	5.1	10.0		
\$4,000-6,999	5.1	2.5	7.1	6.0	10.3	32.7	
\$7,000+	5.2	2.3	7.6	6.3	13.4	33.9	
Unknown	4.9	1.8	8.0	6.6	11.5	(*)	
	"''	1.0	0.0	""	11.5	(")	
55-64 Total			7.0		10.6		
	5.8	2.1	7.9	5.7	10.6	19.4	
Under \$2,000	5.4	1.5	6.8	4.4	7.8	14.5	
\$2,000-3,999	5.9	1.9	7.9	5.8	9.8	20.2	
\$4,000-6,999	5.8	2.0	8.2	6.1	11.4	24.3	
\$7,000+	6.4	2.9	8.7	6.4	16.0	(*)	
Unknown	5.6	1.7	8.5	5.2	12.8	26.0	
<u>65+</u>		•	,				
Total	6.8	2.2	8.1	5.3	8.5	1/- 2	
	0.8	2.2	0.1		. 0.0	14.3	
Under \$2,000	6.5	2.0	7.5	4.9	7.5	12.9	
\$2,000-3,999 \$4,000-6,999	6.6	2.0	8.0	5.3	8.7	14.9	
\$4,000-6,999	6.9	2.5	. 8.4	4.8	9.8	15.0	
\$7,000+	8.7	(*)	11.0	6.9	11.4	21.6	
Unknown	5.9	(*)	7.6	5.7	8.0	11.4	
<u>65-74</u>							
Total	6.5	2.1	8.0	5.2	9.3	15.7	
Under \$2,000	6.3	1.9	7.4	5.0	8.2	13.7	
\$2,000-3,999	6.4	2.0	8.0	5.2	9.8	15.6	
\$4,000-6,999	6.7	2.3	8.4	5.0	9.8	20.4	
\$7,000+	7.9	(*)	10.1	6.5	12.3	22.0	
Unknown	5,4	(*)	7.3	5.3	8.9	. (*)	
<u>75+</u>	* **						
Total	7.3	2.3	8.3	5.4	7.2	13.2	
Under \$2,000	6.9	(*)	7.6	4.9	6.5	12.3	
\$2,000-3,999	7.0	(*)	8.2	5.4	6.2	14.1	
\$4,000-6,999	7.3	(*)	8.5	4.1	10.0	11.5	
\$7,000+	10.5	(*)	12.5	8.0	10.1	21.2	
Unknown	6.7	(*)	8.0	6.6	(*)	10.9	
V	k	(")	٠.٠	· · · · · · · ·	· · · · · · · ·	10.9	
the state of the s						<u> </u>	

NOTE: Rates shown in this table are based on numbers of persons shown in table 5.

Table 36. Average annual number of physician visits for persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

Under \$2,000	are given in Appendix in							
Visits	Ass and faully drawns		visits for	Physicia				
Average number of physician visits in thousands All ages-45+	age and family income		with no	Total	Limita	tion of mo	bility	
Total					None	Partial	Major	
Note Note		Ave	rage number	of physic	ian visits	in thousa	nds	
Total	All ages-45+			[,	v			
Under \$2,000		200 0001	20 211	1 000 000	100 005			
\$2,000-3,999	Total	286,093	38,211	247,882	190,285	37,131	20,466	
\$2,000-3,999		66,374	4,822	61,552	40,902	13,864	6,786	
\$7,000+	\$2,000-3,999			54,075	41,296	8,502	4,277	
Unknown 24,114 3,336 20,778 15,675 2,675 2 Tota1 100,021 20,141 79,880 71,376 5,466 3 Under \$2,000 11,446 1,531 9,916 7,804 1,444 \$2,000-3,999 19,618 4,180 15,438 13,319 1,324 \$4,000-6,999 33,808 7,491 26,318 24,081 1,207 1 \$7,000+ 27,844 5,608 22,237 20,747 1,101 1 Unknown 7,304 1,332 5,972 5,424 (*) Tota1 86,913 10,916 75,997 61,932 8,743 5, Under \$2,000 16,280 1,182 15,098 10,731 2,673 1, \$2,000-3,999 20,353 2,245 18,109 14,914 2,382 \$4,000-6,999 24,051 3,260 20,932 17,367 2,075 1, \$7,000+ 17,817 3,261 14,556 13,365 1,038 Under \$2,000-3,999 22,136 1,608 20,529 13,063 4,796 2, \$7,000+ 13,369 (*) 12,379 7,908 2,242 2	\$4,000-6,999						4,205	
Total	\$7,000+						2,770	
Total	Unknown	24,114	3,336	20,778	15,675	2,675	2,428	
Under \$2,000								
\$2,000-3,999	Tota1	100,021	20,141	79,880	71,376	5,466	3,039	
\$2,000-3,999	Under \$2 000	11 //6	1 521	0.016	7 904	1 444	(*)	
\$4,000-6,999	\$2 000-3 999	10,440		15 / 30	12 210		1 1	
Total	\$4,000-5,999						(*) 1,030	
Unknown	\$7 000+						(*)	
55-64 Total	Inknown						(*)	
Under \$2,000		7,304	1,552	3,,,,	3,424			
Under \$2,000	Total	86 013	10 916	75 007	61 032	9 7/2	5,322	
\$2,000-3,999	1	00,913	10,910	73,337	01,932	0,743	3,322	
\$4,000-6,999	Under \$2,000	16,280	1,182	15,098	10,731	2,673	1,694	
Total	\$2,000-3,999		2,245	18,109	14,914	2,382	(*)	
Total	\$4,000-6,999	24,051	3,120	20,932	17,367	2,075	1,489	
Total	\$7,000+						, (*)	
Under \$2,000	Unknown	8,410	1,108	7,302	-5,555	(*)	1,173	
Under \$2,000	<u>65+</u>		. ,					
\$2,000-3,999	Total	99,159	7,155	92,004	56,977	22,922	12,105	
\$2,000-3,999	Under \$2.000	38,648	2.110	36,538	22,367	9.748	4,424	
\$4,000-6,999	\$2,000-3,999						2,670	
\$7,000+	\$4,000-6,999		1,552			4,425	1,686	
Unknown	\$7,000+		(*)			2,242	2,229	
Total	Unknown	8,400	(*)	7,504	4,696	1,711	1,097	
Under \$2,000	<u>65-74</u>			a [.]				
\$2,000-3,999	Total	62,794	5,235	57,559	40,569	12,353	4,638	
\$2,000-3,999	Hadow \$2 000	22 007	1 474	21 522	1/ 050	5 200	1 /70	
\$4,000-6,999	07 000-2 000						1,470	
Unknown 4,919 (*) 4,291 3,153 (*)	\$4 000-5,999						1,327 (*)	
Unknown 4,919 (*) 4,291 3,153 (*)	\$7 0004						(*) (*)	
75+	Unknown						(*) (*)	
	<u>75+</u>							
Total 36,365 1,920 34,445 16,408 10,569 7,	Total	36,365	1,920	34,445	16,408	10,569	7,468	
	Under \$2,000	15-641	(*)	15,005	7 509	4.542	2,954	
\$2,000-3,999	\$2,000-3,999				2.778		1,343	
\$4,000-6,999 5,635 (*) 5,189 2,339 2,061	\$4.000-6.999						(*)	
\$7,000+ 5,327 (*) 5,056 2,240 1,243 1.	\$7,000+						1,572	
Unknown 3,480 (*) 3,213 1,543 (*)	Unknown						(*)	

Table 37. Number of physician visits per person per year for persons 45 years and older according to chronic limitation of mobility by age and family income: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civillan noninstitutional pop-, ulation of the United States. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

Age and family income	Total physician	Physician visits for persons	T	n visits f	or persons	with 1+
inge und rumriy, ricome	visits	with no	Total	Limita	tion of mo	bility
		chronic condition		None	Partial	Major
	Num	ber of phys	ician visi	ts per per	son per ye	ar
A11 ages-45+		• •		• •		
Total	5.8	2.2	7.7	6.8	12.2	22.3
Under \$2,000	5.8	1.8	7.1	5.9	10.3	18.1
\$2,000-3,999	5.8	2.2	7.6	6.5	13.7	22.9
\$4,000-6,999	5.6	2.3	7.8	6.9	14.4	25.2
\$7,000+ Unknown	6.1	2.5	8.6	7.9	14.4	31.5
Unknown	5.4	1.8	8.0	7.0	11.2	23.8
<u>45-54</u>	·					
Total	5.0	2.3	7.1	6.6	15.6	35.3
Under \$2,000	4.7	1.7	6.3	5.5	12.0	(*)
\$2,000-3,999	5.0	2.6	6.8	6.1	16.8	(*)
\$4,000-6,999	5.1	2.5	7.1	6.7	13.6	42.9
\$7,000+	5.2	2.3	7.6	7.2	24.5	(*)
Unknown	4.9	1.8	8.0	7.5	(*)	(*)
<u>55-64</u>						
Total	5.8	2.1	7.9.	7.1	13.0	27.9
Under \$2,000	5.4	1.5	6.8	5.7	10.0	22.0
\$2,000-3,999	5.9	1.9	7.9	7.2	15.2	(*)
\$4,000-6,999	5.8	2.0	8.2	7.2	15.4	4 2. 5
\$7,000+	. 6.4	2.9	8.7	8.3	15.5	(*)
Unknown	5.6	1.7	8.5	7.1	(*)	48.9
<u>65+</u>						
Tota1	6.8	2.2	8.1	6.6	11.4	18.9
Under \$2,000	6.5	2.0	7.5	6.1	10.2	16.4
\$2,000-3,999	6.6	2.0	8.0	6.4	12.5	21.9
\$4,000-6,999 \$7,000+	6.9	2.5	8.4	6.5	14.2	15.8
\$7,000+	8.7	(*)	11.0	9.1	11.7	32.3
• '	5.9	(*)	7.6	6.3	10.1	15.0
<u>65-74</u>						
Total	. 6.5	2,1	8.0	6.7	13.4	21.2
Under \$2,000	6.3	1.9	7 /	6.3	12.2	16.7
\$2,000-3,999	6.4	2.0	7.4 8.0	6.6	14.5	26.0
\$4,000-6,999	6.7	2.3	8.4	6.6	17.6	(*)
\$7,000+	7.9	(*)	10.1	9.2	(*)	(*)
Unknown	5.4	(*)	7.3	6.4	(*)	(*)
<u>75+</u>						
Total	7.3	2.3	8.3	6.2	9.7	17.7
Under \$2,000	6.9	(*)	7.6	5.9	8.6	16.1
\$2,000-3,999	7.0	(*)	8.2	5.8	10.2	18.9
\$4,000-6,999	7.3	(*)	8.5	6.4	11.6	(*)
\$7,000+ Unknown	10.5 6.7	(*) (*)	12.5 8.0	8.9 6.1	12.0 (*)	32.1 (*)
OIM: IOMIT	0.7	(")	(0.0)	0.1	(~)	(-)

Table 38. Average annual number of dental visits and number of dental visits per person per year for persons 45 years and older by residence, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

				Residenc	e		
Sex and age			Urban	•		Rural	
	All areas	Total	Inside urbanized areas	Other urban places	Total	Nonfarm	Farm
Both sexes	Av	erage nu	mber of den	tal visi	ts in th	ousands	
All ages-45+	63,979	48,564	38,051	10,512	15,415	11,029	4,386
45-6465 +	52,341 11,637	39,665 8,898	31,687 6,364		12,676 2,739	9,219 1,810	3,457 (*)
Male							
All ages-45+	26,859	20,472	15,843	4,629	6,387	4,434	1,952
45-6465+	21,833 5,025	16,645 3,827	13,227 2,616		5,188 1,199	3,612 (*)	1,576 (*)
Female							
All ages-45+	37,120	28,092	22,208	5,883	9,029	6,595	2,434
45-6465+	30,508 6,612	23,020 5,071	18,460 3,748	4,560 1,323		5,607 (*)	1,882 (*)
Both sexes	N	lumber of	dental vis	its per	person p	er year	
All ages-45+	1.3	1.5	1.6	1.2	0.9	1.0	0.7
45-6465+	1.5 0.8	1.7 0.9	1.9 1.0	1.3 0.9	1.1 0.5	1.2 0.5	0.8 (*)
<u>Male</u>							
All ages-45+	1.1	1.4	1.5	1.1	0.7	0.8	0.6
45-64 65+	1.3 0.8	1.5 0.9	1.7 0.9	1.2 1.0	0.9	0.9 (*)	0.7 (*)
Female			·				
All ages-45+	1.4	1.6	1.8	1.2	1.1	1.2	0.8
45-64 65+	1.7 0.8	1.9 0.9	2.1 1.0	1.4	1.3 0.6	1.5 (*)	0.9 (*)

Table 39. Average annual number of dental visits and number of dental visits per person per year for persons 45 years and older by family income and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix 1.

		Family income										
Age	Total	Total Under \$2,000		\$4,000- 6,999	\$7,000+	Unknown						
	Average number of dental visits in thousands											
All ages-45+	63,979	63,979 7,798 10,775 19,344 21,461										
45-64	52,341	4,564	7,787	17.014	19,769	3,208						
65+	11,637	3,234	2,988	2,329	1,692	i,393						
	N	umber of d	ental visi	ts per per	son per ye	ar						
All ages-45+	1.3	0.7	1.0	1.5	2.2	1.0						
45-64	1.5	0.8	1.1	1.6	2.4	1.1						
65+	0.8	0.5	0.9	1.0	1.1	1.0						

POPULATION |

Tables 40-44 contain estimates of the civilian noninstitutional population of the United States. Population estimates in table 40 are for the periods July 1957-June 1958 and July 1958-June 1959 and are appropriate to use for computing rates pertaining to acute conditions, shown separately for the 2 years in this report (text table E and detailed tables 28 and 29). Tables 41-44 show population estimates for the period July 1957-

June 1959 by residence, major activity, and family income.

These population estimates are based on interviews conducted by the National Health Survey, and are not official population estimates. For official estimates for more general use, see Bureau of the Census publications: Current Population Reports, Series P-20, P-25, P-50, P-57, and P-60.

DETAILED TABLES

		Page
40.	Population used in obtaining rates shown in this publication by sex and age: United States, July 1957-June 1958, July 1958-June 1959	59
41.	Population used in obtaining rates shown in this publication by residence, sex, and age: United States, July 1957-June 1959	59
42.	Population used in obtaining rates shown in this publication by major activity, sex, and age: United States, July 1957-June 1959	60
43.	Population used in obtaining rates shown in this publication by family income, sex, and age: United States, July 1957-June 1959	61
44.	Population of "usually working" persons used in obtaining rates shown in this publication by family income, sex, and age: United States, July 1957-June 1959	62
	41. 42. 43.	 41. Population used in obtaining rates shown in this publication by residence, sex, and age: United States, July 1957-June 1959 42. Population used in obtaining rates shown in this publication by major activity, sex, and age: United States, July 1957-June 1959 43. Population used in obtaining rates shown in this publication by family income, sex, and age: United States, July 1957-June 1959 44. Population of "usually working" persons used in obtaining rates shown in this publication by family income, sex, and age: United

Table 40. Population used in obtaining rates shown in this publication by sex and age: United States, July 1957-June 1958, July 1958-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

		Sex	
Age	Both sexes	Male	Female
July 1957-June 1958	Popul	ation in thousa	nds
All ages-45+	48,983	23,380	25,602
.5-64 5+	34,470 14,512	16,739 6,641	17,731 7,871
July 1958-June 1959			
All ages-45+	49,876	23,711	26,165
.5-64 5+	35,055 14,821	16,978 6,733	18,077 8,088

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States, in <u>Current Population Reports</u>: Series P-20, P-25, P-57, and P-60.

Table 41. Population used in obtaining rates shown in this publication by residence, sex, and age: United States, July 1957-June 1959

		(See head	note on table	401		·					
,		<u> </u>	. F	Residence							
	.:		Urban			Rural					
Sex and age	All areas	Total	Total Inside urbanized areas		Total	Nonfarm	Farm				
Both sexes	Population in thousands										
All ages-45+	49,429	32,311	23,415	8,895	17,118	11,023	6,096				
45-54 55-64 65+ 65-74 75+	19,833 14,930 14,667 9,698 4,969	12,944 9,910 9,457 6,330 3,127	9,580 7,303 6,532 4,451 2,081	3,364 2,607 2,924 1,878 1,046	6,888 5,020 5,210 3,368 1,842	4,501 3,173 3,348 2,155 1,193	2,387 1,847 1,862 1,213 649				
<u>Male</u> All ages-45+	23,545	14,863	10,831	4,032	8,683	5,454	3,228				
45-54 55-64 65+ 65-74 75+	9,675 7,183 6,687 4,530 2,157	6,142 4,652 4,069 2,808 1,261	4,539 3,452 2,839 2,002 837	1,603 1,199 1,230 806 424	3,533 2,531 2,618 1,723 896	2,294 1,543 1,617 1,059 558	1,239 988 1,001 664 338				
<u>Female</u> All ages-45+	^{25,884}	17,448	12,585	4,863	8,436	5,569	2,867				
45-54 55-64 65+ 65-74	10,157 7,747 7,980 5,167 2,812	6,802 5,258 5,388 3,522 1,866	5,041 3,851 3,693 2,449 1,244	1,761 1,408 1,695 1,072 622	3,355 2,489 2,592 1,645 946	2,207 1,630 1,732 1,097 635	1,148 859 860 549 311				

Table 42. Population used in obtaining rates shown in this publication by major activity, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

		Major activ	ity (45 year	s and older)	
Sex and age	Total	Usually working	Keeping house	Retired	Other
Both sexes		Popul	ation in tho	usands	
Both sexes		, !	ı	ı	1
All ages-45+	49,429	24,163	16,863	6,134	2,269
45-54	19,833	13,083	5,907	0.2	750
55-64	14,930	8,369	5,907 5,134	83 678	759 750
65+	14,666	2,711	5,134 5,822	5,373	750 760
65-74	9,698	2,312	4,014	2,973	398
75+	4,969	399	1,808	2,400	362
	1,505		1,000	-,-00	
Male	:				•
All ages-45+	23,545	17,178	(*)	4,869	1,471
45-54	0.675	0.000	(45)	,	5.0
55-64	9,675 7,183	9,036 6,047	(*) (*)	67 562	562
65+	6,687	2,095	(*)	4,240 /	568 341
65-74	4,530	1,774	(*)	2,510	240
75+	2,157	321	(*) (*)	1,730	101
Female	,	•,			
All ages-45+	25,884	6,985	16,836	1,265	798
45-54	10,157	4,047	5,897	5 17	197
55-64	7,747	2,322	5,129	115	181
65+	7,980	617	5,811	1,133	420
65-74	5,167	538	4,008	463	158
75+	2,812	79	1,803	669	261
				§ 1	± .*

Table 43. Population used in obtaining rates shown in this publication by family income, sex, and age: United States, July 1957-June 1959

[Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

Garage A. and		_	Family	income								
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown						
	Population in thousands											
Both sexes												
All ages-45+	49,429	11,394	10,713	13,218	9,666	4,438						
45-54	19.833	2,452	3,886	6.662	F 220	1 500						
55-64	14,930	3,004	3,666	6,662	5,330	1,503						
65+	14,667	5,938	3,356	4,144 2,412	2,804	1,507						
65-74	9,698	3,657	2,463	1,643	1,532 1,024	1,428 912						
75+	4,969	2,281	893	769	508	517						
<u>Male</u>												
All ages-45+	23,545	4,781	5,112	6,718	4,998	1,935						
45-54	9,675	999	1,818	3,451	2,738	670						
55-64	7,183	1,176	1,588	2,152	1,573	693						
65+	6,687	2,606	1,707	1,115	687	572						
65-74	4,530	1,584	1,272	788	496	390						
75+	2,157	1,022	435	327	192	182						
<u>Female</u>												
All ages-45+	25,884	6,612	5,601	6,501	4,668	2,502						
45-54	10,157	1,453	2,068	3,212	2,592	832						
55-64	7,747	1,828	1,883	1,992	1,231	814						
65+	7,980	3,332	1,649	1,297	845	856						
65-74	5,167	2,073	1,191	855	528	521						
75+	2,812	1,259	459	443	317	335						

Table 44. Population of "usually working" persons used in obtaining rates shown in this publication by family income, sex, and age: United States, July 1957-June 1959

Data are based on household interviews during July 1957-June 1959. Data refer to the civilian noninstitutional population of the United States. Detailed figures may not add to totals due to rounding. The survey design, general qualifications, and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

	Family income												
Sex and age	Total	Under \$2,000	\$2,000- 3,999	\$4,000- 6,999	\$7,000+	Unknown							
Both sexes	Population in thousands												
All ages-45+	24,163	3,161	5,097	7,802	6,118	1,985							
45-54 55-64 65+ 65-74	13,083 8,369 2,711 2,312 399	1,230 1,166 764 622 142	2,462 1,934 701 603 97	4,640 2,605 557 489 68	3,822 1,879 417 366 50	929 784 273 232 41							
<u>Male</u>	· .					. •							
All ages-45+	17,178	1,943	3,426	5,816	4,546	1,447							
45-54 55-64 65+ 65-74 75+	9,036 6,047 2,095 1,774 321	725 692 526 418 108	1,619 1,282 525 444 81	3,372 1,984 460 405 55	2,694 1,488 365 319 46	627 601 219 188 31							
<u>Female</u>													
All ages-45+	6,985	1,218	1,671	1,986	1,572	539							
45-54 55-64 65+ 65-74	4,047 2,322 617 538 79	506 474 238 204 35	843 652 176 160 16	1,267 621 97 84 13	1,129 391 52 47 5	302 183 54 44 11							

APPENDIX I

TECHNICAL NOTES ON METHODS

Background of This Report

This report on <u>Selected Health Characteristics of Older Persons</u> is one of a series of statistical reports prepared by the U. S. National Health Survey. The report is based on information collected in a continuing nationwide sample of households in the Health Interview Survey, which is a main aspect of the program.

The Health Interview Survey uses a questionnaire which, in addition to personal and demographic characteristics, collects information on illnesses, injuries, chronic conditions, medical care, dental care, and hospitalization. As interview data relating to each of these various broad subject areas are tabulated and analyzed, separate reports are issued covering one or more specific topics. The present report is based on the consolidated sample for 104 weeks of interviewing during the period July 1957-June 1959.

The population covered by the sample for the Health Interview Survey is the civilian noninstitutional population of the United States living at the time of the interview. The sample does not include members of the Armed Forces, United States nationals living in foreign countries, and crews of vessels.

Statistical Design of the Health Interview Survey

General plan.—The sampling plan of the survey follows a multistage probability design which permits a continuous sampling of the civilian population of the United States. The first stage of this design consists of drawing a sample of 500 from the 1,900 geographically defined Primary Sampling Units (PSU's) into which the United States has been divided. A PSU is a county, a group of contiguous counties, or a Standard Metropolitan Statistical Area.

With no loss in general understanding, the remaining stages can be telescoped and treated in this discussion as an ultimate stage. Within PSU's, then, ultimate stage units called segments are defined, also geographically, in such a manner that each segment contains an expected six households in the sample. Each week a random sample of about 120 segments is drawn. In the approximately 700 households in those segments persons are interviewed concerning illnesses, injuries, chronic conditions, disability, and other factors related to health.

The household members included each week are a representative sample of the population so that samples for successive weeks can be combined into larger samples for, say a calendar quarter, or a year. Thus the design permits both continuous measurement of characteristics of high incidence or prevalence in the population, and through the larger consolidated samples, more detailed analysis of less common characteristics and smaller categories. The continuous collection has administrative and operational advantages,

as well as technical assets, since it permits field work to be handled with an experienced, stable staff.

Sample size and geographic detail.—Over the 24-month period ending June 1959, the sample included approximately 235,000 persons from 73,000 households in 12,200 segments. The over-all sample was designed in such a fashion that tabulations can be provided for various geographic sections of the United States and for urban and rural sectors of the Nation.

Collection of data.—The field operations for the household survey are performed by the Bureau of the Census under specifications established by the Public Health Service. In accordance with these specifications the Bureau of the Census designs and selects the sample, conducts the field interviewing acting as collecting agent for the Public Health Service, and edits and codes the questionnaires. Tabulations are prepared by the Public Health Service using the Bureau of the Census electronic computers.

Estimating methods.—Each statistic produced by the survey—for example, the number of work-loss days occurring in a specified period—is the result of two stages of ratio estimation. In the first of these, the ratio factor is the 1950 decennial population count to the estimated population for 1950 for the U.S. National Health Survey's first-stage sample of PSU's. These factors are applied for more than 50 color-residence classes.

Later, ratios of sample-produced estimates of the population to official Bureau of the Census figures for current population in about 60 age-sex-color classes are computed, and serve as second-stage factors for ratio estimating.

The effect of the ratio estimating process is to make the sample more closely representative of the population by age, sex, color, and residence, thus reducing sampling variance.

As noted, each week's sample represents the population living during that week and characteristics of that population. Consolidation of samples over a time period, say a calendar quarter, produces estimates of average characteristics of the U.S. population for that calendar quarter. Similarly, population or prevalence data for a 12-year period are averages of the eight quarterly figures.

For statistics measuring the number of occurrences during a specified time period, such as number of bed-disability days, a similar computational procedure is used, but the statistics have a different interpretation. For the disability-day items, the questionnaire asks for the respondent's experience over the two calendar weeks prior to the week of interview. In such instances the estimated quarterly total for the statistic is simply 6.5 times the average two-week estimate produced by the 13 successive samples taken during the period. The annual total is the sum of the four quarters. Thus, the experience of persons interviewed during a year—experience which actually occurred for each person in a two-calendar-week interval prior to week of

interview—is treated in analysis as though it measured the total of such experience occurring in the year. Such interpretation leads to no significant bias.

General Qualifications

Nonresponse.—Data were adjusted for nonresponse by a procedure which imputed to persons in a household not interviewed the characteristics of interviewed persons in the same segment. The total noninterview rate was 5 percent; 1 percent was refusal, and the remainder was primarily due to the failure to find any eligible household respondent after repeated trials.

The interview process.—The statistics presented in this report are based on replies secured in interviews of persons in the sampled households. Each person 18 years and over, available at the time of interview, was interviewed individually. Proxy respondents within the household were employed for children and for adults not available at the time of the interview, provided the respondent was closely related to the person about whom information was being obtained.

There are limitations to the accuracy of diagnostic and other information collected in household interviews. For diagnostic information, the household respondent can, at best, pass on to the interviewer only the information the physician has given to the family. For conditions not medically attended, diagnostic information is often no more than a description of symptoms. However, other facts such as the number of disability days caused by the condition can be obtained more accurately from household members than from any other source since only the persons concerned are in a position to report information of this type.

Rounding of numbers.—The original tabulations on which the data in this report are based show all estimates to the nearest whole unit. All consolidations were made from the original tabulations using the estimates to the nearest unit. In the final published tables the figures are rounded to the nearest thousand, however, because of sampling error, they are not necessarily accurate to that detail. Derived statistics, such as rates and percent distributions, are computed after the estimates on which they are based have been rounded to the nearest thousand.

Population figures, -- Some of the published tables include population figures for specified categories, Except for certain over-all totals by age and sex (which are adjusted to independent estimates); these figures are based on the sample of households in the U.S. National Health Survey. They are given primarily for the purpose of providing denominators for rate computation. and for this purpose are more appropriate for use with the accompanying measures of health characteristics than other population data that may be available. In some instances they will permit users to recombine published data into classes more suitable to their specific needs. With the exception of the over-all totals by age and sex, mentioned above, the population figures. may in some cases differ from corresponding figures (which are derived from different sources) published in reports of the Bureau of the Census. For population data for general use, see the official estimates presented in Bureau of the Census reports in the P-20, P-25, P-50, P-57, and P-60 series.

Reliability of Estimates

Since the estimates are based on a sample, they will differ somewhat from the figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and interviewing personnel and procedures. As in any survey, the results are also subject to measurement error.

The standard error is primarily a measure of sampling variability, that is, the variations that might occur by chance because only a sample of the population is surveyed. As calculated for this report, the standard error also reflects part of the variation which arises in the measurement process. It does not include estimates of any biases which might lie in the data. The chances are about 68 out of 100 that an estimate from the sample would differ from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference would be less than twice the standard error and about 99 out of 100 that it would be less than 2½ times as large.

In order to derive standard errors which would be applicable to a wide variety of health statistics and which could be prepared at a moderate cost, a number of approximations were required. As a result, the tables of standard errors shown in this Appendix should be interpreted as providing an estimate of approximate standard error rather than as the precise standard error for any specific statistic.

The following rules will enable the reader to determine the sampling errors for the data contained in this report.

1. Estimates of aggregates: Approximate standard errors of estimates of aggregates, such as the number of persons with one or more chronic conditions, the number of physician or dental visits, and the number of disability days, are obtained from appropriate columns of table I.

Example:

The average annual number of bed-disability days for persons 45 years and older was 475,263,000 (table 16). Since the standard error for this estimate is not shown in table I, it is necessary to interpolate between the standard error for 200,000,000 days which is 6,400,000 and the standard error for 500,000,000 days which is 12,000,000. Such interpolation gives 11,538,000 as the standard error for 475,263,000 bed-disability days.

- 2. Estimates of percentages in a percent distribution: Approximate standard errors of percentages in percent distributions of persons, conditions, injuries, and physician and dental visits are given in appropriate columns of table ll. Approximate standard errors of percentages in percent distributions of disability days are obtained from table III. Examples:
 - (A) Approximately 23.9 percent of the 4,969,000 persons 75 years and older had major limitation of activity (tables 1 and 2). Since neither the base nor the percentage is shown in table II, it is necessary to interpolate between 10 percent and 25 percent to obtain 1.37 as the standard error of 23.9 percent with a

base of 3,000,000 and 1,08 as the standard error of 23.9 percent with a base of 5,000,000. A final interpolation between these results yields 1.08, rounded to 1.1, as the standard error for a statistic of 23.9 percent with a base

of 4,969,000.

(B) Of the 475,263,000 days of bed disability reported for persons 45 years and older. 42.1 percent of the days were for males (table 16). Since neither the base nor the percentage is shown in table III, it is necessary to interpolate between 25 percent and 50 percent to obtain 1.14 as the standard error for 42.1 percent with a base of 250,000,000 and 0.84 as the standard error of 42.1 percent with a base of 500,000,000. A final interpolation between these results yields 0.9 as the standard error of 42.1 percent with a base of 475,263,000.

3. Estimates of prevalence rates: Prevalence estimates of a chronic condition per 1,000 persons are obtained from table II. Since table Il is set up for the estimation of the standard error of a rate per 100, the prevalence per 1.000 must first be converted to a percentage; table II is then entered with this percentage and the number of persons in the population category (base of the percentage). The entry in the body of the table must then be multiplied by 10 to apply to the rate per 1,000 persons.

Example:

The prevalence rate of heart disease among persons 65 years and older was 148.8 per 1,000 population (table 21). This rate expressed as a percentage is 14.9 and it is based on 14,667,000 persons in the age group 65 years and older. Since neither the base nor the percentage is shown in table II, it is necessary to interpolate between 10 percent and 25 percent to obtain 0.67 as the standard error for 14.9 percent with a base of 10,000,000 and 0.47 as the standard error of 14.9 percent with a base of 20,000,000. A final interpolation between these results yields 0.58 as the standard error of 14.9 with a base of 14,667,000. Multiplying this standard error by 10 gives 5.8 as the standard error for a rate of 148.8 persons per 1,000 population.

4. Estimates of the number of disability days per person per year, the number of physician or dental visits per person per year, and the incidence rate of acute conditions: Approximate standard errors for these rates are obtained

as follows:

(a) Obtain the standard error of the numerator from table I. Divide the standard error by the numerator itself. Square the result.

(b) Obtain the standard error of the denominator from table I. Divide the standard error by the denominator itself. Square the

(Note: Where the denominator is adjusted to Bureau of the Census figures and therefore is not subject to sampling error, this quantity is

(c) Add the answers from steps (a) and (b)

above and extract the square root,

(d) Multiply the answer from step (c) by the rate. The result is the approximate standard error of the rate. This procedure normally gives an overestimate of the true sampling error. Examples:

(A) There were 5.8 physician visits per person per year for persons 45 years and older with family income less than \$2,000 (table 32). Using Rule 1 we find that the standard error for the numerator of 66,374,000 physician visits is 2,793,000, and the standard error for the denominator of 11,394,000 persons (table 43) is 188,000. Completing the computation as follows:

$$5.8\sqrt{\left(\frac{2,793,000}{66,374,000}\right)^2 + \left(\frac{188,000}{11,394,000}\right)^2}$$

yields 0.3 as the standard error of 5.8 physician visits per person 45 years and older with

family income less than \$2,000.

(B) There were 553.8 upper respiratory conditions per 1,000 persons 45 years and older during the year July 1958-June 1959 (table 28). Using Rule 1 we find that the standard error for the numerator of 27,622,000 conditions is 1,402,000, and the denominator has no sampling error. Completing the computation as follows:

$$553.8\sqrt{\left(\frac{1,402,000}{27,622,000}\right)^2+0}$$

yields a value of 28.1. However, since the data on acute conditions are presented by single years in this report, and the standard error tables are set up for two years of data collection. the value 28.1 must be multiplied by 1.25 to obtain a standard error of 35.1 for a rate of 553.8 upper respiratory conditions per 1,000 persons 45 years and older reported during the year July 1958-June 1959.

	(All num	bers shown in thous	ands)								
	For estimates of the number of:										
Size of estimate	Persons with status or health charac- teristic Chronic conditions by type	Acute conditions Injuries Persons injured	Physician visits Dental visits	Disability days							
	The approximate standard error is:										
100	18	_	· -	-							
500	40	-	-	-							
1,000	60	280	340	400							
2,000	80	400	480	560							
3,000	100	480	600	720							
5,000	130	640	800	960							
10,000	180	880	1,040	1,200							
20,000	240	1,280	1,520	1,760							
30,000	260	1,440	1,800	2,160							
50,000	280	2,000	2,400	2,800							
100,000	320	2,800	3,600	4,400							
200,000	-	4,000	5,200	6,400							
500,000	-	6,000	9,000	12,000							
750,000	-	6,720	11,760	16,800							
1,250,000	-	7,600	16,600	25,600							

NOTE: Since the estimated number of persons in any age-sex class is adjusted to official Bureau of the Census figures, such estimates are not subject to sampling errors.

Table II. Standard errors of percentage distributions of persons, conditions, injuries, and physician and dental visits

When the base of is numb		For estimated percentages of:							
Persons with status or health charac- teristic Chronic conditions by type (In tho	Acute conditions Persons injured Injuries Physician visits Dental visits usands)	2 or 98	5 or 95	10 or 90	25 or 75	50			
		err		d in p	standard in per- is:				
100	2,500	2.9 1.3 0.9 0.6 0.4 0.3 0.2 0.2	1.4 1.0 0.8 0.6 0.5 0.3	2.4 1.7 1.2 1.0 0.8 0.6 0.4 0.3	3.5 2.5 1.8	3.3 2.3 1.9 1.4 1.0 0.7 0.6			

Table III. Standard errors of percentage distributions of disability days

When the base of the percentage is number of:	For estimated percentages of:											
Disability days (In thousands)	2 or 98	5 or 95	10 or 90	25 or 75	50							
				ndard error tage points								
2,500 12,500 25,000 50,000 75,000	3.4 1.5 1.0 0.7 0.6	5.2 2.3 1.7 1.2 1.0	7.2 3.2 2.2 1.6 1.3	10.4 4.6 3.3 2.3 1.9	12.0 5.4 3.8 2.7 2.2							
125,000 250,000 500,000 750,000 1,250,000	0.5 0.3 0.2 0.2 0.2	0.7 0.6 0.4 0.3 0.2	1.0 0.7 0.5 0.4 0.3	1.4 1.0 0.7 0.6 0.5	1.7 1.2 0.9 0.6 0.6							

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Demographic, Social, and Economic Terms

Age.—The age recorded for each person is the age at last birthday. Age is recorded in single years and grouped in a variety of distributions depending upon the purpose of the table.

Marital status.—Marital status is recorded only for persons 14 years of age or older. The categories of marital status are: married, widowed, divorced, separated, and never married. Persons whose only marriage was annulled are counted as "never married." Persons with common-law marriages are considered to be married. "Separated" refers to married persons who have a legal separation or who have parted because of marital discord.

Income of family or of unrelated individuals.—Each member of a family is classified according to the total income of the family of which he is a member. Within the household all persons related to each other by blood, marriage, or adoption constitute a family. Unrelated individuals are classified according to their own income.

The income recorded is the total of all income received by members of the family (or by an unrelated individual) in the 12-month period ending with the week of interview. Income from all sources is included, e.g., wages, salaries, rents from property, pensions, help from relatives, and so forth.

Major activity.—All persons 6 years old or over are classified according to their major activity during the 12-month period prior to the week of interview. The "major" activity, in case more than one is reported, is the one at which the person spent the most time during the 12-month period.

The categories of major activity for adult persons are: usually working, usually keeping house, retired, and other. For several reasons these categories are not comparable with somewhat similarly named categories in official Federal labor force statistics. In the first place, the responses concerning major activity are accepted without detailed questioning, since the objective of the question is not to estimate the numbers of persons in labor force categories but to identify crudely certain population groups which may have differing health problems. In the second place, the figures represent the major activity over the period of an entire year, whereas official labor force statistics relate to a much shorter period, usually one week. Finally, in the definitions of the specific categories which follow, certain marginal groups are classified in a different manner to simplify the procedures.

- I. Usually working includes paid work as an employee for someone else; self-employment in own business, or profession, or in farming; and unpaid work in a family business or farm. Work around the house, or volunteer or unpaid work, such as for church, Red Cross, etc., is not counted as working.
- 2. Usually keeping house includes any activity described as "keeping house" which cannot be clas-

- sified as "working" or "going to school."
- 3. Retired includes persons 50 years old or over who consider themselves to be retired. In case of doubt, a person 50 years old or over is counted as retired if he, or she, has either voluntarily or involuntarily stopped working, is not looking for work, and is not described as "keeping house." A retired person may or may not be unable to work.
- 4. Other includes adults not classed in any of the other categories. Examples of inclusions are: a person who states that he spent most of the past 12 months looking for work, a person doing volunteer work only, a person under 50 years of age who describes himself as "retired" or "taking it easy," a person under 50 years of age who is described as "unable to work," or "unable to go to school" or a person 50 years of age or over who describes himself as "unable to work" and is not "retired."

Location of Residence Terms

Urban residence.—The definition of urban areas used in the U.S. National Health Survey is the same as that used in the 1950 Census. According to this definition, the urban population comprises all persons living in (a) places of 2,500 inhabitants or more incorporated as cities, boroughs, and villages; (b) incorporated towns of 2,500 inhabitants or more except in New England, New York, and Wisconsin, where "Towns" are simply minor civil divisions of counties; (c) the densely settled urban fringe, including both incorporated and unincorporated areas around cities of 50,000 or more; and (d) unincorporated places of 2,500 inhabitants or more outside any urban fringe.

In this report, the urban population has been subdivided into those living "Inside urbanized areas" and those living in "Other urban places."

Inside urbanized areas.—Following the definition used in the 1950 Census, the population in urbanized areas comprises all persons living in (a) cities of 50,000 inhabitants or more in 1940 or according to a special census taken between 1940 and 1950; and (b) the densely settled urban fringe, including both incorporated and unincorporated areas, surrounding these cities.

Other urban places.—The remaining urban population not classified as living "Inside urbanized areas" is classified as living in "Other urban places."

Rural residence.—The remaining population not classified as "Urban" is classified as "Rural." In this report the rural population has been subdivided into "Rural farm" and "Rural nonfarm."

Rural farm.—All rural residents living on farms are classified as "Rural farm." In deciding whether members of a household reside on a farm or ranch.

the statement of the household respondent that the house is on a farm or ranch is accepted, with the following exception. A house occupied by persons who pay cash rent for the house and yard only is not counted as a farm or ranch even though the surrounding area is farm land. This special case does not cover: (1) the living quarters of a tenant farmer who rents farm land as well as house and yard; (2) the quarters of a hired hand who receives living quarters on a farm as part of his compensation; or (3) separate living quarters inside a structure which is classified as on a farm. In all these cases the living quarters are counted as on a farm. Rural nonfarm. - The remaining rural population not classified as "Rural farm" is classified as "Rural nonfarm."

Terms Defining Morbidity Conditions

Condition,—A morbidity condition, or simply a condition, is any entry on the questionnaire which describes a departure from a state of physical or mental wellbeing, it results from a positive response to one of a series of "illness-recall" questions (11-17, Appendix III). In the coding and tabulating process, conditions are selected or classified according to a number of different criteria, such as, whether they were medically attended; whether they resulted in disability; whether they were acute or chronic; or according to the type of disease, injury, impairment, or symptom reported. For the purposes of each published report or set of tables, only those conditions recorded on the questionnaire which satisfy certain stated criteria are included.

Conditions, except impairments, are coded by type according to the International Classification of Diseases, with certain modifications adopted to make the code more suitable for a household-interview-type survey.

Chronic condition.—A condition is considered to be chronic if (1) it is described by the respondent in terms of one of the chronic diseases on the "Check List of Chronic Conditions" or in terms of one of the types of impairments on the "Check List of Impairments" shown as cards A and B in Appendix III, or (2) the condition is described by the respondent as having been first noticed more than 3 months before the week of the interview.

Chronic effect or residual of injury.—A chronic condition resulting from an injury may be either an impairment, such as paralysis, or some other type of late effect of the injury, such as arthritis. Disability from such conditions is included with that resulting directly from the injuries, unless otherwise specified.

With a few exceptions, injuries that are still giving trouble are classified according to the chronic effect of the injury if the injury occurred 3 months or more before the interview week, but to the injury itself if the injury occurred less than 3 months before.

Impairment,—Impairments are chronic or permanent defects, usually static in nature, resulting from disease, injury, or congenital malformation. They represent decrease or loss of ability to perform various functions, particularly those of the musculoskeletal system and the sense organs. All impairments are classified by means of a special supplementary code for impairments. Hence, code numbers for impairments in the International Classification of Diseases are not used. In the Supplementary Code impairments are grouped ac-

cording to the type of functional impairment and etiol-

Acute condition.—All conditions not classed as chronic are considered to be acute, Minor acute conditions, both diseases and injuries, involving neither restricted activity nor medical attendance, are excluded from the statistics. (See definitions of "Restricted-activity day" and "Medically attended condition.")

Injury condition.—An injury condition, or simply an injury, is an acute condition of the type that is classified to the nature of injury code numbers (N800-N999) in the International Classification of Diseases. In addition to fractures, lacerations, contusions, burns, and so forth, which are commonly thought of as injuries, this group of codes include: effects of exposure, such as sunburn; adverse reactions to immunizations and other medical procedures; and poisonings. Unless otherwise specified, the term injury is used to cover all of these.

As in the case of other acute conditions, acute injury conditions involving neither restricted activity nor medical attendance are excluded from the statistics.

Terms Relating to Conditions

Prevalence of conditions.—In general, prevalence of conditions is the estimated number of conditions of a specified type existing at a specified time or the average number existing during a specified interval of time.

The prevalence of chronic conditions denotes the number of chronic cases reported to be present or assumed to be present at the time of interview; those assumed to be present at the time of the interview are cases described by the respondent in terms of one of the chronic conditions on the "Check List of Chronic Conditions" and reported to have been present at some time during the 12-month period prior to the interview.

Estimates of the prevalence of chronic conditions may be restricted to cases that satisfy certain additional stated criteria, such as, for example, cases involving a day or more in bed in the past year, or cases still under medical care.

Incidence of conditions.—Incidence of conditions is the estimated number of conditions of a specified type which had their onset within a specified interval of time. In this survey the interval was one year, or annual incidence.

Onset of condition,—A morbidity condition, whether acute or chronic, is considered to have had its onset when it was first noticed. This could be the time the person first felt "sick," or became injured, or it could be the time the person or his family was first told by a physician that he had a disease of which he was previously unaware. For a chronic condition, episodic in nature, the onset is always considered to be the original onset rather than the start of the most recent episode.

Medically attended condition.—A condition for which a physician was consulted is called a medically attended condition. Consulting a physician includes consultation in person or by telephone for treatment or advice. Advice from the physician transmitted to the patient through the nurse is counted as medical consultation as well as visits to physicians in clinics or hospitals. If at one visit the physician is consulted about more than one condition for each of several patients, each condition is counted as medically attended.

A parent consulting a physician about a child's condition is counted as medical consultation about that condition even if the child was not seen by the physician at that time.

For the purpose of this definition "physician" includes doctors of medicine and osteopathic physicians. The term "doctor" is used in the interview, rather than "physician," because of the need to keep to popular usage. However, the concept toward which all instructions are directed is that which is described here.

A condition is counted as medically attended if a physician was consulted about it at its onset or at any time thereafter. However, the first medical attention for a condition that was present in the 2 calendar weeks before the interview may not occur until after the end of the 2-week period, and, in fact, may not occur until after the interview. Such cases are necessarily treated as though there had been no medical attention.

Activity-restricting condition.—An activity-restricting condition is a condition which has caused at least 1 day of restricted activity during the 2 calendar weeks before the interview week. (See definition of "Restricted-activity day.") The incidence of acute activity-restricting conditions is estimated from the number of such conditions reported as having started in the 2-week period, but a condition starting in the 2-week period which did not result in restricted activity until after the end of that period is not included.

Still under care.—This information is obtained only for chronic conditions. A chronic condition which is "still under care" is one for which the person is still "under instruction" from a physician. By "under instruction" is meant one or more of the following: (1) taking certain medicine or treatment prescribed by a physician, (2) observing a certain systematic course of diet or activity, (3) visiting the physician regularly for checking on the condition, and (4) under instruction from the physician to return if some particular thing happens.

For the purposes of this definition "physician" is defined as in "Medically attended condition."

Terms Relating to Disability

<u>Disability.</u>—Disability is a general term used to describe any temporary or long-term reduction of a person's activity as a result of an acute or chronic condition.

Disability Days

Disability days are classified according to whether they are days of restricted activity, bed-days, hospital days, work-loss days, or school-loss days. All hospital days are, by definition, days of bed disability; all days of bed disability are, by definition, days of restricted activity. The converse form of these statements is, of course, not true, Days lost from work and days lost from school are special terms which apply to the working and school-age populations only, but these, too, are days of restricted activity. Hence, "days of restricted activity" is the most inclusive term used to describe disability days.

Restricted-activity day.—A day of restricted activity is a day when a person cuts down on his usual activities for the whole of that day on account of an illness or an injury. The term "usual activities" for any day means the things that the person would ordinarily

do on that day. For children under school age, "usual activities" depend upon whatever the usual pattern is for the child's day which will, in turn, be affected by the age of the child, weather conditions, and so forth. For retired or elderly persons, "usual activities" might consist of almost no activity, but cutting down on even a small amount for as much as a day would constitute restricted activity. On Sundays or holidays "usual activities" are taken to be the things the person usually does on such days—going to church, playing golf, visiting friends or relatives, or staying at home and listening to the radio, reading, looking at television, and so forth,

Restricted activity does not imply complete inactivity but it does imply only the minimum of "usual activities." A special nap for an hour after lunch does not constitute cutting down on usual activities, nor does the elimination of a heavy chore, such as cleaning ashes out of the furnace or hanging out the wash. If a farmer or housewife carries on only the minimum of the day's chores, however, this is a day of restricted activity.

A day spent in bed or a day home from work or school because of illness or injury is, of course, a restricted-activity day.

Bed-disability day,—A bed-disability day, sometimes for brevity referred to as a "bed-day," is a day on which a person was kept in bed either all or most of the day because of an illness or an injury. "All or most of the day" is defined as more than half of the daylight hours. All hospital days are included as bed-disability days even if the patient was not actually in bed at the hospital.

Work-loss day.—A day is counted as lost from work if the person would have been going to work at a job or business that day but instead lost the entire work day because of an illness or an injury. If the person's regular work day is less than a whole day and the entire work day was lost, it would be counted as a whole work day lost. Work-loss days are determined only for persons 17 years of age and over.

Condition-days of restricted activity, bed disability, etc.—Condition-days of restricted activity, bed disability, and so forth are days of the various forms of disability associated with any one condition. Since any particular day of disability may be associated with more than one condition, the sum of days for all conditions adds to more than the total number of person-days.

Person-days of restricted activity, bed disability, etc.—Person-days of restricted activity, bed disability, and so forth are days of the various forms of disability experienced by any one person. The sum of days for all persons in a group represents an unduplicated count of all days of disability for the group.

Long-Term Disability

Chronic activity limitation.—Chronic activity limitation is ascertained for all persons with one or more chronic conditions. These persons are divided into 4 categories according to the extent to which their activities are limited as a result of the conditions (cards C,D,E, and F, Appendix III). For the purpose of this report categories 2 and 3 have been combined.

Since the major activities of housewives and workers and other persons differ, a different set of criteria is used to determine the amount of reduction of major activity for each group. However, there is a general similarity between the criteria as will be seen in the description of the categories that follows:

Major limitation of activity.—Inability to carry on major activity of the group:

Housewives: Workers and all other adult persons: inability to do any housework.

inability to work at a job or bus-

Partial limitation of activity.—Limitation of amount or kind of participation in activity of the group;

Housewives:

limited in amount or kind of housework or limited in recreational or community activities.

Workers and all other adult persons:

limited in amount of work or kind of employment or limited in recreational or community activities

No limitation of activity.—No limitation as described above.

Chronic mobility limitation.—Persons with one or more chronic conditions are also classified according to limitation of mobility (Card G, Appendix III). The four categories describing the extent of mobility limitation, which have been combined into three for the purpose of this report, are used in the classification of persons regardless of their major activity status.

1. Major limitation of mobility.—Confined to the house except in case of emergencies.

 Partial limitation of mobility.—Limited or in need of help in moving around outside the house.

3. No limitation of mobility.—No limitation as described above.

Terms Relating to Persons Injured

Person injured.—A person injured is one who has sustained an injury in an accident, or in some type of nonaccidental violence. (See definition of "Injury condition," above.) Each time a person is injured he is included in the statistics as a separate "person injured"; hence, one person may be included more than once.

The statistics of persons injured include only persons sustaining injuries which involved at least one full day of restricted activity or medical attendance.

Note that the number of persons injured is not equivalent to the number of "accidents" for several reasons; (1) the term "accident," as commonly used, may not involve injury at all; (2) more than one injured person may be involved in a single accident so that the number of accidents resulting in injury would be less than the number of persons injured in accidents; and (3) the term "accident" ordinarily implies an accidental origin, whereas "persons injured" includes persons whose injury resulted from certain nonaccidental violence.

The number of persons injured in a specified time interval is always equal to or less than the incidence of injury conditions, since one person may incur more than one injury in a single accident or nonaccidental violence.

Class of accident.—Injuries, injured persons, and resulting days of restricted activity may be grouped according to class of accident. This is a broad classification of the types of events which resulted in persons being injured. Most of these events are accidents in the usual sense of the word, but some are other kinds of mishap, such as overexposure to the sun or adverse

reactions to medical procedures, and others are non-accidental violence, such as attempted suicide. The classes of accidents are: (1) motor-vehicle accidents, (2) accidents occurring while at work, (3) home accidents, and (4) other. These categories are not mutually exclusive. For example, a person may be injured in a motor-vehicle accident which occurred while the person was at work.

Motor-vehicle accident,—The class of accident is "motor vehicle" if a motor vehicle was involved in any way. Thus, it is not restricted to moving motor vehicles or to persons riding in motor vehicles. A motor vehicle is any mechanically or electrically powered device, not operated on rails, upon which or by which any person or property may be transported or drawn upon a land highway. Any object, such as a trailer, coaster, sled, or wagon, being towed by a motor vehicle is considered a part of the motor vehicle. Devices used solely for moving persons or materials within the confines of a building and its premises are not counted as motor vehicles.

Accident while at work.—The class of accident is "while at work" if the injured person was 14 years of age or over and was at work at a job or a business at the time the accident happened.

Home accident.—The class of accident is "home" if the injury occurred either inside the house or outside the house. "Outside the house" refers to the yard, buildings, and sidewalks on the property, "Home" includes not only the person's own home but also any other home in which he might have been when he was injured.

Other.—The class of accident is "other" if the occurrence of injury cannot be classified in one or more of the first three class-of-accident categories. This category therefore includes persons injured in public places (e.g., tripping and falling in a store or on a public sidewalk), and also nonaccidental injuries such as homicidal and suicidal attempts. The survey does not cover the military population, but current disability of various types resulting from prior injury occurring while the person was in the Armed Forces is covered and is included in this class. The class also includes mishaps for which the class of accident could not be ascertained.

Medical Care Terms

Physician visit.—A physician visit is defined as consultation with a physician, in person or by telephone, for examination, diagnosis, treatment, or advice. The visit is considered to be a physician visit if the service is provided directly by the physician or by a nurse or other person acting under a physician's supervision. For the purpose of this definition "physician" includes doctors of medicine and osteopathic physicians. The term "doctor" is used in the interview, rather than "physician," because of the need to keep to popular usage. However, the concept toward which all instructions are directed is that which is described here.

Physician visits for services provided on a mass basis are not included in the tabulations. A service received on a mass basis is defined as any service involving only a single test (e.g., test for diabetes) or a single procedure (e.g., smallpox vaccination) when this single service was administered identically to all persons who were at the place for this purpose. Hence, persons passing through a tuberculosis chest X-ray trailer, by this definition, are not included as physician visits. However, a special chest X-ray given in a phy-

sician's office or an outpatient clinic is considered to be a physician visit.

Physician visits to hospital inpatients are not included.

If a physician is called to the house to see more than one person, the call is considered to be a separate physician visit for each person about whom the physician was consulted.

A physician visit is associated with the person about whom the advice was sought, even if that person did not actually see or consult the physician. For example, if a mother consults a physician about one of her children, the physician visit is ascribed to the child.

<u>Place of visit</u>.—The place of visit is a classification of the types of places at which a physician visit took place. (See definition of "Physician visit.") The definitions of the various categories are as follows:

- Home is defined as any place in which the person was staying at the time of the physician's visit. It may be his own home, the home of a friend, a hotel, or any other place the person may be staying (except as an overnight patient in a hospital).
- 2. Office is defined as the office of a physician in private practice only. This may be an office in the physician's home, an individual office in an office building, or a suite of offices occupied by several physicians. For purposes of this survey, physicians connected with prepayment group practice plans are considered to be in private practice.
- 3. Hospital clinic is defined as an outpatient clinic in any hospital.

- Company or industry health unit refers to treatment received from a physician or under a physician's supervision at a place of business (e.g., factory, store, office building). This includes emergency or first-aid rooms located in such places if treatment was received there from a physician or trained nurse.
 Telephone contact refers to advice given in a
- Telephone contact refers to advice given in a telephone call directly by the physician or transmitted through the nurse.
- 6. Other refers to advice or treatment received from a physician or under a physician's general supervision at a school, at an insurance office, at a health department clinic, or any other place at which a physician consultation might take place.

Dental Care Terms

Dental visit.—Each visit to a dentist's office for treatment or advice is considered to be a dental visit. The visit may involve services provided directly by the dentist or by a dental hygienist acting under a dentist's supervision. Services provided while a person was a patient in a hospital for overnight or longer are not considered to be dental visits.

Edentulous persons.—Persons who have lost all of their permanent teeth are classed as edentulous persons. An edentulous person may have dentures but does not have any natural teeth.

APPENDIX III

QUESTIONNAIRE

The items below show the exact content and wording of the questionnaire used in the household survey. The actual questionnaire is designed for a household as a unit and includes additional spaces for reports on more than one person.

CONF	IDENTI	AL: would p	permit	Health Survey is audidentification of the survey, and will a	e iadiv	idual will be	held	strictl	y confidenti	ıl, will l	be use	d only	by per	sons er	ll inform ngaged i	nation whi n and for	ch the	
Form NHS (4-18-58)	-2			BUR	EAU OF	T OF COMMERC THE CENSUS Ling Agent for the BALTH SERVICE	CE						1. 0	Question	osire			
				NATIONAL	. HE	ALTH SUR	VEY	•					-	OI (Toe ation	nairea	· · · · · · · · · · · · · · · · · · ·		
2. (a) Ada	dress or	description of	location	0				3. Idea Code	sample	5. Samp		PSU Number		egment		8. Serial	No.	
weight weight																		
9. la this house on a farm or ranch?																Yes [] No	
			(c) N	ame of Special Dwellin	ng Place	Code		10. Wh	at is the teles				11.	What Is	the best	time to co	IIP	
ing quarters Other No phone No phone 12. Ars there any other living quarters, occupied or vacant, in this building (opartment)?																		
13. Is the	re any o	except apartment ther building of ther occupied o	n this p	cs roperty for people t?	Yes	□ No		If "Y wheel listin	es" to question ser one or mo	us 12, 13 re addition		STRUC apply ducations			welling u	mit to dete	rmine r the	
		· · · · · · · · · · · · · · · · · · ·		,		5. RECORD OF	CAL	<u> </u>										
	-	tem	Date	1	Com.	2		Con.	3	Coc	n.	4		Com.		5	Com.	
Callbac		hold	Time		17.6 17.6		1	-			+						1	
respon	idual ndents	Col. No	Time		Print.	16. REASON	FOR	NON-II	TERVIEW									
TYPE:					8				c	1.				z				
Resson;	Ter	one at home- eated calls nporarily abser- er (Specify)	nt	Vacant - non-s Vacant - seaso Usual residenc Atmed Forces Other (Specify	onal c elsew		00	Elimina semple	shed le by mistake ted in sub- Specity)	۵	tervier ols csuse	mos ob	mained (los:				
Comments 17. Signat	-											-	18.	Code		-		
Special in	struction	18 OF DOTES		·			_					·	<u> </u>					
(b) Who and per: (c) Do (d) Is to awa a he	all pers sons in I ony (oth- here ony ly on bus espital?	o nomes of all of cons staying he the prescribed er) lodgers or no one else who I siness? On a v	other pe re who order.) comers ives he rist?]	re who is now Comporarily in	(List al freside io	Il persons who unce elsewhere. Yes (List)	List t	live b	cre,				→	Last que	me and i	nitial .	(1)	
(f) Do o	my of th	ane else stayir ese people hov eve on question	e a hon	a elsewhere?		Yes (List) Id member, deler				====	_==		-			•		
				o household? (Enter r er-in-law, partner, lod				ple;						Relation	• .	ia la		
3. How old	d wa re y	ou an your last	birthdo	ny?										Age		Und 1 y		
4. Race (C	heck on	e box for each	person)	•		• . •		•					□ Vhi	<u> </u>	Neg	to .	
		box for each p							•	`_				\ \label{Alaberta}		☐ Fe		
6. Where were you born? (Record state or foreign country) If 14 years old or over, ask: 7. Are you now mornlad, wildowed, divorced, separated or never married? (Check one box for each person) Wildowed Wellowed Never married												ars orced arsted						
8. What is	the higi	or over, ask: heat grade you grade complete	-					٠				•		Elem: High:				

9.	(a) Di Ye Ye (b) Ai L	d you s," s e you f "Ye	ever isk: u now is," de	serve in the lete	e Armed F	med Forces or orces, not co o from questi	unting the	reserves?	7				Yes Yes	F		 No	
	(c) Was any of your service during a war or was it peace-time only? [I *War, * ask: (d) During which war did you serve? [I *Peace-time* only, ask: Ww-!1 Korean																
		•				waan 'Juna 27	, 1950 cm	d January 31	, 19557				Yes				
10	(e) Was any of your service between June 27, 1950 and January 31, 1955? If 6 years old or over, sek: Under 6 years Under 6 years Working (For males over 16); working, looking for work, or doing semething else? (For children 6-16); going to school or doing semething else? (For children 6-16); going to school or doing semething else? (For children 6-16); going to school or doing semething else? (b) Are you retired? Yes No																
-		Τ.			ak adale se	reen for him	elf for a	uestions 11-2	5 and Tables	I. 11. and A	v. if		[] Respon		self		
ļ	R	þę	io at	pome	. Enter c	olumn numbe	of respo	ndent in each	column.	-,,			Col. No	w	s respo	No No	
	Were (e) W (b) A	you hat v	sick o ras th ing eli	t any mot	time LAS	of Illness, wi	THE WEI	EK BEFORE	· · · · · · · · · · · · · · · · · · ·			. p					
12	Lost hom	wool or o	may b	om h	ek before : iome?	lid you have	any accid	lente or injur	ies, either at		(a) What ware they? (b) Anything also?		Yes			∐ No	
13	injur (a) W	y tha That v	t bapp vere ti	ened 10 22	offects las before the offects?	st weak or the st time?	week be	fore from on	occident or				Yes			□N∘	
_	Last	wee ition	(best	o wo	which y	did you take on told me al	>out)?				(a) For what conditie (b) Anything else?	ms?	Yes			□ No	
Ľ	for c	THE long	PRES	EMT	TIME do y "No") Eve	ou have any on though the	allments y don't b	er conditions other you all	that have las the time?	ted	(a) What are they? (b) Anything else?		☐ Yes			□ No	
Ē							Table	I - ILLNI	ESSES, IMP	AIRMEN'	TS AND ACCIDENTS	1					
	Did you was?—did the doctor say it was?—did the use any symptom or a condition from q. 13 or q. 17, tak only for: to a doctor on talked to - "No" in col. (c) - record respondable doctor about dust's description. When was the cause of 7, of yes. If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q. 17, ask enly for: If so impairment or graphen to condition from q. 13 or q									ing detail ted below;	OR THE WEEK BE- FORE did couse you to cut						
Ħ		Ques	abou		(Il ill-effe accident, i	cts of earlier record ill effi ill Table A)			or injury, also	over,	rthritis stroke rumor (or cysts) OR Any entry in col. (d-1) or (d-2) of:	Spine - (Upper, or lower) back Arm - (Shoulder,		for as much as a day?			
ě.		tion No.			occurring o weeks, asi	ident or injuduring past 2 k: of the body w				you read ardin- ury news- peper	trouble condition disease coupled with	elbow, lo hand) Leg - (Hip, upp lower, an	wer, wrist, er, knee,	No .	Yea		
					hurt? Who was it? A	nt kind of injusything else l Table A)	-ry			print with plasses?	seeing or bearing; a part of the body; "mental" or any internal organ	ALSO If arm, leg, eye state whether C BOTH.	, or ear, INE or	to Cal. (k))			
┡	(a)	(b)	(c)	\neg		(d-1)	\dashv	(F-2)	(d-3)	(d-4)	(d-5)	x	(e)	(f)	(g)	
1			ים מ	- 1						Yes	<u> </u>					Days	
							able II	- HOSPITA	ALIZATION I	DURING I	PAST 12 MONTHS						-
number	Col. No. of	Qe tic	ies-	Phon you d the h pital (Moni	enter us-	How many days were you in the hespital, not count- ing the	How many of these - days were in	How many of these — days were in	Was this person still in the	did they o (If "they"	they say at the heapital the ca se any medical terms? did't say, ask): the lost doctor you talked to s		Were any on you du the hespi If "Yes" (a) What t	ring thi tal? was the	a stay :	at .	
Line nu	sos		ļ		*	day you left?	the pass 12 months?	2 weeks?	hospital on Sunday night?		ne detail as in cols. (d-1)-(d-5 ion from accident or injury, fil		opera (b) Any e	ther ope	rations	7	
\vdash	(a)	+	(b)	Mo:	(c)	(4)	(e)	(τ)	(g)		. (в)		☐ Yes	(i)		□ No	
				Yr:_			or						,				
ئے	<u>_</u>	Ļ	\perp	_		Days	Days	Days	□ N•				L .				
_	Line from Table	0	1.	What	t part of th	a body was l	ourt? Who		ABLE A (A		and Injuries)		Accid	lent hap 2 weeks		during	,
2. When did it happen? Year (Enter month also if the year is 1957 or 1958) Month past 2 weeks																	
L					n happen?			At home	(inside or on (own home or	staide the l	nouse) [\While in A	armed Services	Some	other pl	ace		
F	invol	ved i	o the	peeid	lent in any			Yee	□ No	•							
*			nt haj			business wh		Yes Yes	□ N	0	Under 14 years at ti	me of accident			,]

16. Has ony THE P.	1& Has onyone in the family - you, your, etchad any of these conditions DURING THE PAST 12 MONTHS?] Yes			No	
	(Read Card A, condition by condition; record any conditions mentioned in the column for the person)																		
17. Does or		the family ha									-			1=	Yes	· · ·		No	
		ad Card B, co			dition; record any c person)	onditions													
					id onyone in the far linic? Anyone else		, your-	-, etctalk							Yes		Ċ] No	
If "Yes"	•	ee during the				-						•					No. of tir	me s	
(c) When	e did you	talk to the	doctor?												lace		Times		
		nes at (hem number of ris												At	At home				
	(Record total number of times for each type of place)															ic ndustry		_	
(*Ho	("Hospital clinic" excludes overnight steys)															Other (Specify)			
if "Yes"	19. (c) East week or the week before did anyone in the family go to a dontist? Anyone else? If "Yea" (b) How many times during the past 2 weeks?															Yes No. of times			
					onths did you ge to	a dentist	7								One Three Four or more				
OVER	night ar l	PAST 12 MG	ONTHS	has any	rone in the family b	ieen o pal	Hent in	a hespital						-	☐ None ☐ Yes (Table II) ☐ No				
If "Yes" (b) How		mes were you	in the	hespita	17			***							No. of times				
seni	torium?	st 12 months	has an	yone in	the family been a p	ootlent in	O RUFS	ing home or							Yes (Table II) No				
If "Yes" (b) How many times were you in a nursing hame ar sanitarium?															No. of times				
					d the total income								_	Gro	p No.				
your's, salaries	your-'s, , rents f	rom property,	pensio	netud me, help	e income from all a from relatives, etc	c.	uch as	wages,		···-									
<u> </u>			L Did.		Table I - I		T i		T			$\overline{}$	<u> </u> -						
How many of these		years old over,ask:	or before that		E PAST J MONTHS		notic	Did you first notice		Hew long since you	Do you still take				ter compli ion for ess				
- dnys ware you in bed	Lost	If "Yes" in col. (i):			Did start	lf col.	PAST			talked ta	any medi- cine ar treatment	days		onse ok at	If 1, 2 or 3 in		IÌ 1. or 2]	
all or	or the low many days did before Durin and days and before would before would before the before before the befo		r—	during the past 2 weeks or	(k) is checked,	hefor	MONTHS or hefore that time? (If during past 12 months, sak):		etor ut?	that the	tha pa	, 16		col. (r): in	in a	or "3" in col.			
the day?			3	before that time?	or the	1			***	prescrib]"	od eoch atement	la this because	(a)	(r) sek:•	ì		
you have		work (going to			(If during past	tion is on				than one month, enter	Or, falls	In bed	~ π	ben teli • which	of the	Which	Piease look at	۱.	
	been working	seen school)?		-	2 weeks, ask): Which week,	either one of	Whic	Which month?		d. 1°	any odv	ice or mes		statement fits you	condi- tions	L. I	this card and	1	
	at a jab or busi-		(n))		last week ar	Cards A or B.			lor '	"Mo.")	ne gove	day?	ь	st. how	have	05	read each statement	43	
	nees except				the week hefore?	continue,						-	c	urds C-	tald me about?	for	Then tell me which		
]	for ? (If 6-16					wise STOP							*F	pro- iste)		condi-	statement fits you	'	
	gra.,]									ľ	- [-		nemed	Show		
	("going to school"	,							1								Card G)	1	
(b) —— Daya	(i)	(j) —— Days	(k)	(1)	(m)	(22)	Mo	(n)	 _	(o) Wos.	(p)	(q)	.y.	(r)	(a)	(t)	(n)	╁	
or None	□ No	or Nose			☐ Week before ☐ Before 2 wks. '		Y	efore Birth	_	Yrs. No Dr.	□ No	_ 01	1		☐ Yes			1	
					Table II - H	IOSPITA	ALIZA	UG HOLTA	RINC	PAST	12 MO	NTHS							
What is th		and address of	f the		For completed			only: If "No" to What part											
					the hespital	If "No" to col. (k), esk:		both cols. (k) and (l)	1	of the hospital	The semina sha			cast of this insurance—that is, who					
(Enter na	me, city	or county and	(State)		hy any kind			Do you exp		bill was bo) take	(will	poys the premium?							
1						kind at plan that		hespital bill to be poid for		of hy insurance									
]						pays for hespital				by lesurance or ony plan									
1		(j)			(k)	costs?		of this kind?		(n)		(6)							
					Yes (Skip	☐ Yes (Skip		(m) Yes				(0) Family member(s) Other (\$pecity)							
		- -			to col.n)	to col.n) to col.n)				⊏%արտ%		1					—.		
<u> </u>					□ No	□ No □ No		□ No (Stop) □ ¾ o		□ ¾ or	more Union, clubs, etc.							_	
			•			F	отно	TES AND C	OMME	NTS			-						
1																			
1																			
1																			
1		•																	
1																			

Card A

1. Asthma or hay fever

3. Chronic bronchitis

4. Repeated attacks of sinus

2. Tuberculosis

NATIONAL HEALTH SURVEY

Check List of Chronic Conditions

14. Any other chronic stomach

15. Kidney stones or chronic

kidney trouble

trouble

Card E

For:

aging to school

present time.

NATIONAL HEALTH SURVEY

Children from 6 years old and others

1. Cannot go to school at all at

Card G

NATIONAL HEALTH SURVEY

1. Confined to the house all the time,

expept in emergencies.

Card C

NATIONAL HEALTH SURVEY

Workers and other persons except

1. Cannot work at all at present.

2. Can work but limited in amount or

Housewives and Children

kind of work.

SELECTED REPORTS FROM THE U.S. NATIONAL HEALTH SURVEY

Series A (Program descriptions, survey designs, concepts, and definitions)

- No. 1. Origin and Program of the U. S. National Health Survey. PHS Pub. No. 584-A1. Price 25 cents.
- No. 2. The Statistical Design of the Health Household-Interview Survey. PHS Pub. No. 584-A2. Price 35 cents.
- No. 3. Concepts and Definitions in the Health Household-Interview Survey. PHS Pub. No. 584-A3. Price 30 cents.

Series 8 (Health Interview Survey results by topics)

- No. 6. Acute Conditions, Incidence and Associated Disability, United States, July 1957-June 1958. PHS Pub. No. 584-86.

 Price 35 cents.
- No. 7. Hospitalization, Patients Discharged From Short-Stay Hospitals, United States, July 1957-June 1958. PHS Pub. No. 584-87. Price 30 cents.
- No. 8. Persons Injured by Class of Accident, United States, July 1957-June 1958. PHS Pub. No. 584-88. Price 40 cents.
- No. 9. Impairments by Type, Age, and Sex, United States, July 1957-June 1958. PHS Pub. No. 584-89. Price 25 cents.
- No. 10. Disability Days, United States, July 1957-June 1958. PHS Pub. No. 584-810. Price 40 cents.
- No. 11. Limitation of Activity and Mobility Due to Chronic Conditions, United States, July 1957-June 1958. PHS Pub.No. 584-811. Price 30 cents.
- No. 12. Chronic Respiratory Conditions Reported in Interviews, United States, July 1957-June 1958. PHS Pub. No. 584-812.

 Price 30 cents.
- No. 13. Heart Conditions and High Blood Pressure Reported in Interviews, United States, July 1957-June 1958. PHS Pub. No. 584-813. Price 30 cents.
- No. 14. Dental Care, Interval and Frequency of Visits, United States, July 1957-June 1959. PHS Pub. No. 584-814. Price 35 cents.
- No. 15. Dental Care, Volume of Visits, United States, July 1957-June 1959. PHS Pub. No. 584-815. Price 35 cents.
- No. 16. Types of Injuries, Incidence and Associated Disability, United States, July 1958-June 1959. PHS Pub. No. 584-B16.
 Price 30 cents.
- No. 17. Peptic Ulcers Reported in Interviews, United States, July 1957-June 1959. PHS Pub. No. 584-817. Price 25 cents.
- No. 18. Acute Conditions, Incidence and Associated Disability, United States, July 1958-June 1959. PHS Pub. No. 584-818.
 Price 30 cents.
- No. 19. Volume of Physician Visits, United States, July 1957-June 1959. PHS Pub. No. 584-819.
- No. 20. Arthritis and Rheumatism Reported in Interviews, United States, July 1957-June 1959. PHS Pub. No. 588-820.
- No. 21. Diabetes Reported in Interviews, United States, July 1957-June 1959. PHS Pub. No. 584-821.
- No. 22. Loss of Teeth, United States, July 1957-June 1958. PHS Pub. No. 584-822.

Series C (Health Interview Survey results for population groups)

- No. 1. Children and Youth, Selected Health Characteristics, United States, July 1957-June 1958. PHS Pub. No. 584-C1.
 Price 35 cents.
- No. 2. Veterans, Health and Medical Care, United States, July 1957-June 1958. PHS Pub. No. 584-C2. Price 40 cents.
- No. 3. The Hawaii Health Survey, Description and Selected Results, Oahu, Hawaii, October 1958-September 1959. PHS Pub. No. 584-C3. Price 40 cents.
- No. 4. Older Persons, Selected Health Characteristics, United States, July 1957-June 1959. PHS Pub. No. 584-C4.

Series D (Developmental and Evaluation Reports)

- No. 1. A Study of Special Purpose Medical-History Techniques. PHS Pub. No. 584-D1. Price 30 cents.
- No. 2. Co-operation in Health Examination Surveys, PHS Pub. No. 584-D2. Price 35 cents.

Catalog Card

U. S. National Health Survey.

Older persons, selected health characteristics, United States, July 1957-June 1959; selected statistics relating to limitation of activity and mobility, disability days, chronic and acute conditions, and physician and dental visits for persons 45 years and older. Based on data collected in household interviews during the period July 1957-June 1959. Washington, U. S. Dept. of Health, Education, and Welfare, Public Health Service, Division of Public Health Methods, 1960.

- 76 p. tables. diagrs. 27cm. (Ite Health statistics, ser. C4)
 - U. S. Public Health Service. Publication no. 584-C4.
 - 1. Aged Statistics. 2. Aged Health services.
- 1. Title. (Series. Series: U. S. Publication no. 584-C4)

Cataloged by U. S. Dept. of Health, Education, and Welfare Library.