



Mortality Trend Tables  
for Leading Causes  
and Selected Components:

United States, 1950-69

U.S. DEPARTMENT OF  
HEALTH, EDUCATION, AND WELFARE  
PUBLIC HEALTH SERVICE  
Health Resources Administration  
National Center for Health Statistics

Table 1.—Death Rates for Diseases of Heart, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 390-398,402,404,410-429 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 400-402,410-443 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	366.1	11.0	1.4	1.0	3.0	11.5	69.5	242.3	667.6	1,634.5	3,644.9	9,162.0
1968-----	372.6	12.1	1.6	1.0	2.9	11.9	71.0	250.2	695.0	1,687.4	3,726.4	9,295.3
1967-----	364.5	11.4	1.1	0.8	2.6	12.4	71.7	252.1	687.4	1,669.5	3,670.6	9,240.5
1966-----	371.2	10.6	1.3	0.8	2.9	13.8	72.0	258.6	706.8	1,719.4	3,805.0	9,486.2
1965-----	367.4	9.6	1.3	0.9	3.1	14.1	72.9	258.8	704.3	1,697.8	3,835.9	9,600.5
1964-----	365.7	9.4	1.4	0.9	3.3	14.2	74.2	260.5	711.5	1,710.9	3,826.4	9,541.1
1963-----	375.2	7.6	1.2	1.0	3.6	14.9	75.3	266.9	724.2	1,758.9	3,992.8	9,964.1
1962-----	370.1	6.8	1.1	1.0	3.6	14.8	73.8	267.9	718.5	1,755.6	3,975.7	9,707.2
1961-----	362.4	6.9	1.1	1.0	3.9	14.9	73.5	267.2	714.7	1,701.3	3,931.7	9,241.6
1960-----	369.0	6.6	1.3	1.3	4.0	15.6	74.6	271.8	737.9	1,740.5	4,089.4	9,317.8
1959-----	365.2	7.1	1.5	1.3	4.3	16.5	73.8	272.6	728.7	1,721.1	4,030.7	9,166.8
1958-----	367.7	7.4	1.4	1.3	4.4	16.8	75.3	276.4	746.8	1,754.6	4,138.3	9,300.0
1957-----	369.4	7.7	1.6	1.5	5.4	16.9	77.3	278.2	761.8	1,783.1	4,155.4	9,314.3
1956-----	361.0	8.1	1.5	1.6	5.5	16.9	75.5	272.8	746.4	1,750.5	4,158.7	9,079.8
1955-----	356.5	7.4	1.6	1.8	5.6	17.2	77.4	275.3	738.9	1,739.7	4,144.3	8,917.2
1954-----	348.3	8.7	1.5	2.2	6.3	17.9	77.6	285.1	740.1	1,725.0	4,016.6	8,472.6
1953-----	361.3	9.5	1.6	2.7	6.9	19.4	84.4	297.1	783.7	1,794.6	4,213.3	8,789.1
1952-----	357.6	9.2	1.7	3.0	7.8	20.0	86.7	301.8	790.7	1,783.9	4,201.6	8,661.8
1951-----	357.0	8.2	1.5	3.2	7.7	20.4	87.5	304.6	793.1	1,798.8	4,258.7	8,973.5
1950 <sup>1</sup> -----	356.8	4.1	1.6	3.9	8.2	20.9	88.3	309.2	804.3	1,857.2	4,311.0	9,152.5
<b>WHITE MALE</b>												
1969-----	443.9	9.9	1.3	0.8	3.0	12.3	98.7	372.5	1,002.0	2,227.5	4,535.6	9,899.3
1968-----	451.4	11.3	1.3	0.8	2.8	12.6	99.9	382.5	1,030.0	2,289.7	4,600.3	10,046.9
1967-----	446.3	11.5	0.8	0.7	2.5	13.0	102.5	388.8	1,028.7	2,270.1	4,511.0	10,105.3
1966-----	453.3	8.6	0.9	0.8	2.8	14.6	102.5	396.3	1,051.8	2,325.5	4,637.1	10,242.3
1965-----	449.7	8.7	1.1	0.7	2.7	15.4	104.1	395.3	1,038.0	2,296.9	4,669.2	10,427.5
1964-----	447.1	7.8	1.1	0.6	3.2	15.9	106.7	395.1	1,033.9	2,306.4	4,616.7	10,215.2
1963 <sup>2</sup> -----	457.0	7.5	0.8	1.0	3.3	16.1	106.9	402.9	1,042.3	2,344.5	4,786.5	10,726.3
1962 <sup>2</sup> -----	450.8	6.0	0.9	0.8	3.5	16.5	105.9	404.4	1,032.1	2,292.5	4,709.5	10,461.4
1961-----	446.3	6.2	0.8	0.8	4.1	16.8	106.5	404.7	1,031.9	2,284.3	4,692.8	10,032.2
1960-----	454.6	6.9	1.0	1.1	3.6	17.6	107.5	413.2	1,056.0	2,297.9	4,839.9	10,135.8
1959-----	446.5	6.5	1.2	1.1	4.1	18.4	105.9	409.3	1,045.8	2,257.1	4,731.0	9,835.6
1958-----	450.0	6.6	1.2	1.1	4.4	18.8	108.7	406.4	1,056.2	2,279.9	4,836.5	9,959.5
1957-----	451.5	7.6	1.2	1.1	5.0	19.2	108.5	406.1	1,070.5	2,302.2	4,836.0	9,841.2
1956-----	443.3	7.0	1.2	1.4	5.1	19.2	106.5	400.0	1,048.1	2,257.0	4,847.1	9,542.1
1955-----	438.5	6.7	1.2	1.5	5.3	19.7	108.4	398.8	1,036.2	2,238.4	4,813.9	9,316.0
1954-----	429.8	8.4	1.3	1.8	6.1	19.9	106.1	405.3	1,031.2	2,206.9	4,637.1	8,854.3
1953-----	443.7	9.4	1.2	2.1	6.5	20.7	112.9	418.2	1,079.3	2,276.2	4,852.6	9,206.3
1952-----	437.9	9.1	1.5	2.2	7.7	21.3	114.6	421.4	1,078.7	2,243.8	4,818.4	9,019.0
1951-----	437.1	8.3	1.3	2.6	7.3	21.5	113.2	419.2	1,071.3	2,268.9	4,901.4	9,620.7
1950-----	434.2	4.6	1.3	3.1	7.1	21.5	112.6	424.1	1,082.6	2,309.4	4,908.0	9,952.3
<b>WHITE FEMALE</b>												
1969-----	315.4	9.6	1.2	0.8	1.9	5.2	24.6	90.8	318.1	1,034.2	3,149.0	9,642.0
1968-----	319.2	8.1	1.3	0.8	1.8	5.7	25.8	95.9	337.1	1,123.3	3,229.0	9,646.5
1967-----	310.2	8.3	0.8	0.6	1.6	5.8	26.1	95.3	332.7	1,123.2	3,198.5	9,517.5
1966-----	314.6	7.4	0.9	0.6	2.0	7.0	26.4	99.7	340.6	1,161.7	3,321.6	9,725.1
1965-----	310.6	7.3	0.9	0.6	2.2	7.1	26.4	100.1	344.6	1,156.6	3,348.5	9,747.7
1964-----	308.5	7.4	1.0	0.7	2.2	6.9	26.9	101.6	354.6	1,170.7	3,369.0	9,702.7
1963 <sup>2</sup> -----	312.0	6.1	0.8	0.7	2.4	7.6	28.4	101.3	362.3	1,195.7	3,470.4	9,893.7
1962 <sup>2</sup> -----	307.5	5.3	0.9	0.8	2.2	7.6	27.2	103.0	360.5	1,196.6	3,496.1	9,534.9
1961-----	301.3	5.2	0.9	0.8	2.7	7.8	27.5	103.8	362.3	1,194.9	3,475.2	9,157.5
1960-----	306.5	4.3	0.9	0.9	2.8	8.2	28.6	103.4	383.0	1,229.8	3,629.7	9,280.8
1959-----	301.7	4.8	0.9	1.0	3.0	8.8	28.3	106.0	377.8	1,230.6	3,614.0	9,194.8
1958-----	305.0	5.4	1.1	1.0	3.1	8.7	28.5	110.2	393.9	1,264.4	3,719.1	9,351.9
1957-----	305.2	5.8	1.1	1.1	3.8	9.6	29.4	111.5	400.0	1,297.5	3,746.6	9,445.7
1956-----	297.2	6.6	0.9	1.3	4.0	9.2	29.0	109.1	394.2	1,278.7	3,743.5	9,275.1
1955-----	293.0	5.6	1.2	1.5	4.1	9.5	31.2	111.4	396.2	1,279.0	3,740.9	9,155.9
1954-----	284.1	6.2	1.0	1.9	4.3	10.1	32.1	118.9	402.9	1,282.7	3,628.6	8,655.7
1953-----	293.9	6.9	1.4	2.3	4.8	11.8	36.6	127.8	431.0	1,344.4	3,820.9	8,928.3
1952-----	290.5	6.6	1.3	2.7	5.3	12.0	38.8	132.0	438.6	1,349.2	3,820.3	8,842.6
1951-----	289.5	6.6	1.4	2.5	5.7	12.7	40.3	139.0	448.1	1,356.3	3,851.5	8,984.6
1950-----	290.5	2.9	1.4	3.7	6.6	13.6	42.4	142.4	460.7	1,401.6	3,926.2	9,086.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 1.—Death Rates for Diseases of Heart, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 390-398,402,404,410-429 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 400-402,410-443 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	316.1	15.9	2.6	1.7	7.3	40.8	180.4	505.2	1,132.6	2,540.3	3,402.0	4,341.0
1968-----	327.1	22.2	3.3	2.1	6.9	39.9	183.1	504.8	1,210.3	2,650.1	3,376.1	4,998.2
1967-----	310.3	19.9	2.9	1.7	6.5	41.1	175.8	487.1	1,120.9	2,487.4	3,245.2	4,896.0
1966-----	323.7	23.6	3.7	1.8	6.9	43.6	180.4	499.3	1,174.1	2,594.6	3,329.2	5,363.0
1965-----	316.5	18.6	2.7	1.8	8.2	42.2	176.7	494.5	1,147.8	2,462.8	3,283.6	5,746.3
1964-----	316.6	20.5	3.8	2.0	6.4	39.2	179.2	492.9	1,180.2	2,432.0	3,182.1	5,835.1
1963 <sup>2</sup> -----	330.1	13.6	3.5	1.9	8.3	40.5	171.0	506.0	1,179.7	2,505.2	3,456.3	6,842.4
1962 <sup>2</sup> -----	324.2	15.1	2.2	2.2	7.9	40.4	170.0	506.7	1,148.0	2,364.7	3,369.2	6,967.7
1961-----	313.9	15.8	2.8	2.3	8.3	39.3	162.3	482.7	1,127.2	2,264.0	3,262.0	6,464.5
1960-----	320.5	13.1	3.8	3.1	8.7	40.1	158.9	487.4	1,175.6	2,216.9	3,323.1	6,128.6
1959-----	317.2	12.0	5.2	2.7	9.7	43.8	161.8	506.7	1,167.7	2,153.0	3,382.6	6,224.1
1958-----	326.5	15.2	2.5	2.4	9.1	44.2	158.4	525.3	1,206.0	2,228.7	3,522.3	6,451.9
1957-----	336.2	15.0	4.4	3.8	11.9	44.8	175.8	532.9	1,297.6	2,211.1	3,445.9	6,692.2
1956-----	322.9	16.5	3.4	3.6	12.3	42.0	160.5	521.5	1,240.3	2,116.2	3,434.6	6,224.0
1955-----	319.4	12.6	4.5	3.9	12.0	40.2	166.6	540.0	1,219.8	2,063.5	3,401.6	5,720.8
1954-----	316.6	19.8	4.0	4.3	12.9	44.7	167.1	548.2	1,232.6	1,981.0	3,305.0	5,854.5
1953-----	343.6	17.9	4.3	5.5	15.9	51.5	186.4	596.6	1,348.7	2,110.2	3,521.1	6,485.7
1952-----	344.7	18.9	4.0	6.7	17.8	49.4	185.5	594.1	1,379.5	2,146.0	3,580.0	6,275.0
1951-----	341.0	15.0	3.3	6.7	15.2	49.4	194.7	591.4	1,367.7	2,112.2	3,571.7	6,315.8
1950 <sup>1</sup> -----	342.0	5.9	4.3	6.3	17.9	50.2	191.4	604.5	1,314.6	2,321.0	3,679.8	6,262.1
ALL OTHER FEMALE												
1969-----	246.7	17.8	2.0	2.3	6.9	24.2	98.2	267.6	728.9	1,870.7	2,532.7	4,570.0
1968-----	258.9	24.0	3.0	1.8	6.2	26.9	105.2	288.1	775.7	1,938.5	2,693.2	4,951.4
1967-----	242.3	17.3	2.0	1.7	6.2	29.3	102.8	287.0	736.7	1,803.9	2,513.1	4,707.6
1966-----	252.8	22.8	2.6	2.0	6.3	30.0	103.8	295.0	787.0	1,850.2	2,743.1	5,113.3
1965-----	249.2	16.1	2.8	2.2	7.1	29.3	110.3	300.9	825.7	1,724.6	2,711.0	5,364.8
1964-----	253.2	16.4	2.9	2.4	9.0	31.3	109.3	317.8	869.5	1,732.8	2,706.0	5,630.6
1963 <sup>2</sup> -----	262.0	11.3	3.5	2.3	9.8	31.3	115.8	331.3	907.4	1,761.2	2,837.2	6,088.6
1962 <sup>2</sup> -----	258.0	10.6	2.3	2.4	10.2	30.3	112.7	325.5	901.8	1,731.0	2,811.1	5,997.6
1961-----	252.9	10.3	2.7	2.2	7.6	30.7	111.6	337.8	899.6	1,613.4	2,805.4	6,039.0
1960-----	255.5	11.7	2.8	2.9	9.6	33.7	118.2	346.8	909.0	1,638.4	2,897.7	5,564.1
1959-----	252.4	18.5	3.1	3.4	9.7	35.4	117.2	365.0	871.7	1,613.0	2,787.0	5,402.6
1958-----	266.9	15.9	3.3	3.1	9.8	39.0	123.3	411.7	970.1	1,668.2	2,835.3	5,431.6
1957-----	273.5	11.1	4.4	4.2	12.3	39.0	138.0	428.9	1,013.8	1,701.9	2,896.6	5,121.6
1956-----	263.8	15.7	4.9	3.6	11.6	37.4	136.5	404.2	1,004.6	1,641.5	2,843.5	4,830.6
1955-----	256.8	16.3	4.1	4.5	12.3	39.2	132.5	424.3	941.2	1,582.2	2,840.9	4,811.8
1954-----	254.0	15.6	3.2	4.7	15.6	42.0	145.6	445.6	922.9	1,555.1	2,667.5	4,659.4
1953-----	271.4	17.9	2.9	6.7	17.1	42.7	161.9	479.5	1,017.1	1,629.5	2,846.3	5,077.4
1952-----	275.5	16.9	3.3	6.7	17.3	50.0	169.9	506.2	1,072.9	1,609.5	2,828.4	4,996.6
1951-----	278.9	10.4	2.6	8.7	18.4	49.8	177.0	513.8	1,101.0	1,606.9	2,920.7	5,167.9
1950 <sup>1</sup> -----	283.0	6.4	3.1	8.9	19.6	50.9	181.7	517.6	1,073.9	1,905.1	3,005.7	5,326.0

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 2.—Death Rates for Malignant Neoplasms, Including Neoplasms of Lymphatic and Hematopoietic Tissues, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 140-209 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 140-205 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	160.0	4.7	7.3	6.1	8.0	16.9	61.4	178.8	419.7	769.4	1,112.0	1,486.5
1968-----	159.4	4.8	7.8	6.3	8.2	17.3	61.2	182.0	418.3	774.2	1,104.4	1,477.6
1967-----	157.2	5.5	8.2	6.6	8.3	17.5	60.3	179.4	412.4	768.8	1,102.1	1,482.9
1966-----	155.1	5.6	8.3	6.4	8.4	18.3	61.1	180.1	406.7	757.6	1,098.7	1,483.7
1965-----	153.5	5.9	8.6	6.5	8.2	19.0	60.7	178.2	406.6	752.2	1,093.9	1,493.3
1964-----	151.3	5.8	9.2	6.5	8.3	19.4	61.0	178.2	401.4	737.5	1,081.8	1,469.8
1963 <sup>2</sup> -----	151.3	6.4	10.0	6.9	8.4	18.8	59.5	176.8	399.7	740.6	1,090.0	1,525.2
1962-----	149.9	6.3	9.7	6.9	8.4	18.8	59.0	176.9	395.6	728.3	1,081.6	1,497.2
1961-----	149.4	6.9	10.4	7.1	8.3	19.4	59.4	175.6	392.6	723.1	1,104.9	1,491.2
1960-----	149.2	7.2	10.9	6.8	8.3	19.5	59.7	177.0	396.8	713.9	1,127.4	1,450.0
1959-----	147.3	6.8	10.6	7.1	8.5	19.9	59.4	174.3	394.4	706.2	1,101.8	1,447.3
1958-----	146.8	6.5	10.9	6.8	8.8	19.4	60.2	173.5	392.4	707.6	1,112.8	1,525.2
1957-----	148.6	7.0	11.0	6.8	8.7	20.1	58.9	176.1	399.3	717.3	1,140.2	1,489.7
1956-----	147.8	7.9	11.0	7.2	8.6	20.1	60.8	173.9	399.5	706.3	1,161.5	1,483.7
1955-----	146.5	7.7	11.1	7.0	9.2	19.3	61.3	174.6	392.3	711.5	1,146.4	1,465.3
1954-----	145.6	8.4	11.3	6.7	9.1	19.7	61.5	176.3	390.3	708.3	1,151.8	1,447.7
1953-----	144.8	8.5	10.9	6.6	8.9	19.5	62.1	176.7	396.1	699.8	1,154.5	1,456.0
1952-----	143.4	8.5	11.6	6.5	9.4	20.3	61.4	175.8	394.3	697.4	1,152.9	1,455.4
1951-----	140.6	8.2	11.8	6.5	9.3	19.7	61.7	175.9	388.7	686.5	1,133.4	1,452.4
1950 <sup>1</sup> -----	139.8	8.7	11.7	6.7	8.6	20.0	62.7	175.1	390.7	698.8	1,153.3	1,451.0
<b>WHITE MALE</b>												
1969-----	181.6	5.2	8.3	7.2	10.3	16.8	51.3	168.8	487.4	992.5	1,507.7	1,941.0
1968-----	180.8	4.8	8.8	7.2	10.3	17.2	50.7	171.5	486.7	1,000.3	1,485.8	1,913.3
1967-----	177.9	5.9	9.3	7.7	10.2	17.1	49.2	169.5	478.5	985.2	1,476.7	1,914.1
1966-----	174.9	5.6	9.5	7.7	10.4	17.5	50.1	169.5	472.5	968.0	1,454.1	1,874.5
1965-----	173.2	6.1	9.8	7.6	10.1	18.0	49.7	167.6	471.4	961.8	1,432.1	1,916.4
1964-----	170.2	6.2	10.7	7.4	9.9	18.6	49.4	166.3	464.4	942.5	1,402.3	1,847.1
1963 <sup>2</sup> -----	168.3	7.9	11.7	7.8	10.1	18.0	46.8	164.4	453.9	931.0	1,390.6	1,919.9
1962 <sup>2</sup> -----	166.5	6.3	11.1	7.7	10.4	17.8	46.7	163.8	451.2	905.8	1,383.5	1,886.7
1961-----	166.8	7.1	12.0	8.4	10.2	18.5	46.1	162.6	445.5	906.1	1,409.7	1,864.9
1960-----	166.1	7.9	13.1	8.0	10.3	18.8	46.3	164.1	450.9	887.3	1,413.7	1,791.4
1959-----	163.4	7.2	12.1	8.6	10.4	18.2	46.4	162.4	443.5	870.7	1,379.1	1,808.6
1958-----	161.6	7.4	12.5	7.6	10.9	17.8	46.5	159.4	438.3	862.5	1,387.5	1,769.0
1957-----	163.8	7.3	12.9	8.0	10.7	18.2	45.1	159.6	443.2	831.1	1,416.1	1,806.8
1956-----	162.4	8.5	12.1	8.1	10.5	18.0	46.2	157.6	442.0	855.3	1,445.0	1,823.8
1955-----	160.0	8.7	13.1	8.6	11.3	17.5	44.3	157.8	429.1	859.8	1,409.1	1,746.9
1954-----	158.2	9.5	12.7	7.8	10.8	18.1	45.4	155.6	424.7	846.0	1,403.9	1,740.8
1953-----	156.7	8.5	12.4	7.9	10.9	17.6	46.9	156.8	431.0	822.4	1,398.0	1,735.1
1952-----	153.8	8.1	12.8	7.2	11.4	17.8	44.4	153.0	422.2	820.6	1,383.1	1,722.9
1951-----	149.5	9.2	13.8	7.6	11.5	17.3	44.2	152.8	409.5	800.2	1,346.4	1,748.9
1950-----	147.2	9.6	13.1	7.6	9.9	17.7	44.5	150.8	409.4	798.7	1,367.6	1,732.7
<b>WHITE FEMALE</b>												
1969-----	146.3	5.1	7.3	5.4	6.0	15.8	64.5	171.8	333.9	560.5	869.9	1,349.4
1968-----	145.6	5.1	7.5	5.9	6.4	16.2	65.1	176.7	332.4	562.1	866.4	1,330.6
1967-----	144.2	5.3	8.3	6.1	6.6	16.5	65.3	174.5	330.2	568.9	866.4	1,321.6
1966-----	142.6	6.3	8.2	5.6	6.6	17.6	64.9	174.8	326.4	564.7	870.9	1,349.7
1965-----	141.9	6.2	8.8	6.0	6.4	18.7	65.2	175.1	327.3	564.2	880.3	1,313.3
1964-----	140.0	5.3	8.3	5.9	7.1	19.4	65.9	175.0	323.2	557.2	875.8	1,313.6
1963 <sup>2</sup> -----	139.2	5.2	9.1	6.3	7.0	18.1	64.3	171.5	323.9	559.7	889.0	1,341.5
1962 <sup>2</sup> -----	138.6	7.2	9.2	6.4	6.6	18.7	64.6	173.9	320.3	559.2	895.2	1,300.6
1961-----	139.4	7.6	10.0	6.4	6.6	19.5	65.7	174.5	323.8	563.1	907.2	1,313.1
1960-----	139.8	6.8	9.7	6.2	6.5	18.8	66.6	175.7	329.0	562.1	939.3	1,304.9
1959-----	138.7	6.5	10.4	6.2	6.4	20.3	66.3	173.7	331.2	563.9	922.0	1,282.1
1958-----	139.4	6.1	10.6	6.4	7.0	19.4	68.1	173.6	332.6	574.5	938.1	1,311.1
1957-----	141.0	7.6	10.6	6.1	7.0	21.2	67.2	178.7	339.8	576.7	964.5	1,360.4
1956-----	140.7	7.3	11.2	6.7	7.0	20.9	69.9	175.8	341.5	583.4	976.4	1,335.3
1955-----	141.0	7.2	10.6	6.0	7.4	19.9	71.8	176.5	343.6	592.3	985.1	1,357.6
1954-----	140.7	7.9	11.2	6.3	7.2	20.0	70.9	181.5	341.8	598.8	994.8	1,327.5
1953-----	140.9	8.6	10.6	5.9	7.2	20.2	71.5	181.7	351.1	603.5	1,008.6	1,350.1
1952-----	141.0	8.7	11.5	6.0	7.7	21.7	72.0	183.6	355.8	603.4	1,014.1	1,360.5
1951-----	139.5	8.0	10.8	6.1	7.4	20.8	73.0	185.0	356.1	601.9	1,012.4	1,340.4
1950-----	139.9	7.8	11.3	6.3	7.5	20.9	74.5	185.8	362.5	616.5	1,026.6	1,348.3

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 2.—Death Rates for Malignant Neoplasms, Including Neoplasms of Lymphatic and Hematopoietic Tissues, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 140-209 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 140-205 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	159.1	2.9	5.1	5.0	7.8	18.2	80.6	285.0	691.2	1,300.2	1,337.7	1,044.3
1968-----	157.7	3.6	6.6	5.3	8.2	18.2	80.3	282.6	671.9	1,311.5	1,300.0	1,087.3
1967-----	153.1	5.6	5.3	6.0	8.8	19.1	79.1	277.4	644.6	1,247.1	1,231.2	1,190.0
1966-----	150.1	3.6	5.4	5.2	7.7	19.5	82.0	278.3	617.6	1,212.9	1,180.5	1,252.2
1965-----	143.4	5.6	5.3	5.2	8.4	18.7	76.4	252.4	603.7	1,150.1	1,145.5	1,209.8
1964-----	142.1	6.0	8.5	6.4	8.4	21.0	77.7	258.4	588.4	1,074.8	1,128.3	1,227.0
1963 <sup>2</sup> -----	142.3	5.6	8.4	5.2	8.7	19.3	74.7	257.3	577.5	1,093.5	1,082.2	1,327.3
1962 <sup>2</sup> -----	135.8	4.0	6.4	6.2	8.4	17.2	69.7	248.2	545.0	1,005.2	1,059.2	1,280.6
1961-----	135.9	4.5	6.8	4.7	9.2	18.4	70.1	237.4	567.7	951.3	1,079.1	1,393.5
1960-----	134.1	6.5	8.2	4.9	9.2	18.2	71.7	233.6	549.8	927.5	1,086.1	1,211.7
1959-----	131.7	7.3	7.1	5.4	9.8	19.9	66.9	229.0	551.7	897.0	1,067.4	1,248.3
1958-----	126.4	3.8	7.0	5.8	8.8	20.9	63.2	227.1	532.3	862.0	988.5	1,018.5
1957-----	129.2	4.9	8.4	5.9	8.7	17.4	60.0	234.9	565.7	860.4	978.5	1,115.4
1956-----	125.4	7.7	7.8	6.1	9.8	19.2	59.6	221.8	562.4	778.4	1,048.5	1,092.0
1955-----	119.9	6.9	6.7	5.9	10.0	17.0	60.5	221.9	530.0	759.1	913.6	1,045.8
1954-----	121.5	8.9	7.1	5.1	12.6	17.8	62.2	226.6	530.6	761.1	942.0	1,022.7
1953-----	115.2	9.0	7.3	5.1	10.1	15.5	57.9	221.6	510.9	725.9	832.5	947.6
1952-----	112.5	9.9	8.2	6.1	9.5	16.1	59.1	214.4	503.6	676.4	889.1	940.0
1951-----	107.2	5.5	8.0	3.6	10.1	18.1	51.5	209.7	495.3	650.2	782.1	942.1
1950 <sup>1</sup> -----	106.1	10.4	8.2	5.7	7.7	17.9	56.0	207.4	457.1	697.8	844.4	916.0
ALL OTHER FEMALE												
1969-----	112.4	1.9	5.0	4.5	6.2	24.3	92.8	229.1	423.9	717.9	641.5	618.8
1968-----	113.4	4.7	4.6	4.2	7.1	25.0	92.0	226.4	429.9	724.5	654.7	715.3
1967-----	111.2	4.7	5.0	4.2	5.7	26.4	88.0	226.9	421.7	699.8	658.6	754.5
1966-----	111.3	4.3	5.6	4.8	7.2	28.3	95.1	228.9	414.8	665.9	695.0	731.7
1965-----	109.4	3.5	4.7	4.2	7.4	28.8	95.3	228.2	421.9	633.8	636.6	855.6
1964-----	108.3	5.2	5.7	4.9	6.3	24.0	95.1	236.8	427.4	599.4	640.5	773.5
1963 <sup>2</sup> -----	108.8	5.7	7.4	5.3	6.4	28.3	100.5	233.0	435.1	578.3	619.6	790.9
1962 <sup>2</sup> -----	109.5	5.0	6.9	6.2	6.9	28.0	99.8	230.4	434.5	576.9	652.4	788.1
1961-----	109.3	4.2	6.6	5.0	6.2	25.9	102.3	241.8	423.5	564.2	648.4	836.6
1960-----	109.8	6.5	7.0	4.8	6.7	29.6	98.3	249.3	427.8	537.6	702.3	727.5
1959-----	106.7	6.0	6.1	4.3	8.1	29.1	100.5	236.3	425.1	519.2	625.5	710.3
1958-----	110.1	5.5	7.5	5.4	7.4	30.3	102.0	250.8	437.2	528.7	639.2	734.2
1957-----	109.3	3.1	4.6	4.3	7.5	28.7	99.4	256.3	435.8	531.1	663.4	683.8
1956-----	111.5	7.3	7.0	5.1	6.8	29.6	113.1	262.7	440.4	513.6	663.0	691.7
1955-----	108.4	5.3	6.3	4.2	6.8	29.5	112.6	266.7	414.4	506.4	623.5	641.2
1954-----	108.7	4.0	6.1	4.5	7.4	30.7	112.6	273.2	434.8	475.1	623.8	634.4
1953-----	107.9	7.2	6.0	3.6	7.2	31.9	111.9	271.9	417.1	486.7	638.0	654.8
1952-----	108.7	8.9	6.6	4.8	7.9	31.4	115.1	278.2	423.4	490.2	615.5	627.6
1951-----	107.8	5.0	7.9	4.4	7.3	31.0	119.3	274.1	435.5	474.3	585.6	617.9
1950 <sup>1</sup> -----	110.1	6.9	7.1	4.0	8.6	33.5	119.3	273.3	437.7	552.7	578.9	716.0

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 3.—Death Rates for Cerebrovascular Diseases, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 430-438 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 330-334 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969	102.6	4.8	0.9	0.7	1.6	5.0	15.9	41.6	117.0	406.5	1,232.3	3,479.0
1968	105.8	5.2	0.8	0.7	1.7	4.9	16.6	44.3	123.1	423.2	1,283.5	3,611.7
1967	102.2	3.0	0.5	0.6	1.4	4.9	15.4	43.5	120.9	412.6	1,261.7	3,592.5
1966	104.6	2.8	0.7	0.7	1.6	4.8	15.6	43.4	124.1	432.0	1,319.7	3,722.5
1965	103.7	4.0	0.7	0.7	1.5	4.8	15.4	45.0	127.9	430.0	1,322.2	3,741.8
1964	103.6	3.1	0.8	0.6	1.7	4.9	15.2	46.3	132.9	434.6	1,337.7	3,678.4
1963	106.6	3.5	0.7	0.6	1.6	4.8	15.2	46.3	138.3	450.5	1,402.7	3,824.1
1962	106.2	3.1	0.8	0.7	1.6	4.7	15.3	47.1	138.5	450.7	1,404.7	3,812.6
1961	105.4	3.4	0.9	0.7	1.7	4.8	14.7	46.9	139.0	451.8	1,421.5	3,709.1
1960	108.0	4.1	0.8	0.7	1.8	4.7	14.7	49.2	147.3	469.2	1,481.3	3,680.5
1959	108.4	4.1	1.0	0.7	1.7	4.8	15.0	49.3	150.8	478.4	1,508.1	3,640.9
1958	110.1	4.6	1.0	0.7	1.8	4.9	15.5	51.7	156.5	495.0	1,539.5	3,676.9
1957	110.2	4.9	1.1	0.8	2.0	4.9	16.6	55.2	161.8	508.2	1,525.4	3,581.9
1956	106.3	4.6	1.1	0.7	1.9	4.8	16.2	54.6	161.1	497.6	1,489.8	3,389.3
1955	106.0	5.6	0.9	0.7	1.9	5.0	16.6	56.2	163.2	506.0	1,497.6	3,290.6
1954	104.1	5.1	0.9	0.7	1.9	4.6	17.9	61.2	172.0	503.8	1,460.0	3,057.3
1953	107.3	5.8	1.0	0.6	1.7	5.1	19.3	67.1	183.8	531.3	1,503.1	3,051.5
1952	106.8	5.5	1.1	0.7	2.0	5.0	19.1	69.4	189.2	534.8	1,500.4	3,060.1
1951	106.7	4.9	0.9	0.6	1.8	4.8	19.5	70.6	192.3	550.9	1,516.4	2,997.1
1950 <sup>1</sup>	104.0	5.1	0.9	0.5	1.6	4.2	18.7	70.4	194.2	554.7	1,499.6	2,990.1
<b>WHITE MALE</b>												
1969	94.4	4.6	1.0	0.6	1.4	3.5	11.3	35.4	120.3	437.3	1,334.0	3,389.7
1968	97.4	5.0	0.7	0.7	1.5	3.6	12.0	37.9	123.9	454.1	1,376.0	3,563.5
1967	95.1	2.7	0.8	0.6	1.2	3.4	11.0	36.4	121.9	443.3	1,363.7	3,576.8
1966	96.9	2.3	0.7	0.8	1.5	3.4	11.3	36.8	125.8	458.1	1,404.7	3,641.6
1965	96.3	4.8	0.6	0.6	1.4	3.5	10.9	37.3	126.9	455.5	1,400.6	3,733.4
1964	96.7	3.2	0.8	0.5	1.8	3.9	10.9	38.3	129.5	462.2	1,418.1	3,689.5
1963 <sup>2</sup>	100.4	3.6	0.8	0.6	1.5	4.0	11.2	39.8	137.3	475.6	1,479.1	3,841.5
1962 <sup>2</sup>	100.3	3.1	0.9	0.7	1.4	3.5	11.1	39.3	131.3	478.7	1,492.7	3,826.8
1961	99.7	3.4	0.9	0.6	1.7	4.0	10.8	39.3	132.1	478.4	1,501.0	3,708.7
1960	102.7	4.3	0.8	0.7	1.7	3.5	11.3	40.9	139.0	501.0	1,564.8	3,744.8
1959	103.1	4.4	0.9	0.8	1.7	3.7	10.9	41.4	142.4	505.9	1,579.6	3,649.1
1958	104.7	4.6	1.1	0.7	1.9	3.7	11.3	42.7	148.5	522.1	1,611.2	3,616.1
1957	105.9	4.7	1.0	0.8	2.0	4.0	12.2	45.4	155.0	535.0	1,609.8	3,560.8
1956	102.2	4.5	0.8	0.7	1.8	3.7	12.2	44.3	153.5	524.3	1,563.3	3,336.1
1955	102.3	5.0	1.0	0.8	1.6	4.2	11.6	45.3	156.4	531.4	1,569.9	3,262.6
1954	100.4	4.8	0.9	0.7	1.7	3.6	13.2	47.6	163.1	530.6	1,530.1	2,941.5
1953	103.6	6.1	0.9	0.7	1.8	4.0	14.2	51.2	175.7	552.7	1,570.4	2,982.5
1952	103.4	5.7	1.2	0.7	1.8	4.2	13.7	54.3	180.2	553.1	1,572.0	3,015.0
1951	103.0	5.6	0.9	0.6	1.4	3.8	14.5	54.2	182.5	575.7	1,558.9	2,984.8
1950	100.5	5.9	1.1	0.5	1.6	3.4	13.1	53.7	182.2	569.7	1,556.3	3,127.1
<b>WHITE FEMALE</b>												
1969	109.5	3.9	0.6	0.7	1.6	3.8	11.8	29.6	78.3	314.1	1,164.0	3,854.7
1968	112.0	4.1	0.8	0.6	1.4	3.6	12.0	32.0	82.6	324.6	1,222.5	3,927.4
1967	107.9	2.8	0.4	0.6	1.3	3.9	11.2	31.0	82.2	320.9	1,201.1	3,885.0
1966	110.0	2.8	0.5	0.6	1.2	3.6	10.7	29.7	82.8	339.8	1,262.3	4,004.9
1965	108.0	2.6	0.6	0.6	1.2	3.8	11.0	30.9	83.4	336.1	1,268.1	3,982.7
1964	107.7	1.9	0.7	0.6	1.3	3.5	10.8	32.4	87.8	350.7	1,283.6	3,888.8
1963 <sup>2</sup>	110.9	2.4	0.6	0.5	1.4	3.7	10.4	32.1	92.0	363.8	1,360.5	4,022.5
1962 <sup>2</sup>	110.3	2.3	0.7	0.7	1.4	3.5	10.9	33.2	96.2	366.2	1,362.3	3,977.9
1961	108.5	2.7	0.7	0.6	1.4	3.3	10.2	32.4	94.3	370.6	1,378.1	3,852.6
1960	110.1	2.6	0.5	0.6	1.4	3.4	10.1	33.8	103.0	383.3	1,444.7	3,795.7
1959	110.6	2.9	1.0	0.5	1.4	3.8	10.1	33.4	108.7	397.6	1,465.5	3,783.7
1958	111.9	3.5	0.7	0.7	1.3	3.6	10.8	36.1	112.4	412.4	1,496.6	3,890.0
1957	110.9	3.8	1.0	0.6	1.5	3.0	11.1	39.1	116.9	430.0	1,479.8	3,756.7
1956	107.0	3.0	1.0	0.7	1.6	3.3	11.3	37.7	117.9	426.4	1,454.3	3,576.3
1955	106.2	4.1	0.7	0.6	1.6	3.5	11.6	40.5	119.9	455.8	1,465.5	3,470.8
1954	103.9	3.9	0.7	0.6	1.8	3.3	12.2	44.7	132.6	436.1	1,424.1	3,276.8
1953	107.1	3.6	0.9	0.5	1.3	3.4	13.9	51.2	143.0	470.9	1,472.8	3,242.3
1952	105.9	2.7	1.0	0.6	1.5	3.6	13.9	53.0	147.7	476.0	1,463.6	3,206.0
1951	106.2	3.7	0.8	0.7	1.5	3.5	14.0	55.7	152.9	487.6	1,499.4	3,139.5
1950	103.3	2.9	0.6	0.4	1.2	2.9	13.6	55.0	156.9	498.1	1,471.3	3,017.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 3.—Death Rates for Cerebrovascular Diseases, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 430-438 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 330-334 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	105.2	7.3	1.3	0.9	2.8	14.4	53.2	136.3	327.7	904.2	1,506.9	1,714.8
1968-----	111.5	8.2	1.3	0.8	3.4	14.1	53.7	139.3	364.7	980.8	1,328.4	1,909.1
1967-----	104.9	4.2	0.5	0.8	2.2	12.8	48.4	138.0	334.9	931.2	1,237.2	1,906.0
1966-----	111.4	4.9	1.0	1.0	2.9	13.1	52.7	145.7	352.2	963.3	1,548.2	2,097.8
1965-----	113.5	6.8	1.4	1.0	2.5	11.7	54.0	150.8	374.8	969.8	1,354.5	2,073.2
1964-----	111.4	6.0	1.0	1.1	2.3	12.8	48.0	147.4	385.7	904.4	1,339.7	2,254.1
1963 <sup>2</sup> -----	116.6	5.2	1.1	1.1	3.0	11.7	46.6	149.0	388.6	946.5	1,420.7	2,630.3
1962 <sup>2</sup> -----	115.6	5.8	1.2	1.1	2.6	11.2	49.5	150.7	398.0	900.2	1,340.8	2,732.3
1961-----	111.0	6.3	1.3	0.9	2.6	11.7	49.3	143.9	378.3	840.5	1,287.7	2,754.8
1960-----	118.2	9.5	1.8	0.8	3.9	12.1	44.4	156.7	411.3	868.2	1,445.0	2,675.2
1959-----	117.3	9.3	1.5	1.1	2.6	11.2	50.3	156.9	417.2	864.4	1,401.4	2,534.5
1958-----	122.5	8.3	2.4	1.2	2.9	13.1	46.0	160.7	447.1	909.0	1,470.5	2,640.7
1957-----	121.9	11.5	1.6	1.0	3.6	13.8	52.9	169.0	449.8	879.4	1,391.9	2,442.3
1956-----	116.5	9.6	2.0	0.3	2.9	14.8	47.2	167.7	435.7	828.6	1,343.8	2,312.0
1955-----	118.0	12.3	1.8	0.9	3.7	13.4	56.7	163.4	451.0	830.9	1,353.6	2,241.7
1954-----	117.3	9.3	1.8	1.4	3.7	12.3	54.9	186.1	450.2	791.5	1,313.4	2,190.9
1953-----	122.0	12.8	1.6	0.7	3.6	14.3	56.5	205.4	477.4	813.1	1,314.0	2,100.0
1952-----	122.9	14.9	1.0	1.0	4.5	11.3	55.3	200.5	497.7	806.9	1,364.5	2,345.0
1951-----	121.1	8.2	1.9	1.0	4.5	13.1	54.8	202.6	482.1	806.4	1,394.0	2,189.5
1950 <sup>1</sup> -----	119.5	10.4	1.0	0.7	3.5	11.9	57.4	204.2	479.3	847.8	1,365.7	2,132.0
ALL OTHER FEMALE												
1969-----	106.9	7.4	0.9	0.5	2.3	15.0	46.7	114.2	283.0	816.3	1,179.4	1,926.3
1968-----	112.8	7.7	1.0	0.7	3.7	14.4	51.8	121.6	301.6	862.3	1,217.0	2,183.3
1967-----	107.3	4.7	0.4	0.8	2.6	15.0	50.5	129.3	303.8	799.2	1,143.8	2,069.4
1966-----	112.9	3.0	1.2	0.9	3.5	15.2	51.9	127.8	312.1	851.4	1,244.8	2,441.7
1965-----	115.9	4.1	1.4	1.0	3.1	14.5	50.1	141.3	354.1	861.8	1,280.4	2,422.2
1964-----	115.1	5.5	1.4	0.8	3.0	15.1	53.2	147.0	381.1	797.7	1,268.4	2,373.5
1963 <sup>2</sup> -----	120.3	6.6	0.8	0.7	2.3	13.6	55.6	147.8	397.0	839.3	1,363.9	2,602.3
1962 <sup>2</sup> -----	120.8	4.7	1.1	0.9	3.2	15.9	55.8	146.9	406.1	821.0	1,367.2	2,776.2
1961-----	118.2	4.2	1.2	1.0	3.6	14.8	51.5	153.9	410.7	802.2	1,300.0	2,607.3
1960-----	121.2	6.5	1.4	1.1	3.2	16.2	54.2	160.3	429.3	804.4	1,389.2	2,510.2
1959-----	123.3	4.3	1.7	0.6	3.7	14.0	57.5	166.9	426.0	820.6	1,423.6	2,548.7
1958-----	124.9	7.9	1.3	1.0	3.1	16.7	61.6	178.8	435.7	833.4	1,402.6	2,334.2
1957-----	126.6	5.9	2.0	1.2	3.7	18.5	67.0	189.5	453.3	831.8	1,335.9	2,402.7
1956-----	123.0	9.5	2.4	0.8	3.3	15.3	61.7	201.0	471.0	763.0	1,279.0	2,330.6
1955-----	121.8	12.9	0.9	0.8	3.9	15.3	62.2	205.2	465.8	779.9	1,268.2	2,073.5
1954-----	122.8	10.0	1.5	1.0	3.2	15.3	70.1	224.7	472.0	764.0	1,265.9	1,978.1
1953-----	125.1	11.5	1.4	0.9	3.1	18.5	73.3	231.5	492.3	781.2	1,287.6	2,009.7
1952-----	127.3	10.2	1.6	0.9	3.8	17.1	74.0	239.5	515.3	788.8	1,264.7	2,175.9
1951-----	127.6	5.4	0.2	0.5	3.9	15.8	74.6	237.1	533.9	796.0	1,355.0	1,960.7
1950 <sup>1</sup> -----	124.9	8.8	1.4	0.6	4.0	15.2	73.7	244.3	504.2	861.4	1,314.0	2,125.1

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 4.—Death Rates for Accidents, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E800-E949 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E800-E962 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	57.6	69.4	31.4	19.8	72.1	53.2	49.4	55.1	65.3	89.0	175.4	489.0
1968-----	57.5	75.6	31.4	20.3	69.5	52.5	47.8	53.3	65.5	95.0	183.9	514.5
1967-----	57.2	77.7	32.4	19.3	67.1	52.0	47.8	54.1	66.0	91.2	189.7	573.8
1966-----	58.0	84.5	33.4	19.8	67.1	51.9	47.9	55.6	67.5	92.5	199.2	591.9
1965-----	55.7	86.0	31.8	18.7	61.7	49.3	46.5	54.4	64.4	90.8	200.0	606.1
1964-----	54.3	84.1	31.2	18.8	59.2	48.6	45.2	52.2	63.2	90.2	197.3	582.1
1963-----	53.4	85.9	31.1	18.3	57.2	45.2	43.0	51.4	62.5	90.0	205.0	654.9
1962-----	52.3	87.8	30.5	18.0	55.1	43.5	42.1	50.1	60.4	90.1	205.9	630.0
1961-----	50.4	84.2	30.7	18.1	54.0	42.4	40.1	47.8	57.7	84.6	199.8	600.5
1960-----	52.3	93.2	31.6	19.3	56.0	43.0	40.9	49.8	59.0	89.1	214.5	624.7
1959-----	52.2	94.7	30.5	18.9	56.5	43.5	40.7	49.4	59.3	87.7	211.2	648.1
1958-----	52.3	96.9	31.0	19.5	56.2	42.5	41.5	47.2	57.6	89.4	221.3	655.1
1957-----	55.9	91.7	30.9	19.8	59.6	46.1	43.3	51.3	63.4	97.8	249.1	746.6
1956-----	56.7	91.3	31.0	20.1	63.6	45.9	43.7	51.2	63.5	101.3	250.7	758.7
1955-----	56.9	88.1	32.6	20.2	60.6	45.6	44.0	52.1	64.4	101.4	262.1	782.9
1954-----	55.9	98.2	33.5	20.4	56.8	44.3	41.9	50.4	62.5	101.4	267.0	780.0
1953-----	60.1	105.0	36.0	22.0	61.8	47.7	45.0	54.6	67.7	107.6	288.6	842.8
1952-----	61.8	109.4	37.2	22.4	61.2	48.3	47.2	55.9	71.1	111.7	300.3	897.7
1951-----	62.5	112.5	36.3	23.5	57.8	48.9	49.3	55.7	71.8	117.1	310.7	944.5
1950 <sup>1</sup> -----	60.6	114.2	36.8	22.7	54.8	45.7	45.7	53.0	70.8	116.9	315.7	972.6
<b>WHITE MALE</b>												
1969-----	78.7	67.4	33.0	24.7	115.4	81.4	70.0	74.7	91.5	119.1	218.6	534.5
1968-----	77.7	66.1	31.8	25.7	111.7	79.5	67.3	75.2	92.2	125.8	227.1	536.2
1967-----	77.6	71.4	34.0	24.8	108.2	79.3	68.8	77.3	93.9	121.6	228.2	602.5
1966-----	78.1	76.3	34.6	25.0	108.2	79.7	68.7	79.1	95.8	121.1	238.9	604.7
1965-----	75.1	81.5	33.5	24.1	99.9	75.6	66.4	77.2	91.6	122.1	238.4	617.5
1964-----	72.9	78.0	32.0	24.2	95.8	74.4	65.1	74.7	89.1	120.9	235.0	562.9
1963 <sup>2</sup> -----	72.1	80.0	32.2	23.4	92.9	70.7	63.5	75.5	90.1	120.0	238.3	655.8
1962 <sup>2</sup> -----	70.4	81.7	31.3	23.4	90.1	69.8	62.2	73.3	86.1	119.2	239.3	638.3
1961-----	68.0	79.9	31.9	23.8	89.3	66.4	59.1	69.8	82.5	113.3	230.2	591.0
1960-----	70.0	85.6	31.8	24.8	92.2	67.7	60.8	72.3	84.2	117.7	238.7	600.5
1959-----	70.5	84.9	31.0	24.8	94.5	68.6	61.1	71.8	84.2	118.0	239.5	651.0
1958-----	70.5	89.5	31.5	25.4	94.0	67.9	63.4	69.4	82.4	116.9	243.7	633.2
1957-----	75.4	82.9	30.8	25.9	101.0	72.7	65.3	75.0	91.1	131.5	273.8	697.4
1956-----	76.9	85.2	32.1	26.5	107.6	72.9	64.6	75.8	91.2	137.0	275.2	706.3
1955-----	77.3	84.2	32.8	26.8	102.5	72.6	66.5	77.0	93.1	137.5	286.0	693.9
1954-----	75.9	93.1	34.7	27.1	97.5	71.4	63.9	73.7	91.0	135.2	289.2	703.9
1953-----	81.9	99.0	37.3	29.2	104.7	76.6	68.2	80.8	99.6	143.8	310.5	759.7
1952-----	85.7	100.5	38.2	29.6	102.6	77.7	71.5	82.0	104.3	150.6	315.1	790.9
1951-----	85.7	103.0	38.3	31.3	97.4	80.0	76.5	83.4	104.5	157.6	319.2	846.0
1950-----	82.5	111.5	39.5	30.1	91.8	74.0	70.1	79.1	103.8	154.8	326.6	853.3
<b>WHITE FEMALE</b>												
1969-----	34.1	51.1	23.4	11.5	28.3	19.6	21.7	25.9	34.7	59.4	149.7	515.2
1968-----	34.3	57.1	23.4	11.7	27.5	18.2	20.4	25.3	33.9	60.6	159.7	553.5
1967-----	34.6	55.8	22.9	10.9	27.0	18.1	20.2	25.9	34.5	61.4	167.4	614.2
1966-----	35.0	59.5	24.2	11.6	26.5	18.1	20.4	25.9	35.0	62.4	174.4	630.3
1965-----	34.0	58.8	22.2	10.5	24.0	17.3	19.8	26.2	33.8	60.5	179.7	647.2
1964-----	33.4	63.1	22.9	10.5	23.3	17.3	19.2	24.6	33.9	60.7	175.3	636.5
1963 <sup>2</sup> -----	33.0	63.1	22.8	10.1	22.5	15.2	18.5	23.9	32.3	60.5	186.0	669.0
1962 <sup>2</sup> -----	32.4	64.6	22.7	9.9	21.0	14.8	18.0	23.4	31.6	60.9	188.1	660.1
1961-----	30.5	62.9	22.2	9.6	18.9	14.0	15.9	21.2	29.0	57.4	182.9	646.4
1960-----	31.5	66.7	23.0	10.4	19.7	14.0	15.7	21.4	29.6	59.3	201.0	675.9
1959-----	31.0	68.0	22.0	9.8	19.0	14.2	15.0	21.6	28.9	59.0	197.1	705.4
1958-----	31.1	64.2	21.7	11.1	18.6	12.8	15.0	20.1	28.9	61.9	209.8	713.4
1957-----	33.5	64.2	23.0	10.8	19.5	13.9	16.2	21.5	31.9	65.4	236.9	833.9
1956-----	33.5	61.7	23.2	10.8	21.1	14.0	16.1	21.0	31.3	67.3	240.1	853.5
1955-----	35.8	61.1	23.6	11.0	20.2	13.9	16.3	20.9	31.5	66.6	252.5	905.5
1954-----	33.2	66.4	24.8	11.4	17.8	12.7	15.2	20.9	30.6	68.6	258.0	889.3
1953-----	35.1	72.7	26.4	12.5	20.4	13.7	15.6	21.4	31.5	73.3	279.0	963.6
1952-----	36.6	79.6	27.7	12.9	21.0	14.1	17.0	22.1	32.8	74.8	297.5	1,042.9
1951-----	36.8	81.4	26.9	12.9	18.8	14.6	16.4	21.2	33.8	79.0	315.0	1,076.8
1950-----	36.3	83.1	27.0	12.5	17.6	14.0	15.7	21.5	33.5	79.9	321.2	1,131.7

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.



Table 4.—Death Rates for Accidents, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E800-E949 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E800-E962 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	105.7	116.2	51.4	39.3	125.0	136.1	137.6	141.4	164.1	192.5	196.6	214.8
1968-----	108.2	138.6	56.5	39.8	120.6	143.1	138.3	148.6	163.0	218.2	195.0	232.7
1967-----	102.4	157.2	59.6	36.8	109.5	136.2	131.9	137.9	155.3	186.2	199.0	236.0
1966-----	104.6	184.5	58.1	38.6	109.3	129.4	132.0	145.3	157.3	198.8	214.4	321.7
1965-----	98.6	168.9	58.6	35.2	99.4	123.9	129.8	136.8	146.5	174.7	193.7	375.6
1964-----	95.3	166.7	55.3	36.9	94.8	119.7	125.0	133.5	137.1	171.6	170.1	275.7
1963 <sup>2</sup> -----	94.4	163.6	57.5	36.1	98.9	114.2	115.7	125.4	138.7	169.2	196.6	339.4
1962 <sup>2</sup> -----	92.2	172.0	52.9	35.4	96.1	109.4	111.6	123.6	136.2	167.2	182.8	341.9
1961-----	88.2	170.1	53.3	34.7	90.3	101.6	108.5	118.6	129.3	139.1	177.3	348.4
1960-----	95.0	204.3	60.0	38.6	98.8	106.1	110.4	127.8	129.1	142.8	206.8	385.8
1959-----	94.5	209.3	59.8	38.5	97.3	106.6	112.2	124.3	134.8	137.4	177.1	286.2
1958-----	94.7	228.4	58.1	35.9	100.9	110.0	107.1	116.8	132.3	141.9	200.7	355.6
1957-----	101.0	213.6	56.8	39.5	101.7	123.6	111.9	130.5	138.5	152.0	219.3	376.9
1956-----	103.6	224.3	56.5	37.6	112.3	121.8	124.1	129.2	143.8	149.9	205.4	416.0
1955-----	100.9	195.8	61.2	36.7	105.5	116.9	115.0	136.7	139.7	155.0	196.8	350.0
1954-----	99.7	221.8	59.9	37.5	105.0	117.7	103.5	130.4	129.4	152.6	223.5	477.3
1953-----	110.9	228.2	68.3	40.4	116.1	130.0	122.3	143.0	149.6	155.8	268.4	414.3
1952-----	113.4	256.8	69.3	38.6	116.5	127.5	127.5	148.8	158.1	156.4	260.9	465.0
1951-----	110.1	229.1	66.6	45.0	104.0	118.2	125.7	140.2	156.2	168.8	250.9	521.1
1950 <sup>1</sup> -----	103.9	245.5	65.0	41.8	97.4	110.1	119.1	129.8	143.5	165.6	223.0	439.3
ALL OTHER FEMALE												
1969-----	37.3	114.2	41.7	17.6	30.2	28.8	32.2	37.5	45.4	85.1	116.9	218.8
1968-----	37.2	118.7	43.7	17.8	24.1	29.6	35.4	34.7	50.4	85.4	120.0	234.7
1967-----	36.7	132.9	44.3	17.2	24.5	28.4	32.5	36.1	45.7	77.3	123.5	260.6
1966-----	40.0	147.5	49.3	17.5	25.1	31.3	33.6	42.3	46.7	91.1	138.5	301.7
1965-----	37.3	159.8	45.5	18.2	22.7	26.5	33.5	36.4	44.9	79.1	106.2	251.9
1964-----	36.6	136.7	46.2	17.9	20.3	28.0	29.8	31.3	46.9	73.9	144.2	318.3
1963 <sup>2</sup> -----	38.0	155.0	46.5	18.9	20.6	28.3	27.8	35.1	47.3	81.5	141.2	365.9
1962 <sup>2</sup> -----	37.9	160.6	49.8	18.6	19.5	23.3	30.0	32.1	45.9	86.0	139.2	357.1
1961-----	36.0	133.5	50.3	17.8	21.0	25.7	29.2	30.0	45.7	68.8	131.5	273.2
1960-----	38.7	173.5	51.5	19.9	19.6	23.1	29.3	34.8	51.0	75.8	146.2	340.2
1959-----	37.6	188.4	49.4	19.1	19.0	24.8	28.7	34.0	40.5	71.4	128.6	320.5
1958-----	39.2	201.7	56.7	19.0	19.6	22.3	27.2	32.4	44.8	81.2	150.3	334.2
1957-----	41.3	184.0	54.3	19.4	22.3	27.1	29.8	38.2	48.9	80.2	167.6	340.5
1956-----	39.5	182.1	47.6	20.2	21.7	24.3	31.0	34.4	52.6	81.2	158.7	294.4
1955-----	40.7	168.8	58.7	19.2	21.8	27.9	29.0	35.5	52.1	80.2	177.3	361.8
1954-----	40.1	204.0	55.4	17.5	21.1	25.1	29.5	36.1	49.1	80.6	181.0	315.6
1953-----	41.8	196.6	57.2	17.9	24.7	28.4	32.1	38.9	48.4	77.2	179.3	406.5
1952-----	45.5	215.1	61.2	20.2	24.9	28.8	28.5	41.0	55.0	86.0	189.7	365.5
1951-----	42.9	222.5	55.9	19.1	25.6	25.7	32.0	38.5	56.5	78.0	188.3	496.4
1950 <sup>1</sup> -----	39.9	207.6	58.3	20.4	22.1	25.8	29.0	30.9	47.8	94.6	166.0	421.2

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 5.—Death Rates for Motor Vehicle Accidents, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E810-E823 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E810-E835 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969	27.6	10.6	11.8	9.8	51.0	32.4	25.8	25.3	28.3	35.2	42.9	34.5
1968	27.5	9.3	11.1	10.0	49.9	32.6	25.0	25.7	28.8	36.2	46.1	39.8
1967	26.7	9.1	11.2	9.4	48.5	31.6	24.2	25.8	28.9	36.7	45.4	40.1
1966	27.1	9.7	11.3	9.6	48.8	32.6	25.0	26.5	29.8	36.5	45.7	43.5
1965	25.4	8.5	10.5	8.9	44.2	29.8	24.6	25.4	29.0	35.5	45.5	40.3
1964	24.5	9.3	10.3	8.7	42.0	29.5	23.6	24.6	28.2	35.9	45.0	38.7
1963	23.1	7.8	10.0	8.0	40.1	27.0	21.9	23.6	27.0	33.4	44.5	39.8
1962	22.0	8.1	9.4	8.1	38.4	25.1	20.9	21.8	26.0	32.8	41.8	38.6
1961	20.8	7.2	9.7	7.6	36.5	24.4	19.5	21.0	24.4	31.0	40.9	36.3
1960	21.5	8.1	10.0	7.9	38.0	24.3	19.3	21.4	25.1	31.4	41.8	37.9
1959	21.5	7.8	9.5	7.9	38.2	24.9	19.5	22.0	24.8	32.3	42.2	36.7
1958	21.3	7.2	9.5	8.1	37.0	24.4	20.2	21.2	24.6	33.5	43.0	37.2
1957	22.7	8.4	9.4	8.0	39.8	26.7	21.2	22.8	27.4	34.6	46.4	38.1
1956	23.7	8.7	9.5	8.4	43.1	27.3	22.0	23.6	27.4	36.5	46.5	42.5
1955	23.4	8.7	10.5	8.0	41.1	27.0	21.9	22.8	28.0	35.3	47.1	41.1
1954	22.1	8.1	11.0	8.0	36.4	25.1	20.0	22.1	26.2	34.1	48.6	43.3
1953	24.0	9.4	12.0	8.5	39.3	26.9	21.8	23.7	28.4	38.0	51.6	49.9
1952	24.3	9.0	11.9	8.6	38.8	26.9	22.4	24.2	29.7	38.7	50.3	44.2
1951	24.1	8.7	11.4	9.2	36.1	26.6	22.6	24.2	29.4	39.5	52.2	47.6
1950 <sup>1</sup>	23.1	8.4	11.5	8.8	34.4	24.6	20.3	22.2	29.0	39.1	52.7	45.1
<b>WHITE MALE</b>												
1969	40.0	10.7	12.4	11.8	82.0	49.5	35.5	34.5	37.9	48.3	67.0	65.7
1968	40.0	10.1	11.2	12.5	80.7	49.5	35.1	35.8	39.6	49.2	73.6	74.2
1967	39.0	9.0	12.1	11.5	78.4	48.5	34.7	36.1	39.6	50.1	70.3	74.5
1966	39.5	9.8	12.1	12.2	78.7	49.9	35.7	36.8	40.9	49.7	72.8	81.3
1965	37.1	9.9	11.1	11.5	71.6	45.8	35.2	35.4	39.5	49.6	72.9	72.8
1964	35.7	9.4	11.1	11.1	67.9	45.8	33.4	34.1	38.0	50.5	70.6	62.6
1963 <sup>2</sup>	34.3	8.0	11.2	10.4	65.4	42.7	32.4	34.0	37.9	47.3	72.0	72.5
1962 <sup>2</sup>	32.5	8.6	10.2	10.6	63.5	39.9	30.3	30.9	36.0	46.4	66.9	73.2
1961	30.8	7.2	10.9	9.8	60.6	36.3	28.6	29.8	33.4	44.9	64.4	67.8
1960	31.5	8.8	11.3	10.3	62.7	38.6	28.4	29.7	34.4	45.5	66.8	61.9
1959	31.9	7.6	10.5	10.3	64.2	39.0	28.7	30.7	34.6	46.9	67.0	64.7
1958	31.8	8.3	10.7	10.7	62.3	38.7	29.9	29.7	34.2	47.8	68.1	71.2
1957	33.9	9.0	10.3	10.4	68.0	42.2	31.1	31.9	38.0	50.8	73.9	70.1
1956	35.3	8.9	9.9	11.2	73.1	42.8	31.8	33.5	38.6	53.4	75.7	75.2
1955	35.0	9.2	11.3	10.6	69.6	42.6	32.8	32.0	39.3	52.3	74.6	74.8
1954	33.1	8.6	12.4	10.4	63.1	40.3	29.6	30.6	36.7	50.4	79.1	69.1
1953	35.8	10.7	13.6	10.9	66.6	42.8	32.2	33.1	40.7	56.3	83.1	83.2
1952	36.5	9.5	13.6	11.2	65.6	43.2	33.4	33.6	42.4	60.3	81.2	70.4
1951	36.5	9.7	13.8	12.0	61.0	42.4	34.0	34.7	41.8	59.4	84.3	85.7
1950	35.1	9.1	13.2	12.0	58.3	39.1	30.9	31.6	41.9	59.1	86.4	79.3
<b>WHITE FEMALE</b>												
1969	15.1	11.3	10.1	6.9	23.1	13.0	13.2	13.8	17.1	23.7	27.6	20.4
1968	14.8	9.0	9.6	6.8	22.7	12.9	12.3	13.6	17.0	24.2	28.3	22.2
1967	14.7	9.1	9.2	6.5	22.1	12.8	12.1	14.0	17.5	24.8	29.1	22.8
1966	14.8	9.3	9.7	6.6	22.0	13.2	12.2	14.3	18.7	24.8	27.4	22.2
1965	13.9	7.8	8.8	6.0	19.6	11.8	12.3	14.1	17.9	23.3	28.0	22.4
1964	13.8	10.0	9.0	5.8	19.3	11.9	12.1	13.7	18.0	23.1	27.8	26.1
1963 <sup>2</sup>	12.7	8.9	8.7	5.4	18.1	10.6	10.9	13.0	16.4	21.9	26.3	20.8
1962 <sup>2</sup>	12.1	7.9	8.6	5.3	16.7	10.1	10.7	12.1	16.2	21.3	25.5	20.3
1961	11.0	7.9	8.1	4.9	14.7	9.3	9.2	11.4	14.8	18.9	23.8	17.5
1960	11.2	7.5	8.3	5.3	15.6	9.0	8.9	11.4	15.3	19.3	23.8	22.2
1959	11.1	8.8	8.2	5.0	14.8	9.6	8.8	11.4	14.6	20.0	25.1	19.4
1958	10.9	6.8	7.9	5.6	14.3	8.6	8.9	11.2	14.5	20.9	25.2	17.6
1957	11.3	8.1	7.9	5.3	15.1	9.2	9.6	11.8	15.5	20.4	24.9	16.8
1956	11.6	7.4	8.5	5.4	16.4	9.6	9.6	11.5	14.8	21.6	24.5	21.9
1955	11.5	8.2	9.2	5.2	15.8	9.8	9.1	11.5	15.8	20.0	26.7	17.6
1954	10.8	6.7	9.1	5.5	13.3	8.4	8.7	11.2	14.5	19.9	25.2	23.9
1953	11.7	7.6	9.9	5.7	15.6	9.2	9.1	12.0	15.3	21.7	26.4	26.7
1952	11.7	8.8	9.8	6.0	15.3	9.0	9.8	12.2	15.9	19.7	25.8	23.9
1951	11.5	7.9	8.9	6.2	13.5	9.5	9.1	11.7	15.8	21.3	26.5	21.1
1950	10.9	7.8	10.1	5.6	12.6	9.0	8.1	10.8	15.0	20.9	25.4	22.3

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 5.—Death Rates for Motor Vehicle Accidents, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E810-E823 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E810-E835 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	47.0	9.6	16.5	15.8	67.2	72.2	65.2	62.9	69.4	68.3	65.2	19.7
1968-----	46.3	6.2	17.0	15.1	66.1	76.2	59.6	62.8	65.1	75.5	65.2	34.5
1967-----	42.6	7.8	17.0	14.7	61.6	68.7	53.9	54.4	59.3	74.6	57.3	36.0
1966-----	43.0	12.3	15.3	13.9	61.3	69.5	57.6	58.3	57.5	70.3	65.1	52.2
1965-----	39.7	5.9	16.4	12.6	55.2	65.7	54.0	52.5	58.2	62.8	50.3	51.2
1964-----	37.9	7.4	14.7	13.4	48.2	59.5	53.8	53.3	52.6	67.5	51.6	45.9
1963 <sup>2</sup> -----	36.6	5.6	14.6	12.2	52.9	57.9	49.6	48.1	51.3	58.5	50.0	63.6
1962 <sup>2</sup> -----	34.6	8.6	12.4	11.5	49.6	53.7	46.6	45.7	51.0	60.2	45.0	25.8
1961-----	32.8	6.0	12.3	11.2	46.5	50.9	42.8	42.3	48.1	53.2	52.1	64.5
1960-----	34.4	7.5	13.1	10.7	50.7	51.7	43.1	49.1	47.6	49.1	56.0	70.2
1959-----	35.4	6.0	12.0	12.5	49.5	53.2	47.2	52.9	49.5	44.4	42.4	65.5
1958-----	35.4	5.9	11.9	11.1	50.1	55.6	47.4	47.3	48.8	54.0	43.2	33.3
1957-----	38.4	8.7	13.3	11.2	50.3	62.1	48.4	51.9	57.0	51.2	73.3	69.2
1956-----	42.6	13.6	13.9	11.2	59.7	67.3	57.3	55.3	58.4	56.9	61.5	64.0
1955-----	40.1	10.0	13.7	10.3	57.4	63.1	51.3	53.1	55.8	54.4	52.8	70.8
1954-----	38.3	13.3	13.2	10.4	50.3	59.1	45.9	55.3	54.8	50.0	60.5	95.5
1953-----	43.4	10.7	17.4	12.7	55.6	66.7	54.5	57.8	56.1	59.9	79.8	81.0
1952-----	42.7	8.6	15.8	11.4	55.5	63.6	53.1	58.9	61.4	54.0	69.1	105.0
1951-----	41.7	6.4	13.5	13.1	50.5	62.1	53.7	52.9	57.6	60.9	73.6	78.9
1950 <sup>1</sup> -----	38.2	8.9	11.7	10.8	44.3	55.4	46.9	52.0	57.0	58.8	66.4	64.3
ALL OTHER FEMALE												
1969-----	14.5	7.8	12.3	8.4	19.2	15.9	15.7	16.6	17.6	21.7	15.8	5.0
1968-----	13.9	10.0	11.0	7.8	15.2	17.1	17.8	15.6	17.1	22.5	16.5	16.7
1967-----	13.8	10.6	10.4	8.7	16.1	15.7	14.1	17.7	17.9	19.5	20.7	10.6
1966-----	14.0	8.3	11.0	7.3	17.8	17.1	15.5	18.6	15.4	20.7	16.3	18.3
1965-----	12.8	6.6	9.8	7.1	15.5	14.8	15.2	16.1	15.7	21.5	11.9	11.1
1964-----	11.6	6.7	8.0	6.5	12.4	14.1	13.9	13.9	17.0	19.2	16.7	10.2
1963 <sup>2</sup> -----	11.0	5.3	8.6	6.0	11.6	14.0	11.3	15.3	16.1	16.7	16.1	11.4
1962 <sup>2</sup> -----	10.8	7.5	9.1	6.5	12.0	10.1	13.6	13.9	14.2	17.0	14.3	4.8
1961-----	10.3	4.8	8.1	5.9	12.4	11.9	11.6	11.4	14.6	14.4	20.1	2.4
1960-----	10.1	7.8	9.4	5.9	11.2	10.3	11.0	12.8	14.4	14.4	11.3	19.9
1959-----	10.2	5.3	8.6	5.7	11.5	12.2	11.3	13.5	12.3	16.2	12.4	5.1
1958-----	9.7	4.8	9.6	4.1	10.7	11.1	11.6	12.8	11.7	15.1	15.7	5.3
1957-----	11.8	6.6	9.1	5.9	13.2	14.4	12.5	15.1	17.7	16.1	18.6	10.8
1956-----	11.9	9.5	9.2	6.6	12.5	12.9	13.9	14.8	20.6	13.8	14.5	2.8
1955-----	12.1	8.4	10.9	5.5	13.5	13.6	13.3	14.4	17.4	18.3	16.9	14.7
1954-----	11.4	8.8	11.2	5.4	11.9	12.0	13.2	13.9	16.8	13.9	19.8	18.8
1953-----	11.9	11.1	9.8	6.2	13.3	13.6	14.5	14.3	13.3	12.5	19.8	19.4
1952-----	11.8	7.6	10.0	4.8	14.5	14.3	10.9	16.8	15.3	14.6	20.7	20.7
1951-----	12.0	9.5	9.1	5.6	13.8	12.0	13.5	15.6	17.7	17.3	18.0	17.9
1950 <sup>1</sup> -----	10.8	6.9	9.4	6.4	12.1	11.4	11.5	11.1	13.9	15.4	22.5	19.1

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 6.—Death Rates for All Other Accidents, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E800-E807,E825-E949 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E800-E802,E840-E962 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	30.0	58.8	19.6	10.0	21.1	20.8	23.6	27.9	36.9	53.8	132.5	454.5
1968-----	30.0	64.2	20.4	10.4	19.5	19.9	22.8	27.6	36.6	56.9	137.8	474.7
1967-----	30.4	69.7	21.3	9.9	18.6	20.3	23.6	28.4	37.1	54.5	144.4	533.6
1966-----	30.9	74.8	22.1	10.2	18.3	19.4	22.9	29.2	37.5	56.1	155.5	548.4
1965-----	30.4	77.5	21.3	9.8	17.5	19.4	21.9	29.0	35.5	55.3	154.5	565.9
1964-----	29.7	74.8	20.9	10.1	17.2	19.1	21.6	27.6	35.0	54.3	152.3	543.4
1963-----	30.3	78.0	21.0	10.2	17.2	18.2	21.0	27.7	35.4	56.6	160.5	595.1
1962 <sup>2</sup> -----	30.3	79.7	21.0	9.9	16.6	18.3	21.2	28.3	34.4	57.3	164.1	591.5
1961-----	29.6	77.0	21.1	10.6	17.5	18.0	20.6	26.8	33.3	53.6	158.8	564.2
1960-----	31.0	85.1	21.6	11.3	18.1	18.6	21.7	28.4	33.9	56.7	172.7	586.8
1959-----	30.7	86.8	21.0	11.0	18.3	18.6	21.2	27.4	33.5	55.4	169.0	611.4
1958-----	30.9	89.6	21.5	11.4	19.2	18.2	21.3	26.0	33.1	55.9	178.3	617.9
1957-----	33.2	85.3	21.5	11.8	19.8	19.3	22.1	28.5	36.0	63.3	202.7	708.4
1956-----	33.0	82.6	21.5	11.7	20.6	18.6	21.7	27.6	36.1	64.8	204.2	716.2
1955-----	33.5	79.4	22.0	12.2	19.4	18.5	22.1	29.4	36.4	66.1	215.0	741.9
1954-----	33.8	90.0	22.6	12.4	20.4	19.2	21.8	28.4	36.3	67.3	218.4	736.6
1953-----	36.1	93.6	24.0	13.5	22.5	20.8	23.1	30.9	39.3	69.5	237.0	792.9
1952-----	37.5	100.3	25.3	13.8	22.3	21.4	24.8	31.7	41.4	75.0	249.9	853.5
1951-----	38.4	103.8	24.9	14.3	21.8	22.3	26.7	31.5	42.4	77.6	258.5	896.9
1950 <sup>1</sup> -----	37.5	105.8	25.3	13.8	20.4	21.1	25.4	30.9	41.8	77.8	263.0	927.5
<b>WHITE MALE</b>												
1969-----	38.7	56.6	20.6	12.9	33.4	32.0	34.5	40.1	53.6	70.8	151.7	468.8
1968-----	37.7	56.0	20.5	13.2	31.0	30.0	32.1	39.4	52.6	74.6	153.5	462.1
1967-----	38.6	62.4	21.9	13.3	29.8	30.8	34.1	41.2	54.3	71.6	157.9	528.0
1966-----	38.6	66.5	22.5	12.7	29.6	29.8	33.0	42.3	55.0	71.4	166.1	523.4
1965-----	38.0	71.6	22.4	12.6	28.3	29.8	31.2	41.8	52.1	72.5	165.5	544.7
1964-----	37.2	69.6	20.9	13.0	27.8	28.7	31.7	40.6	51.1	70.4	164.4	500.3
1963 <sup>2</sup> -----	37.8	72.0	21.0	13.1	27.5	28.0	31.1	41.5	52.2	72.6	166.3	583.3
1962 <sup>2</sup> -----	37.9	73.1	21.1	12.7	26.8	28.9	31.9	42.4	50.1	72.9	172.4	565.2
1961-----	37.2	72.7	21.0	14.0	28.6	28.2	30.5	40.0	49.1	68.3	165.8	523.2
1960-----	38.6	76.8	20.6	14.5	29.5	29.1	32.4	42.6	49.7	72.3	171.9	538.5
1959-----	38.7	77.2	20.5	14.5	30.3	29.6	32.4	41.1	49.6	71.1	172.5	566.3
1958-----	38.7	81.2	20.8	14.7	31.8	29.1	33.5	39.7	48.2	69.1	175.6	562.0
1957-----	41.5	73.9	20.4	15.4	33.0	30.5	34.2	43.1	53.1	80.7	199.9	627.3
1956-----	41.6	74.3	22.2	15.4	34.5	30.2	32.9	42.3	52.7	85.6	199.5	631.1
1955-----	42.3	75.0	21.6	16.2	32.9	30.0	33.7	45.0	53.8	85.3	211.5	619.0
1954-----	42.8	84.5	22.4	16.7	34.4	31.0	34.3	43.1	54.3	84.9	209.1	634.8
1953-----	46.1	89.3	23.7	18.3	39.1	33.8	35.9	47.6	59.9	87.5	227.4	676.5
1952-----	47.2	91.0	24.6	18.4	37.0	34.5	39.1	48.4	62.0	90.3	233.9	720.6
1951-----	49.2	98.3	24.5	19.4	36.4	37.6	42.5	48.7	62.7	98.2	234.9	780.3
1950-----	47.4	102.3	26.3	18.1	33.4	34.9	39.2	47.4	61.9	95.7	240.3	776.0
<b>WHITE FEMALE</b>												
1969-----	19.0	39.8	13.4	4.7	5.3	5.6	8.5	12.1	17.6	34.8	122.2	494.7
1968-----	19.4	48.1	13.7	5.0	4.8	5.2	8.2	11.7	17.0	36.4	130.4	531.3
1967-----	19.9	46.7	13.7	4.5	4.9	5.3	8.0	11.9	16.9	36.6	138.3	591.4
1966-----	20.2	50.1	14.5	5.0	4.5	4.8	8.2	11.4	16.3	37.6	147.0	608.0
1965-----	20.1	51.0	13.4	4.5	4.4	5.5	7.5	12.1	15.9	37.2	151.7	624.8
1964-----	19.6	53.1	13.9	4.7	4.0	5.3	7.1	10.9	16.0	37.6	147.5	610.4
1963 <sup>2</sup> -----	20.3	54.2	14.1	4.7	4.4	4.6	7.6	10.9	16.0	38.6	159.8	648.2
1962 <sup>2</sup> -----	20.3	56.7	14.2	4.5	4.3	4.7	7.3	11.2	15.3	39.6	162.6	639.9
1961-----	19.5	55.1	14.0	4.7	4.1	4.7	6.6	9.8	14.2	38.5	159.1	628.8
1960-----	20.4	59.3	14.7	5.0	4.1	5.0	6.8	10.0	14.4	39.9	177.2	653.7
1959-----	19.9	59.2	13.8	4.8	4.1	4.6	6.2	10.3	14.2	39.0	172.0	685.9
1958-----	20.1	57.4	13.8	5.5	4.3	4.2	6.1	9.0	14.4	41.0	184.6	695.8
1957-----	22.2	56.1	15.1	5.4	4.4	4.6	6.6	9.7	16.3	44.9	212.0	817.1
1956-----	21.9	54.3	14.7	5.4	4.6	4.4	6.5	9.4	16.5	45.6	215.6	831.6
1955-----	22.3	52.9	14.5	5.8	4.5	4.1	7.2	9.4	15.8	46.6	225.9	888.0
1954-----	22.4	59.7	15.6	5.9	4.6	4.2	6.5	9.7	16.1	48.7	232.8	865.4
1953-----	23.5	65.1	16.5	6.8	4.8	4.5	6.5	9.4	16.2	51.5	252.6	936.9
1952-----	24.9	70.8	17.9	6.9	5.6	5.1	7.2	9.9	16.9	55.1	271.8	1,019.0
1951-----	25.3	73.5	18.0	6.7	5.3	5.1	7.2	9.5	18.0	57.7	288.5	1,055.7
1950-----	25.4	75.4	17.0	6.9	5.0	5.0	7.6	10.6	18.5	59.0	295.9	1,109.4

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 6.—Death Rates for All Other Accidents, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E800-E807, E825-E949 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E800-E802, E840-E962 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	58.7	106.7	34.9	23.5	57.9	64.0	72.4	78.5	94.6	124.2	131.4	195.1
1968-----	61.9	132.4	39.5	24.7	54.5	66.9	78.7	85.8	97.9	142.7	129.9	198.2
1967-----	59.8	149.3	42.6	22.1	47.9	67.5	78.1	83.6	95.9	111.6	141.7	200.0
1966-----	61.7	172.2	42.8	24.7	48.0	59.9	74.4	87.0	99.9	128.5	149.2	269.6
1965-----	59.0	163.0	42.1	22.7	44.2	58.2	75.9	84.4	88.3	111.9	143.4	324.4
1964-----	57.4	159.2	40.6	23.5	45.6	60.2	71.1	80.2	84.4	104.2	118.5	229.7
1963 <sup>2</sup> -----	57.8	158.0	42.9	23.9	46.0	56.3	66.1	77.3	87.5	110.7	146.6	275.8
1962 <sup>2</sup> -----	57.6	163.4	40.5	23.9	46.5	55.6	65.0	77.9	85.2	106.9	137.9	316.1
1961-----	55.4	164.2	41.0	23.5	43.7	50.8	65.8	76.3	81.2	85.9	125.2	263.9
1960-----	60.6	196.8	46.9	27.8	48.1	54.4	67.3	78.8	81.5	93.8	150.8	315.7
1959-----	59.1	203.3	47.8	26.0	47.8	53.4	65.0	71.4	85.2	92.9	134.7	220.7
1958-----	59.3	222.5	46.2	24.8	50.8	54.4	59.6	69.5	83.5	87.9	157.6	322.2
1957-----	62.6	204.9	43.4	28.3	51.4	61.6	63.5	78.6	81.5	100.8	145.9	307.7
1956-----	61.1	210.7	42.7	26.4	52.5	54.5	66.8	73.9	85.4	93.0	143.8	352.0
1955-----	60.8	185.8	47.5	26.5	48.0	53.8	63.7	83.6	83.9	100.6	144.0	279.2
1954-----	61.4	208.5	46.7	27.1	54.7	58.5	57.6	75.1	74.6	102.6	163.0	381.8
1953-----	67.5	217.5	50.8	27.7	60.5	63.4	67.8	85.2	93.5	95.9	188.6	333.3
1952-----	70.7	248.2	53.5	27.2	61.0	64.0	74.5	89.9	96.7	102.4	191.8	360.0
1951-----	68.4	222.7	53.1	31.8	53.5	56.1	72.0	87.3	98.6	108.0	177.4	442.1
1950 <sup>1</sup> -----	65.7	236.6	53.3	31.0	53.1	54.7	72.3	77.8	86.4	106.8	156.6	375.0
ALL OTHER FEMALE												
1969-----	22.8	106.5	29.4	9.2	11.0	12.9	16.5	21.0	27.8	63.3	101.1	213.8
1968-----	23.2	108.7	32.6	10.0	8.9	12.5	17.6	19.1	33.3	63.0	101.5	218.1
1967-----	22.9	122.3	34.0	8.5	8.4	12.7	18.4	18.4	27.8	57.8	102.8	250.0
1966-----	26.1	139.3	38.3	10.2	7.3	14.2	18.1	23.8	31.4	70.5	122.2	283.3
1965-----	24.5	153.2	35.7	11.0	7.3	11.7	18.3	20.2	29.2	57.6	94.3	240.7
1964-----	25.0	130.0	38.2	11.4	7.9	13.9	15.9	17.3	29.9	54.7	127.4	306.1
1963 <sup>2</sup> -----	27.1	149.7	37.9	12.9	9.0	14.3	16.4	19.7	31.2	64.6	125.1	354.5
1962 <sup>2</sup> -----	27.1	153.1	40.7	12.1	7.5	13.2	16.4	18.2	31.6	69.0	124.9	352.4
1961-----	25.8	128.7	42.1	11.9	8.5	13.8	17.6	18.7	31.1	54.3	111.4	270.7
1960-----	28.6	165.7	42.2	14.0	8.4	12.8	18.3	22.0	36.7	61.4	134.9	320.3
1959-----	27.4	183.1	40.8	13.3	7.5	12.6	17.4	20.5	28.2	55.1	116.1	315.4
1958-----	29.5	196.9	47.0	14.9	8.9	11.2	15.6	19.5	33.1	66.1	134.6	328.9
1957-----	29.5	177.4	45.2	13.5	9.1	12.7	17.3	23.1	31.2	64.1	149.0	329.7
1956-----	27.6	172.6	38.4	13.6	9.2	11.4	17.1	19.6	32.0	67.4	144.2	291.7
1955-----	28.6	160.5	47.7	13.7	8.3	14.2	15.7	21.1	34.7	61.8	158.3	347.1
1954-----	28.7	195.2	44.2	12.2	9.2	13.1	16.3	22.2	32.3	66.7	161.1	296.9
1953-----	29.9	185.5	47.4	11.7	11.4	14.8	17.6	24.6	35.1	64.8	159.5	387.1
1952-----	31.6	207.6	51.2	15.5	10.5	14.5	17.7	24.3	39.7	71.4	169.0	344.8
1951-----	30.9	213.1	46.8	13.5	11.9	13.7	18.5	22.9	38.8	60.7	170.3	478.6
1950 <sup>1</sup> -----	29.1	200.8	48.9	14.0	10.0	14.4	17.5	19.8	33.9	79.2	143.5	402.0

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 7.—Death Rates for Influenza and Pneumonia, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 470-474,480-486 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 480-483,490-493 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969	33.9	204.3	8.2	1.8	2.5	4.1	9.0	18.0	39.4	101.9	295.4	1,040.6
1968	36.8	230.8	9.6	1.8	2.8	4.4	9.3	19.8	42.4	109.5	322.1	1,169.7
1967	28.8	161.5	9.2	1.7	2.3	3.2	7.1	14.3	29.6	85.2	270.0	971.1
1966	32.5	189.9	10.9	2.0	2.4	3.5	8.3	16.3	34.9	97.6	299.7	1,091.5
1965	31.9	208.3	11.4	2.1	2.2	3.6	7.8	15.1	33.3	94.6	295.1	1,062.6
1964	31.1	205.5	12.3	1.9	2.2	3.6	7.1	15.0	32.3	90.7	286.2	1,029.8
1963 <sup>2</sup>	37.5	217.8	14.4	2.3	2.6	4.2	8.5	18.4	41.4	115.8	350.0	1,263.4
1962 <sup>2</sup>	32.3	215.9	15.7	2.3	2.5	3.5	7.6	15.1	33.0	94.2	297.1	1,054.7
1961	30.1	216.4	15.1	2.1	2.2	3.2	7.2	13.9	30.9	86.3	274.5	952.8
1960	37.3	239.0	16.2	2.6	3.0	4.8	9.3	19.6	43.2	112.1	340.6	1,126.6
1959	31.2	239.2	15.3	2.5	2.7	3.8	7.7	15.5	33.5	86.5	281.6	924.3
1958	33.1	245.3	16.0	2.3	2.9	4.3	7.8	17.3	38.9	96.0	292.8	988.1
1957	35.8	239.9	18.3	3.8	5.5	5.6	10.2	20.6	44.3	106.9	298.6	979.8
1956	28.2	225.3	15.3	2.3	2.4	3.0	7.0	14.0	30.4	80.6	258.0	853.5
1955	27.1	221.6	14.9	2.5	2.7	3.1	6.8	14.2	29.5	76.4	249.8	791.1
1954	25.4	227.2	16.1	2.4	2.4	3.3	6.6	14.2	27.6	70.8	220.6	713.2
1953	33.0	257.5	18.8	3.1	3.2	4.2	8.8	18.6	38.2	96.6	310.7	1,000.7
1952	29.7	266.9	19.9	3.2	3.0	4.2	8.6	17.3	33.7	82.0	266.6	825.2
1951	31.4	252.0	19.0	3.2	3.4	4.0	8.7	18.9	38.3	93.8	299.5	935.1
1950 <sup>1</sup>	31.3	268.1	18.9	3.2	3.2	4.2	9.0	19.3	38.2	97.1	296.5	954.4
<b>WHITE MALE</b>												
1969	36.4	179.5	6.7	1.7	2.6	3.0	7.3	19.1	48.9	135.8	393.7	1,217.5
1968	39.1	194.6	7.5	1.6	2.9	3.4	7.9	21.0	52.3	146.7	416.8	1,364.5
1967	31.1	131.6	7.3	1.6	2.4	2.4	6.3	15.7	37.5	114.2	352.7	1,166.7
1966	34.6	152.0	8.6	1.8	2.6	2.5	7.2	17.3	43.4	128.5	384.8	1,289.9
1965	34.4	171.9	8.4	2.0	2.2	2.8	6.6	16.8	41.0	126.5	386.4	1,280.3
1964	33.2	170.9	9.3	1.6	2.2	2.5	5.8	17.0	39.7	122.3	364.9	1,225.8
1963 <sup>2</sup>	39.3	175.9	10.7	2.0	2.4	3.1	7.3	19.9	48.6	149.0	432.1	1,481.9
1962 <sup>2</sup>	34.1	179.6	10.6	2.0	2.6	2.6	6.9	16.2	39.9	121.7	373.6	1,227.4
1961	31.9	180.8	10.8	2.0	2.3	2.4	6.4	15.3	37.7	113.1	343.9	1,090.7
1960	39.2	191.5	12.4	2.4	3.0	3.9	8.5	21.6	52.2	144.0	416.7	1,303.1
1959	32.7	194.7	12.0	2.2	2.6	3.1	6.7	17.2	40.9	112.5	347.6	1,067.8
1958	34.9	198.2	12.5	2.1	2.9	3.4	7.4	19.2	47.5	123.7	358.6	1,125.3
1957	37.8	200.1	14.4	3.6	4.7	3.9	9.3	23.0	54.3	138.0	359.1	1,117.4
1956	29.4	186.3	11.8	1.9	2.5	2.5	6.2	15.6	37.6	101.1	313.6	949.7
1955	27.7	179.6	11.9	2.3	2.5	2.5	6.4	15.3	35.6	92.4	295.8	846.6
1954	26.2	185.9	12.9	2.2	2.3	2.8	6.4	15.6	33.4	88.7	258.7	782.3
1953	33.8	213.8	15.7	2.6	2.9	3.1	8.4	19.9	44.5	113.1	358.8	1,085.8
1952	30.0	213.3	15.8	2.9	2.7	3.5	8.1	19.8	40.5	95.5	303.4	871.5
1951	31.8	206.7	15.5	2.9	2.8	3.2	8.4	19.6	44.2	109.3	340.1	992.0
1950	31.6	232.1	15.3	2.8	3.0	3.0	8.6	20.3	43.6	111.8	329.2	1,034.1
<b>WHITE FEMALE</b>												
1969	28.9	128.5	6.2	1.7	1.9	2.8	5.8	11.1	22.9	62.6	231.5	1,040.5
1968	31.1	144.9	6.9	1.5	2.2	2.9	5.3	12.1	24.0	66.9	258.8	1,146.8
1967	24.0	97.6	6.2	1.4	1.6	2.1	3.9	7.7	16.3	51.2	215.3	938.4
1966	26.8	108.1	7.4	1.7	1.7	2.2	4.5	10.0	18.8	59.9	240.0	1,045.1
1965	25.9	121.4	8.4	1.6	1.6	2.4	4.6	9.0	18.1	57.0	230.8	1,006.9
1964	25.2	126.8	8.6	1.7	1.6	2.2	3.9	7.8	13.0	55.1	230.3	969.2
1963 <sup>2</sup>	30.3	132.6	10.2	2.0	2.0	3.0	5.1	9.3	21.8	70.0	285.4	1,175.4
1962 <sup>2</sup>	26.3	132.2	9.4	2.1	1.9	2.3	4.5	8.3	18.5	60.1	242.2	1,005.2
1961	24.3	142.3	9.1	1.8	1.5	2.1	4.2	7.4	16.8	54.0	225.7	900.4
1960	29.0	141.1	12.3	2.2	2.3	3.2	5.0	10.0	22.5	69.3	277.1	1,054.7
1959	24.5	147.3	10.6	2.2	2.0	2.5	4.3	8.4	17.6	54.9	232.8	878.4
1958	25.5	156.3	12.1	1.9	2.2	2.8	4.3	9.1	19.7	59.7	237.7	930.8
1957	27.3	146.8	13.6	3.3	5.2	4.4	6.0	10.2	22.5	66.8	249.2	916.4
1956	21.9	145.5	10.9	2.1	1.7	1.7	4.0	7.0	15.5	53.1	215.2	821.4
1955	21.3	145.7	11.1	2.1	2.0	1.8	3.9	6.8	15.5	52.2	211.7	785.1
1954	19.4	150.3	11.8	2.0	1.7	1.9	3.6	6.8	14.2	46.9	189.4	683.2
1953	25.1	173.1	14.7	2.7	2.1	2.6	4.5	7.9	18.2	65.0	262.4	951.5
1952	22.8	180.3	15.5	2.8	2.0	2.3	4.3	7.4	16.8	57.3	235.1	815.9
1951	24.6	171.1	14.9	2.6	2.3	2.5	4.9	8.4	20.6	67.7	263.3	920.2
1950	24.5	189.5	13.9	2.6	2.4	2.7	4.8	8.9	20.2	68.4	264.0	918.6

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 7.—Death Rates for Influenza and Pneumonia, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 470-474,480-486 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 480-483,490-493 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	53.2	464.6	20.0	2.4	4.9	14.6	39.5	65.2	111.0	252.2	392.2	575.4
1968-----	59.6	546.7	22.4	2.9	5.8	17.2	42.2	70.8	125.4	266.4	425.4	732.7
1967-----	47.2	414.4	25.6	2.7	5.1	12.4	31.8	58.1	84.1	222.5	337.2	560.0
1966-----	55.8	521.4	27.1	2.9	4.2	15.5	36.5	59.2	105.2	257.9	392.1	756.7
1965-----	54.7	550.6	27.4	4.0	4.1	14.2	33.3	53.0	98.9	235.3	368.8	634.1
1964-----	55.0	530.7	32.2	3.9	4.7	15.6	33.9	53.8	94.9	210.5	353.3	737.8
1963 <sup>2</sup> -----	70.5	605.2	38.8	4.1	5.0	13.1	36.1	71.4	139.7	310.2	481.6	1,121.2
1962 <sup>2</sup> -----	58.0	574.8	36.9	4.1	4.2	13.0	27.3	57.2	96.4	219.5	353.3	819.4
1961-----	55.6	544.5	32.1	3.6	4.7	13.4	30.4	53.3	94.0	200.7	312.3	848.4
1960-----	74.3	711.7	43.3	4.6	6.0	14.7	38.6	76.0	140.3	276.2	489.0	921.5
1959-----	61.7	685.7	43.6	4.2	5.6	14.4	32.3	55.6	108.1	186.4	321.5	675.9
1958-----	68.5	719.0	39.7	4.0	6.2	17.1	33.2	65.9	130.7	226.1	392.1	935.3
1957-----	71.9	667.9	48.3	6.7	8.8	18.7	39.7	75.4	143.0	227.2	377.8	857.7
1956-----	56.3	628.3	42.5	5.0	4.7	11.2	28.1	50.5	99.2	173.9	288.5	692.0
1955-----	57.6	640.2	40.6	4.7	6.6	12.9	27.9	57.4	98.8	176.0	303.2	704.2
1954-----	56.0	651.2	43.1	5.5	6.4	13.2	25.5	57.3	96.6	158.8	261.3	668.2
1953-----	73.7	714.5	49.6	6.9	9.6	18.2	36.5	79.1	153.5	234.0	417.5	1,068.7
1952-----	67.3	794.1	52.1	6.5	8.1	17.3	38.5	66.3	116.4	189.3	346.4	725.0
1951-----	67.2	726.4	48.5	7.3	10.3	15.6	31.4	78.9	135.9	191.7	354.7	842.1
1950 <sup>1</sup> -----	66.1	714.1	55.1	6.8	7.2	14.7	32.3	73.5	135.0	229.6	358.8	964.2
ALL OTHER FEMALE												
1969-----	33.8	401.9	13.4	2.6	3.7	10.5	19.1	27.8	53.5	128.1	194.1	447.5
1968-----	39.7	476.7	20.3	3.5	3.8	10.7	22.5	32.9	61.7	137.2	221.5	622.2
1967-----	29.7	356.8	18.0	3.0	3.2	8.6	16.6	22.6	40.6	101.6	177.3	451.5
1966-----	37.0	449.5	25.3	3.7	4.2	9.1	21.0	25.1	54.1	121.1	220.1	546.7
1965-----	37.6	482.3	27.3	3.8	3.5	9.0	19.8	21.8	53.3	117.3	208.4	538.9
1964-----	37.0	450.3	28.4	3.0	3.9	10.3	18.3	27.1	47.8	106.9	200.9	553.1
1963 <sup>2</sup> -----	51.0	500.6	34.8	3.8	4.8	12.6	21.7	39.1	85.4	182.0	323.1	997.7
1962 <sup>2</sup> -----	41.9	488.8	32.1	3.6	4.4	8.9	21.4	29.2	57.7	123.4	240.7	654.8
1961-----	37.4	474.0	30.6	3.0	4.3	8.0	17.0	23.2	50.1	96.3	183.7	595.1
1960-----	50.4	589.9	34.3	4.6	5.4	14.1	24.6	37.1	76.3	144.4	282.8	777.1
1959-----	43.2	574.2	34.7	4.2	4.8	10.0	20.9	27.6	57.0	106.4	213.7	502.6
1958-----	46.6	581.0	38.3	4.5	5.0	12.0	17.9	31.6	69.8	121.6	259.5	634.2
1957-----	53.9	598.3	42.7	6.2	10.0	15.6	25.3	43.8	85.3	149.2	252.4	691.9
1956-----	41.6	531.8	38.9	4.6	4.3	10.2	18.4	29.9	54.7	104.0	211.6	544.4
1955-----	40.0	526.6	33.3	4.7	5.3	9.4	14.7	31.7	57.7	102.3	220.5	447.1
1954-----	40.0	546.8	38.2	4.4	5.2	9.2	16.6	29.0	53.2	105.2	186.5	503.1
1953-----	53.7	628.9	36.0	5.6	7.5	13.2	20.3	49.5	94.0	167.2	351.2	809.7
1952-----	49.6	665.3	44.3	5.6	7.2	13.8	22.6	40.6	78.8	129.4	242.2	603.4
1951-----	51.1	589.2	43.1	5.6	9.7	13.1	23.5	52.3	89.7	144.2	286.5	692.9
1950 <sup>1</sup> -----	52.3	589.0	45.6	6.1	7.2	15.7	27.0	55.7	92.5	177.3	314.3	712.2

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 8.—Death Rates for Diabetes Mellitus, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 250 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 260 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969	19.1	0.3	0.1	0.2	0.7	2.5	5.4	13.1	37.1	96.7	185.9	262.0
1968	19.2	0.5	0.1	0.2	0.7	2.6	5.5	13.0	39.2	99.5	183.3	258.5
1967	17.7	0.4	0.1	0.2	0.7	2.5	4.8	12.3	36.2	94.2	169.4	234.8
1966	17.7	0.5	0.2	0.3	0.7	2.6	5.0	12.1	35.7	95.7	171.5	234.4
1965	17.1	0.5	0.2	0.3	0.7	2.6	4.7	11.9	36.1	92.6	166.4	220.0
1964	16.9	0.6	0.2	0.3	0.8	2.6	4.8	12.0	37.3	91.4	161.6	204.9
1963 <sup>2</sup>	17.2	0.4	0.2	0.3	0.9	2.6	5.0	12.1	37.5	95.1	165.9	206.6
1962 <sup>2</sup>	16.8	0.6	0.2	0.3	0.8	2.5	4.6	11.7	36.9	94.0	160.2	206.7
1961	16.4	0.3	0.2	0.3	0.7	2.7	4.7	11.3	35.9	91.4	162.9	184.2
1960	16.7	0.4	0.3	0.4	0.9	2.3	4.5	12.1	37.9	93.4	163.7	181.7
1959	15.9	0.3	0.2	0.4	0.9	2.6	3.8	11.4	37.5	89.9	154.5	158.8
1958	15.9	0.3	0.3	0.3	0.9	2.5	4.1	11.2	37.0	91.6	151.7	169.9
1957	16.0	0.3	0.3	0.4	1.2	2.6	4.5	11.2	38.0	92.5	149.8	158.1
1956	15.7	0.5	0.2	0.3	1.0	2.4	3.8	11.0	37.5	92.1	151.7	162.6
1955	15.5	0.6	0.3	0.4	1.0	2.4	3.9	10.9	37.3	91.7	150.6	165.1
1954	15.6	0.5	0.3	0.4	1.0	2.2	3.9	11.3	38.3	92.3	153.9	155.3
1953	16.3	0.5	0.3	0.4	1.2	2.4	4.1	11.7	41.7	96.2	161.7	156.4
1952	16.4	0.8	0.5	0.6	1.3	2.3	4.0	12.3	40.8	99.7	161.5	152.8
1951	16.3	0.4	0.4	0.5	1.2	2.2	3.9	12.4	42.7	99.8	161.2	151.1
1950 <sup>1</sup>	16.2	0.7	0.3	0.6	1.1	2.2	4.2	12.4	42.1	101.2	166.7	150.3
<b>WHITE MALE</b>												
1969	15.7	0.3	0.1	0.2	0.5	2.5	5.0	11.9	32.0	83.1	170.2	242.2
1968	16.0	0.5	0.2	0.2	0.6	2.6	5.4	12.2	34.8	86.2	161.9	249.8
1967	14.9	0.4	0.1	0.2	0.5	2.6	5.1	11.1	32.6	79.8	153.3	231.1
1966	14.7	0.5	0.2	0.3	0.6	2.8	4.9	11.1	30.1	81.5	154.0	229.6
1965	14.2	0.5	0.3	0.2	0.5	2.9	4.8	11.0	30.1	77.6	148.6	216.2
1964	14.0	0.3	0.3	0.2	0.6	3.0	5.0	10.9	31.2	75.0	145.2	195.5
1963 <sup>2</sup>	14.2	0.3	0.1	0.2	0.6	2.8	5.0	10.9	30.9	77.5	146.6	205.6
1962 <sup>2</sup>	14.1	0.6	0.2	0.3	0.6	2.6	5.4	10.5	30.4	77.0	142.9	220.1
1961	13.5	0.1	0.2	0.2	0.7	2.9	5.0	10.4	29.6	72.8	142.8	175.4
1960	13.7	0.4	0.3	0.2	0.7	2.6	4.7	10.9	29.5	75.1	146.9	174.7
1959	13.0	0.2	0.2	0.3	0.8	2.8	4.1	10.2	29.2	70.9	135.8	161.7
1958	13.0	0.3	0.4	0.2	0.8	2.8	4.4	9.8	29.3	72.9	132.1	165.2
1957	12.8	0.1	0.3	0.3	0.9	2.8	4.7	9.6	28.4	71.7	132.0	148.6
1956	12.8	0.4	0.2	0.2	0.7	2.6	4.0	9.4	29.3	72.0	133.9	157.9
1955	12.8	0.6	0.3	0.3	0.9	2.5	4.1	9.4	29.1	71.7	134.5	150.3
1954	12.7	0.5	0.3	0.3	0.9	2.5	4.3	9.6	28.6	70.9	135.4	147.2
1953	13.0	0.4	0.4	0.3	1.1	2.6	4.2	9.3	30.0	73.5	139.4	155.2
1952	13.3	1.0	0.6	0.4	1.0	2.2	4.1	10.2	30.2	76.8	144.7	154.5
1951	12.8	0.5	0.4	0.3	0.9	2.2	3.7	10.1	31.4	75.5	133.6	141.8
1950	12.8	1.0	0.2	0.5	0.8	2.0	3.9	9.4	30.0	77.0	145.7	147.6
<b>WHITE FEMALE</b>												
1969	21.0	0.3	0.2	0.3	0.7	2.2	3.7	9.5	30.4	93.2	195.7	296.4
1968	21.0	0.6	0.2	0.3	0.7	2.0	3.6	9.0	31.9	95.7	197.7	277.8
1967	19.6	0.6	0.2	0.3	0.7	1.9	2.9	9.2	30.3	94.3	181.6	256.0
1966	19.6	0.3	0.2	0.3	0.7	2.0	3.2	8.3	31.6	96.2	185.2	250.3
1965	19.1	0.5	0.1	0.4	0.6	2.1	3.1	8.3	32.4	95.3	181.6	232.4
1964	18.8	0.5	0.3	0.4	0.7	1.9	3.2	8.4	32.6	97.1	175.3	218.1
1963 <sup>2</sup>	19.2	0.2	0.2	0.4	0.9	2.0	3.2	9.0	33.6	100.4	180.6	212.1
1962 <sup>2</sup>	18.6	0.4	0.3	0.4	0.8	2.1	2.5	8.2	34.0	100.8	173.8	204.8
1961	18.8	0.5	0.2	0.4	0.6	2.1	3.1	8.2	33.6	102.1	182.0	197.3
1960	19.1	0.3	0.3	0.5	0.9	1.8	2.9	8.7	37.2	105.4	179.6	191.2
1959	18.5	0.5	0.3	0.4	0.9	2.0	2.5	8.7	38.2	104.1	172.3	163.1
1958	18.5	0.5	0.3	0.3	0.8	1.8	2.8	9.0	38.6	105.7	170.9	179.1
1957	19.0	0.4	0.2	0.5	1.2	2.1	3.1	8.9	41.8	110.5	169.6	173.5
1956	18.6	0.7	0.2	0.3	1.0	1.8	2.8	9.3	39.7	110.3	172.6	176.7
1955	18.5	0.3	0.3	0.4	1.1	2.1	2.5	9.0	41.1	110.1	169.8	183.4
1954	18.8	0.3	0.3	0.5	1.0	1.8	2.6	9.2	43.6	113.2	177.2	168.4
1953	19.9	0.5	0.3	0.6	1.2	1.8	2.7	10.0	49.5	119.1	186.8	163.9
1952	19.9	0.7	0.4	0.6	1.3	2.1	2.8	11.0	48.0	123.2	183.0	159.9
1951	20.3	0.4	0.5	0.7	1.3	2.0	2.8	11.2	51.3	124.9	192.6	166.0
1950	20.0	0.5	0.3	0.6	1.2	1.9	2.8	11.5	52.0	125.2	192.7	159.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.



Table 8.—Death Rates for Diabetes Mellitus, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 250 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 260 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	17.3	0.3	0.2	0.2	1.0	4.8	16.0	27.9	67.0	133.8	152.9	111.5
1968-----	17.3	0.3	-	0.3	1.1	5.1	10.4	29.0	69.9	142.2	144.3	140.0
1967-----	15.0	-	-	0.1	1.1	4.8	10.6	26.0	54.0	130.2	121.6	96.0
1966-----	15.1	0.3	0.3	0.3	1.1	4.5	11.5	24.1	56.5	118.7	127.7	143.5
1965-----	14.9	0.3	-	0.3	1.5	3.8	10.7	25.6	57.4	115.6	120.6	134.1
1964-----	14.6	2.1	0.2	0.3	1.7	3.7	10.7	24.5	59.3	102.4	126.1	105.4
1963 <sup>2</sup> -----	13.9	0.6	0.2	0.3	1.1	4.7	10.1	21.4	53.4	104.2	108.0	163.6
1962 <sup>2</sup> -----	13.7	0.6	0.2	0.3	1.3	3.8	10.0	20.7	54.0	95.6	118.3	135.5
1961-----	12.8	0.3	0.3	0.3	1.0	3.3	10.8	19.2	47.0	97.6	89.6	116.1
1960-----	13.9	1.0	0.3	0.5	2.0	3.5	9.1	23.2	55.8	86.8	116.1	127.5
1959-----	12.1	0.3	0.3	0.2	0.7	4.1	5.7	22.4	48.6	81.6	104.9	62.1
1958-----	11.2	0.3	0.3	0.4	1.5	3.8	7.8	18.4	38.4	76.0	89.2	148.1
1957-----	10.8	0.3	0.2	0.4	2.0	4.5	7.5	17.4	38.6	74.1	72.6	103.8
1956-----	10.1	0.4	-	0.2	1.4	3.4	6.1	15.0	44.5	66.0	76.2	60.0
1955-----	9.7	1.9	0.3	0.5	1.4	3.4	7.3	14.2	37.8	59.1	88.0	58.3
1954-----	9.8	0.8	0.6	0.8	1.6	2.9	6.4	17.0	38.4	59.9	72.3	90.9
1953-----	9.8	0.4	0.2	0.5	1.4	4.1	6.9	16.0	37.6	52.9	91.2	104.8
1952-----	9.9	-	0.3	0.4	2.0	3.4	5.4	15.9	37.1	64.2	88.2	85.0
1951-----	10.2	-	0.3	0.6	2.3	2.6	6.5	17.6	40.4	64.8	72.6	94.7
1950 <sup>1</sup> -----	10.0	1.0	0.1	0.3	1.7	3.0	8.3	18.1	33.3	67.1	88.2	58.9
ALL OTHER FEMALE												
1969-----	29.5	0.3	0.1	0.3	1.3	3.2	13.0	40.6	122.3	240.6	220.2	173.8
1968-----	30.0	0.7	-	0.3	1.1	5.0	13.1	41.5	123.2	240.9	217.0	212.5
1967-----	26.2	-	0.1	0.2	1.2	4.2	11.1	38.9	112.6	203.1	194.4	153.0
1966-----	26.8	1.3	0.2	0.2	1.7	3.6	13.4	45.5	108.7	209.1	190.0	163.3
1965-----	25.0	0.6	0.2	0.5	1.4	3.7	11.3	39.6	110.1	187.9	172.7	175.9
1964-----	25.3	0.6	0.1	0.3	0.9	3.5	11.0	42.9	119.7	175.4	170.7	198.0
1963 <sup>2</sup> -----	25.4	1.3	0.3	0.3	2.1	3.9	14.1	42.8	117.6	169.8	170.4	145.5
1962 <sup>2</sup> -----	23.7	0.9	-	0.3	1.5	2.5	10.8	45.9	104.8	159.6	166.7	166.7
1961-----	23.3	0.6	0.2	0.2	1.7	5.0	10.7	40.2	108.8	148.5	173.9	139.0
1960-----	23.4	0.3	0.3	0.4	1.5	3.3	12.9	42.4	110.1	144.7	161.7	156.4
1959-----	21.2	0.3	0.4	0.5	1.5	3.4	11.3	37.7	100.7	127.0	151.6	151.3
1958-----	21.1	-	0.5	0.8	1.7	3.9	9.8	39.1	96.8	134.1	150.3	113.2
1957-----	20.4	1.0	0.4	0.7	2.0	3.5	13.3	41.7	95.6	114.7	131.0	86.5
1956-----	19.1	-	0.2	0.4	2.2	3.8	9.4	38.5	93.2	113.1	118.1	102.8
1955-----	18.5	1.5	0.1	0.5	1.5	2.9	10.4	39.7	84.9	115.3	112.9	100.0
1954-----	18.1	1.2	0.2	0.4	1.5	3.0	10.3	41.8	86.9	103.4	99.2	109.4
1953-----	19.4	0.4	-	0.5	1.9	3.5	12.0	45.4	90.2	108.9	116.5	112.9
1952-----	16.1	0.4	0.1	1.3	2.2	3.1	11.5	40.4	85.5	103.1	108.6	96.6
1951-----	17.9	-	0.4	0.7	2.2	3.7	13.1	41.5	79.9	99.4	113.5	92.9
1950 <sup>1</sup> -----	18.7	-	0.8	1.1	2.6	4.2	14.5	43.0	80.2	113.5	108.4	122.5

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 9.—Death Rates for Arteriosclerosis, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 440 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 450 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	16.4	-	-	0.0	0.0	0.0	0.2	1.1	6.7	40.1	206.8	1,075.6
1968-----	16.8	0.0	0.0	0.0	0.0	0.0	0.2	1.2	6.9	42.1	215.0	1,123.5
1967-----	19.0	-	-	-	0.0	0.0	0.2	1.3	8.3	48.4	249.6	1,298.5
1966-----	19.9	0.0	-	-	0.0	0.0	0.2	1.6	9.3	53.1	268.0	1,359.0
1965-----	19.7	-	0.0	-	0.0	0.1	0.3	1.6	9.0	52.1	270.5	1,396.3
1964-----	19.4	-	-	-	0.0	0.0	0.2	1.6	9.8	51.9	271.9	1,398.3
1963-----	19.8	0.0	-	-	0.0	0.1	0.2	1.7	9.6	54.3	286.0	1,425.8
1962-----	19.8	0.0	0.0	0.0	0.0	0.0	0.2	1.9	9.3	52.7	289.8	1,449.8
1961-----	19.3	0.0	-	0.0	0.0	0.1	0.2	1.6	10.0	53.3	284.7	1,402.5
1960-----	20.0	-	-	-	-	0.1	0.3	1.8	10.9	56.9	307.4	1,423.3
1959-----	19.6	0.0	-	-	0.0	0.0	0.3	2.1	10.8	54.8	306.6	1,410.5
1958-----	19.9	-	0.0	0.0	0.0	0.1	0.3	1.7	10.4	58.7	313.9	1,465.7
1957-----	19.5	-	-	-	0.0	0.0	0.2	1.9	10.4	58.8	312.0	1,448.9
1956-----	19.1	-	0.0	0.0	0.0	0.0	0.2	1.8	9.8	56.7	319.1	1,436.4
1955-----	19.8	-	0.0	-	0.0	0.1	0.3	1.8	10.1	58.7	340.2	1,491.4
1954-----	18.8	0.0	0.0	-	0.0	0.1	0.3	1.9	10.3	57.6	324.0	1,444.0
1953-----	20.4	-	-	-	0.0	0.1	0.3	1.8	11.1	66.0	361.0	1,594.4
1952-----	20.4	-	-	0.0	0.0	0.1	0.3	2.1	11.8	66.4	370.8	1,598.2
1951-----	20.8	-	0.0	-	-	0.1	0.3	2.1	12.0	70.3	391.0	1,664.0
1950 <sup>1</sup> -----	20.4	-	-	-	0.0	0.1	0.3	2.0	12.3	70.0	392.0	1,727.7
<b>WHITE MALE</b>												
1969-----	15.2	-	-	-	0.0	0.0	0.2	1.2	8.2	46.9	237.0	1,079.3
1968-----	15.7	0.1	-	0.0	-	0.0	0.2	1.3	8.1	48.5	240.4	1,163.8
1967-----	18.0	-	-	-	-	0.0	0.3	1.6	9.5	56.1	282.0	1,342.4
1966-----	19.0	0.1	-	-	0.0	0.0	0.3	1.8	10.7	61.7	304.3	1,397.9
1965-----	19.0	-	0.0	-	0.0	0.1	0.3	1.9	10.5	61.1	302.0	1,462.5
1964-----	18.5	-	-	-	0.0	0.0	0.2	1.8	11.2	58.4	298.2	1,448.2
1963-----	19.3	0.1	-	-	-	0.1	0.2	2.0	11.0	62.7	316.9	1,490.9
1962 <sup>2</sup> -----	19.2	-	-	0.0	-	0.0	0.2	2.4	10.9	58.7	321.1	1,496.8
1961-----	19.0	0.1	-	-	0.0	0.0	0.3	1.8	11.9	61.1	316.6	1,468.4
1960-----	20.0	-	-	-	-	0.1	0.3	2.3	12.5	65.9	346.6	1,502.2
1959-----	19.6	0.1	-	-	0.0	0.0	0.3	2.4	12.7	62.9	342.6	1,474.8
1958-----	20.1	-	-	0.0	0.0	0.1	0.3	1.8	11.8	67.7	354.9	1,539.6
1957-----	19.8	-	-	-	0.0	0.0	0.2	2.2	11.7	69.2	347.7	1,509.3
1956-----	19.6	-	0.0	-	0.0	0.0	0.2	2.2	10.9	65.3	362.9	1,494.7
1955-----	20.5	-	0.0	-	0.0	0.0	0.2	2.1	11.2	68.4	392.0	1,559.9
1954-----	19.3	0.1	0.0	-	0.0	0.1	0.2	2.0	11.3	66.7	368.8	1,461.7
1953-----	21.3	-	-	-	0.0	0.1	0.3	1.7	12.1	77.5	412.7	1,641.4
1952-----	21.1	-	-	0.0	0.0	0.0	0.3	2.4	13.1	76.7	412.6	1,662.8
1951-----	21.9	-	-	-	-	0.0	0.2	2.0	13.1	82.9	438.7	1,797.9
1950-----	21.7	-	-	-	-	0.1	0.3	2.0	13.9	80.9	445.3	1,937.6
<b>WHITE FEMALE</b>												
1969-----	19.3	-	-	0.0	0.0	0.0	0.1	0.7	4.0	29.9	192.4	1,195.1
1968-----	19.7	-	0.0	-	0.0	0.0	0.1	0.7	4.0	31.4	202.8	1,224.0
1967-----	22.1	-	-	-	0.0	0.0	0.1	0.8	5.5	36.8	235.0	1,405.0
1966-----	22.7	-	-	-	0.0	0.0	0.2	1.0	5.5	40.2	250.0	1,455.2
1965-----	22.3	-	-	-	0.0	0.1	0.1	0.8	5.3	39.3	253.8	1,470.2
1964-----	22.2	-	-	-	0.0	0.0	0.2	0.9	6.1	40.7	258.7	1,470.9
1963 <sup>2</sup> -----	22.4	-	-	-	0.0	0.0	0.1	1.0	5.7	41.6	270.6	1,481.0
1962 <sup>2</sup> -----	22.5	0.1	0.0	-	0.0	-	0.1	0.9	5.7	42.5	274.4	1,502.0
1961-----	21.5	-	-	0.0	-	0.1	0.1	0.8	5.8	41.1	267.8	1,445.8
1960-----	21.8	-	-	-	-	0.0	0.1	0.8	6.4	43.4	285.1	1,452.8
1959-----	21.3	-	-	-	0.0	0.0	0.1	1.0	6.2	42.5	285.3	1,458.9
1958-----	21.3	-	-	-	0.0	0.0	0.1	1.0	6.1	44.6	286.5	1,504.6
1957-----	20.9	-	-	-	0.0	0.0	0.1	1.0	6.2	44.1	291.5	1,492.6
1956-----	20.4	-	-	0.0	0.0	0.0	0.2	0.9	5.6	43.4	294.7	1,493.8
1955-----	20.7	-	-	-	0.0	0.0	0.1	0.8	5.8	45.4	308.0	1,539.0
1954-----	19.8	-	-	-	0.0	0.0	0.2	0.9	6.2	44.9	296.7	1,518.6
1953-----	21.2	-	-	-	0.0	0.1	0.2	0.9	6.4	50.7	328.6	1,670.4
1952-----	21.4	-	-	-	0.0	0.1	0.1	0.9	7.2	53.0	349.6	1,653.7
1951-----	21.6	-	0.0	-	-	0.1	0.2	1.0	7.4	55.2	367.1	1,691.0
1950-----	20.9	-	-	-	0.0	-	0.1	0.9	7.4	54.4	362.2	1,706.0

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 9.—Death Rates for Arteriosclerosis, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 440 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 450 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	9.7	-	-	0.0	-	-	0.7	3.6	15.1	83.2	161.4	439.3
1968-----	10.2	-	-	-	-	0.1	0.4	3.3	18.4	93.5	190.0	440.0
1967-----	11.1	-	-	-	0.0	0.1	0.7	2.9	19.5	96.9	209.5	544.0
1966-----	12.3	-	-	-	0.1	0.1	0.8	4.0	25.1	96.1	240.0	634.8
1965-----	12.4	-	-	-	-	0.1	1.0	5.2	23.1	100.0	242.3	685.4
1964-----	12.3	-	-	-	-	-	0.7	4.8	25.3	98.8	245.1	678.4
1963 <sup>2</sup> -----	13.4	-	-	-	-	0.3	0.8	4.4	24.8	104.2	268.4	857.6
1962 <sup>2</sup> -----	13.7	-	-	-	-	0.3	0.8	5.2	24.2	107.4	253.8	964.5
1961-----	12.9	-	-	-	0.1	-	0.7	3.8	26.2	102.6	252.8	803.2
1960-----	14.0	-	-	-	-	0.2	1.3	4.6	31.6	101.7	269.8	937.5
1959-----	13.6	-	-	-	-	0.2	1.0	6.9	26.2	105.8	286.8	734.5
1958-----	14.2	-	0.1	-	0.1	0.2	0.9	5.7	25.8	107.2	303.6	940.7
1957-----	13.5	-	-	-	0.1	0.2	0.5	4.4	27.9	107.1	267.4	900.0
1956-----	13.0	-	-	-	0.1	0.1	1.0	5.1	32.0	100.3	249.2	812.0
1955-----	14.2	-	-	-	0.2	0.1	1.3	7.8	32.1	110.2	276.8	854.2
1954-----	13.4	-	-	-	-	0.2	1.2	6.7	30.6	96.3	284.9	886.4
1953-----	15.1	-	-	-	0.1	0.4	1.1	7.9	39.4	114.0	304.4	923.8
1952-----	14.4	-	-	-	0.1	0.3	1.6	6.5	37.9	106.3	304.5	925.0
1951-----	14.2	-	-	-	-	0.3	0.7	9.4	37.3	106.7	308.5	852.6
1950 <sup>1</sup> -----	14.4	-	-	-	-	0.2	0.9	7.1	37.3	124.5	315.2	905.3
ALL OTHER FEMALE												
1969-----	9.6	-	-	-	-	0.1	0.4	1.9	11.3	62.9	150.7	467.5
1968-----	10.1	-	-	-	-	0.1	0.3	2.4	15.3	67.8	166.4	456.9
1967-----	10.7	-	-	-	-	-	0.1	2.2	14.4	75.0	177.3	559.1
1966-----	12.1	-	-	-	0.0	0.1	0.3	3.6	19.6	89.3	188.3	625.0
1965-----	11.9	-	-	-	-	-	0.7	3.2	18.9	78.5	211.0	674.1
1964-----	12.1	-	-	-	-	-	0.3	3.6	19.3	83.6	213.5	732.7
1963 <sup>2</sup> -----	12.5	-	-	-	-	0.1	0.4	3.7	22.7	83.7	223.6	768.2
1962 <sup>2</sup> -----	12.2	-	-	-	-	0.1	0.6	4.0	17.8	77.7	232.3	823.8
1961-----	11.4	-	-	-	-	0.2	0.3	3.4	20.2	74.6	210.9	756.1
1960-----	12.0	-	-	-	-	0.1	0.7	3.1	21.4	83.7	227.4	767.2
1959-----	12.0	-	-	-	-	0.1	0.7	4.8	24.9	73.0	229.8	748.7
1958-----	13.2	-	-	-	0.1	0.1	1.0	4.1	27.0	88.0	256.2	773.7
1957-----	12.5	-	-	-	-	0.1	0.6	5.6	25.5	81.9	239.3	786.5
1956-----	11.8	-	-	-	0.1	0.1	0.6	5.0	23.6	83.7	226.8	688.9
1955-----	11.7	-	-	-	-	0.1	1.1	3.6	26.1	65.6	260.6	767.6
1954-----	11.6	-	-	-	-	0.1	1.2	5.5	27.6	74.3	221.4	756.3
1953-----	12.1	-	-	-	-	0.3	0.6	5.7	29.6	77.8	260.3	732.3
1952-----	11.7	-	-	-	-	0.2	1.1	6.4	27.9	77.3	227.6	824.1
1951-----	11.4	-	-	-	-	0.1	1.4	5.4	31.8	78.6	219.8	760.7
1950 <sup>1</sup> -----	11.3	-	-	-	0.1	0.4	1.2	6.6	26.6	91.1	220.6	823.2

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 10.—Death Rates for Bronchitis, Emphysema, and Asthma, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 490-493 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 501,502,527.1, and 241 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	15.4	4.5	0.8	0.3	0.5	0.9	2.7	10.1	37.5	90.3	126.8	146.5
1968-----	16.6	5.1	0.9	0.3	0.5	1.0	2.9	10.8	41.3	97.6	135.5	154.7
1967-----	15.3	5.5	0.9	0.3	0.5	1.0	3.0	10.8	38.9	90.2	123.3	139.3
1966-----	15.2	5.9	0.7	0.4	0.5	1.0	3.0	10.9	39.1	90.3	122.3	134.3
1965-----	14.4	7.7	0.8	0.3	0.5	1.0	2.9	10.8	37.0	86.0	114.1	130.9
1964-----	12.9	6.3	1.0	0.3	0.5	0.8	2.9	10.4	33.6	75.3	100.1	115.6
1963 <sup>1</sup> -----	13.0	7.5	1.0	0.3	0.5	1.0	2.7	10.3	34.5	76.7	97.5	121.7
1962-----	11.2	7.7	0.9	0.3	0.5	1.0	2.7	9.2	29.6	64.8	85.0	99.6
1961-----	9.9	8.0	1.0	0.3	0.5	1.0	2.5	8.0	27.1	56.0	73.4	87.2
1960-----	9.9	8.7	1.2	0.3	0.5	0.9	2.9	8.9	26.9	55.2	71.0	82.0
1959-----	8.7	9.4	1.2	0.3	0.4	1.0	2.4	7.8	24.4	47.3	60.4	76.5
1958-----	8.3	9.2	1.4	0.3	0.4	0.9	2.3	7.1	23.4	44.4	58.7	71.5
1957-----	8.7	10.3	1.5	0.3	0.5	0.9	2.5	7.6	24.3	46.1	63.5	73.4
1956-----	7.5	11.3	1.4	0.2	0.4	0.7	1.9	6.3	21.1	38.9	55.6	74.2
1955-----	7.0	9.2	1.5	0.3	0.5	0.8	2.1	6.2	19.4	36.6	51.1	72.5
1954 <sup>1</sup> -----	6.8	10.5	1.5	0.2	0.5	0.9	2.0	6.2	18.8	34.7	51.3	68.9
1953 <sup>1</sup> -----	7.2	10.9	1.6	0.3	0.4	0.8	1.9	5.1	14.3	27.4	41.6	61.1
1952 <sup>1</sup> -----	6.9	11.3	1.6	0.3	0.4	0.9	2.0	5.6	14.8	28.2	41.1	60.2
1951 <sup>1</sup> -----	6.9	9.9	1.8	0.3	0.4	0.8	1.9	5.8	14.9	27.9	44.6	74.7
1950 <sup>1,2</sup> -----	5.0	10.4	1.8	0.3	0.4	0.8	1.6	4.5	10.3	17.8	30.3	65.0
<b>WHITE MALE</b>												
1969-----	26.7	3.9	0.7	0.2	0.4	0.6	2.8	13.3	64.2	179.2	270.2	281.6
1968-----	28.6	5.0	0.8	0.3	0.3	0.6	2.5	14.3	70.6	194.0	284.3	305.1
1967-----	26.7	6.3	0.8	0.3	0.4	0.6	2.7	14.6	68.0	179.1	258.9	274.0
1966-----	26.6	6.1	0.7	0.4	0.3	0.6	2.6	14.7	69.2	179.6	257.0	268.1
1965-----	25.0	7.4	0.6	0.3	0.3	0.5	2.7	14.8	65.2	172.4	231.2	253.1
1964-----	22.2	6.1	0.9	0.3	0.4	0.5	2.6	14.7	59.2	149.1	199.0	217.7
1963 <sup>1</sup> -----	22.5	7.5	0.8	0.3	0.3	0.5	2.4	15.0	62.4	153.0	191.9	223.7
1962 <sup>3</sup> -----	19.2	7.4	0.8	0.4	0.2	0.5	2.6	12.9	53.7	128.4	160.3	179.4
1961-----	16.6	8.8	1.0	0.2	0.2	0.6	2.4	11.0	48.3	107.3	138.8	146.7
1960-----	16.5	9.8	1.0	0.2	0.2	0.5	2.5	12.7	47.4	105.6	131.3	136.3
1959-----	14.5	9.6	1.0	0.2	0.3	0.6	2.1	11.2	44.3	89.7	110.1	131.9
1958-----	13.7	9.2	1.4	0.3	0.2	0.6	2.3	10.0	41.9	85.8	107.3	111.4
1957-----	13.9	10.0	1.5	0.3	0.3	0.6	2.4	10.7	42.3	85.6	108.6	112.5
1956-----	11.9	10.9	1.5	0.2	0.3	0.5	1.8	9.0	36.9	71.1	92.3	98.0
1955-----	10.9	9.3	1.5	0.3	0.2	0.4	2.0	8.9	35.1	65.2	81.5	98.6
1954 <sup>1</sup> -----	10.4	11.3	1.5	0.2	0.4	0.7	2.0	8.3	31.6	59.7	80.8	88.0
1953 <sup>1</sup> -----	10.9	11.0	1.7	0.3	0.2	0.4	1.9	6.3	22.1	43.3	60.5	79.3
1952 <sup>1</sup> -----	10.2	10.8	1.8	0.3	0.3	0.4	1.8	7.3	22.5	44.0	59.8	70.7
1951 <sup>1</sup> -----	10.0	9.3	1.8	0.3	0.4	0.5	1.6	7.3	23.4	43.1	61.2	92.5
1950 <sup>1</sup> -----	6.8	10.9	1.8	0.3	0.2	0.4	1.3	5.7	15.0	25.6	39.8	86.2
<b>WHITE FEMALE</b>												
1969-----	6.5	3.2	0.5	0.2	0.3	0.6	1.9	6.0	14.7	25.2	39.6	84.0
1968-----	6.9	4.0	0.6	0.3	0.4	0.8	2.0	6.1	16.1	27.0	42.6	82.9
1967-----	6.1	3.0	0.7	0.2	0.4	0.8	2.2	6.4	14.0	24.0	37.2	71.9
1966-----	6.0	3.4	0.5	0.2	0.4	0.9	2.2	6.2	13.4	24.2	36.6	66.1
1965-----	5.7	5.2	0.7	0.2	0.5	0.8	2.2	5.9	12.5	21.8	37.6	66.1
1964-----	5.2	4.6	0.8	0.3	0.4	0.7	2.2	5.6	10.7	19.7	34.9	63.0
1963 <sup>1</sup> -----	5.1	5.4	0.7	0.3	0.4	0.9	2.1	5.4	10.6	18.9	35.4	67.0
1962 <sup>3</sup> -----	4.6	5.5	0.6	0.2	0.4	0.7	2.0	5.1	9.0	16.5	32.1	56.5
1961-----	4.1	5.7	0.6	0.2	0.4	0.8	1.7	4.4	8.1	15.5	28.3	52.2
1960-----	4.2	5.5	1.0	0.2	0.5	0.7	2.3	4.4	8.4	14.6	29.5	50.9
1959-----	3.6	6.3	1.0	0.2	0.3	0.8	1.7	3.9	6.3	13.3	25.6	45.9
1958-----	3.5	6.8	1.2	0.2	0.5	0.7	1.4	3.6	6.4	12.1	24.6	51.5
1957-----	4.0	7.6	1.2	0.2	0.4	0.7	1.8	3.6	7.2	15.3	31.7	50.0
1956-----	3.6	8.4	1.0	0.2	0.4	0.7	1.4	3.0	6.5	12.6	29.8	63.2
1955-----	3.7	6.5	1.4	0.2	0.5	0.8	1.6	3.1	6.5	13.7	30.1	56.6
1954 <sup>1</sup> -----	3.8	6.2	1.3	0.2	0.4	0.8	1.7	3.6	6.7	14.7	30.7	59.4
1953 <sup>1</sup> -----	4.0	7.7	1.2	0.2	0.4	0.8	1.5	3.5	6.4	14.7	28.9	51.7
1952 <sup>1</sup> -----	4.0	8.6	1.2	0.3	0.5	1.0	1.7	3.2	7.2	15.0	28.2	53.4
1951 <sup>1</sup> -----	4.2	8.6	1.4	0.2	0.4	0.9	1.8	3.6	6.5	15.5	33.4	62.8
1950 <sup>1</sup> -----	3.3	7.8	1.4	0.2	0.5	0.9	1.4	3.2	5.6	11.0	23.8	53.5

<sup>1</sup>Excludes data for Emphysema without mention of bronchitis (ICD No. 527.1) because for these years data were not available.

<sup>2</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>3</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 10.—Death Rates for Bronchitis, Emphysema, and Asthma, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 490-493 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 501,502,527.1, and 241 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	13.2	8.9	1.8	0.8	1.7	1.8	6.6	21.1	51.5	103.3	112.3	90.2
1968-----	14.7	8.5	1.6	0.7	1.4	2.9	8.9	26.7	56.0	107.7	133.3	90.9
1967-----	13.3	9.2	1.2	0.7	1.2	2.8	7.8	21.0	49.9	105.3	113.1	108.0
1966-----	13.1	12.6	1.4	0.9	1.3	2.9	8.3	23.8	49.1	97.6	91.3	97.8
1965-----	13.0	15.2	1.6	0.8	1.4	2.7	8.0	23.9	48.2	86.6	100.0	126.8
1964-----	11.8	11.0	2.3	0.6	1.4	2.4	8.1	18.0	48.6	84.8	78.3	64.9
1963 <sup>3</sup> -----	12.3	13.0	2.4	0.8	1.5	2.3	7.0	19.9	44.6	89.6	84.5	127.3
1962 <sup>3</sup> -----	10.6	15.7	2.7	0.6	1.3	3.2	7.3	17.2	38.0	72.3	58.6	71.0
1961-----	10.3	15.2	2.9	0.8	1.2	3.5	6.8	17.1	36.2	61.2	62.6	96.8
1960-----	10.0	13.1	2.2	0.5	1.2	2.7	7.1	16.9	37.7	60.9	54.0	79.7
1959-----	9.0	19.0	2.7	0.9	0.7	3.4	7.1	15.4	31.6	47.2	45.8	51.7
1958-----	8.5	19.4	2.4	0.7	0.8	2.6	6.1	11.8	33.1	49.9	44.6	35.3
1957-----	10.0	23.3	2.9	0.5	0.9	2.3	7.0	17.2	36.0	50.9	60.0	59.3
1956-----	7.5	27.6	2.2	0.4	1.0	2.0	3.8	12.4	29.5	35.3	39.2	47.8
1955-----	7.0	21.1	2.4	0.5	1.3	2.0	4.1	10.1	26.5	33.7	38.4	77.3
1954-----	6.6	24.6	2.0	0.3	0.7	2.4	4.6	11.1	25.1	29.0	32.8	30.0
1953 <sup>1</sup> -----	6.9	24.4	2.6	0.6	1.0	1.7	3.5	8.6	20.6	21.8	23.7	38.9
1952 <sup>1</sup> -----	6.9	27.0	3.0	0.4	0.9	2.0	3.6	11.6	17.6	27.8	24.5	29.4
1951 <sup>1</sup> -----	6.3	18.6	3.3	0.5	0.6	1.5	3.5	9.8	16.4	23.5	32.1	46.7
1950 <sup>1,2</sup> -----	5.4	22.7	3.8	0.9	0.5	1.4	3.9	6.2	11.4	19.5	22.8	42.9
ALL OTHER FEMALE												
1969-----	4.7	8.7	1.7	0.5	1.5	3.3	4.9	8.5	13.4	16.9	19.5	27.5
1968-----	5.3	7.3	1.5	0.7	1.6	3.6	6.8	9.7	15.1	17.8	20.0	36.1
1967-----	5.0	10.0	1.6	0.6	1.4	3.3	6.0	8.8	13.6	19.7	17.9	31.8
1966-----	4.5	9.9	0.9	0.7	1.7	2.3	6.4	9.2	11.7	16.9	13.4	21.7
1965-----	4.5	14.2	1.5	0.4	1.7	3.7	5.4	8.3	11.7	13.8	15.4	25.9
1964-----	4.3	11.8	1.1	0.4	1.2	2.7	6.5	8.7	12.1	9.1	17.7	32.7
1963 <sup>3</sup> -----	5.2	12.3	2.3	0.6	1.5	3.7	6.0	9.5	13.0	16.9	16.6	50.0
1962 <sup>3</sup> -----	4.8	13.1	1.5	0.7	2.1	4.2	6.0	7.7	11.0	12.9	18.5	28.6
1961-----	4.4	8.8	1.7	0.6	2.1	3.3	5.5	6.2	11.1	11.6	18.5	48.8
1960-----	4.9	15.3	2.4	0.7	1.8	3.5	6.9	7.8	9.6	12.6	17.3	44.7
1959-----	4.6	15.9	2.4	0.5	1.8	2.9	6.5	6.4	10.9	10.1	19.3	26.3
1958-----	4.4	13.4	1.8	0.6	1.3	2.9	6.6	8.8	8.7	12.2	16.3	16.2
1957-----	4.7	15.3	1.3	0.6	1.8	4.1	5.4	7.7	12.0	13.3	11.0	37.1
1956-----	3.7	14.2	2.4	0.6	0.9	2.1	3.8	6.3	7.7	12.3	19.6	18.2
1955-----	3.6	13.7	1.4	0.5	1.3	2.8	4.7	5.3	10.0	6.9	11.4	33.3
1954-----	3.6	18.4	2.0	0.3	1.2	2.0	3.3	5.8	8.1	8.9	19.8	35.7
1953 <sup>1</sup> -----	4.1	17.4	2.5	0.4	0.8	2.4	3.3	6.8	11.4	10.8	16.5	24.0
1952 <sup>1</sup> -----	4.5	16.0	2.1	0.3	0.9	2.7	4.8	6.2	12.0	13.7	19.8	73.9
1951 <sup>1</sup> -----	4.4	13.5	2.6	0.5	1.0	2.5	3.5	6.9	12.9	12.1	16.2	85.7
1950 <sup>1,2</sup> -----	3.6	11.8	2.8	0.5	1.0	2.5	3.6	4.8	9.1	9.8	12.7	42.1

<sup>1</sup>Excludes data for Emphysema without mention of bronchitis (ICD No. 527.1) because for these years data were not available.

<sup>2</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>3</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 11.—Death Rates for Cirrhosis of Liver, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 571 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 581 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	14.8	1.2	0.2	0.1	0.5	4.3	19.0	36.5	47.7	40.9	28.8	21.8
1968-----	14.6	1.5	0.2	0.1	0.4	4.2	18.1	36.6	46.7	41.6	28.8	22.8
1967-----	14.1	1.0	0.1	0.1	0.4	4.0	17.2	34.8	45.7	41.1	27.4	22.8
1966-----	13.6	0.7	0.2	0.1	0.4	3.8	16.6	33.8	43.3	39.8	29.8	24.3
1965-----	12.8	0.8	0.2	0.1	0.3	3.5	15.0	31.9	40.1	38.0	29.9	21.7
1964-----	12.1	0.7	0.1	0.1	0.3	3.3	13.9	29.5	38.2	36.8	29.4	25.4
1963 <sup>2</sup> -----	11.9	0.9	0.1	0.1	0.3	3.3	12.7	29.9	36.4	37.6	29.5	26.1
1962 <sup>2</sup> -----	11.7	0.6	0.2	0.1	0.3	3.2	13.0	28.9	35.4	37.1	29.4	26.2
1961-----	11.3	0.7	0.1	0.1	0.3	3.1	12.3	27.5	34.2	35.7	30.1	25.9
1960-----	11.3	0.8	0.2	0.1	0.3	2.9	11.8	27.6	32.7	37.4	32.1	28.8
1959-----	10.9	0.9	0.2	0.1	0.3	2.8	11.5	26.1	31.8	34.9	33.0	28.6
1958-----	10.8	1.0	0.1	0.1	0.3	2.5	11.1	24.9	31.3	36.6	34.0	31.2
1957-----	11.3	1.0	0.2	0.1	0.3	2.9	12.1	25.6	33.0	38.2	35.3	32.3
1956-----	10.7	1.0	0.2	0.1	0.3	2.4	11.2	23.8	30.9	36.8	37.3	32.7
1955-----	10.2	1.0	0.2	0.1	0.3	2.1	10.5	22.8	28.8	36.1	35.9	37.0
1954-----	10.1	1.2	0.1	0.1	0.3	2.2	10.4	21.5	29.1	35.5	35.7	37.4
1953-----	10.4	1.2	0.1	0.1	0.3	2.3	10.7	23.0	29.9	34.7	38.8	37.8
1952-----	10.2	1.3	0.2	0.2	0.4	2.4	10.6	22.4	29.9	34.4	37.6	37.9
1951-----	9.8	1.3	0.1	0.1	0.3	2.3	10.1	21.1	28.7	34.0	38.7	40.6
1950 <sup>1</sup> -----	9.2	1.2	0.1	0.1	0.3	2.1	9.3	19.1	27.2	33.8	36.9	39.0
<b>WHITE MALE</b>												
1969-----	19.1	1.3	0.2	0.1	0.3	3.3	19.2	45.0	66.6	62.5	44.0	35.2
1968-----	18.8	1.4	0.1	0.1	0.2	3.0	18.8	44.9	66.8	63.3	44.6	31.4
1967-----	18.2	1.0	0.0	0.1	0.1	3.0	18.1	43.1	65.1	62.7	41.2	32.3
1966-----	17.8	0.5	0.2	0.1	0.2	2.5	17.3	41.9	63.3	61.3	45.2	37.4
1965-----	16.6	0.7	0.2	0.1	0.2	2.5	15.1	39.5	57.9	58.7	47.1	31.8
1964-----	16.0	0.6	0.1	0.1	0.2	2.5	14.3	37.7	55.1	58.3	44.8	35.5
1963 <sup>2</sup> -----	15.7	0.8	0.1	0.1	0.2	2.5	13.2	37.8	52.4	59.1	43.6	37.1
1962 <sup>2</sup> -----	15.7	0.5	0.2	0.1	0.2	2.5	13.6	38.0	51.6	57.2	44.2	38.9
1961-----	15.4	0.5	0.1	0.1	0.2	2.5	13.3	35.8	50.3	56.9	46.7	34.5
1960-----	15.6	0.7	0.2	0.1	0.2	2.3	13.4	36.5	49.2	59.7	46.8	41.7
1959-----	14.9	1.0	0.1	0.1	0.2	2.2	13.3	34.7	47.0	54.9	48.3	38.7
1958-----	14.9	1.0	0.1	0.1	0.1	2.1	13.0	34.0	46.2	57.9	48.0	40.8
1957-----	15.8	1.0	0.1	0.1	0.2	2.6	14.1	35.7	48.7	59.6	50.0	44.1
1956-----	14.7	0.5	0.2	0.1	0.3	2.0	13.2	32.4	44.7	56.7	50.5	44.0
1955-----	14.2	1.2	0.2	0.1	0.2	1.8	12.7	31.4	42.4	54.3	51.7	44.6
1954-----	14.1	1.1	0.2	0.1	0.3	1.9	12.6	30.4	43.0	53.5	49.0	48.9
1953-----	14.3	0.8	0.1	0.1	0.2	2.2	13.0	31.2	44.2	51.7	53.5	47.0
1952-----	14.3	1.0	0.1	0.1	0.2	2.2	13.3	30.6	44.5	52.3	51.9	50.2
1951-----	13.5	1.4	0.1	0.1	0.2	2.1	12.6	28.5	40.2	50.2	54.9	43.5
1950-----	12.6	1.3	0.1	0.1	0.2	2.0	11.9	26.3	38.8	48.9	47.2	47.2
<b>WHITE FEMALE</b>												
1969-----	9.5	1.4	0.3	0.2	0.4	1.8	10.6	23.5	27.7	23.8	20.3	15.4
1968-----	9.6	1.4	0.3	0.1	0.4	1.8	10.4	23.6	27.9	25.0	19.9	19.9
1967-----	9.4	0.6	0.2	0.1	0.3	1.8	10.0	23.7	27.9	24.2	19.4	19.4
1966-----	8.9	0.9	0.1	0.1	0.3	1.8	9.9	22.4	24.6	23.4	20.6	17.8
1965-----	8.5	0.6	0.2	0.1	0.3	1.8	9.3	22.0	23.4	22.7	19.3	16.8
1964-----	8.2	0.7	0.1	0.1	0.3	1.7	9.3	20.2	23.4	20.9	19.6	21.8
1963 <sup>2</sup> -----	8.0	0.7	0.2	0.1	0.3	1.6	8.8	20.9	21.2	20.8	20.5	21.6
1962 <sup>2</sup> -----	7.6	0.4	0.1	0.1	0.2	1.7	9.0	18.9	19.7	20.6	19.6	19.9
1961-----	7.5	0.5	0.1	0.1	0.3	1.6	8.6	18.7	20.1	19.5	19.6	22.3
1960-----	7.4	0.8	0.1	0.2	0.2	1.8	8.2	18.1	18.0	20.8	22.7	22.2
1959-----	7.3	0.8	0.2	0.1	0.2	1.8	8.2	17.2	18.3	19.4	23.4	24.6
1958-----	7.2	1.1	0.1	0.1	0.3	1.6	8.1	16.0	18.5	20.2	25.4	25.9
1957-----	7.5	0.8	0.1	0.1	0.3	2.2	9.0	15.9	18.3	22.0	25.3	26.9
1956-----	7.4	1.1	0.1	0.1	0.3	1.8	8.1	15.7	18.2	21.7	29.2	27.9
1955-----	7.0	0.6	0.2	0.1	0.3	1.7	7.7	14.5	16.9	22.2	25.0	34.2
1954-----	6.8	1.4	0.1	0.2	0.3	1.8	7.8	13.3	16.6	21.7	27.2	32.1
1953-----	7.2	0.9	0.2	0.1	0.3	1.8	8.4	15.1	17.4	21.7	29.8	33.7
1952-----	7.0	1.6	0.2	0.1	0.4	2.0	7.8	14.7	16.8	20.6	28.3	30.4
1951-----	6.9	1.0	0.1	0.2	0.3	2.1	7.4	13.9	18.0	21.1	27.5	41.3
1950-----	6.5	0.9	0.2	0.1	0.5	1.8	6.7	12.4	16.4	21.0	29.9	36.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 11.—Death Rates for Cirrhosis of Liver, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category number 571 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 581 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	23.9	0.6	0.2	0.0	1.9	20.2	66.8	75.7	76.7	61.5	22.1	18.0
1968-----	22.8	1.6	0.3	0.0	1.1	21.2	55.7	61.0	70.2	56.4	23.9	10.9
1967-----	21.1	2.3	0.2	0.0	1.7	18.2	56.3	64.6	62.4	59.4	27.1	24.0
1966-----	20.7	0.3	0.2	0.1	1.5	20.3	51.7	65.5	61.0	55.5	24.6	23.9
1965-----	18.3	1.2	0.2	0.1	0.9	15.3	46.3	59.3	57.4	42.1	25.4	19.5
1964-----	15.6	0.9	-	0.1	0.7	12.3	37.8	49.6	49.3	37.7	28.3	8.1
1963 <sup>2</sup> -----	13.9	1.9	0.2	0.2	0.4	11.3	31.9	39.4	46.9	37.6	23.6	15.2
1962 <sup>2</sup> -----	13.8	1.5	0.2	0.0	1.1	10.4	31.6	36.6	45.1	37.3	24.3	29.0
1961-----	13.3	2.1	0.5	0.2	0.8	12.1	27.4	36.6	41.0	35.0	20.9	9.7
1960-----	12.6	1.3	0.1	0.1	0.5	9.3	23.8	39.1	41.5	30.9	23.3	22.3
1959-----	11.7	1.3	0.3	0.2	0.5	8.9	21.4	34.5	37.2	34.3	19.4	24.1
1958-----	10.7	1.0	0.3	0.1	0.8	6.6	17.1	33.1	37.2	35.4	21.6	44.4
1957-----	11.6	2.4	0.3	0.2	0.2	7.4	19.6	31.5	42.5	31.4	35.6	19.2
1956-----	10.2	2.2	0.4	0.1	0.6	5.7	17.8	27.0	36.1	29.9	23.8	28.0
1955-----	9.0	1.9	0.2	0.1	0.3	4.4	14.1	26.0	29.3	30.4	29.6	33.3
1954-----	8.6	1.6	0.1	0.1	0.5	5.0	13.5	21.6	32.1	27.6	26.9	27.3
1953-----	8.9	3.8	-	0.2	0.4	4.7	11.6	27.7	30.2	28.8	15.8	19.0
1952-----	9.4	1.4	0.1	0.2	0.6	5.0	13.8	25.8	33.8	27.2	27.3	40.0
1951-----	8.7	0.9	-	0.2	0.2	3.4	12.7	24.8	32.3	29.7	24.5	26.3
1950 <sup>1</sup> -----	7.8	2.5	0.1	0.3	0.2	2.7	10.0	19.4	30.2	36.9	28.7	5.4
ALL OTHER FEMALE												
1969-----	14.4	1.0	0.2	0.1	1.3	15.0	39.2	45.5	33.7	24.1	12.5	10.0
1968-----	13.9	2.0	0.3	0.3	1.4	14.2	38.8	44.0	31.0	22.3	9.1	9.7
1967-----	12.3	1.3	0.2	0.2	1.3	14.7	31.3	36.8	31.8	19.9	6.4	-
1966-----	12.4	1.0	0.2	0.3	1.1	12.7	32.9	38.0	31.0	18.0	10.9	8.3
1965-----	11.1	0.9	0.2	0.2	1.4	12.3	30.9	30.7	26.9	16.5	9.7	9.3
1964-----	9.5	1.5	0.4	0.2	1.2	11.9	26.3	25.2	19.8	13.9	10.2	8.2
1963 <sup>2</sup> -----	9.3	1.9	0.1	0.3	0.7	12.3	21.0	27.1	21.8	15.6	8.5	13.6
1962 <sup>2</sup> -----	9.3	1.3	0.3	0.1	1.4	11.4	21.3	26.8	21.0	17.0	11.1	7.1
1961-----	8.4	1.2	-	0.2	1.0	9.9	21.3	23.4	18.7	12.7	8.7	14.6
1960-----	7.9	1.3	0.3	0.1	0.7	9.0	19.5	23.1	17.1	11.5	13.1	14.9
1959-----	7.9	0.7	0.4	0.1	1.0	9.4	16.5	22.1	20.2	11.7	11.8	-
1958-----	6.5	1.0	-	0.0	0.9	7.8	15.0	16.8	15.1	9.6	11.1	7.9
1957-----	7.4	1.0	0.4	0.3	1.4	7.0	15.5	19.0	21.5	12.3	11.0	8.1
1956-----	6.9	1.8	0.2	0.2	0.9	7.0	14.9	18.3	18.6	10.1	13.0	-
1955-----	6.0	1.9	0.3	0.2	0.4	5.5	12.0	18.1	13.6	11.2	15.2	8.8
1954-----	5.7	1.2	-	0.2	0.7	4.3	11.7	16.5	15.8	11.5	8.7	9.4
1953-----	5.7	2.1	0.2	0.2	1.0	4.2	11.1	16.7	13.3	9.8	14.9	19.4
1952-----	5.1	1.8	0.1	0.2	1.0	4.3	9.5	15.0	11.8	9.2	10.3	20.7
1951-----	5.6	2.7	0.2	-	0.7	4.1	10.1	15.9	15.4	11.8	17.1	17.9
1950 <sup>1</sup> -----	5.2	1.0	0.3	0.3	0.5	3.6	9.2	14.6	12.9	14.7	17.6	19.1

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 12.—Death Rates for Suicide, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E950-E959 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E963, E970-E979 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	11.1	...	...	0.3	8.0	12.9	16.6	19.4	21.3	20.7	22.1	20.7
1968-----	10.7	...	...	0.3	7.1	12.1	16.2	19.6	21.8	19.6	20.8	22.1
1967-----	10.8	...	...	0.3	7.0	12.4	16.6	19.5	22.4	19.8	21.0	22.7
1966-----	10.9	...	...	0.3	6.4	12.3	15.8	20.0	22.9	21.6	24.3	23.8
1965-----	11.1	...	...	0.3	6.2	12.3	16.7	20.7	23.8	22.0	24.2	24.4
1964-----	10.8	...	...	0.2	6.0	11.9	15.6	20.5	22.7	22.1	23.9	25.3
1963-----	11.0	...	...	0.3	6.0	11.8	16.0	21.1	23.6	22.4	25.4	24.6
1962-----	10.9	...	...	0.3	5.7	11.3	15.0	21.0	23.7	22.2	27.2	26.7
1961-----	10.4	...	...	0.2	5.1	10.3	14.4	20.3	23.1	22.0	26.0	24.9
1960-----	10.6	...	...	0.3	5.2	10.0	14.2	20.7	23.7	23.0	27.9	26.0
1959-----	10.6	...	...	0.2	4.9	9.9	13.6	19.8	24.3	24.8	27.8	25.7
1958-----	10.7	...	...	0.2	4.8	9.8	13.7	20.7	24.1	25.1	27.7	25.8
1957-----	9.8	...	...	0.2	4.0	8.7	12.7	18.2	22.4	23.6	26.5	26.4
1956-----	10.0	...	...	0.2	4.0	8.5	12.1	18.5	24.2	25.2	28.0	23.5
1955-----	10.2	...	...	0.1	4.1	8.4	12.3	19.6	24.8	25.0	27.5	27.9
1954-----	10.1	...	...	0.1	4.2	8.7	12.5	19.3	23.9	25.1	25.6	25.1
1953-----	10.1	...	...	0.2	4.4	8.5	12.6	18.7	22.4	25.4	28.0	27.9
1952-----	10.0	...	...	0.2	4.2	8.5	12.4	18.3	22.5	25.6	27.9	31.2
1951-----	10.4	...	...	0.2	4.4	8.6	13.3	18.8	25.2	27.1	27.7	30.5
1950-----	11.4	...	...	0.2	4.5	9.1	14.3	20.9	26.8	29.6	31.1	28.8
<b>WHITE MALE</b>												
1969-----	17.2	...	...	0.5	12.6	18.3	22.4	28.6	34.9	37.0	46.7	52.2
1968-----	16.9	...	...	0.5	11.3	17.2	23.2	28.8	36.1	36.2	44.2	56.8
1967-----	16.8	...	...	0.5	10.8	17.2	23.8	29.2	36.5	34.6	43.8	56.3
1966-----	17.2	...	...	0.5	9.9	17.2	22.7	30.1	38.4	38.8	51.1	59.0
1965-----	17.4	...	...	0.5	9.6	17.7	23.4	30.8	39.7	39.3	49.5	57.7
1964-----	17.2	...	...	0.5	9.3	17.0	22.2	31.8	39.5	39.1	50.5	65.1
1963 <sup>2</sup> -----	17.8	...	...	0.5	9.2	16.9	23.6	33.0	40.2	40.3	52.5	59.4
1962 <sup>2</sup> -----	17.8	...	...	0.5	8.7	16.5	22.8	33.4	40.5	41.3	57.0	61.4
1961-----	17.1	...	...	0.4	7.9	14.7	22.3	32.8	39.7	39.9	53.2	60.3
1960-----	17.6	...	...	0.5	8.6	14.9	21.9	33.7	40.2	42.0	55.7	61.3
1959-----	17.7	...	...	0.5	7.9	14.4	21.5	32.8	41.8	45.0	56.6	63.8
1958-----	18.0	...	...	0.4	7.6	14.3	22.1	34.0	41.9	45.8	58.3	58.5
1957-----	16.5	...	...	0.4	6.5	12.7	20.1	30.3	38.1	43.5	55.3	62.7
1956-----	16.9	...	...	0.3	6.4	12.9	19.1	30.2	41.5	46.4	58.1	54.3
1955-----	17.2	...	...	0.2	6.2	12.7	19.8	31.7	42.8	45.6	54.7	61.2
1954-----	17.5	...	...	0.2	6.9	13.5	19.9	33.0	41.6	46.2	52.6	54.3
1953-----	17.2	...	...	0.4	7.0	13.0	20.5	31.0	38.4	47.5	55.9	63.4
1952-----	16.9	...	...	0.2	6.9	12.6	19.6	29.8	39.4	47.2	56.7	70.0
1951-----	17.3	...	...	0.3	6.8	12.8	20.5	30.3	39.7	50.3	53.8	70.9
1950-----	19.0	...	...	0.3	6.6	13.8	22.4	34.1	45.9	53.2	61.9	61.9
<b>WHITE FEMALE</b>												
1969-----	6.8	...	...	0.2	3.8	8.1	12.7	13.3	11.8	9.8	7.5	4.5
1968-----	6.3	...	...	0.1	3.4	7.5	11.5	13.4	11.6	8.1	6.5	4.9
1967-----	6.5	...	...	0.1	3.4	7.9	11.5	13.0	12.3	9.9	6.9	5.7
1966-----	6.3	...	...	0.1	3.2	7.8	11.3	12.9	11.5	9.7	7.4	4.3
1965-----	6.6	...	...	0.1	3.0	7.6	12.0	13.6	12.3	10.0	7.9	6.7
1964-----	6.1	...	...	0.1	2.9	7.3	10.9	12.5	10.9	10.4	6.7	4.4
1963 <sup>2</sup> -----	6.3	...	...	0.1	3.1	7.5	10.9	12.9	11.6	9.5	7.7	5.1
1962 <sup>2</sup> -----	5.9	...	...	0.1	2.9	7.3	9.5	12.4	11.1	8.8	7.6	7.6
1961-----	5.3	...	...	0.1	2.3	6.1	8.3	10.8	10.4	9.1	7.6	4.9
1960-----	5.3	...	...	0.1	2.3	5.8	8.1	10.9	10.9	8.8	9.2	6.1
1959-----	5.0	...	...	0.0	2.1	5.7	7.5	9.5	10.6	9.8	7.6	3.8
1958-----	5.1	...	...	0.1	2.4	5.9	7.1	10.4	10.0	9.6	6.6	6.7
1957-----	4.6	...	...	0.1	1.8	5.0	7.1	8.8	10.1	8.3	6.2	3.9
1956-----	4.8	...	...	0.1	2.0	4.7	6.7	9.6	10.5	9.4	6.8	4.6
1955-----	4.9	...	...	0.1	2.0	4.9	6.6	10.3	10.4	9.4	8.3	7.2
1954-----	4.5	...	...	0.1	1.8	4.4	6.8	8.2	9.5	8.7	6.3	7.1
1953-----	4.6	...	...	0.1	2.3	4.8	6.5	9.0	9.3	8.3	7.5	5.4
1952-----	4.7	...	...	0.1	2.1	5.0	6.9	9.4	9.2	9.0	7.0	6.3
1951-----	5.0	...	...	0.1	2.3	5.0	7.7	9.8	9.2	9.1	8.3	5.7
1950-----	5.5	...	...	0.1	2.7	5.2	8.2	10.5	10.7	10.6	8.4	8.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.



Table 12.—Death Rates for Suicide, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E950-E959 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E963, E970-E979 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	8.1	...	...	0.2	9.9	17.8	15.8	12.1	10.6	14.5	16.2	13.1
1968-----	7.3	...	...	0.2	8.2	16.0	12.6	12.3	12.4	14.9	11.9	5.5
1967-----	7.6	...	...	0.2	8.2	17.7	14.7	11.6	13.3	12.3	12.6	8.0
1966-----	7.8	...	...	0.1	8.6	17.9	11.4	14.1	14.4	14.1	15.4	23.9
1965-----	7.7	...	...	0.2	8.5	14.2	15.0	13.5	14.2	15.3	15.9	7.3
1964-----	7.2	...	...	0.1	8.0	16.2	12.8	11.8	12.3	13.4	12.5	2.7
1963 <sup>2</sup> -----	7.9	...	...	0.3	7.5	15.9	14.9	13.6	12.7	18.2	19.0	12.1
1962 <sup>2</sup> -----	7.2	...	...	0.1	7.5	12.8	12.8	12.4	14.6	16.8	13.6	12.9
1961-----	7.6	...	...	0.1	7.6	16.3	11.5	14.0	15.0	13.2	16.6	12.9
1960-----	7.2	...	...	0.1	5.3	12.9	13.5	12.8	16.9	12.6	11.3	15.9
1959-----	7.5	...	...	0.2	6.6	14.4	11.0	13.9	14.8	14.9	22.9	10.3
1958-----	7.1	...	...	0.2	5.4	12.6	12.3	12.8	15.3	14.2	11.5	14.8
1957-----	6.8	...	...	0.0	5.6	13.0	10.6	11.8	11.9	17.4	15.6	11.5
1956-----	6.1	...	...	0.1	5.8	10.8	10.5	9.8	12.2	10.0	11.5	12.0
1955-----	6.1	...	...	0.1	6.5	9.6	9.6	10.5	12.7	10.8	12.8	16.7
1954-----	6.8	...	...	0.1	5.5	12.8	10.6	11.4	12.2	13.6	13.4	4.5
1953-----	6.4	...	...	0.1	4.2	10.1	10.3	12.6	12.0	12.2	21.9	14.3
1952-----	6.1	...	...	0.3	4.5	10.7	8.9	9.9	14.3	11.0	10.9	20.0
1951-----	6.6	...	...	0.3	5.2	10.7	10.8	11.4	12.0	10.1	15.1	5.3
1950 <sup>1</sup> -----	7.0	...	...	0.1	5.3	10.1	11.3	11.7	16.8	15.0	7.9	16.1
ALL OTHER FEMALE												
1969-----	2.8	...	...	0.2	4.1	6.1	4.8	3.7	2.3	2.8	2.6	3.8
1968-----	2.4	...	...	0.1	3.1	4.6	4.5	3.6	2.8	3.0	3.8	2.8
1967-----	2.7	...	...	0.1	4.0	5.9	4.2	3.5	3.3	3.3	2.4	3.0
1966-----	2.4	...	...	0.1	2.9	6.2	3.1	3.5	3.6	3.7	2.5	3.3
1965-----	2.5	...	...	0.1	3.1	6.0	4.2	4.1	2.8	2.5	3.5	7.4
1964-----	2.2	...	...	0.0	2.0	4.7	4.1	3.4	4.0	3.2	2.8	-
1963 <sup>2</sup> -----	2.2	...	...	0.2	2.6	5.0	4.0	2.7	2.3	5.0	1.5	4.5
1962 <sup>2</sup> -----	2.2	...	...	0.0	3.0	4.6	3.9	3.3	3.9	2.0	3.2	2.4
1961-----	1.9	...	...	0.2	2.0	3.5	3.9	2.7	3.2	1.9	1.6	4.9
1960-----	2.0	...	...	0.0	1.5	3.5	3.7	3.2	3.4	3.8	4.2	5.0
1959-----	1.9	...	...	0.1	2.2	3.7	2.7	2.9	4.3	3.0	1.9	2.6
1958-----	1.8	...	...	0.0	1.7	3.6	2.8	3.1	2.9	2.4	3.3	2.6
1957-----	1.4	...	...	-	1.4	2.5	2.3	1.3	4.0	1.9	2.1	8.1
1956-----	1.6	...	...	0.1	1.2	2.6	3.0	3.0	3.3	2.7	1.4	-
1955-----	1.5	...	...	-	1.8	2.8	1.7	2.8	2.6	2.5	1.5	-
1954-----	1.5	...	...	-	1.5	2.8	2.0	2.8	2.9	2.6	0.8	3.1
1953-----	1.3	...	...	0.2	2.0	2.1	1.6	1.8	1.9	2.2	0.8	-
1952-----	1.3	...	...	0.1	1.3	2.7	2.5	1.6	2.2	0.3	0.9	3.4
1951-----	1.7	...	...	-	2.4	2.6	2.7	2.3	2.9	2.3	1.8	-
1950 <sup>1</sup> -----	1.7	...	...	0.1	1.7	2.8	2.2	4.0	1.2	2.5	2.9	-

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 13.—Death Rates for Homicide, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E960-E978 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E964, E980-E985 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	7.7	4.3	1.6	0.7	11.0	15.9	13.4	8.7	6.5	4.7	2.9	2.9
1968-----	7.3	4.8	1.4	0.7	10.1	15.3	12.9	9.0	6.3	4.5	3.3	3.7
1967-----	6.8	6.4	1.1	0.7	9.1	14.6	12.1	8.2	5.6	4.2	3.2	3.8
1966-----	5.9	5.8	1.2	0.6	7.6	12.3	10.6	7.7	5.4	3.4	2.8	3.1
1965-----	5.5	5.5	1.1	0.6	6.8	11.9	9.9	6.9	5.0	3.4	2.7	2.4
1964-----	5.1	5.4	1.2	0.6	6.3	10.5	9.2	6.7	4.6	2.9	2.5	3.1
1963-----	4.9	5.0	1.1	0.6	5.8	10.2	8.8	6.2	4.5	2.9	2.8	2.4
1962-----	4.8	5.1	0.9	0.6	5.9	9.9	8.6	6.3	4.5	3.1	2.6	3.4
1961-----	4.7	4.6	1.0	0.5	5.7	9.6	8.3	6.3	3.9	2.9	2.9	2.8
1960-----	4.7	4.8	0.7	0.5	5.9	9.7	8.1	6.2	4.2	2.8	2.4	2.4
1959-----	4.6	3.7	0.9	0.5	5.9	9.3	8.0	5.9	4.0	2.9	2.5	2.6
1958-----	4.5	3.5	0.7	0.5	5.8	8.7	7.4	6.3	4.0	2.9	2.6	3.3
1957-----	4.5	3.1	0.6	0.4	5.8	9.0	7.5	5.8	4.0	3.0	2.2	1.4
1956-----	4.6	3.3	0.7	0.4	6.1	9.0	7.6	5.6	4.1	2.8	2.2	2.3
1955-----	4.5	3.4	0.5	0.4	5.5	8.8	7.7	5.9	4.0	2.4	2.5	3.0
1954-----	4.8	3.7	0.6	0.5	6.0	9.1	8.2	6.5	3.5	2.8	2.7	1.9
1953-----	4.8	3.1	0.7	0.5	5.9	9.0	8.7	6.1	3.1	2.8	2.3	3.6
1952-----	5.2	3.7	0.6	0.5	6.2	10.0	9.1	6.1	3.8	3.0	2.7	2.0
1951-----	4.9	4.3	0.6	0.4	5.9	9.1	8.2	5.9	3.9	2.7	2.4	2.1
1950 <sup>1</sup> -----	5.3	4.4	0.6	0.5	6.3	9.9	8.8	6.1	4.0	3.2	2.6	2.3
<b>WHITE MALE</b>												
1969-----	6.0	3.5	1.4	0.5	7.6	11.6	10.1	7.1	6.9	5.2	4.1	3.5
1968-----	5.9	3.9	1.0	0.5	7.5	11.0	9.8	8.0	6.4	5.6	4.3	4.8
1967-----	5.3	5.6	0.9	0.5	6.3	10.4	8.8	6.9	5.7	5.0	3.8	4.3
1966-----	4.5	5.2	0.8	0.5	5.0	8.2	7.6	6.5	5.4	3.7	2.8	4.7
1965-----	4.4	5.3	1.0	0.5	4.9	7.8	7.3	6.0	5.2	3.5	3.5	3.0
1964-----	3.9	4.6	1.1	0.6	4.4	7.0	6.1	5.7	4.7	2.9	3.0	4.4
1963 <sup>2</sup> -----	3.9	4.2	1.1	0.5	4.2	7.3	6.4	5.1	4.4	3.4	3.1	2.9
1962 <sup>2</sup> -----	3.8	4.3	0.7	0.5	3.9	6.9	6.0	5.7	4.6	3.4	3.1	2.9
1961-----	3.6	3.9	0.8	0.5	3.9	6.4	5.8	5.4	4.1	3.4	2.9	2.6
1960-----	3.6	3.8	0.6	0.4	4.4	6.2	5.5	5.0	4.3	3.4	2.7	2.7
1959-----	3.5	3.2	0.8	0.5	4.2	5.5	5.4	5.1	4.3	3.4	3.0	2.1
1958-----	3.4	2.5	0.5	0.5	4.0	5.5	4.8	5.1	4.1	3.5	2.6	4.4
1957-----	3.2	2.1	0.3	0.4	3.9	5.6	4.8	4.3	4.0	3.4	2.5	1.3
1956-----	3.3	3.5	0.6	0.5	4.0	5.4	4.8	4.3	4.3	3.3	2.7	2.3
1955-----	3.4	3.3	0.4	0.4	3.5	5.2	5.7	4.6	4.1	3.0	3.4	3.4
1954-----	3.5	3.0	0.5	0.4	3.9	5.4	5.7	5.1	3.6	3.1	3.5	2.5
1953-----	3.5	2.7	0.6	0.4	3.8	5.4	5.7	5.2	3.7	3.1	3.3	3.4
1952-----	3.7	3.7	0.7	0.5	3.9	5.6	5.8	5.3	4.0	3.9	3.6	3.6
1951-----	3.6	3.6	0.4	0.4	3.6	5.5	5.6	4.9	4.3	3.5	3.0	2.1
1950-----	3.9	4.3	0.4	0.4	3.7	5.4	6.4	5.5	4.4	4.1	3.5	1.8
<b>WHITE FEMALE</b>												
1969-----	2.0	3.6	1.3	0.5	2.4	3.2	2.6	2.5	1.8	1.8	1.6	2.4
1968-----	1.9	4.2	1.1	0.5	2.2	2.8	2.9	2.2	1.8	1.5	1.9	2.6
1967-----	1.9	4.7	0.9	0.5	2.2	2.8	3.2	2.3	1.5	1.5	2.3	3.2
1966-----	1.8	4.7	0.9	0.4	2.0	3.0	2.4	2.5	1.9	1.6	1.8	2.1
1965-----	1.6	3.6	0.8	0.4	1.8	2.5	2.3	2.0	1.6	1.3	1.3	1.3
1964-----	1.6	4.0	0.9	0.4	1.6	2.7	2.4	1.9	1.5	1.5	1.4	1.9
1963 <sup>2</sup> -----	1.5	4.1	0.9	0.4	1.4	2.3	2.2	1.9	1.4	1.3	1.7	1.8
1962 <sup>2</sup> -----	1.6	4.2	0.9	0.5	1.7	2.2	2.5	1.8	1.4	1.5	1.6	3.3
1961-----	1.5	3.3	0.8	0.4	1.5	2.4	2.5	2.1	1.2	1.1	2.0	1.8
1960-----	1.4	3.5	0.5	0.3	1.5	2.0	2.2	1.9	1.5	1.1	1.2	1.5
1959-----	1.4	2.9	0.7	0.4	1.4	2.4	2.2	1.8	1.1	1.1	1.3	1.6
1958-----	1.4	2.8	0.7	0.3	1.6	1.7	2.2	1.9	1.2	1.2	2.1	2.1
1957-----	1.3	2.2	0.5	0.3	1.4	1.9	2.2	1.7	1.5	0.9	1.1	1.3
1956-----	1.3	2.6	0.6	0.2	1.4	1.9	1.8	1.8	1.2	1.0	1.3	0.9
1955-----	1.2	2.5	0.4	0.3	1.3	1.7	1.9	1.9	1.1	0.7	1.4	1.2
1954-----	1.4	3.0	0.6	0.4	1.5	1.9	2.1	2.0	1.0	1.0	1.1	1.0
1953-----	1.4	3.1	0.6	0.3	1.2	2.0	2.1	1.7	1.1	1.2	0.8	3.0
1952-----	1.3	3.0	0.5	0.3	1.4	1.8	2.3	1.7	1.1	0.9	0.8	1.1
1951-----	1.4	4.2	0.5	0.4	1.3	1.9	2.0	1.7	1.1	1.0	1.3	1.5
1950-----	1.4	3.9	0.6	0.4	1.3	1.9	2.2	1.6	1.3	1.1	1.2	1.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 13.—Death Rates for Homicide, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers E960-E978 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers E964,E980-E985 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	58.1	11.5	3.7	2.7	88.0	138.4	122.9	83.5	50.1	39.9	9.3	8.2
1968-----	54.6	7.8	3.5	2.9	77.1	137.3	114.6	81.8	52.7	34.3	15.9	10.9
1967-----	49.6	12.4	2.0	2.4	68.6	126.5	107.5	71.7	45.2	32.1	13.6	8.0
1966-----	43.5	11.7	2.7	1.8	61.1	104.2	95.4	67.8	40.7	27.5	15.4	6.5
1965-----	40.1	11.2	2.4	2.2	51.8	104.1	86.5	58.7	36.7	30.7	16.4	12.2
1964-----	37.4	9.8	1.9	1.3	49.6	89.6	86.9	56.1	34.6	21.5	14.7	8.1
1963 <sup>2</sup> -----	35.7	9.3	1.8	1.3	45.2	87.7	81.1	54.7	37.5	17.9	12.6	12.1
1962 <sup>2</sup> -----	35.5	10.5	1.7	1.2	48.2	85.2	76.3	56.1	36.0	21.0	11.8	6.5
1961-----	33.6	9.3	2.1	1.5	44.2	81.3	71.0	51.3	29.9	21.8	15.3	22.6
1960-----	34.5	10.1	1.6	1.4	43.7	84.7	72.3	51.5	29.1	18.6	16.7	12.8
1959-----	35.0	9.7	1.5	1.3	45.3	84.2	73.4	51.3	30.8	18.4	15.3	17.2
1958-----	34.9	8.3	1.5	0.9	44.2	83.4	68.8	54.7	31.8	18.1	10.8	14.8
1957-----	36.5	4.9	1.7	1.6	45.8	86.7	69.6	56.3	31.5	23.7	14.8	7.7
1956-----	37.1	6.6	0.9	1.7	48.8	88.0	73.7	49.0	30.5	18.3	10.8	24.0
1955-----	36.9	6.1	1.1	1.4	45.3	89.1	67.8	53.0	35.2	16.0	8.8	16.7
1954-----	40.6	10.1	1.4	1.6	49.8	94.8	76.2	57.6	33.2	19.9	15.1	4.5
1953-----	41.3	5.6	1.1	1.5	53.7	90.6	85.6	54.2	27.0	19.2	12.3	-
1952-----	45.4	5.4	1.3	1.2	55.8	106.1	90.1	53.1	32.7	19.1	18.2	-
1951-----	41.3	5.9	1.5	0.8	50.8	91.5	79.8	51.6	33.9	15.0	11.3	15.8
1950 <sup>1</sup> -----	45.5	5.9	0.9	1.8	56.5	105.6	80.4	52.8	32.3	20.9	11.9	10.7
ALL OTHER FEMALE												
1969-----	11.7	4.5	2.7	1.0	15.8	27.4	23.8	13.9	9.8	6.0	3.3	2.5
1968-----	11.6	8.0	2.9	1.3	16.1	26.2	23.6	13.3	8.5	5.9	4.2	2.8
1967-----	11.9	12.3	2.5	1.5	16.0	27.1	23.0	15.2	9.2	6.8	1.2	4.5
1966-----	10.6	7.6	2.9	1.3	12.6	24.7	22.1	13.3	7.7	3.9	7.5	1.7
1965-----	10.0	10.4	2.4	1.0	11.6	24.0	21.3	12.5	7.0	3.3	3.5	3.7
1964-----	9.2	12.4	2.4	1.4	11.1	20.5	19.4	10.5	6.8	4.2	0.9	4.1
1963 <sup>2</sup> -----	9.1	8.5	1.9	1.3	10.9	20.4	19.6	10.5	5.5	6.1	3.5	-
1962 <sup>2</sup> -----	8.9	8.4	1.3	0.8	11.0	21.2	20.0	8.5	5.9	3.3	4.2	7.1
1961-----	8.9	10.0	2.3	0.8	10.6	20.4	19.1	9.5	5.6	2.6	4.3	2.4
1960-----	9.9	13.0	1.7	1.0	11.3	22.9	19.6	12.3	6.8	3.1	3.0	2.5
1959-----	9.4	4.6	1.2	0.7	12.4	22.9	19.2	9.2	6.0	3.9	2.5	7.7
1958-----	9.3	6.6	1.1	0.8	12.5	20.5	17.7	12.4	5.3	3.8	3.3	-
1957-----	9.2	6.3	1.5	0.5	12.0	21.3	17.4	11.0	4.9	4.1	2.1	-
1956-----	10.3	8.4	1.6	0.6	13.0	23.1	19.0	11.1	7.6	4.9	1.4	2.8
1955-----	9.5	6.8	1.2	0.6	12.3	21.5	17.2	10.2	6.0	2.5	3.8	11.8
1954-----	9.5	5.6	0.9	0.7	12.6	20.8	16.4	12.5	2.5	4.7	7.9	6.3
1953-----	9.6	3.8	1.0	0.8	12.2	21.6	17.2	9.7	5.6	3.5	2.5	16.1
1952-----	10.8	7.1	0.6	0.7	13.2	25.2	19.4	10.4	4.9	3.4	4.3	-
1951-----	10.7	7.2	1.2	0.7	15.3	24.3	16.6	10.4	4.5	2.6	4.5	-
1950 <sup>1</sup> -----	11.2	6.4	1.5	1.2	15.8	25.8	17.7	8.7	3.2	3.9	3.9	3.8

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 14.—Death Rates for Nephritis and Nephrosis, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 580-584 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 590-594 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	4.7	1.1	0.3	0.4	1.0	1.6	3.0	4.8	9.6	16.5	32.5	69.7
1968-----	4.7	0.8	0.2	0.4	1.0	1.5	3.4	5.2	9.3	16.4	31.3	74.6
1967-----	5.5	1.2	0.4	0.5	1.4	2.1	3.8	6.5	10.6	19.2	36.2	93.8
1966-----	5.9	1.4	0.4	0.5	1.6	2.6	4.0	6.5	11.0	20.5	40.7	99.6
1965-----	6.2	1.3	0.6	0.6	1.6	2.6	3.9	7.2	11.5	20.9	43.0	108.3
1964-----	6.5	1.4	0.6	0.6	1.8	2.8	4.2	7.2	12.3	21.5	47.4	118.2
1963 <sup>2</sup> -----	6.7	0.9	0.6	0.7	1.8	3.0	4.2	7.7	12.7	23.3	48.0	118.9
1962 <sup>2</sup> -----	7.0	2.1	0.7	0.8	1.9	3.1	4.6	7.4	12.5	23.4	52.8	127.9
1961-----	7.2	1.4	0.7	0.8	2.1	3.2	4.5	7.7	13.1	24.2	55.1	135.4
1960-----	7.6	1.6	0.7	0.8	2.1	3.3	4.8	8.4	13.8	26.7	58.0	141.8
1959-----	8.0	2.0	0.6	0.9	1.9	3.3	5.1	8.6	14.6	27.2	64.6	151.9
1958-----	9.2	2.0	0.9	0.9	2.3	3.5	5.5	9.8	16.3	33.2	76.8	186.2
1957-----	10.2	1.8	1.1	1.1	2.7	4.1	6.3	10.2	18.8	36.6	82.0	198.7
1956-----	10.6	2.1	1.0	1.1	2.4	3.9	6.2	10.7	19.1	38.8	91.7	215.1
1955-----	11.1	1.9	1.3	1.1	2.7	3.9	6.3	11.3	20.2	41.1	99.5	217.2
1954-----	12.2	2.5	1.5	1.2	2.8	4.4	6.5	12.5	22.6	45.3	111.7	235.9
1953-----	13.6	2.2	1.7	1.4	3.2	4.8	7.5	14.2	25.0	51.4	124.2	279.9
1952-----	15.3	2.7	1.9	1.7	3.2	4.9	8.5	16.5	29.0	57.9	145.7	309.0
1951-----	16.8	2.7	2.2	1.8	3.4	5.0	8.8	16.8	32.0	65.5	164.4	383.4
1950-----	18.7	2.7	2.5	1.7	3.5	5.5	9.9	18.8	34.9	77.0	189.3	442.2
<b>WHITE MALE</b>												
1969-----	4.5	1.2	0.3	0.3	0.9	1.5	2.3	4.2	9.5	16.6	39.5	80.4
1968-----	4.6	0.9	0.2	0.3	0.9	1.4	2.7	4.8	9.2	18.1	36.9	87.7
1967-----	5.5	1.3	0.3	0.4	1.4	1.9	3.3	6.0	11.1	20.5	44.0	111.4
1966-----	5.9	1.1	0.4	0.5	1.7	2.4	3.6	6.1	10.8	22.9	46.5	122.3
1965-----	6.1	1.3	0.5	0.5	1.8	2.6	3.5	7.2	11.3	22.3	49.5	128.6
1964-----	6.5	1.4	0.5	0.5	1.9	2.6	3.9	6.9	12.2	22.7	55.2	139.3
1963 <sup>2</sup> -----	6.7	0.8	0.4	0.6	1.7	3.2	3.9	7.3	13.1	23.4	55.5	145.3
1962 <sup>2</sup> -----	7.1	1.8	0.7	0.6	2.2	3.3	4.4	7.1	12.5	25.6	59.7	156.3
1961-----	7.2	1.1	0.7	0.6	2.4	3.6	4.5	7.7	12.7	25.2	62.6	146.7
1960-----	7.6	1.7	0.5	0.8	2.5	3.7	4.5	7.9	13.4	26.4	64.1	165.3
1959-----	7.9	2.3	0.6	0.7	2.1	3.6	4.8	8.5	14.4	26.5	71.4	167.8
1958-----	9.1	1.9	0.8	0.9	2.5	3.5	5.4	8.9	15.3	33.9	85.2	205.4
1957-----	9.9	1.6	1.0	1.1	3.0	4.3	5.7	9.9	17.6	38.1	85.5	210.3
1956-----	10.2	1.7	1.0	1.0	2.6	4.2	5.8	9.6	17.9	37.6	97.7	240.1
1955-----	11.0	2.1	1.4	1.1	3.2	4.1	5.9	10.5	18.7	41.6	109.9	235.1
1954-----	11.9	2.9	1.6	1.2	3.1	4.3	6.0	11.2	21.3	45.2	120.3	232.8
1953-----	13.4	2.1	1.8	1.5	3.7	4.5	6.8	13.0	23.2	50.7	135.5	306.7
1952-----	15.0	3.0	2.1	1.8	3.5	4.8	7.3	15.1	26.6	57.8	156.1	326.2
1951-----	16.1	2.3	2.1	1.7	3.6	4.8	7.8	15.3	29.9	64.3	173.3	385.4
1950-----	18.1	2.4	2.5	1.8	4.0	5.3	8.8	17.1	32.5	73.3	200.2	484.5
<b>WHITE FEMALE</b>												
1969-----	3.6	0.7	0.3	0.4	0.7	0.9	1.7	2.6	5.4	11.5	24.5	62.3
1968-----	3.4	0.6	0.3	0.3	0.6	0.9	1.8	2.7	5.9	10.1	23.1	62.3
1967-----	4.0	0.8	0.4	0.4	1.0	1.0	2.0	3.6	5.8	13.2	26.5	76.7
1966-----	4.4	1.4	0.3	0.5	1.1	1.7	2.1	3.7	6.7	13.1	31.6	78.4
1965-----	4.6	1.3	0.6	0.6	1.3	1.4	2.4	4.0	7.2	13.9	32.6	87.7
1964-----	4.9	1.1	0.5	0.5	1.3	1.8	2.5	4.1	7.4	15.0	36.3	97.6
1963 <sup>2</sup> -----	5.1	0.9	0.4	0.7	1.4	1.7	2.5	4.4	7.9	16.1	36.6	94.7
1962 <sup>2</sup> -----	5.2	1.8	0.4	0.7	1.4	2.1	2.7	4.1	7.7	15.6	42.3	97.0
1961-----	5.6	1.3	0.4	0.7	1.4	2.0	2.6	4.4	8.7	17.5	42.6	115.9
1960-----	5.9	1.2	0.6	0.7	1.4	2.0	2.9	4.9	8.9	19.8	45.1	114.1
1959-----	6.2	1.2	0.4	0.9	1.4	2.1	2.9	4.7	9.7	19.7	51.1	128.3
1958-----	7.3	1.9	0.6	0.8	1.7	2.4	3.0	6.1	10.4	24.6	61.4	157.3
1957-----	8.1	1.5	0.8	1.0	1.9	2.4	3.8	6.1	13.3	26.2	68.8	172.8
1956-----	8.5	1.9	0.8	1.0	1.8	2.2	3.7	6.5	13.5	30.0	76.4	180.5
1955-----	8.8	1.3	0.8	1.1	2.0	2.4	3.7	6.9	14.3	31.6	80.4	187.8
1954-----	9.9	1.5	1.3	1.1	2.1	3.1	4.2	8.3	16.3	34.4	92.8	201.0
1953-----	10.9	1.7	1.2	1.3	2.3	3.4	4.7	9.0	18.1	39.8	99.3	235.0
1952-----	12.0	2.1	1.4	1.5	2.3	3.3	5.6	9.9	20.0	44.0	117.1	255.3
1951-----	13.6	2.3	1.8	1.7	2.7	3.3	5.6	10.4	23.0	49.6	137.2	336.0
1950-----	15.2	1.4	2.1	1.5	2.5	3.8	6.1	12.0	24.8	60.3	156.9	388.9

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 14.—Death Rates for Nephritis and Nephrosis, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 580-584 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 590-594 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	10.3	1.3	0.4	0.5	2.8	5.4	13.0	20.9	32.5	54.1	73.0	83.6
1968-----	10.3	0.7	0.2	0.6	3.0	4.7	13.0	21.2	28.6	57.3	80.6	101.8
1967-----	11.9	1.6	0.9	0.8	2.7	6.0	13.5	23.7	37.3	60.4	91.5	158.0
1966-----	13.0	3.6	0.8	0.9	3.2	7.7	13.6	24.9	41.9	64.7	94.9	178.3
1965-----	13.2	2.2	1.5	0.9	2.8	8.0	13.8	26.7	35.4	66.9	106.9	219.5
1964-----	13.4	2.7	1.1	1.2	3.0	7.6	13.1	25.8	41.3	68.0	107.6	183.8
1963 <sup>2</sup> -----	14.6	1.9	1.6	1.3	3.9	7.6	13.9	27.9	38.7	90.3	113.2	239.4
1962 <sup>2</sup> -----	14.4	3.1	1.7	1.5	2.6	6.7	15.0	25.7	38.8	76.0	108.3	312.9
1961-----	14.7	3.3	1.8	1.2	3.7	7.3	13.5	24.7	40.7	78.4	121.5	261.3
1960-----	16.2	4.3	1.9	1.7	3.4	6.9	16.4	26.3	44.1	83.9	146.1	264.7
1959-----	17.0	3.7	2.1	1.2	2.8	7.7	16.7	29.9	44.4	91.4	142.4	524.1
1958-----	19.1	3.8	2.3	1.7	3.8	6.9	16.5	32.8	56.5	100.0	158.3	400.0
1957-----	20.3	3.1	2.6	1.3	4.3	11.0	19.0	29.8	59.7	100.8	183.7	432.0
1956-----	21.7	3.7	2.7	2.2	4.3	9.0	17.2	34.9	62.2	115.1	195.4	478.3
1955-----	21.4	5.0	3.2	1.3	3.7	8.5	19.3	35.2	61.4	111.0	212.8	350.0
1954-----	24.8	4.4	2.7	1.6	5.1	10.3	18.5	38.2	68.5	155.8	268.9	555.0
1953-----	27.3	3.8	3.9	1.5	3.7	11.1	20.9	42.7	80.2	148.3	293.9	600.0
1952-----	32.8	5.4	3.7	2.1	5.1	11.0	24.8	54.3	101.7	186.9	374.5	741.2
1951-----	34.3	6.8	4.6	2.5	4.5	11.4	25.6	52.3	101.2	190.8	382.1	1,033.3
1950 <sup>1</sup> -----	37.8	9.4	4.8	1.8	5.2	12.1	27.3	56.7	112.3	241.1	435.1	782.1
ALL OTHER FEMALE												
1969-----	8.6	2.3	0.7	0.7	1.8	3.6	9.4	14.4	30.3	44.0	42.6	67.5
1968-----	9.0	1.7	0.5	0.7	1.8	4.8	11.6	16.0	26.8	42.3	47.5	98.6
1967-----	10.3	1.7	0.5	0.6	2.3	6.6	12.3	20.5	31.2	46.3	47.0	112.1
1966-----	10.2	0.3	0.3	0.6	1.9	5.5	13.0	18.8	28.7	49.4	61.1	115.0
1965-----	10.6	0.9	0.9	1.0	1.9	7.1	10.2	18.0	34.0	52.8	65.2	118.5
1964-----	11.5	1.2	0.9	0.8	2.9	7.3	11.4	21.0	36.0	50.1	67.4	161.2
1963 <sup>2</sup> -----	11.6	1.3	1.8	1.0	2.9	6.4	12.2	20.7	33.1	56.8	70.4	143.2
1962 <sup>2</sup> -----	12.0	4.4	0.9	0.9	2.2	6.3	12.4	22.1	35.6	51.7	87.8	169.0
1961-----	12.0	0.9	1.2	1.3	2.9	6.4	12.3	20.9	36.0	45.0	91.3	209.8
1960-----	13.3	1.0	1.4	0.9	2.6	6.4	11.9	27.0	38.6	58.5	97.3	216.0
1959-----	14.7	3.0	1.0	1.9	3.2	6.5	15.0	25.3	41.3	67.0	116.1	197.4
1958-----	16.8	1.7	2.3	0.9	2.6	9.0	17.2	30.2	51.7	70.6	133.3	235.1
1957-----	19.1	3.5	2.0	1.8	4.0	9.1	20.4	34.1	52.1	88.7	146.9	262.9
1956-----	19.6	3.6	1.2	1.1	3.3	10.0	21.0	37.3	55.4	88.1	157.2	257.6
1955-----	20.3	1.5	1.6	1.4	3.9	9.4	19.2	37.7	66.5	86.3	159.8	353.3
1954-----	20.6	4.0	1.6	1.3	3.9	10.1	19.1	39.5	67.4	92.7	150.0	332.1
1953-----	25.4	4.3	2.9	1.8	6.7	12.0	25.2	46.8	76.2	110.3	208.3	448.0
1952-----	29.5	1.3	2.2	1.9	6.2	12.9	28.5	54.2	100.2	125.5	242.2	634.8
1951-----	32.4	4.5	3.9	2.5	6.9	14.8	30.1	56.9	105.3	152.3	268.5	681.0
1950 <sup>1</sup> -----	34.9	6.9	2.3	2.1	4.9	15.2	35.9	61.4	106.8	197.0	326.0	486.3

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 15.—Death Rates for Peptic Ulcer, by Age, Color, and Sex: United States, 1950-1969

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 531-533 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 540,541 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
<b>BOTH SEXES</b>												
1969-----	4.6	1.1	0.1	0.0	0.1	0.5	1.8	4.2	10.3	20.7	40.2	65.7
1968-----	4.7	1.6	0.1	0.0	0.1	0.5	1.7	4.5	10.9	22.2	38.5	69.1
1967-----	5.0	1.6	0.1	0.0	0.2	0.5	1.8	4.9	11.5	22.3	41.5	80.2
1966-----	5.3	1.5	0.0	0.0	0.2	0.5	1.9	5.3	11.9	24.7	45.5	79.3
1965-----	5.4	1.9	0.1	0.0	0.2	0.5	1.9	5.6	12.7	24.7	46.3	81.8
1964-----	5.7	1.6	0.1	0.0	0.1	0.6	2.2	5.8	13.6	26.6	50.0	85.6
1963-----	6.4	1.7	0.1	0.0	0.2	0.8	2.5	7.0	14.8	30.2	55.9	93.0
1962 <sup>2</sup> -----	6.6	1.6	0.1	0.1	0.2	0.7	2.6	7.0	15.4	31.1	56.8	94.5
1961-----	6.5	1.3	0.1	0.0	0.2	0.8	2.4	7.4	15.1	29.7	53.3	82.6
1960-----	6.3	1.9	0.1	0.0	0.1	0.7	2.5	7.3	15.7	29.9	53.6	77.9
1959-----	6.0	1.6	0.1	0.0	0.2	0.7	2.7	7.1	14.9	28.2	52.1	66.8
1958-----	6.2	1.9	0.1	0.0	0.2	0.9	2.8	7.5	16.1	28.7	51.7	70.7
1957-----	6.1	1.7	0.1	0.0	0.1	0.7	2.8	7.6	16.0	28.6	47.4	72.0
1956-----	6.0	1.3	0.1	0.0	0.2	0.8	2.6	7.5	15.6	28.4	48.5	68.5
1955-----	5.9	1.5	0.1	0.0	0.2	0.8	2.9	7.8	16.0	28.2	44.9	55.8
1954-----	5.8	1.9	0.1	0.0	0.2	0.8	3.2	8.1	15.9	27.3	41.4	53.9
1953-----	5.8	2.0	0.1	0.0	0.2	0.9	3.1	8.4	16.6	26.6	41.9	50.1
1952-----	5.6	1.4	0.1	0.0	0.2	0.8	3.3	8.4	16.5	25.9	37.3	44.6
1951-----	5.5	1.2	0.1	0.0	0.2	1.0	3.3	8.4	16.5	25.6	35.3	43.2
1950 <sup>1</sup> -----	5.5	1.6	0.0	0.1	0.3	0.9	3.4	8.6	16.7	24.8	34.6	38.3
<b>WHITE MALE</b>												
1969-----	6.5	0.9	0.1	0.0	0.2	0.7	2.2	6.3	15.9	32.6	62.1	91.6
1968-----	6.8	1.3	0.1	0.0	0.1	0.6	2.0	6.4	16.8	35.7	61.8	98.6
1967-----	7.2	1.5	0.1	0.0	0.2	0.6	2.3	6.8	18.7	37.1	64.6	118.4
1966-----	7.7	1.3	0.0	0.0	0.2	0.6	2.5	7.5	18.6	40.2	74.1	116.1
1965-----	8.0	1.3	0.1	0.0	0.2	0.6	2.5	8.1	20.6	41.6	73.7	125.9
1964-----	8.6	1.2	0.1	0.1	0.2	0.7	2.9	8.3	22.3	44.6	78.8	134.9
1963 <sup>2</sup> -----	9.8	1.5	0.2	0.0	0.2	1.0	3.3	10.7	24.0	51.0	90.3	142.7
1962 <sup>2</sup> -----	10.0	1.6	0.1	0.1	0.2	0.9	3.6	10.3	24.7	52.1	91.2	142.2
1961-----	9.7	1.1	0.1	0.0	0.2	1.1	3.5	11.0	25.0	49.8	85.8	128.1
1960-----	10.0	1.8	0.1	0.0	0.2	1.0	3.6	11.3	26.8	50.4	88.5	120.9
1959-----	9.5	1.6	0.1	0.0	0.2	1.0	3.8	11.1	25.6	47.1	85.7	102.1
1958-----	9.9	1.4	0.1	0.0	0.2	1.3	3.9	11.8	27.3	48.5	84.4	107.0
1957-----	9.7	1.2	0.1	0.0	0.2	0.9	4.3	12.3	26.7	48.8	76.5	104.8
1956-----	9.8	1.2	0.1	0.0	0.2	1.2	4.1	12.4	26.7	49.2	78.3	106.3
1955-----	9.7	1.7	0.1	0.0	0.3	1.1	4.5	12.9	27.6	47.9	73.7	80.3
1954-----	9.7	1.7	0.0	0.1	0.3	1.2	4.9	13.6	27.4	47.1	69.4	83.0
1953-----	9.8	1.5	0.1	0.0	0.2	1.3	5.0	14.0	28.3	46.6	70.2	81.0
1952-----	9.5	1.5	0.1	0.0	0.2	1.3	5.2	14.2	28.7	45.2	60.7	69.6
1951-----	9.3	1.4	0.0	0.0	0.3	1.5	5.5	14.1	28.0	44.0	59.1	67.5
1950-----	9.1	1.9	0.1	0.1	0.3	1.4	5.4	14.1	28.6	42.1	57.3	61.4
<b>WHITE FEMALE</b>												
1969-----	3.1	0.3	0.1	0.0	0.1	0.3	1.0	1.9	4.9	11.0	26.7	59.0
1968-----	3.1	0.9	0.1	0.0	0.1	0.2	0.8	2.3	5.1	11.2	24.8	58.1
1967-----	3.1	0.9	0.0	0.0	0.1	0.2	0.9	2.6	4.8	10.9	26.9	64.9
1966-----	3.3	0.7	0.0	0.0	0.1	0.3	0.9	2.5	5.4	12.1	27.8	64.2
1965-----	3.2	1.0	0.1	0.0	0.1	0.3	1.0	2.7	5.2	11.7	28.6	63.7
1964-----	3.3	0.6	0.0	0.0	0.1	0.3	1.1	2.8	5.3	12.3	31.4	62.9
1963 <sup>2</sup> -----	3.7	0.8	0.1	0.0	0.1	0.3	1.3	3.0	5.7	13.7	34.3	71.7
1962 <sup>2</sup> -----	3.8	0.7	0.1	0.0	0.1	0.3	1.2	3.3	6.1	14.3	34.8	72.5
1961-----	3.4	0.7	0.0	0.1	0.1	0.3	1.1	3.3	5.7	13.3	31.6	58.9
1960-----	3.3	0.6	0.1	0.1	0.1	0.3	1.0	3.0	5.5	13.1	30.4	57.3
1959-----	3.1	0.7	0.1	0.1	0.1	0.3	1.1	2.8	5.2	12.9	30.9	49.6
1958-----	3.1	1.2	0.1	0.0	0.1	0.3	1.2	3.0	5.5	12.3	28.5	53.6
1957-----	3.0	1.5	0.1	0.0	0.1	0.4	1.0	2.8	5.7	12.3	27.0	55.1
1956-----	2.8	0.6	0.1	0.0	0.1	0.3	0.9	2.6	5.0	11.4	27.7	49.5
1955-----	2.7	0.6	0.1	0.0	0.1	0.3	1.1	2.4	5.1	11.8	24.0	45.9
1954-----	2.4	1.2	0.1	0.0	0.1	0.3	1.1	2.4	4.7	10.9	21.6	38.2
1953-----	2.3	1.7	0.1	0.0	0.1	0.3	1.0	2.4	4.6	9.8	21.6	31.5
1952-----	2.2	0.5	0.1	0.0	0.1	0.3	1.1	2.3	4.8	10.0	20.0	29.5
1951-----	2.1	0.8	0.1	-	0.1	0.3	0.9	2.2	5.0	9.8	19.0	28.3
1950-----	2.1	0.8	0.0	0.0	0.1	0.2	1.1	2.6	4.9	9.6	16.9	26.8

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 15.—Death Rates for Peptic Ulcer, by Age, Color, and Sex: United States, 1950-1969—Con.

[Rates per 100,000 population in specified age-color-sex group. For 1968 and 1969 rates are based on deaths assigned to category numbers 531-533 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category numbers 540,541 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
ALL OTHER MALE												
1969-----	4.9	5.4	0.1	0.1	0.2	1.3	4.6	8.9	18.3	31.9	49.5	16.4
1968-----	5.5	4.6	0.2	0.1	0.4	2.4	6.3	10.7	19.9	38.8	30.8	29.1
1967-----	5.3	2.9	0.2	0.1	0.4	1.5	5.7	12.1	18.8	30.0	44.2	48.0
1966-----	6.0	4.2	0.2	0.1	0.3	1.4	6.1	12.1	22.7	39.4	43.6	47.8
1965-----	5.6	6.5	-	0.1	0.6	1.6	5.0	12.8	19.5	32.8	44.4	31.7
1964-----	6.2	5.4	-	-	0.5	1.5	5.6	13.2	24.1	38.6	45.1	40.5
1963 <sup>2</sup> -----	6.7	5.9	0.2	0.1	0.5	2.3	6.1	14.0	25.1	41.0	45.4	33.3
1962 <sup>2</sup> -----	7.0	4.6	0.2	0.1	0.4	3.2	6.9	13.9	26.1	43.2	39.1	54.8
1961-----	6.7	4.8	0.1	-	0.5	3.3	5.7	14.8	21.9	39.3	44.8	54.8
1960-----	6.9	7.8	0.2	0.1	0.1	2.2	6.6	15.0	23.8	38.7	40.7	44.6
1959-----	6.5	7.0	0.3	0.1	0.3	1.7	7.1	14.0	23.6	35.6	39.6	37.9
1958-----	6.5	5.2	-	0.0	0.6	1.9	6.6	13.0	24.1	34.6	46.8	22.2
1957-----	6.6	4.9	-	0.0	0.2	2.3	6.2	14.6	26.3	30.9	41.5	46.2
1956-----	5.9	5.1	-	0.1	0.3	2.3	5.5	12.6	24.7	27.0	36.9	24.0
1955-----	6.3	5.0	0.1	-	0.6	2.3	6.9	14.2	22.9	28.7	41.6	25.0
1954-----	6.5	5.2	0.1	-	0.7	2.3	7.7	14.6	26.0	27.6	29.4	31.8
1953-----	6.9	6.4	-	0.1	0.5	2.9	6.0	17.2	28.3	29.1	31.6	35.3
1952-----	6.4	7.7	0.1	-	0.9	2.2	7.2	15.2	22.9	26.0	33.6	25.0
1951-----	6.4	2.7	0.2	-	0.8	2.7	7.3	16.6	24.9	29.1	7.5	47.4
1950 <sup>1</sup> -----	7.0	4.4	0.1	0.1	0.8	3.1	8.4	16.0	27.8	26.8	35.7	16.1
ALL OTHER FEMALE												
1969-----	2.1	1.6	0.1	-	0.0	0.7	2.4	2.8	5.8	14.1	17.3	25.0
1968-----	2.2	3.3	-	-	-	0.5	2.6	2.9	7.0	11.3	14.3	37.5
1967-----	2.2	3.7	0.2	0.1	0.3	0.7	2.2	3.0	6.8	9.8	18.7	30.3
1966-----	2.3	4.0	-	0.1	0.2	0.4	2.1	5.5	6.3	13.0	12.1	25.0
1965-----	2.2	5.1	0.1	0.0	-	0.6	1.9	4.0	7.3	10.6	18.1	22.2
1964-----	2.6	4.5	-	-	0.1	1.1	3.0	5.7	6.2	11.8	19.1	30.6
1963 <sup>2</sup> -----	2.5	2.5	0.1	0.1	0.2	0.9	1.9	4.9	8.9	14.5	13.1	29.5
1962 <sup>2</sup> -----	2.3	1.9	0.1	-	0.2	0.4	2.4	3.6	9.7	11.8	15.3	28.6
1961-----	2.5	2.1	0.2	-	0.2	0.7	1.7	5.2	9.2	12.1	17.9	39.0
1960-----	2.2	3.6	0.2	0.0	0.1	0.8	2.3	4.2	6.6	11.0	14.3	19.9
1959-----	2.0	1.7	-	-	0.2	0.6	2.7	4.5	4.5	9.2	16.1	15.4
1958-----	2.7	5.2	-	0.1	0.3	1.5	3.2	4.7	7.5	10.1	21.6	21.1
1957-----	2.0	3.1	-	-	0.3	0.7	1.8	3.7	7.3	8.2	15.2	21.6
1956-----	2.2	2.6	-	-	0.1	1.1	1.9	5.0	7.2	10.9	16.7	11.1
1955-----	2.2	3.0	0.1	-	0.4	0.9	1.7	4.6	8.1	10.7	15.2	11.8
1954-----	2.2	3.6	0.1	-	0.3	0.9	2.5	5.1	7.3	9.4	11.1	6.3
1953-----	2.0	2.6	0.1	-	0.2	0.4	1.8	5.7	7.5	8.4	9.1	16.1
1952-----	1.7	0.9	0.1	-	0.1	0.9	2.4	4.2	5.5	4.5	11.2	24.1
1951-----	2.1	1.4	-	-	0.3	0.8	2.0	5.4	9.2	9.0	8.1	10.7
1950 <sup>1</sup> -----	1.9	1.5	-	-	0.6	0.4	2.8	5.4	3.6	10.2	10.7	-

<sup>1</sup>Based on enumerated population adjusted for age bias in the population of races other than white.

<sup>2</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

Table 16.—Death Rates for Malignant Neoplasms of Breast, for the Female Population, by Age and Color: United States, 1950-1969

[Rates per 100,000 population in specified age-color group. For 1968 and 1969 rates are based on deaths assigned to category number 174 of the Eighth Revision of the International Classification of Diseases, Adapted for Use in the United States; and for 1950-67, are based on deaths assigned to category number 170 of the Sixth and Seventh Revisions adopted, respectively, in 1948 and 1955]

Year, color, and sex	Total	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
WHITE FEMALE												
1969-----	29.1	-	-	0.0	0.1	3.1	19.7	51.5	79.5	96.2	120.7	192.8
1968-----	29.3	0.1	0.0	0.0	0.1	3.6	20.6	52.6	78.6	97.9	124.5	194.1
1967-----	28.8	-	0.0	-	0.1	3.5	20.2	52.3	78.6	94.8	123.6	198.8
1966-----	28.6	-	-	-	0.1	3.4	20.2	52.3	78.7	94.8	124.3	192.1
1965-----	28.4	-	-	0.0	0.1	3.5	20.2	51.9	76.6	95.0	128.5	192.8
1964-----	27.8	-	-	-	0.2	4.1	20.0	52.9	73.2	93.5	121.9	200.2
1963 <sup>1</sup> -----	26.9	0.1	-	-	0.1	3.3	19.0	50.6	72.5	93.3	120.1	190.4
1962 <sup>1</sup> -----	27.0	-	-	0.0	0.1	3.9	18.7	51.1	71.1	93.0	127.1	190.8
1961-----	27.3	0.1	-	0.0	0.1	3.9	19.5	50.9	70.4	93.0	133.1	201.6
1960-----	27.2	0.1	-	0.0	0.1	3.6	19.7	51.2	71.8	91.6	132.8	199.7
1959-----	26.6	0.1	0.0	0.0	0.1	4.1	19.3	48.8	71.5	91.1	129.3	197.8
1958-----	26.7	0.1	-	0.0	0.1	3.6	20.3	49.4	69.9	92.7	132.9	188.9
1957-----	27.1	0.1	0.0	-	0.1	4.0	19.3	50.1	71.3	95.1	139.1	208.8
1956-----	27.0	0.3	0.0	0.0	0.1	3.7	20.7	48.6	71.7	95.9	139.8	205.5
1955-----	27.2	0.1	0.0	0.0	0.2	3.6	21.0	50.0	71.8	94.9	144.8	208.9
1954-----	26.3	0.1	-	0.0	0.1	3.7	19.3	48.4	70.0	95.4	140.2	202.8
1953-----	26.4	-	-	0.0	0.1	3.6	20.5	48.5	70.2	96.4	144.1	205.7
1952-----	26.0	0.1	0.1	0.0	0.1	3.9	20.4	47.6	72.4	93.8	136.6	193.8
1951-----	25.5	-	-	0.0	0.1	3.5	19.4	45.6	69.9	95.7	148.0	203.9
1950-----	25.7	-	-	0.0	0.2	3.7	20.8	47.1	70.9	96.3	143.6	204.2
ALL OTHER FEMALE												
1969-----	19.4	-	-	-	0.3	5.3	25.7	52.6	71.7	97.4	77.9	71.3
1968-----	19.8	-	-	-	0.1	5.8	23.9	55.0	70.9	98.4	85.8	113.9
1967-----	19.4	-	-	-	0.2	5.7	21.6	55.7	76.2	88.3	79.3	101.5
1966-----	18.1	-	-	-	0.3	6.0	21.4	50.9	71.0	80.6	74.5	91.7
1965-----	18.3	-	-	-	0.2	6.1	24.2	50.8	69.2	75.4	81.1	127.8
1964-----	19.1	-	-	-	0.2	5.7	21.6	52.2	68.9	76.4	80.9	112.2
1963 <sup>1</sup> -----	18.2	0.3	-	0.1	0.1	5.6	25.1	50.0	68.9	74.0	80.4	118.2
1962 <sup>1</sup> -----	17.5	0.6	-	0.0	0.2	5.3	24.1	47.1	62.3	72.7	86.2	123.8
1961-----	17.8	-	-	-	0.2	4.6	26.0	47.7	64.1	69.4	91.3	119.5
1960-----	17.9	0.6	-	-	0.3	5.9	24.1	53.0	61.1	70.0	87.1	89.4
1959-----	17.0	-	-	0.0	0.1	5.5	21.7	45.8	66.9	65.7	75.8	107.7
1958-----	16.8	0.7	-	-	0.3	5.3	23.6	43.7	64.9	61.9	79.1	105.3
1957-----	17.2	-	-	0.0	0.1	4.6	22.3	49.5	63.1	66.7	89.7	89.2
1956-----	17.7	-	-	-	0.2	4.1	25.3	50.9	65.1	65.4	82.6	122.2
1955-----	17.1	-	0.1	-	0.1	5.5	24.5	45.8	60.7	67.9	89.4	67.6
1954-----	17.2	-	-	-	0.4	6.7	24.9	46.1	62.0	65.6	77.0	84.4
1953-----	16.3	-	-	-	0.4	6.3	21.6	46.7	55.0	60.2	84.3	116.1
1952-----	16.2	-	-	0.1	0.6	5.0	23.5	41.9	64.0	58.8	91.4	75.9
1951-----	16.1	-	-	-	0.2	4.5	21.8	47.1	62.0	59.5	74.8	110.7
1950 <sup>2</sup> -----	16.0	-	-	-	0.1	4.7	20.8	45.3	58.1	76.4	78.1	91.9

<sup>1</sup>Figures by color exclude data for residents of New Jersey because this State did not require reporting the item.

<sup>2</sup>Based on enumerated population adjusted for age bias in the population of races other than white.



Table 17.—Death Rates for Malignant Neoplasms of Female Genital Organs, by Site, for White Women, by Birth Cohort and Age at Death: United States, 1949, 1954, 1959, 1964, and 1969

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers]

Cause of death and age	Period of birth of cohort																	
	1950-1954	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
Malignant neoplasm of genital organs (ICD No. 180-184) <sup>1</sup>	0.6	0.5	0.7	0.6	0.6													
15-19-----																		
20-24-----		0.9	1.1	1.1	1.2	1.3												
25-29-----			1.3	2.3	2.4	3.0	3.1											
30-34-----				4.2	5.9	6.4	8.4	8.7										
35-39-----					9.1	10.5	13.3	14.6	18.3									
40-44-----						17.8	20.3	22.2	26.5	30.8								
45-49-----							27.0	31.0	34.6	40.1	48.4							
50-54-----								40.4	44.9	49.5	57.2	62.1						
55-59-----									51.3	55.8	61.9	69.7	80.9					
60-64-----										61.9	69.9	78.7	83.8	87.6				
65-69-----											78.5	87.9	92.6	94.6	101.9			
70-74-----												89.7	100.7	106.0	109.9	120.9		
75-79-----													101.3	105.0	114.0	127.8	128.8	
80-84-----														107.9	120.2	126.0	136.5	141.9
Malignant neoplasm of cervix uteri (ICD No. 180) <sup>2</sup>	0.0	0.0	0.0	0.0	0.0	0.3												
15-19-----																		
20-24-----		0.2			0.4	1.6	1.9											
25-29-----			1.1	1.2	1.1	4.1	4.1	5.5										
30-34-----					4.9	6.1	8.1	8.2	9.8									
35-39-----						8.0	10.9	11.4	13.7	14.9								
40-44-----							10.5	12.9	15.2	16.1	21.5							
45-49-----								12.2	16.6	18.4	23.0	23.1						
50-54-----									14.5	17.8	19.9	23.9	25.5					
55-59-----										15.6	19.6	22.5	25.2	26.6				
60-64-----											16.9	22.2	25.2	27.4	28.3			
65-69-----												19.9	23.1	27.0	30.0	32.5		
70-74-----													21.0	26.5	30.4	37.5	31.2	
75-79-----														20.9	30.3	34.5	33.9	38.7
80-84-----																		
Malignant neoplasm of other parts of uterus (ICD No. 181,182) <sup>2</sup>	0.1	0.1	0.1	0.1	0.1	0.5	0.5											
15-19-----																		
20-24-----		0.2	0.2	0.2	0.2	0.6	0.6	1.4										
25-29-----			0.2	0.4	0.5	0.6	0.8	1.4	3.9									
30-34-----				0.3	0.5	0.7	0.8	1.4	3.9	7.5								
35-39-----					0.8	1.0	1.6	2.3	4.4	7.5								
40-44-----						2.0	3.3	3.0	4.4									
45-49-----							4.4	4.0	5.1	7.1	12.3							
50-54-----								5.8	7.7	9.3	13.1	18.9						
55-59-----									9.5	12.1	14.6	19.9	28.2					
60-64-----										14.7	18.6	23.0	28.0	33.1				
65-69-----											23.0	26.6	29.9	32.8	41.4			
70-74-----												27.2	31.3	33.3	38.2	53.6		
75-79-----													32.8	35.3	38.9	45.8	57.8	
80-84-----														36.4	39.0	41.8	55.1	70.6

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 171-176 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>2</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 171 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>3</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 172-174 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

Table 17.—Death Rates for Malignant Neoplasms of Female Genital Organs, by Site, for White Women, by Birth Cohort and Age at Death: United States, 1940, 1954, 1959, 1964, and 1969—Con.

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers]

Cause of death and age	Period of birth of cohort																	
	1930-1934	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
Malignant neoplasm of ovary, fallopian tube, and broad ligament (ICDA No. 183) <sup>1/</sup>																		
15-19	0.4																	
20-24	0.4	0.4																
25-29		0.6	0.5															
30-34		0.6	0.7	0.8														
35-39			1.2	1.3	1.6													
40-44				3.3	3.5	3.9												
45-49					7.2	7.5	8.2	4.3										
50-54						13.5	15.6	15.8	14.2	13.8	15.8	18.7						
55-59							21.7	19.3	20.8	23.4	24.4	25.3						
60-64								26.0	25.8	26.3	24.4	25.3	25.5					
65-69									30.0	30.2	31.0	30.2	28.7					
70-74										36.0	35.9	35.3	31.3	28.7				
75-79											39.4	35.6	25.6	25.7	28.7			
80-84												41.8	37.5	35.7	35.0	31.2		
													40.9	40.0	39.2	38.3	22.4	
Malignant neoplasm of other and unspecified female genital organs (ICDA No. 184) <sup>2/</sup>																		
15-19	0.0																	
20-24		-	0.0	0.0	0.0													
25-29		-	0.0	-	-	0.0												
30-34			-	0.0	0.1	0.0	0.1	0.1										
35-39				0.0	0.1	0.1	0.2	0.2	0.3	0.5								
40-44					0.1	0.1	0.4	0.3	0.3	0.5								
45-49						0.3	0.4	0.5	0.5	0.7	0.3							
50-54							0.6	0.6	1.0	0.6	1.4							
55-59									1.2	1.0	1.1	1.5	1.3					
60-64										1.6	2.2	2.5	2.4	2.4				
65-69											2.7	3.2	3.0	3.4				
70-74												4.2	4.7	4.8	6.0	6.1		
75-79													5.7	6.6	8.9	9.5	8.8	
80-84														9.6	10.9	11.5	11.0	10.2

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1/</sup>For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 175 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.  
<sup>2/</sup>For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 176 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

Table 18.—Death Rates for Malignant Neoplasms of Female Genital Organs, by Site, for Women of Races Other Than White, by Birth Cohort and Age at Death: United States, 1949, 1954, 1959, 1964, and 1969

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers]

Cause of death and age	Period of birth of cohort																	
	1950-1954	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
Malignant neoplasm of genital organs (ICDA Nos. 180-184) <sup>1/</sup>	0.6																	
15-19-----		1.1	1.1	0.9	1.5													
20-24-----		1.3		2.0	1.8	3.0												
25-29-----			3.0	4.1	7.9	7.4	3.0											
30-34-----				11.3	11.9	12.9	16.2	25.5										
35-39-----					19.9	24.4	20.8	33.6	48.6									
40-44-----						31.2	28.5	47.8	58.7	74.3								
45-49-----							48.4	57.8	65.4	85.5	104.7							
50-54-----								61.8	85.2	97.3	112.8	119.8						
55-59-----									92.1	91.5	117.2	120.8	154.6					
60-64-----										112.2	126.6	124.0	154.6	144.9				
65-69-----											153.8	155.1	131.3	111.4	159.5			
70-74-----												111.4	131.3	117.7	133.1	154.0		
75-79-----													121.7	123.0	126.0	152.6	150.8	
80-84-----														108.2	116.3	147.2	116.0	117.9
Malignant neoplasm of cervix uteri (ICDA No. 180) <sup>2/</sup>																		
15-19-----		0.1	0.3	-	0.3													
20-24-----		0.3	0.6	0.7	1.5	1.4												
25-29-----			2.5	2.8	5.1	4.9	5.5											
30-34-----				7.1	8.6	9.1	11.4	16.1										
35-39-----					15.1	15.2	23.4	21.9	28.8									
40-44-----						21.3	26.0	31.2	35.1	42.2								
45-49-----							28.7	35.3	42.8	48.3	53.7							
50-54-----								32.0	42.2	46.7	59.6	55.9						
55-59-----									45.0	42.5	60.5	56.0	58.6					
60-64-----										45.3	61.0	48.9	67.5	55.6				
65-69-----											61.1	70.9	64.5	47.5	53.0			
70-74-----												44.3	52.0	49.7	55.9	62.8		
75-79-----													36.8	47.4	56.5	55.3	65.6	
80-84-----														41.8	50.0	55.6	52.0	38.5
Malignant neoplasm of other parts of uterus (ICDA Nos. 181,182) <sup>3/</sup>																		
15-19-----	0.2				0.3	0.3												
20-24-----		0.2	0.5	0.3	0.7	0.9												
25-29-----		0.3	0.5	0.4	1.0	1.8	2.3											
30-34-----			0.4	0.7	1.6	2.1	2.9	7.1										
35-39-----				1.9	3.7	4.2	7.8	14.4										
40-44-----					2.2	4.8	5.0	8.7	10.4	23.9								
45-49-----						4.8	6.3	7.8	12.1	22.4	36.2							
50-54-----								13.0	20.1	19.4	34.2	48.3						
55-59-----									22.6	23.3	31.3	42.5	56.8					
60-64-----										32.6	43.8	46.3	58.3	56.8				
65-69-----											50.2	49.2	43.9	44.1	69.3			
70-74-----												46.2	46.3	59.3	60.5			
75-79-----													53.3	44.4	77.6	62.3		
80-84-----														45.9	35.0	67.9	48.0	74.4

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 171-176 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>2/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 171 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>3/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 172-174 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

Table 18.—Death Rates for Malignant Neoplasms of Female Genital Organs, by Site, for Women of Races Other Than White, by Birth Cohort and Age at Death: United States, 1949, 1954, 1959, 1964, and 1969—Con.

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers]

Cause of death and age	Period of birth of cohort																	
	1950-1954	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
Malignant neoplasms of ovary, fallopian tube, and broad ligament (ICDA No. 153) <sup>1/2</sup>																		
15-19-----	0.5	0.7	0.4	0.6	0.9													
20-24-----		0.6	0.3	0.6	0.6													
25-29-----			1.0	0.6	1.3	0.9												
30-34-----				2.1	1.4	1.7	1.6	1.5										
35-39-----					2.4	4.5	2.5	3.5	3.9									
40-44-----						4.2	6.6	7.2	8.3	7.1								
45-49-----							12.0	13.5	12.7	13.0	12.8							
50-54-----								15.7	17.6	19.0	16.1	13.4						
55-59-----									24.3	23.1	22.6	21.4	16.3					
60-64-----										30.7	27.4	27.4	22.5	15.4				
65-69-----											32.2	29.6	17.4	15.4	15.5			
70-74-----												25.9	20.8	20.0	15.2	9.7		
75-79-----													20.7	16.3	20.4	17.1	13.7	
80-84-----														17.5	15.0	17.0	10.0	2.6
Malignant neoplasms of other and unspecified female genital organs (ICDA No. 154) <sup>1/2</sup>																		
15-19-----		0.1	-	-	-													
20-24-----		0.1	-	-	0.1	0.1												
25-29-----			-	-	0.4	0.1	0.3	0.8										
30-34-----			0.1	-	0.3	-	0.3	0.8	1.6									
35-39-----				0.1	0.1	1.0	0.7	0.5	1.9	1.2								
40-44-----					0.8		1.3	1.2	1.8	1.7	2.0							
45-49-----								1.1	2.3	2.9	3.9	2.2						
50-54-----									2.6	2.8	0.9	2.5						
55-59-----										3.6	3.4	2.7	2.8					
60-64-----											4.3	4.3	2.5	1.8				
65-69-----												2.7	1.7	2.8	0.9			
70-74-----												4.6	5.9	3.7	2.6	3.3		
75-79-----													3.1	6.3	5.7	6.0		
80-84-----																		2.6

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1/2</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 175 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

<sup>2/2</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 176 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

Table 19.—Death Rates for Malignant Neoplasms of Male Genital Organs, by Site, for White Men, by Birth Cohort and Age at Death: United States, 1949, 1954, 1959, 1964, and 1969

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers]

Cause of death and age	Period of birth of cohort																	
	1960-1954	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
<b>Malignant neoplasm of genital organs</b> (ICDA Nos. 185-187) <sup>1/</sup>	0.7	0.6	0.7	0.7	0.5													
15-19		1.7	1.3	1.2	1.8	1.6												
20-24			2.3	2.2	1.7	2.3	2.1											
25-29				1.7	1.8	1.8	2.0	2.0										
30-34					1.8	1.8	2.1	1.8	1.8									
35-39					1.8	1.9	2.1	1.8	1.6	1.8								
40-44						1.3	1.4	1.4	1.6	1.8								
45-49							1.9	2.0	1.9	2.4	2.6							
50-54								4.2	4.3	4.1	4.3	6.3						
55-59									11.0	12.0	12.8	13.9	14.1					
60-64										30.0	29.6	29.6	32.6	34.2				
65-69											66.9	67.5	73.2	73.2	71.0			
70-74												132.9	136.5	130.1	142.3	144.5		
75-79													231.2	228.5	226.9	251.8	259.3	
80-84														337.2	349.7	373.9	402.3	368.4
<b>Malignant neoplasm of prostate</b> (ICDA No. 185) <sup>2/</sup>	0.0	0.0	0.1	0.1	0.1	0.0												
15-19					0.1													
20-24					0.0													
25-29					0.0													
30-34					0.0													
35-39					0.1		0.1	0.0	0.1									
40-44							0.2	0.3	0.3	0.4								
45-49								0.8	0.9	0.9	1.3	1.4						
50-54									3.2	3.2	3.2	3.2	4.8					
55-59										10.3	10.9	11.5	12.5	12.8				
60-64											28.7	28.2	28.1	30.8	32.2			
65-69												64.9	65.1	65.9	70.6	68.1		
70-74													130.1	133.8	127.6	139.4	139.4	
75-79														228.0	225.1	221.8	256.9	253.4
80-84															333.4	344.3	369.5	360.0
<b>Malignant neoplasm of all other and unspecified male genital organs</b> (ICDA Nos. 186,187) <sup>3/</sup>	0.7	0.6	0.6	0.7	0.4													
15-19		1.7	1.3	1.2	1.8	1.5												
20-24			2.2	2.2	1.7	2.2	2.1											
25-29				1.6	1.8	1.8	2.0	2.0										
30-34					1.8	1.8	2.0	1.7	1.7									
35-39					1.8	1.8	2.0	1.7	1.3									
40-44						1.1	1.2	1.5	1.3	1.5								
45-49							1.1	1.0	1.0	1.0	1.2							
50-54								1.0	1.0	0.9	1.1	1.5						
55-59									0.8	1.1	1.3	1.4	1.3					
60-64										1.3	1.3	1.5	1.8	2.0				
65-69											1.7	1.8	2.0	2.6	2.9			
70-74												2.9	2.7	2.5	2.9	5.1		
75-79													3.2	3.5	5.1	4.8	5.8	
80-84														3.8	5.4	5.4	8.8	8.4

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 177-179 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

<sup>2/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 177 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

<sup>3/</sup> For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 178-179 of the Sixth and Seventh Revisions, adopted respectively, in 1949 and 1955.

Table 20.—Death Rates for Malignant Neoplasms of Male Genital Organs, by Site, for Men of Races Other Than White, by Birth Cohort and Age at Death: United States, 1949, 1954, 1959, 1964, and 1969

[Numbers after causes of death are category numbers of the Eighth Revision of the International Lists, 1965. For data year 1969, rates are based on deaths assigned to these Eighth Revision category numbers.]

Cause of death and age	Period of birth of cohort																	
	1950-1954	1945-1949	1940-1944	1935-1939	1930-1934	1925-1929	1920-1924	1915-1919	1910-1914	1905-1909	1900-1904	1895-1899	1890-1894	1885-1889	1880-1884	1875-1879	1870-1874	1865-1869
Rates per 100,000 population in age group																		
<b>Malignant neoplasm of genital organs</b> (ICM No. 153-157) <sup>1</sup>	0.1	0.3	0.3	0.5	0.3	0.2	0.5	1.1	2.0	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
15-19		0.5	0.1	0.3	-	0.2	0.5	1.1	2.0	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
20-24			0.7	0.2	0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
25-29					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
30-34					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
35-39					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
40-44					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
45-49					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
50-54					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
55-59					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
60-64					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
65-69					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
70-74					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
75-79					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
80-84					0.7	1.3	1.3	1.4	2.1	4.3	6.1	21.0	46.0	75.1	103.8	129.9	256.5	242.9
<b>Malignant neoplasm of prostate</b> (ICM No. 155) <sup>2</sup>	-	-	-	0.2	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
15-19				0.2	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
20-24		0.1		-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
25-29			0.1	0.2	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
30-34				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
35-39				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
40-44				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
45-49				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
50-54				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
55-59				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
60-64				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
65-69				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
70-74				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
75-79				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
80-84				-	-	-	-	0.2	0.5	2.6	4.6	19.8	46.4	70.2	105.3	124.6	230.2	231.4
<b>Malignant neoplasm of all other and unspecified male genital organs</b> (ICM Nos. 126,137) <sup>3</sup>	0.1	0.3	0.3	0.3	0.3	0.2	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
15-19		0.4	0.1	0.3	-	-	-	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
20-24			0.5	0.2	0.3	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
25-29				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
30-34				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
35-39				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
40-44				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
45-49				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
50-54				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
55-59				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
60-64				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
65-69				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
70-74				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
75-79				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3
80-84				0.2	0.7	0.5	0.5	0.9	1.4	1.6	1.4	2.2	2.7	2.2	2.9	3.4	4.4	6.3

NOTE: The first figure in each row, moving from right to left, is the death rate for data year 1949, the second figure for 1954, the third figure for 1959, the fourth figure for 1964, and the fifth figure for 1969.

<sup>1</sup>For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category numbers 177-179 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>2</sup>For data years 1949, 1954, 1959, 1964, death rates are based on deaths assigned to category number 177 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

<sup>3</sup>For data years 1949, 1959, 1964, death rates are based on deaths assigned to category numbers 178-179 of the Sixth and Seventh Revisions, adopted respectively, in 1948 and 1955.

## Explanatory Notes

These 20 tables present age-specific death rates by color and sex for 13 of the 15 leading causes and selected components of these causes for each year during 1950-69. The two causes not included are **Certain causes of mortality in early infancy**, which ranked as the sixth leading cause in 1969, and **Congenital anomalies**, which ranked as the twelfth leading cause in that year. Mortality trends for these causes, based on the number of live births for each year during 1950-69, are shown in a report of the series *Vital and Health Statistics* (National Center for Health Statistics: Mortality trends for leading causes of death, United States, 1950-69. *Vital and Health Statistics*. Series 20-No. 16. Health Resources Administration, DHEW, Rockville, Md.).

During 1950-69 causes of death were classified according to three revisions of the International Classification of Diseases. The Sixth Revision was in use during 1949-57; the Seventh Revision during 1958-67; and the Eighth Revision was introduced in 1968. These revisions are made in order that the disease classification may be consistent with advances in medical science and changes in diagnostic practice.

Each revision has produced some breaks in the comparability of cause-of-death statistics. For the breaks between the Sixth and Seventh Revision, see *Vital Statistics—Special Reports*, Vol. 51, No. 4, March 1965; for the breaks between the Seventh and Eighth Revision, see the above mentioned *Vital and Health Statistics* report.

SYMBOLS	
Category not applicable-----	...
Quantity zero-----	-
Quantity more than 0 but less than 0.05----	0.0