

Healthy People 2010 Operational Definition

3-4. Reduce the death rate from cancer of the uterine cervix.

National Data Source	National Vital Statistics System-Mortality (NVSS-M), CDC, NCHS.
State Data Source	National Vital Statistics System-Mortality (NVSS-M), CDC, NCHS.
Healthy People 2000 Objective	16.4 (Cancer).
Changes since the 2000 Publication	Revised baseline (see Comments). Revised target (see Comments).
Measure	Female rate per 100,000 standard population (age adjusted-see Comments).
Baseline (Year)	2.8 (1999)
Target	2.0
Target-Setting Method	Better than the best racial/ethnic subgroup. For a discussion of target-setting methods, see Part A, section 4.
Numerator	Number of female deaths due to cancer of the uterine cervix (ICD-10 code C53).
Denominator	Number of females.
Population Targeted	U.S. resident population.
Questions Used To Obtain the National Baseline Data	Not applicable.
Expected Periodicity	Annual.
Comments	The original baseline was revised from 3.0 deaths per 100,000 population to 2.8 per 100,000 due to changes in the International Classification of Diseases (ICD) system. The original baseline (1998) was based on ICD-9 code 180. The revised baseline (1999) is based on the ICD-10 system. The baseline year for all Healthy People 2010 mortality objectives was changed to 1999 so that a consistent trend based on a single ICD classification

scheme could be tracked throughout the decade. See Part A, section 8 for a discussion of the International Classification of Diseases (ICD). Information on comparability between ICD-9 and ICD-10 has been published by NCHS.¹

Although the original target of 2.0 deaths per 100,000 population was proportionally adjusted to reflect the revised baseline using the original target-setting method, the target value did not change.

Data (except those by education status) are age adjusted to the 2000 standard population using the age groups under 1 year, 1-4, 5-14, 15-24, 25-34, 35-44, 45-54, 55-64, 65-74, 75-84, 85 years and over. Data by education status are adjusted using the age groups 25-34, 35-44, 45-54, and 55-64. Age-adjusted rates are weighted sums of age-specific rates. For a discussion of age adjustment, see Part A, section 7.

This objective differs from Healthy People 2000 objective 16.4, which age adjusted the death rates using the 1940 standard population.

For some measures, data do not meet the criteria for statistical reliability, data quality, or confidentiality and have been suppressed. Information on suppression of data for the major Healthy People 2010 data systems has been published in a *Healthy People Statistical Note*.²

See Part C for a description of NVSS-M and Appendix A for focus area contact information.

References

1. Anderson, R.N.; Minino, A.M.; Hoyert, D.L.; Rosenberg, H.M.; Comparability of cause of death between ICD-9 and ICD-10: Preliminary estimates. *National Vital Statistics Reports*. Vol. 49 No.2. Hyattsville, MD: National Center for Health Statistics, 2001.
2. Klein, R.J.; Proctor, S.E.; Boudreault, M.A.; Turczyn, K.M. Healthy People 2010 Criteria for Data Suppression. *Statistical Notes* No. 24. Hyattsville, MD: National Center for Health Statistics. 2002.