Persons with disabilities experiencing problems accessing PDF file should contact nchsed@cdc.gov, or call 301-458-4688

HEALTH STATISTICS

FROM THE U.S. NATIONAL HEALTH SURVEY

Acute Conditions seasonal variations

United States July 1957 - June 1961

Statistics on incidence of acute conditions and number of associated days of disability in each calendar quarter according to condition group and age. Based on data collected in household interviews during the period July 1957-June 1961.

U. S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Abraham Ribicoff, Secretary

PUBLIC HEALTH SERVICE Luther L. Terry, Surgeon General

Washington, D. C.

NATIONAL CENTER FOR HEALTH STATISTICS

Forrest E. Linder, Ph.D., Director Theodore D. Woolsey, Assistant Director O. K. Sagen, Ph.D., Assistant Director

U. S. NATIONAL HEALTH SURVEY

Theodore D. Woolsey, Chief
Alice M. Waterhouse, M.D., Chief Medical Advisor
James E. Kelly, D.D.S., Dental Advisor
Walt R. Simmons, Statistical Advisor
Arthur J. Mc Dowell, Chief, Health Examination Survey
Philip S. Lawrence, Sc.D., Chief, Health Interview Survey
Robert T. Little, Chief, Automatic Data Processing

The U. S. National Health Survey is a continuing program under which the Public Health Service makes studies to determine the extent of illness and disability in the population of the United States and to gather related information. It is authorized by Public Law 652, 84th Congress.

CO-OPERATION OF THE BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Health Survey, the Bureau of the Census, under a contractual arrangement, participates in most aspects of survey planning, selects the sample, collects the data, and carries out certain parts of the statistical processing.

CONTENTS

| · · · · · · · · · · · · · · · · · · · | Page |
|--|------|
| Introduction | 1 |
| Source of Data | 1 |
| Incidence of Acute Conditions and Associated Days of Disability by Quarter | 2 |
| Detailed Tables | 8 |
| Appendix I. Technical Notes on Methods | 46 |
| Background of This ReportStatistical Design of the Health Interview | 46 |
| SurveyGeneral Qualifications | 46 |
| General Qualifications | 47 |
| Reliability of EstimatesGuide to Use of Relative Standard Error | 47 |
| Charts | 49 |
| Appendix II. Definitions of Certain Terms | |
| Used in This Report | 52 |
| Terms Relating to Conditions | 52 |
| Terms Relating to Disability | 53 |
| Demographic Terms | 53 |
| Appendix III. Questionnaire | 54 |

ACUTE CONDITIONS

SEASONAL VARIATIONS

INTRODUCTION

During the 16 calendar quarters from July 1957-June 1961 an estimated average of 95 million acute conditions per quarter involving either activity restriction or medical attention were experienced by members of the civilian, noninstitutional population of the United States. This estimate is based on data collected in household interviews conducted by the U. S. National Health Survey, National Center for Health Statistics. The largest number of acute conditions occurring in any quarter was 160 million conditions in October-December 1957, the peak of the epidemic of Asian infleunza. The smallest estimated number of acute conditions was the 60 million cases which occurred in July-September 1959.

The average of 95 million acute conditions per quarter resulted in restriction of activity for the average person of 2.3 days per quarter. The largest number of restricted-activity days per quarter was experienced in the October-December 1957 quarter with 4.5 days per person and the lowest number occurred in July-September 1959 with 1.4 days per person.

This report presents an analysis of the incidence of and associated disability from acute conditions for 16 quarters of data collection to show seasonal variations. A companion report, Acute Conditions, Geographic Distribution, B-34, presents annual estimates of incidence of acute conditions for each of the four years of data collection to show trends in the geographic distribution of acute conditions and to present detailed information for the year, July 1960-June 1961. Similar data, reported from the National Health Survey, Series B, No. 24, presented an analysis of seasonal variations for the 12 quarters of reporting of acute conditions from July 1957-June

1960. Other reports from the National Health Survey containing information about acute conditions are Series B, Nos. 5, 6, 18, and 23, and Series C Nos. 1 and 4.

SOURCE OF DATA

The information contained in this publication is derived from household interviews conducted by the National Health Survey in co-operation with the Bureau of the Census of a probability sample of the civilian, noninstitutional population of the United States. The sample is designed so that interviews are conducted during every week of the year. During the 16-quarter period from July 1957-June 1961 the sample was composed of approximately 149,000 households containing about 485,000 persons living at the time of the interview.

A description of the design of the survey, the methods used in estimation, and general qualifications of the data obtained from surveys is presented in Appendix 1. Since the estimates shown in this report are based on a sample of the population rather than on the entire population, they are subject to sampling error. Therefore, particular attention should be paid to the section entitled "Reliability of Estimates." Since the estimates based on a three-month-interviewing period are subject to higher sampling errors than estimates based on a 12-month period, the quarterly estimates should be interpreted with caution, especially for those estimates of relatively low magnitude.

Certain terms used in this report are defined in Appendix II. Some of the terms have specialized meanings for the purpose of the survey. For example, an acute condition is defined, with certain exceptions, as a condition which has lasted less than three months and which has involved either medical attention or restricted activity. The exceptions, which are listed in Appendix II, are certain conditions such as heart trouble and diabetes which are always considered to be chronic.

This report was prepared by Charles S. Wilder of the U. S. National Realth Survey staff.

The questionnaire used by the Health Interview Survey during July 1960-June 1961 is reproduced in Appendix III. The questions (II-I7 and Table I) designed to elicit information about acute conditions are substantially the same as those used in the earlier versions of the questionnaire.

Each condition reported in response to the illness-recall questions (11-17) was entered on a line of Table I. Further questions were asked to obtain a better description of the specific condition and to obtain information about the actions taken by the person, such as seeking medical attention or restricting his usual activities.

INCIDENCE OF ACUTE CONDITIONS AND ASSOCIATED DAYS OF DISABILITY BY QUARTER

During the 16 quarters in the period from July 1957-June 1961 there was a marked seasonal variation in the incidence rates of acute conditions per 100 persons (tables 1 and 2). As shown in figure 1, the seasonal pattern had peaks in the January-March quarter of three of the four years; seasonal lows were reached in the July-September quarter of each year. In the first year of reporting the seasonal high was in the October-December 1957 quarter as a result of the epidemic of Asian influenza. Figure 2 shows that other respiratory conditions. including influenza. reached the high point in October-December 1957, while upper respiratory conditions reached a peak during the January-March 1958 quarter in conforming with the usual pattern. During July 1960-June 1961 the incidence of acute conditions showed the least amount of seasonal variation of any of the four years.

The influence of the incidence of acute respiratory conditions on the rates for all acute conditions is shown in figure 1. Since about three of each five acute conditions are respiratory conditions, it is not surprising that the pattern of the seasonal distribution of all acute conditions is quite similar to the pattern for acute respiratory conditions (fig. 1).

There is less seasonal variation in the incidence of injuries and of infective and parasitic diseases than for all acute conditions as evidenced by repetition of cyclical movement. However, the rise and fall of these curves is well within sampling variability and may be due to

chance alone. The high and low points for the incidence of infective and parasitic diseases correspond to the pattern for all acute conditions in the last three of the four years of data. The low rates for infective and parasitic conditions during July-December 1957 may have been due to the influence of the Asian influenza epidemic on the diagnostic information obtained in interviews. Illnesses which would have ordinarily been reported as "virus infection," "the virus," or other ill-defined conditions classifiable to infective and parasitic diseases may have been reported during the epidemic period as Asian influenza. The classification of such conditions as respiratory would result in the unusually low rate for infective and parasitic conditions.

The curve for injuries exhibits a seasonal pattern that is the reverse of that for all conditions—high when the curve for all conditions is low, and vice versa. A small seasonal pattern of variation is shown for digestive system conditions; however, it is quite irregular since it disappears during the four quarters of 1958 and part of 1959.

Tables 3 and 4 show the age distribution of the incidence of acute conditions during each of the 16 quarters. The effect of age on seasonal variation as shown in figure 3 is such that for the youngest age group there is less relative rise and fall in the age-specific incidence rates than is present for the older age groups. The age group, 5-14, has the largest relative difference between high and low points in each cycle. Age-specific incidence rates for the acute respiratory and injury condition groups are shown in tables 17 and 20, respectively.

The distribution of days of restricted activity and bed disability per 100 persons exhibits about the same type of seasonal pattern as for the incidence rates (tables 5-12). In 14 of the 16 quarters respiratory conditions contributed the largest proportion of days to the total number of restricted-activity days. The exceptions were July-September of 1958 and 1960 during which injuries had the largest number of restricted-activity days (table 5). The proportionate contribution to the total by the other condition groups varied from quarter to quarter.

In terms of the number of restricted-activity days per condition, a rate which may be considered as a measure of severity among the persons exposed to the risk of incurring such disability, the most interesting curve in figures 4 and 5 is that for injuries. The curve for a particular condition group was obtained by dividing the number of restricted-activity days for the group by the number of cases in the condition group. In figure 1 the incidence curve for injuries

¹The rates shown in all of the figures depicting seasonal trends are plotted on semilogarithmic scale so that visual comparisons can be made of relative amounts of seasonal variation within and between individual curves of condition groups or age groups.




Figure 1. Incidence of acute conditions per 100 persons per quarter.

had seasonal high points in the July-September quarter while the curve for restricted-activity days per condition had high points in the January-March quarter. Tables 20-22 have been included to explain this seeming discrepancy. Comparing these two quarters in each of the four years (table 20) shows that in July-September an average of 37.4 percent of the injuries were sustained by young people under 15 and 22.9 percent by persons 45 years of age and over. During January-

March an average of 33.4 percent of the injuries were sustained by young people under 15 and 25.2 percent by older persons 45 years and over. Since older persons have a longer recuperation time per injury (fig. 6), this explains the apparent discrepancy in seasonal peaks.

The gradual rise in the duration of restricted activity per injury during the 16 quarters, as shown in figures 5 and 6, should not, at present be taken to indicate that the severity of injuries

is changing. The period under study is too short to judge the significance of this rise until further data have been accumulated.

An indication of the effect of acute conditions on work-loss days is given in figure 7 and tables 13 and 14. Figure 7 shows the average number of persons absent from work each day because of acute conditions. More males than females were absent from work each day, but since the working population is heavily weighted with males, such an effect is expected. Respiratory conditions and

injuries were the leading causes of work loss, but injuries exerted a relatively greater influence as a cause of work loss than they did as a cause of restricted activity among all persons. The proportion of work-loss days resulting from injuries is 26 percent of the total number, while only 17 percent of all restricted-activity days may be attributed to injuries.

Seasonal variations in the average number of persons with work loss indicate no sharp rise during the winters of 1958-59 and 1960-61. The




Figure 2. Incidence of acute upper and other respiratory conditions per 100 persons per quarter.




Figure 3. Incidence of acute conditions per 100 persons per quarter by age.




Figure 4. Number of restricted-activity days per condition per quarter by condition group.




Figure 5. Number of restricted activity days per condition per quorter by condition group.




Figure 6. Number of restricted-activity days per injury per quarter by age.




Figure 7. Fersons absent from work each day because of acute conditions by sex.

epidemics of Asian influenza during the winters of 1957-58 and 1959-60 account for the high rate of work loss during these years.

13

School-loss days associated with acute conditions per 100 children aged 6-16 years of age are shown in tables 15 and 16. The seasonal peaks coincide with those shown in figure 1 for the incidence of acute conditions. Respiratory

conditions were the primary cause of days lost from school resulting from acute conditions.

Since acute respiratory conditions and injuries represent such a large proportion of the total number of acute conditions, separate tables showing incidence, restricted-activity days, and bed-disability days by age are shown in tables 17-22.

DETAILED TABLES

| | | | Page |
|------|-----|---|------|
| | | ACUTE CONDITIONS | |
| able | 1. | Incidence of acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | . 10 |
| | 2. | Incidence of acute conditions per 100 persons per quarter by sex and condition group: United States, July 1957-June 1961 | 12 |
| | | Incidence of acute conditions per quarter by sex and age: United States, July 1957-June 1961 | 14 |
| | 4. | Incidence of acute conditions per 100 persons per quarter by sex and age: United States, July 1957-June 1961 | 16 |
| | | DAYS OF RESTRICTED ACTIVITY ASSOCIATED WITH ACUTE CONDITIONS | |
| | 5. | Days of restricted activity associated with acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | 18 |
| | 6. | Days of restricted activity associated with acute conditions per 100 persons per quarter by sex and condition group: United States, July 1957-June 1961 | 20 |
| | 7. | Days of restricted activity associated with acute conditions per quarter by sex and age: United States, July 1957-June 1961 | 22 |
| | 8. | Days of restricted activity associated with acute conditions per 100 persons per quarter by sex and age: United States, July 1957-June 1961 | 24 |
| | | DAYS OF BED DISABILITY ASSOCIATED WITH ACUTE CONDITIONS | |
| | 9. | Days of bed disability associated with acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | 26 |
| | 10. | Days of bed disability associated with acute conditions per 100 persons per quarter by sex and condition group: United States, July 1957-June 1961 | 28 |
| | 11. | Days of bed disability associated with acute conditions per quarter by sex and age: United States, July 1957-June 1961 | 30 |
| | 12. | Days of bed disability associated with acute conditions per 100 persons per quarter by sex and age: United States, July 1957-June 1961 | 32 |
| | | DAYS OF TIME LOST ASSOCIATED WITH ACUTE CONDITIONS | |
| | 13. | Days lost from work associated with acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | 34 |
| | 14. | Persons absent from work each day because of acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | 34 |
| | 15. | Days lost from school associated with acute conditions per quarter by sex and condition group: United States, July 1957-June 1961 | 36 |
| | 16. | Days lost from school associated with acute conditions per 100 children per quarter by sex and condition group: United States, July 1957-June 1961 | 36 |

DETAILED TABLES—Continued

| | | | Page |
|--|--------------|--|---------|
| ACUTE RESPIRATORY CONDITIONS Table 17. Incidence of acute respiratory conditions per quarter by age: United States, July 1957-June 1961 | | | |
| Table | 17. | | 38 |
| | 18. | | 38 |
| | 19., | | 40 |
| | , | INJURIES | , |
| | 20. | Incidence of injuries per quarter by age: United States, July 1957-June 1961- | 40 |
| | 21. | | - 42 |
| | ,2 2. | Days of bed disability associated with injuries per quarter by age: United States, July 1957-June 1961 | 42 |
| | | POPULATION | |
| | 23. | Population used in obtaining rates shown in this publication by quarter, sex, | 44 |

Table 1. Incidence of acute conditions per quarter by sex Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| 1 | | | · · · · · · · · · · · · · · · · · · · | | • 0 | | |
|--|----------------------------|-----------------------------|---------------------------------------|----------------------------|---------------------------|----------------------------|--|
| | 195 | 57 | 1958 | | | | |
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December | |
| Both sexes | | | | Ir | ncidence of | acute con- | |
| All acute conditions | 69,704 | 160,312 | 125,437 | 82,433 | 63,196 | 93,202 | |
| Infective and parasitic diseases | 7,946 | 6,817 | 10,034 | 13,815 | 7,131 | 9,178 | |
| Respiratory conditions Upper respiratory Other respiratory | 30,612 20,123 10,489 | 126,340 51,445 74,895 | 88,727 53,524 35,203 | 38,854 28,506 10,348 | 25,458 17,745 7,713 | 59,246 42,031 17,215 | |
| Digestive system conditions | 6,915 | 4,662 | 5,884 | 6,565 | 5,748 | 5,284 | |
| Injuries | 13,910 | 11,123 | 10,192 | 12,383 | - 15,480 | 11,063 | |
| All other acute conditions | 10,322 | 11,370 | 10,601 | 10,816 | 9,380 | 8,431 | |
| Male | | | | | - | | |
| All acute conditions | 31,587 | 74,119 | 59,707 | 37,279 | 29,566 | 42,875 | |
| Infective and parasitic diseases | 3,322 | 3,109 | 5,422 | 6,019 | 2,979 | 4,579 | |
| Respiratory conditions Upper respiratory Other respiratory | 12,778 8,360 4,418 | 58,866 23,268 35,598 | 41,514 24,262 17,252 | 16,249 11,867 4,382 | 11,886 8,198 3,688 | 26,473 18,827 7,646 | |
| Digestive system conditions | 2,554 | 2,552 | 2,763 | 2,943 | 2,475 | 2,556 | |
| Injuries | 8,527 | 5,607 | 5,840 | 7,808 | 8,725 | 6,330 | |
| All other acute conditions | 4,406 | 3,985 | 4,168 | 4,260 | 3,502 | 2,937 | |
| Female | | .) . | | | | | |
| All acute conditions | 38,117 | 86,193 | 65,730 | 45,154 | 33,630 | 50,327 | |
| Infective and parasitic diseases | 4,624 | 3,708 | 4,612 | 7,796 | 4,152 | 4,599 | |
| Respiratory conditions Upper respiratory Other respiratory | 17,834 11,763 6,071 | 67,474 28,177 39,297 | 47,212 29,261 17,951 | 22,605 16,639 5,966 | 13,572 9,547 4,025 | 23,203 | |
| Digestive system conditions | 4,361 | 2,110 | 3,121 | 3,623 | 3,273 | 2,728 | |
| Injuries | 5,383 | 5,516 | 4,351 | 4,575 | 6,755 | 4,732 | |
| All other acute conditions | 5,916 | 7,385 | 6,434 | 6,556 | 5,877 | 5,495 | |
| | | <u> </u> | | | · | | |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

and condition group: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | | | 1961 | | |
|-------------------|------------------|--------------------|----------------------|-------------------|------------------|--------------------|----------------------|-------------------|------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| litions in | thousand | 8 | | | | | | | |
| 118,305 | 93,238 | 59,742 | 92,178 | 130,572 | 72,659 | 65,719 | 100,308 | 112,359 | 80,954 |
| 14,530 | 13,401 | 7,227 | 10,263 | 13,854 | 11,359 | 7,552 | 10,108 | 16,435 | 14,962 |
| 78,101 | 52,543 | 22,137 | - 57,216 | 93,656 | 34,896 | 24,866 | 63,576 | 71,075 | 36,759 |
| 52,532 25,568 | 30,073 22,469 | 16,258 5,879 | 39,733 17,483 | 45,911 47,745 | 23,992 10,904 | 17,219 7,647 | 44,612 18,964 | 50,406 20,670 | 26,499 10,260 |
| 5,044 | 4,286 | 5,385 | 5,083 | 3,942 | ·4,180 | 6,217 | 4,797 | 5,346 | 6,279 |
| 9,986 | 12,941 | 13,751 | 9,992 | 9,927 | 12,017 | 15,434 | ` 12,078` | 10,066 | 12,129 |
| 10,645 | 10,067 | 11,243 | 9,623 | 9,192 | 10,207 | 11,649 | 9,748 | 9,437 | 10,824 |
| | | · | | | | | , | | |
| 53,455 | 44,684 | 27,627 | 40,662 | 59,779 | 33,496 | 31,524 | 47,195 | . 52,609 | 36,211 |
| 7,788 | 7,062 | 3,258 | 5,272 | 6,537 | 4,444 | 3,831 | 4,492 | 8,080 | 6,296 |
| 33,844 | 23,594 | 10,533 | 24,729 | 42,824 | 16,898 | 10,834 | 29,137 | 32,437 | 16,096 |
| 22,674 11,169 | 13,317 10,277 | 7,783 2,750 | 17,722 7,007 | 20,804 22,020 | 11,577 5,321 | 7,596 3,238 | 20,628 8,509 | 23,305 9,132 | 11,410 4,686 |
| 2,087 | 1,887 | 2,523 | 1,942 | 1,989 | 1,823 | 3,173 | 2,364 | 2,507 | 2,819 |
| 5,760 | 8,102 | 7,363 | 5,440 | . 5,260 | 6,917 | 9,056 | 7,477 | 5,605 | 7,210 |
| 3,976 | 4,039 | 3,949 | 3,279 | 3,169 | 3,413 | 4,630 | 3,724 | 3,980 | 3,790 |
| • | • | | • | | | | | | |
| 64,851 | 48,554 | 32,115 | 51,516 | 70,793 | 39,163 | 34,194 | 53,113 | 59,750 | 44,742 |
| 6,742 | 6,339 | 3,968 | 4,991 | 7,318 | 6,914 | 3,721 | 5,616 | 8,355 | 8,665 |
| 44,257 | 28,949 | 11,604 | 32,487 | 50,833 | 17,998 | 14,032 | 34,439 | 38,638 | 20,663 |
| 29,858 14,399 | 16,756 12,192 | 8,474 3,130 | 22,011 10,476 | 25,107 25,726 | 12,415 5,583 | 9,623 4,408 | 23,984 10,455 | 27,100 11,538 | 15,088 5,575 |
| 2,957 | 2,399 | 2,861 | 3,142 | 1,954 | 2,356 | 3,044 | 2,432 | 2,839 | 3,460 |
| 4,225 | 4,839 | 6,388 | 4,552 | 4,667 | 5,100 | 6,378 | 4,601 | 4,461 | 4,919 |
| 6,669 | 6,029 | 7,294 | 6,345 | 6,022 | 6,794 | 7,020 | 6,024 | 5,457 | 7,034 |

Table 2. Incidence of acute conditions per 100 persons per quarter [Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | 1958 | | | | |
|--|-----------------------------|----------------------------|----------------------|---------------------|---------------------|----------------------|--|
| Sex and condition group | Jul <u>y</u> - September | October- December | January- March | April- June | July- September | October- December | |
| Both sexes | , | Incidence of acute conditi | | | | | |
| All acute conditions | 41.7 | 95.4 | 74.3 | 48.6 | 37,1 | 54.5 | |
| Infective and parasitic diseases | 4.8 | 4.1 | 5.9 | 8.2 | 4.2 | 5.4 | |
| Respiratory conditions Upper respiratory Other respiratory | 18.3 12.0 6.3 | 75.2 30.6 44.6 | 52.6 31.7 20.9 | 22.9 16.8 6.1 | 15.0 10.4 4.5 | 34.7 24.6 10.1 | |
| Digestive system conditions | 4.1 | 2.8 | 3.5 | 3.9 | 3.4 | 3.1 | |
| Injuries | 8.3 | 6.6 | 6.0 | 7.3 | 9.1 | 6.5 | |
| All other acute conditions | 6.2 | 6.8 | 6.3 | 6.4 | 5.5 | 4.9 | |
| Male | | _ | | , | | | |
| All acute conditions | 38.9 | 90.7 | 72.7 | 45.2 | 35.7 | 51.5 | |
| Infective and parasitic diseases | 4.1 | 3.8 | 6.6 | 7.3 | 3.6 | 5.5 | |
| Respiratory conditions Upper respiratory Other respiratory | 15.7 10.3 5.4 | 72.0 28.5 43.5 | 50.5 29.5 21.0 | 19.7 14.4 5.3 | 14.4 9.9 4.5 | 31.8 22.6 9.2 | |
| Digestive system conditions | 3.1 | 3.1 | 3.4 | 3.6 | 3.0 | 3 71 | |
| Injuries | 10.5 | 6.9 | 7.1 | 9.5 | 10.5 | 7.6 | |
| All other acute conditions | 5.4 | 4.9 | 5.1 | 5.2 | 4.2 | 3.5 | |
| Female | | | | | | | |
| All acute conditions | 44.4 | 99.9 | 75.8 | 51.9 | 38.5 | 57.3 | |
| Infective and parasitic diseases | 5.4 | . 4.3 | 5.3 | 9.0 | 4.8 | 5.2 | |
| Respiratory conditions Upper respiratory Other respiratory | 20.8 13.7 7.1 | 78.2 32.7 45.5 | 54.5 33.8 20.7 | 26.0 19.1 6.9 | 15.5 10.9 4.6 | 37.3 26.4 10.9 | |
| Digestive system conditions | 5.1 | 2.4 | 3.6 | 4.2 | 3.7 | 3.1 | |
| Injuries | 6.3 | 6.4 | 5.0 | 5.3 | 7.7 | 5.4 | |
| All other acute conditions | 6.9 | 8.6 | 7.4 | 7.5 | 6.7 | 6.3 | |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

by sex and condition group: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | - 1 | 959 | | | 1 | | 1961 | | |
|----------------------|----------------------|--------------------|----------------------|----------------------|---------------------|---------------------|----------------------|----------------------|---------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| per 100 p | ersons pe | r quarter | | | | | | | <u> </u> |
| 68.9 | 54.1 | 34.5 | 53.0 | 74.5 | 41.3 | 37.2 | 56.5 | 63.0 | 45.2 |
| 8.5 | 7.8 | 4.2 | 5.9 | 7.9 | 6.5 | 4.3 | 5.7 | 9.2 | 8.4 |
| 45.5 30.6 | 30.5 17.4 | 12.8 9.4 | 32.9 22.8 | 53.4 26.2 | 19.8 13.6 | 14.1 9.7 | 35.8 25.1 | 39.8 28.3 | 20.5 14.8 |
| 14.9 | 13.0 | 3.4 | 10.1 | 27.2 | 6.2 | 4.3 | 10.7 | 11.6 | 5.7 |
| 2.9 | 2.5 | 3.1 | 2.9 | 2.2 | 2.4 | 3.5 | 2.7 | 3.0 | 3.5 |
| 5.8 | 7.5 | 7.9 | 5.7 | 5.7 | 6.8 | 8.7 | 6.8 | 5.6 | 6.8 |
| 6.2 | 5.8 | 6.5 | 5.5 | 5.2 | 5.8 | 6.6 | 5.5 | 5.3 | 6.0 |
| 64.0 | 53.3 | 32.8 | 48.0 | 70.0 | 39.1 | 36.7 | 54.6 | 60.6 | 41.6 |
| 9.3 | 8.4 | 3.9 | 6.2 | 7.7 | 5.2 | 4.5 | 5.2 | 9.3 | 7.2 |
| 40.5 27.1 13.4 | 28.1 15.9 12.2 | 12.5 9.2 3.3 | 29.2 20.9 8.3 | 50.2 24.4 25.8 | 19.7 13.5 6.2 | 12.6 8.8 3.8 | 33.7 23.9 9.8 | 37.4 26.9 10.5 | 18.5 13.1 5.4 |
| 2.5 | 2.2 | 3.0 | 2.3 | 2.3 | 2.1 | 3.7 | 2.7 | 2.9 | 3.2 |
| 6.9 | 9.7 | 8.7 | 6.4 | 6.2 | 8.1 | 10.5 | 8.7 | 6.5 | 8.3 |
| 4.8 | 4.8 | 4.7 | 3.9 | 3.7 | 4.0 | 5.4 | 4.3 | 4.6 | 4.4 |
| 73.6 | 54.9 | 36.1 | 57.7 | 78.6 | 43.3 | 37.7 | 58.2 | 65.2 | 48.6 |
| 7.6 | 7.2 | 4.5 | 5.6 | 8.1 | 7.7 | 4.1 | 6.2 | 9.1 | 9.4 |
| 50.2 33.9 16.3 | 32.7 18.9 13.8 | 13.1 9.5 3.5 | 36.4 24.7 11.7 | 56.5 27.9 28.6 | 19.9 13.7 6.2 | 15.5 10.6 4.9 | 37.8 26.3 11.5 | 42.2 29.6 12.6 | 22.5 16.4 6.1 |
| 3.4 | 2.7 | 3.2 | 3.5 | 2.2 | 2.6 | 3.4 | 2.7 | 3.1 | 3.8 |
| 4.8 | 5.5 | 7.2 | 5.1 | 5.2 | 5.6 | 7.0 | 5.0 | 4.9 | 5.3 |
| 7.6 | 6.8 | 8.2 | 7.1 | 6.7 | 7.5 | 7.7 | 6.6 | 6.0 | 7.6 |

Table 3. Incidence of acute conditions per quarter Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | .958 | | | |
|-------------|-----------------------------|----------------------|-------------------|----------------|--------------------|----------------------|--|--|
| Sex and age | July- September | October- December | January- March | April- June | July- September | October- December | | |
| Both sexes | Both sexes Incidence of acc | | | | | | | |
| All ages | 69,704 | 160,312 | 125,437 | 82,433 | 63,196 | 93,202 | | |
| Under 5 | 12,466 | 26,119 | 22,991 | 16,571 | 13,142 | 17,192 | | |
| 5-14 | 16,044 | 46,246 | 32,245 | 22,329 | 14,986 | 24,214 | | |
| 15-24 | 10,787 | 22,512 | 14,286 | 9,841 | 7,919 | 11,382 | | |
| 25-44 | 16,886 | 36,916 | 29,261 | 18,069 | 15,850 | 21,554 | | |
| 45-64 | 9,374 | 21,658 | 18,980 | 10,714 | 7,982 | 13,507 | | |
| 65+ | 4,148 | 6,861 | 7,674 | 4,909 | 3,317 | 5,353 | | |
| Male | | | ٠ | | | | | |
| All ages | 31,587 | 74,119 | 59,707 | 37,279 | 29,566 | 42,875 | | |
| Under 5 | 6,320 | 12,946 | 12,403 | 8,336 | 7,464 | 9,371 | | |
| 5-14 | 8,216 | 23,304 | 16,673 | 10,788 | 7,081 | 12,601 | | |
| 15-24 | 5,077 | 9,638 | 5,748 | 4,117 | 3,173 | 4,045 | | |
| 25-44 | 6,609 | 15,071 | 13,192 | 7,610 | 6,998 | 8,533 | | |
| 45-64 | 3,704 | 9,972 | 8,444 | 4,226 | 3,717 | 6,312 | | |
| 65+ | 1,660 | 3,188 | 3,247 | 2,203 | 1,133 | 2,012 | | |
| Female | · | | | | | | | |
| All ages | 38,117 | 86,193 | 65,730 | 45,154 | 33,630 | 50,327 | | |
| Under 5 | 6,146 | 13,172 | 10,587 | 8,235 | 5,678 | 7,820 | | |
| 5-14 | 7,827 | 22,942 | 15,572 | 11,542 | 7,904 | 11,613 | | |
| 15-24 | 5,710 | 12,874 | 8,538 | 5,724 | 4,746 | 7,337 | | |
| 25-44 | 10,277 | 21,846 | 16,069 | 10,460 | 8,852 | 13,021 | | |
| 45-64 | 5,670 | 11,686 | 10,537 | 6,488 | 4,265 | 7,195 | | |
| 65+ | 2,488 | 3,673 | 4,427 | 2,706 | 2,184 | 3,340 | | |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

by sex and age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II]

| | | 1959 | | | | | 1961 | | |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| ditions in | thousan | ds | | | | | | | |
| 118,305 | 93,238 | 59,742 | 92,178 | 130,572 | 72,659 | 65,719 | 100,308 | 112,359 | 80,954 |
| 22,502 | 16,472 | 11,765 | 20,567 | 22,713 | 16,817 | 15,365 | 18,795 | 25,151 | 15,915 |
| 33,503 | 28,617 | 14,623 | 24,727 | 32,575 | 18,799 | 15,255 | 27,891 | 29,771 | 21,748 |
| 13,319 | 11,077 | 7,198 | 11,003 | 15,733 | 9,104 | 8,262 | 12,188 | 14,149 | 9,810 |
| 26,504 | 19,857 | 15,209 | 20,214 | 30,389 | 13,934 | 15,274 | 23,132 | 21,532 | 18,037 |
| 15,220 | 13,296 | 7,845 | 11,096 | 20,693 | 10,400 | 8,074 | 13,276 | 15,820 | 11,447 |
| 7,257 | 3,919 | 3,102 | 4,570 | 8,469 | 3,606 | 3,489 | 5,026 | 5,936 | 3,996 |
| | | | | | | | | | |
| 53,455 | 44,684 | 27,627 | 40,662 | 59,779 | 33,496 | 31,524 | 47,195 | 52,609 | 36,211 |
| 11,884 | 8,519 | 5,766 | 10,621 | 11,716 | 8,862 | 8,225 | 9,972 | 12,865 | 7,348 |
| 17,075 | 14,877 | 7,563 | 11,187 | 15,526 | 9,040 | 7,718 | 14,488 | 15,105 | 11,593 |
| 5,612 | 4,071 | 2,807 | 4,331 | 7,055 | 4,001 | 3,429 | 5,041 | 6,198 | 4,264 |
| 10,041 | 9,150 | 6,507 | 8,511 | 12,395 | 5,633 | 6,637 | 9,499 | 9,202 | 6,506 |
| 6,224 | 6,745 | 3,772 | 4,378 | 9,500 | 4,540 | 3,900 | 6,022 | 6,728 | 4,984 |
| 2,618 | 1,321 | 1,212 | 1,635 | 3,587 | 1,420 | 1,615 | 2,173 | 2,511 | 1,516 |
| | | | | | | | | | : |
| 64,851 | 48,554 | 32,115 | 51,516 | 70,793 | 39,163 | 34,194 | 53,113 | 59,750 | 44,742 |
| 10,618 | 7,953 | 6,000 | 9,947 | 10,997 | 7,955 | 7,140 | 8,823 | 12,286 | 8,567 |
| 16,428 | 13,740 | 7,059 | 13,541 | 17,050 | 9,759 | 7,538 | 13,403 | 14,666 | 10,155 |
| 7,707 | 7,006 | 4,391 | 6,673 | 8,678 | 5,103 | 4,833 | 7,148 | 7,951 | 5,546 |
| 16,463 | 10,707 | 8,701 | 11,703 | 17,994 | 8,301 | 8,636 | 13,633 | 12,330 | 11,531 |
| 8,996 | 6,551 | 4,073 | 6,718 | 11,192 | 5,859 | 4,174 | 7,253 | 9,092 | 6,463 |
| 4,639 | 2,597 | 1,890 | 2,935 | 4,881 | 2,186 | 1,874 | 2,853 | 3,425 | 2,480 |

Table 4. Incidence of acute conditions per 100 persons per [Data are based on household interviews of the civilian noninstitutional population. The survey design, general qualifica-

| | 1957 | | 1958 | | | | | |
|-------------|-------------------------------|----------|-------------------|----------------|--------------------|----------------------|--|--|
| Sex and age | July- | October- | Tanuama | | | 0-1-1 | | |
| bek and age | September | December | January- March | April- June | July- September | October- December | | |
| Both sexes | Incidence of acute conditions | | | | | | | |
| All ages | 41.7 | 95.4 | 74.3 | 48.6 | 37.1 | 54.5 | | |
| Under 5 | 65.0 | 135.0 | 118.4 | 85.2 | 67.3 | 87.5 | | |
| 5-14 | 48.7 | 139.4 | 96.5 | 66.4 | 44.2 | 70.9 | | |
| 15-24 | 52.1 | 107.3 | 67.3 | 45.9 | 36.6 | 52.1 | | |
| 25-44 | 37.0 | 80.8 | 64.1 | 39.6 | 34.8 | 47.4 | | |
| 45-64 | 27.4 | 63.0 | 55.0 | 30.9 | 22.9 | 38.6 | | |
| 65+ | 28.8 | 47.3 | 52.8 | 33.6 | 22.6 | 36.2 | | |
| Male | | | | | | | | |
| All ages | 38.9 | 90.7 | 72.7 | 45.2 | 35.7 | 51.5 | | |
| Under 5 | 64.7 | 131.4 | 125.3 | 84.2 | 75.1 | 93.7 | | |
| 5-14 | 48.9 | 137.7 | 97.8 | 62.9 | 41.0 | 72.3 | | |
| 15-24 | 53.1 | 99.0 | 58.1 | 41.1 | 31.4 | 39.5 | | |
| 25-44 | 30.2 | 68.9 | 60.2 | 34.8 | 32.0 | 39.1 | | |
| 45-64 | 22.3 | 59.7 | 50.4 | 25.1 | 22.0 | 37.2 | | |
| 65+ | 25.1 | 48.0 | 48.8 | 33.1 | 16.9 | 29.9 | | |
| Female | | | | | | | | |
| All ages | 44.4 | 99.9 | 75.8 | 51.9 | 38.5 | 57.3 | | |
| Under 5 | 65.3 | 138.8 | 111.1 | 86.3 | 59.3 | 81.1 | | |
| 5-14 | 48.5 | 141.2 | 95.2 | 70.1 | 47.6 | 69.4 | | |
| 15-24 | 51.3 | 114.5 | 75.3 | 50.0 | 41.2 | 63.2 | | |
| 25-44 | 43.2 | 91.9 | 67.6 | 44.1 | 37.3 | 55.0 | | |
| 45-64 | 32.2 | 66.1 | 59.3 | 36.3 | 23.8 | 39.9 | | |
| 65+ | 31.9 | 46.8 | 56.1 | 34.1 | 27.3 | 41.4 | | |
| | L | L | L | L | L | L | | |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

quarter by sex and age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | : | 1961 | | |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June | |
| per 100 p | ersons pe | r quarter | | | | | | | | |
| 68.9 | 54.1 | 34.5 | 53.0 | 74.5 | 41.3 | 37.2 | 56.5 | 63.0 | 45.2 | |
| 114.3 | 83.5 | 59.5 | 103.4 | 113.4 | 84.1 | 76.7 | 93.2 | 124.5 | 78.8 | |
| 97.2 | 82.3 | 41.7 | 69.9 | 90.8 | 51.9 | 41.8 | 75.6 | 80.0 | 57.9 | |
| 60.4 | 49.8 | 32.1 | 48.5 | 68.6 | 39.4 | 35.5 | 52.0 | 59.8 | 41.2 | |
| 58.3 | 43.7 | 33.5 | 44.6 | 66.8 | 30.7 | 33.6 | 50.9 | 47.4 | 39.7 | |
| 43.3 | 37.7 | 22.1 | 31.2 | 57.8 | 28.9 | 22.4 | 36.6 | 43.5 | 31.4 | |
| 48.8 | 26.2 | 20.6 | 30.2 | 55.6 | 23.6 | 22.7 | 32.5 | 38.2 | 25.6 | |
| | | | | | | | | | 2 | |
| 64.0 | 53.3 | 32.8 | 48.0 | 70.0 | 39.1 | 36.7 | 54.6 | 60.6 | 41.6 | |
| 118.5 | 84.8 | 57.2 | 104.8 | 114.9 | 87.0 | 80.7 | 97.2 | 125.1 | 71.4 | |
| 97.1 | 83.9 | 42.3 | 61.9 | 84.7 | 48.9 | 41.4 | 77.0 | 79.5 | 60.5 | |
| 54.2 | 38.9 | 26.6 | 40.2 | 64.8 | 36.3 | 30.9 | 45.2 | 55.0 | 37.5 | |
| 46.0 | 42.0 | 29.8 | 39.3 | 57.0 | 25.9 | 30.5 | 43.7 | 42.3 | 29.9 | |
| 36.6 | 39.5 | 22.0 | 25.5 | 54.9 | 26.2 | 22.4 | 34.5 | 38.4 | 28.4 | |
| 38.8 | 19.5 | 17.8 | 23.9 | 52.2 | 20.6 | 23.4 | 31.3 | 36.1 | 21.7 | |
| | | | | | | | | | | |
| 73.6 | 54.9 | 36.1 | 57.7 | 78.6 | 43.3 | 37.7 | 58.2 | 65.2 | 48.6 | |
| 109.9 | 82.2 | 61.8 | 101.9 | 111.9 | 81.0 | 72.6 | 89.2 | 123.9 | 86.4 | |
| 97.3 | 80.7 | 41.1 | 78.2 | 97.0 | 55.1 | 42.2 | 74.3 | 80.5 | 55.3 | |
| 65.9 | 59.5 | 37.0 | 55.9 | 72.0 | 42.1 | 39.6 | 58.2 | 64.2 | 44.5 | |
| 69.6 | 45.3 | 36.8 | 49.6 | 75.9 | 35.0 | 36.5 | 57.5 | 52.0 | 48.7 | |
| 49.6 | 36.0 | 22.3 | 36.6 | 60.5 | 31.5 | 22.3 | 38.6 | 48.2 | 34.1 | |
| 57.1 | 31.8 | 23.0 | 35.4 | 58.4 | 26.0 | 22.2 | 33.4 | 39.9 | 28.7 | |

Table 5. Days of restricted activity associated with acute conditions per [Data are based on household interviews of the civilian, noninstitutional repulation. The survey design, general qualifica-

| | 195 | 57 | | 1 | 958 | |
|----------------------------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | D | ays of rest | ricted ac- |
| All acute conditions | 282,240 | 753,492 | 535,499 | 351,489 | 249,991 | 338,319 |
| Infective and parasitic diseases | 32,307 | 29,148 | 54,445 | 74,421 | 26,609 | 32,876 |
| Respiratory conditions | 100,721 | 584,101 | 339,869 | 147,355 | 65,776 | 183,662 |
| Digestive system conditions | 22,828 | 17,873 | 21,545 | 16,912 | 16,881 | 17,510 |
| Injuries | 69,144 | 59,087 | 61,637 | 56,985 | 91,038 | 61,320 |
| All other acute conditions | 57,239 | 63,284 | 58,005 | 55,816 | 49,688 | 42,950 |
| Male | | | | | | |
| All acute conditions | 111,481 | 332,804 | 237,740 | 151,701 | 112,142 | 139,335 |
| Infective and parasitic diseases | 14,444 | 9,622 | 28,437 | 35,497 | 11,175 | 15,469 |
| Respiratory conditions | 38,276 | 263,801 | 148,799 | 62,079 | 30,747 | 73,068 |
| Digestive system conditions | 9,896 | 7,959 | 7,106 | 7,187 | 7,216 | 7,909 |
| Injuries | 31,522 | 32,486 | 33,868 | 30,776 | 44,894 | 33,165 |
| All other acute conditions | 17,343 | 18,936 | 19,532 | 16,163 | 18,110 | 9,724 |
| Female | | | : | | | [|
| All acute conditions | 170,759 | 420,688 | 297,758 | 199,787 | 137,849 | 198,984 |
| Infective and parasitic diseases | 17,864 | 19,525 | 26,008 | 38,924 | 15,434 | 17,406 |
| Respiratory conditions | 62,445 | 320,299 | 191,070 | 85,276 | 35,029 | 110,594 |
| Digestive system conditions | 12,933 | 9,913 | 14,439 | 9,725 | 9,665 | 9,602 |
| Injuries | 37,622 | 26,602 | 27,769 | 26,208 | 46,143 | 28,156 |
| All other acute conditions | 39,896 | 44,348 | 38,473 | 39,654 | 31,578 | 33,226 |

quarter by sex and condition group: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Pefinitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | .960 | | 196 | 1 |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| tivity in | thousand | s | | | | | | | |
| 463,654 | 394,349 | 246,721 | 356,285 | 619,276 | 323,169 | 254,411 | 382,081 | 516,912 | 371,815 |
| 69,954 | 74,281 | 30,095 | 48,216 | 69,203 | 61,873 | 28,578 | 38,453 | 81,553 | 76,789 |
| 260,941 | 189,693 | 68,870 | 169,165 | 402,446 | 114,803 | 69,265 | 187,019 | 250,753 | 124,350 |
| 16,590 | 13,584 | 19,705 | 21,922 | 17,668 | 18,420 | 18,067 | 16,503 | 19,503 | 23,301 |
| 59,985 | 58,219 | 63,844 | 59,720 | 70,166 | 70,784 | 79,126 | 81,340 | 96,540 | 74,604 |
| 56,184 | 58,573 | 64,207 | 57,263 | .59,793 | 57,289 | 59,375 | 58,766 | 68,564 | 72,771 |
| | | , | | | | | ļ | | |
| 200,536 | 170,560 | 108,724 | 148,613 | 266,408 | 141,932 | 119,781 | 175,596 | 221,813 | 160,612 |
| 40,624 | 36,967 | 14,169 | 20,869 | 34,246 | 24,078 | 12,789 | 17,053 | 41,457 | 33,339 |
| 108,807 | 79,411 | 33,169 | 73,417 | 171,148 | 52,417 | 30,701 | 82,685 | 105,700 | 53,917 |
| 4,406 | 5,911 | 9,322 | 7,070 | 11,609 | 9,223 | 7,891 | 6,092 | 7,715 | 10,126 |
| 29,879 | 27,161 | 32,144 | 31,117 | 32,764 | 39,126 | 47,090 | 53,260 | 49,460 | 43,417 |
| 16,820 | 21,110 | 19,921 | 16,140 | 16,642 | 17,089 | 21,309 | 16,506 | 17,481 | 19,813 |
| | | | | | | | | | |
| 263,119 | 223,789 | 137,998 | 207,672 | 352,868 | 181,237 | 134,630 | 206,485 | 295,099 | 211,203 |
| 29,331 | 37,314 | 15,927 | 27,347 | 34,957 | 37,795 | 15,788 | 21,400 | 40,096 | 43,451 |
| 152,134 | 110,282 | 35,701 | 95,747 | 231,298 | 62,386 | 38,564 | 104,334 | 145,053 | 70,433 |
| 12,184 | 7,673 | 10,383 | 14,852 | 6,060 | 9,198 | 10,176 | 10,411 | 11,787 | 13,175 |
| 30,106 | 31,058 | 31,700 | 28,603 | 37,402 | 31,658 | 32,036 | 28,080 | 47,079 | 31,187 |
| 39,364 | 37,463 | 44,287 | 41,122 | 43,151 | 40,199 | 38,066 | 42,259 | 51,083 | 52,957 |

Table 6. Days of restricted activity associated with acute conditions per 100 per[Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 57 | | 1 | .958 | |
|----------------------------------|----------------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and condition group | Jul y- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | Days o | f restricte | d activity |
| All acute conditions | 168.9 | 448.4 | 317.2 | 207.4 | 146.9 | 197.9 |
| Infective and parasitic diseases | 19.3 | 17.3 | 32.3 | 43.9 | 15.6 | 19.2 |
| Respiratory conditions | 60.3 | 347.6 | 201.3 | 87.0 | 38.7 | 107.4 |
| Digestive system conditions | 13.7 | 10.6 | 12.8 | 10.0 | 9.9 | 10.2 |
| Injuries | 41.4 | 35.2 | 36.5 | 33.6 | 53.5 | 35.9 |
| All other acute conditions | 34.3 | 37.7 | 34.4 | 32.9 | 29.2 | 25.1 |
| Male | | | | | | |
| All acute conditions | 137.2 | 407.1 | 289.4 | 184.0 | 135.5 | 167.5 |
| Infective and parasitic diseases | 17.8 | 11.8 | 34.6 | 43.1 | 13.5 | 18.6 |
| Respiratory conditions | 47.1 | 322.7 | 181.1 | 75.3 | 37.1 | 87.8 |
| Digestive system conditions | 12.2 | 9.7 | 8.7 | 8.7 | 8.7 | 9.5 |
| Injuries | 38.8 | 39.7 | 41.2 | 37.3 | 54.2 | 39.9 |
| All other acute conditions | 21.3 | 23.2 | 23.8 | 19.6 | 21.9 | 11.7 |
| Female | | 1 | i | | | |
| All acute conditions | 198.9 | 487.5 | 343.6 | 229.7 | 157.8 | 226.7 |
| Infective and parasitic diseases | 20.8 | 22.6 | 30.0 | 44.7 | 17.7 | 19.8 |
| Respiratory conditions | 72.7 | 371.2 | 220.5 | 98.0 | 40.1 | 126.0 |
| Digestive system conditions | 15.1 | 11.5 | 16.7 | 11.2 | 11.1 | 10.9 |
| Injuries | 43.8 | 30.8 | 32.0 | 30.1 | 52.8 | 32.1 |
| All other acute conditions | 46.5 | 51.4 | 44.4 | 45.6 | 36.1 | 37.9 |

sons per quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | .960 | | 196 | 1 |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| per 100 per | sons per | quarter | | | | | | | |
| 270.1 | 228.8 | 142.5 | 204.8 | 353.1 | 183.6 | 144.0 | 215.1 | 289.7 | 207.6 |
| 40.7 | 43.1 | 17.4 | 27.7 | 39.5 | 35.2 | 16.2 | 21.7 | 45.7 | 42.9 |
| 152.0 | 110.0 | 39.8 | 97.3 | 229.5 | 65.2 | 39.2 | 105.3 | 140.5 | 69.4 |
| 9.7 | 7.9 | 11.4 | 12.6 | 10.1 | 10.5 | 10.2 | 9.3 | 10.9 | 13.0 |
| 34.9 | 33.8 | 36.9 | 34.3 | 40.0 | 40.2 | 44.8 | 45.8 | 54.1 | 41.6 |
| 32.7 | 34.0 | 37.1 | 32.9 | 34.1 | 32.6 | 33.6 | 33.1 | 38.4 | 40.6 |
| | | | | | : | | | | |
| 240.0 | 203.3 | 129.0 | 175.6 | 312.2 | 165.7 | 139.3 | 203.3 | 255.7 | 184.4 |
| 48.6 | 44.1 | 16.8 | 24.7 | 40.1 | 28.1 | 14.9 | 19.7 | 47.8 | 38.3 |
| 130.2 | 94.6 | 39.4 | 86.7 | 200.5 | 61.2 | 35.7 | 95.7 | 121.8 | 61.9 |
| 5.3 | 7.0 | 11.1 | 8.4 | 13.6 | 10.8 | 9.2 | 7.1 | 8.9 | 11.6 |
| 35.8 | 32.4 | 38.1 | 36.8 | 38.4 | 45.7 | 54.8 | 61.7 | 57.0 | 49.8 |
| 20.1 | 25.2 | 23.6 | 19.1 | 19.5 | 20.0 | 24.8 | 19.1 | 20.1 | 22.7 |
| | | | | | | | | | |
| 298.6 | 252.9 | 155.3 | 232.6 | 392.0 | 200.6 | 148.4 | 226.4 | 322.0 | 229.5 |
| 33.3 | 42.2 | 17.9 | 30.6 | 38.8 | 41.8 | 17.4 | 23.5 | 43.7 | 47.2 |
| 172.6 | 124.6 | 40.2 | 107.2 | 256.9 | 69.0 | 42.5 | 114.4 | 158.3 | 76.5 |
| 13.8 | 8.7 | 11.7 | 16.6 | 6.7 | 10.2 | 11.2 | 11.4 | 12.9 | 14.3 |
| 34.2 | 35.1 | 35.7 | 32.0 | 41.5 | 35.0 | 35.3 | 30.8 | 51.4 | 33.9 |
| 44.7 | 42.3 | 49.8 | 46.1 | 47.9 | 44.5 | 42.0 | 46.3 | 55.7 | 57.5 |

Table 7. Days of restricted activity associated with acute condi-[Pata are based on household interviews of the civilian noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | • |
|-------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and age | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | D | ays of rest | ricted ac- |
| All ages | 282,240 | 753,492 | 535,499 | 351,489 | 249,991 | 338,319 |
| Under 5 | 29,769 | 85,078 | 72,969 | 54,030 | 32,433 | 44,185 |
| 5-14 | 50,464 | 228,985 | 122,340 | 91,311 | 49,904 | 72,787 |
| 15-24 | 44,500 | 97,541 | 47,838 | 37,420 | 43,491 | 39,863 |
| 25-44 | 75,576 | 153,950 | 127,826 | 66,665 | 57,369 | 78,624 |
| 45-64 | 49,113 | 131,667 | 108,801 | 57,435 | 42,728 | 59,404 |
| 65+ | 32,818 | 56,272 | 55,724 | 44,628 | 24,066 | 43,455 |
| Male | | | | | | |
| All ages | 111,481 | 332,804 | 237,740 | 151,701 | 112,142 | 139,335 |
| Under 5 | 14,922 | 39,779 | 38,020 | 24,775 | 18,250 | 21,864 |
| 5-14 | 24,269 | 109,844 | 60,843 | 46,732 | 26,883 | 38,280 |
| 15-24 | 13,454 | 38,411 | 17,377 | 17,040 | 16,134 | 11,650 |
| 25-44 | 28,897 | 59,662 | 50,243 | 21,811 | 21,729 | 24,851 |
| 45-64 | 19,915 | 58,107 | 48,104 | 20,807 | 18,987 | 28,020 |
| 65+ | 10,025 | 27,000 | 23,154 | 20,535 | 10,159 | 14,670 |
| Female | | | | | | |
| All ages | 170,759 | 420,688 | 297,758 | 199,787 | 137,849 | 198,984 |
| Under 5 | 14,847 | 45,299 | 34,949 | 29,255 | 14,183 | 22,321 |
| 5-14 | 26,196 | 119,141 | 61,498 | 44,579 | 23,021 | 34,507 |
| 15-24 | 31,046 | 59,130 | 30,461 | 20,379 | 27,357 | 28,213 |
| 25-44 | 46,679 | 94,287 | 77,583 | 44,853 | 35,640 | 53,773 |
| 45-64 | 29,198 | 73,560 | 60,698 | 36,627 | 23,741 | 31,384 |
| 65+ | 22,793 | 29,272 | 32,570 | 24,093 | 13,907 | 28,785 |

tions per quarter by sex and age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| tivity in | thousand | s | | | | | | | |
| 463,654 | 394,349 | 246,721 | 356,285 | 619,276 | 323,169 | 254,411 | 382,081 | 516,912 | 371,815 |
| 67,610 | 53,348 | 31,042 | 56,161 | 66,569 | 46,176 | 33,865 | 47,613 | 84,776 | 46,093 |
| 128,272 | 115,638 | 53,593 | 88,127 | 142,553 | 79,306 | 45,504 | 86,854 | 121,960 | 79,800 |
| 41,058 | 35,004 | 27,790 | 32,665 | 65,440 | 39,181 | 30,960 | 43,099 | 51,178 | 45,257 |
| 94,280 | 75,884 | 59,617 | 83,967 | 133,644 | 64,002 | 66,114 | 90,603 | 108,005 | 92,772 |
| 85,582 | 76,874 | 49,937 | 57,131 | 139,966 | 60,084 | 47,483 | 74,280 | 94,125 | 59,680 |
| 46,852 | 37,601 | 24,743 | 38,234 | 71,104 | 34,420 | 30,485 | 39,631 | 56,868 | 48,214 |
| | | | | | | | | | |
| 200,536 | 170,560 | 108,724 | 148,613 | 266,408 | 141,932 | 119,781 | 175,596 | 221,813 | 160,612 |
| 37,713 | 28,474 | 13,935 | 29,332 | 35,812 | 24,428 | 16,808 | 26,673 | 42,070 | 19,041 |
| 65,654 | 54,360 | 29,007 | 44;780 | 66,189 | 38,456 | 26,802 | 45,446 | 63,906 | 42,748 |
| 15,535 | 9,256 | 12,319 | 11,031 | 26,080 | 15,333 | 12,748 | 20,784 | 19,523 | 19,233 |
| 33,201 | 27,644 | 25,634 | 28,237 | 49,641 | 25,809 | 26,707 | 36,490 | 39,078 | 33,585 |
| 32,434 | 34,163 | 19,879 | 19,855 | 56,633 | 24,387 | 21,073 | 30,938 | 35,513 | 22,882 |
| 15,998 | 16,664 | 7,950 | 15,379 | 32,053 | 13,520 | 15,643 | 15,266 | 21,725 | 23,123 |
| | | | | | | | | | |
| 263,119 | 223,789 | 137,998 | 207,672 | 352,868 | 181,237 | 134,630 | 206,485 | 295,099 | 211,203 |
| 29,897 | 24,875 | 17,107 | 26,829 | 30,757 | 21,748 | 17,057 | 20,940 | 42,707 | 27,052 |
| 62,618 | 61,278 | 24,586 | 43,346 | 76,364 | 40,851 | 18,702 | 41,408 | 58,054 | 37,052 |
| 25,522 | 25,748 | 15,471 | 21,635 | 39,360 | 23,848 | 18,212 | 22,315 | 31,655 | 26,024 |
| 61,079 | 48,240 | 33,983 | 55,731 | 84,004 | 38,193 | 39,407 | 54,114 | 68,928 | 59,187 |
| 53,148 | 42,711 | 30,057 | 37,277 | 83,333 | 35,697 | 26,410 | 43,342 | 58,612 | 36,797 |
| 30,855 | 20,938 | 16,793 | 22,855 | 39,051 | 20,900 | 14,842 | 24,366 | 35,143 | 25,091 |

Table 8. Days of restricted activity associated with acute conditions per [Data are based on household interviews of the civilian noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|-------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and age | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | Days o | of restricte | d activity |
| All ages | 168.9 | 448.4 | 317.2 | 207.4 | 146.9 | 197.9 |
| Under 5 | 155.1 | 439.8 | 375.6 | 277.9 | 166.2 | 224.9 |
| 5-14 | 153.2 | 690.4 | 366.3 | 271.5 | 147.3 | 213.0 |
| 15-24 | 215.0 | 465.0 | 225.2 | 174.4 | 200.9 | 182.4 |
| 25-44 | 165.6 | 337.2 | 279.9 | 146.1 | 125.9 | 172.8 |
| 45-64 | 143.4 | 382.7 | 315.0 | 165.6 | 122.7 | 169.8 |
| 65+ | 227.9 | 388.2 | 383.1 | 305.5 | 163.9 | 293.8 |
| Male | | | | | | |
| All ages | 137.2 | 407.1 | 289.4 | 184.0 | 135.5 | 167.5 |
| Under 5 | 152.7 | 403.6 | 384.2 | 250.2 | 183.5 | 218.5 |
| 5-14 | 144.5 | 649.2 | 357.1 | 272.3 | 155.5 | 219.5 |
| 15-24 | 140.7 | 394.5 | 175.6 | 170.2 | 159.4 | 113.7 |
| 25-44 | 132.2 | 272.6 | 229.4 | 99.6 | 99.4 | 113.9 |
| 45-64 | 119.6 | 347.7 | 286.9 | 123.7 | 112.4 | 165.3 |
| 65+ | 151.8 | 406.6 | 348.2 | 308.1 | 151.9 | 218.1 |
| Female | · | | | | | j |
| All ages | 198.9 | 487.5 | 343.6 | 229.7 | 157.8 | 226.7 |
| Under 5 | 157.7 | 477.3 | 366.7 | 306.7 | 148.1 | 231.6 |
| 5-14 | 162.4 | 733.4 | 376.0 | 270.7 | 138.8 | 206.2 |
| 15-24 | 278.8 | 526.1 | 268.6 | 178.1 | 237.3 | 243.0 |
| 25-44 | 196.2 | 396.5 | 326.5 | 188.9 | 150.3 | 227.0 |
| 45-64 | 165.9 | 415.9 | 341.5 | 205.1 | 132.3 | 174.0 |
| 65+ | 292.5 | 372.7 | 412.6 | 303.4 | 173.9 | 356.9 |

100 persons per quarter by sex and age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | | 1959 | | | 1 | 960 | | 196 | 1 |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| per 100 pe | ersons pe | r quarter | | | - | | | | |
| 270.1 | 228.8 | 142.5 | 204.8 | 353.1 | 183.6 | 144.0 | 215.1 | 289.7 | 207.6 |
| 343.3 | 270.6 | 156.9 | 282.3 | 332.5 | 230.8 | 169.1 | 236.2 | 419.7 | 228.2 |
| 372.2 | 332.6 | 152.8 | 249.1 | 397.2 | 219.0 | 124.6 | 235.6 | 327.8 | 212.6 |
| 186.2 | 157.3 | 123.9 | 144.0 | 285.2 | 169.4 | 132.9 | 183.8 | 216.5 | 190.0 |
| 207.3 | 167.0 | 131.2 | 185.4 | 293.9 | 140.9 | 145.5 | 199.5 | 237.7 | 204.2 |
| 243.6 | 218.0 | 141.0 | 160.7 | 390.9 | 167.2 | 131.6 | 205.0 | 258.8 | 163.5 |
| 315.2 | 251.7 | 164.7 | 252.9 | 467.0 | 225.3 | 198.4 | 256.1 | 365.8 | 308.4 |
| ļ | | · | | | | j | | | |
| 240.0 | 203.3 | 129.0 | 175.6 | 312.2 | 165.7 | 139.3 | 203.3 | 255.7 | 184.4 |
| 375.9 | 283.5 | 138.3 | 289.5 | 351.2 | 239.8 | 164.8 | 259.9 | 409.2 | 185.1 |
| 373.2 | 306.4 | 162.1 | 248.0 | 361.3 | 208.0 | 143.7 | 241.5 | 336.5 | 223.1 |
| 149.9 | 88.4 | 116.5 | 102.5 | 239.4 | 139.3 | 114.8 | 186.3 | 173.4 | 169.3 |
| 152.2 | 126.8 | 117.6 | 130.2 | 228.1 | 118.7 | 122.8 | 167.9 | 179.7 | 154.4 |
| 190.7 | 200.2 | 116.1 | 115.6 | 327.5 | 140.6 | 121.1 | 1 7 7.2 | 202.7 | 130.2 |
| 237.1 | 246.3 | 117.0 | 225.1 | 466.4 | 196.5 | 226.4 | 219.8 | 312.0 | 330.8 |
| | | | | | | | | | |
| 298.6 | 252.9 | 155.3 | 232.6 | 392.0 | 200.6 | 148.4 | 226.4 | 322.0 | 229.5 |
| 309.4 | 257.1 | 176.3 | 274.8 | 313.0 | 221.5 | 173.5 | 211.6 | 430.6 | 272.8 |
| 371.0 | 360.0 | 143.2 | 250.3 | 434.6 | 230.5 | 104.6 | 229.4 | 318.8 | 201.7 |
| 218.3 | 218.7 | 130.5 | 181.4 | 326.6 | 196.7 | 149.3 | 181.6 | 255.6 | 208.9 |
| 258.1 | 204.0 | 143.9 | 236.0 | 354.3 | 161.2 | 166.4 | 228.4 | 290.8 | 249.8 |
| 293.3 | 234.6 | 164.3 | 202.8 | 450.1 | 192.0 | 141.4 | 231.0 | 310.9 | 194.3 |
| 380.1 | 256.3 | 204.0 | 275.9 | 467.5 | 248.9 | 175.5 | 285.7 | 409.4 | 290.2 |

Table 9. Days of bed disability associated with acute conditions per Data are based on household interviews of the civilian noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|----------------------------------|--------------------|----------------------|-------------------|----------------|--------------------|---------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October December |
| Both sexes | | | | | Days of | bed disa- |
| All acute conditions | 103,737 | 378,836 | 251,570 | 138,877 | 101,613 | 145,125 |
| Infective and parasitic diseases | 15,462 | 15,056 | 29,234 | 29,940 | 13,806 | 14,646 |
| Respiratory conditions | 42,400 | 312,251 | 173,929 | 64,521 | 33,474 | 79,975 |
| Digestive system conditions | 9,207 | 6,942 | 7,986 | 8,996 | 6,955 | 8,503 |
| Injuries | 19,917 | 16,429 | 20,206 | 15,695 | 26,790 | 22,046 |
| All other acute conditions | 16,751 | 28,159 | 20,215 | 19,725 | 20,588 | 19,954 |
| Male | | | | | | |
| All acute conditions | 42,253 | 165,026 | 108,522 | 55,830 | 47,292 | 59,679 |
| Infective and parasitic diseases | 6,312 | 4,971 | 14,317 | 13,762 | 5,525 | 7,538 |
| Respiratory conditions | 18,148 | 141,119 | 77,904 | 26,061 | 16,581 | 30,699 |
| Digestive system conditions | 3,344 | 3,386 | 1,867 | 3,160 | 2,624 | 3,420 |
| Injuries | 9,686 | 8,881 | 8,845 | 7,677 | 13,810 | 13,406 |
| All other acute conditions | 4,761 | 6,668 | 5,590 | 5,171 | 8,752 | 4,617 |
| <u>Female</u> | | | ! | | | |
| All acute conditions | 61,484 | 213,811 | 143,048 | 83,046 | 54,321 | 85,446 |
| Infective and parasitic diseases | 9,150 | 10,085 | 14,917 | 16,178 | 8,281 | 7,109 |
| Respiratory conditions | 24,252 | 171,133 | 96,025 | 38,460 | 16,894 | 49,276 |
| Digestive system conditions | 5,862 | 3,555 | 6,119 | 5,836 | 4,330 | 5,083 |
| Injuries | 10,230 | 7,547 | 11,361 | 8,018 | 12,980 | 8,641 |
| All other acute conditions | 11,990 | 21,490 | 14,625 | 14,555 | 11,836 | 15,337 |

quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | | 1961 | |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|----------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| bility in | thousand | ls | | | | | | | |
| 195,661 | 173,448 | 96,706 | 139,668 | 282,239 | 126,603 | 85,061 | 146,013 | 212,429 | 147,851 |
| 30,663 | 31,654 | 14,795 | 22,908 | 33,114 | 27,265 | 15,233 | 19,711 | 38,279 | 34,492 |
| 117,505 | 95,069 | . 26,756 | 70,753 | 195,607 | 51,617 | 26,041 | 76,329 | 111,196 | 54,379 |
| 7,007 | 6,322 | 10,592 | 10,136 | 7,054 | 7,348 | 8,309 | 7,176 | 7,146 | 11,757 |
| 17,662 | 17,263 | 19,402 | 14,525 | 18,042 | 18,881 | 14,315 | 19,578 | 29,236 | 20,374 |
| 22,823 | 23,139 | 25,162 | 21,347 | 28,422 | 21,492 | 21,162 | 23,219 | 26,573 | 26,850 |
| | | | | | | | | | |
| 84,054 | 75,502 | 39,761 | 59,229 | 120,421 | 55,690 | 39,162 | 63,938 | 86,250 | 59,642 |
| 17,579 | 16,162 | 7,053 | 9,552 | 16,130 | 11,307 | 5,890 | 9,009 | 18,783 | 14,747 |
| 49,609 | 41,178 | 13,490 | 30,482 | 84,499 | 24,515 | 12,742 | 33,738 | 44,151 | 23,472 |
| 1,714 | 2,394 | 4,511 | 3,045 | 5,014 | 3,557 | 2,698 | 2,953 | 3,040 | 4,971 |
| 8,602 | 7,896 | 8,059 | 9,418 | 8,042 | 10,383 | 10,264 | 11,954 | 12,918 | 11,611 |
| 6,550 | 7,872 | 6,648 | 6,732 | 6,737 | 5,928 | 7 ¹ , 567 | 6,284 | 7,359 | 4,842 |
| | - | | | : | | | i | | |
| 111,607 | 97,946 | 56,945 | 80,439 | 161,818 | 70,913 | 45,899 | 82,075 | 126,179 | 88,209 |
| 13,085 | 15,492 | 7,742 | 13,355 | 16,984 | 15,957 | 9,343 | 10,702 | 19,497 | 19,746 |
| 67,896 | 53,891 | 13,266 | 40,271 | 111,109 | 27,103 | 13,299 | 42,591 | 67,045 | 30,907 |
| 5,292 | 3,928 | 6,081 | 7,091 | 2,040 | 3,791 | 5,611 | 4,223 | 4,105 | 6,785 |
| 9,060 | 9,367 | 11,342 | 5,107 | 10,001 | 8,498 | 4,052 | 7,624 | 16,317 | 8,763 |
| 16,274 | 15,267 | 18,513 | 14,614 | 21,685 | 15,564 | 13,595 | 16,935 | 19,214 | 22,008 |

Table 10. Days of bed disability associated with acute conditions per 100 per-Pata are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|----------------------------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | Days o | f bed disab | ility per |
| All acute conditions | 62.1 | 225.4 | 149.0 | 82.0 | 59.7 | 84.9 |
| Infective and parasitic diseases | 9.3 | 9.0 | 17.3 | 17.7 | 8.1 | 8.6 |
| Respiratory conditions | 25.4 | 185.8 | 103.0 | 38.1 | 19.7 | 46.8 |
| Digestive system conditions | 5.5 | 4.1 | 4.7 | 5.3 | 4.1 | 5.0 |
| Injuries | 11.9 | 9.8 | 12.0 | 9.3 | 15.7 | 12.9 |
| All other acute conditions | 10.0 | 16.8 | 12.0 | 11.6 | 12.1 | 11.7 |
| Male | | | i | • | · | |
| All acute conditions | 52.0 | 201.9 | 132.1 | 67.7 | 57.1 | 71.7 |
| Infective and parasitic diseases | 7.8 | 6.1 | 17.4 | 16.7 | 6.7 | 9.1 |
| Respiratory conditions | 22.3 | 172.6 | 94.8 | 31.6 | 20.0 | 36.9 |
| Digestive system conditions | 4.1 | 4.1 | 2.3 | 3.8 | 3.2 | 4.1 |
| Injuries | 11.9 | 10.9 | 10.8 | 9.3 | 16.7 | 16.1 |
| All other acute conditions | 5.9 | 8.2 | 6.8 | 6.3 | 10.6 | 5.6 |
| Female | | | | | li . | - |
| All acute conditions | 71.6 | 247.8 | 165.1 | 95.5 | 62.2 | 97.4 |
| Infective and parasitic diseases | 10.7 | 11.7 | 17.2 | 18.6 | 9.5 | 8.1 |
| Respiratory conditions | 28.2 | 198.3 | 110.8 | 44.2 | 19.3 | 56.1 |
| Digestive system conditions | 6.8 | 4.1 | 7.1 | 6.7 | 5.0 | 5.8 |
| Injuries | 11.9 | 8.7 | 13.1 | 9.2 | 14.9 | 9.8 |
| All other acute conditions | 14.0 | 24.9 | 16.9 | 16.7 | 13.5 | 17.5 |

sons per quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix I. Pefinitions of terms are given in Appendix II]

| ſ | | . 1 | 959 | | | 1 | 960 | | 1961 | |
|---|----------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| [| March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| 1 | 00 perso | ns per qu | arter | | | | - | | | |
| L | 114.0 | 100.6 | 55.9 | 80.3 | 160.9 | 71.9 | 48.1 | 82.2 | 119.1 | 82.5 |
| | 17.9 | 18.4 | 8.5 | 13.2 | 18.9 | 15.5 | 8.6 | 11.1 | 21.5 | 19.3 |
| | 68.4 | 55.1 | 15.5 | 40.7 | 111.5 | 29.3 | 14.7 | 43.0 | 62.3 | 30.4 |
| | 4.1 | 3.7 | 6.1 | 5.8 | 4.0 | 4.2 | 4.7 | 4.0 | 4.0 | 6.6 |
| | 10.3 | 10.0 | 11.2 | 8.4 | 10.3 | 10.7 | 8.1 | 11.0 | 16.4 | 11.4 |
| | 13.3 | 13.4 | 14.5 | 12.3 | 16.2 | 12.2 | 12.0 | 13.1 | 14.9 | 15.0 |
| | | | | | | | | | | |
| | 100.6 | 90.0 | 47.2 | 70.0 | 141.1 | 65.0 | 45.5 | 74.0 | 99.4 | 68.5 |
| | 21.0 | 19.3 | 8.4 | 11.3 | 18.9 | 13.2 | 6.8 | 10.4 | 21.6 | 16.9 |
| | 59.4 | 49.1 | 16.0 | 36.0 | 99.0 | 28.6 | 14.8 | 39.1 | 50.9 | 26.9 |
| | 2.1 | 2.9 | 5.4 | 3.6 | 5.9 | 4.2 | 3.1 | 3.4 | 3.5 | 5.7 |
| | 10.3 | 9.4 | 9.6 | 11.1 | 9.4 | 12.1 | 11.9 | 13.8 | 14.9 | 13.3 |
| | 7.8 | 9.4 | 7.9 | 8.0 | 7.9 | 6.9 | . 8.8 | 7.3 | 8.5 | 5.6 |
| | | | | | | | | | | |
| | 126.6 | 110.7 | 64.1 | 90.1 | 179.8 | 78.5 | 50.6 | 90.0 | 137.7 | 95.9 |
| | 14.8 | 17.5 | 8.7 | 15.0 | 18.9 | 17.7 | 10.3 | 11.7 | 21.3 | 21.5 |
| | 77.0 | 60.9 | 14.9 | 45.1 | 123.4 | 30.0 | 14.7 | 46.7 | 73.2 | 33.6 |
| | 6.0 | 4.4 | 6.8 | 7.9 | 2.3 | 4.2 | 6.2 | 4.6 | 4.5 | 7.4 |
| | 10.3 | 10.6 | 12.8 | 5.7 | 11.1 | 9.4 | 4.5 | 8.4 | 17.8 | 9.5 |
| | 18.5 | 17.3 | 20.8 | 16.4 | 24.1 | 17.2 | 15.0 | 18.6 | 21.0 | 23.9 |

Table 11. Days of bed disability associated with acute conditions [Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 57 | | 1 | 958 | |
|---------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and age | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | | Days of | bed disa- |
| All ages | 103,737 | 378,836 | 251,570 | 138,877 | 101,613 | 145,125 |
| Under 5 | 14,385 | 37,753 | 31,736 | 22,395 | 15,473 | 16,523 |
| 5-14 | 20,929 | 126,383 | 60,665 | 35,633 | 17,745 | 33,715 |
| 15-24 | 18,076 | 55,532 | 23,579 | 19,069 | 17,095 | 17,117 |
| 25-44 | 25,136 | 71,672 | 59,751 | 23,147 | 23,398 | 33,898 |
| 45-64 | 16,831 | 62,774 | 49,168 | 24,620 | 15,980 | 26,630 |
| 65+ | 8,379 | 24,722 | 26,670 | 14,012 | 11,923 | 17,241 |
| <u>Male</u> | | | | | | |
| All ages | 42,253 | 165,026 | 108,522 | 55,830 | 47,292 | 59,679 |
| Under 5 | 6,747 | 17,352 | 15,849 | 7,870 | 8,027 | 9,035 |
| 5-14 | 10,087 | 60,343 | 30,370 | 17,287 | 8,866 | 16,647 |
| 15-24 | 5,564 | 20,575 | 8,421 | 9,118 | 5,699 | 4,706 |
| 25-44 | 9,078 | 28,347 | 23,022 | 7,427 | 9,759 | 11,029 |
| 45-64 | 8,174 | 25,252 | 20,038 | 8,917 | 9,423 | 14,069 |
| 65+ | 2,603 | 13,157 | 10,822 | 5,211 | 5,519 | 4,194 |
| <u>Female</u> | | | | | | |
| All ages | 61,484 | 213,811 | 143,048 | 83,046 | 54,321 | 85,446 |
| Under 5 | 7,638 | 20,401 | 15,887 | 14,526 | 7,446 | 7,488 |
| 5-14 | 10,842 | 66,040 | 30,296 | 18,346 | 8,879 | 17,068 |
| 15-24 | 12,512 | 34,957 | 15,158 | 9,951 | 11,395 | 12,412 |
| 25-44 | 16,058 | 43,325 | 36,729 | 15,719 | 13,639 | 22,869 |
| 45-64 | 8,658 | 37,523 | 29,130 | 15,703 | 6,558 | 12,561 |
| 65+ | 5,776 | 11,565 | 15,848 | 8,801 | 6,404 | 13,048 |

per quarter by sex and age: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix II. Definitions of terms are given in Appendix II.

1959 1960 1961 January-April-July-October-April-October-January-April-January-July-March June September December March September December March June June bility in thousands 126,603 195,661 | 173,448 96,706 139,668 282,239 85,061 146,013 212,429 147,851 26,905 24,317 13,034 20,880 28,269 22,199 13,294 18,614 37,212 20,880 56,532 20,679 37,549 69,522 30,940 52,824 15,662 40,421 51,785 33,690 21,695 15,381 12,227 14,395 32,129 15,925 9,561 17,466 23,835 18,138 40,229 35,064 22,441 29,393 61,072 24,413 22,583 35,159 40,060 37,072 31,013 33,278 17,286 21,585 65,435 21,301 15,120 23,694 35,296 18,974 19,288 12,583 11,039 15,866 25,812 11,824 8,840 10,659 24,242 19,097 84,054 75,502 39,761 59,229 120,421 55,690 39,162 63,938 86,250 59,642 12,493 14,423 6,756 12,171 14,805 11,381 6,285 10,310 17,926 8,032 27,188 25,474 9,390 18,119 31,598 15,979 9,152 20,628 24,704 17,754 7,994 3,856 13,946 4,607 3,561 4,420 2,867 6,290 9,301 7,824 15,960 13,501 9,841 12,047 22,582 10,646 9,107 13,960 12,480 12,100 10,452 15,202 7,203 6,684 27,321 8,965 7,588 6,937 12,403 6,745 8,037 4,225 2,715 6,647 10,169 4,299 5,813 4,162 9,435 7,189 97,946 111,607 56,945 80,439 161,818 70,913 45,899 82,075 126,179 88,209 12,481 11,824 8,709 6,277 13,464 10,818 7,009 8,304 19,285 12,848 29,344 19,430 27,351 11,289 37,924 14,961 6,510 19,793 27,081 15,937 13,701 10,774 8,371 10,835 18,183 11,506 6,694 11,176 14,534 10,315 24,269 21,563 12,600 17,345 38,490 13,767 13,476 21,199 27,580 24,972 10,083 20,561 18,076 14,902 38,114 12,337 22,893 7,532 16,757 12,229 11,251 8,358 8,324 9,219 15,643 7,525 4,677 4,846 14,806 11,909

Table 12. Days of bed disability associated with acute conditions per 100 Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|---------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|
| Sex and age | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | Days o | f bed disab | ility per |
| All ages | 62.1 | 225.4 | 149.0 | 82.0 | 59.7 | 84.9 |
| Under 5 | 75.0 | 195.1 | 163.4 | 115.2 | 79.3 | 84.1 |
| 5-14 | 63.6 | 381.1 | 181.6 | 106.0 | 52.4 | 98.7 |
| 15-24 | 87.3 | 264.7 | 111.0 | 88.9 | 79.0 | 78.3 |
| 25-44 | 55.1 | 157.0 | 130.9 | 50.7 | 51.3 | 74.5 |
| 45-64 | 49.1 | 182.5 | 142.4 | 71.0 | 45.9 | 76.1 |
| 65+ | 58.2 | 170.5 | 183.4 | 95.9 | 81.2 | 116.6 |
| Male | | | | | | |
| All ages | 52.0 | 201.9 | 132.1 | 67.7 | 57.1 | 71.7 |
| Under 5 | 69.0 | 176.1 | 160.2 | 79.5 | 80.7 | 90.3 |
| 5-14 | 60.0 | 356.6 | 178.2 | 100.7 | 51.3 | 95.5 |
| 15-24 | 58.2 | 211.3 | 85.1 | 91.1 | 56.3 | 45.9 |
| 25-44 | 41.5 | 129.5 | 105.1 | 33.9 | 44.6 | 50.6 |
| 45-64 | 49.1 | 151.1 | 119.5 | 53.0 | 55.8 | 83.0 |
| 65+ | 39.4 | 198.1 | 162.7 | 78.2 | 82.5 | 62.4 |
| <u>Female</u> | | | | | | |
| All ages | 71.6 | 247.8 | 165.1 | 95.5 | 62.2 | 97.4 |
| Under 5 | 81.1 | 215.0 | 166.7 | 152.3 | 77.7 | 77.7 |
| 5-14 | 67.2 | 406.5 | 185.2 | 111.4 | 53.5 | 102.0 |
| 15-24 | 112.3 | 311.0 | 133.6 | 87.0 | 98.8 | 106.9 |
| 25-44 | 67.5 | 182.2 | 154.6 | 66.2 | 57.5 | 96.5 |
| 45-64 | 49.2 | 212.1 | 163.9 | 87.9 | 36.5 | 69.7 |
| 65+ | 74.1 | 147.2 | 200.8 | 110.8 | 80.1 | 161.8 |

persons per quarter by sex and age: United States, $\,$ July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|-------------------|----------------|--------------------|----------------------|-------------------|----------------|--------------------|----------------------|-------------------|----------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| 100 perso | ns per qu | arter | | | | | | | |
| 114.0 | 100.6 | 55.9 | 80.3 | 160.9 | 71.9 | 48.1 | 82.2 | 119.1 | 82.5 |
| 136.6 | 123.3 | 65.9 | 104.9 | 141.2 | 111.0 | 66.4 | 92.3 | 184.2 | 103.4 |
| 164.0 | 152.0 | 59.0 | 106.1 | 193.7 | 85.4 | 42.9 | 109.6 | 139.2 | 89.8 |
| 98.4 | 69.1 | 54.5 | 63.4 | 140.0 | 68.8 | 41.0 | 74.5 | 100.8 | 76.1 |
| 88.5 | 77.1 | 49.4 | 64.9 | 134.3 | 53.7 | 49.7 | 77.4 | 88.1 | 81.6 |
| 88.3 | 94.3 | 48.8 | 60.7 | 182.8 | 59.3 | 41.9 | 65.4 | 97.0 | 52.0 |
| 129.8 | 84.2 | 73.5 | 105.0 | 169.5 | 77.4 | 57.5 | 68.9 | 155.9 | 122.1 |
| , | | | | | | | | | |
| 100.6 | 90.0 | 47.2 | 70.0 | 141.1 | 65.0 | 45.5 | 74.0 | 99.4 | 68.5 |
| 143.8 | 124.4 | 67.1 | 120.1 | 145.2 | 111.7 | 61.6 | 100.4 | 174.3 | 78.1 |
| 154.6 | 143.6 | 52.5 | 100.3 | 172.5 | 86.4 | 49.1 | 109.6 | 130.1 | 92.7 |
| 77.2 | 44.0 | 36.5 | 33.1 | 128.0 | 40.1 | 25.8 | 56.4 | 82.6 | 68.9 |
| 73.2 | 61.9 | 45.1 | 55.6 | 103.8 | 49.0 | 41.9 | 64.2 | 57.4 | 55.6 |
| 61.5 | 89.1 | 42.1 | 38.9 | 158.0 | 51.7 | 43.6 | 39.7 | 70.8 | 38.4 |
| 119.1 | 62.4 | 40.0 | 97.3 | 148.0 | 62.5 | 60.2 | 83.7 | 135.5 | 102.8 |
| | | | | | | | | | |
| 126.6 | 110.7 | 64.1 | 90.1 | 179.8 | 78.5 | 50.6 | 90.0 | 137.7 | 95.9 |
| 129.2 | 122.2 | 64.7 | 89.2 | 137.0 | 110.2 | 71.3 | 83.9 | 194.4 | 129.6 |
| 173.9 | 160.7 | 65.8 | 112.2 | 215.8 | 84.4 | 36.4 | 109.7 | 148.7 | 86.7 |
| 117.2 | 91.5 | 70.6 | 90.8 | 150.9 | 94.9 | 54.9 | 91.0 | 117.4 | 82.8 |
| 102.6 | 91.2 | 53.3 | 73.5 | 162.4 | 58.1 | 56.9 | 89.5 | 116.4 | 105.4 |
| 113.5 | 99.3 | 55.1 | 81.1 | 205.9 | 66.4 | 40.3 | 89.3 | 121.4 | 64.6 |
| 138.6 | 102.3 | 101.1 | 111.3 | 187.3 | 89.6 | 55.3 | 56.8 | 172.5 | 137.7 |

Table 13. Days lost from work associated with acute conditions per Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|---|--|---|---|---|---|---|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | - <u>-</u> | | | Days | lost from |
| All acute conditions | 57,426 | 146,008 | 104,996 | 48,111 | 45,221 | 61,538 |
| Infective and parasitic diseases Respiratory conditions Digestive system conditions Injuries All other acute conditions | 5,959 16,506 5,371 18,686 10,905 | 4,517 109,057 2,598 19,275 10,561 | 5,098 70,281 3,076 17,671 8,870 | 4,066 22,833 2,595 11,893 6,723 | 3,398 8,255 3,324 18,943 11,301 | 3,446 25,385 4,620 20,286 7,800 |
| Male | | | | | | |
| All acute conditions | 29,729 | 89,913 | 66,402 | 30,230 | 30,273 | 37,147 |
| Respiratory conditions | 7,926 11,814 9,990 | 65,231 16,223 8,460 | 42,491 12,926 10,985 | 13,301 9,584 7,346 | 4,269 14,851 11,153 | 12,898 16,688 7,560 |
| All acute conditions | 27,697 | 56,095 | 38,593 | 17,881 | 14,948 | 24,391 |
| Respiratory conditions InjuriesAll other acute conditions | 8,580 6,872 12,245 | 43,826 3,052 9,217 | 27,790 4,744 6,059 | 9,533 2,310 6,038 | 3,986 4,091 6,871 | 12,487 3,598 8,306 |

Table 14. Persons absent from work each day because of acute conditions per [Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | · | 1 | 958 | |
|---|-------------------------------|---------------------------------|---------------------------------|-------------------------------|-------------------------------|-------------------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | Per | sons absent | from work |
| All acute conditions | 938 | 2,384 | 1,714 | 785 | 738 | 1,005 |
| Infective and parasitic diseases Respiratory conditions Digestive system conditions Injuries All other acute conditions | 97 269 88 305 178 | 74 1,781 42 315 172 | 83 1,147 50 289 145 | 66 373 42 194 110 | 55 135 54 309 185 | 56 414 75 331 127 |
| Male | : | | | · | | |
| All acute conditions | 485 | 1,468 | 1,084 | 494 | 494 | 606 |
| Respiratory conditions InjuriesAll other acute conditions | 129 193 163 | 1,065 265 138 | 694 211 179 | 217 156 120 | 70 242 182 | 211 272 123 |
| Female | | | | | | |
| All acute conditions | 452 | 916 | 630 | 292 | 244 | 398 |
| Respiratory conditions Injuries All other acute conditions | 140 112 200 | 716 50 150 | 454 77 99 | 156 38 99 | 65 67 112 | 204 59 136 |

quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|---|---|--|---|---|---|--|---|---|---|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| work in t | housands | | | | | | | | |
| 65,552 | 57,290 | 41,289 | 52,175 | 103,211 | 44,755 | 43,516 | 58,796 | 64,443 | 54,349 |
| 6,387 36,017 2,385 15,134 5,630 | 8,246 25,924 1,891 14,122 7,107 | 3,338 9,912 3,784 16,585 7,670 | 3,650 23,017 3,250 14,656 7,603 | 6,601 67,966 6,019 16,154 6,472 | 3,660 17,048 4,426 12,073 7,549 | 3,689 7,628 2,912 21,685 7,603 | 3,175 25,215 3,329 19,755 7,322 | 6,177 27,130 2,150 21,450 7,536 | 5,703 17,132 5,669 17,663 8,182 |
| 21,388 10,070 7,845 26,250 | 16,970 10,490 11,317 | 6,054 9,532 10,261 | 13,388 9,150 9,080 20,557 | 41,257 11,269 12,317 38,368 | 10,173 9,571 9,271 | 4,603 16,776 7,354 | 15,317 15,999 5,339 | 15,916 16,763 9,018 | 10,786 13,038 9,222 21,304 |
| 14,628 5,064 6,557 | 8,954 3,632 5,927 | 3,858 7,053 4,531 | 9,629 5,506 5,423 | 26,709 4,885 6,774 | 6,875 2,502 6,364 | 3,024 4,908 6,850 | 9,898 3,755 8,488 | 11,214 4,687 6,845 | 6,346 4,625 10,332 |

quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|-------------------------------|--------------------------------|-------------------------------|-------------------------------|----------------------------------|-------------------------------|-------------------------------|-------------------------------|--------------------------------|-------------------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| each day | in thousa | nds | | | | | | | |
| 1,070 | 935 | 674 | 852 | 1,685 | 731 | 710 | 960 | 1,052 | 887 |
| 104 588 39 247 92 | 135 423 31 231 116 | 54 162 62 271 125 | 60 376 53 239 124 | 108 1,110 98 264 106 | 60 278 72 197 123 | 60 125 48 354 124 | 52 412 54 323 120 | 101 443 35 350 123 | 93 280 93 288 134 |
| 642 | 633 | 422 | 516 | 1,059 | 474 | 469 | 598 | 681 | 540 |
| 349 164 128 | 277 171 185 | 99 156 168 | 219 149 148 | 674 184 201 | 166 156 151 | 75 274 120 | 250 261 87 | 260 274 147 | 176 213 151 |
| 429 | 302 | 252 | 336 | 626 | 257 | 241 | 361 | 371 | 348 |
| 239 83 107 | 146 59 97 | 63 115 74 | 157 90 89 | 436 80 111 | 112 41 104 | 49 80 112 | 162 61 139 | 183 77 112 | 104 76 169 |

Table 15. Days lost from school associated with acute conditions per Pata are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|--|--------------------|----------------------------|----------------------------|---------------------------|--------------------|--------------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | | 33 | Days | lost from |
| All acute conditions | (*) | 143,956 | 71,530 | 49,232 | (*) | 43,210 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 6,525 126,327 11,105 | 14,578 46,457 10,495 | 20,884 19,407 8,941 | (*) (*) (*) | 8,711 27,438 7,061 |
| Male | | 1 | · | | | |
| All acute conditions | (*) | 68,763 | 35,524 | 22,389 | (*) | 22,146 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 2,719 60,815 5,230 | 6,592 23,265 5,667 | 9,831 7,999 4,558 | (*) (*) (*) | 5,754 11,842 4,550 |
| Female | | | | | | : |
| All acute conditions | (*) | 75,193 | 36,006 | 26,844 | (*) | 21,063 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 3,806 65,512 5,875 | 7,986 23,192 4,828 | 11,053 11,407 4,383 | (*) (*) (*) | 2,956 15,595 2,511 |

Table 16. Days lost from school associated with acute conditions per 100 chil-Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | i7 | | 1 | 958 | |
|--|--------------------|-----------------------|-----------------------|----------------------|--------------------|----------------------|
| Sex and condition group | July- September | October- December | January- March | April- June | July- September | October- December |
| Both sexes | | | - | Days lost | from schoo | ol per 100 |
| All acute conditions | (*) | 417.3 | 205.3 | 136.1 | (*) | 120.6 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 18.9 366.2 32.2 | 41.8 133.3 30.1 | 57.7 53.6 24.7 | (*) (*) (*) | 24.3 76.6 19.7 |
| <u>Male</u> | | | | | | |
| All acute conditions | (*) | 391.1 | 200.1 | 124.9 | (*) | 121.2 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 15.5 345.9 29.7 | 37.1 131.0 31.9 | 54.8 44.6 25.4 | (*) (*) (*) | 31.5 64.8 24.9 |
| <u>Female</u> | | | | | | |
| All acute conditions | (*) | 444.4 | 210.8 | 147.1 | (*) | 120.0 |
| Infective and parasitic diseases Respiratory conditions All other acute conditions | (*) (*) (*) | 22.5 387.2 34.7 | 46.7 135.8 28.3 | 60.6 62.5 24.0 | (*) (*) (*) | 16.8 88.8 14.3 |

quarter by sex and condition group: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | | 1961 | |
|---------------------------|----------------------------|--------------------|---------------------------|----------------------------|----------------------------|--------------------|--------------------------|---------------------------|---------------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| school in | thousand | ls | | | | | | | <u> </u> |
| 75,365 | 67 <u>,</u> 072 | (*) | 49,659 | 81,741 | 41,454 | (*) | 48,822 | 63,910 | 44,182 |
| 21,523 43,988 9,854 | 18,206 37,076 11,790 | (*) (*) (*) | 10,743 29,545 9,370 | 15,679 55,251 10,811 | 14,988 15,155 11,311 | (*) (*) (*) | 9,340 30,462 9,020 | 19,521 35,867 8,522 | 17,705 16,767 9,710 |
| 39,822 | 30,607 | (*) | 24,356 | 38,115 | 21,060 | (*) | 25,708 | 33,770 | 24,000 |
| 14,384 20,800 4,638 | 9,129 15,520 5,958 | (*) (*) (*) | 5,125 14,944 4,288 | 7,933 24,139 6,043 | 5,517 8,070 7,472 | (*) (*) (*) | 5,590 13,877 6,241 | 11,928 17,331 4,511 | 10,003 8,889 5,108 |
| 35,544 | 36,465 | (*) | 25,302 | 43,625 | 20,395 | (*) | 23,114 | 30,140 | 20,183 |
| 7,139 23,188 5,216 | 9,077 21,557 5,832 | (*) (*) (*) | 5,618 14,601 5,083 | 7,746 31,111 4,768 | 9,471 7,085 3,838 | (*) (*) (*) | 3,751 16,584 2,779 | 7,593 18,536 4,011 | 7,703 7,878 4,602 |

dren per quarter by sex and condition group: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix II. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|-----------------------|-----------------------|--------------------|----------------------|-----------------------|----------------------|--------------------|----------------------|----------------------|----------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| children | aged 6-16 | per quarte | r | | | | | | |
| 208.3 | 184.3 | (*) | 134.4 | 218.3 | 110.1 | (*) | 127.1 | 165.6 | 113.7 |
| 59.5 121.6 27.2 | 50.0 101.9 32.4 | (*) (*) (*) | 29.1 80.0 25.4 | 41.9 147.5 28.9 | 39.8 40.2 30.0 | (*) (*) (*) | 24.3 79.3 23.5 | 50.6 92.9 22.1 | 45.6 43.1 25.0 |
| 216.0 | 164.8 | (*) | 129.3 | 199.9 | 109.0 | (*) | 131.9 | 170.9 | 121.1 |
| 78.0 112.8 25.2 | 49.1 83.5 32.1 | (*) (*) (*) | 27.2 79.3 22.8 | 41.6 126.6 31.7 | 28.5 41.8 38.7 | (*) (*) (*) | 28.7 71.2 32.0 | 60.4 87.7 22.8 | 50.5 44.9 25.8 |
| 200.2 | 204.8 | (*) | 139.8 | 237.4 | 111.3 | (*) | 122.2 | 159.9 | 105.9 |
| 40.2 130.6 29.4 | 51.0 121.0 32.7 | (*) (*) (*) | 31.0 80.7 28.1 | 42.1 169.3 25.9 | 51.7 38.7 20.9 | (*) (*) (*) | 19.8 87.7 14.7 | 40.3 98.4 21.3 | 40.4 41.3 24.2 |

Table 17. Incidence of acute respiratory conditions Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|----------|--|---|--|--|--|---|
| Age | July- September | October- December | January- March | April- June | July- September | October- December |
| 1 | | | | Nu | mber of acu | te condi- |
| All ages | 30,612 | 126,340 | 88,727 | 38,854 | 25,458 | 59,246 |
| Under 5 | 5,689 6,708 4,678 7,873 3,988 1,676 | 21,030 38,174 17,079 27,774 16,989 5,293 | | 2,782 | 5,436 6,125 3,433 6,002 3,020 1,442 | 12,206 15,900 6,357 12,802 8,704 3,277 |
| 411 | 10.2 | 1 75 0 | l soci | 22.9 | • | |
| Under 5 | 29.6 20.4 22.6 17.2 11.6 11.6 | 75.2 108.7 115.1 81.4 60.8 49.4 36.5 | 52.6 86.4 64.9 43.9 45.8 41.8 38.2 | 41.2 27.5 17.4 20.7 16.3 19.0 | 27.8 18.1 15.9 13.2 8.7 9.8 | 34.7 62.1 46.5 29.1 28.1 24.9 22.2 |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

Table 18. Days of restricted activity associated with acute respiratory Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|----------|---|---|--|--|---|--|
| Age | July- September | October- December | January- March | April- June | July- September | October- December |
| | | | | Da | ys of restr | icted ac- |
| All ages | 100,721 | 584,101 | 339,869 | 147,355 | 65,776 | 183,662 |
| Under 5 | 13,891 21,711 11,891 24,973 20,293 7,960 | 69,945 193,606 68,926 111,242 101,776 38,606 | | 11,439 28,255 29,366 | 13,337 19,153 6,993 13,025 8,331 4,938 | 32,057 43,549 18,909 36,079 30,160 22,909 |
| | <u> </u> | | | | Days per | 100 per- |
| All ages | 60.3 | 347.6 | 201.3 | 87.0 | 38.7 | 107.4 |
| Under 5 | 72.4 65.9 57.5 54.7 59.3 55.3 | 361.5 583.7 328.6 243.6 295.8 266.3 | 244.8 228.5 129.4 175.0 211.7 244.0 | 122.4 96.1 53.3 61.9 84.7 151.8 | 68.3 56.5 32.3 28.6 23.9 33.6 | 163.2 127.4 86.5 79.3 86.2 154.9 |

per quarter by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II]

| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
|---|--|--|---|---|--|--|---|--|--|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| tions in thousands | | | | | | | | | |
| 78,101 | 52,543 | 22,137 | 57,216 | 93,656 | 34,896 | 24,866 | 63,576 | 71,075 | 36,759 |
| 14,857 21,438 8,248 17,879 10,609 5,070 per 100 p | | 5,821 5,339 2,113 4,857 2,728 1,278 | 13,558 16,190 6,402 11,969 6,894 2,204 | 15,436 22,707 10,472 22,638 15,905 6,499 | 9,291 8,254 3,787 6,318 5,202 2,044 | 6,758 5,686 3,211 5,930 2,264 1,016 | 12,680 17,074 7,925 15,360 7,188 3,350 | 16,087 18,269 9,444 12,788 10,315 4,173 | 7,419 9,415 4,086 8,188 5,529 2,123 |
| 45.5 | 30.5 | 12.8 | 32.9 | 53.4 | 19.8 | 14.1 | 35.8 | 39.8 | 20.5 |
| 75.4 62.2 37.4 39.3 30.2 34.1 | 45.4 48.3 26.1 25.1 20.6 15.5 | 29.4 15.2 9.4 10.7 7.7 8.5 | 68.1 45.8 28.2 26.4 19.4 14.6 | 77.1 63.3 45.6 49.8 44.4 42.7 | 46.4 22.8 16.4 13.9 14.5 13.4 | 33.7 15.6 13.8 13.1 6.3 6.6 | 62.9 46.3 33.8 33.8 19.8 21.6 | 79.6 49.1 39.9 28.1 28.4 26.8 | 36.7 25.1 17.2 18.0 15.1 13.6 |

conditions per quarter by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | | · · · · · · · · · · · · · · · · · · · | | | | | | | |
|--|---|--|--|--|--|---|--|--|---|
| | 1 | 959 | | | 1 | 960 | | 196 | 1 |
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| tivity in | thousand | is | | | <u> </u> | | | | |
| 260,941 | 189,693 | 68,870 | 169,165 | 402,446 | 114,803 | 69,265 | 187,019 | 250,753 | 124,350 |
| 46,353 70,246 20,825 47,111 48,525 27,880 sons per | 25,749 61,582 14,887 34,895 35,187 17,393 quarter | 15,018 16,226 5,055 13,012 12,630 6,930 | 35,456 48,429 15,046 32,911 25,800 11,524 | 43,319 89,775 36,746 87,233 93,908 51,464 | 21,768 25,771 11,128 20,389 21,280 14,467 | 14,085 14,531 8,594 17,532 8,006 6,517 | 31,007 49,237 19,951 41,872 27,797 17,154 | 49,460 62,828 23,105 42,366 40,886 32,108 | 24,739 22,703 |
| 152.0 | 110.0 | 39.8 | 97.3 | 229.5 | 65.2 | 39.2 | 105.3 | 140.5 | 69.4 |
| 235.4 203.8 94.4 103.6 138.1 187.6 | 130.6 177.1 66.9 76.8 99.8 116.4 | 75.9 46.3 22.5 28.6 35.7 46.1 | 178.2 136.9 66.3 72.7 72.6 76.2 | 216.4 250.1 160.1 191.8 262.3 338.0 | 108.8 71.2 48.1 44.9 59.2 94.7 | 70.3 39.8 36.9 38.6 22.2 42.4 | 153.8 133.5 85.1 92.2 76.7 110.9 | 244.8 168.9 97.7 93.2 112.4 206.5 | 93.7 74.1 49.7 54.4 62.2 117.3 |

Table 19. Days of bed disability associated with acute respiratory Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|-----------------|-----------|----------|----------|--------|-------------|-----------|
| Age | July- | October- | January- | April- | July- | October- |
| | September | December | March | June | September | December |
| | | | | | Days of | bed disa- |
| All ages | 42,400 | 312,251 | 173,929 | 64,521 | 33,474 | 79,975 |
| Under 55-14 | 6,634 | 32,008 | 23,262 | 11,299 | 7,441 | 10,918 |
| | 10,043 | 112,966 | 39,312 | 14,665 | 8,706 | 22,533 |
| 15-24 | 6,806 | 41,879 | 14,121 | 7,024 | 3,265 | 8,250 |
| 25-44 | 10,301 | 55,364 | 40,323 | 12,587 | 7,776 | 15,114 |
| 45-64 | 6,888 | 52,522 | 37,777 | 13,712 | 3,441 | 11,872 |
| 65+ | 1,729 | 17,512 | 19,133 | 5,234 | 2,847 | 11,289 |
| | | | • | Days o | f bed disab | ility per |
| All ages | 25.4 | 185.8 | 103.0 | 38.1 | 19.7 | 46.8 |
| Under 55-14 | 34.6 | 165.5 | 119.8 | 58.1 | 38.1 | 55.6 |
| | 30.5 | 340.6 | 117.7 | 43.6 | 25.7 | 65.9 |
| 15-24 | 32.9 | 199.7 | 66.5 | 32.7 | 15.1 | 37.7 |
| 25-44 | 22.6 | 121.2 | 88.3 | 27.6 | 17.1 | 33.2 |
| 45-64 | 20.1 | 152.7 | 109.4 | 39.5 | 9.9 | 33.9 |
| 65 + | 12.0 | 120.8 | 131.5 | 35.8 | 19.4 | 76.3 |

Table 20. Incidence of injuries per quarter Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | <u></u> | 1 | 958 | |
|----------|-------------------------|-------------------------|-------------------|-------------------------|-------------------------|-------------------------|
| Age | July- September | October- December | January- March | April- June | July- September | October- December |
| | | | | | | er of in- |
| All ages | 13,910 | 11,123 | 10,192 | 12,383 | 15,480 | 11,063 |
| Under 15 | 5,008 5,289 3,613 | 3,134 4,852 3,137 | 3,950 | 4,602 4,709 3,072 | 5,755 6,144 3,581 | 3,067 4,792 3,204 |
| | | | | Nu | mber of inj | uries per |
| All ages | 8.3 | 6.6 | 6.0 | 7.3 | 9.1 | 6.5 |
| Under 15 | 9.6 8.0 7.4 | 6.0 7.3 6.4 | 6.7 5.9 5.5 | 8.7 7.0 6.2 | 10.8 9.1 7.2 | 5.7 7.1 6.4 |

Excluded from these statistics are all conditions involving neither restricted activity nor medical attention.

conditions per quarter by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

| | 1 | .959 | | | 1 | 960 | | 1961 | |
|-----------|-----------|-----------|----------|----------|--------|-----------|----------|----------|--------|
| January- | April- | July- | October- | January- | April- | July- | October- | January- | April- |
| March | June | September | December | March | June | September | December | March | June |
| bility in | thousand | ls | | | | | | | |
| 117,505 | 95,069 | 26,756 | 70,753 | 195,607 | 51,617 | 26,041 | 76,329 | 111,196 | 54,379 |
| 18,795 | 12,492 | 5,508 | 13,244 | 18,744 | 9,943 | 5,761 | 10,658 | 20,028 | 8,479 |
| 34,994 | 33,192 | 6,228 | 21,553 | 45,847 | 12,704 | 6,073 | 24,882 | 27,288 | 11,638 |
| 12,081 | 8,088 | 2,868 | 7,092 | 21,315 | 5,837 | 2,138 | 10,343 | 11,916 | 6,409 |
| 20,389 | 17,804 | 5,269 | 15,202 | 41,723 | 7,913 | 6,680 | 16,768 | 20,808 | 11,328 |
| 19,199 | 16,116 | 4,905 | 9,537 | 47,614 | 8,379 | 3,370 | 10,061 | 17,727 | 8,871 |
| 12,048 | 7,377 | 1,977 | 4,126 | 20,363 | 6,842 | 2,018 | 3,617 | 13,429 | 7,654 |
| 100 perso | ns per qu | arter | | | | | | | |
| 68.4 | 55.1 | 15.5 | 40.7 | 111.5 | 29.3 | 14.7 | 43.0 | 62.3 | 30.4 |
| 95.4 | 63.4 | 27.8 | 66.6 | 93.6 | 49.7 | 28.8 | 52.9 | 99.1 | 42.0 |
| 101.5 | 95.5 | 17.8 | 60.9 | 127.7 | 35.1 | 16.6 | 67.5 | 73.3 | 31.0 |
| 54.8 | 36.4 | 12.8 | 31.3 | 92.9 | 25.2 | 9.2 | 44.1 | 50.4 | 26.9 |
| 44.8 | 39.2 | 11.6 | 33.6 | 91.8 | 17.4 | 14.7 | 36.9 | 45.8 | 24.9 |
| 54.7 | 45.7 | 13.8 | 26.8 | 133.0 | 23.3 | 9.3 | 27.8 | 48.7 | 24.3 |
| 81.0 | 49.4 | 13.2 | 27.3 | 133.7 | 44.8 | 13.1 | 23.4 | 86.4 | 49.0 |

by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix 1. Definitions of terms are given in Appendix II

| | 1 | .959 | | | 1 | .960 | | 1961 | |
|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| juries in thousands | | | | | | | | | |
| 9,986 | 12,941 | 13,751 | 9,992 | 9,927 | 12,017 | 15,434 | 12,078 | 10,066 | 12,129 |
| 3,432 4,231 2,322 | 5,183 4,988 2,771 | 5,167 5,900 2,684 | 3,844 3,597 2,550 | 3,250 4,112 2,565 | 4,996 4,229 2,792 | 5,952 5,933 3,549 | 4,785 3,918 3,376 | 3,188 4,342 2,536 | 4,841 4,348 2,940 |
| 100 perso | ns per qu | arter | | | | | | | |
| 5.8 | 7.5 | 7.9 | 5.7 | 5.7 | 6.8 | 8.7 | 6.8 | , 5.6 | 6.8 |
| 6.3 6.3 4.6 | 9.5 7.4 5.5 | 9.4 8.7 5.3 | 7.0 5.3 5.0 | 5.8 6.0 5.0 | 8.9 6.2 5.5 | 10.5 8.6 6.9 | 8.4 5.7 6.5 | 5.6 6.3 4.9 | 8.4 6.3 5.6 |

Table 21. Days of restricted activity associated with in-Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 57 | | 1 | .958 | |
|--------------------------|----------------------------|----------------------------|---------------------------|---------------------------|----------------------------|---------------------------|
| Age | July- September | October- December | January- March | April- June | July- September | October- December |
| | = | | | Da | ys of restr | icted ac- |
| All ages | 69,144 | 59,087 | 61,637 | 56,985 | 91,038 | 61,320 |
| Under 15 | 10,763 31,992 26,389 | 11,477 22,751 24,860 | 7,906 24,619 29,111 | 9,664 21,522 25,799 | 19,304 42,771 28,963 | 8,515 24,250 28,555 |
| | | | | Days of | restricted | activity |
| All ages | 41.4 | 35.2 | 36.5 | 33.6 | 53.5 | 35.9 |
| Under 15 15-44 45+ | 20.6 48.2 54.2 | 21.9 34.1 50.8 | 15.0 36.8 59.3 | 18.2 32.1 52.3 | 36.2 63.6 58.5 | 15.8 36.0 57.4 |

Table 22. Days of bed disability associated with injuries Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|-------------------|-------------------------|-------------------------|-------------------------|-------------------------|---------------------------|--------------------------|
| Age | July- September | October- December | January- March | April- June | July- September | October December |
| | | | | | Days of | bed disa- |
| All ages | 19,917 | 16,429 | 20,206 | 15,695 | 26,790 | 22,046 |
| Under 15 15-44 | 3,497 7,853 8,567 | 1,976 6,807 7,646 | 2,296 8,348 9,562 | 2,086 5,050 8,559 | 4,013 10,904 11,873 | 1,907 8,173 11,966 |
| | | | • | Da | ys of bed d | isability |
| All ages | 11.9 | 9.8 | 12.0 | 9.3 | 15.7 | 12.9 |
| Under 15 | 6.7 11.8 17.6 | 3.8 10.2 15.6 | 4.3 12.5 19.5 | 3.9 7.5 17.4 | 7.5 16.2 24.0 | 3.5 12.1 24.0 |

juries per quarter by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | 1 | 959 | | | 1 | 960 | | 1961 | |
|---------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| tivity in thousands | | | | | | | | | |
| 59,985 | 58,219 | 63,844 | 59,720 | 70,166 | 70,784 | 79,126 | 81,340 | 96,540 | 74,604 |
| 7,397 20,517 32,070 | 13,579 18,593 26,046 | 13,965 24,962 24,916 | 11,827 18,494 29,399 | 11,770 27,854 30,541 | 16,181 27,067 27,536 | 13,893 32,430 32,804 | 14,429 30,279 36,632 | 14,869 40,605 41,066 | 11,172 38,246 25,186 |
| per 100 p | ersons pe | r quarter | | | | • | | | |
| 34.9 | 33.8 | 36.9 | 34.3 | 40.0 | 40.2 | 44.8 | 45.8 | 54.1 | 41.6 |
| 13.7 30.4 64.2 | 24.9 27.5 51.9 | 25.5 36.8 49.4 | 21.4 27.2 58.0 | 21.0 40.7 59.8 | 28.8 39.5 53.8 | 24.6 47.2 63.8 | 25.3 44.0 70.9 | 25.9 58.8 79.1 | 19.4 55.2 48.3 |

per quarter by age: United States, July 1957-June 1961

tions and information on the reliability of the estimates are given in Appendix I. Definitions of terms are given in Appendix II

| | 1 | 959 | ====== | | 1 | 960 | | 1961 | |
|-------------------------|--------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|--------------------------|-------------------------|
| January- March | April- June | July- September | October- December | January- March | April- June | July- September | October- December | January- March | April- June |
| bility in thousands | | | | | | | | | |
| 17,662 | 17,263 | 19,402 | 14,525 | 18,042 | 18,881 | 14,315 | 19,578 | 29,236 | 20,374 |
| 2,449 5,795 9,418 | ·3,578 4,783 8,902 | 3,854 6,519 9,029 | 3,412 2,216 8,897 | 2,496 7,142 8,404 | 4,207 6,554 8,120 | 1,955 6,172 6,189 | 4,166 6,860 8,553 | 6,991 8,677 13,568 | 4,857 9,425 6,092 |
| per 100 p | ersons pe | r quarter | | | | | | | |
| 10.3 | 10.0 | 11.2 | 8.4 | 10.3 | 10.7 | 8.1 | 11.0 | 16.4 | 11.4 |
| 4.5 8.6 18.8 | 6.6 7.1 17.7 | 7.0 9.6 17.9 | 6.2 3.3 17.6 | 4.5 10.4 16.5 | 7.5 9.6 15.9 | 3.5 9.0 12.0 | 7.3 10.0 16.5 | 12.2 12.6 26.1 | 8.4 13.6 11.7 |

Table 23. Population used in obtaining rates shown in this publica-Data are based on household interviews of the civilian, noninstitutional population. The survey design, general qualifica-

| | 195 | 7 | | 1 | 958 | |
|---------------|-----------|----------|----------|---------|-----------|-----------|
| Sex and age | July- | October- | January- | April- | July- | October- |
| | September | December | March | June | September | December |
| Both sexes | | | | | P | opulation |
| All ages | 167,111 | 168,048 | 168,807 | 169,448 | 170,135 | 170,942 |
| Under 15 | 52,122 | 52,512 | 52,823 | 53,071 | 53,397 | 53,814 |
| 0-4 | 19,189 | 19,346 | 19,425 | 19,441 | 19,519 | 19,643 |
| 5-14 | 32,933 | 33,166 | 33,398 | 33,630 | 33,878 | 34,171 |
| 15-44 | 66,345 | 66,638 | 66,901 | 67,089 | 67,224 | 67,355 |
| 15-24 | 20,696 | 20,976 | 21,239 | 21,453 | 21,650 | 21,856 |
| 25-44 | 45,649 | 45,662 | 45,662 | 45,636 | 45,574 | 45,499 |
| 45+ | 48,644 | 48,898 | 49,084 | 49,288 | 49,515 | 49,773 |
| 45-64 | 34,246 | 34,402 | 34,539 | 34,682 | 34,830 | 34,982 |
| 65+ | 14,398 | 14,496 | 14,545 | 14,606 | 14,684 | 14,791 |
| Male | | | | | | |
| All ages | 81,243 | 81,749 | 82,149 | 82,451 | 82,781 | 83,177 |
| Under 55-14 | 9,774 | 9,855 | 9,895 | 9,903 | 9,943 | 10,006 |
| | 16,800 | 16,920 | 17,040 | 17,160 | 17,287 | 17,438 |
| 15-24 | 9,559 | 9,736 | 9,897 | 10,009 | 10,120 | 10,248 |
| 25-44 | 21,855 | 21,884 | 21,901 | 21,891 | 21,859 | 21,812 |
| 45-64 | 16,647 | 16,713 | 16,766 | 16,823 | 16,885 | 16,948 |
| 65+ | 6,606 | 6,641 | 6,650 | 6,665 | 6,688 | 6,726 |
| <u>Female</u> | | | | | | |
| All ages | 85,868 | 86,299 | 86,658 | 86,996 | 87,354 | 87,764 |
| Under 55-14 | 9,414 | 9,491 | 9,530 | 9,538 | 9,577 | 9,637 |
| | 16,133 | 16,246 | 16,358 | 16,471 | 16,591 | 16,733 |
| 15-24 | 11,137 | 11,240 | 11,342 | 11,444 | 11,530 | 11,608 |
| 25-44 | 23,794 | 23,778 | 23,761 | 23,745 | 23,715 | 23,687 |
| 45-64 | 17,599 | 17,689 | 17,773 | 17,859 | 17,946 | 18,034 |
| 65+ | 7,792 | 7,855 | 7,894 | 7,940 | 7,996 | 8,065 |
| Children 6-16 | | | | | | |
| Both sexes | 34,162 | 34,501 | 34,840 | 36,178 | 35,505 | 35,824 |
| MaleFemale | 17,407 | 17,582 | 17,757 | 17,932 | 18,100 | 18,265 |
| | 16,755 | 16,919 | 17,083 | 18,247 | 17,404 | 17,559 |

NOTE: For official population estimates for more general use, see Bureau of the Census reports on the civilian population of the United States,

tion by quarter, sex, and age: United States, July 1957-June 1961 tions and information on the reliability of the estimates are given in Appendix II

| | 1959 | | | | 1 | 960 | | 1961 | |
|-----------|----------|-----------|----------|----------|---------|-----------|----------|----------|---------|
| January- | April- | July- | October- | January- | April- | July- | October- | January- | April- |
| March | June | September | December | March | June | September | December | March | June |
| in thousa | ınds | . • | | | | · . | | | |
| 171,676 | 172,389 | 173,136 | 173,926 | 175,361 | 175,992 | 176,727 | 177,601 | 178,417 | 179,134 |
| 54,162 | 54,481 | 54,846 | 55,277 | 55,915 | 56,219 | 56,560 | 57,027 | 57,408 | 57,733 |
| 19,695 | 19,718 | 19,779 | 19,896 | 20,022 | 20,007 | 20,027 | 20,158 | 20,201 | 20,201 |
| 34,467 | 34,763 | 35,067 | 35,382 | 35,892 | 36,212 | 36,533 | 36,869 | 37,207 | 37,533 |
| 67,524 | 67,700 | 67,847 | 67,987 | 68,417 | 68,565 | 68,725 | 68,871 | 69,089 | 69,261 |
| 22,051 | 22,250 | 22,423 | 22,690 | 22,946 | 23,134 | 23,299 | 23,446 | 23,642 | 23,822 |
| 45,473 | 45,450 | 45,424 | 45,297 | 45,471 | 45,431 | 45,426 | 45,425 | 45,447 | 45,439 |
| 49,991 | 50,209 | 50,442 | 50,662 | 51,030 | 51,208 | 51,442 | 51,703 | 51,919 | 52,140 |
| 35,126 | 35,271 | 35,418 | 35,545 | 35,805 | 35,933 | 36,076 | 36,229 | 36,371 | 36,503 |
| 14,865 | 14,938 | 15,024 | 15,117 | 15,225 | 15,275 | 15,366 | 15,474 | 15,548 | 15,636 |
| | - | r | , | • | | | | | |
| 83,545 | 83,904 | 84,268 | 84,638 | 85,340 | 85,639 | 85,999 | 86,386 | 86,763 | 87,112 |
| 10,032 | 10,043 | 10,074 | 10,131 | 10,196 | 10,187 | 10,197 | 10,264 | 10,282 | 10,285 |
| 17,590 | 17,742 | 17,898 | 18,060 | 18,322 | 18,486 | 18,651 | 18,822 | 18,993 | 19,159 |
| 10,361 | 10,476 | 10,571 | 10,762 | 10,894 | 11,009 | 11,100 | 11,158 | 11,260 | 11,363 |
| 21,809 | 21,808 | 21,802 | 21,684 | 21,764 | 21,738 | 21,745 | 21,736 | 21,746 | 21,745 |
| 17,007 | 17,067 | 17,128 | 17,169 | 17,292 | 17,340 | 17,399 | 17,462 | | 17,569 |
| 6,747 | ,6,767 | 6,794 | 6,831 | 6,872 | 6,879 | 6,908 | 6,945 | | 6,990 |
| 88,131 | 88,485 | 88,867 | 89,288 | 90,021 | 90,353 | 90,728 | 91,214 | 91,654 | 92,022 |
| 9,663 | 9,675 | 9,705 | 9,764 | 9,826 | 9,820 | 9,830 | 9,894 | 9,919 | 9,916 |
| 16,877 | 17,021 | 17,169 | 17,321 | 17,570 | 17,726 | 17,882 | 18,047 | 18,213 | 18,373 |
| 11,690 | 11,773 | 11,852 | 11,928 | 12,052 | 12,126 | 12,199 | 12,288 | 12,383 | 12,459 |
| 23,664 | - 23,642 | 23,622 | 23,613 | 23,707 | 23,693 | 23,681 | 23,689 | 23,701 | 23,694 |
| 18,119 | 18,204 | 18,290 | 18,377 | 18,513 | 18,593 | 18,677 | 18,766 | 18,853 | 18,935 |
| 8,118 | 8,170 | 8,230 | 8,285 | 8,353 | 8,396 | 8,458 | 8,530 | 8,585 | 8,646 |
| 36 100 | 36 300 | 24 700 | 36 044 | 37 /50 | 27 450 | 20 020 | 30 /03 | 20 500 | 20 047 |
| 36,188 | 36,386 | ,36,789 | 36,944 | 37,450 | 37,658 | 38,028 | 38,403 | | 38,867 |
| 18,433 | 18,577 | 18,796 | 18,842 | 19,071 | 19,329 | 19,383 | 19,486 | 19,756 | 19,815 |
| 17,756 | 17,809 | 17,993 | 18,101 | 18,380 | 18,329 | 18,645 | 18,917 | 18,844 | 19,052 |

in Current Population Reports: Series P-20, P-25 and P-60.

APPENDIX I

TECHNICAL NOTES ON METHODS

Background of This Report

This report, <u>Acute Conditions</u>; <u>Seasonal Variations</u>, is one of a series of statistical reports prepared by the U. S. National Health Survey. It is based on information collected in a continuing nationwide sample of households in the Health Interview Survey, which is one of the major parts of the program.

The Health Interview Survey utilizes a questionnaire which, in addition to personal and demographic characteristics, obtains information on illnesses, injuries, chronic conditions and impairments, health insurance coverage, and other health topics. As data relating to each of these various broad topics are tabulated and analyzed, separate reports are issued which cover one or more of the specific topics. The present report is based on the consolidated sample for 208 weeks of interviewing ending June 1961.

The population covered by the sample for the Health Interview Survey is the civilian, noninstitutional population of the United States living at the time of the interview. The sample does not include members of the Armed Forces, U. S. nationals living in foreign countries, or crews of vessels. It should also be noted that the estimates shown do not represent a complete inventory of acute conditions for the specified calendar period since no adjustment has been made for persons who incurred acute conditions during the two-week-recall period but who died prior to the interview.

Statistical Design of the Health Interview Survey

General plan.—The sampling plan of the survey follows a multistage probability design which permits a continuous sampling of the civilian population of the United States. The first stage of this design consists of drawing a sample of 500 from the 1,900 geographically defined Primary Sampling Units (PSU's) into which the United States has been divided. A PSU is a county, a group of contiguous counties, or a Standard Metropolitan Statistical Area.

With no loss in general understanding, the remaining stages can be telescoped and treated in this discussion as an ultimate stage. Within PSU's, then, ultimate stage units called segments are defined, also geographically, in such a manner that each segment contains an expected six households in the sample. Each week a random sample of about 120 segments is drawn. In the approximately 700 households in those segments, household members are interviewed concerning factors related to health.

Since the household members interviewed each week are a representative sample of the population, samples for successive weeks can be combined into

larger samples. Thus the design permits both continuous measurement of characteristics of high incidence or prevalence in the population, and through the larger consolidated samples, more detailed analysis of less common characteristics and smaller categories. The continuous collection has administrative and operational advantages as well as technical assets, since it permits field work to be handled with an experienced, stable staff.

Sample size and geographic detail.—The sample plan for each 13-week quarter from July 1957-June 1961 included about 30,000 persons from 9,000 households. Over the 12-month period ending June 1961, the sample included approximately 125,000 persons from 38,000 households in 6,400 segments. The over-all sample was designed in such a fashion that tabulations can be provided for each of the major geographic regions and for urban and rural sectors of the United States.

Collection of data.—The field operations for the household survey are performed by the Bureau of the Census under specifications established by the Public Health Service. In accordance with these specifications the Bureau of the Census designs and selects the sample; conducts the field interviewing, acting as the collecting agent for the Public Health Service; and edits and codes the questionnaires. Tabulations are prepared by the Public Health Service using the Bureau of the Census electronic computers.

Estimating methods.—Each statistic produced by the survey—for example, the number of work-loss days occurring in a specified period—is the result of two stages of ratio estimation. In the first of these, the factor is the ratio of the 1950 decennial population count to the 1950 estimated population in the U. S. National Health Survey's first-stage sample of PSU's. These factors are applied for some 50 color-residence classes.

Later, ratios of sample-produced estimates of the population to official Bureau of the Census figures for current population in about 60 age-sex-color classes are computed, and serve as second-stage factors for ratio estimating.

The effect of the ratio estimating process is to make the sample more closely representative of the population by age, sex, color, and residence, thus reducing sampling variance.

As noted, each week's sample represents the population living during that week and characteristics of that population. Consolidation of samples over a time period, say a calendar quarter, produces estimates of average characteristics of the U. S. population for that calendar quarter. Similarly, population data for a year are averages of the four quarterly figures.

For statistics measuring the number of occurrences during a specified time period, such as the number of bed-disability days, or incidence of acute conditions, a similar computational procedure is used, but the statistics have a different interpretation. For these items, the questionnaire asks for the respondent's experience over the two calendar weeks prior to the week of interview. In such instances the estimated quarterly total for the statistic is simply 6.5 times the average two-week estimate produced by the 13 successive samples taken during the period. The annual total is the sum of the four quarters. Thus, the experience of persons interviewed during a year-experience which actually occurred for each person in a twocalendar-week interval prior to week of interview-is treated as though it measured the total of such experience during the year. Such interpretation leads to no significant bias.

General Qualifications

Nonresponse.—Data were adjusted for nonresponse by a procedure which imputes to persons in a household which was not interviewed the characteristics of persons in households in the same segment which were interviewed. The total noninterview rate was 5 percent; 1 percent was refusal, and the remainder was primarily due to the failure to find any eligible household respondent after repeated trials.

The interview process.—The statistics presented in this report are based on replies secured in interviews of persons in the sampled households. Each person 18 years of age and over, available at the time of interview, was interviewed individually. Proxy respondents within the household were employed for children and for adults not available at the time of the interview, provided the respondent was closely related to the person about whom information was being obtained.

There are limitations to the accuracy of diagnostic and other information collected in household interviews. For diagnostic information, the household respondent can, at best, pass on to the interviewer only the information the physician has given to the family. For conditions not medically attended, diagnostic information is often no more than a description of symptoms. However, other facts, such as the number of disability days caused by the condition, can be obtained more accurately from household members than from any other source since only the persons concerned are in a position to report information of this type.

Rounding of numbers.—The original tabulations on which the data in this report are based show all estimates to the nearest whole unit. All consolidations were made from the original tabulations using the estimates to the nearest unit. In the final published tables the figures are rounded to the nearest thousand, although they are not necessarily accurate to that detail. Devised statistics, such as rates and percent distributions, are computed after the estimates on which they are based have been rounded to the nearest thousand.

Population figures.—Some of the published tables include population figures for specified categories. Except for certain over-all totals by age and sex, which are adjusted to independent estimates, these figures are based on the sample of households in the U. S. National Health Survey. They are given primarily for the purpose of providing denominators for rate computation,

and for this purpose are more appropriate for use with the accompanying measures of health characteristics than other population data that may be available. In some instances they will permit users to recombine published data into classes more suitable to their specific needs. With the exception of the over-all totals by age and sex, mentioned above, the population figures differ from corresponding figures (which are derived from different sources) published in reports of the Bureau of the Census. For population data for general use, see the official estimates presented in Bureau of the Census reports in the P-20, P-25, and P-60 series.

Reliability of Estimates

Since the estimates are based on a sample, they will differ somewhat from the figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and interviewing personnel and procedures. As in any survey, the results are also subject to measurement error.

The standard error is primarily a measure of sampling variability, that is, the variations that might occur by chance because only a sample of the population is surveyed. As calculated for this report, the standard error also reflects part of the variation which arises in the measurement process. It does not include estimates of any biases which might lie in the data. The chances are about 68 out of 100 that an estimate from the sample would differ from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference would be less than twice the standard error and about 99 out of 100 that it would be less than 2½ times as large.

The relative standard error of an estimate is obtained by dividing the standard error of the estimate by the estimate itself, and is expressed as a percentage of the estimate. Included in this Appendix are charts from which the relative standard errors can be determined for estimates shown in the report. In order to derive relative errors which would be applicable to a wide variety of health statistics and which could be prepared at a moderate cost, a number of approximations were required. As a result, the charts provide an estimate of the approximate relative standard error rather than the precise error for any specific aggregate or percentage.

Three classes of statistics for the health survey are identified for purposes of estimating variances.

Narrow range.—This class consists of (1) statistics which estimate a population attribute, e.g., the number of persons in a particular income group, and (2) statistics for which the measure for a single individual for the period of reference is usually either 0 or 1, on occasion may take on the value 2, and very rarely, 3.

Medium range.—This class consists of other statistics for which the measure for a single individual for the period of reference will rarely lie outside the range 0 to 5.

Wide range.—This class consists of statistics for which the measure for a single individual for the period of reference frequently will range from 0 to a number in excess of 5, e.g., the number of days of work loss experienced during the year.

In addition to classifying variables according to whether they are narrow-, medium-, or wide-range, statistics in the survey are further defined as:

<u>Type A.</u>—Statistics on prevalence, and incidence data for which the period of reference in the questionnaire is 12 months.

Type B.—Incidence-type statistics for which the period of reference in the questionnaire is two weeks.

Only the charts on sampling error applicable to data contained in this report are presented. Those shown are charts for aggregates based on one calendar quarter of data collection.

General rules for determining relative sampling errors.—The ''guide'' on page 49, together with the following rules, will enable the reader to determine approximate relative standard errors from the charts for estimates presented in this report.

Rule 1. Estimates of aggregates: Approximate relative standard errors of estimates of aggregates, such as the number of persons with a given characteristic, or the number of disability days are obtained from appropriate curves on pages 50 and 51. The number of persons in the total U.S. population or in an age-sex class of the total population is adjusted to official Bureau of the Census figures and is not subject to sampling error.


- Rule 2. Estimates of rates where the numerator is not a subclass of the denominator: This rule applies where a unit of the numerator often occurs more than once for any one unit in the denominator. For example, in the computation of the number of days of work loss per person per year, several of the days included in the numerator could be assigned to a person (one unit) in the denominator. Approximate relative standard errors for rates of this kind may be computed as follows:
 - (a) Where the denominator is the total U. S. population, or includes all persons in one or more of the age-sex groups of the total population, the relative error of the rate is equivalent to the relative error of the numerator which can be obtained directly from the appropriate chart.
 - (b) In other cases, obtain the relative standard error of the numerator and of the denominator from the appropriate curve. Square each of these relative errors, add the resulting values, and extract the square root of the sum. This procedure will result in an upper bound, and often will overstate the error.

Guide to Use of Relative Standard Error Charts

The code shown below identifies the appropriate curve to be used in estimating the relative standard error of the statistic described. The four components of each code describe the statistic as follows: (1)


A = aggregate, P = percentage; (2) the number of calendar quarters of data collection; (3) the type of the statistic as described on page 48; and (4) the range of the statistic as described on page 47.

| | | Use: | |
|--|----------|---|----------|
| Statistic | Rule | Code on | page |
| Number of: Acute conditions per quarter | 1 | Albn | 50 |
| Disability days per quarter | 1 , | A1BW | 51 |
| Persons in the U. S. population, or total number of persons in any age-sex category | Not subj | ect to sampling error | , 1 |
| Incidence rates for acute conditions: Per 100 total U. S. population or per 100 persons in any age-sex group of the total U. S. population | 2(a) | A1BN | , 50 |
| Number of disability days: Per case for acute conditions | 2(b) | Numer : A1BW Denom : A1BN | 51 50 |
| Average number of persons absent from work each day because of an acute condition | 1 | AlBW (with value for number of work days lost because of the condition read from the curve) | 51 |



Size of estimate (in thousands)

Example of use of chart: An aggregate of 6,000,000 (on scale at bottom of chart) for a Narrow range Type B statistic has a relative standard error of 19.3 percent, read from scale at left side of chart, or a standard error of 1,158,000 (19.3 percent of 6,000,000).



Example of use of chart: An aggregate of 20,000,000 (on scale at bottom of chart) for a wide range type B statistic has a relative standard error of 16.0 percent, read from scale at left side of chart, or a standard error of 3,200,000 (16.0 percent of 20,000,000).

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Terms Relating to Conditions

Condition.—A morbidity condition, or simply a condition, is any entry on the questionnaire which describes a departure from a state of physical or mental wellbeing. It results from a positive response to one of a series of "illness-recall" questions. In the coding and tabulating process, conditions are selected or classified according to a number of different criteria, such as, whether they were medically attended; whether they resulted in disability; whether they were acute or chronic; or according to the type of disease, injury, impairment, or symptoms reported. For the purposes of each published report or set of tables, only those conditions recorded on the questionnaire which satisfy certain stated criteria are included.

Acute conditions are classified by type according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the code more suitable for a household-interview-type survey.

Acute condition.—An acute condition is defined as a condition which has lasted less than three months and which has involved either medical attention or restricted activity. Because of the procedures used to estimate incidence, the acute conditions included in this report are the conditions which had their onset during the two weeks prior to the interview week and which involved either medical attention or restricted activity during that two-week period. However, it excludes certain conditions which are always classified as chronic (listed below) even though the onset occurred within three months.

Conditions always classified as chronic:

Asthma Hav fever Tuberculosis Chronic bronchitis Repeated attacks of sinus trouble Rheumatic fever Hardening of the arteries High blood pressure Heart trouble Stroke Trouble with varicose veins Hemorrhoids or piles Tumor, cyst, or growth Chronic gallbladder or liver trouble Stomach ulcer Any other chronic stomach trouble Kidney stones or chronic kidney trouble

Arthritis or rheumatism Mental illness Diabetes Thyroid trouble or goiter Any allergy Epilepsy Chronic nervous trouble Cancer Chronic skin trouble Hernia or rupture Prostate trouble Deafness or serious trouble with hearing Serious trouble with seeing, even when wearing glasses Cleft palate Any speech defect Missing fingers, hand, or arm-toes, foot, or leg Palsy

Paralysis of any kind Repeated trouble with back or spine Club foot Permanent stiffness or

V Other conditions

deformity of the foot, leg, fingers, arm, or back Condition present since birth

All other acute code

numbers

Condition groups.—Conditions are classified according to the International Classification of Diseases, 1955 Revision, with certain modifications adopted to make the code more suitable for a household interview survey. In this report, all tables which have data classified by type of condition employ a 5 category regrouping plus several selected subgroups. The International Classification code numbers included in each category are shown below.

| Condition Groups | International Classification Code Numbers |
|--|---|
| I Infective and parasits diseases | 020-138 |
| II Respiratory condition Upper respiratory Other respiratory | ns 470-501, 511, 514 517-525, 527, 783 470-475, 511, 514, 517 480-501, 518-525, 527 783 |
| III Digestive system conditions | 530-539, 543-553, 570 571, 573-587, 784, 785 |
| IV Injuries | N800-N885, N890-N895 N900-N994, N996-N999 |

Onset of condition.—A condition is considered to have had its onset when it was first noticed. This could be the time the person first felt sick or became injured, or it could be the time when the person or his family was first told by a physician that he had a condition of which he was previously unaware.

Incidence of conditions.—The incidence of conditions is the estimated number of conditions having their onset in a specified time period. As previously mentioned, minor acute conditions, involving neither restricted activity nor medical attention are excluded from the statistics. The incidence data shown in some reports are further limited to various subclasses of conditions, such as, "incidence of conditions involving bed disability."

Activity-restricting condition.—An activity restricting condition is a condition which has caused at least one day of restricted activity during the two calendar weeks before the interview week. (See def-

inition of "Restricted-activity day.") The incidence of acute activity-restricting conditions is estimated from the number of such conditions reported as having started in the two-week period, but a condition which did not result in restricted activity until after the end of the two-week period in which it had its onset is not included.

Bed-disabling condition.—A condition involving at least one day of bed disability is called a bed-disabling condition. (See definition of "Bed-disability day.") The incidence of acute bed-disabling conditions is defined in a manner analogous to the incidence of acute activity-restricting conditions.

Medically attended condition.—A condition is considered to be medically attended if a physician has been consulted about it either at its onset or at any time thereafter. Medical attention includes consultation either in person or by telephone for treatment or advice. Advice from the physician transmitted to the patient through the nurse is counted as well as visits to physicians in clinics or hospitals. If during the course of a single visit the physician is consulted about more than one condition for each of several patients, each condition of each patient is counted as medically attended.

Discussions of a child's condition by the physician and a responsible member of the household are considered as medical attention even if the child was not seen at that time.

For the purpose of this definition, the term ''physician'' includes doctors of medicine and osteopathic physicians.

Terms Relating to Disability

<u>Disability</u>.—Disability is the general term used to describe any temporaty or long-term reduction of a person's activity as a result of an acute or chronic condition.

Disability days are classified according to whether they are days of restricted activity, bed-days, workloss days, or school-loss days. All days of bed disability are, by definition, days of restricted activity. The converse form of this statement is, of course, not true. Days lost from work and days lost from school are also days of restricted activity for the working and school-age populations. Hence, restricted activity is the most inclusive term used in describing disability days.

Condition-days of restricted activity, bed disability, etc.—Condition-days of restricted activity, bed disability, and so forth are days of the various forms of disability associated with any one condition. Since any particular day of disability may be associated with more than one condition, the sum of days for all con-

ditions adds to more than the total number of persondays of disability.

Restricted-activity day.—A day of restricted activity is one on which a person substantially reduces the amount of activity normal for that day because of a specific illness or injury. The type of reduction varies with the age and occupation of the individual as well as with the day of the week or season of the year. Restricted activity covers the range from substantial reduction to complete inactivity for the entire day.

Bed-disability day.—A day of bed disability is one on which a person stays in bed for all or most of the day because of a specific illness or injury. It is considered to be a day only if the period of bed disability includes more than half of the daylight hours. All hospital days for inpatients are considered to be days of bed disability even if the patient was not actually in bed at the hospital.

Work-loss day.—A day lost from work is a normal working day on which a person did not work at his job or business because of a specific illness or injury. If the person's regular work day is less than a whole day and the entire work day was lost, it would be counted as a whole work day lost. The number of days lost from work is determined only for persons 17 years of age or over who reported that at any time during the two-week period covered by the interview they either worked at, or had a job or business.

School-loss day.—A day lost from school is a normal school day on which a child did not attend school because of a specific illness or injury. The number of days lost from school is determined only for children 6-16 years of age.

Demographic Terms

Age.—The age recorded for each person is the age at last birthday. Age is recorded in single years and grouped in a variety of distributions depending upon the purpose of the table.

Quarter.—The quarters used by the U. S. National Health Survey are actually 13-week periods rather than 3 calendar months. Since each 13-week period begins on a Monday and ends on a Sunday, the actual dates of the beginning and end of each 13-week period may overlap into another calendar quarter. Therefore, the time periods in the table headings are the approximate rather than the precise periods during which the interviewing was conducted.

In order to compute the number of persons absent from work each day, the assumption was made that there are 245 work days in the year or 61.25 work days in each quarter.

APPENDIX III

QUESTIONNAIRE

The items below show the exact content and wording of the basic questionnaire used in the nationwide household survey of the U. S. National Health Survey. The actual questionnaire is designed for a household as a unit and includes additional spaces for reports on more than one person, condition, accident or hospitalization. Such repetitive spaces are omitted in this illustration. CONFIDENTIAL - The National Health Survey is authorized by Public Law 652 of the 84th Congress (70 Stat 489; 42 U.S.C. 305). All information which would permit identification of the individual will be held strictly confidential, will be used only by persons engaged in and for the purposes of the survey, and will not be disclosed or released to others for any other purposes (22 FR 1687). U.S. DEPARTMENT OF COMMERCE BUREAU OF THE CENSUS ACTING AS COLLECTING AGENT FOR THE U.S. PUBLIC HEALTH SERVICE FORM NHS-4 1 Overtionnaire NATIONAL HEALTH SURVEY Questionnaires 2. (a) Address or description of location a. Reg. · PSU 7. Segment No. 8. Serial N (b) Mailing address if not shown in (a) Type of | Housing unit | (d) Name of Special Dwelling Place | diving | Other 9. Is this house on a form ar ranch?..... 🗀 Yès 10. Do you awn or rent this place? Ask items 10 and 11 only, if "rural" box is checked: Rent Own Rent free 11. If "Own" or "rent free" in question 10, ask: (c) During the past 12 months did soles □ No (d) During the past 12 m (a) Does this place have 10 or more acres? of crops, livestock, and other form products from the place amount to of crops, livestock, and other fam products from the place amount to If "rent" in question 10, ask: products from \$50 or more? \$250 or more? (h) Does the place you rent have 10 or more acres? ☐.Yes ☐ No □ Yes □ No □ No INSTRUCTIONS FOR O 12 13 AND 14 If "Yes," to questions 12, 13 or 14 apply definition of a housing unit to determine whether one or more additional questionnaires should be filled and whether the 13. Does anyone else living in this building use YOUR ENTRANCE to get to his living quarters?..... Yes ☐ No 15. What is the telephone number here? 16. In case I've overlooked anything, what is the best time to call? Ask at all units except apartment houses; □ No No phone 17. RECORD OF CALLS AT HOUSEHOLDS Com Date Entire household Time Callbacks for individual respondents Date 18. REASON FOR NON-INTERVIEW TYPE Demolished Refusal (Fill Hom 19) Vacant - non-seasonal Interview not obtained for: No one at home-repeated calls Vacant - seasonal In sample by mistake Fill item Usual residence elsewhere Eliminated in sub-sample Temporarily absent Cols. Armed Forces Other (Specify) because: Other (Specify) Other (Specify) 19. Reason for refusal 20. TYPE A FOLLOW-UP PROCEDURE If final call results in a Type A non-interview (except Refusals) take the following steps: 1. Contact neighbors (caretakers, etc.) until you find someone who knows the family. 2. Find out the number of people in the household, their names and approximate ages; if names of all members not known, ascertain relationships. Record this information in the regular spaces inside the questionaire. 3. Find out if anyone in the bousing unit is now in a hospital as a patient; if so, which person it is. This is done by asking the following question: 4. Is anyone in the household now in the hespital? □No Don't know No contact made (a) If "Yes,"--Who? (Enter name)_ (Col. No.) 1. (a) What is the name of the head of this household? (Enter name in first column) ast name (I) Last name (b) What are the names of all other persons who live here? (List all persons who usually live here, and all persons staying here who have no usual place of residence elsewere. List these persons in the prescribed order.) Yes (List) (c) Do any (other) ladgers or roomers live here? ☐ No (d) Is there anyone else who lives here who is now temporarily in a hospital? □ No Yes (List) Yes (List) (e) Away on business? No No First name and initial First name and initial Yes (List) (f) On a visit? ☐ No □ No Yes (List) (a) Is there anyone else stoying here now? (h) Do any of the people in this household have a home elsewhere? Yes (apply household membership rules; if not a member, delete) No (leave on questionnaire) 2. How are you related to the head of the household? (Enter relationship to head, for example: Relationship Relazionship head, wife, daughter, grandson, mother-in-law, partner, lodger, lodger's wife, etc.) Head

| • | Age Under 1 year |
|---|---|
| 3. How old were you on your last birthday? | |
| • | |
| | White Negro |
| 4. Race (Check one box for each person) | ا |
| | Other |
| • | |
| 5. Sex (Check one box for each person) | Male Female |
| 16 17 | |
| If 17 years old or over, ask: | Under 17 years |
| 6. Are you now married, widowed, divorced, separated as never married? | Married Divorced |
| (Check one hox for each person) | ☐ Widowed ☐ Separated |
| | . Never married |
| If 17 years old or over, ask: | Under 17 years |
| 7. (a) What is the highest grade you attended in school? | Elem: 1 2 3 4 5 6 7 8 |
| (Circle highest grade attended or check "None") | High: 1 2 3 4 |
| • | College: 1 2 3 4 5+ |
| (b) Did you finish thegrade (year)? | |
| | ☐ Yes ☐ No |
| If Male and 17 years old or over, ask: | Fem.or und. 17 yrs |
| 8. (a) Did you ever serve in the Armed Forces of the United States? | Yes No |
| If "Yes," ask: | |
| (b) Are you now in the Armed Forces, not counting the reserves? | |
| (If "Yes," delete this person from questionnaire) | Yes No |
| (c) Was any of your service during a war or was it peace-time only? | Var □ Peace- |
| | time only |
| If "War," ask: | ₩W II Korean |
| (d) During which war did you aerve? If "Peace-time" only, ask: | Other |
| (e) Was any of your service between June 27, 1950 and January 31, 1955? | L |
| | Yes No |
| If 17 years old or over, ask: | Under 17 years |
| 9. (a) What were you doing most of the past 12 months. | ☐ Working |
| (For males): working, or doing samething else? | Keeping house |
| (For females): working, keeping house, or doing something else? | Somerhing else |
| If "Something else" checked, and person is 45 years old or over, ask: | ☐ Yes · ☐ No |
| (b) Are you ratired? | |
| If "Working," in q. 9(a), ask: | Under 17 years |
| 10. (a) Were you warking last week or the week before? | _ ; |
| If "Keeping bouse" or "Something else" in q. 9(a), ask: | Yes No |
| (b) Did you work at a job or business at any time last week or the week before? | |
| If "No," in q. 10(a) or 10(h), ask: | |
| (c) Even though you did not wark last week or the week before, do you have a job at business? | Yes . No |
| NOTE: Determine which adults are at home and record this information. Beginning with question 11 you | Under 17 years |
| sre to interview for himself or herself, each adult person who is at home. | At home Not st |
| 11. Were you sick at any time LAST WEEK OR THE WEEK BEFORE? (That is, the 2-week period | ☐ Yes ☐ No |
| which ended last Sunday)? | |
| (a) What was the matter? | |
| (b) Anything else? | |
| 12. Last week or the week before did you take any medicine or treatment for any condition (besideswhich you told me about)? | ☐ Yes ☐ No |
| (a) For what conditions? | |
| (b) Anything else? | |
| 13. Lost week or the week before did you have any accidents or injuries? | Yes No |
| (a) What were they? | |
| (b) Anything else? | |
| 14. Did you ever have an (any other) accident or injury that was still bothering you last week or the week before? | Yes No |
| (a) In what way did it bother you? | |
| (b) Anything else? | |
| 15. AT THE PRESENT TIME do you have any ailments or conditions that have lasted for a | Yes No |
| long time? (If "No") Even though they dan't bother you all the time? | |
| (v) What are they? | |
| (b) Anything else? | • |
| 16. Nos anyone in the family - you, your, etc had any of these conditions DURING | Yes No |
| THE PAST 12 MONTHS? (Read Card A, condition by condition; record any conditions | |
| mentioned in the column for the person) | |
| 17. Does anyone in the family have any of these conditions? | Yes No |
| (Read Card B, condition by condition; record any conditions | · · |
| mentioned in the column for the person) | |
| For persons 17 years old or over, show who responded for(or was present during the asking of) questions 11-17. If person responded for self, show whether entirely or partly. For persons | Responded for self-entirely Responded for self-partly |
| R questions 11-17. It person responded for sell, show whether entirely or partly. For persons under 17, show who responded for them. | Col. Nowas respondent |
| 18. (a) Hos anyone in the family been in a hospital DURING THE PAST 12 MONTHS? | Yes No |
| If "Yes," | |
| (b) How many different times were you in the hospital overnight or langer? | |
| | No. of times |
| 19 (a) During the past 12 months has anyone in the family baen a patient in a nursing home or sanitarium? | Yes No |
| If "Yes," | |
| (b) How many times were you in a nursing hame or sanitarium? | No. of times |
| 20. If baby under one year listed as a bousehold member, ask: | Hospital Home |
| (a) Wasboby born in a hospital or at hame? If "hospital" in q. 20(a) and 1 or more in q. 18(b), ask: | Yes No |
| | |

| Toble 1 - ILLNESSES, IMPAIRMENTS AND INJURIES | | | | | | | | | | | | | | | | | | |
|---|---|---------------------|--|---|---|---|---|---|---|---|--|---|--|--|--|--|---|-------------|
| Line number | Col. No. of person | E . Question number | Did you EVER at any time talk to a doctor about? | and prese old injurie (a) If doc What did t say it was give it o r name? (b) If doc to: Re entry (d-2)- requir Ask for al during pas | tor talked to: the doctor et? did he medical tor not talke: cord original and ask (d-5) as ed. Il injuries st 2 weeks of the bedy When kind was it? alse? I Table A uries) | of? (This cc asked if (d-1) is lm; a 5 If entry is from o (If "Cau injury, a A) | s the coust | If eye trouble of any kind and be old or over, ask: Can you see well enough to read or dinory news-print with the control of | What Ask of Any e or (d-the w Asthur Cysts Growt Tumo) For a ask: | kind of is it? mly for: nry in Col. (d-1) 2) that includes ords: "condition" "disease" ts "trouble" n allergy or stroke | What pan offected Ask only Impairment Assess infect mation Aches, 1 weaks Bleedin, Cancer, growth Neuralgi Virus Show de Eer or ey Head - (U Bock - (U Bock - (U Back - (U Bock | rt of the body is 7 y for: ents; lojuries; nd foe: es, boils, ions, inflam- a, sores, ulcers pains, soreness, ess g or blood clots tumor, cysts or as a or neutitis | OR TH WEEK FORE coi to cut to ry ou activit as mu day? | BE- did use you down ir usual ries for ch as a | Satur- | How many of these days were you in bed all or most of the day? (b) Days of None | H 6-16 years old ask: How many days did year school last week or the week before? | - |
| Щ. | _ | L | | | • | 11 | | | - I | | | | L | L | I | T | Node | |
| | | | | | | - | | | | | | | | | | | | |
| _ | Table 11 - HOSPITALIZATION DURING PAST 12 MONTHS | | | | | | | | | | | | | | | | | |
| H | Т | | | | | | Table To Inte | | ALIZAT | , | | ONTHS | tion wo | w | Onv | operations p | arformed on | ł |
| Line number | Co No of pe so | r- m | No. | When did you enter the hos- pitol? Month, year) | nights were you in the hospital? | How many of these nights were in the past 12 months? | Will you need to ask cols. (f) and (g)? | of thesenights were last week or the week before? | Vas this person still in the bos- pital on ast Sunday night? | did they give it (If "they" dida" What did the los (Show same deta | o medicolo csay, asl t doctory il as in c n acciden | nome? | it was? f T.I) | p If | ou durin ital? ''Yes,' i) What opera | g this stay of was the nam tion? other operation | of the | |
| Н | +" | a) | (P) | (c) Mo: | (d) | (e) | (x) | (f) | (g) | | (h) | | | + | Yes Yes | (i) | ☐ No | 1 |
| 1 | | | 1 I | Yr: | Nights | Of Nights | □ No | Nights 1 | □ No | | | | | | | | | |
| 2 | T | | | Mo: Yr: | Nights | Or Nights | ☐ Yes | | Yes | | | | | | Yes Yes | | □ No | |
| 3 | 1 | | 1 1 | Mo: | Nights | Or All | ☐ Yes | Nights | Yès | | | | | | Yes | | □ No | |
| L | 1. | - | | Yr: | | Nights | | None | | L | | | | | | | | |
| | | | | | | | | | | | | ٠, | | , | - | | | |
| | | | | | | | | | | | | | | | | | | |
| 21. | (a) | Wed | are intere | sted in all | kinds of X-r | oys - Did | you have yo | our teeth X-r | | Y QUESTIONS | T □ Y | es | ☐ No | | Yes | | □ No | |
| | 3 months (that is, fram through last Sunday)? If "Yes," No. of times No. of times No. | | | | | | | | | | | | | | | | | |
| 22 | | _ | | | id vou have | CHEST | X-ray? | | - | | + | es-Chest | ∏ No | - | Yes-C | | □ No | |
| | 22. During the post 3 months did you have a CHEST X-ray? 23. (a) Did you have any (other) kind of X-ray at all during the post 3 months? Yes No Yes No | | | | | | | | | | | | | | | | | |
| 23, | | Did 'Yes | | e any (other | r) kind of X- | ray at oll | during the | post 3 mont | 18? | | Part(| es s) of body: | ☐ No | | ∐Yes art(s) o | f body: | ☐ No | |
| | (Ъ) | Who | t part of | the body w | as X-rayed? | | | | | | 1 | | | | | | | |
| | | | | | | | | | • | | | | | | | | | |
| | _ | 7 | | | | | | | 1 | RT OF BODY E | | | 1. | | | | | |
| Line number | B Col. No. of | person | (d Question No. | Part of bo | | How many different times did have your X-rayed d ing the pa 3 months? | you How | e did you the X-ray(s) many X-rays at the (hos- doctor's e, etc.)? | | was this X-ray(s) k-up or an examinateatment? | | If "both" in col (f) ask: How mony of theseX-roy(s were for treat- ment? (g) | | | andition | tment" in cr were you b | ol. (f) ask: | |
| | , ·a | + | | (0) | | (4) | Hosp | | | Check-up/examinati | on | 18/ | \top | | 78 | ·-/ | | |
| I | | | | | | | | ffice | | Treatment | | | | | | | | - |
| 2 | | \dagger | | | | | Hosp | | | Check-up/examinati Treatment | оп | | | | | | | |
| 3 | T | + | - | | | | Other Hosp Dr. o | | | Both Check-np/examinati Freatment | on . | | 1 | | | | . ` | |
| | Ļ | \perp | | | | | Othe | | | Both | T - | <u></u> | ' | | | | | |
| 26. | you | ···· | s, etc.? | (Show Card | | income fr | om all sour | | | , that is, your's, solaries, rents | Group | No | | Gr | oup No. | • | | |
| | | | | | | | | | | | 1 . | | | - 1 | | | | |

| | | | | | | Table I | - ILLNESS | ES. IMPAI | RMENTS | ULNI DNA | RIES | | | • | | | \neg |
|---|---|--|------------------|---|---|---|---|---|---|--|---|--|--|--|--|--|-------------|
| | If 17 year old or over and if "Yes", in | THE | it hop | st notice pen) DURING 3 MONTHS or time? | To Inter- viewer: | Did you first notice DURING THE PAST 12 | How long since you last talked to a doctor | Do you still take any medicine | About how many days | If lor more days in col. (q-1) and | A | for | each per | last condi | tion, · | If "1," or "2" or "3" in col. | |
| , | q. 10(a), 10(b) or 10(c), as How man days did | k: Cbec | k one | Didstart during the past 2 weeks | CON- TINUE if col. | MONTHS or before that time? | (If less than one month, | or treatment that the doctor prescrib- ed for? | during the past 12 months, has kept you | col: (e) is check- ed, ask: How many of these | Please look at this cord and read each state- | If "1," "2" or "3" in col. (r): | If "Yes" in col (s); Which? | If "1" col. (r | or "2" in) ask: - | (r) ask: | |
| | keep you from work lost week or the week before? | 3 mos | ing 3 mos. | or before that time? (If during past 2 weeks, ask) Which week, last week or the week before? | (k) is check- ed, or the condi- tion is on Card A or is an im- ment; or her- wise, STOP | | month ('Und enter ('Und I'' for ''Mo.'') | Or, follow any advice he gave? | for all or most of | days were during last week or the week before? | ment. Then toll me which state- ment fits you best, in terms of health. (Show Cards C- F, as appro- priate) | because of any of the condi- tians you have told me obout? | (Enter X on line for each condi- tion named) | How long have you been? (Insert the words of the statement aelect- | If 17 years old or over, ask: Were you working of a job or business up to that time? | Please look at this card and read each state- ment. Then tell me which atate- ment | Line number |
| | (j) Dny: | (k) | (1) | (m) | (38) | (n) During past | (o) Mos. | (p) | (q-1) Days | (q-2) Days | (r) | (s) | (t) | ed) (u) Mos. | (v) | fits you best. (Show Card G) (w) | |
| | or None | | | ☐ Week before ☐ Before 2 wh | | 12 montha Before Birth | Yrs. | No Dr. | or | or None | | □No | | Yrs. | No Und.17 | | l L |
| | | | , | | | | | | | | | | | | | 4. | |
| | `` | | | | | Table II - H | | ZATION D | URING PA | | | | | | | _ | 4 |
| | over who | sbow an | perat | ioo, a setting of | a fracture, | of persons 6 year ora delivery in Co | ls.(h)or(i): | | - | What is th | e name ond | oddrass | of the ho | spital you v | were in? | | |
| ` | How many you in the fore you ! tion (deli | hospita ad your | l, be- opera- | After you le pital, how n was it befor returned to activities for | e you your usual | If "still unsi asi How long hos since you let hospital? | it been | | | Enter nam | e, city and | State; if | city not k | nowa, eate | r county) | - | |
| | No. of n | ights | | . (k No. of days ☐ Still on | | (I) Over 6 mo | onths uths: Months: | , | | | (| m) | | | | : | - |
| - | Nn. of n | in. of nights No. of days If under 6 months: | | | | | onths | ıths: | | | | | | | | | |
| | No. of n | ights | | No. of days | | Over 6 mo | nths nths;. Months: | | | ••••• | | ••••• | | | | | |
| , | | | | ¥., | | , · | | | | • | | | | | | | |
| | 24 (-) 0 | · | 7 | ر المالية المالية المالي | in ab | e family have any | | RAY QUES | | ПП | | | - I | Yes | | | |
| | If "Ye | condition s,'' at part c | ? if the ! | ody was treate | d? | • | | | | | of body: | |] No | Part(s) | | . ــــــــــــــــــــــــــــــــــــ | 10 |
| ļ | (e) W | sthis in | clude | I in the X-ray(s | you told | me obout before? | | | • | Y | | |] No | Yes | , | | _ |
| | If "Ye | :s," | | e family have a | | e during the past | 3 months? | | | Part(| es s) of body: | |]No | Tart(s) | of body: | | No |
| | (c) W | s this ir | clude | d in the X-ray(s | ı) you told | me about before? | 1 , | | | □ Y | es | |] No | Yes | | | Νο |
| , | | | | | | ٠. | ٠ | | ` | | . * | s. | | | | | |
| | | | | | able X - I | FILL ONE LINE | FOR EAC | | F BODY | | ROM QUE | STIONS | 22-25 | | | | \Box |
| | (Ask after | | | | | re lines in Table ols.(a)-(b) of Tab | | | COTNOTES | , | • | | | | | | |
| | If "Yes, Which X | " | e thes | -rays you told (| ne about ta | ken at the same ti | me? | | | | | | | | | | |
| | (Stop) Ye | | | f body: | for X-rays No. | Part(s) of body | | No. | | | , | | } | ٠ | | | · |
| | | | | | | | | | | | | - | | . | | | |
| | | Pa | t(s) α | f body: | No. | Parr(s) of body | | No. | | | | 1 | | ٠. | | | |
| | | Pa | τ(s) α | f body: | No. | Part(s) of body | | No. | | • | | | | | | | |
| | Group N | . <u> </u> | | | Group No. | 1 | Group N | 0. | | Grou | No. | | | Group No. | | | \dashv |
| | 1 | - | | | | | | | | | | | | | | | - 1 |

| | | Table A - (Accidents and Injuries) | | | | | | |
|---------------------------------|--|--|---|--|--|--|--|--|
| Line No. | I. When did the accident happen? | 2. At the time of the accident, what part of the be Anything else? | ody was hurt? What kind of injury was it? | | | | | |
| Table I | Year: | Part(s) of body | . Kind of injury(s) | | | | | |
| | (If 1960 or 1961 also enter the month) | · | • | | | | | |
| Accident happened | ĺ | | , | | | | | |
| last week or | Month: | | | | | | | |
| week before (Go to q. 3) | | , | * | | | | | |
| | ruck, bus or other motor vehicle involved in the o | | No (Go to Section B) | | | | | |
| \$ | on one mater vehicle involved? or one) maving at the time? | Yes (more than on | e) No No (Go to Section B) | | | | | |
| ı | the vehicle, getting in or out of it, a passenger | | 2. Getting in or out 3. Passenger Qo to Section A q. 6) | | | | | |
| Same | n A - (Motor Vehicle Accidents) | A q.5) | 4. Driver ptor Vohicle Accidents) | | | | | |
| 36010 | <u> </u> | | 1 | | | | | |
| 5. (a) How did the | If "Outside" in q. 4, ask: accident happen? | 7. How did the occident happen? A.1. Any injury involving an uncontrolled f | · | | | | | |
| I. Accide | ent between motor vehicle and person riding | 2. Any injury involving the discharge of | | | | | | |
| on bic | yele, in streetcar, on milrord train, on horse- vehicle | | a non-motor vehicle in morion (streetcar, railroad | | | | | |
| 1 | ent between moror vehicle and person who | train, airplane, boat, bicycle, horse-di | | | | | | |
| ₩as ₩ | alking, running, or standing | B.4. Any injury caused by machinery (belt | or motor driven) while in operation | | | | | |
| 3. Other | (Specify how the eccident happened) | (Specify kind of mechinery) | - | | | | | |
| · - | : | 5. Any injury caused by edge or point of piercing implement | knife, scissors, nail or orber cutting or | | | | | |
| (b) What kind(s) | of motor vehicle was involved? | 6. Any injury caused by foreign body in e | eye, windpipe, or orher orifices | | | | | |
| l. Car | 2 Taxi 3 Bus | 7. Any injury caused by animal or insect | • | | | | | |
| 4. 🔲 Truck | 5. Motorcycle 6. Other (Specify) | 8. Any injury caused by poisonous aubstr | ance swallowed (Specify substance) | | | | | |
| | | C.9. Fell on stairs or steps or from a beigh | or , | | | | | |
| | | 10. All other fails | • | | | | | |
| # "Center" | out" "Passenger" or "Driver," in q. 4, ask: | Bumped into object or person (covers punching, kicking, etc.) | all collisions between persons including striking, | | | | | |
| 6. (a) How did the | • | l ' ' | cts beld in own hand or hand of other petson, also | | | | | |
| | ent between two or more motor vehicles on | falling, flying, or thrown objects) | | | | | | |
| roadw | ay | Handling or stepping on sharp or rough objects such as stones, splinters, broken glass, rope, etc. | | | | | | |
| | ent between motor webicle and some other t on roadwsy | I4. Caught in, pinched or crushed between stationary object | n two moving objects or between a moving and a | | | | | |
| · · | ity object) | 15. Came in contact with hot object or so | batance or open flame | | | | | |
| _ | vehicle came to sudden stop on roadway | 16. Oue-time lifting or other one-time exer | rtion | | | | | |
| _ | vehicle ran off roadway | 17. Twisting, stumbling, etc. | | | | | | |
| o. Uther | (Specify how the eccident happened) | D.18. Other (Specify how accident happened | 0 | | | | | |
| | Acc. on roadway | | | | | | | |
| | Acc. not ou roadway | | · | | | | | |
| | motor vehicle were you in (getting in) (getting | , | | | | | | |
| out of) wher | n the accident happened? 2. [] Taxi 3. [] Bus | | | | | | | |
| 4. Truck | | | | | | | | |
| | - 11/2 | 100 100 100 100 100 100 100 100 100 100 | | | | | | |
| | | ASK FOR ALL ACCIDENTS | | | | | | |
| 1 | e accident happenat home or some other place? ne (inside house) 2. 🔲 At | | ome other place | | | | | |
| If "Some other p | place,'' ask: | | | | | | | |
| (b) What kind of 3. ☐ Street | | hool (includes school premises) | | | | | | |
| 4. 🔲 Farm | 7. 🔲 PI | ace of recreation and sports, except at school | | | | | | |
| | | her (Specify the place where eccident happened) | | | | | | |
| 9. Were you at wo | rk at your job or business when the accident happ | | nder 17 at time of accident | | | | | |
| <u> </u> | | FOOTHOTES AND COMMENTS | | | | | | |
| | | | • | | | | | |
| | | | , | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| 1 | | | | | | | | |
| | | | ` . | | | | | |
| | | ` | | | | | | |

| Gard A | Card C | Card E | Card G |
|--|--|--|--|
| | | | · |
| NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY |
| Check List of Chronic Conditions | For: Workers and other persons except Housewives and Children | For: Children from 6 through 16 years old | |
| 1. Asthma 2. Tuberculesis 3. Chronic brenchitis 4. Repeated attacks of sinus 5. Rheumatic fever 6. Hardening of the arteries 7. High blood pressure 8. Heart trouble 9. Stroke 10. Trouble with varicose veins 12. Hay fever 13. Tumor, cyst or growth 14. Chronic gallbladder or liver trouble 2 Touble vith vericose veins 2 Chronic skin trouble 2 Chronic skin trouble 2 Tennia or rupture trouble 2 Prostate trouble 2 Prostate trouble | 2. Able to work but limited in amount of work or kind of work. atism 3. Able to work but limited in kind or amount of other activities. goiter 4. /Not limited in any of these ways. | Not able to go to school at all. Able to go to school but limited to certain types of schools or in school attendance. Able to go to school but limited in other activities. Not limited in any of these ways. | 1. Confined to the house all the time except in emergencies. 2. Able to go outside but need the he of another person in getting around outside. 3. Able to go outside alone but have trouble in getting around freely. 4. Not limited in any of these ways. |
| Card B | Card D | Card F | Cord H |
| NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY | NATIONAL HEALTH SURVEY |
| Check List of Selected Impairments 1. Deafness or serious trouble with hearing | For: Housewife | For: Children under 6 years old 1. Not able to take part at all in | Family income during past 12 months Group 1. Under \$500 (Including loss) |
| Serious trouble with seeing, even when wearing glass Cleft palate Any speech defect Missing fingers, hand, or arm——toes, foot, or leg Palsy Paralysis of any kind | 2. Able to keep house but limited in amount or kind of housework. 3. Able to keep house but limited in kind or amount of other activities. | ordinary play with other children. 2. Able to play with other children but limited in amount or kind of play. 4. Not limited in any of these ways. | Group 2. \$500- \$999 Group 3. \$1,000- \$1,999 |
| Repeated trouble with back or spine Club foot Permanent stiffness or any deformity of the foot, lefingers, arm or back Any condition present since birth | 4. Not limited in any of these ways. | The state of the services and services are services and services and services and services are services are services and services are services are services and services are services are services are services are services are s | Group 5. \$3,000- \$3,999 Group 6. \$4,000- \$4,999 |
| | | • | |

SELECTED REPORTS FROM THE U.S. NATIONAL HEALTH SURVEY

Public Health Service Publication No. 584

Series A (Program descriptions, survey designs, concepts, and definitions)

- No. 1. Origin and Program of the U. S. National Health Survey. 25 cents.
- No. 2. The Statistical Design of the Health Household-Interview Survey. 35 cents.
- No. 3. Concepts and Definitions in the Health Household-Interview Survey. 30 cents.
- No. 4. Plan and Initial Program of the Health Examination Survey.

Series B (Health Interview Survey results by topics)

- No. 8. Persons Injured by Class of Accident, United States, July 1957-June 1958. 40 cents.
- No. 9. Impairments by Type, Age, and Sex, United States, July 1957-June 1958. 25 cents.
- No. 10. Disability Days, United States, July 1957-June 1958. 40 cents.
- No. 11. Limitation of Activity and Mobility Due to Chronic Conditions, United States, July 1957-June 1958. 30 cents.
- No. 12. Chronic Respiratory Conditions Reported in Interviews, United States, July 1957-June 1958. 30 cents.
- No. 13. Heart Conditions and High Blood Pressure Reported i., Interviews, United States, July 1957-June 1958. 30 cents.
- No. 14. Dental Care, Interval and Frequency of Visits, United States, July 1957-June 1959. 35 cents.
- No. 15. Dental Care, Volume of Visits, United States, July 1957-June 1959. 35 cents.
- No. 16. Types of Injuries, Incidence and Associated Disability, United States, July 1958-June 1959. 30 cents.
- No. 17. Peptic Ulcers Reported in Interviews, United States, July 1957-June 1959. 25 cents.
- No. 18. Acute Conditions, Incidence and Associated Disability, United States, July 1958-June 1959. 30 cents.
- No. 19. Volume of Physician Visits, United States, July 1957-June 1959. 40 cents.
- No. 20. Arthritis and Rheumatism Reported in Interviews, United States, July 1957-June 1959. 25 cents.
- No. 21. Diabetes Reported in Interviews, United States, July 1957-June 1959. 25 cents.
- No. 22. Loss of Teeth, United States, July 1957-June 1958. 25 cents.
- No. 23. Acute Conditions, Geographic Distribution, United States, July 1958-June 1959. 30 cents.
- No. 24. Acute Conditions, Seasonal Variations, United States, July 1957-June 1960. 35 cents.
- No. 25. Hernias Reported in Interviews, United States, July 1957-June 1959. 25 cents.
- No. 26. Interim Report on Health Insurance, United States, July-December 1959. 45 cents.
- No. 27. Distribution and Use of Hearing Aids, Wheel Chairs, Braces, and Artificial Limbs, United States, July 1958-June 1959. 25 cents.
- No. 28. Persons Receiving Care at Home, United States, July 1958-June 1959. 30 cents.
- No. 29. Disability Days, United States, July 1959-June 1960. 40 cents.
- No. 30. Proportion of Hospital Bill Paid By Insurance, Discharged From Short-Stay Hospitals, United States, July 1958-June 1960. 40 cents.
- No. 31. Duration of Limitation of Activity Due to Chronic Conditions, United States, July 1959-June 1960. 30 cents.
- No. 32. Hospital Discharges and Length of Stay: Short-Stay Hospitals, United States, 1958-1960.
- No. 33. Acute Conditions, Seasonal Variations, United States, July 1957-June 1961.
- No. 34. Acute Conditions, Geographic Distribution, United States, July 1960-June 1961.

Series C (Health Interview Survey results for population groups)

- No. 1. Children and Youth, Selected Health Characteristics, United States, July 1957-June 1958. 35 cents.
- No. 2. Veterans, Health and Medical Care, United States, July 1957-June 1958. 40 cents.
- No. 3. The Hawaii Health Survey, Description and Selected Results, Oahu, Hawaii, October 1958-September 1959. 40 cents.
- No. 4. Older Persons, Selected Health Characteristics, United States, July 1957-June 1959. 45 cents.
- No. 5. Selected Health Characteristics by Area, Regions and Urban-Rural Residence, United States, July 1957-June 1959. 35 cents.
- No. 6. Selected Health Characteristics by Area, Divisions and Large Metropolitan Areas, United States, July 1957-June 1959. 35 cents.
- No. 7. Currently Employed Persons, Illness and Work-Loss Days, United States, July 1959-June 1960.

Series D (Developmental and Evaluation Reports)

- No. 1. A Study of Special Purpose Medical-History Techniques. 30 cents.
- No. 2. Co-operation in Health Examination Surveys. 35 cents.
- No. 3. Hospital Utilization in the Last Year of Life. 30 cents.
- No. 4. Reporting of Hospitalization in the Health Interview Survey. 50 cents.
- No. 5. Health Interview Responses Compared With Medical Records. 45 cents.
- No. 6. Attitudes Toward Co-operation in a Health Examination Survey. 35 cents.
- No. 7. Evaluation of a Single-Visit Cardiovascular Examination. 30 cents.

Catalog Card

U. S. National Health Survey

Acute conditions, seasonal variations, United States, July 1957-June 1961; statistics on incidence of acute conditions and number of associated days of disability in each calendar quarter according to condition group and age. Based on data collected in household interviews during the period July 1957-June 1961. Washington, U.S. Dept. of Health, Education, and Welfare, Public Health Service, 1962.

59 p. tables. diagrs. 27cm. (11s Health statistics, ser. B33)

- U. S. Public Health Service, Publication no. 584-B33
- 1. Sickness U.S. Statistics. 2. Accidents U.S. Statistics.
- I. Title.

Cataloged by Department of Health, Education, and Welfare Library.