

**Profile of Chronic Illness
in Nursing Homes
United States:
National Nursing Home Survey
August 1973-April 1974**

Statistics on reported chronic conditions and impairments; primary diagnosis at last examination and at admission; behavioral problems; impairments in vision, hearing, and speech; and use of dentures and special aids. Based on information collected in the 1973-74 National Nursing Home Survey. Comparisons made with data from previous nursing home surveys and from surveys of the noninstitutional population.

DHEW Publication No. (PHS) 78-1780

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Center for Health Statistics
Hyattsville, Md. December 1977



Library of Congress Cataloging in Publication Data

Ingram, Donald K

Profile of chronic illness in nursing homes, United States, August 1973-April 1974.

(Vital and health statistics: Series 13, Data from the National Health Survey, Data on health resources utilization; no. 29) (DHEW publication; no. (PHS) 78-1780)

Bibliography: p.

1. Nursing home patients—United States—Statistics. 2. Chronic diseases—United States—Statistics. I. United States. National Center for Health Statistics. II. Title. III. Series: United States. National Center for Health Statistics. Vital and health statistics: Series 13, Data from the National Health Survey, Data on health resources utilization; no. 29. IV. Series: United States. Dept. of Health, Education, and Welfare. DHEW publication; no. (PHS) 78-1780.

RA407.3.A349

no.29

[RA997]

362.1'1'0973s

ISBN 0-8496-0096-8

[312.3]

77-22564

NATIONAL CENTER FOR HEALTH STATISTICS

DOROTHY P. RICE, *Director*

ROBERT A. ISRAEL, *Deputy Director*

JACOB J. FELDMAN, Ph.D., *Associate Director for Analysis*

GAIL F. FISHER, *Associate Director for the Cooperative Health Statistics System*

ELIJAH L. WHITE, *Associate Director for Data Systems*

JAMES T. BAIRD, JR., Ph.D., *Associate Director for International Statistics*

ROBERT C. HUBER, *Associate Director for Management*

MONROE G. SIRKEN, Ph.D., *Associate Director for Mathematical Statistics*

PETER L. HURLEY, *Associate Director for Operations*

JAMES M. ROBEY, Ph.D., *Associate Director for Program Development*

PAUL E. LEAVERTON, Ph.D., *Associate Director for Research*

ALICE HAYWOOD, *Information Officer*

DIVISION OF HEALTH RESOURCES UTILIZATION STATISTICS

SIEGFRIED A. HOERMANN, *Director*

JOAN F. VAN NOSTRAND, *Chief, Long-Term Care Statistics Branch*

W. EDWARD BACON, Ph.D., *Chief, Hospital Care Statistics Branch*

JAMES E. DeLOZIER, *Chief, Ambulatory Care Statistics Branch*

MANOOCH EHR K. NOZARY, *Chief, Technical Services Branch*

Vital and Health Statistics-Series 13-No. 29

DHEW Publication No. (PHS) 78-1780
Library of Congress Catalog Card Number 77-22564

CONTENTS

The Survey	1
Overview of Findings	1
Overview of the Report	2
Sources and Qualifications of Data	2
The Population	3
Age and Sex	3
Certification Status of Facility	4
The Profile of Chronic Illness	5
Reported Conditions	5
Age and Sex	7
Certification Status of Facility	9
Primary Diagnosis	9
Primary Diagnosis at Last Examination	9
Primary Diagnosis at Admission	13
Behavioral Problems	14
Age and Sex	15
Certification Status of Facility	16
Impairments in Vision, Hearing, and Speech and Use of Dentures.....	16
Vision	16
Hearing	17
Speech	19
Use of Dentures	19
Special Aids	20
Age and Sex	20
Certification Status of Facility	20
List of Detailed Tables	24
Appendixes	
I. Technical Notes on Method	48
II. Definitions of Certain Terms Used in This Report	56
III. Resident Questionnaire Used in the 1973-74 National Nursing Home Survey	58

LIST OF FIGURES

1. Percent distribution of nursing home residents by age: United States, August 1973-April 1974	3
2. Prevalence rates per 1,000 nursing home residents for selected reported chronic conditions, by age: United States, August 1973-April 1974	8
3. Rate per 1,000 nursing home residents for selected primary diagnoses at last examination, by age: United States, August 1973-April 1974	12
4. Comparison of rates for those primary diagnoses showing significant changes between admission and last examination: United States, August 1973-April 1974	14
5. Percent of nursing home residents with behavioral problems by problem and frequency: United States, August 1973-April 1974	15

6. Percent distribution of nursing home residents by vision status, according to age and sex: United States, August 1973-April 1974	16
7. Percent of nursing home residents using eyeglasses, by age and sex: United States, August 1973-April 1974	17
8. Percent distribution of nursing home residents by hearing status, according to age and sex: United States, August 1973-April 1974	18
9. Percent of nursing home residents using a hearing aid, by age and sex: United States, August 1973-April 1974	18
10. Percent distribution of nursing home residents by speech status, according to age and sex: United States, August 1973-April 1974	19
11. Percent distribution of nursing home residents by use of dentures, according to age and sex: United States, August 1973-April 1974	19

LIST OF TEXT TABLES

A. Number, percent distribution, and median age of nursing home residents by age, according to sex: United States, August 1973-April 1974	4
B. Number of nursing homes and number and percent distribution of nursing home residents by sex and age, according to certification status of home: United States, August 1973-April 1974	4
C. Number and percent of nursing home residents by selected reported chronic conditions and impairments: United States, 1973-74, 1969, and 1964	6
D. Number and percent distribution of nursing home residents by primary diagnosis at last examination and rank in August 1973-April 1974 and June-August 1969: United States	10
E. Number and percent distribution of nursing home residents by sex and age, according to special aids used: United States, August 1973-April 1974	21

SYMBOLS

Data not available-----	---
Category not applicable-----	...
Quantity zero-----	-
Quantity more than 0 but less than 0.05-----	0.0
Figure does not meet standards of reliability or precision-----	*

PROFILE OF CHRONIC ILLNESS IN NURSING HOMES

Donald K. Ingram, *Formerly of Division of Health Resources Utilization Statistics*

THE SURVEY

OVERVIEW OF FINDINGS

Presented in this report are descriptive statistics which provide a profile of the health of the over 1 million persons in the United States who reside in 15,700 nursing homes. Based on the results of a nationwide survey conducted between August 1973 and April 1974, the size of this institutionalized population is estimated at 1,075,800 persons—approximately 1 person out of every 200 living in the country. If only persons 65 years and over are counted, it is estimated that 5 percent of the Nation's older population resided in a nursing home at the time of the survey. About 11 percent of the nursing home population were younger than 65 years, 82 years being the median age.

It is the inability to care adequately for chronic illness outside an institution which has been assumed to be the primary reason motivating the great number of admissions to nursing homes. Statistics on chronic illness among persons living outside institutions support this assumption.¹ During 1972 approximately 26 million persons, representing 13 percent of the noninstitutionalized population, were reported to be limited in their daily activities as a result of a chronic condition. The impact of chronic illness was felt most severely in the elderly segment of the noninstitutionalized population. In particular, around 9 million elderly persons, or 43 percent of those 65 years and older, had a chronic health problem that limited their daily

activity in some way. For nearly 4 million of the elderly, chronic illness was severe enough to cause a limitation in their mobility, and over 1 million were confined to the house. With such a great prevalence of chronic illness among the elderly in general it is understandable that many of those who found it difficult, if not impossible, to be cared for at home eventually were admitted to nursing homes.

Inside nursing homes the health profile of the residents is dominated by chronic illness. The current survey found that practically all residents had to cope with at least one type of chronic condition or impairment. Fifty-eight percent reportedly suffered from senility. Two other major chronic conditions were heart trouble and arthritis or rheumatism, both of which affected about 34 percent of the nursing home population. In addition, sensory impairments were very prevalent. For about 13 percent of all residents vision was either severely impaired or completely lost; around 32 percent had impaired hearing. The speech of about 26 percent of the population was considered impaired. Around 64 percent had dentures. Behavioral problems were also associated with chronic illness. Approximately 42 percent of the residents were described as agitated or nervous, while 39 percent were reported to act depressed at least once a week. The high level of chronic illness was also reflected in the use of special aids. About half of the population used at least one type of special aid. For about 35 percent of

all residents this aid was a wheelchair, and around 15 percent used a walker for moving about.

OVERVIEW OF THE REPORT

The health status of the nursing home population during August 1973-April 1974 is the focus of this report. In effect, to examine the health status of this institutionalized population was to examine the profile of chronic illness among its members. The survey data provided several different perspectives through which the profile of chronic illness could be analyzed.

The prevalence of various chronic conditions and impairments provided the primary perspective. One source of prevalence data was the information on 11 major chronic conditions and impairments reported by the nurse or other staff member most familiar with the resident's health condition. Another source was drawn from the medical records of the resident. Of interest was the primary diagnosis noted at the last examination as well as at the time of admission. In addition to these prevalence data on medical problems, the profile of chronic illness was also viewed from the perspective of the behavioral problems affecting the residents. Finally, the relative extent of chronic disability was assessed by two other measures of chronic impairments. One such source was derived from the detailed information which dealt with the extent of visual, hearing, speech, and dental impairments. This included data on the use of eyeglasses, hearing aids, and dentures. The last source concerned data on the use of special aids such as wheelchairs, walkers, canes, braces, and crutches.

The analysis of these main variables provided the statistical profile of chronic illness which constitutes the scope of this report. Secondary variables providing additional detail to the analysis include the age and sex of residents. The one facility variable analyzed in detail was certification status. Certification status refers to whether or not the facility was certified for Medicare (Title XVIII of the Social Security Act), Medicaid (Title XIX of the Social Security

Act), or both. A detailed description of this classification is provided in appendix II.

SOURCES AND QUALIFICATIONS OF DATA

The data presented in this report were obtained in a sample survey conducted by the National Center for Health Statistics (NCHS) between August 1973 and April 1974. This survey is referred to in the report as the 1973-74 National Nursing Home Survey (1973-74 NNHS). Detailed information concerning the development of the survey and its methodology are presented in appendix I.

The 1973-74 NNHS represents an expansion of previous NCHS surveys of nursing homes and related facilities which were on an *ad hoc* basis. Background on the establishment of these surveys has been reported previously.² The first was a mail-out survey conducted during 1963. It was succeeded by two interview surveys conducted during 1964 and 1969.

Since the information herein was derived from a sample survey, the reader should be aware of the exact sources of the data and the qualifications which bear upon their interpretation. There are three appendixes provided to aid in understanding the more technical aspects of the survey operation and methodology.

Contained in appendix I is a general description of the survey, including the sample design and the survey procedures. Since the statistics in this report are estimates based on a sample of nursing homes, they are of course subject to sampling variability. Therefore the reader should pay particular attention to the tables of standard errors located in appendix I. The survey terminology is found in appendix II. Definitions of various terms should be noted as they are essential to the proper interpretation of the data. In particular the reader is directed to the criteria for classifying nursing homes. The information on the facilities was obtained in a personal interview with the administrator. Responses to items contained in the Facility Questionnaire were used to determine the certification status of the institution. The Resident Questionnaire was used to gather information on a sample of resi-

dents within each facility. This was administered during a personal interview with a nurse, usually the one who provided care for the sample resident. During the interview the respondent was asked to refer to the resident's medical record. A facsimile of the Resident Questionnaire is exhibited in appendix III.

This report represents a followup to previous NCHS reports on the health of nursing home residents.³⁻⁶ Wherever appropriate, comparisons were made with data from the 1964 and 1969 nursing home surveys previously described. Special care should be taken in analyzing these comparisons, however, since there were differences in scope. The 1964 and 1969 surveys included all nursing homes, whereas the 1973-74 NNHS included only those homes providing

some level of nursing care (i.e., personal care homes were excluded). This exclusion should have minimal impact on comparisons, however, because residents of personal care homes comprised only about 5-6 percent of the total resident population of the 1964 and 1969 surveys.

The Health Interview Survey (HIS) provides the main source of data for comparisons with the noninstitutionalized population. This national household survey, also conducted by NCHS, relies upon the respondent's knowledge of his own health condition. Differences in survey material and method of data collection should be taken into account, especially since data are not supplied by medically trained personnel from the medical record.

THE POPULATION

In order to provide a backdrop to the chronic illness profile of nursing home residents, this section will highlight some demographic features of this institutionalized population. There were an estimated 15,700 nursing homes in operation during the survey period. The size of the resident population within these facilities totaled 1,075,800 persons.

AGE AND SEX

The age distribution of the nursing home population has always been a very skewed one. As shown in figure 1, the population for the 1973-74 survey period was no exception. The median age of the population was 82 years. The modal age class was 75-84 years, which accounted for 36 percent of all residents.

In addition to the skewness in the age distribution, the nursing home population has also been traditionally characterized by its preponderance of female residents. An estimated 70 percent of the population was female, compared to 30 percent who were male. The sex ratio was




Figure 1. Percent distribution of nursing home residents by age: United States, August 1973-April 1974

42 men for every 100 women. In contrast, the proportion of male residents under 65 was twice that of female residents.

In general, female residents were also older than male residents (table A). The median age of women was 82 years, while the median age of men was 78 years.

Table A. Number, percent distribution, and median age of nursing home residents by age, according to sex: United States, August 1973-April 1974

Sex	Number of residents	All ages	Percent distribution				Median age
			Under 65 years	65-74 years	75-84 years	85 years and over	
Both sexes.....	1,075,800	100.0	10.6	15.2	35.8	38.4	82
Male.....	318,100	100.0	16.5	20.5	32.2	30.9	78
Female.....	757,700	100.0	8.2	12.9	37.3	41.6	82

CERTIFICATION STATUS OF FACILITY

The nursing homes included in the 1973-74 NNHS were:

- Those certified as extended care facilities by Medicare (Title XVIII of the Social Security Act).^a
- Those certified as skilled nursing homes by Medicaid (Title XIX of the Social Security Act).^a
- Those certified as intermediate care facilities by Medicaid.

- Those not certified by either program but providing some level of nursing care.

Of the homes which were certified, some were certified under *both* the Medicare and the Medicaid programs. Other homes certified by the Medicaid program were certified to participate as both a skilled nursing home *and* an intermediate care facility. In order to provide detailed data by certification status, some small certification subgroups were combined with larger ones when both provided a similar level of care. Thus those 406,900 residents classified in table B as residing in facilities certified by *both* Medicare and Medicaid (BM&M) include 23,500 which were in facilities certified by Medicare only. Similarly, the 292,500 residents in facilities certified by Medicaid as skilled

^aThe extended care facility and skilled nursing home designations are used in this report because most of the survey was conducted prior to the legislation which created the skilled nursing facility.

Table B. Number of nursing homes and number and percent distribution of nursing home residents by sex and age, according to certification status of home: United States, August 1973-April 1974

Certification status	Number of nursing homes	Number of residents	Sex			Age				
			Both sexes	Male	Female	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
All homes.....	15,700	1,075,800	100.0	29.6	70.4	100.0	10.6	15.2	35.8	38.4
Both Medicare & Medicaid ..	4,200	¹ 406,900	100.0	27.7	72.3	100.0	9.3	15.1	37.1	38.4
Medicaid only										
Skilled nursing home	3,500	² 292,500	100.0	29.7	70.3	100.0	9.8	15.3	35.6	39.3
Intermediate care facility.....	4,400	236,700	100.0	32.4	67.6	100.0	12.5	15.8	34.0	37.8
Not certified	3,600	139,800	100.0	29.8	70.2	100.0	12.9	13.9	35.2	37.9

¹23,500 of the residents were in facilities certified by Medicare only.

²129,600 of the residents were in facilities certified as both SNH and ICF.

nursing homes (SNH's) included 129,600 which were in homes also certified as intermediate care facilities (ICF's).

Although the homes certified by both Medicare and Medicaid or by Medicare only (the BM&M group in this report) comprised about 27 percent of all homes, they housed about 38 percent of all residents. Similarly, those certified as skilled nursing homes by Medicaid (the SNH's in this report) comprised only about 22 percent of the Nation's nursing homes but had 27 percent of the population in residence. Those homes certified only by Medicaid as intermediate care

facilities (the ICF's in this report) comprised 28 percent of all homes but had only about 22 percent of all residents. Finally, homes not certified for Medicare or Medicaid comprised 23 percent of all homes but housed about 13 percent of the residents.

Sex and age of the residents did not vary greatly across the facility types (table B). The only significant differences were found in ICF's and not certified homes, which had significantly greater proportions of younger residents than found in the other two types of facilities.

THE PROFILE OF CHRONIC ILLNESS

This section provides a quantitative index of chronic illness in the nursing home population during the 1973-74 survey period. Prevalence patterns by age and sex of the residents and by certification status of the facilities can be drawn directly from the data collected on the reported chronic conditions and impairments, on the condition identified as the primary diagnosis at admission and at the last examination, and on the reported behavioral problems affecting the residents. In addition, an assessment of chronic disabilities can be made by analyzing data on particular impairments related to vision, hearing, and speech and on use of dentures and special aids.

REPORTED CONDITIONS

Listed in table C are the chronic conditions for which reported prevalence was sought from the nurse respondents. This information was obtained from item 9 of the Resident Questionnaire. Based on the findings of previous NCHS surveys of nursing homes, it was thought that these 11 conditions were the most relevant to the population under study, particularly in terms of their prevalence. About 95 percent of the residents had at least one of these

conditions, with the average number of conditions per resident at 2.2.

Previous reports have shown that residents are likely to have multiple conditions.^{3,6} For the 1969 survey the mean was computed to be 3.4 conditions, however, that survey used a more extensive and open-ended listing of reported conditions. An estimated 98 percent of all residents were reported to have at least one of the conditions in that listing. Because the 1973-74 NNHS did not include all the chronic conditions used in the previous surveys, the comparison of averages for these two time periods is not valid.

By far, senility was found to be the most prevalent condition among nursing home residents. Well over half the population was reported to have this chronic illness. As observed in table C, the rate of 58 percent was similar to that reported in the 1969 survey and over twice the rate reported 10 years earlier in the 1964 survey. For the noninstitutionalized population during July 1966-June 1968, senility was reported in HIS as the major reason that 154,000 persons, or 9 percent of those persons aged 55 years and over, received home care.⁷

As discussed in a previous report, the classification of senility as a disease entity presents many nosological difficulties.⁵ Senility may refer specifically to advanced types which in-

Table C. Number and percent of nursing home residents by selected reported chronic conditions and impairments: United States, 1973-74, 1969, and 1964

Reported chronic conditions and impairments	1973-74		1969 ²	1964 ³
	Number of residents	Percent ¹		
			Percent ¹	
Senility.....	627,000	58.3	56.4	26.8
Arthritis or rheumatism.....	368,500	34.3	33.2	22.1
Heart trouble.....	360,500	33.5	36.3	28.2
Mental illness ⁴	200,400	18.6	18.1	18.2
Amputation of extremities or limbs, or permanent stiffness or any deformity of foot, leg, fingers, arm, or back.....	150,000	13.9	23.8	15.6
Diabetes.....	142,700	13.3	11.9	8.0
Paralysis due to stroke.....	122,100	11.3	25.1	12.0
Glaucoma or cataracts.....	110,900	10.3	---	6.2
Any chronic trouble with back or spine.....	106,100	9.9	10.0	---
Mental retardation.....	72,800	6.8	---	---
Paralysis or palsy not related to stroke, arthritis, or rheumatism.....	66,200	6.2	9.8	4.7
None of these conditions.....	57,700	5.4	---	---

¹ Figures may not total to 100 percent because residents may have had more than one condition.

² See reference 4 for source of estimate.

³ See reference 5 for source of estimate.

⁴ Estimates for 1969 and 1964 include mentally retarded residents; for 1973-74 the estimate for mental retardation is presented separately.

volve psychosis, such as chronic brain syndrome or senile dementia, and consequently are classified as mental disorders under the *Eighth Revision International Classification of Diseases, Adapted for Use in the United States (ICDA)*.⁸ Under more generalized lay terminology, senility may refer simply to symptoms such as a decline in intellect, memory, and judgment, a loss of orientation, difficulty in speaking, or feebleness. In the 1973-74 NNHS this later definition was used. Thus senility was considered as a very broad category encompassing disease etiology and symptomology generally associated with deteriorating effect of the aging process on mental functioning. Since senility has been linked with other diseases, most notably with cerebral arteriosclerosis, and with impairments in perception and sensation, there may have been many physical causes leading to a prevalence rate of 58 percent in the nursing home population.⁵ There is also the possibility that the term senility as defined in the survey represented a classification so broad that the respondent may have equated this condition with old age when responding.

After senility the most frequently reported conditions were arthritis or rheumatism (34 percent) and heart trouble (34 percent). Arthritis or rheumatism is a category which represents a variety of diseases that are generally considered to be associated with the degeneration and resultant inflammation of joint tissue.

For noninstitutionalized persons 65 years and over during 1969-70 it was the leading cause of limitation of activity, affecting 21 percent of the persons limited in activity.⁹ As seen in table C, the proportion of the current nursing home population reported to have arthritis or rheumatism (34 percent) was around the level reported in the 1969 survey. In the 5 years between the 1964 and 1969 surveys, however, the proportion increased from 22 to 33 percent, although the age distribution of the population remained relatively constant.

During 1973-74 heart trouble was third in the ranking of prevalence. The rate of 34 percent from the current survey was nearly the same as the 1969 rate, but higher than the 1964 rate (28 percent). Like senility, heart trouble rep-

resents a very broad often ill-defined disease category. This classification can include rheumatic heart disease, hypertensive heart disease, ischemic heart disease, and other diseases of the circulatory system. Thus some respondents may have included residents with arteriosclerosis, or hardening of the arteries, and possibly some of those suffering from cerebrovascular disease, or stroke. Nevertheless, heart trouble would undoubtedly remain among the top three conditions for reported prevalence in the nursing home population even if these miscellaneous categories were eliminated. During 1972 heart conditions were reported by the Health Interview Survey (HIS) as the most prevalent condition among noninstitutionalized persons with limitations of activity.¹ It was second only to arthritis or rheumatism as an activity limiting condition among persons 65 years and over during 1969-70.⁹

At 19 percent, mental illness was the fourth most reported condition in the 1973-74 nursing home population. Although the prevalence rate remained constant since 1964, this comparison ignores the fact that in the current survey mental retardation was a distinct category separate from mental illness, whereas in the previous two surveys both mental illness and mental retardation were combined in one category (labeled mental illness). Although in the 1973-74 NNHS only 7 percent of the residents were reported as mentally retarded, the addition of mentally retarded residents to those with mental illness gives a total of 25 percent with mental disorders. This represents a substantial increase from the 1969 prevalence rate of 18 percent. Among the 1969-70 noninstitutionalized population only about 4 percent of all persons with activity limitations reported mental and nervous conditions as the chronic condition causing their limitation.⁹

The fifth most prevalent chronic condition among nursing home residents was another broad category—amputation of extremities or limbs; or permanent stiffness or any deformity of the foot, leg, fingers, arm, or back (to be abbreviated amputation, stiffness, or deformity). As shown in table C, 14 percent of all residents were included under this category as having at least one of these orthopedic impairments. This

rate is similar to that for 1964 but represents a 10 percent drop since 1969. The factors influencing this decline were not apparent.

An estimated 13 percent of the nursing home population were reported to be diabetic. Among those with activity limitations in the noninstitutionalized population during 1969-70, HIS reported diabetes as the cause of the limitation for 4 percent of those persons 65 years and over.⁹

Eleven percent or less of the nursing home population were reported to have the following conditions: paralysis or palsy due to stroke (11 percent), glaucoma or cataracts (10 percent), chronic trouble with back or spine (10 percent), mental retardation (7 percent), and paralysis or palsy not related to stroke, arthritis, or rheumatism (6 percent).

Age and Sex

Comparisons across age and sex groups in the nursing home population yielded some interesting patterns in the prevalence of the 11 reported conditions. As illustrated in Figure 2, the prevalence of several of the conditions increased markedly with age. The top three in overall prevalence—senility, arthritis or rheumatism, and heart trouble—all approximated a fourfold increase when rates for residents under 65 years of age were compared with those residents 85 years and over. Particularly noteworthy was the fact that around 7 out of every 10 of the older residents were reported to be senile. Glaucoma or cataracts also approximated a fourfold increase between the youngest and the oldest groups. As a result of these trends, the four conditions mentioned above rank-ordered as the most prevalent among older residents.

In contrast to the pattern of increased prevalence, several conditions markedly decreased with age—mental illness; mental retardation; paralysis not related to stroke, arthritis, or rheumatism; and amputation, stiffness, or deformity. In general these were the ones with the highest prevalence rates among younger residents, or those under 65 years. Mental illness and mental retardation were the most prevalent conditions reported for younger residents. The prevalence of mental retardation showed the steepest




Figure 2. Prevalence rates per 1,000 nursing home residents for selected reported chronic conditions, by age: United States, August 1973-April 1974

decline with age. Figure 2 depicts a drop in rate of nearly twentyfold between the youngest and the oldest age groups.

Comparing male and female residents as groups, differences were found in the prevalence

of reported conditions; but these were relatively small in comparison to the age differences. An overall sex-age comparison on prevalence of reported conditions can be found in table 1.

Mental retardation was the only condition

for which the prevalence rate among male residents was distinctly higher than that reported for females. The rate of 100 per 1,000 males was nearly twice the rate for females (54 per 1,000). Other conditions for which the prevalence rate among men exceeded that for women included amputation, stiffness, or deformity, and paralysis or palsy both due to stroke and other causes.

Female residents experienced higher prevalence rates on five of the reported conditions—arthritis or rheumatism, senility, heart trouble, glaucoma or cataracts, and chronic back or spine trouble. The greatest sex differential was estimated for arthritis or rheumatism, 379 cases per 1,000 women as compared to 256 per 1,000 males.

Certification Status of Facility

Provided in table 2 are the prevalence patterns of reported conditions according to the certification status of the home. For several conditions these patterns varied considerably among the different types of facilities. Regardless of certification status senility remained as the most frequently reported condition.

Both Medicare and Medicaid (BM&M) certified homes accounted for the highest rates of residents with paralysis or palsy due to stroke, heart trouble, and paralysis or palsy not due to stroke. At 14 percent the rate of stroke after-effects among residents of BM&M's was slightly higher than that for the SNH's or ICF's, while it was considerably higher than the rate of 8 percent reported for residents of not certified facilities. In general this pattern was repeated for residents with heart trouble and those with other types of paralysis or palsy. In these cases, however, the differences in rates between BM&M's and SNH's were not statistically significant.

Conversely, residents in BM&M's accounted for the lowest rates of mental illness and mental retardation. With rates over 9 percent, mentally retarded residents were much more prevalent in ICF's and homes not certified than they were in other facilities. Other conditions which varied to any significant degree by certification status were diabetes and amputation, stiffness, or de-

formity. The lowest rates for both these conditions were reported among residents of not certified homes, with estimates of about 10 percent.

Although not statistically significant for several conditions, the figures suggest that BM&M's and SNH's tended to provide care for residents with more medically serious conditions such as heart trouble or paralysis or palsy; whereas ICF's and not certified homes had larger shares of residents with conditions such as mental illness and mental retardation which require custodial care.

PRIMARY DIAGNOSIS

To complement the data on reported chronic conditions and impairments, the nurse respondent was asked to consult the medical record of each sample resident to obtain medical information based on a physician's diagnosis. The respondent was asked to extract the *primary* diagnosis from both the most current examination and from the examination given when the resident was admitted. For each of these specific time references the primary diagnosis was then classified according to the categories listed in item 8 of the Resident Questionnaire. This classification was fashioned to correspond to the major disease categories of the ICDA.⁸ Exception to this format was made in the major category Diseases of the Circulatory System. As noted in table D, this category was subdivided according to the ICDA classification into heart attack, stroke, hardening of the arteries, and other diseases of the circulatory system.

Primary Diagnosis at Last Examination

A primary diagnosis at last examination was obtained for 99 percent of the population. Since by definition only one condition was noted, these estimates are not to be considered actual prevalence rates. Instead, this question was intended to determine the condition which was currently the most serious one affecting the resident. It is assumed that the difference between the rates of a reported condition and its corresponding primary diagnosis is a rough

Table D. Number and percent distribution of nursing home residents by primary diagnosis at last examination and rank in August 1973- April 1974 and June-August 1969: United States

Primary diagnosis at last examination in rank order for 1973-74	1973-74			1969 ¹		
	Rank	Number of residents	Percent distribution	Rank	Number of residents	Percent distribution
Diseases of the circulatory system ²	---	---	---	1	298,400	39.1
Hardening of the arteries.....	1	241,800	22.5	---	---	---
Senility, old age, or other symptoms and ill-defined conditions.....	2	146,800	13.6	3	77,200	10.1
Mental disorders ³	3	115,800	10.8	2	84,600	11.1
Stroke.....	4	113,400	10.5	---	---	---
Diseases of the musculoskeletal system and connective tissue.....	5	73,100	6.8	5	49,500	6.5
Diseases of the nervous system and sense organs.....	6	64,200	6.0	4	57,700	7.6
Heart attack.....	7	55,700	5.2	---	---	---
Accidents, poisonings, and violence.....	8	49,300	4.6	7	36,700	4.8
Endocrine, nutritional, and metabolic diseases.....	9	48,100	4.5	6	39,400	5.2
Diseases of the circulatory system other than hardening of the arteries, stroke or heart attack.....	10	39,400	3.7	---	---	---
Neoplasms.....	11	25,600	2.4	11	16,900	2.2
Diseases of the respiratory system.....	12	22,200	2.1	12	14,900	2.0
Diseases of the digestive system.....	13	20,500	1.9	10	18,100	2.4
Other diagnoses ⁴	14	16,100	1.5	---	---	---
Diseases of the genitourinary system.....	15	15,600	1.5	13	8,900	1.2
Unknown diagnoses.....	16	9,600	0.9	9	24,400	3.2
Diseases of the blood and blood-forming organs.....	17	7,600	0.7	14	3,900	0.5
Diseases of the skin and subcutaneous tissue.....	18	6,000	0.6	15	2,800	0.4
Congenital anomalies.....	19	3,100	0.3	16	1,900	0.3
Infective and parasitic diseases.....	20	*	*	8	28,700	3.8

¹See reference 11 for source of estimates.

²In the 1973-74 National Nursing Home Survey, this category was subdivided into specific categories of hardening of the arteries, stroke, heart attack, and other diseases of the circulatory system. These categories were not reported separately in the 1969 survey.

³Includes mental retardation and mental illness.

⁴Includes certain causes of perinatal morbidity, complications of pregnancy and childbirth, and other diagnoses not listed.

indication of overall severity. For example, the more severe the condition the less difference there should be between its reported prevalence rate and its frequency as a primary diagnosis. Table 3 provides a cross-tabulation of primary diagnosis at last examination and reported conditions to permit this comparison.

Table D provides a rank ordering of primary diagnosis at last examination and a comparison with data from the 1969 survey.¹⁰ Hardening of the arteries was the primary diagnosis for 23 percent of all the residents and was highly related to reported conditions (table 3). It was the most frequent primary diagnosis for the following conditions: senility; arthritis or rheumatism; glaucoma or cataracts; amputation, stiffness, or deformity; and heart trouble. For all other con-

ditions (except mental retardation) it was the second most frequent primary diagnosis.

Although senility was reported to affect 58 percent of the nursing home population, its frequency as a primary diagnosis was far less. At 14 percent it was the second most frequent primary diagnosis. (Actually, the percent of residents whose primary diagnosis was senility was slightly less than 14 percent because the classification also included old age, other symptoms, and ill-defined conditions such as coma and uremia.) The difference in the reported rate (58 percent) and the frequency of primary diagnosis (14 percent) is due to the fact that senility had been considered a secondary rather than a primary condition at the last examination. There is also the possibility, mentioned

earlier, that the respondents could have over-reported this condition because it was so broadly defined.

Mental disorder was the major diagnosis for an estimated 11 percent of the residents. This rate is only about half of that reported as a chronic condition (19 percent). Table 3 shows that only about 38 percent of those residents with reported mental illness and 53 percent of those with reported mental retardation had mental disorders as their primary diagnosis. Apparently there were more severe conditions affecting residents with reported mental problems.

The fourth most prevalent primary diagnosis was stroke. Nearly 11 percent of all residents had this condition as the primary diagnosis at their last examination. This rate was nearly comparable to its reported prevalence rate of 11 percent, a fact indicating the severity of this chronic condition. Diseases of the musculo-skeletal system and connective tissue comprised a category that included many other chronic conditions besides arthritis or rheumatism. As seen in table D, the diagnosed rate for this ICDA category was 7 percent, far below that of the reported rate for arthritis or rheumatism at 34 percent. The category of diseases of the nervous system and sense organs represented a primary diagnostic rate of 6 percent. This diagnosis should have encompassed two major reported conditions—glaucoma or cataracts and paralysis and palsy not related to stroke, arthritis or rheumatism.

Although approximately 34 percent of all residents reportedly had heart trouble, only an estimated 5 percent had heart attack as their primary diagnosis. With an overall ranking of seventh in prevalence, this diagnosis specifically referred to those residents with ischemic heart disease. Tenth in overall ranking, residents with diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack accounted for about 4 percent of all primary diagnoses. Adding this to the rate for heart attack would yield better comparability with the classification for reported heart trouble, but the total of around 9 percent would still fall far short of the reported prevalence figure (34 percent). An examination of table 3

reveals the reason for this discrepancy. Hardening of the arteries was by far the most prevalent diagnosis of residents with reported heart trouble.

The category of accidents, poisonings, and violence was eighth in the rank ordering of primary diagnoses, with a rate of 5 percent. It was assumed that the great majority of the residents in this category had suffered from some type of bodily injury, most particularly a fracture resulting from accidents. In support of this assumption, a fracture of the femur had been reported for about 3 percent of the 1964 nursing home population.⁴

In ninth position in overall ranking of primary diagnoses, the category of endocrine, nutritional, and metabolic diseases yielded a rate of nearly 5 percent. Based on table 3, there was little doubt residents with diabetes comprised the largest proportion of this rate. The reported rate of diabetes (13 percent) was almost 3 times greater than the diagnosed rate for the broader category of endocrine, nutritional, and metabolic diseases, which possibly included other conditions such as goiter.

The remaining 10 categories of primary diagnoses account for only 12 percent of the total. Each was estimated to have a rate of about 2 percent or less.

With regard to comparisons between the 1973-74 NNHS and the 1969 survey, table D shows that relatively little change occurred in the pattern of primary diagnoses. About the only conspicuous difference was that for the rate of infective and parasitic diseases. This category, which includes primarily acute diseases, occupied a significant position among primary diagnoses in 1969; it ranked eighth and included 4 percent of the residents. In 1973-74, however, the rate was not even reportable because so few residents were in this category. A comparison to the 1969 survey can be made for a primary diagnosis involving diseases of the circulatory system by combining the four sub-categories used in the 1973-74 survey—hardening of the arteries, stroke, heart attack, and other diseases of the circulatory system. The resulting total rate of 42 percent for 1973-74 was only slightly higher than the 39 percent obtained in the 1969 survey.




Figure 3. Rate per 1,000 nursing home residents for selected primary diagnoses at last examination, by age: United States, August 1973-April 1974

Age and sex.—The frequency of many primary diagnoses varied greatly with age. Figure 3 illustrates the age trends for selected diagnoses. Mental disorders were the most prevalent conditions affecting residents under 65 years. As depicted in figure 3, the primary diagnosis rate of mental disorders dropped dramatically as age increased, from 396 per 1,000 under 65 years to 31 per 1,000 for 85 years or older. The rate for diseases of the nervous system and sense organs at 156 cases per 1,000 was the second most frequent primary diagnosis for the younger residents. This classification, too, showed an unmistakable decline in rate with age to 35 cases per 1,000 for those 85 years and over.

Stroke was the third most frequent primary diagnosis among residents under 65 years of age

and the third ranked diagnosis for those over 65. The highest rates were for residents 65-74 and 75-84 years of age; the lowest, for residents under 65 years and those 85 years and over.

Hardening of the arteries ranked seventh among the young residents and showed the greatest increase in rates with age to rank first among older residents. Nearly 30 percent of those 85 years and over had this disease as their primary diagnosis at last examination. The next largest increase with age was in the category of senility, old age, other symptoms, and ill-defined conditions. With a rate of 185 per 1,000, these conditions were the second most prevalent primary diagnoses for residents 85 years and over. In contrast, the rate was 21 per 1,000 for those under 65 years.

Other diagnoses with increased rates with age were diseases of the musculoskeletal system and connective tissue; heart attack; and diseases of the circulatory system other than heart attack, stroke, and hardening of the arteries.

In contrast, diseases of the respiratory system showed a small but significant decrease with age. Primary diagnosis by sex and age is presented in table 1. One of the diagnoses representing the greatest overall sex differential was hardening of the arteries. The rate for males was 190 per 1,000; the rate for females, 239 per 1,000. This difference appeared mainly among residents at least 65 years old. Female residents also experienced the highest rate of senility, old age, or other symptoms and ill-defined conditions. In addition, women had a higher rate of musculoskeletal diseases and accidents, poisonings, and violence than men had.

In contrast, men had higher rates than females for mental disorders and stroke. The rate for mental disorders was 143 per 1,000 males versus 93 per 1,000 females. When examined across age groups, this difference was not consistent. The sex difference in rates for stroke was more evident among residents over 65 years of age.

Among the less prevalent diagnoses there were two significant sex differences that were clearly evident. The rate for male residents with either diseases of the respiratory system or diseases of the genitourinary system was 3 times higher than that for female residents. Sex comparison on these conditions by age was not possible because of the small numbers in the younger age groups.

Certification status of facility.—Comparisons of how the current primary diagnosis varied according to certification status can be made from data presented in table 2. A primary diagnosis of stroke was observed to occur more frequently among residents of BM&M's than among residents of the other types of facilities. In fact, with a rate of 13 percent it was the second most prevalent primary diagnosis in these institutions, whereas it was the fourth or fifth leading diagnosis in other homes. Hardening of the arteries, which was the most frequent primary diagnosis for all facilities, was at its greatest rate of 25 percent among residents of

SNH's. Accidents, poisonings, and violence constituted a primary diagnosis for which the rate among residents of BM&M homes was unmistakably greater than that for residents in the other facilities. The rate of 7 percent was 2 times higher than that in other types of homes.

In contrast, the pattern of primary diagnosis for mental disorders showed the lowest rate of 8 percent among residents of BM&M's. The highest rate of 16 percent was found in not certified homes, where mental disorders were the second most frequent primary diagnosis. Not certified homes also had the highest rate for diseases of the musculoskeletal system and connective tissue.

An overall rate for diseases of the circulatory system can be obtained by combining rates for heart attack, stroke, hardening of the arteries, and other diseases of the circulatory system. This procedure is possible since only one diagnosis per resident was given. Examining this overall rate by certification status, it is evident that diseases of the circulatory system were most prevalent among residents of BM&M's and SNH's, with rates of 45 percent and 44 percent, respectively. The rate was 40 percent in ICF's and only 33 percent in not certified homes. An opposite pattern is obtained when the categories for mental disorders and senility, old age, or other symptoms and ill-defined conditions are combined to provide a category for "mental problems." In this case, 32 percent of the residents in not certified homes are included in this category, whereas about 28 percent in ICF's are included, 24 in SNH's, and only 20 percent in BM&M's.

Primary Diagnosis at Admission

Information abstracted from medical records permitted classification of the primary diagnosis at the time of admission for over 99 percent of the residents. These data were considered to indicate the primary medical reasons for persons initially seeking institutionalization. Presented in figure 4 are the few primary diagnoses which showed significant differences between the rate at admission and that at last examination. Tables 1 and 2 provide additional data on the pattern of primary diagnosis at admission according to




Figure 4. Comparison of rates for those primary diagnoses showing significant changes between admission and last examination: United States, August 1973-April 1974

the age and sex of the residents and the certification status of the facility. The patterns emerging from an analysis of these variables generally reaffirmed the findings established from the data on primary diagnosis at last examination.

Comparing the admitting diagnoses to the current, the greatest difference in rates occurred in the category of accidents, poisonings, and violence. The difference showed that the number of residents in this category at admission was nearly twice the number at last examination. Nearly 9 percent of persons in nursing homes during the 1973-74 survey period were admitted as a result of injuries received from accidents, poisonings, and violence. Examined across age groups, this category showed the greatest difference between admission and current diagnosis among women, among residents 85 years and over, and among residents in BM&M's.

Rates for stroke represented another notable difference between diagnosis at admission and at last examination. Stroke was more likely to be the primary diagnosis at admission than at the most current examination. As shown in table 2, the prominence of stroke as an admission diagnosis was noted more particularly among

residents of BM&M's. Taken together, stroke and accidents, poisonings, and violence apparently accounted for over a fifth of all admissions to these facilities.

In contrast to this pattern, the difference in rates for residents whose primary diagnosis was senility, old age, or other symptoms and ill-defined conditions was in the opposite direction. This primary diagnosis was more frequent at the most recent examination than at admission. The same trend occurred for the primary diagnoses hardening of the arteries and diseases of the musculoskeletal system. When these three diagnoses were examined by age groups, the greatest differences in rates between the most recent and the admission diagnosis occurred in the eldest age group, 85 years and over.

Any remaining differences in rates between diagnosis at time of admission and time of last examination were not significant. For example, the rates for mental disorders were practically the same.

BEHAVIORAL PROBLEMS

Information on specific behavioral problems listed in item 15 of the Resident Questionnaire provided an additional source of prevalence data related to the state of the resident's mental health. The respondent reported how frequently the resident was depressed, agitated or nervous, abusive or aggressive, or confused or senile, had disturbed sleep, or exhibited other behavioral problems.

As shown in figure 5, the most prevalent problem was being confused or senile. This was the only behavioral problem affecting a majority of the residents, 57 percent being the estimated rate of occurrence. This figure was comparable to the rate of 58 percent for reported senility (see previous section on reported conditions for details). Despite the high prevalence rate, only about 10 percent were reported to exhibit this behavior more often than once a week. Agitation or nervousness was the second most prevalent behavioral problem. It affected over 42 percent of all residents, with 13 percent being affected more often than once a week. Depression was reported for 39 percent of the nurs-




Figure 5. Percent of nursing home residents with behavioral problems by problem and frequency: United States, August 1973-April 1974

ing home population. In terms of the "severity" of the behavioral problems (i.e., those occurring more often than once a week), depression had the highest reported frequency of 14 percent. Disturbed sleep affected a substantially smaller proportion of the population. About 19 percent of all residents apparently had problems affecting their sleep. For around 5 percent this condition occurred more often than once a week. The least prevalent of these problems was abusive or aggressive behavior. Only 17 percent of all residents were reported to display this type of behavior.

Besides the specific problems mentioned, about 5 percent of all residents had other behavioral problems. Included in this category were behavioral problems due to alcoholism, drug addiction, and unspecified reasons. These problems, however, affected only 1 percent of the population more often than once a week.

For purposes of comparison over time, the only available data on behavioral problems from nursing home surveys were from a 1956 survey of 13 States.¹¹ An estimated 56 percent of the residents in that survey were reported to act confused part or most of the time. This rate nearly matched that from the current survey. For data on behavioral problems in the noninsti-

tutionalized population, the only available source was the 1960-62 Health Examination Survey.¹² An estimated 51 percent of all persons aged 65 years and over suffered feelings of nervousness. This rate compared with the rate of 42 percent of the nursing home population reported to be agitated or nervous.

Age and Sex

There were three behavioral problems that occurred more often among residents under 65 years of age than among older residents. In table 4 it can be seen that the youngest residents were more likely to be depressed (46 percent), to be agitated or nervous (52 percent), and to be abusive or aggressive (23 percent). The rate of occurrence of all three problems appeared to decline at every age level, although this trend was not statistically significant throughout. In contrast to the youngest residents, the eldest residents were reported as the least depressed (34 percent), the least agitated or nervous (37 percent), and the least abusive or aggressive (15 percent). Ranging from a rate of 10 percent in the youngest group to 4 percent in the eldest group, the category of other behavioral problems showed a similar trend. Residents 85 years and over were more often described as confused or senile. In fact, about 54 percent of this age group were reported to exhibit this behavior once a week or less. In comparison, only around 27 percent of those under 65 years of age fell into this category. The rate of occurrence for disturbed sleep did not reveal a significant age pattern.

There were significant sex differences in the prevalence of only three behavioral problems. Analogous to the pattern of reported senility, the frequency of confused or senile behavior was 59 percent among female residents, while it was 53 percent among male residents. Depression was also slightly more prevalent among women, 40 percent compared to 36 percent for men. On the other hand, it was very clear that males tended toward more frequent occurrence of abusive or aggressive behavior. Their rate for this problem was 22 percent compared to 14 percent for females. This difference was significant for both younger and older residents.

Certification Status of Facility

The certification status of a facility was related to the prevalence pattern of behavioral problems (table 5). The lowest rate of occurrence for every one of the specified problems was reported for residents in not certified homes. (The difference between the rates for abusive or aggressive behavior in not certified homes as compared to BM&M's was not statistically significant, however.) Although residents of SNH's appeared to have the greatest levels of depression (41 percent), abusive or aggressive behavior (18 percent), confused or senile behavior (60 percent), and disturbed sleep (20 percent) the differences among ICF's, SNH's, and BM&M's were relatively small and often insignificant compared to the decidedly lower occurrence for not certified homes. Residents of ICF's had the greatest rate of agitation or nervousness (45 percent). Table 6 presents data on conditions and primary diagnoses of residents with behavioral problems.

IMPAIRMENTS IN VISION, HEARING, AND SPEECH AND USE OF DENTURES

To complement the data on reported conditions and primary diagnosis, information was also collected on sensory impairments. Referring to questions 17, 18, and 19 of the Resident Questionnaire (appendix III), the respondent was asked whether visual, hearing, and speech impairments existed and if so whether they were partial, severe, or complete. To provide as reliable a status rating as possible, definitions were provided to the respondent. For instance, in order to rate a resident's vision status as partially impaired, the respondent had to judge whether the resident was able to watch television from 8 to 12 feet across the room. To rate a resident's speech as severely impaired, the respondent had to judge that the resident could not carry on a normal conversation and was understood only with difficulty. Questions 17, 18, and 19 of the Resident Questionnaire (appendix III) should be consulted for the definitions used to rate the extent of the resident's sensory impairments. In addition to the ratings of sensory impairments,

information was also obtained on the use of eyeglasses, hearing aids, and dentures.

Vision

As shown in figure 6, the majority of residents (54 percent) were considered to have unimpaired vision, i.e., they could read ordinary newspaper print. About 34 percent of the population had partially impaired vision, while 10 percent had a vision impairment that was considered severe. An estimated 30,400 residents, or about 3 percent of the population, were judged to be blind.

Overall the impairment pattern was similar for males and females. In contrast, differences in the extent of impairment for several age groups were distinctly significant. While the proportion of blind residents was nearly equal for all age groups, the proportion of residents 85 years and over with severe visual impairments (14 percent) was nearly triple that for residents younger than 65 (5 percent). Table 7 provides additional data on visual impairments according to whether or not the resident wore eyeglasses.




Figure 6. Percent distribution of nursing home residents by vision status, according to age and sex: United States, August 1973-April 1974




Figure 7. Percent of nursing home residents using eyeglasses, by age and sex: United States, August 1973-April 1974

Eyeglasses were reportedly used by 62 percent of the population. Figure 7 illustrates how the utilization of eyeglasses varied by age and sex. Since visual impairments are typically associated with the aging process, it was not surprising to find the utilization rate increased with age up to the 75-84 year group; but no difference was observed between those aged 75-84 and those aged 85 years and older. Utilization differed between the sexes also, with female residents having the higher utilization rate. This pattern was probably due in part to the preponderance of older females in the resident population.

Table 8 provides the prevalence pattern of visual impairments according to certification status. Among residents in BM&M's, SNH's, and ICF's, the differences in the degree of impairment were either relatively small or insignificant. The degree of visual impairments in not certified homes, however, appeared to be considerably less than that found in the other types of facilities. The proportion of residents in not certified homes with unimpaired vision was 62 percent, compared to 55 percent in ICF's, 51 percent in SNH's, and 52 percent in BM&M's.

Compared to the 1969 nursing home population, overall vision status for the 1973-74 survey period population appeared to have improved somewhat.⁵ A greater proportion of the 1969 population was counted as severely im-

paired (17 percent versus 10 percent). The proportion of blind residents was similar for both surveys (3-4 percent), as was the proportion of eyeglass users (60-62 percent).

Utilization of eyeglasses by nursing home residents has generally been at a much lower rate than that reported for the older noninstitutionalized population. In three different surveys conducted during 1960-62, 1965-66, and 1971 the rate was estimated to be over 90 percent for noninstitutionalized persons 65 years and over.¹³⁻¹⁵ When comparing vision status, the older noninstitutionalized population has not reported the prevalence of visual impairments anywhere near the 47 percent prevalence of the nursing home population. In 1971 only 5 percent of the noninstitutionalized population 65 years and over had a visual impairment considered serious enough to prevent the person from reading newspaper.¹⁶

Hearing

Impairments to hearing were less common among nursing home residents than were impairments to sight. Figure 8 illustrates that about 68 percent of the population were considered unimpaired in audition, i.e., they could hear a telephone conversation on an ordinary telephone. For around 26 percent of all residents, hearing was judged to be partially impaired. Severe impairments affected only about 5 percent of the population. And only 10,800 persons, or around 1 percent, were judged to be deaf.

As seen in figure 8, sex differentials in the degree of hearing impairment were insignificant. In contrast, age differences in the degree of impairment were well defined. Only about 12 percent of the residents under 65 years of age were impaired, whereas about 45 percent of those 85 years and over were reported to have some type of hearing impairment.

Although about 32 percent of the population was rated as impaired in hearing, only a comparatively small proportion, 4 percent, were reported to use a hearing aid. As depicted in figure 9, there was a pattern of increased utilization of hearing aids with age. There was no significant difference in utilization by sex, however.




Figure 8. Percent distribution of nursing home residents by hearing status, according to age and sex: United States, August 1973-April 1974




Figure 9. Percent of nursing home residents using a hearing aid, by age and sex: United States, August 1973-April 1974

Comparison of audition status on the basis of the use of hearing aids reveals some interesting patterns as shown in table 7. For about 31 percent of the residents using hearing aids, the use of the aid apparently resulted in eliminating the auditory impairment. For the remaining 69 percent, some impairment was evident in spite of the use of the hearing aid. Among those residents not using a hearing aid, 69 percent did not need one since they were considered unimpaired in hearing. As for the remaining 30 percent who had an auditory impairment, the reason for a hearing aid not being used could not be determined from the data. Whether or not the impairment could have been alleviated by the use of an aid remains unknown.

As shown in table 8, when the prevalence of auditory impairments was compared according to the certification status of the facilities, the pattern which emerged was similar to one previously discussed for visual impairments. Little difference in the degree of impairment appeared among residents in BM&M's, SNH's, and ICF's. The prevalence of auditory impairments among the residents of not certified homes was noticeably less than that found among the residents of the other types of homes. An estimated 75 percent of the residents in not certified homes were unimpaired in hearing, while 65 percent in SNH's, 67 percent in BM&M's, and 68 percent in ICF's were assigned this status.

Based on comparisons with data from the 1969 survey, it seemed that audition status, like vision status, improved in the 1973-74 nursing home population.⁵ Proportionately, there were 10 percent more residents in 1969 who had hearing impairments than reported by the current survey (43 percent versus 32 percent). The greatest difference was among those residents with severe impairments, including deafness. In 1969 this proportion was 13 percent, twice that of the more current estimate. This apparent improvement in audition status occurred in spite of any distinct increases in the use rate of hearing aids.

When compared to findings from surveys of the noninstitutionalized population, nursing home residents had more hearing impairments. During 1971 hearing problems were reported by 27 percent of those persons in the non-

stitutionalized population aged 65 years and over.¹⁷

Speech

For the first time in an NCHS survey of nursing homes data were collected on the extent to which a resident's speech was impaired. Figure 10 illustrates that 74 percent of all residents had no speech impairment reported. Of those remaining, the ability to speak was judged to be partially impaired for an estimated 14 percent of the population and to be severely impaired for about 8 percent. Mute residents numbered 32,100 and comprised 3 percent of the population.

As depicted in figure 10, the pattern of speech defects varied appreciably with age. Unlike vision and audition impairments, speech impairments were more prevalent among younger residents, particularly those under 65 years. The proportion of younger residents who had severely impaired speech or were mute was nearly 3 times that for residents 85 years and over. Although less pronounced, there was also a

sex difference in the pattern of speech defects. Twenty-nine percent of the males had some level of impairment as compared to 24 percent of the females.

Similar to the pattern found for vision and auditory impairments across facility types, speech impairments were less prevalent in not certified homes. Table 8 reveals that 80 percent of the residents in not certified homes were not impaired in speech, while 74 percent of residents in ICF's and 73 percent in BM&M's and SNH's were included in this category.

According to the 1971 HIS, about 10 percent of all persons living outside institutions were reported to have a speech defect, which included cleft palate, harelip, stammering, stuttering, and other speech problems.¹⁶

Use of Dentures

Item 20 of the Resident Questionnaire pertained to the use of dentures. This information had not been collected in any previous NCHS survey of nursing homes. The resulting data are presented in figure 11. A sizable majority of




Figure 10. Percent distribution of nursing home residents by speech status, according to age and sex: United States, August 1973-April 1974




Figure 11. Percent distribution of nursing home residents by use of dentures, according to age and sex: United States, August 1973-April 1974

residents (59 percent) had dentures and used them, while 36 percent did not have dentures. An estimated 5 percent of the population had dentures but did *not* use them.

As expected, the use of dentures was related to age. Whereas only 26 percent of those residents under 65 years had dentures, 74 percent of those 85 years and over had dentures. Utilization also varied by sex; about 15 percent more women than men had dentures. According to the data provided in table 8, there was little discernable difference in the use of dentures across the various certification statuses.

SPECIAL AIDS

Data on the use of special aids were analyzed to provide an indication of the extent of chronic disabilities within the nursing home population. Each respondent was asked to specify from among the aids listed in table E those which the resident regularly used.

As shown in table 9, the total number of residents using one or more aids was estimated to be 558,300, 52 percent of the population. The total number of aids in use was estimated to be 662,700, an indication that some residents used more than one special aid.

As shown in table E, some aids were in much greater use than others. The wheelchair was the most frequently used special aid. Approximately 35 percent of all residents were reported to use this device regularly. About 15 percent of the population used a walker for aid in ambulation, while about 6 percent relied upon a cane for assistance. The remaining aids for which specific information was collected all had utilization rates of less than 2 percent. The category "other aids" (geriatric chairs, special shoes, triangle bars, and unclassified aids) included 2 percent of all residents.

Compared to previous nursing home surveys, the utilization pattern of special aids has not changed appreciably.^{6,18} The utilization rate of wheelchairs changed relatively little between 1969 (31 percent) and the 1973-74 survey period (35 percent) following a large increase from 1964, when only 21 percent of the population were reported to use this device. The walker was

the one aid which continued to show a steady and considerable increase in utilization. In 1964 only about 9 percent of the population used walkers, in 1969 only 12 percent used this aid, and in the 1973-74 survey period the rate rose to 15 percent. Utilization of the remaining special aids has remained about the same over the years.

Comparison of the utilization of special aids between nursing home residents and the noninstitutionalized population indicates the high degree of chronic disability of the nursing home population. Rates for wheelchairs and walkers showed the greatest differences in utilization and, thus, in the indicated level of disability for the two populations. According to the 1969 HIS, 3 percent of the noninstitutionalized population 65 years or older used either wheelchairs or walkers.¹⁹ In contrast, 35 percent of the nursing home population used wheelchairs and 15 percent used walkers.

Age and Sex

The pattern of utilization for some special aids varied as a function of age. For those using wheelchairs, table E illustrates that the rate remained practically unchanged (32-34 percent) until the oldest age group where the rate increased significantly (38 percent). There were clear and consistent increases, though, in the rates for walkers (from 6 to 17 percent) and canes (from 3 to 8 percent). In contrast, the use of braces shows a marked decrease between the under 65 and the 85 and over age group (from 4 down to 1 percent). Although the use of crutches appeared to decrease with age, the trend was not significant.

There were some sex differences in the utilization of special aids. Although female and male residents did not differ significantly in their utilization of the wheelchair, females were considerably more likely to be using walkers (17 versus 10 percent). Their utilization of the self-feeder was also significantly higher. On the other hand, males had significantly higher utilization of crutches and canes.

Certification Status of Facility

By examining the utilization of special aids according to certification status in table 10, it is

Table E. Number and percent distribution of nursing home residents by sex and age, according to special aids used: United States, August 1973-April 1974

Sex and age	Number of residents ¹	No aid used	1 aid or more used	Special aid used ²							
				Wheel-chair	Walker	Cane	Crutches	Braces	Artificial limb	Self feeder	Other aids
Number of residents.....	1,075,800	517,500	558,300	374,800	156,400	68,800	9,300	16,700	4,400	11,400	20,900
Both sexes	Percent of residents ³										
All ages.....	1,075,800	48.1	51.9	34.8	14.5	6.4	0.9	1.6	0.4	1.1	1.9
Under 65 years.....	114,300	58.8	41.2	32.2	6.2	2.8	*	4.0	*	*	2.6
65-74 years.....	163,100	55.1	44.9	31.2	10.4	4.9	1.4	2.2	*	*	1.6
75-84 years.....	384,900	48.4	51.6	33.9	15.5	6.8	0.8	1.5	*	1.1	1.9
85 years and over ..	413,600	42.1	57.9	37.9	17.5	7.6	*	0.6	*	1.0	2.0
Male											
All ages.....	318,100	50.2	49.8	33.7	10.0	8.6	1.5	1.6	0.8	0.8	1.8
Under 65 years.....	52,400	60.3	39.7	30.2	5.1	*	*	*	*	*	*
65-74 years.....	65,100	56.4	43.6	31.3	7.4	7.0	*	*	*	*	*
75-84 years.....	102,300	47.4	52.6	35.1	11.0	9.6	*	*	*	*	*
85 years and over ..	98,300	43.7	56.3	35.8	13.2	11.7	*	*	*	*	*
Female											
All ages.....	757,700	47.2	52.8	35.3	16.5	5.5	0.6	1.6	*	1.2	2.0
Under 65 years.....	61,900	57.5	42.5	34.0	7.2	*	*	4.1	*	*	*
65-74 years.....	98,000	54.3	45.7	31.1	12.4	3.6	*	2.5	*	*	*
75-84 years.....	282,600	48.8	51.2	33.5	17.2	5.8	*	1.6	*	1.1	1.9
85 years and over ..	315,300	41.6	58.4	38.5	18.9	6.3	*	0.7	*	1.2	2.0

¹Includes all residents whether or not a special aid is used.

²Excludes eyeglasses and hearing aids.

³Figures may not total to 100 percent because residents may have more than one special aid.

evident that differences existed in the utilization of special aids across the facility types. The greatest difference appeared between BM&M's, where about 60 percent of the residents used one or more special aids, and not certified homes, where only 40 percent of the residents used an aid.

A look at the pattern by type of aid revealed that most of the difference in overall utilization was attributable to the use of wheelchairs. In BM&M's approximately 44 percent of all resi-

dents used a wheelchair. This rate was 10 percent higher than for SNH's, about 14 percent higher than for ICF's, and 21 percent higher than for residents of not certified homes. Little difference, most of it insignificant, appeared among the certification types when the utilization of any of the other types of aids was examined. Data on the chronic conditions and primary diagnoses of residents using special aids are presented in table 11.

REFERENCES

- ¹National Center for Health Statistics: Limitations of activity and mobility due to chronic conditions, United States, 1972. *Vital and Health Statistics*. Series 10-No. 96. DHEW Pub. No. (HRA) 75-1523. Health Resources Administration. Washington, U.S. Government Printing Office, Nov. 1974.
- ²National Center for Health Statistics: Design and methodology for a national survey for nursing homes. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 7. Public Health Service. Washington. U.S. Government Printing Office, Sept. 1968.
- ³National Center for Health Statistics: Chronic illness among residents of nursing and personal care homes, United States, May-June, 1964. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 12-No. 7. Public Health Service. Washington. U.S. Government Printing Office, Mar. 1967.
- ⁴National Center for Health Statistics: Prevalence of chronic conditions and impairments among residents of nursing and personal care homes, United States, May-June 1964. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 12-No. 8. Public Health Service. Washington. U.S. Government Printing Office, July 1967.
- ⁵National Center for Health Statistics: Chronic conditions and impairments of nursing home residents, United States, 1969. *Vital and Health Statistics*. Series 12-No. 22. DHEW Pub. No. (HRA) 74-1707. Health Resources Administration. Washington. U.S. Government Printing Office, Dec. 1973.
- ⁶National Center for Health Statistics: Measures of chronic illness among residents of nursing and personal care homes, United States, June-August 1969. *Vital and Health Statistics*. Series 12-No. 24. DHEW Pub. No. (HRA) 74-1709. Health Resources Administration. Washington. U.S. Government Printing Office, Mar. 1974.
- ⁷National Center for Health Statistics: Home care for persons 55 years and over, United States, July 1966-June 1968. *Vital and Health Statistics*. Series 10-No. 73. DHEW Pub. No. (HSM) 72-1062. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, July 1972.
- ⁸National Center for Health Statistics: *Eighth Revision International Classification of Diseases, Adapted for Use in the United States*. PHS Pub. No. 1693. Public Health Service. Washington. U.S. Government Printing Office, 1967.
- ⁹National Center for Health Statistics: Limitation of activity due to chronic conditions, United States, 1969 and 1970. *Vital and Health Statistics*. Series 10-No. 80. DHEW Pub. No. (HSM) 73-1506. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Apr. 1973.
- ¹⁰National Center for Health Statistics: Charges for care and sources of payment for residents in nursing homes, United States, June-August 1969. *Vital and Health Statistics*. Series 12-No. 21. DHEW Pub. No. (HRA) 74-1706. Health Resources Administration. Washington. U.S. Government Printing Office, July 1973.
- ¹¹Commission on Chronic Illness: Nursing homes, their patients and their care. *Monograph 46*. PHS Pub. No. 503. Public Health Service. Washington. U.S. Government Printing Office, 1956.
- ¹²National Center for Health Statistics: Selected symptoms of psychological distress, United States, 1960-62. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 11-No. 37. Public Health Service. Washington. U.S. Government Printing Office, Aug. 1970.
- ¹³National Center for Health Statistics: History and examination findings related to visual activity among adults, United States, 1960-62. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 11-No. 28. Public Health Service. Washington. U.S. Government Printing Office, Mar. 1968.
- ¹⁴National Center for Health Statistics: Characteristics of persons with corrective lenses, United States, July 1965-June 1966. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 10-No. 53. Public Health Service. Washington. U.S. Government Printing Office, June 1969.
- ¹⁵National Center for Health Statistics: Current estimates from the Health Interview Survey, United States, 1971. *Vital and Health Statistics*. Series 10-No. 79. DHEW Pub. No. (HSM) 73-1505. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Feb. 1973.
- ¹⁶National Center for Health Statistics: Prevalence of selected impairments, United States, 1971. *Vital and Health Statistics*. Series 10-No. 99. DHEW Pub. No. (HRA) 75-1526. Health Resources Administration. Washington. U.S. Government Printing Office, May 1975.
- ¹⁷National Center for Health Statistics: Persons with impaired hearing, United States, 1971. *Vital and Health Statistics*. Series 10-No. 101. DHEW Pub. No. (HRA)

76-1528. Health Resources Administration. Washington. U.S. Government Printing Office, Nov. 1975.

¹⁸National Center for Health Statistics: Use of special aids in homes for the aged and chronically ill, United States, May-June 1964. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 12-No. 11. Public Health Service. Washington. U.S. Government Printing Office. Dec. 1968.

¹⁹National Center for Health Statistics: Use of special aids, United States, 1969. *Vital and Health Statistics*. Series 10-No. 78. DHEW Pub. No. (HSM) 73-1504. Health Services and Mental Health Administration. Washington. U.S. Government Printing Office, Dec. 1972.

²⁰Van Nostrand, J. F.: Development of Survey Methodology to Measure Cost and Quality of Care in Nursing Homes. Paper presented at 101st Annual Meeting of American Public Health Association, San Francisco, Nov. 8, 1973.

²¹National Center for Health Statistics: Development and maintenance of a national inventory of hospitals and institutions. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Feb. 1965.

²²National Center for Health Statistics: Design and methodology of the 1967 Master Facility Inventory Survey. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 9. Public Health Service. Washington. U.S. Government Printing Office, Feb. 1965.

²³National Center for Health Statistics: The Agency Reporting System for Maintaining the National Inventory of Hospitals and Institutions. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 6. Public

Health Service. Washington. U.S. Government Printing Office, Apr. 1968.

²⁴National Center for Health Statistics: Institutions for the aged and chronically ill, United States, April-June 1963. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 12-No. 1. Public Health Service. Washington. U.S. Government Printing Office, July 1965.

²⁵National Center for Health Statistics: Arrangements for physicians services to residents in nursing and personal care homes, United States, May-June 1969. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 12-No. 13. Public Health Service. Washington. U.S. Government Printing Office, Feb. 1970.

²⁶National Center for Health Statistics: Selected characteristics of nursing homes for the aged and chronically ill, United States, June-August 1969. *Vital and Health Statistics*. Series 12-No. 23. DHEW Pub. No. (HRA) 74-1708. Health Resources Administration. Washington. U.S. Government Printing Office, Jan. 1974.

²⁷National Center for Health Statistics: Selected operating and financial characteristics of nursing homes, United States, 1973-74 National Nursing Home Survey. *Vital and Health Statistics*. Series 13-No. 22. DHEW Pub. No. (HRA) 76-1773. Health Resources Administration. Washington. U.S. Government Printing Office, Dec. 1975.

²⁸National Center for Health Statistics: Inpatient health facilities as reported from the 1971 Master Facility Inventory Survey. *Vital and Health Statistics*. Series 14-No. 12. DHEW Pub. No. (HRA) 74-1807. Health Resources Administration. Washington. U.S. Government Printing Office, Feb. 1970.

LIST OF DETAILED TABLES

1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974	25
2. Number and percent distribution of nursing home residents by reported chronic conditions and impairments and primary diagnosis at last examination and at admission, according to certification status of home: United States, August 1973-April 1974	30
3. Percent distribution of nursing home residents by primary diagnosis at last examination, according to reported chronic conditions and impairments: United States, August 1973-April 1974	32
4. Number and percent of nursing home residents with behavioral problems by type of problem, according to sex and age: United States, August 1973-April 1974	33
5. Number and percent of nursing home residents with behavioral problems by type of problem, according to certification status of home: United States, August 1973-April 1974	35
6. Number and percent of nursing home residents with behavioral problems by type of problem, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974	36
7. Number and percent distribution of nursing home residents by vision, hearing, speech, and use of denture status, according to sex and age: United States, August 1973-April 1974	37
8. Number and percent distribution of nursing home residents by vision, hearing, speech, and use of denture status, according to certification status of home: United States, August 1973-April 1974	38
9. Number and percent of nursing home residents by special aids used, according to sex and age: United States, August 1973-April 1974	40
10. Number and percent of nursing home residents by special aids used, according to certification status of home: United States, August 1973-April 1974	42
11. Number and percent distribution of nursing home residents by special aids used, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974	43

Table 1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974

Sex, reported chronic conditions and impairments, and primary diagnosis at last examination and at admission	Number of residents	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
<u>BOTH SEXES</u>						
Number of residents-----	1,075,800	1,075,800	114,300	163,100	384,900	413,600
<u>Reported chronic conditions and impairments</u>		Rate per 1,000 residents				
Senility-----	627,200	583.0	189.1	482.2	623.2	694.3
Mental illness-----	200,400	186.3	348.5	293.0	171.9	112.8
Mental retardation-----	72,800	67.7	345.7	94.5	27.6	17.5
Arthritis or rheumatism-----	368,500	342.5	109.7	232.4	363.2	431.1
Paralysis or palsy due to stroke-----	122,100	113.5	106.6	166.1	129.1	80.1
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	66,200	61.6	161.5	75.5	52.6	36.8
Glaucoma or cataracts-----	110,900	103.1	38.5	62.8	95.6	143.9
Diabetes-----	142,700	132.6	112.0	168.1	148.8	109.4
Any chronic trouble with back or spine-----	106,100	98.6	74.7	86.6	100.8	107.9
Amputation of extremities or limbs, or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	150,000	139.4	189.2	160.8	133.0	123.2
Heart trouble-----	360,500	335.1	110.2	271.9	354.7	403.9
None of the above-----	57,700	53.6	64.5	62.2	53.9	47.0
<u>Primary diagnosis at last examination</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	146,800	136.5	20.6	85.2	140.9	184.6
Heart attack-----	55,700	51.8	*	41.1	55.3	64.1
Stroke-----	113,400	105.4	94.0	138.0	120.6	81.5
Hardening of the arteries-----	241,800	224.7	36.6	151.7	237.2	293.9
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	39,400	36.6	*	31.4	39.9	40.4
Accidents, poisonings, and violence-----	49,300	45.8	39.6	35.8	45.8	51.4
Mental disorders-----	115,800	107.6	395.8	185.1	72.1	30.5
Diseases of the musculoskeletal system and connective tissue-----	73,100	67.9	48.3	58.5	70.7	74.4
Endocrine, nutritional, and metabolic diseases-----	48,100	44.7	44.0	59.5	46.9	37.1
Diseases of the respiratory system-----	22,200	20.6	*	33.3	22.9	13.7
Neoplasms-----	25,600	23.8	27.9	29.4	23.6	20.7
Diseases of the nervous system and sense organs-----	64,200	59.7	156.0	78.4	49.3	35.4
Diseases of the digestive system-----	20,500	19.0	*	18.6	17.9	20.1
Infective and parasitic diseases-----	*	*	*	*	*	*
Diseases of the genitourinary system-----	15,600	14.5	*	*	16.7	15.1
Diseases of the skin and subcutaneous tissue-----	6,000	5.6	*	*	6.1	*
Diseases of the blood and blood-forming organs-----	7,600	7.1	*	*	7.6	8.8
Congenital anomalies-----	3,100	2.9	19.1	*	*	*
Other diagnoses ¹ -----	16,100	15.0	22.1	15.9	15.0	12.6
Unknown diagnoses-----	9,600	8.9	*	*	9.3	9.8

See footnote at end of table.

Table 1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974—Con.

Sex, reported chronic conditions and impairments, and primary diagnosis at last examination and at admission	Number of residents	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
<u>BOTH SEXES—Con.</u>						
<u>Primary diagnosis at admission</u>		Rate per 1,000 residents				
Senility, old age, or other symptoms and ill-defined conditions-----	110,700	102.9	15.0	68.7	104.0	139.7
Heart attack-----	59,600	55.4	*	46.4	58.9	68.6
Stroke-----	132,800	123.4	104.9	165.9	142.3	94.2
Hardening of the arteries-----	219,900	204.4	27.4	138.3	212.4	272.1
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	34,300	31.8	*	23.9	36.5	35.4
Accidents, poisonings, and violence-----	95,700	89.0	55.0	54.6	89.9	111.0
Mental disorders-----	122,100	113.5	421.0	194.6	77.3	30.2
Diseases of the musculoskeletal system and connective tissue-----	62,600	58.2	47.0	51.3	63.7	58.8
Endocrine, nutritional, and metabolic diseases-----	44,700	41.6	34.9	50.9	45.0	36.6
Diseases of the respiratory system-----	23,100	21.5	18.6	31.9	22.9	16.8
Neoplasms-----	24,200	22.5	28.1	28.0	23.0	18.3
Diseases of the nervous system and sense organs-----	60,500	56.2	147.9	74.3	46.3	32.9
Diseases of the digestive system-----	21,800	20.3	*	18.2	19.4	22.6
Infective and parasitic diseases-----	2,300	2.2	*	*	*	*
Diseases of the genitourinary system-----	15,300	14.2	*	*	15.8	15.8
Diseases of the skin and subcutaneous tissue-----	5,000	4.7	*	*	6.4	*
Diseases of the blood and blood-forming organs-----	6,700	6.2	*	*	*	8.2
Congenital anomalies-----	2,800	2.6	*	*	*	*
Other diagnoses ¹ -----	24,800	23.0	22.7	21.5	22.7	24.1
Unknown diagnoses-----	6,900	6.4	*	*	5.3	9.7
<u>MALE</u>						
All residents-----	318,100	318,100	52,400	65,100	102,300	98,300
<u>Reported chronic conditions and impairments</u>		Rate per 1,000 residents				
Senility-----	169,900	534.0	170.4	482.4	615.8	677.0
Mental illness-----	63,700	200.1	334.8	274.8	164.1	116.3
Mental retardation-----	31,600	99.5	362.4	104.1	37.9	*
Arthritis or rheumatism-----	81,300	255.6	79.2	180.2	298.1	355.6
Paralysis or palsy due to stroke-----	42,000	131.9	104.9	193.4	147.9	88.8
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	23,600	74.3	167.4	78.7	61.6	34.9
Glaucoma or cataracts-----	28,500	89.6	40.4	*	85.5	141.5
Diabetes-----	39,100	122.9	101.6	147.5	137.3	102.8
Any chronic trouble with back or spine-----	23,800	74.8	71.0	68.0	77.6	78.5
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	48,600	152.8	203.2	176.5	151.6	111.6
Heart trouble-----	100,400	315.5	104.7	280.3	350.0	415.2
None of the above-----	19,400	60.9	70.2	69.0	57.0	54.5

See footnote at end of table.

Table 1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974—Con.

Sex, reported chronic conditions and impairments, and primary diagnosis at last examination and at admission	Number of residents	All ages	Rate per 1,000 residents			
			Under 65 years	65-74 years	75-84 years	85 years and over
<u>MALE—Con.</u>						
<u>Primary diagnosis at last examination</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	35,900	112.8	*	67.0	125.4	179.9
Heart attack-----	14,500	45.5	*	41.4	49.5	60.9
Stroke-----	38,000	119.5	96.7	156.0	132.2	94.4
Hardening of the arteries-----	60,600	190.4	*	130.1	226.9	275.7
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	11,000	34.6	*	*	41.2	42.1
Accidents, poisonings, and violence-----	9,700	30.6	48.0	*	27.1	27.9
Mental disorders-----	45,400	142.7	419.4	201.9	73.6	28.0
Diseases of the musculoskeletal system and connective tissue-----	16,300	51.2	*	45.3	56.1	57.5
Endocrine, nutritional, and metabolic diseases-----	12,900	40.4	*	53.2	39.0	38.4
Diseases of the respiratory system-----	12,100	38.0	*	57.0	43.9	23.4
Neoplasms-----	9,700	30.6	*	*	30.9	32.8
Diseases of the nervous system and sense organs-----	22,400	70.4	152.8	73.7	56.3	39.0
Diseases of the digestive system-----	6,700	21.1	*	23.5	*	*
Infective and parasitic diseases-----	*	*	*	*	*	*
Diseases of the genitourinary system-----	8,900	28.0	*	*	35.6	35.6
Diseases of the skin and subcutaneous tissue-----	*	*	*	*	*	*
Diseases of the blood and blood-forming organs-----	*	*	*	*	*	*
Congenital anomalies-----	*	*	*	*	*	*
Other diagnoses ¹ -----	5,200	16.5	*	*	*	*
Unknown diagnoses-----	2,900	9.0	*	*	*	*
<u>Primary diagnosis at admission</u>						
Senility, old age, other symptoms and ill-defined conditions-----	27,300	85.9	*	54.9	86.6	145.2
Heart attack-----	17,200	54.0	*	51.9	60.9	72.0
Stroke-----	44,900	141.2	108.2	188.9	159.4	108.4
Hardening of the arteries-----	54,800	172.3	*	112.4	196.9	264.8
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	1,000	31.3	*	*	40.4	39.7
Accidents, poisonings, and violence-----	16,700	52.5	60.5	41.6	52.0	56.0
Mental disorders-----	49,400	155.2	454.4	207.7	86.8	32.0
Diseases of the musculoskeletal system and connective tissue-----	13,300	41.7	34.9	37.3	45.7	44.0
Endocrine, nutritional, and metabolic diseases-----	11,400	35.7	*	43.3	39.2	33.9
Diseases of the respiratory system-----	12,200	38.3	*	50.4	44.9	28.7
Neoplasms-----	9,200	29.1	*	*	29.9	28.3
Diseases of the nervous system and sense organs-----	20,900	65.7	141.7	70.9	54.4	33.6
Diseases of the digestive system-----	7,000	22.0	*	23.3	*	25.9
Infective and parasitic diseases-----	*	*	*	*	*	*
Diseases of the genitourinary system-----	9,500	29.8	*	*	36.6	41.7
Diseases of the skin and subcutaneous tissue-----	*	*	*	*	*	*
Diseases of the blood and blood-forming organs-----	*	*	*	*	*	*
Congenital anomalies-----	*	*	*	*	*	*
Other diagnoses ¹ -----	7,400	23.2	*	*	25.6	22.2
Unknown diagnoses-----	*	*	*	*	*	*

See footnote at end of table.

Table 1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974—Con.

Sex, reported chronic conditions and impairments, and primary diagnosis at last examination and at admission	Number of residents	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
<u>FEMALE</u>						
All residents-----	757,700	757,700	61,900	98,000	282,600	315,300
<u>Reported chronic conditions and impairments</u>			Rate per 1,000 residents			
Senility-----	457,300	603.6	205.0	482.1	625.8	699.7
Mental illness-----	136,800	180.5	360.0	305.0	174.8	111.7
Mental retardation-----	41,100	54.3	331.6	88.1	23.8	16.7
Arthritis or rheumatism-----	287,200	379.0	135.4	267.1	386.8	454.7
Paralysis or palsy due to stroke-----	80,100	105.8	108.1	147.9	122.3	77.4
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	42,600	56.2	156.5	73.4	49.3	37.4
Glaucoma or cataracts-----	82,400	108.8	*	66.6	99.3	144.6
Diabetes-----	103,600	136.8	120.9	181.8	152.9	111.4
Any chronic trouble with back or spine-----	82,300	108.6	77.9	99.0	109.3	117.1
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	101,400	133.8	177.2	150.3	126.3	126.8
Heart trouble-----	260,100	343.3	114.9	266.3	356.4	400.4
None of the above-----	38,300	50.6	59.7	57.6	52.7	44.7
<u>Primary diagnosis at last examination</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	11,100	146.4	*	97.3	146.6	186.1
Heart attack-----	41,200	54.4	*	40.9	57.4	65.1
Stroke-----	75,400	99.5	91.7	126.1	116.5	77.5
Hardening of the arteries-----	181,200	239.1	38.8	166.1	240.9	299.6
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	28,400	37.5	*	32.1	39.4	39.8
Accidents, poisonings, and violence-----	39,500	52.1	32.4	42.1	52.6	58.8
Mental disorders-----	70,400	92.9	375.8	173.9	71.5	31.2
Diseases of the musculoskeletal system and connective tissue-----	56,800	74.9	57.7	67.3	76.0	79.7
Endocrine, nutritional, and metabolic diseases-----	35,300	46.5	55.0	63.7	49.8	36.6
Diseases of the respiratory system-----	10,100	13.3	*	*	15.3	10.7
Neoplasms-----	15,900	21.0	*	28.6	21.0	16.9
Diseases of the nervous system and sense organs-----	41,800	55.2	158.8	81.4	46.8	34.2
Diseases of the digestive system-----	13,800	18.2	*	*	17.8	19.6
Infective and parasitic diseases-----	*	*	*	*	*	*
Diseases of the genitourinary system-----	6,700	8.9	*	*	9.8	8.7
Diseases of the skin and subcutaneous tissue-----	4,000	5.3	*	*	*	*
Diseases of the blood and blood-forming organs-----	5,400	7.1	*	*	7.8	8.1
Congenital anomalies-----	*	*	*	*	*	*
Other diagnoses ¹ -----	10,900	14.3	*	*	14.4	11.2
Unknown diagnoses-----	6,800	8.9	*	*	8.9	10.2

See footnote at end of table.

Table 1. Number of nursing home residents and rate per 1,000 residents by sex, reported chronic conditions and impairments and primary diagnosis at last examination and at admission, and age: United States, August 1973-April 1974—Con.

Sex, reported chronic conditions and impairments, and primary diagnosis at last examination and at admission	Number of residents	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
FEMALE—Con.		Rate per 1,000 residents				
<u>Primary diagnosis at admission</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	83,400	110.1	*	77.9	110.3	138.0
Heart attack-----	42,500	56.0	*	42.7	58.1	67.5
Stroke-----	87,900	115.9	102.0	150.6	136.1	89.8
Hardening of the arteries-----	165,100	217.9	*	155.6	218.0	274.4
Diseases of the circulatory system other than hardening of the arteries, stroke, or heart attack-----	24,300	32.1	*	26.1	35.1	34.0
Accidents, poisonings, and violence-----	79,000	104.3	50.3	63.2	103.7	128.2
Mental disorders-----	72,800	96.0	392.8	185.9	73.9	29.6
Diseases of the musculoskeletal system and connective tissue-----	49,300	65.1	57.2	60.7	70.2	63.4
Endocrine, nutritional, and metabolic diseases-----	33,400	44.0	45.2	55.8	47.1	37.4
Diseases of the respiratory system-----	10,900	14.4	*	*	14.9	13.0
Neoplasms-----	14,900	19.7	*	25.5	20.5	15.2
Diseases of the nervous system and sense organs-----	39,600	52.2	153.2	76.5	43.4	32.7
Diseases of the digestive system-----	14,800	19.5	*	*	19.3	21.6
Infective and parasitic diseases-----	*	*	*	*	*	*
Diseases of the genitourinary system-----	5,800	7.7	*	*	8.3	7.7
Diseases of the skin and subcutaneous tissue-----	3,300	4.3	*	*	*	*
Diseases of the blood and blood-forming organs-----	4,700	6.2	*	*	*	*
Congenital anomalies-----	*	*	*	*	*	*
Other diagnoses ¹ -----	17,400	23.0	*	*	21.6	24.7
Unknown diagnoses-----	5,500	7.2	*	*	*	10.5

¹Includes certain causes of perinatal morbidity, complications of pregnancy and childbirth, and other diagnoses not listed above.

Table 2. Number and percent distribution of nursing home residents by reported chronic conditions and impairments and primary diagnosis at last examinations and at admission, according to certification status of home: United States, August 1973-April 1974

Reported chronic conditions and impairments and primary diagnoses at last examination and at admission	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing home only ²	Intermediate care facility only	
Number of residents					
All residents-----	1,075,800	406,900	292,500	236,700	139,800
<u>Reported chronic conditions and impairments³</u>					
Senility-----	627,200	235,900	179,300	141,400	70,700
Mental illness-----	200,400	58,900	60,100	52,800	28,800
Mental retardation-----	72,800	18,300	20,100	21,500	12,900
Arthritis or rheumatism-----	368,500	125,600	106,100	90,100	46,700
Paralysis or palsy due to stroke-----	122,100	55,400	32,200	23,700	10,900
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	66,200	27,100	18,300	14,000	6,800
Glaucoma or cataracts-----	110,900	40,900	34,300	23,200	12,600
Diabetes-----	142,700	57,400	43,400	28,700	13,300
Any chronic trouble with back or spine-----	106,100	38,700	30,700	24,100	12,500
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arms, or back-----	150,000	56,700	44,900	34,400	14,000
Heart trouble-----	360,500	149,600	101,100	71,400	38,400
None of the above-----	57,700	23,900	13,700	9,600	10,600
<u>Primary diagnosis at last examination</u>					
Senility, old age, or other symptoms and ill-defined conditions-----	146,800	50,300	39,000	34,900	22,500
Heart attack-----	55,700	19,400	16,400	14,200	5,700
Stroke-----	113,400	53,700	28,000	21,500	10,200
Hardening of the arteries-----	241,800	92,900	74,400	50,700	23,700
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	39,400	15,200	8,800	8,800	6,700
Accidents, poisonings, and violence-----	49,300	26,900	10,100	7,800	4,500
Mental disorders-----	115,800	30,500	30,900	31,700	22,700
Diseases of the musculoskeletal system and connective tissues-----	73,100	22,000	20,200	18,200	12,600
Endocrine, nutritional, and metabolic diseases-----	48,100	18,100	14,400	10,700	4,900
Diseases of the respiratory system-----	22,200	9,500	6,200	3,800	2,700
Neoplasms-----	25,600	12,400	6,100	4,300	2,900
Diseases of the nervous system, sense organs-----	64,200	24,200	17,300	15,600	7,200
Diseases of the digestive system-----	20,500	7,700	6,000	3,600	3,200
Diseases of the genitourinary system-----	15,600	6,700	4,800	2,500	*
Other diagnoses ⁴ -----	34,700	15,100	7,600	5,500	6,600
Unknown diagnoses-----	9,600	*	2,400	3,000	*
<u>Primary diagnosis at admission</u>					
Senility, old age, or other symptoms and ill-defined conditions-----	110,700	34,700	30,600	26,400	19,000
Heart attack-----	59,600	21,600	18,300	14,400	5,300
Stroke-----	132,800	61,100	33,800	26,000	11,900
Hardening of the arteries-----	219,900	77,500	67,800	50,600	24,000
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	34,300	14,300	7,400	7,400	5,300
Accidents, poisonings, and violence-----	95,700	50,400	22,000	15,500	7,800
Mental disorders-----	122,100	29,800	31,800	36,000	24,600
Diseases of the musculoskeletal system and connective tissue-----	62,600	20,500	18,000	14,600	9,500
Endocrine, nutritional, and metabolic diseases-----	44,700	15,200	14,400	10,400	4,700
Diseases of the respiratory system-----	23,100	10,500	5,600	4,600	*
Neoplasms-----	24,200	12,400	5,300	3,400	3,000
Diseases of the nervous system and sense organs-----	60,500	21,900	16,800	15,200	6,600
Diseases of the digestive system-----	21,800	9,300	6,500	3,000	3,000
Diseases of the genitourinary system-----	15,300	7,900	4,200	*	*
Other diagnoses ⁴ -----	41,600	17,400	8,200	5,700	10,400
Unknown diagnoses-----	6,900	*	*	*	*

See footnotes at end of table.

Table 2. Number and percent distribution of nursing home residents by reported chronic conditions and impairments and primary diagnosis at last examinations and at admission, according to certification status of home: United States, August 1973-April 1974—Con.

Reported chronic conditions and impairments and primary diagnoses at last examination and at admission	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing home only ²	Intermediate care facility only	
<u>Reported chronic conditions and impairments³</u>		Percent distribution ³			
Senility-----	58.3	58.0	61.3	59.8	50.6
Mental illness-----	18.6	14.5	20.5	22.3	20.6
Mental retardation-----	6.8	4.5	6.9	9.1	9.2
Arthritis and rheumatism-----	34.3	30.9	36.3	38.1	33.4
Paralysis or palsy due to stroke-----	11.3	13.6	11.0	10.0	7.8
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	6.2	6.7	6.3	5.9	4.9
Glaucoma or cataracts-----	10.3	10.0	11.7	9.8	9.0
Diabetes-----	13.3	14.1	14.8	12.1	9.5
Any chronic trouble with back or spine-----	9.9	9.5	10.5	10.2	9.0
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arms, or back-----	13.9	13.9	15.4	14.5	10.0
Heart trouble-----	33.5	36.8	34.6	30.2	27.5
None of the above-----	5.4	5.9	4.7	4.0	7.6
<u>Primary diagnosis at last examination</u>					
All primary diagnoses-----	100.0	100.0	100.0	100.0	100.0
Senility, old age, or other symptoms and ill-defined conditions-----	13.6	12.4	13.3	14.8	16.1
Heart attack-----	5.2	4.8	5.6	6.0	4.1
Stroke-----	10.5	13.2	9.6	9.1	7.3
Hardening of the arteries-----	22.5	22.8	25.4	21.4	17.0
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	3.7	3.7	3.0	3.7	4.7
Accidents, poisonings, and violence-----	4.6	6.6	3.4	3.3	3.2
Mental disorders-----	10.8	7.5	10.6	13.4	16.3
Diseases of the musculoskeletal system and connective tissue-----	6.8	5.4	6.9	7.7	9.0
Endocrine, nutritional, and metabolic diseases-----	4.5	4.5	4.9	4.5	3.5
Diseases of the respiratory system-----	2.1	2.3	2.1	1.6	2.0
Neoplasms-----	2.4	3.0	2.1	1.8	2.1
Diseases of the nervous system and sense organs-----	6.0	5.9	5.9	6.6	5.1
Diseases of the digestive system-----	1.9	1.9	2.0	1.5	2.3
Diseases of the genitourinary system-----	1.5	1.6	1.7	1.0	*
Other diagnoses ⁴ -----	3.2	3.7	2.6	2.3	4.7
Unknown diagnoses-----	0.9	*	0.8	1.3	*
<u>Primary diagnosis at admission</u>					
All primary diagnoses-----	100.0	100.0	100.0	100.0	100.0
Senility, old age, or other symptoms and ill-defined conditions-----	10.3	8.5	10.5	11.1	13.6
Heart attack-----	5.5	5.3	6.3	6.1	3.8
Stroke-----	12.3	15.0	11.6	11.0	8.5
Hardening of the arteries-----	20.4	19.1	23.2	21.4	17.2
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	3.2	3.5	2.5	3.1	3.8
Accidents, poisonings, and violence-----	8.9	12.4	7.5	6.5	5.6
Mental disorders-----	11.4	7.3	10.9	15.2	17.6
Diseases of the musculoskeletal system and connective tissues-----	5.8	5.0	6.2	6.2	6.8
Endocrine, nutritional, and metabolic diseases-----	4.2	3.7	4.9	4.4	3.4
Diseases of the respiratory system-----	2.1	2.6	1.9	2.0	*
Neoplasms-----	2.2	3.1	1.8	1.4	2.2
Diseases of the nervous system and sense organs-----	5.6	5.4	5.7	6.4	4.7
Diseases of the digestive system-----	2.0	2.3	2.2	1.2	2.2
Diseases of the genitourinary system-----	1.4	1.9	1.4	*	*
Other diagnoses ⁴ -----	3.9	4.3	2.8	2.4	7.4
Unknown diagnoses-----	0.6	*	*	*	*

¹23,500 of the residents were in facilities certified by Medicare only.

²129,600 of the residents were in facilities certified as both SNH and ICF.

³Figures may not add to 100 percent because residents may have had more than one reported condition or impairment.

⁴Includes diseases of the blood and blood-forming organs, diseases of the skin and subcutaneous tissue, congenital anomalies, infective and parasitic diseases, certain causes of perinatal morbidity, complications of pregnancy and childbirth, and other diagnoses not listed above.

Table 3. Percent distribution of nursing home residents by primary diagnosis at last examination, according to reported chronic conditions and impairments: United States, August 1973-April 1974

Primary diagnosis at last examination	Reported chronic conditions and impairments ¹					
	Senility	Mental illness	Mental retardation	Arthritis or rheumatism	Paralysis or palsy due to stroke	Paralysis or palsy, etc. ²
Number of residents-----	627,200	200,400	72,800	368,500	122,100	66,200
	Percent distribution					
All diagnoses-----	100.0	100.0	100.0	100.0	100.0	100.0
Senility, old age, or other symptoms and ill-defined conditions-----	22.0	10.6	4.8	13.1	6.2	6.1
Heart attack-----	4.9	3.4	*	6.1	2.3	*
Stroke-----	10.5	6.5	3.9	7.5	60.5	*
Hardening of the arteries-----	26.2	17.0	8.2	25.6	12.3	16.5
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	3.2	2.2	*	3.5	2.3	*
Accidents, poisonings, and violence-----	4.2	2.3	*	4.7	*	6.4
Mental disorders-----	7.5	37.9	52.8	4.5	*	9.9
Diseases of the musculoskeletal system and connective tissue-----	4.7	3.1	*	17.3	2.4	7.4
Endocrine, nutritional, and metabolic diseases-----	3.7	3.8	3.6	3.7	2.9	3.5
Diseases of the respiratory system-----	1.7	1.6	*	1.6	*	*
Neoplasms-----	1.8	1.4	*	1.7	*	*
Diseases of the nervous system and sense organs-----	4.4	5.6	10.2	4.2	2.6	32.6
Diseases of the digestive system-----	1.5	1.8	*	2.0	*	*
Diseases of the genitourinary system-----	1.4	*	*	1.4	*	*
Other diagnoses ² -----	1.9	1.7	5.0	2.5	*	4.0
Unknown diagnoses-----	*	*	*	*	*	*

Primary diagnosis at last examination	Reported chronic conditions and impairments ¹					
	Glaucoma or cataracts	Diabetes	Any chronic trouble with back or spine	Amputation, etc. ¹	Heart trouble	None of these
Number of residents-----	110,900	142,700	106,100	150,000	360,500	57,700
	Percent distribution					
All diagnoses-----	100.0	100.0	100.0	100.0	100.0	100.0
Senility, old age, or other symptoms and ill-defined conditions-----	12.7	8.6	10.1	8.3	10.1	4.6
Heart attack-----	6.1	4.6	5.3	3.9	14.0	*
Stroke-----	8.0	10.3	7.7	12.7	10.6	5.8
Hardening of the arteries-----	26.8	18.6	21.3	19.4	30.8	16.6
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	3.0	3.1	2.7	3.3	5.5	5.4
Accidents, poisonings, and violence-----	3.1	3.4	5.3	7.5	3.0	10.9
Mental disorders-----	5.3	7.0	5.4	7.9	4.1	*
Diseases of the musculoskeletal system and connective tissue-----	8.2	4.0	21.6	11.4	4.7	*
Endocrine, nutritional, and metabolic diseases-----	4.7	29.3	3.0	6.1	4.2	*
Diseases of the respiratory system-----	*	*	*	*	2.5	5.4
Neoplasms-----	2.3	1.6	3.1	1.9	1.7	8.7
Diseases of the nervous system and sense organs-----	11.5	3.5	6.7	8.0	3.1	8.2
Diseases of the digestive system-----	*	*	*	*	1.6	5.4
Diseases of the genitourinary system-----	*	*	*	*	1.5	*
Other diagnoses ² -----	2.9	2.3	3.0	4.6	2.0	12.2
Unknown diagnoses-----	*	*	*	*	*	6.3

¹For complete category titles, see table 1.

²Includes diseases of blood and blood-forming organs, diseases of the skin and subcutaneous tissue, congenital anomalies, infective and parasitic diseases, certain causes of perinatal morbidity, complications of pregnancy and childbirth, and other diagnoses not listed above.

Table 4. Number and percent of nursing home residents with behavioral problems by type of problem, according to sex and age: United States, August 1973-April 1974

Behavioral problems	Both sexes				
	All ages	Under 65 years	65-74 years	75-84 years	85 years and over
	Number of residents				
All residents ¹ -----	1,075,800	114,300	163,100	384,900	413,600
Depressed-----	418,800	52,500	70,900	154,600	140,700
Once a week or less-----	263,900	31,600	44,600	100,200	87,500
More often than once a week-----	154,900	20,900	26,300	54,500	53,200
Agitated or nervous-----	455,400	59,900	75,800	166,200	153,400
Once a week or less-----	318,100	42,800	52,000	117,500	105,800
More often than once a week-----	137,300	17,100	23,800	48,800	47,600
Abusive or aggressive-----	177,800	26,200	29,500	60,800	61,300
Once a week or less-----	114,100	16,000	19,300	39,400	39,400
More often than once a week-----	63,700	10,200	10,200	21,400	21,900
Confused or senile-----	613,200	38,000	80,800	228,300	266,200
Once a week or less-----	510,800	30,500	65,600	190,900	223,800
More often than once a week-----	102,400	7,400	15,200	37,400	42,400
Disturbed sleep-----	201,400	20,100	32,200	74,900	74,200
Once a week or less-----	144,500	14,600	22,700	54,800	52,400
More often than once a week-----	56,900	5,500	9,600	20,000	21,800
Other behavioral problems-----	55,800	11,200	10,700	19,700	14,300
Once a week or less-----	45,300	9,100	8,200	16,300	11,700
More often than once a week-----	10,500	*	2,500	3,400	2,600
	Percent of residents ²				
Depressed-----	38.9	46.0	43.4	40.2	34.0
Once a week or less-----	24.5	27.6	27.3	26.0	21.2
More often than once a week-----	14.4	18.3	16.1	14.1	12.9
Agitated or nervous-----	42.3	52.4	46.5	43.2	37.1
Once a week or less-----	29.6	37.4	31.9	30.5	25.6
More often than once a week-----	12.8	15.0	14.6	12.7	11.5
Abusive or aggressive-----	16.5	22.9	18.1	15.8	14.8
Once a week or less-----	10.6	14.0	11.8	10.2	9.5
More often than once a week-----	5.9	8.9	6.3	5.6	5.3
Confused or senile-----	57.0	33.2	49.5	59.3	64.4
Once a week or less-----	47.5	26.7	40.2	49.6	54.1
More often than once a week-----	9.5	6.5	9.3	9.7	10.2
Disturbed sleep-----	18.7	17.6	19.8	19.5	17.9
Once a week or less-----	13.4	12.8	13.9	14.2	12.7
More often than once a week-----	5.3	4.8	5.9	5.2	5.3
Other behavioral problems-----	5.2	9.8	6.5	5.1	3.5
Once a week or less-----	4.2	7.9	5.0	4.2	2.8
More often than once a week-----	1.0	*	1.5	0.9	0.6

See footnotes at end of table.

Table 4. Number and percent of nursing home residents with behavioral problems by type of problem, according to sex and age: United States, August 1973-April 1974—Con.

Behavioral problems	Male			Female		
	All ages	Under 65 years	65 years and over	All ages	Under 65 years	65 years and over
Number of residents						
All residents ¹ -----	318,100	52,400	265,700	757,700	61,900	695,800
Depressed-----	114,000	20,900	93,200	304,700	31,700	273,100
Once a week or less-----	71,200	13,100	58,100	192,700	18,500	174,200
More often than once a week---	42,900	7,800	35,100	112,000	13,200	98,900
Agitated or nervous-----	133,700	26,900	106,800	321,700	33,000	288,600
Once a week or less-----	91,300	19,200	72,000	226,800	23,600	203,200
More often than once a week---	42,400	7,600	34,800	94,900	9,500	85,400
Abusive or aggressive-----	70,900	14,600	56,300	106,900	11,600	95,300
Once a week or less-----	42,700	8,600	34,100	71,400	7,400	64,000
More often than once a week---	28,200	6,000	22,200	35,500	4,200	31,300
Confused or senile-----	170,000	16,200	153,800	443,200	21,700	421,500
Once a week or less-----	137,900	13,500	124,400	372,900	17,000	355,900
More often than once a week---	32,100	2,700	29,400	70,300	4,700	65,600
Disturbed sleep-----	57,000	8,000	49,000	144,300	12,000	132,300
Once a week or less-----	41,200	6,200	35,000	103,300	8,400	94,900
More often than once a week---	15,800	*	14,000	41,000	3,600	37,400
Other behavioral problems-----	18,600	5,800	12,800	37,200	5,300	31,800
Once a week or less-----	14,600	4,800	9,800	30,700	4,300	26,400
More often than once a week---	4,000	*	3,000	6,500	*	5,400
Percent of residence ²						
Depressed-----	35.9	39.8	35.1	40.2	51.2	39.3
Once a week or less-----	22.4	25.0	21.9	25.4	29.9	25.0
More often than once a week---	13.5	14.9	13.2	14.8	21.3	14.2
Agitated or nervous-----	42.0	51.3	40.2	42.5	53.4	41.5
Once a week or less-----	28.7	36.7	27.1	29.9	38.1	29.2
More often than once a week---	13.3	14.6	13.1	12.5	15.3	12.3
Abusive or aggressive-----	22.3	27.8	21.2	14.1	18.7	13.7
Once a week or less-----	13.4	16.4	12.8	9.4	11.9	9.2
More often than once a week---	8.9	11.4	8.4	4.7	6.8	4.5
Confused or senile-----	53.4	31.0	57.9	58.5	35.1	60.6
Once a week or less-----	43.4	25.8	46.8	49.2	27.5	51.2
More often than once a week---	10.1	5.2	11.1	9.3	7.6	9.4
Disturbed sleep-----	17.9	15.4	18.4	19.1	19.4	19.0
Once a week or less-----	13.0	11.8	13.2	13.6	13.6	13.6
More often than once a week---	5.0	*	5.3	5.4	5.8	5.4
Other behavioral problems-----	5.9	11.1	4.8	4.9	8.6	4.6
Once a week or less-----	4.6	9.1	3.7	4.1	6.9	3.8
More often than once a week---	1.3	*	1.1	0.9	*	0.8

¹Includes all residents whether or not they had behavioral problems.

²Includes only those residents with behavioral problems. Figures may not total to 100 because residents may have had more than one behavioral problem.

Table 5. Number and percent of nursing home residents with behavioral problems by type of problem, according to certification status of home: United States, August 1973-April 1974

Behavioral problems	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing homes ²	Intermediate care facilities	
Number of residents					
All residents ³ -----	1,075,800	406,900	292,500	236,700	139,800
Depressed-----	418,800	163,600	120,900	92,200	42,100
Once a week or less-----	263,900	106,700	79,900	54,300	23,000
More often than once a week-----	154,900	56,900	40,900	37,900	19,200
Agitated or nervous-----	455,400	170,100	127,500	107,400	50,300
Once a week or less-----	318,100	118,800	90,800	73,800	34,700
More often than once a week-----	137,300	51,400	36,700	33,600	15,600
Abusive or aggressive-----	177,800	65,600	53,300	39,700	19,100
Once a week or less-----	114,100	42,300	36,200	23,800	11,700
More often than once a week-----	63,700	23,400	17,000	15,900	7,400
Confused or senile-----	613,200	234,900	175,700	135,900	66,800
Once a week or less-----	510,800	198,700	148,600	109,300	54,200
More often than once a week-----	102,400	36,200	27,100	26,600	12,600
Disturbed sleep-----	201,400	75,000	59,300	45,400	21,600
Once a week or less-----	144,500	53,900	43,300	32,300	15,000
More often than once a week-----	56,900	21,100	16,000	13,100	6,600
Other behavioral problems-----	55,800	22,000	16,900	10,200	6,700
Once a week or less-----	45,300	18,500	13,800	7,900	5,100
More often than once a week-----	10,500	3,600	3,100	*	*
Percent of residents with behavioral problems ⁴					
Depressed-----	38.9	40.2	41.3	39.0	30.1
Once a week or less-----	24.5	26.2	27.3	22.9	16.4
More often than once a week-----	14.4	14.0	14.0	16.0	13.7
Agitated or nervous-----	42.3	41.8	43.6	45.4	36.0
Once a week or less-----	29.6	29.2	31.0	31.2	24.8
More often than once a week-----	12.8	12.6	12.5	14.2	11.2
Abusive or aggressive-----	16.5	16.1	18.2	16.8	13.7
Once a week or less-----	10.6	10.4	12.4	10.1	8.4
More often than once a week-----	5.9	5.7	5.8	6.7	5.3
Confused or senile-----	57.0	57.7	60.1	57.4	47.8
Once a week or less-----	47.5	48.8	50.8	46.2	38.8
More often than once a week-----	9.5	8.9	9.3	11.2	9.0
Disturbed sleep-----	18.7	18.4	20.3	19.2	15.5
Once a week or less-----	13.4	13.2	14.8	13.6	10.8
More often than once a week-----	5.3	5.2	5.5	5.6	4.7
Other behavioral problems-----	5.2	5.4	5.8	4.3	4.8
Once a week or less-----	4.2	4.5	4.7	3.4	3.7
More often than once a week-----	1.0	0.9	1.1	*	*

¹23,500 of the residents were in facilities certified by Medicare only.

²129,600 of the residents were in facilities certified as both SNH and ICF.

³Includes all residents whether or not they had behavioral problems.

⁴Figures may not total to 100 because residents may have had more than one behavioral problem.

Table 6. Number and percent of nursing home residents with behavioral problems by type of problem, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974

Reported chronic conditions and impairments and primary diagnosis at last examination	Number of residents ¹	Behavioral problems					
		De-pressed	Agita-ted or nervous	Abusive or aggres-sive	Con-fused or senile	Dis-turbed sleep	Other behav-ioral prob-lems
		Percent of residents with behavioral problems ²					
All residents-----	1,075,800	38.9	42.3	16.5	57.0	18.7	5.2
<u>Reported chronic conditions and impairments</u>							
Senility-----	627,200	41.8	48.3	19.6	82.7	22.6	5.3
Mental illness-----	200,400	54.4	65.4	31.2	69.5	29.2	11.3
Mental retardation-----	72,800	37.7	54.9	25.2	48.2	18.3	9.4
Arthritis or rheumatism-----	368,500	43.0	44.3	16.2	59.5	22.3	4.2
Paralysis or palsy due to stroke-----	122,100	46.9	41.2	15.4	57.0	18.8	4.1
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	66,200	46.1	49.0	17.1	52.3	20.2	7.0
Glaucoma or cataracts-----	110,900	42.7	43.7	17.5	59.8	23.3	4.9
Diabetes-----	142,700	42.7	43.0	16.4	56.7	18.8	4.3
Any chronic trouble with back or spine-----	106,100	49.8	50.8	18.0	58.5	26.4	6.6
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arms, or back-----	150,000	46.9	46.7	20.7	58.9	22.9	6.6
Heart trouble-----	360,500	42.2	43.5	16.2	61.1	22.0	4.5
None of the above-----	57,700	23.5	21.5	7.0	12.4	8.9	*
<u>Primary diagnosis at last examination</u>							
Senility, old age, or other symptoms and ill-defined conditions-----	146,800	34.2	42.4	17.2	79.7	19.4	4.2
Heart attack-----	55,700	40.6	42.0	13.5	54.3	21.2	*
Stroke-----	113,400	43.7	39.4	14.5	57.1	17.1	4.5
Hardening of the arteries-----	241,800	35.8	41.2	15.7	64.9	18.7	4.4
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	39,400	43.2	42.4	16.6	49.4	18.2	6.1
Accidents, poisonings, and violence-----	49,300	35.6	36.5	14.2	50.3	17.0	*
Mental disorders-----	115,800	44.1	56.8	26.4	55.0	20.0	11.2
Diseases of the musculoskeletal system and connective tissue-----	73,100	37.2	34.6	12.8	39.1	17.4	4.3
Endocrine, nutritional, and metabolic diseases-----	48,100	40.1	38.7	16.5	48.7	15.8	*
Diseases of the respiratory system-----	22,200	41.8	43.7	15.1	44.1	23.1	*
Neoplasms-----	25,600	49.5	39.3	12.1	48.5	24.2	*
Diseases of the nervous system and sense organs-----	64,200	42.9	49.9	18.5	45.3	17.9	5.8
Diseases of the digestive system-----	20,500	43.2	38.9	12.6	47.6	20.3	*
Diseases of the genitourinary system-----	15,600	40.4	42.9	21.0	53.0	21.7	*
Other diagnoses ³ -----	34,700	30.9	34.7	10.5	34.5	16.5	*
Unknown diagnoses-----	9,600	25.1	27.4	*	33.1	*	*

¹Includes all residents whether or not they had behavioral problems.

²Figures may not total to 100 percent because residents may have had more than one behavioral problem.

³Includes diseases of the blood and blood-forming organs, diseases of the skin and subcutaneous tissue, congenital anomalies, infective and parasitic diseases, certain causes of perinatal morbidity, complications of pregnancy and childbirth, and other diagnoses not listed above.

Table 7. Number and percent distribution of nursing home residents by vision, hearing, speech, and use of denture status, according to sex and age: United States, August 1973-April 1974

Vision, hearing, speech, and use of dentures status	Number of residents	Sex			Age			
		Both sexes	Male	Female	Under 65 years	65-74 years	75-84 years	85 years and over
Number of residents-----	1,075,800	1,075,800	318,100	757,700	114,300	163,100	384,900	413,600
<u>Vision status</u>		Percent distribution						
All residents-----	1,075,800	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	575,500	53.5	56.5	52.2	70.6	62.3	53.8	45.0
Sight partially impaired-----	362,000	33.7	31.5	34.6	21.7	28.8	35.0	37.6
Sight severely impaired-----	107,900	10.0	9.0	10.5	5.0	6.3	8.9	14.0
Sight completely lost-----	30,400	2.8	3.0	2.7	2.7	2.6	2.3	3.5
All residents using eyeglasses-----	670,400	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	371,400	55.4	57.0	54.9	65.6	63.3	57.1	49.9
Sight partially impaired-----	226,600	33.8	32.0	34.4	28.8	29.3	33.8	35.9
Sight severely impaired-----	66,500	9.9	9.9	9.9	5.4	6.5	8.4	13.0
Sight completely lost-----	6,000	0.9	1.0	0.8	*	*	0.7	1.1
All residents not using eyeglasses-----	405,400	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	204,100	50.3	55.9	47.0	73.2	61.2	47.0	34.5
Sight partially impaired-----	135,500	33.4	30.9	34.9	18.0	28.3	37.5	41.2
Sight severely impaired-----	41,400	10.2	8.0	11.5	4.7	6.0	9.9	15.9
Sight completely lost-----	24,500	6.0	5.2	6.5	4.0	4.5	5.6	8.4
<u>Hearing status</u>								
All residents-----	1,075,800	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	729,600	67.8	66.9	68.2	88.4	80.3	70.0	55.2
Hearing partially impaired-----	280,300	26.1	26.7	25.8	9.2	17.0	25.5	34.8
Hearing severely impaired-----	55,200	5.1	5.2	5.1	1.6	1.9	3.8	8.7
Hearing completely lost-----	10,800	1.0	1.2	0.9	*	*	0.7	1.4
All residents using hearing aids-----	45,900	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	14,100	30.6	32.1	30.0	*	45.1	32.6	27.1
Hearing partially impaired-----	20,400	44.4	42.6	45.0	*	*	42.2	48.0
Hearing severely impaired-----	10,400	22.7	23.7	22.3	*	*	23.1	22.5
Hearing completely lost-----	*	*	*	*	*	*	*	*
All residents not using hearing aids-----	1,029,900	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	715,500	69.4	68.3	70.0	88.9	81.1	71.4	57.2
Hearing partially impaired-----	259,900	25.2	26.0	24.9	9.0	16.7	24.9	33.8
Hearing severely impaired-----	44,700	4.3	4.5	4.3	*	1.5	3.0	7.7
Hearing completely lost-----	9,700	0.9	1.1	0.9	*	*	0.7	1.3
<u>Speech status</u>								
All residents-----	1,075,800	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Speech not impaired-----	799,100	74.3	70.6	75.8	58.6	70.7	76.2	78.2
Speech partially impaired-----	155,200	14.4	16.8	13.4	18.2	15.9	13.9	13.3
Speech severely impaired-----	89,500	8.3	9.3	7.9	15.5	9.7	7.4	6.7
Speech completely lost-----	32,100	3.0	3.3	2.8	7.6	3.8	2.5	1.8
<u>Use of dentures status</u>								
All residents-----	1,075,800	100.0	100.0	100.0	100.0	100.0	100.0	100.0
No dentures-----	388,800	36.1	46.8	31.7	74.2	48.6	30.5	25.9
Dentures and used them-----	632,800	58.8	48.7	63.1	22.3	47.7	64.4	68.1
Dentures but did not use them--	54,200	5.0	4.5	5.3	3.4	3.7	5.1	5.9

Table 8. Number and percent distribution of nursing home residents by vision, hearing, speech, and use of dentures status, according to certification status of home: United States, August 1973-April 1974

Vision, hearing, speech, and use of dentures status	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing homes ²	Intermediate care facilities	
<u>Vision status</u>		Number of residents			
All residents-----	1,075,800	406,900	292,500	236,700	139,800
Sight not impaired-----	575,500	210,900	148,400	129,600	86,600
Sight partially impaired-----	362,000	145,300	101,900	75,400	39,400
Sight severely impaired-----	107,900	39,700	33,600	23,500	11,200
Sight completely lost-----	30,400	10,900	8,700	8,300	2,600
All residents using eyeglasses-----	670,400	253,700	184,000	140,600	92,200
Sight not impaired-----	371,400	132,700	97,000	82,400	59,300
Sight partially impaired-----	226,600	93,700	64,700	43,900	24,300
Sight severely impaired-----	66,500	25,100	21,000	12,800	7,600
Sight completely lost-----	6,000	*	*	*	*
All residents not using eyeglasses-----	405,400	153,200	108,500	96,100	47,600
Sight not impaired-----	204,100	78,200	51,400	47,200	27,300
Sight partially impaired-----	135,500	51,700	37,200	31,500	15,100
Sight severely impaired-----	41,400	14,600	12,600	10,700	3,500
Sight completely lost-----	24,500	8,600	7,300	6,800	*
<u>Hearing status</u>		Number of residents			
All residents-----	1,075,800	406,900	292,500	236,700	139,800
Hearing not impaired-----	729,600	272,700	190,800	161,400	104,700
Hearing partially impaired-----	280,300	109,600	82,000	60,200	28,500
Hearing severely impaired-----	55,200	20,600	16,800	12,200	5,500
Hearing completely lost-----	10,800	3,900	2,800	2,900	*
All residents using hearing aids-----	45,900	16,500	13,200	9,200	7,000
Hearing not impaired-----	14,100	5,000	3,700	2,700	2,600
Hearing partially impaired-----	20,400	7,100	6,100	4,400	2,800
Hearing severely impaired-----	10,400	3,900	3,300	*	*
Hearing completely lost-----	*	*	*	*	*
All residents not using hearing aids-----	1,029,900	390,400	279,300	227,400	132,700
Hearing not impaired-----	715,500	267,700	187,100	158,700	102,100
Hearing partially impaired-----	259,900	102,500	75,900	55,800	25,700
Hearing severely impaired-----	44,700	16,800	13,500	10,400	4,100
Hearing completely lost-----	9,700	3,500	2,800	2,600	*
<u>Speech status</u>		Number of residents			
All residents-----	1,075,800	406,900	292,500	236,700	139,800
Speech not impaired-----	799,100	297,700	212,800	176,100	112,500
Speech partially impaired-----	155,200	61,200	44,300	33,700	15,900
Speech severely impaired-----	89,500	36,400	25,700	18,700	8,700
Speech completely lost-----	32,100	11,600	9,700	8,100	2,600
<u>Use of dentures status</u>		Number of residents			
All residents-----	1,075,800	406,900	292,500	236,700	139,800
No dentures-----	388,800	139,600	102,200	95,500	51,400
Dentures and used them-----	632,800	243,700	175,700	131,300	82,200
Dentures but did not use them-----	54,200	23,500	14,600	9,900	6,200

See footnotes at end of table.

Table 8. Number and percent distribution of nursing home residents by vision, hearing, speech, and use of dentures status, according to certification status of home: United States, August 1973-April 1974—Con.

Vision, hearing, speech, and use of dentures status	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing homes ²	Intermediate care facilities	
<u>Vision status</u>		Percent distribution			
All residents-----	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	53.5	51.8	50.7	54.7	61.9
Sight partially impaired-----	33.7	35.7	34.8	31.8	28.2
Sight severely impaired-----	10.0	9.8	11.5	9.9	8.0
Sight completely lost-----	2.8	2.7	3.0	3.5	1.9
All residents using eyeglasses-----	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	55.4	52.3	52.7	58.6	64.1
Sight partially impaired-----	33.8	36.9	35.2	31.2	26.3
Sight severely impaired-----	9.9	9.9	11.4	9.1	8.3
Sight completely lost-----	0.9	*	*	*	*
All residents not using eyeglasses-----	100.0	100.0	100.0	100.0	100.0
Sight not impaired-----	50.3	51.1	47.4	49.1	57.2
Sight partially impaired-----	33.4	33.7	34.3	32.8	31.7
Sight severely impaired-----	10.2	9.5	11.6	11.1	7.4
Sight completely lost-----	6.0	5.6	6.7	7.1	*
<u>Hearing status</u>					
All residents-----	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	67.8	67.0	65.2	68.2	74.9
Hearing partially impaired-----	26.1	26.9	28.0	25.4	20.4
Hearing severely impaired-----	5.1	5.1	5.8	5.1	4.0
Hearing completely lost-----	1.0	1.0	1.0	1.2	*
All residents using hearing aids-----	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	30.6	30.3	28.3	29.7	36.9
Hearing partially impaired-----	44.4	43.3	46.2	47.2	39.7
Hearing severely impaired-----	22.7	23.5	25.0	*	*
Hearing completely lost-----	*	*	*	*	*
All residents not using hearing aids-----	100.0	100.0	100.0	100.0	100.0
Hearing not impaired-----	69.4	68.6	67.0	69.8	76.9
Hearing partially impaired-----	25.2	26.3	27.2	24.5	19.4
Hearing severely impaired-----	4.3	4.3	4.8	4.6	3.1
Hearing completely lost-----	0.9	0.9	1.0	1.1	*
<u>Speech status</u>					
All residents-----	100.0	100.0	100.0	100.0	100.0
Speech not impaired-----	74.3	73.2	72.7	74.4	80.4
Speech partially impaired-----	14.4	15.0	15.1	14.3	11.4
Speech severely impaired-----	8.3	8.9	8.8	7.8	6.3
Speech completely lost-----	3.0	2.8	3.3	3.4	1.9
<u>Use of dentures status</u>					
All residents-----	100.0	100.0	100.0	100.0	100.0
No dentures-----	36.1	34.3	35.0	40.4	36.8
Dentures and used them-----	58.8	59.9	60.0	55.5	58.8
Dentures but did not use them-----	5.0	5.8	5.0	4.2	4.4

¹23,500 of the residents were in facilities certified by Medicare only.
²129,600 of the residents were in facilities certified as both SNH and ICF.

Table 9. Number and percent of nursing home residents by special aids used, according to sex and age: United States, August 1973-April 1974

Sex and age	No aid used	1 aid or more used	Special aid used			
			Wheel-chair	Walker	Cane	Crutches
<u>Both sexes</u>			Number of residents			
All ages-----	517,500	558,300	374,800	156,400	68,800	9,300
Under 65 years-----	67,200	47,100	36,800	7,100	3,200	*
65-74 years-----	89,900	73,200	50,900	17,000	8,000	2,400
75-84 years-----	186,300	198,500	130,500	59,800	26,200	3,000
85 years and over-----	174,100	239,500	156,600	72,500	31,400	*
<u>Male</u>						
All ages-----	159,800	158,400	107,300	31,700	27,400	4,600
Under 65 years-----	31,600	20,800	15,800	2,700	*	*
65-74 years-----	36,700	28,400	20,400	4,800	4,600	*
75-84 years-----	48,500	53,800	35,900	11,200	9,800	*
85 years and over-----	42,900	55,400	35,200	13,000	11,500	*
<u>Female</u>						
All ages-----	357,700	400,000	267,500	124,700	41,400	4,700
Under 65 years-----	35,600	26,300	21,000	4,400	*	*
65-74 years-----	53,200	44,800	30,500	12,200	3,500	*
75-84 years-----	137,800	144,700	94,500	48,600	16,400	*
85 years and over-----	131,100	184,100	121,400	59,500	19,800	*
<u>Both sexes</u>			Percent of residents			
All ages-----	48.1	51.9	34.8	14.5	6.4	0.9
Under 65 years-----	58.8	41.2	32.2	6.2	2.8	*
65-74 years-----	55.1	44.9	31.2	10.4	4.9	1.4
75-84 years-----	48.4	51.6	33.9	15.5	6.8	0.8
85 years and over-----	42.1	57.9	37.9	17.5	7.6	*
<u>Male</u>						
All ages-----	50.2	49.8	33.7	10.0	8.6	1.5
Under 65 years-----	60.3	39.7	30.2	5.1	*	*
65-74 years-----	56.4	43.6	31.3	7.4	7.0	*
75-84 years-----	47.4	52.6	35.1	11.0	9.6	*
85 years and over-----	43.7	56.3	35.8	13.2	11.7	*
<u>Female</u>						
All ages-----	47.2	52.8	35.3	16.5	5.5	0.6
Under 65 years-----	57.5	42.5	34.0	7.2	*	*
65-74 years-----	54.3	45.7	31.1	12.4	3.6	*
75-84 years-----	48.8	51.2	33.5	17.2	5.8	*
85 years and over-----	41.6	58.4	38.5	18.9	6.3	*

Table 9. Number and percent of nursing home residents by special aids used, according to sex and age: United States, August 1973-April 1974—Con.

Sex and age	Special aid used			
	Braces	Artificial limb	Self-feeder	Other aids
<u>Both sexes</u>				
Number of residents				
All ages-----	16,700	4,400	11,400	20,900
Under 65 years-----	4,600	*	*	3,000
65-74 years-----	3,600	*	*	2,700
75-84 years-----	5,700	*	4,300	7,100
85 years and over-----	2,700	*	4,300	8,100
<u>Male</u>				
All ages-----	5,000	2,500	2,500	5,600
Under 65 years-----	*	*	*	*
65-74 years-----	*	*	*	*
75-84 years-----	*	*	*	*
85 years and over-----	*	*	*	*
<u>Female</u>				
All ages-----	11,700	*	8,900	15,300
Under 65 years-----	2,500	*	*	*
65-74 years-----	2,400	*	*	*
75-84 years-----	4,400	*	3,200	5,500
85 years and over-----	2,300	*	3,700	6,300
<u>Both sexes</u>				
Percent of residents				
All ages-----	1.6	0.4	1.1	1.9
Under 65 years-----	4.0	*	*	2.6
65-74 years-----	2.2	*	*	1.6
75-84 years-----	1.5	*	1.1	1.9
85 years and over-----	0.6	*	1.0	2.0
<u>Male</u>				
All ages-----	1.6	0.8	0.8	1.8
Under 65 years-----	*	*	*	*
65-74 years-----	*	*	*	*
75-84 years-----	*	*	*	*
85 years and over-----	*	*	*	*
<u>Female</u>				
All ages-----	1.6	*	1.2	2.0
Under 65 years-----	4.1	*	*	*
65-74 years-----	2.5	*	*	*
75-84 years-----	1.6	*	1.1	1.9
85 years and over-----	0.7	*	1.2	2.0

Table 10. Number and percent of nursing home residents by special aids used, according to certification status of home: United States, August 1973-April 1974

Special aid used	All homes	Certification status			
		Both Medicare and Medicaid ¹	Medicaid only		Not certified
			Skilled nursing homes ²	Intermediate care facilities	
Number of residents					
All residents ³ -----	1,075,800	406,900	292,500	236,700	139,800
No aid-----	517,500	161,900	142,900	128,300	84,400
1 aid or more-----	558,300	244,900	149,600	108,400	55,400
Wheelchair-----	374,800	178,500	98,700	67,300	30,300
Walker-----	156,400	65,200	39,700	33,500	18,000
Cane-----	68,800	22,700	20,300	14,800	10,900
Crutches-----	9,300	2,500	2,300	2,800	*
Braces-----	16,700	7,700	4,800	2,800	*
Artificial limb-----	4,400	*	*	*	*
Self-feeder-----	11,400	3,700	*	4,400	*
Other aid-----	20,900	7,600	7,500	4,400	*
Percent of residents ⁴					
No aid-----	48.1	39.8	48.9	54.2	60.4
1 aid or more-----	51.9	60.2	51.1	45.8	39.6
Wheelchair-----	34.8	43.9	33.8	28.4	21.7
Walker-----	14.5	16.0	13.6	14.2	12.9
Cane-----	6.4	5.6	7.0	6.3	7.8
Crutches-----	0.9	0.6	0.8	1.2	*
Braces-----	1.6	1.9	1.7	1.2	*
Artificial limb-----	0.4	*	*	*	*
Self-feeder-----	1.1	0.9	*	1.9	*
Other aid-----	1.9	1.9	2.6	1.8	*

¹23,500 of the residents were in facilities certified by Medicare only.

²129,600 of the residents were in facilities certified as both SNH and ICF.

³Includes all residents whether or not a special aid was used.

⁴Figures may not total to 100 percent because residents may have used more than one special aid.

Table 11. Number and percent distribution of nursing home residents by special aids used, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974

Reported chronic conditions and impairments and primary diagnosis at last examination	Number of residents	Special aid used				
		No aid used	1 aid or more used	Wheelchair	Walker	Cane
Number of residents ¹						
All residents-----	1,075,800	517,500	558,300	374,800	156,400	68,800
<u>Reported chronic conditions and impairments²</u>						
Senility-----	627,200	294,700	332,500	235,700	83,700	34,300
Mental illness-----	200,400	126,700	73,700	51,100	17,800	7,500
Mental retardation-----	72,800	48,700	24,000	18,300	4,300	*
Arthritis or rheumatism-----	368,500	141,100	227,400	143,800	77,200	32,700
Paralysis or palsy due to stroke---	122,100	35,500	86,600	68,400	17,700	7,100
Paralysis or palsy not related to stroke, arthritis, or rheumatism---	66,200	23,600	42,600	33,900	7,900	3,100
Glaucoma or cataracts-----	110,900	49,900	61,100	37,600	19,300	11,600
Diabetes-----	142,700	63,400	79,300	53,500	23,800	10,000
Any chronic trouble with back or spine-----	106,100	35,400	70,700	46,400	24,300	9,000
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	150,000	41,300	108,700	80,900	23,900	10,000
Heart trouble-----	360,500	161,300	199,200	132,600	56,800	28,400
None of the above-----	57,700	35,400	22,300	11,400	8,800	3,400
<u>Primary diagnosis at last examination</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	146,800	74,500	72,300	48,600	18,900	7,300
Heart attack-----	55,700	26,500	29,200	16,300	9,900	5,600
Stroke-----	113,400	38,200	75,200	58,000	16,900	6,600
Hardening of the arteries-----	241,800	120,300	121,400	78,900	32,700	17,900
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	39,400	20,200	19,300	11,000	5,300	4,000
Accidents, poisonings, or violence-----	49,300	10,700	38,600	27,400	15,100	*
Mental disorders-----	115,800	90,200	25,600	18,400	4,900	2,500
Diseases of the musculoskeletal system and connective tissue-----	73,100	22,300	50,800	31,200	18,500	7,100
Endocrine, nutritional, and metabolic diseases-----	48,100	21,700	26,400	17,500	9,000	3,700
Diseases of the respiratory system-----	22,200	12,700	9,500	6,100	2,800	*
Neoplasms-----	25,600	12,700	130,000	8,200	3,600	*
Diseases of the nervous system and sense organs-----	64,200	26,800	37,400	27,000	8,400	4,000
Diseases of the digestive system---	20,500	10,300	10,200	6,400	3,100	*
Diseases of the genitourinary system-----	15,600	6,700	8,900	6,500	*	*
Other diagnoses ³ -----	34,700	16,900	16,000	10,600	4,400	*
Unknown diagnoses-----	9,600	6,000	3,600	*	*	*

See footnotes at end of table.

Table 11. Number and percent distribution of nursing home residents by special aids used, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974—Con.

Reported chronic conditions and impairments and primary diagnosis at last examination	Special aid used				
	Crutches	Braces	Artificial limb	Self-feeder	Other aids
	Number of residents ¹				
All residents-----	9,300	16,700	4,400	11,400	20,900
<u>Reported chronic conditions and impairments²</u>					
Senility-----	3,600	6,000	*	7,700	14,600
Mental illness-----	*	2,400	*	2,400	4,400
Mental retardation-----	*	*	*	*	*
Arthritis or rheumatism-----	4,200	5,000	*	5,100	7,500
Paralysis or palsy due to stroke-----	*	7,400	*	*	3,400
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	*	2,400	*	*	*
Glaucoma or cataracts-----	*	*	*	*	*
Diabetes-----	*	*	*	*	2,300
Any chronic trouble with back or spine-----	*	3,600	*	2,900	3,000
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	3,800	6,200	4,400	2,400	5,000
Heart trouble-----	3,000	4,400	*	3,600	6,800
None of the above-----	*	*	-	*	*
<u>Primary diagnosis at last examination</u>					
Senility, old age, or other symptoms and ill-defined conditions-----	*	*	*	2,700	3,400
Heart attack-----	*	*	*	*	*
Stroke-----	*	6,300	*	*	2,800
Hardening of the arteries-----	*	*	*	*	4,400
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	*	*	*	*	*
Accidents, poisonings, or violence-----	*	*	*	*	*
Mental disorders-----	*	*	*	*	*
Diseases of the musculoskeletal system and connective tissue-----	*	*	*	*	*
Endocrine, nutritional, and metabolic diseases-----	*	*	*	*	*
Diseases of the respiratory system-----	*	*	*	*	*
Neoplasms-----	*	*	*	*	*
Diseases of the nervous system and sense organs-----	*	*	*	*	*
Diseases of the digestive system-----	*	*	*	*	*
Diseases of the genitourinary system-----	*	*	*	*	*
Other diagnoses ³ -----	*	*	*	*	*
Unknown diagnoses-----	*	*	*	*	-

See footnotes at end of table.

Table 11. Number and percent distribution of nursing home residents by special aids used, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974—Con.

Reported chronic conditions and impairments and primary diagnosis at last examination	Total	Special aid used				
		No aid used	1 aid or more used	Wheelchair	Walker	Cane
Percent distribution ¹						
All residents-----	100.0	48.1	51.9	34.8	14.5	6.4
<u>Reported chronic conditions and impairments²</u>						
Senility-----	100.0	47.0	53.0	37.6	13.3	5.5
Mental illness-----	100.0	63.2	36.8	25.5	8.9	3.8
Mental retardation-----	100.0	67.0	33.0	25.2	5.9	*
Arthritis or rheumatism-----	100.0	38.3	61.7	39.0	21.0	8.9
Paralysis or palsy due to stroke-----	100.0	29.1	70.9	56.0	14.5	5.8
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	100.0	35.6	64.4	51.1	11.9	4.7
Glaucoma or cataracts-----	100.0	45.0	55.1	33.9	17.4	10.5
Diabetes-----	100.0	44.4	55.6	37.5	16.7	7.0
Any chronic trouble with back or spine-----	100.0	33.3	66.7	43.8	22.9	8.5
Amputation of extremities or limbs; or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	100.0	27.5	72.5	54.0	15.9	6.7
Heart trouble-----	100.0	44.8	55.3	36.8	15.8	7.9
None of the above-----	100.0	61.4	38.6	19.8	15.3	5.9
<u>Primary diagnosis at last examination</u>						
Senility, old age, or other symptoms and ill-defined conditions-----	100.0	50.7	49.3	33.1	12.9	5.0
Heart attack-----	100.0	47.6	52.4	29.3	17.7	10.1
Stroke-----	100.0	33.7	66.3	51.2	14.9	5.9
Hardening of the arteries-----	100.0	49.8	50.2	32.6	13.5	7.4
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	100.0	51.1	48.9	27.9	13.5	9.9
Accidents, poisonings, or violence-----	100.0	21.7	78.3	55.6	30.7	*
Mental disorders-----	100.0	77.9	22.1	15.9	4.2	2.2
Diseases of the musculoskeletal system and connective tissue-----	100.0	30.1	69.5	42.8	25.4	9.8
Endocrine, nutritional, and metabolic diseases-----	100.0	45.0	55.0	36.3	18.7	7.8
Diseases of the respiratory system-----	100.0	57.4	42.6	27.4	12.6	*
Neoplasms-----	100.0	49.4	50.6	32.2	14.1	*
Diseases of the nervous system and sense organs-----	100.0	41.7	58.3	42.0	13.1	6.3
Diseases of the digestive system-----	100.0	50.4	49.6	31.2	15.1	*
Diseases of the genitourinary system-----	100.0	43.0	57.0	41.4	*	*
Other diagnoses ³ -----	100.0	51.4	48.6	32.3	13.3	*
Unknown diagnoses-----	100.0	62.2	37.8	*	*	*

See footnotes at end of table.

Table 11. Number and percent distribution of nursing home residents by special aids used, according to reported chronic conditions and impairments and primary diagnosis at last examination: United States, August 1973-April 1974—Con.

Reported chronic conditions and impairments and primary diagnosis at last examination and at admission	Special aid used				
	Crutches	Braces	Artificial limb	Self-feeder	Other aids
All residents-----	0.9	1.6	0.4	1.1	1.9
Percent distribution ¹					
Reported chronic conditions and impairments ²					
Senility-----	0.6	1.0	*	1.2	2.3
Mental illness-----	*	1.2	*	1.2	2.2
Mental retardation-----	*	*	*	*	*
Arthritis or rheumatism-----	1.1	1.4	*	1.4	2.0
Paralysis or palsy due to stroke-----	*	6.1	*	*	2.8
Paralysis or palsy not related to stroke, arthritis, or rheumatism-----	*	3.6	*	*	*
Glaucoma or cataracts-----	*	*	*	*	*
Diabetes-----	*	*	*	*	1.6
Any chronic trouble with back or spine-----	*	3.4	*	2.7	2.8
Amputation of extremities or limbs, or permanent stiffness or any deformity of foot, leg, fingers, arm, or back-----	2.5	4.2	2.9	1.6	3.3
Heart trouble-----	0.8	1.2	*	1.0	1.9
None of the above-----	*	*	---	*	*
Primary diagnosis at last examination					
Senility, old age, or other symptoms and ill-defined conditions-----	*	*	*	1.1	2.3
Heart attack-----	*	*	*	*	*
Stroke-----	*	5.5	*	*	2.5
Hardening of the arteries-----	*	*	*	*	1.8
Diseases of the circulatory system other than hardening of the arteries, stroke, and heart attack-----	*	*	*	*	*
Accidents, poisonings, or violence-----	*	*	*	*	*
Mental disorders-----	*	*	*	*	*
Diseases of the musculoskeletal system and connective tissue-----	*	*	*	*	*
Endocrine, nutritional, and metabolic diseases-----	*	*	*	*	*
Diseases of the respiratory system-----	*	*	*	*	*
Neoplasms-----	*	*	*	*	*
Diseases of the nervous system and sense organs-----	*	*	*	*	*
Diseases of the digestive system-----	*	*	*	*	*
Diseases of the genitourinary system-----	*	*	*	*	*
Other diagnoses ³ -----	*	*	*	*	*
Unknown diagnoses-----	*	*	*	*	---

¹Figures may not sum to total because residents may have used more than one special aid.

²Figures may not total to 100 percent because residents may have had more than one reported condition or impairment.

³Includes diseases of the blood and blood-forming organs, diseases of the skin and subcutaneous tissue, congenital anomalies, infective and parasitic diseases, certain causes of perinatal morbidity, complications of pregnancy and childbirth and other diagnoses not listed above.

APPENDIXES

CONTENTS

I.	Technical Notes on Method	48
	Survey Design	48
	General	48
	Sampling Frame	48
	Sampling Design	50
	Data Collection Procedures for 1973-74 National Nursing Home Survey	50
	General Qualifications	51
	Nonresponse and Imputation of Missing Data	51
	Rounding of Numbers	52
	Data Processing	52
	Estimation Procedures	52
	Reliability of Estimates	52
II.	Definitions of Certain Terms Used in This Report	56
	Terms Relating to Residents and Health Status	56
	Terms Related to Facilities	56
	Types of Facilities Included in The Survey	56
	Certification Status	57
III.	Resident Questionnaire Used in The 1973-74 National Nursing Home Survey	58

LIST OF APPENDIX TABLES

I.	Distribution of facilities in the 1973-74 National Nursing Home Survey universe and disposition of sample facilities according to primary sampling strata and bed size of facility: conterminous United States	49
II.	Data collection procedures	51
III.	Standard errors of percentages of residents	54

LIST OF APPENDIX FIGURES

I.	Relative standard error of estimated number of residents	53
----	--	----

APPENDIX I

TECHNICAL NOTES ON METHOD

SURVEY DESIGN

General

From August 1973 to April 1974 the Division of Health Resources Utilization Statistics (DHRUS) conducted the National Nursing Home Survey (NNHS)—a sample survey of nursing homes and their residents and staff in the conterminous United States. The survey was designed and developed by DHRUS in conjunction with a group of experts in various fields encompassing the broad area of long-term care.²⁰ It was specifically designed as the first of a series of surveys to satisfy the diverse data needs of those who establish standards for, plan, provide, and assess long-term care services.

Sampling Frame

The 1973-74 NNHS focused on nursing homes which *provided some level of nursing care*. Only facilities providing nursing care were included because detailed questions on facility services and resident health status were relevant only to these facilities. They included both nursing care homes and personal care with nursing homes, while personal care homes and domiciliary care homes were excluded. Facilities were either freestanding establishments or nursing care units of hospitals, retirement centers, and similar institutions. A definition of nursing care and detailed criteria for classifying facilities providing such care are presented in appendix II.

The survey universe consisted of two groups of facilities: those providing some level of nursing care as classified in the 1971 Master

Facility Inventory (MFI) and those opening for business in 1972. The major group (93 percent) was composed of all nursing homes providing some level of nursing care as classified by the 1971 MFI. The MFI is a census of all inpatient health facilities conducted every 2 years by mail by the National Center for Health Statistics. A detailed description of how the MFI was developed, its content, and procedures for updating and assessing its coverage has been published.²¹⁻²³

In order for data collection to begin in August, the sampling frame was "frozen" in the spring of 1973 so that the sample could be selected in ample time to permit the scheduling of nationwide data collection. To obtain as current a sample frame as possible, all nursing homes which opened for business during 1972 were also included in the universe. (Facilities opening in early 1973 could not be included since data about them were not yet available.) The facilities which opened in 1972 comprised the second, and smaller (7 percent), group of facilities in the universe. Although the universe included only facilities providing nursing care, all facilities opened in 1972 were included because the level of nursing care they provided was unknown prior to survey. Once the NNHS was conducted, facilities not meeting the criteria were classified as out of scope (see table I for details).

Although the NNHS was conducted in 1973-74, it should be noted that estimates will not correspond precisely to figures from the 1973 MFI census for several reasons. In comparison to the MFI, the NNHS universe excluded the following facilities: (1) personal care homes

Table I. Distribution of facilities in the 1973-74 National Nursing Home Survey universe and disposition of sample facilities according to primary sampling strata and bed size of facility: conterminous United States

Certification status and size of home	Universe (sampling frame) ¹	Number of facilities in sample			
		Total facilities	Out of scope or out of business	In scope and in business	
				Non-responding	Responding
All types of strata	17,685	2,118	147	63	1,908
Both Medicare and Medicaid and Medicare only	4,099	803	20	26	757
Unknown number of beds.....	2	0	0	0	0
Less than 25 beds.....	149	4	0	1	3
25-49 beds	538	35	0	1	34
50-99 beds	1,713	228	7	7	214
100-199 beds	1,385	370	8	11	351
200-299 beds	224	100	4	3	93
300-499 beds	68	46	1	2	43
500 beds or more	20	20	0	1	19
Medicaid only	7,473	790	34	24	732
Unknown number of beds.....	3	0	0	0	0
Less than 15 beds.....	250	5	1	2	2
15-24 beds	967	36	5	1	30
25-49 beds	2,253	123	11	3	109
50-99 beds	2,688	293	4	8	281
100-199 beds	1,108	241	3	6	232
200-299 beds	145	52	5	3	44
300-499 beds	43	24	3	1	20
500 beds or more	16	16	2	0	14
Not certified	6,113	525	93	13	419
Unknown number of beds.....	19	0	0	0	0
Less than 15 beds.....	1,279	23	10	0	13
15-24 beds	1,062	38	9	0	29
25-49 beds	1,575	87	13	3	71
50-99 beds	1,334	145	19	5	121
100-199 beds	652	141	21	4	116
200-299 beds	120	43	12	0	31
300-499	52	28	4	1	23
500 beds or more	20	20	5	0	15

¹The universe consisted of nursing homes providing some level of nursing care as classified in the 1971 MFI and those opened for business in 1972

and domiciliary care homes, (2) facilities which opened in 1973, and (3) facilities which, between 1971 and 1973, upgraded the level of care they provided, thereby meeting the "nursing care" criteria when surveyed in the 1973 MFI. Data from the NNHS are also subject to sampling variability, while data from the MFI are not since the MFI is a census.

Sampling Design

The sampling was a stratified two-stage probability design: The first stage was a selection of establishments and the second stage was a selection of residents and employees of the sample establishments. In preparation for the first-stage sample selection, establishments listed in the MFI were sorted into the following types of strata based on Medicare and Medicaid certification: (1) Both Medicare and Medicaid and Medicare only, (2) Medicaid only, and (3) Not certified. Facilities in each of these three strata were sorted into bed size groups, producing 26 primary strata as shown in table I. The nursing homes in the universe were ordered by type of ownership, geographic region, State and county. The sample was then selected systematically after a random start within each primary stratum. Table I shows the distribution of establishments in the sampling frame and the final disposition of the sample with regard to response and in-scope status. The number of facilities estimated in the survey (15,749) is less than the universe figure (17,685) because some facilities went out of business or out of scope between the time the universe was "frozen" and the survey was conducted. Differences ranging from 2,100-2,900 between survey estimates and universe figures occurred in the 1963,²⁴ 1964²⁵ and 1969²⁶ nursing home surveys for the same reason.

The second-stage selection of residents and employees was carried out by the interviewers at the time of their visits to the establishments in accordance with specific instructions given for each sample establishment. The sample frame for residents was the total number of residents on the register of the establishment on the evening prior to the day of the survey. Residents who were physically absent from the facility due

to overnight leave or a hospital visit but had a bed maintained for them at the establishment were included in the sample frame. An average of 10 residents per facility were in the sample.

The sampling frame for employees was the Staff Control Record, on which the interviewer listed the names of all staff employees (including those employed by contract) and sampled professional, semiprofessional, and nursing staff. Those generally *not* involved in direct patient care, such as office staff and food service, house-keeping, and maintenance personnel were excluded from the sample. The interviewer used predesignated sampling instructions that appeared at the head of each column of this form. An average of 14 persons per facility were in the sample.

Data Collection Procedures for 1973-74 National Nursing Home Survey

The 1973-74 NNHS utilized the following questionnaires and forms:

- Administrator Letter and worksheet
- Facility Questionnaire
- Expense Questionnaire
- Resident Control Record
- Resident Questionnaire
- Staff Questionnaire—Parts I and II
- Staff Control Record

(See appendix III for a copy of the Resident Questionnaire. For all other data collection instruments, see reference 27.)

Data were collected according to the following procedures:

1. A letter was sent to the administrators of sample facilities informing them of the survey and the fact that an interviewer would contact them for an appointment. On the back of the letter was a worksheet which the administrator was requested to fill out prior to the interviewer's visit. This worksheet asked for those data that required access to records and some time for compiling (such as total admission and discharges, and inpatient days of care). Included with this introductory letter were letters of endorsement from the American

Nursing Home Association and the American Association of Homes for the Aging urging the administrators to participate in the survey.

2. Several days to 1 week after the mailing of the letters, the interviewer telephoned the sample facility and made an appointment with the administrator.
3. At the time of the appointment the Facility Questionnaire was completed by the interviewer who interviewed the administrator or owner of the facility. After completing this form, the interviewer secured the administrator's permission to send the Expense Questionnaire to the facility's accountant. (If financial records were not kept by an outside firm, the Expense Questionnaire was filled in by the administrator with the interviewer present.) The interviewer completed the Staff Control Record (a list of all current employees both full- and part-time), selected the sample from it and prepared Staff Questionnaires, Parts I and II, which were left for each sample person to complete, seal in addressed and franked envelopes (one for each part of the questionnaire), and either return to the interviewer or mail. The interviewer then completed the Resident Control Record (a list of all residents currently in the facility), selected the sample of residents from it, and filled in a Resident Questionnaire for each sample person by interviewing the member of the nursing staff familiar with care provided to the resident. The nurse referred to the resident's medical records. No resident was interviewed directly.

If the Expense Questionnaire was not returned within 2 weeks, the interviewer telephoned the accountant, requesting its prompt return. If the Staff Questionnaires were not returned in 1 week, the interviewer contacted the staff member and requested the return of the form.

Table II presents a summary of the data collection procedures.

Table II. Data collection procedures

Questionnaire	Respondent	Interview situation
Facility	Administrator	Personal interview
Expense	Facility's accountant	Self-enumerated questionnaire
Resident.....	Member of nursing staff familiar with care provided to the resident and with access to resident's medical records (10 samples residents per facility)	Personal interview
Staff.....	Sampled staff member (14 per facility)	Self-enumerated questionnaire

GENERAL QUALIFICATIONS

Nonresponse and Imputation of Missing Data

Response rates differed for each type of questionnaire as indicated below:

<i>Questionnaire</i>	<i>Response rate</i>
Facility	97percent
Expense	88 percent
Resident	98 percent
Staff	82 percent

Generally response rates were higher for questionnaires administered in a personal interview situation (facility and resident) as compared to those which were self-enumerated (expense and staff). Statistics presented in this report were adjusted for failure of a facility to respond. Data were also adjusted for non-response which resulted from failure to complete one of the questionnaires (expense, resident, staff) or from failure to complete an item on a questionnaire. Those items left unanswered on a partially completed questionnaire (facility, expense, resident, staff) were generally

imputed by assigning a value from a responding unit with major characteristics identical to those of the nonresponding unit.

Rounding of Numbers

Estimates of residents have been rounded to the nearest hundred. For this reason detailed figures within tables do not always add to totals. Percents were calculated on the basis of unrounded figures and will not necessarily agree precisely with percents which may be calculated from rounded data.

Data Processing

A series of checks were performed during the course of the survey. This included field followups for missing and inconsistent data, some manual editing of the questionnaires, extensive editing conducted by computer to assure that all responses were accurate, consistent, logical, and complete. Once the data base was edited, the computer was used to calculate and assign weights, ratio adjustments, and recodes and for other related procedures necessary to produce national estimates from the sample data.

Estimation Procedures

Statistics reported in this publication are derived by a ratio estimating procedure. The purpose of ratio estimation is to take into account all relevant information in the estimation process, thereby reducing the variability of the estimate. The estimation of number of establishments and establishment data not related to size are inflated by the reciprocal of the probability of selecting the sample establishment and adjusted for the nonresponding establishments within primary certification-size strata. Two ratio adjustments, one at each stage of selection, were also used in the estimation process. The first-stage ratio adjustment (along with the above inflation factors) was included in the estimation of establishment data related to size, resident data, and staff data for all primary certification-size strata from which a sample of facilities was drawn. The numerator was the total beds according to the Master Facility Inventory data for all facilities in the stratum. The denom-

inator was the estimate of the total beds obtained through a simple inflation of the Master Facility Inventory data for the sample homes in the stratum. The effect of the first-stage ratio adjustment was to bring the sample in closer agreement with the known universe of beds. The second-stage ratio adjustment was included in the estimation of resident and staff data within establishments. The second-stage ratio adjustment is the product of two fractions—the first is the universe of the sampling fraction for residents (or staff) upon which the selection is based; the second is the ratio of the number of sample residents (or staff) in the establishment to the number of residents (or staff) for whom questionnaires were completed within the facility.

RELIABILITY OF ESTIMATES

As in any survey, the results are subject to reporting and processing errors and errors due to nonresponse. To the extent possible, these types of errors were kept to a minimum by methods built into survey procedures.

Since statistics presented in this report are based on a sample, they will differ somewhat from figures that would have been obtained if a complete census had been taken using the same schedules, instructions, and procedures.

The standard error is primarily a measure of the variability that occurs by chance because only a sample is surveyed rather than the entire universe. The standard error also reflects part of the measurement error, but it does not measure any systematic biases in the data. It is inversely proportional to the square root of the number of observations in the sample. Thus as the sample size increases, the standard error generally decreases.

The relative standard error of an estimate is obtained by dividing the standard error of the estimate by the estimate itself and is expressed as a percentage of the estimate. According to NCHS standards, reliable estimates are those which have a relative standard error of 25 percent or less. For example, figure I shows the relative standard errors for the estimated number of residents. For a relative standard

Figure I. RELATIVE STANDARD ERROR OF ESTIMATED NUMBER OF RESIDENTS




Illustration of use of figure I. As shown in table 9, a walker was used by an estimated 156,400 residents in 1973-74. The relative standard error of this estimate as read in figure I is approximately 2.90 percent, and the standard error is 4536 (2.90 percent of 156,400).

error of 25 percent or less, the minimum number of residents is 2,300. Thus resident estimates must be 2,300 or larger in order to meet the standards of reliability. In tables in this report asterisks are shown for any cell with a number of less than 2,300 or a percentage which represents a number less than 2,300, i.e., with more than a 25-percent relative standard error.

Because of the relationship between the relative standard error and the estimate, the standard error of an estimate can be found by multiplying the estimate by its relative standard error. Both values can be determined from the curve in figure I. For example, table 2 shows that the total number of residents with a primary diagnosis of stroke was 113,400. The relative standard error corresponding to this estimate in figure I is approximately 3.40 percent. The standard error is $113,400 \times (.0340) = 3856$. The chances are about 68 out of 100 that an estimate from the sample differs from the

value which would have been obtained from a complete census by less than the standard error. The chances are about 95 out of 100 that the difference is less than twice the standard error and about 99 out of 100 that it is less than 2 1/2 times as large.

The standard errors of *percentages* of residents are presented in table III. For example, about 36 percent of the 318,100 male residents were depressed once a week or less (table 4). Linear interpolation between values in table III yields an approximate standard error of 1.03 percent. Thus the chances are 95 out of 100 that the estimate is contained in the interval 36 percent ± 2.0 (1.03 percent) (i.e., between 33.9 percent and 38.1 percent) and 99 out of 100 for the interval 36 percent ± 2.5 (1.03 percent) (i.e., between 33.4 percent and 38.6 percent).

Statistical tests to determine whether the differences between selected characteristics are statistically significant can be performed by

Table III. Standard errors of percentages of residents

Base of estimated percent (number of residents)	Estimated percent							
	1 or 99	2 or 98	5 or 95	10 or 90	20 or 80	30 or 70	40 or 60	50
2,000.....	2.69	3.79	5.90	8.12	10.83	12.41	13.27	13.54
4,000.....	1.90	2.68	4.17	5.75	7.66	8.78	9.38	9.58
6,000.....	1.56	2.19	3.41	4.69	6.25	7.17	7.66	7.82
8,000.....	1.35	1.90	2.95	4.06	5.42	6.21	6.63	6.77
10,000.....	1.21	1.70	2.64	3.63	4.84	5.55	5.93	6.06
20,000.....	0.85	1.20	1.87	2.57	3.43	3.92	4.20	4.28
30,000.....	0.70	0.98	1.52	2.10	2.80	3.20	3.43	3.50
40,000.....	0.60	0.85	1.32	1.82	2.42	2.78	2.97	3.03
50,000.....	0.54	0.76	1.18	1.62	2.17	2.48	2.65	2.71
60,000.....	0.49	0.69	1.08	1.48	1.98	2.27	2.42	2.47
70,000.....	0.46	0.64	1.00	1.37	1.83	2.10	2.24	2.29
80,000.....	0.43	0.60	0.93	1.28	1.71	1.96	2.10	2.14
90,000.....	0.40	0.57	0.88	1.21	1.61	1.85	1.98	2.02
100,000.....	0.38	0.54	0.83	1.15	1.53	1.76	1.88	1.92
200,000.....	0.27	0.38	0.59	0.81	1.08	1.24	1.33	1.35
300,000.....	0.22	0.31	0.48	0.66	0.88	1.01	1.08	1.11
400,000.....	0.19	0.27	0.42	0.57	0.77	0.88	0.94	0.96
500,000.....	0.17	0.24	0.37	0.51	0.69	0.78	0.84	0.86
600,000.....	0.16	0.22	0.34	0.47	0.63	0.72	0.77	0.78
700,000.....	0.14	0.20	0.32	0.43	0.58	0.66	0.71	0.72
800,000.....	0.13	0.19	0.30	0.41	0.54	0.62	0.66	0.68
900,000.....	0.13	0.18	0.28	0.38	0.51	0.59	0.63	0.64
1,000,000.....	0.12	0.17	0.26	0.36	0.48	0.56	0.59	0.61
1,100,000.....	0.10	0.17	0.24	0.35	0.46	0.53	0.57	0.57

Illustration of use of table II. In the text it was noted that about 36 percent of the 318,100 male residents in the United States were reported to act depressed at least once a week. Linear interpolation between values shown in table II will yield an approximate standard error of 1.03 percent for an estimate of 36 percent with a base of 318,100.

examining the confidence interval for the difference of the estimates in question. If the confidence interval does not overlap with zero, the estimates are considered statistically different. This test can be used, for example, to determine if the rate of depression is greater for females than for males. Of the 757,700 female residents, 40.2 percent were reported to act depressed at least once a week. The approximate standard error of this estimate is 0.68 percent (from table III). Similarly, the standard error of the 35.9 percent of the 318,000 males reported to act depressed at least once a week was 1.02 percent.

The standard error of a difference is approximately the square root of the sum of the squares of the standard errors of each of the estimates. In other words, letting X and Y be the two estimates,

$$\text{S.E. } (X - Y) = \sqrt{\text{S.E.}^2 (X) + \text{S.E.}^2 (Y)}$$

(This formula will represent the actual standard error quite accurately for the difference between separate and uncorrelated characteristics, although it is only a rough approximation in most other cases.)

Thus the standard error of the difference of the two percentages is approximately

$$\begin{aligned} \text{S.E. } (40.2\% - 35.9\%) &= \sqrt{(1.02)^2 + (.68)^2} \\ \text{S.E. } (4.3\%) &= 1.23 \end{aligned}$$

Since the confidence interval of the difference (4.3 percent \pm 1.96 (1.23)) does not overlap with zero, the chances are 95 out of 100 that the rate of depression is greater for females than for males. In 95 out of 100 cases the difference between the percentages will be positive and lie between 1.9 and 6.7 percentage points.

— o o o —

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

TERMS RELATING TO RESIDENTS AND HEALTH STATUS

Resident.—A person who had been formally admitted but not discharged from an establishment was a resident. All such persons were included in the survey whether or not they were physically present in the facility at the time of the survey.

Age.—Age at date of survey was age of resident.

Reported chronic conditions and impairments.—A reported condition was considered to be the affirmative response by the respondent to any and all categories of item 9 of the Resident Questionnaire. The respondent, who was the nurse most familiar with the care provided to the resident, reported the existence of these chronic conditions and impairments from knowledge of the resident's health and a check of the resident's medical record.

Primary diagnosis.—The primary diagnosis was the one condition reported by the respondent in response to item 8 of the Resident Questionnaire. The list of conditions corresponds to the ICDA.⁸ With the assistance of the interviewer, the respondent was instructed to extract from the resident's medical record the primary diagnosis recorded at the last examination and at admission.

Special aid.—A special aid is a device used to compensate for defects resulting from disease, injury, impairment, or congenital malformation. Aids included in the survey were wheelchairs, walkers, crutches, braces, artificial limbs, self-feeders, and other aids as were specified in item 10 of the Resident Questionnaire.

Behavioral problem.—A behavioral problem referred to the respondent's report that the resident exhibited any of the following behaviors: depressed, agitated or nervous, abusive or aggressive, confused or senile, disturbed sleep, and other problems.

TERMS RELATED TO FACILITIES

Types of Facilities Included in the Survey

Institutions included in the 1973-74 National Nursing Home Survey were those classified as either nursing care homes or personal care homes with nursing according to data collected in the 1971 Master Facility Inventory Survey²⁹ conducted by the National Center for Health Statistics.

Definitions for these two classes of nursing homes were as follows:

Nursing Care Home

- Fifty percent or more of the residents received nursing care during the week prior to the survey. (Nursing care is defined as the provision of one or more of the following services: taking temperature, pulse, respiration; or blood pressure; full bed bath; application of dressings or bandages; catheterization; intravenous, intramuscular, or hypodermic injection; nasal feeding; irrigation; bowel and bladder retraining; oxygen therapy; and enema.)

- At least one full-time (35 or more hours per week) registered nurse (RN) or licensed practical nurse (LPN) was employed.

Personal Care Home with Nursing

- Some, but less than 50 percent, of the residents received nursing care during the week prior to the survey.
- At least one full-time RN or LPN was employed.

or

- Some of the residents received nursing care during the week prior to the survey.
- No full-time RN or LPN was employed.
- The institution either:

Provided administration of medicines or supervision over self-administered medicines,

or

Provided assistance with three or more activities for daily living (such as help with tub bath or shower; help with dres-

sing, correspondence, or shopping; help with walking or getting about; and help with eating).

Certification Status

Medicare.—Refers to the medical assistance provided in Title XVIII of the Social Security Act. Medicare is a health insurance program administered by the Social Security Administration for persons aged 65 and older who are eligible for benefits.

Medicaid.—Refers to the medical assistance provided in Title XIX of the Social Security Act. Medicaid is a State-administered program for the medically indigent.

Skilled nursing home (SNH).—Refers to certification as a skilled nursing home under Medicaid.

Intermediate care facility (ICF).—Refers to certification as an intermediate care facility under Medicaid.

Not certified.—Refers to facilities which are not certified as providers of care either by Medicare or Medicaid.

APPENDIX III

RESIDENT QUESTIONNAIRE USED
IN THE 1973-74 NATIONAL NURSING HOME SURVEY

RESIDENT QUESTIONNAIRE

OMB #068-S-72172
Expires 7-31-74

1973 Nursing Home Survey
National Center for Health Statistics
Health Resources Administration
Rockville, Maryland

1-7

ASSURANCE OF CONFIDENTIALITY — All information which would permit identification of the individual will be held in strict confidence, will be used only by persons engaged in and for the purposes of the survey, and will not be disclosed or released to others for any purposes.	ESTABLISHMENT NO.									

cc2

LINE NO.

--	--	--

cc11

cc14-1

1. WHAT IS — DATE OF BIRTH? or

Month Day Year Age

cc15,16 17, 18 19-21 cc22-24

2. WHAT IS — SEX? Male Female

25-1 -2

3. WHAT IS — ETHNIC BACK- 26-1 Caucasian -2 Negro -3 Oriental

GROUND? (Mark (X) Only -4 Spanish American -5 American Indian -6 Other

one box)

4. WHAT IS — CURRENT MARITAL 27-1 Married -2 Widowed -3 Divorced

STATUS? (Mark (X) only -4 Separated -5 Never Married

one box)

5. WHAT WAS THE DATE OF — CURRENT ADMISSION TO THIS PLACE?

Month Day Year

cc28, 29 30-31 32-34

6a. WHERE DID — LIVE AT THE TIME OF ADMISSION? (Mark (X) only one box)

(1) In a boarding home 35-1

(2) In another nursing home or related facility -2

(3) In a mental hospital or other long-term specialty hospital -3

(4) In a general or short-stay hospital -4

(5) In a private apartment or house -5

(6) Other place, (Specify) _____ -6

(7) Don't know -7

36-

6b. AT THE TIME OF ADMISSION DID — LIVE WITH: (Mark (X) all that apply)

	Yes	No
(1) Spouse? 37-1 <input type="checkbox"/>	<input type="checkbox"/>	-2 <input type="checkbox"/>
(2) Children? 38-1 <input type="checkbox"/>	<input type="checkbox"/>	-2 <input type="checkbox"/>
(3) Other relatives? 39-1 <input type="checkbox"/>	<input type="checkbox"/>	-2 <input type="checkbox"/>
(4) Unrelated persons? 40-1 <input type="checkbox"/>	<input type="checkbox"/>	-2 <input type="checkbox"/>
(5) Lived alone? 41-1 <input type="checkbox"/>	<input type="checkbox"/>	-2 <input type="checkbox"/>
(6) Don't know? 42-1 <input type="checkbox"/>		

7. WHAT IS THE PRIMARY REASON FOR — ADMISSION TO THE HOME? (Enter "1" in box for primary reason; if secondary reason given, enter "2".)

43- Physical reasons (e.g., illness or need for treatments)

44- Social reasons (e.g., no family, or lack of family interest)

45- Behavioral reasons (e.g., disruptive behavior, mental deterioration)

46- Economic reasons (e.g., no money and/or resources)

8. WHAT WAS THIS RESIDENT'S PRIMARY DIAGNOSIS: (Mark (X) only one box in each column)

- | | | | |
|-------|------------------------------|-------------------------------------|--|
| | a. AT ADMISSION? | b. AT THE TIME OF—LAST EXAMINATION? | |
| 49,50 | <input type="checkbox"/> a01 | 51,52 | <input type="checkbox"/> b01 Senility, old age, and other symptoms and ill-defined conditions (e.g., coma, uremia) |
| | <input type="checkbox"/> a02 | <input type="checkbox"/> b02 | Heart attack (e.g., ischemic heart disease) |
| | <input type="checkbox"/> a03 | <input type="checkbox"/> b03 | Stroke (e.g., cerebrovascular diseases) |
| | <input type="checkbox"/> a04 | <input type="checkbox"/> b04 | Hardening of arteries (e.g., arteriosclerosis, diseases of the arteries, arterioles, capillaries) |
| | <input type="checkbox"/> a05 | <input type="checkbox"/> b05 | Other diseases of the circulatory system (e.g., NOT heart attack, stroke, or hardening of the arteries) |
| | <input type="checkbox"/> a06 | <input type="checkbox"/> b06 | Accidents, poisonings, and violence (e.g., fracture of hip, other broken bones, burns, concussion) |
| | <input type="checkbox"/> a07 | <input type="checkbox"/> b07 | Mental disorders (e.g., mental retardation, psychoses, neuroses, mental illness, emotional problems) |
| | <input type="checkbox"/> a08 | <input type="checkbox"/> b08 | Diseases of the musculoskeletal system and connective tissue (e.g., arthritis, rheumatism, back pain) |
| | <input type="checkbox"/> a09 | <input type="checkbox"/> b09 | Endocrine, nutritional, and metabolic diseases (e.g., goiter, diabetes, gout) |
| | <input type="checkbox"/> a10 | <input type="checkbox"/> b10 | Diseases of the respiratory system (e.g., pneumonia, emphysema) |
| | <input type="checkbox"/> a11 | <input type="checkbox"/> b11 | Neoplasms (e.g., cancer, tumors) |
| | <input type="checkbox"/> a12 | <input type="checkbox"/> b12 | Diseases of the nervous system and sense organs (e.g., Parkinson's disease, glaucoma, cataracts, blindness, multiple sclerosis, spastic paralysis, epilepsy) |
| | <input type="checkbox"/> a13 | <input type="checkbox"/> b13 | Diseases of the digestive system (e.g., cirrhosis of liver, ulcer, intestinal obstruction) |
| | <input type="checkbox"/> a14 | <input type="checkbox"/> b14 | Infective and parasitic diseases (e.g., T.B., polio, syphilis) |
| | <input type="checkbox"/> a15 | <input type="checkbox"/> b15 | Diseases of the genitourinary system (e.g., nephrosis, chronic pelvic infection, hyperplasia of prostate) |
| | <input type="checkbox"/> a16 | <input type="checkbox"/> b16 | Diseases of the skin and subcutaneous tissue (e.g., cellulitis, abscess, chronic ulcer) |
| | <input type="checkbox"/> a17 | <input type="checkbox"/> b17 | Diseases of the blood and blood-forming organs (e.g., anemia) |
| | <input type="checkbox"/> a18 | <input type="checkbox"/> b18 | Congenital anomalies (e.g., hydrocephalus) |
| | <input type="checkbox"/> a19 | <input type="checkbox"/> b19 | Complications of pregnancy, childbirth and the puerperium (e.g., infections, hemorrhage, toxemias) |
| | <input type="checkbox"/> a20 | <input type="checkbox"/> b20 | Certain causes of perinatal morbidity and mortality (e.g., birth injury or immaturity of infant) |
| | <input type="checkbox"/> a21 | <input type="checkbox"/> b21 | Don't know |
| | <input type="checkbox"/> a22 | <input type="checkbox"/> b22 | Other (Specify) _____ 54- |
- Specify: _____ 53-

9. DOES — HAVE ANY OF THE FOLLOWING CONDITIONS OR IMPAIRMENTS? (Mark (X) all that apply)

- cc55-65 -1 a. Senility (includes decline in intellect, memory, and judgement, loss of orientation, difficulty in speaking; feableness.)
- 2 b. Mental illness (Psychiatric or emotional problems)
- 3 c. Mental retardation
- 4 d. Arthritis or rheumatism
- 5 e. Paralysis or palsy other than arthritis
- e. (1) IS THIS THE RESULT OF A STROKE? 66-1 Yes -2 No
- 6 f. Glaucoma or cataracts
- 7 g. Diabetes
- 8 h. Any CHRONIC trouble with back or spine
- 9 i. Amputation of extremities or limbs; or permanent stiffness or any deformity of the foot, leg, fingers, arm, or back
- 0 j. Heart trouble
- OR
- 8 k. Resident has none of the above conditions or impairments

10. DOES THIS RESIDENT REGULARLY USE ANY OF THE FOLLOWING AIDS?

CARD 2
14-2

		No	Yes
a.	Walker	15-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
b.	Crutches	16-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
c.	Braces	17-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
d.	Wheelchair	18-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
e.	Artificial Limb	19-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
f.	Self-feeder	20-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
g.	Any other aids (do not count glasses or hearing aids)	21-2 <input type="checkbox"/>	-1 <input type="checkbox"/>

Specify _____ 22-

11. DURING THE LAST MONTH, HOW MANY TIMES DID—RECEIVE ANY OF THE FOLLOWING THERAPY SERVICES? (INCLUDE ONLY SERVICES PROVIDED BY A LICENSED OR REGISTERED PROFESSIONAL WHETHER INSIDE OR OUTSIDE THE HOME.)

		NUMBER OF TIMES
a.	Physical therapy <input type="checkbox"/> None or	<input type="text"/> cc23
b.	Recreational therapy <input type="checkbox"/> None or	<input type="text"/> cc25
c.	Occupational therapy <input type="checkbox"/> None or	<input type="text"/> cc27
d.	Speech therapy <input type="checkbox"/> None or	<input type="text"/> cc29
e.	Hearing therapy <input type="checkbox"/> None or	<input type="text"/> cc31
f.	Professional counseling by social worker, psychologist or other mental health worker <input type="checkbox"/> None or	<input type="text"/> cc33

12. DURING THE PAST 7 DAYS, WHICH OF THESE SERVICES DID—RECEIVE? (Mark (X) all that apply)

- cc35-62
- 01 a. Rub or massage
 - 02 b. Administration of treatment by staff
 - 03 c. Special diet
 - 04 d. Application of sterile dressings or bandages
 - 05 e. Temperature-pulse-respiration
 - 06 f. Full bed-bath
 - 07 g. Enema
 - 08 h. Catheterization
 - 09 i. Blood pressure reading
 - 10 j. Irrigation
 - 11 k. Oxygen therapy
 - 12 l. Intravenous injection
 - 13 m. Hypodermic injection
- OR
- 14 n. None of the above services received

13. DURING THE PAST 7 DAYS, DID – RECEIVE ANY MEDICATIONS?

CARD 3
14-3

15-2

No (Skip to Question 14)

-1

Yes

WHICH TYPES OF MEDICATIONS DID – RECEIVE? (Mark (X) All That Apply)

- cc16-45 -01 a. Tranquilizers (e.g., Thorazine, Mellaril)
- 02 b. Hypnotics – Sedatives (e.g., Nembutal, Seconal, Phenobarbital, Butisol, Placidyl, Chloral Hydrate)
- 03 c. Stool softeners (e.g., Peri-Colace)
- 04 d. Anti-Depressant (e.g., Elavil)
- 05 e. Anti-Hypertensives (e.g., Ismelin)
- 06 f. Diuretics (e.g., Diuril, Esidrex)
- 07 g. Analgesics (e.g., Aspirin, Darvon, Demerol, Percodan, Empirin with Codeine)
- 08 h. Diabetic agents (e.g., Orinase, Insulin)
- 09 i. Anti-inflammatory agents (e.g., Cortisone, Sodium Salicylate, Butazolidin, Indocin)
- 10 j. Anti-infectives (i.e., antibiotics)
- 11 k. Anti-Anginal drugs (e.g., Nitroglycerin, Peritrate)
- 12 l. Cardiac Glycosides (e.g., Digitalis, Lanoxin)
- 13 m. Anti-Coagulants (e.g., Dicumarol, Warfarin)
- 14 n. Vitamins or iron
- 15 o. Other types of medications not listed above

14. THE FOLLOWING ACTIVITIES FOR DAILY LIVING LIST VARIOUS LEVELS OF CARE THAT MAY BE NEEDED BY A RESIDENT. PLEASE INDICATE THE ONE THAT BEST DESCRIBES THE LEVEL OF CARE NEEDED BY THIS RESIDENT. FOR EACH ACTIVITY, THE LEVELS ARE GIVEN IN ASCENDING ORDER: IN OTHER WORDS, THE LEVEL DESCRIBING THE MINIMUM CARE IS FIRST AND THE LEVEL DESCRIBING THE MOST CARE IS LAST. IF YOU ARE UNDECIDED WHICH OF TWO LEVELS TO INDICATE, CHOOSE THE ONE DESCRIBING THE LESSER AMOUNT OF CARE:

a. CONSIDERING THE FOLLOWING FOUR HYGIENE ACTIVITIES (WASHING FACE AND HANDS, BRUSHING TEETH OR DENTURES, COMBING HAIR, AND SHAVING OR APPLYING MAKE-UP) DOES THIS RESIDENT:

(Mark (X) Only One Box)

- 46-1 Perform all four with no assistance?
- 2 Perform all four with no assistance, but needs help in getting and/or putting away equipment?
- 3 Perform three or four with no assistance, but requires help with a complete bath?
- 4 Require assistance with one or two of these hygiene activities?
- 5 Require assistance with all four of these hygiene activities?

b. CONCERNING DRESSING, DOES THIS RESIDENT:

(Mark (X) Only One Box)

- 47-1 Get clothes from closets and drawers and completely dress without assistance?
- 2 Get clothes from closets and drawers and completely dress with some assistance (tying shoes, fastening braces, closing buttons or zippers in back of garments)?
- 3 Receive assistance in getting clothes, or in dressing (do not count tying shoes, fastening braces, closing buttons or zippers in back of garments as assistance)?
- 4 Stay partly or completely undressed?

c. CONCERNING FEEDING, DOES THIS RESIDENT:

(Mark (X) Only One Box)

- 48-1 Feed self without assistance?
- 2 Feed self with minor assistance (cutting meat or buttering bread)?
- 3 Receive major assistance in feeding (do not count cutting meat or buttering bread)?
- 4 Require intravenous feeding?
- 5 Require tube feeding?

d. CONCERNING AMBULATION TO REACH THE TOILET ROOM, IS THIS RESIDENT:

(Mark (X) Only One Box)

- 51-1 Able to go to the toilet room without nurses' assistance (may use cane, walker, wheelchair, or other object of support), may manage bedpan or commode at night?
- 2 Receiving nurses' assistance in going to the toilet room (do not count use of cane, walker, or other object of support), using bedpan or commode at night, or cleaning self or arranging clothes after elimination?
- 3 Unable to go to the toilet room for the elimination process?

e. CONCERNING MOVING IN AND OUT OF A BED OR CHAIR, IS THIS RESIDENT:

(Mark (X) Only One Box)

- Receiving no assistance? 52-1
- Walking with assistance of one person? -2
- Walking with assistance of two persons? -3
- Up in a chair with assistance once in 8 hours? -4
- Up in a chair with assistance twice in 8 hours? -5
- Bedfast with assistance in turning every two hours? -6
- Bedfast with assistance in turning every hour? -7

e.(1) DOES - HAVE BED SORES?

53-1 Yes No
 (continue with part f.)

f. CONCERNING CONTINENCE, IS THIS RESIDENT:

(Mark (X) Only One Box)

- In control of both bowels and bladder? 54-1
- An ostomy patient? -2
- In control of bladder only? -3
- In control of bowels only? -4
- Not in control of bowels or bladder? -5

f.(1) IS - RECEIVING BOWEL AND/OR BLADDER RETRAINING?

55-1 Yes (Skip to Question 15a.)
 -2 No

f.(2) WOULD RETRAINING GIVE THIS RESIDENT CONTROL OVER BOWELS AND/OR BLADDER?

Yes No Doubtful
 56-1 -2 -3

15a. DOES THIS RESIDENT EXHIBIT ANY OF THE FOLLOWING BEHAVIOR?

b. DOES THIS RESIDENT EXHIBIT THIS BEHAVIOR MORE OFTEN THAN ONCE A WEEK OR ONCE A WEEK OR LESS?

	No	Yes		More often than once a week	Once a week or less
(1) Depressed	57-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→	58-1 <input type="checkbox"/>	-2 <input type="checkbox"/>
(2) Agitated, nervous	59-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→	60-1 <input type="checkbox"/>	-2 <input type="checkbox"/>
(3) Abusive, aggressive	61-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→	62-1 <input type="checkbox"/>	-2 <input type="checkbox"/>
(4) Confused, senile	63-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→	64-1 <input type="checkbox"/>	-2 <input type="checkbox"/>
(5) Disturbed sleep	65-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→	66-1 <input type="checkbox"/>	-2 <input type="checkbox"/>
(6) Other problem behavior	67-2 <input type="checkbox"/>	-1 <input type="checkbox"/>	→ (Specify) _____ 68-	→ 69-1 <input type="checkbox"/>	-2 <input type="checkbox"/>

16a. DURING THIS RESIDENT'S STAY HERE, WHEN DID -- LAST SEE A PHYSICIAN FOR TREATMENT, MEDICATION, OR FOR AN EXAMINATION?

CARD 4
14-4

Month Day Year

 cc15,16 17,18 19,20

OR 21-1 Has Never Seen A Doctor While Here (Skip to Question 17a.)

b. AT THAT TIME, DID -- RECEIVE :

- | | | Yes | No |
|---------------------|------|--------------------------|-----------------------------|
| (1) An examination? | 22-1 | <input type="checkbox"/> | -2 <input type="checkbox"/> |
| (2) Treatment? | 23-1 | <input type="checkbox"/> | -2 <input type="checkbox"/> |
| (3) Prescription? | 24-1 | <input type="checkbox"/> | -2 <input type="checkbox"/> |
| (4) Other? | 25-1 | <input type="checkbox"/> | -2 <input type="checkbox"/> |

Specify _____ 26-

c. DID THE PHYSICIAN ATTEND THE RESIDENT: (Mark (X) Only One Box)

- 27-1 as a private physician?
 -2 for the home itself which furnishes the medical care?
 -3 temporarily as a replacement for the resident's private physician who was unable to attend the resident?
 -4 under some other arrangement? (Specify) _____ 28-

d. DOES A PHYSICIAN EXAMINE THIS RESIDENT: (Mark (X) Only One Box)

- 29-1 only when called?
 -2 irregularly, but without being called?
 -3 on a scheduled basis?

d. (1) HOW OFTEN DOES THE PHYSICIAN EXAMINE THE RESIDENT?

(Mark (X) Only One Box.)

- 30-1 once a week
 -2 every 2 weeks
 -3 once a month
 -4 every three months
 -5 once a year
 -6 other (Specify) _____ 31-

17a. DOES -- WEAR EYE GLASSES?

Yes
 32-1

No
 -2

b. IS -- SIGHT WITH GLASSES: (Mark (X) Only One Box)

- 33-1 not impaired? (e.g., can read ordinary newspaper print)
 -2 partially impaired? (e.g., can watch television 8 to 12 feet across the room)
 -3 severely impaired? (e.g., can recognize the features of familiar persons if they are within 2 to 3 feet)
 -4 completely lost? (e.g., blind)

c. IS -- SIGHT: (Mark (X) Only One Box)

- 33-1 not impaired? (e.g., can read ordinary newspaper print without glasses)
 -2 partially impaired? (e.g., can watch television 8 to 12 feet across the room)
 -3 severely impaired? (e.g., can recognize the features of familiar persons if they are within 2 to 3 feet)
 -4 completely lost? (e.g., blind)

18a. DOES -- USE A HEARING AID?

Yes
 34-1

No
 -2

b. IS -- HEARING WITH A HEARING AID: (Mark (X) Only One Box)

- 35-1 not impaired? (e.g., can hear a telephone conversation on an ordinary telephone)
 -2 partially impaired? (e.g., can hear most of the things a person says)
 -3 severely impaired? (e.g., can hear only a few words a person says or loud noises)
 -4 completely lost? (e.g., deaf)

c. IS -- HEARING: (Mark (X) Only One Box)

- 35-1 not impaired? (e.g., can hear a telephone conversation on an ordinary telephone)
 -2 partially impaired? (e.g., can hear most of the things a person says)
 -3 severely impaired? (e.g., can hear only a few words a person says or loud noises)
 -4 completely lost? (e.g., deaf)

19. IS — ABILITY TO SPEAK: (Mark (X) Only One Box)

- 38-1 not impaired? (e.g., is able to be understood; can carry on a normal conversation)
- 2 partially impaired? (e.g., is able to be understood but has difficulty pronouncing some words)
- 3 severely impaired? (e.g., cannot carry on a normal conversation; is understood only with difficulty)
- 4 completely lost? (e.g., is mute)

20a. DOES THIS RESIDENT HAVE DENTURES?

- Yes 39-1
- No (Skip to Question 21a.) -2

b. DOES — USE THE DENTURES?

- Yes 40-1
- No -2

21a. DURING THE LAST MONTH, DID — LEAVE THE HOME FOR ANY RECREATIONAL OR LEISURE ACTIVITIES?

- Yes 41-1
- No -2

b. FOR WHICH OF THE FOLLOWING ACTIVITIES DID — LEAVE THE HOME? (Mark (X) All That Apply)

- cc42-52 -1 Get books, etc., from the library
- 2 Attend plays, movies, concerts, etc.
- 3 Attend arts and crafts classes outside the home
- 4 Visit museums, parks, fairs, etc.
- 5 Go on shopping trips organized by the home
- 6 Go on independent shopping trips organized by the resident or visitors
- 7 Visit a beauty shop or barber shop
- 8 Visit community clubs (such as community centers, senior citizen clubs, service clubs, bridge clubs, unions, etc.)
- 9 Attend religious services or other religious activities
- 0 Go for a walk
- 8 Other, (Specify) _____ 53-

c. WHY DIDN'T — LEAVE THE HOME TO PARTICIPATE IN ANY ACTIVITIES DURING THE LAST MONTH? (Mark (X) All That Apply)

- cc42-52 -1 Resident was too ill or was not able to move well enough to participate
- 2 Resident was not interested
- 3 Staff was unable to determine resident's interests at this point
- 4 Staff feels that the resident's behavior will not be tolerated outside the home
- 5 No one was available to accompany the resident
- 6 Resident cannot afford these activities
- 7 Lack of transportation
- 8 Other, (Specify) _____ 53-

22a. DURING THE PAST YEAR, HAS THIS RESIDENT BEEN ON ANY KIND OF LEAVE OVERNIGHT OR LONGER, EXCLUDING LEAVE FOR MEDICAL REASONS?

- Yes 54-1
- No (Skip to Question 23a.) -2
- Don't know (Skip to Question 23a.) -3

b. WHERE DID — USUALLY GO WHEN ON LEAVE? (Mark (X) Only One Box)

- 55-1 To own home or apartment
- 2 To home of family or relatives
- 3 To home of unrelated friends
- 4 To foster home
- 5 To boardinghouse or room
- 6 To another place, (Specify) _____ 56-
- 7 Don't know

c. ABOUT HOW OFTEN DID THIS RESIDENT GO ON LEAVE? (Mark (X) Only One Box)

- 57-1 Nearly every week
- 2 About once a month
- 3 About once every two months
- 4 Several times a year
- 5 About once a year or less
- 6 Other (Specify) _____ 58-
- 7 Don't know

CARD 5
14-5

23a. DOES — HAVE ANY VISITORS?

- Yes 15-1
- No -2 (Skip to Question 24)
- Don't know -3

b. HOW FREQUENTLY DO VISITORS SEE THE RESIDENT? (Mark (X) Only One Box)

- 16-1 Nearly every week
- 2 About once a month
- 3 About once every two months
- 4 Several times a year
- 5 About once a year or less
- 6 Other (Specify) _____ 17-
- 7 Don't know

24. HOW MANY BEDS ARE IN — ROOM? (Mark (X) Only One Box)

- 18-1 One bed (i.e., the resident's own bed)
- 2 Two beds
- 3 Three beds
- 4 Four beds
- 5 Five or more beds

25a. HAS THIS RESIDENT LIVED IN THIS FACILITY FOR ONE FULL MONTH OR LONGER?

- Yes 19-1
 - No -2
- Stop; go on to next questionnaire.

b. LAST MONTH, WHAT WAS THE BASIC CHARGE FOR THIS RESIDENT'S LODGING, MEALS, AND NURSING CARE NOT INCLUDING PRIVATE DUTY NURSING OR OTHER SPECIAL CHARGES?

No charge is made for care (Skip to Question 26a.) \$ _____ cc20-25

c. LAST MONTH, WHAT WAS THE TOTAL CHARGE FOR THIS RESIDENT'S CARE, INCLUDING ALL CHARGES FOR SPECIAL SERVICES, DRUGS, AND SPECIAL MEDICAL SUPPLIES?

No charge is made for care (Skip to Question 26a.) \$ _____ cc26-31

(1) DID THIS AMOUNT INCLUDE SPECIAL CHARGES FOR

	No	Yes
(a) physician services?	32-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(b) private duty nursing?	33-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(c) therapy?	34-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(d) drugs?	35-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(e) special medical supplies?	36-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(f) special diet?	37-2 <input type="checkbox"/>	-1 <input type="checkbox"/>
(g) other?	38-2 <input type="checkbox"/>	-1 <input type="checkbox"/>

Specify

-39

26a. WHAT WERE ALL THE SOURCES OF PAYMENT FOR THIS RESIDENT'S CARE LAST MONTH? (Mark (X) All That Apply)

- cc40-48 (1) Own income or family support (private plans, retirement funds, social security, etc.)
- (2) Medicare (Title XVIII)
- (3) Medicaid (Title XIX)
- (4) Other public assistance or welfare
- (5) Church support
- (6) VA contract
- (7) Initial payment-life care
- (8) No charge is made for care
- (9) Other (Specify) _____ 49-

b. WHAT WAS THE PRIMARY SOURCE OF PAYMENTS FOR — CARE LAST MONTH? (Mark (X) Only One Box.)

- 50-1 Own income or family support (private plans, retirement funds, social security, etc.)
- 2 Medicare (Title XVIII)
- 3 Medicaid (Title XIX)
- 4 Other public assistance or welfare
- 5 Church support
- 6 VA contract
- 7 Initial payment-life care
- 8 No charge is made for care
- 9 Other (Specify) _____ 51-

Form 73NHS-7

RESIDENT CONTROL RECORD

RESIDENTS IN NURSING HOME					
SAMPLE DESIGNATION	NAME OF RESIDENT	LINE NO.	SAMPLE DESIGNATION	NAME OF RESIDENT	LINE NO.
SW _____			SW _____		
TE _____			TE _____		
a	b	c	a	b	c
		51			76
		52			77
		53			78
		54			79
		55			80
		56			81
		57			82
		58			83
		59			84
		60			85
		61			86
		62			87
		63			88
		64			89
		65			90
		66			91
		67			92
		68			93
		69			94
		70			95
		71			96
		72			97
		73			98
		74			99
		75			00

VITAL AND HEALTH STATISTICS PUBLICATIONS SERIES

Formerly Public Health Service Publication No. 1000

- Series 1. Programs and Collection Procedures.*—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data Evaluation and Methods Research.*—Studies of new statistical methodology including experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical Studies.*—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and Committee Reports.*—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.*—Statistics on illness; accidental injuries; disability; use of hospital, medical, dental, and other services; and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.*—Data from direct examination, testing, and measurement of national samples of the civilian, noninstitutionalized population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics; and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutionalized Population Surveys.*—Discontinued effective 1975. Future reports from these surveys will be in Series 13.
- Series 13. Data on Health Resources Utilization.*—Statistics on the utilization of health manpower and facilities providing long-term care, ambulatory care, hospital care, and family planning services.
- Series 14. Data on Health Resources: Manpower and Facilities.*—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on Mortality.*—Various statistics on mortality other than as included in regular annual or monthly reports. Special analyses by cause of death, age, and other demographic variables; geographic and time series analyses; and statistics on characteristics of deaths not available from the vital records, based on sample surveys of those records.
- Series 21. Data on Natality, Marriage, and Divorce.*—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports. Special analyses by demographic variables; geographic and time series analyses; studies of fertility; and statistics on characteristics of births not available from the vital records, based on sample surveys of those records.
- Series 22. Data from the National Mortality and Natality Surveys.*—Discontinued effective 1975. Future reports from these sample surveys based on vital records will be included in Series 20 and 21, respectively.
- Series 23. Data from the National Survey of Family Growth.*—Statistics on fertility, family formation and dissolution, family planning, and related maternal and infant health topics derived from a biennial survey of a nationwide probability sample of ever-married women 15-44 years of age.

For a list of titles of reports published in these series, write to: Scientific and Technical Information Branch
National Center for Health Statistics
Public Health Service
Hyattsville, Md. 20782

NCHS

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
National Center for Health Statistics
Center Building
3700 East-West Highway
Hyattsville, Maryland 20782

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

For publications in the
Vital and Health Statistics
Series call 301-436-NCHS.

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF HEW
HEW 390

THIRD CLASS
BLK. RATE

