

SAFER • HEALTHIER • PEOPLE™



Vital and Health Statistics

November 2002

Series 13, Number 153

National Hospital Discharge Survey: 2000 Annual Summary With Detailed Diagnosis and Procedure Data



DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Trade name disclaimer

The use of trade names is for identification only and does not imply endorsement by the Public Health Service, U.S. Department of Health and Human Services.

Suggested citation

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Copyright information

Kozak LJ, Hall MJ, Owings MF. National Hospital Discharge Survey: 2000 Annual Summary with detailed diagnosis and procedure data. National Center for Health Statistics. Vital Health Stat 13(153). 2002.

Library of Congress Catalog Card Number 90-657616

For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.

Vital and Health Statistics

Series 13, Number 153

National Hospital Discharge Survey: 2000 Annual Summary With Detailed Diagnosis and Procedure Data

Data From the National
Hospital Discharge Survey

DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
November 2002
DHHS Publication No. (PHS) 2003-1724

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for International Statistics*

Jennifer H. Madans, Ph.D., *Associate Director for Science*

Lawrence H. Cox, Ph.D., *Associate Director for Research and Methodology*

Diane M. Makuc, Dr.P.H., *Acting Associate Director for Analysis, Epidemiology, and Health Promotion*

P. Douglas Williams, *Acting Associate Director for Data Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Vital and Health Statistics Systems*

Douglas L. Zinn, *Acting Associate Director for Management and Operations*

Charles J. Rothwell, *Associate Director for Information Technology and Services*

Division of Health Care Statistics

Linda K. Demlo, Ph.D., *Division Director*

Thomas McLemore, *Deputy Director*

Irma Arispe, Ph.D., *Associate Director for Science*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

Cooperation of the U.S. Bureau of the Census

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the U.S. Census Bureau, under a contractual arrangement, participated in planning the surveys and collecting the data for the National Hospital Discharge Survey.

Contents

Abstract	1
Highlights.....	1
Patient Characteristics.....	1
Hospital Characteristics.....	1
Diagnoses.....	1
Procedures.....	2
Deliveries and Newborn Infants.....	2
Introduction	2
Methods.....	3
Data Source.....	3
Estimation.....	4
Use of Tables.....	4
References	5
Appendix	181
Technical Notes	181
Abstract Form	181
Population Estimates	181
Reliability of Estimates	181
Definitions of Selected Terms	184
Changes in ICD–9–CM Codes.....	186

Text Figures

1. Number of discharges by age and sex: United States, 2000	2
2. Average length of stay in days by region: United States, selected years 1970–2000	2
3. Number of discharges for selected diagnoses by age: United States, 2000	3
4. Rate of discharges with a diagnosis of human immunodeficiency virus by region: United States, 2000	3
5. Rate of discharges with coronary artery procedures by sex: United States, 2000	4
6. Percent distribution of discharges with deliveries by length of stay: United States, selected years 1980–2000.....	4

Detailed Tables

Tables on Patient and Hospital Characteristics

1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2000	7
2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2000	8
3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2000	9
4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2000	11
5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age, and geographic region: United States, 2000	14

6.	Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age, and geographic region: United States, 2000	16
7.	Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 2000	18

Diagnosis Tables

8.	Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2000	18
9.	Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2000	19
10.	Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000	21
11.	Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000	22
12.	Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000	23
13.	Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000	24
14.	Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000	25
15.	Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000	26
16.	Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000	27
17.	Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000	28
18.	Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000	29
19.	Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2000	30
20.	Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2000	31
21.	Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2000	32
22.	Number and rate of discharges from short-stay hospitals with human immunodeficiency virus diagnoses, by selected characteristics: United States, selected years 1990–2000	33
23.	Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnoses: United States, 2000	34

Procedure Tables

24.	Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 2000	34
25.	Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 2000	35
26.	Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2000	36
27.	Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures by sex, age, and geographic region: United States, 2000	37
28.	Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2000	38
29.	Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2000	39
30.	Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2000	40
31.	Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2000	41
32.	Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2000	42
33.	Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2000	43

Tables on Deliveries and Newborn Infants

34.	Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2000	44
35.	Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–2000	45
36.	Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2000	46
37.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–2000	46
38.	Number and rate of females with cesarean deliveries, by type of delivery, and of females with vaginal birth after cesarean delivery, discharged from short-stay hospitals, by age and geographic region: United States, 2000	47
39.	Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2000	48
40.	Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay by health status: United States, selected years 1980–2000	48
41.	Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2000	49
42.	Number and percent of male newborn infants circumcised during hospitalization by geographic region: United States, selected years 1980–2000	49

Diagnosis and Procedure Tables by ICD–9–CM Code

43.	Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2000	50
44.	Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000	77
45.	Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2000	104
46.	Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2000	162

Appendix Tables

I.	Civilian population by selected characteristics: United States, 2000	181
II.	Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics, 2000	184
III.	Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey by ICD–9–CM category, 2000	186
IV.	Changes in diagnosis codes	187
V.	Changes in procedure codes	192

Appendix Figure

I.	Medical abstract form for the National Hospital Discharge Survey, 2000	182
----	--	-----

Objectives

This report presents national estimates for 2000 and selected trend data on the use of non-Federal short-stay hospitals in the United States. Estimates are provided by demographic characteristics of patients discharged, geographic region, ownership and bed size of hospitals, principal expected source of payment, conditions diagnosed, and surgical and nonsurgical procedures performed.

Methods

The estimates are based on data collected through the National Hospital Discharge Survey. The survey has been conducted annually by the National Center for Health Statistics since 1965. In 2000, data were collected for approximately 313,000 discharges. Of the 481 eligible non-Federal short-stay hospitals in the sample, 434 (90 percent) responded to the survey. Measurements of hospital use include number and rate of discharges and days of care, and the average length of stay. The population estimates used to compute rates were based on the 1990 Census. Estimates of first-listed diagnoses, days of care, all-listed diagnoses, and all-listed procedures are presented according to their code number in the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM).

Results

An estimated 31.7 million inpatients were discharged from non-Federal short-stay hospitals in 2000. These patients used an estimated 155.9 million days of care. Females had 19.2 million discharges compared with 12.5 million for males. Fifty-four percent of all discharges under age 65 years had a principal expected source of payment from a private source, and 22 percent were covered by Medicaid. Discharge rates per 1,000 population ranged from 85.4 in the West to 135.5 in the Northeast. The leading discharge diagnosis was delivery for patients under age 65 years and heart disease for those 65 years of age and over. The rate of discharges with coronary artery bypass graft for males was 2.4 times the rate for females, and males had almost twice the female rate for discharges with removal of coronary artery obstruction. Hospital stays were 1 day or less for 15 percent of women with deliveries in 2000, compared with 37 percent in 1995.

Keywords: *hospitalization • inpatient • diagnoses • procedures*

National Hospital Discharge Survey: 2000 Annual Summary With Detailed Diagnosis and Procedure Data

by *Lola J. Kozak, Ph.D.; Margaret J. Hall, Ph.D.; and Maria F. Owings, Ph.D., Division of Health Care Statistics*

Highlights

Patient Characteristics

- An estimated 31.7 million inpatients were discharged from non-Federal short-stay hospitals in 2000. These patients used an estimated 155.9 million days of care.
- During 2000, 19.2 million of the discharges were for females, compared with 12.5 million for males. Females had 3.8 million more discharges among those 15–34 years of age and 2.0 million more discharges among those 75 years of age and over than did males ([figure 1](#)).
- In 1970, average lengths of stay ranged from 6.7 days (West) to 9.1 days (Northeast). In 2000, the range was from 4.5 days (Midwest and West) to 5.7 days (Northeast) ([figure 2](#)).
- The principal expected source of payment for 54 percent of discharges under age 65 years was a private source (HMO/PPO, Blue Cross/Blue Shield, or another private source), and for 22 percent it was Medicaid.
- Deaths accounted for 2.4 percent of all hospital discharges, 1.0 percent of discharges under 65 years and 4.7 percent of discharges 65 years of age and over.

Hospital Characteristics

- Discharge rates per 1,000 population ranged from 85.4 in the West to 135.5 in the Northeast.
- In 2000, half of inpatients were discharged from hospitals with less than 200 beds, while about 10 percent were discharged from hospitals with 500 or more beds.
- Nonprofit hospitals accounted for more than three-fourths of hospital discharges in 2000. The proportion ranged from 93 percent in the Northeast to 61 percent in the South.

Diagnoses

- The leading discharge diagnosis was delivery for patients under 65 years of age and heart disease for those 65 years of age and over ([figure 3](#)).
- Other diagnoses that were reported for more than a million discharges in 2000 were psychoses, pneumonia, and malignant neoplasms.
- The discharge rate per 100,000 population for patients with HIV diagnoses was highest in the Northeast (129.5) and higher in the South (64.9) than in the Midwest (31.1) or West (33.4) ([figure 4](#)).

This report was prepared in the *Division of Health Care Statistics*. Rong Cai of Lockheed Martin Information Technology did the computer programming for the report and produced the estimated parameters for approximate relative standard error equations. The report was edited by Klaudia M. Cox and typeset by Zung T.N. Le of the Publications Branch, Division of Data Services.




Figure 1. Number of discharges by age and sex: United States, 2000




Figure 2. Average length of stay in days by region: United States, selected years 1970-2000

Procedures

- In 2000, 23.2 million surgical and 16.7 million nonsurgical procedures were performed on hospitalized patients.
- Frequent procedures were spinal tap and respiratory therapy for children under 15 years, obstetrical procedures for those 15-44 years of age, and coronary artery procedures for patients 45 years of age and over.
- The rate of discharges with a coronary artery bypass graft for

males was 2.4 times the rate for females, and males had almost twice the female rate for discharges with removal of coronary artery obstruction (figure 5).

Deliveries and Newborn Infants

- An estimated 3.7 million women were hospitalized for delivery in 2000, and they had average stays of 2.5 days.

- Hospital stays were 1 day or less for 15 percent of women with deliveries in 2000, compared with 37 percent in 1995 (figure 6).
- In 2000, 38 percent of newborn infants had at least one illness or risk-related diagnosis.
- More than half the diagnoses for sick newborn infants were hemolytic disease of fetus or newborn, respiratory distress syndrome and other respiratory conditions, disorders relating to short gestation and unspecified low birthweight, or congenital anomalies.

Introduction

This report presents data from the 2000 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report provides a summary of 2000 data, including estimates of diagnoses and procedures by *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) code (1).

The NHDS is the principal source for national data on the characteristics of patients discharged from non-Federal short-stay hospitals. Data from the NHDS are used to examine important topics in public health (2-6) and for a variety of planning, administrative, and evaluation activities by governmental, scientific, academic, and commercial institutions.

Estimates of the number of diagnoses and procedures shown in this report are for hospital inpatients only. Data on ambulatory medical care services are collected in other NCHS surveys. The National Ambulatory Medical Care Survey obtains information on patients' visits to physicians' offices (7). The National Hospital Ambulatory Medical Care Survey collects data on visits to hospital outpatient and emergency departments (8,9). Data on ambulatory surgery performed in hospitals and in

freestanding ambulatory surgery centers were collected by the National Survey of Ambulatory Surgery from 1994–96 (10).

The NHDS and the ambulatory care surveys are components of the National Health Care Survey. Other components are the National Nursing Home Survey (11) and the National Home and Hospice Care Survey (12).

Methods

Data Source

NHDS data are collected from a sample of inpatient records acquired from a national probability sample of hospitals. Because persons with multiple discharges during the year can be sampled more than once, the NHDS produces estimates for discharges, not persons. Only hospitals with an average length of stay of fewer than 30 days for all patients, general hospitals, or children's general hospitals are included in the survey. Federal, military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals), and hospitals with fewer than six beds staffed for patient use, are excluded. Beginning in 1988, the sampling frame for hospitals has been the SMG Hospital Market Database (13). The 1988 sample was updated in 1991, 1994, 1997, and 2000 to allow the inclusion of hospitals that opened or changed their eligibility status since the previous sample update.

The NHDS originally used a two-stage design, but it was redesigned in 1988 and implemented a stratified three-stage design (14). Both the original and the redesigned samples include the largest hospitals with certainty and sample noncertainty hospitals. The first stage of the current three-stage design consists of the selection of 112 primary sampling units (PSUs) that comprise a probability sample of PSUs used in the 1985–94 National Health Interview Survey (15). The second stage consists of the selection of noncertainty hospitals from the sample PSUs. At the third stage, a systematic random




Figure 3. Number of discharges for selected diagnoses by age: United States, 2000




Figure 4. Rate of discharges with a diagnosis of human immunodeficiency virus by region: United States, 2000

sample of discharges within hospitals is selected.

For 2000, the sample consisted of 509 hospitals, of which 28 were out of scope (ineligible) because they had gone out of business or otherwise failed to meet the criteria for the NHDS universe. Of the 481 in-scope (eligible) hospitals, 434 (90 percent) responded to the survey. Data were collected for approximately 313,000 discharges from the 434 responding hospitals.

Two data collection procedures are used for the survey. One is a manual system in which sample selection and transcription of information from

medical records are performed by hospital staff or by staff of the U.S. Bureau of the Census on behalf of NCHS. Completed forms are sent to NCHS for medical coding, data entry, editing, and estimation.

For the other data collection procedure, NCHS purchases electronic files containing medical record data from commercial organizations, State data systems, hospitals, or hospital associations. Discharges from these files are systematically sampled. Approximately 40 percent of respondent hospitals provided data through this automated system for the 2000 survey.




Figure 5. Rate of discharges with coronary artery procedures by sex: United States, 2000




Figure 6. Percent distribution of discharges with deliveries by length of stay: United States, selected years 1980-2000

The NHDS includes data elements relating to patient demographic characteristics and medical information, as well as administrative and hospital-related data elements. The medical abstract form shown in figure 1 in the appendix collects patients' date of birth or age, ZIP code, dates of admission and discharge, discharge status, sex, race, ethnicity, marital status, expected source of payment, final medical diagnoses, procedures performed, and dates of surgery. Medical data are coded according to the *International Classification of Diseases, 9th Revision, Clinical Modification*

(ICD-9-CM) (1). Administrative and hospital data collected include patient medical record numbers, number of hospital beds, type of ownership, and geographic region of the hospital. The patient medical record number, date of birth, and ZIP code are confidential information that are not available to the public.

For the manual data collection, an ongoing quality control program is undertaken for the medical coding and data entry from abstracts to electronic form. Approximately 10 percent of abstracts are independently recoded, with discrepancies resolved by a chief

coder. The overall error rate for records manually coded for the 2000 survey year was 0.6 percent for medical (ICD-9-CM) coding and data entry, and 0.2 percent for demographic coding and data entry.

Estimation

Because they are sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components:

- (1) inflation by reciprocals of the probabilities of sample selection,
- (2) adjustment for nonresponse, and
- (3) population weighting ratio adjustments.

These three components of the final weight are described in more detail in another report (14). Information about the standard errors of statistics for the 2000 NHDS is in the appendix of this report.

Use of Tables

Sociodemographic information about patients and utilization estimates by hospital characteristics are shown in tables 1-7. Diagnostic data are included in tables 8-23, and information about procedures is found in tables 24-33. Tables 34-42 present data on women with deliveries and newborn infants. Estimates of diagnoses, days of care by diagnosis, and procedures are shown by ICD-9-CM code in tables 43-46. Figures 1-6 illustrate highlights from the survey data. Definitions of terms used in these tables are presented in the appendix.

The diagnostic and procedure data in this report are often presented by chapter of the ICD-9-CM. Subcategories of diagnoses or procedures are also shown. These specific categories are selected because of their large estimates or because they were of special interest. Although diagnoses assigned ICD-9-CM codes E800-E999 (supplementary classification of external causes of injury and poisoning) are included in the NHDS, these diagnoses are not well reported in the survey and, therefore, are

excluded from this report. A detailed discussion of injury diagnoses and E-codes has been published (16).

Table 27 shows data for coronary artery bypass graft (CABG) surgery and removal of coronary artery obstruction and insertion of stent(s). These procedures are often assigned more than one code for a single operation. CABG surgery is frequently assigned multiple codes to indicate the number and type of bypasses. Beginning with the 1996 NHDS data, changes to the ICD-9-CM coding system allowed patients to have a separate code for insertion of stent(s) along with a code for removal of coronary artery obstruction. As shown in table 27, the number of these procedures exceeds the number of inpatients who had one or more of the procedures.

Tables 7 and 23 show rates of death per 100 discharges. In contrast, mortality rates are ratios of deaths to the population. Mortality rates include deaths that occur outside of hospitals and are for the underlying cause of death, which might not be the same as the first-listed diagnosis for a patient discharged dead. Mortality rates are published regularly by NCHS in *National Vital Statistics Reports* (17).

Data for newborn infants are included only in tables 39–42. Because estimates of newborn infants are based on a sample and because they do not include out-of-hospital births, these estimates may not agree with data on births published in *National Vital Statistics Reports* (18), which are based on a complete count of births.

The final four tables consist of use measures for detailed lists of ICD-9-CM codes. Tables 43–45 provide information by three-digit, four-digit, and five-digit ICD-9-CM diagnostic codes. Table 46 shows two-, three-, and four-digit procedure codes. Only ICD-9-CM code numbers appear in tables 43–46. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1). A specific diagnosis or procedure code appears in tables 43, 45, and 46 if the estimate of the total for the code is 5,000 or more. The codes in table 44 for days of care are identical to those in table 43 for first-listed diagnoses.

Estimates in all tables are rounded to the nearest thousand. Totals may include data for categories that are not shown individually in the tables. For these reasons, figures within tables do not always add to the totals. Rates, percents, and average lengths of stay are calculated from unrounded figures and may not precisely agree with measures calculated from rounded data.

Because of low reliability, estimates in tables 1–42 with relative standard errors of more than 30 percent or that are based on fewer than 30 records are not presented; only an asterisk (*) appears in the tables. Estimates based on 30–59 records are preceded by an asterisk (*) to indicate that they also have low reliability.

Only an asterisk is shown in tables 43, 45, and 46 if the estimate of the diagnosis or procedure is less than 5,000, and only an asterisk is shown in table 44 for estimates of days of care derived from fewer than 5,000 estimated discharges. These estimates generally have a relative standard error of more than 30 percent or a sample size of less than 30. Estimates of diagnoses and procedures of less than 9,000 are preceded by an asterisk because they are generally based on fewer than 60 records. Days of care estimates in table 44 that are derived from fewer than 9,000 estimated discharges are also preceded by asterisks.

The population estimates used to compute rates are shown in table I in the appendix. These estimates are based on the 1990 census. Population estimates for the civilian population from the 2000 census were not available when this report was prepared. When 2000-based civilian population estimates are received, NCHS will provide them on request. A comparison of populations based on the 1990 and 2000 census will be included in the 2001 Annual Summary. The population estimates used in this report have been adjusted for the net underenumeration in the 1990 census. Populations used to compute rates in the annual summaries of 1990–96 NHDS data were not adjusted for the underenumeration.

References

1. International Classification of Diseases, 9th Revision, Clinical Modification, 6th edition. U.S. Department of Health and Human Services, National Center for Health Statistics, Health Care Financing Administration. 1998. <http://www.cdc.gov/nchs/icd9.htm#ICD-9-CM>.
2. Wang G, Dietz WH. Economic burden of obesity in youths aged 6 to 17 years: 1979–1999. *Pediatrics* 109(5): E81. 2002. <http://www.pediatrics.org/cgi/content/full/109/5/e81>.
3. Kennedy BS, Kasl SV, Brass LM, Vaccarino V. Trends in hospitalized stroke for blacks and whites in the United States, 1980–1999. *Neuroepidemiology* 21(3): 131–41. 2002.
4. Kozak LJ, Hall MJ, Owings MF. Trends in avoidable hospitalizations, 1980–1998. *Health Affairs* 20(2): 225–32. 2001.
5. Heidenreich PA, McClellan M. Trends in treatment and outcomes for acute myocardial infarction: 1975–1995. *American Journal of Medicine* 110(3): 165–74. 2001.
6. Weeks JD, Kozak LJ. Trends in the use of episiotomy in the United States: 1980–1998. *Birth* 28(3): 152–60. 2001.
7. Cherry DK, Woodwell DA. National Ambulatory Medical Care Survey: 2000 summary. Advance data from vital and health statistics; no 328. Hyattsville, Maryland: National Center for Health Statistics. 2002. <http://www.cdc.gov/nchs/data/ad/ad328.pdf>.
8. Ly N, McCaig LF. National Hospital Ambulatory Medical Care Survey: 2000 outpatient department summary. Advance data from vital and health statistics; no 327. Hyattsville, Maryland: National Center for Health Statistics. 2002. <http://www.cdc.gov/nchs/data/ad/ad327.pdf>.
9. McCaig LF, Ly N. National Hospital Ambulatory Medical Care Survey: 2000 emergency department summary. Advance data from vital and health statistics; no 326. Hyattsville, Maryland: National Center for Health Statistics. 2002. <http://www.cdc.gov/nchs/data/ad/ad326.pdf>.
10. Hall MJ, Lawrence L. Ambulatory surgery in the United States, 1996. Advance data from vital and health statistics; no 300. Hyattsville, Maryland: National Center for Health

- Statistics. 1998. <http://www.cdc.gov/nchs/data/ad/ad300.pdf>.
11. Jones A. The National Nursing Home Survey: 1999 summary. National Center for Health Statistics. Vital Health Stat 13(152). 2002. http://www.cdc.gov/nchs/data/series/sr_13/sr13_152.pdf.
 12. National Center for Health Statistics. National Home and Hospice Care Survey. 2002. <http://www.cdc.gov/nchs/about/major/nhhcsd/nhhcsd.htm>.
 13. SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1987, 1991, 1994, 1997, 2000.
 14. Dennison CF, Pokras R. Design and operation of the National Hospital Discharge Survey: 1988 redesign. National Center for Health Statistics. Vital Health Stat 1(39). 2000. http://www.cdc.gov/nchs/data/series/sr_01/sr1_39.pdf.
 15. Massey JT, Moore TF, Parsons VL, Tadros W. Design and estimation for the National Health Interview Survey, 1985–94. National Center for Health Statistics. Vital Health Stat 2(110). 1989. http://www.cdc.gov/nchs/data/series/sr_02/sr02_110.pdf.
 16. Hall MJ, Owings MF. Hospitalizations for injury: United States, 1996. Advance data from vital and health statistics; no 318. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/data/ad/ad318.pdf>.
 17. Minino AM, Smith BL. Deaths: Preliminary data for 2000. National vital statistics reports vol 49 no 12. Hyattsville, Maryland: National Center for Health Statistics. 2001. http://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_12.pdf.
 18. Martin JA, Hamilton BE, Ventura SJ, et al. Births: Final data for 2000. National vital statistics reports vol 50 no 5. Hyattsville, Maryland: National Center for Health Statistics. 2002. http://www.cdc.gov/nchs/data/nvsr/nvsr50/nvsr50_05.pdf.
 19. Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual, Release 7.0. Research Triangle Park, NC: Research Triangle Institute. 1996.
 20. Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. New York: John Wiley & Sons. Wiley Classics Library Edition. 1993.
 21. Cochran WG. Sampling techniques, 3rd ed. New York: John Wiley & Sons, 1977.

Table 1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	1970	1975	1980	1985	1990	1995	2000
Number of discharges in thousands							
All ages	29,127	34,043	37,832	35,056	30,788	30,722	31,706
Under 15 years	3,873	3,826	3,672	2,972	2,412	2,405	2,383
15–44 years	12,664	14,171	15,635	13,966	11,799	10,593	9,969
45–64 years	6,693	8,391	8,660	7,610	6,244	6,168	6,958
65 years and over	5,897	7,654	9,864	10,508	10,333	11,556	12,396
65–74 years	3,163	4,057	4,943	5,011	4,689	4,832	4,678
75–84 years	2,099	2,728	3,611	3,969	3,949	4,590	5,119
85 years and over	635	869	1,310	1,528	1,694	2,134	2,599
Rate of discharges per 1,000 population ¹							
All ages	144.3	159.2	167.7	148.4	122.3	115.7	114.0
Under 15 years	66.8	70.4	71.6	57.7	43.1	40.4	39.4
15–44 years	154.6	152.4	150.1	125.0	99.3	87.8	81.6
45–64 years	159.6	191.8	194.8	170.8	135.5	118.5	114.2
65 years and over	293.3	337.3	383.7	369.8	334.1	347.7	359.6
65–74 years	253.2	291.5	315.8	297.3	261.6	260.0	260.0
75–84 years	339.5	392.1	463.9	446.4	395.7	415.6	419.1
85 years and over	443.7	477.1	576.4	572.9	560.6	592.5	608.8
Number of days of care in thousands							
All ages	226,445	262,389	274,508	226,217	197,422	164,627	155,857
Under 15 years	18,189	17,554	16,191	13,554	11,655	10,715	10,734
15–44 years	71,605	80,687	81,951	67,397	54,062	40,825	36,593
45–64 years	62,419	75,393	71,008	53,541	42,153	34,207	34,443
65 years and over	74,232	88,754	105,358	91,726	89,552	78,880	74,086
65–74 years	37,873	44,963	49,253	41,090	37,422	31,310	26,438
75–84 years	27,633	32,906	40,355	36,024	35,926	31,974	31,525
85 years and over	8,726	10,886	15,750	14,612	16,204	15,597	16,123
Rate of days of care per 1,000 population ¹							
All ages	1,121.6	1,227.3	1,216.7	957.7	784.0	620.2	560.4
Under 15 years	313.9	322.9	315.7	263.0	208.4	179.9	177.4
15–44 years	874.0	867.9	786.8	603.3	454.9	338.4	299.5
45–64 years	1,488.8	1,723.0	1,596.9	1,201.6	914.4	657.5	565.2
65 years and over	3,691.9	3,910.6	4,098.4	3,228.0	2,895.6	2,373.7	2,148.9
65–74 years	3,031.6	3,230.8	3,147.0	2,437.3	2,087.8	1,684.7	1,469.3
75–84 years	4,469.2	4,729.2	5,183.7	4,052.2	3,599.5	2,894.7	2,581.1
85 years and over	6,102.1	5,978.1	6,932.1	5,478.7	5,361.7	4,330.6	3,776.6
Average length of stay in days							
All ages	7.8	7.7	7.3	6.5	6.4	5.4	4.9
Under 15 years	4.7	4.6	4.4	4.6	4.8	4.5	4.5
15–44 years	5.7	5.7	5.2	4.8	4.6	3.9	3.7
45–64 years	9.3	9.0	8.2	7.0	6.8	5.5	5.0
65 years and over	12.6	11.6	10.7	8.7	8.7	6.8	6.0
65–74 years	12.0	11.1	10.0	8.2	8.0	6.5	5.7
75–84 years	13.2	12.1	11.2	9.1	9.1	7.0	6.2
85 years and over	13.7	12.5	12.0	9.6	9.6	7.3	6.2

¹Population estimates used to compute rates for 1990, 1995, and 2000 are based on the 1990 census. See appendix.

Table 2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex and age	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	
Both sexes							
All ages	31,706	100.0	114.0	155,857	100.0	560.4	4.9
Under 15 years	2,383	7.5	39.4	10,734	6.9	177.4	4.5
Under 1 year	782	2.5	195.7	4,444	2.9	1,112.5	5.7
1–4 years	703	2.2	44.9	2,276	1.5	145.2	3.2
5–14 years	898	2.8	22.0	4,014	2.6	98.3	4.5
15–44 years	9,969	31.4	81.6	36,593	23.5	299.5	3.7
15–19 years	1,138	3.6	56.1	4,241	2.7	209.1	3.7
20–24 years	1,681	5.3	89.7	5,154	3.3	275.0	3.1
25–34 years	3,717	11.7	97.7	12,468	8.0	327.7	3.4
35–44 years	3,433	10.8	76.1	14,731	9.5	326.5	4.3
45–64 years	6,958	21.9	114.2	34,443	22.1	565.2	5.0
45–54 years	3,500	11.0	94.2	16,629	10.7	447.5	4.8
55–64 years	3,458	10.9	145.4	17,815	11.4	749.1	5.2
65 years and over	12,396	39.1	359.6	74,086	47.5	2,148.9	6.0
65–74 years	4,678	14.8	260.0	26,438	17.0	1,469.3	5.7
75–84 years	5,119	16.1	419.1	31,525	20.2	2,581.1	6.2
85 years and over	2,599	8.2	608.8	16,123	10.3	3,776.6	6.2
Male							
All ages	12,514	100.0	92.0	66,797	100.0	491.2	5.3
Under 15 years	1,333	10.7	43.1	6,241	9.3	201.7	4.7
Under 1 year	450	3.6	220.6	2,701	4.0	1,324.4	6.0
1–4 years	388	3.1	48.4	1,241	1.9	154.9	3.2
5–14 years	495	4.0	23.7	2,299	3.4	110.0	4.6
15–44 years	2,680	21.4	43.9	13,084	19.6	214.2	4.9
15–19 years	302	2.4	29.0	1,593	2.4	153.3	5.3
20–24 years	293	2.3	31.0	1,354	2.0	143.5	4.6
25–34 years	768	6.1	40.8	3,637	5.4	192.9	4.7
35–44 years	1,317	10.5	58.8	6,500	9.7	290.2	4.9
45–64 years	3,424	27.4	115.8	17,325	25.9	585.9	5.1
45–54 years	1,679	13.4	92.3	8,410	12.6	462.2	5.0
55–64 years	1,745	13.9	153.5	8,914	13.3	784.0	5.1
65 years and over	5,077	40.6	352.8	30,147	45.1	2,095.2	5.9
65–74 years	2,199	17.6	269.7	12,366	18.5	1,516.6	5.6
75–84 years	2,107	16.8	426.1	13,060	19.6	2,641.1	6.2
85 years and over	771	6.2	597.4	4,721	7.1	3,658.5	6.1
Female							
All ages	19,192	100.0	135.0	89,059	100.0	626.7	4.6
Under 15 years	1,050	5.5	35.5	4,493	5.0	152.1	4.3
Under 1 year	332	1.7	169.8	1,742	2.0	891.5	5.3
1–4 years	315	1.6	41.2	1,035	1.2	135.0	3.3
5–14 years	402	2.1	20.2	1,716	1.9	86.1	4.3
15–44 years	7,289	38.0	119.3	23,509	26.4	384.8	3.2
15–19 years	836	4.4	84.5	2,647	3.0	267.6	3.2
20–24 years	1,388	7.2	149.2	3,800	4.3	408.5	2.7
25–34 years	2,948	15.4	153.6	8,830	9.9	460.1	3.0
35–44 years	2,116	11.0	93.2	8,231	9.2	362.3	3.9
45–64 years	3,534	18.4	112.6	17,118	19.2	545.6	4.8
45–54 years	1,822	9.5	96.1	8,218	9.2	433.3	4.5
55–64 years	1,712	8.9	137.9	8,900	10.0	717.0	5.2
65 years and over	7,319	38.1	364.4	43,939	49.3	2,187.4	6.0
65–74 years	2,479	12.9	251.9	14,072	15.8	1,430.1	5.7
75–84 years	3,013	15.7	414.4	18,466	20.7	2,540.3	6.1
85 years and over	1,828	9.5	613.7	11,402	12.8	3,827.7	6.2

¹Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
Both sexes					
All ages:					
United States	31,706	114.0	155,857	560.4	4.9
Northeast	7,103	135.5	40,366	770.1	5.7
Midwest	7,207	112.8	32,600	510.4	4.5
South	12,016	121.6	58,730	594.6	4.9
West	5,380	85.4	24,160	383.4	4.5
Under 15 years:					
United States	2,383	39.4	10,734	177.4	4.5
Northeast	472	44.0	2,538	236.8	5.4
Midwest	503	36.6	1,930	140.4	3.8
South	806	37.3	3,568	165.2	4.4
West	602	41.8	2,698	187.0	4.5
15–44 years:					
United States	9,969	81.6	36,593	299.5	3.7
Northeast	2,130	93.8	9,465	416.9	4.4
Midwest	2,087	74.8	7,142	255.8	3.4
South	3,922	90.9	14,210	329.2	3.6
West	1,830	64.4	5,777	203.4	3.2
45–64 years:					
United States	6,958	114.2	34,443	565.2	5.0
Northeast	1,592	135.5	8,774	746.7	5.5
Midwest	1,497	106.3	6,970	494.8	4.7
South	2,770	126.7	13,548	619.5	4.9
West	1,099	83.0	5,152	389.1	4.7
65 years and over:					
United States	12,396	359.6	74,086	2,148.9	6.0
Northeast	2,909	401.3	19,589	2,702.2	6.7
Midwest	3,121	383.8	16,559	2,036.8	5.3
South	4,517	372.0	27,405	2,257.2	6.1
West	1,849	265.9	10,533	1,514.3	5.7
Male					
All ages:					
United States	12,514	92.0	66,797	491.2	5.3
Northeast	2,979	117.4	18,006	709.2	6.0
Midwest	2,857	91.5	13,756	440.8	4.8
South	4,621	96.4	24,372	508.6	5.3
West	2,057	65.4	10,663	338.9	5.2
Under 15 years:					
United States	1,333	43.1	6,241	201.7	4.7
Northeast	266	48.6	1,552	283.0	5.8
Midwest	284	40.3	1,098	156.1	3.9
South	438	39.7	1,910	173.1	4.4
West	345	46.7	1,681	227.6	4.9
15–44 years:					
United States	2,680	43.9	13,084	214.2	4.9
Northeast	684	60.6	3,753	332.4	5.5
Midwest	565	40.5	2,479	177.5	4.4
South	1,030	48.3	4,988	234.0	4.8
West	401	27.6	1,865	128.4	4.7
45–64 years:					
United States	3,424	115.8	17,325	585.9	5.1
Northeast	805	142.5	4,504	796.9	5.6
Midwest	734	107.1	3,472	506.5	4.7
South	1,353	128.6	6,756	642.2	5.0
West	531	81.2	2,592	396.4	4.9

See footnote at end of table.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
65 years and over:					
United States	5,077	352.8	30,147	2,095.2	5.9
Northeast	1,224	412.9	8,197	2,766.3	6.7
Midwest	1,273	379.3	6,708	1,998.4	5.3
South	1,800	356.2	10,718	2,121.7	6.0
West	780	258.6	4,524	1,499.3	5.8
Female					
All ages:					
United States	19,192	135.0	89,059	626.7	4.6
Northeast	4,123	152.5	22,361	827.2	5.4
Midwest	4,351	133.2	18,844	576.8	4.3
South	7,395	145.4	34,358	675.6	4.6
West	3,323	105.3	13,497	427.7	4.1
Under 15 years:					
United States	1,050	35.5	4,493	152.1	4.3
Northeast	205	39.2	986	188.5	4.8
Midwest	219	32.6	832	124.0	3.8
South	368	34.8	1,658	156.9	4.5
West	257	36.6	1,017	144.4	3.9
15–44 years:					
United States	7,289	119.3	23,509	384.8	3.2
Northeast	1,446	126.7	5,712	500.4	4.0
Midwest	1,522	109.0	4,663	334.1	3.1
South	2,893	132.4	9,222	422.0	3.2
West	1,429	103.0	3,912	281.9	2.7
45–64 years:					
United States	3,534	112.6	17,118	545.6	4.8
Northeast	787	129.0	4,270	700.2	5.4
Midwest	763	105.5	3,497	483.8	4.6
South	1,417	124.8	6,792	598.4	4.8
West	568	84.8	2,560	382.0	4.5
65 years and over:					
United States	7,319	364.4	43,939	2,187.4	6.0
Northeast	1,685	393.2	11,392	2,657.9	6.8
Midwest	1,847	387.0	9,852	2,063.8	5.3
South	2,717	383.3	16,686	2,353.7	6.1
West	1,069	271.5	6,009	1,525.7	5.6

¹Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO ²	Blue Cross/ Blue Shield and other private insurance	Self-pay	Other payments and no charge
Number of discharges in thousands								
United States								
All	31,706	12,329	4,451	581	4,864	6,863	1,506	813
Male	12,514	5,131	1,414	330	1,769	2,706	704	321
Female	19,192	7,199	3,038	251	3,096	4,157	803	491
Under 15 years	2,383	*6	894	41	407	711	109	*
15–44 years	9,969	548	2,446	263	2,297	3,008	925	359
45–64 years	6,958	1,214	880	199	1,488	2,452	422	208
65 years and over	12,396	10,561	231	78	672	692	51	62
Northeast	7,103	2,763	903	97	1,158	1,635	350	*
Midwest	7,207	3,247	746	106	1,071	1,672	241	74
South	12,016	4,695	1,825	235	1,768	2,355	710	*
West	5,380	1,624	978	142	867	1,201	205	324
Northeast								
Male	2,979	1,166	355	65	432	697	175	*
Female	4,123	1,597	548	32	726	938	175	*
Under 15 years	472	*	122	*	63	*	25	*
15–44 years	2,130	126	503	46	496	670	221	34
45–64 years	1,592	270	230	29	345	586	93	*
65 years and over	2,909	2,366	47	*	254	191	11	*
Midwest								
Male	2,857	1,348	241	60	370	662	123	33
Female	4,351	1,899	505	*	701	1,010	118	41
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	2,087	110	437	*	540	750	146	37
45–64 years	1,497	257	144	31	370	589	68	19
65 years and over	3,121	2,879	21	*	86	101	5	*
South								
Male	4,621	1,925	517	129	633	913	317	*
Female	7,395	2,771	1,308	106	1,135	1,442	393	*
Under 15 years	806	*	350	*11	125	176	46	*
15–44 years	3,922	252	1,011	114	892	1,047	429	*
45–64 years	2,770	552	355	84	534	913	210	*
65 years and over	4,517	3,888	*	27	218	220	25	11
West								
Male	2,057	692	301	75	334	434	88	113
Female	3,323	932	676	67	534	767	117	211
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	1,830	60	495	55	369	541	129	161
45–64 years	1,099	135	150	56	239	365	50	96
65 years and over	1,849	1,428	54	*	114	180	*	*
Number of days of care in thousands								
United States								
All	155,857	74,707	21,211	2,746	19,269	27,033	6,009	3,509
Male	66,797	31,366	8,614	1,670	7,930	11,774	3,175	1,648
Female	89,059	43,341	12,597	1,076	11,339	15,259	2,834	1,861
Under 15 years	10,734	*	4,711	*	1,717	2,792	369	*
15–44 years	36,593	3,616	9,551	1,068	7,372	9,911	3,295	1,343
45–64 years	34,443	7,542	5,415	1,038	6,486	10,395	2,053	1,005
65 years and over	74,086	63,519	1,535	458	3,695	3,936	292	350

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO ²	Blue Cross/ Blue Shield and other private insurance	Self-pay	Other payments and no charge
Number of days of care in thousands								
Northeast	40,366	19,006	5,637	*	5,106	7,360	1,729	*
Midwest	32,600	17,464	3,193	*	3,888	6,047	932	358
South	58,730	28,710	8,128	1,084	7,203	9,141	2,681	*
West	24,160	9,528	4,253	639	3,072	4,485	667	1,363
Northeast								
Male	18,006	8,124	2,704	*	2,057	3,418	888	*
Female	22,361	10,882	2,933	*	3,049	3,942	841	*
Under 15 years	2,538	*	*	*	255	*	100	*
15–44 years	9,465	880	2,737	*	1,724	2,500	1,062	172
45–64 years	8,774	1,882	1,676	139	1,582	2,776	493	77
65 years and over	19,589	16,237	330	*	1,545	1,169	73	*
Midwest								
Male	13,756	7,316	1,262	*	1,514	2,592	538	178
Female	18,844	10,148	1,932	*	2,375	3,455	394	180
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	7,142	682	1,641	*	1,619	2,305	484	159
45–64 years	6,970	1,525	766	*	1,569	2,378	341	114
65 years and over	16,559	15,247	123	62	461	533	*	*
South								
Male	24,372	11,715	2,910	661	3,011	3,826	1,407	*
Female	34,358	16,994	5,217	423	4,193	5,314	1,274	*
Under 15 years	3,568	*	1,749	*33	*	586	150	*
15–44 years	14,210	1,738	3,561	439	2,972	3,488	1,376	*
45–64 years	13,548	3,331	2,060	456	2,444	3,695	1,020	*
65 years and over	27,405	23,629	*	157	1,150	1,371	135	63
West								
Male	10,663	4,211	1,738	372	1,349	1,938	342	628
Female	13,497	5,317	2,515	266	1,723	2,547	325	735
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	5,777	317	1,612	196	1,058	1,618	373	542
45–64 years	5,152	804	913	280	891	1,545	199	491
65 years and over	10,533	8,406	324	*	539	863	*	*
United States								
Average length of stay								
All	4.9	6.1	4.8	4.7	4.0	3.9	4.0	4.3
Male	5.3	6.1	6.1	5.1	4.5	4.4	4.5	5.1
Female	4.6	6.0	4.1	4.3	3.7	3.7	3.5	3.8
Under 15 years	4.5	*5.0	5.3	4.5	4.2	3.9	3.4	4.4
15–44 years	3.7	6.6	3.9	4.1	3.2	3.3	3.6	3.7
45–64 years	5.0	6.2	6.2	5.2	4.4	4.2	4.9	4.8
65 years and over	6.0	6.0	6.6	5.9	5.5	5.7	5.8	5.6
Northeast	5.7	6.9	6.2	5.5	4.4	4.5	4.9	4.9
Midwest	4.5	5.4	4.3	4.6	3.6	3.6	3.9	4.8
South	4.9	6.1	4.5	4.6	4.1	3.9	3.8	4.1
West	4.5	5.9	4.4	4.5	3.5	3.7	3.3	4.2
Northeast								
Male	6.0	7.0	7.6	5.5	4.8	4.9	5.1	5.5
Female	5.4	6.8	5.3	5.6	4.2	4.2	4.8	4.5
Under 15 years	5.4	*	7.3	11.4	4.1	4.9	3.9	4.9
15–44 years	4.4	7.0	5.4	5.1	3.5	3.7	4.8	5.1
45–64 years	5.5	7.0	7.3	4.7	4.6	4.7	5.3	3.8
65 years and over	6.7	6.9	7.0	*6.2	6.1	6.1	6.9	*7.7

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO/PPO ²	Blue Cross/ Blue Shield and other private insurance	Self-pay	Other payments and no charge
	Average length of stay							
Midwest								
Male	4.8	5.4	5.2	4.6	4.1	3.9	4.4	5.4
Female	4.3	5.3	3.8	4.5	3.4	3.4	3.3	4.4
Under 15 years	3.8	*	4.6	4.0	3.2	3.6	3.4	3.6
15–44 years	3.4	6.2	3.8	4.1	3.0	3.1	3.3	4.3
45–64 years	4.7	5.9	5.3	5.3	4.2	4.0	5.0	5.9
65 years and over	5.3	5.3	5.8	5.0	5.4	5.3	6.7	6.0
South								
Male	5.3	6.1	5.6	5.1	4.8	4.2	4.4	4.5
Female	4.6	6.1	4.0	4.0	3.7	3.7	3.2	3.8
Under 15 years	4.4	*	5.0	*3.0	5.1	3.3	3.3	4.2
15–44 years	3.6	6.9	3.5	3.8	3.3	3.3	3.2	3.7
45–64 years	4.9	6.0	5.8	5.5	4.6	4.0	4.9	4.5
65 years and over	6.1	6.1	7.0	5.9	5.3	6.2	5.5	5.6
West								
Male	5.2	6.1	5.8	4.9	4.0	4.5	3.9	5.5
Female	4.1	5.7	3.7	4.0	3.2	3.3	2.8	3.5
Under 15 years	4.5	*	5.0	3.3	4.0	4.0	2.9	4.6
15–44 years	3.2	5.3	3.3	3.6	2.9	3.0	2.9	3.4
45–64 years	4.7	5.9	6.1	5.0	3.7	4.2	4.0	5.1
65 years and over	5.7	5.9	6.0	6.0	4.7	4.8	4.8	5.1

*Figure does not meet standard of reliability or precision.

¹Includes discharges for which no expected source of payment was provided.²HMO/PPO is health maintenance organization and preferred provider organization.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All bed sizes	6-99	100-199	200-299	300-499	500 or more
Number of discharges in thousands						
All	31,706	7,353	8,590	5,437	7,253	3,072
Sex						
Male	12,514	2,873	3,205	2,180	2,962	1,295
Female	19,192	4,480	5,385	3,257	4,291	1,778
Age						
Under 15 years	2,383	413	746	423	597	203
15-44 years	9,969	2,170	2,747	1,550	2,424	1,078
45-64 years	6,958	1,461	1,744	1,230	1,694	829
65 years and over	12,396	3,309	3,353	2,234	2,538	962
Region						
Northeast	7,103	1,382	1,266	1,393	2,182	880
Midwest	7,207	2,428	1,547	1,206	1,364	662
South	12,016	2,798	3,655	1,860	2,405	1,297
West	5,380	745	2,123	978	1,301	233
Percent distribution of patients discharged						
All	100.0	23.2	27.1	17.2	22.9	9.7
Sex						
Male	100.0	23.0	25.6	17.4	23.7	10.3
Female	100.0	23.3	28.1	17.0	22.4	9.3
Age						
Under 15 years	100.0	17.4	31.3	17.8	25.1	8.6
15-44 years	100.0	21.8	27.6	15.6	24.3	10.8
45-64 years	100.0	21.0	25.1	17.7	24.4	11.9
65 years and over	100.0	26.7	27.1	18.0	20.5	7.8
Region						
Northeast	100.0	19.5	17.8	19.6	30.7	12.4
Midwest	100.0	33.7	21.5	16.7	18.9	9.2
South	100.0	23.3	30.4	15.5	20.0	10.8
West	100.0	13.9	39.5	18.2	24.2	4.3
Number of days of care in thousands						
All	155,857	32,568	41,062	28,112	37,015	17,100
Sex						
Male	66,797	13,232	16,934	11,973	16,686	7,972
Female	89,059	19,336	24,129	16,139	20,329	9,127
Age						
Under 15 years	10,734	1,504	3,342	1,919	2,895	1,074
15-44 years	36,593	8,056	9,249	5,621	9,093	4,574
45-64 years	34,443	6,234	8,122	6,229	9,143	4,716
65 years and over	74,086	16,774	20,350	14,343	15,884	6,735
Region						
Northeast	40,366	6,877	8,336	7,865	12,218	5,070
Midwest	32,600	9,433	7,105	5,984	6,561	3,517
South	58,730	13,195	16,660	9,479	12,034	7,362
West	24,160	3,063	8,961	4,783	6,201	1,151

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All bed sizes	6–99	100–199	200–299	300–499	500 or more
			Percent distribution of days of care			
All	100.0	20.9	26.3	18.0	23.7	11.0
Sex						
Male	100.0	19.8	25.4	17.9	25.0	11.9
Female	100.0	21.7	27.1	18.1	22.8	10.2
Age						
Under 15 years	100.0	14.0	31.1	17.9	27.0	10.0
15–44 years	100.0	22.0	25.3	15.4	24.8	12.5
45–64 years	100.0	18.1	23.6	18.1	26.5	13.7
65 years and over	100.0	22.6	27.5	19.4	21.4	9.1
Region						
Northeast	100.0	17.0	20.7	19.5	30.3	12.6
Midwest	100.0	28.9	21.8	18.4	20.1	10.8
South	100.0	22.5	28.4	16.1	20.5	12.5
West	100.0	12.7	37.1	19.8	25.7	4.8
			Average length of stay in days			
All	4.9	4.4	4.8	5.2	5.1	5.6
Sex						
Male	5.3	4.6	5.3	5.5	5.6	6.2
Female	4.6	4.3	4.5	5.0	4.7	5.1
Age						
Under 15 years	4.5	3.6	4.5	4.5	4.8	5.3
15–44 years	3.7	3.7	3.4	3.6	3.8	4.2
45–64 years	5.0	4.3	4.7	5.1	5.4	5.7
65 years and over	6.0	5.1	6.1	6.4	6.3	7.0
Region						
Northeast	5.7	5.0	6.6	5.6	5.6	5.8
Midwest	4.5	3.9	4.6	5.0	4.8	5.3
South	4.9	4.7	4.6	5.1	5.0	5.7
West	4.5	4.1	4.2	4.9	4.8	4.9

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All	Proprietary	Government	Nonprofit
Number of discharges in thousands				
All	31,706	3,312	3,979	24,414
Sex				
Male	12,514	1,260	1,579	9,676
Female	19,192	2,053	2,401	14,738
Age				
Under 15 years	2,383	122	304	1,956
15-44 years	9,969	1,123	1,369	7,477
45-64 years	6,958	677	952	5,329
65 years and over	12,396	1,390	1,354	9,653
Region				
Northeast	7,103	115	373	6,614
Midwest	7,207	236	580	6,391
South	12,016	2,312	2,368	7,335
West	5,380	649	658	4,073
Percent distribution of patients discharged				
All	100.0	10.4	12.6	77.0
Sex				
Male	100.0	10.1	12.6	77.3
Female	100.0	10.7	12.5	76.8
Age				
Under 15 years	100.0	5.2	12.8	82.1
15-44 years	100.0	11.3	13.7	75.0
45-64 years	100.0	9.7	13.7	76.6
65 years and over	100.0	11.2	10.9	77.9
Region				
Northeast	100.0	1.6	5.3	93.1
Midwest	100.0	3.3	8.1	88.7
South	100.0	19.2	19.7	61.0
West	100.0	12.1	12.2	75.7
Number of days of care in thousands				
All	155,857	17,155	18,717	119,985
Sex				
Male	66,797	6,956	8,259	51,582
Female	89,059	10,198	10,459	68,403
Age				
Under 15 years	10,734	618	1,144	8,972
15-44 years	36,593	4,354	5,257	26,983
45-64 years	34,443	3,480	4,779	26,184
65 years and over	74,086	8,703	7,537	57,846
Region				
Northeast	40,366	484	2,374	37,508
Midwest	32,600	1,117	2,529	28,954
South	58,730	12,705	10,987	35,038
West	24,160	2,848	2,827	18,484

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

	All	Proprietary	Government	Nonprofit
Percent distribution of days of care				
All	100.0	11.0	12.0	77.0
Sex				
Male	100.0	10.4	12.4	77.2
Female	100.0	11.5	11.7	76.8
Age				
Under 15 years	100.0	5.8	10.7	83.6
15–44 years	100.0	11.9	14.4	73.7
45–64 years	100.0	10.1	13.9	76.0
65 years and over	100.0	11.7	10.2	78.1
Region				
Northeast	100.0	1.2	5.9	92.9
Midwest	100.0	3.4	7.8	88.8
South	100.0	21.6	18.7	59.7
West	100.0	11.8	11.7	76.5
Average length of stay in days				
All	4.9	5.2	4.7	4.9
Sex				
Male	5.3	5.5	5.2	5.3
Female	4.6	5.0	4.4	4.6
Age				
Under 15 years	4.5	5.0	3.8	4.6
15–44 years	3.7	3.9	3.8	3.6
45–64 years	5.0	5.1	5.0	4.9
65 years and over	6.0	6.3	5.6	6.0
Region				
Northeast	5.7	4.2	6.4	5.7
Midwest	4.5	4.7	4.4	4.5
South	4.9	5.5	4.6	4.8
West	4.5	4.4	4.3	4.5

Table 7. Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	Number of deaths in thousands			Rate per 100 discharges		
	Both sexes	Male	Female	Both sexes	Male	Female
All ages	776	366	411	2.4	2.9	2.1
Under 65 years	189	107	83	1.0	1.4	0.7
Under 15 years	14	10	4	0.6	0.7	0.4
15–44 years	44	21	22	0.4	0.8	0.3
45–64 years	132	76	56	1.9	2.2	1.6
65 years and over	587	259	328	4.7	5.1	4.5
65–74 years	158	88	70	3.4	4.0	2.8
75–84 years	238	112	126	4.7	5.3	4.2
85 years and over	190	59	132	7.3	7.6	7.2

Table 8. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)*]

Diagnostic category and ICD–9–CM code	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 10,000 population ¹	Number in thousands	Rate per 10,000 population ¹	
All conditions ²	31,706	1,140.1	155,857	5,604.4	4.9
Heart disease 391–392.0,393–398,402,404,410–416,420–429	4,385	157.7	20,514	737.7	4.7
Acute myocardial infarction 410	781	28.1	4,411	158.6	5.7
Coronary atherosclerosis 414.0	1,102	39.6	4,274	153.7	3.9
Other ischemic heart disease 411–413,414.1–414.9	283	10.2	766	27.5	2.7
Cardiac dysrhythmias 427	716	25.7	2,640	94.9	3.7
Congestive heart failure 428.0	999	35.9	5,448	195.9	5.5
Females with deliveries V27	3,738	134.4	9,271	333.4	2.5
Psychoses 290–299	1,445	52.0	11,750	422.5	8.1
Pneumonia 480–486	1,282	46.1	7,527	270.7	5.9
Malignant neoplasms 140–208,230–234	1,156	41.6	7,801	280.5	6.7
Malignant neoplasm of large intestine and rectum 153–154,197.5	159	5.7	1,365	49.1	8.6
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	139	5.0	925	33.3	6.7
Malignant neoplasm of breast 174–175,198.81	108	3.9	319	11.5	2.9
Fractures, all sites ³ 800–829	982	35.3	5,658	203.5	5.8
Fracture of neck of femur ³ 820	321	11.6	2,246	80.8	7.0
Cerebrovascular disease 430–438	981	35.3	5,276	189.7	5.4
Diabetes mellitus 250	557	20.0	2,877	103.5	5.2
Chronic bronchitis 491	495	17.8	2,563	92.1	5.2
Volume depletion 276.5	485	17.4	1,947	70.0	4.0
Asthma 493	465	16.7	1,415	50.9	3.0
Osteoarthritis and allied disorders 715	441	15.9	1,994	71.7	4.5
Benign neoplasms 210–229	390	14.0	1,228	44.2	3.2
Cellulitis and abscess 681–682	353	12.7	1,856	66.7	5.3
Cholelithiasis 574	344	12.4	1,316	47.3	3.8
Intervertebral disc disorder 722	329	11.8	971	34.9	3.0
Septicemia 038	326	11.7	2,597	93.4	8.0

¹Population estimates used to compute rates are based on the 1990 census. See appendix.²Includes data for diagnostic conditions not shown in table.³Excludes fractures coded as 733.1, pathologic fracture.

Table 9. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
	Discharges in thousands			
All conditions ¹	12,396	4,678	5,119	2,599
Heart disease 391-392.0,393-398,402,404,410-416,420-429	2,854	1,111	1,185	558
Acute myocardial infarction 410	499	184	207	108
Coronary atherosclerosis 414.0	609	310	247	51
Other ischemic heart disease 411-413,414.1-414.9	157	70	63	24
Cardiac dysrhythmias 427	508	188	219	100
Congestive heart failure 428.0	767	239	323	204
Pneumonia 480-486	763	219	323	220
Cerebrovascular disease 430-438	711	233	317	161
Malignant neoplasms 140-208,230-234	617	292	241	84
Malignant neoplasm of large intestine and rectum 153-154,197.5	103	42	40	21
Fractures, all sites ² 800-829	521	116	211	194
Fracture of neck of femur ² 820	290	49	123	118
Chronic bronchitis 491	328	150	135	44
Osteoarthritis and allied disorders 715	303	173	110	19
Volume depletion 276.5	261	75	101	85
Psychoses 290-299	244	97	95	53
Septicemia 038	216	65	85	66
Diabetes mellitus 250	192	85	79	28
Diverticula of intestine 562	164	57	69	38
Cholelithiasis 574	127	61	49	17
Cellulitis and abscess 681-682	124	47	46	30
	Discharge rate per 10,000 population ³			
All conditions ¹	3,595.5	2,599.7	4,191.5	6,087.6
Heart disease 391-392.0,393-398,402,404,410-416,420-429	827.8	617.5	970.3	1,306.1
Acute myocardial infarction 410	144.9	102.4	169.6	252.7
Coronary atherosclerosis 414.0	176.5	172.4	202.5	119.9
Other ischemic heart disease 411-413,414.1-414.9	45.5	38.7	51.3	57.3
Cardiac dysrhythmias 427	147.3	104.7	179.7	233.8
Congestive heart failure 428.0	222.4	133.1	264.5	478.8
Pneumonia 480-486	221.2	122.0	264.8	514.6
Cerebrovascular disease 430-438	206.3	129.8	259.7	376.0
Malignant neoplasms 140-208,230-234	178.8	162.4	197.2	195.7
Malignant neoplasm of large intestine and rectum 153-154,197.5	29.8	23.1	32.9	49.4
Fractures, all sites ² 800-829	151.2	64.6	173.0	453.7
Fracture of neck of femur ² 820	84.0	27.0	100.6	276.7
Chronic bronchitis 491	95.2	83.2	110.3	102.8
Osteoarthritis and allied disorders 715	87.9	96.3	90.3	45.4
Volume depletion 276.5	75.8	41.7	82.8	199.9
Psychoses 290-299	70.7	53.7	77.4	123.3
Septicemia 038	62.7	36.4	69.4	154.8
Diabetes mellitus 250	55.7	47.4	64.4	66.0
Diverticula of intestine 562	47.7	31.9	56.2	89.5
Cholelithiasis 574	36.9	34.1	40.3	39.4
Cellulitis and abscess 681-682	35.8	26.4	38.0	69.5
	Average length of stay in days			
All conditions ¹	6.0	5.7	6.2	6.2
Heart disease 391-392.0,393-398,402,404,410-416,420-429	5.1	4.8	5.3	5.2
Acute myocardial infarction 410	6.2	5.9	6.2	6.7
Coronary atherosclerosis 414.0	4.4	4.1	4.8	3.9
Other ischemic heart disease 411-413,414.1-414.9	3.1	3.1	3.1	2.9
Cardiac dysrhythmias 427	4.1	3.8	4.2	4.4
Congestive heart failure 428.0	5.6	5.5	5.9	5.3
Pneumonia 480-486	6.5	6.4	6.3	7.0
Cerebrovascular disease 430-438	5.3	4.7	5.9	5.3

See footnotes at end of table.

Table 9. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Diagnostic category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
	Average length of stay in days			
Malignant neoplasms 140-208,230-234	7.2	7.0	7.2	7.5
Malignant neoplasm of large intestine and rectum 153-154,197.5	9.2	9.1	9.0	9.9
Fractures, all sites ² 800-829	6.6	5.9	7.4	6.1
Fracture of neck of femur ² 820	7.1	7.1	7.7	6.5
Chronic bronchitis 491	5.2	5.1	5.1	5.9
Osteoarthritis and allied disorders 715	4.7	4.8	4.6	4.8
Volume depletion 276.5	5.0	5.1	5.0	5.1
Psychoses 290-299	9.6	10.5	9.4	8.5
Septicemia 038	7.8	8.6	7.9	6.9
Diabetes mellitus 250	5.9	5.9	6.0	5.5
Diverticula of intestine 562	6.0	5.1	6.2	6.8
Cholelithiasis 574	5.0	4.4	5.3	5.8
Cellulitis and abscess 681-682	6.5	6.1	6.9	6.5

¹Includes data for diagnostic conditions not shown in table.

²Excludes fractures coded as 733.1, pathologic fracture.

³Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 10. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	31,706	2,383	9,969	6,958	12,396
Infectious and parasitic diseases 001-139	787	160	173	150	305
Septicemia 038	326	16	32	62	216
Neoplasms 140-239	1,587	37	289	566	695
Malignant neoplasms 140-208,230-234	1,156	27	120	393	617
Malignant neoplasm of large intestine and rectum 153-154,197.5	159	*	*7	49	103
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	139	*	8	43	87
Malignant neoplasm of breast 174-175,198.81	108	*	12	40	56
Benign neoplasms 210-229	390	*	163	161	58
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	1,455	152	289	381	634
Diabetes mellitus 250	557	22	136	207	192
Volume depletion 276.5	485	111	54	59	261
Diseases of the blood and blood-forming organs 280-289	392	70	91	83	149
Mental disorders 290-319	2,147	103	1,255	495	293
Psychoses 290-299	1,445	*	803	343	244
Alcohol dependence syndrome 303	146	*	92	49	6
Diseases of the nervous system and sense organs 320-389	471	84	112	109	167
Diseases of the circulatory system 390-459	6,294	30	389	1,813	4,062
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,385	16	244	1,271	2,854
Acute myocardial infarction 410	781	*	38	242	499
Coronary atherosclerosis 414.0	1,102	*	50	442	609
Other ischemic heart disease 411-413,414.1-414.9	283	*	21	105	157
Cardiac dysrhythmias 427	716	8	43	157	508
Congestive heart failure 428.0	999	*	35	195	767
Cerebrovascular disease 430-438	981	*	37	229	711
Diseases of the respiratory system 460-519	3,444	699	394	648	1,703
Acute bronchitis and bronchiolitis 466	285	201	13	19	52
Pneumonia 480-486	1,282	173	128	218	763
Chronic bronchitis 491	495	*	18	147	328
Asthma 493	465	203	111	84	68
Diseases of the digestive system 520-579	3,143	217	758	852	1,317
Appendicitis 540-543	273	68	145	43	16
Noninfectious enteritis and colitis 555-558	277	49	81	56	91
Diverticula of intestine 562	264	*	30	70	164
Cholelithiasis 574	344	*	108	107	127
Diseases of the genitourinary system 580-629	1,743	83	574	436	650
Calculus of kidney and ureter 592	177	*	80	62	31
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	504	*	503	*	...
Diseases of the skin and subcutaneous tissue 680-709	514	*	132	133	179
Cellulitis and abscess 681-682	353	28	96	106	124
Diseases of the musculoskeletal system and connective tissue 710-739	1,530	37	322	497	674
Osteoarthritis and allied disorders 715	441	*	13	125	303
Intervertebral disc disorders 722	329	*	140	132	56
Congenital anomalies 740-759	165	113	33	11	9
Certain conditions originating in the perinatal period 760-779	156	155	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	267	56	93	74	44
Injury and poisoning 800-999	2,466	228	712	514	1,012
Fractures, all sites ² 800-829	982	74	223	164	521
Fracture of neck of femur ² 820	321	*	*6	24	290
Poisonings 960-989	192	21	108	39	25
Supplementary classifications V01-V82	4,640	89	3,851	196	505
Females with deliveries V27	3,738	12	3,722	*3	...

*Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 11. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Rate per 10,000 population			
		Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	1,140.1	393.9	815.9	1,141.7	3,595.5
Infectious and parasitic diseases 001-139	28.3	26.5	14.1	24.6	88.3
Septicemia 038	11.7	2.6	2.6	10.1	62.7
Neoplasms 140-239	57.1	6.1	23.7	92.9	201.7
Malignant neoplasms 140-208,230-234	41.6	4.5	9.8	64.4	178.8
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.7	*	*0.6	8.0	29.8
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.0	*	0.6	7.0	25.1
Malignant neoplasm of breast 174-175,198.81	3.9	*	1.0	6.6	16.2
Benign neoplasms 210-229	14.0	*	13.3	26.5	16.9
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	52.3	25.1	23.6	62.4	183.8
Diabetes mellitus 250	20.0	3.7	11.1	33.9	55.7
Volume depletion 276.5	17.4	18.3	4.4	9.6	75.8
Diseases of the blood and blood-forming organs 280-289	14.1	11.5	7.4	13.6	43.1
Mental disorders 290-319	77.2	17.0	102.7	81.3	85.1
Psychoses 290-299	52.0	*	65.7	56.3	70.7
Alcohol dependence syndrome 303	5.3	*	7.5	8.0	1.6
Diseases of the nervous system and sense organs 320-389	16.9	13.9	9.1	17.9	48.3
Diseases of the circulatory system 390-459	226.3	5.0	31.8	297.5	1,178.3
Heart disease 391-392.0,393-398,402,404,410-416,420-429	157.7	2.6	20.0	208.6	827.8
Acute myocardial infarction 410	28.1	*	3.1	39.7	144.9
Coronary atherosclerosis 414.0	39.6	*	4.1	72.6	176.5
Other ischemic heart disease 411-413,414.1-414.9	10.2	*	1.8	17.2	45.5
Cardiac dysrhythmias 427	25.7	1.3	3.5	25.7	147.3
Congestive heart failure 428.0	35.9	*	2.9	32.0	222.4
Cerebrovascular disease 430-438	35.3	*	3.0	37.6	206.3
Diseases of the respiratory system 460-519	123.8	115.6	32.2	106.3	493.8
Acute bronchitis and bronchiolitis 466	10.3	33.3	1.1	3.1	15.2
Pneumonia 480-486	46.1	28.6	10.5	35.8	221.2
Chronic bronchitis 491	17.8	*	1.5	24.2	95.2
Asthma 493	16.7	33.6	9.1	13.7	19.6
Diseases of the digestive system 520-579	113.0	35.8	62.0	139.8	381.9
Appendicitis 540-543	9.8	11.3	11.9	7.1	4.7
Noninfectious enteritis and colitis 555-558	10.0	8.0	6.6	9.2	26.5
Diverticula of intestine 562	9.5	*	2.5	11.4	47.7
Cholelithiasis 574	12.4	*	8.8	17.5	36.9
Diseases of the genitourinary system 580-629	62.7	13.7	47.0	71.5	188.6
Calculus of kidney and ureter 592	6.3	*	6.6	10.1	9.1
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.1	*	41.1	*	...
Diseases of the skin and subcutaneous tissue 680-709	18.5	*	10.8	21.8	51.9
Cellulitis and abscess 681-682	12.7	4.6	7.9	17.4	35.8
Diseases of the musculoskeletal system and connective tissue 710-739	55.0	6.2	26.3	81.6	195.4
Osteoarthritis and allied disorders 715	15.9	*	1.1	20.5	87.9
Intervertebral disc disorders 722	11.8	*	11.4	21.7	16.3
Congenital anomalies 740-759	5.9	18.7	2.7	1.8	2.6
Certain conditions originating in the perinatal period 760-779	5.6	25.6	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	9.6	9.3	7.6	12.1	12.8
Injury and poisoning 800-999	88.7	37.7	58.3	84.4	293.5
Fractures, all sites ² 800-829	35.3	12.3	18.3	26.8	151.2
Fracture of neck of femur ² 820	11.6	*	*0.5	3.9	84.0
Poisonings 960-989	6.9	3.4	8.8	6.5	7.2
Supplementary classifications V01-V82	166.8	14.6	315.1	32.2	146.3
Females with deliveries V27	134.4	2.0	304.6	*0.5	...

*Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 12. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	4.9	4.5	3.7	5.0	6.0
Infectious and parasitic diseases 001-139	6.3	3.3	6.6	6.5	7.5
Septicemia 038	8.0	5.5	9.6	8.3	7.8
Neoplasms 140-239	5.8	6.5	4.2	5.3	7.0
Malignant neoplasms 140-208,230-234	6.7	7.6	6.4	6.2	7.2
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	*	*7.2	7.4	9.2
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.7	*	7.3	6.2	6.9
Malignant neoplasm of breast 174-175,198.81	2.9	*	2.3	2.1	3.7
Benign neoplasms 210-229	3.2	3.0	2.5	3.1	5.2
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	4.5	2.8	3.6	4.8	5.1
Diabetes mellitus 250	5.2	3.1	3.8	5.6	5.9
Volume depletion 276.5	4.0	2.2	3.1	3.8	5.0
Diseases of the blood and blood-forming organs 280-289	4.4	4.0	4.7	4.5	4.5
Mental disorders 290-319	7.3	12.4	6.5	7.5	8.8
Psychoses 290-299	8.1	11.5	7.3	8.3	9.6
Alcohol dependence syndrome 303	5.7	*	5.8	5.7	4.6
Diseases of the nervous system and sense organs 320-389	4.8	3.9	4.5	4.3	5.7
Diseases of the circulatory system 390-459	4.8	5.6	3.9	4.2	5.2
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	5.4	3.6	3.9	5.1
Acute myocardial infarction 410	5.7	*	3.8	4.8	6.2
Coronary atherosclerosis 414.0	3.9	*	2.9	3.3	4.4
Other ischemic heart disease 411-413,414.1-414.9	2.7	*	2.1	2.3	3.1
Cardiac dysrhythmias 427	3.7	3.5	2.4	2.9	4.1
Congestive heart failure 428.0	5.5	*	4.4	4.9	5.6
Cerebrovascular disease 430-438	5.4	*	6.4	5.3	5.3
Diseases of the respiratory system 460-519	5.4	3.3	4.5	5.6	6.4
Acute bronchitis and bronchiolitis 466	3.4	3.1	3.8	3.0	4.8
Pneumonia 480-486	5.9	3.4	5.3	5.9	6.5
Chronic bronchitis 491	5.2	*	4.6	5.1	5.2
Asthma 493	3.0	2.2	2.9	3.9	4.9
Diseases of the digestive system 520-579	4.7	3.3	3.8	4.7	5.5
Appendicitis 540-543	3.3	3.0	3.1	3.6	5.1
Noninfectious enteritis and colitis 555-558	4.6	2.3	4.5	4.7	5.8
Diverticula of intestine 562	5.5	*	4.4	4.8	6.0
Cholelithiasis 574	3.8	*	2.9	3.4	5.0
Diseases of the genitourinary system 580-629	3.8	3.3	2.9	3.4	4.8
Calculus of kidney and ureter 592	2.4	*	2.0	2.7	2.8
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.5	*	2.5	*	...
Diseases of the skin and subcutaneous tissue 680-709	5.5	3.1	4.4	6.0	7.0
Cellulitis and abscess 681-682	5.3	3.2	4.1	5.4	6.5
Diseases of the musculoskeletal system and connective tissue 710-739	4.1	4.4	3.0	3.8	4.9
Osteoarthritis and allied disorders 715	4.5	*	3.7	4.1	4.7
Intervertebral disc disorders 722	3.0	*	2.3	2.8	4.9
Congenital anomalies 740-759	5.7	6.4	3.9	4.7	5.0
Certain conditions originating in the perinatal period 760-779	10.1	10.2	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	2.4	2.1	1.9	2.1	4.2
Injury and poisoning 800-999	5.4	4.1	4.4	5.5	6.4
Fractures, all sites ² 800-829	5.8	3.1	4.8	5.6	6.6
Fracture of neck of femur ² 820	7.0	*	*5.1	6.3	7.1
Poisonings 960-989	2.9	1.8	2.5	3.0	*
Supplementary classifications V01-V82	3.9	5.5	2.6	8.5	11.6
Females with deliveries V27	2.5	2.4	2.5	*2.9	...

*Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 13. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Number in thousands	
		Male	Female
All conditions	31,706	12,514	19,192
Infectious and parasitic diseases 001-139	787	389	398
Septicemia 038	326	154	172
Neoplasms 140-239	1,587	586	1,002
Malignant neoplasms 140-208,230-234	1,156	528	629
Malignant neoplasm of large intestine and rectum 153-154,197.5	159	80	79
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	139	72	66
Malignant neoplasm of breast 174-175,198.81	108	*	108
Benign neoplasms 210-229	390	40	350
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	1,455	625	830
Diabetes mellitus 250	557	279	278
Volume depletion 276.5	485	206	279
Diseases of the blood and blood-forming organs 280-289	392	163	229
Mental disorders 290-319	2,147	1,104	1,043
Psychoses 290-299	1,445	712	733
Alcohol dependence syndrome 303	146	107	40
Diseases of the nervous system and sense organs 320-389	471	196	275
Diseases of the circulatory system 390-459	6,294	3,115	3,179
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,385	2,243	2,142
Acute myocardial infarction 410	781	453	328
Coronary atherosclerosis 414.0	1,102	676	425
Other ischemic heart disease 411-413,414.1-414.9	283	145	139
Cardiac dysrhythmias 427	716	336	380
Congestive heart failure 428.0	999	418	581
Cerebrovascular disease 430-438	981	428	553
Diseases of the respiratory system 460-519	3,444	1,603	1,841
Acute bronchitis and bronchiolitis 466	285	145	140
Pneumonia 480-486	1,282	583	699
Chronic bronchitis 491	495	215	280
Asthma 493	465	198	267
Diseases of the digestive system 520-579	3,143	1,376	1,767
Appendicitis 540-543	273	156	117
Noninfectious enteritis and colitis 555-558	277	105	172
Diverticula of intestine 562	264	105	159
Cholelithiasis 574	344	111	233
Diseases of the genitourinary system 580-629	1,743	533	1,211
Calculus of kidney and ureter 592	177	104	73
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	504	...	504
Diseases of the skin and subcutaneous tissue 680-709	514	259	255
Cellulitis and abscess 681-682	353	184	169
Diseases of the musculoskeletal system and connective tissue 710-739	1,530	666	865
Osteoarthritis and allied disorders 715	441	175	267
Intervertebral disc disorders 722	329	175	154
Congenital anomalies 740-759	165	94	71
Certain conditions originating in the perinatal period 760-779	156	86	69
Symptoms, signs, and ill-defined conditions 780-799	267	127	139
Injury and poisoning 800-999	2,466	1,214	1,252
Fractures, all sites ² 800-829	982	413	569
Fracture of neck of femur ² 820	321	81	240
Poisonings 960-989	192	72	120
Supplementary classifications V01-V82	4,640	378	4,261
Females with deliveries V27	3,738	...	3,738

*Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 14. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Rate per 10,000 population	
		Male	Female
All conditions	1,140.1	920.2	1,350.5
Infectious and parasitic diseases 001-139	28.3	28.6	28.0
Septicemia 038	11.7	11.3	12.1
Neoplasms 140-239	57.1	43.1	70.5
Malignant neoplasms 140-208,230-234	41.6	38.8	44.2
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.7	5.9	5.5
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.0	5.3	4.7
Malignant neoplasm of breast 174-175,198.81	3.9	*	7.6
Benign neoplasms 210-229	14.0	2.9	24.6
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	52.3	46.0	58.4
Diabetes mellitus 250	20.0	20.5	19.6
Volume depletion 276.5	17.4	15.2	19.6
Diseases of the blood and blood-forming organs 280-289	14.1	12.0	16.1
Mental disorders 290-319	77.2	81.2	73.4
Psychoses 290-299	52.0	52.4	51.6
Alcohol dependence syndrome 303	5.3	7.8	2.8
Diseases of the nervous system and sense organs 320-389	16.9	14.4	19.4
Diseases of the circulatory system 390-459	226.3	229.1	223.7
Heart disease 391-392.0,393-398,402,404,410-416,420-429	157.7	165.0	150.7
Acute myocardial infarction 410	28.1	33.3	23.1
Coronary atherosclerosis 414.0	39.6	49.7	29.9
Other ischemic heart disease 411-413,414.1-414.9	10.2	10.6	9.8
Cardiac dysrhythmias 427	25.7	24.7	26.8
Congestive heart failure 428.0	35.9	30.7	40.9
Cerebrovascular disease 430-438	35.3	31.5	38.9
Diseases of the respiratory system 460-519	123.8	117.9	129.5
Acute bronchitis and bronchiolitis 466	10.3	10.7	9.9
Pneumonia 480-486	46.1	42.9	49.2
Chronic bronchitis 491	17.8	15.8	19.7
Asthma 493	16.7	14.5	18.8
Diseases of the digestive system 520-579	113.0	101.2	124.4
Appendicitis 540-543	9.8	11.5	8.3
Noninfectious enteritis and colitis 555-558	10.0	7.7	12.1
Diverticula of intestine 562	9.5	7.7	11.2
Cholelithiasis 574	12.4	8.2	16.4
Diseases of the genitourinary system 580-629	62.7	39.2	85.2
Calculus of kidney and ureter 592	6.3	7.6	5.1
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.1	...	35.5
Diseases of the skin and subcutaneous tissue 680-709	18.5	19.1	17.9
Cellulitis and abscess 681-682	12.7	13.6	11.9
Diseases of the musculoskeletal system and connective tissue 710-739	55.0	49.0	60.8
Osteoarthritis and allied disorders 715	15.9	12.8	18.8
Intervertebral disc disorders 722	11.8	12.9	10.9
Congenital anomalies 740-759	5.9	6.9	5.0
Certain conditions originating in the perinatal period 760-779	5.6	6.3	4.9
Symptoms, signs, and ill-defined conditions 780-799	9.6	9.4	9.8
Injury and poisoning 800-999	88.7	89.3	88.1
Fractures, all sites ² 800-829	35.3	30.4	40.0
Fracture of neck of femur ² 820	11.6	6.0	16.9
Poisonings 960-989	6.9	5.3	8.5
Supplementary classifications V01-V82	166.8	27.8	299.9
Females with deliveries V27	134.4	...	263.0

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 15. Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Average length of stay in days	
		Male	Female
All conditions	4.9	5.3	4.6
Infectious and parasitic diseases 001-139	6.3	6.2	6.3
Septicemia 038	8.0	7.6	8.3
Neoplasms 140-239	5.8	7.0	5.2
Malignant neoplasms 140-208,230-234	6.7	7.2	6.4
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	9.0	8.1
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.7	6.3	7.0
Malignant neoplasm of breast 174-175,198.81	2.9	*	2.9
Benign neoplasms 210-229	3.2	4.5	3.0
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	4.5	4.6	4.4
Diabetes mellitus 250	5.2	5.4	5.0
Volume depletion 276.5	4.0	3.9	4.1
Diseases of the blood and blood-forming organs 280-289	4.4	4.3	4.5
Mental disorders 290-319	7.3	7.2	7.5
Psychoses 290-299	8.1	8.2	8.1
Alcohol dependence syndrome 303	5.7	5.4	6.7
Diseases of the nervous system and sense organs 320-389	4.8	5.2	4.5
Diseases of the circulatory system 390-459	4.8	4.7	5.0
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	4.5	4.9
Acute myocardial infarction 410	5.7	5.2	6.3
Coronary atherosclerosis 414.0	3.9	3.8	4.0
Other ischemic heart disease 411-413,414.1-414.9	2.7	2.6	2.8
Cardiac dysrhythmias 427	3.7	3.6	3.8
Congestive heart failure 428.0	5.5	5.6	5.4
Cerebrovascular disease 430-438	5.4	5.2	5.5
Diseases of the respiratory system 460-519	5.4	5.3	5.5
Acute bronchitis and bronchiolitis 466	3.4	3.2	3.7
Pneumonia 480-486	5.9	5.6	6.1
Chronic bronchitis 491	5.2	5.2	5.2
Asthma 493	3.0	2.6	3.4
Diseases of the digestive system 520-579	4.7	4.7	4.8
Appendicitis 540-543	3.3	3.3	3.2
Noninfectious enteritis and colitis 555-558	4.6	4.0	4.9
Diverticula of intestine 562	5.5	4.9	5.9
Cholelithiasis 574	3.8	4.4	3.6
Diseases of the genitourinary system 580-629	3.8	4.1	3.6
Calculus of kidney and ureter 592	2.4	2.0	2.9
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.5	...	2.5
Diseases of the skin and subcutaneous tissue 680-709	5.5	5.5	5.6
Cellulitis and abscess 681-682	5.3	5.1	5.4
Diseases of the musculoskeletal system and connective tissue 710-739	4.1	4.0	4.3
Osteoarthritis and allied disorders 715	4.5	4.5	4.5
Intervertebral disc disorders 722	3.0	2.6	3.3
Congenital anomalies 740-759	5.7	5.4	6.1
Certain conditions originating in the perinatal period 760-779	10.1	11.5	8.4
Symptoms, signs, and ill-defined conditions 780-799	2.4	2.3	2.5
Injury and poisoning 800-999	5.4	5.4	5.4
Fractures, all sites ² 800-829	5.8	5.9	5.6
Fracture of neck of femur ² 820	7.0	7.3	6.9
Poisonings 960-989	2.9	2.7	3.0
Supplementary classifications V01-V82	3.9	9.2	3.4
Females with deliveries V27	2.5	...	2.5

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 16. Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions	31,706	7,103	7,207	12,016	5,380
Infectious and parasitic diseases 001-139	787	189	169	288	140
Septicemia 038	326	87	76	114	49
Neoplasms 140-239	1,587	383	333	578	295
Malignant neoplasms 140-208,230-234	1,156	287	245	411	213
Malignant neoplasm of large intestine and rectum 153-154,197.5	159	35	40	56	28
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	139	32	33	49	24
Malignant neoplasm of breast 174-175,198.81	108	29	25	34	19
Benign neoplasms 210-229	390	87	81	148	73
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	1,455	318	350	574	213
Diabetes mellitus 250	557	129	120	228	80
Volume depletion 276.5	485	100	130	193	62
Diseases of the blood and blood-forming organs 280-289	392	92	87	154	59
Mental disorders 290-319	2,147	692	446	*	185
Psychoses 290-299	1,445	459	312	*	113
Alcohol dependence syndrome 303	146	28	25	*	18
Diseases of the nervous system and sense organs 320-389	471	108	115	167	80
Diseases of the circulatory system 390-459	6,294	1,507	1,509	2,382	895
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,385	1,085	1,062	1,627	610
Acute myocardial infarction 410	781	190	185	283	122
Coronary atherosclerosis 414.0	1,102	254	274	426	147
Other ischemic heart disease 411-413,414.1-414.9	283	86	61	101	36
Cardiac dysrhythmias 427	716	181	176	252	106
Congestive heart failure 428.0	999	235	257	384	122
Cerebrovascular disease 430-438	981	217	230	384	151
Diseases of the respiratory system 460-519	3,444	765	824	1,286	569
Acute bronchitis and bronchiolitis 466	285	43	64	99	*
Pneumonia 480-486	1,282	263	336	480	203
Chronic bronchitis 491	495	126	111	198	59
Asthma 493	465	110	109	167	78
Diseases of the digestive system 520-579	3,143	693	732	1,150	569
Appendicitis 540-543	273	56	62	84	70
Noninfectious enteritis and colitis 555-558	277	57	68	103	48
Diverticula of intestine 562	264	63	67	95	39
Cholelithiasis 574	344	79	73	131	61
Diseases of the genitourinary system 580-629	1,743	367	392	692	292
Calculus of kidney and ureter 592	177	38	45	78	15
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	504	99	93	211	102
Diseases of the skin and subcutaneous tissue 680-709	514	115	115	210	74
Cellulitis and abscess 681-682	353	93	87	116	57
Diseases of the musculoskeletal system and connective tissue 710-739	1,530	304	376	565	285
Osteoarthritis and allied disorders 715	441	81	117	161	82
Intervertebral disc disorders 722	329	62	80	130	57
Congenital anomalies 740-759	165	*	*	57	*
Certain conditions originating in the perinatal period 760-779	156	26	*	42	*
Symptoms, signs, and ill-defined conditions 780-799	267	63	53	119	32
Injury and poisoning 800-999	2,466	527	580	907	453
Fractures, all sites ² 800-829	982	195	228	374	186
Fracture of neck of femur ² 820	321	63	74	127	57
Poisonings 960-989	192	40	49	69	34
Supplementary classifications V01-V82	4,640	814	974	1,808	1,043
Females with deliveries V27	3,738	632	771	1,439	895

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 17. Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
All conditions	1,140.1	1,354.9	1,128.3	1,216.4	853.7
Infectious and parasitic diseases 001-139	28.3	36.1	26.5	29.2	22.3
Septicemia 038	11.7	16.6	11.9	11.6	7.8
Neoplasms 140-239	57.1	73.0	52.1	58.5	46.7
Malignant neoplasms 140-208,230-234	41.6	54.7	38.4	41.6	33.9
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.7	6.6	6.2	5.7	4.5
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.0	6.2	5.1	5.0	3.9
Malignant neoplasm of breast 174-175,198.81	3.9	5.6	3.9	3.5	3.0
Benign neoplasms 210-229	14.0	16.7	12.7	15.0	11.6
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	52.3	60.6	54.8	58.1	33.8
Diabetes mellitus 250	20.0	24.7	18.8	23.1	12.7
Volume depletion 276.5	17.4	19.0	20.4	19.5	9.8
Diseases of the blood and blood-forming organs 280-289	14.1	17.5	13.7	15.5	9.4
Mental disorders 290-319	77.2	132.0	69.8	*	29.3
Psychoses 290-299	52.0	87.5	48.9	*	18.0
Alcohol dependence syndrome 303	5.3	5.4	3.9	*	2.8
Diseases of the nervous system and sense organs 320-389	16.9	20.6	18.1	16.9	12.8
Diseases of the circulatory system 390-459	226.3	287.5	236.3	241.1	142.1
Heart disease 391-392.0,393-398,402,404,410-416,420-429	157.7	207.1	166.3	164.8	96.8
Acute myocardial infarction 410	28.1	36.3	28.9	28.7	19.4
Coronary atherosclerosis 414.0	39.6	48.5	42.9	43.2	23.3
Other ischemic heart disease 411-413,414.1-414.9	10.2	16.4	9.5	10.2	5.7
Cardiac dysrhythmias 427	25.7	34.6	27.6	25.5	16.9
Congestive heart failure 428.0	35.9	44.8	40.3	38.9	19.4
Cerebrovascular disease 430-438	35.3	41.3	36.0	38.9	23.9
Diseases of the respiratory system 460-519	123.8	146.0	128.9	130.2	90.3
Acute bronchitis and bronchiolitis 466	10.3	8.3	10.1	10.0	*
Pneumonia 480-486	46.1	50.1	52.7	48.6	32.2
Chronic bronchitis 491	17.8	24.1	17.4	20.1	9.4
Asthma 493	16.7	21.1	17.1	16.9	12.4
Diseases of the digestive system 520-579	113.0	132.2	114.6	116.4	90.2
Appendicitis 540-543	9.8	10.7	9.8	8.5	11.2
Noninfectious enteritis and colitis 555-558	10.0	10.9	10.6	10.5	7.7
Diverticula of intestine 562	9.5	12.0	10.4	9.6	6.2
Cholelithiasis 574	12.4	15.0	11.5	13.3	9.7
Diseases of the genitourinary system 580-629	62.7	70.0	61.4	70.1	46.4
Calculus of kidney and ureter 592	6.3	7.2	7.1	7.9	2.4
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	18.1	18.8	14.6	21.4	16.1
Diseases of the skin and subcutaneous tissue 680-709	18.5	22.0	18.0	21.2	11.7
Cellulitis and abscess 681-682	12.7	17.8	13.6	11.8	9.0
Diseases of the musculoskeletal system and connective tissue 710-739	55.0	58.0	58.8	57.2	45.3
Osteoarthritis and allied disorders 715	15.9	15.5	18.3	16.3	13.0
Intervertebral disc disorders 722	11.8	11.9	12.5	13.2	9.0
Congenital anomalies 740-759	5.9	*	*	5.8	*
Certain conditions originating in the perinatal period 760-779	5.6	5.0	*	4.2	*
Symptoms, signs, and ill-defined conditions 780-799	9.6	12.0	8.3	12.1	5.0
Injury and poisoning 800-999	88.7	100.5	90.7	91.8	71.9
Fractures, all sites ² 800-829	35.3	37.1	35.6	37.9	29.6
Fracture of neck of femur ² 820	11.6	12.0	11.6	12.9	9.0
Poisonings 960-989	6.9	7.7	7.7	7.0	5.4
Supplementary classifications V01-V82	166.8	155.3	152.5	183.0	165.5
Females with deliveries V27	134.4	120.7	120.8	145.7	142.0

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 18. Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Average length of stay in days				
All conditions	4.9	5.7	4.5	4.9	4.5
Infectious and parasitic diseases 001-139	6.3	7.7	5.3	6.2	5.6
Septicemia 038	8.0	8.9	6.4	8.4	7.6
Neoplasms 140-239	5.8	6.5	5.4	5.9	5.3
Malignant neoplasms 140-208,230-234	6.7	7.5	6.2	6.9	6.1
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	10.0	8.3	8.4	7.5
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.7	8.3	5.9	6.7	5.6
Malignant neoplasm of breast 174-175,198.81	2.9	3.0	2.4	*	1.7
Benign neoplasms 210-229	3.2	3.4	3.0	3.0	3.3
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	4.5	5.3	4.0	4.5	4.2
Diabetes mellitus 250	5.2	6.0	4.5	5.0	5.2
Volume depletion 276.5	4.0	5.0	3.5	4.1	3.3
Diseases of the blood and blood-forming organs 280-289	4.4	5.1	4.2	4.4	4.0
Mental disorders 290-319	7.3	8.5	6.4	7.1	6.5
Psychoses 290-299	8.1	9.3	7.1	7.9	7.1
Alcohol dependence syndrome 303	5.7	6.8	6.5	4.9	6.2
Diseases of the nervous system and sense organs 320-389	4.8	5.1	4.6	4.7	4.8
Diseases of the circulatory system 390-459	4.8	5.5	4.4	4.8	4.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.7	5.4	4.2	4.7	4.2
Acute myocardial infarction 410	5.7	6.4	5.4	5.6	5.0
Coronary atherosclerosis 414.0	3.9	4.2	3.5	4.0	3.7
Other ischemic heart disease 411-413,414.1-414.9	2.7	3.1	2.4	2.6	2.6
Cardiac dysrhythmias 427	3.7	4.2	3.2	3.9	3.1
Congestive heart failure 428.0	5.5	6.7	4.8	5.3	4.8
Cerebrovascular disease 430-438	5.4	6.0	5.0	5.3	5.1
Diseases of the respiratory system 460-519	5.4	5.9	4.8	5.5	5.4
Acute bronchitis and bronchiolitis 466	3.4	4.0	3.1	3.5	3.4
Pneumonia 480-486	5.9	6.2	5.4	6.2	5.6
Chronic bronchitis 491	5.2	5.6	4.6	5.2	5.1
Asthma 493	3.0	3.5	2.6	3.2	2.6
Diseases of the digestive system 520-579	4.7	5.5	4.3	4.7	4.5
Appendicitis 540-543	3.3	3.0	3.2	3.3	3.5
Noninfectious enteritis and colitis 555-558	4.6	5.1	4.6	4.2	4.7
Diverticula of intestine 562	5.5	6.5	5.3	5.1	5.1
Cholelithiasis 574	3.8	4.6	3.5	3.6	3.7
Diseases of the genitourinary system 580-629	3.8	4.6	3.5	3.6	3.4
Calculus of kidney and ureter 592	2.4	2.9	2.7	2.1	1.9
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.5	2.6	2.6	2.4	2.6
Diseases of the skin and subcutaneous tissue 680-709	5.5	5.6	5.1	5.9	5.2
Cellulitis and abscess 681-682	5.3	5.5	4.8	5.6	4.8
Diseases of the musculoskeletal system and connective tissue 710-739	4.1	4.6	4.1	4.0	3.9
Osteoarthritis and allied disorders 715	4.5	5.1	4.3	4.6	4.0
Intervertebral disc disorders 722	3.0	3.0	3.2	2.6	3.3
Congenital anomalies 740-759	5.7	6.4	6.4	4.9	5.6
Certain conditions originating in the perinatal period 760-779	10.1	9.9	9.9	12.4	8.7
Symptoms, signs, and ill-defined conditions 780-799	2.4	2.4	2.1	2.5	2.4
Injury and poisoning 800-999	5.4	5.9	4.8	5.6	5.3
Fractures, all sites ² 800-829	5.8	6.7	5.0	5.8	5.6
Fracture of neck of femur ² 820	7.0	7.4	6.1	7.2	7.1
Poisonings 960-989	2.9	3.3	2.2	2.5	*
Supplementary classifications V01-V82	3.9	4.2	4.0	4.0	3.4
Females with deliveries V27	2.5	2.8	2.4	2.5	2.3

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 19. Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Number in thousands			
All conditions	138,032	6,324	34,149	31,372	66,187
Infectious and parasitic diseases 001-139	3,441	437	840	752	1,411
Septicemia 038	637	43	78	136	380
Neoplasms 140-239	4,237	139	649	1,395	2,054
Malignant neoplasms 140-208,230-234	3,236	115	318	1,031	1,772
Malignant neoplasm of large intestine and rectum 153-154,197.5	254	*	11	77	166
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	443	*	27	137	272
Malignant neoplasm of breast 174-175,198.81	186	*	21	66	98
Benign neoplasms 210-229	871	*	316	336	207
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	14,727	486	1,843	4,236	8,162
Diabetes mellitus 250	4,658	28	485	1,553	2,592
Volume depletion 276.5	2,176	304	293	332	1,248
Diseases of the blood and blood-forming organs 280-289	4,158	188	877	921	2,172
Mental disorders 290-319	9,688	300	4,256	2,493	2,639
Psychoses 290-299	3,400	*	1,266	676	1,385
Alcohol dependence syndrome 303	717	*	367	270	79
Diseases of the nervous system and sense organs 320-389	3,494	405	557	739	1,792
Diseases of the circulatory system 390-459	29,894	115	1,558	7,321	20,900
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,916	71	740	4,048	13,057
Acute myocardial infarction 410	941	*	45	280	616
Coronary atherosclerosis 414.0	4,144	*	112	1,119	2,908
Other ischemic heart disease 411-413,414.1-414.9	2,375	*	96	758	1,520
Cardiac dysrhythmias 427	3,748	27	153	615	2,954
Congestive heart failure 428.0	3,226	11	85	561	2,569
Cerebrovascular disease 430-438	1,984	*	61	406	1,508
Diseases of the respiratory system 460-519	10,262	1,198	1,266	2,199	5,598
Acute bronchitis and bronchiolitis 466	424	239	40	41	104
Pneumonia 480-486	2,055	258	211	357	1,228
Chronic bronchitis 491	1,024	*	45	270	707
Asthma 493	1,391	314	402	334	342
Diseases of the digestive system 520-579	9,131	452	1,795	2,527	4,357
Appendicitis 540-543	305	71	164	50	20
Noninfectious enteritis and colitis 555-558	606	109	152	123	223
Diverticula of intestine 562	546	*	39	111	396
Cholelithiasis 574	604	*	166	177	256
Diseases of the genitourinary system 580-629	7,451	167	1,955	1,852	3,477
Calculus of kidney and ureter 592	258	*5	107	84	62
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,547	27	8,508	*12	...
Diseases of the skin and subcutaneous tissue 680-709	1,822	153	344	477	848
Cellulitis and abscess 681-682	799	47	179	241	332
Diseases of the musculoskeletal system and connective tissue 710-739	5,499	74	823	1,430	3,172
Osteoarthritis and allied disorders 715	1,461	*	45	285	1,130
Intervertebral disc disorders 722	490	*	182	189	118
Congenital anomalies 740-759	626	353	128	75	70
Certain conditions originating in the perinatal period 760-779	444	440	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	7,289	624	1,520	1,926	3,219
Injury and poisoning 800-999	5,770	465	1,772	1,241	2,291
Fractures, all sites ² 800-829	1,542	102	416	262	763
Fracture of neck of femur ² 820	371	*	8	28	333
Poisonings 960-989	343	28	191	75	49
Supplementary classifications V01-V82	11,553	302	5,453	1,774	4,024
Females with deliveries V27	3,738	12	3,722	*3	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹The diagnosis of female with delivery is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 20. Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	Both sexes	Number in thousands	
		Male	Female
All conditions	138,032	55,138	82,894
Infectious and parasitic diseases 001-139	3,441	1,538	1,903
Septicemia 038	637	316	321
Neoplasms 140-239	4,237	1,752	2,485
Malignant neoplasms 140-208,230-234	3,236	1,565	1,671
Malignant neoplasm of large intestine and rectum 153-154,197.5	254	133	121
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	443	240	203
Malignant neoplasm of breast 174-175,198.81	186	*	185
Benign neoplasms 210-229	871	126	744
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	14,727	5,992	8,735
Diabetes mellitus 250	4,658	2,056	2,602
Volume depletion 276.5	2,176	901	1,275
Diseases of the blood and blood-forming organs 280-289	4,158	1,626	2,532
Mental disorders 290-319	9,688	4,702	4,985
Psychoses 290-299	3,400	1,554	1,846
Alcohol dependence syndrome 303	717	514	203
Diseases of the nervous system and sense organs 320-389	3,494	1,517	1,977
Diseases of the circulatory system 390-459	29,894	13,960	15,933
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,916	8,755	9,161
Acute myocardial infarction 410	941	533	408
Coronary atherosclerosis 414.0	4,144	2,296	1,848
Other ischemic heart disease 411-413,414.1-414.9	2,375	1,317	1,058
Cardiac dysrhythmias 427	3,748	1,781	1,967
Congestive heart failure 428.0	3,226	1,334	1,893
Cerebrovascular disease 430-438	1,984	880	1,104
Diseases of the respiratory system 460-519	10,262	4,766	5,496
Acute bronchitis and bronchiolitis 466	424	207	217
Pneumonia 480-486	2,055	956	1,099
Chronic bronchitis 491	1,024	448	576
Asthma 493	1,391	490	901
Diseases of the digestive system 520-579	9,131	3,901	5,231
Appendicitis 540-543	305	166	138
Noninfectious enteritis and colitis 555-558	606	232	374
Diverticula of intestine 562	546	199	347
Cholelithiasis 574	604	205	399
Diseases of the genitourinary system 580-629	7,451	2,464	4,987
Calculus of kidney and ureter 592	258	135	122
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,547	...	8,547
Diseases of the skin and subcutaneous tissue 680-709	1,822	882	940
Cellulitis and abscess 681-682	799	407	392
Diseases of the musculoskeletal system and connective tissue 710-739	5,499	1,928	3,571
Osteoarthritis and allied disorders 715	1,461	482	979
Intervertebral disc disorders 722	490	249	241
Congenital anomalies 740-759	626	317	309
Certain conditions originating in the perinatal period 760-779	444	261	183
Symptoms, signs, and ill-defined conditions 780-799	7,289	3,227	4,063
Injury and poisoning 800-999	5,770	2,945	2,825
Fractures, all sites ² 800-829	1,542	705	837
Fracture of neck of femur ² 820	371	96	275
Poisonings 960-989	343	126	218
Supplementary classifications V01-V82	11,553	3,360	8,192
Females with deliveries V27	3,738	...	3,738

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹The diagnosis of female with delivery is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 21. Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions	138,032	31,635	32,815	50,272	23,309
Infectious and parasitic diseases 001-139	3,441	799	782	1,246	614
Septicemia 038	637	169	136	233	99
Neoplasms 140-239	4,237	1,061	923	1,500	754
Malignant neoplasms 140-208,230-234	3,236	831	720	1,129	557
Malignant neoplasm of large intestine and rectum 153-154,197.5	254	54	69	86	44
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	443	95	107	173	68
Malignant neoplasm of breast 174-175,198.81	186	49	43	60	34
Benign neoplasms 210-229	871	197	175	328	171
Endocrine, nutritional, and metabolic diseases, and immunity disorders 240-279	14,727	3,411	3,586	5,442	2,287
Diabetes mellitus 250	4,658	1,149	1,064	1,781	664
Volume depletion 276.5	2,176	459	532	831	354
Diseases of the blood and blood-forming organs 280-289	4,158	954	969	1,547	687
Mental disorders 290-319	9,688	2,635	2,252	3,536	1,266
Psychoses 290-299	3,400	890	750	1,380	380
Alcohol dependence syndrome 303	717	232	*	218	95
Diseases of the nervous system and sense organs 320-389	3,494	810	863	1,189	632
Diseases of the circulatory system 390-459	29,894	7,372	7,266	10,891	4,365
Heart disease 391-392.0,393-398,402,404,410-416,420-429	17,916	4,576	4,399	6,344	2,598
Acute myocardial infarction 410	941	234	223	333	151
Coronary atherosclerosis 414.0	4,144	1,061	1,045	1,477	561
Other ischemic heart disease 411-413,414.1-414.9	2,375	641	569	838	326
Cardiac dysrhythmias 427	3,748	964	924	1,249	611
Congestive heart failure 428.0	3,226	747	843	1,211	425
Cerebrovascular disease 430-438	1,984	432	463	787	302
Diseases of the respiratory system 460-519	10,262	2,379	2,435	3,720	1,728
Acute bronchitis and bronchiolitis 466	424	72	97	151	105
Pneumonia 480-486	2,055	430	515	772	338
Chronic bronchitis 491	1,024	250	225	411	139
Asthma 493	1,391	387	325	434	246
Diseases of the digestive system 520-579	9,131	2,093	2,177	3,318	1,543
Appendicitis 540-543	305	62	72	94	76
Noninfectious enteritis and colitis 555-558	606	133	141	234	99
Diverticula of intestine 562	546	141	130	195	80
Cholelithiasis 574	604	142	135	217	110
Diseases of the genitourinary system 580-629	7,451	1,621	1,722	2,862	1,246
Calculus of kidney and ureter 592	258	61	60	108	29
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	8,547	1,519	1,776	3,182	2,069
Diseases of the skin and subcutaneous tissue 680-709	1,822	422	459	664	276
Cellulitis and abscess 681-682	799	193	203	280	124
Diseases of the musculoskeletal system and connective tissue 710-739	5,499	1,212	1,438	1,946	903
Osteoarthritis and allied disorders 715	1,461	310	404	537	209
Intervertebral disc disorders 722	490	94	117	193	86
Congenital anomalies 740-759	626	152	128	195	*
Certain conditions originating in the perinatal period 760-779	444	75	*	122	*
Symptoms, signs, and ill-defined conditions 780-799	7,289	1,543	1,900	2,634	1,213
Injury and poisoning 800-999	5,770	1,232	1,373	2,058	1,107
Fractures, all sites ² 800-829	1,542	291	358	586	306
Fracture of neck of femur ² 820	371	71	85	150	64
Poisonings 960-989	343	66	99	120	59
Supplementary classifications V01-V82	11,553	2,345	2,674	4,221	2,313
Females with deliveries V27	3,738	632	772	1,439	895

* Figure does not meet standard of reliability or precision.

¹The diagnosis of female with delivery is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 22. Number and rate of discharges from short-stay hospitals with human immunodeficiency virus diagnoses, by selected characteristics: United States, selected years 1990–2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Data are for discharges with at least one of the following *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) codes as a first-listed or secondary diagnosis: 042–044, 279.19, 795.8 (in 1990); 042, V08 (beginning in 1995)]

Characteristic	1990	1995	1996	1997	1998	1999	2000
Number in thousands							
All HIV discharges ¹	146	249	227	178	189	180	173
Sex							
Male	114	183	165	123	132	121	115
Female	32	66	62	55	57	59	58
Age							
Under 25 years	14	17	12	11	9	8	7
25–29 years	20	26	18	17	15	13	11
30–34 years	38	48	44	32	30	41	28
35–39 years	36	59	52	40	42	34	35
40–44 years	17	49	49	38	43	34	34
45 years and over	21	51	53	40	51	49	58
Region							
Northeast	66	95	86	69	69	70	68
Midwest	20	27	25	*	23	20	20
South	37	87	77	72	71	67	64
West	24	41	39	19	26	23	21
Rate per 100,000 population ²							
All HIV discharges ¹	58.1	93.9	84.7	65.8	69.1	65.3	62.2
Sex							
Male	93.4	141.6	125.7	93.1	98.8	89.5	84.8
Female	24.7	48.4	45.4	39.7	40.7	42.1	40.5
Age							
Under 25 years	14.8	17.6	12.6	11.3	9.1	8.1	7.4
25–29 years	95.1	132.2	91.3	86.7	77.0	70.5	59.0
30–34 years	172.8	216.2	203.6	153.6	145.2	208.3	139.2
35–39 years	179.0	262.7	227.8	176.5	185.4	149.1	154.8
40–44 years	94.8	241.4	233.1	175.2	194.4	152.0	150.7
45 years and over	27.4	60.2	60.6	45.5	55.6	52.9	61.2
Region							
Northeast	128.8	183.3	164.9	132.3	133.4	135.0	129.5
Midwest	33.6	43.5	40.5	*	36.2	30.7	31.1
South	42.1	93.1	82.0	75.1	73.1	68.4	64.9
West	44.1	69.5	65.2	30.9	42.0	37.0	33.4

* Figure does not meet standard of reliability or precision.

¹HIV is human immunodeficiency virus.

²Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 23. Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnoses: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

First-listed diagnosis and ICD-9-CM code	All ages	Under 65 years	65 years and over	All ages	Under 65 years	65 years and over
	Number of deaths in thousands			Rate per 100 discharges		
All conditions ¹	776	189	587	2.4	1.0	4.7
Heart disease 391-392.0,393-398,402,404,410-416,420-429	163	29	134	3.7	1.9	4.7
Acute myocardial infarction 410	77	*	63	9.9	*	12.5
Cardiac dysrhythmias 427	15	*5	11	2.1	*2.2	2.1
Congestive heart failure 428.0	41	*	38	4.1	*	5.0
Malignant neoplasms 140-208,230-234	94	36	58	8.1	6.6	9.4
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	20	7	13	14.1	13.3	14.7
Pneumonia 480-486	63	8	54	4.9	1.6	7.1
Cerebrovascular disease 430-438	63	16	47	6.4	5.8	6.6
Septicemia 038	45	10	36	13.9	9.0	16.4
Fractures, all sites ² 800-829	16	*	14	1.6	*	2.6
Volume depletion 276.5	12	*	11	2.5	*	4.3
Chronic bronchitis 491	10	*	9	2.1	*	2.6

* Figure does not meet standard of reliability or precision.

¹Includes data for deaths not shown in table.

²Excludes fractures coded as 733.1, pathologic fracture.

Table 24. Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Characteristic	All discharges	Discharges without procedures	Discharges with procedures			
			Total	With surgical procedures	Total	With surgical procedures
				Number in thousands		Percent
All patients	31,706	12,455	19,251	14,127	60.7	44.6
Age						
Under 15 years	2,383	1,316	1,067	580	44.8	24.4
15-44 years	9,969	2,894	7,074	5,564	71.0	55.8
45-64 years	6,958	2,620	4,338	3,295	62.3	47.4
65 years and over	12,396	5,624	6,772	4,688	54.6	37.8
65-74 years	4,678	1,812	2,866	2,113	61.3	45.2
75-84 years	5,119	2,373	2,746	1,890	53.6	36.9
85 years and over	2,599	1,439	1,160	684	44.6	26.3
Sex						
Male	12,514	5,302	7,212	5,050	57.6	40.4
Female	19,192	7,152	12,039	9,077	62.7	47.3
Region						
Northeast	7,103	2,593	4,509	3,001	63.5	42.3
Midwest	7,207	3,107	4,100	3,117	56.9	43.2
South	12,016	5,057	6,959	5,288	57.9	44.0
West	5,380	1,697	3,683	2,720	68.5	50.6

NOTE: See appendix for definition of surgical procedures.

Table 25. Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Number in thousands	Rate per 10,000 population ¹
Surgical procedures ²	23,244	835.8
Cardiac catheterization 37.21-37.23	1,221	43.9
Repair of current obstetric laceration ³ 75.5-75.6	1,136	40.9
Removal of coronary artery obstruction and insertion of stent(s) ⁴ 36.0	1,025	36.9
Episiotomy with or without forceps or vacuum extraction ³ 72.1,72.21,72.31,72.71,73.6	944	33.9
Cesarean section ³ 74.0-74.2,74.4,74.99	855	30.8
Artificial rupture of membranes ³ 73.0	833	30.0
Hysterectomy 68.3-68.7,68.9	633	22.8
Reduction of fracture 76.7,79.0-79.3	628	22.6
Open reduction of fracture with internal fixation 79.3	423	15.2
Coronary artery bypass graft ⁴ 36.1	519	18.7
Endoscopy of small intestine with biopsy 45.14,45.16	512	18.4
Oophorectomy and salpingo-oophorectomy 65.3-65.6	494	17.8
Cholecystectomy 51.2	419	15.1
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	327	11.8
Debridement of wound, infection, or burn 86.22,86.28	325	11.7
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	315	11.3
Lysis of peritoneal adhesions 54.5	307	11.0
Appendectomy, excluding incidental 47.0	303	10.9
Total knee replacement 81.54	299	10.8
Excision or destruction of intervertebral disc 80.5	296	10.6
Partial excision of large intestine 45.7	236	8.5
Partial excision of bone 76.2-76.3,77.6-77.8	209	7.5
Nonsurgical procedures ⁵	16,737	601.8
Arteriography and angiocardiography using contrast material 88.4-88.5	2,005	72.1
Respiratory therapy 93.9,96.7	991	35.6
Diagnostic ultrasound 88.7	886	31.9
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	754	27.1
Hemodialysis 39.95	456	16.4
Insertion of endotracheal tube 96.04	429	15.4
Endoscopy of small intestine without biopsy 45.11-45.13	403	14.5
Endoscopy of large intestine without biopsy 45.21-45.24	352	12.7
Spinal tap 03.31	296	10.6

¹Population estimates used to compute rates are based on the 1990 census. See appendix.²Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.³See table 36 for rate per 100 deliveries or per 100 vaginal deliveries.⁴See table 27 for comparison of number of procedures with number of discharges with procedures.⁵Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.

Table 26. Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2000[Discharges of inpatients from non-Federal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
Surgical procedures ¹	7,801	3,665	3,131	1,005
Cardiac catheterization 37.21-37.23	597	321	237	38
Removal of coronary artery obstruction and insertion of stent(s) ² 36.0	524	297	197	30
Endoscopy of small intestine with biopsy 45.14,45.16	304	115	132	57
Coronary artery bypass graft ² 36.1	286	163	114	9
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	274	75	132	67
Reduction of fracture 76.7,79.0-79.3	265	73	103	89
Open reduction of fracture with internal fixation 79.3	203	54	79	71
Total knee replacement 81.54	211	125	76	10
Cholecystectomy 51.2	149	77	54	18
Prostatectomy 60.2-60.6	134	71	48	14
Debridement of wound, infection, or burn 86.22,86.28	120	51	46	23
Nonsurgical procedures ³	6,937	2,772	2,896	1,269
Arteriography and angiocardiography using contrast material 88.4-88.5	998	511	411	77
Diagnostic ultrasound 88.7	474	189	188	98
Respiratory therapy 93.9,96.7	440	174	178	88
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	383	133	171	78
Endoscopy of small intestine without biopsy 45.11-45.13	250	84	109	57
Endoscopy of large intestine without biopsy 45.21-45.24	233	81	105	47
Insertion of endotracheal tube 96.04	226	89	96	40
Hemodialysis 39.95	226	118	91	16
		Rate per 10,000 population ⁴		
Surgical procedures ¹	2,262.7	2,037.2	2,563.1	2,353.3
Cardiac catheterization 37.21-37.23	173.0	178.6	194.2	89.0
Removal of coronary artery obstruction and insertion of stent(s) ² 36.0	152.0	165.1	161.1	70.7
Endoscopy of small intestine with biopsy 45.14,45.16	88.1	63.7	108.2	133.2
Coronary artery bypass graft ² 36.1	82.9	90.7	93.2	20.6
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	79.4	41.7	107.8	156.5
Reduction of fracture 76.7,79.0-79.3	76.8	40.5	84.6	207.6
Open reduction of fracture with internal fixation 79.3	58.9	29.9	64.3	165.6
Total knee replacement 81.54	61.2	69.7	62.0	23.1
Cholecystectomy 51.2	43.3	43.0	44.4	41.3
Prostatectomy 60.2-60.6	38.9	39.7	39.4	33.8
Debridement of wound, infection, or burn 86.22,86.28	34.7	28.3	37.4	53.7
Nonsurgical procedures ³	2,012.0	1,540.4	2,371.3	2,971.9
Arteriography and angiocardiography using contrast material 88.4-88.5	289.5	283.8	336.2	180.0
Diagnostic ultrasound 88.7	137.6	104.8	154.2	228.4
Respiratory therapy 93.9,96.7	127.7	96.8	145.7	206.5
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	111.0	74.1	140.2	183.0
Endoscopy of small intestine without biopsy 45.11-45.13	72.5	46.6	89.1	134.2
Endoscopy of large intestine without biopsy 45.21-45.24	67.5	45.0	85.6	110.5
Insertion of endotracheal tube 96.04	65.5	49.6	78.8	94.3
Hemodialysis 39.95	65.5	65.7	74.9	37.8

¹Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.²See table 27 for a comparison of number of procedures with number of discharges with procedures.³Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.⁴Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 27. Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures by sex, age, and geographic region: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Sex, age, and region	Removal of coronary artery obstruction and insertion of stents(s) ¹		Coronary artery bypass graft ²	
	Number in thousands	Rate per 10,000 population ³	Number in thousands	Rate per 10,000 population ³
Discharges with procedures				
Total	554	19.9	314	11.3
Sex				
Male	357	26.2	219	16.1
Female	198	13.9	94	6.6
Age				
Under 15 years	*	*	*	*
15-44 years	34	2.8	10	0.8
45-64 years	237	38.9	126	20.8
65 years and over	283	82.1	177	51.4
65-74 years	161	89.3	99	55.0
75-84 years	107	87.5	72	58.6
85 years and over	16	36.6	6	15.2
Region				
Northeast	106	20.2	72	13.7
Midwest	157	24.6	78	12.2
South	194	19.6	111	11.2
West	98	15.5	53	8.4
Procedures				
Total	1,025	36.9	519	18.7
Sex				
Male	655	48.2	371	27.3
Female	370	26.0	148	10.4
Age				
Under 15 years	*	*	*	*
15-44 years	62	5.1	17	1.4
45-64 years	437	71.7	216	35.4
65 years and over	524	152.0	286	82.9
65-74 years	297	165.1	163	90.7
75-84 years	197	161.1	114	93.2
85 years and over	30	70.7	9	20.6
Region				
Northeast	194	37.0	123	23.4
Midwest	290	45.4	133	20.9
South	360	36.4	174	17.6
West	181	28.7	89	14.1

* Figure does not meet standard of reliability or precision.

¹ICD-9-CM code 36.0.²ICD-9-CM code 36.1.³Population estimates used to compute rates are based on the 1990 census. See appendix.

NOTE: The number of procedures exceeds the number of discharges because these procedures are often assigned more than one code for a single operation. See Methods section.

Table 28. Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over	
						Number in thousands
All procedures	39,981	1,951	13,518	9,775	14,737	
Operations on the nervous system	01-05	1,000	210	309	238	242
Spinal tap	03.31	296	131	70	47	48
Operations on the endocrine system	06-07	90	*	27	37	23
Operations on the eye	08-16	80	9	20	20	32
Operations on the ear	18-20	37	22	8	*4	*4
Operations on the nose, mouth, and pharynx	21-29	261	56	89	57	59
Operations on the respiratory system	30-34	969	57	164	271	477
Bronchoscopy with or without biopsy	33.21-33.24,33.27	248	*	42	73	116
Operations on the cardiovascular system	35-39	5,939	154	549	2,113	3,123
Removal of coronary artery obstruction and insertion of stent(s) ¹	36.0	1,025	*	62	437	524
Coronary artery bypass graft ¹	36.1	519	*	17	216	286
Cardiac catheterization	37.21-37.23	1,221	8	96	520	597
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7-37.8	327	*	*5	45	274
Hemodialysis	39.95	456	*	65	164	226
Operations on the hemic and lymphatic system	40-41	315	*	49	108	135
Operations on the digestive system	42-54	5,145	208	1,136	1,404	2,397
Endoscopy of small intestine with or without biopsy	45.11-45.14,45.16	915	11	120	231	553
Endoscopy of large intestine with or without biopsy	45.21-45.25	541	*	65	125	346
Partial excision of large intestine	45.7	236	*	27	73	134
Appendectomy, excluding incidental	47.0	303	71	167	50	16
Cholecystectomy	51.2	419	*	138	129	149
Lysis of peritoneal adhesions	54.5	307	4	123	94	85
Operations on the urinary system	55-59	962	35	235	291	401
Cystoscopy with or without biopsy	57.31-57.33	181	*	33	47	97
Operations on the male genital organs	60-64	258	19	14	68	157
Prostatectomy	60.2-60.6	184	*	*	48	134
Operations on the female genital organs	65-71	2,061	6	1,192	638	225
Oophorectomy and salpingo-oophorectomy	65.3-65.6	494	*	205	230	57
Bilateral destruction or occlusion of fallopian tubes	66.2-66.3	315	*	315	*	*
Hysterectomy	68.3-68.7,68.9	633	*	317	254	62
Obstetrical procedures	72-75	6,209	19	6,185	*5	...
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	944	*	940	*	...
Artificial rupture of membranes	73.0	833	*	830	*	...
Cesarean section	74.0-74.2,74.4,74.99	855	*	853	*	...
Repair of current obstetric laceration	75.5-75.6	1,136	*4	1,130	*	...
Operations on the musculoskeletal system	76-84	3,171	172	836	918	1,245
Partial excision of bone	76.2-76.3,77.6-77.8	209	11	70	77	51
Reduction of fracture	76.7,79.0-79.3	628	51	187	125	265
Open reduction of fracture with internal fixation	79.3	423	15	117	87	203
Excision or destruction of intervertebral disc	80.5	296	*	126	128	41
Total hip replacement	81.51	152	*	13	51	88
Total knee replacement	81.54	299	*	*6	82	211
Operations on the integumentary system	85-86	1,264	*	355	378	407
Debridement of wound, infection, or burn	86.22,86.28	325	17	84	105	120
Miscellaneous diagnostic and therapeutic procedures	87-99	12,223	837	2,349	3,226	5,812
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	754	39	150	182	383
Arteriography and angiocardiology using contrast material	88.4-88.5	2,005	11	175	821	998
Diagnostic ultrasound	88.7	886	37	143	232	474
Respiratory therapy	93.9,96.7	991	206	129	215	440
Insertion of endotracheal tube	96.04	429	40	61	103	226
Injection or infusion of cancer chemotherapeutic substance	99.25	199	48	37	68	46

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 29. Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All procedures	1,437.7	322.5	1,106.3	1,603.9	4,274.7
Operations on the nervous system 01-05	35.9	34.8	25.3	39.1	70.3
Spinal tap 03.31	10.6	21.6	5.8	7.8	13.8
Operations on the endocrine system 06-07	3.2	*	2.2	6.1	6.5
Operations on the eye 08-16	2.9	1.5	1.6	3.2	9.3
Operations on the ear 18-20	1.3	3.6	0.6	*0.6	*1.1
Operations on the nose, mouth, and pharynx 21-29	9.4	9.2	7.3	9.4	17.0
Operations on the respiratory system 30-34	34.8	9.4	13.4	44.4	138.3
Bronchoscopy with or without biopsy 33.21-33.24,33.27	8.9	*	3.5	12.1	33.7
Operations on the cardiovascular system 35-39	213.5	25.4	44.9	346.8	905.9
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	36.9	*	5.1	71.7	152.0
Coronary artery bypass graft ¹ 36.1	18.7	*	1.4	35.4	82.9
Cardiac catheterization 37.21-37.23	43.9	1.4	7.8	85.4	173.0
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	11.8	*	*0.4	7.4	79.4
Hemodialysis 39.95	16.4	*	5.3	26.9	65.5
Operations on the hemic and lymphatic system 40-41	11.3	*	4.0	17.7	39.1
Operations on the digestive system 42-54	185.0	34.4	93.0	230.3	695.2
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	32.9	1.8	9.8	37.9	160.5
Endoscopy of large intestine with or without biopsy 45.21-45.25	19.4	*	5.3	20.5	100.5
Partial excision of large intestine 45.7	8.5	*	2.2	11.9	38.8
Appendectomy, excluding incidental 47.0	10.9	11.7	13.6	8.2	4.7
Cholecystectomy 51.2	15.1	*	11.3	21.2	43.3
Lysis of peritoneal adhesions 54.5	11.0	0.7	10.1	15.5	24.8
Operations on the urinary system 55-59	34.6	5.8	19.3	47.7	116.3
Cystoscopy with or without biopsy 57.31-57.33	6.5	*	2.7	7.8	28.0
Operations on the male genital organs 60-64	9.3	3.2	1.2	11.1	45.5
Prostatectomy 60.2-60.6	6.6	*	*	7.9	38.9
Operations on the female genital organs 65-71	74.1	0.9	97.5	104.7	65.3
Oophorectomy and salpingo-oophorectomy 65.3-65.6	17.8	*	16.8	37.8	16.6
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	11.3	*	25.8	*	*
Hysterectomy 68.3-68.7,68.9	22.8	*	26.0	41.7	17.9
Obstetrical procedures 72-75	223.3	3.1	506.2	*0.8	...
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	33.9	*	76.9	*	...
Artificial rupture of membranes 73.0	30.0	*	67.9	*	...
Cesarean section 74.0-74.2,74.4,74.99	30.8	*	69.8	*	...
Repair of current obstetric laceration 75.5-75.6	40.9	*0.7	92.5	*	...
Operations on the musculoskeletal system 76-84	114.0	28.4	68.4	150.6	361.1
Partial excision of bone 76.2-76.3,77.6-77.8	7.5	1.9	5.7	12.7	14.7
Reduction of fracture 76.7,79.0-79.3	22.6	8.5	15.3	20.5	76.8
Open reduction of fracture with internal fixation 79.3	15.2	2.5	9.5	14.3	58.9
Excision or destruction of intervertebral disc 80.5	10.6	*	10.3	21.0	12.0
Total hip replacement 81.51	5.5	*	1.0	8.3	25.7
Total knee replacement 81.54	10.8	*	*0.5	13.5	61.2
Operations on the integumentary system 85-86	45.4	*	29.1	62.0	118.0
Debridement of wound, infection, or burn 86.22,86.28	11.7	2.9	6.8	17.2	34.7
Miscellaneous diagnostic and therapeutic procedures 87-99	439.5	138.3	192.3	529.3	1,685.7
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	27.1	6.5	12.3	29.8	111.0
Arteriography and angiocardiology using contrast material 88.4-88.5	72.1	1.8	14.3	134.6	289.5
Diagnostic ultrasound 88.7	31.9	6.1	11.7	38.1	137.6
Respiratory therapy 93.9,96.7	35.6	34.1	10.6	35.2	127.7
Insertion of endotracheal tube 96.04	15.4	6.5	5.0	16.9	65.5
Injection or infusion of cancer chemotherapeutic substance 99.25	7.2	8.0	3.0	11.2	13.3

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 30. Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
All procedures	39,981	15,654	24,328
Operations on the nervous system 01-05	1,000	458	542
Spinal tap 03.31	296	156	141
Operations on the endocrine system 06-07	90	23	67
Operations on the eye 08-16	80	43	37
Operations on the ear 18-20	37	24	13
Operations on the nose, mouth, and pharynx 21-29	261	158	103
Operations on the respiratory system 30-34	969	526	443
Bronchoscopy with or without biopsy 33.21-33.24,33.27	248	145	103
Operations on the cardiovascular system 35-39	5,939	3,420	2,519
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	1,025	655	370
Coronary artery bypass graft ¹ 36.1	519	371	148
Cardiac catheterization 37.21-37.23	1,221	732	490
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	327	162	165
Hemodialysis 39.95	456	224	232
Operations on the hemic and lymphatic system 40-41	315	160	155
Operations on the digestive system 42-54	5,145	2,180	2,965
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	915	408	508
Endoscopy of large intestine with or without biopsy 45.21-45.25	541	210	331
Partial excision of large intestine 45.7	236	107	129
Appendectomy, excluding incidental 47.0	303	163	140
Cholecystectomy 51.2	419	134	284
Lysis of peritoneal adhesions 54.5	307	61	246
Operations on the urinary system 55-59	962	450	511
Cystoscopy with or without biopsy 57.31-57.33	181	96	85
Operations on the male genital organs 60-64	258	258	...
Prostatectomy 60.2-60.6	184	184	...
Operations on the female genital organs 65-71	2,061	...	2,061
Oophorectomy and salpingo-oophorectomy 65.3-65.6	494	...	494
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	315	...	315
Hysterectomy 68.3-68.7,68.9	633	...	633
Obstetrical procedures 72-75	6,209	...	6,209
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	944	...	944
Artificial rupture of membranes 73.0	833	...	833
Cesarean section 74.0-74.2,74.4,74.99	855	...	855
Repair of current obstetric laceration 75.5-75.6	1,136	...	1,136
Operations on the musculoskeletal system 76-84	3,171	1,546	1,624
Partial excision of bone 76.2-76.3,77.6-77.8	209	111	98
Reduction of fracture 76.7,79.0-79.3	628	285	344
Open reduction of fracture with internal fixation 79.3	423	177	246
Excision or destruction of intervertebral disc 80.5	296	162	134
Total hip replacement 81.51	152	69	83
Total knee replacement 81.54	299	115	185
Operations on the integumentary system 85-86	1,264	567	697
Debridement of wound, infection, or burn 86.22,86.28	325	189	136
Miscellaneous diagnostic and therapeutic procedures 87-99	12,223	5,841	6,382
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	754	345	409
Arteriography and angiocardiology using contrast material 88.4-88.5	2,005	1,157	848
Diagnostic ultrasound 88.7	886	385	501
Respiratory therapy 93.9,96.7	991	507	484
Insertion of endotracheal tube 96.04	429	221	209
Injection or infusion of cancer chemotherapeutic substance 99.25	199	115	85

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 31. Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Rate per 10,000 population	
		Male	Female
All procedures	1,437.7	1,151.1	1,711.8
Operations on the nervous system	35.9	33.7	38.1
Spinal tap	10.6	11.4	9.9
Operations on the endocrine system	3.2	1.7	4.7
Operations on the eye	2.9	3.2	2.6
Operations on the ear	1.3	1.7	0.9
Operations on the nose, mouth, and pharynx	9.4	11.6	7.2
Operations on the respiratory system	34.8	38.7	31.2
Bronchoscopy with or without biopsy	8.9	10.6	7.3
Operations on the cardiovascular system	213.5	251.5	177.2
Removal of coronary artery obstruction and insertion of stent(s) ¹	36.0	48.2	26.0
Coronary artery bypass graft ¹	18.7	27.3	10.4
Cardiac catheterization	43.9	53.8	34.4
Insertion, replacement, removal, and revision of pacemaker leads or device	11.8	11.9	11.6
Hemodialysis	16.4	16.4	16.3
Operations on the hemic and lymphatic system	11.3	11.7	10.9
Operations on the digestive system	185.0	160.3	208.6
Endoscopy of small intestine with or without biopsy	32.9	30.0	35.7
Endoscopy of large intestine with or without biopsy	19.4	15.5	23.3
Partial excision of large intestine	8.5	7.9	9.1
Appendectomy, excluding incidental	10.9	12.0	9.8
Cholecystectomy	15.1	9.9	20.0
Lysis of peritoneal adhesions	11.0	4.5	17.3
Operations on the urinary system	34.6	33.1	36.0
Cystoscopy with or without biopsy	6.5	7.1	6.0
Operations on the male genital organs	9.3	18.9	...
Prostatectomy	6.6	13.5	...
Operations on the female genital organs	74.1	...	145.0
Oophorectomy and salpingo-oophorectomy	17.8	...	34.7
Bilateral destruction or occlusion of fallopian tubes	11.3	...	22.2
Hysterectomy	22.8	...	44.6
Obstetrical procedures	223.3	...	436.9
Episiotomy with or without forceps or vacuum extraction	33.9	...	66.4
Artificial rupture of membranes	30.0	...	58.6
Cesarean section	30.8	...	60.2
Repair of current obstetric laceration	40.9	...	79.9
Operations on the musculoskeletal system	114.0	113.7	114.3
Partial excision of bone	7.5	8.2	6.9
Reduction of fracture	22.6	20.9	24.2
Open reduction of fracture with internal fixation	15.2	13.0	17.3
Excision or destruction of intervertebral disc	10.6	11.9	9.4
Total hip replacement	5.5	5.1	5.8
Total knee replacement	10.8	8.4	13.0
Operations on the integumentary system	45.4	41.7	49.0
Debridement of wound, infection, or burn	11.7	13.9	9.6
Miscellaneous diagnostic and therapeutic procedures	439.5	429.6	449.1
Computerized axial tomography	27.1	25.3	28.8
Arteriography and angiocardiology using contrast material	72.1	85.1	59.6
Diagnostic ultrasound	31.9	28.3	35.3
Respiratory therapy	35.6	37.3	34.0
Insertion of endotracheal tube	15.4	16.2	14.7
Injection or infusion of cancer chemotherapeutic substance	7.2	8.4	6.0

... Category not applicable.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 32. Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures	39,981	9,689	8,756	13,852	7,683
Operations on the nervous system 01-05	1,000	206	273	297	223
Spinal tap 03.31	296	60	62	103	72
Operations on the endocrine system 06-07	90	22	14	37	18
Operations on the eye 08-16	80	19	16	34	11
Operations on the ear 18-20	37	9	6	13	*8
Operations on the nose, mouth, and pharynx 21-29	261	69	56	88	48
Operations on the respiratory system 30-34	969	249	198	358	164
Bronchoscopy with or without biopsy 33.21-33.24,33.27	248	68	45	96	38
Operations on the cardiovascular system 35-39	5,939	1,315	1,412	2,167	1,045
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	1,025	194	290	360	181
Coronary artery bypass graft ¹ 36.1	519	123	133	174	89
Cardiac catheterization 37.21-37.23	1,221	244	302	489	186
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	327	98	73	113	43
Hemodialysis 39.95	456	97	83	179	97
Operations on the hemic and lymphatic system 40-41	315	77	66	108	64
Operations on the digestive system 42-54	5,145	1,212	1,158	1,814	961
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	915	212	220	336	147
Endoscopy of large intestine with or without biopsy 45.21-45.25	541	130	124	197	90
Partial excision of large intestine 45.7	236	57	61	74	45
Appendectomy, excluding incidental 47.0	303	61	71	94	76
Cholecystectomy 51.2	419	86	83	169	81
Lysis of peritoneal adhesions 54.5	307	71	63	123	51
Operations on the urinary system 55-59	962	244	202	375	141
Cystoscopy with or without biopsy 57.31-57.33	181	49	45	63	24
Operations on the male genital organs 60-64	258	57	70	85	46
Prostatectomy 60.2-60.6	184	39	52	59	35
Operations on the female genital organs 65-71	2,061	376	441	846	397
Oophorectomy and salpingo-oophorectomy 65.3-65.6	494	87	111	206	89
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	315	45	57	150	63
Hysterectomy 68.3-68.7,68.9	633	112	150	255	117
Obstetrical procedures 72-75	6,209	1,173	1,401	2,155	1,479
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	944	183	201	371	190
Artificial rupture of membranes 73.0	833	122	246	234	232
Cesarean section 74.0-74.2,74.4,74.99	855	155	156	358	186
Repair of current obstetric laceration 75.5-75.6	1,136	208	263	366	299
Operations on the musculoskeletal system 76-84	3,171	637	728	1,178	628
Partial excision of bone 76.2-76.3,77.6-77.8	209	38	46	76	49
Reduction of fracture 76.7,79.0-79.3	628	133	141	234	120
Open reduction of fracture with internal fixation 79.3	423	85	97	167	74
Excision or destruction of intervertebral disc 80.5	296	52	65	125	54
Total hip replacement 81.51	152	32	39	53	28
Total knee replacement 81.54	299	52	87	109	52
Operations on the integumentary system 85-86	1,264	312	274	506	172
Debridement of wound, infection, or burn 86.22,86.28	325	74	86	125	40
Miscellaneous diagnostic and therapeutic procedures 87-99	12,223	3,712	2,442	3,791	2,278
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	754	338	*	149	157
Arteriography and angiocardiology using contrast material 88.4-88.5	2,005	409	499	825	271
Diagnostic ultrasound 88.7	886	316	146	232	193
Respiratory therapy 93.9,96.7	991	294	166	245	286
Insertion of endotracheal tube 96.04	429	114	83	149	83
Injection or infusion of cancer chemotherapeutic substance 99.25	199	59	28	77	35

* Figure does not meet standard of reliability or precision.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

Table 33. Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures	1,437.7	1,848.4	1,370.8	1,402.3	1,219.2
Operations on the nervous system 01-05	35.9	39.3	42.8	30.0	35.5
Spinal tap 03.31	10.6	11.4	9.8	10.4	11.4
Operations on the endocrine system 06-07	3.2	4.2	2.1	3.7	2.8
Operations on the eye 08-16	2.9	3.7	2.4	3.5	1.8
Operations on the ear 18-20	1.3	1.7	0.9	1.3	*1.3
Operations on the nose, mouth, and pharynx 21-29	9.4	13.2	8.7	8.9	7.6
Operations on the respiratory system 30-34	34.8	47.4	31.0	36.2	26.0
Bronchoscopy with or without biopsy 33.21-33.24,33.27	8.9	12.9	7.1	9.7	6.1
Operations on the cardiovascular system 35-39	213.5	250.9	221.0	219.4	165.8
Removal of coronary artery obstruction and insertion of stent(s) ¹ 36.0	36.9	37.0	45.4	36.4	28.7
Coronary artery bypass graft ¹ 36.1	18.7	23.4	20.9	17.6	14.1
Cardiac catheterization 37.21-37.23	43.9	46.5	47.3	49.5	29.5
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	11.8	18.8	11.4	11.5	6.8
Hemodialysis 39.95	16.4	18.5	13.0	18.1	15.3
Operations on the hemic and lymphatic system 40-41	11.3	14.6	10.3	11.0	10.1
Operations on the digestive system 42-54	185.0	231.1	181.3	183.6	152.5
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	32.9	40.4	34.4	34.0	23.4
Endoscopy of large intestine with or without biopsy 45.21-45.25	19.4	24.8	19.5	19.9	14.2
Partial excision of large intestine 45.7	8.5	10.8	9.5	7.5	7.1
Appendectomy, excluding incidental 47.0	10.9	11.7	11.2	9.6	12.1
Cholecystectomy 51.2	15.1	16.4	13.0	17.1	12.8
Lysis of peritoneal adhesions 54.5	11.0	13.5	9.8	12.4	8.1
Operations on the urinary system 55-59	34.6	46.5	31.6	38.0	22.3
Cystoscopy with or without biopsy 57.31-57.33	6.5	9.3	7.0	6.3	3.9
Operations on the male genital organs 60-64	9.3	10.8	11.0	8.6	7.3
Prostatectomy 60.2-60.6	6.6	7.5	8.1	5.9	5.5
Operations on the female genital organs 65-71	74.1	71.8	69.0	85.7	63.1
Oophorectomy and salpingo-oophorectomy 65.3-65.6	17.8	16.6	17.4	20.9	14.1
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	11.3	8.6	8.9	15.2	10.1
Hysterectomy 68.3-68.7,68.9	22.8	21.4	23.4	25.8	18.6
Obstetrical procedures 72-75	223.3	223.9	219.4	218.2	234.7
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	33.9	34.9	31.4	37.5	30.1
Artificial rupture of membranes 73.0	30.0	23.2	38.5	23.7	36.7
Cesarean section 74.0-74.2,74.4,74.99	30.8	29.5	24.5	36.2	29.5
Repair of current obstetric laceration 75.5-75.6	40.9	39.7	41.1	37.1	47.4
Operations on the musculoskeletal system 76-84	114.0	121.6	114.0	119.2	99.6
Partial excision of bone 76.2-76.3,77.6-77.8	7.5	7.2	7.2	7.7	7.8
Reduction of fracture 76.7,79.0-79.3	22.6	25.3	22.1	23.7	19.1
Open reduction of fracture with internal fixation 79.3	15.2	16.2	15.1	16.9	11.8
Excision or destruction of intervertebral disc 80.5	10.6	9.9	10.1	12.7	8.6
Total hip replacement 81.51	5.5	6.1	6.1	5.3	4.4
Total knee replacement 81.54	10.8	9.9	13.6	11.0	8.2
Operations on the integumentary system 85-86	45.4	59.5	42.9	51.2	27.3
Debridement of wound, infection, or burn 86.22,86.28	11.7	14.1	13.4	12.7	6.4
Miscellaneous diagnostic and therapeutic procedures 87-99	439.5	708.2	382.3	383.8	361.5
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	27.1	64.4	*	15.0	24.9
Arteriography and angiocardiology using contrast material 88.4-88.5	72.1	78.0	78.2	83.5	43.0
Diagnostic ultrasound 88.7	31.9	60.2	22.8	23.5	30.6
Respiratory therapy 93.9,96.7	35.6	56.0	26.0	24.8	45.4
Insertion of endotracheal tube 96.04	15.4	21.7	13.0	15.1	13.2
Injection or infusion of cancer chemotherapeutic substance 99.25	7.2	11.3	4.4	7.8	5.6

* Figure does not meet standard of reliability or precision.

¹See table 27 for comparison of number of procedures with number of discharges with procedures.

NOTE: Population estimates used to compute rates are based on the 1990 census. See appendix.

Table 34. Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2000[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	
All deliveries							
10–54 years	3,738	100.0	41.5	9,271	100.0	103.0	2.5
10–19 years	453	12.1	22.8	1,094	11.8	55.1	2.4
20–24 years	954	25.5	102.5	2,303	24.8	247.6	2.4
25–29 years	984	26.3	106.7	2,378	25.6	257.8	2.4
30–34 years	851	22.8	85.4	2,150	23.2	215.7	2.5
35–39 years	415	11.1	36.9	1,128	12.2	100.1	2.7
40–54 years	80	2.1	2.6	217	2.3	7.2	2.7
All regions	3,738	100.0	41.5	9,271	100.0	103.0	2.5
Northeast	632	16.9	37.4	1,754	18.9	103.6	2.8
Midwest	771	20.6	37.3	1,853	20.0	89.6	2.4
South	1,439	38.5	44.8	3,571	38.5	111.2	2.5
West	895	23.9	44.0	2,092	22.6	103.0	2.3
Vaginal deliveries²							
10–54 years	2,882	100.0	32.0	6,113	100.0	67.9	2.1
10–19 years	379	13.1	19.1	838	13.7	42.2	2.2
20–24 years	768	26.6	82.5	1,639	26.8	176.2	2.1
25–29 years	762	26.4	82.6	1,578	25.8	171.1	2.1
30–34 years	636	22.1	63.8	1,332	21.8	133.6	2.1
35–39 years	285	9.9	25.3	610	10.0	54.2	2.1
40–54 years	53	1.8	1.7	116	1.9	3.8	2.2
All regions	2,882	100.0	32.0	6,113	100.0	67.9	2.1
Northeast	478	16.6	28.2	1,110	18.2	65.6	2.3
Midwest	615	21.3	29.8	1,294	21.2	62.6	2.1
South	1,081	37.5	33.7	2,295	37.6	71.5	2.1
West	709	24.6	34.9	1,414	23.1	69.6	2.0
Cesarean deliveries³							
10–54 years	855	100.0	9.5	3,158	100.0	35.1	3.7
10–19 years	75	8.7	3.8	256	8.1	12.9	3.4
20–24 years	186	21.7	20.0	665	21.0	71.5	3.6
25–29 years	222	26.0	24.1	800	25.3	86.7	3.6
30–34 years	215	25.2	21.6	818	25.9	82.0	3.8
35–39 years	130	15.2	11.5	518	16.4	46.0	4.0
40–54 years	27	3.2	0.9	102	3.2	3.3	3.8
All regions	855	100.0	9.5	3,158	100.0	35.1	3.7
Northeast	155	18.1	9.1	644	20.4	38.1	4.2
Midwest	156	18.3	7.6	559	17.7	27.0	3.6
South	358	41.9	11.1	1,276	40.4	39.7	3.6
West	186	21.8	9.2	679	21.5	33.4	3.6

¹ Population estimates used to compute rates are based on the 1990 census and are for females 10–54 years of age. See appendix.² Discharges of females with deliveries except those with cesarean deliveries.³ Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 35. Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–2000[Discharges of inpatients from non-Federal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)* code V27, females with deliveries]

Length of stay and type of delivery	1980	1985	1990	1995	2000
Length of stay					
Number in thousands					
All deliveries	3,762	3,854	4,025	3,766	3,738
Less than 1 day	23	34	37	36	16
1 day	216	373	623	1,368	551
2 days	747	1,037	1,649	1,418	1,863
3 days	1,205	1,140	900	606	863
4 days	691	586	468	193	292
5–7 days	726	598	282	100	112
8 days or more	154	86	67	46	41
Vaginal deliveries ¹	3,143	2,977	3,080	2,981	2,882
Less than 1 day	23	33	35	34	14
1 day	215	371	620	1,362	546
2 days	743	1,030	1,624	1,271	1,726
3 days	1,185	1,054	618	228	500
4 days	618	338	106	39	53
5–7 days	317	125	55	27	25
8 days or more	42	26	23	21	17
Cesarean deliveries ²	619	877	945	785	855
Less than 1 day	*	*	*	*	*
1 day	*	*	*	*6	*
2 days	*	*	25	147	137
3 days	20	86	283	379	363
4 days	73	249	361	154	239
5–7 days	409	473	227	73	87
8 days or more	112	60	44	25	24
Percent distribution					
All deliveries	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.6	0.9	0.9	1.0	0.4
1 day	5.7	9.7	15.5	36.3	14.7
2 days	19.9	26.9	41.0	37.7	49.9
3 days	32.0	29.6	22.4	16.1	23.1
4 days	18.4	15.2	11.6	5.1	7.8
5–7 days	19.3	15.5	7.0	2.7	3.0
8 days or more	4.1	2.2	1.7	1.2	1.1
Vaginal deliveries ¹	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.7	1.1	1.1	1.1	0.5
1 day	6.8	12.5	20.1	45.7	19.0
2 days	23.6	34.6	52.7	42.6	59.9
3 days	37.7	35.4	20.1	7.6	17.3
4 days	19.7	11.4	3.4	1.3	1.8
5–7 days	10.1	4.2	1.8	0.9	0.9
8 days or more	1.3	0.9	0.7	0.7	0.6
Cesarean deliveries ²	100.0	100.0	100.0	100.0	100.0
Less than 1 day	*	*	*	*	*
1 day	*	*	*	*0.8	*
2 days	*	*	2.6	18.7	16.0
3 days	3.2	9.8	29.9	48.3	42.4
4 days	11.8	28.4	38.2	19.6	27.9
5–7 days	66.1	53.9	24.0	9.3	10.2
8 days or more	18.1	6.8	4.7	3.2	2.8
Average length of stay in days					
All deliveries	3.8	3.3	2.8	2.1	2.5
Vaginal deliveries ¹	3.2	2.7	2.3	1.7	2.1
Cesarean deliveries ²	6.5	5.2	4.5	3.6	3.7

* Figure does not meet standard of reliability or precision.

¹Discharges of females with deliveries except those with cesarean deliveries.²Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 36. Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2000[Discharges from non-Federal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Data are for selected procedures performed on discharges with code V27, females with deliveries]

Procedure category and ICD–9–CM code	1980	1985	1990	1995	2000
Number of procedures in thousands					
All obstetrical procedures 72–75	3,526	4,239	6,660	6,282	6,145
Repair of current obstetric laceration 75.5–75.6	350	543	789	961	1,134
Episiotomy 72.1,72.21,72.31,72.71,73.6	2,012	1,820	1,714	1,410	944
Manually assisted delivery 73.5	70	193	750	866	898
Cesarean section 74.0–74.2,74.4,74.99	619	877	945	785	855
Artificial rupture of membranes 73.0	120	315	691	752	833
Medical induction of labor 73.4	42	66	225	366	435
Vacuum extraction ¹ 72.7	22	66	190	279	245
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	555	376	269	176	117
Rate per 100 deliveries ²					
All obstetrical procedures 72–75	93.7	110.0	165.5	166.8	164.4
Cesarean section 74.0–74.2,74.4,74.99	16.5	22.7	23.5	20.8	22.9
Artificial rupture of membranes 73.0	3.2	8.2	17.2	20.0	22.3
Medical induction of labor 73.4	1.1	1.7	5.6	9.7	11.7
Rate per 100 vaginal deliveries ³					
Repair of current obstetric laceration 75.5–75.6	11.1	18.1	25.5	32.0	39.2
Episiotomy 72.1,72.21,72.31,72.71,73.6	64.0	61.1	55.6	47.2	32.7
Manually assisted delivery 73.5	2.2	6.5	24.3	29.0	31.1
Vacuum extraction ¹ 72.7	0.7	2.2	6.1	9.2	8.4
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	17.6	12.5	8.6	5.8	4.0

¹Forceps delivery and vacuum extraction with episiotomy are also included in the episiotomy category.²Procedures per 100 discharges of females with deliveries.³Procedures for females with vaginal deliveries per 100 discharges with vaginal deliveries. Vaginal deliveries are all deliveries except those with a cesarean section.**Table 37. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–2000**[Discharges of inpatients from non-Federal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Type of delivery	1980	1985	1990	1995	2000
Number in thousands					
Females with deliveries	3,762	3,854	4,025	3,766	3,738
Previous cesarean delivery	192	324	426	413	447
No previous cesarean delivery	3,571	3,530	3,600	3,353	3,290
Vaginal deliveries	3,143	2,977	3,080	2,981	2,882
Vaginal birth after cesarean delivery (VBAC)	*6	21	87	147	123
Other vaginal deliveries	3,137	2,956	2,993	2,835	2,759
Cesarean deliveries	619	877	945	785	855
No previous cesarean delivery (primary)	434	574	606	519	531
Previous cesarean delivery (repeat)	185	303	339	266	324
Rate per 100 deliveries					
Cesarean deliveries per 100 deliveries	16.5	22.7	23.5	20.8	22.9
Primary cesarean deliveries per 100 deliveries without previous cesarean delivery	12.1	16.3	16.8	15.5	16.1
Repeat cesarean deliveries per 100 cesarean deliveries	29.9	34.6	35.9	33.9	37.9
Vaginal birth after cesarean deliveries (VBAC) per 100 deliveries with previous cesarean delivery	*3.4	6.6	20.4	35.5	27.6

* Figure does not meet standard of reliability or precision.

NOTE: Previous cesarean deliveries are discharges with ICD–9–CM diagnostic code 654.2. Vaginal deliveries are all females with deliveries except those with cesarean deliveries. Cesarean deliveries are discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 38. Number and rate of females with cesarean deliveries, by type of delivery, and of females with vaginal birth after cesarean delivery, discharged from short-stay hospitals, by age and geographic region: United States, 2000[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Cesarean deliveries			VBAC ^{4,5}
	Total ¹	Primary ²	Repeat ³	
Number in thousands				
10–54 years	855	531	324	123
10–19 years	75	67	8	*
20–24 years	186	121	65	22
25–29 years	222	136	87	35
30–34 years	215	125	91	38
35–39 years	130	67	63	20
40–54 years	27	15	12	*4
All regions	855	531	324	123
Northeast	155	98	57	22
Midwest	156	95	61	28
South	358	228	130	43
West	186	110	76	31
Rate per 100 deliveries				
10–54 years	22.9	16.1	37.9	27.6
10–19 years	16.5	15.2	10.1	*
20–24 years	19.5	14.0	34.8	25.8
25–29 years	22.6	15.7	39.0	28.6
30–34 years	25.3	17.3	42.1	29.5
35–39 years	31.3	20.3	48.1	24.2
40–54 years	33.8	23.5	44.5	*25.8
All regions	22.9	16.1	37.9	27.6
Northeast	24.5	17.7	36.6	28.2
Midwest	20.3	13.9	39.3	31.0
South	24.9	18.0	36.2	24.8
West	20.8	13.9	41.0	28.7

* Figure does not meet standard of reliability or precision.

¹Discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99. Rate per 100 total deliveries.²Cesarean deliveries without ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries without previous cesarean delivery.³Cesarean deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 cesarean deliveries.⁴VBAC is vaginal birth after cesarean delivery.⁵Vaginal deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries with previous cesarean delivery.

Table 39. Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2000

[Infants born in non-Federal hospitals]

Sex, region, and health status	Discharges		Average length of stay in days
	Number in thousands	Percent distribution	
All newborn infants	3,643	100.0	3.2
Sex			
Male	1,869	51.3	3.2
Female	1,773	48.7	3.2
Region			
Northeast	755	20.7	3.4
Midwest	792	21.8	2.9
South	1,344	36.9	3.3
West	751	20.6	3.2
Health status			
Well ¹	2,256	61.9	2.1
Male	1,112	30.5	2.1
Female	1,144	31.4	2.0
Sick ²	1,386	38.1	5.1
Male	758	20.8	4.9
Female	629	17.3	5.4

¹ Without any illness or risk-related diagnoses.

²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

Table 40. Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay by health status: United States, selected years 1980-2000

[Infants born in non-Federal hospitals]

Length of stay and health status	1980	1985	1990	1995	2000
Length of stay					
All newborn infants	3,824	3,794	3,869	3,631	3,643
Less than 1 day	69	84	79	87	45
1 day	270	428	693	1,523	722
2 days	842	1,057	1,624	1,145	1,877
3 days	1,175	1,034	636	486	546
4 days	618	528	447	134	205
5-7 days	633	497	232	106	94
8 days or more	218	167	157	149	153
Percent distribution					
All newborn infants	100.0	100.0	100.0	100.0	100.0
Less than 1 day	1.8	2.2	2.0	2.4	1.2
1 day	7.1	11.3	17.9	41.9	19.8
2 days	22.0	27.9	42.0	31.5	51.5
3 days	30.7	27.3	16.4	13.4	15.0
4 days	16.2	13.9	11.6	3.7	5.6
5-7 days	16.5	13.1	6.0	2.9	2.6
8 days or more	5.7	4.4	4.1	4.1	4.2
Health status					
All newborn infants	4.3	3.8	3.3	2.8	3.2
Well ¹	3.2	2.8	2.3	1.7	2.1
Sick ²	7.1	5.7	4.8	4.4	5.1

¹Without any illness or risk-related diagnoses.

²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

Table 41. Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2000[Infants born in non-Federal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Diagnostic category and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
Sick newborn diagnoses 001-999	2,626	1,454	1,173
Hemolytic disease of fetus or newborn, due to isoimmunization, and other perinatal jaundice 773-774	479	246	233
Neonatal jaundice associated with preterm delivery 774.2	102	52	51
Unspecified fetal and neonatal jaundice 774.6	315	164	152
Respiratory distress syndrome and other respiratory conditions of fetus and newborn 769-770	372	216	156
Transitory tachypnea of newborn 770.6	107	66	41
Disorders relating to short gestation and unspecified low birthweight (prematurity) 765	271	140	130
Congenital anomalies 740-759	246	140	107
Disorders relating to long gestation and high birthweight 766	204	127	77
Endocrine and metabolic disturbances specific to the fetus and newborn 775	123	72	51
Birth trauma 767	104	55	49
Injuries to scalp 767.1	61	34	27
Infections specific to the perinatal period 771	95	48	47

Table 42. Number and percent of male newborn infants circumcised during hospitalization by geographic region: United States, selected years 1980-2000[Infants born in non-Federal hospitals with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) code 64.0]

	1980	1985	1990	1995	2000
	Number circumcised in thousands				
United States	1,261	1,162	1,169	1,200	1,166
Northeast	253	221	238	249	256
Midwest	403	331	341	317	330
South	379	393	396	454	438
West	226	218	195	180	143
	Percent circumcised				
United States	64.7	59.5	59.0	64.1	62.4
Northeast	67.4	65.2	62.6	68.3	64.6
Midwest	75.9	70.5	76.0	79.8	81.4
South	56.0	56.0	57.1	66.1	63.9
West	61.8	49.0	42.4	42.6	37.3

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	31,706	12,514	19,192	2,383	9,969	6,958	12,396	7,103	7,207	12,016	5,380
001-139	787	389	398	160	173	150	305	189	169	288	140
005	*5	*	*	*	*	*	*	*	*	*	*
008	119	49	70	51	18	17	32	21	25	48	25
008.4	31	12	18	*	*5	*5	18	*9	*8	*9	*
008.45	25	*9	17	*	*	*	16	*8	*6	*8	*
008.6	29	16	13	29	*	*	*	*	*6	10	11
008.61	28	16	13	28	*	-	*	*	*6	10	10
008.8	58	20	38	21	12	12	13	10	11	28	9
009	12	*5	*7	*	*	*	*	*	*	*	*
009.0	*5	*	*	*	*	*	*	*	*	*	*
011	*6	*	*	*	*	*	*	*	*	*	*
034	*6	*	*	*	*	*	*	*	*	*	*
034.0	*6	*	*	*	*	*	*	*	*	*	*
038	326	154	172	16	32	62	216	87	76	114	49
038.0	17	10	*7	*	*	*	12	*	*5	*	*
038.1	56	34	22	*	*7	13	33	12	11	23	10
038.10	*8	*6	*	*	*	*	*5	*	*	*	*
038.11	32	19	13	*	*	10	17	*6	*6	13	*7
038.19	15	10	*6	*	*	*	11	*	*	*5	*
038.2	15	*6	*9	*	*	*	*7	*	*	*6	*
038.4	82	34	48	*	*	16	60	24	19	24	15
038.40	*8	*	*	*	*	*	*	*	*	*	*
038.42	52	18	34	*	*	9	39	15	12	16	9
038.43	*6	*	*	-	*	*	*	*	*	*	*
038.49	16	*8	*9	*	*	*	12	*	*6	*	*
038.8	10	*	*6	*	*	*	*7	*	*	*	*
038.9	145	66	79	*8	16	24	97	40	34	53	18
041	14	*8	*6	*	*	*	*6	*	*	*7	*
042	69	48	20	*	47	20	*	26	*9	23	11
047	32	18	13	16	12	*	*	*	*6	12	11
047.9	29	18	12	15	11	*	*	*	*5	12	9
053	19	*8	11	*	*	*	13	*	*6	*6	*
053.9	10	*	*6	*	*	*	*6	*	*	*	*
054	*7	*	*	*	*	*	*	*	*	*	*
070	20	12	*8	*	*8	*8	*	*	*	*8	*
070.5	*8	*	*	-	*	*	*	*	*	*	*
075	9	*	*6	*	*5	*	-	*	*	*	*
079	65	33	32	39	12	*5	*8	16	14	22	13
079.6	*6	*	*	*6	-	-	-	*	*	*	*
079.9	58	29	28	33	11	*5	*8	15	11	21	11
079.99	58	29	28	33	11	*5	*8	15	11	21	11
112	17	*8	*9	*	*	*5	*6	*	*	*7	*
135	*8	*	*5	-	*	*	*	*	*	*	*
136	*5	*	*	*	*	*	*	*	*	*	*
140-239	1,587	586	1,002	37	289	566	695	383	333	578	295
150	11	*8	*	-	*	*	*7	*	*	*	*
151	24	14	9	-	*	*9	12	*5	*	*	*9
151.0	*8	*7	*	-	*	*	*	*	*	*	*
151.9	*6	*	*	-	*	*	*	*	*	*	*
153	117	56	61	-	*5	36	76	24	31	42	20
153.0	*6	*	*	-	*	*	*	*	*	*	*
153.1	10	*	*6	-	*	*	*6	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
153.2	*7	*	*	—	*	*	*	*	*	*	*
153.3	29	14	15	—	*	10	18	*6	*9	10	*
153.4	19	*9	11	—	*	*6	14	*	*	*7	*
153.6	21	9	12	—	*	*	16	*	*	9	*
153.7	*7	*5	*	—	*	*	*	*	*	*	*
153.9	14	*6	*8	—	*	*	*9	*	*6	*	*
154	38	22	16	*	*	12	24	10	*8	13	*7
154.0	11	*7	*	—	*	*	*6	*	*	*	*
154.1	25	14	11	*	*	*7	17	*6	*	9	*
155	11	*7	*	*	*	*6	*	*	*	*	*
155.0	*5	*	*	*	*	*	*	*	*	*	*
156	*6	*	*	—	—	*	*	*	*	*	*
157	26	12	14	*	*	*7	17	*6	*5	10	*
157.0	11	*	*6	*	*	*	*7	*	*	*	*
157.9	*9	*	*	—	*	*	*6	*	*	*	*
161	*7	*	*	—	*	*	*	*	*	*	*
162	122	67	55	*	*5	38	78	28	29	43	22
162.2	*8	*	*	—	*	*	*5	*	*	*	*
162.3	38	22	16	—	*	13	24	10	*9	11	*8
162.5	17	*9	*9	—	*	*	12	*	*	*	*
162.8	12	*6	*6	—	*	*	*8	*6	*	*	*
162.9	42	23	19	*	*	12	27	*7	10	20	*5
170	*6	*	*	*	*	*	*	*	*	*	*
171	*6	*	*	*	*	*	*	*	*	*	*
174	108	—	108	—	12	40	55	29	25	34	19
174.3	*5	—	*5	—	*	*	*	*	*	*	*
174.4	21	—	21	—	*	*6	13	*6	*5	*	*5
174.8	26	—	26	—	*	12	11	10	*	*6	*5
174.9	40	—	40	—	*	16	20	*8	*9	18	*
180	16	—	16	—	*8	*	*	*	*	*6	*
180.9	10	—	10	—	*5	*	*	*	*	*	*
182	30	—	30	—	*	12	16	*7	*6	11	*5
182.0	30	—	30	—	*	12	16	*7	*6	11	*5
183	26	—	26	*	*	9	12	10	*	*7	*
183.0	25	—	25	*	*	*9	12	9	*	*7	*
185	80	80	—	—	*	29	50	17	20	28	14
188	37	28	10	*	*	*7	30	14	*7	10	*5
188.8	11	*8	*	—	*	*	*8	*	*	*	*
188.9	13	10	*	*	*	*	11	*	*	*	*
189	34	19	15	*	*	15	16	*8	*8	15	*
189.0	30	17	13	*	*	14	13	*6	*7	15	*
191	27	15	12	*	*6	11	*7	*6	*	9	*7
191.8	*5	*	*	*	*	*	*	*	*	*	*
191.9	*6	*	*	*	*	*	*	*	*	*	*
193	16	*	12	*	*6	*7	*	*	*	*6	*
196	13	*5	*7	—	*	*6	*5	*	*	*	*
196.0	*6	*	*	—	*	*	*	*	*	*	*
197	93	33	60	*	11	28	53	26	20	31	15
197.0	16	*5	11	*	*	*	*9	*	*	*5	*
197.2	28	*6	21	—	*	*8	16	9	*	11	*
197.6	11	*	*7	—	*	*	*7	*	*	*	*
197.7	28	12	15	*	*	*8	16	*7	*7	*9	*
198	83	37	46	*	*8	31	43	23	20	26	15
198.3	33	14	19	*	*	15	14	9	*8	*8	*7
198.5	24	11	13	*	*	*8	16	*5	*8	*7	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
198.8	18	*9	10	—	*	*7	9	*	*	*9	*
198.89	16	*7	9	—	*	*6	9	*	*	*7	*
199	35	16	20	*	*6	12	18	10	*	17	*
199.0	26	10	16	—	*	10	12	*7	*	13	*
199.1	9	*5	*	*	*	*	*5	*	*	*	*
200	12	*6	*6	*	*	*	*6	*	*	*	*
200.0	*9	*	*	*	*	*	*	*	*	*	*
202	36	20	16	*	*	13	17	*9	*8	15	*
202.8	31	18	12	—	*	12	15	*8	*7	13	*
202.80	14	*8	*6	—	*	*7	*6	*	*	*6	*
202.83	*6	*	*	—	*	*	*	*	*	*	*
203	14	*7	*7	—	*	*5	*7	*	*	*6	*
203.0	13	*7	*7	—	*	*5	*7	*	*	*6	*
203.00	13	*7	*7	—	*	*5	*7	*	*	*6	*
204	17	10	*7	10	*	*	*	*	*	*	*8
204.0	11	*6	*	10	*	*	*	*	*	*	*6
204.00	*7	*	*	*6	*	*	*	*	*	*	*
204.1	*6	*	*	—	*	*	*	*	*	*	*
204.10	*6	*	*	—	*	*	*	*	*	*	*
205	17	11	*6	*	*	*6	*5	*	*	*5	*
205.0	12	*8	*	*	*	*5	*	*	*	*	*
205.00	11	*7	*	*	*	*	*	*	*	*	*
210	*5	*	*	*	*	*	*	*	*	*	*
211	35	12	22	*	*	*7	25	*7	*6	15	*7
211.3	22	9	13	*	*	*	16	*	*	9	*5
211.4	*7	*	*5	—	*	*	*6	*	*	*	*
218	240	—	240	—	121	114	*5	48	51	99	42
218.0	31	—	31	—	14	15	*	*8	*	12	*6
218.1	77	—	77	—	39	37	*	16	14	35	13
218.2	27	—	27	—	14	13	*	*8	*	*7	*8
218.9	105	—	105	—	54	49	*	16	29	45	15
220	35	—	35	*	18	11	*5	*9	10	10	*7
225	22	*8	14	*	*	9	*9	*6	*	*6	*5
225.2	14	*	10	—	*	*6	*6	*	*	*	*
226	10	*	*9	—	*	*	*	*	*	*	*
227	15	*	11	*	*	*6	*6	*	*	*	*
227.1	*5	*	*	—	*	*	*	*	*	*	*
227.3	*6	*	*	*	*	*	*	*	*	*	*
228	*6	*	*	*	*	*	*	*	*	*	*
233	25	*	24	—	10	*9	*6	*	*5	*9	*7
233.0	12	*	12	—	*	*5	*5	*	*	*	*
233.1	12	—	12	—	*8	*	*	*	*	*	*
235	*8	*	*	*	*	*	*	*	*	*5	*
237	*6	*	*	*	*	*	*	*	*	*	*
238	13	*6	*7	*	*	*	*9	*	*	*	*
238.7	11	*	*6	*	*	*	*8	*	*	*	*
239	11	*	*6	*	*	*	*6	*	*	*6	*
240-279	1,455	625	830	152	289	381	634	318	350	574	213
241	15	*	13	—	*	*7	*	*	*	*7	*
241.1	10	*	*9	—	*	*5	*	*	*	*5	*
242	14	*	9	*	*	*	*	*	*	*	*
242.0	*6	*	*	*	*	*	*	*	*	*	*
242.00	*5	*	*	*	*	*	*	*	*	*	*
242.9	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
244	18	*	16	—	*	*6	*9	*6	*	*6	*
244.9	17	*	15	—	*	*6	*9	*6	*	*6	*
250	557	279	278	22	136	207	192	129	120	228	80
250.0	179	83	96	*8	38	74	59	39	34	79	27
250.00	79	35	44	*	13	36	29	14	15	40	10
250.01	29	14	15	*6	*5	10	*8	*8	*7	10	*
250.02	52	22	31	*	12	22	18	12	*8	21	12
250.03	18	12	*7	*	*8	*6	*	*	*	*9	*
250.1	103	58	45	12	60	23	*8	21	27	41	12
250.10	18	11	*6	*	*9	*6	*	*	*	*8	*
250.11	38	19	19	*7	24	*6	*	*8	12	11	*6
250.12	15	9	*6	—	*7	*5	*	*	*	*6	*
250.13	32	18	14	*	20	*6	*	*	*9	16	*
250.2	13	*6	*7	*	*	*	*7	*	*	*	*
250.22	*6	*	*	—	*	*	*	*	*	*	*
250.4	33	17	16	—	*	18	10	*8	*6	9	9
250.40	19	10	9	—	*	10	*7	*	*	*	*6
250.41	*8	*	*	—	*	*	*	*	*	*	*
250.6	52	22	30	—	11	25	16	13	11	24	*
250.60	24	11	13	—	*	10	10	*5	*6	11	*
250.61	10	*	*	—	*	*5	*	*	*	*6	*
250.62	12	*	*7	—	*	*7	*	*	*	*	*
250.63	*6	*	*	—	*	*	*	*	*	*	*
250.7	52	30	23	—	*	21	29	12	11	20	9
250.70	28	17	11	—	*	9	18	*6	*7	11	*
250.71	15	*9	*7	—	*	*7	*8	*	*	*7	*
250.72	*6	*	*	—	*	*	*	*	*	*	*
250.8	109	57	52	*	15	37	56	28	25	44	13
250.80	60	32	28	—	*5	19	35	16	12	21	11
250.81	18	12	*7	*	*	*8	*7	*7	*	*8	*
250.82	20	*7	12	—	*	*7	10	*	*7	*8	*
250.83	11	*6	*	*	*	*	*	*	*	*7	*
250.9	*8	*	*	*	*	*	*	*	*	*	*
251	17	*7	10	*	*	*	10	*	*	*8	*
251.2	13	*7	*6	*	*	*	*8	*	*	*7	*
253	12	*	*7	*	*	*	*7	*	*	*	*
253.6	*8	*	*6	*	*	*	*6	*	*	*	*
255	*6	*	*	*	*	*	*	*	*	*	*
263	15	*	11	*	*	*	10	*	*	*6	*
263.9	13	*	10	*	*	*	9	*	*	*5	*
272	34	19	15	*	*8	20	*6	9	*7	10	*8
272.0	17	9	*8	—	*	11	*	*	*	*5	*
272.4	15	*9	*6	—	*	*7	*	*	*	*	*
274	11	*6	*6	—	*	*	*7	*	*	*	*
274.0	*7	*	*	—	*	*	*	*	*	*	*
275	19	*6	13	*	*5	*	*9	*	*6	*7	*
275.4	15	*	11	*	*	*	*8	*	*5	*	*
275.42	*9	*	*7	*	*	*	*6	*	*	*	*
276	646	261	385	115	75	92	363	128	172	256	89
276.0	*8	*	*	*	*	*	*5	*	*	*	*
276.1	69	18	51	*	*6	12	51	13	18	24	15
276.2	*7	*	*5	*	*	*	*	*	*	*	*
276.5	485	206	279	111	54	59	261	100	130	193	62
276.6	*6	*	*	*	*	*	*	*	*	*	*
276.7	28	13	15	*	*	*7	19	*	*8	12	*
276.8	35	12	23	*	*9	*9	17	*7	*9	15	*
276.9	*6	*	*	*	*	*	*5	*	*	*	*
277	18	*7	11	*5	9	*	*	*	*7	*6	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
277.0	12	*6	*7	*	*7	*	—	*	*	*5	*
277.00	12	*6	*7	*	*7	*	—	*	*	*5	*
278	45	11	34	*	29	15	*	*9	13	13	10
278.0	42	11	31	*	27	14	*	*8	12	12	10
278.00	*8	*	*	—	*	*	*	*	*	*	*
278.01	34	*6	27	*	24	10	*	*6	*9	10	*9
280-289	392	163	229	70	91	83	149	92	87	154	59
280	57	18	39	*	*8	14	34	11	14	23	*8
280.0	27	*8	19	*	*	*7	16	*5	*	13	*
280.9	29	10	19	*	*	*7	18	*6	10	10	*
282	74	31	42	24	46	*	*	26	16	26	*6
282.6	67	30	37	21	43	*	*	24	15	23	*
282.62	60	27	34	17	41	*	—	23	13	21	*
283	*6	*	*	*	*	*	*	*	*	*	*
284	28	16	12	*7	*	*8	*9	*9	*5	9	*
284.8	21	10	11	*	*	*7	*8	*5	*5	*6	*
284.9	*6	*	*	*	*	*	*	*	*	*	—
285	82	28	53	*	*9	17	55	16	13	40	13
285.1	14	*7	*7	*	*	*	9	*	*	*5	*
285.9	64	20	45	*	*7	14	43	13	9	33	9
286	20	*8	12	*	*	*	11	*	*5	*9	*
286.7	*6	*	*	—	*	*	*	*	*	*	*
287	40	19	21	10	*6	13	10	*8	10	14	*8
287.3	13	*7	*6	*	*	*	*	*	*	*	*
287.4	*6	*	*	*	*	*	*	*	*	*	*
287.5	16	*7	*8	*	*	*6	*	*	*	*6	*
288	64	29	36	17	9	15	23	14	19	22	10
288.0	49	21	28	13	*7	14	14	12	14	16	*7
288.8	13	*7	*6	*	*	*	*7	*	*	*	*
289	18	9	9	*5	*5	*	*	*	*	*8	*
290-319	2,147	1,104	1,043	103	1,255	495	293	692	446	825	185
290	44	15	29	—	—	*	41	9	10	20	*
290.0	*6	*	*	—	—	—	*6	*	*	*	*
290.2	*8	*	*6	—	—	*	*8	*	*	*	*
290.3	*6	*	*	—	—	—	*6	*	*	*	*
290.4	22	*8	14	—	—	*	20	*	*	13	*
290.40	*7	*	*	—	—	*	*6	*	*	*	*
290.41	*8	*	*5	—	—	*	*8	*	*	*	*
291	126	93	33	*	75	43	*7	37	48	28	12
291.0	20	16	*	—	12	*8	*	*7	*5	*6	*
291.8	97	72	26	*	62	31	*	28	40	19	9
291.81	89	66	23	*	55	29	*	27	39	15	*8
291.89	*8	*6	*	—	*7	*	—	*	*	*	*
292	89	48	40	*	59	17	12	48	13	19	*9
292.0	51	31	20	*	40	10	*	34	*7	*8	*
292.8	29	13	15	—	17	*	*8	12	*	*9	*
292.81	15	*	12	—	*5	*	*6	*	*	*5	*
292.84	*9	*7	*	—	*7	*	*	*	*	*	*
292.9	*5	*	*	—	*	*	*	*	*	*	*
293	39	19	20	*	13	9	17	18	*9	*8	*
293.0	18	*6	12	*	*	*	12	*	*5	*	*
293.8	19	12	*7	*	11	*	*	13	*	*	*
293.83	16	11	*	*	10	*	*	11	*	*	*
294	36	15	21	—	*	*	30	12	*8	10	*7
294.8	33	13	19	—	*	*	28	10	*8	9	*6

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
295	264	158	106	*	162	80	19	95	50	96	22
295.3	78	52	26	—	45	26	*7	33	17	18	*9
295.30	25	18	*8	—	17	*7	*	*7	10	*	*
295.32	15	10	*	—	*7	*5	*	*6	*	*	*
295.34	37	24	13	—	21	13	*	21	*	*9	*
295.6	19	10	*9	—	12	*6	*	*6	*	*8	*
295.64	11	*6	*5	—	*7	*	*	*	*	*	*
295.7	132	75	56	*	86	38	*6	43	22	59	*8
295.70	87	53	35	*	56	26	*	18	17	47	*6
295.74	41	21	20	—	28	11	*	25	*	10	*
295.9	28	17	10	*	16	*7	*	11	*5	*8	*
295.90	16	9	*7	*	*9	*	*	*	*	*7	*
295.94	*8	*7	*	—	*6	*	*	*7	*	*	*
296	759	321	437	48	451	168	92	200	160	355	43
296.2	163	63	100	16	96	26	25	35	43	74	11
296.20	92	39	53	10	55	14	13	20	29	40	*
296.22	*8	*	*	*	*	*	*	*	*	*5	*
296.23	32	12	20	*	18	*7	*	*6	*6	16	*
296.24	31	*8	22	*	19	*	*6	*9	*6	13	*
296.3	264	99	165	*9	136	73	47	78	59	113	15
296.30	70	22	48	*	38	19	10	18	19	32	*
296.32	13	*6	*7	*	*7	*	*	*	*	*	*
296.33	123	49	74	*	68	34	17	44	23	49	*8
296.34	56	20	35	*	21	17	17	14	12	27	*
296.4	62	28	34	*	35	20	*6	23	12	22	*5
296.40	32	15	17	*	19	9	*	*7	*7	15	*
296.43	*5	*	*	—	*	*	*	*	*	*	*
296.44	23	11	12	*	12	*9	*	13	*	*	*
296.5	73	31	41	*	46	18	*7	15	15	40	*
296.50	44	20	25	*	27	12	*	*8	*9	27	*
296.53	14	*	10	*	10	*	*	*	*	*7	*
296.54	13	*7	*6	—	*8	*	*	*	*	*5	*
296.6	86	44	43	*8	62	12	*	16	12	55	*
296.60	43	24	19	*7	30	*	*	*5	*7	29	*
296.63	13	*6	*7	*	10	*	*	*6	*	*	*
296.64	29	13	16	*	23	*5	*	*	*	21	*
296.7	51	26	26	*	33	11	*	16	15	16	*
296.8	36	19	17	*7	27	*	*	*	*	28	*
296.80	18	14	*	*	14	*	*	*	*	17	—
296.89	17	*	12	*	13	*	*	*	*	11	*
296.9	20	10	10	*	12	*6	*	12	*	*5	*
296.90	19	9	9	*	11	*6	*	12	*	*	*
297	*9	*	*6	—	*	*	*6	*	*	*	*
298	79	38	40	*	40	18	18	35	11	23	10
298.9	75	36	39	*	39	16	17	33	11	22	9
300	80	30	50	*	42	19	14	19	22	31	*8
300.0	38	12	25	*	18	9	11	10	10	15	*
300.00	26	10	17	*	12	*8	*6	*6	*8	11	*
300.01	9	*	*7	—	*	*	*	*	*	*	*
300.1	*7	*	*	*	*	*	*	*	*	*	*
300.4	23	11	12	*	13	*	*	*5	*7	*8	*
300.9	*5	*	*	*	*	*	*	*	*	*	*
301	9	*	*5	*	*7	*	*	*	*	*	*
303	146	107	40	*	92	49	*6	28	25	75	18
303.0	27	20	*7	—	14	12	*	*	*9	*8	*5
303.00	10	*7	*	—	*5	*	*	*	*	*	*
303.01	17	13	*	—	*8	*7	*	*	*	*	*
303.9	119	87	33	*	78	37	*	23	16	67	13
303.90	61	45	16	*	37	22	*	*	*6	47	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
303.91	57	41	16	*	40	14	*	20	9	20	*8
304	93	57	36	*	76	14	*	37	10	40	*6
304.0	46	27	19	—	35	9	*	24	*	16	*
304.00	*8	*	*5	—	*5	*	*	*	*	*6	*
304.01	38	24	14	—	30	*8	*	23	*	10	*
304.2	14	10	*	*	13	*	*	*6	*	*6	*
304.21	*9	*5	*	—	*9	*	*	*	*	*	*
304.3	*7	*6	*	*	*7	*	—	*	*	*	*
304.4	*5	*	*	—	*	*	—	—	*	*	*
304.8	10	*6	*	*	9	*	—	*	*	*8	*
304.80	*8	*5	*	*	*7	*	—	*	*	*8	*
305	87	52	35	*	51	30	*6	15	12	45	15
305.0	38	25	14	*	22	12	*	*7	*5	20	*6
305.00	26	16	11	*	15	*9	*	*	*	15	*
305.01	*9	*7	*	—	*	*	*	*	*	*	*
305.1	20	12	*8	—	11	*8	*	*	*	9	*
305.6	11	*6	*5	—	*7	*	*	*	*	*6	*
305.60	*8	*5	*	—	*	*	*	*	*	*	*
305.9	13	*6	*7	*	*8	*	*	*	*	*8	*
305.90	12	*5	*7	*	*8	*	*	*	*	*8	*
307	*8	*	*5	*	*	*	*	*	*	*	*
308	*5	*	*	—	*	*	*	*	*	*	*
309	93	44	49	*6	73	9	*	40	23	19	11
309.0	34	16	18	*	29	*	*	17	*5	*7	*
309.2	14	*7	*8	*	10	*	*	*	*6	*	*
309.28	13	*6	*7	*	10	*	*	*	*6	*	*
309.4	24	12	12	*	18	*	*	11	*6	*	*
309.8	11	*	*8	*	*8	*	*	*7	*	*	*
309.81	10	*	*7	*	*7	*	*	*6	*	*	*
309.9	*8	*5	*	*	*6	*	*	*	*	*	*
310	*9	*	*5	*	*	*	*5	*	*	*	*
311	110	56	54	*8	72	22	*8	57	19	28	*6
312	29	16	13	*9	19	*	*	17	*	*	*
312.3	24	13	11	*6	16	*	*	16	*	*	*
312.30	18	9	9	*	13	*	—	15	*	*	*
313	11	*5	*6	*5	*5	—	—	*	*	*	*
313.8	11	*5	*6	*5	*5	—	—	*	*	*	*
313.81	10	*	*	*	*5	—	—	*	*	*	*
314	13	10	*	10	*	—	—	*	*	*	*
314.0	12	9	*	10	*	—	—	*	*	*	*
314.01	11	*8	*	9	*	—	—	*	*	*	*
320-389	471	196	275	84	112	109	167	108	115	167	80
320	*6	*	*	*	*	*	*	*	*	*	*
331	59	29	31	*	*	*	49	11	18	21	*9
331.0	42	18	24	—	—	*	40	*8	14	15	*
331.3	*6	*	*	*	*	*	*5	*	*	*	*
331.4	*9	*5	*	*	*	*	*	*	*	*	*
332	19	*7	12	—	*	*	17	*7	*	*5	*
332.0	19	*7	12	—	*	*	17	*7	*	*5	*
333	13	*7	*5	*	*	*	*7	*	*	*	*
337	*9	*	*5	*	*	*	*	*	*	*	*
337.2	*6	*	*	*	*	*	*	*	*	*	*
340	25	*6	19	—	11	11	*	*	*9	9	*
343	*8	*	*	*5	*	*	—	*	*	*	*
343.9	*5	*	*	*	*	—	—	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
345	56	29	27	22	16	9	*9	15	13	17	11
345.1	*8	*5	*	*	*	*	*	*	*	*	*
345.10	*7	*	*	*	*	*	*	*	*	*	*
345.3	16	9	*6	*6	*	*	*	*	*	*5	*
345.4	*5	*	*	*	*	*	*	*	*	*	—
345.5	*7	*	*5	*	*	*	*	*	*	*	*
345.9	14	*8	*6	*6	*	*	*	*	*	*5	*
345.90	12	*7	*	*	*	*	*	*	*	*5	*
346	43	*7	36	*	24	15	*	*7	11	19	*5
346.8	*7	*	*5	*	*	*	*	*	*	*	*
346.9	27	*	23	*	16	*9	*	*	*6	13	*
346.90	20	*	17	*	10	*7	*	*	*	10	*
346.91	*7	*	*6	*	*5	*	*	*	*	*	*
348	25	10	14	*	*6	*7	*8	*6	*8	*8	*
348.3	*8	*	*5	*	*	*	*	*	*	*	*
349	*7	*	*	*	*	*	*	*	*	*	*
350	*5	*	*	—	*	*	*	*	*	*	*
350.1	*5	*	*	—	*	*	*	*	*	*	*
351	*9	*	*	*	*	*	*	*	*	*	*
351.0	*8	*	*	*	*	*	*	*	*	*	*
354	*7	*	*	*	*	*	*	*	*	*	*
356	*6	*	*	*	*	*	*	*	*	*	*
357	*6	*	*	*	*	*	*	*	*	*	*
359	*5	*	*	*	*	*	*	*	*	*	*
376	*8	*	*	*6	*	*	*	*	*	*	*
376.0	*8	*	*	*6	*	*	*	*	*	*	*
376.01	*7	*	*	*6	*	*	*	*	*	*	*
380	*7	*	*	*	*	*	*	*	*	*	*
380.1	*5	*	*	*	*	*	*	*	*	*	*
382	16	*8	*8	15	*	*	*	*	*	*5	*
382.9	15	*7	*8	13	*	*	*	*	*	*5	*
386	38	10	28	—	*	14	22	11	11	13	*
386.1	14	*	10	—	*	*	*9	*6	*	*	*
386.11	*8	*	*7	—	*	*	*	*	*	*	*
386.3	20	*6	13	—	*	*9	10	*	*	*9	*
386.30	17	*5	12	—	*	*6	9	*	*	*8	*
390-459	6,294	3,115	3,179	30	389	1,813	4,062	1,507	1,509	2,382	895
394	*6	*	*5	—	*	*	*	*	*	*	*
396	12	*	*8	—	*	*	10	*	*	*5	*
396.2	*5	*	*	—	—	*	*	*	*	*	*
398	29	11	18	—	*	*	27	15	*	*7	*
398.9	29	11	18	—	*	*	27	15	*	*7	*
398.91	28	11	18	—	*	*	27	15	*	*6	*
401	267	105	162	*	42	119	106	58	58	112	38
401.0	28	10	17	—	*7	11	10	*6	*5	14	*
401.9	235	92	143	*	34	106	94	51	51	98	36
402	83	32	50	*	*	20	58	23	14	35	10
402.9	80	32	49	*	*	20	56	23	13	35	10
402.90	16	*7	9	*	*	*7	*8	*5	*	*7	*
402.91	64	25	39	—	*	13	48	17	11	27	*9
403	80	36	44	*	15	27	37	14	15	39	12
403.9	75	34	41	*	13	26	36	13	15	36	12
403.90	*7	*	*	*	*	*	*	*	*	*	—
403.91	68	31	37	*	12	24	32	10	14	33	12

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
404	27	13	15	—	*	*8	18	*6	*	13	*
404.9	25	11	14	—	*	*6	17	*	*	12	*
404.93	18	*8	9	—	*	*	12	*	*	*9	*
410	781	453	328	*	38	242	499	190	185	283	122
410.0	30	18	11	—	*	*9	20	*6	*7	13	*
410.01	29	18	11	—	*	*9	19	*5	*7	13	*
410.1	101	60	41	—	*8	32	61	22	23	39	17
410.11	95	56	39	—	*7	30	59	19	23	37	17
410.2	20	10	10	—	*	*7	12	*6	*	*6	*
410.21	19	10	9	—	*	*7	11	*6	*	*5	*
410.3	14	9	*	—	*	*6	*6	*	*	*	*
410.31	13	*9	*	—	*	*6	*6	*	*	*	*
410.4	117	74	43	—	*7	53	57	30	29	40	18
410.41	113	73	40	—	*7	50	56	29	29	37	18
410.5	12	9	*	*	*	*	*7	*	*	*	*
410.51	12	*9	*	*	*	*	*7	*	*	*	*
410.7	350	195	154	*	15	98	236	92	83	120	54
410.71	343	192	151	*	15	97	230	91	82	116	54
410.8	*8	*	*	*	*	*	*6	*	*	*	*
410.81	*7	*	*	*	*	*	*6	*	*	*	*
410.9	125	68	56	*	*	29	91	27	29	52	17
410.90	*9	*	*	—	*	*	*7	*	*	*	—
410.91	116	63	52	*	*	28	83	23	28	48	17
411	171	90	81	*	14	67	90	53	40	58	20
411.1	153	82	71	—	13	63	77	45	36	55	17
411.8	15	*7	*8	*	*	*	11	*7	*	*	*
411.89	14	*6	*8	*	*	*	10	*7	*	*	*
412	*6	*	*	—	*	*	*	*	*	*	*
413	82	34	48	—	*6	29	48	23	15	31	13
413.9	81	34	47	—	*5	28	48	23	15	31	13
414	1,125	692	433	*	51	450	624	262	279	435	150
414.0	1,102	676	425	*	50	442	609	254	274	426	147
414.00	187	109	78	—	*7	68	111	41	55	74	17
414.01	874	540	335	*	42	362	471	205	208	336	125
414.02	31	21	10	—	*	9	22	*6	10	11	*
414.05	*7	*	*	*	*	*	*	*	*	*	*
414.8	11	*7	*	—	*	*	9	*	*	*	*
414.9	12	*8	*	—	*	*6	*6	*	*	*	*
415	83	31	52	—	12	27	44	20	20	31	13
415.1	82	31	51	—	12	27	43	19	20	30	13
415.11	*6	*	*	—	*	*	*	*	*	*	*
415.19	76	29	47	—	10	24	42	18	18	28	12
416	11	*	*8	*	*	*	*6	*	*	*	*
416.0	*7	*	*6	*	*	*	*	*	*	*	*
420	12	*7	*	*	*	*	*	*	*	*	*
420.9	12	*7	*	*	*	*	*	*	*	*	*
420.90	10	*6	*	*	*	*	*	*	*	*	*
421	17	11	*6	*	*	*5	*8	*6	*	*6	*
421.0	16	10	*6	*	*	*5	*8	*6	*	*5	*
423	13	*	*8	*	*	*	*6	*	*	*5	*
423.9	11	*	*7	—	*	*	*	*	*	*	*
424	93	40	53	*	11	28	53	25	22	29	18
424.0	39	13	26	*	*7	15	16	9	*9	14	*7
424.1	51	26	25	—	*	12	35	15	13	13	10
425	31	15	16	*	*5	10	14	*7	*9	12	*
425.4	26	13	14	*	*	9	12	*6	*8	10	*
426	53	27	26	*	*	*7	42	16	13	16	*8

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
426.0	27	13	14	*	*	*	22	*7	*7	9	*
426.1	11	*6	*	*	*	*	*8	*	*	*	*
427	716	336	380	*8	43	157	508	181	176	252	106
427.0	29	11	18	*	*	*7	17	*5	9	*9	*6
427.1	50	38	12	*	*	15	31	17	10	17	*7
427.3	399	180	219	*	18	90	291	99	105	142	52
427.31	357	153	203	—	17	75	265	91	95	125	45
427.32	42	27	15	*	*	15	25	*8	10	17	*7
427.4	*6	*	*	*	*	*	*	*	*	*	*
427.41	*6	*	*	*	*	*	*	*	*	*	*
427.5	11	*5	*5	*	*	*	*7	*	*	*	*
427.6	14	*8	*6	—	*	*	*9	*	*	*6	*
427.69	10	*7	*	—	*	*	*6	*	*	*	*
427.8	185	78	108	*	10	33	137	45	45	64	31
427.81	72	32	40	*	*	*8	63	22	16	26	*7
427.89	113	46	67	*	10	26	74	23	28	38	23
427.9	21	11	10	*	*	*5	11	*8	*	*8	*
428	1,008	422	586	*	35	196	775	238	260	387	123
428.0	999	418	581	*	35	195	767	235	257	384	122
428.9	*6	*	*	*	—	*	*5	*	*	*	—
429	17	*8	9	*	*	*6	10	*	*	*8	*
429.2	*8	*	*	—	—	*	*6	*	*	*	*
430	22	*9	14	—	*6	9	*7	*5	*	9	*
431	66	29	38	*	*	21	41	17	15	21	13
432	21	13	*8	*	*	*	14	*	*	*7	*6
432.1	19	12	*7	*	*	*	13	*	*	*7	*5
433	164	87	78	*	*	41	121	37	40	66	21
433.1	128	66	62	—	*	30	97	28	31	53	16
433.10	108	54	54	—	*	27	81	22	27	47	13
433.11	19	12	*7	—	*	*	16	*6	*	*6	*
433.3	31	19	12	—	*	*9	22	*8	*7	11	*
433.30	25	16	9	—	*	*7	17	*7	*6	*9	*
433.31	*6	*	*	—	*	*	*	*	*	*	*
434	287	125	162	*	*9	68	209	61	65	116	45
434.0	12	*6	*6	—	*	*	*8	*	*	*	*
434.01	10	*	*5	—	*	*	*7	*	*	*	*
434.1	39	16	24	—	*	10	28	10	*8	12	9
434.10	*6	*	*	—	—	*	*6	*	*	*	*
434.11	33	14	19	—	*	9	21	*8	*6	9	9
434.9	236	103	133	*	*6	55	173	49	53	101	33
434.90	*6	*	*	—	*	*	*	*	*	*	*
434.91	230	101	129	*	*6	53	169	48	52	98	33
435	224	91	133	*	10	43	171	66	47	83	28
435.3	*7	*	*	—	*	*	*6	*	*	*	*
435.8	*5	*	*	—	*	*	*	*	*	*	*
435.9	209	84	125	*	9	40	159	63	44	79	24
436	146	56	91	*	*	32	112	18	42	60	26
437	24	*8	16	*	*	*7	15	*	*7	*9	*
437.3	*7	*	*	—	*	*	*	*	*	*	*
438	26	12	14	*	*	*	22	*	*5	13	*5
438.2	*9	*	*	—	*	*	*7	*	*	*	*
438.20	*8	*	*	—	*	*	*7	*	*	*	*
438.8	11	*6	*6	—	*	*	10	*	*	*	*
438.89	*8	*	*	—	*	*	*6	*	*	*	*
440	125	72	53	*	*	35	87	32	31	43	19
440.1	*6	*	*	—	*	*	*	*	*	*	*
440.2	108	63	45	*	*	30	75	28	25	39	16

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
440.20	12	*8	*	*	*	*	*7	*	*	*5	*
440.21	27	16	10	—	*	11	16	*7	*6	9	*
440.22	12	*6	*6	—	*	*	*8	*	*	*	*
440.23	20	11	*8	—	*	*	18	*5	*6	*	*
440.24	36	20	16	—	*	*8	27	*8	*8	14	*5
440.3	*5	*	*	—	—	*	*	*	*	*	*
441	60	43	17	*	*	12	46	16	12	21	11
441.0	10	*6	*	*	*	*	*	*	*	*	*
441.01	*6	*	*	*	*	*	*	*	*	*	*
441.3	*7	*	*	—	—	*	*6	*	*	*	*
441.4	36	29	*7	—	*	*6	30	10	*8	12	*6
442	*8	*	*	*	*	*	*	*	*	*	*
443	27	12	15	*	*	*7	17	*	10	11	*
443.9	22	10	12	—	*	*6	16	*	*8	9	*
444	37	21	17	—	*	12	24	*8	*5	19	*5
444.2	27	15	13	—	*	*8	18	*6	*	13	*
444.22	23	12	10	—	*	*6	15	*5	*	11	*
446	13	*6	*7	*6	*	*	*	*	*	*	*
446.1	*6	*	*	*6	*	—	—	*	*	*	*
447	14	*6	*8	*	*	*6	*6	*	*	*6	*
447.1	*8	*	*	—	*	*	*	*	*	*	*
451	23	*8	15	*	*	*7	12	*	*7	11	*
451.0	*7	*	*5	—	*	*	*	*	*	*	*
451.1	11	*	*7	—	*	*	*6	*	*	*	*
451.19	*7	*	*5	—	*	*	*	*	*	*	*
453	136	62	74	*	19	41	76	32	39	47	18
453.8	118	53	66	*	16	37	65	26	35	41	16
453.9	16	9	*7	—	*	*	10	*6	*	*5	*
454	*8	*	*	—	*	*	*	*	*	*	*
455	29	16	13	—	*6	11	12	*6	*6	13	*
455.2	19	10	*8	—	*	*7	*8	*	*	*9	*
456	*5	*	*	—	*	*	*	*	*	*	*
458	66	30	36	*	*	14	47	12	16	27	11
458.0	31	15	16	*	*	*7	23	*6	*8	13	*
458.9	29	12	17	*	*	*6	20	*6	*7	11	*6
459	18	10	*9	*	*	*6	11	*	*7	*	*
459.8	10	*6	*	—	*	*	*8	*	*	*	*
459.81	*8	*	*	—	*	*	*5	*	*	*	*
460-519	3,444	1,603	1,841	699	394	648	1,703	765	824	1,286	569
461	14	*7	*7	*	*6	*	*	*	*	*7	*
462	11	*	*7	*	*6	*	*	*	*	*	*
463	*8	*	*	*	*6	*	*	*	*	*	*
464	28	19	9	25	*	*	*	*6	*7	*8	*8
464.4	22	16	*6	22	*	—	—	*	*6	*6	*6
465	31	15	15	18	*	*	*6	*7	*6	10	*8
465.9	29	15	14	18	*	*	*6	*6	*6	10	*8
466	285	145	140	201	13	19	52	43	64	99	79
466.0	97	40	57	16	13	19	49	18	24	43	11
466.1	189	105	84	185	*	*	*	25	40	56	68
466.11	106	58	48	105	*	*	*	14	22	25	45
466.19	82	47	36	80	*	*	*	11	18	31	23
470	*6	*5	*	—	*	*	*	*	*	*	*
473	11	*6	*5	*	*	*	*	*	*	*6	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
474	15	*7	*8	12	*	*	*	*6	*	*6	*
474.0	*6	*	*	*	*	*	*	*	*	*	*
474.1	*9	*5	*	*8	*	*	—	*	*	*	*
474.10	*6	*	*	*6	*	—	—	*	*	*	*
475	15	*8	*7	*	*8	*	*	*	*	*	*
478	19	*8	11	*7	*	*	*	*6	*	*6	*5
478.2	*8	*	*	*	*	*	*	*	*	*	*
480	27	14	13	17	*	*	*7	*	*7	10	*7
480.1	12	*9	*	10	*	—	*	*	*	*5	*
480.9	13	*	*9	*6	*	*	*	*	*	*	*
481	36	17	18	*	*	10	21	*6	10	11	*9
482	141	71	70	*7	13	22	99	25	41	59	17
482.0	*7	*	*	—	*	*	*	*	*	*	*
482.1	27	13	14	*	*5	*	15	*6	*8	*8	*
482.2	19	10	*9	*	*	*	16	*	*7	*7	*
482.3	*8	*	*	*	*	*	*5	*	*	*	*
482.4	34	18	16	*	*	*	26	*	*7	20	*
482.41	29	17	12	*	*	*	23	*	*6	16	*
482.8	28	12	16	*	*	*	22	*	12	11	*
482.83	22	*8	14	*	*	*	19	*	11	*7	*
482.9	19	11	*8	*	*	*5	10	*7	*	*7	*
483	*7	*	*	*	*	*	*	*	*	*	*
483.0	*6	*	*	*	*	*	*	*	*	*	—
485	39	15	25	17	*	*5	14	*	*9	22	*
486	1,032	463	569	128	107	177	620	224	268	373	167
487	39	16	23	*7	*	*6	23	*6	14	15	*
487.0	13	*5	*8	*	*	*	*9	*	*	*5	*
487.1	25	11	15	*6	*	*	15	*	10	9	*
490	22	9	12	*5	*	*	10	*	*	11	*
491	495	215	280	*	18	147	328	126	111	198	59
491.2	493	214	279	*	18	146	328	126	111	198	59
491.20	17	10	*7	—	*	*	12	*	*	*8	*
491.21	477	205	272	*	17	142	316	121	107	190	59
492	21	12	*9	*	*	*5	14	*7	*6	*6	*
492.8	19	11	*8	—	*	*	14	*6	*5	*6	*
493	465	198	267	203	111	84	68	110	109	167	78
493.0	46	28	18	36	*7	*	*	*7	*	*7	29
493.00	23	13	10	17	*	*	*	*	*	*5	14
493.01	24	15	*8	19	*	*	*	*	*	*	15
493.2	58	20	37	*	10	22	25	19	12	21	*6
493.20	45	16	29	*	*9	16	19	15	10	15	*
493.21	13	*	*8	—	*	*6	*6	*	*	*5	*
493.9	360	149	211	166	93	59	42	85	93	139	43
493.90	237	90	147	91	68	43	35	56	56	97	28
493.91	123	59	64	76	25	15	*6	29	37	42	15
494	*8	*	*5	*	*	*	*6	*	*	*	*
496	115	64	51	*	*	36	74	29	26	43	17
507	155	89	66	*7	*9	15	125	32	35	65	23
507.0	153	88	66	*6	*9	14	124	32	34	65	22
510	10	*6	*	*	*	*	*	*	*	*	*
510.9	10	*6	*	*	*	*	*	*	*	*	*
511	53	23	29	*	*7	13	32	13	11	20	*9
511.0	*8	*	*	*	*	*	*	*	*	*	*
511.8	*6	*	*	*	*	*	*	*	*	*	*
511.9	38	17	21	*	*	9	24	*8	*7	16	*7

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
512	44	27	18	*	15	10	18	11	*9	18	*7
512.1	12	*5	*7	*	*	*	*7	*	*	*	*
512.8	29	19	10	*	14	*	10	*6	*6	13	*
513	*5	*	*	*	*	*	*	*	*	*	*
513.0	*5	*	*	*	*	*	*	*	*	*	*
514	*7	*5	*	*	*	*	*	*	*	*	*
515	24	*8	16	*	*	*	15	*6	*5	*9	*
516	15	*7	*8	*	*	*	9	*	*	*6	*
516.3	*6	*	*	*	*	*	*	*	*	*	*
518	211	89	122	10	20	52	129	59	43	73	36
518.0	*6	*	*	*	*	*	*	*	*	*	*
518.4	*8	*	*	*	*	*	*	*	*	*	*
518.8	188	78	110	*8	16	46	118	53	38	66	31
518.81	147	61	86	*	11	34	99	42	30	52	24
518.82	11	*	*6	*	*	*	*	*	*	*5	*
518.84	17	*7	10	*	*	*7	9	*6	*	*	*
518.89	10	*5	*5	*	*	*	*	*	*	*	*
519	21	9	11	*8	*	*	*5	*7	*6	*	*
519.1	10	*	*6	*	*	*	*	*	*	*	*
520-579	3,143	1,376	1,767	217	758	852	1,317	693	732	1,150	569
522	*7	*	*	*	*	*	*	*	*	*	*
522.5	*5	*	*	*	*	*	*	*	*	*	*
524	11	*	*8	*	9	*	*	*	*	*	*
524.0	*5	*	*	*	*	*	—	*	*	*	*
527	*7	*	*	*	*	*	*	*	*	*	*
527.2	*5	*	*	*	*	*	*	*	*	*	*
528	*9	*	*	*	*	*	*	*	*	*	*
530	222	101	121	28	44	66	84	46	54	79	43
530.1	35	21	15	*	*9	13	10	*6	12	9	*8
530.10	13	*6	*6	*	*	*5	*	*	*	*	*
530.11	15	*9	*6	*	*	*6	*	*	*	*	*
530.19	*8	*5	*	*	*	*	*	*	*	*	*
530.2	18	10	*8	*	*	*	11	*5	*	*5	*
530.3	*8	*	*	*	*	*	*6	*	*	*	*
530.5	*5	*	*	*	*	*	*	*	*	*	*
530.6	*5	*	*	*	*	*	*	*	*	*	*
530.7	11	*6	*6	*	*	*	*6	*	*	*	*
530.8	135	54	81	25	27	40	43	28	29	50	27
530.81	127	50	76	25	26	38	38	26	27	49	25
530.82	*5	*	*	—	*	*	*	*	*	*	*
531	82	37	45	*	*8	17	56	14	25	26	17
531.0	16	*9	*8	*	*	*	12	*	*6	*6	*
531.00	15	*8	*7	*	*	*	12	*	*6	*6	*
531.4	45	22	23	*	*	10	31	*8	12	14	11
531.40	44	22	23	*	*	10	30	*7	12	14	10
531.9	12	*	*8	*	*	*	*7	*	*	*	*
531.90	10	*	*7	*	*	*	*6	*	*	*	*
532	71	44	27	*	14	16	41	15	15	25	16
532.0	15	*8	*7	*	*	*	10	*	*	*6	*
532.00	15	*8	*7	*	*	*	10	*	*	*6	*
532.4	40	27	13	—	*7	10	23	*9	*7	13	11
532.40	38	26	12	—	*7	9	22	*9	*7	12	10
532.5	*5	*	*	—	*	*	*	*	*	*	*
532.50	*5	*	*	—	*	*	*	*	*	*	*
532.9	*7	*	*	—	*	*	*	*	*	*	*
532.90	*5	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
533	16	*6	9	*	*	*	*9	*	*	*8	*
533.4	*6	*	*	—	*	*	*	*	*	*	*
533.40	*6	*	*	—	*	*	*	*	*	*	*
533.9	*5	*	*	*	*	*	*	*	*	*	*
533.90	*5	*	*	*	*	*	*	*	*	*	*
535	151	55	96	*	30	38	78	34	37	60	19
535.0	30	11	19	*	*7	*6	14	*7	*8	12	*
535.00	21	*	16	*	*7	*	*8	*	*5	10	*
535.01	*9	*6	*	*	*	*	*6	*	*	*	*
535.1	15	*	12	*	*	*	*9	*	*	*6	*
535.10	10	*	*9	*	*	*	*	*	*	*	*
535.11	*5	*	*	*	*	*	*	*	*	*	*
535.3	*6	*5	*	—	*	*	*	*	*	*	*
535.4	34	14	20	*	*	*9	20	*7	13	10	*
535.40	19	*6	14	*	*	*	11	*	*7	*5	*
535.41	15	*8	*7	*	*	*	9	*	*5	*	*
535.5	53	14	39	*	9	13	29	12	*9	27	*
535.50	37	*8	29	*	*8	11	17	*7	*6	20	*
535.51	16	*6	10	*	*	*	12	*5	*	*7	*
535.6	12	*7	*	*	*	*	*	*	*	*	*
535.60	*7	*	*	*	*	*	*	*	*	*	*
536	50	22	28	*7	12	12	19	*8	10	22	9
536.2	11	*5	*6	*	*	*	*	*	*	*5	*
536.3	10	*	*9	*	*	*	*	*	*	*	*
536.4	13	*6	*6	*	*	*	*9	*	*	*	*
536.42	*5	*	*	*	*	*	*	*	*	*	*
536.8	13	*8	*	*	*	*	*	*	*	*6	*
537	28	10	18	*	*	*	20	*8	*8	*7	*6
537.0	10	*	*6	*	*	*	*7	*	*	*	*
537.8	14	*5	*9	*	*	*	11	*	*	*	*
537.83	12	*	*8	—	*	*	10	*	*	*	*
540	255	146	109	62	136	41	16	54	60	77	65
540.0	53	30	23	16	23	10	*	10	14	15	14
540.1	33	17	16	*6	15	*8	*5	*6	9	11	*7
540.9	169	99	70	41	98	23	*7	38	37	51	44
541	10	*6	*	*	*6	*	*	*	*	*5	*
543	*5	*	*	*	*	*	*	*	*	*	*
550	47	40	*7	*5	*7	11	24	13	10	13	10
550.1	22	18	*	*	*	*	11	*8	*	*	*5
550.10	17	14	*	*	*	*	*8	*6	*	*	*
550.9	24	21	*	*	*	*6	13	*6	*5	*8	*5
550.90	15	12	*	*	*	*	*8	*	*	*	*
550.92	*6	*6	*	*	*	*	*	*	*	*	*
552	49	15	34	*	*6	19	24	15	11	17	*7
552.1	10	*	*7	*	*	*	*	*	*	*	*
552.2	30	9	20	—	*	12	15	10	*6	10	*
552.20	*8	*	*6	—	*	*	*7	*	*	*	*
552.21	20	*7	13	—	*	10	*8	*6	*	*7	*
553	75	29	45	*	16	27	30	15	16	35	9
553.1	*5	*	*	*	*	*	*	*	*	*	*
553.2	51	21	30	*	9	20	22	10	11	23	*7
553.20	9	*5	*	*	*	*	*	*	*	*	*
553.21	40	15	24	*	*6	14	19	*6	10	18	*5
553.3	18	*5	13	*	*	*7	*7	*	*	*8	*
555	45	19	27	*	27	11	*6	13	14	10	*8
555.0	15	*6	9	*	*9	*	*	*	*5	*	*
555.1	*8	*	*	*	*	*	*	*	*	*	*
555.2	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
555.9	16	*7	9	*	10	*	*	*	*6	*	*
556	28	14	14	*	13	*7	*8	*7	*8	*8	*
556.9	14	*5	*8	*	*7	*	*	*	*	*	*
557	44	10	35	*	*	*9	30	*8	13	16	*8
557.0	22	*6	16	*	*	*	15	*	*7	*6	*
557.9	20	*	16	*	*	*	13	*	*	*9	*
558	160	63	96	45	37	30	47	30	33	69	28
558.1	*5	*	*	—	*	*	*	*	*	*	*
558.9	154	61	93	45	37	29	42	28	32	67	27
560	275	100	175	10	35	74	156	63	63	101	48
560.1	45	20	26	*	*8	10	26	12	*9	21	*
560.2	11	*5	*6	*	*	*	*8	*	*	*	*
560.3	34	10	25	*	*	*	25	*8	*8	15	*
560.39	30	9	21	*	*	*	22	*6	*8	12	*
560.8	75	27	48	*	14	21	38	13	19	26	17
560.81	52	17	35	*	9	17	25	*9	13	19	12
560.89	23	10	13	*	*	*	13	*	*6	*7	*
560.9	107	38	69	*	10	37	57	26	24	34	22
562	264	105	159	—	30	70	164	63	67	95	39
562.1	261	104	157	—	29	69	162	63	65	95	39
562.10	36	12	23	—	*	*9	23	*8	*7	17	*
562.11	142	59	83	—	23	51	67	34	40	44	23
562.12	67	26	41	—	*	*7	60	18	12	27	10
562.13	16	*6	10	—	*	*	12	*	*5	*7	*
564	63	22	40	9	13	14	26	12	20	18	13
564.0	36	14	22	*7	*	*6	18	*6	11	10	*8
564.1	17	*	14	*	*7	*	*5	*	*	*5	*
566	17	12	*	*	*7	*6	*	*	*	*7	*
567	19	11	*8	*	*	*7	*6	*	*	*8	*
567.2	12	*8	*	*	*	*6	*	*	*	*	*
568	15	*	13	*	*5	*5	*	*	*	*6	*
568.0	11	*	10	*	*	*	*	*	*	*5	*
569	96	42	54	*	11	21	62	24	18	33	21
569.1	*7	*	*6	*	*	*	*	*	*	*	*
569.3	16	9	*7	*	*	*	11	*	*	*6	*
569.4	10	*	*5	*	*	*	*6	*	*	*	*
569.49	*5	*	*	*	*	*	*	*	*	*	*
569.6	16	*7	9	*	*	*	*7	*	*	*6	*
569.69	10	*	*6	*	*	*	*6	*	*	*	*
569.8	40	16	24	*	*	*9	28	11	*7	13	*8
569.81	*7	*	*	*	*	*	*	*	*	*	*
569.82	*6	*	*	*	*	*	*	*	*	*	*
569.83	*7	*	*5	*	*	*	*	*	*	*	*
569.85	16	*7	*9	—	—	*	14	*	*	*	*
570	*5	*	*	*	*	*	*	*	*	*	*
571	82	52	31	*	19	43	20	19	17	32	14
571.1	13	*9	*	*	*5	*7	*	*	*	*7	*
571.2	36	27	9	—	*8	21	*7	9	*8	12	*7
571.5	27	13	14	*	*	12	11	*7	*6	10	*
572	35	19	16	*	*	19	12	12	*	12	*8
572.2	23	11	12	—	*	12	*8	*8	*	*9	*
572.3	*5	*	*	—	*	*	*	*	*	*	*
573	11	*5	*6	*	*5	*	*	*	*	*	*
573.3	*7	*	*	*	*	*	*	*	*	*	*
574	344	111	233	*	108	107	127	79	73	131	61
574.0	108	44	64	*	34	33	40	27	25	39	17

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
574.00	100	40	60	*	32	31	36	26	22	38	14
574.01	*8	*	*	*	*	*	*	*	*	*	*
574.1	144	37	107	*	48	49	46	34	24	60	26
574.10	139	35	104	*	47	47	44	33	23	59	24
574.2	37	12	25	*	10	13	14	*7	9	15	*6
574.20	35	10	25	*	9	13	13	*6	*9	14	*6
574.5	16	*	11	*	*	*	10	*	*	*	*
574.50	11	*	*8	*	*	*	*7	*	*	*	*
574.7	15	*	11	*	*7	*	*5	*	*6	*	*
574.70	*8	*	*6	—	*	*	*	*	*	*	*
574.71	*7	*	*	*	*	*	*	*	*	*	*
574.8	*8	*	*	—	*	*	*	*	*	*	*
574.80	*6	*	*	—	*	*	*	*	*	*	*
575	68	26	42	*	25	21	22	13	15	33	*8
575.0	23	12	11	—	*7	*8	*9	*	*	10	*
575.1	36	11	25	*	14	11	11	*7	*6	20	*
575.10	*6	*	*	—	*	*	*	*	*	*	*
575.11	23	*5	18	*	10	*8	*5	*5	*	12	*
575.12	*7	*	*	—	*	*	*	*	*	*	*
575.8	*5	*	*	*	*	*	*	*	*	*	—
576	31	13	18	*	*	9	16	*9	*9	*8	*6
576.1	12	*6	*6	*	*	*	*5	*	*	*	*
576.2	10	*	*6	*	*	*	*5	*	*	*	*
576.8	*7	*	*	*	*	*	*	*	*	*	*
577	226	117	110	*5	82	82	57	43	51	90	42
577.0	195	102	93	*	74	67	49	38	44	77	37
577.1	23	11	12	*	*7	12	*	*	*	12	*
578	195	93	102	*	15	41	135	45	50	68	33
578.0	23	12	11	*	*	*6	13	*7	*6	*6	*
578.1	33	17	17	*	*	*8	23	*9	*9	9	*7
578.9	139	64	74	*	10	28	99	29	35	52	22
579	*6	*	*	*	*	*	*	*	*	*	*
580-629	1,743	533	1,211	83	574	436	650	367	392	692	292
581	*6	*	*	*	*	*	*	*	*	*	*
584	116	60	56	*	9	28	78	39	19	35	23
584.5	*7	*	*	*	*	*	*	*	*	*	*
584.9	106	53	53	*	*8	25	73	36	18	33	19
585	31	17	14	*	*5	9	16	*7	*7	14	*
586	*8	*	*	—	*	*	*5	*	*	*	*
590	130	21	109	26	54	24	26	22	31	46	31
590.1	75	11	64	12	33	14	16	14	20	24	17
590.10	75	11	64	12	33	14	16	14	20	24	17
590.8	51	*9	42	13	19	10	9	*8	11	20	13
590.80	51	*9	42	13	19	10	9	*8	11	20	13
591	*7	*	*	*	*	*	*	*	*	*	*
592	177	104	73	*	80	62	31	38	45	78	15
592.0	47	21	26	*	22	15	9	12	11	20	*
592.1	129	83	46	*	59	46	22	26	34	58	11
593	41	16	25	*8	10	10	14	*8	*8	18	*7
593.4	10	*	*5	*	*	*	*	*	*	*6	*
593.7	*6	*	*5	*6	*	*	—	*	*	*	*
593.9	15	*7	*8	*	*	*	*8	*	*	*8	*
594	*8	*7	*	*	*	*	*6	*	*	*	*
594.1	*7	*7	*	*	*	*	*5	*	*	*	*
595	19	*7	11	*	*	*	12	*5	*	*6	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
595.9	*8	*	*	*	*	*	*5	*	*	*	*
596	22	14	*8	*	*	*	13	*	*	11	*
596.0	*9	*8	*	—	—	*	*7	*	*	*	*
596.5	*6	*	*	*	*	*	*	*	*	*	*
596.54	*5	*	*	*	*	*	*	*	*	*	*
598	*7	*6	*	*	*	*	*	*	*	*	*
599	384	131	253	31	38	56	260	85	77	158	64
599.0	339	106	233	30	30	43	235	74	67	142	55
599.7	31	19	13	*	*	*8	17	*9	*8	10	*
599.8	*6	*	*5	—	*	*	*	*	*	*	*
600	105	105	—	—	*	18	86	22	26	39	18
601	*6	*6	—	—	*	*	*	*	*	*	*
604	*7	*7	—	*	*	*	*	*	*	*	*
604.9	*7	*7	—	*	*	*	*	*	*	*	*
604.90	*5	*5	—	*	*	*	*	*	*	*	*
608	*7	*7	—	*	*	*	*	*	*	*	*
611	28	*	28	*	18	*7	*	11	*	*8	*5
611.0	*5	*	*	*	*	*	*	*	*	*	*
611.1	16	*	16	*	12	*	*	*8	*	*	*
614	50	—	50	*	42	*6	*	10	10	25	*
614.2	*6	—	*6	*	*	*	*	*	*	*	*
614.6	15	—	15	—	12	*	*	*	*	*8	*
614.9	18	—	18	*	18	*	*	*	*6	*8	*
616	13	—	13	*	*8	*	*	*	*	*6	*
616.0	*9	—	*9	—	*6	*	*	*	*	*	*
617	74	—	74	—	59	15	*	15	19	26	14
617.0	42	—	42	—	31	10	*	*7	11	16	*8
617.1	13	—	13	—	10	*	*	*	*	*	*
617.3	13	—	13	—	11	*	*	*	*	*	*
618	151	—	151	—	33	65	53	26	39	61	25
618.0	35	—	35	—	*5	18	12	*6	*9	15	*5
618.1	20	—	20	—	10	*7	*	*	*	*9	*
618.2	17	—	17	—	*5	*8	*	*	*	*	*
618.3	17	—	17	—	*	*7	*7	*	*	*8	*
618.4	21	—	21	—	*	9	*7	*	*5	9	*
618.5	26	—	26	—	*	10	15	*	*7	11	*
618.6	*6	—	*6	—	*	*	*	*	*	*	*
618.8	9	—	9	—	*	*	*	*	*	*	*
619	*5	—	*5	—	*	*	*	*	*	*	*
620	74	—	74	*	48	17	*6	12	18	30	13
620.0	*6	—	*6	*	*	*	—	*	*	*	*
620.1	14	—	14	*	12	*	—	*	*	*	*
620.2	45	—	45	*	25	12	*6	*9	11	18	*7
620.8	*5	—	*5	*	*	*	*	*	*	*	*
621	17	—	17	—	*7	*7	*	*	*	*5	*
621.3	*6	—	*6	—	*	*	*	*	*	*	*
622	*9	—	*9	—	*5	*	*	*	*	*	*
622.1	*8	—	*8	—	*	*	*	*	*	*	*
625	97	—	97	*	53	32	12	14	17	45	20
625.3	14	—	14	—	10	*	—	*	*	*8	*
625.6	45	—	45	—	13	20	12	*7	10	19	10
625.9	32	—	32	*	25	*7	*	*6	*	16	*6
626	101	—	101	*	68	33	*	12	27	46	16
626.2	72	—	72	*	49	23	—	9	21	29	13
626.8	23	—	23	*	17	*6	*	*	*5	14	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
627	13	—	13	—	*	*9	*	*	*	*	*
627.1	10	—	10	—	—	*8	*	*	*	*	*
630-677	504	—	504	*	503	*	—	99	93	211	102
632	14	—	14	—	14	*	—	*	*	*5	*
633	34	—	34	—	34	*	—	9	*5	11	*9
633.1	30	—	30	—	29	*	—	*7	*	10	*8
634	26	—	26	*	26	*	—	*7	*	12	*6
634.9	19	—	19	*	19	—	—	*5	*	*7	*
634.91	14	—	14	—	14	—	—	*5	*	*5	*
637	*7	—	*7	—	*7	—	—	*	*	*	*
637.9	*7	—	*7	—	*7	—	—	*	*	*	*
640	*7	—	*7	—	*7	—	—	*	*	*	*
641	13	—	13	—	13	*	—	*	*	*	*
641.1	*6	—	*6	—	*6	*	—	*	*	*	*
641.13	*6	—	*6	—	*6	*	—	*	*	*	*
642	27	—	27	*	26	—	—	*7	*	12	*
642.3	10	—	10	—	10	—	—	*	*	*5	*
642.33	10	—	10	—	10	—	—	*	*	*5	*
642.4	*8	—	*8	*	*8	—	—	*	*	*	*
642.43	*7	—	*7	*	*7	—	—	*	*	*	*
643	49	—	49	—	49	—	—	*6	15	21	*7
643.0	22	—	22	—	22	—	—	*	*7	12	*
643.03	22	—	22	—	22	—	—	*	*7	12	*
643.1	19	—	19	—	19	—	—	*	*	*7	*
643.13	19	—	19	—	19	—	—	*	*	*7	*
643.2	*6	—	*6	—	*6	—	—	*	*	*	*
643.23	*6	—	*6	—	*6	—	—	*	*	*	*
644	114	—	114	—	114	—	—	23	23	45	23
644.0	94	—	94	—	94	—	—	20	20	33	20
644.03	93	—	93	—	93	—	—	20	20	33	20
644.1	20	—	20	—	20	—	—	*	*	12	*
644.13	20	—	20	—	20	—	—	*	*	12	*
646	57	—	57	*	57	—	—	*8	*9	28	11
646.6	32	—	32	*	32	—	—	*	*	14	*8
646.63	29	—	29	*	29	—	—	*	*	13	*7
646.8	20	—	20	—	20	—	—	*	*	12	*
646.83	19	—	19	—	19	—	—	*	*	12	*
647	*7	—	*7	*	*7	—	—	*	*	*	*
648	73	—	73	*	73	*	—	14	13	32	15
648.0	*9	—	*9	—	*9	—	—	*	*	*	*
648.03	*8	—	*8	—	*8	—	—	*	*	*	*
648.8	11	—	11	—	11	*	—	*	*	*7	*
648.83	11	—	11	—	11	*	—	*	*	*7	*
648.9	44	—	44	*	44	—	—	9	*8	17	10
648.93	43	—	43	*	43	—	—	*9	*8	17	9
654	12	—	12	—	12	*	—	*	*	*6	*
654.5	*5	—	*5	—	*5	*	—	*	*	*	*
654.53	*5	—	*5	—	*5	*	—	*	*	*	*
658	*7	—	*7	—	*7	*	—	*	*	*	*
659	*8	—	*8	—	*8	*	—	*	*	*	*
670	*7	—	*7	—	*7	—	—	*	*	*	*
670.0	*7	—	*7	—	*7	—	—	*	*	*	*
670.04	*7	—	*7	—	*7	—	—	*	*	*	*
674	*7	—	*7	—	*7	—	—	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
680-709	514	259	255	70	132	133	179	115	115	210	74
681	20	13	*7	*	*6	*	*6	*	*7	*7	*
681.0	11	*8	*	*	*5	*	*	*	*	*	*
681.00	10	*7	*	*	*	*	*	*	*	*	*
681.1	9	*5	*	*	*	*	*	*	*	*	*
681.10	9	*5	*	*	*	*	*	*	*	*	*
682	333	171	161	24	90	101	118	90	80	110	52
682.0	20	12	*8	*6	*6	*	*	*5	*	*7	*
682.2	25	14	12	*	*8	*7	*7	*6	*5	9	*
682.3	34	15	19	*	12	10	10	11	*8	10	*5
682.4	23	13	9	*	*9	*7	*	*8	*6	*6	*
682.5	*6	*	*	*	*	*	*	*	*	*	*
682.6	172	85	87	*5	40	54	73	45	43	56	28
682.7	41	22	19	*	*8	15	16	9	10	14	*8
682.9	*5	*	*	*	*	*	*	*	*	*	*
693	*8	*	*	*	*	*	*	*	*	*	*
693.0	*7	*	*	*	*	*	*	*	*	*	*
695	*7	*	*	*	*	*	*	*	*	*	*
695.1	*5	*	*	*	*	*	*	*	*	*	*
707	71	31	40	*	12	14	44	12	18	32	9
707.0	37	15	22	*	*7	*6	24	*6	*8	19	*
707.1	29	14	15	*	*	*7	18	*6	*8	11	*
709	37	18	19	27	*8	*	*	*	*	35	*
709.2	30	15	15	21	*8	*	*	*	*	29	*
709.8	*7	*	*	*5	*	*	*	*	*	*6	*
710-739	1,530	666	865	37	322	497	674	304	376	565	285
710	23	*	19	*	11	*8	*	*	*5	*8	*
710.0	17	*	14	*	10	*	*	*	*	*6	*
711	23	14	*9	*	*5	*6	*8	*6	*6	*8	*
711.0	22	13	*8	*	*5	*6	*8	*5	*6	*8	*
711.06	*8	*6	*	*	*	*	*	*	*	*	*
714	19	*	15	*	*	*8	*9	*	*	*7	*
714.0	15	*	12	—	*	*6	*7	*	*	*6	*
715	441	175	267	*	13	125	303	81	117	161	82
715.1	39	15	24	*	*	13	25	9	*9	13	*8
715.15	9	*	*6	—	*	*	*7	*	*	*	*
715.16	27	9	18	*	*	10	17	*6	*7	9	*5
715.2	*8	*	*5	—	*	*	*	*	*	*	*
715.3	205	81	124	*	*7	62	137	33	58	79	36
715.35	56	23	33	—	*	20	33	11	15	18	11
715.36	142	54	88	*	*	38	101	21	41	58	23
715.8	*5	*	*	—	*	*	*	*	*	*	*
715.89	*5	*	*	—	*	*	*	*	*	*	*
715.9	182	74	108	—	*	47	132	35	47	67	34
715.90	*8	*	*	—	*	*	*6	*	*	*6	*
715.95	54	24	30	—	*	16	35	13	13	18	9
715.96	113	44	69	—	*	27	86	19	31	41	22
716	20	10	11	*	*	10	*7	*	*	9	*
716.1	*7	*	*	—	*	*	*	*	*	*	*
716.9	11	*	*7	*	*	*5	*	*	*	*	*
717	12	*6	*6	*	9	*	*	*	*	*	*
717.8	*7	*	*	*	*7	—	*	*	*	*	*
717.83	*7	*	*	*	*6	—	—	*	*	*	*
718	15	*6	*9	*	10	*	*	*	*	*	*
718.3	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
718.8	*5	*	*	*	*	*	*	*	*	*	*
719	42	16	27	*	*9	13	19	*8	*8	15	12
719.4	33	11	22	*	*7	10	16	*6	*7	11	9
719.41	9	*	*6	—	*	*	*	*	*	*	*
719.45	14	*	11	*	*	*	11	*	*	*	*
719.46	*5	*	*	*	*	*	*	*	*	*	*
721	67	31	36	—	10	24	32	11	17	23	15
721.0	23	10	13	—	*5	11	*6	*	*	11	*5
721.1	12	*8	*	—	*	*5	*6	*	*	*	*
721.3	24	*9	15	—	*	*6	16	*6	*6	*7	*
722	329	175	154	*	140	132	56	62	80	130	57
722.0	56	29	27	—	25	29	*	*9	11	25	11
722.1	188	104	84	*	85	69	34	39	44	75	29
722.10	187	104	83	*	85	68	33	39	44	74	29
722.2	*6	*	*	—	*	*	*	*	*	*	—
722.4	*7	*	*	—	*	*	*	*	*	*	*
722.5	34	15	19	—	11	13	11	*5	12	10	*7
722.52	34	15	19	—	11	13	11	*5	12	10	*7
722.7	18	9	9	—	*7	*8	*	*	*	*6	*
722.71	11	*6	*5	—	*	*6	*	*	*	*	*
722.73	*5	*	*	—	*	*	*	*	*	*	*
722.8	*7	*	*	*	*	*	*	*	*	*	*
722.83	*7	*	*	*	*	*	*	*	*	*	*
722.9	11	*6	*	*	*	*	*	*	*	*	*
722.93	*8	*	*	*	*	*	*	*	*	*	*
723	21	9	12	*	*5	11	*	*	*	*7	*6
723.0	*6	*	*	—	*	*	*	*	*	*	*
723.1	*6	*	*	*	*	*	*	*	*	*	*
723.4	*7	*	*	—	*	*6	*	*	*	*	*
724	134	59	75	*	23	45	65	26	34	46	29
724.0	65	33	33	*	*	25	36	13	13	24	15
724.02	61	31	29	*	*	23	34	12	13	21	15
724.2	35	12	23	*	9	10	15	*	12	11	*8
724.3	*5	*	*	—	*	*	*	*	*	*	*
724.4	9	*5	*	*	*	*	*	*	*	*	*
724.5	13	*	10	*	*	*	*7	*	*	*	*
726	45	21	24	*	10	19	15	*7	10	22	*6
726.1	18	*8	10	—	*	*7	*9	*	*	*7	*
726.10	16	*8	*9	—	*	*6	*9	*	*	*7	*
726.2	13	*	*8	—	*	*7	*	*	*	10	*
726.3	*5	*	*	—	*	*	*	*	*	*	*
727	22	10	12	*	*	*8	*9	*	*	11	*
727.0	*6	*	*	*	*	*	*	*	*	*	*
727.6	10	*	*6	—	*	*	*	*	*	*8	*
727.61	9	*	*5	—	*	*	*	*	*	*7	*
728	19	10	10	*	*	*	10	*	*	*6	*
728.8	12	*6	*6	*	*	*	*7	*	*	*	*
728.89	9	*5	*	*	*	*	*6	*	*	*	*
729	43	20	23	*	12	14	15	9	10	18	*6
729.1	12	*	9	*	*	*	*	*	*	*	*
729.5	14	*8	*6	*	*	*5	*	*	*	*6	*
729.8	13	*6	*7	*	*	*	*	*	*	*5	*
729.81	*7	*	*	—	*	*	*	*	*	*	*
729.89	*6	*	*	*	*	*	*	*	*	*	*
730	51	31	20	*	10	17	20	14	11	17	*9
730.0	19	12	*7	*	*	*7	*6	*6	*	*6	*
730.07	11	*8	*	*	*	*	*5	*	*	*	*
730.1	*9	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
730.2	23	13	10	*	*6	*7	10	*7	*5	*7	*
730.27	15	*6	*8	*	*	*	*7	*5	*	*	*
732	*6	*	*	*	*	*	*	*	*	*	*
733	142	44	98	*	28	32	79	29	37	52	24
733.1	69	16	53	*	*	12	54	15	20	23	11
733.13	44	*8	35	—	*	*	38	9	13	13	*8
733.14	13	*	9	*	*	*	*8	*	*	*6	*
733.19	*6	*	*5	—	—	*	*	*	*	*	*
733.4	23	12	10	*	*8	*8	*7	*	*5	11	*
733.42	21	12	9	*	*7	*7	*7	*	*	10	*
733.6	17	*	14	*	*	*	*8	*	*5	*7	*
733.8	24	10	14	*	10	*7	*7	*	*5	*7	*7
733.81	*5	*	*	*	*	*	*	*	*	*	*
733.82	19	*7	12	*	*7	*6	*6	*	*	*6	*
736	10	*5	*	*	*	*	*	*	*	*	*
737	13	*	11	*7	*	*	*	*	*	*	*
737.3	12	*	11	*7	*	*	*	*	*	*	*
737.30	*9	*	*8	*	*	*	*	*	*	*	*
738	22	*7	15	*	*8	*9	*	*5	*	*8	*
738.4	16	*	11	—	*	*7	*	*5	*	*	*
740-759	165	94	71	113	33	11	*9	41	28	57	38
742	*7	*	*	*5	*	*	—	*	*	*	*
745	17	*8	9	14	*	*	*	*5	*	*	*6
746	12	9	*	*7	*	*	*	*	*	*	*
747	15	*8	*7	*7	*	*	*	*	*	*6	*
747.6	*6	*	*	*	*	*	*	*	*	*	*
749	13	*8	*5	12	*	—	—	*	*	*5	*
749.0	*6	*	*	*6	*	—	—	*	*	*	*
750	13	11	*	13	*	—	*	*	*	*	*
750.5	11	9	*	11	—	—	—	*	*	*	*
751	12	*7	*5	10	*	*	*	*	*	*	*
752	*5	*	*	*	*	*	*	*	*	*	*
753	9	*	*	*	*	*	*	*	*	*	*
754	9	*	*	*8	*	*	—	*	*	*	*
756	20	12	*9	9	*5	*	*	*	*	*8	*6
756.1	10	*7	*	*	*	*	*	*	*	*	*
756.12	*7	*	*	—	*	*	*	*	*	*	*
757	11	*	*7	*6	*	*	—	*	*	10	—
757.3	10	*	*6	*6	*	—	—	*	*	9	—
757.32	10	*	*6	*6	*	—	—	*	—	9	—
760-779	156	86	69	155	*	—	—	26	31	42	56
765	21	12	9	21	—	—	—	*	*	10	*
765.1	17	9	*7	17	—	—	—	*	*	*8	*
769	12	*7	*	12	—	—	—	*	*	*	*
770	32	17	15	32	—	—	—	*6	*6	9	12
770.6	*7	*	*	*7	—	—	—	*	*	*	*
770.8	17	*9	*8	17	—	—	—	*	*	*	*7
771	14	*9	*5	14	—	—	—	*	*	*	*6
771.8	11	*8	*	11	—	—	—	*	*	*	*
774	37	20	17	37	—	—	—	*5	*6	*8	18
774.6	29	15	14	29	—	—	—	*	*	*6	14

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
775	*5	*	*	*5	—	—	—	*	*	*	*
777	*6	*	*	*6	—	—	—	*	*	*	*
778	*6	*	*	*6	—	—	—	*	*	*	*
779	12	*7	*	12	—	—	—	*	*	*	*
779.3	*7	*	*	*7	—	—	—	*	*	*	*
780-799	267	127	139	56	93	74	44	63	53	119	32
780	64	30	34	25	20	11	*7	17	13	27	*7
780.2	12	*	*7	*	*	*	*	*	*	*7	*
780.3	28	14	13	15	10	*	*	*7	*	12	*
780.39	25	13	12	12	10	*	*	*7	*	11	*
780.6	14	*7	*7	9	*	*	*	*5	*	*	*
784	*8	*	*	*	*	*	*	*	*	*	*
786	81	43	38	*	27	39	12	18	18	38	*8
786.5	75	40	35	*	27	38	11	17	17	35	*7
786.50	46	23	22	*	14	24	*7	10	11	21	*
786.59	26	16	11	*	10	13	*	*7	*	12	*
787	*5	*	*	*	*	*	*	*	*	*	*
788	*6	*	*	*	*	*	*	*	*	*	*
789	45	17	28	12	22	*7	*	11	10	18	*6
789.0	45	17	27	12	21	*7	*	11	10	17	*6
789.00	10	*	*6	*	*	*	*	*	*	*6	*
789.01	*5	*	*	*	*	*	*	*	*	*	*
789.03	15	*6	*9	*	10	*	*	*5	*	*	*
799	49	24	25	*8	14	11	16	10	*6	27	*6
799.9	49	24	25	*8	14	11	16	10	*6	27	*6
800-999	2,466	1,214	1,252	228	712	514	1,012	527	580	907	453
800	10	*7	*	*7	*	*	*	*	*	*	*
801	25	18	*7	*7	11	*	*	*5	*	*6	*8
801.0	*9	*5	*	*	*	*	*	*	*	*	*
801.1	*6	*	*	*	*	*	*	*	*	*	*
801.2	*6	*	*	*	*	*	*	*	*	*	*
802	31	24	*7	*	21	*	*	*6	*5	13	*7
802.2	11	*9	*	*	*8	*	*	*	*	*	*
802.4	*8	*7	*	*	*6	*	*	*	*	*5	*
805	67	31	36	*	21	9	35	11	18	24	14
805.0	14	*7	*6	*	*5	*	*5	*	*	*5	*
805.02	*5	*	*	*	*	*	*	*	*	*	*
805.2	18	*7	11	*	*	*	11	*	*	*7	*
805.4	32	15	17	*	10	*	17	*6	10	10	*6
807	48	24	24	*	10	11	25	*7	12	22	*7
807.0	39	19	21	*	*8	9	21	*5	9	18	*7
807.01	*8	*	*	*	*	*	*	*	*	*	*
807.02	*6	*	*	—	*	*	*	*	*	*	*
807.03	*7	*	*	—	*	*	*	*	*	*	*
807.05	*5	*	*	*	*	*	*	*	*	*	*
807.2	*5	*	*	—	*	*	*	*	*	*	*
808	54	18	36	*	9	*	39	10	14	24	*6
808.0	11	*8	*	*	*	*	*	*	*	*	*
808.2	35	*7	28	*	*	*	30	*6	*8	17	*
810	*7	*	*	*	*	*	*	*	*	*	*
810.0	*6	*	*	*	*	*	*	*	*	*	*
812	59	22	38	12	10	*9	29	13	15	19	12

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
812.0	28	*7	21	*	*	*6	21	*7	*9	*9	*
812.00	9	*	*7	*	*	*	*6	*	*	*	*
812.01	11	*	*9	—	*	*	*9	*	*	*	*
812.09	*6	*	*	*	*	*	*	*	*	*	*
812.2	12	*	*8	*	*	*	*6	*	*	*5	*
812.21	*8	*	*6	*	*	*	*	*	*	*	*
812.4	15	*8	*7	11	*	*	*	*	*	*	*
812.41	12	*6	*6	*9	*	*	*	*	*	*	*
813	64	31	33	14	20	13	17	14	12	24	14
813.0	15	9	*6	*	*6	*	*	*5	*	*6	*
813.01	*8	*6	*	*	*	*	*	*	*	*	*
813.2	*8	*5	*	*	*	*	*	*	*	*	*
813.4	27	10	17	*	*8	*5	9	*	*7	11	*5
813.41	*7	*	*5	*	*	*	*	*	*	*	*
813.42	*9	*	*5	*	*	*	*	*	*	*	*
813.44	11	*	*6	*	*	*	*	*	*	*	*
813.5	*5	*	*	*	*	*	*	*	*	*	*
816	9	*8	*	*	*	*	*	*	*	*	*
816.1	*7	*7	*	*	*	*	*	*	*	*	*
820	321	81	240	*	*6	24	290	63	74	127	57
820.0	92	19	73	*	*	*8	81	17	24	32	19
820.00	11	*	9	—	*	*	11	*	*	*	*
820.02	15	*	12	—	*	*	12	*	*	10	*
820.03	*7	*	*	*	*	*	*6	*	*	*	*
820.09	57	11	47	*	*	*	53	12	17	16	12
820.2	155	43	111	*	*	9	142	33	35	61	25
820.20	10	*	*8	—	*	*	9	*	*	*	*
820.21	134	39	95	*	*	*7	125	26	31	57	20
820.22	11	*	*8	*	*	*	*8	*	*	*	*
820.8	72	18	54	*	*	*7	63	13	14	33	12
821	56	25	30	11	14	*8	23	10	13	19	15
821.0	28	17	11	*9	*9	*	*7	*	*7	*8	*9
821.00	*6	*	*	*	*	*	*	*	*	*	*
821.01	22	14	*9	*8	*8	*	*	*	*6	*7	*7
821.2	22	*6	17	*	*	*	16	*	*	*9	*5
821.20	*7	*	*	*	*	*	*	*	*	*	*
821.23	9	*	*8	*	*	*	*7	*	*	*	*
822	15	*7	*8	—	*	*6	*6	*	*	*	*
822.0	14	*6	*8	—	*	*6	*6	*	*	*	*
823	69	41	28	*	27	23	14	15	15	26	13
823.0	25	13	12	*	*9	*8	*7	*	*7	*8	*5
823.00	19	10	*9	*	*7	*6	*6	*	*6	*5	*
823.2	19	11	*8	*	*7	*8	*	*	*	*7	*6
823.22	15	*9	*6	*	*5	*6	*	*	*	*	*5
823.3	*5	*	*	*	*	*	*	*	*	*	*
823.8	13	*8	*5	*	*7	*	*	*5	*	*6	*
823.82	9	*	*	*	*	*	*	*	*	*	*
824	111	47	63	*	46	34	26	24	25	43	18
824.0	*6	*	*	*	*	*	*	*	*	*	*
824.2	*9	*6	*	*	*	*	*	*	*	*	*
824.4	33	13	20	*	13	12	*7	*	9	14	*
824.6	30	*8	22	*	10	*9	11	9	*6	12	*
824.8	25	14	11	*	13	*6	*	*6	*	*9	*6
825	17	13	*	*	*7	*6	*	*	*	*7	*
825.0	*7	*5	*	—	*	*	*	*	*	*	*
825.2	*8	*6	*	*	*	*	*	*	*	*	*
825.25	*5	*	*	*	*	*	*	*	*	*	*
836	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
840	20	11	9	—	*	*6	12	*	*8	*6	*
840.4	18	9	*9	—	*	*5	12	*	*7	*5	*
844	16	12	*	*	12	*	*	*	*	*6	*
844.2	*7	*	*	—	*7	*	—	*	*	*	*
844.8	*6	*6	*	—	*	*	*	*	*	*	*
847	15	*6	*9	*	*5	*	*7	*	*	*6	*
847.2	*7	*	*	—	*	*	*	*	*	*	*
850	36	23	12	10	14	*	*8	12	*7	10	*7
850.0	*7	*	*	*	*	*	*	*	*	*	*
850.1	13	*9	*	*6	*	*	*	*	*	*	*
850.5	*8	*6	*	*	*	*	*	*	*	*	*
850.9	*6	*	*	*	*	*	*	*	*	*	*
851	17	12	*6	*	*8	*	*6	*	*	*6	*
851.8	12	*7	*	*	*6	*	*	*	*	*5	*
852	33	20	14	*	*9	*6	16	*7	*8	9	9
852.0	*7	*	*	*	*	*	*	*	*	*	*
852.2	24	14	10	*	*5	*	13	*5	*	*6	*8
852.20	*8	*	*	*	*	*	*6	*	*	*	*
852.21	*7	*	*	*	*	*	*	*	*	*	*
853	*8	*	*	*	*	*	*	*	*	*	*
853.0	*8	*	*	*	*	*	*	*	*	*	*
854	10	*6	*	*	*	*	*	*	*	*	*
854.0	*9	*6	*	*	*	*	*	*	*	*	*
860	25	16	9	*	14	*	*7	*	*6	10	*6
860.0	16	*9	*7	*	*9	*	*	*	*	*6	*
861	*8	*5	*	*	*	*	*	*	*	*	*
861.2	*6	*	*	*	*	*	*	*	*	*	*
861.21	*6	*	*	*	*	*	*	*	*	*	*
863	10	*7	*	*	*6	*	*	*	*	*	*
864	*8	*6	*	*	*	*	*	*	*	*	*
864.0	*6	*	*	*	*	*	*	*	*	*	*
865	15	11	*	*	9	*	*	*	*	*	*
865.0	14	11	*	*	*9	*	*	*	*	*	*
868	*7	*5	*	*	*	*	*	*	*	*	*
873	22	13	9	*	10	*	*6	*6	*	*5	*6
873.0	*6	*	*	*	*	*	*	*	*	*	*
873.4	11	*6	*5	*	*	*	*	*	*	*	*
873.42	*7	*	*	*	*	*	*	*	*	*	*
881	*8	*6	*	*	*5	*	*	*	*	*	*
883	*8	*5	*	*	*6	*	*	*	*	*	*
886	*6	*5	*	*	*	*	*	*	*	*	*
886.0	*5	*	*	*	*	*	*	*	*	*	*
891	12	*8	*	*	*5	*	*	*	*	*6	*
891.0	*6	*	*	*	*	*	*	*	*	*	*
920	*9	*	*	*	*	*	*	*	*	*	*
922	15	*7	*8	*	*	*	*8	*	*	*	*
922.1	*8	*	*	*	*	*	*	*	*	*	*
924	20	*7	14	*	*	*	14	*	*5	*7	*6
924.0	11	*	*7	*	*	*	*8	*	*	*	*
924.01	10	*	*7	*	*	*	*7	*	*	*	*
933	*6	*	*	*	*	*	*	*	*	*	*
935	*8	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
935.1	*7	*	*	*	*	*	*	*	*	*	*
941	*8	*7	*	*	*	*	*	*	*	*	*
941.2	*6	*6	*	*	*	*	*	*	*	*	*
942	10	*7	*	*	*	*	*	*	*	*	*
942.2	*6	*	*	*	*	*	*	*	*	*	*
943	*5	*	*	*	*	*	*	*	*	*	*
944	*6	*5	*	*	*	*	*	*	*	*	*
945	14	11	*	*	*5	*	*	*	*	*9	*
945.2	*7	*	*	*	*	*	*	*	*	*	*
945.3	*6	*	*	*	*	*	*	*	*	*	*
948	13	*8	*5	11	*	*	*	*	—	12	—
958	*6	*	*	*	*	*	*	*	*	*	*
959	36	22	14	*	16	*5	10	*7	*7	13	9
959.0	27	17	10	*	13	*	*7	*	*	10	*7
959.01	25	16	10	*	12	*	*7	*	*	10	*7
962	*8	*	*	*	*	*	*	*	*	*	*
962.3	*7	*	*	*	*	*	*	*	*	*	*
965	49	14	35	*	34	*8	*	*7	14	17	10
965.0	12	*5	*7	*	*7	*	*	*	*	*5	*
965.09	*8	*	*5	*	*	*	*	*	*	*	*
965.1	*7	*	*	*	*	*	*	*	*	*	*
965.4	22	*	18	*	18	*	*	*	*8	*7	*
966	9	*	*6	*	*	*	*	*	*	*	*
967	*8	*	*6	*	*	*	*	*	*	*	*
968	9	*6	*	*	*7	*	*	*	*	*	*
968.5	*7	*	*	*	*5	*	*	*	*	*	*
969	51	18	33	*	34	12	*	10	12	20	*8
969.0	18	*7	11	*	13	*	*	*	*5	*8	*
969.4	20	*7	14	*	13	*6	*	*	*	*9	*
972	*8	*	*5	*	*	*	*	*	*	*	*
977	11	*	*7	*	*6	*	*	*	*	*	*
977.8	*5	*	*	*	*	*	*	*	*	*	*
977.9	*5	*	*	*	*	*	*	*	*	*	*
989	*9	*	*	*	*	*	*	*	*	*	*
989.5	*5	*	*	*	*	*	*	*	*	*	*
995	34	11	23	*	*9	11	12	9	11	11	*
995.0	*7	*	*	*	*	*	*	*	*	*	*
995.1	10	*	*7	*	*	*	*	*	*	*	*
995.2	12	*	*9	*	*	*	*6	*	*	*5	*
996	392	206	186	31	64	119	178	88	89	139	76
996.0	20	11	*9	*	*	*	13	*6	*	*6	*
996.01	13	*6	*6	*	*	*	10	*	*	*	*
996.04	*6	*	*	—	*	*	*	*	*	*	*
996.1	*9	*	*	*	*	*	*	*	*	*	*
996.2	24	11	13	14	*	*	*	*5	*8	*6	*
996.3	*6	*	*	*	*	*	*	*	*	*	*
996.4	78	36	43	*	*8	27	43	17	21	26	14
996.5	*8	*	*	*	*	*	*	*	*	*	*
996.59	*5	*	*	*	*	*	*	*	*	*	*
996.6	122	70	52	11	19	36	55	26	30	44	21
996.62	59	32	27	*7	11	18	22	13	16	22	*9
996.64	12	*7	*	—	*	*	*8	*	*	*	*
996.66	16	10	*6	—	*	*	11	*	*	*7	*
996.67	12	*7	*	*	*	*	*6	*	*	*5	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
996.68	*6	*	*	*	*	*	*	*	*	*	*
996.69	*6	*	*	*	*	*	*	*	*	*	*
996.7	106	54	52	*	18	33	53	21	21	37	27
996.73	37	15	21	*	*5	12	20	*5	*5	14	12
996.74	31	18	14	*	*	10	16	*9	*5	11	*7
996.77	9	*6	*	*	*	*	*	*	*	*	*
996.78	11	*	*7	*	*	*	*	*	*	*	*
996.8	19	10	*9	*	*8	*8	*	*5	*	12	*
996.81	12	*6	*5	*	*5	*6	*	*	*	*7	*
997	93	48	45	*	18	29	42	23	22	33	15
997.0	*6	*	*	*	*	*	*	*	*	*	*
997.1	10	*5	*	*	*	*	*7	*	*	*	*
997.2	*7	*	*	*	*	*	*	*	*	*	*
997.3	*7	*	*	*	*	*	*	*	*	*	*
997.4	33	13	20	*	11	10	12	9	*9	10	*6
997.5	*8	*5	*	*	*	*	*	*	*	*	*
997.6	20	13	*7	*	*	*8	11	*6	*	*8	*
997.62	13	*9	*	*	*	*	*8	*	*	*6	*
998	188	92	96	10	52	54	72	48	45	67	28
998.1	47	27	19	*	14	12	18	13	10	16	*8
998.11	32	21	11	*	*8	*8	13	*8	*7	12	*5
998.12	13	*6	*7	*	*5	*	*	*	*	*	*
998.3	13	*6	*7	*	*	*	*	*	*	*7	*
998.5	99	44	54	*5	26	28	39	26	25	34	14
998.59	95	42	53	*5	25	27	39	25	24	32	14
998.6	*7	*	*	*	*	*	*	*	*	*	*
998.8	16	*7	9	*	*6	*	*	*	*	*7	*
998.89	11	*5	*6	*	*5	*	*	*	*	*	*
999	*6	*	*	*	*	*	*	*	*	*	*
V01-V82	4,640	378	4,261	89	3,851	196	505	814	974	1,808	1,043
V10	*7	*	*5	*	*	*	*	*	*	*	*
V12	14	*	*9	*	*	*	*7	*	*	*8	*
V12.5	9	*	*6	*	*	*	*5	*	*	*5	*
V22	*8	—	*8	—	*8	*	—	*	*	*6	*
V22.2	*6	—	*6	—	*6	*	—	*	*	*5	*
V27	3,738	—	3,738	12	3,722	*	—	632	771	1,439	895
V27.0	3,499	—	3,499	12	3,484	*	—	581	686	1,369	863
V27.1	18	—	18	*	18	—	—	*	*	*9	*
V27.2	54	—	54	—	54	*	—	*8	11	22	12
V27.9	163	—	163	*	163	*	—	40	70	39	15
V30	*9	*	*	*9	—	—	—	*	*	*	*
V30.1	*9	*	*	*9	—	—	—	*	*	*	*
V42	*5	*	*	*	*	*	*	*	*	*	*
V43	*5	*	*	—	*	*	*	*	*	*	*
V45	19	11	*8	*	*	*6	*9	*5	*	*9	*
V45.8	12	*7	*6	*	*	*5	*	*	*	*6	*
V45.89	*7	*	*	*	*	*	*	*	*	*	*
V53	*8	*	*	*	*	*	*6	*	*	*	*
V53.3	*8	*	*	*	*	*	*6	*	*	*	*
V53.31	*6	*	*	*	*	*	*	*	*	*	*
V54	9	*	*	*	*	*	*	*	*	*	*
V55	35	19	16	*	*7	13	12	10	*7	11	*7
V55.2	*6	*	*	*	*	*	*	*	*	*	*
V55.3	24	14	10	*	*	9	*9	*8	*5	*7	*

See footnotes at end of table.

Table 43. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
V57	513	187	326	*	31	84	394	82	141	210	80
V57.1	134	45	89	*	10	22	101	34	14	64	22
V57.8	365	138	227	*	19	60	283	41	126	141	57
V57.89	365	138	227	*	19	60	283	41	126	141	57
V57.9	13	*	*9	-	*	*	10	*7	*	*	*
V58	188	103	85	43	38	57	50	52	29	72	36
V58.0	18	*6	11	*	*	*6	*8	*	*	10	*
V58.1	134	82	51	39	27	43	24	39	22	49	24
V58.4	25	10	15	*	*6	*7	9	*9	*	*6	*8
V58.49	25	10	15	*	*6	*7	9	*9	*	*5	*8
V58.8	10	*	*6	*	*	*	*8	*	*	*7	*
V58.89	9	*	*6	*	*	*	*8	*	*	*7	*
V59	*7	*	*	*	*	*	*	*	*	*	*
V70	11	*5	*6	*	*	*	*	*	*	*5	*
V70.7	10	*	*6	*	*	*	*	*	*	*	*
V71	10	*	*5	*	*	*	*	*	*	*	*
V72	*5	*	*	*	*	*	*	-	*	*	-

* Figure does not meet standard of reliability or precision.

- Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
All codes	155,857	66,797	89,059	10,734	36,593	34,443	74,086	40,366	32,600	58,730	24,160
001-139	4,929	2,423	2,505	529	1,136	976	2,288	1,460	892	1,788	789
005	*10	*	*	*	*	*	*	*	*	*	*
008	437	152	285	110	55	61	210	110	100	158	70
008.4	233	76	158	*	*26	*30	172	*80	*61	*65	*
008.45	216	*62	153	*	*	*	165	*77	*54	*64	*
008.6	64	36	28	63	*	*	*	*	*11	23	25
008.61	62	35	28	62	*	-	*	*	*11	22	25
008.8	135	38	97	41	26	30	37	23	26	67	18
009	42	*16	*26	*	*	*	*	*	*	*	*
009.0	*15	*	*	*	*	*	*	*	*	*	*
011	*58	*	*	*	*	*	*	*	*	*	*
034	*12	*	*	*	*	*	*	*	*	*	*
034.0	*12	*	*	*	*	*	*	*	*	*	*
038	2,597	1,169	1,428	87	309	511	1,690	776	485	966	370
038.0	146	92	*53	*	*	*	94	*	*37	*	*
038.1	622	316	307	*	*126	125	358	162	88	275	98
038.10	*96	*68	*	*	*	*	*37	*	*	*	*
038.11	393	170	223	*	*	85	230	*87	*53	185	*67
038.19	134	77	*57	*	*	*	90	*	*	*56	*
038.2	122	*57	*65	*	*	*	*47	*	*	*53	*
038.4	607	252	355	*	*	122	442	204	114	180	109
038.40	*70	*	*	*	*	*	*	*	*	*	*
038.42	342	110	232	*	*	60	262	112	62	98	70
038.43	*38	*	*	-	*	*	*	*	*	*	*
038.49	155	*75	*79	*	*	*	107	*	*39	*	*
038.8	89	*	*48	*	*	*	*57	*	*	*	*
038.9	973	397	576	*33	101	158	680	302	179	377	115
041	97	*48	*48	*	*	*	*57	*	*	*62	*
042	602	445	157	*	415	156	*	271	*58	189	83
047	82	46	37	38	33	*	*	*	*15	30	29
047.9	75	43	32	36	32	*	*	*	*14	30	24
053	83	*37	46	*	*	*	60	*	*28	*22	*
053.9	47	*	*27	*	*	*	*33	*	*	*	*
054	*68	*	*	*	*	*	*	*	*	*	*
070	100	61	*39	*	*34	*45	*	*	*	*30	*
070.5	*33	*	*	-	*	*	*	*	*	*	*
075	36	*	*25	*	*18	*	-	*	*	*	*
079	178	90	88	111	24	*13	*31	53	35	53	37
079.6	*22	*	*	*22	-	-	-	*	*	*	*
079.9	151	76	75	85	23	*13	*31	51	24	50	26
079.99	151	76	75	85	23	*13	*31	51	24	50	26
112	103	*59	*45	*	*	*29	*58	*	*	*42	*
135	*41	*	*31	-	*	*	*	*	*	*	*
136	*35	*	*	*	*	*	*	*	*	*	*
140-239	9,264	4,081	5,183	239	1,205	2,977	4,843	2,505	1,799	3,387	1,574
150	78	*61	*	-	*	*	*39	*	*	*	*
151	302	163	138	-	*	*73	202	*54	*	*	*109
151.0	*115	*95	*	-	*	*	*	*	*	*	*
151.9	*94	*	*	-	*	*	*	*	*	*	*
153	997	504	493	-	*36	251	709	228	266	361	142
153.0	*62	*	*	-	*	*	*	*	*	*	*
153.1	101	*	*64	-	*	*	*64	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
153.2	*62	*	*	—	*	*	*	*	*	*	*
153.3	229	109	120	—	*	71	153	*45	*74	81	*
153.4	174	*78	95	—	*	*39	133	*	*	*61	*
153.6	197	103	94	—	*	*	156	*	*	95	*
153.7	*52	*39	*	—	*	*	*	*	*	*	*
153.9	87	*51	*35	—	*	*	*62	*	*39	*	*
154	324	190	134	*	*	105	202	92	*62	104	*67
154.0	95	*50	*	—	*	*	*64	*	*	*	*
154.1	200	122	77	*	*	*66	122	*54	*	71	*
155	78	*57	*	*	*	*42	*	*	*	*	*
155.0	*38	*	*	*	*	*	*	*	*	*	*
156	*52	*	*	—	—	*	*	*	*	*	*
157	233	107	126	*	*	*78	142	*67	*42	71	*
157.0	93	*	*55	*	*	*	*53	*	*	*	*
157.9	*79	*	*	—	*	*	*49	*	*	*	*
161	*64	*	*	—	*	*	*	*	*	*	*
162	818	425	393	*	*34	237	539	237	165	287	129
162.2	*52	*	*	—	*	*	*32	*	*	*	*
162.3	265	154	112	—	*	80	178	72	*58	79	*56
162.5	125	*57	*68	—	*	*	78	*	*	*	*
162.8	113	*45	*68	—	*	*	*82	*67	*	*	*
162.9	231	123	107	*	*	61	153	*55	40	110	*25
170	*54	*	*	*	*	*	*	*	*	*	*
171	*31	*	*	*	*	*	*	*	*	*	*
174	317	—	317	—	27	83	207	87	60	137	32
174.3	*19	—	*19	—	*	*	*	*	*	*	*
174.4	43	—	43	—	*	*12	27	*15	*10	*	*8
174.8	62	—	62	—	*	24	30	28	*	*12	*10
174.9	161	—	161	—	*	36	117	*29	*26	98	*
180	78	—	78	—	*41	*	*	*	*	*32	*
180.9	51	—	51	—	*32	*	*	*	*	*	*
182	127	—	127	—	*	48	69	*31	*26	51	*19
182.0	125	—	125	—	*	48	67	*31	*26	50	*19
183	220	—	220	*	*	59	119	84	*	*85	*
183.0	218	—	218	*	*	*58	118	83	*	*85	*
185	331	331	—	—	*	93	235	91	71	118	51
188	245	176	69	*	*	*36	203	120	*34	35	*55
188.8	73	*53	*	—	*	*	*58	*	*	*	*
188.9	101	80	*	*	*	*	80	*	*	*	*
189	213	115	98	*	*	82	114	*48	*52	94	*
189.0	181	98	83	*	*	75	90	*30	*48	87	*
191	160	100	60	*	*35	63	*36	*37	*	47	*47
191.8	*30	*	*	*	*	*	*	*	*	*	*
191.9	*30	*	*	*	*	*	*	*	*	*	*
193	31	*	20	*	*8	*16	*	*	*	*11	*
196	53	*22	*30	—	*	*20	*26	*	*	*	*
196.0	*17	*	*	—	*	*	*	*	*	*	*
197	743	256	487	*	102	196	443	269	139	243	92
197.0	105	*32	73	*	*	*	*56	*	*	*39	*
197.2	228	*46	182	—	*	*63	126	83	*	83	*
197.6	80	*	*55	—	*	*	*51	*	*	*	*
197.7	233	92	141	*	*	*63	142	*87	*46	*70	*
198	549	264	285	*	*54	204	288	158	118	174	99
198.3	225	104	121	*	*	89	112	65	*47	*68	*45
198.5	164	82	82	*	*	*63	92	*44	*41	*41	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
198.8	115	*57	58	—	*	*43	58	*	*	*52	*
198.89	110	*55	55	—	*	*40	57	*	*	*49	*
199	277	128	149	*	*35	125	118	69	*	154	*
199.0	229	98	131	—	*	117	86	*53	*	132	*
199.1	48	*30	*	*	*	*	*32	*	*	*	*
200	96	*33	*63	*	*	*	*54	*	*	*	*
200.0	*71	*	*	*	*	*	*	*	*	*	*
202	306	165	141	*	*	111	149	*73	*48	156	*
202.8	253	155	98	—	*	106	115	*70	*41	118	*
202.80	126	*82	*44	—	*	*67	*44	*	*	*71	*
202.83	*37	*	*	—	*	*	*	*	*	*	*
203	95	*43	*53	—	*	*34	*57	*	*	*39	*
203.0	95	*43	*52	—	*	*34	*56	*	*	*39	*
203.00	94	*42	*52	—	*	*34	*56	*	*	*39	*
204	110	69	*41	44	*	*	*	*	*	*	*27
204.0	67	*41	*	44	*	*	*	*	*	*	*19
204.00	*51	*	*	*37	*	*	*	*	*	*	*
204.1	*43	*	*	—	*	*	*	*	*	*	*
204.10	*43	*	*	—	*	*	*	*	*	*	*
205	250	160	*90	*	*	*100	*47	*	*	*74	*
205.0	212	*134	*	*	*	*93	*	*	*	*	*
205.00	184	*112	*	*	*	*	*	*	*	*	*
210	*8	*	*	*	*	*	*	*	*	*	*
211	203	70	132	*	*	*38	153	*52	*31	82	*38
211.3	124	58	66	*	*	*	96	*	*	50	*30
211.4	*42	*	*37	—	*	*	*38	*	*	*	*
218	643	—	643	—	304	319	*20	136	134	246	127
218.0	78	—	78	—	34	39	*	*22	*	31	*15
218.1	207	—	207	—	102	100	*	44	38	89	37
218.2	86	—	86	—	33	53	*	*24	*	*15	*34
218.9	272	—	272	—	135	128	*	47	73	111	41
220	96	—	96	*	42	30	*23	*25	24	29	*18
225	111	*41	70	*	*	53	*40	*31	*	*35	*23
225.2	70	*	48	—	*	*39	*27	*	*	*	*
226	13	*	*12	—	*	*	*	*	*	*	*
227	61	*	44	*	*	*19	*33	*	*	*	*
227.1	*16	*	*	—	*	*	*	*	*	*	*
227.3	*25	*	*	*	*	*	*	*	*	*	*
228	*23	*	*	*	*	*	*	*	*	*	*
233	53	*	50	—	20	*19	*13	*	*11	*18	*14
233.0	24	*	24	—	*	*10	*11	*	*	*	*
233.1	26	—	26	—	*16	*	*	*	*	*	*
235	*56	*	*	*	*	*	*	*	*	*36	*
237	*29	*	*	*	*	*	*	*	*	*	*
238	88	*56	*32	*	*	*	*60	*	*	*	*
238.7	72	*	*28	*	*	*	*53	*	*	*	*
239	51	*	*30	*	*	*	*31	*	*	*26	*
240-279	6,570	2,891	3,678	426	1,044	1,845	3,254	1,685	1,384	2,609	892
241	26	*	21	—	*	*10	*	*	*	*12	*
241.1	19	*	*15	—	*	*7	*	*	*	*10	*
242	46	*	30	*	*	*	*	*	*	*	*
242.0	*15	*	*	*	*	*	*	*	*	*	*
242.00	*13	*	*	*	*	*	*	*	*	*	*
242.9	*20	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
244	60	*	51	—	*	*12	*42	*17	*	*22	*
244.9	59	*	50	—	*	*11	*42	*17	*	*22	*
250	2,877	1,494	1,384	69	516	1,165	1,128	779	537	1,145	417
250.0	562	247	314	*23	100	233	206	140	94	248	79
250.00	242	115	127	*	36	110	94	48	40	125	30
250.01	77	37	40	*15	*11	25	*26	*26	*18	26	*
250.02	185	61	124	*	34	80	70	46	*25	72	41
250.03	58	34	*24	*	*19	*18	*	*	*	*25	*
250.1	402	233	169	37	190	129	*46	103	82	173	44
250.10	68	43	*24	*	*31	*26	*	*	*	*34	*
250.11	125	67	59	*21	71	*28	*	*28	34	49	*15
250.12	90	56	*34	—	*28	*41	*	*	*	*26	*
250.13	119	67	52	*	61	*34	*	*	*24	65	*
250.2	76	*39	*37	*	*	*	*48	*	*	*	*
250.22	*32	*	*	—	*	*	*	*	*	*	*
250.4	208	108	100	—	*	106	72	*58	*46	64	40
250.40	115	62	52	—	*	59	*47	*	*	*	*24
250.41	*51	*	*	—	*	*	*	*	*	*	*
250.6	307	146	161	—	63	149	94	79	66	146	*
250.60	155	80	75	—	*	58	67	*38	*40	66	*
250.61	54	*	*	—	*	*31	*	*	*	*29	*
250.62	45	*	*23	—	*	*31	*	*	*	*	*
250.63	*52	*	*	—	*	*	*	*	*	*	*
250.7	572	326	246	—	*	263	286	158	122	191	102
250.70	298	192	106	—	*	130	157	*90	*70	106	*
250.71	165	*90	*74	—	*	*63	*93	*	*	*68	*
250.72	*86	*	*	—	*	*	*	*	*	*	*
250.8	673	370	302	*	90	242	337	184	103	273	113
250.80	364	204	159	—	*29	102	234	95	46	118	105
250.81	130	75	*55	*	*	*71	*33	*58	*	*56	*
250.82	107	*53	54	—	*	*49	49	*	*33	*48	*
250.83	71	*38	*	*	*	*	*	*	*	*50	*
250.9	*35	*	*	*	*	*	*	*	*	*	*
251	54	*25	29	*	*	*	33	*	*	*27	*
251.2	42	*23	*19	*	*	*	*25	*	*	*22	*
253	70	*	*38	*	*	*	*38	*	*	*	*
253.6	*47	*	*29	*	*	*	*33	*	*	*	*
255	*21	*	*	*	*	*	*	*	*	*	*
263	99	*	70	*	*	*	67	*	*	*44	*
263.9	91	*	67	*	*	*	64	*	*	*40	*
272	57	31	26	*	*13	34	*8	19	*11	15	*12
272.0	28	15	*13	—	*	19	*	*	*	*8	*
272.4	22	*14	*8	—	*	*11	*	*	*	*	*
274	43	*18	*25	—	*	*	*31	*	*	*	*
274.0	*25	*	*	—	*	*	*	*	*	*	*
275	93	*26	66	*	*14	*	*55	*	*17	*30	*
275.4	79	*	60	*	*	*	*53	*	*17	*	*
275.42	*59	*	*49	*	*	*	*46	*	*	*	*
276	2,625	1,051	1,574	253	224	373	1,775	631	622	1,042	330
276.0	*73	*	*	*	*	*	*61	*	*	*	*
276.1	311	85	226	*	*23	71	214	57	96	102	57
276.2	*25	*	*18	*	*	*	*	*	*	*	*
276.5	1,947	814	1,133	239	166	224	1,318	497	455	788	206
276.6	*29	*	*	*	*	*	*	*	*	*	*
276.7	108	51	57	*	*	*20	83	*	*27	44	*
276.8	106	43	63	*	*20	*27	58	*23	*20	49	*
276.9	*19	*	*	*	*	*	*18	*	*	*	*
277	146	*53	94	*47	73	*	*	*	*54	*44	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
277.0	104	*45	*59	*	*58	*	—	*	*	*39	*
277.00	101	*44	*57	*	*58	*	—	*	*	*39	*
278	178	35	143	*	112	64	*	*41	56	46	35
278.0	164	32	132	*	108	55	*	*37	52	41	35
278.00	*15	*	*	—	*	*	*	*	*	*	*
278.01	149	*25	124	*	102	46	*	*33	*47	36	*33
280-289	1,741	704	1,036	277	424	371	669	468	365	675	233
280	256	67	189	*	*19	69	165	55	62	100	*39
280.0	137	*34	103	*	*	*39	88	*23	*	65	*
280.9	117	33	84	*	*	*28	76	*31	40	35	*
282	371	137	233	94	260	*	*	132	65	152	*21
282.6	353	132	220	87	254	*	*	127	64	145	*
282.62	331	122	208	75	245	*	—	125	58	134	*
283	*33	*	*	*	*	*	*	*	*	*	*
284	168	83	85	*41	*	*58	*46	*51	*33	68	*
284.8	144	64	80	*	*	*53	*44	*38	*33	*56	*
284.9	*24	*	*	*	*	*	*	*	*	*	—
285	260	95	165	*	*29	48	179	68	43	106	43
285.1	54	*31	*23	*	*	*	39	*	*	*28	*
285.9	194	57	137	*	*25	38	130	57	31	73	33
286	85	*37	49	*	*	*	53	*	*19	*41	*
286.7	*20	*	*	—	*	*	*	*	*	*	*
287	157	87	70	29	*24	52	51	*39	40	51	*27
287.3	56	*34	*21	*	*	*	*	*	*	*	*
287.4	*25	*	*	*	*	*	*	*	*	*	*
287.5	59	*37	*22	*	*	*24	*	*	*	*20	*
288	324	144	180	68	42	86	129	83	84	112	45
288.0	265	113	152	57	*37	84	88	64	75	96	*30
288.8	46	*27	*19	*	*	*	*36	*	*	*	*
289	74	40	35	*21	*14	*	*	*	*	*33	*
290-319	15,762	7,986	7,776	1,277	8,188	3,703	2,595	5,879	2,866	5,820	1,198
290	430	145	285	—	—	*	399	91	83	210	*
290.0	*44	*	*	—	—	—	*44	*	*	*	*
290.2	*95	*	*62	—	—	—	*93	*	*	*	*
290.3	*60	*	*	—	—	—	*60	*	*	*	*
290.4	217	*76	140	—	—	*	194	*	*	137	*
290.40	*74	*	*	—	—	*	*56	*	*	*	*
290.41	*57	*	*34	—	—	*	*57	*	*	*	*
291	594	437	158	*	344	203	*46	157	278	122	38
291.0	116	97	*	—	48	*60	*	*37	*36	*34	*
291.8	432	312	121	*	285	119	*	114	216	78	24
291.81	392	282	109	*	248	115	*	102	211	58	*22
291.89	*41	*29	*	—	*36	*	—	*	*	*	*
292	418	238	180	*	273	84	61	213	60	91	*53
292.0	193	118	75	*	152	39	*	129	*24	*32	*
292.8	175	91	84	—	109	*	*43	66	*	*53	*
292.81	77	*	61	—	*32	*	*28	*	*	*32	*
292.84	*60	*46	*	—	*44	*	*	*	*	*	*
292.9	*40	*	*	—	*	*	*	*	*	*	*
293	289	121	167	*	92	87	108	125	*65	*67	*
293.0	114	*31	82	*	*	*	78	*	*38	*	*
293.8	126	81	*45	*	69	*	*	64	*	*	*
293.83	104	78	*	*	62	*	*	56	*	*	*
294	263	123	140	—	*	*	217	97	*60	61	*45
294.8	228	103	125	—	*	*	199	76	*55	58	*39

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
295	3,004	1,730	1,274	*	1,776	954	232	1,290	517	1,007	191
295.3	908	602	305	—	509	292	*107	448	185	190	*85
295.30	268	181	*87	—	160	*77	*	*89	109	*	*
295.32	167	110	*	—	*73	*42	*	*57	*	*	*
295.34	460	309	151	—	269	167	*	295	*	*91	*
295.6	177	96	*81	—	104	*55	*	*68	*	*51	*
295.64	93	*50	*43	—	*56	*	*	*	*	*	*
295.7	1,567	858	709	*	991	463	*71	630	206	651	*80
295.70	1,026	585	441	*	638	295	*	266	167	558	*35
295.74	513	257	255	—	342	152	*	356	*	82	*
295.9	252	142	111	*	137	*81	*	126	*57	*62	*
295.90	134	83	*51	*	*72	*	*	*	*	*55	*
295.94	*60	*43	*	—	*45	*	*	*53	*	*	*
296	6,007	2,673	3,334	555	3,036	1,326	1,091	1,938	1,061	2,707	302
296.2	1,100	438	662	177	490	162	271	313	220	479	88
296.20	517	221	296	81	241	76	119	163	141	197	*
296.22	*82	*	*	*	*	*	*	*	*	*66	*
296.23	230	95	134	*	97	*37	*	*73	*27	103	*
296.24	267	*74	194	*	129	*	*83	*70	*43	110	*
296.3	2,215	966	1,248	*163	984	533	534	810	399	923	84
296.30	502	184	319	*	230	115	136	133	106	239	*
296.32	122	*85	*38	*	*35	*	*	*	*	*	*
296.33	1,078	497	581	*	526	235	191	476	131	444	*28
296.34	498	198	301	*	186	121	184	170	92	212	*
296.4	563	226	337	*	286	184	*86	232	125	161	*46
296.40	255	109	146	*	142	67	*	*62	*78	101	*
296.43	*47	*	*	—	*	*	*	*	*	*	*
296.44	247	95	153	*	112	*98	*	146	*	*	*
296.5	573	233	340	*	281	197	*67	137	113	310	*
296.50	367	166	201	*	168	131	*	*84	*68	211	*
296.53	89	*	62	*	60	*	*	*	*	*39	*
296.54	113	*37	*76	—	*49	*	*	*	*	*58	*
296.6	670	315	355	*44	447	113	*	172	75	402	*
296.60	276	153	122	*39	162	*	*	*48	*45	176	*
296.63	112	*36	*76	*	74	*	*	*78	*	*	*
296.64	274	119	155	*	208	*48	*	*	*	198	*
296.7	428	204	225	*	213	98	*	147	90	154	*
296.8	319	212	107	*57	249	*	*	*	*	240	*
296.80	157	140	*	*	115	*	*	*	*	148	—
296.89	158	*	86	*	132	*	*	*	*	89	*
296.9	111	55	56	*	70	*23	*	68	*	*28	*
296.90	91	49	42	*	53	*21	*	62	*	*	*
297	*95	*	*67	—	*	*	*68	*	*	*	*
298	634	309	325	*	357	128	127	296	89	151	98
298.9	620	303	317	*	352	125	123	291	88	147	95
300	458	158	300	*	163	159	78	119	85	137	*116
300.0	231	49	182	*	63	115	52	41	31	57	*
300.00	182	36	146	*	44	*111	*27	*20	*25	37	*
300.01	41	*	*30	—	*	*	*	*	*	*	*
300.1	*20	*	*	*	*	*	*	*	*	*	*
300.4	108	54	54	*	48	*	*	*32	*26	*41	*
300.9	*15	*	*	*	*	*	*	*	*	*	*
301	112	*	*33	*	*76	*	*	*	*	*	*
303	836	570	266	*	529	277	*26	192	162	371	112
303.0	88	63	*25	—	37	48	*	*	*25	*36	*15
303.00	31	*17	*	—	*13	*	*	*	*	*	*
303.01	57	46	*	—	*24	*30	*	*	*	*	*
303.9	748	507	241	*	492	230	*	180	137	335	97
303.90	323	205	118	*	196	119	*	*	*33	238	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
303.91	419	296	123	*	292	109	*	162	104	95	*59
304	503	303	199	*	420	64	*	186	68	187	*61
304.0	239	139	100	—	194	38	*	106	*	83	*
304.00	*46	*	*34	—	*35	*	*	*	*	*38	*
304.01	193	127	66	—	159	*31	*	102	*	45	*
304.2	74	48	*	*	68	*	*	*31	*	*24	*
304.21	*54	*33	*	—	*51	*	*	*	*	*	*
304.3	*50	*36	*	*	*47	*	—	*	*	*	*
304.4	*21	*	*	—	*	*	—	—	*	*	*
304.8	60	*41	*	*	54	*	—	*	*	*31	*
304.80	*31	*23	*	*	*27	*	—	*	*	*29	*
305	267	134	133	*	169	77	*21	45	24	172	26
305.0	141	73	68	*	94	30	*	*27	*10	95	*10
305.00	104	42	62	*	75	*23	*	*	*	79	*
305.01	*32	*29	*	—	*	*	*	*	*	*	*
305.1	32	20	*11	—	17	*11	*	*	*	14	*
305.6	38	*13	*25	—	*15	*	*	*	*	*24	*
305.60	*28	*9	*	—	*	*	*	*	*	*	*
305.9	39	*18	*21	*	*29	*	*	*	*	*29	*
305.90	37	*17	*20	*	*28	*	*	*	*	*29	*
307	*62	*	*32	*	*	*	*	*	*	*	*
308	*12	*	*	—	*	*	*	*	*	*	*
309	444	186	257	*86	298	38	*	275	67	69	33
309.0	116	45	71	*	98	*	*	79	*11	*17	*
309.2	62	*22	*40	*	45	*	*	*	*23	*	*
309.28	59	*20	*39	*	44	*	*	*	*22	*	*
309.4	99	50	50	*	65	*	*	59	*19	*	*
309.8	141	*	*83	*	*74	*	*	*106	*	*	*
309.81	138	*	*81	*	*72	*	*	*106	*	*	*
309.9	*14	*8	*	*	*11	*	*	*	*	*	*
310	*45	*	*26	*	*	*	*29	*	*	*	*
311	662	360	302	*56	406	158	*43	426	70	147	*20
312	229	138	91	*113	102	*	*	159	*	*	*
312.3	174	93	81	*73	86	*	*	130	*	*	*
312.30	134	63	71	*	64	*	—	108	*	*	*
313	191	*60	*131	*156	*34	—	—	*	*	*	*
313.8	190	*60	*131	*156	*34	—	—	*	*	*	*
313.81	87	*	*	*	*32	—	—	*	*	*	*
314	155	103	*	146	*	—	—	*	*	*	*
314.0	153	101	*	145	*	—	—	*	*	*	*
314.01	151	*100	*	143	*	—	—	*	*	*	*
320-389	2,253	1,018	1,236	331	498	468	958	551	528	786	388
320	*94	*	*	*	*	*	*	*	*	*	*
331	429	204	225	*	*	*	371	91	125	167	*47
331.0	334	143	191	—	—	*	312	*74	102	128	*
331.3	*43	*	*	*	*	*	*39	*	*	*	*
331.4	*40	*22	*	*	*	*	*	*	*	*	*
332	94	*34	60	—	*	*	86	*43	*	*26	*
332.0	94	*34	60	—	*	*	86	*43	*	*26	*
333	76	*44	*33	*	*	*	*35	*	*	*	*
337	*24	*	*17	*	*	*	*	*	*	*	*
337.2	*18	*	*	*	*	*	*	*	*	*	*
340	129	*36	94	—	44	70	*	*	*44	43	*
343	*51	*	*	*15	*	*	—	*	*	*	*
343.9	*17	*	*	*	*	—	—	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
345	224	121	103	78	57	39	*50	59	43	72	50
345.1	*32	*19	*	*	*	*	*	*	*	*	*
345.10	*23	*	*	*	*	*	*	*	*	*	*
345.3	58	37	*21	*24	*	*	*	*	*	*21	*
345.4	*22	*	*	*	*	*	*	*	*	*	—
345.5	*29	*	*22	*	*	*	*	*	*	*	*
345.9	54	*35	*20	*16	*	*	*	*	*	*23	*
345.90	46	*31	*	*	*	*	*	*	*	*22	*
346	142	*19	123	*	85	46	*	*22	58	48	*14
346.8	*30	*	*24	*	*	*	*	*	*	*	*
346.9	69	*	62	*	46	*16	*	*	*21	29	*
346.90	47	*	43	*	28	*13	*	*	*	21	*
346.91	*22	*	*19	*	*18	*	*	*	*	*	*
348	165	88	77	*	*39	*47	*62	*51	*38	*54	*
348.3	*56	*	*33	*	*	*	*	*	*	*	*
349	*33	*	*	*	*	*	*	*	*	*	*
350	*15	*	*	—	*	*	*	*	*	*	*
350.1	*15	*	*	—	*	*	*	*	*	*	*
351	*20	*	*	*	*	*	*	*	*	*	*
351.0	*18	*	*	*	*	*	*	*	*	*	*
354	*14	*	*	*	*	*	*	*	*	*	*
356	*24	*	*	*	*	*	*	*	*	*	*
357	*64	*	*	*	*	*	*	*	*	*	*
359	*34	*	*	*	*	*	*	*	*	*	*
376	*23	*	*	*17	*	*	*	*	*	*	*
376.0	*22	*	*	*17	*	*	*	*	*	*	*
376.01	*21	*	*	*16	*	*	*	*	*	*	*
380	*19	*	*	*	*	*	*	*	*	*	*
380.1	*15	*	*	*	*	*	*	*	*	*	*
382	45	*22	*22	42	*	*	*	*	*	*13	*
382.9	37	*16	*21	34	*	*	*	*	*	*12	*
386	114	47	67	—	*	26	82	49	20	35	*
386.1	33	*	21	—	*	*	*22	*16	*	*	*
386.11	*19	*	*15	—	*	*	*	*	*	*	*
386.3	68	*34	34	—	*	*15	50	*	*	*22	*
386.30	62	*31	30	—	*	*10	48	*	*	*21	*
390-459	30,407	14,637	15,770	168	1,516	7,610	21,113	8,317	6,662	11,468	3,959
394	*69	*	*60	—	*	*	*	*	*	*	*
396	79	*	*50	—	*	*	61	*	*	*23	*
396.2	*27	*	*	—	—	*	*	*	*	*	*
398	168	62	106	—	*	*	159	74	*	*52	*
398.9	168	62	106	—	*	*	159	74	*	*52	*
398.91	162	61	101	—	*	*	153	74	*	*50	*
401	645	217	428	*	75	264	303	152	135	292	66
401.0	102	27	76	—	*15	32	56	*24	*19	55	*
401.9	534	187	347	*	60	229	242	125	111	236	61
402	421	163	258	*	*	80	326	138	66	174	44
402.9	413	161	252	*	*	79	320	137	64	170	43
402.90	51	*24	27	*	*	*22	*22	*16	*	*25	*
402.91	361	137	225	—	*	56	297	120	59	145	*37
403	468	223	245	*	89	146	206	90	69	205	105
403.9	440	207	233	*	77	140	196	84	68	185	104
403.90	*42	*	*	*	*	*	*	*	*	*	—
403.91	398	179	219	*	72	130	168	56	66	172	104

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
404	217	101	116	—	*	*62	141	*54	*	91	*
404.9	198	89	109	—	*	*48	136	*	*	86	*
404.93	153	*66	86	—	*	*	108	*	*	*66	*
410	4,411	2,349	2,062	*	143	1,165	3,091	1,215	998	1,584	615
410.0	145	75	70	—	*	*38	105	*27	*30	67	*
410.01	142	74	68	—	*	*37	102	*24	*30	67	*
410.1	569	299	269	—	*31	163	374	117	127	237	88
410.11	546	286	260	—	*28	156	361	108	125	230	83
410.2	105	50	55	—	*	*25	77	*41	*	*21	*
410.21	99	50	49	—	*	*25	71	*37	*	*20	*
410.3	67	35	*	—	*	*22	*41	*	*	*	*
410.31	65	*34	*	—	*	*22	*40	*	*	*	*
410.4	622	334	289	—	*21	232	370	210	130	217	65
410.41	607	330	276	—	*21	220	366	208	130	204	65
410.5	72	51	*	*	*	*	*42	*	*	*	*
410.51	70	*50	*	*	*	*	*41	*	*	*	*
410.7	2,113	1,098	1,014	*	55	531	1,527	623	486	686	317
410.71	2,069	1,078	991	*	54	526	1,489	612	483	660	314
410.8	*54	*	*	*	*	*	*41	*	*	*	*
410.81	*48	*	*	*	*	*	*41	*	*	*	*
410.9	635	352	283	*	*	121	488	143	141	284	67
410.90	*51	*	*	—	*	*	*39	*	*	*	—
410.91	576	320	255	*	*	112	440	124	134	251	67
411	453	234	219	*	25	154	274	150	96	151	57
411.1	396	210	186	—	22	142	232	126	84	142	44
411.8	48	*21	*26	*	*	*	36	*21	*	*	*
411.89	43	*18	*25	*	*	*	33	*21	*	*	*
412	*7	*	*	—	*	*	*	*	*	*	*
413	194	79	114	—	*13	51	130	64	32	71	27
413.9	191	79	113	—	*11	50	130	63	32	70	26
414	4,386	2,618	1,768	*	149	1,509	2,726	1,113	964	1,755	553
414.0	4,274	2,561	1,713	*	143	1,479	2,650	1,062	947	1,722	543
414.00	614	333	281	—	*15	175	424	138	132	288	56
414.01	3,496	2,118	1,378	*	121	1,262	2,112	888	760	1,376	471
414.02	131	90	41	—	*	28	101	*27	50	41	*
414.05	*26	*	*	*	*	*	*	*	*	*	*
414.8	61	*33	*	—	*	*	45	*	*	*	*
414.9	46	*23	*	—	*	*15	*26	*	*	*	*
415	523	205	318	—	71	164	288	132	114	207	70
415.1	513	199	315	—	71	162	280	126	113	204	70
415.11	*38	*	*	—	*	*	*	*	*	*	*
415.19	476	187	288	—	60	145	270	116	101	191	67
416	77	*	*55	*	*	*	*37	*	*	*	*
416.0	*45	*	*32	*	*	*	*	*	*	*	*
420	70	*32	*	*	*	*	*	*	*	*	*
420.9	70	*32	*	*	*	*	*	*	*	*	*
420.90	55	*23	*	*	*	*	*	*	*	*	*
421	242	141	*101	*	*	*76	*105	*94	*	*84	*
421.0	223	121	*101	*	*	*76	*98	*92	*	*67	*
423	51	*	*26	*	*	*	*28	*	*	*22	*
423.9	37	*	*20	—	*	*	*	*	*	*	*
424	520	260	260	*	44	133	341	150	128	142	100
424.0	185	82	103	*	*23	59	103	47	*53	52	*33
424.1	310	161	149	—	*	69	222	92	73	82	63
425	149	66	82	*	*24	37	80	*46	*33	49	*
425.4	121	53	68	*	*	31	68	*37	*29	42	*
426	214	112	103	*	*	*24	178	67	49	73	*25

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
426.0	133	71	62	*	*	*	115	*38	*29	50	*
426.1	41	*20	*	*	*	*	*32	*	*	*	*
427	2,640	1,195	1,444	*29	102	448	2,060	769	566	979	325
427.0	77	27	50	*	*	*17	50	*15	24	*23	*14
427.1	234	184	50	*	*	61	159	94	35	75	*30
427.3	1,466	613	853	*	38	251	1,177	431	357	540	138
427.31	1,311	515	796	—	34	202	1,075	392	318	475	127
427.32	155	98	57	*	*	49	102	*39	39	65	*11
427.4	*52	*	*	*	*	*	*	*	*	*	*
427.41	*52	*	*	*	*	*	*	*	*	*	*
427.5	69	*37	*33	*	*	*	*62	*	*	*	*
427.6	40	*19	*21	—	*	*	*24	*	*	*15	*
427.69	25	*16	*	—	*	*	*17	*	*	*	*
427.8	634	233	401	*	24	92	499	176	130	233	96
427.81	289	111	178	*	*	*26	257	93	56	109	*31
427.89	346	122	223	*	22	66	242	83	74	124	65
427.9	65	44	21	*	*	*15	43	*23	*	*25	*
428	5,505	2,357	3,148	*	157	964	4,374	1,611	1,245	2,067	583
428.0	5,448	2,328	3,120	*	157	958	4,323	1,575	1,237	2,055	581
428.9	*47	*	*	*	—	*	*42	*	*	*	—
429	66	*29	37	*	*	*20	42	*	*	*33	*
429.2	*33	*	*	—	—	*	*26	*	*	*	*
430	301	*98	203	—	*100	95	*106	*75	*	104	*
431	547	283	264	*	*	191	329	107	173	135	133
432	143	93	*50	*	*	*	109	*	*	*41	*25
432.1	137	88	*49	*	*	*	105	*	*	*39	*23
433	598	312	286	*	*	140	445	145	125	258	70
433.1	455	235	220	—	*	101	347	114	95	197	50
433.10	322	141	180	—	*	84	234	65	73	151	33
433.11	134	94	*40	—	*	*	113	*49	*	*46	*
433.3	119	70	48	—	*	*29	85	*28	*25	48	*
433.30	78	50	28	—	*	*19	55	*18	*16	*35	*
433.31	*41	*	*	—	*	*	*	*	*	*	*
434	1,806	736	1,069	*	*42	409	1,346	505	360	713	228
434.0	79	*35	*44	—	*	*	*61	*	*	*	*
434.01	67	*	*38	—	*	*	*54	*	*	*	*
434.1	273	120	153	—	*	59	202	101	*49	70	53
434.10	*35	*	*	—	—	*	*34	*	*	*	*
434.11	238	110	128	—	*	59	169	*92	*41	55	51
434.9	1,454	581	873	*	*29	333	1,082	386	285	621	162
434.90	*37	*	*	—	*	*	*	*	*	*	*
434.91	1,416	569	847	*	*29	315	1,063	384	278	595	160
435	787	302	485	*	31	156	600	296	155	263	73
435.3	*30	*	*	—	*	*	*26	*	*	*	*
435.8	*20	*	*	—	*	*	*	*	*	*	*
435.9	725	272	453	*	28	147	549	284	136	249	56
436	789	289	500	*	*	139	636	104	176	372	136
437	145	*42	103	*	*	*49	89	*	*38	*58	*
437.3	*37	*	*	—	*	*	*	*	*	*	*
438	160	78	82	*	*	*	139	*	*20	96	*26
438.2	*54	*	*	—	*	*	*46	*	*	*	*
438.20	*46	*	*	—	*	*	*39	*	*	*	*
438.8	74	*46	*28	—	*	*	66	*	*	*	*
438.89	*45	*	*	—	*	*	*38	*	*	*	*
440	877	515	362	*	*	243	614	229	224	315	110
440.1	*17	*	*	—	*	*	*	*	*	*	*
440.2	803	484	319	*	*	223	566	210	205	294	94

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
440.20	54	*38	*	*	*	*	*31	*	*	*24	*
440.21	118	76	42	—	*	46	68	*43	*21	34	*
440.22	91	*46	*45	—	*	*	*58	*	*	*	*
440.23	143	84	*60	—	*	*	131	*33	*50	*	*
440.24	394	239	154	—	*	*113	274	*92	*109	154	*38
440.3	*38	*	*	—	—	*	*	*	*	*	*
441	548	349	198	*	*	98	439	201	97	153	97
441.0	69	*42	*	*	*	*	*	*	*	*	*
441.01	*42	*	*	*	*	*	*	*	*	*	*
441.3	*73	*	*	—	—	*	*53	*	*	*	*
441.4	309	231	*79	—	*	*35	274	102	*62	93	*53
442	*55	*	*	*	*	*	*	*	*	*	*
443	178	82	97	*	*	*61	106	*	59	78	*
443.9	157	74	83	—	*	*57	99	*	*50	69	*
444	285	177	109	—	*	95	179	*64	*36	139	*46
444.2	219	143	76	—	*	*65	145	*51	*	106	*
444.22	190	126	64	—	*	*54	129	*48	*	91	*
446	72	*22	*50	*18	*	*	*	*	*	*	*
446.1	*18	*	*	*18	*	—	—	*	*	*	*
447	64	*29	*35	*	*	*29	*23	*	*	*23	*
447.1	*26	*	*	—	*	*	*	*	*	*	*
451	111	*41	70	*	*	*36	58	*	*35	52	*
451.0	*25	*	*20	—	*	*	*	*	*	*	*
451.1	52	*	*31	—	*	*	*33	*	*	*	*
451.19	*33	*	*25	—	*	*	*	*	*	*	*
453	725	327	398	*	97	214	411	190	188	269	78
453.8	622	270	351	*	83	186	349	144	174	231	73
453.9	95	56	*39	—	*	*	57	*41	*	*35	*
454	*43	*	*	—	*	*	*	*	*	*	*
455	94	58	36	—	*21	27	47	*16	*18	51	*
455.2	59	36	*23	—	*	*20	*33	*	*	*32	*
456	*28	*	*	—	*	*	*	*	*	*	*
458	298	113	186	*	*	83	204	87	59	109	43
458.0	139	61	78	*	*	*19	116	*31	*30	59	*
458.9	140	39	101	*	*	*59	74	*53	*23	42	*23
459	93	48	*45	*	*	*23	63	*	*35	*	*
459.8	59	*32	*	—	*	*	*44	*	*	*	*
459.81	*45	*	*	—	*	*	*31	*	*	*	*
460-519	18,518	8,449	10,069	2,279	1,763	3,613	10,863	4,524	3,919	7,012	3,063
461	50	*29	*21	*	*21	*	*	*	*	*23	*
462	24	*	*17	*	*11	*	*	*	*	*	*
463	*19	*	*	*	*16	*	*	*	*	*	*
464	56	34	22	46	*	*	*	*11	*13	*16	*17
464.4	34	24	*10	34	*	—	—	*	*10	*9	*8
465	79	36	43	44	*	*	*18	*17	*15	24	*22
465.9	77	35	41	44	*	*	*18	*16	*15	24	*22
466	983	457	526	623	49	57	254	175	196	342	270
466.0	370	132	238	36	48	56	230	67	95	169	39
466.1	613	324	289	587	*	*	*	107	102	173	231
466.11	394	206	188	393	*	*	*	72	62	82	177
466.19	219	118	101	195	*	*	*	35	39	91	54
470	*10	*10	*	—	*	*	*	*	*	*	*
473	38	*22	*16	*	*	*	*	*	*	*20	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
474	26	*16	*10	22	*	*	*	*13	*	*8	*
474.0	*8	*	*	*	*	*	*	*	*	*	*
474.1	*18	*13	*	*17	*	*	—	*	*	*	*
474.10	*14	*	*	*14	*	—	—	*	*	*	*
475	36	*18	*18	*	*20	*	*	*	*	*	*
478	80	*36	44	*25	*	*	*	*40	*	*18	*15
478.2	*50	*	*	*	*	*	*	*	*	*	*
480	128	59	69	78	*	*	*37	*	*36	37	*28
480.1	59	*39	*	54	*	—	*	*	*	*15	*
480.9	61	*	*46	*19	*	*	*	*	*	*	*
481	220	109	112	*	*	57	132	*36	59	77	*49
482	1,200	613	586	*71	116	194	819	220	297	503	180
482.0	*46	*	*	—	*	*	*	*	*	*	*
482.1	281	123	158	*	*68	*	137	*59	*77	*84	*
482.2	109	54	*55	*	*	*	92	*	*41	*31	*
482.3	*54	*	*	*	*	*	*40	*	*	*	*
482.4	362	213	149	*	*	*	280	*	*71	203	*
482.41	333	200	133	*	*	*	259	*	*66	181	*
482.8	246	120	126	*	*	*	174	*	73	113	*
482.83	176	*69	107	*	*	*	142	*	66	*61	*
482.9	101	59	*42	*	*	*26	58	*51	*	*29	*
483	*35	*	*	*	*	*	*	*	*	*	*
483.0	*32	*	*	*	*	*	*	*	*	*	—
485	160	49	112	49	*	*21	82	*	*32	82	*
486	5,783	2,442	3,342	372	530	989	3,893	1,308	1,376	2,234	865
487	180	79	100	*21	*	*20	124	*36	60	64	*
487.0	87	*44	*43	*	*	*	*63	*	*	*36	*
487.1	90	35	55	*12	*	*	60	*	33	26	*
490	81	37	44	*12	*	*	49	*	*	38	*
491	2,563	1,111	1,451	*	85	755	1,719	709	509	1,041	304
491.2	2,559	1,109	1,449	*	85	753	1,718	706	508	1,040	304
491.20	78	48	*30	—	*	*	56	*	*	*39	*
491.21	2,481	1,061	1,419	*	77	739	1,662	679	496	1,001	304
492	95	57	*38	*	*	*23	63	*33	*26	*30	*
492.8	81	45	*36	—	*	*	59	*31	*21	*23	*
493	1,415	505	910	445	317	322	331	387	284	539	205
493.0	111	55	56	78	*23	*	*	*15	*	*18	65
493.00	52	23	29	34	*	*	*	*	*	*14	28
493.01	59	32	*27	44	*	*	*	*	*	*	37
493.2	255	78	178	*	35	89	131	86	44	100	*25
493.20	189	63	126	*	*30	59	100	66	35	72	*
493.21	66	*	*52	—	*	*30	*31	*	*	*27	*
493.9	1,045	372	673	366	258	224	197	286	224	420	115
493.90	719	236	483	200	176	169	174	196	137	309	76
493.91	326	136	190	166	82	55	*24	90	87	111	39
494	*50	*	*29	*	*	*	*28	*	*	*	*
496	627	448	179	*	*	116	272	118	79	147	282
507	1,428	882	546	*42	*110	166	1,110	364	252	585	226
507.0	1,404	861	543	*40	*109	148	1,107	363	251	584	206
510	108	*67	*	*	*	*	*	*	*	*	*
510.9	99	*60	*	*	*	*	*	*	*	*	*
511	375	127	248	*	*38	81	254	86	67	137	*85
511.0	*26	*	*	*	*	*	*	*	*	*	*
511.8	*42	*	*	*	*	*	*	*	*	*	*
511.9	302	97	205	*	*	63	212	*58	*47	116	*80

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
512	291	170	121	*	90	88	109	74	*46	130	*41
512.1	66	*25	*42	*	*	*	*34	*	*	*	*
512.8	193	118	75	*	84	*	65	*37	*31	89	*
513	*57	*	*	*	*	*	*	*	*	*	*
513.0	*57	*	*	*	*	*	*	*	*	*	*
514	*32	*25	*	*	*	*	*	*	*	*	*
515	135	*43	92	*	*	*	91	*37	*23	*51	*
516	118	*51	*67	*	*	*	70	*	*	*32	*
516.3	*42	*	*	*	*	*	*	*	*	*	*
518	1,935	770	1,165	82	134	504	1,215	615	351	672	297
518.0	*31	*	*	*	*	*	*	*	*	*	*
518.4	*37	*	*	*	*	*	*	*	*	*	*
518.8	1,793	714	1,079	*69	121	456	1,147	567	326	633	267
518.81	1,464	576	888	*	88	346	986	461	280	533	190
518.82	70	*	*32	*	*	*	*	*	*	*28	*
518.84	175	*67	108	*	*	*77	87	*72	*	*	*
518.89	45	*23	*22	*	*	*	*	*	*	*	*
519	78	40	39	*22	*	*	*25	*20	*28	*	*
519.1	30	*	*17	*	*	*	*	*	*	*	*
520-579	14,856	6,446	8,411	723	2,861	4,005	7,267	3,793	3,173	5,349	2,542
522	*19	*	*	*	*	*	*	*	*	*	*
522.5	*13	*	*	*	*	*	*	*	*	*	*
524	20	*	*16	*	17	*	*	*	*	*	*
524.0	*11	*	*	*	*	*	—	*	*	*	*
527	*25	*	*	*	*	*	*	*	*	*	*
527.2	*20	*	*	*	*	*	*	*	*	*	*
528	*35	*	*	*	*	*	*	*	*	*	*
530	708	338	370	112	95	168	334	169	166	249	124
530.1	136	93	43	*	*19	35	65	*49	34	25	*29
530.10	38	*17	*20	*	*	*16	*	*	*	*	*
530.11	41	*25	*16	*	*	*13	*	*	*	*	*
530.19	*58	*51	*	*	*	*	*	*	*	*	*
530.2	76	44	*32	*	*	*	48	*19	*	*31	*
530.3	*51	*	*	*	*	*	*36	*	*	*	*
530.5	*13	*	*	*	*	*	*	*	*	*	*
530.6	*21	*	*	*	*	*	*	*	*	*	*
530.7	69	*28	*41	*	*	*	*47	*	*	*	*
530.8	319	131	189	88	48	82	101	72	74	108	65
530.81	287	115	172	88	47	70	82	64	66	99	59
530.82	*19	*	*	—	*	*	*	*	*	*	*
531	432	199	234	*	*37	106	287	92	124	148	68
531.0	116	*70	*46	*	*	*	66	*	*29	*58	*
531.00	111	*66	*45	*	*	*	64	*	*27	*58	*
531.4	208	94	114	*	*	41	149	*41	60	64	42
531.40	201	93	108	*	*	41	142	*40	59	64	38
531.9	48	*	*30	*	*	*	*30	*	*	*	*
531.90	43	*	*28	*	*	*	*27	*	*	*	*
532	505	316	189	*	51	141	313	173	80	148	104
532.0	99	*58	*41	*	*	*	74	*	*	*40	*
532.00	98	*57	*40	*	*	*	72	*	*	*40	*
532.4	287	198	89	—	*23	110	155	*118	*35	63	71
532.40	276	192	85	—	*22	106	148	*118	*32	59	67
532.5	*50	*	*	—	*	*	*	*	*	*	*
532.50	*50	*	*	—	*	*	*	*	*	*	*
532.9	*29	*	*	—	*	*	*	*	*	*	*
532.90	*19	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
533	48	*19	29	*	*	*	*30	*	*	*23	*
533.4	*19	*	*	—	*	*	*	*	*	*	*
533.40	*19	*	*	—	*	*	*	*	*	*	*
533.9	*12	*	*	*	*	*	*	*	*	*	*
533.90	*12	*	*	*	*	*	*	*	*	*	*
535	518	180	338	*	84	119	308	143	121	196	58
535.0	83	35	48	*	*16	*14	50	*21	*21	32	*
535.00	46	*	37	*	*15	*	*19	*	*12	21	*
535.01	*37	*26	*	*	*	*	*31	*	*	*	*
535.1	75	*	60	*	*	*	*54	*	*	*22	*
535.10	41	*	*36	*	*	*	*	*	*	*	*
535.11	*34	*	*	*	*	*	*	*	*	*	*
535.3	*20	*16	*	—	*	*	*	*	*	*	*
535.4	130	50	80	*	*	*28	79	*28	42	40	*
535.40	67	*21	46	*	*	*	41	*	*24	*20	*
535.41	63	*29	*34	*	*	*	39	*	*19	*	*
535.5	175	44	130	*	23	42	108	50	*32	81	*
535.50	105	*21	84	*	*18	25	59	*25	*14	56	*
535.51	70	*24	46	*	*	*	48	*25	*	*25	*
535.6	36	*20	*	*	*	*	*	*	*	*	*
535.60	*23	*	*	*	*	*	*	*	*	*	*
536	225	96	129	*27	51	40	107	*47	32	97	50
536.2	41	*17	*24	*	*	*	*	*	*	*23	*
536.3	48	*	*40	*	*	*	*	*	*	*	*
536.4	102	*53	*49	*	*	*	*68	*	*	*	*
536.42	*39	*	*	*	*	*	*	*	*	*	*
536.8	26	*15	*	*	*	*	*	*	*	*13	*
537	166	61	105	*	*	*	107	*50	*38	*43	*35
537.0	62	*	*39	*	*	*	*38	*	*	*	*
537.8	62	*26	*37	*	*	*	54	*	*	*	*
537.83	49	*	*31	—	*	*	42	*	*	*	*
540	846	491	355	193	424	149	80	157	194	261	235
540.0	266	155	111	78	115	48	*	47	68	75	76
540.1	202	102	101	*37	85	*49	*31	*34	57	74	*37
540.9	378	234	144	77	224	52	*24	76	69	111	122
541	29	*15	*	*	*15	*	*	*	*	*11	*
543	*11	*	*	*	*	*	*	*	*	*	*
550	148	124	*24	*12	*12	30	94	48	26	43	31
550.1	83	68	*	*	*	*	59	*30	*	*	*16
550.10	58	49	*	*	*	*	*41	*27	*	*	*
550.9	63	54	*	*	*	*17	33	*18	*13	*17	*15
550.90	40	32	*	*	*	*	*26	*	*	*	*
550.92	*16	*14	*	*	*	*	*	*	*	*	*
552	295	79	216	*	*18	99	177	83	68	103	*41
552.1	59	*	*39	*	*	*	*	*	*	*	*
552.2	159	37	122	—	*	55	98	53	*28	57	*
552.20	*48	*	*44	—	*	*	*43	*	*	*	*
552.21	105	*31	74	—	*	48	*52	*24	*	*45	*
553	254	97	158	*	48	87	114	53	46	130	26
553.1	*14	*	*	*	*	*	*	*	*	*	*
553.2	174	72	103	*	36	59	77	32	36	85	*21
553.20	33	*16	*	*	*	*	*	*	*	*	*
553.21	135	53	82	*	*24	47	63	*22	32	66	*15
553.3	65	*17	49	*	*	*24	*32	*	*	*34	*
555	256	101	154	*	142	58	*46	84	76	58	*38
555.0	92	*36	56	*	*50	*	*	*	*30	*	*
555.1	*45	*	*	*	*	*	*	*	*	*	*
555.2	*43	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
555.9	76	*29	48	*	41	*	*	*	*28	*	*
556	195	81	114	*	97	*39	*57	*61	*50	*42	*
556.9	96	*35	*61	*	*46	*	*	*	*	*	*
557	359	82	277	*	*	*78	221	*57	99	134	*69
557.0	200	*54	146	*	*	*	112	*	*62	*55	*
557.9	133	*	111	*	*	*	93	*	*	*70	*
558	458	155	304	89	74	89	207	91	91	196	80
558.1	*29	*	*	—	*	*	*	*	*	*	*
558.9	424	147	277	89	73	84	179	81	86	181	76
560	1,650	580	1,070	43	197	411	998	404	369	610	266
560.1	218	97	121	*	*29	54	133	66	*37	96	*
560.2	126	*63	*62	*	*	*	*109	*	*	*	*
560.3	144	38	107	*	*	*	108	*37	*33	55	*
560.39	122	36	86	*	*	*	87	*32	*32	41	*
560.8	656	225	431	*	111	161	368	113	171	244	128
560.81	493	163	330	*	81	129	268	*81	126	183	103
560.89	163	62	101	*	*	*	100	*	*45	*61	*
560.9	494	154	339	*	41	167	273	138	97	167	91
562	1,455	512	943	—	133	336	985	412	355	490	198
562.1	1,431	507	925	—	127	333	972	408	341	485	198
562.10	174	52	122	—	*	*35	121	*40	*37	77	*
562.11	810	303	507	—	96	263	452	236	227	218	130
562.12	357	114	243	—	*	*24	331	113	54	154	36
562.13	90	*38	52	—	*	*	68	*	*23	*36	*
564	236	76	159	29	36	53	118	77	57	54	48
564.0	93	32	61	*13	*	*19	52	*18	29	30	*17
564.1	94	*	78	*	*21	*	*48	*	*	*17	*
566	72	46	*	*	*20	*34	*	*	*	*38	*
567	152	87	*65	*	*	*59	*60	*	*	*58	*
567.2	113	*76	*	*	*	*48	*	*	*	*	*
568	78	*	68	*	*25	*23	*	*	*	*25	*
568.0	57	*	53	*	*	*	*	*	*	*24	*
569	640	314	326	*	60	144	415	196	110	251	83
569.1	*37	*	*26	*	*	*	*	*	*	*	*
569.3	60	31	*29	*	*	*	44	*	*	*25	*
569.4	39	*	*17	*	*	*	*27	*	*	*	*
569.49	*19	*	*	*	*	*	*	*	*	*	*
569.6	114	*59	55	*	*	*	*67	*	*	*36	*
569.69	79	*	*35	*	*	*	*55	*	*	*	*
569.8	347	167	179	*	*	*90	226	111	*61	141	*34
569.81	*85	*	*	*	*	*	*	*	*	*	*
569.82	*43	*	*	*	*	*	*	*	*	*	*
569.83	*89	*	*72	*	*	*	*	*	*	*	*
569.85	84	*45	*39	—	—	*	78	*	*	*	*
570	*40	*	*	*	*	*	*	*	*	*	*
571	541	322	218	*	113	279	146	138	96	231	76
571.1	95	*65	*	*	*38	*54	*	*	*	*56	*
571.2	241	167	75	—	*41	137	*63	74	*46	79	*42
571.5	173	77	96	*	*	76	73	*44	*27	84	*
572	222	128	94	*	*	120	67	79	*	70	*54
572.2	131	61	70	—	*	71	*46	*53	*	*46	*
572.3	*24	*	*	—	*	*	*	*	*	*	*
573	61	*23	*39	*	*23	*	*	*	*	*	*
573.3	*43	*	*	*	*	*	*	*	*	*	*
574	1,316	486	830	*	308	365	632	364	257	467	228
574.0	435	193	242	*	91	115	227	145	82	152	56

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
574.00	396	172	224	*	85	106	204	137	72	142	45
574.01	*39	*	*	*	*	*	*	*	*	*	*
574.1	472	130	342	*	124	160	181	121	76	180	95
574.10	456	120	336	*	122	152	174	120	70	177	89
574.2	104	41	63	*	21	36	47	*25	22	41	*15
574.20	92	31	60	*	20	33	38	*16	*21	39	*15
574.5	65	*	42	*	*	*	40	*	*	*	*
574.50	45	*	*30	*	*	*	*26	*	*	*	*
574.7	76	*	52	*	*26	*	*37	*	*31	*	*
574.70	*38	*	*29	—	*	*	*	*	*	*	*
574.71	*38	*	*	*	*	*	*	*	*	*	*
574.8	*56	*	*	—	*	*	*	*	*	*	*
574.80	*34	*	*	—	*	*	*	*	*	*	*
575	289	131	159	*	80	89	119	74	60	131	*24
575.0	84	47	37	—	*17	*23	*44	*	*	33	*
575.1	162	66	95	*	47	52	63	*43	*27	82	*
575.10	*16	*	*	—	*	*	*	*	*	*	*
575.11	105	*41	64	*	35	*41	*28	*36	*	48	*
575.12	*40	*	*	—	*	*	*	*	*	*	*
575.8	*23	*	*	*	*	*	*	*	*	*	—
576	152	62	91	*	*	38	85	*50	*33	*39	*31
576.1	61	*31	*30	*	*	*	*28	*	*	*	*
576.2	44	*	*30	*	*	*	*22	*	*	*	*
576.8	*38	*	*	*	*	*	*	*	*	*	*
577	1,344	692	652	*25	453	511	356	284	273	509	278
577.0	1,172	604	567	*	399	428	324	252	232	435	253
577.1	123	62	61	*	*42	59	*	*	*	61	*
578	900	398	502	*	46	195	652	224	186	342	148
578.0	94	48	46	*	*	*18	68	*34	*25	*21	*
578.1	132	62	70	*	*	*27	97	*40	*32	38	*22
578.9	674	288	386	*	34	149	487	151	130	283	111
579	*51	*	*	*	*	*	*	*	*	*	*
580-629	6,566	2,205	4,361	271	1,679	1,490	3,126	1,674	1,376	2,526	990
581	*30	*	*	*	*	*	*	*	*	*	*
584	929	495	434	*	61	215	642	350	151	263	165
584.5	*74	*	*	*	*	*	*	*	*	*	*
584.9	835	434	401	*	*53	174	602	306	133	253	143
585	167	85	82	*	*30	42	92	*33	*32	77	*
586	*54	*	*	—	*	*	*29	*	*	*	*
590	466	86	379	79	164	88	135	90	107	176	92
590.1	264	37	227	38	91	49	86	58	68	93	46
590.10	264	37	227	38	91	49	86	58	68	93	46
590.8	178	*39	139	40	61	37	40	*27	36	72	43
590.80	178	*39	139	40	61	37	40	*27	36	72	43
591	*34	*	*	*	*	*	*	*	*	*	*
592	422	211	212	*	161	164	87	109	120	163	130
592.0	127	45	82	*	51	37	33	40	29	48	*
592.1	294	165	128	*	110	126	52	68	90	114	21
593	147	56	91	*22	33	42	49	*25	*35	65	*22
593.4	38	*	*21	*	*	*	*	*	*	*19	*
593.7	*20	*	*16	*16	*	*	—	*	*	*	*
593.9	57	*24	*32	*	*	*	*31	*	*	*30	*
594	*25	*16	*	*	*	*	*17	*	*	*	*
594.1	*17	*15	*	*	*	*	*12	*	*	*	*
595	115	*52	62	*	*	*	86	*41	*	*26	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
595.9	*47	*	*	*	*	*	*31	*	*	*	*
596	110	61	*50	*	*	*	78	*	*	44	*
596.0	*25	*21	*	—	—	*	*20	*	*	*	*
596.5	*24	*	*	*	*	*	*	*	*	*	*
596.54	*21	*	*	*	*	*	*	*	*	*	*
598	*16	*15	*	*	*	*	*	*	*	*	*
599	1,846	653	1,194	108	135	265	1,338	422	345	818	262
599.0	1,689	572	1,117	107	114	229	1,239	378	303	771	238
599.7	109	60	49	*	*	*28	68	*35	*29	31	*
599.8	*14	*	*13	—	*	*	*	*	*	*	*
600	299	299	—	—	*	38	257	66	81	107	45
601	*21	*21	—	—	*	*	*	*	*	*	*
604	*26	*26	—	*	*	*	*	*	*	*	*
604.9	*22	*22	—	*	*	*	*	*	*	*	*
604.90	*17	*17	—	*	*	*	*	*	*	*	*
608	*32	*32	—	*	*	*	*	*	*	*	*
611	169	*	168	*	142	*17	*	134	*	*14	*13
611.0	*18	*	*	*	*	*	*	*	*	*	*
611.1	140	*	140	*	132	*	*	*126	*	*	*
614	171	—	171	*	139	*24	*	32	27	97	*
614.2	*39	—	*39	*	*	*	*	*	*	*	*
614.6	44	—	44	—	35	*	*	*	*	*24	*
614.9	49	—	49	*	47	*	*	*	*12	*23	*
616	39	—	39	*	*23	*	*	*	*	*20	*
616.0	*19	—	*19	—	*13	*	*	*	*	*	*
617	189	—	189	—	150	38	*	37	48	68	36
617.0	104	—	104	—	77	26	*	*16	30	41	*17
617.1	35	—	35	—	27	*	*	*	*	*	*
617.3	33	—	33	—	28	*	*	*	*	*	*
618	362	—	362	—	78	151	133	67	94	141	60
618.0	81	—	81	—	*17	36	28	*13	*24	30	*14
618.1	44	—	44	—	21	*15	*	*	*	*18	*
618.2	48	—	48	—	*12	*23	*	*	*	*	*
618.3	45	—	45	—	*	*18	*19	*	*	*23	*
618.4	48	—	48	—	*	22	*19	*	*14	20	*
618.5	61	—	61	—	*	25	34	*	*16	25	*
618.6	*11	—	*11	—	*	*	*	*	*	*	*
618.8	22	—	22	—	*	*	*	*	*	*	*
619	*26	—	*26	—	*	*	*	*	*	*	*
620	198	—	198	*	115	53	*24	43	46	82	27
620.0	*13	—	*13	*	*	*	—	*	*	*	*
620.1	33	—	33	*	30	*	—	*	*	*	*
620.2	128	—	128	*	60	43	*22	*33	29	53	*14
620.8	*12	—	*12	*	*	*	*	*	*	*	*
621	45	—	45	—	*22	*17	*	*	*	*14	*
621.3	*18	—	*18	—	*	*	*	*	*	*	*
622	*22	—	*22	—	*11	*	*	*	*	*	*
622.1	*20	—	*20	—	*	*	*	*	*	*	*
625	233	—	233	*	125	73	34	43	38	112	40
625.3	29	—	29	—	21	*	—	*	*	*16	*
625.6	101	—	101	—	29	40	33	*23	21	40	16
625.9	89	—	89	*	66	*22	*	*14	*	48	*16
626	241	—	241	*	164	77	*	33	70	107	32
626.2	173	—	173	*	117	56	—	23	57	69	25
626.8	57	—	57	*	42	*15	*	*	*12	34	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
627	39	—	39	—	*	*25	*	*	*	*	*
627.1	33	—	33	—	—	*23	*	*	*	*	*
630-677	1,267	—	1,267	*	1,264	*	—	258	243	498	267
632	23	—	23	—	23	*	—	*	*	*9	*
633	71	—	71	—	71	*	—	18	*11	26	*17
633.1	63	—	63	—	62	*	—	*14	*	24	*15
634	43	—	43	*	43	*	—	*18	*	14	*8
634.9	31	—	31	*	31	—	—	*16	*	*8	*
634.91	24	—	24	—	24	—	—	*15	*	*5	*
637	*12	—	*12	—	*12	—	—	*	*	*	*
637.9	*12	—	*12	—	*12	—	—	*	*	*	*
640	*15	—	*15	—	*15	—	—	*	*	*	*
641	59	—	59	—	58	*	—	*	*	*	*
641.1	*31	—	*31	—	*30	*	—	*	*	*	*
641.13	*31	—	*31	—	*30	*	—	*	*	*	*
642	62	—	62	*	62	—	—	*16	*	28	*
642.3	21	—	21	—	21	—	—	*	*	*12	*
642.33	21	—	21	—	21	—	—	*	*	*12	*
642.4	*22	—	*22	*	*22	—	—	*	*	*	*
642.43	*21	—	*21	*	*21	—	—	*	*	*	*
643	117	—	117	—	117	—	—	*20	34	48	*15
643.0	53	—	53	—	53	—	—	*	*12	30	*
643.03	53	—	53	—	53	—	—	*	*12	30	*
643.1	44	—	44	—	44	—	—	*	*	*14	*
643.13	44	—	44	—	44	—	—	*	*	*14	*
643.2	*18	—	*18	—	*18	—	—	*	*	*	*
643.23	*18	—	*18	—	*18	—	—	*	*	*	*
644	277	—	277	—	277	—	—	55	67	94	61
644.0	251	—	251	—	251	—	—	51	64	79	57
644.03	250	—	250	—	250	—	—	51	64	78	57
644.1	25	—	25	—	25	—	—	*	*	14	*
644.13	24	—	24	—	24	—	—	*	*	14	*
646	135	—	135	*	135	—	—	*23	*20	65	28
646.6	80	—	80	*	80	—	—	*	*	36	*20
646.63	75	—	75	*	74	—	—	*	*	32	*19
646.8	39	—	39	—	39	—	—	*	*	24	*
646.83	38	—	38	—	38	—	—	*	*	23	*
647	*22	—	*22	*	*21	—	—	*	*	*	*
648	208	—	208	*	208	*	—	39	47	80	42
648.0	*43	—	*43	—	*43	—	—	*	*	*	*
648.03	*42	—	*42	—	*42	—	—	*	*	*	*
648.8	28	—	28	—	28	*	—	*	*	*19	*
648.83	28	—	28	—	28	*	—	*	*	*19	*
648.9	102	—	102	*	102	—	—	25	*21	34	22
648.93	98	—	98	*	98	—	—	*24	*21	33	21
654	33	—	33	—	33	*	—	*	*	*15	*
654.5	*14	—	*14	—	*14	*	—	*	*	*	*
654.53	*14	—	*14	—	*14	*	—	*	*	*	*
658	*28	—	*28	—	*27	*	—	*	*	*	*
659	*11	—	*11	—	*10	*	—	*	*	*	*
670	*24	—	*24	—	*24	—	—	*	*	*	*
670.0	*24	—	*24	—	*24	—	—	*	*	*	*
670.04	*24	—	*24	—	*24	—	—	*	*	*	*
674	*23	—	*23	—	*23	—	—	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
680-709	2,848	1,418	1,430	219	581	790	1,257	643	588	1,234	383
681	82	52	*30	*	*17	*	*37	*	*27	*34	*
681.0	35	*26	*	*	*15	*	*	*	*	*	*
681.00	33	*24	*	*	*	*	*	*	*	*	*
681.1	47	*27	*	*	*	*	*	*	*	*	*
681.10	47	*27	*	*	*	*	*	*	*	*	*
682	1,774	895	879	80	377	554	762	501	390	623	260
682.0	70	42	*28	*15	*26	*	*	*18	*	*27	*
682.2	122	61	61	*	*32	*32	*44	*38	*21	44	*
682.3	175	73	102	*	50	58	63	53	*46	58	*17
682.4	87	46	41	*	*31	*23	*	*28	*16	*31	*
682.5	*32	*	*	*	*	*	*	*	*	*	*
682.6	988	491	497	*19	171	313	486	276	221	348	144
682.7	226	120	106	*	*28	84	104	49	55	78	*44
682.9	*31	*	*	*	*	*	*	*	*	*	*
693	*21	*	*	*	*	*	*	*	*	*	*
693.0	*19	*	*	*	*	*	*	*	*	*	*
695	*42	*	*	*	*	*	*	*	*	*	*
695.1	*26	*	*	*	*	*	*	*	*	*	*
707	689	314	375	*	106	169	407	98	126	377	88
707.0	453	204	249	*	*69	*116	264	*55	*72	264	*
707.1	212	101	112	*	*	*49	130	*39	*46	102	*
709	116	57	58	79	*30	*	*	*	*	109	*
709.2	94	45	49	63	*26	*	*	*	*	92	*
709.8	*21	*	*	*16	*	*	*	*	*	*18	*
710-739	6,324	2,644	3,679	166	972	1,885	3,300	1,407	1,549	2,258	1,109
710	136	*	107	*	56	*45	*	*	*32	*51	*
710.0	91	*	69	*	50	*	*	*	*	*34	*
711	160	89	*71	*	*26	*48	*66	*48	*36	*57	*
711.0	147	86	*61	*	*26	*43	*65	*45	*35	*57	*
711.06	*62	*46	*	*	*	*	*	*	*	*	*
714	95	*	66	*	*	*44	*41	*	*	*48	*
714.0	77	*	49	—	*	*39	*33	*	*	*39	*
715	1,994	787	1,208	*	49	515	1,430	417	503	746	329
715.1	173	67	106	*	*	51	117	44	*45	54	*30
715.15	37	*	*23	—	*	*	*26	*	*	*	*
715.16	128	47	80	*	*	39	85	*29	*39	41	*19
715.2	*34	*	*21	—	*	*	*	*	*	*	*
715.3	929	370	558	*	*23	261	643	186	238	352	152
715.35	267	102	165	—	*	96	159	80	63	73	51
715.36	642	261	382	*	*	157	475	104	171	270	98
715.8	*25	*	*	—	*	*	*	*	*	*	*
715.89	*25	*	*	—	*	*	*	*	*	*	*
715.9	828	325	504	—	*	189	627	167	198	330	133
715.90	*53	*	*	—	*	*	*44	*	*	*42	*
715.95	251	104	147	—	*	65	177	61	51	97	42
715.96	506	198	308	—	*	110	393	94	142	184	86
716	69	30	38	*	*	29	*30	*	*	31	*
716.1	*26	*	*	—	*	*	*	*	*	*	*
716.9	35	*	*20	*	*	*13	*	*	*	*	*
717	21	*11	*10	*	15	*	*	*	*	*	*
717.8	*11	*	*	*	*10	—	*	*	*	*	*
717.83	*10	*	*	*	*10	—	—	*	*	*	*
718	32	*12	*20	*	19	*	*	*	*	*	*
718.3	*13	*	*	*	*	*	*	*	*	*	*
718.8	*9	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
719	152	53	99	*	*26	37	85	*30	*23	64	35
719.4	112	37	75	*	*20	24	67	*21	*19	45	27
719.41	19	*	*13	—	*	*	*	*	*	*	*
719.45	59	*	48	*	*	*	53	*	*	*	*
719.46	*21	*	*	*	*	*	*	*	*	*	*
721	227	84	144	—	25	63	139	44	65	68	50
721.0	56	17	39	—	*9	20	*26	*	*	20	*19
721.1	52	*27	*	—	*	*16	*32	*	*	*	*
721.3	95	*31	64	—	*	*23	66	*24	*27	*28	*
722	971	459	513	*	315	377	278	190	255	340	186
722.0	122	58	64	—	40	69	*	*16	32	56	18
722.1	482	244	238	*	184	165	132	108	110	182	82
722.10	475	241	234	*	182	162	131	107	109	180	79
722.2	*12	*	*	—	*	*	*	*	*	*	—
722.4	*15	*	*	—	*	*	*	*	*	*	*
722.5	151	61	90	—	38	63	50	*24	46	45	*36
722.52	151	61	90	—	38	63	50	*24	46	45	*36
722.7	60	24	36	—	*20	*19	*	*	*	*14	*
722.71	27	*13	*15	—	*	*11	*	*	*	*	*
722.73	*18	*	*	—	*	*	*	*	*	*	*
722.8	*38	*	*	*	*	*	*	*	*	*	*
722.83	*37	*	*	*	*	*	*	*	*	*	*
722.9	85	*37	*	*	*	*	*	*	*	*	*
722.93	*77	*	*	*	*	*	*	*	*	*	*
723	53	25	28	*	*9	28	*	*	*	*19	*13
723.0	*19	*	*	—	*	*	*	*	*	*	*
723.1	*18	*	*	*	*	*	*	*	*	*	*
723.4	*13	*	*	—	*	*10	*	*	*	*	*
724	510	205	305	*	54	168	285	119	122	166	103
724.0	251	116	134	*	*	84	155	52	50	92	56
724.02	239	114	126	*	*	79	150	50	49	86	54
724.2	153	43	110	*	21	50	80	*	49	39	*25
724.3	*13	*	*	—	*	*	*	*	*	*	*
724.4	26	*11	*	*	*	*	*	*	*	*	*
724.5	43	*	28	*	*	*	*22	*	*	*	*
726	98	46	53	*	20	39	39	*14	27	41	*17
726.1	34	*11	22	—	*	*15	*15	*	*	*16	*
726.10	32	*11	*21	—	*	*15	*15	*	*	*15	*
726.2	16	*	*11	—	*	*9	*	*	*	12	*
726.3	*22	*	*	—	*	*	*	*	*	*	*
727	46	20	26	*	*	*15	*16	*	*	19	*
727.0	*21	*	*	*	*	*	*	*	*	*	*
727.6	12	*	*7	—	*	*	*	*	*	*8	*
727.61	10	*	*6	—	*	*	*	*	*	*8	*
728	149	89	60	*	*	*	78	*	*	*54	*
728.8	104	*74	*30	*	*	*	*44	*	*	*	*
728.89	78	*57	*	*	*	*	*40	*	*	*	*
729	149	62	87	*	39	52	53	33	31	72	*14
729.1	43	*	35	*	*	*	*	*	*	*	*
729.5	48	*26	*21	*	*	*21	*	*	*	*24	*
729.8	37	*15	*21	*	*	*	*	*	*	*17	*
729.81	*18	*	*	—	*	*	*	*	*	*	*
729.89	*18	*	*	*	*	*	*	*	*	*	*
730	513	315	198	*	112	161	205	156	119	165	*72
730.0	200	132	*68	*	*	*64	*70	*75	*	*60	*
730.07	119	*78	*	*	*	*	*45	*	*	*	*
730.1	*79	*59	*	*	*	*	*	*	*	*	*
730.2	233	123	109	*	*54	*65	105	*61	*67	*78	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
730.27	135	*50	*85	*	*	*	*87	*47	*	*	*
732	*15	*	*	*	*	*	*	*	*	*	*
733	669	212	458	*	93	152	417	153	154	233	130
733.1	404	96	308	*	*	84	302	99	103	135	67
733.13	246	*56	190	—	*	*	209	60	66	71	*48
733.14	76	*	61	*	*	*	*37	*	*	*32	*
733.19	*32	*	*29	—	—	*	*	*	*	*	*
733.4	111	64	47	*	*27	*32	*52	*	*20	44	*
733.42	102	60	41	*	*25	*28	*49	*	*	39	*
733.6	36	*	29	*	*	*	*20	*	*8	*17	*
733.8	88	37	51	*	31	*26	*29	*	*19	*27	*24
733.81	*17	*	*	*	*	*	*	*	*	*	*
733.82	71	*27	44	*	*22	*21	*27	*	*	*24	*
736	39	*24	*	*	*	*	*	*	*	*	*
737	92	*	73	*54	*	*	*	*	*	*	*
737.3	86	*	72	*53	*	*	*	*	*	*	*
737.30	*56	*	*48	*	*	*	*	*	*	*	*
738	101	*27	74	*	*33	*39	*	*27	*	*34	*
738.4	81	*	58	—	*	*36	*	*25	*	*	*
740-759	946	514	432	724	128	51	*44	265	181	284	215
742	*43	*	*	*27	*	*	—	*	*	*	*
745	147	*86	61	135	*	*	*	*32	*	*	*51
746	81	55	*	*64	*	*	*	*	*	*	*
747	112	*69	*44	*71	*	*	*	*	*	*21	*
747.6	*15	*	*	*	*	*	*	*	*	*	*
749	23	*14	*9	21	*	—	—	*	*	*9	*
749.0	*13	*	*	*12	*	—	—	*	*	*	*
750	46	32	*	40	*	—	*	*	*	*	*
750.5	24	21	*	24	—	—	—	*	*	*	*
751	124	*65	*59	107	*	*	*	*	*	*	*
752	*12	*	*	*	*	*	*	*	*	*	*
753	33	*	*	*	*	*	*	*	*	*	*
754	29	*	*	*23	*	*	—	*	*	*	*
756	157	84	*73	101	*24	*	*	*	*	*43	*43
756.1	52	*35	*	*	*	*	*	*	*	*	*
756.12	*38	*	*	—	*	*	*	*	*	*	*
757	12	*	*7	*7	*	*	—	*	*	10	—
757.3	10	*	*6	*6	*	—	—	*	*	10	—
757.32	10	*	*6	*6	*	—	—	*	—	10	—
760-779	1,578	994	583	1,576	*	—	—	260	312	517	489
765	626	424	202	626	—	—	—	*	*	324	*
765.1	414	302	*112	414	—	—	—	*	*	*220	*
769	312	*203	*	312	—	—	—	*	*	*	*
770	217	128	89	217	—	—	—	*49	*40	52	75
770.6	*44	*	*	*44	—	—	—	*	*	*	*
770.8	95	*56	*39	95	—	—	—	*	*	*	*34
771	90	*45	*45	90	—	—	—	*	*	*	*32
771.8	66	*40	*	66	—	—	—	*	*	*	*
774	115	48	66	115	—	—	—	*11	*12	*19	71
774.6	85	32	53	85	—	—	—	*	*	*15	54
775	*22	*	*	*22	—	—	—	*	*	*	*
777	*58	*	*	*58	—	—	—	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
778	*15	*	*	*15	—	—	—	*	*	*	*
779	55	*36	*	55	—	—	—	*	*	*	*
779.3	*26	*	*	*26	—	—	—	*	*	*	*
780-799	632	288	344	116	172	158	186	149	110	299	75
780	130	56	74	52	34	20	*25	34	23	62	*12
780.2	32	*	*23	*	*	*	*	*	*	*23	*
780.3	51	25	27	27	16	*	*	*14	*	23	*
780.39	48	23	25	24	16	*	*	*14	*	22	*
780.6	31	*17	*13	22	*	*	*	*11	*	*	*
784	*18	*	*	*	*	*	*	*	*	*	*
786	119	63	57	*	39	54	20	25	28	57	*10
786.5	107	55	52	*	37	53	17	22	26	50	*8
786.50	68	34	33	*	21	34	*13	13	18	32	*
786.59	35	20	15	*	13	18	*	*9	*	16	*
787	*8	*	*	*	*	*	*	*	*	*	*
788	*9	*	*	*	*	*	*	*	*	*	*
789	81	30	51	19	36	*14	*	17	15	39	*11
789.0	80	30	50	19	35	*14	*	17	14	37	*11
789.00	21	*	*15	*	*	*	*	*	*	*15	*
789.01	*10	*	*	*	*	*	*	*	*	*	*
789.03	23	*11	*12	*	15	*	*	*8	*	*	*
799	241	109	132	*25	44	49	123	54	*33	120	*33
799.9	238	109	129	*25	43	49	121	54	*31	120	*33
800-999	13,373	6,610	6,763	923	3,138	2,834	6,478	3,084	2,803	5,070	2,415
800	66	*52	*	*20	*	*	*	*	*	*	*
801	179	139	*40	*34	86	*	*	*42	*	*38	*69
801.0	*40	*26	*	*	*	*	*	*	*	*	*
801.1	*51	*	*	*	*	*	*	*	*	*	*
801.2	*52	*	*	*	*	*	*	*	*	*	*
802	153	127	*26	*	119	*	*	*18	*16	86	*33
802.2	37	*26	*	*	*26	*	*	*	*	*	*
802.4	*55	*50	*	*	*47	*	*	*	*	*45	*
805	361	169	191	*	95	62	192	49	86	150	76
805.0	74	*42	*32	*	*23	*	*25	*	*	*24	*
805.02	*36	*	*	*	*	*	*	*	*	*	*
805.2	87	*35	52	*	*	*	53	*	*	*40	*
805.4	185	86	99	*	51	*	108	*24	49	78	*33
807	228	97	131	*	39	43	144	*39	54	95	*40
807.0	179	67	112	*	*23	33	122	*32	37	76	*35
807.01	*29	*	*	*	*	*	*	*	*	*	*
807.02	*24	*	*	—	*	*	*	*	*	*	*
807.03	*20	*	*	—	*	*	*	*	*	*	*
807.05	*21	*	*	*	*	*	*	*	*	*	*
807.2	*20	*	*	—	*	*	*	*	*	*	*
808	316	129	187	*	72	*	201	66	76	147	*27
808.0	96	*72	*	*	*	*	*	*	*	*	*
808.2	174	*42	132	*	*	*	143	*28	*34	95	*
810	*32	*	*	*	*	*	*	*	*	*	*
810.0	*31	*	*	*	*	*	*	*	*	*	*
812	230	84	146	16	31	*55	128	60	51	88	32
812.0	131	*47	84	*	*	*38	91	*33	*31	*50	*
812.00	47	*	*28	*	*	*	*27	*	*	*	*
812.01	50	*	*37	—	*	*	*41	*	*	*	*
812.09	*24	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
812.2	52	*	*41	*	*	*	*27	*	*	*23	*
812.21	*25	*	*20	*	*	*	*	*	*	*	*
812.4	29	*13	*16	12	*	*	*	*	*	*	*
812.41	20	*8	*12	*10	*	*	*	*	*	*	*
813	327	240	86	20	52	34	221	195	34	65	33
813.0	41	21	*20	*	*11	*	*	*12	*	*13	*
813.01	*22	*12	*	*	*	*	*	*	*	*	*
813.2	*170	*166	*	*	*	*	*	*	*	*	*
813.4	63	26	36	*	*18	*10	29	*	*18	25	*12
813.41	*18	*	*12	*	*	*	*	*	*	*	*
813.42	*24	*	*14	*	*	*	*	*	*	*	*
813.44	19	*	*10	*	*	*	*	*	*	*	*
813.5	*15	*	*	*	*	*	*	*	*	*	*
816	20	*17	*	*	*	*	*	*	*	*	*
816.1	*17	*16	*	*	*	*	*	*	*	*	*
820	2,246	594	1,651	*	*29	153	2,056	465	456	921	404
820.0	651	132	520	*	*	*38	605	125	147	219	161
820.00	73	*	65	—	*	*	72	*	*	*	*
820.02	175	*	140	—	*	*	158	*	*	65	*
820.03	*38	*	*	*	*	*	*36	*	*	*	*
820.09	362	74	288	*	*	*	340	81	106	112	63
820.2	1,087	314	772	*	*	67	1,000	239	220	464	164
820.20	54	*	*39	—	*	*	44	*	*	*	*
820.21	942	283	659	*	*	*49	882	188	190	437	127
820.22	91	*	*75	*	*	*	*74	*	*	*	*
820.8	485	136	349	*	*	*48	428	97	81	231	75
821	403	186	217	68	119	*63	153	76	72	161	94
821.0	204	135	69	*60	*64	*	*45	*	*35	*70	*56
821.00	*35	*	*	*	*	*	*	*	*	*	*
821.01	169	117	*52	*58	*62	*	*	*	*30	*58	*52
821.2	142	*30	112	*	*	*	108	*	*	*64	*29
821.20	*36	*	*	*	*	*	*	*	*	*	*
821.23	63	*	*53	*	*	*	*49	*	*	*	*
822	46	*18	*28	—	*	*16	*25	*	*	*	*
822.0	44	*16	*28	—	*	*16	*25	*	*	*	*
823	378	224	154	*	150	137	81	89	81	152	56
823.0	137	60	76	*	*32	*56	*45	*	*40	*49	*19
823.00	106	48	*58	*	*21	*49	*35	*	*33	*37	*
823.2	78	47	*31	*	*23	*35	*	*	*	*24	*24
823.22	58	*39	*19	*	*17	*28	*	*	*	*	*21
823.3	*38	*	*	*	*	*	*	*	*	*	*
823.8	88	*57	*31	*	*57	*	*	*30	*	*51	*
823.82	62	*	*	*	*	*	*	*	*	*	*
824	452	196	256	*	120	186	135	111	82	158	101
824.0	*34	*	*	*	*	*	*	*	*	*	*
824.2	*21	*11	*	*	*	*	*	*	*	*	*
824.4	109	41	67	*	30	40	*38	*	20	49	*
824.6	110	*32	77	*	28	*35	47	43	*16	42	*
824.8	82	33	49	*	32	*15	*	*28	*	*28	*14
825	75	50	*	*	*31	*27	*	*	*	*29	*
825.0	*25	*17	*	—	*	*	*	*	*	*	*
825.2	*31	*17	*	*	*	*	*	*	*	*	*
825.25	*25	*	*	*	*	*	*	*	*	*	*
836	*9	*	*	*	*	*	*	*	*	*	*
840	29	14	15	—	*	*10	17	*	*10	*10	*
840.4	27	12	*15	—	*	*9	16	*	*9	*9	*
844	37	22	*	*	25	*	*	*	*	*18	*
844.2	*10	*	*	—	*10	*	—	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
844.8	*21	*14	*	—	*	*	*	*	*	*	*
847	52	*19	*33	*	*14	*	*26	*	*	*15	*
847.2	*29	*	*	—	*	*	*	*	*	*	*
850	91	58	33	14	29	*	*39	27	*13	26	*25
850.0	*17	*	*	*	*	*	*	*	*	*	*
850.1	25	*17	*	*9	*	*	*	*	*	*	*
850.5	*23	*15	*	*	*	*	*	*	*	*	*
850.9	*13	*	*	*	*	*	*	*	*	*	*
851	133	96	*37	*	*59	*	*45	*	*	*49	*
851.8	82	*56	*	*	*47	*	*	*	*	*34	*
852	271	162	109	*	*80	*41	134	*56	*70	95	50
852.0	*65	*	*	*	*	*	*	*	*	*	*
852.2	189	108	81	*	*46	*	119	*48	*	*61	*44
852.20	*59	*	*	*	*	*	*48	*	*	*	*
852.21	*48	*	*	*	*	*	*	*	*	*	*
853	*54	*	*	*	*	*	*	*	*	*	*
853.0	*52	*	*	*	*	*	*	*	*	*	*
854	44	*29	*	*	*	*	*	*	*	*	*
854.0	*40	*29	*	*	*	*	*	*	*	*	*
860	146	92	54	*	72	*	*45	*	*32	62	*30
860.0	72	*38	*34	*	*41	*	*	*	*	*28	*
861	*54	*43	*	*	*	*	*	*	*	*	*
861.2	*43	*	*	*	*	*	*	*	*	*	*
861.21	*42	*	*	*	*	*	*	*	*	*	*
863	80	*52	*	*	*42	*	*	*	*	*	*
864	*47	*41	*	*	*	*	*	*	*	*	*
864.0	*33	*	*	*	*	*	*	*	*	*	*
865	127	63	*	*	77	*	*	*	*	*	*
865.0	124	61	*	*	*75	*	*	*	*	*	*
868	*46	*36	*	*	*	*	*	*	*	*	*
873	63	35	28	*	31	*	*18	*17	*	*20	*15
873.0	*20	*	*	*	*	*	*	*	*	*	*
873.4	23	*10	*13	*	*	*	*	*	*	*	*
873.42	*14	*	*	*	*	*	*	*	*	*	*
881	*32	*21	*	*	*15	*	*	*	*	*	*
883	*17	*11	*	*	*13	*	*	*	*	*	*
886	*15	*12	*	*	*	*	*	*	*	*	*
886.0	*12	*	*	*	*	*	*	*	*	*	*
891	58	*39	*	*	*32	*	*	*	*	*28	*
891.0	*20	*	*	*	*	*	*	*	*	*	*
920	*16	*	*	*	*	*	*	*	*	*	*
922	44	*14	*30	*	*	*	*34	*	*	*	*
922.1	*21	*	*	*	*	*	*	*	*	*	*
924	69	*19	50	*	*	*	49	*	*16	*29	*17
924.0	40	*	*29	*	*	*	*29	*	*	*	*
924.01	35	*	*28	*	*	*	*27	*	*	*	*
933	*17	*	*	*	*	*	*	*	*	*	*
935	*15	*	*	*	*	*	*	*	*	*	*
935.1	*12	*	*	*	*	*	*	*	*	*	*
941	*52	*48	*	*	*	*	*	*	*	*	*
941.2	*28	*26	*	*	*	*	*	*	*	*	*
942	121	*97	*	*	*	*	*	*	*	*	*
942.2	*54	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
943	*50	*	*	*	*	*	*	*	*	*	*
944	*23	*15	*	*	*	*	*	*	*	*	*
945	95	68	*	*	*40	*	*	*	*	*58	*
945.2	*35	*	*	*	*	*	*	*	*	*	*
945.3	*51	*	*	*	*	*	*	*	*	*	*
948	177	*104	*73	146	*	*	*	*	—	171	—
958	*33	*	*	*	*	*	*	*	*	*	*
959	135	78	58	*	42	*22	62	*31	*19	63	23
959.0	85	52	33	*	29	*	*39	*	*	47	*14
959.01	83	50	33	*	28	*	*39	*	*	47	*13
962	*17	*	*	*	*	*	*	*	*	*	*
962.3	*17	*	*	*	*	*	*	*	*	*	*
965	116	36	80	*	76	*25	*	*18	29	44	25
965.0	36	*14	*22	*	*15	*	*	*	*	*17	*
965.09	*16	*	*10	*	*	*	*	*	*	*	*
965.1	*10	*	*	*	*	*	*	*	*	*	*
965.4	55	*	42	*	46	*	*	*	*15	*17	*
966	28	*	*21	*	*	*	*	*	*	*	*
967	*27	*	*23	*	*	*	*	*	*	*	*
968	24	*11	*	*	*14	*	*	*	*	*	*
968.5	*19	*	*	*	*11	*	*	*	*	*	*
969	134	50	84	*	89	31	*	43	25	43	*23
969.0	38	*13	25	*	24	*	*	*	*11	*13	*
969.4	40	*11	29	*	21	*14	*	*	*	*16	*
972	*69	*	*62	*	*	*	*	*	*	*	*
977	20	*	*10	*	*10	*	*	*	*	*	*
977.8	*8	*	*	*	*	*	*	*	*	*	*
977.9	*12	*	*	*	*	*	*	*	*	*	*
989	*26	*	*	*	*	*	*	*	*	*	*
989.5	*19	*	*	*	*	*	*	*	*	*	*
995	125	35	90	*	*21	59	38	34	54	32	*
995.0	*17	*	*	*	*	*	*	*	*	*	*
995.1	18	*	*13	*	*	*	*	*	*	*	*
995.2	69	*	*57	*	*	*	*26	*	*	*17	*
996	2,456	1,349	1,107	195	357	674	1,230	662	495	821	477
996.0	78	41	*37	*	*	*	49	*28	*	*21	*
996.01	33	*14	*18	*	*	*	27	*	*	*	*
996.04	*26	*	*	—	*	*	*	*	*	*	*
996.1	*31	*	*	*	*	*	*	*	*	*	*
996.2	113	46	66	61	*	*	*	*31	*46	*23	*
996.3	*91	*	*	*	*	*	*	*	*	*	*
996.4	426	165	261	*	*34	107	285	96	99	104	127
996.5	*71	*	*	*	*	*	*	*	*	*	*
996.59	*54	*	*	*	*	*	*	*	*	*	*
996.6	1,025	612	413	103	145	306	471	256	224	398	147
996.62	520	289	231	*63	74	164	220	149	113	191	*67
996.64	67	*36	*	—	*	*	*45	*	*	*	*
996.66	111	67	*44	—	*	*	74	*	*	*42	*
996.67	82	*45	*	*	*	*	*49	*	*	*39	*
996.68	*68	*	*	*	*	*	*	*	*	*	*
996.69	*35	*	*	*	*	*	*	*	*	*	*
996.7	481	241	240	*	79	135	255	105	89	168	119
996.73	167	70	97	*	*29	45	93	*24	*15	71	58
996.74	171	96	75	*	*	51	82	*45	*33	59	*34
996.77	41	*22	*	*	*	*	*	*	*	*	*
996.78	41	*	*26	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
996.8	141	82	*60	*	*57	*65	*	*47	*	75	*
996.81	71	*35	*36	*	*25	*41	*	*	*	*44	*
997	546	279	266	*	79	167	277	139	131	196	79
997.0	*38	*	*	*	*	*	*	*	*	*	*
997.1	45	*29	*	*	*	*	*28	*	*	*	*
997.2	*34	*	*	*	*	*	*	*	*	*	*
997.3	*52	*	*	*	*	*	*	*	*	*	*
997.4	197	84	113	*	49	59	80	60	*51	57	*28
997.5	*35	*22	*	*	*	*	*	*	*	*	*
997.6	140	83	*57	*	*	*52	80	*36	*	*47	*
997.62	95	*56	*	*	*	*	*57	*	*	*35	*
998	1,223	542	681	41	204	414	563	296	286	418	223
998.1	150	85	66	*	31	37	77	41	28	59	*22
998.11	98	69	30	*	*21	*23	52	*26	*16	46	*10
998.12	41	*13	*27	*	*10	*	*	*	*	*	*
998.3	119	*44	*75	*	*	*	*	*	*	*46	*
998.5	779	332	447	*32	115	245	387	200	175	245	159
998.59	753	321	432	*30	108	231	384	199	165	231	158
998.6	*67	*	*	*	*	*	*	*	*	*	*
998.8	65	*26	39	*	*22	*	*	*	*	*25	*
998.89	37	*16	*21	*	*19	*	*	*	*	*	*
999	*31	*	*	*	*	*	*	*	*	*	*
V01-V82	18,024	3,488	14,536	491	10,022	1,666	5,845	3,444	3,850	7,150	3,579
V10	*24	*	*19	*	*	*	*	*	*	*	*
V12	57	*	*40	*	*	*	*40	*	*	*35	*
V12.5	30	*	*18	*	*	*	*22	*	*	*21	*
V22	*17	—	*17	—	*16	*	—	*	*	*14	*
V22.2	*14	—	*14	—	*13	*	—	*	*	*12	*
V27	9,271	—	9,271	30	9,231	*	—	1,754	1,853	3,571	2,092
V27.0	8,450	—	8,450	29	8,413	*	—	1,573	1,580	3,334	1,962
V27.1	43	—	43	*	43	—	—	*	*	*22	*
V27.2	198	—	198	—	198	*	—	*34	42	75	47
V27.9	561	—	561	*	560	*	—	134	221	136	70
V30	*102	*	*	*102	—	—	—	*	*	*	*
V30.1	*102	*	*	*102	—	—	—	*	*	*	*
V42	*12	*	*	*	*	*	*	*	*	*	*
V43	*23	*	*	—	*	*	*	*	*	*	*
V45	69	34	*34	*	*	*24	*37	*23	*	*36	*
V45.8	46	*17	*29	*	*	*23	*	*	*	*24	*
V45.89	*34	*	*	*	*	*	*	*	*	*	*
V53	*12	*	*	*	*	*	*9	*	*	*	*
V53.3	*12	*	*	*	*	*	*9	*	*	*	*
V53.31	*8	*	*	*	*	*	*	*	*	*	*
V54	56	*	*	*	*	*	*	*	*	*	*
V55	221	111	110	*	*40	88	80	68	*50	71	*32
V55.2	*37	*	*	*	*	*	*	*	*	*	*
V55.3	169	87	82	*	*	69	*62	*57	*40	*44	*
V57	6,801	2,509	4,292	*	468	1,129	5,098	1,179	1,740	2,787	1,095
V57.1	1,775	600	1,176	*	155	280	1,310	519	189	832	235
V57.8	4,902	1,881	3,021	*	308	827	3,690	609	1,543	1,892	858
V57.89	4,902	1,881	3,021	*	308	827	3,690	609	1,543	1,892	858
V57.9	117	*	*89	—	*	*	94	*50	*	*	*
V58	1,034	576	458	174	160	318	382	223	143	421	247
V58.0	117	*53	64	*	*	*41	*66	*	*	62	*
V58.1	711	454	257	162	121	229	198	185	92	277	157

See footnotes at end of table.

Table 44. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
V58.4	140	51	89	*	*29	*42	64	*16	*	*42	*63
V58.49	137	50	87	*	*28	*40	64	*14	*	*41	*62
V58.8	65	*	*48	*	*	*	*53	*	*	*40	*
V58.89	65	*	*48	*	*	*	*53	*	*	*40	*
V59	*23	*	*	*	*	*	*	*	*	*	*
V70	30	*18	*11	*	*	*	*	*	*	*16	*
V70.7	28	*	*11	*	*	*	*	*	*	*	*
V71	18	*	*8	*	*	*	*	*	*	*	*
V72	*11	*	*	*	*	*	*	—	*	*	—

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 43; see Methods section.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	138,032	55,138	82,894	6,324	34,149	31,372	66,187	31,635	32,815	50,272	23,309
001-139	3,441	1,538	1,903	437	840	752	1,411	799	782	1,246	614
003	*7	*	*5	*	*	*	*	*	*	*	*
003.0	*5	*	*	*	*	*	*	*	*	*	*
005	*7	*	*	*	*	*	*	*	*	*5	*
005.9	*6	*	*	-	*	*	*	*	*	*	*
008	247	111	136	85	34	37	91	51	53	89	54
008.4	106	47	59	*5	16	19	65	30	27	29	20
008.45	98	41	57	*	15	17	61	29	24	28	17
008.6	46	27	19	45	*	*	*	*	*8	15	18
008.61	45	26	19	44	*	*	*	*	*8	15	18
008.8	90	34	55	32	16	17	24	15	16	43	16
009	18	*8	10	*	*6	*	*6	*	*6	*	*
009.0	10	*	*6	*	*	*	*	*	*	*	*
011	12	*8	*	*	*	*	*	*	*	*6	*
011.9	10	*6	*	*	*	*	*	*	*	*5	*
031	*9	*	*	*	*	*	*	*	*	*	*
034	22	13	*9	11	*9	*	*	*8	*6	*6	*
034.0	21	13	*9	10	*9	*	*	*8	*6	*6	*
038	637	316	321	43	78	136	380	169	136	233	99
038.0	42	24	18	*	*5	11	24	11	11	13	*8
038.1	127	74	53	*8	20	30	69	34	26	46	21
038.10	18	10	*7	*	*	*	11	*	*	*8	*
038.11	73	42	31	*	12	19	37	19	15	27	13
038.19	36	22	14	*	*	*8	21	11	*8	12	*6
038.2	21	*9	12	*	*	*	9	*5	*5	*9	*
038.4	142	67	75	*7	14	28	93	39	29	45	29
038.40	15	*7	*8	*	*	*	*8	*5	*	*6	*
038.42	72	26	46	*	*6	13	50	21	15	20	15
038.43	16	12	*	*	*	*	*7	*	*	*	*
038.49	36	20	16	*	*	*7	24	*8	*9	12	*8
038.8	20	*7	13	*	*	*	13	*7	*	*6	*
038.9	281	133	148	21	33	56	172	72	60	114	36
041	1,195	447	748	102	274	221	598	236	313	444	201
041.0	249	78	170	16	93	38	101	40	72	98	38
041.00	29	14	15	*	*	*6	15	*	*8	12	*
041.01	15	10	*5	*	*	*	*	*	*5	*5	*
041.02	70	*	65	*	61	*	*	14	19	29	*8
041.04	92	31	61	*	11	16	62	12	29	38	13
041.09	37	16	21	*	10	*7	14	*6	10	12	10
041.1	231	119	111	20	48	59	104	50	63	82	35
041.10	26	16	10	*	*6	*5	12	*8	*7	*8	*
041.11	168	86	82	14	37	47	71	35	44	63	26
041.19	36	16	20	*	*5	*6	22	*7	13	12	*6
041.2	*7	*	*	*	*	*	*	*	*	*	*
041.3	75	25	51	*	11	14	47	15	19	29	12
041.4	325	75	249	35	61	49	180	71	78	114	62
041.5	12	*6	*7	*	*	*	*5	*	*	*	*
041.6	45	15	29	*	*6	*6	33	*9	14	18	*
041.7	90	46	44	9	17	17	48	15	20	37	18
041.8	143	69	74	11	27	32	73	31	35	50	27
041.84	11	*	*6	*	*	*	*	*	*	*	*
041.85	49	24	25	*6	*7	9	26	10	11	18	10
041.86	58	29	29	*	13	14	30	15	15	19	10
041.89	21	9	12	*	*	*	11	*	*	*8	*
041.9	18	*9	9	*	*6	*	*6	*	*	10	*
042	130	88	41	*	84	42	*	45	15	50	20

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
047	33	19	14	16	12	*	*	*	*6	12	11
047.9	31	18	13	15	12	*	*	*	*6	12	10
052	*6	*	*	*	*	*	*	*	*	*	*
053	58	23	35	*	*5	*8	42	15	19	15	10
053.1	18	*7	11	*	*	*	16	*	*7	*	*
053.19	15	*6	9	*	*	*	14	*	*6	*	*
053.9	33	13	20	*	*	*6	22	9	11	*8	*
054	47	14	32	*7	24	*7	*9	*8	*9	20	10
054.1	16	*	13	*	14	*	*	*	*	*9	*
054.10	11	*	*9	—	10	*	*	*	*	*6	*
054.2	*6	*	*	*	*	*	*	*	*	*	*
054.9	16	*	12	*	*8	*	*	*	*	*6	*
070	231	132	99	*	98	108	24	67	38	85	42
070.1	*7	*	*	*	*	*	*	*	*	*	*
070.3	38	23	14	—	16	18	*	12	*7	15	*
070.30	21	14	*7	—	10	*9	*	*	*	9	*
070.32	17	9	*7	—	*6	*9	*	*7	*	*6	*
070.4	*8	*	*	—	*	*5	*	*	*	*	*
070.44	*7	*	*	—	*	*	*	*	*	*	*
070.5	171	95	76	*	74	81	17	50	27	60	34
070.51	86	50	36	—	39	39	*8	23	16	31	17
070.54	84	45	39	—	34	42	*8	26	11	29	17
070.9	*5	*	*	*	*	*	*	*	*	*	*
075	14	*5	*9	*	*9	*	*	*	*	*	*
078	20	*8	12	*	11	*	*	*	*	9	*
078.1	10	*	*7	*	*7	*	*	*	*	*	*
078.10	*5	*	*	*	*	*	*	*	*	*	*
078.5	*7	*	*	*	*	*	*	*	*	*	*
079	165	77	88	88	35	15	26	37	32	54	42
079.6	13	*6	*7	12	*	*	*	*	*	*	*
079.8	*6	*	*	*	*	*	*	*	*	*	*
079.9	140	67	72	70	30	14	25	33	25	47	35
079.99	136	67	69	70	27	14	25	32	25	46	34
094	*6	*	*	—	*	*	*	*	*	*	*
098	*5	*	*	*	*	*	—	*	*	*	*
110	56	32	24	*	13	12	29	20	15	14	*8
110.1	30	16	14	*	*	*6	21	12	9	*6	*
110.3	*6	*	*	*	*	*	*	*	*	*	*
110.4	13	*9	*	*	*7	*	*	*	*	*	*
111	*6	*	*	*	*	*	*	*	*	*	*
112	298	119	179	35	61	76	125	71	65	112	50
112.0	161	68	94	21	31	45	65	43	34	57	28
112.1	20	—	20	*	10	*	*	*	*	10	*
112.2	29	14	16	*	*	*7	18	*6	*5	13	*
112.3	22	*6	16	10	*	*	*8	*	*8	*6	*6
112.4	*8	*5	*	*	*	*	*	*	*	*	*
112.5	*9	*5	*	*	*	*	*	*	*	*	*
112.8	40	18	22	*	*7	13	19	11	*9	15	*6
112.84	28	14	14	*	*5	*9	14	*8	*	12	*
112.89	*8	*	*	*	*	*	*	*	*	*	*
112.9	*8	*	*5	*	*	*	*	*	*	*	*
117	28	17	12	*	9	*7	10	*6	*	13	*5
117.3	11	*6	*6	*	*	*	*	*	*	*	*
117.5	*6	*	*	*	*	*	*	*	*	*	*
117.9	11	*6	*	*	*	*	*5	*	*	*6	*
131	11	*	11	*	9	*	*	*	*	*6	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
131.0	*9	*	*8	—	*7	*	*	*	*	*5	*
131.01	*5	—	*5	—	*	*	—	*	*	*	*
135	40	10	30	—	10	22	*9	13	*9	15	*
136	21	14	*7	*	11	*6	*	*6	*	*7	*
136.3	15	11	*	*	9	*	*	*	*	*5	*
136.9	*5	*	*	*	*	*	*	*	*	*	*
138	18	*8	10	*	*	*6	11	*5	*5	*	*5
140-239	4,237	1,752	2,485	139	649	1,395	2,054	1,061	923	1,500	754
140	*6	*	*	—	—	*	*5	*	*6	*	—
141	*7	*6	*	—	*	*	*	*	*	*	*
146	*8	*8	*	—	*	*	*	*	*	*	*
150	25	19	*6	—	*	*8	16	*7	*	11	*
150.5	*6	*	*	—	*	*	*	*	*	*	*
150.9	14	11	*	—	*	*5	*9	*	*	*8	*
151	40	24	16	—	*6	13	21	9	*8	9	14
151.0	13	11	*	—	*	*	*7	*	*	*	*
151.8	*5	*	*	—	*	*	*	*	*	*	*
151.9	10	*5	*5	—	*	*	*	*	*	*	*
152	*8	*	*	—	*	*	*	*	*	*	*
153	177	91	87	*	*7	51	120	36	52	61	29
153.0	*7	*	*	—	*	*	*5	*	*	*	*
153.1	13	*5	*8	—	*	*	*8	*	*	*	*
153.2	*9	*5	*	—	*	*	*5	*	*	*	*
153.3	35	17	17	—	*	12	21	*7	11	12	*5
153.4	24	11	14	—	*	*8	17	*	*	9	*6
153.6	26	11	14	—	*	*6	19	*	*7	11	*
153.7	*7	*5	*	—	*	*	*	*	*	*	*
153.8	*5	*	*	—	*	*	*	*	*	*	*
153.9	50	27	23	*	*	12	36	10	19	14	*6
154	65	37	28	*	*	21	41	15	15	21	14
154.0	16	11	*6	—	*	*5	10	*	*	*	*
154.1	44	24	19	*	*	13	28	10	10	15	*9
155	26	16	11	*	*	13	11	10	*	*9	*
155.0	14	*8	*6	*	*	*6	*6	*	*	*	*
155.1	*7	*	*	—	*	*	*	*	*	*	*
155.2	*6	*	*	—	*	*	*	*	*	*	*
156	10	*	*6	—	*	*	*8	*	*	*	*
157	50	25	26	*	*	14	35	14	10	16	9
157.0	14	*6	*8	*	*	*	*9	*	*	*	*
157.8	*5	*	*	*	*	*	*	*	*	*	*
157.9	25	12	14	—	*	*5	19	*6	*6	*9	*5
158	*6	*	*	*	*	*	*	*	*	*	*
161	17	10	*7	—	*	*7	*8	*6	*	*7	*
161.9	9	*6	*	—	*	*	*	*	*	*	*
162	338	193	145	*	11	101	222	69	82	139	48
162.2	13	*7	*6	*	*	*	9	*	*	*6	*
162.3	69	42	28	—	*	21	46	18	16	22	13
162.4	*8	*	*	*	*	*	*	*	*	*	*
162.5	26	13	12	—	*	*8	17	*5	*7	*7	*6
162.8	28	13	15	—	*	10	17	13	*7	*	*
162.9	193	113	80	*	*7	56	128	30	47	94	21
170	39	30	*9	22	10	*	*	17	*6	12	*
170.7	23	20	*	18	*	*	*	15	*	*	*
171	19	*8	11	*	*	*6	*7	*	*	*9	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
171.9	*5	*	*	*	*	*	*	*	*	*	*
172	15	*6	*9	—	*	*7	*6	*5	*	*6	*
172.9	*6	*	*	—	*	*	*	*	*	*	*
173	19	10	10	*	*	*	14	*7	*	*6	*
174	183	—	183	—	21	66	96	48	43	59	33
174.2	*5	—	*5	—	*	*	*	*	*	*	*
174.3	*6	—	*6	—	*	*	*	*	*	*	*
174.4	23	—	23	—	*	*7	14	*6	*6	*5	*6
174.8	34	—	34	—	*	15	15	14	*6	*7	*7
174.9	102	—	102	—	12	36	54	23	23	41	15
176	*6	*	*	*	*	*	*	*	*	*	*
179	*7	—	*7	—	*	*	*	*	*	*	*
180	36	—	36	—	14	12	9	*7	*6	16	*7
180.8	*6	—	*6	—	*	*	*	*	*	*	*
180.9	27	—	27	—	11	9	*7	*	*5	14	*
182	36	—	36	—	*	14	18	*9	*8	13	*7
182.0	35	—	35	—	*	14	18	*9	*8	12	*7
183	54	—	54	*	*9	22	24	16	12	18	*9
183.0	52	—	52	*	*9	21	23	14	11	18	*8
184	*7	—	*7	*	*	*	*	*	*	*	*
185	181	181	—	*	*	38	142	49	40	64	28
186	*6	*6	—	*	*5	*	*	*	*	*	*
186.9	*6	*6	—	*	*5	*	*	*	*	*	*
188	63	47	16	*	*	*9	53	22	14	20	*7
188.8	15	11	*	*	*	*	11	*6	*	*	*
188.9	29	21	*8	*	*	*	25	*9	*8	11	*
189	56	31	25	*	*	22	29	12	13	24	*6
189.0	50	27	23	*	*	21	25	10	13	22	*6
191	50	26	25	10	12	16	13	12	*8	17	13
191.1	*7	*	*	*	*	*	*	*	*	*	*
191.8	*7	*	*	*	*	*	*	*	*	*	*
191.9	13	*7	*7	*	*	*	*	*	*	*6	*
192	*6	*	*	*	*	*	*	*	*	*	*
193	20	*	16	*	*7	9	*	*5	*	*9	*
194	*6	*	*	*	*	*	*	*	*	*	*
195	16	*8	*8	*	*	*8	*6	*6	*	*7	*
195.0	9	*7	*	*	*	*	*	*	*	*	*
196	163	66	97	*	18	61	82	45	36	47	35
196.0	22	9	13	*	*	9	*7	*7	*	*7	*
196.1	21	11	11	*	*	*8	11	*7	*	*6	*
196.2	63	32	31	*	*	21	37	14	16	17	16
196.3	32	*	30	—	*5	14	12	10	*7	*9	*6
196.6	12	*6	*7	*	*	*	*8	*	*	*	*
196.9	*8	*	*	—	*	*	*	*	*	*	*
197	456	202	254	*8	44	157	248	126	99	145	86
197.0	101	45	56	*	16	34	48	25	24	32	20
197.1	15	9	*6	*	*	*	*6	*	*	*	*6
197.2	58	19	39	*	*7	17	34	16	11	22	9
197.4	10	*5	*5	—	*	*	*6	*	*	*	*
197.5	11	*	*6	*	*	*5	*5	*	*	*	*
197.6	56	19	37	—	*6	19	31	16	10	15	14
197.7	182	90	92	*	12	65	105	51	42	59	30
197.8	19	*9	10	*	*	*6	12	*6	*	*5	*
198	400	193	208	*8	47	147	198	106	93	133	68

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
198.1	16	10	*6	—	*	*	11	*6	*	*	*
198.2	*7	*	*	*	*	*	*	*	*	*	*
198.3	92	43	50	*	13	46	33	27	20	29	16
198.4	*9	*	*6	*	*	*	*	*	*	*	*
198.5	164	84	80	*	15	49	96	38	42	52	32
198.7	12	9	*	—	*	*6	*5	*	*	*	*
198.8	94	39	55	*	13	36	44	26	18	36	14
198.82	14	*	10	*	*	*	*8	*	*	*6	*
198.89	78	35	44	*	11	32	35	22	15	30	11
199	161	65	96	*	17	53	89	42	32	63	25
199.0	71	26	45	*	9	28	33	20	10	31	11
199.1	90	39	51	*	*8	25	56	22	22	32	14
200	25	13	12	*	*	*8	12	*7	*	*8	*
200.0	16	*8	*8	*	*	*6	*9	*	*	*7	*
200.00	*7	*	*	*	*	*	*	*	*	*	*
201	16	*8	*8	*	*5	*	*7	*	*	*6	*
201.9	14	*6	*7	*	*	*	*7	*	*	*	*
201.90	10	*	*6	*	*	*	*6	*	*	*	*
202	109	58	50	*	11	37	59	28	25	39	16
202.0	*6	*	*	—	*	*	*	*	*	*	*
202.1	*5	*	*	*	*	*	*	*	*	*	*
202.8	94	51	43	*	9	33	52	25	22	35	12
202.80	51	28	23	*	*	19	27	15	12	18	*6
202.81	*6	*	*	*	*	*	*	*	*	*	*
202.82	*7	*5	*	—	*	*	*	*	*	*	*
202.83	16	*7	9	—	*	*5	9	*	*	*7	*
202.88	*8	*	*	—	*	*	*	*	*	*	*
203	63	32	30	—	*	20	40	17	13	22	11
203.0	62	32	30	—	*	20	39	16	13	22	11
203.00	61	31	30	—	*	20	39	16	13	22	11
204	89	50	39	31	*6	*7	45	19	22	27	21
204.0	37	21	15	31	*	*	*	*8	*6	11	12
204.00	23	13	10	19	*	*	*	*5	*	10	*
204.01	13	*8	*5	12	*	*	*	*	*	*	*7
204.1	51	28	23	—	*	*6	44	11	15	15	9
204.10	49	27	22	—	*	*6	42	11	14	15	9
205	48	31	17	*7	*9	16	17	14	*9	15	10
205.0	33	22	10	*5	*7	13	*8	11	*6	*9	*7
205.00	28	18	9	*	*5	11	*8	10	*5	*8	*
205.01	*5	*	*	*	*	*	*	*	*	*	*
205.1	13	*6	*6	*	*	*	*7	*	*	*	*
205.10	12	*6	*6	*	*	*	*7	*	*	*	*
208	11	*6	*	*	*	*	*6	*	*	*6	*
208.9	*6	*	*	—	*	*	*	*	*	*5	*
210	*5	*	*	*	*	*	*	*	*	*	*
211	164	68	96	*	11	37	116	40	35	62	27
211.1	15	*6	9	—	*	*	10	*	*	*6	*
211.3	124	53	70	*	*6	28	90	30	27	46	21
211.4	15	*5	10	—	*	*	12	*	*	*7	*
212	*6	*	*	*	*	*	*	*	*	*	*
213	9	*6	*	*	*6	*	*	*	*	*	*
214	19	*8	10	*	*	*6	*7	*	*	*8	*
214.1	*6	*	*	*	*	*	*	*	*	*	*
214.8	*5	*	*	*	*	*	*	*	*	*	*
215	*8	*	*5	*	*	*	*	*	*	*	*
216	*8	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
217	*8	*	*8	*	*	*	*	*	*	*	*
218	474	—	474	—	233	221	21	96	97	191	91
218.0	55	—	55	—	25	28	*	14	*8	20	12
218.1	135	—	135	—	61	68	*6	29	24	57	25
218.2	73	—	73	—	34	37	*	17	11	25	20
218.9	212	—	212	—	112	88	12	35	54	89	34
219	*6	—	*6	—	*	*	*	*	*	*	*
220	62	—	62	*	29	22	10	15	14	21	12
225	35	13	22	*	*	13	18	11	*8	9	*7
225.0	*5	*	*	*	*	*	*	*	*	*	*
225.2	25	*8	17	—	*	9	14	*7	*	*7	*5
226	14	*	12	—	*	*5	*	*	*	*6	*
227	25	*7	18	*	*6	10	9	*7	*	*7	*8
227.0	*6	*	*	*	*	*	*	*	*	*	*
227.1	9	*	*7	—	*	*	*	*	*	*	*
227.3	9	*	*5	*	*	*	*	*	*	*	*
228	17	9	*8	*6	*	*5	*	*	*	*8	*
228.0	15	*7	*8	*	*	*	*	*	*	*7	*
230	*6	*	*	—	*	*	*	*	*	*	*
233	41	*	39	—	13	15	14	*7	10	15	9
233.0	23	*	23	—	*	10	11	*6	*	*8	*5
233.1	14	—	14	—	10	*	*	*	*	*6	*
235	15	*6	*9	*	*	*	*9	*	*	*7	*
235.2	*7	*	*	*	*	*	*	*	*	*	*
236	*7	*	*	*	*	*	*	*	*	*	*
237	13	*7	*6	*	*	*	*	*	*	*	*
237.7	*6	*	*	*	*	*	*	*	*	*	*
238	69	33	36	*7	*	11	46	16	18	20	15
238.4	14	10	*	*	*	*	*7	*	*5	*	*
238.7	49	21	28	*6	*	*	36	11	12	15	11
239	26	12	14	*	*	*6	16	*8	*	10	*
240-279	14,727	5,992	8,735	486	1,843	4,236	8,162	3,411	3,586	5,442	2,287
240	11	*	10	*	*	*	*6	*	*	*	*
240.9	11	*	10	*	*	*	*6	*	*	*	*
241	29	*	25	*	*7	11	11	*6	*	14	*
241.0	*9	*	*8	*	*	*	*	*	*	*	*
241.1	17	*	15	—	*	*7	*6	*	*	*9	*
242	67	18	49	*	18	20	29	17	14	23	13
242.0	18	*	14	*	*7	*6	*	*	*	*5	*
242.00	18	*	13	*	*6	*6	*	*	*	*5	*
242.9	43	13	30	*	*9	13	21	11	9	15	*7
242.90	40	13	28	*	*8	13	19	11	*8	14	*7
244	1,360	259	1,102	*	140	297	919	334	365	437	224
244.0	46	*5	41	*	*5	14	27	*8	9	19	10
244.1	11	*	10	—	*	*	*6	*	*	*	*
244.9	1,296	250	1,045	*	131	277	883	320	353	412	210
245	14	*	12	*	*6	*	*	*	*	*6	*
245.2	10	*	*9	*	*	*	*	*	*	*	*
246	*9	*	*7	*	*	*	*	*	*	*	*
246.9	*6	*	*	*	*	*	*	*	*	*	*
250	4,658	2,056	2,602	28	485	1,553	2,592	1,149	1,064	1,781	664
250.0	3,337	1,417	1,921	13	283	1,051	1,990	821	758	1,324	435
250.00	2,582	1,104	1,478	*	184	795	1,601	644	602	987	349

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
250.01	438	177	262	*8	53	146	231	114	98	184	42
250.02	243	103	140	*	28	87	127	51	44	112	37
250.03	74	33	40	*	18	24	31	12	14	41	*6
250.1	129	69	60	13	71	29	16	26	34	52	17
250.10	27	15	11	*	11	*7	*8	*6	*6	12	*
250.11	45	22	23	*7	28	*7	*	10	13	12	*9
250.12	20	13	*7	—	*9	*8	*	*	*	*9	*
250.13	38	20	18	*	24	*7	*	*5	11	19	*
250.2	19	10	*9	*	*	*6	10	*6	*	*6	*
250.20	*7	*	*	*	*	*	*	*	*	*	*
250.22	*8	*	*	—	*	*	*	*	*	*	*
250.3	*7	*	*	*	*	*	*	*	*	*	*
250.4	289	129	159	—	29	118	141	66	61	87	74
250.40	173	80	92	—	*8	66	99	38	34	51	49
250.41	74	33	41	—	15	32	27	18	18	22	15
250.42	28	*9	19	—	*	14	12	*7	*	*9	*8
250.43	14	*7	*7	—	*	*6	*	*	*	*5	*
250.5	138	61	77	—	19	70	49	42	33	37	25
250.50	60	28	32	—	*	32	25	19	14	15	12
250.51	55	22	33	—	11	26	18	17	13	16	9
250.52	14	*6	*8	—	*	*8	*5	*	*	*	*
250.53	*9	*	*	—	*	*	*	*	*	*	*
250.6	355	175	180	*	45	143	166	97	87	116	55
250.60	185	92	93	—	12	72	101	51	44	59	32
250.61	94	47	47	—	19	39	37	32	19	31	12
250.62	47	23	24	—	*	22	22	11	16	14	*7
250.63	28	13	16	*	11	11	*7	*	*8	13	*
250.7	132	69	64	—	*7	47	78	35	26	49	22
250.70	78	38	40	—	*	24	52	19	17	29	13
250.71	36	20	16	—	*	16	18	13	*5	14	*
250.72	13	*7	*6	—	*	*5	*7	*	*	*	*
250.73	*5	*	*	—	*	*	*	*	*	*	*
250.8	215	108	107	*	21	72	121	50	51	86	28
250.80	118	63	56	—	*7	38	74	29	27	41	21
250.81	41	20	21	*	*5	17	18	13	*7	17	*
250.82	38	16	22	—	*	12	22	*6	11	18	*
250.83	18	10	*8	*	*6	*	*7	*	*5	10	*
250.9	37	16	21	*	*6	13	17	*	9	21	*
250.90	*7	*	*	—	*	*	*	*	*	*5	*
250.92	23	10	14	*	*	10	10	*	*7	12	*
250.93	*5	*	*	*	*	*	*	*	*	*	*
251	81	30	50	*5	12	20	45	21	15	34	11
251.2	40	13	27	*	*5	*7	24	*7	*8	18	*6
251.8	32	13	19	*	*6	*9	16	12	*5	11	*
252	28	*8	20	*	*6	*6	16	*5	*5	13	*
252.0	20	*7	13	—	*5	*	11	*	*	9	*
252.1	*8	*	*6	*	*	*	*	*	*	*	*
253	64	26	38	*6	13	17	28	17	11	23	12
253.2	13	*8	*5	*	*	*	*	*	*	*	*
253.5	12	*	*7	*	*	*	*	*	*	*	*
253.6	31	10	20	*	*	*8	19	*8	*6	11	*6
255	57	20	37	*	13	20	22	14	12	18	13
255.0	12	*	10	*	*	*	*	*	*	*	*
255.4	34	13	21	*	*7	10	16	*9	*7	11	*7
256	15	—	15	*	11	*	*	*	*	*6	*
256.4	10	—	10	*	*8	*	—	*	*	*	*
259	*8	*	*6	*	*	*	*	*	*	*	*
260	*8	*	*6	*	*	*	*5	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
261	21	10	11	*	*	*	13	*5	*5	*7	*
263	375	161	213	10	35	76	253	76	82	167	49
263.0	*5	*	*	*	*	*	*	*	*	*	*
263.1	11	*6	*5	*	*	*	*8	*	*	*5	*
263.8	19	10	9	*	*	*	13	*	*7	*7	*
263.9	340	143	196	9	33	70	228	68	73	154	45
266	38	14	23	*	*6	*	27	13	*5	15	*
266.2	33	13	20	*	*6	*	23	12	*5	12	*
269	*8	*	*	*	*	*	*6	*	*	*	*
269.9	*5	*	*	*	*	*	*	*	*	*	*
270	9	*7	*	*	*	*	*	*	*	*	*
271	25	*9	16	*	*8	*6	*9	10	*6	*6	*
271.3	23	*8	15	*	*8	*5	*8	*9	*6	*5	*
272	1,945	998	948	*	140	778	1,024	494	501	651	300
272.0	899	453	447	*	63	371	465	275	222	282	121
272.1	37	22	15	*	*8	19	*9	*5	10	14	*7
272.2	9	*	*	—	*	*	*	*	*	*	*
272.4	960	495	465	*	63	371	525	199	259	340	162
272.8	*6	*	*	*	*	*	*	*	*	*	*
272.9	28	17	11	*	*	*9	18	11	*6	*5	*5
273	59	20	39	*	*7	14	35	17	14	19	*9
273.1	*6	*	*	—	*	*	*	*	*	*	*
273.8	45	15	30	*	*6	11	25	12	10	16	*7
274	188	133	55	*	10	55	123	49	46	62	31
274.0	50	30	20	—	*	13	32	10	11	22	*7
274.9	137	103	34	*	*6	41	89	39	34	39	24
275	216	82	134	*5	45	61	106	45	65	78	29
275.0	11	*	*8	*	*6	*	*	*	*	*	*
275.2	89	35	55	*	19	28	42	16	32	35	*7
275.3	17	*7	9	*	*5	*5	*6	*	*	*6	*
275.4	96	34	61	*	14	25	54	20	28	31	17
275.41	42	13	29	*	9	12	18	*9	13	13	*7
275.42	43	15	28	*	*	11	28	*9	11	16	*6
275.49	*8	*	*	*	*	*	*7	*	*	*	*
276	4,558	1,825	2,733	378	598	908	2,675	940	1,145	1,745	727
276.0	108	53	55	*7	*8	15	78	24	25	42	17
276.1	646	253	393	17	54	144	432	129	169	240	108
276.2	213	97	116	21	36	55	101	38	45	88	42
276.3	46	20	25	*	*6	15	22	9	14	16	*7
276.4	16	*7	9	*	*	*	9	*	*	*	*
276.5	2,176	901	1,275	304	293	332	1,248	459	532	831	354
276.6	82	34	48	*	11	25	44	*7	22	39	15
276.7	321	155	166	*5	29	70	218	72	82	110	57
276.8	860	274	586	18	148	224	471	190	215	343	113
276.9	89	31	58	*	11	24	52	9	39	31	10
277	59	24	35	16	22	*9	12	13	15	18	13
277.0	33	13	21	13	16	*	*	*6	*9	11	*8
277.00	33	13	20	13	16	*	*	*6	*8	11	*8
277.3	*8	*	*6	*	*	*	*6	*	*	*	*
278	782	256	526	*7	244	344	187	156	187	287	152
278.0	767	253	514	*7	240	335	185	153	183	281	150
278.00	514	176	338	*6	142	224	142	101	127	183	104
278.01	253	77	176	*	97	111	43	52	56	98	46
278.1	*8	*	*7	—	*	*	*	*	*	*	*
278.8	*6	*	*	*	*	*	*	*	*	*	*
279	18	*8	10	*	*	*7	*	*	*	*8	*
279.0	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
280-289	4,158	1,626	2,532	188	877	921	2,172	954	969	1,547	687
280	652	234	418	17	114	132	390	146	175	235	97
280.0	382	139	242	*	61	81	239	82	103	136	60
280.9	264	92	172	15	51	49	148	62	70	96	36
281	95	37	59	*	*6	17	72	21	27	33	14
281.0	35	9	26	—	*	*5	30	*7	10	13	*
281.1	*9	*	*	—	*	*	*8	*	*	*	*
281.9	48	21	27	*	*5	10	33	11	14	17	*7
282	133	51	83	34	79	13	*7	48	26	48	11
282.4	18	*	14	*	9	*	*5	*9	*	*6	*
282.5	19	*	15	*	11	*	*	*6	*	*8	*
282.6	88	39	49	27	55	*	*	31	19	31	*6
282.60	12	*5	*7	*	*6	*	*	*	*	*6	*
282.61	*7	*	*	*	*	*	*	*	*	*	*
282.62	68	29	39	18	48	*	—	27	14	23	*
283	20	*8	12	*	*	*	11	*	*	*6	*
283.0	*8	*	*6	*	*	*	*	*	*	*	*
283.9	*8	*	*	*	*	*	*5	*	*	*	*
284	168	90	78	15	23	52	78	52	35	56	25
284.8	143	77	67	9	20	46	69	41	34	46	23
284.9	24	13	11	*5	*	*6	9	11	*	10	*
285	2,155	766	1,389	43	460	450	1,202	454	490	828	383
285.1	609	234	375	*7	119	132	350	121	174	204	109
285.8	37	15	22	*	*7	*7	22	*6	*8	16	*8
285.9	1,507	516	991	34	334	311	828	326	307	607	266
286	181	96	85	*7	36	56	83	45	40	66	30
286.0	*8	*6	*	*	*	*	*	*	*	*	*
286.3	10	*	*5	*	*	*	*	*	*	*	*
286.4	*7	*	*	*	*	*	*	*	*	*	*
286.5	15	*5	9	*	*	*5	*7	*	*	*8	*
286.6	18	*9	10	*	*	*	*9	*	*	*7	*
286.7	35	17	18	*	*	11	20	9	9	11	*
286.9	83	48	35	*	15	27	39	23	15	29	17
287	357	175	182	24	69	99	166	103	72	125	57
287.0	*5	*	*	*	*	*	—	*	*	*	*
287.3	39	16	23	*	10	9	16	12	*9	13	*
287.4	61	31	29	*	14	20	23	20	14	13	15
287.5	248	124	124	13	43	68	124	68	48	95	37
288	308	128	180	36	67	72	133	63	78	118	49
288.0	172	73	99	25	38	48	62	35	41	68	28
288.3	*6	*	*	*	*	*	*	*	*	*	*
288.8	120	48	72	9	25	22	64	24	36	43	17
288.9	*8	*	*	*	*	*	*	*	*	*	*
289	89	41	48	10	22	26	30	18	20	34	17
289.0	*8	*	*	*	*	*5	*	*	*	*	*
289.2	*6	*	*	*	*	*	*	*	*	*	*
289.3	10	*5	*	*	*	*	*	*	*	*	*
289.5	12	*6	*6	*	*	*	*	*	*	*	*
289.59	10	*5	*5	*	*	*	*	*	*	*	*
289.8	26	12	14	*	*6	*6	13	*	*7	10	*
289.9	21	*7	14	*	*	*7	10	*	*	10	*
290-319	9,688	4,702	4,985	300	4,256	2,493	2,639	2,635	2,252	3,536	1,266
290	248	84	164	—	*	9	237	63	54	98	34
290.0	117	33	83	—	*	*	112	39	27	40	10
290.1	*6	*	*	—	—	*	*5	*	*	*	*
290.2	17	*6	11	—	—	*	17	*	*7	*5	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
290.20	*8	*	*	—	—	*	*8	*	*	*	*
290.21	9	*	*6	—	—	*	9	*	*	*	*
290.3	21	*6	15	—	—	*	21	*	*	10	*
290.4	86	36	50	—	—	*	81	15	13	40	17
290.40	60	23	36	—	—	*	56	12	10	25	13
290.41	17	9	*8	—	—	*	17	*	*	9	*
290.42	*6	*	*	—	—	*	*6	*	*	*5	*
291	261	194	67	*	129	100	31	76	79	68	38
291.0	52	42	10	—	19	25	*7	16	11	19	*6
291.2	15	13	*	—	*	*7	*6	*	*5	*	*
291.3	*8	*6	*	—	*	*	*	*	*	*	*
291.8	181	131	51	*	104	62	15	54	59	42	26
291.81	166	122	44	*	93	57	15	50	56	36	24
291.89	15	*9	*7	—	10	*	*	*	*	*7	*
292	191	102	89	*	103	40	46	72	38	55	25
292.0	94	54	40	*	68	21	*	49	16	21	*8
292.1	11	*	*7	*	*	*	*6	*	*	*	*
292.12	10	*	*6	*	*	*	*	*	*	*	*
292.8	75	39	36	*	28	14	33	20	17	26	11
292.81	47	18	28	*	*7	*9	30	*8	13	17	*9
292.84	21	16	*	*	15	*	*	*8	*	*8	*
292.89	*7	*	*	—	*5	—	*	*	*	*	*
292.9	11	*6	*5	*	*	*	*	*	*	*	*
293	123	50	73	*	20	21	81	33	32	34	23
293.0	70	24	46	*	*	11	56	14	20	19	17
293.8	35	17	18	*	16	*7	12	16	*8	9	*
293.83	27	14	13	*	13	*	*9	14	*	*7	*
293.9	15	*8	*7	—	*	*	12	*	*	*5	*
294	684	251	433	*	*9	27	648	158	181	245	99
294.1	258	96	162	*	*	*7	248	64	75	89	30
294.8	395	140	255	—	*	13	380	87	101	144	64
294.9	30	13	16	*	*	*6	19	*7	*5	13	*5
295	439	245	195	*	232	137	68	133	84	177	46
295.2	*8	*7	*	—	*8	*	—	*	*	*8	—
295.20	*8	*7	*	—	*7	*	—	*	*	*7	—
295.3	115	77	38	*	62	38	15	41	24	35	15
295.30	54	38	16	*	31	16	*8	13	16	17	*8
295.32	19	13	*7	—	*8	*7	*	*7	*	*7	*
295.34	39	25	14	—	22	14	*	21	*	11	*
295.4	*7	*	*5	*	*6	*	*	*	*	*5	*
295.40	*6	*	*	*	*5	*	*	*	*	*5	*
295.6	32	17	15	—	16	11	*5	9	*7	12	*
295.60	*7	*	*	—	*	*	*	*	*	*	*
295.62	13	*8	*5	—	*5	*5	*	*	*	*6	*
295.64	11	*6	*5	—	*8	*	*	*	*	*	*
295.7	173	92	82	*	108	51	12	50	31	81	12
295.70	124	67	57	*	76	36	*9	23	25	67	*9
295.74	43	23	20	—	28	12	*	25	*	11	*
295.9	97	48	49	*	31	31	35	31	19	34	13
295.90	82	39	43	*	23	27	32	21	17	32	11
295.92	*5	*	*	—	*	*	*	*	*	*	*
295.94	*8	*7	*	—	*6	*	*	*7	*	*	*
296	1,237	526	711	59	704	305	169	291	236	624	86
296.0	*5	*	*	—	*	*	*	*	*	*	*
296.2	293	118	175	17	150	69	56	55	65	153	20
296.20	185	81	104	12	92	46	35	36	45	93	11
296.22	10	*	*7	*	*5	*	*	*	*	*6	*
296.23	42	16	26	*	24	*9	*6	*6	*8	23	*
296.24	52	16	36	*	28	12	11	11	*9	30	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
296.3	347	137	211	10	173	103	62	96	74	157	21
296.30	103	36	67	*	51	30	18	26	26	45	*6
296.32	19	*8	11	*	10	*	*	*	*6	*6	*
296.33	145	59	86	*	81	39	21	50	25	61	*9
296.34	73	32	42	*	28	27	17	16	13	42	*
296.4	95	45	50	*	57	26	*8	24	14	51	*7
296.40	48	24	24	*	31	10	*	*8	*8	29	*
296.43	*5	*	*	—	*	*	*	*	*	*	*
296.44	37	16	21	*	21	12	*	13	*	17	*
296.5	124	60	64	*	88	26	*7	19	18	83	*
296.50	77	35	42	*	57	14	*	10	11	54	*
296.53	21	*6	14	*	16	*	*	*	*	13	*
296.54	23	16	*7	—	14	*7	*	*	*	15	*
296.6	100	49	50	*8	74	13	*	17	13	66	*
296.60	49	26	24	*7	35	*	*	*7	*8	33	*
296.63	19	*9	10	*	15	*	*	*6	*	10	*
296.64	29	13	16	*	23	*5	*	*	*	22	*
296.7	171	70	101	*7	91	47	25	53	42	50	26
296.8	71	29	42	*9	48	12	*	9	*6	53	*
296.80	46	21	25	*6	29	9	*	*	*	38	*
296.89	22	*6	16	*	17	*	*	*	*	13	*
296.9	31	16	15	*	18	9	*	16	*	9	*
296.90	28	15	14	*	16	*8	*	15	*	*8	*
297	34	13	21	—	*5	*	24	*8	*7	16	*
297.1	12	*	*8	—	*	*	*8	*	*	*	*
297.9	20	*9	12	—	*	*	15	*	*	10	*
298	175	83	92	*	57	34	80	52	37	62	25
298.9	167	79	89	*	55	32	77	49	35	59	23
299	*9	*7	*	*	*	*	*	*	*	*	*
299.0	*5	*	*	*	*	*	—	*	*	*	*
300	861	265	596	18	272	256	315	231	210	315	105
300.0	473	144	329	*	116	149	204	127	121	173	53
300.00	393	117	276	*	78	127	186	108	104	139	43
300.01	54	16	38	*	27	16	10	14	13	21	*7
300.02	25	10	15	*	11	*6	*7	*	*	13	*
300.1	23	*5	18	*	12	*8	*	*	9	*6	*
300.11	10	*	*7	*	*	*	*	*	*	*	*
300.14	*6	*	*6	—	*	*	—	*	*	*	*
300.2	25	*5	20	*	18	*	*	*7	*5	10	*
300.21	18	*	16	*	14	*	*	*5	*	*8	*
300.3	24	10	14	*	13	*	*	*9	*5	*7	*
300.4	242	71	171	*6	64	72	99	73	50	88	31
300.8	*9	*	*7	*	*	*	*	*	*	*	*
300.9	64	28	36	*	44	12	*5	9	17	27	11
301	348	150	198	*7	242	75	25	145	73	107	23
301.0	9	*7	*	—	*6	*	*	*	*	*7	—
301.2	*9	*7	*	*	*6	*	*	*	*	*	*
301.5	*7	*	*6	*	*	*	*	*	*	*	*
301.50	*7	*	*5	*	*	*	*	*	*	*	*
301.6	31	9	22	*	16	*9	*6	*7	*9	15	*
301.7	24	21	*	*	22	*	*	*9	*8	*6	*
301.8	134	41	93	*	96	30	*	61	35	27	10
301.81	*9	*6	*	*	*5	*	*	*	*	*	*
301.83	85	11	74	*	68	14	*	30	26	22	*7
301.84	*5	*	*	*	*	*	*	*	*	*	—
301.89	35	20	15	*	20	12	*	28	*	*	*
301.9	123	59	64	*	86	28	9	62	14	39	*8
303	717	514	203	*	367	270	79	232	172	218	95
303.0	101	72	29	*	51	40	9	22	46	19	14

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
303.00	28	21	*7	*	12	13	*	*5	12	*6	*6
303.01	70	49	21	—	36	27	*6	17	32	13	*8
303.9	616	442	174	*	316	230	70	210	126	199	81
303.90	252	184	68	*	112	106	34	43	55	114	40
303.91	293	212	81	*	170	96	26	141	50	70	32
303.92	16	10	*7	*	12	*	*	*7	*7	*	*
303.93	55	36	19	—	21	24	*9	19	13	15	*8
304	461	258	204	*	346	99	13	203	77	135	47
304.0	157	91	66	*	100	50	*6	82	17	41	17
304.00	35	17	18	*	16	14	*	*6	*6	18	*
304.01	117	71	46	—	81	35	*	73	10	23	12
304.1	25	11	14	*	18	*5	*	*7	*5	11	*
304.10	13	*	*8	*	*9	*	*	*	*	*6	*
304.11	12	*6	*6	—	*8	*	*	*	*	*	*
304.2	91	53	37	*	69	21	*	40	13	30	*7
304.20	29	13	16	*	20	*9	—	*	*5	17	*
304.21	56	37	19	—	44	12	*	33	*6	13	*
304.3	57	40	17	*	53	*	*	16	22	15	*5
304.30	15	10	*5	*	14	*	—	*	*6	*6	*
304.31	37	27	10	*	34	*	*	13	13	*8	*
304.4	17	*6	11	—	14	*	*	*	*	*	*7
304.40	*5	*	*	—	*	*	—	*	*	*	*
304.41	*9	*	*6	—	*7	*	*	*	*	*	*
304.6	10	*	*7	*	*6	*	*	*	*	*	*
304.61	*6	*	*	*	*	*	*	*	*	*	*
304.7	34	17	16	*	28	*5	*	20	*	*7	*
304.70	*6	*	*	*	*	*	—	*	*	*	*
304.71	27	13	13	—	24	*	*	18	*	*5	*
304.8	37	22	16	*	35	*	—	18	*	14	*
304.80	19	10	*9	*	17	*	—	*	*	12	*
304.81	12	*7	*	*	11	*	—	*9	*	*	*
304.83	*6	*	*	—	*5	*	—	*	*	*	*
304.9	32	14	18	*	24	*6	*	15	*	9	*
304.90	17	*8	10	*	11	*	*	*	*	*8	*
304.91	11	*5	*6	—	9	*	*	*9	*	*	*
305	2,235	1,255	980	13	1,113	763	345	443	533	861	398
305.0	417	296	121	*	222	140	51	101	82	169	65
305.00	271	195	76	*	142	91	35	46	66	112	47
305.01	95	69	27	*	52	34	*9	27	11	47	11
305.02	24	16	*8	—	19	*	*	18	*	*	*
305.03	27	16	10	*	*8	11	*7	9	*	*8	*6
305.1	1,375	700	674	*	547	535	289	219	360	531	266
305.2	100	61	39	*	86	10	—	33	23	31	13
305.20	60	36	24	*	52	*6	—	14	19	19	*8
305.21	26	18	*9	*	24	*	—	11	*	9	*
305.22	*8	*	*	*	*7	*	—	*6	*	*	*
305.23	*6	*	*	*	*	*	—	*	*	*	*
305.4	15	*7	*8	—	12	*	*	*	*	10	*
305.40	11	*	*6	—	*8	*	*	*	*	*7	*
305.5	42	25	17	*	29	12	*	14	10	13	*5
305.50	24	15	9	—	17	*7	*	*7	*7	*7	*
305.51	13	*8	*6	*	9	*	*	*6	*	*	*
305.6	144	89	54	*	107	36	*	39	32	56	17
305.60	99	63	37	*	73	25	*	16	27	44	12
305.61	30	17	12	—	22	*8	—	16	*	*8	*
305.62	*6	*	*	—	*5	*	*	*	*	*	*
305.63	*9	*	*	—	*6	*	—	*	*	*	*
305.7	20	*9	11	*	18	*	—	*	*	*5	11
305.70	15	*7	*8	*	13	*	—	*	*	*	*9
305.9	118	64	54	*	89	25	*	33	22	45	19

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see Methods section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
305.90	82	44	38	*	59	19	*	16	17	35	15
305.91	17	10	*7	*	13	*	*	*6	*	*7	*
305.92	*7	*	*	–	*6	*	*	*6	*	*	–
305.93	13	*6	*6	*	10	*	–	*5	*	*	*
306	11	*	*7	*	*	*	*	*	*	*	*
307	71	26	45	*8	27	11	24	21	16	27	*7
307.1	*8	*	*7	*	*6	*	*	*	*	*	*
307.2	10	*7	*	*	*	*	*	*	*	*	*
307.23	9	*7	*	*	*	*	*	*	*	*	*
307.5	14	*	13	*	*9	*	*	*	*	*	*
307.50	*7	*	*7	*	*	*	*	*	*	*	*
307.51	*5	*	*5	–	*	*	*	*	*	*	*
307.8	15	*	11	*	*6	*	*	*	*	*6	*
307.81	13	*	10	*	*6	*	*	*	*	*6	*
307.9	20	13	*8	*	*	*	17	*6	*	*9	*
308	23	*8	15	*	9	*7	*6	*	*	11	*
308.3	9	*	*	*	*	*	*	*	*	*	*
308.9	*8	*	*5	*	*	*	*	*	*	*	*
309	233	91	142	16	154	38	26	81	67	61	25
309.0	66	30	36	*	45	10	11	25	18	18	*5
309.2	30	10	20	*	16	*5	*7	*7	12	*8	*
309.24	*6	*	*	*	*	*	*	*	*	*	*
309.28	23	*8	15	*	14	*	*	*5	9	*5	*
309.4	31	14	17	*	23	*	*	13	*7	*	*6
309.8	81	24	57	10	55	15	*	31	22	20	*8
309.81	79	24	55	10	53	14	*	31	21	19	*7
309.9	17	*9	*8	*	10	*	*	*	*7	*5	*
310	79	29	50	*	*6	*9	63	20	*9	38	13
310.1	*8	*	*	–	*	*	*5	*	*	*	*6
310.2	*8	*	*	*	*	*	*	*	*	*	*
310.9	62	20	42	*	*	*	55	16	*7	35	*
311	867	307	560	16	281	243	328	240	228	280	119
312	82	56	26	28	44	*6	*	38	20	17	*7
312.3	51	34	18	14	31	*6	*	28	13	*7	*
312.30	36	22	14	10	22	*	*	24	*7	*	*
312.34	13	11	*	*	*8	*	*	*	*	*	*
312.8	*8	*6	*	*	*	*	*	*	*	*	*
312.9	20	15	*6	9	*7	*	*	9	*	*	*
313	45	26	19	31	13	*	–	12	13	16	*
313.8	44	26	18	30	13	*	–	11	13	16	*
313.81	42	24	17	29	13	*	–	11	13	15	*
314	78	57	21	40	36	*	*	19	30	21	*8
314.0	75	55	20	39	34	*	*	18	29	20	*8
314.00	20	13	*7	*7	12	*	*	*	*8	*	*
314.01	55	42	13	32	23	*	*	13	22	16	*
315	29	18	12	21	*6	*	*	*	12	*	*9
315.8	*8	*	*	*8	*	*	*	*	*	*	*
315.9	10	*6	*	*5	*	*	*	*	*5	*	*
317	41	18	23	*	28	*7	*	19	*8	10	*
318	29	18	11	*	14	*8	*	*6	13	*6	*
318.0	9	*6	*	*	*	*	*	*	*5	*	*
318.1	13	*9	*	*	*7	*	*	*	*	*	*
318.2	*7	*	*	*	*	*	*	*	*	*	*
319	71	45	25	16	25	18	11	23	13	24	11
320–389	3,494	1,517	1,977	405	557	739	1,792	810	863	1,189	632
320	10	*	*5	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
322	13	*6	*7	*	*	*	*	*	*	*	*
322.9	10	*	*6	*	*	*	*	*	*	*	*
323	*7	*	*	*	*	*	*	*	*	*	*
324	10	*7	*	*	*	*	*	*	*	*	*
324.1	*6	*	*	*	*	*	*	*	*	*	*
330	*5	*	*	*	*	*	*	*	*	*	*
331	445	165	280	24	12	24	385	107	114	169	55
331.0	349	121	228	*	*	*6	343	88	91	136	34
331.3	15	*8	*6	*	*	*	10	*	*	*5	*
331.4	60	30	30	22	9	14	15	11	14	19	16
331.9	16	*	11	*	*	*	13	*	*	*6	*
332	270	138	132	—	*	19	250	66	65	105	35
332.0	267	137	130	—	*	18	248	65	63	104	34
333	83	37	46	*	14	19	48	16	25	25	17
333.0	*8	*5	*	—	*	*	*7	*	*	*	*
333.1	20	10	10	*	*	*	15	*	*7	*7	*
333.2	*8	*	*	*	*	*	*	*	*	*	*
333.8	*8	*	*	—	*	*	*5	*	*	*	*
333.82	*7	*	*	—	*	*	*	*	*	*	*
333.9	30	10	20	*	*5	*8	16	*	10	*8	*8
333.90	*7	*	*	*	*	*	*	*	*	*	*
333.99	21	*6	15	*	*	*6	12	*	*8	*	*6
334	*7	*	*	*	*	*	*	*	*	*	*
335	12	*6	*6	*	*	*	*6	*	*	*	*
335.2	9	*5	*	*	*	*	*6	*	*	*	*
335.20	*6	*	*	—	*	*	*	*	*	*	*
336	21	12	*9	*	*	*6	10	*7	*	*	*6
336.3	*7	*	*	—	*	*	*	*	*	*	*
336.9	*7	*	*	*	*	*	*	*	*	*	*
337	72	34	39	*	23	23	26	17	15	27	13
337.1	48	24	25	—	11	17	21	9	10	21	*9
337.2	16	*5	11	*	11	*	*	*	*	*	*
337.20	*5	*	*	—	*	*	*	*	*	*	*
337.22	*6	*	*	*	*	*	*	*	*	*	*
340	90	20	70	—	27	42	22	21	32	22	16
341	*6	*	*	*	*	*	*	*	*	*	*
342	192	81	111	*	15	44	128	35	49	62	46
342.0	11	*	*6	—	*	*	*8	*	*	*	*
342.00	*9	*	*	—	*	*	*6	*	*	*	*
342.8	*9	*	*6	*	*	*	*5	*	*	*	*
342.9	170	72	98	*	13	40	114	32	44	56	38
342.90	147	62	85	*	12	35	97	29	38	49	30
342.91	10	*	*	*	*	*	*7	*	*	*	*
342.92	14	*6	*8	*	*	*	10	*	*	*	*
343	72	43	29	37	24	*8	*	18	16	18	19
343.2	14	*8	*6	*9	*	*	*	*5	*	*	*
343.9	52	31	21	25	16	*8	*	11	12	13	15
344	131	69	62	*5	44	37	45	23	31	52	25
344.0	46	30	16	*	18	13	12	*8	12	14	11
344.00	38	25	13	*	14	11	11	*6	11	13	*8
344.1	41	24	17	*	16	16	*8	*8	*7	18	*8
344.3	13	*	11	—	*	*	10	*	*	12	—
344.30	13	*	11	—	*	*	10	*	*	12	—
344.6	*9	*	*	*	*	*	*	*	*	*	*
344.8	*5	*	*	*	*	*	*	*	*	*	*
344.9	16	*6	10	*	*	*	*8	*	*6	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
345	108	51	56	34	31	22	22	25	25	32	26
345.0	*5	*	*	*	*	*	*	*	*	*	*
345.1	13	*9	*	*	*	*	*	*	*	*	*
345.10	12	*8	*	*	*	*	*	*	*	*	*
345.3	21	12	*9	*7	*5	*5	*	*	*6	*6	*
345.4	*7	*	*	*	*	*	*	*	*	*	*
345.40	*5	*	*	*	*	*	*	*	*	*	*
345.5	11	*	*8	*	*	*	*	*	*	*	*
345.50	*9	*	*6	*	*	*	*	*	*	*	*
345.9	43	19	24	13	13	*9	*8	*8	*9	14	12
345.90	40	18	22	12	12	*9	*8	*7	*8	14	11
346	171	29	143	*	83	70	17	39	42	64	27
346.0	*6	*	*6	*	*	*	*	*	*	*	*
346.8	10	*	*8	*	*	*	*	*	*	*	*
346.80	*8	*	*6	*	*	*	*	*	*	*	*
346.9	147	24	123	*	71	60	14	33	35	56	23
346.90	138	23	115	*	65	58	14	31	32	52	22
346.91	*9	*	*8	*	*7	*	*	*	*	*	*
347	*7	*	*	*	*	*	*	*	*	*	*
348	153	79	75	16	26	40	71	37	34	51	32
348.0	*6	*	*	*	*	*	*	*	*	*	*
348.1	56	33	23	*	*6	17	29	15	10	18	12
348.3	39	18	22	*	*6	*7	23	*6	10	16	*8
348.4	12	*7	*5	*	*	*	*	*	*	*	*
348.5	16	*8	*7	*	*	*	*8	*	*	*	*
348.8	13	*7	*6	*	*	*5	*	*	*	*	*
348.9	*5	*	*	*	*	*	*	*	*	*	*
349	42	14	28	*	23	10	*8	11	*9	14	*8
349.0	22	*	17	*	19	*	*	*5	*6	*6	*
349.2	*6	*	*	*	*	*	*	*	*	*	*
349.8	10	*	*6	*	*	*	*	*	*	*	*
349.82	*6	*	*	—	*	*	*	*	*	*	*
350	17	*	12	*	*	*5	10	*	*	*6	*
350.1	16	*	12	*	*	*5	10	*	*	*5	*
351	30	13	17	*	*7	11	11	*8	*5	11	*6
351.0	27	10	17	*	*7	10	10	*7	*5	*9	*6
353	11	*	*7	*	*	*	*	*	*	*	*
353.0	*6	*	*	*	*	*	*	*	*	*	*
354	31	14	18	*	*8	13	9	*9	*8	11	*
354.0	17	*6	11	*	*5	*8	*	*	*	*7	*
354.2	*5	*	*	*	*	*	*	*	*	*	*
355	34	15	19	*	*	12	18	*7	*8	15	*
355.8	14	*8	*7	*	*	*	*8	*	*	*6	*
355.9	15	*5	10	—	*	*5	*8	*	*	*8	*
356	72	30	42	*	*8	16	47	13	17	29	13
356.1	*6	*	*	—	*	*	*	*	*	*	*
356.9	60	25	35	*	*6	14	40	9	14	24	12
357	232	118	114	*	26	96	109	63	57	75	37
357.0	11	*6	*6	*	*	*	*	*	*	*	*
357.2	209	106	103	—	20	88	101	57	51	68	32
358	27	10	17	*	*	*	20	*	10	*7	*5
358.0	24	9	15	*	*	*	18	*	9	*6	*5
359	27	14	13	*	10	*6	*8	10	*7	*	*
359.1	11	*8	*	*	*	*	*	*	*	*	*
360	*6	*	*	*	*	*	*	*	*	*	*
361	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
362	170	66	104	*7	17	55	91	47	40	49	34
362.0	100	44	56	—	14	48	38	31	21	29	19
362.01	94	41	53	—	13	45	36	29	20	28	17
362.02	*6	*	*	—	*	*	*	*	*	*	*
362.2	*7	*	*	*6	*	*	*	*	*	*	*
362.21	*6	*	*	*6	—	—	*	*	*	*	*
362.3	13	*6	*7	—	*	*	9	*	*	*	*
362.34	*7	*	*	—	*	*	*	*	*	*	*
362.5	43	10	33	—	*	*	40	10	12	*9	11
362.50	40	*8	32	—	*	*	38	10	12	*7	11
365	161	59	102	*	*	23	135	52	44	45	20
365.9	150	54	96	*	*	21	126	48	41	42	18
366	28	11	17	*	*	*	22	*9	*8	*7	*
366.9	23	9	14	*	*	*	19	*8	*7	*5	*
368	60	24	36	*	15	15	28	9	17	18	16
368.1	10	*	*6	*	*	*	*5	*	*	*	*
368.16	*5	*	*	—	*	*	*	*	*	*	*
368.2	13	*6	*7	*	*	*	*	*	*	*	*
368.4	20	*8	12	—	*	*	12	*	*5	*	*6
368.46	17	*7	10	—	*	*	11	*	*	*	*
368.8	11	*	*7	*	*	*	*	*	*	*	*
369	72	29	43	*	10	15	45	13	19	28	11
369.0	25	*8	17	*	*	*	17	*	*6	12	*
369.00	25	*8	17	*	*	*	17	*	*6	11	*
369.4	19	*7	12	*	*	*	12	*6	*6	*5	*
369.6	16	11	*6	*	*	*	*8	*	*	*6	*
369.60	15	10	*	*	*	*	*7	*	*	*6	*
369.9	*6	*	*	*	*	*	*	*	*	*	*
372	48	28	20	17	*6	*5	20	*8	12	17	10
372.0	*8	*	*	*	*	*	*	*	*	*	*
372.00	*8	*	*	*	*	*	*	*	*	*	*
372.3	29	17	12	12	*	*	12	*	*8	9	*7
372.30	26	15	11	9	*	*	12	*	*8	*9	*5
372.7	*9	*5	*	*	*	*	*	*	*	*	*
372.72	*7	*	*	*	*	*	*	*	*	*	*
373	*6	*	*	*	*	*	*	*	*	*	*
374	10	*	*6	*	*	*	*	*	*	*6	*
374.3	*7	*	*6	*	*	*	*	*	*	*	*
376	15	9	*6	*7	*	*	*	*	*	*5	*
376.0	10	*6	*	*7	*	*	*	*	*	*	*
376.01	10	*6	*	*6	*	*	*	*	*	*	*
377	10	*	*6	*	*	*	*	*	*	*	*
378	16	*8	*8	*	*	*	*	*	*	*7	*
378.5	*9	*	*	*	*	*	*	*	*	*	*
379	22	10	11	*	*5	*5	*7	*	*5	*8	*
379.2	*6	*	*	*	*	*	*	*	*	*	*
379.5	*5	*	*	*	*	*	*	*	*	*	*
379.9	*5	*	*	*	*	*	*	*	*	*	*
380	31	10	21	*5	11	*6	*9	*6	*7	12	*5
380.1	19	*5	13	*	*9	*	*	*	*	*7	*
380.10	18	*	13	*	*8	*	*	*	*	*6	*
380.4	*8	*	*6	*	*	*	*	*	*	*	*
381	18	10	*8	13	*	*	*	*6	*	*	*
382	192	115	77	166	16	*	*7	29	37	76	49
382.9	186	112	73	161	15	*	*6	28	35	75	47
383	11	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
386	64	19	45	*	*	21	37	18	18	20	*8
386.0	12	*	*9	—	*	*	*7	*	*	*	*
386.00	12	*	*9	—	*	*	*7	*	*	*	*
386.1	20	*5	15	—	*	*6	13	*8	*7	*	*
386.11	13	*	10	—	*	*	*8	*	*	*	*
386.3	29	10	19	*	*	10	15	*6	*7	11	*
386.30	26	*9	17	*	*	*8	14	*6	*7	11	*
388	15	*6	*9	*	*	*	*7	*	*	*5	*
389	86	44	42	*5	13	13	55	22	25	26	13
389.1	11	*7	*	*	*	*	*8	*	*8	*	*
389.10	*9	*6	*	*	*	*	*7	*	*7	*	*
389.7	*6	*5	*	*	*	*	*	*	*	*	*
389.9	61	29	32	*	*8	11	39	18	14	20	*9
390-459	29,894	13,960	15,933	115	1,558	7,321	20,900	7,372	7,266	10,891	4,365
394	27	*8	19	*	*	*8	17	*7	9	*6	*
394.0	*7	*	*6	*	*	*	*5	*	*	*	*
394.1	*9	*	*6	—	*	*	*6	*	*	*	*
394.2	*8	*	*6	—	*	*	*	*	*	*	*
395	*5	*	*	—	*	*	*	*	*	*	*
396	144	57	87	*	*	14	125	44	31	51	18
396.0	*6	*	*	*	*	*	*	*	*	*	*
396.2	41	17	25	*	*	*	40	15	10	11	*7
396.3	72	30	42	*	*	*8	61	21	15	28	*8
396.8	15	*6	*9	—	*	*	13	*	*	*	*
396.9	*7	*	*5	—	*	*	*5	*	*	*	*
397	116	42	74	*	*6	20	89	36	25	44	11
397.0	113	41	73	*	*6	19	88	35	24	43	11
398	69	26	43	—	*	11	56	32	13	17	*7
398.9	69	26	43	—	*	11	56	32	13	17	*7
398.90	11	*5	*5	—	*	*5	*	*	*	*	*
398.91	58	21	37	—	*	*6	51	29	10	13	*6
401	6,943	2,806	4,137	11	467	2,049	4,416	1,647	1,681	2,630	985
401.0	60	20	40	*	*9	20	32	14	11	30	*5
401.1	151	65	86	—	9	51	90	62	20	22	47
401.9	6,732	2,720	4,012	11	449	1,978	4,294	1,571	1,650	2,577	933
402	360	137	223	*	11	72	277	102	71	134	53
402.0	*7	*	*	—	*	*	*	*	*	*	*
402.9	351	132	219	*	*9	70	272	101	70	128	52
402.90	192	72	120	*	*5	45	143	55	43	72	23
402.91	159	59	99	—	*	26	129	46	27	56	29
403	661	329	332	*	74	203	381	135	134	268	124
403.0	11	*	*6	—	*	*	*	*	*	*6	*
403.01	10	*	*5	—	*	*	*	*	*	*5	*
403.1	14	*8	*6	*	*	*	*7	*6	*	*	*
403.11	13	*7	*6	*	*	*	*6	*5	*	*	*
403.9	636	316	319	*	67	197	370	128	131	259	117
403.90	58	29	29	*	*	15	38	16	*8	22	12
403.91	578	287	291	*	63	181	332	112	123	237	106
404	70	30	40	—	*	18	49	17	13	27	13
404.9	65	29	37	—	*	16	47	16	11	26	11
404.90	*6	*	*	—	*	*	*	*	*	*	*
404.91	*8	*	*	—	*	*	*7	*	*	*	*
404.92	12	*5	*6	—	*	*	*9	*	*	*	*
404.93	39	16	23	—	*	9	28	*8	*7	17	*7
405	9	*	*	*	*	*	*	*	*	*	*
405.9	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
410	941	533	408	*	45	280	616	234	223	333	151
410.0	32	19	12	—	*	10	21	*7	*7	13	*
410.01	30	18	12	—	*	9	20	*6	*7	13	*
410.1	113	67	46	*	*8	35	70	25	26	41	21
410.11	101	60	42	*	*7	31	64	20	24	39	18
410.12	*8	*	*	—	*	*	*	*	*	*	*
410.2	22	11	11	—	*	*8	13	*7	*	*7	*
410.21	20	10	10	—	*	*8	12	*6	*	*6	*
410.3	15	10	*5	—	*	*7	*7	*	*	*	*
410.31	14	9	*	—	*	*6	*6	*	*	*	*
410.4	133	84	50	—	*9	57	67	33	32	44	24
410.41	118	77	42	—	*8	51	60	31	30	39	19
410.42	12	*7	*5	—	*	*	*6	*	*	*	*5
410.5	14	10	*	*	*	*	*8	*	*	*5	*
410.51	13	9	*	*	*	*	*8	*	*	*	*
410.6	*6	*	*	—	*	*	*	*	*	*	*
410.7	421	229	193	*	17	114	290	113	98	145	65
410.70	*6	*	*	—	*	*	*5	*	*	*	*
410.71	390	213	177	*	16	107	267	105	92	132	60
410.72	26	12	13	—	*	*7	18	*6	*	9	*5
410.8	15	*7	*8	*	*	*5	10	*	*	*6	*
410.81	13	*6	*7	*	*	*	9	*	*	*	*
410.9	169	91	78	*	*6	37	126	39	42	65	24
410.90	12	*6	*6	—	*	*	11	*5	*	*	—
410.91	146	78	69	*	*5	31	109	31	35	58	21
410.92	11	*8	*	—	*	*	*6	*	*	*	*
411	851	486	365	*	44	322	484	238	178	311	124
411.0	*9	*5	*	—	*	*	*5	*	*	*	*
411.1	808	465	343	*	41	310	456	225	170	298	116
411.8	34	16	18	*	*	9	23	12	*5	*9	*7
411.81	*7	*	*	—	*	*	*	*	*	*	*
411.89	27	12	15	*	*	*7	19	11	*	*8	*
412	661	387	274	*	29	205	426	179	163	212	107
413	543	255	288	—	19	155	368	132	141	205	64
413.9	538	252	286	—	17	153	368	132	140	203	64
414	4,465	2,484	1,980	*5	116	1,194	3,149	1,153	1,132	1,588	592
414.0	4,144	2,296	1,848	*5	112	1,119	2,908	1,061	1,045	1,477	561
414.00	1,265	689	576	*	22	264	978	307	430	412	116
414.01	2,730	1,504	1,226	*	85	811	1,830	716	579	1,007	427
414.02	95	68	27	—	*	32	61	20	29	31	15
414.04	12	*8	*	—	*	*	*8	*	*	*6	*
414.05	42	27	15	*	*	*9	31	17	*	20	*
414.1	*7	*	*	*	*	*	*	*	*	*	*
414.10	*6	*	*	—	*	*	*	*	*	*	*
414.8	214	135	80	—	*	47	165	57	64	71	22
414.9	99	50	49	—	*	26	72	33	21	38	*7
415	133	51	82	*	18	40	74	37	28	47	20
415.0	*5	*	*	*	*	*	*	*	*	*	*
415.1	127	48	79	*	17	39	72	35	28	45	20
415.11	12	*	*7	—	*	*	*5	*	*	*	*
415.19	116	43	73	*	14	35	66	31	24	42	18
416	286	113	173	*	17	62	203	67	68	99	53
416.0	163	59	104	*	10	33	116	40	40	61	21
416.8	43	15	28	*	*	*9	31	11	*9	*9	16
416.9	80	40	40	*	*	20	55	16	19	29	16
420	24	14	10	*	*6	*8	*8	*6	*6	*7	*5
420.9	22	13	9	*	*6	*7	*8	*6	*6	*6	*
420.90	17	9	*8	*	*5	*	*6	*5	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
421	25	16	*9	*	*5	*9	11	*8	*5	*8	*
421.0	23	15	*9	*	*	*9	10	*8	*	*8	*
423	37	16	20	*	*6	10	20	9	*7	14	*6
423.9	28	12	16	*	*6	*7	15	*8	*5	11	*
424	959	362	597	*6	88	178	688	248	217	350	144
424.0	547	184	363	*	74	118	351	136	124	213	74
424.1	335	147	188	*	9	41	283	92	81	103	60
424.2	41	17	24	*	*	11	27	*7	*6	21	*7
424.3	*6	*	*	*	*	*	*	*	*	*	*
424.9	30	11	19	*	*	*5	23	12	*	10	*
424.90	27	9	17	*	*	*5	20	9	*	9	*
425	534	273	261	*	44	145	342	130	122	200	83
425.1	11	*	*7	*	*	*	*8	*	*	*	*
425.4	478	245	232	*	39	128	309	117	109	178	74
425.5	14	10	*	—	*	*8	*	*	*	*5	*
425.8	20	*9	11	*	*	*	13	*	*	*8	*
425.9	*7	*	*	*	*	*	*	*	*	*	*
426	385	197	188	*	19	60	304	96	108	112	69
426.0	65	35	30	*	*	13	48	17	16	22	9
426.1	97	50	47	*	*	13	79	29	24	27	17
426.10	*8	*	*	—	*	*	*6	*	*	*	*
426.11	55	26	29	*	*	*6	45	15	15	14	10
426.12	*9	*	*	—	*	*	*7	*	*	*	*
426.13	26	16	9	*	*	*	20	*9	*	*8	*5
426.2	10	*6	*	—	*	*	*8	*	*	*	*
426.3	81	33	48	—	*	11	68	22	21	21	17
426.4	63	32	30	*	*	10	49	15	20	19	9
426.5	34	22	12	—	*	*	29	*	12	9	*8
426.52	*6	*	*	—	*	*	*	*	*	*	*
426.53	21	13	*7	—	*	*	18	*	*9	*	*6
426.6	10	*5	*	—	*	*	*8	*	*	*	*
426.7	11	*	*7	*	*	*	*	*	*	*	*
426.9	10	*7	*	*	*	*	*7	*	*	*	*
427	3,748	1,781	1,967	27	153	615	2,954	964	924	1,249	611
427.0	95	37	57	*	*8	19	65	20	22	31	21
427.1	233	157	76	*	12	66	155	68	51	73	42
427.3	2,400	1,107	1,293	*	40	325	2,032	636	599	782	384
427.31	2,227	1,010	1,217	*	35	282	1,907	600	554	721	352
427.32	173	97	76	*	*5	43	125	36	45	61	32
427.4	48	29	19	*	*	17	27	11	10	16	11
427.41	47	29	19	*	*	17	27	11	10	16	10
427.5	97	46	51	*	*8	20	67	17	21	39	20
427.6	119	62	57	*	10	28	80	30	30	40	20
427.60	*7	*	*	*	*	*	*	*	*	*	*
427.61	28	11	17	*	*	*	20	*7	*8	*8	*5
427.69	85	47	38	*	*6	22	56	21	21	29	14
427.8	627	283	345	15	57	114	441	140	167	219	101
427.81	163	74	89	*	*	18	142	45	40	59	18
427.89	465	209	256	14	55	96	299	95	127	160	83
427.9	124	57	67	*	14	24	85	41	24	47	12
428	3,255	1,347	1,907	11	86	567	2,591	761	849	1,218	426
428.0	3,226	1,334	1,893	11	85	561	2,569	747	843	1,211	425
428.1	*8	*	*	*	*	*	*5	*	*	*	*
428.9	21	10	11	*	*	*	17	13	*	*6	*
429	274	133	142	*	16	57	198	70	62	110	33
429.2	114	55	59	—	*	12	99	38	19	47	9
429.3	97	43	54	*	9	24	61	19	18	43	16
429.4	9	*	*6	*	*	*5	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
429.9	39	22	17	*	*	11	27	*8	16	12	*
430	27	10	17	—	*6	12	*9	*6	*5	12	*
431	84	38	45	*	*5	26	52	19	18	31	16
432	37	19	18	*	*	*7	26	9	*6	10	11
432.1	32	17	15	*	*	*6	23	*9	*	9	9
433	300	150	150	*	*	63	234	68	76	120	37
433.1	218	111	107	—	*	43	174	49	56	87	26
433.10	192	96	96	—	*	38	152	41	52	78	22
433.11	26	15	11	—	*	*	21	*8	*	*9	*
433.3	69	35	34	—	*	16	52	17	16	28	*8
433.30	60	30	30	—	*	14	45	15	14	24	*7
433.31	*8	*	*	—	*	*	*7	*	*	*	*
434	354	156	198	*	12	83	257	75	80	144	54
434.0	13	*6	*7	*	*	*	9	*	*	*	*
434.01	11	*	*6	*	*	*	*8	*	*	*	*
434.1	50	21	29	*	*	12	36	14	11	15	11
434.10	*9	*	*6	—	—	*	*8	*	*	*	*
434.11	41	18	23	*	*	11	27	12	*8	12	10
434.9	291	128	162	*	*9	68	212	58	65	126	41
434.90	11	*5	*5	*	*	*	*7	*	*	*5	*
434.91	280	123	157	*	*8	65	205	57	63	121	39
435	282	114	167	*	11	53	217	83	60	104	34
435.3	12	*	*7	—	*	*	9	*	*	*	*
435.8	*7	*	*	—	*	*	*5	*	*	*	*
435.9	255	103	153	*	10	47	197	77	54	95	29
436	197	78	119	*	*	40	155	28	55	83	32
437	150	56	93	*	*	22	122	24	33	69	23
437.0	63	26	38	—	*	*	60	11	11	30	11
437.1	13	*	*8	*	*	*	11	*	*	*7	*
437.2	11	*	*8	*	*	*	*7	*	*	*5	*
437.3	15	*5	10	—	*	*6	*7	*	*	*7	*
437.9	38	15	23	—	*	*5	33	*7	11	16	*
438	553	259	295	*	14	100	438	119	129	213	92
438.0	15	*7	*8	*	*	*	12	*	*5	*	*
438.1	94	43	52	*	*	18	74	22	25	31	16
438.10	10	*	*5	—	*	*	*9	*	*	*	*
438.11	64	27	37	*	*	12	51	16	17	20	11
438.12	10	*	*6	—	*	*	*8	*	*	*	*
438.19	10	*7	*	—	*	*	*7	*	*	*	*
438.2	250	112	138	*	*7	45	197	63	56	90	42
438.20	224	100	123	*	*6	40	177	57	49	84	34
438.21	13	*6	*8	*	*	*	10	*	*	*	*
438.22	13	*6	*7	*	*	*	10	*	*	*	*
438.5	*6	*	*	*	*	*	*	*	*	*	*
438.8	148	74	74	*	*	24	121	23	33	65	27
438.82	36	22	14	—	*	*	32	*6	*6	17	*8
438.89	109	51	58	*	*	19	88	17	26	47	18
438.9	39	20	19	*	*	*8	29	*8	*7	21	*
440	364	196	168	*	*8	80	274	89	98	126	51
440.0	18	*8	10	*	*	*	13	*5	*	*	*
440.1	35	15	20	*	*	*8	26	*8	12	11	*
440.2	224	128	97	*	*6	56	162	51	58	79	35
440.20	56	31	25	*	*	14	40	12	15	23	*6
440.21	43	28	15	—	*	14	28	10	11	14	*8
440.22	16	*9	*7	—	*	*6	10	*	*	*6	*
440.23	36	19	17	—	*	*5	31	*9	9	9	*9
440.24	68	39	29	—	*	16	49	16	17	25	10
440.29	*5	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
440.3	10	*	*	—	—	*	*7	*	*	*	*
440.31	*5	*	*	—	—	*	*	*	*	*	*
440.8	*7	*	*	—	—	*	*	*	*	*	*
440.9	70	35	35	—	*	*7	62	22	17	26	*
441	168	107	61	*	*	23	142	45	40	58	25
441.0	14	*8	*6	*	*	*6	*6	*	*	*	*
441.01	*7	*	*	*	*	*	*	*	*	*	*
441.2	13	*6	*6	*	*	*	9	*	*	*5	*
441.3	*7	*5	*	—	—	*	*6	*	*	*	*
441.4	123	82	42	—	*	12	111	33	32	42	17
441.7	*6	*	*	—	*	*	*5	*	*	*	*
442	27	18	9	*	*	*8	18	*	*6	11	*7
442.2	10	*8	*	—	*	*	*8	*	*	*	*
442.3	11	*7	*	—	*	*	*8	*	*	*	*
443	475	240	235	*	18	114	342	127	108	183	56
443.0	12	*	9	*	*	*5	*	*	*	*	*
443.8	61	28	33	*	*	20	37	17	11	22	11
443.81	50	24	27	—	*	18	30	15	*7	19	9
443.89	11	*	*6	*	*	*	*7	*	*	*	*
443.9	397	205	192	*	*8	88	301	106	94	155	41
444	83	46	37	*	*	25	53	20	15	33	16
444.0	9	*	*	*	*	*	*5	*	*	*	*
444.2	56	32	24	*	*	17	35	12	10	23	10
444.21	*7	*	*	—	*	*	*	*	*	*	*
444.22	49	29	20	*	*	15	32	11	9	20	*8
444.8	13	*6	*6	—	*	*	*8	*	*	*	*
444.81	9	*	*	—	*	*	*6	*	*	*	*
446	37	12	25	*6	*	*7	20	10	*7	11	9
446.1	*6	*	*	*6	*	—	—	*	*	*	*
446.4	*7	*	*	—	*	*	*	*	*	*	*
446.5	14	*	13	—	—	*	13	*	*	*	*
447	55	25	30	*	*6	18	29	18	13	19	*6
447.1	28	15	13	*	*	10	17	10	*	12	*
447.6	13	*	*8	*	*	*5	*	*	*	*	*
448	*7	*	*	*	*	*	*5	*	*	*	*
451	63	28	36	*	13	18	31	14	15	28	*5
451.0	10	*	*7	—	*	*	*	*	*	*5	*
451.1	19	*9	11	—	*	*6	11	*6	*	9	*
451.11	*6	*	*	—	*	*	*	*	*	*	*
451.19	13	*6	*8	—	*	*5	*7	*	*	*7	*
451.2	12	*6	*6	*	*	*	*8	*	*	*	*
451.8	19	10	9	*	*6	*5	*8	*	*	11	*
451.82	*6	*	*	*	*	*	*	*	*	*	*
451.84	9	*6	*	*	*	*	*	*	*	*6	—
453	296	132	163	*	42	86	167	70	77	108	41
453.2	*5	*	*	*	*	*	*	*	*	*	*
453.8	245	109	137	*	34	73	137	56	69	86	35
453.9	43	21	22	—	*5	10	28	12	*7	19	*5
454	30	14	16	—	*	12	15	9	*6	11	*
454.0	*6	*	*	—	*	*	*	*	*	*	*
454.2	*7	*	*	—	*	*	*	*	*	*	—
454.9	15	*6	*9	—	*	*	*7	*6	*	*	*
455	161	68	93	*	37	49	74	38	37	61	24
455.0	51	20	31	*	9	15	27	12	10	19	11
455.2	30	15	15	—	*	10	15	*6	*6	12	*6
455.3	18	*	14	—	*	*5	*9	*	*6	*6	*
455.5	12	*6	*5	—	*	*	*	*	*	*	*
455.6	28	10	18	—	11	*7	10	*7	*8	11	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
455.8	12	*6	*6	—	*	*	*5	*	*	*	*
456	60	38	22	*	13	31	15	17	12	16	15
456.1	*8	*	*	*	*	*	*	*	*	*	*
456.2	37	27	10	*	*7	22	*9	12	*8	9	*9
456.20	21	15	*7	*	*	13	*	*7	*	*	*6
456.21	16	12	*	*	*	*9	*	*5	*	*5	*
456.8	*6	*	*	—	*	*	*	*	*	*	*
457	28	12	17	*	*6	11	10	*8	*7	9	*
457.1	19	*6	13	*	*	*8	*7	*6	*	*5	*
458	385	183	202	*5	38	92	250	73	104	141	68
458.0	106	49	57	*	10	23	72	17	27	42	21
458.1	*6	*	*	—	—	*	*	*	*	*	*
458.2	37	16	20	*	*	*9	24	10	10	*8	*9
458.8	23	12	11	*	*	*	15	*	*6	*7	*
458.9	214	102	112	*	23	54	133	40	57	83	33
459	134	61	72	*	12	36	85	33	39	42	20
459.0	18	*9	*9	*	*	*	11	*	*	*6	*
459.2	11	*	*7	*	*	*	*6	*	*	*5	*
459.8	87	39	49	*	*7	21	59	21	30	22	14
459.81	79	36	42	—	*6	20	52	19	25	21	13
459.89	*9	*	*6	*	*	*	*6	*	*	*	*
459.9	14	*8	*6	—	*	*	*8	*	*	*6	*
460-519	10,262	4,766	5,496	1,198	1,266	2,199	5,598	2,379	2,435	3,720	1,728
460	*6	*	*	*	*	*	*	*	*	*	*
461	58	24	34	10	24	10	14	10	16	24	*8
461.0	15	*5	10	*	9	*	*	*	*6	*6	*
461.8	*6	*	*	*	*	*	*	*	*	*	*
461.9	29	13	17	*5	11	*7	*6	*	*7	13	*6
462	52	21	31	15	24	*7	*6	10	*9	24	*9
463	18	11	*8	*	12	*	*	*	*6	*7	*
464	47	28	19	36	*6	*	*	11	9	14	13
464.1	*8	*	*	*6	*	*	*	*	*	*	*
464.10	*8	*	*	*6	*	*	*	*	*	*	*
464.4	30	21	9	29	*	—	*	*	*7	9	9
465	156	71	85	64	35	20	36	33	35	55	33
465.9	155	70	84	63	35	20	36	32	34	55	33
466	424	207	217	239	40	41	104	72	97	151	105
466.0	201	82	119	21	39	40	100	44	50	82	26
466.1	223	125	98	218	*	*	*	28	47	69	79
466.11	117	66	51	116	*	*	*	15	26	29	48
466.19	105	59	47	101	*	*	*	13	21	40	31
470	12	10	*	*	*6	*	*	*	*	*	*
472	11	*	*6	*	*	*	*	*	*	*	*
472.0	10	*	*6	*	*	*	*	*	*	*	*
473	148	57	91	18	43	47	40	28	30	72	18
473.0	20	*7	13	*	*	*5	*7	*	*	9	*
473.2	10	*	*6	*	*	*	*	*	*	*	*
473.8	11	*6	*	*	*	*	*	*	*	*5	*
473.9	103	38	65	10	32	34	27	19	21	52	12
474	27	15	12	18	*7	*	*	*8	*	11	*
474.0	11	*5	*6	*7	*	*	*	*	*	*5	*
474.00	*6	*	*	*	*	*	*	*	*	*	*
474.02	*5	*	*	*	*	—	—	*	*	*	*
474.1	15	9	*6	11	*	*	*	*6	*	*5	*
474.10	*8	*5	*	*8	*	*	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
474.11	*5	*	*	*	*	*	*	*	*	*	*
475	16	*8	*8	*	9	*	*	*5	*	*5	*
477	45	17	28	*	*7	19	17	*5	11	21	*8
477.9	44	17	27	*	*6	18	17	*5	11	20	*7
478	77	37	40	16	21	21	19	20	18	24	15
478.1	12	*6	*6	*	*	*	*	*	*	*	*
478.2	22	13	9	*	*6	*7	*	*5	*5	*7	*
478.29	14	*9	*6	*	*	*6	*	*	*	*	*
478.3	14	*	*9	*	*	*	*8	*	*	*6	*
478.6	*5	*	*	*	*	*	*	*	*	*	*
478.7	13	*6	*7	*7	*	*	*	*	*	*	*
478.75	*6	*	*	*	*	*	*	*	*	*	*
480	39	20	19	26	*	*	*8	*	*8	14	12
480.1	16	10	*6	14	*	—	*	*	*	*7	*
480.9	21	*8	12	11	*	*	*5	*	*	*6	*7
481	60	28	32	*	*8	17	32	11	16	21	13
482	240	125	115	14	27	39	161	47	64	91	38
482.0	16	*8	*9	*	*	*	11	*	*	*7	*
482.1	43	21	22	*	*8	*8	24	10	12	13	*8
482.2	26	15	12	*	*	*	20	*	10	9	*
482.3	15	*7	*8	*	*	*	10	*	*	*	*
482.30	*8	*	*	*	*	*	*	*	*	*	*
482.4	69	37	33	*	*8	*7	52	13	16	32	9
482.40	*6	*	*	*	*	*	*	*	*	*	*
482.41	59	33	26	*	*6	*6	45	12	12	27	*9
482.8	43	20	23	*	*	*7	29	*	16	16	*5
482.82	*5	*	*	*	*	*	*	*	*	*	*
482.83	31	13	18	*	*	*	24	*	14	11	*
482.9	28	17	11	*	*	*7	15	11	*	*9	*
483	10	*5	*	*	*	*	*	*	*	*	*
483.0	*9	*	*	*	*	*	*	*	*	*	*
484	*7	*5	*	*	*	*	*	*	*	*	*
485	55	21	34	27	*	*6	18	*6	12	32	*5
486	1,645	752	893	184	168	287	1,005	359	411	607	268
487	69	27	42	12	*7	12	39	12	25	24	*8
487.0	18	*8	10	*	*	*	12	*	*6	*7	*
487.1	49	19	30	10	*5	*8	26	9	20	16	*
490	92	38	55	12	18	18	43	20	23	35	14
491	1,024	448	576	*	45	270	707	250	225	411	139
491.0	*8	*	*	—	*5	*	*	*	*	*	*
491.2	987	432	555	*	33	258	694	244	212	398	132
491.20	110	54	56	*	*	24	80	32	24	40	15
491.21	877	378	499	*	28	234	613	213	188	359	117
491.9	27	12	15	*	*6	10	11	*	*9	10	*
492	200	112	89	*	9	50	141	51	52	72	25
492.0	14	11	*	*	*	*5	*	*	*	*	*
492.8	186	101	85	*	*	44	136	46	50	68	22
493	1,391	490	901	314	402	334	342	387	325	434	246
493.0	74	41	34	56	11	*5	*	11	*7	10	47
493.00	47	23	24	33	*8	*	*	*5	*	*8	29
493.01	27	17	10	23	*	*	*	*6	*	*	18
493.2	226	77	149	*	21	74	130	73	53	68	31
493.20	206	70	136	*	19	65	121	67	50	61	28
493.21	20	*7	12	*	*	9	*9	*7	*	*7	*
493.9	1,089	372	717	257	369	254	210	302	265	355	167
493.90	939	302	637	170	334	234	201	265	221	304	149

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
493.91	151	70	81	87	35	20	9	37	44	50	19
494	38	15	24	*	*	*6	28	13	*5	13	*7
496	1,851	975	876	*	38	417	1,392	443	485	656	267
500	10	10	*	*	*	*	9	*8	*	*	*
501	20	18	*	—	*	*	17	*6	*	*9	*
507	294	170	124	*9	26	38	221	69	64	110	51
507.0	291	168	124	*8	26	37	220	68	64	109	50
508	*6	*	*	*	*	*	*	*	*	*	*
510	22	14	*9	*	*6	*8	*7	*	*	*8	*7
510.9	20	12	*8	*	*5	*7	*6	*	*	*7	*7
511	375	164	211	13	39	89	233	90	85	137	62
511.0	30	14	16	*	*8	*8	13	*8	*6	10	*5
511.8	29	12	18	*	*	*7	16	*8	*6	*9	*7
511.9	313	137	176	9	28	73	203	74	71	118	50
512	114	65	49	*	23	29	58	24	21	50	18
512.0	*5	*	*	*	*	*	*	*	*	*	*
512.1	56	29	27	*	*5	16	33	12	11	23	9
512.8	54	33	21	*	17	12	22	10	9	26	*9
513	11	*8	*	*	*	*	*5	*	*	*	*
513.0	10	*7	*	*	*	*	*	*	*	*	*
514	34	15	19	*	*	*6	23	*7	11	12	*
515	151	64	87	*	*8	21	119	31	46	47	26
516	33	16	17	*	*	*8	21	*8	*8	11	*5
516.3	14	*7	*8	*	*	*	11	*	*5	*	*
516.8	12	*7	*5	*	*	*	*6	*	*	*	*
516.9	*6	*	*	*	*	*	*	*	*	*	*
517	20	*	16	*	*	10	*	*	*	*7	*
517.8	18	*	14	*	*	9	*	*	*	*7	*
518	1,251	594	657	111	154	308	677	283	250	458	260
518.0	295	141	154	22	49	78	146	56	63	115	61
518.3	17	*8	*9	*	*	*	*8	*	*	*7	*
518.4	23	*8	14	*	*6	*6	9	*	*6	*9	*
518.5	98	60	38	*	17	29	47	24	22	33	19
518.8	814	375	439	81	78	189	466	194	155	294	171
518.81	548	249	299	15	53	129	350	145	102	213	88
518.82	122	59	63	57	12	18	34	12	19	35	55
518.83	28	12	16	*	*	*5	20	*8	*8	9	*
518.84	31	13	17	*	*	11	18	9	*6	*9	*7
518.89	86	41	44	*8	10	25	44	20	20	28	17
519	89	42	47	19	17	24	29	21	27	26	14
519.0	12	*8	*	*	*	*	*	*	*	*	*
519.09	*6	*	*	*	*	*	*	*	*	*	*
519.1	36	14	22	10	*6	9	12	*7	14	11	*
519.4	*7	*	*	*	*	*	*	*	*	*	*
519.8	22	12	10	*	*	*6	*7	*	*8	*5	*
519.9	9	*	*6	*	*	*	*	*	*	*	*
520-579	9,131	3,901	5,231	452	1,795	2,527	4,357	2,093	2,177	3,318	1,543
520	*5	*	*	*	*	*	*	*	*	*	*
521	22	11	12	*	10	*	*	*6	*	10	*
521.0	21	10	11	*	10	*	*	*6	*	10	*
522	21	10	11	*	*8	*8	*	*	*6	11	*
522.5	18	*8	9	*	*7	*6	*	*	*	*9	*
523	12	*7	*	*	*	*	*	*	*	*6	*
524	30	10	20	*	21	*	*	*6	*	10	9

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
524.0	10	*	*8	*	*8	*	*	*	*	*	*
524.6	11	*	*7	*	*5	*	*	*	*	*	*
524.60	*8	*	*	*	*	*	*	*	*	*	*
525	*6	*	*	*	*	*	*	*	*	*	*
526	*9	*	*	*	*	*	*	*	*	*	*
527	16	*7	*9	*	*	*	*7	*	*	*6	*
527.2	10	*	*6	*	*	*	*	*	*	*	*
528	41	21	20	*8	12	9	12	10	*8	15	*8
528.0	22	11	11	*5	*	*6	*7	*7	*	*7	*
528.3	*5	*	*	*	*	*	*	*	*	*	*
528.9	10	*6	*	*	*	*	*	*	*	*	*
530	1,477	620	857	88	206	450	733	358	364	524	231
530.0	*7	*5	*	*	*	*	*	*	*	*	*
530.1	210	102	108	*5	37	65	103	50	57	71	32
530.10	91	44	47	*	16	27	46	25	24	34	*9
530.11	78	36	41	*	12	28	35	17	25	23	14
530.19	42	22	20	*	*8	10	22	*8	9	15	10
530.2	70	42	27	*	*9	18	41	17	17	21	14
530.3	43	21	22	*	*	11	30	*7	*9	16	12
530.5	19	*6	12	*	*	*6	11	*	*5	*8	*
530.6	9	*	*5	*	*	*	*8	*	*	*6	*
530.7	20	12	*8	*	*	*	10	*	*7	*7	*
530.8	1,095	424	671	79	149	340	526	277	262	393	163
530.81	1,062	408	654	79	146	331	506	268	253	385	157
530.82	15	*8	*8	—	*	*	*9	*	*	*	*
530.89	16	*8	*9	*	*	*	11	*6	*5	*	*
531	157	69	87	*	15	36	106	29	44	51	33
531.0	23	11	11	*	*	*	17	*	*8	*6	*
531.00	21	11	11	*	*	*	16	*	*7	*6	*
531.3	*7	*	*	*	*	*	*6	*	*	*	*
531.30	*7	*	*	*	*	*	*6	*	*	*	*
531.4	62	30	33	*	*5	14	44	13	15	20	15
531.40	62	30	32	*	*5	14	43	13	14	20	15
531.7	*6	*	*	—	*	*	*	*	*	*	*
531.70	*5	*	*	—	*	*	*	*	*	*	*
531.9	53	21	32	*	*7	14	32	*9	16	20	*8
531.90	52	20	32	*	*6	14	31	*9	15	19	*8
532	112	65	47	*	18	29	65	24	24	40	24
532.0	17	9	*8	*	*	*	11	*	*	*7	*
532.00	17	*9	*8	*	*	*	11	*	*	*7	*
532.4	51	34	17	*	*8	12	31	12	9	16	13
532.40	49	32	16	*	*8	11	30	12	9	16	12
532.5	*5	*	*	—	*	*	*	*	*	*	*
532.50	*5	*	*	—	*	*	*	*	*	*	*
532.9	28	15	13	—	*6	*7	15	*5	*7	10	*6
532.90	26	14	12	—	*6	*7	13	*5	*6	10	*5
533	157	65	92	*	21	36	100	52	31	53	22
533.4	12	*6	*7	—	*	*	*9	*	*	*	*
533.40	12	*6	*7	—	*	*	*9	*	*	*	*
533.7	*8	*	*6	—	*	*	*	*	*	*	*
533.70	*8	*	*6	—	*	*	*	*	*	*	*
533.9	130	54	76	*	18	30	82	45	24	42	19
533.90	130	54	76	*	18	30	82	45	24	42	19
534	*5	*	*	—	*	*	*	*	*	*	*
535	607	260	347	12	110	164	321	147	151	226	83
535.0	67	24	43	*	13	17	34	17	15	28	*7
535.00	50	15	35	*	12	12	23	11	10	24	*5
535.01	16	9	*7	*	*	*	11	*6	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
535.1	82	34	48	*	11	21	49	23	18	31	11
535.10	68	27	41	*	10	19	38	20	16	24	*8
535.11	14	*7	*7	*	*	*	11	*	*	*7	*
535.3	21	17	*	*	11	*8	*	*	*6	*7	*
535.30	16	13	*	*	*9	*6	*	*	*	*	*
535.31	*5	*	*	—	*	*	*	*	*	*	*
535.4	127	53	75	*	18	35	73	23	37	44	24
535.40	94	34	60	*	16	26	51	15	28	34	18
535.41	33	19	15	*	*	9	21	*8	9	10	*6
535.5	244	96	148	*	43	67	129	66	54	99	24
535.50	211	83	128	*	41	58	108	56	49	85	22
535.51	32	13	20	*	*	9	20	11	*6	13	*
535.6	66	36	30	*	13	17	35	14	21	18	13
535.60	54	30	24	*	12	13	28	11	17	16	9
535.61	12	*6	*6	*	*	*	*7	*	*	*	*
536	254	103	151	18	58	72	107	39	72	105	38
536.2	33	15	18	*9	*9	*8	*8	*	*	15	9
536.3	81	29	52	*	24	28	28	14	18	40	9
536.4	34	15	19	*	*	*	22	*7	*6	15	*6
536.41	10	*5	*	*	*	*	*7	*	*	*	*
536.42	13	*6	*7	*	*	*	*7	*	*	*5	*
536.49	*8	*	*	*	*	*	*5	*	*	*	*
536.8	84	35	49	*	16	27	40	11	38	24	12
536.9	21	*8	12	*	*	*6	9	*	*6	11	*
537	67	31	36	*	*7	17	40	17	18	19	13
537.0	21	9	12	*	*	*6	11	*	*6	*8	*
537.3	*7	*	*	*	*	*	*	*	*	*	*
537.8	35	16	19	*	*	*9	24	10	*9	10	*7
537.83	15	*6	*9	—	*	*	13	*	*	*	*
537.89	14	*8	*7	*	*	*6	*6	*	*	*	*
540	271	151	120	64	143	45	19	57	65	82	67
540.0	57	32	25	16	25	12	*	11	16	15	15
540.1	36	18	18	*6	15	*9	*7	*7	10	13	*7
540.9	178	101	77	42	103	24	*8	39	39	54	45
541	13	*6	*7	*	*8	*	*	*	*	*7	*
542	*5	*	*	*	*	*	*	*	*	*	*
543	15	*7	*9	*	*8	*	*	*	*	*	*
543.9	12	*	*7	*	*7	*	*	*	*	*	*
550	85	70	15	*8	11	19	47	22	16	29	19
550.1	26	21	*5	*	*	*5	14	*8	*	*6	*7
550.10	21	16	*	*	*	*	11	*7	*	*	*
550.9	59	49	9	*5	*7	14	32	13	11	23	12
550.90	41	34	*7	*	*5	9	25	9	*7	16	*8
550.92	13	11	*	*	*	*	*	*	*	*	*
552	65	21	44	*	10	25	30	19	14	23	9
552.1	17	*7	10	*	*	*8	*5	*	*	*7	*
552.2	35	11	24	—	*	14	18	11	*6	13	*
552.20	10	*	*8	—	*	*	*8	*	*	*	*
552.21	23	*8	15	—	*	12	9	*7	*	*8	*
553	524	188	337	*7	68	144	306	118	136	201	70
553.1	47	21	26	*	11	14	20	*8	13	17	10
553.2	96	35	61	*	18	34	43	20	23	38	14
553.20	26	9	17	*	*	10	12	*	*6	12	*
553.21	65	23	42	*	12	23	30	14	17	25	9
553.3	378	130	248	*	38	95	241	89	99	144	46
555	110	43	67	*	45	36	28	30	27	38	15
555.0	19	*7	12	*	11	*6	*	*5	*6	*5	*
555.1	15	*5	9	*	*7	*	*	*6	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
555.2	*8	*	*	*	*	*	*	*	*	*	*
555.9	69	27	42	*	23	24	20	17	17	27	*8
556	54	26	27	*	22	12	19	15	12	15	11
556.6	*6	*	*	*	*	*	*	*	*	*	*
556.8	*5	*	*	*	*	*	*	*	*	*	*
556.9	34	16	18	*	13	*6	15	10	*6	10	*7
557	76	20	55	*	*6	16	50	15	19	28	14
557.0	37	13	24	*	*	*7	25	*8	10	11	*9
557.1	*6	*	*	*	*	*	*	*	*	*	*
557.9	32	*6	26	*	*	*8	21	*6	*7	15	*
558	367	142	224	104	79	59	125	72	83	153	59
558.1	11	*	*9	—	*	*	*8	*	*	*	*
558.9	353	139	214	104	78	56	116	67	79	149	58
560	674	273	401	21	101	172	381	138	162	251	123
560.1	283	132	151	*8	54	72	149	52	70	111	51
560.2	15	*6	*9	*	*	*	10	*	*	*	*
560.3	86	28	58	*	*5	11	68	17	17	42	10
560.30	9	*	*6	*	*	*	*7	*	*	*6	*
560.39	76	25	51	*	*	*9	60	14	16	36	9
560.8	121	47	73	*	22	34	62	24	31	37	29
560.81	68	25	43	*	11	22	32	12	16	24	15
560.89	53	22	30	*	10	12	29	12	15	13	13
560.9	166	58	108	*	19	53	91	40	41	54	31
562	546	199	347	—	39	111	396	141	130	195	80
562.0	*7	*	*	—	*	*	*6	*	*	*	*
562.1	539	195	344	—	38	111	390	140	127	193	79
562.10	243	81	161	—	9	40	194	68	55	86	34
562.11	189	74	115	—	26	59	104	48	50	62	29
562.12	88	33	55	—	*	9	78	21	15	37	15
562.13	19	*7	12	—	*	*	15	*	*7	*7	*
564	454	146	308	30	83	108	234	89	120	166	78
564.0	303	111	192	25	42	60	176	61	77	114	51
564.1	116	19	97	*	35	38	42	23	32	40	21
564.2	*6	*	*	*	*	*	*	*	*	*	*
564.8	13	*6	*7	*	*	*	*6	*	*	*6	*
564.89	10	*	*6	*	*	*	*5	*	*	*5	*
564.9	*6	*	*	—	*	*	*	*	*	*	*
565	13	*7	*6	*	*	*	*	*	*	*	*
565.0	*8	*	*	*	*	*	*	*	*	*	*
565.1	*6	*	*	*	*	*	*	*	*	*	*
566	23	17	*7	*	*8	*9	*	*	*5	10	*
567	84	45	39	*	20	30	30	19	17	29	19
567.2	54	31	23	*	12	21	17	14	12	16	13
567.8	11	*6	*	*	*	*	*	*	*	*	*
567.9	19	*8	12	*	*	*5	*9	*	*	*9	*
568	161	42	120	*	55	47	57	39	35	59	28
568.0	132	34	97	*	39	41	51	33	26	51	22
568.8	28	*6	22	*	16	*6	*5	*6	*8	*8	*6
568.81	18	*	14	*	11	*	*	*	*	*5	*
568.89	9	*	*7	*	*5	*	*	*	*	*	*
569	271	128	143	*6	32	65	167	61	70	93	47
569.0	9	*6	*	*	*	*	*	*	*	*	*
569.1	10	*	*8	*	*	*	*5	*	*	*	*
569.2	*5	*	*	—	*	*	*	*	*	*	*
569.3	54	27	27	*	*5	14	35	13	15	20	*6
569.4	27	14	14	*	*6	*5	15	*7	*6	10	*
569.41	*8	*	*	*	*	*	*	*	*	*	*
569.49	16	*8	*7	*	*	*	*9	*	*	*6	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
569.5	21	10	11	*	*	*7	11	*5	*5	*5	*5
569.6	31	16	15	*	*	*9	16	*6	*6	13	*6
569.62	*6	*	*	*	*	*	*	*	*	*	*
569.69	19	10	9	*	*	*6	10	*	*	*7	*
569.8	104	44	60	*	10	23	68	25	26	33	20
569.81	15	*8	*7	*	*	*5	*8	*6	*	*	*
569.82	16	*6	10	*	*	*6	*9	*	*	*6	*
569.83	22	*7	14	*	*	*6	13	*	*6	*8	*
569.84	*9	*	*	*	*	*	*8	*	*	*	*
569.85	26	12	14	*	*	*	22	*7	*7	*6	*6
569.89	17	*6	11	*	*	*	*8	*	*	*5	*
569.9	*9	*	*	—	*	*	*8	*	*	*	*
570	20	10	10	*	*	*7	*8	*7	*	*6	*
571	360	211	149	*	81	187	90	84	83	129	64
571.1	44	27	16	*	18	22	*	10	11	16	*7
571.2	135	94	41	—	28	82	26	31	27	52	25
571.3	25	19	*7	—	*9	13	*	*	11	*5	*
571.4	11	*	*7	*	*	*	*	*	*	*	*
571.40	*6	*	*	*	*	*	*	*	*	*	*
571.5	114	53	61	*	14	52	48	28	27	40	18
571.8	17	*8	*8	*	*	9	*	*	*	*6	*
571.9	*7	*	*	*	*	*	*	*	*	*	*
572	130	72	58	*	24	65	39	37	22	44	26
572.0	*7	*	*	*	*	*	*	*	*	*	*
572.2	60	33	27	*	11	29	19	18	*8	23	11
572.3	42	23	18	*	*8	21	12	10	10	12	10
572.4	*8	*	*5	*	*	*5	*	*	*	*	*
572.8	13	*8	*	*	*	*7	*	*	*	*	*
573	70	36	34	*	20	22	25	20	13	27	10
573.3	30	14	16	*	13	*6	*9	*8	*5	13	*
573.8	18	10	*9	*	*	*8	*8	*6	*	*	*
573.9	16	*9	*7	*	*	*5	*7	*	*	*7	*
574	604	205	399	*	166	177	256	142	135	217	110
574.0	129	51	77	*	41	39	49	30	29	48	21
574.00	118	47	72	*	39	35	44	28	26	45	19
574.01	11	*	*6	*	*	*	*5	*	*	*	*
574.1	257	77	180	*	77	86	93	57	51	102	48
574.10	246	73	173	*	74	82	88	56	48	98	44
574.11	11	*	*7	—	*	*	*	*	*	*	*
574.2	133	46	87	*	28	34	69	34	33	45	21
574.20	126	43	83	*	26	33	66	31	33	41	21
574.21	*7	*	*	*	*	*	*	*	*	*	*
574.4	*8	*	*	—	*	*	*5	*	*	*	*
574.5	32	13	19	*	*6	*7	19	12	*7	*6	*8
574.50	20	*8	12	*	*	*	12	*8	*	*	*
574.51	12	*5	*7	—	*	*	*7	*	*	*	*
574.6	*6	*	*	—	*	*	*	*	*	*	*
574.7	18	*	14	*	*8	*	*6	*	*6	*	*
574.70	11	*	*8	—	*	*	*	*	*	*	*
574.71	*8	*	*5	*	*	*	*	*	*	*	*
574.8	*9	*	*	—	*	*	*5	*	*	*	*
574.80	*6	*	*	—	*	*	*	*	*	*	*
574.9	*8	*	*5	*	*	*	*	*	*	*	*
574.90	*5	*	*	*	*	*	*	*	*	*	*
575	128	50	78	*	43	39	45	23	28	55	22
575.0	30	14	16	—	*8	10	12	*5	*7	11	*6
575.1	57	19	38	*	19	17	21	10	10	31	*6
575.10	14	*6	*8	—	*	*	*5	*	*	*8	*
575.11	35	10	25	*	13	11	11	*8	*6	18	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
575.12	*8	*	*	—	*	*	*	*	*	*5	*
575.3	*7	*	*5	*	*	*	*	*	*	*	*
575.6	*6	*	*	—	*	*	*	*	*	*	*
575.8	17	*7	11	*	*7	*	*	*	*6	*5	*
576	107	50	56	*	17	32	55	32	22	30	22
576.1	38	23	16	*	*	13	20	13	*7	11	*7
576.2	29	12	17	*	*	*8	17	9	*	*9	*7
576.8	32	12	20	*	*7	*7	17	*9	*8	9	*6
577	385	198	186	10	132	140	103	79	84	150	72
577.0	285	145	140	*7	103	96	78	58	62	111	54
577.1	61	33	28	*	21	28	11	12	11	26	11
577.2	17	11	*6	*	*5	*7	*	*	*	*7	*
577.8	16	*7	*9	*	*	*	*7	*	*6	*	*
577.9	*7	*	*	*	*	*	*	*	*	*	*
578	468	223	245	11	49	95	313	104	120	163	81
578.0	58	33	25	*	12	12	31	14	17	18	*9
578.1	113	52	61	*	12	26	70	25	28	39	21
578.9	297	138	159	*	24	57	211	65	75	106	51
579	45	17	28	12	*5	15	12	10	11	18	*6
579.0	*6	*	*	*	*	*	*	*	*	*	*
579.3	19	*5	13	*5	*	*6	*	*	*5	*8	*
579.8	12	*8	*	*5	*	*	*	*	*	*	*
579.9	*7	*	*	*	*	*	*	*	*	*	*
580-629	7,451	2,464	4,987	167	1,955	1,852	3,477	1,621	1,722	2,862	1,246
580	11	*6	*5	*	*	*	*	*	*	*	*
580.9	*6	*	*	*	*	*	*	*	*	*	*
581	42	18	24	*	*7	15	16	11	*9	13	9
581.8	21	10	11	*	*	*8	10	*	*	*5	*7
581.81	20	9	11	*	*	*8	10	*	*	*	*7
581.9	16	*7	10	*	*	*	*5	*5	*	*6	*
582	30	14	17	*	*6	9	15	*9	*	13	*
582.9	23	12	11	*	*	*7	14	*7	*	10	*
583	145	59	86	*	23	59	62	35	29	49	33
583.8	132	53	79	*	19	54	58	31	26	44	31
583.81	125	49	76	*	16	53	57	30	25	42	29
583.89	*7	*	*	*	*	*	*	*	*	*	*
583.9	11	*6	*	*	*	*	*	*	*	*	*
584	468	244	224	*	39	111	313	136	95	158	79
584.5	50	26	23	*	*	14	29	16	10	13	11
584.8	*5	*	*	*	*	*	*	*	*	*	*
584.9	413	215	198	*	33	96	281	119	84	143	66
585	303	148	156	*	27	81	191	78	68	102	55
586	51	27	24	*	*	12	36	9	14	21	*7
588	14	*5	9	*	*	*	*5	*	*	*6	*
588.8	10	*	*6	*	*	*	*	*	*	*	*
590	210	35	175	30	97	36	46	35	50	75	50
590.1	110	16	94	14	53	18	25	18	30	36	25
590.10	110	16	94	14	53	18	25	18	30	36	25
590.8	93	16	77	15	41	18	19	15	18	36	23
590.80	93	16	77	15	41	18	19	15	18	36	23
591	159	73	86	*7	46	36	70	42	34	57	27
592	258	135	122	*5	107	84	62	61	60	108	29
592.0	102	43	59	*	39	31	30	25	22	41	14
592.1	154	91	63	*	67	52	32	35	37	67	14
593	569	287	282	19	45	101	404	121	144	214	91

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
593.2	17	*9	*8	*	*	*7	9	*6	*	*	*
593.3	10	*	*6	*	*	*	*	*	*	*	*
593.4	27	10	17	*	*7	*7	12	*	*7	12	*
593.5	*7	*	*	*	*	*	*	*	*	*	*
593.7	14	*	9	12	*	*	*	*	*	*	*
593.70	11	*	*8	10	*	*	*	*	*	*	*
593.8	20	10	10	*	*5	*5	*8	*	*	*7	*
593.89	14	*7	*7	*	*	*	*	*	*	*	*
593.9	471	244	226	*	25	76	367	95	123	179	74
594	20	17	*	*	*	*	15	*5	*	*8	*
594.1	18	17	*	*	*	*	13	*	*	*7	*
595	66	23	43	*	9	15	40	17	18	21	*9
595.0	12	*	*8	*	*	*	*8	*	*	*	*
595.1	*6	*	*6	—	*	*	*	*	*	*	*
595.2	10	*	*5	—	*	*	*7	*	*	*	*
595.8	*7	*	*	*	—	*	*6	*	*	*	*
595.82	*5	*	*	—	—	*	*	*	*	*	*
595.9	30	10	20	*	*6	*8	16	*7	*8	10	*
596	205	136	69	9	26	35	135	40	62	64	39
596.0	67	64	*	*	*	9	57	9	20	24	14
596.1	*7	*	*	*	*	*	*	*	*	*	*
596.3	*8	*6	*	*	*	*	*6	*	*	*	*
596.4	*8	*	*	*	*	*	*6	*	*	*	*
596.5	73	36	37	*6	13	16	37	19	21	19	14
596.54	64	32	32	*6	12	14	32	16	19	17	12
596.59	*8	*	*	*	*	*	*	*	*	*	*
596.8	29	13	16	*	*6	*	17	*6	10	9	*
598	34	23	11	*	*5	*6	22	*7	*6	16	*
598.9	29	19	10	*	*	*5	19	*	*	15	*
599	2,018	672	1,346	60	248	328	1,382	474	455	786	304
599.0	1,711	473	1,238	55	201	259	1,196	392	388	677	254
599.6	34	30	*	*	*	*8	23	*8	*9	13	*
599.7	248	164	85	*	41	52	152	67	56	86	39
599.8	13	*	11	*	*	*6	*	*	*	*	*
599.82	*5	*	*	—	*	*	*	*	*	*	*
599.9	*6	*	*	—	*	*	*5	*	*	*	*
600	398	398	—	—	*	63	332	103	97	133	65
601	65	65	—	—	*5	18	41	15	15	24	11
601.0	18	18	—	—	*	*6	10	*	*	*6	*
601.1	26	26	—	—	*	*5	19	*7	*	10	*5
601.9	18	18	—	—	*	*6	10	*	*7	*7	*
603	*8	*8	—	*	*	*	*	*	*	*	*
603.9	*5	*5	—	*	*	*	*	*	*	*	*
604	12	12	—	*	*	*	*	*	*	*	*
604.9	11	11	—	*	*	*	*	*	*	*	*
604.90	9	9	—	*	*	*	*	*	*	*	*
605	12	12	—	*	*	*	*6	*	*	*	*
607	19	19	—	*	*	*6	10	*	*	*8	*
607.8	12	12	—	*	*	*	*6	*	*	*6	*
607.84	*7	*7	—	—	*	*	*	*	*	*	*
608	18	18	—	*	*5	*7	*	*	*	*9	*
608.4	*7	*7	—	*	*	*	*	*	*	*	*
608.8	*7	*7	—	*	*	*	*	*	*	*	*
610	26	—	26	—	*8	11	*8	11	*	*8	*
610.1	19	—	19	—	*	*8	*7	*6	*	*7	*
611	56	*	54	*	23	18	14	20	*9	18	*9
611.0	11	*	11	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
611.1	21	*	20	*	14	*	*	10	*	*	*
611.7	15	*	14	—	*	*	*7	*	*	*6	*
611.72	11	*	11	—	*	*	*6	*	*	*	*
611.8	*7	—	*7	—	*	*	*	*	*	*	*
614	260	—	260	*	171	77	10	45	53	126	37
614.0	*7	—	*7	*	*	*	*	*	*	*	*
614.1	28	—	28	—	15	11	*	*	*	15	*
614.2	11	—	11	*	*7	*	*	*	*	*6	*
614.3	*5	—	*5	*	*	*	*	*	*	*	*
614.4	*7	—	*7	*	*	*	*	*	*	*	*
614.6	173	—	173	*	109	57	*6	29	35	83	26
614.9	28	—	28	*	26	*	*	*	*7	14	*
615	20	—	20	*	17	*	*	*	*5	*8	*
615.9	12	—	12	*	11	*	*	*	*	*6	*
616	168	—	168	*	87	61	19	30	39	74	25
616.0	133	—	133	—	67	55	11	23	29	60	20
616.1	25	—	25	*	15	*	*	*	*8	11	*
616.10	25	—	25	*	15	*	*	*	*8	11	*
617	288	—	288	—	173	103	12	54	65	115	55
617.0	172	—	172	—	87	75	11	31	39	66	37
617.1	36	—	36	—	26	10	*	*8	*8	11	*8
617.2	*9	—	*9	—	*6	*	*	*	*	*	*
617.3	42	—	42	—	33	*8	*	10	10	16	*7
617.9	22	—	22	—	15	*6	*	*	*	14	*
618	279	—	279	—	74	117	88	44	70	111	54
618.0	80	—	80	—	19	37	25	14	17	33	17
618.1	39	—	39	—	22	11	*6	*8	10	16	*
618.2	25	—	25	—	*7	13	*	*6	*6	*8	*6
618.3	20	—	20	—	*	*8	*8	*	*	10	*
618.4	33	—	33	—	*9	15	9	*	*9	14	*6
618.5	31	—	31	—	*	11	18	*	*9	13	*7
618.6	30	—	30	—	*5	13	12	*	10	11	*6
618.8	21	—	21	—	*7	*9	*	*	*5	*7	*7
619	10	—	10	*	*	*	*	*	*	*	*
619.1	*7	—	*7	*	*	*	*	*	*	*	*
620	322	—	322	*	191	109	18	56	85	126	55
620.0	50	—	50	*	31	18	*	*8	14	20	*7
620.1	68	—	68	*	41	26	*	10	17	29	11
620.2	143	—	143	*	86	42	12	26	33	57	26
620.5	*7	—	*7	*	*	*	*	*	*	*	*
620.8	47	—	47	*	26	17	*	10	16	13	*9
621	93	—	93	*	38	41	14	19	22	32	21
621.0	38	—	38	—	14	19	*5	9	*7	14	*8
621.2	9	—	9	—	*6	*	*	*	*	*	*
621.3	17	—	17	—	*	9	*	*	*	*6	*
621.5	*6	—	*6	—	*	*	*	*	*	*	*
621.8	15	—	15	—	*7	*	*	*	*	*	*5
622	38	—	38	—	20	14	*	*7	9	15	*7
622.1	23	—	23	—	15	*7	*	*	*5	11	*
622.7	*8	—	*8	—	*	*	*	*	*	*	*
623	21	—	21	*	11	*	*6	*	*	9	*
623.8	15	—	15	*	*8	*	*	*	*	*7	*
625	370	—	370	*	220	112	38	45	75	180	71
625.0	32	—	32	—	29	*	—	*	*7	16	*6
625.3	81	—	81	*	61	20	—	9	14	43	14
625.6	149	—	149	*	46	67	36	18	30	69	32
625.8	*7	—	*7	*	*	*	*	*	*	*	*
625.9	92	—	92	*	72	19	*	12	21	44	15

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
626	287	—	287	*	179	105	*	47	72	118	50
626.2	205	—	205	*	131	74	*	34	51	83	37
626.4	*5	—	*5	*	*	*	—	*	*	*	*
626.6	*5	—	*5	—	*	*	*	*	*	*	*
626.8	56	—	56	*	34	21	*	*9	14	23	9
626.9	11	—	11	*	*	*6	*	*	*	*6	*
627	53	—	53	—	*7	32	14	9	14	17	14
627.0	*6	—	*6	—	*	*	—	*	*	*	*
627.1	27	—	27	—	*	19	*8	*8	*6	*6	*8
627.2	*8	—	*8	—	*	*	*	*	*	*	*
627.9	*6	—	*6	—	*	*	*	*	*	*	*
628	*6	—	*6	*	*5	*	—	*	*	*	*
630-677	8,547	—	8,547	27	8,508	12	—	1,519	1,776	3,182	2,069
632	15	—	15	—	15	*	—	*	*	*6	*
633	35	—	35	—	35	*	—	10	*5	11	9
633.1	30	—	30	—	30	*	—	*8	*	10	*8
634	27	—	27	*	27	*	—	*7	*	12	*7
634.9	19	—	19	*	19	—	—	*5	*	*7	*
634.91	14	—	14	—	14	—	—	*5	*	*5	*
635	*5	—	*5	*	*	*	—	*	*	*	*
637	*7	—	*7	—	*7	—	—	*	*	*	*
637.9	*7	—	*7	—	*7	—	—	*	*	*	*
639	*9	—	*9	—	*9	*	—	*	*	*	*
640	11	—	11	—	11	*	—	*	*	*	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*6	—	*6	—	*6	—	—	*	*	*	*
641	81	—	81	—	81	*	—	15	15	26	25
641.0	*5	—	*5	—	*5	*	—	*	*	*	*
641.1	18	—	18	—	17	*	—	*	*	*6	*
641.11	11	—	11	—	11	—	—	*	*	*	*
641.13	*7	—	*7	—	*6	*	—	*	*	*	*
641.2	39	—	39	—	39	*	—	*6	*7	14	12
641.21	35	—	35	—	35	—	—	*6	*6	13	10
641.9	13	—	13	—	13	—	—	*	*	*	*
641.91	*8	—	*8	—	*8	—	—	*	*	*	*
641.93	*5	—	*5	—	*5	—	—	*	*	*	*
642	325	—	325	*	323	*	—	55	54	154	63
642.0	33	—	33	*	32	*	—	*6	*6	16	*
642.01	27	—	27	*	27	*	—	*	*6	14	*
642.2	*5	—	*5	—	*5	—	—	*	*	*	*
642.3	134	—	134	*	133	*	—	25	22	58	29
642.31	117	—	117	*	116	*	—	20	21	51	25
642.33	14	—	14	—	14	—	—	*	*	*6	*
642.4	82	—	82	*	81	*	—	13	17	35	17
642.41	71	—	71	*	70	*	—	10	14	31	16
642.43	*8	—	*8	*	*8	—	—	*	*	*	*
642.5	33	—	33	*	33	*	—	*	*	18	*6
642.51	31	—	31	*	31	*	—	*	*	17	*6
642.6	*5	—	*5	—	*5	—	—	*	*	*	*
642.7	13	—	13	—	13	*	—	*	*	9	*
642.71	12	—	12	—	12	*	—	*	*	*8	*
642.9	20	—	20	—	20	—	—	*	*	11	*
642.91	14	—	14	—	14	—	—	*	*	*7	*
643	60	—	60	—	60	—	—	*8	17	26	*9
643.0	26	—	26	—	26	—	—	*	*7	15	*
643.03	25	—	25	—	25	—	—	*	*7	14	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
643.1	22	—	22	—	22	—	—	*	*	*8	*
643.13	21	—	21	—	21	—	—	*	*	*8	*
643.2	*9	—	*9	—	*9	—	—	*	*	*	*
643.23	*7	—	*7	—	*7	—	—	*	*	*	*
644	367	—	367	*	366	*	—	66	64	167	70
644.0	108	—	108	*	108	*	—	21	23	41	24
644.03	108	—	108	*	108	*	—	21	22	40	24
644.1	21	—	21	—	21	—	—	*	*	13	*
644.13	20	—	20	—	20	—	—	*	*	13	*
644.2	238	—	238	*	237	*	—	41	39	114	44
644.21	238	—	238	*	237	*	—	41	39	114	44
645	117	—	117	*	116	*	—	21	16	38	41
645.0	117	—	117	*	116	*	—	21	16	38	41
645.01	115	—	115	*	114	*	—	20	15	38	41
646	258	—	258	*	257	*	—	46	52	110	50
646.1	23	—	23	*	23	—	—	*	*5	*7	9
646.11	21	—	21	*	21	—	—	*	*	*7	*8
646.2	12	—	12	—	12	—	—	*	*	*	*
646.21	*5	—	*5	—	*5	—	—	*	*	*	*
646.23	*6	—	*6	—	*6	—	—	*	*	*	*
646.6	162	—	162	*	161	—	—	31	33	69	28
646.61	90	—	90	*	90	—	—	17	19	41	13
646.62	*7	—	*7	—	*7	—	—	*	*	*	*
646.63	61	—	61	*	61	—	—	12	12	25	12
646.8	53	—	53	*	53	—	—	*8	11	24	10
646.81	27	—	27	*	26	—	—	*	*6	10	*6
646.83	24	—	24	—	24	—	—	*	*	13	*
647	95	—	95	*	95	—	—	14	16	50	15
647.6	41	—	41	*	41	—	—	*	*6	22	*9
647.61	35	—	35	—	35	—	—	*	*	19	*8
647.8	46	—	46	—	46	—	—	*9	*8	22	*7
647.81	35	—	35	—	35	—	—	*5	*7	18	*5
647.83	*9	—	*9	—	*9	—	—	*	*	*	*
648	1,107	—	1,107	*	1,103	*	—	211	232	418	246
648.0	38	—	38	*	38	*	—	*	*8	15	11
648.01	23	—	23	*	23	*	—	*	*5	9	*7
648.03	14	—	14	—	14	—	—	*	*	*5	*
648.1	42	—	42	—	42	—	—	9	10	14	*9
648.11	37	—	37	—	37	—	—	*8	9	11	*9
648.2	291	—	291	*	289	*	—	49	60	118	65
648.21	178	—	178	*	177	*	—	28	34	80	36
648.22	84	—	84	*	84	*	—	16	20	25	22
648.23	23	—	23	—	23	—	—	*	*	10	*5
648.24	*6	—	*6	—	*6	—	—	*	*	*	*
648.3	*8	—	*8	*	*8	—	—	*	*	*	*
648.4	121	—	121	*	120	—	—	19	34	41	28
648.41	98	—	98	*	98	—	—	15	27	32	24
648.43	17	—	17	*	17	—	—	*	*5	*6	*
648.6	35	—	35	—	35	*	—	10	*6	14	*5
648.61	29	—	29	—	29	—	—	*7	*6	12	*
648.7	*5	—	*5	—	*5	—	—	*	*	*	*
648.8	155	—	155	—	154	*	—	35	27	55	38
648.81	132	—	132	—	132	*	—	32	25	42	34
648.83	20	—	20	—	20	*	—	*	*	11	*
648.9	411	—	411	*	410	*	—	81	85	158	87
648.91	329	—	329	*	329	*	—	63	71	126	69
648.92	*8	—	*8	—	*8	—	—	*	*	*	*
648.93	72	—	72	*	71	—	—	15	13	29	15
650	475	—	475	*	475	—	—	72	89	208	107

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
651	65	—	65	*	65	*	—	14	15	26	11
651.0	61	—	61	*	61	*	—	12	13	25	11
651.01	43	—	43	—	43	*	—	*7	10	17	*8
651.03	18	—	18	*	18	*	—	*5	*	*7	*
652	259	—	259	*	258	*	—	49	58	88	64
652.1	*6	—	*6	—	*6	—	—	*	*	*	*
652.11	*6	—	*6	—	*6	—	—	*	*	*	*
652.2	116	—	116	—	116	*	—	20	24	45	27
652.21	110	—	110	—	110	*	—	19	22	44	25
652.23	*6	—	*6	—	*6	—	—	*	*	*	*
652.3	24	—	24	—	24	*	—	*	*	*7	*8
652.31	24	—	24	—	24	*	—	*	*	*7	*8
652.5	36	—	36	*	36	—	—	9	*8	12	*7
652.51	36	—	36	*	36	—	—	9	*8	12	*7
652.6	16	—	16	—	16	*	—	*	*	*	*
652.61	15	—	15	—	15	*	—	*	*	*	*
652.8	49	—	49	*	49	*	—	10	13	13	14
652.81	49	—	49	*	49	*	—	10	13	13	14
652.9	*5	—	*5	—	*5	—	—	*	*	*	*
652.91	*5	—	*5	—	*5	—	—	*	*	*	*
653	111	—	111	*	110	*	—	18	19	49	25
653.4	85	—	85	*	85	*	—	11	16	40	18
653.41	85	—	85	*	84	*	—	11	16	40	18
653.5	18	—	18	—	18	—	—	*	*	*5	*5
653.51	18	—	18	—	18	—	—	*	*	*5	*5
654	564	—	564	*	562	*	—	105	110	215	133
654.0	14	—	14	—	14	—	—	*	*	*	*
654.01	12	—	12	—	12	—	—	*	*	*	*
654.1	25	—	25	—	25	*	—	*5	*	*9	*8
654.11	21	—	21	—	21	*	—	*	*	*6	*7
654.2	459	—	459	—	458	*	—	81	91	177	110
654.21	447	—	447	—	446	*	—	79	89	172	107
654.23	12	—	12	—	12	*	—	*	*	*	*
654.4	22	—	22	*	22	—	—	*	*	*8	*5
654.41	16	—	16	*	16	—	—	*	*	*6	*
654.43	*5	—	*5	—	*5	—	—	*	*	*	*
654.5	23	—	23	—	22	*	—	*6	*	9	*
654.51	12	—	12	—	12	*	—	*	*	*5	*
654.53	10	—	10	—	10	*	—	*	*	*	*
654.6	13	—	13	—	13	—	—	*	*	*6	*
654.61	10	—	10	—	10	—	—	*	*	*6	*
655	33	—	33	*	33	*	—	*5	*7	11	10
655.7	15	—	15	—	15	—	—	*	*	*6	*5
655.71	13	—	13	—	13	—	—	*	*	*	*
655.8	11	—	11	*	11	—	—	*	*	*	*
655.81	10	—	10	*	10	—	—	*	*	*	*
656	393	—	393	*	391	*	—	59	88	156	90
656.1	72	—	72	*	72	*	—	*9	18	28	17
656.11	70	—	70	*	70	*	—	*9	18	27	17
656.3	31	—	31	—	31	*	—	*	*7	13	*6
656.31	31	—	31	—	31	*	—	*	*7	13	*6
656.4	17	—	17	—	17	—	—	*	*	*8	*
656.41	17	—	17	—	17	—	—	*	*	*8	*
656.5	57	—	57	—	57	—	—	12	10	20	15
656.51	54	—	54	—	54	—	—	11	10	19	14
656.6	90	—	90	*	90	—	—	12	22	33	22
656.61	88	—	88	*	88	—	—	12	22	33	21
656.7	26	—	26	—	26	—	—	*	12	*	*7
656.71	26	—	26	—	26	—	—	*	12	*	*7

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
656.8	91	—	91	*	90	—	—	15	12	45	19
656.81	90	—	90	*	89	—	—	15	11	45	18
657	21	—	21	—	21	—	—	*	*	*8	*
657.0	21	—	21	—	21	—	—	*	*	*8	*
657.01	19	—	19	—	19	—	—	*	*	*8	*
658	358	—	358	*	357	*	—	76	66	138	79
658.0	86	—	86	*	86	*	—	22	11	36	17
658.01	80	—	80	*	79	—	—	21	10	33	15
658.03	*6	—	*6	—	*6	*	—	*	*	*	*
658.1	146	—	146	*	145	*	—	28	33	54	31
658.11	140	—	140	*	140	*	—	27	32	50	31
658.2	57	—	57	—	57	*	—	13	13	18	13
658.21	55	—	55	—	55	*	—	12	13	17	13
658.4	64	—	64	*	64	—	—	12	*8	29	14
658.41	62	—	62	*	62	—	—	12	*8	27	14
659	706	—	706	*8	695	*	—	116	132	247	211
659.1	34	—	34	—	34	*	—	*	*8	13	*9
659.11	25	—	25	—	25	*	—	*	*6	10	*6
659.13	9	—	9	—	9	*	—	*	*	*	*
659.2	43	—	43	*	43	—	—	*7	*8	15	13
659.21	43	—	43	*	43	—	—	*7	*8	14	13
659.4	14	—	14	—	14	—	—	*	*	*6	*
659.41	14	—	14	—	14	—	—	*	*	*6	*
659.5	22	—	22	*	22	*	—	*5	*	*7	*7
659.51	21	—	21	*	21	*	—	*	*	*7	*6
659.6	133	—	133	—	131	*	—	19	19	40	54
659.61	126	—	126	—	125	*	—	18	18	37	53
659.63	*6	—	*6	—	*6	*	—	*	*	*	*
659.7	359	—	359	*	357	*	—	64	71	133	91
659.71	357	—	357	*	354	*	—	64	71	132	91
659.8	68	—	68	*5	62	*	—	*8	15	14	30
659.81	65	—	65	*5	59	*	—	*8	15	13	30
659.9	29	—	29	—	29	—	—	*	*	18	*
659.91	29	—	29	—	29	—	—	*	*	18	*
660	202	—	202	*	202	*	—	31	51	67	53
660.0	38	—	38	—	38	—	—	*6	*8	11	13
660.01	38	—	38	—	38	—	—	*6	*8	11	13
660.1	37	—	37	*	37	*	—	*5	*5	13	13
660.11	37	—	37	*	36	*	—	*5	*5	13	13
660.3	41	—	41	—	41	*	—	*7	11	15	*8
660.31	41	—	41	—	41	*	—	*7	11	15	*8
660.4	62	—	62	—	62	—	—	9	21	18	13
660.41	62	—	62	—	62	—	—	9	21	18	13
660.6	*8	—	*8	—	*8	—	—	*	*	*	*
660.61	*8	—	*8	—	*8	—	—	*	*	*	*
660.7	10	—	10	*	10	—	—	*	*	*	*
660.71	10	—	10	*	10	—	—	*	*	*	*
661	346	—	346	*	345	*	—	71	74	124	77
661.0	77	—	77	*	77	*	—	15	19	23	19
661.01	76	—	76	*	75	*	—	15	18	23	18
661.1	74	—	74	—	74	*	—	20	12	27	15
661.11	74	—	74	—	74	*	—	20	12	27	15
661.2	83	—	83	*	82	*	—	19	15	34	15
661.21	82	—	82	*	82	*	—	19	15	34	15
661.3	97	—	97	—	97	—	—	13	23	34	27
661.31	97	—	97	—	97	—	—	13	23	34	27
661.9	10	—	10	—	10	—	—	*	*	*	*
661.91	*9	—	*9	—	*9	—	—	*	*	*	*
662	41	—	41	*	40	*	—	12	*8	*7	14

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
662.0	*9	—	*9	—	*9	—	—	*	*	*	*
662.01	*9	—	*9	—	*9	—	—	*	*	*	*
662.1	*8	—	*8	—	*8	—	—	*	*	*	*
662.11	*8	—	*8	—	*8	—	—	*	*	*	*
662.2	25	—	25	*	24	*	—	*7	*6	*	*7
662.21	25	—	25	*	24	*	—	*7	*6	*	*7
663	746	—	746	*	744	*	—	124	162	260	200
663.0	13	—	13	—	13	—	—	*	*	*6	*
663.01	13	—	13	—	13	—	—	*	*	*6	*
663.1	87	—	87	—	87	*	—	15	18	28	26
663.11	87	—	87	—	87	*	—	15	18	28	26
663.2	31	—	31	—	31	*	—	*5	*7	10	*8
663.21	31	—	31	—	31	*	—	*5	*7	10	*8
663.3	575	—	575	*	575	*	—	93	126	203	153
663.31	575	—	575	*	574	*	—	93	126	203	153
663.4	15	—	15	*	14	—	—	*	*	*	*
663.41	15	—	15	*	14	—	—	*	*	*	*
663.8	18	—	18	*	18	—	—	*	*	*6	*6
663.81	18	—	18	*	17	—	—	*	*	*5	*6
664	1,133	—	1,133	*	1,127	*	—	209	266	353	305
664.0	493	—	493	*	491	*	—	92	115	154	132
664.01	491	—	491	*	489	*	—	92	115	153	131
664.1	473	—	473	*	469	*	—	83	114	141	136
664.11	471	—	471	*	468	*	—	82	113	140	135
664.2	104	—	104	*	104	*	—	23	25	34	22
664.21	104	—	104	*	104	*	—	23	25	33	22
664.3	38	—	38	*	38	*	—	*7	*8	17	*7
664.31	38	—	38	*	38	*	—	*7	*8	17	*7
664.4	*7	—	*7	—	*7	—	—	*	*	*	*
664.41	*7	—	*7	—	*7	—	—	*	*	*	*
664.8	*6	—	*6	—	*6	—	—	*	*	*	*
664.81	*6	—	*6	—	*6	—	—	*	*	*	*
664.9	10	—	10	—	10	—	—	*	*	*	*5
664.91	10	—	10	—	10	—	—	*	*	*	*5
665	156	—	156	*	156	—	—	22	43	57	35
665.3	*8	—	*8	—	*8	—	—	*	*	*	*
665.31	*8	—	*8	—	*8	—	—	*	*	*	*
665.4	53	—	53	—	53	—	—	*7	12	23	11
665.41	53	—	53	—	53	—	—	*7	12	23	11
665.5	83	—	83	*	83	—	—	11	24	29	19
665.51	83	—	83	*	83	—	—	11	24	28	19
666	111	—	111	*	111	*	—	18	27	33	33
666.0	13	—	13	—	13	—	—	*	*	*	*
666.02	13	—	13	—	13	—	—	*	*	*	*
666.1	80	—	80	*	80	*	—	12	21	24	24
666.12	80	—	80	*	80	*	—	12	21	24	24
666.2	12	—	12	—	12	—	—	*	*	*	*
666.22	10	—	10	—	10	—	—	*	*	*	*
666.3	*5	—	*5	—	*5	—	—	*	*	*	*
667	22	—	22	—	22	—	—	*	*6	*6	*5
667.0	15	—	15	—	15	—	—	*	*	*	*
667.02	14	—	14	—	14	—	—	*	*	*	*
667.1	*7	—	*7	—	*7	—	—	*	*	*	*
667.12	*6	—	*6	—	*6	—	—	*	*	*	*
668	14	—	14	—	14	—	—	*	*	*	*
668.8	12	—	12	—	12	—	—	*	*	*	*
668.81	*5	—	*5	—	*5	—	—	*	*	*	*
668.82	*6	—	*6	—	*6	—	—	*	*	*	*
669	159	—	159	*	159	*	—	20	39	64	37

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
669.4	*7	—	*7	—	*7	—	—	*	*	*	*
669.5	80	—	80	*	80	—	—	9	20	38	14
669.51	80	—	80	*	80	—	—	9	20	38	14
669.7	10	—	10	—	10	—	—	*	*	*	*
669.71	10	—	10	—	10	—	—	*	*	*	*
669.8	54	—	54	—	54	*	—	*8	13	17	16
669.81	53	—	53	—	52	*	—	*8	12	17	16
670	21	—	21	*	21	—	—	*	*5	*9	*
670.0	21	—	21	*	21	—	—	*	*5	*9	*
670.02	13	—	13	*	13	—	—	*	*	*6	*
670.04	*8	—	*8	—	*8	—	—	*	*	*	*
671	28	—	28	—	28	—	—	*8	*7	*6	*7
671.8	17	—	17	—	17	—	—	*6	*	*	*
671.81	11	—	11	—	11	—	—	*	*	*	*
672	15	—	15	*	15	—	—	*	*	*	*
672.0	15	—	15	*	15	—	—	*	*	*	*
672.02	15	—	15	*	15	—	—	*	*	*	*
674	31	—	31	*	31	—	—	*6	*9	*9	*7
674.3	13	—	13	—	13	—	—	*	*	*	*
674.32	10	—	10	—	10	—	—	*	*	*	*
674.8	14	—	14	*	14	—	—	*	*6	*	*
674.82	10	—	10	*	10	—	—	*	*	*	*
676	*7	—	*7	*	*7	—	—	*	*	*	*
680-709	1,822	882	940	153	344	477	848	422	459	664	276
681	50	27	23	*	10	14	22	*7	16	19	*8
681.0	20	12	*8	*	*7	*	*6	*	*8	*6	*
681.00	18	11	*7	*	*6	*	*	*	*7	*5	*
681.1	30	15	15	*	*	9	16	*5	*8	12	*
681.10	28	14	14	*	*	*9	15	*	*8	11	*
682	749	380	369	42	169	227	310	185	186	261	116
682.0	37	20	17	9	12	*8	*7	9	9	12	*7
682.1	12	*8	*	*	*6	*	*	*	*	*6	*
682.2	82	36	46	*7	24	23	28	17	19	33	14
682.3	75	35	40	*	24	22	25	18	16	29	12
682.4	36	21	16	*	13	10	10	12	11	10	*
682.5	10	*5	*	*	*	*	*	*	*	*	*
682.6	335	165	171	*7	58	105	165	86	86	113	51
682.7	139	79	60	*6	23	48	61	33	35	47	23
682.8	*8	*	*	*	*	*	*	*	*	*	*
682.9	15	*7	*8	*	*	*	*6	*	*	*5	*
683	*6	*	*	*	*	*	*	*	*	*	*
684	*9	*5	*	*	*	*	*	*	*	*	*
686	18	9	9	*	*	*5	*8	*	*5	*8	*
686.8	*5	*	*	*	*	*	*	*	*	*	*
686.9	*9	*5	*	*	*	*	*	*	*	*	*
690	10	*7	*	*	*	*	*	*	*	*	*
690.1	*9	*7	*	*	*	*	*	*	*	*	*
690.10	*6	*	*	*	*	*	*	*	*	*	*
691	33	17	16	29	*	*	*	*	*6	*8	14
691.0	27	14	13	26	*	—	*	*	*	*5	14
691.8	*6	*	*	*	*	*	*	*	*	*	*
692	50	24	26	11	12	11	16	9	15	14	12
692.9	41	21	19	10	*9	*9	13	*9	12	11	*9
693	40	18	22	*	12	10	13	*8	11	12	*9
693.0	38	16	22	*	12	10	13	*8	10	11	*8

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
694	*7	*	*	—	*	*	*6	*	*	*	*
695	31	13	18	*	*9	*8	10	*6	9	11	*
695.1	*8	*	*	*	*	*	*	*	*	*	*
695.4	*6	*	*	—	*	*	*	*	*	*	*
695.8	*7	*	*	*	*	*	*	*	*	*	—
695.89	*7	*	*	*	*	*	*	*	*	*	—
695.9	*5	*	*	*	*	*	*	*	*	*	*
696	42	17	24	*	*7	17	17	15	*9	11	*7
696.1	36	14	22	*	*6	15	15	14	*6	9	*7
698	27	*7	20	*	12	*7	*7	*	11	*7	*
698.9	19	*5	13	*	*9	*	*	*	*7	*	*
701	21	*6	15	*	*6	*7	*6	*6	*	*7	*
701.8	*5	*	*	*	*	*	*	*	*	*	*
701.9	*6	*	*	*	*	*	*	*	*	*	*
703	13	*6	*7	*	*	*	9	*	*	*5	*
703.0	*6	*	*	*	*	*	*	*	*	*	*
703.8	*5	*	*	*	*	*	*	*	*	*	—
704	13	*	*8	*	*6	*	*	*	*	*7	*
705	*8	*	*	*	*	*	*	*	*	*	*
705.8	*6	*	*	*	*	*	*	*	*	*	*
706	16	*9	*7	*	*7	*5	*	*	*6	*	*
706.2	*9	*	*	*	*	*	*	*	*	*	*
707	581	275	306	*	55	134	388	144	143	224	70
707.0	306	130	176	*	25	45	235	74	72	127	33
707.1	252	135	117	*	28	82	141	64	64	89	36
707.8	11	*	*6	*	*	*	*	*	*	*	*
707.9	12	*5	*6	*	*	*	*8	*	*	*	*
708	17	*8	9	*	*6	*	*	*	*5	*	*
708.0	10	*	*6	*	*	*	*	*	*	*	*
709	65	32	33	29	16	10	10	*7	11	44	*
709.2	37	17	20	22	11	*	*	*	*	32	*
709.8	16	9	*7	*6	*	*	*5	*	*	*9	*
709.9	*8	*	*	*	*	*	*	*	*	*	*
710-739	5,499	1,928	3,571	74	823	1,430	3,172	1,212	1,438	1,946	903
710	152	20	132	*	50	58	44	33	42	53	24
710.0	104	13	90	*	41	38	24	20	27	39	19
710.1	22	*	19	*	*	9	*8	*6	*5	*7	*
710.2	*7	*	*7	*	*	*	*	*	*	*	*
710.4	*7	*	*	*	*	*	*	*	*	*	*
710.9	*6	*	*6	—	*	*	*	*	*	*	*
711	44	28	16	*	*8	15	16	11	10	16	*7
711.0	41	26	15	*	*8	14	15	9	9	16	*6
711.06	15	11	*	*	*	*6	*5	*	*	*6	*
711.07	*7	*	*	*	*	*	*	*	*	*	*
712	*7	*5	*	—	—	*	*6	*	*	*	*
712.3	*6	*	*	—	—	*	*6	*	*	*	*
713	*8	*	*5	*	*	*	*	*	*	*	*
714	232	69	163	*	17	65	147	55	63	78	37
714.0	217	65	152	*	14	63	140	51	61	74	32
714.3	*5	*	*	*	*	*	*	*	*	*	*
715	1,461	482	979	*	45	285	1,130	310	404	537	209
715.0	40	13	27	—	*	*5	34	16	*9	10	*
715.00	*9	*	*5	—	*	*	*8	*	*	*	*
715.09	31	9	21	—	*	*	26	15	*7	*7	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
715.1	52	18	34	*	*	16	33	12	12	18	11
715.15	11	*	*7	—	*	*	*8	*	*	*	*
715.16	34	11	23	*	*	11	21	*7	*8	12	*6
715.2	10	*	*6	—	*	*	*	*	*	*	*
715.3	302	116	186	*	13	83	205	49	88	114	51
715.30	*6	*	*	—	*	*	*	*	*	*	—
715.31	15	11	*	—	*	*	*9	*	*5	*	*
715.35	77	29	47	—	*	23	50	14	20	27	14
715.36	194	70	124	*	*6	50	137	29	58	75	32
715.8	54	13	41	—	*	*8	45	*8	23	16	*7
715.89	54	13	41	—	*	*8	45	*8	23	16	*7
715.9	1,003	316	686	*	24	170	808	222	272	378	131
715.90	683	195	488	*	15	96	572	154	190	266	73
715.91	19	*6	12	—	*	*	14	*	*5	*6	*
715.95	86	38	48	—	*	23	59	22	24	25	16
715.96	184	69	115	*	*	39	141	33	48	72	31
715.97	*6	*	*	—	*	*	*	*	*	*	*
715.98	18	*	14	—	*	*	14	*6	*	*6	*
716	263	90	173	*	23	74	166	57	67	110	29
716.1	11	*7	*	—	*	*5	*	*	*	*	*
716.5	*9	*	*6	*	*	*	*7	*	*	*	*
716.59	*8	*	*6	*	*	*	*6	*	*	*	*
716.9	239	78	160	*	17	66	155	52	62	101	24
716.90	180	56	124	*	*9	48	123	44	45	75	16
716.91	11	*	*7	—	*	*	*	*	*	*8	*
716.95	*6	*	*	*	*	*	*	*	*	*	*
716.96	16	*8	*8	*	*	*	9	*	*	*7	*
716.99	11	*	*9	*	*	*	*5	*	*	*6	*
717	33	17	16	*	18	*	9	*6	*	9	14
717.7	*8	*	*	*	*	*	*	*	*	*	*
717.8	10	*6	*	*	*8	*	*	*	*	*	*
717.83	*7	*	*	*	*7	*	—	*	*	*	*
718	51	25	26	*	15	11	22	10	13	18	11
718.3	13	*6	*7	*	*	*	*6	*	*	*	*
718.4	16	*8	*9	*	*	*	10	*	*	*5	*
718.46	*6	*	*	*	*	*	*	*	*	*	*
718.8	*8	*	*	*	*	*	*	*	*	*	*
719	217	90	128	*5	42	53	117	53	74	55	35
719.0	19	13	*6	*	*	*	9	*	*	*8	*
719.06	16	11	*	*	*	*	*8	*	*	*6	*
719.4	121	45	76	*	30	33	56	26	35	38	23
719.40	*6	*	*	*	*	*	*	*	*	*	*
719.41	39	16	23	*	11	13	15	10	13	11	*5
719.45	41	12	29	*	*6	*	29	*8	11	15	*8
719.46	22	9	12	*	*8	*6	*7	*	*	*8	*
719.49	*6	*	*	*	*	*	*	*	*	*	*
719.7	59	23	36	*	*	11	45	19	32	*	*
719.70	27	13	13	*	*	*5	21	*6	17	*	*
719.75	16	*	11	—	*	*	12	*6	*8	*	*
719.76	14	*	10	*	*	*	11	*7	*5	*	*
720	13	*8	*	*	*	*7	*	*	*	*5	*
720.0	*9	*6	*	—	*	*5	*	*	*	*	*
721	229	87	142	—	25	73	131	40	60	86	43
721.0	60	24	36	—	11	23	27	9	14	23	15
721.1	18	11	*7	—	*	*7	*8	*	*6	*	*
721.2	*8	*	*6	—	*	*	*8	*	*	*	*
721.3	95	32	63	—	*7	29	59	20	24	36	15
721.9	39	12	27	—	*	9	26	*7	12	16	*
721.90	38	12	26	—	*	9	26	*7	12	16	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
722	490	249	241	*	182	189	118	94	117	193	86
722.0	68	34	35	—	30	34	*	11	13	29	14
722.1	226	121	105	*	97	81	48	49	54	87	36
722.10	222	119	102	*	96	79	46	48	53	85	35
722.2	*9	*	*5	—	*	*	*	*	*	*7	—
722.4	21	10	11	—	*6	12	*	*	*6	*7	*
722.5	90	43	47	—	24	30	35	16	25	32	17
722.52	86	41	46	—	23	29	34	15	24	32	16
722.6	12	*	*7	—	*	*	*6	*	*	*6	*
722.7	22	11	11	—	*7	10	*5	*	*5	*8	*
722.71	14	*7	*6	—	*	*7	*	*	*	*5	*
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	15	*8	*6	*	*	*7	*	*	*	*5	*
722.83	13	*7	*6	*	*	*6	*	*	*	*	*
722.9	27	13	14	*	*7	9	11	*5	*	12	*
722.93	18	*9	9	*	*	*6	*6	*	*	*7	*
723	60	27	33	*	18	27	14	14	13	20	15
723.0	14	*8	*6	*	*	*6	*	*	*	*	*5
723.1	28	11	17	*	11	10	*6	*6	*	10	*6
723.4	16	*7	*8	—	*	10	*	*	*	*5	*
724	451	180	271	*	98	142	207	94	118	156	83
724.0	129	54	75	*	*7	36	86	29	29	46	25
724.00	27	9	17	—	*	*	21	*7	*6	11	*
724.02	98	44	54	*	*5	31	62	21	22	34	21
724.2	148	57	91	*	44	47	55	25	43	51	29
724.3	21	*8	13	—	*	*5	11	*7	*5	*5	*
724.4	29	11	17	*	*6	*7	15	*7	*6	11	*
724.5	102	40	62	*	33	36	32	21	26	37	18
724.6	*5	*	*	—	*	*	*	*	*	*	*
724.8	10	*	*7	*	*	*	*	*	*5	*	*
725	35	*8	27	—	—	*	31	*7	11	12	*5
726	114	48	66	*	22	43	48	20	25	47	21
726.0	9	*	*7	*	*	*	*	*	*	*	*
726.1	40	16	25	—	*5	13	22	*7	10	16	*7
726.10	36	13	23	—	*	11	21	*6	9	15	*6
726.2	26	10	16	*	*6	11	*9	*	*	14	*5
726.3	12	*8	*	*	*	*	*	*	*	*	*
726.33	11	*7	*	*	*	*	*	*	*	*	*
726.5	*9	*	*5	—	*	*	*5	*	*	*	*
726.6	*6	*	*	*	*	*	*	*	*	*	*
726.9	*8	*	*5	—	*	*	*	*	*	*	*
726.91	*7	*	*	—	*	*	*	*	*	*	*
727	64	27	37	*	12	21	27	15	13	25	10
727.0	24	10	15	*	*6	*6	9	*7	*6	*7	*5
727.00	14	*	10	*	*	*	*6	*	*	*	*
727.05	*6	*	*	*	*	*	*	*	*	*	*
727.4	*6	*	*	*	*	*	*	*	*	*	*
727.6	14	*	*9	—	*	*7	*6	*	*	11	*
727.61	12	*	*7	—	*	*5	*5	*	*	10	*
727.8	*9	*	*	*	*	*	*	*	*	*	*
728	107	53	54	*	25	27	51	28	24	37	18
728.2	11	*	*9	*	*	*	*9	*6	*	*	*
728.3	*6	*	*	*	*	*	*5	*	*	*	*
728.8	63	34	29	*	21	17	24	11	14	25	13
728.85	13	*5	*8	*	*	*	*	*	*	*	*
728.86	*7	*	*	*	*	*	*	*	*	*	*
728.89	38	25	13	*	13	*7	18	*7	*7	17	*7
728.9	19	11	*9	*	*	*5	10	*5	*6	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
729	305	96	209	*6	72	96	132	56	76	116	57
729.1	114	20	94	*	31	52	30	25	31	35	24
729.2	*9	*	*	—	*	*	*	*	*	*	*
729.5	62	26	36	*	18	15	26	14	18	20	11
729.8	108	41	67	*	15	23	68	13	23	52	19
729.81	17	*7	9	*	*	*5	*6	*5	*	*	*
729.82	*6	*	*	*	*	*	*	*	*	*	*
729.89	85	31	54	*	9	17	58	*7	18	45	15
730	136	81	55	*6	28	46	56	38	31	45	22
730.0	42	27	16	*	9	14	15	12	10	12	*8
730.07	27	18	*9	*	*6	*9	11	*9	*8	*9	*
730.1	28	18	10	*	*	12	10	*8	*6	9	*
730.17	15	12	*	*	*	*6	*6	*	*	*6	*
730.2	64	35	29	*	13	20	30	17	15	23	9
730.25	*5	*	*	*	*	*	*	*	*	*	*
730.26	*5	*	*	*	*	*	*	*	*	*	*
730.27	38	20	17	*	*6	14	18	12	*8	12	*5
730.28	10	*6	*	*	*	*	*5	*	*	*	*
731	48	31	17	—	*5	14	29	14	15	15	*
731.0	19	11	*8	—	*	*	18	*6	*8	*	*
731.8	29	20	9	—	*	13	11	*7	*6	11	*
732	10	*5	*	*5	*	*	*	*	*	*	*
733	760	138	622	*5	54	120	581	173	202	259	126
733.0	441	37	403	*	*	52	383	109	126	138	68
733.00	416	34	382	*	*	49	362	105	119	132	60
733.01	12	*	10	—	—	*	12	*	*	*	*5
733.09	12	*	10	—	*	*	9	*	*	*	*
733.1	155	43	112	*	*5	23	126	36	40	52	26
733.10	*6	*	*	—	—	*	*	*	*	*	*
733.13	114	31	84	*	*	12	99	26	29	37	22
733.14	15	*	11	*	*	*5	10	*	*	*6	*
733.15	*6	*	*	*	*	*	*	*	*	*	*
733.19	9	*	*8	—	*	*	*8	*	*	*	*
733.2	*5	*	*	*	*	*	*	*	*	*	*
733.4	39	18	21	*	15	12	11	*8	10	17	*
733.42	31	14	17	*	12	9	9	*6	*8	14	*
733.6	30	*7	23	*	*8	*9	13	*7	*9	10	*
733.8	39	16	23	*	12	10	16	*6	*8	12	13
733.81	*6	*	*	*	*	*	*	*	*	*	*
733.82	33	12	20	*	9	*8	14	*6	*7	10	10
733.9	50	15	35	*	*8	13	27	*6	9	25	10
733.90	41	10	31	*	*	10	26	*	*8	22	*8
733.99	*7	*	*	*	*	*	*	*	*	*	*
735	11	*	*8	*	*	*	*5	*	*	*	*
736	41	18	23	*5	*6	15	15	10	9	11	11
736.4	10	*	*7	*	*	*5	*	*	*	*	*
736.42	*6	*	*	*	*	*	*	*	*	*	*
736.6	*5	*	*	—	*	*	*	*	*	*	*
736.7	16	*8	*8	*	*	*	*6	*	*	*	*
736.79	12	*6	*6	*	*	*	*	*	*	*	*
736.8	*5	*	*	*	*	*	*	*	*	*	*
737	74	16	59	12	15	*7	39	19	23	17	15
737.1	12	*	10	*	*	*	*7	*	*	*	*
737.10	11	*	10	*	*	*	*7	*	*	*	*
737.3	54	11	42	12	11	*5	25	14	17	13	10
737.30	46	10	37	10	10	*	22	13	15	11	*7
737.39	*6	*	*	*	*	*	*	*	*	*	*
737.4	*7	*	*5	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
738	45	16	29	*	13	15	17	11	12	14	*9
738.4	36	13	23	*	*8	11	16	9	10	9	*7
739	36	10	25	—	19	*7	10	32	*	*	*
739.2	12	*6	*6	—	*	*	*7	11	*	*	—
739.3	18	*	16	—	17	*	*	18	*	*	—
740-759	626	317	309	353	128	75	70	152	128	195	152
741	27	12	15	12	11	*	*	*6	*9	*7	*6
741.0	13	*	*8	*8	*	*	*	*	*	*	*
741.00	*6	*	*	*	*	*	*	*	*	*	*
741.9	14	*7	*7	*	*7	*	*	*	*	*	*
741.90	12	*6	*6	*	*7	*	*	*	*	*	*
742	34	14	20	27	*	*	*	9	*8	9	*7
742.1	*8	*	*	*7	*	*	*	*	*	*	*
742.3	13	*5	*7	11	*	*	*	*	*	*	*
744	*6	*	*	*	*	*	*	*	*	*	*
745	74	38	36	55	10	*5	*	19	12	19	25
745.1	*7	*	*	*6	*	*	—	*	*	*	*
745.2	*7	*	*	*6	*	*	*	*	*	*	*
745.4	22	11	12	17	*	*	*	*7	*	*	*7
745.5	28	14	14	18	*	*	*	*7	*6	*6	10
745.6	*6	*	*	*5	*	*	*	*	*	*	*
746	55	33	22	28	13	*8	*5	15	11	18	11
746.4	*8	*6	*	*	*	*	*	*	*	*	*
746.8	26	16	10	12	*7	*5	*	*6	*	9	*6
746.85	*9	*5	*	*	*	*	*	*	*	*5	*
746.89	11	*6	*	*8	*	*	*	*	*	*	*
747	61	34	27	35	*6	*8	11	15	13	21	12
747.0	19	12	*7	18	*	*	*	*	*	*5	*6
747.1	*5	*	*	*	*	*	*	*	*	*	*
747.2	*5	*	*	*	*	*	*	*	*	*	*
747.3	*8	*	*5	*7	*	*	*	*	*	*	*
747.6	13	*7	*6	*	*	*	*6	*	*	*6	*
747.61	*8	*	*	*	*	*	*6	*	*	*	*
747.8	*8	*6	*	*	*	*	*	*	*	*	*
747.81	*6	*	*	*	*	*	*	*	*	*	*
748	18	10	*8	15	*	*	*	*7	*	*	*
748.3	11	*7	*	10	—	*	*	*	*	*	*
749	18	11	*7	16	*	—	*	*	*	*6	*
749.0	*9	*	*	*8	*	—	*	*	*	*	*
749.00	*6	*	*	*5	*	—	*	*	*	*	*
749.2	*7	*5	*	*6	*	—	—	*	*	*	*
750	18	13	*6	15	*	*	*	*	*5	*	*
750.5	11	9	*	11	—	*	—	*	*	*	*
751	42	21	21	21	*9	*6	*6	*7	11	13	10
751.0	*8	*	*	*	*	*	*	*	*	*	*
751.2	*6	*	*	*6	*	—	—	*	*	*	*
751.5	11	*	*7	*	*	*	*	*	*	*	*
752	29	10	19	10	14	*	*	*7	*9	*7	*7
752.1	*8	—	*8	*	*5	*	*	*	*	*	*
752.11	*8	—	*8	*	*	*	*	*	*	*	*
752.3	*6	—	*6	*	*	*	*	*	*	*	*
752.5	*6	*6	—	*5	*	—	*	*	*	*	*
752.51	*6	*6	—	*5	*	—	*	*	*	*	*
753	68	33	35	22	14	14	19	17	13	26	12
753.0	12	*7	*	*	*	*	*	*	*	*6	*
753.1	29	12	17	*	*8	*8	11	*6	*	14	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
753.10	9	*	*6	*	*	*	*	*	*	*	*
753.12	11	*	*6	*	*	*	*	*	*	*	*
753.2	13	*8	*	10	*	*	*	*	*	*	*
753.29	*7	*	*	*6	*	*	—	*	*	*	*
753.3	*6	*	*	*	*	*	*	*	*	*	*
754	21	11	10	15	*	*	*	*	*	*7	*6
754.7	*5	*	*	*	*	*	*	*	*	*	*
754.8	*6	*	*	*	*	*	*	*	*	*	*
755	16	*8	*9	*7	*	*	*	*	*	*	*5
755.6	10	*	*5	*	*	*	*	*	*	*	*
755.63	*8	*	*	*	*	*	*	*	*	*	*
756	56	28	28	23	12	10	11	11	*9	21	16
756.0	9	*	*	*9	*	*	*	*	*	*	*
756.1	24	13	11	*	*7	*7	*9	*	*	11	*6
756.11	*5	*	*	—	*	*	*	*	*	*	*
756.12	14	*7	*7	—	*	*	*6	*	*	*6	*
756.5	*5	*	*	*	*	*	*	*	*	*	*
757	16	*7	9	*8	*7	*	*	*	*	11	*
757.3	14	*6	*8	*8	*6	*	*	*	*	10	*
757.32	10	*	*6	*6	*	—	—	*	*	10	*
758	40	18	22	25	*8	*6	*	10	9	*7	14
758.0	33	15	18	20	*7	*5	*	*9	*8	*	11
759	24	12	12	13	*	*	*	*9	*	*6	*
759.8	13	*7	*6	*8	*	*	*	*	*	*	*
759.89	*9	*	*	*6	*	*	*	*	*	*	*
760-779	444	261	183	440	*	*	*	75	94	122	154
760	*5	*	*	*	*	*	—	*	*	*	*
764	*5	*	*	*5	—	—	—	*	*	*	*
765	49	27	22	49	—	—	—	*7	12	16	14
765.0	10	*6	*	10	—	—	—	*	*	*	*
765.1	39	21	18	39	—	—	—	*5	11	13	10
765.17	*6	*	*	*6	—	—	—	*	*	*	*
765.18	*8	*	*	*8	—	—	—	*	*	*	*
765.19	13	*6	*6	13	—	—	—	*	*	*	*
767	*6	*	*	*6	—	—	—	*	*	*	*
769	25	17	*8	25	—	—	—	*	*7	*8	*7
770	87	51	37	87	*	*	*	17	18	27	25
770.6	13	*9	*	13	—	—	—	*	*	*6	*
770.7	18	12	*6	18	*	*	*	*	*5	*	*
770.8	41	22	19	41	—	—	*	*9	*7	11	14
771	46	30	16	46	—	*	*	*9	*8	11	18
771.7	*5	*	*	*5	—	—	—	*	*	*	*
771.8	34	23	12	34	—	—	*	*7	*6	9	12
772	*5	*	*	*5	*	—	—	*	*	*	*
773	*7	*	*	*7	*	—	—	*	*	*	*
774	82	49	33	82	—	*	*	12	15	21	34
774.2	31	20	11	31	—	—	—	*	*7	9	11
774.3	*6	*	*	*6	—	—	*	*	*	*	*
774.6	44	25	19	44	—	*	*	*7	*7	10	20
775	31	17	15	31	—	—	—	*6	*	*7	13
775.5	15	*8	*8	15	—	—	—	*	*	*	*7
775.6	*5	*	*	*5	—	—	—	*	*	*	*
776	16	11	*	16	*	—	*	*	*	*	*
776.6	*8	*5	*	*8	—	—	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
777	10	*	*5	10	—	—	—	*	*	*	*
778	15	10	*5	15	—	—	*	*	*	*	*5
778.4	*8	*5	*	*8	—	—	—	*	*	*	*
779	40	22	18	40	*	*	—	*7	*8	10	16
779.0	*6	*	*	*6	—	—	—	*	*	*	*
779.3	27	13	13	27	—	*	—	*	*	*7	12
780-799	7,289	3,227	4,063	624	1,520	1,926	3,219	1,543	1,900	2,634	1,213
780	2,104	957	1,147	209	415	501	979	497	532	739	336
780.0	99	43	56	*7	19	13	59	18	26	29	26
780.01	17	*6	11	*	*	*	12	*6	*	*	*
780.02	*7	*	*	*	*	*	*	*	*	*	*
780.03	*5	*	*	*	*	*	*	*	*	*	*
780.09	70	34	36	*	17	*8	39	11	17	23	18
780.1	13	*6	*7	*	*	*	*9	*	*	*	*
780.2	393	162	231	*	41	78	272	123	79	144	48
780.3	761	368	393	95	194	209	263	195	193	266	107
780.31	16	*9	*7	15	*	*	*	*	*	*6	*
780.39	745	359	386	80	194	208	263	192	189	260	104
780.4	110	37	73	*	11	23	74	24	35	35	16
780.5	182	101	81	10	47	74	51	40	52	60	30
780.50	*7	*	*5	*	*	*	*	*	*	*	*
780.52	24	10	14	—	*6	*6	12	*	10	*9	*
780.57	142	84	58	*8	37	61	36	34	40	45	23
780.6	285	138	147	85	62	55	82	50	69	102	64
780.7	159	60	99	*5	20	25	108	29	42	59	29
780.79	154	59	95	*5	18	23	108	28	41	58	27
780.8	*6	*	*	*	*	*	*	*	*	*	*
780.9	95	39	57	*	15	18	59	15	29	38	14
781	223	88	135	12	26	34	151	43	65	79	36
781.0	26	11	15	*	*7	*7	9	*	*6	11	*
781.2	130	48	83	*	*8	17	103	27	41	46	17
781.3	53	23	30	*6	*7	*8	32	*8	14	19	13
781.9	*8	*	*	*	*	*	*	*	*	*	*
782	198	91	107	20	41	54	83	43	57	57	41
782.0	47	21	26	*	15	19	11	*8	15	16	*7
782.1	33	17	16	11	*8	*6	*8	*7	*9	*9	*9
782.3	83	33	51	*	9	21	52	20	25	21	17
782.4	23	14	*9	*	*7	*6	*9	*5	*6	*6	*
782.5	*5	*	*	*	*	*	*	*	*	*	*
783	171	85	86	62	23	20	66	38	38	59	36
783.0	38	18	20	*	*	*7	24	*	*8	22	*5
783.2	41	19	22	*	*8	*7	21	*8	11	15	*7
783.3	12	*5	*7	*9	*	*	*	*	*	*	*
783.4	72	37	34	46	*7	*	16	23	14	16	19
784	326	139	187	17	78	91	140	49	100	121	56
784.0	148	52	96	*9	56	47	36	20	46	63	19
784.2	12	*7	*	*	*	*	*	*	*	*5	*
784.3	56	27	30	*	*	13	39	*	19	19	14
784.4	*7	*	*	*	*	*	*	*	*	*	*
784.49	*7	*	*	*	*	*	*	*	*	*	*
784.5	55	27	28	*	*6	15	33	*6	17	18	14
784.7	39	21	19	*	*	*8	24	12	10	11	*7
785	323	135	188	22	59	87	155	79	88	109	47
785.0	71	29	42	*6	21	16	28	13	20	26	13
785.1	35	10	24	*	*7	13	14	12	*	13	*
785.2	29	12	17	*8	10	*6	*6	*5	*8	11	*5
785.4	72	33	39	*	*7	21	44	17	21	26	*8

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
785.5	77	35	42	*5	*7	21	45	22	24	20	12
785.50	*9	*	*6	*	*	*	*	*	*	*	*
785.51	28	12	16	*	*	*9	18	*7	12	*5	*
785.59	41	21	20	*	*	10	22	12	10	13	*6
785.6	26	13	13	*	*7	11	*6	*8	*6	*9	*
785.9	12	*	9	—	*	*	10	*	*	*	*
786	1,362	632	730	74	247	537	504	283	310	538	230
786.0	198	92	107	55	23	43	77	32	52	78	37
786.00	*6	*	*	*	*	*	*	*	*	*	*
786.03	20	12	*8	18	*	*	*	*5	*	*	*8
786.05	56	23	33	*	*6	19	29	11	9	26	11
786.06	*9	*	*	*6	*	*	*	*	*	*	*
786.07	10	*	*5	*6	*	*	*	*	*	*7	*
786.09	91	41	49	22	14	18	37	12	32	36	11
786.1	*6	*	*	*	*	*	*	*	*	*	*
786.2	28	13	15	*9	*7	*5	*7	*	*7	*8	10
786.3	53	31	21	*	12	14	26	12	11	21	9
786.5	1,039	470	569	*	200	467	370	224	233	415	166
786.50	532	246	286	*	98	234	200	99	118	235	81
786.51	31	14	17	*	*7	12	12	*6	*7	14	*
786.52	52	18	34	*	10	19	23	10	15	17	10
786.59	424	191	232	*	86	202	135	110	93	149	71
786.6	29	17	11	*	*	*6	21	*8	*	12	*
786.8	*6	*5	*	*	*	*	*	*	*	*	*
787	698	284	415	97	151	154	296	121	200	249	128
787.0	362	133	229	64	94	83	121	56	104	136	66
787.01	224	80	144	18	66	59	81	35	63	90	37
787.02	64	17	48	*6	18	16	25	11	18	21	14
787.03	73	36	37	40	11	*8	14	10	22	26	14
787.1	*7	*	*	*	*	*	*	*	*	*	*
787.2	119	61	58	*	13	23	80	23	30	47	19
787.3	14	*	*9	*	*	*	*5	*	*	*	*
787.6	11	*	*7	*	*	*	*5	*	*	*	*
787.9	185	78	106	27	36	39	84	37	56	58	34
787.91	182	78	104	27	35	38	81	36	55	57	34
788	406	209	197	*9	61	75	261	94	118	127	67
788.0	17	*6	11	*	11	*	*	*	*	*6	*
788.1	10	*	*7	*	*	*	*5	*	*	*	*
788.2	234	148	86	*	31	46	155	63	64	71	36
788.20	189	117	73	*	26	36	126	51	53	55	30
788.29	41	28	12	*	*	*9	27	11	10	14	*6
788.3	99	29	70	*	11	17	68	18	30	29	22
788.30	70	21	48	*	*7	10	51	12	23	21	14
788.31	11	*	*8	*	*	*	*8	*	*	*	*
788.33	*9	—	*9	—	*	*	*	*	*	*	*5
788.39	*7	*	*	*	*	*	*	*	*	*	*
788.4	12	*7	*	*	*	*	*8	*	*	*	*
788.41	*8	*	*	—	*	*	*6	*	*	*	*
788.9	25	11	14	*	*	*	20	*	*7	11	*
789	549	204	345	45	183	159	162	104	140	214	91
789.0	401	129	272	41	153	99	109	70	102	164	66
789.00	147	42	105	16	51	34	46	23	29	69	26
789.01	45	15	30	*	18	12	10	*6	10	20	*8
789.02	*8	*	*5	*	*	*	*	*	*	*	*
789.03	59	19	40	12	29	11	*8	11	15	23	*9
789.04	25	*7	18	*	*9	*7	9	*7	*7	9	*
789.05	*7	*	*6	*	*	*	*	*	*	*	*
789.06	46	18	28	*	17	14	15	*7	16	13	10
789.07	18	*7	11	*	*5	*6	*	*	*	*8	*
789.09	47	17	30	*	19	11	13	10	16	16	*6

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
789.1	10	*5	*5	*	*	*5	*	*	*	*6	*
789.2	11	*6	*	*	*	*	*	*	*	*	*
789.3	17	*	13	*	*	*6	*6	*	*	*6	*
789.30	*6	*	*5	—	*	*	*	*	*	*	*
789.5	107	58	49	*	20	46	40	27	29	34	17
790	330	150	180	22	79	76	154	79	79	122	50
790.2	44	*8	36	*	28	*	11	13	*6	17	*9
790.4	13	*6	*7	*	*	*	*	*	*5	*	*
790.5	23	13	10	*	*	*5	12	*	*8	*7	*
790.6	128	59	69	*	19	33	70	33	30	48	18
790.7	78	41	36	13	15	20	29	19	18	28	12
790.9	41	21	20	*	*7	*8	25	*8	10	17	*6
790.92	24	10	14	*	*	*	16	*	*5	9	*
790.93	*8	*8	—	—	—	*	*6	*	*	*	*
790.99	*7	*	*	*	*	*	*	*	*	*	*
791	35	15	20	*	*8	9	13	9	11	11	*
791.0	15	*6	*9	*	*	*	*6	*	*6	*	*
791.9	15	*	10	*	*	*	*6	*	*	*6	*
792	76	12	64	*	50	*5	18	14	21	24	18
792.1	27	11	16	*	*	*5	18	*7	*5	10	*
792.3	47	—	47	—	47	—	—	*7	15	13	13
793	18	*8	10	*	*	*6	*8	*	*5	*6	*
793.1	*9	*	*	—	*	*	*	*	*	*	*
794	106	57	49	*	24	38	41	21	30	39	16
794.3	48	29	19	*	*6	20	21	*7	15	20	*5
794.31	30	17	13	*	*5	11	13	*5	*8	12	*
794.39	14	9	*5	—	*	*8	*6	*	*6	*6	*
794.8	50	25	24	*	17	17	13	12	11	16	10
795	32	11	21	*	19	*6	*5	10	*7	*8	*7
795.5	20	9	11	*	12	*	*	*7	*	*	*5
795.7	*7	*	*6	*	*	*	*	*	*	*	*
795.79	*7	*	*6	*	*	*	*	*	*	*	*
796	46	19	28	*	16	11	17	*8	15	17	*7
796.0	*8	*	*	*	*	*	*6	*	*	*	*
796.2	27	11	16	*	11	*8	*7	*	*9	*8	*6
797	*6	*	*	—	—	—	*6	*	*	*	*
799	281	128	153	23	37	61	159	47	80	113	40
799.0	103	46	57	14	12	23	54	17	34	37	16
799.1	16	9	*7	*	*	*	11	*	*	*7	*
799.3	74	29	44	*	*	13	58	*9	30	29	*6
799.4	34	19	15	*	*5	10	18	*9	*6	10	9
799.9	49	24	25	*8	14	11	16	10	*6	27	*6
800-999	5,770	2,945	2,825	465	1,772	1,241	2,291	1,232	1,373	2,058	1,107
800	13	*9	*	*8	*	*	*	*	*	*	*
801	37	28	9	10	17	*5	*	*7	*7	10	12
801.0	15	10	*	*	*5	*	*	*	*	*7	*
801.1	*7	*5	*	*	*	*	*	*	*	*	*
801.2	9	*7	*	*	*	*	*	*	*	*	*
802	94	69	26	*5	58	15	16	18	20	35	21
802.0	18	12	*6	*	*8	*	*5	*	*5	*6	*
802.2	21	15	*6	*	16	*	*	*5	*	*6	*6
802.25	*5	*	*	*	*	*	—	*	*	*	*
802.4	21	16	*	*	13	*	*	*	*	9	*
802.6	12	*8	*	*	*7	*	*	*	*	*6	*
802.8	14	10	*	*	*8	*	*	*	*	*5	*
803	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
805	134	63	71	*	42	21	66	21	31	50	33
805.0	30	17	13	*	13	*5	9	*	*	10	11
805.01	*6	*	*	*	*	*	*	*	*	*	*
805.02	9	*	*6	*	*	*	*	*	*	*	*
805.2	34	16	18	*	*8	*5	20	*	*9	12	*9
805.4	57	23	34	*	16	*8	32	10	15	22	9
805.6	10	*5	*5	*	*	*	*	*	*	*	*
806	*6	*	*	—	*	*	*	*	*	*	*
807	131	74	57	*	33	27	66	23	33	51	23
807.0	115	63	52	*	29	24	59	20	29	45	21
807.00	17	*8	*8	*	*	*	13	*	*7	*	*
807.01	28	17	11	*	*7	*6	13	*6	*6	10	*5
807.02	16	*8	*8	*	*	*	9	*	*6	*5	*
807.03	13	*8	*5	*	*	*	*6	*	*	*5	*
807.04	10	*	*	*	*	*	*6	*	*	*	*
807.05	*8	*	*5	*	*	*	*	*	*	*	*
807.09	17	11	*6	*	*	*6	*6	*	*	*6	*
807.2	*7	*	*	—	*	*	*	*	*	*	*
808	94	40	53	*	20	11	59	17	23	38	16
808.0	20	15	*5	*	*7	*	*8	*	*	*7	*
808.2	57	17	40	*	*7	*5	43	*9	14	25	10
808.4	10	*7	*	*	*	*	*5	*	*	*	*
808.8	*6	*	*	*	*	*	*	*	*	*	*
810	28	17	12	*	*7	*	16	*6	*7	*6	*9
810.0	27	16	12	*	*6	*	16	*6	*7	*6	*8
810.00	18	10	*8	*	*	*	10	*5	*	*	*
810.02	*6	*	*	*	*	*	*	*	*	*	*
811	*9	*6	*	*	*5	*	*	*	*	*	*
811.0	*9	*6	*	*	*5	*	*	*	*	*	*
812	95	34	61	14	18	12	52	18	24	35	17
812.0	50	15	35	*	*	*8	37	10	14	18	*8
812.00	18	*5	12	*	*	*	12	*	*	*9	*
812.01	18	*6	12	*	*	*	16	*	*7	*	*
812.03	*6	*	*	—	*	*	*	*	*	*	*
812.09	*8	*	*6	*	*	*	*6	*	*	*	*
812.2	18	*	13	*	*6	*	*9	*	*	*9	*
812.20	*6	*	*	*	*	*	*	*	*	*	*
812.21	12	*	*9	*	*	*	*5	*	*	*	*
812.4	22	11	11	12	*	*	*5	*5	*	*7	*5
812.41	12	*6	*6	*9	*	*	*	*	*	*	*
813	102	47	54	17	32	22	31	20	22	37	22
813.0	23	14	9	*	*8	*	*7	*6	*	*8	*6
813.01	12	*8	*	*	*	*	*	*	*	*	*
813.2	9	*6	*	*	*	*	*	*	*	*	*
813.4	50	18	31	*6	15	11	18	*8	14	19	9
813.41	11	*	*8	*	*	*	*6	*	*	*	*
813.42	18	*8	11	*	*7	*	*5	*	*5	*7	*
813.44	17	*7	11	*	*7	*	*	*	*5	*6	*
813.5	*6	*	*	*	*	*	*	*	*	*	*
813.8	*7	*	*	*	*	*	*	*	*	*	*
814	*6	*	*	*	*	*	*	*	*	*	*
814.0	*6	*	*	*	*	*	*	*	*	*	*
815	16	*8	*8	*	*8	*	*	*	*	*8	*
815.0	12	*	*8	*	*5	*	*	*	*	*6	*
816	24	17	*6	*	11	*	*7	*6	*6	*8	*
816.0	12	*7	*5	*	*	*	*	*	*	*	*
816.01	*6	*	*	*	*	*	*	*	*	*	*
816.1	12	10	*	*	*6	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
816.11	*7	*6	*	*	*	*	*	*	*	*	*
820	371	96	275	*	*8	28	333	71	85	150	64
820.0	99	22	77	*	*	*8	88	18	25	35	21
820.00	12	*	9	—	*	*	11	*	*	*	*
820.02	16	*	12	—	*	*	12	*	*	10	*
820.03	*8	*	*5	*	*	*	*7	*	*	*	*
820.09	62	11	51	*	*	*	57	12	18	18	14
820.2	181	51	131	*	*	11	166	38	42	73	29
820.20	13	*	*9	—	*	*	12	*5	*	*	*
820.21	148	42	106	*	*	*9	137	28	33	64	22
820.22	21	*	16	*	*	*	17	*	*7	*	*
820.8	87	22	65	*	*	*9	76	15	18	41	13
821	73	35	38	13	21	11	28	11	16	27	18
821.0	39	23	15	10	13	*6	10	*6	9	13	11
821.00	9	*6	*	*	*	*	*	*	*	*	*
821.01	30	18	12	*8	11	*	*6	*	*7	10	*9
821.2	26	*7	19	*	*	*	17	*	*	10	*6
821.20	*7	*	*5	*	*	*	*	*	*	*	*
821.23	11	*	*9	*	*	*	*8	*	*	*	*
822	24	12	12	*	*6	*7	10	*6	*6	10	*
822.0	21	9	12	*	*	*7	10	*5	*5	*7	*
823	97	54	44	*6	39	30	22	19	22	38	18
823.0	39	19	20	*	13	10	14	*7	11	15	*7
823.00	29	15	14	*	10	*8	10	*5	*8	11	*
823.02	*6	*	*	*	*	*	*	*	*	*	*
823.2	24	13	11	*	10	*9	*	*	*	*8	*8
823.20	*5	*	*	*	*	*	*	*	*	*	*
823.22	17	10	*8	*	*7	*6	*	*	*	*6	*6
823.3	*6	*5	*	*	*	*	*	*	*	*	*
823.8	19	10	9	*	*8	*7	*	*6	*	*8	*
823.80	*6	*	*	*	*	*	*	*	*	*	*
823.82	11	*5	*6	*	*	*	*	*	*	*	*
824	136	59	78	*6	58	39	34	29	31	53	24
824.0	9	*	*	*	*	*	*	*	*	*	*
824.2	12	*7	*5	*	*6	*	*	*	*	*	*
824.4	36	14	23	*	14	13	*9	*6	9	15	*6
824.6	34	11	23	*	12	*9	12	10	*6	14	*
824.8	36	18	18	*	18	*7	*8	*7	*7	13	9
825	36	23	13	*	17	12	*	*5	*8	15	*8
825.0	13	9	*	*	*	*6	*	*	*	*6	*
825.2	20	11	9	*	11	*5	*	*	*	*8	*
825.21	*5	*	*	*	*	*	*	*	*	*	*
825.25	12	*6	*6	*	*5	*	*	*	*	*	*
826	*8	*	*	*	*	*	*	*	*	*	*
831	9	*5	*	*	*	*	*	*	*	*	*
831.0	9	*5	*	*	*	*	*	*	*	*	*
836	21	11	10	*	14	*	*	*	*	10	*5
836.0	10	*	*	*	*	*	*	*	*	*	*
836.1	*8	*5	*	—	*7	*	*	*	*	*	*
839	*7	*	*	*	*	*	*	*	*	*	*
840	36	18	18	*	*5	10	21	*6	13	13	*
840.4	27	13	15	—	*	*8	17	*5	*9	10	*
840.9	*5	*	*	*	*	*	*	*	*	*	*
842	*5	*	*	*	*	*	*	*	*	*	*
843	*5	*	*	*	*	*	*	*	*	*	*
844	33	20	12	*	21	*6	*6	*6	*	12	10
844.2	13	*8	*	*	12	*	*	*	*	*5	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
844.8	*9	*7	*	*	*	*	*	*	*	*	*
844.9	*7	*	*	*	*	*	*	*	*	*	*
845	21	10	11	*	*8	*	*7	*	*	*7	*
845.0	20	10	10	*	*8	*	*7	*	*	*7	*
845.00	14	*6	*7	*	*6	*	*5	*	*	*	*
846	10	*	*7	*	*	*	*	*	*	*	*
846.0	*8	*	*6	*	*	*	*	*	*	*	*
847	34	15	19	*	14	*	16	*5	*8	12	*8
847.0	14	*6	*8	*	*7	*	*6	*	*	*5	*
847.2	14	*7	*7	—	*5	*	*6	*	*	*5	*
848	*8	*	*6	—	*	*	*	*	*	*	*
848.8	*5	*	*	—	*	*	*	*	*	*	*
850	58	36	22	11	25	*9	14	19	11	16	12
850.0	12	*6	*6	*	*	*	*5	*	*	*	*
850.1	21	14	*7	*7	9	*	*	*7	*	*	*6
850.5	12	*9	*	*	*7	*	*	*	*	*	*
850.9	11	*6	*6	*	*	*	*	*	*	*	*
851	23	15	*7	*	10	*	*7	*	*5	*8	*6
851.8	16	10	*6	*	*8	*	*6	*	*	*7	*
852	43	26	16	*5	10	*8	19	*9	10	13	11
852.0	11	*6	*	*	*	*	*	*	*	*	*
852.2	30	18	12	*	*6	*5	16	*6	*6	9	*8
852.20	9	*5	*	*	*	*	*6	*	*	*	*
852.21	*9	*6	*	*	*	*	*	*	*	*	*
853	11	*5	*6	*	*	*	*5	*	*	*	*
853.0	11	*5	*6	*	*	*	*5	*	*	*	*
853.00	*6	*	*	*	*	*	*	*	*	*	*
854	16	11	*5	*	*7	*	*	*	*	*6	*
854.0	15	11	*	*	*7	*	*	*	*	*5	*
854.02	*6	*	*	*	*	*	*	*	*	*	*
860	52	36	16	*	29	10	11	*8	11	19	14
860.0	30	19	11	*	16	*6	*6	*	*7	10	*8
860.4	11	*8	*	*	*6	*	*	*	*	*	*
861	36	23	13	*	15	*8	12	*7	9	14	*6
861.0	*6	*	*	*	*	*	*	*	*	*	*
861.2	28	19	9	*	12	*7	*8	*5	*8	10	*5
861.21	27	19	*9	*	12	*7	*8	*	*8	10	*5
862	10	*7	*	*	*	*	*	*	*	*	*
863	20	13	*6	*	11	*	*	*	*6	*6	*
864	15	11	*	*	*8	*	*	*	*	*	*
864.0	12	*9	*	*	*6	*	*	*	*	*	*
865	27	19	*8	*	15	*	*	*6	*6	10	*5
865.0	25	18	*8	*	14	*	*	*6	*6	*9	*5
865.00	*6	*	*	*	*	*	*	*	*	*	*
865.04	*6	*	*	*	*	*	*	*	*	*	*
866	12	9	*	*	*5	*	*	*	*	*	*
866.0	11	*9	*	*	*	*	*	*	*	*	*
866.01	*5	*	*	*	*	*	*	*	*	*	*
867	15	14	*	*	*	*	*8	*	*	*	*
867.0	13	12	*	*	*	*	*8	*	*	*	*
868	19	12	*7	*	9	*	*6	*	*	*8	*
868.0	14	*8	*6	*	*	*	*6	*	*	*5	*
870	14	11	*	*	9	*	*	*	*	*	*
870.8	*5	*	*	*	*	*	*	*	*	*	*
871	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
872	10	*9	*	*	*7	*	*	*	*	*	*
872.0	*6	*5	*	*	*	*	*	*	*	*	*
873	152	96	57	12	65	26	50	35	38	45	34
873.0	43	26	17	*	15	*6	20	*7	12	14	10
873.2	*7	*	*	*	*	*	*	*	*	*	*
873.20	*6	*	*	*	*	*	*	*	*	*	*
873.4	82	51	31	*8	34	14	26	22	19	22	18
873.40	*8	*	*	*	*	—	*	*	*	*	*
873.41	*6	*	*	*	*	*	*	*	*	*	*
873.42	40	24	16	*	15	*8	16	*9	11	11	10
873.43	15	10	*5	*	*6	*	*	*5	*	*	*
873.44	*8	*6	*	*	*	*	*	*	*	*	*
873.49	*5	*	*	*	*	*	*	*	*	*	*
873.6	10	*8	*	*	*5	*	*	*	*	*	*
873.63	*6	*	*	*	*	*	*	*	*	*	*
874	*6	*	*	*	*	*	*	*	*	*	*
875	*8	*7	*	*	*7	*	*	*	*	*	*
875.0	*7	*6	*	*	*6	*	*	*	*	*	*
879	12	*8	*	*	*8	*	*	*	*	*5	*
880	10	*7	*	*	*7	*	*	*	*	*6	*
880.0	*8	*6	*	*	*6	*	*	*	*	*	*
881	38	22	16	*	25	*	*7	*6	*9	13	10
881.0	30	16	14	*	20	*	*6	*5	*7	11	*7
881.00	10	*7	*	*	*7	*	*	*	*	*	*
881.01	*6	*	*	*	*	*	*	*	*	*	*
881.02	13	*6	*7	*	10	*	*	*	*	*6	*
881.2	*5	*	*	*	*	*	*	*	*	*	*
882	20	13	*7	*	*8	*7	*	*6	*7	*	*
882.0	11	*6	*	*	*	*	*	*	*	*	*
882.1	*8	*5	*	*	*	*	*	*	*	*	*
883	24	17	*7	*	13	*5	*	*	9	*	*5
883.0	*8	*	*	*	*	*	*	*	*	*	*
883.1	*6	*	*	*	*	*	*	*	*	*	*
883.2	10	*9	*	*	*6	*	*	*	*	*	*
886	*6	*5	*	*	*	*	*	*	*	*	*
886.0	*5	*	*	*	*	*	*	*	*	*	*
890	10	*7	*	*	*6	*	*	*	*	*	*
890.0	*7	*6	*	*	*	*	*	*	*	*	*
891	31	22	9	*	16	*6	*5	*5	*7	14	*
891.0	17	12	*6	*	9	*	*	*	*	*7	*
891.1	11	*8	*	*	*5	*	*	*	*	*5	*
892	11	*7	*	*	*6	*	*	*	*	*	*
892.1	*7	*	*	*	*	*	*	*	*	*	*
903	11	*9	*	*	*7	*	*	*	*	*5	*
904	*5	*	*	*	*	*	*	*	*	*	*
905	56	25	32	*	16	11	28	14	19	*9	14
905.1	*9	*	*5	*	*	*	*	*	*	*	*
905.2	*7	*	*	*	*	*	*	*	*	*	*
905.3	14	*	10	*	*	*	13	*	*8	*	*
905.4	17	*8	*9	*	*	*6	*5	*	*	*	*6
906	44	23	20	23	15	*	*	*	*	36	*
906.9	21	11	10	15	*6	—	*	*	—	21	—
907	46	28	18	*	23	17	*	*7	13	13	13
907.0	27	16	11	*	13	*9	*	*5	*8	*6	*8
907.2	18	11	*7	*	10	*7	*	*	*	*6	*5

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
908	20	14	*6	*	*9	*7	*	*	*	*7	*
908.9	15	11	*	*	*6	*6	*	*	*	*6	*
909	*8	*	*	*	*	*	*	*	*	*	*
910	30	20	10	*8	*9	*6	*7	*8	*8	*7	*7
910.0	27	18	*9	*7	*8	*5	*7	*7	*8	*6	*6
911	12	10	*	*	*7	*	*	*	*	*	*
911.0	11	10	*	*	*6	*	*	*	*	*	*
912	*7	*6	*	*	*	*	*	*	*	*	*
912.0	*6	*6	*	*	*	*	*	*	*	*	*
913	16	*8	*8	*	*7	*	*	*	*	*	*
913.0	13	*7	*6	*	*5	*	*	*	*	*	*
914	*7	*	*	*	*	*	*	*	*	*	*
914.0	*5	*	*	*	*	*	*	*	*	*	*
916	24	14	10	*	*6	*	9	*6	*5	*6	*7
916.0	18	11	*7	*	*6	*	*7	*	*	*	*6
917	*5	*	*	*	*	*	*	*	*	*	*
919	18	11	*7	*	*8	*	*	*	*5	*7	*
919.0	16	10	*6	*	*7	*	*	*	*5	*5	*
920	60	22	38	*8	13	*6	32	13	14	18	15
921	13	*7	*7	*	*7	*	*	*	*	*	*
922	65	31	34	*	22	*9	32	12	18	22	14
922.1	31	15	15	*	11	*	15	*7	*7	11	*
922.2	17	*8	9	*	*6	*	*7	*	*6	*5	*
922.3	13	*	*8	*	*	*	*9	*	*	*	*
922.31	*7	*	*	*	*	*	*	*	*	*	*
922.32	*6	*	*	*	*	*	*	*	*	*	*
923	28	11	17	*	9	*	16	*6	11	*7	*
923.0	13	*7	*6	*	*5	*	*7	*	*6	*	*
923.00	11	*7	*	*	*	*	*5	*	*	*	*
923.1	*7	*	*	*	*	*	*	*	*	*	*
923.2	*5	*	*	*	*	*	*	*	*	*	*
924	85	32	54	*	22	14	49	17	23	31	14
924.0	25	*8	17	*	*	*	17	*6	*7	*7	*6
924.00	*6	*	*	*	*	*	*	*	*	*	*
924.01	19	*6	13	*	*	*	13	*	*	*	*5
924.1	21	*6	15	*	*6	*	10	*	*5	*8	*
924.10	*6	*	*	*	*	*	*	*	*	*	*
924.11	15	*	11	*	*	*	*8	*	*	*6	*
924.8	29	13	16	*	10	*	14	*	*8	14	*
933	14	*8	*6	*	*	*	*8	*	*	*5	*
933.1	11	*6	*	*	*	*	*7	*	*	*	*
934	*8	*	*	*	*	*	*6	*	*	*	*
935	10	*6	*	*	*	*	*	*	*	*	*
935.1	*8	*5	*	*	*	*	*	*	*	*	*
941	18	14	*	*5	*6	*	*	*	*8	*7	*
941.2	11	10	*	*	*	*	*	*	*6	*	*
941.29	*5	*	*	*	*	*	*	*	*	*	*
942	22	15	*7	*9	*8	*	*	*	*6	*8	*
942.2	14	11	*	*6	*6	*	*	*	*	*6	*
942.3	*6	*	*	*	*	*	*	*	*	*	*
943	22	17	*5	*6	12	*	*	*	*7	10	*
943.2	14	11	*	*	*8	*	*	*	*	*7	*
943.3	*5	*	*	*	*	*	*	*	*	*	*
944	20	16	*	*6	*7	*5	*	*	*7	*9	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
944.2	11	9	*	*	*	*	*	*	*	*	*
944.3	*7	*6	*	*	*	*	*	*	*	*	*
945	27	20	*6	*7	9	*6	*	*5	*6	13	*
945.2	14	10	*	*	*	*	*	*	*	*6	*
945.22	*5	*	*	*	*	*	*	*	*	*	*
945.3	9	*7	*	*	*	*	*	*	*	*5	*
948	48	34	13	21	17	*6	*	*9	*7	29	*
948.0	25	19	*6	*8	11	*	*	*6	*	13	*
948.00	25	19	*6	*8	11	*	*	*6	*	13	*
948.1	10	*9	*	*	*	*	*	*	*	*6	*
948.10	*8	*7	*	*	*	*	*	*	*	*5	*
948.2	*5	*	*	*	*	*	*	*	*	*	*
952	*7	*6	*	*	*	*	*	*	*	*	*
955	12	9	*	*	*6	*	*	*	*	*	*
958	34	23	11	*	17	*8	*5	*7	*8	14	*5
958.3	9	*7	*	*	*	*	*	*	*	*	*
958.7	*6	*	*	*	*	*	*	*	*	*	*
958.8	11	*7	*	*	*6	*	*	*	*	*	*
959	82	40	41	*6	35	11	30	16	24	26	16
959.0	47	25	22	*	19	*5	19	*7	12	16	12
959.01	43	22	21	*	17	*	18	*7	11	15	10
959.1	11	*6	*5	*	*	*	*	*	*	*	*
959.7	*7	*	*	*	*	*	*	*	*	*	*
962	10	*	*6	*	*	*	*	*	*	*6	*
962.3	*9	*	*5	*	*	*	*	*	*	*5	*
963	12	*	*8	*	10	*	*	*	*	*	*
963.0	10	*	*8	*	*8	*	*	*	*	*	*
964	*7	*	*	*	*	*	*	*	*	*	*
964.2	*6	*	*	*	*	*	*	*	*	*	*
965	84	26	58	*	54	19	*6	14	24	28	17
965.0	21	*9	12	*	12	*7	*	*	*	*8	*
965.09	11	*	*7	*	*6	*	*	*	*	*	*
965.1	14	*	9	*	*7	*5	*	*	*	*	*
965.4	31	*7	24	*	23	*	*	*5	11	9	*6
965.6	9	*	*7	*	*7	*	*	*	*	*	*
965.61	*8	*	*6	*	*6	*	*	*	*	*	*
965.8	*6	*	*	*	*	*	*	*	*	*	*
966	16	*5	11	*	*8	*	*	*	*	*	*
966.1	*7	*	*5	*	*	*	*	*	*	*	*
966.3	9	*	*6	*	*7	*	*	*	*	*	*
967	14	*	11	*	*7	*	*	*	*	*	*
967.8	*6	*	*	*	*	*	*	*	*	*	*
968	14	*8	*6	*	10	*	*	*	*	*5	*
968.5	*9	*6	*	*	*7	*	*	*	*	*	*
969	87	28	60	*	57	22	*	16	27	31	14
969.0	33	11	22	*	22	*7	*	*6	12	11	*
969.4	33	10	23	*	19	11	*	*6	9	13	*5
969.7	*7	*	*	*	*	*	*	*	*	*	*
972	21	*8	13	*	*	*	11	*	*	*8	*
972.1	10	*	*7	—	*	*	*8	*	*	*	*
972.6	*6	*	*	*	*	*	*	*	*	*	*
977	16	*6	10	*	*9	*	*	*	*	*7	*
977.8	*7	*	*	*	*	*	*	*	*	*	*
977.9	9	*	*6	*	*	*	*	*	*	*	*
980	15	*8	*6	*	11	*	*	*	*6	*5	*
980.0	11	*5	*6	*	*8	*	*	*	*5	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
987	*7	*	*	*	*	*	*	*	*	*	*
989	13	*7	*6	*	*	*	*	*	*	*6	*
989.5	*7	*	*	*	*	*	*	*	*	*	*
990	*5	*	*	*	*	*	*	*	*	*	*
991	11	*7	*	*	*5	*	*	*	*	*	*
991.6	*8	*	*	*	*	*	*	*	*	*	*
994	*8	*	*	*	*	*	*	*	*	*	*
995	112	41	70	10	26	26	50	32	24	45	11
995.0	*8	*	*	*	*	*	*	*	*	*	*
995.1	12	*	*8	*	*	*	*	*	*	*	*
995.2	68	27	41	*	11	17	39	22	12	27	*7
995.3	*5	*	*	*	*	*	*	*	*	*	*
995.5	10	*	*6	*8	*	*	—	*	*	*7	*
995.8	*7	*	*6	—	*	*	*	*	*	*	*
996	653	351	302	50	115	194	294	155	139	236	124
996.0	30	16	15	*	*	*7	21	10	*7	9	*
996.01	19	9	10	*	*	*	15	*6	*	*6	*
996.04	*6	*	*	—	*	*	*	*	*	*	*
996.1	22	12	10	*	*	*7	9	*7	*	*7	*
996.2	29	15	14	17	*7	*	*	*6	10	*9	*5
996.3	10	*8	*	*	*	*	*6	*	*	*	*
996.39	*6	*	*	*	*	*	*	*	*	*	*
996.4	99	46	53	*	12	34	53	23	26	33	18
996.5	20	11	9	*	*	*6	*8	*	*	*9	*
996.59	14	*7	*7	*	*	*	*6	*	*	*6	*
996.6	206	117	89	21	38	58	89	43	45	78	40
996.61	*6	*	*	*	*	*	*	*	*	*	*
996.62	110	60	50	15	22	35	38	23	25	44	18
996.63	*6	*	*	*	*	*	*	*	*	*	*
996.64	26	16	9	*	*	*	19	*6	*5	*8	*7
996.66	20	12	*8	—	*	*5	14	*	*	*8	*
996.67	14	*8	*6	*	*	*	*6	*	*	*7	*
996.68	*7	*	*	*	*	*	*	*	*	*	*
996.69	10	*	*6	*	*	*	*	*	*	*	*
996.7	193	103	91	*	30	58	102	46	40	68	40
996.72	14	10	*	*	*	*	*8	*5	*	*	*
996.73	58	25	32	*	*9	19	31	11	*9	23	15
996.74	60	32	28	*	*8	19	32	17	12	22	9
996.76	12	10	*	*	*	*	*7	*	*	*	*
996.77	13	*7	*6	*	*	*	*7	*	*	*5	*
996.78	16	*8	*8	*	*	*6	*6	*	*	*5	*
996.79	11	*6	*5	*	*	*	*6	*	*	*	*
996.8	42	24	19	*	19	17	*	12	*	22	*
996.81	25	13	12	*	12	11	*	*6	*	13	*
996.82	*5	*	*	*	*	*	*	*	*	*	*
997	552	270	282	17	88	153	294	140	128	167	117
997.0	29	14	15	*	*	*8	16	*7	*6	*6	10
997.02	14	*7	*7	*	*	*	11	*	*	*	*
997.09	12	*	*8	*	*	*5	*	*	*	*	*
997.1	120	65	55	*	*8	30	79	38	26	36	20
997.2	17	*7	10	*	*	*	11	*5	*	*6	*
997.3	105	47	58	*6	16	31	53	26	20	32	26
997.4	172	76	95	*	44	47	76	36	46	52	38
997.5	64	35	29	*	11	18	34	17	14	20	14
997.6	33	21	12	*	*	11	19	9	*9	11	*
997.62	22	13	*9	*	*	*7	13	*7	*5	*7	*
997.69	*8	*	*	—	*	*	*	*	*	*	*
997.9	12	*	*8	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
997.99	*8	*	*	*	*	*	*5	*	*	*	*
998	632	296	336	23	144	185	280	157	139	219	117
998.1	207	104	103	*6	42	56	103	54	45	66	42
998.11	109	61	48	*	25	29	50	25	24	37	23
998.12	89	41	49	*	15	24	49	26	19	28	17
998.13	9	*	*6	*	*	*	*	*	*	*	*
998.2	77	28	49	*	20	24	32	17	16	26	19
998.3	49	25	24	*	10	14	23	11	11	22	*5
998.5	177	83	94	10	42	53	72	42	43	61	31
998.59	172	80	92	10	40	51	71	41	41	59	31
998.6	16	*9	*7	*	*	*7	*6	*	*	*6	*
998.8	92	41	51	*	25	27	37	22	19	34	17
998.83	13	*7	*6	*	*	*	*5	*	*	*	*
998.89	78	34	44	*	22	22	32	19	16	29	14
998.9	9	*	*7	*	*	*	*	*6	*	*	*
999	42	19	23	*	11	11	15	12	*7	15	*7
999.2	12	*	*7	*	*	*	*	*	*	*6	*
999.3	11	*	*7	*	*	*	*5	*	*	*	*
999.8	12	*5	*7	*	*	*	*	*	*	*	*
999.9	*5	*	*	*	*	*	*	*	*	*	*
V01-V82	11,553	3,360	8,192	302	5,453	1,774	4,024	2,345	2,674	4,221	2,313
V02	281	11	270	*	269	*8	*	58	58	107	58
V02.5	262	*	260	*	256	*	*	50	54	105	54
V02.51	253	—	253	*	252	*	—	48	52	101	53
V02.59	*9	*	*6	*	*	*	*	*	*	*	*
V02.6	18	*8	9	—	11	*5	*	*8	*	*	*
V02.61	*8	*	*5	—	*6	*	*	*	*	*	*
V02.62	10	*6	*	—	*	*	*	*	*	*	*
V04	13	*	13	—	13	*	*	*	*	*6	*
V04.3	12	—	12	—	12	—	—	*	*	*6	*
V05	*5	*	*	*	*	*	*	*	*	*	*
V07	44	*	42	*	33	*	*7	*9	14	9	12
V07.2	32	*	31	*	30	*	*	*8	9	*7	*8
V07.4	*8	—	*8	—	*	*	*5	*	*	*	*
V08	43	27	16	*	29	13	*	23	*	14	*
V09	58	25	33	*	10	14	30	11	12	28	*7
V09.0	51	22	29	*	*8	13	27	10	11	24	*6
V10	967	394	572	*8	64	227	668	262	241	306	158
V10.0	213	107	106	*	10	46	156	62	58	61	33
V10.02	11	*7	*	*	*	*	*6	*	*	*	*
V10.03	*5	*	*	—	*	*	*	*	*	*	*
V10.04	*9	*	*6	—	*	*	*7	*	*	*	*
V10.05	149	72	77	*	*6	25	118	43	39	42	24
V10.06	20	*8	12	*	*	*7	12	*7	*	*7	*
V10.09	*9	*	*	*	*	*	*6	*	*	*	*
V10.1	63	32	31	*	*	16	43	20	15	20	*8
V10.11	62	31	31	*	*	16	43	20	15	20	*8
V10.2	19	15	*	—	*	*	13	*	*	10	*
V10.21	16	12	*	—	*	*	12	*	*	*8	*
V10.3	256	*	255	*	18	62	176	81	58	78	38
V10.4	218	129	90	—	13	46	159	50	56	67	45
V10.41	36	—	36	—	*6	15	15	*6	10	12	*7
V10.42	29	—	29	—	*	*8	18	*7	*6	10	*6
V10.43	16	—	16	—	*	*6	*7	*	*	*	*
V10.44	*6	—	*6	—	*	*	*	*	*	*	*
V10.46	122	122	—	—	*	13	109	28	33	34	27
V10.5	72	43	29	*	*	15	54	19	19	26	*9

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V10.51	40	26	13	*	*	*7	31	9	12	14	*
V10.52	32	16	16	*	*	*8	23	9	*7	12	*
V10.7	25	14	11	*	*	*7	14	*5	9	*8	*
V10.79	21	11	10	*	*	*6	13	*	*7	*7	*
V10.8	92	50	42	*	11	29	50	20	21	34	17
V10.82	20	11	9	*	*	9	*9	*	*5	*7	*
V10.83	41	27	14	—	*	10	29	*8	*9	18	*6
V10.85	*7	*	*	*	*	*	*	*	*	*	*
V10.87	13	*	11	—	*	*	*7	*	*	*	*
V10.9	*6	*	*	*	*	*	*	*	*	*	*
V11	37	22	15	*	13	12	11	*	13	11	*8
V11.3	16	12	*	—	*	*8	*	*	*5	*	*
V11.8	13	*6	*6	*	*6	*	*	*	*6	*	*
V12	773	321	451	*6	80	180	507	164	222	257	129
V12.0	41	19	22	*	12	*9	20	*7	11	13	11
V12.01	16	*9	*8	*	*	*	10	*	*	*5	*
V12.09	18	*7	11	*	*9	*	*7	*	*	*6	*5
V12.2	19	*6	13	*	*	*	15	*	*	*6	*6
V12.4	25	*8	18	*	*6	*9	10	*5	*8	*8	*
V12.49	19	*6	13	*	*	*	9	*	*7	*7	*
V12.5	518	214	304	*	37	112	367	114	152	174	78
V12.50	40	15	25	*	*	12	27	*9	10	19	*
V12.51	102	33	69	*	19	28	55	18	36	29	20
V12.52	*7	*	*	—	*	*	*	*	*	*	*
V12.59	369	162	207	*	15	71	281	85	105	125	54
V12.6	23	11	13	*	*	*6	12	*	*6	*9	*
V12.7	141	62	79	*	19	42	80	28	39	46	27
V12.70	*7	*	*	*	*	*	*	*	*	*	—
V12.71	75	35	40	*	10	21	44	9	22	28	16
V12.72	22	11	11	—	*	*5	15	*	*7	*7	*
V12.79	37	13	24	*	*8	11	17	11	9	*9	*8
V13	74	35	39	17	20	15	21	12	21	18	23
V13.0	34	19	15	*	10	11	11	*6	*8	11	*8
V13.01	21	14	*7	*	*7	*8	*6	*	*	*6	*6
V13.09	10	*	*6	*	*	*	*	*	*	*	*
V13.2	*8	*	*7	—	*6	*	*	*	*	*	*
V13.5	*7	*	*	—	*	*	*	*	*	*	*
V13.7	13	*8	*	13	—	—	—	—	*5	*	*7
V13.8	*5	*	*	*	*	*	*	*	*	*	*
V14	54	14	40	*	19	14	19	10	*7	24	12
V14.0	28	*8	20	*	10	*8	10	*8	*	11	*5
V14.2	*8	*	*6	*	*	*	*	*	*	*	*
V14.5	*7	*	*5	—	*	*	*	*	*	*	*
V15	636	355	281	21	143	217	255	112	158	201	164
V15.0	*6	*	*	*	*	*	*	*	*	*	*
V15.1	16	*8	*9	10	*	*	*	*	*	*	*
V15.3	25	16	*8	*	*	*7	12	*	*	*8	11
V15.4	11	*	*6	*	*7	*	—	*	*	*5	*
V15.41	*9	*	*6	*	*6	*	—	*	*	*	*
V15.5	21	12	*9	*	10	*5	*	*	*6	*8	*
V15.8	552	309	243	*5	115	199	233	97	138	174	143
V15.81	158	85	72	*	68	56	30	31	47	55	25
V15.82	372	211	161	—	42	139	192	62	84	113	114
V15.84	*6	*5	*	—	*	*	*	*	*	*	*
V15.89	16	*7	9	*	*5	*	*6	*	*6	*	*
V16	29	*7	22	*	*6	16	*6	*	*6	*6	12
V16.0	9	*	*6	—	*	*5	*	*	*	*	*
V16.3	*6	*	*6	*	*	*	*	*	*	*	*
V16.4	*7	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V16.41	*6	—	*6	*	*	*	*	*	*	*	*
V17	116	68	49	*	30	58	26	20	25	42	29
V17.3	85	53	32	*	20	46	19	17	20	31	18
V17.4	22	10	11	*	*8	10	*	*	*	*8	*8
V18	13	*	*9	*	*8	*	*	*	*	*	*7
V18.0	10	*	*6	*	*6	*	*	*	*	*	*5
V22	25	—	25	*	24	*	—	*	*	13	*6
V22.2	18	—	18	*	18	*	—	*	*	10	*
V23	135	—	135	*	132	*	—	13	20	71	30
V23.2	*6	—	*6	—	*6	—	—	*	*	*	*
V23.4	12	—	12	—	12	—	—	*	*	*	*
V23.7	84	—	84	*	82	*	—	*5	13	48	17
V23.8	24	—	24	*	23	*	—	*	*	13	*5
V23.82	*7	—	*7	—	*6	*	—	*	*	*	*
V23.89	14	—	14	—	14	—	—	*	*	*9	*
V25	306	*	305	*	305	*	—	41	53	154	58
V25.0	9	—	9	*	9	—	—	—	*	*8	*
V25.2	294	*	294	—	293	*	—	41	53	144	56
V26	10	*	10	—	*9	*	*	*	*	*6	*
V26.5	*6	*	*6	—	*	*	*	*	*	*	*
V26.51	*6	—	*6	—	*	*	*	*	*	*	*
V27	3,738	—	3,738	12	3,722	*	—	632	772	1,439	895
V27.0	3,499	—	3,499	12	3,484	*	—	581	686	1,369	863
V27.1	18	—	18	*	18	—	—	*	*	*9	*
V27.2	54	—	54	—	54	*	—	*8	11	22	12
V27.9	163	—	163	*	163	*	—	40	70	39	15
V29	*7	*	*	*7	—	—	—	*	*	*	*
V29.0	*7	*	*	*7	—	—	—	*	*	*	*
V30	*9	*	*	*9	—	—	*	*	*	*	*
V30.1	*9	*	*	*9	—	—	*	*	*	*	*
V42	110	62	48	20	29	35	26	33	19	40	17
V42.0	39	22	17	*	14	18	*	*8	*8	18	*5
V42.1	*7	*	*	*	*	*	*	*	*	*	*
V42.2	19	*7	11	*	*	*	15	*5	*	*	*6
V42.7	14	*7	*6	*	*	*6	*	*	*	*7	*
V42.8	26	18	*8	15	*8	*	*	12	*	*6	*
V42.81	10	*7	*	*6	*	*	*	*	*	*	*
V42.82	*7	*	*	*	*	*	*	*6	*	*	—
V42.83	*8	*6	*	*	*	*	*	*	*	*	*
V43	428	160	268	*	18	87	323	98	125	138	67
V43.1	*5	*	*	*	*	*	*5	*	*	*	*
V43.3	124	66	58	*	*8	34	82	37	35	37	16
V43.6	292	92	200	*	9	51	232	59	84	97	51
V43.64	129	39	90	*	*6	23	101	28	37	41	23
V43.65	155	51	104	—	*	27	125	31	45	53	27
V44	168	83	85	40	25	38	65	47	40	50	31
V44.0	34	16	18	13	*	10	*8	10	*9	10	*5
V44.1	65	34	31	21	*7	11	26	17	15	22	11
V44.2	19	10	9	*	*6	*	*8	*	*	*6	*
V44.3	38	18	20	*	*	10	21	12	9	10	*6
V44.6	*6	*	*	*	*	*	*	*	*	*	*
V45	1,744	1,009	736	34	117	461	1,132	397	484	594	269
V45.0	294	142	151	*	*8	33	252	85	73	102	34
V45.01	254	116	138	*	*	21	227	70	64	89	31
V45.02	33	23	10	*	*	10	20	11	*7	11	*
V45.1	79	39	40	*	16	26	37	17	15	33	14
V45.2	32	16	16	15	*7	*	*6	*6	*7	11	*8

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V45.3	12	*	*9	*	*	*	*	*	*	*	*
V45.4	38	16	21	*	10	14	13	*	11	12	12
V45.6	*7	*	*	*	*	*	*6	*	*	*	*
V45.61	*6	*	*	*	*	*	*5	*	*	*	*
V45.7	49	12	37	*	*6	13	29	11	15	12	11
V45.71	20	*	20	—	*	*5	14	*	*8	*	*
V45.72	*9	*	*	*	*	*	*5	*	*	*	*
V45.73	20	*8	12	*	*	*5	10	*	*	*6	*5
V45.8	1,233	774	458	16	64	366	786	273	356	418	185
V45.81	663	454	210	*	10	167	486	160	193	226	85
V45.82	364	236	129	*	19	147	197	70	108	123	63
V45.89	205	85	119	15	36	52	102	43	56	69	37
V46	17	*7	9	*	*	*	*5	*6	*	*	*
V46.1	14	*7	*7	*	*	*	*5	*6	*	*	*
V49	85	52	33	*	10	23	51	15	20	36	14
V49.7	81	50	31	*	9	21	50	15	19	35	12
V49.72	*7	*	*	—	*	*	*	*	*	*	*
V49.75	40	25	15	*	*	10	26	*8	*9	18	*
V49.76	24	15	9	*	*	*6	14	*	*6	9	*
V50	*7	*	*5	*	*	*	*	*	*	*	*
V53	13	*7	*6	*	*	*	*8	*	*	*	*
V53.3	11	*6	*5	*	*	*	*8	*	*	*	*
V53.31	*8	*	*	*	*	*	*6	*	*	*	*
V54	45	16	29	*	*6	*6	29	*7	9	19	9
V54.0	*7	*	*	*	*	*	*	*	*	*	*
V54.8	25	*8	18	*	*	*	19	*	*7	10	*6
V54.9	13	*	*9	*	*	*	10	*	*	*6	*
V55	48	24	23	*6	*9	15	17	13	9	14	11
V55.1	*5	*	*	*	*	*	*	*	*	*	*
V55.2	9	*	*	*	*	*	*	*	*	*	*
V55.3	27	15	12	*	*	10	11	*9	*6	*7	*6
V56	78	35	43	*	11	29	37	13	*7	37	21
V56.0	70	30	39	*	10	26	34	13	*5	33	18
V56.8	*7	*	*	*	*	*	*	*	*	*	*
V57	688	259	429	11	71	109	498	140	162	281	106
V57.1	184	62	122	*	13	30	138	49	19	85	31
V57.2	71	27	45	*	15	10	46	19	11	29	13
V57.21	71	27	45	*	15	10	46	19	11	29	13
V57.3	*7	*	*	*	*	*	*	*	*	*	*
V57.8	401	159	242	*6	37	64	294	60	127	153	61
V57.89	401	159	242	*6	37	64	294	60	127	153	61
V57.9	23	*8	15	*	*	*	15	11	*	11	*
V58	376	187	189	49	58	104	166	92	72	137	76
V58.0	31	13	18	*	*6	11	13	*8	*	16	*
V58.1	150	90	60	42	30	50	28	41	22	59	29
V58.4	30	13	17	*	*7	*7	12	*9	*	*7	10
V58.49	29	13	17	*	*7	*7	12	*9	*	*7	10
V58.6	152	66	87	*	13	33	105	33	41	47	32
V58.61	137	61	76	*	11	30	96	30	35	43	30
V58.69	13	*	9	*	*	*	*8	*	*6	*	*
V58.8	11	*	*7	*	*	*	*8	*	*	*7	*
V58.89	10	*	*6	*	*	*	*8	*	*	*7	*
V59	*7	*	*	*	*	*	*	*	*	*	*
V60	26	16	10	*	14	*7	*6	*	*6	9	*7
V60.0	15	11	*	*	*8	*6	*	*	*5	*	*5
V61	51	20	31	*9	38	*	*	*	15	23	9
V61.0	10	*	*5	*	*8	*	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V61.1	11	*	*6	—	*9	*	*	*	*	*	*
V61.10	10	*	*6	—	*9	*	*	*	*	*	*
V61.2	14	10	*	*7	*7	*	—	*	*5	*8	*
V61.20	13	*9	*	*6	*7	*	—	*	*	*8	*
V61.5	9	—	9	—	9	—	—	*	*	*	*
V62	27	13	14	*	14	*	*6	*7	*5	12	*
V62.8	19	*8	11	*	10	*	*	*6	*	*7	*
V62.81	*6	*	*	*	*	*	*	*	*	*	*
V62.89	12	*7	*6	*	*7	*	*	*	*	*	*
V63	*5	*	*	*	*	*	*	*	*	*	*
V64	99	42	56	*	37	30	29	20	22	39	19
V64.1	20	10	10	*	*	*6	*9	*	*	*8	*
V64.2	16	*7	9	—	*5	*	*8	*	*	*7	*
V64.3	15	*9	*6	*	*6	*	*	*	*	*7	*
V64.4	47	17	31	*	21	16	*9	9	12	17	*9
V65	22	*8	14	*	13	*	*	*7	*5	*	*6
V65.3	12	*	*7	*	*	*	*	*	*	*	*
V65.4	*7	*	*	—	*7	*	—	*5	*	—	*
V66	42	20	23	*	*	*9	30	10	9	16	*7
V66.7	32	14	17	*	*	*7	22	*	*7	14	*7
V70	13	*6	*6	*	*6	*	*	*	*	*6	*
V70.7	11	*5	*6	*	*6	*	*	*	*	*	*
V71	19	9	10	*	9	*	*	*	*5	*8	*
V71.0	*6	*	*	*	*	*	*	—	*	*	—
V71.09	*6	*	*	*	*	*	*	—	*	*	—
V72	17	*7	10	*	*	*	9	*9	*	*6	—
V72.8	12	*	*7	—	*	*	*8	*8	*	*	—
V72.83	*5	*	*	—	*	*	*	*	*	*	—

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
All codes	39,981	15,654	24,328	1,951	13,518	9,775	14,737	9,689	8,756	13,852	7,683
01-05	1,000	458	542	210	309	238	242	206	273	297	223
01	122	71	51	19	33	37	33	24	29	33	35
01.0	13	*7	*6	*8	*	*	*	*	*	*	*5
01.02	9	*	*	*6	*	*	*	*	*	*	*
01.1	20	13	*7	*	*6	*5	*	*	*6	*	*6
01.13	*6	*	*	*	*	*	*	*	*	*	*
01.18	12	*8	*	*	*5	*	*	*	*	*	*
01.2	16	*7	9	*	*6	*	*	*	*	*5	*
01.24	11	*	*7	*	*	*	*	*	*	*	*
01.3	30	20	10	*	*7	*7	15	*8	*6	*6	10
01.31	20	15	*5	*	*	*	12	*	*	*	*7
01.39	10	*5	*5	*	*	*	*	*	*	*	*
01.5	40	21	19	*	11	18	*8	*9	*8	13	10
01.51	*8	*	*6	*	*	*	*	*	*	*	*
01.59	31	18	13	*	9	14	*	*7	*7	*9	*8
02	109	56	52	40	29	22	18	23	28	33	25
02.0	18	11	*7	*7	*6	*	*	*	*	*6	*
02.02	*5	*	*	*	*	*	*	*	*	*	*
02.06	*7	*	*	*	*	*	*	*	*	*	*
02.1	*9	*	*	*	*	*	*	*	*	*	*
02.12	*8	*	*	*	*	*	*	*	*	*	*
02.2	23	11	12	*	*7	*8	*	*6	*	*8	*5
02.3	18	10	*8	*7	*	*	*6	*	*	*5	*
02.34	15	*9	*7	*6	*	*	*	*	*	*	*
02.4	30	15	15	19	*7	*	*	*5	12	*9	*
02.42	25	12	13	16	*	*	*	*	10	*7	*
02.43	*6	*	*	*	*	*	*	*	*	*	*
02.9	10	*	*	*	*	*	*	*	*	*	*
02.94	*6	*	*	*	*	*	*	*	*	*	*
03	711	305	406	147	229	161	175	144	204	212	150
03.0	132	67	66	*	19	54	58	26	31	44	31
03.09	128	64	64	*	17	53	57	25	31	43	30
03.3	298	156	141	131	70	48	48	60	63	103	72
03.31	296	156	141	131	70	47	48	60	62	103	72
03.4	10	*6	*	*	*	*	*	*	*	*	*
03.5	21	11	10	*	*8	*6	*	*	*6	*5	*7
03.53	*8	*	*	*	*	*	*	*	*	*	*
03.59	11	*6	*5	*	*	*	*	*	*	*	*
03.6	*8	*5	*	*	*	*	*	*	*	*	*
03.8	*6	*	*	*	*	*	*	*	*	*	*
03.9	232	55	177	*	125	45	58	52	97	49	34
03.90	93	25	67	*	51	17	22	26	35	17	15
03.91	76	10	66	*	52	10	13	10	42	15	*9
03.92	40	13	28	*	*5	13	22	12	14	*9	*6
03.93	*8	*	*	—	*	*	*	*	*	*	*
03.95	11	*	10	*	10	*	—	*	*	*	*
04	50	24	26	*	15	15	15	13	10	17	10
04.0	*8	*	*	*	*	*	*	*	*	*	*
04.3	*7	*	*	*	*	*	*	*	*	*	*
04.4	15	*6	*9	*	*	*7	*	*	*	*6	*
04.43	*7	*	*	*	*	*	*	*	*	*	*
04.49	*6	*	*	*	*	*	*	*	*	*	*
04.8	*9	*	*5	*	*	*	*5	*	*	*	*
04.81	*8	*	*	*	*	*	*5	*	*	*	*
05	*8	*	*6	*	*	*	*	*	*	*	*
05.3	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
06-07	90	23	67	*	27	37	23	22	14	37	18
06	74	17	57	*	22	31	18	17	11	32	13
06.2	19	*	15	*	*5	11	*	*	*	*8	*
06.3	12	*	11	*	*	*5	*	*	*	*6	*
06.39	11	*	10	*	*	*	*	*	*	*6	*
06.4	14	*	12	—	*5	*6	*	*	*	*5	*
06.8	15	*	11	*	*	*6	*5	*	*	*6	*
06.89	14	*	10	*	*	*5	*	*	*	*5	*
07	16	*6	10	*	*	*6	*	*	*	*5	*
07.2	*5	*	*	*	*	*	*	*	*	*	*
07.6	*7	*	*	*	*	*	*	*	*	*	*
08-16	80	43	37	*9	20	20	32	19	16	34	11
08	24	16	*9	*	11	*8	*5	*7	*5	9	*
08.8	19	13	*6	*	*9	*5	*	*6	*	*6	*
08.81	14	11	*	*	*8	*	*	*	*	*	*
12	*6	*	*	*	*	*	*	*	*	*	*
13	*8	*	*5	*	*	*	*6	*	*	*	*
14	26	13	13	*	*	*7	12	*	*5	14	*
14.7	10	*5	*5	*	*	*	*5	*	*	*6	*
16	*7	*	*	*	*	*	*	*	*	*	*
18-20	37	24	13	22	*8	*	*	9	*6	13	*8
18	12	9	*	*	*5	*	*	*	*	*	*
20	23	13	9	18	*	*	*	*7	*	*9	*
20.0	17	11	*6	15	*	*	*	*	*	*7	*
20.01	16	10	*6	15	*	*	*	*	*	*7	*
21-29	261	158	103	56	89	57	59	69	56	88	48
21	63	41	22	*	23	14	22	15	17	17	13
21.0	24	15	9	*	*	*5	15	*6	*8	*	*6
21.01	*8	*	*	*	*	*	*	*	*	*	*
21.02	*7	*5	*	*	*	*	*	*	*	*	*
21.03	*8	*5	*	*	*	*	*5	*	*	*	*
21.6	*5	*	*	*	*	*	*	*	*	*	*
21.7	*6	*	*	*	*	*	*	*	*	*	*
21.8	19	14	*6	*	11	*	*	*	*5	*7	*
21.81	*6	*	*	*	*	*	*	*	*	*	*
21.88	*8	*7	*	*	*	*	*	*	*	*	*
22	28	17	11	*	11	9	*	11	*6	*8	*
22.6	13	*7	*5	*	*	*	*	*	*	*	*
22.63	*8	*	*	*	*	*	*	*	*	*	*
23	23	12	11	*6	11	*	*	*6	*	10	*
23.0	*8	*	*	*	*	*	*	*	*	*	*
23.09	*8	*	*	*	*	*	*	*	*	*	*
23.1	12	*6	*6	*	*8	*	*	*	*	*	*
23.19	12	*6	*6	*	*7	*	*	*	*	*	*
24	*5	*	*	*	*	*	*	*	*	*	*
25	9	*6	*	*	*	*	*	*	*	*	*
26	16	*8	*8	*	*	*	*8	*	*	*7	*
26.3	12	*6	*5	*	*	*	*6	*	*	*	*
26.31	*7	*	*	*	*	*	*	*	*	*	*
27	50	31	19	17	15	11	*7	13	9	16	12
27.0	*6	*	*	*	*	*	*	*	*	*	*
27.5	19	13	*6	*8	*6	*	*	*5	*	*	*5
27.51	10	*8	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
27.6	17	11	*5	*8	*	*	*	*	*	*	*
27.62	*6	*	*	*6	*	—	—	*	*	*	*
27.69	*9	*7	*	*	*	*	*	*	*	*	*
28	41	23	18	20	13	*6	*	12	*8	16	*5
28.0	*9	*	*	*	*	*	*	*	*	*	*
28.2	11	*7	*	*	*6	*	*	*	*	*	*
28.3	15	*8	*6	14	*	*	—	*6	*	*5	*
29	26	16	10	*	*6	*7	9	*5	*7	*9	*5
29.1	9	*	*5	*	*	*	*	*	*	*	*
29.11	*7	*	*	*	*	*	*	*	*	*	*
29.3	*6	*	*	*	*	*	*	—	*	*	*
29.4	10	*8	*	*	*	*	*	*	*	*	*
30-34	969	526	443	57	164	271	477	249	198	358	164
30	*7	*	*	*	*	*	*	*	*	*	*
31	136	75	61	16	27	38	55	36	29	50	22
31.1	67	40	27	*	16	19	30	17	13	27	10
31.2	13	*7	*6	*	*	*5	*6	*	*	*	*
31.29	13	*7	*6	*	*	*5	*6	*	*	*	*
31.4	44	23	21	11	*6	11	15	13	10	14	*7
31.42	40	20	20	11	*6	10	13	12	9	12	*7
32	62	36	26	*	9	23	27	19	12	18	12
32.2	28	17	11	*	*6	11	10	10	*	*8	*
32.29	26	16	10	*	*5	11	*9	9	*	*8	*
32.3	*6	*	*	*	*	*	*	*	*	*	*
32.4	22	11	11	*	*	9	11	*5	*5	*7	*
33	302	174	129	18	48	89	147	82	54	118	49
33.2	291	166	125	17	46	84	143	77	53	113	48
33.22	52	31	21	*	10	16	21	14	9	16	12
33.23	39	19	20	*8	*	12	13	13	*6	15	*6
33.24	98	57	41	*	17	27	52	27	20	39	12
33.26	25	14	11	*	*	*	19	*	*	10	*7
33.27	54	35	19	*	*8	18	28	12	10	25	*8
33.28	18	*7	11	*	*	*6	*8	*	*	*8	*
33.3	*6	*	*	*	*	*	*	*	*	*	*
34	461	238	224	19	79	118	245	110	102	169	81
34.0	178	102	76	11	42	47	78	39	39	66	34
34.03	*8	*5	*	*	*	*	*6	*	*	*	*
34.04	138	79	59	*8	33	38	58	33	28	51	25
34.09	26	14	12	*	*8	*6	11	*	*8	*9	*6
34.2	35	20	15	*	*7	11	17	12	*6	11	*
34.21	12	*8	*	*	*	*	*6	*	*	*	*
34.22	*8	*	*	—	*	*	*	*	*	*	*
34.24	*8	*6	*	*	*	*	*	*	*	*	*
34.5	16	9	*6	*	*5	*	*	*	*	*5	*
34.51	13	*7	*5	*	*	*	*	*	*	*	*
34.6	*7	*	*	*	*	*	*	*	*	*	*
34.7	*6	*	*	*	*	*	*	*	*	*	*
34.9	207	92	116	*	18	46	140	48	50	75	34
34.91	181	78	103	*	16	40	122	41	46	64	30
34.92	25	13	13	*	*	*6	17	*7	*	11	*
35-39	5,939	3,420	2,519	154	549	2,113	3,123	1,315	1,412	2,167	1,045
35	118	66	52	20	12	31	54	33	25	33	27
35.1	14	*7	*7	*	*	*	*7	*	*	*	*
35.12	11	*6	*	*	*	*	*5	*	*	*	*
35.2	73	41	32	*	*7	23	42	19	18	21	14
35.21	17	*9	*9	*	*	*	13	*7	*	*	*
35.22	28	20	*9	*	*	11	14	*7	*8	*9	*5

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
35.23	*7	*	*	—	—	*	*6	*	*	*	*
35.24	16	*7	*9	*	*	*7	*7	*	*	*6	*
35.3	*7	*	*	*	*	*	*	*	*	*	*
35.33	*7	*	*	*	*	*	*	*	*	*	*
35.6	*5	*	*	*	*	*	*	*	*	*	*
35.7	*6	*	*	*	*	*	*	*	*	*	*
36	1,549	1,029	519	*	80	654	813	319	424	535	271
36.0	1,025	655	370	*	62	437	524	194	290	360	181
36.01	470	300	170	*	30	200	240	92	137	162	78
36.02	11	*8	*	—	*	*	*	*	*	*	*
36.04	*6	*	*	—	*	*	*	*	*	*	*
36.05	80	52	28	*	*	36	43	15	22	26	16
36.06	456	290	166	*	28	195	233	85	127	163	82
36.1	519	371	148	*	17	216	286	123	133	174	89
36.10	13	9	*	—	*	*	9	*	*	*7	*
36.11	51	34	17	*	*	23	26	13	14	17	*7
36.12	98	72	27	*	*	37	57	23	24	32	20
36.13	89	62	27	—	*	36	52	20	25	29	14
36.14	48	33	15	—	*	19	28	*7	11	23	*8
36.15	202	147	55	—	*7	86	109	49	54	63	36
36.16	11	9	*	—	*	*7	*	*	*	*	*
36.19	*6	*	*	—	*	*	*	*	*	*	*
37	1,782	1,039	743	15	135	639	993	409	431	671	271
37.0	*9	*	*	*	*	*	*	*	*	*	*
37.1	10	*	*6	*	*	*	*	*	*	*	*
37.12	*9	*	*5	*	*	*	*	*	*	*	*
37.2	1,318	793	525	9	113	550	646	274	324	516	202
37.21	29	16	13	*	*	*9	15	11	*	12	*
37.22	1,035	626	409	*	82	463	489	191	262	423	159
37.23	157	89	68	*7	9	48	93	43	36	54	24
37.26	71	48	23	*	11	22	38	22	19	19	11
37.27	14	*7	*7	*	*5	*	*	*	*	*5	*
37.29	*6	*	*	*	*	*	*	*	*	*	*
37.3	34	16	18	*	*8	10	14	*8	*7	11	*7
37.34	25	12	13	*	*7	*7	11	*6	*5	*8	*6
37.6	37	23	13	*	*	12	22	11	12	*8	*
37.61	32	21	12	—	*	11	20	10	11	*8	*
37.7	175	88	86	*	*	25	145	52	38	61	24
37.71	30	15	15	*	*	*	27	*9	*5	11	*
37.72	100	51	49	*	*	14	85	31	24	33	12
37.75	*6	*	*	—	*	*	*	*	*	*	*
37.76	*8	*	*	*	*	*	*6	*	*	*	*
37.78	24	12	12	*	*	*5	17	*7	*	*8	*6
37.8	152	74	79	*	*	20	129	47	35	52	19
37.80	*8	*	*5	—	—	*	*7	*	*	*	*
37.81	11	*6	*5	*	*	*	10	*	*	*5	*
37.82	19	*9	10	—	*	*	16	*6	*	*6	*
37.83	99	50	49	*	*	15	83	31	23	33	13
37.87	*8	*	*	*	*	*	*6	*	*	*	*
37.9	45	34	11	*	*	14	28	11	10	15	9
37.94	34	27	*7	*	*	10	22	9	*8	11	*7
38	1,300	637	663	91	192	363	653	291	278	494	237
38.0	34	16	18	*	*	10	21	*8	*7	12	*7
38.03	*7	*	*	*	*	*	*	*	*	*	*
38.08	17	9	*8	*	*	*6	10	*	*	*7	*
38.1	144	79	65	*	*	38	104	32	35	57	20
38.12	124	67	57	—	*	32	91	29	30	50	16
38.18	13	*8	*5	—	*	*	*9	*	*	*	*
38.2	*6	*	*	—	*	*	*5	*	*	*	*
38.4	49	32	17	*	*	10	36	13	10	16	10

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
38.44	30	22	*8	*	*	*	26	*8	*6	10	*5
38.45	10	*5	*	*	*	*	*	*	*	*	*
38.5	*6	*	*	—	*	*	*	*	*	*	*
38.59	*6	*	*	—	*	*	*	*	*	*	*
38.6	*9	*	*	*	*	*	*6	*	*	*5	*
38.7	52	22	30	—	*	14	34	19	10	18	*
38.8	32	17	15	*8	*9	*8	*8	9	*	13	*6
38.85	9	*	*	*7	*	*	*	*	*	*	*
38.86	*8	*	*	*	*	*	*	*	*	*	*
38.9	966	459	507	82	169	277	438	202	209	368	186
38.91	86	45	41	17	12	21	36	16	27	28	15
38.92	10	*6	*	10	—	—	*	*	*	*	*5
38.93	682	312	371	51	130	190	312	147	149	252	134
38.94	*8	*	*	*	*	*	*	*	*	*	*
38.95	136	65	70	*	22	51	62	32	24	54	26
38.98	35	22	13	*	*	11	21	*	*5	27	*
38.99	*9	*6	*	*	*	*	*	*	*	*	*
39	1,190	648	542	25	130	425	609	263	253	434	239
39.2	151	92	60	*	*9	50	91	36	31	57	27
39.25	24	15	9	—	*	10	12	*	*8	9	*
39.27	33	17	16	*	*	14	16	*6	*	16	*6
39.29	86	56	31	*	*	24	60	25	18	29	14
39.3	20	12	*8	*	9	*	*5	*	*	*9	*
39.31	15	9	*6	*	*7	*	*	*	*	*8	*
39.4	72	36	35	*	10	25	36	15	10	28	19
39.42	20	*9	11	*	*6	*6	*7	*	*	*9	*6
39.43	*7	*	*	—	*	*	*	*	*	*	*
39.49	43	22	20	*	*	16	24	9	*7	15	11
39.5	126	63	63	*	14	38	71	26	38	44	18
39.50	81	41	40	*	*6	26	49	17	26	28	10
39.51	11	*	*8	*	*	*	*	*	*	*	*
39.52	14	10	*	—	*	*	10	*	*	*6	*
39.56	*6	*	*	*	*	*	*	*	*	*	*
39.59	*7	*	*	*	*	*	*	*	*	*	*
39.6	297	187	110	15	16	117	147	73	65	95	64
39.61	271	173	98	14	15	104	137	70	62	85	53
39.62	*9	*5	*	*	*	*	*	*	*	*	*6
39.63	*6	*	*	*	*	*	*	*	*	*	*
39.64	9	*5	*	*	*	*	*	*	*	*6	*
39.9	516	254	262	*	70	187	257	110	103	198	105
39.90	35	16	19	*	*	13	21	*7	12	12	*
39.93	*6	*	*	*	*	*	*	*	*	*	*
39.95	456	224	232	*	65	164	226	97	83	179	97
39.98	10	*7	*	*	*	*	*	*	*	*	*
40-41	315	160	155	22	49	108	135	77	66	108	64
40	187	88	100	*	25	75	84	47	39	64	38
40.1	48	21	27	*	*8	17	21	12	9	17	9
40.11	44	20	24	*	*7	16	19	11	*8	17	*7
40.2	47	19	29	*	*7	18	21	12	12	14	*9
40.23	11	*	11	—	*	*	*6	*	*	*	*
40.29	28	14	14	*	*	11	13	*7	*8	11	*
40.3	64	30	33	*	*7	28	29	15	13	22	14
40.4	13	*7	*6	*	*	*5	*5	*	*	*6	*
40.41	11	*6	*	*	*	*	*	*	*	*6	*
40.5	15	10	*	*	*	*6	*7	*	*	*	*
40.59	*6	*	*	*	*	*	*	*	*	*	*
41	127	72	55	18	25	33	51	30	27	45	26
41.0	*6	*	*	*	*	*	*	*	*	*	*
41.3	89	50	39	14	12	20	43	19	20	32	18

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
41.31	89	50	39	14	12	20	42	19	20	31	18
41.5	27	15	11	*	*9	9	*7	*5	*6	9	*6
42-54	5,145	2,180	2,965	208	1,136	1,404	2,397	1,212	1,158	1,814	961
42	180	96	84	*6	26	59	90	46	39	58	37
42.2	111	59	52	*	16	37	54	35	24	35	17
42.23	106	55	51	*	14	37	52	33	23	34	16
42.3	24	15	*9	*	*	11	*8	*	*6	*6	*9
42.33	22	15	*7	*	*	11	*7	*	*6	*5	*8
42.9	34	15	19	*	*	*7	23	*5	*7	13	*8
42.92	33	14	19	*	*	*7	23	*	*7	13	*8
43	223	111	112	22	17	43	141	58	45	84	36
43.1	171	80	91	10	15	29	116	45	32	68	27
43.11	140	64	76	*	10	23	103	36	25	59	20
43.19	30	16	15	*6	*	*6	13	*8	*7	*9	*7
43.3	12	10	*	11	*	*	*	*	*	*	*
43.4	15	*9	*	*	*	*	10	*6	*	*	*
43.41	12	*7	*5	*	*	*	*9	*5	*	*	*
43.7	*8	*	*	—	*	*	*	*	*	*	*
43.8	*6	*	*	—	*	*	*5	*	*	*	*
43.89	*6	*	*	—	*	*	*5	*	*	*	*
44	211	95	117	12	53	58	89	46	46	74	44
44.0	*7	*	*	—	*	*	*	*	*	*	*
44.1	27	11	16	*	*	*7	16	*	*5	15	*
44.13	10	*	*7	*	*	*	*6	*	*	*	*
44.14	16	*7	*9	*	*	*5	*8	*	*	10	*
44.2	10	*	*5	*	*	*	*5	*	*	*	*
44.29	*7	*	*	*	*	*	*	*	*	*	*
44.3	38	10	28	*	21	11	*5	*7	11	10	10
44.31	22	*	18	—	16	*6	*	*	*7	*6	*6
44.39	16	*7	10	*	*5	*	*5	*	*	*	*
44.4	71	40	31	*	10	12	49	20	14	21	16
44.42	11	*8	*	—	*	*	*8	*5	*	*	*
44.43	54	29	25	*	*7	10	37	12	12	18	12
44.6	54	23	31	10	14	21	10	*8	13	19	14
44.66	43	20	24	*7	11	17	*8	*6	11	15	12
44.69	*6	*	*	*	*	*	*	*	*	*	*
45	1,992	864	1,128	23	250	507	1,212	477	471	710	334
45.1	921	410	511	11	120	233	557	216	220	337	148
45.13	402	182	220	*	50	99	249	97	95	145	65
45.16	507	222	285	*7	68	129	303	113	123	190	81
45.2	545	211	334	*	65	126	349	132	125	197	91
45.23	282	120	162	*	29	60	193	73	63	97	49
45.24	66	23	42	*	11	17	38	11	16	30	*9
45.25	189	65	124	*	26	47	114	44	45	69	30
45.3	18	10	*8	*	*	*	9	*	*	*7	*
45.33	12	*7	*5	*	*	*	*5	*	*	*5	*
45.4	131	60	71	*	*6	30	95	31	30	48	22
45.42	98	41	57	*	*	21	73	23	23	37	15
45.43	28	18	11	—	*	*8	19	*6	*5	10	*7
45.6	58	27	31	*	11	14	30	15	17	17	*9
45.62	54	26	29	*	11	13	28	14	15	17	*8
45.7	236	107	129	*	27	73	134	57	61	74	45
45.72	16	*7	*9	*	*	*	*7	*	*	*6	*
45.73	78	33	45	*	*7	21	50	15	20	25	18
45.74	12	*5	*7	*	*	*	*8	*	*	*	*
45.75	29	14	16	*	*	11	15	*8	*8	9	*
45.76	77	37	41	*	*8	27	42	20	19	23	15
45.79	21	10	11	*	*	*7	10	*6	*	*6	*
45.8	12	*6	*6	*	*	*	*	*	*	*5	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
45.9	64	27	36	*	10	20	32	19	10	23	12
45.91	13	*5	*7	*	*	*	*5	*5	*	*	*
45.93	26	12	14	*	*5	*6	14	*6	*6	*8	*5
45.94	18	*8	10	*	*	*7	10	*6	*	*7	*
46	201	100	101	13	37	57	94	49	48	63	40
46.0	14	*8	*7	*	*	*	*7	*	*	*	*
46.01	*5	*	*	*	*	*	*	*	*	*	*
46.03	*9	*	*	*	*	*	*5	*	*	*	*
46.1	45	21	23	*	*	13	25	11	*9	14	11
46.10	22	10	12	*	*	*6	13	*	*5	*8	*
46.11	13	*8	*	*	*	*	*5	*	*	*	*
46.13	9	*	*7	*	*	*	*6	*	*	*	*
46.2	15	*7	*7	*	*5	*	*	*	*	*	*
46.20	*5	*	*	*	*	*	*	*	*	*	*
46.3	28	12	16	*	*	*7	16	*7	*6	11	*
46.32	*5	*	*	*	*	*	*	*	*	*	*
46.39	23	10	13	*	*	*6	13	*5	*5	*9	*
46.4	15	*6	*9	*	*	*	*8	*	*	*	*
46.41	*5	*	*	*	*	*	*	*	*	*	*
46.42	*5	*	*	—	*	*	*	*	*	*	*
46.5	37	20	17	*	*8	13	13	12	*8	11	*6
46.51	9	*5	*	*	*	*	*	*	*	*	*
46.52	27	14	13	*	*	10	10	*8	*6	*8	*5
46.7	32	14	18	*	10	*8	13	*6	*9	11	*7
46.73	14	*	9	*	*	*	*6	*	*	*	*
46.75	*7	*	*	*	*	*	*	*	*	*	*
46.8	13	9	*	*	*	*	*9	*	*	*	*
46.85	*9	*6	*	*	*	*	*7	*	*	*	*
47	355	179	177	77	186	66	26	71	90	109	85
47.0	303	163	140	71	167	50	16	61	71	94	76
47.01	86	41	45	23	48	13	*	11	20	28	28
47.09	217	123	94	47	119	37	14	51	52	67	48
47.1	49	14	35	*5	19	16	*9	10	17	14	*8
47.11	*5	*	*	*	*	*	*	*	*	*	*
47.19	44	12	32	*	16	16	*8	10	14	13	*7
48	143	70	72	*7	29	38	69	36	30	50	27
48.2	48	21	27	*	10	14	21	11	13	15	10
48.23	22	11	11	*	*6	*8	*9	*	*	*8	*6
48.24	25	9	15	*	*	*6	12	*5	*8	*7	*
48.3	24	12	12	*	*	*	17	*	*6	*9	*
48.36	17	*8	*9	*	*	*	13	*	*	*7	*
48.5	13	*7	*6	—	*	*	*7	*	*	*	*
48.6	30	16	15	*	*	9	16	11	*	12	*
48.63	19	11	*9	—	*	*5	11	*8	*	*7	*
48.69	*8	*	*	*	*	*	*	*	*	*	*
48.7	*9	*	*	*	*	*	*	*	*	*	*
48.8	13	*8	*5	*	*5	*	*	*	*	*7	*
48.81	11	*8	*	*	*	*	*	*	*	*	*
49	46	23	23	*	13	14	17	*7	12	16	10
49.0	*7	*6	*	*	*	*	*	*	*	*	*
49.01	*6	*5	*	*	*	*	*	*	*	*	*
49.4	19	9	10	—	*6	*6	*8	*	*6	*7	*
49.46	14	*7	*7	—	*	*6	*	*	*	*5	*
49.7	*6	*	*	*	*	*	*	*	*	*	*
50	99	51	47	*6	18	35	40	28	17	36	17
50.1	77	38	39	*	12	28	33	22	14	27	14
50.11	52	25	27	*	*9	19	21	14	10	18	10
50.12	24	13	11	*	*	*8	12	*7	*	*9	*
50.2	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
50.6	*5	*	*	*	*	*	*	*	*	*	*
50.61	*5	*	*	*	*	*	*	*	*	*	*
50.9	*6	*	*	*	*	*	*	*	*	*	*
51	667	233	434	*	195	192	275	154	137	240	136
51.0	*5	*	*	*	*	*	*	*	*	*	*
51.1	65	26	39	*	15	19	31	19	14	20	12
51.10	51	21	30	*	14	16	22	14	12	16	10
51.14	*6	*	*	—	*	*	*	*	*	*	*
51.2	419	134	284	*	138	129	149	86	83	169	81
51.22	101	41	60	*	22	33	45	20	25	38	18
51.23	317	93	224	*	115	96	103	65	58	130	63
51.3	9	*	*	*	*	*	*	*	*	*	*
51.4	10	*	*5	—	*	*	*6	*	*	*	*
51.41	*7	*	*	—	*	*	*	*	*	*	*
51.5	*8	*	*	—	*	*	*	*	*	*	*
51.8	138	49	89	*	35	32	71	33	30	39	36
51.85	65	22	43	*	18	16	31	14	16	18	18
51.87	31	12	19	—	*	9	17	10	*	9	*7
51.88	36	13	23	*	11	*	20	*7	*9	10	10
51.9	12	*5	*6	*	*	*	*6	*7	*	*	*
51.98	10	*	*5	*	*	*	*5	*6	*	*	*
52	39	16	24	*	*8	14	16	*8	10	13	*8
52.1	11	*	*7	*	*	*	*5	*	*	*	*
52.7	*5	*	*	—	*	*	*	*	*	*	*
52.9	10	*	*7	*	*	*	*	*	*	*	*
52.93	*7	*	*5	*	*	*	*	*	*	*	*
53	230	112	119	10	42	79	99	53	49	87	41
53.0	46	37	*9	*	*7	10	26	13	*8	16	10
53.00	*8	*6	*	*	*	*	*	*	*	*	*
53.01	*6	*	*	*	*	*	*	*	*	*	*
53.03	*9	*8	*	—	*	*	*6	*	*	*	*
53.04	*8	*7	*	—	*	*	*	*	*	*	*
53.05	12	*8	*	—	*	*	*7	*	*	*	*
53.1	11	10	*	*	*	*	*	*	*	*	*
53.4	50	20	30	*	12	18	20	9	12	19	10
53.49	45	19	26	*	11	16	18	*8	11	18	*9
53.5	41	11	30	*	*7	14	18	*8	*9	16	*8
53.51	24	*6	18	*	*	10	9	*	*	*9	*6
53.59	17	*5	11	*	*	*	*8	*	*	*7	*
53.6	65	27	38	—	10	28	27	16	14	27	*8
53.61	52	21	31	—	*7	22	23	12	12	22	*6
53.69	13	*6	*7	—	*	*6	*	*	*	*5	*
53.7	*6	*	*	*	*	*	*	*	*	*	*
54	759	231	528	25	263	241	230	179	163	274	143
54.0	15	*6	*8	*	*	*	*6	*	*	*5	*
54.1	62	25	37	*	22	18	19	14	14	24	10
54.11	31	12	19	*	12	*9	9	*6	*6	13	*6
54.12	13	*	*8	*	*5	*	*	*	*	*	*
54.19	18	*8	10	*	*5	*7	*6	*	*	*7	*
54.2	113	25	88	*	51	33	27	28	25	37	22
54.21	69	12	57	*	41	16	10	14	17	24	14
54.23	20	*7	13	*	*	*8	*7	*7	*	*	*
54.24	*8	*	*5	*	*	*	*	*	*	*	*
54.25	12	*	10	*	*	*	*	*	*	*	*
54.3	13	*	*8	*	*	*6	*	*	*	*	*
54.4	59	12	46	*	17	19	21	14	12	20	13
54.5	307	61	246	*	123	94	85	71	63	123	51
54.51	41	*7	34	*	15	14	12	*7	*9	16	*9
54.59	266	55	211	*	108	81	73	63	54	106	42
54.6	15	*7	*7	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
54.61	*7	*	*	*	*	*	*	*	*	*	*
54.7	*9	*	*6	*	*	*	*	*	*	*	*
54.9	168	86	82	10	36	63	59	40	38	54	36
54.91	106	50	56	*	22	43	40	25	27	37	17
54.93	12	*7	*	*	*	*	*	*	*	*	*
54.95	15	*8	*6	*5	*	*5	*	*	*	*	*
54.98	31	18	13	*	*7	9	12	*6	*7	*8	10
55-59	962	450	511	35	235	291	401	244	202	375	141
55	126	64	62	*7	39	46	34	31	22	55	18
55.0	28	13	14	*	10	*7	*9	9	*	10	*
55.03	18	*9	10	*	*7	*	*7	*7	*	*7	*
55.2	22	12	10	*	*7	9	*	*	*	*8	*
55.23	20	11	*9	*	*7	*8	*	*	*	*8	*
55.5	47	25	22	*	10	20	16	11	9	22	*
55.51	44	23	21	*	*9	18	16	11	9	20	*
55.6	10	*6	*	*	*5	*	*	*	*	*6	*
55.69	10	*6	*	*	*5	*	*	*	*	*6	*
55.8	*5	*	*	*	*	*	*	*	*	*	*
56	111	60	51	12	35	31	32	30	22	46	13
56.0	51	30	22	*	21	16	14	13	9	27	*
56.2	*6	*	*	*	*	*	*	*	*	*	*
56.3	24	11	13	*	9	*7	*8	*7	*	10	*
56.31	20	10	10	*	*9	*6	*	*6	*	10	*
56.5	*9	*6	*	*	*	*	*6	*	*	*	*
56.51	*9	*6	*	*	*	*	*5	*	*	*	*
56.7	13	*	*8	*9	*	*	*	*	*	*	*
56.74	12	*	*8	*9	*	*	*	*	*	*	*
57	392	216	177	9	67	99	217	111	88	138	55
57.0	25	22	*	*	*	*6	18	*6	*9	*8	*
57.1	44	15	29	*	*8	17	18	*8	*9	21	*6
57.17	16	*6	10	*	*	*6	*8	*5	*	*	*
57.18	23	*8	15	*	*6	10	*7	*	*	15	*
57.3	183	97	86	*	33	48	97	50	45	63	24
57.32	167	89	79	*	32	44	87	44	42	59	22
57.33	13	*7	*6	—	*	*	9	*	*	*	*
57.4	42	32	10	—	*	9	31	14	10	15	*
57.49	42	32	10	—	*	9	31	14	10	15	*
57.7	*8	*7	*	*	*	*	*	*	*	*	*
57.71	*7	*6	*	*	*	*	*	*	*	*	*
57.8	22	*5	17	*	*8	*	*8	*	*6	*8	*5
57.81	11	*	9	*	*5	*	*	*	*	*	*
57.9	61	33	28	*	12	9	38	23	*7	20	10
57.91	*5	*	*	—	*	*	*	*	*	*	*
57.94	49	23	27	*	12	*7	30	21	*5	14	9
58	58	44	14	*	*7	13	34	13	15	22	*8
58.4	*7	*	*	*	*	*	*	*	*	*	*
58.5	*8	*7	*	—	*	*	*6	*	*	*	*
58.6	31	22	*8	*	*	*5	22	*7	10	11	*
59	274	67	208	*	87	102	83	58	56	114	46
59.0	*7	*	*6	*	*	*	*	*	*	*	*
59.02	*6	*	*	*	*	*	*	*	*	*	*
59.3	*8	*	*7	—	*	*	*	*	*	*	*
59.4	11	—	11	*	*	*	*	*	*	*	*
59.5	50	—	50	—	19	23	*8	*	11	26	9
59.7	50	*	50	*	11	23	16	*6	11	18	14
59.79	48	*	48	—	11	22	16	*6	11	17	14
59.8	135	59	76	*	45	40	48	38	26	56	14
59.9	*7	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
60-64	258	258	—	19	14	68	157	57	70	85	46
60	200	200	—	—	*	52	146	44	55	64	37
60.1	12	12	—	—	—	*	10	*	*	*	*
60.11	12	12	—	—	—	*	10	*	*	*	*
60.2	132	132	—	—	*	22	109	30	39	40	23
60.29	129	129	—	—	*	21	106	29	37	40	23
60.5	46	46	—	—	*	26	20	*8	12	17	10
61	10	10	—	*	*	*5	*	*	*	*	*
61.0	*5	*5	—	*	*	*	*	*	*	*	*
62	10	10	—	*	*	*	*	*	*	*	*
63	*6	*6	—	*	*	*	*	*	*	*	*
64	30	30	—	15	*	*7	*5	*6	10	11	*
64.0	18	18	—	14	*	*	*	*	*6	*6	*
64.9	*8	*8	—	*	*	*	*	*	*	*	*
65-71	2,061	—	2,061	*6	1,192	638	225	376	441	846	397
65	586	—	586	*	279	244	60	107	130	241	108
65.2	43	—	43	*	38	*	*	10	9	13	11
65.25	*8	—	*8	*	*7	*	—	*	*	*	*
65.29	32	—	32	*	29	*	*	*8	*7	11	*7
65.3	20	—	20	*	12	*	*	*	*	*7	*
65.39	18	—	18	*	11	*	*	*	*	*6	*
65.4	86	—	86	*	57	20	*8	15	17	40	14
65.41	11	—	11	*	*8	*	*	*	*	*7	*
65.49	75	—	75	*	49	18	*8	14	16	33	12
65.5	9	—	9	—	*	*	*	*	*	*	*
65.51	*8	—	*8	—	*	*	*	*	*	*	*
65.6	378	—	378	*	132	201	45	67	86	156	70
65.61	337	—	337	*	115	180	42	60	75	143	59
65.62	10	—	10	—	*5	*	*	*	*	*	*
65.63	30	—	30	—	12	16	*	*	*8	10	*9
65.8	33	—	33	—	24	*8	*	*	*7	17	*
65.81	*5	—	*5	—	*	*	*	*	*	*	*
65.89	28	—	28	—	20	*7	*	*	*5	14	*
65.9	*5	—	*5	*	*	*	*	*	*	*	*
66	389	—	389	*	382	*6	*	61	71	179	78
66.0	*7	—	*7	—	*7	—	—	*	*	*	*
66.02	*6	—	*6	—	*6	—	—	*	*	*	*
66.2	*8	—	*8	—	*8	—	—	*	*	*	*
66.3	307	—	307	—	307	*	—	44	56	145	62
66.32	180	—	180	—	180	*	—	23	33	81	42
66.39	126	—	126	—	126	*	—	20	22	64	20
66.4	9	—	9	*	*7	*	*	*	*	*6	*
66.5	*9	—	*9	—	*6	*	*	*	*	*5	*
66.51	*7	—	*7	—	*5	*	*	*	*	*	*
66.6	36	—	36	*	34	*	*	10	*7	11	*8
66.61	*8	—	*8	*	*8	*	*	*	*	*	*
66.62	21	—	21	—	21	—	—	*7	*	*7	*
66.69	*6	—	*6	*	*	*	*	*	*	*	*
66.7	*8	—	*8	*	*8	—	—	*	*	*	*
66.79	*6	—	*6	*	*6	—	—	*	*	*	*
67	20	—	20	—	14	*	*	*6	*	*9	*
67.5	*7	—	*7	—	*7	*	—	*	*	*	*
68	683	—	683	*	354	263	66	126	156	276	125
68.1	11	—	11	—	*5	*	*	*	*	*	*
68.12	*5	—	*5	—	*	*	*	*	*	*	*
68.2	37	—	37	—	30	*6	*	10	*	16	*5
68.29	34	—	34	—	28	*5	*	10	*	15	*5

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
68.3	25	—	25	—	12	11	*	*7	*	*7	*8
68.4	401	—	401	—	193	172	36	70	98	172	61
68.5	198	—	198	—	107	70	22	32	48	74	45
68.51	58	—	58	—	34	22	*	*9	17	21	11
68.59	141	—	141	—	73	48	20	23	31	52	35
68.6	*6	—	*6	—	*	*	*	*	*	*	*
69	106	—	106	*	90	11	*6	29	12	37	28
69.0	62	—	62	—	50	*7	*5	16	*7	20	19
69.02	32	—	32	—	32	—	—	*6	*	11	11
69.09	28	—	28	—	15	*7	*5	*8	*	*9	*7
69.1	*8	—	*8	*	*6	*	*	*	*	*5	*
69.19	*8	—	*8	*	*6	*	*	*	*	*5	*
69.5	23	—	23	*	22	*	*	*9	*	*6	*6
69.52	19	—	19	*	19	—	—	*8	*	*6	*
69.9	9	—	9	*	*9	*	—	*	*	*	*
69.96	*7	—	*7	—	*7	—	—	*	*	*	*
70	246	—	246	*	61	104	79	42	62	95	47
70.3	10	—	10	*	*	*	*	*	*	*	*
70.33	*5	—	*5	*	*	*	*	*	*	*	*
70.5	141	—	141	—	27	67	48	26	35	54	27
70.50	70	—	70	—	13	33	25	15	17	27	10
70.51	37	—	37	—	*6	17	14	*7	10	13	*8
70.52	34	—	34	—	*8	17	9	*	*8	14	*9
70.7	39	—	39	*	13	13	12	*	*7	17	9
70.77	25	—	25	—	*5	9	11	*	*5	11	*7
70.79	*7	—	*7	*	*	*	*	*	*	*	*
70.9	43	—	43	—	11	19	13	*6	15	15	*7
70.92	43	—	43	—	11	19	13	*6	15	15	*7
71	29	—	29	*	13	*7	*9	*	*7	10	*7
71.3	*7	—	*7	*	*5	*	*	*	*	*	*
71.7	12	—	12	*	*	*	*	*	*	*	*
71.79	*8	—	*8	—	*	*	*	*	*	*	*
72-75	6,209	—	6,209	19	6,185	*5	—	1,173	1,401	2,155	1,479
72	366	—	366	*	362	*	—	59	82	132	93
72.0	24	—	24	—	24	—	—	*	*6	11	*
72.1	75	—	75	*	75	*	—	10	16	42	*8
72.2	*6	—	*6	—	*6	—	—	*	*	*	*
72.4	*7	—	*7	—	*7	—	—	*	*	*	*
72.5	*5	—	*5	—	*5	—	—	*	*	*	*
72.7	245	—	245	*	242	*	—	42	54	70	79
72.71	147	—	147	*	145	*	—	30	33	38	46
72.79	98	—	98	*	97	—	—	11	21	32	34
73	2,911	—	2,911	*6	2,903	*	—	479	674	1,080	678
73.0	833	—	833	*	830	*	—	122	246	234	232
73.01	131	—	131	*	131	*	—	13	34	44	39
73.09	702	—	702	*	700	*	—	109	211	190	192
73.1	*9	—	*9	—	*9	—	—	*	*	*	*
73.4	444	—	444	*	443	*	—	80	96	146	122
73.5	899	—	899	*	896	*	—	132	177	404	186
73.51	9	—	9	—	9	—	—	*	*	*	*
73.59	890	—	890	*	887	*	—	131	174	402	183
73.6	718	—	718	*	717	*	—	142	151	289	136
74	856	—	856	*	853	*	—	155	156	358	186
74.0	12	—	12	—	12	*	—	*	*	*6	*
74.1	824	—	824	*	822	*	—	153	148	346	177
74.9	17	—	17	*	16	—	—	*	*5	*5	*6
74.99	17	—	17	*	16	—	—	*	*5	*5	*6
75	2,077	—	2,077	*8	2,067	*	—	481	489	585	522

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
75.1	15	—	15	—	15	—	—	*	*	*	*
75.3	854	—	854	*	851	*	—	257	205	191	201
75.32	193	—	193	*	191	*	—	52	59	28	54
75.34	557	—	557	*	555	*	—	185	119	134	118
75.35	66	—	66	—	65	*	—	16	22	14	14
75.37	37	—	37	*	37	—	—	*	*	14	15
75.4	47	—	47	—	47	*	—	9	11	13	14
75.5	*7	—	*7	—	*7	—	—	*	*	*	*
75.51	*5	—	*5	—	*5	—	—	*	*	*	*
75.6	1,129	—	1,129	*	1,124	*	—	206	261	363	299
75.61	33	—	33	—	33	—	—	*5	*7	13	*8
75.62	109	—	109	*	108	—	—	23	25	37	24
75.69	987	—	987	*	982	*	—	178	229	313	267
75.7	19	—	19	—	19	—	—	*	*	9	*
76-84	3,171	1,546	1,624	172	836	918	1,245	637	728	1,178	628
76	74	47	27	*5	55	*8	*7	16	9	29	19
76.6	16	*	11	*	14	*	*	*	*	*6	*5
76.7	38	29	*8	*	28	*	*	9	*	15	*9
76.72	*5	*	*	*	*	*	*	*	*	*	*
76.75	*7	*	*	*	*5	*	*	*	*	*	*
76.76	13	11	*	*	10	*	*	*	*	*	*
76.79	*5	*	*	*	*	*	*	*	*	*	*
76.9	12	*8	*	*	9	*	*	*	*	*5	*
76.92	*5	*	*	*	*	*	*	*	*	*	*
77	268	139	129	22	83	93	69	53	56	95	64
77.1	*5	*	*	*	*	*	*	*	*	*	*
77.3	17	*9	*8	*6	*	*	*	*	*	*	*
77.37	*6	*	*	*	*	*	*	*	*	*	*
77.4	18	*8	10	*	*	*	10	*	*	*6	*
77.49	10	*	*6	*	*	*	*6	*	*	*	*
77.5	*7	*	*	*	*	*	*	*	*	*	*
77.6	43	26	16	*	12	15	13	11	*9	13	10
77.61	*6	*	*	*	*	*	*	*	*	*	*
77.65	*6	*	*	*	*	*	*	*	*	*	*
77.67	*7	*	*	*	*	*	*	*	*	*	*
77.68	*6	*	*	*	*	*	*	*	*	*	*
77.69	11	*6	*	*	*	*	*	*	*	*	*
77.7	109	55	54	*8	42	43	16	20	24	37	28
77.70	*8	*5	*	*	*	*5	*	*	*	*	*
77.79	91	46	45	*5	37	36	13	14	22	32	24
77.8	54	28	26	*	15	18	20	*7	12	25	10
77.81	24	9	14	*	*5	*9	10	*	*6	12	*
77.88	*6	*	*	*	*	*	*	*	*	*	*
77.89	14	9	*5	*	*5	*	*	*	*	*7	*
77.9	*9	*	*	*	*	*	*	*	*	*	*
78	214	107	107	25	64	56	70	38	53	76	47
78.0	44	23	21	*	13	12	16	9	13	12	10
78.03	*5	*	*	*	*	*	*	*	*	*	*
78.05	13	*5	*8	*	*	*	*8	*	*6	*	*
78.07	13	*7	*5	*	*	*	*	*	*	*	*6
78.09	*7	*6	*	*	*	*	*	*	*	*	*
78.1	31	18	13	*	14	*9	*	*	*7	15	*
78.13	*8	*	*6	*	*	*	*	*	*	*	*
78.17	15	12	*	*	*9	*	*	*	*	10	*
78.4	10	*	*7	*	*	*	*5	*	*	*6	*
78.5	55	27	28	*9	14	11	22	11	13	17	14
78.55	34	15	19	*	*5	*6	18	*6	10	10	*8
78.57	*7	*	*	*	*	*	*	*	*	*	*
78.59	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
78.6	64	32	32	*6	20	20	19	10	18	22	14
78.65	21	*8	13	*	*	*	12	*	*7	*7	*
78.67	14	*7	*6	*	*	*7	*	*	*	*	*
78.69	18	9	*9	*	9	*6	*	*	*	*6	*
78.9	*6	*	*	*	*	*	*	*	*	*	*
79	675	304	370	60	191	140	284	137	157	251	129
79.0	79	38	41	20	20	17	22	21	18	24	15
79.01	10	*	*7	*	*	*	*6	*	*	*	*
79.02	22	*9	13	*8	*5	*	*5	*	*5	*6	*6
79.05	*7	*	*	*6	*	*	*	*	*	*	*
79.06	31	16	15	*	*9	10	*8	*9	*7	11	*
79.1	79	35	44	12	18	14	35	17	18	24	20
79.11	11	*6	*	*7	*	*	*	*	*	*	*
79.12	11	*	*6	*	*	*	*5	*	*	*	*
79.15	37	14	23	*	*5	*	26	*8	11	10	*7
79.16	13	*7	*7	*	*	*6	*	*	*	*6	*
79.2	10	*5	*	*	*	*	*	*	*	*	*
79.3	423	177	246	15	117	87	203	85	97	167	74
79.31	27	10	17	*	*6	*7	11	*9	*6	10	*
79.32	36	19	17	*	15	9	*8	10	*8	12	*7
79.34	9	*7	*	*	*6	*	*	*	*	*	*
79.35	185	53	131	*	15	18	149	32	43	77	32
79.36	139	69	71	*	63	42	31	27	32	54	27
79.37	10	*8	*	*	*	*	*	*	*	*	*
79.39	11	*7	*	*	*5	*	*	*	*	*	*
79.6	43	28	14	*6	21	*9	*7	*6	11	18	*9
79.62	*9	*	*6	*	*	*	*	*	*	*	*
79.65	*6	*	*	*	*	*	*	*	*	*	*
79.66	15	11	*	*	*7	*	*	*	*	*6	*
79.7	27	14	13	*	*	9	12	*7	*7	9	*
79.75	18	*9	9	*	*	*7	*9	*	*	*7	*
79.8	*6	*	*	*	*	*	*	*	*	*	*
80	452	247	206	11	174	175	94	87	95	185	85
80.0	13	*8	*	*	*	*	*7	*	*	*	*
80.05	*5	*	*	—	*	*	*	*	*	*	*
80.06	*6	*	*	—	*	*	*	*	*	*	*
80.1	24	15	*9	*	*7	*6	*7	*6	*5	10	*
80.16	9	*6	*	*	*	*	*	*	*	*	*
80.2	27	15	12	*	10	*7	*9	*	*	10	*7
80.21	11	*7	*	—	*	*6	*	*	*	*	*
80.26	15	*8	*7	*	*8	*	*	*	*	*5	*
80.4	15	*	11	*	*	*	*	*	*	*6	*
80.46	*7	*	*6	*	*	*	*	*	*	*	*
80.5	296	162	134	*	126	128	41	52	65	125	54
80.50	15	*7	*9	—	*5	*7	*	*7	*	*7	—
80.51	279	154	124	*	119	120	38	45	63	118	54
80.6	16	*9	*7	*	9	*	*	*	*	*6	*
80.7	15	*8	*6	*	*	*	*7	*	*	*6	*
80.76	11	*6	*5	*	*	*	*6	*	*	*	*
80.8	41	21	20	*	12	14	15	*7	13	15	*7
80.81	10	*	*6	—	*	*	*	*	*	*	*
80.86	15	*8	*6	*	*6	*	*	*	*	*7	*
80.88	*5	*	*	*	*	*	*	*	*	*	*
81	1,057	455	602	20	169	315	554	205	262	379	212
81.0	244	115	129	12	89	106	37	41	55	92	56
81.02	97	50	47	*	37	50	10	12	22	40	22
81.03	*6	*	*	*	*	*	*	*	*	*	*
81.04	*7	*	*5	*	*	*	*	*	*	*	*
81.05	14	*6	*9	*6	*	*	*	*	*	*	*
81.06	20	10	10	*	11	*7	*	*	*5	*5	*5

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
81.07	13	*6	*7	—	*	*5	*	*	*5	*	*
81.08	76	33	44	*	24	33	16	15	12	28	20
81.09	9	*	*5	*	*5	*	*	*	*	*5	*
81.1	15	*7	*8	*	*	*6	*5	*	*	*	*6
81.11	*5	*	*	*	*	*	*	*	*	*	*
81.2	*8	*	*	*	*	*	*	*	*	*	*
81.4	45	26	19	*	31	*5	*7	*6	*7	20	13
81.40	*6	*	*	*	*	*	*	*	*	*	*
81.45	22	13	*8	*	20	*	—	*	*	*9	*8
81.47	13	*7	*6	*	*9	*	*	*	*	*6	*
81.5	618	241	377	*	25	160	432	121	168	217	111
81.51	152	69	83	—	13	51	88	32	39	53	28
81.52	106	24	82	—	*	*8	95	21	25	39	21
81.53	31	16	15	—	*	10	19	*7	10	*7	*7
81.54	299	115	185	*	*6	82	211	52	87	109	52
81.55	28	17	11	—	*	*8	18	*8	*7	9	*
81.8	47	19	28	*	*8	14	25	*8	*9	20	10
81.80	*8	*	*	—	*	*	*	*	*	*	*
81.81	12	*	10	—	*	*	9	*	*	*	*
81.83	22	10	12	*	*	*8	9	*	*	10	*
81.9	77	43	34	*	*9	17	46	22	19	23	13
81.91	50	32	18	*	*6	10	30	15	11	13	10
81.92	20	*7	13	*	*	*	15	*5	*7	*5	*
82	25	19	*6	*	11	*7	*6	*6	*8	*5	*6
82.4	11	10	*	*	*7	*	*	*	*	*	*
82.45	*5	*	*	*	*	*	*	*	*	*	*
83	237	126	111	26	66	73	72	51	52	89	45
83.0	24	15	9	*	10	*9	*	*6	*5	*7	*6
83.09	16	*9	*7	*	*6	*	*	*	*	*5	*
83.1	26	12	14	*7	*7	*6	*6	*7	*6	*7	*6
83.14	14	*7	*7	*	*6	*	*	*	*	*	*
83.2	13	*7	*6	*	*	*	*	*	*	*5	*
83.21	13	*7	*6	*	*	*	*	*	*	*5	*
83.3	15	*6	9	*	*	*7	*6	*	*	*8	*
83.39	12	*	*7	*	*	*5	*	*	*	*6	*
83.4	31	19	12	*	11	*8	10	*8	*7	11	*5
83.45	19	12	*7	*	*8	*	*7	*6	*	*7	*
83.6	72	35	37	*	18	26	28	13	15	32	12
83.63	48	21	27	—	*	19	24	*8	11	23	*6
83.64	11	*8	*	*	*6	*	*	*	*	*	*
83.65	12	*6	*7	*	*7	*	*	*	*	*5	*
83.7	*7	*	*	*	*	*	*	*	*	*	*
83.8	33	20	12	*7	*8	9	*8	9	*7	11	*5
83.82	*7	*	*	*	*	*	*	*	*	*	*
83.85	10	*6	*	*6	*	*	*	*	*	*	*
83.88	10	*6	*	*	*	*	*	*	*	*	*
83.9	10	*	*5	*	*	*	*	*	*	*	*
84	168	102	67	*	23	53	90	43	36	68	21
84.0	10	*8	*	*	*6	*	*	*	*	*	*
84.01	9	*7	*	*	*5	*	*	*	*	*	*
84.1	137	78	59	*	12	43	81	36	30	53	18
84.11	46	28	17	*	*7	15	24	16	*8	16	*5
84.12	16	11	*5	*	*	*7	*7	*5	*	*	*
84.15	38	22	16	*	*	12	22	*8	9	15	*6
84.17	36	16	20	*	*	*9	28	*7	10	17	*
84.3	19	15	*	*	*	*7	*8	*	*	10	*
85-86	1,264	567	697	124	355	378	407	312	274	506	172
85	207	*	204	*	44	83	79	63	42	61	41
85.0	*6	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
85.1	10	—	10	*	*	*	*7	*	*	*	*
85.2	38	*	38	*	*	15	19	15	*6	10	*8
85.21	27	*	27	*	*	12	13	12	*	*6	*5
85.23	*8	—	*8	—	*	*	*	*	*	*	*
85.3	19	*	19	*	14	*	*	9	*	*	*
85.32	17	*	17	*	13	*	*	*8	*	*	*
85.4	93	*	92	—	10	37	46	20	24	28	20
85.41	21	*	20	—	*	10	*8	*	*5	*6	*6
85.43	65	*	64	—	*7	23	35	15	18	20	12
85.5	*5	—	*5	—	*	*	*	*	*	*	*
85.7	9	*	9	—	*	*6	*	*	*	*	*
85.8	*9	*	*8	—	*	*	*	*	*	*	*
85.9	14	*	14	*	*	*8	*	*6	*	*	*
85.95	*8	—	*8	—	*	*	*	*	*	*	*
86	1,057	564	493	123	312	295	327	249	232	445	131
86.0	276	134	142	29	80	88	80	69	57	110	40
86.01	14	*7	*7	*	*	*	*	*8	*	*	*
86.04	108	59	49	9	37	33	28	23	27	39	20
86.05	33	16	17	*8	*8	*9	*9	10	*7	13	*
86.06	11	*	*6	*	*	*	*	*	*	*6	*
86.07	81	31	50	*7	16	31	27	21	13	38	*9
86.09	27	17	10	*	12	*6	*7	*5	*6	12	*
86.1	20	*8	12	*	*	*8	*6	*6	*	*7	*
86.11	19	*7	12	*	*	*8	*6	*6	*	*7	*
86.2	373	210	164	19	89	113	151	93	97	139	45
86.22	260	150	111	14	70	83	94	62	63	103	32
86.23	*7	*	*	*	*	*	*	*	*	*	*
86.27	36	16	21	—	*	*6	28	17	*9	*8	*
86.28	65	40	25	*	13	22	26	12	22	22	*8
86.3	73	34	39	12	22	18	21	17	13	35	*8
86.4	*6	*	*	*	*	*	*	*	*	*	*
86.5	135	79	56	*8	60	28	39	33	36	42	24
86.59	135	79	56	*8	60	28	39	33	36	42	24
86.6	105	64	42	42	32	16	15	16	14	70	*6
86.62	*6	*	*	*	*	*	*	*	*	*	*
86.69	82	50	32	38	22	12	10	11	10	58	*
86.7	31	19	12	*	*8	*9	10	*7	*5	16	*
86.70	*7	*	*	*	*	*	*	*	*	*6	*
86.74	17	10	*6	*	*	*6	*7	*	*	*7	*
86.8	33	11	22	*5	12	13	*	*6	*5	18	*
86.83	14	*	13	*	*6	*7	*	*	*	*5	*
86.84	*9	*	*	*	*	*	*	*	*	*7	*
86.89	*6	*	*	*	*	*	*	*	*	*	*
87-99	12,223	5,841	6,382	837	2,349	3,226	5,812	3,712	2,442	3,791	2,278
87	1,072	466	606	67	197	266	543	267	238	329	238
87.0	349	161	188	19	54	75	201	157	52	67	73
87.03	346	158	187	18	53	75	200	157	51	67	71
87.2	26	15	11	*	*7	*7	11	*	*9	10	*6
87.21	14	*7	*7	*	*	*	*	*	*	*7	*
87.4	354	149	205	31	47	79	197	42	107	130	75
87.41	92	45	47	*	10	25	53	40	13	16	22
87.44	255	101	154	27	35	52	142	*	92	111	52
87.49	*6	*	*	*	*	*	*	*	*	*	*
87.5	133	46	87	*	40	44	48	20	27	55	31
87.51	10	*6	*	*	*	*	*7	*	*	*	*
87.53	113	36	78	*	36	37	38	14	23	51	25
87.54	*6	*	*	*	*	*	*	*	*	*	*
87.6	58	23	35	*7	*7	14	30	12	*7	10	29
87.61	15	*6	*8	*	*	*	*9	*	*	*5	*5
87.62	20	*7	12	*	*	*	10	*	*	*	10

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
87.63	12	*5	*6	*	*	*	*6	*	*	*	*7
87.64	*9	*	*6	*	*	*	*5	*	*	*	*5
87.7	143	70	73	*7	40	43	53	32	34	54	24
87.71	*5	*	*	—	*	*	*	*	*	*	*
87.73	29	13	15	*	11	11	*6	*5	*6	10	*8
87.74	86	40	46	*	20	28	35	20	21	36	*8
87.77	*6	*	*	*	*	*	*	*	*	*	*
87.79	*8	*	*	*	*	*	*	*	*	*	*
88	3,539	1,823	1,716	90	469	1,219	1,761	967	767	1,222	583
88.0	235	106	129	11	64	64	97	103	30	48	54
88.01	232	104	128	11	63	62	96	102	30	47	53
88.1	38	*9	29	*	*6	*8	20	*	16	13	*8
88.19	37	*9	28	*	*6	*8	19	*	16	13	*8
88.2	29	15	15	*	*	*8	17	*	*9	11	*7
88.26	*9	*	*	*	*	*	*7	*	*	*	*
88.27	10	*6	*	*	*	*	*	*	*5	*	*
88.3	91	38	52	*9	23	22	37	38	17	24	12
88.38	80	34	46	*7	21	20	32	37	14	18	10
88.39	*7	*	*6	—	*	*	*	—	*	*	*
88.4	421	222	200	*	46	132	239	83	104	183	52
88.40	*8	*	*	—	*	*	*	*	*	*	*
88.41	120	58	62	*	19	38	61	25	24	56	14
88.42	129	70	59	*	*8	42	77	20	37	56	16
88.43	11	*	*7	*	*	*	*	*	*	*	*
88.45	24	14	10	*	*	*8	14	*	*7	10	*
88.47	20	*9	11	*	*	*7	10	*	*	10	*
88.48	82	50	32	*	*5	22	54	17	22	33	10
88.49	25	13	12	*	*6	*7	12	*7	*	11	*
88.5	1,583	935	648	*6	129	689	759	326	395	643	219
88.50	*7	*	*	*	*	*	*	*6	*	*	*
88.51	11	*5	*6	*	*	*	*5	*5	*	*	*
88.53	621	362	259	*	54	278	288	133	151	263	73
88.54	36	22	14	*	*	14	15	*	18	*9	*
88.55	17	10	*6	—	*	*7	*9	*	*6	*6	*
88.56	825	489	336	*	63	356	404	156	201	333	135
88.57	63	40	23	*	*	26	32	17	12	27	*6
88.6	23	13	10	*	*	*7	11	10	*	*5	*
88.67	10	*5	*	*	*	*	*	*	*	*	*
88.7	886	385	501	37	143	232	474	316	146	232	193
88.71	74	30	44	*	*5	15	50	37	*6	11	19
88.72	497	230	268	14	47	135	302	170	70	165	92
88.74	20	*7	13	*	*5	*6	*6	*	*6	*7	*
88.75	51	26	25	*6	10	12	23	19	*9	9	14
88.76	108	43	64	*6	30	33	38	36	21	16	36
88.77	80	35	45	*	10	23	46	31	20	13	17
88.78	25	—	25	—	25	*	*	*6	*7	*	*8
88.79	27	13	15	*	10	*8	*7	11	*6	*7	*
88.9	231	100	131	18	49	59	105	87	46	63	35
88.91	151	67	84	13	25	41	72	52	29	47	23
88.93	41	18	24	*	15	*9	16	17	12	*6	*7
88.94	12	*	*7	*	*	*	*6	*6	*	*	*
88.97	21	*8	13	*	*	*7	*8	10	*	*6	*
89	944	436	508	57	113	255	519	282	205	232	225
89.0	50	26	24	*	*8	15	27	20	*	*	28
89.04	18	*8	10	*	*	*	11	18	—	—	—
89.06	13	*7	*7	*	*	*5	*6	—	—	*	13
89.07	*8	*	*	—	*	*	*	—	—	—	*8
89.09	*7	*5	*	—	*	*	*	—	*	—	*7
89.1	99	51	48	31	15	17	35	43	10	23	22
89.14	73	39	34	19	*9	13	33	29	*7	18	19

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
89.19	15	*6	*8	*7	*	*	*	*9	*	*	*
89.2	27	*9	18	*	*6	*	16	11	*	9	*
89.22	11	*6	*	—	*	*	*9	*7	*	*	—
89.26	*9	—	*9	*	*	*	*	*	*	*	*
89.3	49	20	30	13	*7	10	19	9	13	16	11
89.37	18	*7	11	*	*	*7	*6	*	*	*6	*
89.38	*6	*	*	*	*	*	*	*	*	*	*
89.39	21	*8	13	*7	*	*	11	*	*6	9	*
89.4	143	68	74	*	19	58	65	52	20	38	32
89.41	45	24	21	*	*8	21	16	19	*5	*9	13
89.44	95	42	52	*	11	38	46	32	14	29	19
89.5	434	191	243	*6	44	111	274	111	121	99	103
89.50	*9	*	*	*	*	*	*6	*6	*	*	*
89.52	275	114	161	*	29	69	175	31	108	72	65
89.54	131	63	68	*	13	34	80	69	9	19	34
89.59	15	*7	*8	*	*	*	11	*5	*	*6	*
89.6	141	71	70	*	14	39	84	35	36	45	25
89.61	10	*6	*	*	*	*	*5	*	*	*	*
89.62	11	*6	*5	*	*	*	*7	*	*	*	*
89.64	49	25	23	—	*	15	32	10	18	13	*7
89.65	61	30	31	*	*9	15	36	17	*8	26	10
89.68	*7	*	*	—	*	*	*	*	*	*	*
90	16	*7	9	*	*	*6	*7	*	*8	*7	*
90.5	*6	*	*	*	*	*	*	*	*	*	—
90.8	*8	*	*5	—	*	*	*	*	*6	*	—
90.81	*8	*	*5	—	*	*	*	*	*6	*	—
91	*6	*	*	*	*	*	*	*	*	*	*
92	279	123	156	*7	36	88	148	105	40	79	54
92.0	106	46	60	*	12	35	57	40	16	28	22
92.02	*8	*	*5	*	*	*	*	*	*	*	*
92.03	*6	*	*	*	*	*	*	*	*	*	*
92.04	*8	*	*	*	*	*	*	*	*	*	*
92.05	83	38	46	*	*7	30	46	30	13	24	16
92.1	76	30	46	*	10	20	44	35	*8	16	17
92.14	24	12	12	*	*	*6	13	10	*	*7	*6
92.15	44	15	29	*	*	11	27	22	*6	*8	*8
92.2	88	43	45	*	13	29	45	28	15	34	11
92.24	22	9	12	*	*	*8	9	10	*	*5	*
92.27	14	*6	*8	—	*	*	*7	*6	*	*5	*
92.29	44	24	20	*	*	14	25	10	*8	20	*6
92.3	*8	*	*	—	*	*	*	*	*	*	*
92.32	*5	*	*	—	*	*	*	*	*	*	*
93	1,478	621	858	182	177	264	855	462	217	445	355
93.0	56	21	35	*	*9	12	32	25	10	*5	15
93.01	50	19	31	*	*7	11	31	24	*9	*	15
93.1	25	*8	17	*	*	*6	16	23	*	*	*
93.11	*7	*	*	*	*	*	*5	*7	—	—	*
93.12	12	*	*8	*	*	*	*8	12	—	—	—
93.2	40	18	22	*	*7	10	22	28	*	*	*6
93.22	28	13	15	*	*	*6	17	21	*	*	*6
93.24	*5	*	*	*	*	*	*	*	*	*	—
93.3	385	134	252	*7	29	60	290	92	51	171	71
93.32	*7	*	*	*	*	*	*	*	*	*	*
93.38	16	*6	10	*	*	*	12	*	*	*7	*7
93.39	357	121	236	*6	25	54	272	86	48	161	62
93.4	11	*6	*6	*	*	*	*	*5	*	*	*
93.46	*6	*	*	*	*	*	*	*	*	*	*
93.5	90	48	43	16	26	20	29	24	24	25	18
93.53	18	*9	9	*	*	*	*6	*	*9	*	*
93.54	19	*9	10	*	*6	*	*7	*	*6	*5	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
93.55	*5	*	*	*	*	*	*	*	*	*	*
93.57	19	13	*6	*6	*6	*	*	*5	*6	*6	*
93.59	25	11	15	*	*7	*7	11	*8	*	*9	*7
93.7	33	14	20	*	*	*8	22	*6	10	11	*7
93.75	32	14	19	*	*	*8	21	*6	10	11	*7
93.8	395	151	244	10	51	63	271	114	59	164	58
93.81	26	11	15	*	*6	*5	13	*6	10	*6	*
93.83	271	97	174	*	35	45	187	82	39	113	37
93.89	97	43	55	*	9	13	71	26	9	46	17
93.9	438	220	217	141	47	81	169	145	58	59	176
93.90	60	26	33	*8	*	17	31	20	17	12	10
93.94	193	100	92	82	22	29	60	74	15	26	78
93.96	156	80	76	47	17	29	63	40	22	16	78
93.99	20	*8	12	*	*	*	*9	10	*	*	*6
94	809	467	342	40	528	179	62	451	156	166	35
94.0	*9	*5	*	*	*6	*	*	*	*	*6	*
94.2	203	101	103	*9	121	48	26	145	20	35	*
94.22	*6	*	*	*	*	*	*	*5	*	*	—
94.23	36	18	18	*	22	*7	*	35	—	*	*
94.25	113	61	52	*6	77	25	*6	97	11	*	*
94.27	48	18	30	—	19	14	14	*8	*9	28	*
94.3	114	54	60	11	72	21	9	69	25	19	—
94.37	*8	*6	*	*	*	—	—	*8	—	—	—
94.38	44	19	26	*	30	11	*	44	—	*	—
94.39	55	28	28	*7	34	10	*	12	25	18	—
94.4	141	73	69	16	90	23	13	83	30	28	*
94.42	13	*7	*7	*	*7	*	*	*8	*5	*	—
94.44	118	58	60	12	74	21	12	65	25	28	*
94.46	*6	*	*	—	*	*	*	*6	—	—	—
94.6	338	233	106	*	238	86	13	151	81	77	30
94.61	14	10	*	*	9	*	*	*7	*	*	*
94.62	149	111	38	*	91	48	9	56	44	34	15
94.63	19	14	*	—	10	*7	*	*	*9	*	*
94.64	11	*8	*	*	10	*	*	*6	*	*	*
94.65	64	35	29	*	50	14	*	37	*8	17	*
94.66	*6	*	*	—	*	*	*	*	*	*	*
94.67	*8	*	*	*	*7	*	*	*	*	*	*
94.68	60	43	17	*	50	10	*	43	*6	10	*
94.69	*7	*	*	*	*6	*	*	*	*	*	*
95	*6	*	*	*	*	*	*	*	*	*	*
96	1,335	642	693	126	272	295	642	369	265	427	275
96.0	503	255	247	45	72	125	261	130	102	167	103
96.04	429	221	209	40	61	103	226	114	83	149	83
96.07	63	30	34	*5	10	19	30	13	17	15	18
96.2	*5	*	*	*	*	*	*	*	*	*	*
96.3	21	*7	14	*	9	*	*7	*7	*	*7	*
96.33	11	*	*7	*	*8	*	*	*	*	*	*
96.38	*7	*	*	*	*	*	*	*	*	*	*
96.4	86	*	83	*	82	*	*	10	17	25	35
96.49	82	*	82	*	81	*	*	10	15	24	33
96.5	49	26	23	*5	14	9	20	11	13	17	*8
96.56	26	16	9	*	*6	*5	11	*	*	10	*7
96.59	17	9	*8	*	*6	*	*6	*	*7	*6	*
96.6	115	62	53	*7	10	21	77	60	18	23	14
96.7	553	287	266	65	83	134	272	148	108	186	110
96.70	23	*9	14	*	*	9	9	*8	*	9	*
96.71	356	185	171	41	53	83	178	91	71	120	73
96.72	174	93	81	23	26	41	84	49	33	56	36
97	97	47	49	13	21	22	40	24	15	41	16
97.0	33	18	15	*	*	*7	20	*8	*	16	*

See footnotes at end of table.

Table 46. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2000—Con.

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see Methods section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
97.02	23	13	10	*	*	*	14	*	*	13	*
97.05	*7	*	*	*	*	*	*	*	*	*	*
97.2	*8	*	*	*	*	*	*	*	*	*	*
97.23	*7	*	*	*	*	*	*	*	*	*	*
97.4	11	*5	*5	*	*	*	*	*	*	*	*
97.49	*9	*	*	*	*	*	*	*	*	*	*
97.5	*7	*	*	*	*	*	*	*	*	*	*
97.6	17	*6	10	*	*6	*	*7	*	*	*9	*
97.62	10	*	*7	*	*	*	*	*	*	*5	*
97.8	14	*7	*6	*	*	*	*	*	*	*	*
98	28	17	10	*	*6	*7	10	*8	*	10	*6
98.0	10	*7	*	*	*	*	*5	*	*	*	*
98.02	*8	*5	*	*	*	*	*	*	*	*	*
98.5	*9	*7	*	—	*	*	*	*	*	*	*
98.51	*9	*7	*	—	*	*	*	*	*	*	*
99	2,615	1,185	1,430	247	526	623	1,219	772	526	827	490
99.0	1,157	502	655	48	148	267	695	230	237	466	223
99.00	10	*5	*	*	*	*	*5	*	*	*	*
99.02	50	20	30	*	*	16	29	10	12	15	13
99.03	20	10	10	*	*	*	14	10	*	*7	*
99.04	932	387	545	31	119	203	579	182	186	382	181
99.05	60	32	28	*9	*9	19	23	11	11	28	10
99.07	75	43	32	*	10	21	40	12	20	29	14
99.1	359	152	207	29	118	81	131	145	69	79	66
99.10	63	41	22	*	*8	22	34	19	12	20	11
99.11	68	*	68	*	67	*	*	14	16	22	16
99.14	*8	*	*	*	*	*	*	*	*	*	*
99.15	106	55	51	19	16	27	44	43	23	22	19
99.17	28	14	15	*	*9	*8	9	16	*5	*	*5
99.18	39	14	25	*	10	*7	17	30	*	*7	*
99.19	46	24	22	*	*7	15	24	20	10	*	12
99.2	851	405	446	120	217	227	287	333	167	207	144
99.20	82	57	25	—	*	38	39	24	26	18	14
99.21	244	102	142	36	81	46	81	110	48	43	44
99.23	77	30	48	*7	14	19	38	29	21	18	9
99.25	199	115	85	48	37	68	46	59	28	77	35
99.29	237	98	139	28	76	52	82	109	43	47	38
99.4	20	*	20	*	20	*	—	*	*	*8	*6
99.47	15	—	15	—	15	—	—	*	*	*6	*
99.5	*7	*	*	*	*	*	*	*	*	*	*
99.6	140	79	60	*	9	34	94	41	36	37	25
99.60	43	25	18	*	*	10	29	12	10	11	10
99.61	29	17	12	*	*	*6	20	*7	*7	*9	*6
99.62	55	30	25	*	*	15	36	17	15	16	*7
99.69	13	*8	*	*	*	*	*8	*	*	*	*
99.7	14	*6	*8	*	*	*	*	*	*	*	*
99.71	*8	*	*	*	*	*	*	*	*	*	*
99.8	44	25	19	40	*	*	*	*6	*	14	19
99.83	37	20	16	37	*	*	—	*6	*	10	17
99.9	18	10	*9	*	*	*	*6	*	*	10	*
99.99	18	10	*9	*	*	*	*6	*	*	10	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see Methods section.

Appendix

Technical Notes

Abstract Form

Figure I shows the medical abstract form used for the 2000 NHDS. The hospital number, medical record number, date of birth, and patient ZIP code are confidential information and are not released to the public.

Population Estimates

Population estimates used to compute rates are shown in table I. These estimates are for the civilian population of the United States, including institutionalized persons, as of July 1, 2000. The data are based on the 1990 Census and are from unpublished tabulations provided by the U.S. Census Bureau. Population estimates from the 2000 Census were not available for the civilian population when this report was prepared. NCHS will provide 2000-based estimates of the civilian population on request when they are received. The 2001 NHDS Annual Summary will include a comparison of populations based on 1990 and 2000 Censuses.

The population estimates in table I have been adjusted for the net underenumeration using the 1990 National Population Adjustment Matrix. Population data provided in annual summaries of 1990–96 NHDS data were not adjusted for the underenumeration.

Reliability of Estimates

The statistics presented in this report are based on a sample survey, and therefore may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from the NHDS were calculated with SUDAAN software, which takes into account the complex sample design. A description of

Table I. Civilian population by selected characteristics: United States, 2000

Age and geographic region	Both sexes	Male	Female
Number in thousands			
All ages			
Total	278,099	135,986	142,113
Region:			
Northeast	52,420	25,388	27,032
Midwest	63,877	31,208	32,668
South	98,781	47,924	50,857
West	63,021	31,465	31,556
Under 15 years			
Total	60,493	30,943	29,549
Under 1 year	3,994	2,040	1,955
1–4 years	15,678	8,012	7,666
5–14 years	40,820	20,892	19,929
Region:			
Northeast	10,716	5,483	5,233
Midwest	13,745	7,035	6,710
South	21,604	11,039	10,564
West	14,427	7,386	7,041
15–44 years			
Total	122,186	61,086	61,100
15–24 years	39,025	19,831	19,193
25–34 years	38,046	18,855	19,191
35–44 years	45,115	22,399	22,716
Region:			
Northeast	22,704	11,289	11,414
Midwest	27,917	13,961	13,956
South	43,166	21,312	21,854
West	28,399	14,523	13,876
45–64 years			
Total	60,944	29,567	31,377
45–54 years	37,162	18,198	18,964
55–64 years	23,783	11,370	12,413
Region:			
Northeast	11,751	5,652	6,098
Midwest	14,084	6,856	7,229
South	21,870	10,521	11,349
West	13,239	6,539	6,700
65 years and over			
Total	34,476	14,389	20,087
65–74 years	17,993	8,153	9,840
75–84 years	12,214	4,945	7,269
85 years and over	4,269	1,290	2,979
Region:			
Northeast	7,249	2,963	4,286
Midwest	8,130	3,356	4,773
South	12,141	5,052	7,089
West	6,956	3,017	3,938

NOTE: Population estimates have been adjusted for net underenumeration using the 1990 National Population Adjustment Matrix. Estimates are consistent with annual population estimates from the Bureau of the Census that are based on the 1990 census.

the software and the approach it uses has been published (19).

The chances are about 32 in 100 that an estimate from the sample would differ from a complete census by more than the standard error. The chances are

5 in 100 that the difference would be more than twice the standard error and about 1 in 100 that the difference would be more than 2.5 times as large as the standard error. The relative standard error (RSE) expresses the standard error

Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics, 2000

Characteristic	Discharges or first-listed diagnoses		Days of care		All-listed diagnoses		All-listed procedures	
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>
Total	0.002287	307.96962	0.003503	824.08156	0.002846	279.63119	0.003840	309.28733
Sex								
Male	0.002492	330.84852	0.004031	1,524.79149	0.004294	453.83201	0.004677	284.75142
Female	0.002296	300.21724	0.003590	758.48227	0.002176	277.66480	0.002855	296.90783
Age group								
Under 15 years	0.021332	284.11415	0.035212	520.36022	0.023027	249.07632	0.031270	242.65411
15–44 years	0.002306	309.63184	0.003788	846.80014	0.002161	303.96264	0.003001	307.89369
45–64 years	0.002495	294.03933	0.003990	1,245.18684	0.003046	312.39495	0.004791	287.43793
65 years and over	0.003283	334.40841	0.003590	2,018.69120	0.003000	362.13321	0.004734	311.25612
Geographic region								
Northeast	0.013516	176.21901	0.017328	581.83012	0.018498	189.26179	0.017236	293.50715
Midwest	0.009542	213.25373	0.012172	428.23450	0.010369	231.86476	0.013317	120.09107
South	0.003554	363.25616	0.006166	1,223.94608	0.004845	352.50429	0.005879	261.57132
West	0.005469	351.03280	0.010168	1,046.78781	0.005785	346.80421	0.008474	316.77742
Principal expected source of payment								
Medicare	0.003730	347.59293	0.004077	1,885.28816	0.003649	315.73286	0.004973	325.03095
Medicaid	0.006329	323.72265	0.009505	1,113.96754	0.005271	363.82164	0.007488	305.73437
Worker's compensation and other government payment	0.010432	337.34361	0.022387	1,100.21733	0.017122	259.22962	0.015112	274.59336
HMO/PPO ¹	0.004596	267.09076	0.008327	467.29188	0.004956	235.70957	0.005690	231.02811
Blue Cross/Blue Shield and other private insurance	0.005157	295.04612	0.007509	828.24439	0.005760	296.04960	0.009506	334.25082
Self pay	0.004822	314.42800	0.007746	1,005.64303	0.004554	312.44459	0.006655	260.43225
No charge and other	0.038141	171.15139	0.051189	506.79249	0.037292	200.24927	0.043448	144.74118

¹HMO/PPO is health maintenance organization and preferred provider organization.

NOTE: The relative standard error (RSE) can be determined from the equation $RSE(x) = \sqrt{a + b/X}$.

as a percent of the estimate and can be multiplied by the estimate to obtain the standard error.

To derive approximate standard errors for the estimates in this report, first calculate the approximate RSE of the estimate using the parameters shown in table II. Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate *X*, may be estimated from the formula:

$$RSE(X) = \sqrt{a + b/X}$$

where *a* and *b* are defined in table II. When multiplied by 100, RSE(*X*) is expressed as a percent of *X*.

The approximate RSE for a percent, 100*p*, in which the numerator is a subclass of the denominator, may be calculated directly using the formula:

$$RSE(p) = \sqrt{b(1-p) / (pX)}$$

where 100*p* is the percent of interest, *X* is the denominator of the percent, and *b* is the parameter *b* in the formula for

approximating the RSE(*X*). The values for *b* are given in table II. When multiplied by 100, RSE(*p*) is expressed as a percent of *p*. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (20,21).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained by the method described above using figures from table II), if the denominator is an estimate produced by the U.S. Bureau of the Census for the total population or one or more of the age-sex groups of the total population.

The approximate RSE of a ratio (*X/Y*) in which the numerator (*X*) and the denominator (*Y*) are both estimated from the same survey, but the numerator is not a subclass of the denominator, is calculated using the formula:

$$RSE(X/Y) = \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

When multiplied by 100, the RSE(*X/Y*) is expressed as a percent of the ratio *X/Y*. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of both the numerator and denominator are both less than 10 percent (20,21).

Definitions of Selected Terms

Age—Patient's age at the birthday prior to admission to the hospital.

Average length of stay—The mean length of stay for discharges. It is calculated by dividing the number of days of care by the number of discharges.

Bed size of hospital—The number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for patients, not including bassinets for newborn infants. In this report, the classification of hospitals by bed size is based on the number of beds at or near midyear as reported by the hospital.

Days of care—The total number of days a patient spent in the hospital. A stay of less than 1 day (patient admission and discharge on the same day) is counted as 1 day. For patients admitted and discharged on different days, the number of days of care is computed by counting all the days from (and including) the day of admission to (but not including) the day of discharge.

Death rate—Calculated by dividing the number of deaths for a category by the total number of discharges in that category and multiplying by 100. Discharges whose status is unknown (0.7 percent) were assumed to be discharged alive.

Diagnosis—A disease or injury (or other factor that influences a patient's health status or causes contact with health services) that is listed on the medical record of the patient. The diagnosis established after study to be chiefly responsible for the admission of the patient to the hospital is the principal diagnosis. The first-listed diagnosis is the one specified as the principal diagnosis on the face sheet or discharge summary of the medical record. However, if the principal diagnosis is not specified, the first one listed on the face sheet or discharge summary of the medical record is used. The number of first-listed diagnoses is the same as the number of discharges. Data for all-listed diagnoses include first-listed diagnoses and other diagnoses appearing on the medical record. A minimum of one and a maximum of seven diagnoses are coded per discharge in the NHDS. The HIV data (table 22) are for discharges with an HIV diagnosis listed in any position among the coded diagnoses.

Discharge—A completed inpatient hospitalization. A hospitalization may be completed by death or by releasing the patient to the customary place of residence, a nursing home, another hospital, or other location.

Hospital—Hospitals with an average length of stay for all inpatients of less than 30 days (short stay) or those whose speciality is general (medical or surgical) or children's general. Federal hospitals, hospital units of institutions,

and hospitals with less than six beds staffed for patient use are excluded.

Newborn infant—Inpatients admitted to a hospital by birth. Sick newborn infants are those who had at least one of the ICD-9-CM (1) diagnostic codes 001-999.

Ownership of hospital—The type of ownership is determined by the organization that controls and operates the hospital. Hospitals are grouped as follows:

Proprietary—Hospitals operated by individuals, partnerships, or corporations for profit.

Government—Hospitals operated by State or local governments.

Nonprofit—Hospitals operated by a nonprofit organization or church.

Principal expected source of payment—The primary expected method of payment for the hospitalization. Types of payment are as follows:

Medicare—The health insurance program for the aged and disabled administered by the Centers for Medicare and Medicaid Services (formerly the Health Care Financing Administration).

Medicaid—A jointly funded Federal-State health insurance program providing medical care to those unable to afford it.

Worker's compensation—A state or municipal disability insurance or industrial accident insurance.

Other government—This includes payments under the Title V Program, the State-funded Maternal and Child Health Program that provides coverage for children up to 21 years of age with specific chronic medical conditions (e.g., Crippled Children's Programs), CHAMPUS (Civilian Health and Medical Program of the Uniformed Medical Services) or TRICARE (the Defense Department's health care program for service families), casualty insurance paid by the State, vocational rehabilitation, Federal or State research grant (medical research), or legal hold (prisoner in medical detention).

HMO/PPO—Any Health Maintenance Organization (HMO) or Preferred Provider Organization (PPO) sponsored by consumers, communities, physicians, or hospitals.

Blue Cross/Blue Shield and other private—A private insurance plan not specified as an HMO/PPO. This includes Blue Cross/Blue Shield plans, medical coverage provided by life insurance companies, casualty insurance companies, health insurance companies, and independent plans such as employer/union-sponsored plans and/or self-funded plans (partial or total).

Self-pay—The majority of the costs for the hospitalization were expected to be paid by the patient, spouse, family, or next-of-kin.

No charge—Patients admitted with the understanding that the medical services would be free. This includes hospital-sponsored special research or "teaching" patients.

Other—Includes any nonprofit source of payment (such as church welfare, United Way (United Appeal), or Shriner's Hospitals for Children).

Procedure—Surgical procedures (such as appendectomy), diagnostic procedures (such as spinal tap), and therapeutic treatments (such as injection or infusion of cancer chemotherapeutic substance) reported on a patient's medical record. Data are for all-listed procedures, which include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded per discharge in the NHDS. The procedures classified as nonsurgical in this report are listed in table III by ICD-9-CM code numbers. All other procedures were considered surgical. This classification was begun in 1979 for the presentation of NHDS data. The distinction between surgical and nonsurgical procedures has become less meaningful in the last 2 decades with the development of minimally invasive and noninvasive procedures. The procedures classified as nonsurgical may

Table III. Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey by ICD-9-CM category, 2000

[Procedure categories and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Nonsurgical codes
Operations on the nervous system 01-05	01.18-01.19,03.31,03.39,04.19,05.19
Operations on the endocrine system 06-07	06.19,07.19
Operations on the eye 08-16	08.19,09.19,09.41-09.49,10.29,11.29,12.29,14.19,15.09,16.21,16.29
Operations on the ear 18-20	18.01,18.11,18.19,20.31,20.39
Operations on the nose, mouth, and pharynx 21-29	21.00-21.02,21.21,21.29,22.19,24.19,25.09,26.19,27.29,28.19,29.11,29.19
Operations on the respiratory system 30-34	31.41-31.42,31.48-31.49,33.21-33.23,33.29,34.21-34.22,34.28-34.29
Operations on the cardiovascular system 35-39	37.26-37.27,37.29,38.29,39.95
Operations on the hemic and lymphatic system 40-41	40.19,41.38-41.39
Operations on the digestive system 42-54	42.22-42.23,42.29,44.11-44.13,44.19,45.11-45.13,45.19,45.21-45.24, 45.28-45.29,48.21-48.23,48.29,49.21,49.29,50.19,51.10-51.11, 51.19,52.19,54.21,54.29
Operations on the urinary system 55-59	55.21-55.22,55.29,56.31,56.35,56.39,57.31-57.32,57.39,57.94-57.95, 58.21-58.22,58.29,59.29
Operations on the male genital organs 60-64	60.18-60.19,61.19,62.19,63.09,64.19,64.94
Operations on the female genital organs 65-71	65.19,66.19,67.19,68.11,68.19,69.92,70.21-70.22,70.29,71.19
Obstetrical procedures 72-75	73.4-73.59,73.91-73.92,75.31-75.32,75.34-75.35,75.94
Operations on the musculoskeletal system 76-84	76.19,78.80-78.89,80.20-80.29,81.98,83.29,84.41-84.43,84.45-84.47
Operations on the integumentary system 85-86	85.19,86.19,86.92
Miscellaneous diagnostic and therapeutic procedures 87-99	87.01-99.99

have as much operative or anesthetic risk and require highly trained personnel or special equipment similar to those classified as surgical.

Region—Hospitals are classified by location in one of the four geographic regions of the United States that correspond to those used by the U.S. Bureau of the Census.

Region	States included
Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania
Midwest	Michigan, Ohio, Illinois, Indiana, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
South	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas
West	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada,

Washington, Oregon, California, Hawaii, and Alaska

Changes in ICD-9-CM Codes

Although the ICD-9-CM has been used for coding NHDS data since 1979, the classification system undergoes annual updating. Assignment of new diagnostic and procedure codes, fourth- and fifth-digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Centers for Medicare and Medicaid Services (formerly Health Care Financing Administration). Addenda to the ICD-9-CM have become effective October 1 of each year since 1986, except in 1999. No addendum was released in 1999 because of concerns about instituting coding changes just before the millennium crossover.

The data in this report were coded using the addenda for October 1, 1986-98. When hospitals reported data using code changes introduced in the October 1, 2000, addendum, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

Tables IV and V show the new ICD-9-CM codes from the 1986-1998 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned before development of the new code. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1).

Table IV. Changes in diagnosis codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81	1995	005.8	205.01	1991	V10.62
005.89	1995	005.8	205.10	1991	205.1
007.4	1997	007.8	205.11	1991	V10.62
008.00–008.09	1992	008.0	205.20	1991	205.2
008.43–008.47	1992	008.49	205.21	1991	V10.62
008.61–008.69	1992	008.6	205.30	1991	205.3
031.2	1997	031.8	205.31	1991	V10.62
038.10	1997	038.1	205.80	1991	205.8
038.11	1997	038.1	205.81	1991	V10.62
038.19	1997	038.1	205.90	1991	205.9
041.00–041.09	1992	041.0	205.91	1991	V10.62
041.04 (code title restated)	1997	041.04	206.00	1991	206.0
041.10–041.19	1992	041.1	206.01	1991	V10.63
041.81–041.85	1992	041.8	206.10	1991	206.1
041.86	1995	041.84	206.11	1991	V10.63
041.89	1992	041.8	206.20	1991	206.2
042	1994	042.0–042.2, 042.9, 043.0–043.3, 043.9, 044.0, 044.9 (codes deleted)	206.21	1991	V10.63
042.0–042.9	1986	279.19	206.80	1991	206.8
043.0–043.9	1986	279.19	206.81	1991	V10.63
044.0–044.9	1986	279.19	206.90	1991	206.9
070.20–070.21	1991	070.2	206.91	1991	V10.63
070.22	1994	070.20	207.00	1991	207.0
070.23	1994	070.21	207.01	1991	V10.69
070.30–070.31	1991	070.3	207.10	1991	207.1
070.32	1994	070.30	207.11	1991	V10.69
070.33	1994	070.31	207.20	1991	207.2
070.41–070.43	1991	070.4	207.21	1991	V10.69
070.44	1994	070.41	207.80	1991	207.8
070.49	1991	070.4	207.81	1991	V10.69
070.51–070.53	1991	070.5	208.00	1991	208.0
070.54	1994	070.51	208.01	1991	V10.60
070.59	1991	070.5	208.10	1991	208.1
077.98–077.99	1993	077.9	208.11	1991	V10.60
078.10–078.11, 078.19	1993	078.1	208.20	1991	208.2
078.88	1993	078.89	208.21	1991	V10.60
079.4	1993	079.8	208.80	1991	208.8
079.50–079.53, 079.59	1993	079.8	208.81	1991	V10.60
079.6	1996	79.89	208.90	1991	208.9
079.81	1995	079.89	208.91	1991	V10.60
079.88–079.89	1993	79.8	237.70–237.72	1990	237.7
079.98–079.99	1993	079.9	250.02	1993	250.90
088.81, 088.89	1989	088.8	250.03	1993	250.91
088.82	1993	088.89	250.12	1993	250.10
099.40–099.49	1992	099.4	250.13	1993	250.11
099.50–099.59	1992	078.89	250.22	1993	250.20
112.84–112.85	1992	112.89	250.23	1993	250.21
114.4–114.5	1993	114.3	250.32	1993	250.30
176.0–176.9	1991	173.0–173.9	250.33	1993	250.31
203.00	1991	203.0	250.42	1993	250.40
203.01	1991	V10.79	250.43	1993	250.41
203.10	1991	203.1	250.52	1993	250.50
203.11	1991	V10.79	250.53	1993	250.51
203.80	1991	203.8	250.62	1993	250.60
203.81	1991	V10.79	250.63	1993	250.61
204.00	1991	204.0	250.72	1993	250.70
204.01	1991	V10.61	250.73	1993	250.71
204.10	1991	204.1	250.82	1993	250.80
204.11	1991	V10.61	250.83	1993	250.81
204.20	1991	204.2	250.92	1993	250.90
204.21	1991	V10.61	250.93	1993	250.91
204.80	1991	204.8	275.40–275.42	1997	275.4
204.81	1991	V10.61	275.49	1997	275.4
204.90	1991	204.9	278.00–278.01	1995	278.0
204.91	1991	V10.61	283.10–283.11, 283.19	1993	283.1
205.00	1991	205.0	291.81	1996	291.8
			291.89	1996	291.8
			293.84	1996	293.89

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
300.82	1996	300.81	434.90–434.91	1993	434.9
305.1	1994	305.10, 305.11, 305.12, 305.13 (codes deleted)	435.3	1995	435.0 & 435.1
312.81–312.82, 312.89	1994	312.8	437.7	1992	780.9
315.32	1996	315.39	438.0	1997	294.9 & 438
320.81–320.89	1992	320.8	438.10	1997	784.5 & 438
333.92–333.93	1994	333.99	438.11	1997	784.3 & 438
337.20–337.22, 337.29	1993	337.9	438.12	1997	784.4 & 438
337.3	1998	337.9	438.19	1997	784.5 & 438
342.00–342.02	1994	342.0	438.20	1997	342.90 & 438
342.10–342.12	1994	342.1	438.21	1997	342.91 & 438
342.80–342.82	1994	342.9	438.22	1997	342.92 & 438
342.90–342.92	1994	342.9	438.30	1997	344.40 & 438
344.00–344.04, 344.09	1994	344.0	438.31	1997	344.41 & 438
344.30–344.32	1994	344.3	438.32	1997	344.42 & 438
344.40–344.42	1994	344.4	438.40	1997	344.30 & 438
344.81, 344.89	1993	344.8	438.41	1997	344.31 & 438
345.00–345.01	1989	345.0	438.42	1997	344.32 & 438
345.10–345.11	1989	345.1	438.50–438.52	1997	344.89 & 438
345.40–345.41	1989	345.4	438.53	1998	438.50
345.50–345.51	1989	345.5	438.81	1997	784.69 & 438
345.60–345.61	1989	345.6	438.82	1997	787.2 & 438
345.70–345.71	1989	345.7	438.89	1997	438
345.80–345.81	1989	345.8	438.9	1997	438
345.90–345.91	1989	345.9	440.20–440.22	1992	440.2
346.00–346.01	1992	346.0	440.23	1993	440.20 & 707.1 or 707.8 or 707.9
346.10–346.11	1992	346.1	440.24	1993	440.20 & 785.4
346.20–346.21	1992	346.2	440.29	1993	440.2
346.80–346.81	1992	346.8	440.30–440.32	1994	996.1
346.90–346.91	1992	346.9	441.00–441.03	1994	441.0
355.71	1993	354.4	441.6	1993	441.1 & 441.3
355.79	1993	355.7	441.7	1993	441.2 & 441.4
371.82	1992	371.89	446.20–446.21, 446.29	1990	446.2
374.87	1990	374.89	451.82–451.84	1993	451.89
403.00–403.01	1989	403.0	458.2	1995	997.9 & 458.9
403.10–403.11	1989	403.1	458.8	1997	458.9
403.90–403.91	1989	403.9	466.11	1996	466.1
404.00–404.03	1989	404.0	466.19	1996	466.1
404.10–404.13	1989	404.1	474.0 (code title restated)	1997	474.0
404.90–404.93	1989	404.9	474.00–474.02	1997	474.0
410.00–410.02	1989	410.0	482.30–482.39	1992	482.3
410.10–410.12	1989	410.1	482.4	1998	482.4
410.20–410.22	1989	410.2	482.41	1998	482.4
410.30–410.32	1989	410.3	482.49	1998	482.4
410.40–410.42	1989	410.4	482.81–482.89	1992	482.8
410.50–410.52	1989	410.5	482.84	1997	482.83
410.60–410.62	1989	410.6	483.0	1992	483
410.70–410.72	1989	410.7	483.1	1996	078.88 & 484.8
410.80–410.82	1989	410.8	483.8	1992	483
410.90–410.92	1989	410.9	491.20–491.21	1991	491.2
411.81	1989	410.9	493.20	1989	493.90
411.89	1989	411.8	493.21	1989	493.91
414.00–414.01	1994	414.0	512.1	1994	997.3
414.02–414.03	1994	996.03	518.6	1997	518.89
414.04–414.05	1996	414.00	518.81	1987	799.1
415.11	1995	997.3 & 415.1	518.82–518.89	1987	518.8
415.19	1995	415.1	518.83	1998	518.81
429.71	1989	410.0–410.9	518.84	1998	518.81
429.79	1989	410.0–410.9	519.00–519.02, 519.09	1998	519.0
433.00–433.01	1993	433.0	524.00–524.09	1992	524.0
433.10–433.11	1993	433.1	524.10–524.19	1992	524.1
433.20–433.21	1993	433.2	524.60–524.69	1991	524.6
433.30–433.31	1993	433.3	524.70–524.79	1992	524.8
433.80–433.81	1993	433.8	530.10–530.11, 530.19	1993	530.1
433.90–433.91	1993	433.9	530.81	1993	530.1
434.00–434.01	1993	434.0	530.82–530.84, 530.89	1993	530.8
434.10–434.11	1993	434.1	535.00–535.01	1991	535.0
			535.10–535.11	1991	535.1

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
535.20–535.21	1991	535.2	690.11	1995	691.8 & 704.8
535.30–535.31	1991	535.3	690.12	1995	691.8
535.40–535.41	1991	535.4	690.18	1995	690
535.50–535.51	1991	535.5	690.8	1995	690
535.60–535.61	1991	535.6	692.72–692.74	1992	692.79
536.3	1994	536.8	692.82–692.83	1992	692.89
536.40–536.42, 536.49	1998	997.4	702.0–702.8	1991	702
537.82	1990	537.89	702.11, 702.19	1994	702.1
537.83	1991	537.82	704.02	1993	704.09
556.0–556.6	1994	556	709.00–709.01, 709.09	1994	709.0
556.8–556.9	1994	556	710.5	1992	288.3, 729.1
562.02	1991	562.00	728.86	1995	729.4
562.03	1991	562.01	733.10–733.16, 733.19	1993	733.1
562.12	1991	562.10	738.10–738.19	1992	738.1
562.13	1991	562.11	747.60–747.64, 747.69	1993	747.6
564.81	1998	564.8	747.82	1993	747.89
564.89	1998	564.8	752.51–752.52	1996	752.5
569.60–569.61	1995	569.6	752.61–752.63	1996	752.6
569.62	1998	569.69	752.64–752.65	1996	752.8
569.69	1995	569.6	752.69	1996	752.8
569.84	1990	557.1	753.10–753.17, 753.19	1990	753.1
569.85	1991	569.84	753.20–753.23	1996	753.2
574.60	1996	574.00 & 574.30	753.29	1996	753.2
574.61	1996	574.01 & 574.31	756.70–756.71	1997	756.7
574.70	1996	574.10 & 574.40	756.79	1997	756.7
574.71	1996	574.11 & 574.41	758.81	1996	758.8
574.80	1996	574.00 & 574.10 574.30 & 574.40	758.89	1996	758.9
574.81	1996	574.01 & 574.11 574.31 & 574.41	759.81–759.82	1989	759.8
574.90	1996	574.20 & 574.50	759.83	1994	759.89
574.91	1996	574.21 & 574.51	759.89	1989	759.8
575.10–575.11	1996	575.1	760.75	1991	760.79
575.12	1996	575.0 & 575.1	760.76	1994	760.79
593.70–593.73	1994	593.7	763.81–763.83, 763.89	1998	763.8
596.51–596.53	1992	596.5	764.00–764.09	1988	764.0
596.54	1992	344.61	764.10–764.19	1988	764.1
596.55–596.59	1992	596.5	764.20–764.29	1988	764.2
599.81–599.89	1992	599.8	764.90–764.99	1988	764.9
645.0	1991	645	765.00–765.09	1988	765.0
651.30–651.31, 651.33	1989	651.00–651.01, 651.03	765.10–765.19	1988	765.1
651.40–651.41, 651.43	1989	651.10–651.11, 651.13	780.01–780.02	1992	780.0
651.50–651.51, 651.53	1989	651.20–651.21, 651.23	780.03	1993	780.01
651.60–651.61, 651.63	1989	651.80–651.81, 651.83	780.09	1992	780.0
654.20–654.21, 654.23	1990	654.2, 654.9	780.31	1997	780.3
654.90–654.94	1990	654.2, 654.9	780.39	1997	780.3
655.70 & 655.71	1997	655.8	780.57	1992	780.51, 780.53
655.73	1997	655.8	780.71	1998	780.7
657.0	1991	657	780.79	1998	780.7
659.60, 659.61, 659.63	1992	659.80–659.81, 659.83	781.8	1994	781.9
659.70, 659.71, 659.73	1998	656.30, 656.31, 656.33	786.03	1998	786.09
665.10, 665.11	1992	665.10, 665.11, 665.12, 665.14 ¹	786.04	1998	786.09
670.0	1991	670	786.05	1998	786.09
672.0	1991	672	786.06	1998	786.09
677	1994	No previous code assignment	786.07	1998	786.09
686.00 & 686.01	1997	686.0	787.01–787.03	1994	787.0
686.09	1997	686.0	787.91	1995	558.9
690.10	1995	690	787.99	1995	787.9
			788.20–788.21, 788.29	1993	788.2
			788.30–788.39	1992	788.3
			788.41–788.43	1993	788.4
			788.61–788.62, 788.69	1993	788.6
			789.00–789.07, 789.09	1994	789.0
			789.30–789.37, 789.39	1994	789.3
			789.40–789.47, 789.49	1994	789.4
			789.60–789.67, 789.69	1994	789.6
			790.91	1993	790.9
			790.92	1993	286.9
			790.93, 790.99	1993	790.9

¹The title for the subcategory 665.1 has been changed, making the fifth-digit subclassifications 665.12 and 665.14 invalid.

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
790.94	1997	790.99	V12.70–V12.72, V12.79	1994	V12.7
795.71	1994	785.8 (code deleted)	V13.00–V13.01, V13.09	1994	V13.0
795.79	1994	795.7	V13.61	1998	V13.6
795.8	1986	795.7	V13.69	1998	V13.6
796.5	1997	796.9	V15.41–V15.42	1996	V15.4
864.05	1992	864.09	V15.49	1996	V15.4
864.15	1992	864.19	V15.82	1994	305.13 (codes deleted)
909.5	1994	909.9	V15.84–V15.86	1995	V15.89
922.31–922.33	1996	922.3	V16.40–V16.43	1997	V16.4
925.1–925.2	1993	925	V16.49	1997	V16.4
959.0 (code title restated)	1997	959.0	V16.51	1998	V16.5
959.01	1997	854.00	V16.59	1998	V16.5
959.09	1997	959.0	V18.61	1998	V18.6
965.61	1998	965.6	V18.69	1998	V18.6
965.69	1998	965.6	V23.7	1989	V23.8
989.81–989.84	1995	989.8	V23.81	1998	V23.8
989.89	1995	989.8	V23.82	1998	V23.8
995.50–995.55	1996	995.5	V23.83	1998	V23.8
995.59	1996	995.5	V23.84	1998	V23.8
995.60–995.69	1993	995.0	V23.89	1998	V23.8
995.80	1996	995.81	V25.43	1992	V25.49
995.81 (code title restated)	1996	995.81	V25.5	1992	V25.8
995.82–995.85	1996	995.81	V26.51	1998	No previous code assignment
995.86	1998	995.89	V26.52	1998	No previous code assignment
996.04	1994	996.09	V28.6	1997	V28.8
996.51–996.59	1987	996.5	V29.0–V29.1, V29.8	1992	V71.8
996.55	1998	996.52	V29.2	1994	V29.8
996.56	1998	996.59	V29.3	1998	V29.8
996.60–996.69	1989	996.6	V29.8	1992	V71.8
996.68	1998	996.69	V29.9	1992	V71.9
996.70–996.79	1989	996.7	V30.00–V30.01	1989	V30.0
996.80–996.84	1987	996.8	V31.00–V31.01	1989	V31.0
996.85	1990	996.8	V32.00–V32.01	1989	V32.0
996.86	1987	996.8	V33.00–V33.01	1989	V33.0
996.89	1987	996.8	V34.00–V34.01	1989	V34.0
997.00–997.01	1995	997.0	V35.00–V35.01	1989	V35.0
997.02	1995	997.9 & 430–434, 436	V36.00–V36.01	1989	V36.0
997.09	1995	997.0	V37.00–V37.01	1989	V37.0
997.91	1995	997.9	V39.00–V39.01	1989	V39.0
997.99	1995	997.9	V42.81–V42.83	1997	V42.8
998.11–998.12	1996	998.1	V42.89	1997	V42.8
998.13	1996	998.89	V43.60–V43.66, V43.69	1994	V43.6
998.51	1996	998.5	V43.81–V43.82	1995	V43.8
998.59	1996	998.5	V43.83	1998	V43.89
998.81–998.82, 998.89	1994	998.8	V43.89	1995	V43.8
998.83	1996	998.89	V44.50	1998	V44.5
V02.51	1998	V02.5	V44.51	1998	V44.5
V02.52	1998	V02.5	V44.52	1998	V44.5
V02.59	1998	V02.5	V44.59	1998	V44.5
V02.60–V02.62	1997	V02.6	V45.00	1994	V45.89
V02.69	1997	V02.6	V45.01	1994	V45.0
V03.81–V03.82, V03.89	1994	V03.8	V45.02, V45.09	1994	V45.89
V05.3–V05.4	1993	V05.8	V45.51	1994	V45.5
V06.5–V06.6	1994	V06.8	V45.52, V45.59	1994	V45.89
V07.31, V07.39	1994	V07.3	V45.61	1997	V45.6
V07.4	1992	V07.8	V45.63	1997	V45.6
V08	1994	044.9, 795.8 (codes deleted)	V45.71	1997	611.8
V09.0–V09.91	1993	No previous code assignments	V45.72	1997	569.89
V10.48	1998	V10.49	V45.73	1997	593.89
V12.00–V12.03, V12.09	1994	V12.0	V45.82	1994	V45.89
V12.40–V12.41	1997	V12.4	V45.83	1995	V45.89
V12.49	1997	V12.4	V49.60–V49.67	1994	V49.5
V12.50–V12.52	1995	V12.5	V49.70–V49.77	1994	V49.5
V12.59	1995	V12.5	V50.41–V50.42, V50.49	1994	V50.8
			V53.01–V53.02	1997	V53.0

Table IV. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
V53.09	1997	V53.0	V72.81–V72.85	1993	V72.8
V53.31	1994	V53.3	V73.88–V73.89	1993	V73.8
V53.32, V53.39	1994	V53.9	V73.98–V73.99	1993	V73.9
V56.1	1995	V58.89	V76.10–V76.12	1997	V76.1
V56.1 (code title restated)	1998	V56.1	V76.19	1997	V76.1
V56.2	1998	V56.1	V76.44	1998	V76.49
V57.21–V57.22	1994	V57.2	V76.45	1998	V76.49
V58.41, V58.49	1994	V58.4	E854.8	1995	E858.8
V58.61	1995	V67.51	E869.4	1994	E869.8
V58.62	1998	V58.69	E880.1	1995	E884.9
V58.69	1995	V67.51	E884.3–E884.4	1995	E884.2
V58.81, V58.89	1994	V58.8	E884.5–E884.6	1995	E884.9
V58.82	1995	V58.89	E906.5	1995	E906.3
V59.01–V59.02	1995	V59.0	E908.0–E908.4	1995	E908
V59.09	1995	V59.0	E908.8–E908.9	1995	E908
V59.6	1995	V59.8	E909.0–E909.4	1995	E909
V61.10–V61.12	1996	V61.1	E909.8–E909.9	1995	E909
V61.22	1996	V61.21	E920.5	1995	E920.4
V62.83	1996	V65.49	E922.4	1997	E917.9
V64.4	1997	No previous code assignment	E924.2	1995	E924.0
V65.40–V65.45, V65.49	1994	V65.4	E955.6	1997	E955.9
V66.7	1996	No previous code assignment	E967.2	1996	E967.0
V69.0–V69.3	1994	No previous code assignments	E967.3	1996	No previous code assignment
V69.8–V69.9	1994	No previous code assignments	E967.4–E967.8	1996	E967.1
			E968.5	1995	E968.8
			E968.6	1997	E968.8
			E985.6	1997	E985.4

Table V. Changes in procedure codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
02.96	1992	89.19	37.83 (device only)	1987	37.73–37.77 (leads/device)
03.90	1987	03.99 (insertion of catheter)	37.85–37.87	1987	37.85
05.25	1995	39.7	37.89	1987	37.86 + 37.89
11.75	1989	11.79	37.94–37.98	1986	37.99
11.76	1989	11.62	38.22	1986	38.29
20.96–20.98	1986	20.95	38.44 (abdominal aorta only)	1986	38.44 (entire aorta)
22.12	1988	22.11	38.45 (thoracic aorta added)	1986	38.44–38.45
26.12	1988	26.11	38.95	1989	38.93
29.31	1991	83.02	39.28	1991	39.29
29.32	1991	29.3	39.50	1995	39.59
29.33	1991	29.3	39.65	1988	39.61
29.39	1991	29.3	39.66	1990	39.65
31.45	1988	31.43–31.44	39.90	1996	39.50
31.95	1989	31.75	41.00–41.03	1988	41.0
32.01	1989	32.0	41.04	1994	99.79
32.09	1989	32.0	41.05	1997	No previous code assignment
32.22	1995	32.29, 32.9	41.06	1997	No previous code assignment
32.28	1989	32.29	42.25	1988	42.24
33.27	1987	33.22 + 33.27	42.33	1989	42.32, 42.39
33.28	1987	33.27	42.33	1990	42.91
33.29	1987	33.28–33.29	43.11	1989	43.1
33.50	1995	33.5	43.19	1989	43.1, 43.2
33.51	1995	33.5	43.41	1989	43.41, 43.49
33.52	1995	33.5	44.21	1986	44.2
33.6	1990	33.5 + 37.5	44.22	1986	44.99
34.05	1994	34.99	44.29	1986	44.2
35.84	1988	35.82	44.43	1989	43.49, 45.32
35.96	1986	35.03	44.44	1989	38.86
36.00–36.03	1986	36.0	44.49	1989	43.0
36.04	1986	39.97	44.93–44.94	1986	44.99
36.05	1987	36.01	45.16	1988	45.14 (45.15 before 1987)
36.05	1986	36.01 ¹ , 36.02	45.30	1989	45.31, 45.32
36.06	1995	36.01, 36.02, 36.03, 36.05	45.42	1988	45.41
36.09	1986	36.0	45.43	1989	45.49
36.09	1991	36.00 (code deleted)	45.75 (Hartmann resection added)	1988	48.66 (code deleted)
36.17	1996	36.19	45.95	1987	45.93
36.31	1998	36.3	46.13	1992	46.12 (code deleted)
36.32	1998	36.3	46.32	1989	46.39
36.39	1998	36.3	46.85	1989	46.99
37.26–37.27	1988	37.29	47.01	1996	47.0
37.34	1988	37.33	47.09	1996	47.0
37.35	1997	37.33	47.11	1996	47.1
37.65	1995	37.62	47.19	1996	47.1
37.66	1995	37.62	48.36	1995	45.42
37.67	1998	37.4	49.31	1989	49.3
37.70 (leads only)	1987	37.70 (leads/device)	49.39	1989	49.3
37.71–37.72 (leads only)	1987	37.74 (leads/device)	51.10	1989	51.97
37.73 (leads only)	1987	37.73 (leads/device)	51.11	1989	51.11, 51.97
37.74 (leads only)	1987	37.76 (leads/device)	51.14	1989	51.12
37.75 (leads only)	1987	37.89 (leads/device)	51.15	1989	51.97
37.76 (leads only)	1987	37.81 (leads/device)	51.21	1996	51.22, 51.23
37.77 (leads only)	1987	37.83–37.84 (leads/device)	51.22	1991	51.21 (code deleted), 51.22
37.78	1987	37.71–37.72	51.23	1991	51.22
37.79	1987	86.09	51.24	1996	51.22, 51.23
37.80–37.87	1992	89.49 (code deleted. This procedure is included in the code for pacemaker insertion/replacement.)	51.64	1989	51.69
37.80 (device only)	1987	37.73–37.77 (leads/device)	51.84–51.88	1989	51.97
37.81 (device only)	1987	37.73–37.77 (leads/device)	51.97	1986	52.91, 51.99, or 51.82
37.82 (device only)	1987	37.73–37.77 (leads/device)	51.98	1986	51.99
			52.13	1989	51.97, 52.91
			52.14	1989	52.11
			52.21	1989	52.2
			52.22	1989	52.2
			52.84	1996	99.29

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
52.85	1996	99.29	78.15	1991	78.45
52.86	1996	99.29	78.16	1991	78.46
52.93	1989	52.93 + 52.91	78.17	1991	78.47
52.94	1989	52.09	78.18	1991	78.48
52.97	1989	52.91	78.19	1991	78.49
52.98	1989	52.91	78.20	1991	78.10, 78.20, 78.30
52.99	1989	52.93, 52.94, 52.99	78.21 (invalid code)	1991	78.11, 78.31
54.24	1987	54.23	78.22	1991	78.12, 78.22, 78.32
54.25	1993	54.98	78.23	1991	78.13, 78.23, 78.33
54.51	1996	54.5	78.24	1991	78.14, 78.34
54.59	1996	54.5	78.25	1991	78.15, 78.25, 78.35
55.03–55.04	1986	55.02	78.27	1991	78.17, 78.27, 78.37
56.33–56.34	1987	56.33	78.28	1991	78.18, 78.38
56.35	1987	45.12	78.29	1991	78.11, 78.16, 78.19, 78.29, 78.39
57.17–57.18	1989	57.21	78.39	1991	78.31
57.22	1989	57.22, 57.82	78.90 ²	1987	78.40
58.31	1990	58.3	78.91 ²	1987	78.41
58.39	1990	58.3	78.92 ²	1987	78.42
58.93	1986	57.99	78.93 ²	1987	78.43
59.03	1996	59.02	78.94 ²	1987	78.44
59.12	1996	59.11	78.95 ²	1987	78.45
59.72	1995	59.79	78.96 ²	1987	78.46
59.96	1986	59.95	78.97 ²	1987	78.47
60.21	1995	60.2	78.98 ²	1987	78.48
60.29	1995	60.2	78.99 ²	1987	78.49
60.95	1991	60.99	80.50–80.59	1986	80.5
64.97	1986	64.95	81.03	1989	81.02
65.01	1996	65.0	81.04–81.05	1989	81.03, 81.04, 81.05
65.09	1996	65.0	81.06–81.07	1989	81.06, 81.07
65.13	1996	65.12	81.08	1989	81.06, 81.07, 81.08
65.14	1996	65.19	81.09	1989	81.08
65.23	1996	65.21	81.40	1989	81.69
65.24	1996	65.22	81.51	1989	81.51, 81.59
65.25	1996	65.29	81.52	1989	81.61, 81.62, 81.63, 81.64
65.31	1996	65.3	81.53	1989	81.51, 81.59, 81.61, 81.62, 81.63, 81.64
65.39	1996	65.3	81.54–81.55	1989	81.41 (code deleted)
65.41	1996	65.4	81.56	1989	81.48
65.49	1996	65.4	81.57	1989	81.31, 81.39
65.53	1996	65.51	81.59	1989	81.39
65.54	1996	65.52	81.72	1989	81.79
65.63	1996	65.61	81.73–81.74	1989	81.86 (code deleted)
65.64	1996	65.62	81.75	1989	81.87 (code deleted)
65.74	1996	65.71	81.79	1989	81.79, 81.87
65.75	1996	65.72	81.80	1989	81.81
65.76	1996	65.73	81.97	1992	81.59
65.81	1996	65.8	85.95	1987	85.99
65.89	1996	65.8	85.96	1987	85.99
66.01	1992	66.0	86.06	1987	86.09
66.02	1992	66.73	86.07	1990	86.09
68.15	1987	68.14	86.27	1986	86.22–86.23
68.16	1987	68.13	86.28	1988	86.22
68.23	1996	68.29	86.67	1998	86.65
68.51	1996	68.5	86.93	1987	86.89
68.59	1996	68.5	88.90	1986	88.39
68.9	1992	68.4	88.91	1986	89.15
74.3	1992	69.11 (code deleted)	88.92	1986	89.39
75.37	1998	99.29	88.93	1986	89.15
77.56	1989	77.89, 78.49, 81.18	88.94	1986	89.39
77.57	1989	77.89, 80.48, 81.18, 83.85	88.95	1986	89.29
77.58	1989	77.59, 81.18	88.97	1989	88.99
78.10	1991	78.40	88.98	1989	88.90
78.11	1991	78.41	88.99	1986	89.39
78.12	1991	78.42	89.10	1989	89.15
78.13	1991	78.43	89.17–89.18	1988	89.15
78.14	1991	78.44			

See footnotes at end of table.

Table V. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
89.19	1989	89.15	96.71	1991	93.92 (code deleted)
89.50	1991	89.54	96.72	1991	93.92 (code deleted)
92.3	1995	01.59, 04.07, 07.63, 07.68	97.05	1989	51.97
92.30	1998	92.3	98.51–98.52	1989	59.96 (code deleted)
92.31	1998	92.3	98.59	1989	59.96 (code deleted)
92.32	1998	92.3	99.00	1995	99.02
92.33	1998	92.3	99.10	1998	99.29
92.39	1998	92.3	99.15	1986	99.29
93.90	1988	93.92	99.20	1998	99.29
94.61–94.69	1989	94.25	99.28	1994	99.25
96.29	1998	96.39	99.71–99.74, 99.79 ³	1988	99.07
96.6	1986	96.35	99.85	1987	93.35
96.70	1991	93.92 (code deleted)	99.86	1987	93.39
			99.88	1988	99.83

¹Before October 1986 contents of current code 36.05 would have been assigned to 36.0.

²Codes 78.90–78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90–78.99 were reassigned to codes 78.40–78.49.

³Codes 99.71–99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71–99.79 were reclassified as "Therapeutic apheresis." Codes 99.75–99.78 have not yet been reassigned.

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *National Vital Statistics Report* (NVSR). These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the NVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Data Dissemination Branch
National Center for Health Statistics
Centers for Disease Control and Prevention
6525 Belcrest Road, Room 1064
Hyattsville, MD 20782-2003
(301) 458-4636
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs

**DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
6525 Belcrest Road
Hyattsville, Maryland 20782-2003

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

MEDIA MAIL
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284