

SAFER • HEALTHIER • PEOPLE™



Vital and Health Statistics

Series 13, Number 158

March 2005

# National Hospital Discharge Survey: 2002 Annual Summary With Detailed Diagnosis and Procedure Data



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics



**Copyright information**

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

---

**Suggested citation**

Kozak LJ, Owings MF, Hall MJ. National Hospital Discharge Survey: 2002 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. Vital Health Stat 13(158). 2005.

---

**Library of Congress Catalog Card Number 90-657616**

---

For sale by the U.S. Government Printing Office  
Superintendent of Documents  
Mail Stop: SSOP  
Washington, DC 20402-9328  
Printed on acid-free paper.

---

# Vital and Health Statistics

---

Series 13, Number 158

## National Hospital Discharge Survey: 2002 Annual Summary With Detailed Diagnosis and Procedure Data

Data From the National Health Care  
Survey

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics

Hyattsville, Maryland  
March 2005  
DHHS Publication No. (PHS) 2005-1729

## **National Center for Health Statistics**

Edward J. Sondik, Ph.D., *Director*

Jennifer H. Madans, Ph.D., *Acting Co-Deputy Director*

Michael H. Sadagursky, *Acting Co-Deputy Director*

Jennifer H. Madans, Ph.D., *Associate Director for Science*

Lawrence H. Cox, Ph.D., *Associate Director for Research and Methodology*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Michael H. Sadagursky, *Associate Director for Management and Operations*

Margot A. Palmer, *Director for Information Technology*

Phillip R. Beattie, *Director for Information Services*

Diane M. Makuc, Dr.P.H., *Acting Associate Director for Analysis and Epidemiology*

Charles J. Rothwell, *Director for Vital Statistics*

Jane E. Sisk, Ph.D., *Director for Health Care Statistics*

Jane F. Gentleman, Ph.D., *Director for Health Interview Statistics*

Clifford L. Johnson, *Director for Health and Nutrition Examination Surveys*

## **Division of Health Care Statistics**

Jane E. Sisk, *Director*

Robin Remsburg, Dr.N., *Deputy Director*

Irma Arispe, Ph.D., *Associate Director for Science*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

# Contents

---

Abstract .....	1
Highlights.....	1
Patient and Hospital Characteristics .....	1
Diagnoses.....	1
Procedures .....	2
Deliveries and Newborn Infants .....	2
Introduction .....	2
Methods .....	3
Data Source .....	3
Estimation .....	5
Use of Tables.....	5
References .....	6
Appendix .....	184
Technical Notes .....	184
Abstract Form .....	184
Population Estimates .....	184
Reliability of Estimates.....	184
Definitions of Selected Terms.....	187
Changes in ICD-9-CM Codes .....	191

## Text Figures

1. Number of elderly discharges from short-stay hospitals by age: United States, selected years, 1970–2002.....	2
2. Selected first-listed diagnoses for short-stay hospital discharges by age: United States, 2002 .....	2
3. Selected secondary diagnoses for short-stay hospital discharges by age: United States, 2002.....	3
4. Annual average number of discharges with first-listed injury diagnosis by selected Barell injury diagnosis matrix categories: United States, 2000–2002 .....	3
5. Rate of selected cardiovascular procedures by sex: United States, 2002 .....	4
6. Number of primary and repeat cesarean deliveries: United States, 1995–2002 .....	4

## Detailed Tables

### Tables on Patient and Hospital Characteristics

1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2002 .....	7
2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2002 .....	8
3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2002 .....	9
4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2002 .....	11
5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by bed size of hospital, according to sex, age, and geographic region: United States, 2002.....	14

6.	Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by hospital ownership, according to sex, age, and geographic region: United States, 2002. ....	16
7.	Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2002. ....	18
8.	Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 2002. ....	18

**Diagnosis Tables**

9.	Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2002. ....	19
10.	Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 2002. ....	20
11.	Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002. ....	22
12.	Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002. ....	23
13.	Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002. .	24
14.	Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002. ....	25
15.	Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002. ....	26
16.	Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002. .	27
17.	Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002. .	28
18.	Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002. ....	29
19.	Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002. ....	30
20.	Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2002. ....	31
21.	Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2002. ....	32
22.	Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2002. ....	33
23.	Number and rate of discharges from short-stay hospitals with human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years 1990–2002. ....	34
24.	Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2000–2002. ....	35
25.	Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2002. ....	35

**Procedure Tables**

26.	Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 2002. ....	36
27.	Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 2002. ....	37
28.	Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2002. ....	38
29.	Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures by sex, age, and geographic region: United States, 2002. ....	39
30.	Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2002. ....	40
31.	Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2002. ....	41
32.	Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2002. ....	42
33.	Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2002. ....	43
34.	Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2002. ....	44
35.	Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2002. ....	45

**Tables on Deliveries and Newborn Infants**

36.	Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals by age and geographic region, according to type of delivery: United States, 2002. ....	46
-----	--	----

37.	Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–2002 .....	47
38.	Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2002.....	48
39.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–2002.....	48
40.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 2002.....	49
41.	Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2002 .....	50
42.	Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay, by health status: United States, selected years 1980–2002 .....	50
43.	Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2002 .....	51
44.	Number and percent of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years 1980–2002.....	51

**Diagnosis, Days of Care, and Procedure Tables by ICD–9–CM Code**

45.	Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2002 .....	52
46.	Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002 .....	79
47.	Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2002 .....	106
48.	Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2002 .....	165

**Appendix Tables**

I.	Civilian population by age, geographic region, and sex: United States, 2002 .....	184
II.	Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics, 2002 .....	187
III.	Code numbers for Barell injury diagnosis matrix categories in table 24.....	189
IV.	Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD–9–CM category, 2002.....	190
V.	Changes in diagnosis codes .....	192
VI.	Changes in procedure codes .....	197

**Appendix Figure**

I.	Medical abstract form for the National Hospital Discharge Survey, 2002 .....	185
----	--	-----

**Objectives**

This report presents 2002 national estimates and selected trend data on the use of non-Federal short-stay hospitals in the United States. Estimates are provided by selected patient and hospital characteristics, diagnoses, and surgical and nonsurgical procedures performed. Estimates of diagnoses and procedures are presented according to *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) codes.

**Methods**

The estimates are based on data collected through the National Hospital Discharge Survey (NHDS). The survey has been conducted annually since 1965. In 2002, data were collected for approximately 327,000 discharges. Of the 474 eligible non-Federal short-stay hospitals in the sample, 445 (94 percent) responded to the survey.

**Results**

An estimated 33.7 million inpatients were discharged from non-Federal short-stay hospitals in 2002. They used 164.2 million days of care and had an average length of stay of 4.9 days. Common first-listed discharge diagnoses included delivery, ischemic heart disease, psychoses, pneumonia, and malignant neoplasms. Inpatients had 6.8 million cardiovascular procedures and 6.6 million obstetric procedures. Males had higher rates for cardiac procedures such as cardiac catheterization and coronary artery bypass graft, but males and females had similar rates of pacemaker procedures. The number and rate of all cesarean deliveries, primary and repeat, rose from 1995 to 2002; the rate of vaginal birth after cesarean delivery dropped from 35.5 in 1995 to 15.8 in 2002.

**Keywords:** hospitalization • inpatient • diagnoses • procedures

# National Hospital Discharge Survey: 2002 Annual Summary With Detailed Diagnosis and Procedure Data

by *Lola J. Kozak, Ph.D.; Maria F. Owings, Ph.D.; and Margaret J. Hall, Ph.D., Division of Health Care Statistics*

## Highlights

### Patient and Hospital Characteristics

- An estimated 33.7 million inpatients were discharged from non-Federal short-stay hospitals in 2002. They used an estimated 164.2 million days of care and were hospitalized for an average of 4.9 days.
- As shown in [figure 1](#), the number of discharges from short-stay hospitals steadily increased for patients 75–84 years and 85 years and over between 1970 and 2002, but did not change significantly for the 65–74 years age group between 1980 and 2002. Discharge rates per 1,000 population also increased for the 75–84 and 85 and over age groups between 1970 and 2002, but the rate for the 65–74 years group decreased between 1980 and 2002.
- The average length of a hospital stay ranged from 4.4 days in the Midwest to 5.6 days in the Northeast.
- Discharges 15–44 years of age made up 60 percent of discharges in the self-pay category, but only 32 percent of total discharges. Hospitals in the South accounted for 49 percent of self-pay discharges compared with 39 percent of total discharges.
- Nonprofit hospitals accounted for 75 percent of hospital discharges in

2002. The proportion ranged from 94 percent in the Northeast to 56 percent in the South.

- Deaths accounted for 2.4 percent of all hospital discharges in 2002: 3.0 percent for males and 2.0 percent for females.

### Diagnoses

- The first-listed discharge diagnosis (generally the cause of hospitalization) for 4.0 million discharges under 65 years of age was delivery ([figure 2](#)).
- Ischemic heart disease, which includes acute myocardial infarction and coronary atherosclerosis, was the first-listed diagnosis for 1.2 million discharges 65 years of age and over ([figure 2](#)).
- Other first-listed diagnoses that were reported for more than a million discharges for all ages in 2002 included psychoses (1.7 million), pneumonia (1.3 million), and malignant neoplasms (1.2 million).
- [Figure 3](#) shows five frequently occurring secondary diagnoses for discharges under age 65 and age 65 and over. Essential hypertension was reported as a secondary diagnosis 2.9 million times for those under age 65 and 4.8 million times for those age 65 years and over.
- The average annual number of discharges with first-listed diagnoses of all types of injuries was 1,802,000 in 2000–2002. Using the Barell injury diagnosis matrix






Figure 1. Number of elderly discharges from short-stay hospitals by age: United States, selected years, 1970–2002




Figure 2. Selected first-listed diagnoses for short-stay hospital discharges by age: United States, 2002

(figure 4), fracture (992,000) was the most common type of injury reported and lower extremity, including the hip (688,000), was the most common site of fracture and of all injuries.

### Procedures

- In 2002, 25.3 million surgical and 17.3 million nonsurgical procedures were performed on hospitalized patients.
- Females had 6.6 million obstetrical procedures, including 1.2 million

repairs of current obstetric laceration and 1.1 million cesarean sections.


- A total of 6.8 million operations on the cardiovascular system, including 1.3 million cardiac catheterizations and 1.2 million removal of coronary artery obstruction and insertion of stents, were performed on inpatients in 2002.
- Males had higher rates of cardiac procedures such as cardiac catheterization, removal of coronary artery obstruction and insertion of stents (coronary angioplasty), and coronary artery bypass graft. However, the rate of pacemaker procedures was similar for males and females (figure 5).

### Deliveries and Newborn Infants


- Approximately 4 million women were hospitalized for delivery in 2002, and their average stay was 2.6 days.
- Cesarean sections per 100 deliveries rose from 20.8 in 1995 to 26.8 in 2002.
- The number and rate of both primary and repeat cesarean deliveries increased from 1995 through 2002, but repeat cesareans expanded from 266,000 (34 per 100 cesareans) to 420,000 (40 per 100 cesareans) (figure 6).
- The rate of vaginal birth after cesarean delivery (VBAC) dropped from 35.5 in 1995 to 15.8 in 2002.
- In 2002, 40 percent of newborn infants had at least one illness or risk-related diagnosis. Perinatal jaundice, respiratory conditions, and disorders related to prematurity were common diagnoses.
- The proportion of male newborn infants circumcised during the birth hospitalization was 60 percent nationally, but ranged from 33 percent in the West to 81 percent in the Midwest.

### Introduction

This report presents data from the 2002 National Hospital Discharge Survey (NHDS). The survey has



**Figure 3. Selected secondary diagnoses for short-stay hospital discharges by age: United States, 2002**



**Figure 4. Annual average number of discharges with first-listed injury diagnosis by selected Barell injury diagnosis matrix categories: United States, 2000–2002**

been conducted continuously by the National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report provides a summary of the 2002 data, including estimates of diagnoses and procedures by

*International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM) codes (1).*

The NHDS is the principal source for national data on the characteristics of patients discharged from non-Federal short-stay hospitals. Data from the NHDS are used for evaluating the health

status of the population, planning programs to improve health status, studying trends in morbidity, and carrying out research activities in the health field (2–6). The Department of Health and Human Services uses NHDS data in the development and monitoring of goals for the 2010 Health Objectives. Other CDC centers, including the National Center for Infectious Diseases, the National Center for Injury Prevention and Control, and the National Center for Chronic Disease Prevention and Health Promotion, track a variety of conditions, including human immunodeficiency virus (HIV), injuries, and chronic diseases with NHDS data. The National Institutes of Health makes extensive use of NHDS data on heart disease and cancer. The Veterans Administration and Department of Defense compare NHDS data with statistics on the inpatient care provided in their hospitals (7–8).

The NHDS produces estimates of utilization by inpatients only in non-Federal short-stay hospitals. Data on ambulatory medical care services are collected in other NCHS surveys. The National Ambulatory Medical Care Survey obtains information on visits to physicians' offices (9). The National Hospital Ambulatory Medical Care Survey collects data on visits to hospital outpatient and emergency departments (10,11). Information about long-term care is collected in the National Nursing Home Survey (12) and the National Home and Hospice Care Survey (13).

## Methods

### Data Source

NHDS data are collected from a sample of inpatient records acquired from a national probability sample of hospitals. Because persons with multiple discharges during the year can be sampled more than once, the NHDS produces estimates for discharges, not persons. Only general hospitals, children's general hospitals, or hospitals with an average length of stay of fewer than 30 days for all patients are eligible for inclusion in the survey. Federal,




Figure 5. Rate of selected cardiovascular procedures by sex: United States, 2002




Figure 6. Number of primary and repeat cesarean deliveries: United States, 1995–2002

military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals) and hospitals with fewer than six beds staffed for patient use, are excluded. Beginning in 1988, the sampling frame for hospitals has been the SMG Hospital Market Database

(14). The 1988 sample was updated in 1991, 1994, 1997, and 2000 to include hospitals that opened or changed their eligibility status since the previous sample update.

Since 1988, the NHDS has used a modified three-stage design (15). Units selected at the first stage consist of

either hospitals or geographic areas such as counties, groups of counties, or metropolitan statistical areas in the 50 States and the District of Columbia. Within sampled geographic areas, additional hospitals are selected. Finally, at the last stage, discharges are selected within sampled hospitals using systematic random sampling.

For 2002, the sample consisted of 504 hospitals, of which 30 were out of scope (ineligible) because they had gone out of business or were otherwise ineligible to be in the NHDS universe. Of the 474 in-scope (eligible) hospitals, 445 (94 percent) responded to the survey. Data were collected for approximately 327,000 discharges from the 445 responding hospitals.

Two data collection procedures are used for the survey. One is a manual system in which sample selection and transcription of information from medical records are performed by hospital staff or by staff of the U.S. Census Bureau on behalf of NCHS. Completed forms are sent to NCHS for medical coding, data entry, editing, and estimation.

For the other data collection procedure, NCHS purchases electronic files containing medical record data from commercial organizations, State data systems, hospitals, or hospital associations. Discharges from these files are systematically sampled. Approximately 40 percent of respondent hospitals provided data through this automated system for the 2002 survey.

The NHDS medical abstract form (figure 1) and the automated data contain items that relate to the personal characteristics of the patient. These include birth date (converted to age), sex, race, ethnicity, marital status, ZIP Code, and expected sources of payment. Administrative items such as admission and discharge dates, admission type and source, discharge status, and medical record number are also included. The medical information includes final medical diagnoses, procedures performed, and dates of surgery. Medical data are coded according to the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) (1). Date of birth and ZIP



Code are confidential information not available to the public.

For the manual data collection in the NHDS, an ongoing quality control program is undertaken for the medical coding and data entry from abstracts to electronic form. Approximately 10 percent of abstracts are independently recoded, with discrepancies resolved by a chief coder. The overall error rate for records manually coded for the 2002 survey year was 0.1 percent for demographic and medical (ICD-9-CM) coding and data entry.

## Estimation

Because they are sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components:

- (1) inflation by reciprocals of the probabilities of sample selection,
- (2) adjustment for nonresponse, and
- (3) population weighting ratio adjustments.

These three components of the final weight are described in more detail in another report (15). Information about the standard errors of statistics for the 2002 NHDS is in [table II](#).

## Use of Tables

Sociodemographic information about patients and utilization estimates by hospital characteristics are shown in [tables 1–8](#). Diagnostic data are included in [tables 9–25](#), and information about procedures is in [tables 26–35](#). [Tables 36–44](#) present data on women with deliveries and newborn infants. Estimates of diagnoses, days of care by diagnosis, and procedures are shown by ICD-9-CM code in [tables 45–48](#). [Figures 1–6](#) illustrate highlights from the survey data. Definitions of terms used in these tables and figures are presented in the “Appendix.”

Because there is no widely accepted method of grouping diagnoses or procedures, the data in this report are generally presented by chapter of the ICD-9-CM with frequently occurring broad and specific subcategories within

chapters also shown (see [tables 11–22](#) and [30–35](#)). The first-listed diagnostic categories shown in [tables 9](#) and [10](#) and [figure 2](#) are based on the categories with relatively large estimates in [tables 11–18](#). Likewise, the procedures shown in [tables 27](#) and [28](#) are based on the categories with relatively large estimates in [tables 30–35](#). The absence of standard categories makes it inadvisable to refer to the leading or most common diagnoses or procedures or to rank categories in order of frequency because the order of magnitude would vary based on what is or is not included in the various categories.

Diagnoses assigned ICD-9-CM codes E800–E999 (supplementary classification of external causes of injury and poisoning) are not included in this report. However, a detailed discussion of injury diagnoses and E-codes has been published (16).

Injury data are presented in this report using the recently developed Barell injury diagnosis matrix ([table 24](#), [figure 4](#)). This is a two-dimensional array of ICD-9-CM codes (excluding external cause codes) that groups injuries by body site and type of injury. The codes for the matrix are shown in [table III](#). The matrix provides a standard format for reporting injury data that improves the usefulness and comparability of epidemiological and clinical analyses (17). Data categorized using the Barell matrix are included in this report to provide a national benchmark for States and localities.

[Table 29](#) shows data for coronary artery bypass graft (CABG) surgery and removal of coronary artery obstruction and insertion of stent(s). These procedures are often assigned more than one code for a single operation. Coronary artery bypass graft surgery often is assigned multiple codes to indicate the number and type of bypasses. Beginning with the 1996 NHDS data, changes to the ICD-9-CM coding system allowed patients to have a separate code for insertion of stent(s) along with a code for removal of coronary artery obstruction. As shown in [table 29](#), the number of these procedures exceeds the number of inpatients who had one or more of the procedures.

[Tables 8](#) and [25](#) show rates of death per 100 discharges. In contrast, mortality rates are ratios of deaths to the population. Mortality rates include deaths that occur outside of hospitals and are for the underlying cause of death, which might not be the same as the first-listed diagnosis for a patient discharged dead. Mortality rates are published regularly by NCHS in *National Vital Statistics Reports* (18).

Data for newborn infants are included only in [tables 41–44](#). Because estimates of newborn infants are based on a sample and because they do not include out-of-hospital births, these estimates may not agree with data on births published in *National Vital Statistics Reports* (19), which are based on a complete count of births.

[Tables 45–47](#) provide information by three-digit, four-digit, and five-digit ICD-9-CM diagnostic codes. [Table 48](#) shows two-, three-, and four-digit procedure codes. Only ICD-9-CM code numbers appear in [tables 45–48](#). Specific titles and more detailed information about the coding system are in appropriate volumes of the ICD-9-CM (1). A specific diagnosis or procedure code appears in [tables 45](#), [47](#), and [48](#) if the estimate of the total for the code is 5,000 or more. The codes that appear in [table 46](#) for days of care are identical to those in [table 45](#) for first-listed diagnoses.

Estimates in all tables are rounded to the nearest thousand. Totals may include data for categories that are not shown individually in the tables. For these reasons, figures within tables do not always add to the totals. Rates, percents, and average lengths of stay are calculated from unrounded figures and may not precisely agree with measures calculated from rounded data.

Because of low reliability, estimates in [tables 1–44](#) with relative standard errors of more than 30 percent or that are based on fewer than 30 records are not presented; only an asterisk (\*) appears in the tables. Estimates based on 30–59 records are preceded by an asterisk (\*) to indicate that they also have low reliability.

Due to the large number of estimates in [tables 45–48](#), more generalized criteria for reliability were

used. Only an asterisk (\*) is shown in tables 45, 47, and 48 if the estimate of the diagnosis or procedure is less than 5,000, and only an asterisk (\*) is shown in table 46 for estimates of days of care derived from fewer than 5,000 estimated discharges. These estimates generally have a relative standard error of more than 30 percent or a sample size of less than 30. Estimates of diagnoses and procedures of less than 9,000 are preceded by an asterisk (\*) because they are generally based on fewer than 60 records. Days of care estimates in table 46 that are derived from fewer than 9,000 estimated discharges are also preceded by asterisks.

The population estimates used to compute rates are shown in table I. These estimates are based on the 2000 census.

A weighted least squares regression method was used to test for trends (20). Differences between NHDS estimates were tested using the two-sided *t*-test. Terms that express differences such as higher, highest, lower, lowest, longer, shorter, largest, smallest, more than, most, less than, leading, increased or decreased, were only used when the differences were statistically significant at the 0.05 level.

## References

- International Classification of Diseases, 9th Revision, Clinical Modification, 6th edition. U.S. Department of Health and Human Services, National Center for Health Statistics, Health Care Financing Administration. 1998. <http://www.cdc.gov/nchs/icd9.htm#ICD-9-CM>.
- Liu JH, Etzioni DA, O'Connell JB, et al. The increasing workload of general surgery. *Arch Surg* 139:423–28. 2004.
- Wattigney WA, Mensah GA, Croft JB. Increasing trends in hospitalization for atrial fibrillation in the United States, 1985 through 1999: implications for primary prevention. *Circulation* 108:711–16. 2003.
- Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Eng J Med* 348:1546–54. 2003.
- Danel I, Berg C, Johnson CH, Atrash H. Magnitude of maternal morbidity during labor and delivery: United States, 1993–1997. *Am J Public Health* 93(4):631–4. 2003.
- Wang G, Dietz WH. Economic burden of obesity in youths aged 6 to 17 years: 1979–1999. *Pediatrics* 109(5):E81. 2002. <http://www.pediatrics.org/cgi/content/full/109/5/e81>.
- Rosenthal GE, Kaboli PJ, Barnett MJ. Differences in length of stay in Veterans Health Administration and other United States hospitals: is the gap closing? *Med Care* 41(8):882–94. 2003.
- Jackson JL, Cheng EY, Jones DL, Meyer G. Comparison of discharge diagnoses and inpatient procedures between military and civilian health care systems. *Mil Med* 164(10):701–4. 1999.
- Woodwell DA, Cherry DK. National Ambulatory Medical Care Survey: 2002 summary. Advance data from vital and health statistics; no 346. Hyattsville, Maryland: National Center for Health Statistics. 2004. <http://www.cdc.gov/nchs/data/ad/ad346.pdf>.
- Hing E, Middleton K. National Hospital Ambulatory Medical Care Survey: 2002 outpatient department summary. Advance data from vital and health statistics; no 345. Hyattsville, Maryland: National Center for Health Statistics. 2004. <http://www.cdc.gov/nchs/data/ad/ad345.pdf>.
- McCaig LF, Burt CW. National Hospital Ambulatory Medical Care Survey: 2002 emergency department summary. Advance data from vital and health statistics; no 340. Hyattsville, Maryland: National Center for Health Statistics. 2004. <http://www.cdc.gov/nchs/data/ad/ad340.pdf>.
- Jones A. The National Nursing Home Survey: 1999 summary. National Center for Health Statistics. *Vital Health Stat* 13(152). 2002. [http://www.cdc.gov/nchs/data/series/sr\\_13/sr13\\_152.pdf](http://www.cdc.gov/nchs/data/series/sr_13/sr13_152.pdf).
- Haupt BJ. Characteristics of hospice care discharges and their length of service: United States, 2000. National Center for Health Statistics. *Vital Health Stat* 13(154). 2003. [http://www.cdc.gov/nchs/data/series/sr\\_13/sr13\\_154.pdf](http://www.cdc.gov/nchs/data/series/sr_13/sr13_154.pdf).
- SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1987, 1991, 1994, 1997, 2000.
- Dennison CF, Pokras R. Design and Operation of the National Hospital Discharge Survey: 1988 Redesign. National Center for Health Statistics. *Vital Health Stat* 1(39). 2000. [http://www.cdc.gov/nchs/data/series/sr\\_01/sr01\\_039.pdf](http://www.cdc.gov/nchs/data/series/sr_01/sr01_039.pdf).
- Hall MJ, Owings MF. Hospitalizations for injury: United States, 1996. Advance data from vital and health statistics; no 318. Hyattsville, Maryland: National Center for Health Statistics. 2000. <http://www.cdc.gov/nchs/data/ad/ad318.pdf>.
- Barell V, Aharonson-Daniel L, Fingerhut LA, et al. An introduction to the Barell body region by nature of injury diagnosis matrix. *Injury Prevention* 8:91–6. 2002.
- Kochanek KD, Smith BL. Deaths: preliminary data for 2002. National vital statistics reports vol 52 no 13. Hyattsville, Maryland: National Center for Health Statistics. 2004. [http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52\\_13.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52_13.pdf).
- Martin JA, Hamilton BE, Sutton PD, et al. Births: final data for 2002. National vital statistics reports vol 52 no 10. Hyattsville, Maryland: National Center for Health Statistics. 2003. [http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52\\_10.pdf](http://www.cdc.gov/nchs/data/nvsr/nvsr52/nvsr52_10.pdf).
- Gillum BS, Graves EJ, Kozak LJ. Trends in hospital utilization: United States, 1988–92. National Center for Health Statistics. *Vital Health Stat* 13(124). 1996. [http://www.cdc.gov/nchs/data/series/sr\\_13/sr13\\_124.pdf](http://www.cdc.gov/nchs/data/series/sr_13/sr13_124.pdf).
- Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual, Release 7.0. Research Triangle Park, NC: Research Triangle Institute. 1996.
- Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. New York: John Wiley & Sons. Wiley Classics Library Edition. 1993.
- Cochran WG. Sampling techniques, 3rd ed. New York: John Wiley & Sons. 1977.

**Table 1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	1970	1980	1985	1990	1995	2000	2002
Number of discharges in thousands							
All ages . . . . .	29,127	37,832	35,056	30,788	30,722	31,706	33,727
Under 15 years . . . . .	3,873	3,672	2,972	2,412	2,405	2,383	2,540
15–44 years . . . . .	12,664	15,635	13,966	11,799	10,593	9,969	10,736
45–64 years . . . . .	6,693	8,660	7,610	6,244	6,168	6,958	7,723
65 years and over . . . . .	5,897	9,864	10,508	10,333	11,556	12,396	12,727
65–74 years . . . . .	3,163	4,943	5,011	4,689	4,832	4,678	4,642
75–84 years . . . . .	2,099	3,611	3,969	3,949	4,590	5,119	5,271
85 years and over . . . . .	635	1,310	1,528	1,694	2,134	2,599	2,814
Rate of discharges per 1,000 population <sup>1</sup>							
All ages . . . . .	144.3	167.7	148.4	122.3	115.7	112.8	117.5
Under 15 years . . . . .	66.8	71.6	57.7	43.1	40.4	39.5	41.9
15–44 years . . . . .	154.6	150.1	125.0	99.3	87.8	80.9	86.4
45–64 years . . . . .	159.6	194.8	170.8	135.5	118.5	111.4	115.9
65 years and over . . . . .	293.3	383.7	369.8	334.1	347.7	353.4	357.5
65–74 years . . . . .	253.2	315.8	297.3	261.6	260.0	254.6	254.0
75–84 years . . . . .	339.5	463.9	446.4	395.7	415.6	412.4	413.9
85 years and over . . . . .	443.7	576.4	572.9	560.6	592.5	605.1	612.8
Number of days of care in thousands							
All ages . . . . .	226,445	274,508	226,217	197,422	164,627	155,857	164,152
Under 15 years . . . . .	18,189	16,191	13,554	11,655	10,715	10,734	11,531
15–44 years . . . . .	71,605	81,951	67,397	54,062	40,825	36,593	40,091
45–64 years . . . . .	62,419	71,008	53,541	42,153	34,207	34,443	38,298
65 years and over . . . . .	74,232	105,358	91,726	89,552	78,880	74,086	74,233
65–74 years . . . . .	37,873	49,253	41,090	37,422	31,310	26,438	25,801
75–84 years . . . . .	27,633	40,355	36,024	35,926	31,974	31,525	31,497
85 years and over . . . . .	8,726	15,750	14,612	16,204	15,597	16,123	16,934
Rate of days of care per 1,000 population <sup>1</sup>							
All ages . . . . .	1,121.6	1,216.7	957.7	784.0	620.2	554.6	571.7
Under 15 years . . . . .	313.9	315.7	263.0	208.4	179.9	178.0	190.1
15–44 years . . . . .	874.0	786.8	603.3	454.9	338.4	297.1	322.7
45–64 years . . . . .	1,488.8	1,596.9	1,201.6	914.4	657.5	551.5	574.7
65 years and over . . . . .	3,691.9	4,098.4	3,228.0	2,895.6	2,373.7	2,111.9	2,085.1
65–74 years . . . . .	3,031.6	3,147.0	2,437.3	2,087.8	1,684.7	1,439.0	1,411.9
75–84 years . . . . .	4,469.2	5,183.7	4,052.2	3,599.5	2,894.7	2,539.8	2,473.4
85 years and over . . . . .	6,102.1	6,932.1	5,478.7	5,361.7	4,330.6	3,753.8	3,686.9
Average length of stay in days							
All ages . . . . .	7.8	7.3	6.5	6.4	5.4	4.9	4.9
Under 15 years . . . . .	4.7	4.4	4.6	4.8	4.5	4.5	4.5
15–44 years . . . . .	5.7	5.2	4.8	4.6	3.9	3.7	3.7
45–64 years . . . . .	9.3	8.2	7.0	6.8	5.5	5.0	5.0
65 years and over . . . . .	12.6	10.7	8.7	8.7	6.8	6.0	5.8
65–74 years . . . . .	12.0	10.0	8.2	8.0	6.5	5.7	5.6
75–84 years . . . . .	13.2	11.2	9.1	9.1	7.0	6.2	6.0
85 years and over . . . . .	13.7	12.0	9.6	9.6	7.3	6.2	6.0

<sup>1</sup>Rates were calculated using the U.S. Census Bureau estimates of the civilian population. Rates for 1990 and 1995 were based on population estimates adjusted for the net underenumeration in the 1990 census. Rates for 2000 and 2002 were calculated using estimates based on the 2000 census. Rates for 2000 differ from the ones shown in earlier reports because they were calculated using revised population estimates.



**Table 2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex and age	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population <sup>1</sup>	Number in thousands	Percent distribution	Rate per 1,000 population <sup>1</sup>	
<b>Both sexes</b>							
All ages . . . . .	33,727	100.0	117.5	164,152	100.0	571.7	4.9
Under 15 years . . . . .	2,540	7.5	41.9	11,531	7.0	190.1	4.5
Under 1 year . . . . .	810	2.4	200.8	4,575	2.8	1,134.1	5.6
1–4 years . . . . .	713	2.1	45.8	2,473	1.5	158.8	3.5
5–14 years . . . . .	1,016	3.0	24.8	4,483	2.7	109.3	4.4
15–44 years . . . . .	10,736	31.8	86.4	40,091	24.4	322.7	3.7
15–19 years . . . . .	1,239	3.7	61.1	4,817	2.9	237.7	3.9
20–24 years . . . . .	1,844	5.5	93.2	5,857	3.6	295.9	3.2
25–34 years . . . . .	3,897	11.6	98.6	12,965	7.9	328.1	3.3
35–44 years . . . . .	3,757	11.1	84.1	16,452	10.0	368.4	4.4
45–64 years . . . . .	7,723	22.9	115.9	38,298	23.3	574.7	5.0
45–54 years . . . . .	3,826	11.3	95.6	18,289	11.1	456.7	4.8
55–64 years . . . . .	3,897	11.6	146.5	20,009	12.2	752.2	5.1
65 years and over . . . . .	12,727	37.7	357.5	74,233	45.2	2,085.1	5.8
65–74 years . . . . .	4,642	13.8	254.0	25,801	15.7	1,411.9	5.6
75–84 years . . . . .	5,271	15.6	413.9	31,497	19.2	2,473.4	6.0
85 years and over . . . . .	2,814	8.3	612.8	16,934	10.3	3,686.9	6.0
<b>Male</b>							
All ages . . . . .	13,389	100.0	95.2	71,076	100.0	505.5	5.3
Under 15 years . . . . .	1,412	10.5	45.5	6,282	8.8	202.4	4.4
Under 1 year . . . . .	463	3.5	224.4	2,495	3.5	1,209.1	5.4
1–4 years . . . . .	401	3.0	50.3	1,318	1.9	165.5	3.3
5–14 years . . . . .	548	4.1	26.1	2,469	3.5	117.5	4.5
15–44 years . . . . .	2,906	21.7	46.6	14,362	20.2	230.3	4.9
15–19 years . . . . .	335	2.5	32.3	1,759	2.5	169.4	5.2
20–24 years . . . . .	347	2.6	34.7	1,661	2.3	166.2	4.8
25–34 years . . . . .	783	5.9	39.5	3,648	5.1	183.8	4.7
35–44 years . . . . .	1,441	10.8	65.1	7,294	10.3	329.5	5.1
45–64 years . . . . .	3,755	28.0	115.8	19,359	27.2	597.0	5.2
45–54 years . . . . .	1,812	13.5	92.3	9,276	13.1	472.2	5.1
55–64 years . . . . .	1,943	14.5	152.0	10,083	14.2	788.8	5.2
65 years and over . . . . .	5,315	39.7	359.8	31,074	43.7	2,103.6	5.8
65–74 years . . . . .	2,205	16.5	265.6	12,341	17.4	1,486.7	5.6
75–84 years . . . . .	2,234	16.7	439.7	13,284	18.7	2,614.4	5.9
85 years and over . . . . .	877	6.5	630.9	5,448	7.7	3,920.2	6.2
<b>Female</b>							
All ages . . . . .	20,338	100.0	138.8	93,076	100.0	635.2	4.6
Under 15 years . . . . .	1,128	5.5	38.1	5,249	5.6	177.3	4.7
Under 1 year . . . . .	347	1.7	176.1	2,079	2.2	1,055.5	6.0
1–4 years . . . . .	313	1.5	41.1	1,155	1.2	151.7	3.7
5–14 years . . . . .	468	2.3	23.4	2,015	2.2	100.6	4.3
15–44 years . . . . .	7,830	38.5	126.6	25,729	27.6	415.9	3.3
15–19 years . . . . .	903	4.4	91.4	3,058	3.3	309.3	3.4
20–24 years . . . . .	1,498	7.4	152.9	4,195	4.5	428.3	2.8
25–34 years . . . . .	3,114	15.3	158.3	9,318	10.0	473.8	3.0
35–44 years . . . . .	2,316	11.4	102.8	9,158	9.8	406.7	4.0
45–64 years . . . . .	3,968	19.5	116.0	18,939	20.3	553.4	4.8
45–54 years . . . . .	2,014	9.9	98.7	9,014	9.7	441.8	4.5
55–64 years . . . . .	1,954	9.6	141.4	9,925	10.7	718.3	5.1
65 years and over . . . . .	7,412	36.4	355.8	43,159	46.4	2,072.0	5.8
65–74 years . . . . .	2,437	12.0	244.4	13,460	14.5	1,349.6	5.5
75–84 years . . . . .	3,037	14.9	396.8	18,213	19.6	2,379.7	6.0
85 years and over . . . . .	1,938	9.5	604.9	11,486	12.3	3,585.6	5.9

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.

**Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population <sup>1</sup>	Number in thousands	Rate per 1,000 population <sup>1</sup>	
Both sexes					
All ages:					
United States . . . . .	33,727	117.5	164,152	571.7	4.9
Northeast . . . . .	6,990	129.0	39,371	726.8	5.6
Midwest . . . . .	7,503	115.4	33,277	511.6	4.4
South . . . . .	12,994	126.7	63,403	618.0	4.9
West . . . . .	6,239	95.5	28,101	430.3	4.5
Under 15 years:					
United States . . . . .	2,540	41.9	11,531	190.1	4.5
Northeast . . . . .	512	48.0	2,585	242.1	5.0
Midwest . . . . .	487	35.8	1,926	141.6	4.0
South . . . . .	931	43.0	4,290	198.0	4.6
West . . . . .	610	41.5	2,730	185.7	4.5
15–44 years:					
United States . . . . .	10,736	86.4	40,091	322.7	3.7
Northeast . . . . .	2,008	87.1	9,355	405.8	4.7
Midwest . . . . .	2,119	75.8	7,146	255.6	3.4
South . . . . .	4,492	101.4	16,654	376.1	3.7
West . . . . .	2,117	73.1	6,936	239.6	3.3
45–64 years:					
United States . . . . .	7,723	115.9	38,298	574.7	5.0
Northeast . . . . .	1,584	121.7	8,794	675.5	5.5
Midwest . . . . .	1,711	112.6	7,578	498.8	4.4
South . . . . .	3,094	129.3	15,486	647.4	5.0
West . . . . .	1,334	91.9	6,441	443.7	4.8
65 years and over:					
United States . . . . .	12,727	357.5	74,233	2,085.1	5.8
Northeast . . . . .	2,885	388.5	18,638	2,510.2	6.5
Midwest . . . . .	3,187	384.0	16,627	2,003.7	5.2
South . . . . .	4,478	351.5	26,973	2,117.4	6.0
West . . . . .	2,178	305.1	11,994	1,680.0	5.5
Male					
All ages:					
United States . . . . .	13,389	95.2	71,076	505.5	5.3
Northeast . . . . .	2,962	113.0	18,020	687.8	6.1
Midwest . . . . .	3,025	94.9	14,230	446.3	4.7
South . . . . .	4,947	98.9	26,670	533.1	5.4
West . . . . .	2,455	75.6	12,157	374.1	5.0
Under 15 years:					
United States . . . . .	1,412	45.5	6,282	202.4	4.4
Northeast . . . . .	289	52.8	1,539	281.7	5.3
Midwest . . . . .	279	40.0	1,090	156.5	3.9
South . . . . .	502	45.3	2,295	207.1	4.6
West . . . . .	343	45.5	1,358	180.3	4.0
15–44 years:					
United States . . . . .	2,906	46.6	14,362	230.3	4.9
Northeast . . . . .	650	56.7	3,872	337.6	6.0
Midwest . . . . .	588	41.8	2,438	173.3	4.1
South . . . . .	1,158	52.5	5,755	260.6	5.0
West . . . . .	509	34.5	2,298	155.8	4.5
45–64 years:					
United States . . . . .	3,755	115.8	19,359	597.0	5.2
Northeast . . . . .	806	128.5	4,634	739.0	5.8
Midwest . . . . .	839	112.7	3,783	507.8	4.5
South . . . . .	1,460	126.2	7,590	656.0	5.2
West . . . . .	650	91.1	3,353	469.9	5.2

See footnotes at end of table.

**Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population <sup>1</sup>	Number in thousands	Rate per 1,000 population <sup>1</sup>	
65 years and over:					
United States . . . . .	5,315	359.8	31,074	2,103.6	5.8
Northeast . . . . .	1,217	406.1	7,975	2,660.4	6.6
Midwest . . . . .	1,319	387.3	6,920	2,032.3	5.2
South . . . . .	1,826	345.1	11,030	2,084.3	6.0
West . . . . .	953	309.8	5,149	1,673.1	5.4
Female					
All ages:					
United States . . . . .	20,338	138.8	93,076	635.2	4.6
Northeast . . . . .	4,028	144.0	21,352	763.4	5.3
Midwest . . . . .	4,478	135.0	19,047	574.4	4.3
South . . . . .	8,048	153.1	36,733	698.7	4.6
West . . . . .	3,784	115.3	15,944	485.8	4.2
Under 15 years:					
United States . . . . .	1,128	38.1	5,249	177.3	4.7
Northeast . . . . .	223	42.9	1,046	200.6	4.7
Midwest . . . . .	208	31.3	836	125.9	4.0
South . . . . .	429	40.6	1,995	188.5	4.6
West . . . . .	267	37.2	1,373	191.3	5.1
15–44 years:					
United States . . . . .	7,830	126.6	25,729	415.9	3.3
Northeast . . . . .	1,358	117.3	5,483	473.3	4.0
Midwest . . . . .	1,531	110.2	4,708	339.0	3.1
South . . . . .	3,334	150.2	10,899	491.1	3.3
West . . . . .	1,608	113.2	4,638	326.6	2.9
45–64 years:					
United States . . . . .	3,968	116.0	18,939	553.4	4.8
Northeast . . . . .	779	115.4	4,160	616.5	5.3
Midwest . . . . .	872	112.6	3,795	490.2	4.4
South . . . . .	1,633	132.2	7,895	639.2	4.8
West . . . . .	684	92.8	3,088	418.5	4.5
65 years and over:					
United States . . . . .	7,412	355.8	43,159	2,072.0	5.8
Northeast . . . . .	1,667	376.6	10,663	2,408.4	6.4
Midwest . . . . .	1,868	381.8	9,708	1,983.8	5.2
South . . . . .	2,652	356.0	15,943	2,140.8	6.0
West . . . . .	1,225	301.6	6,845	1,685.2	5.6

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.



**Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment <sup>1</sup>	Medicare	Medicaid	Workers' compensation and other government payments	HMO/PPO <sup>2</sup>	Blue Cross/Blue Shield and other private insurance	Self-pay	Other payments and no charge
United States								
Number of discharges in thousands								
All	33,727	13,191	4,933	680	5,795	6,360	1,429	753
Male	13,389	5,625	1,527	387	2,044	2,525	710	338
Female	20,338	7,566	3,407	293	3,751	3,834	719	415
Under 15 years	2,540	*	966	45	545	669	88	*
15-44 years	10,736	622	2,790	314	2,759	2,814	860	332
45-64 years	7,723	1,443	968	245	1,912	2,360	425	201
65 years and over	12,727	11,114	209	75	580	517	56	51
Northeast								
All	6,990	2,863	922	91	1,141	1,492	238	*
Male	2,962	1,232	376	57	427	626	132	*
Female	4,028	1,631	546	35	715	865	106	*
Under 15 years	512	*	135	*	108	*	18	*
15-44 years	2,008	138	505	41	502	590	146	52
45-64 years	1,584	304	222	32	366	539	63	33
65 years and over	2,885	2,421	60	*	165	182	11	*
Midwest								
All	7,503	3,373	859	*	1,165	1,616	232	54
Male	3,025	1,413	283	68	417	662	119	27
Female	4,478	1,960	576	*	748	954	113	27
Under 15 years	*	*	*	*	*	*	*	*
15-44 years	2,119	128	499	*	554	680	140	29
45-64 years	1,711	319	182	35	460	595	70	16
65 years and over	3,187	2,924	27	*	70	*	*	*
South								
All	12,994	4,831	2,045	288	2,115	2,393	706	*
Male	4,947	2,032	546	147	716	913	341	*
Female	8,048	2,799	1,499	141	1,398	1,480	365	137
Under 15 years	931	*	417	*	183	180	32	*
15-44 years	4,492	265	1,174	145	1,063	1,155	427	113
45-64 years	3,094	610	393	111	673	931	225	54
65 years and over	4,478	3,948	62	19	196	127	22	11
West								
All	6,239	2,124	1,107	189	1,374	859	253	*
Male	2,455	948	322	115	484	324	118	*
Female	3,784	1,176	785	74	890	534	135	*
Under 15 years	*	*	*	*	*	*	*	*
15-44 years	2,117	91	612	78	641	390	147	*
45-64 years	1,334	211	171	67	413	294	68	*
65 years and over	2,178	1,820	60	*	149	78	17	*
United States								
Number of days of care in thousands								
All	164,152	77,880	23,511	3,216	22,074	25,429	5,631	3,429
Male	71,076	33,819	9,310	1,917	8,875	11,048	3,091	1,745
Female	93,076	44,061	14,201	1,299	13,199	14,382	2,540	1,684
Under 15 years	11,531	*48	5,055	255	2,109	2,826	302	*
15-44 years	40,091	3,877	10,956	1,360	8,843	9,683	3,080	1,286
45-64 years	38,298	8,791	6,098	1,143	8,218	9,998	1,952	1,195
65 years and over	74,233	65,164	1,402	458	2,904	2,922	297	242

See footnotes at end of table.

**Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment <sup>1</sup>	Medicare	Medicaid	Workers' compensation and other government payments	HMO/PPO <sup>2</sup>	Blue Cross/Blue Shield and other private insurance	Self-pay	Other payments and no charge
Northeast				Number of days of care in thousands				
All	39,371	19,036	6,144	596	4,684	6,519	1,096	*
Male	18,020	8,574	2,981	355	1,925	2,931	639	*
Female	21,352	10,462	3,163	*	2,759	3,588	456	*
Under 15 years	2,585	*	*	*	384	*	*	*
15–44 years	9,355	1,071	3,061	*	1,722	2,237	640	230
45–64 years	8,794	2,181	1,648	173	1,696	2,398	319	170
65 years and over	18,638	15,783	472	*	882	1,121	64	*
Midwest								
All	33,277	17,691	3,532	*	4,152	5,952	774	272
Male	14,230	7,512	1,371	*	1,665	2,644	420	*
Female	19,047	10,179	2,161	*	2,487	3,309	354	136
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	7,146	651	1,797	*	1,632	2,150	411	151
45–64 years	7,578	1,729	976	*	1,879	2,317	260	89
65 years and over	16,627	15,294	125	*	347	*	*	*
South								
All	63,403	29,173	9,235	1,230	8,476	9,654	2,749	*
Male	26,670	12,434	3,264	695	3,374	4,174	1,428	*
Female	36,733	16,739	5,971	535	5,102	5,480	1,321	500
Under 15 years	4,290	*	2,146	*	*	771	98	*
15–44 years	16,654	1,706	4,159	569	3,608	4,077	1,510	410
45–64 years	15,486	3,563	2,501	491	3,052	4,071	1,031	281
65 years and over	26,973	23,881	429	118	1,037	736	109	44
West								
All	28,101	11,980	4,600	889	4,762	3,304	1,013	*
Male	12,157	5,300	1,694	548	1,911	1,299	604	*
Female	15,944	6,680	2,906	341	2,851	2,005	409	*
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	6,936	450	1,939	339	1,881	1,220	519	*
45–64 years	6,441	1,318	972	301	1,592	1,212	342	*
65 years and over	11,994	10,206	376	*	638	411	84	*
United States				Average length of stay				
All	4.9	5.9	4.8	4.7	3.8	4.0	3.9	4.6
Male	5.3	6.0	6.1	5.0	4.3	4.4	4.4	5.2
Female	4.6	5.8	4.2	4.4	3.5	3.8	3.5	4.1
Under 15 years	4.5	*	5.2	5.7	3.9	4.2	3.4	4.2
15–44 years	3.7	6.2	3.9	4.3	3.2	3.4	3.6	3.9
45–64 years	5.0	6.1	6.3	4.7	4.3	4.2	4.6	5.9
65 years and over	5.8	5.9	6.7	6.1	5.0	5.7	5.3	4.7
Northeast								
All	5.6	6.6	6.7	6.5	4.1	4.4	4.6	4.9
Male	6.1	7.0	7.9	6.2	4.5	4.7	4.8	5.2
Female	5.3	6.4	5.8	7.0	3.9	4.1	4.3	4.5
Under 15 years	5.0	*	7.1	15.8	3.6	4.2	4.0	4.8
15–44 years	4.7	7.8	6.1	5.8	3.4	3.8	4.4	4.4
45–64 years	5.5	7.2	7.4	5.4	4.6	4.4	5.1	5.2
65 years and over	6.5	6.5	7.8	7.8	5.3	6.1	5.9	6.1

See footnotes at end of table.

**Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment <sup>1</sup>	Medicare	Medicaid	Workers' compensation and other government payments	HMO/PPO <sup>2</sup>	Blue Cross/Blue Shield and other private insurance	Self-pay	Other payments and no charge
Midwest				Average length of stay				
All . . . . .	4.4	5.2	4.1	4.5	3.6	3.7	3.3	5.1
Male . . . . .	4.7	5.3	4.9	4.7	4.0	4.0	3.5	5.1
Female . . . . .	4.3	5.2	3.8	4.2	3.3	3.5	3.1	5.0
Under 15 years . . . . .	4.0	*	4.2	3.3	3.6	3.9	3.6	3.3
15–44 years . . . . .	3.4	5.1	3.6	4.3	2.9	3.2	2.9	5.3
45–64 years . . . . .	4.4	5.4	5.4	5.1	4.1	3.9	3.7	5.6
65 years and over . . . . .	5.2	5.2	4.7	4.6	5.0	5.1	*	*
South								
All . . . . .	4.9	6.0	4.5	4.3	4.0	4.0	3.9	4.1
Male . . . . .	5.4	6.1	6.0	4.7	4.7	4.6	4.2	4.6
Female . . . . .	4.6	6.0	4.0	3.8	3.6	3.7	3.6	3.6
Under 15 years . . . . .	4.6	*	5.2	4.0	4.3	4.3	3.0	4.0
15–44 years . . . . .	3.7	6.4	3.5	3.9	3.4	3.5	3.5	3.6
45–64 years . . . . .	5.0	5.8	6.4	4.4	4.5	4.4	4.6	5.2
65 years and over . . . . .	6.0	6.0	6.9	6.4	5.3	5.8	4.9	3.9
West								
All . . . . .	4.5	5.6	4.2	4.7	3.5	3.8	4.0	4.7
Male . . . . .	5.0	5.6	5.3	4.8	3.9	4.0	5.1	*
Female . . . . .	4.2	5.7	3.7	4.6	3.2	3.8	3.0	4.0
Under 15 years . . . . .	4.5	*	5.0	5.1	3.8	4.8	3.4	3.9
15–44 years . . . . .	3.3	5.0	3.2	4.3	2.9	3.1	3.5	3.6
45–64 years . . . . .	4.8	6.2	5.7	4.5	3.9	4.1	5.0	*
65 years and over . . . . .	5.5	5.6	6.2	6.0	4.3	5.3	4.9	4.1

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Includes discharges for which no expected source of payment was provided.<sup>2</sup>HMO/PPO is defined as health maintenance organization and preferred provider organization.



**Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by bed size of hospital, according to sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Number of discharges in thousands						
All . . . . .	33,727	7,903	8,353	6,767	7,446	3,257
Sex						
Male . . . . .	13,389	3,011	3,182	2,770	3,011	1,413
Female . . . . .	20,338	4,891	5,172	3,997	4,434	1,844
Age						
Under 15 years . . . . .	2,540	502	469	678	688	203
15–44 years . . . . .	10,736	2,450	2,725	2,019	2,431	1,111
45–64 years . . . . .	7,723	1,633	1,834	1,535	1,810	911
65 years and over . . . . .	12,727	3,317	3,326	2,535	2,517	1,032
Region						
Northeast . . . . .	6,990	1,599	1,680	1,064	1,798	849
Midwest . . . . .	7,503	2,530	1,582	1,364	1,292	735
South . . . . .	12,994	2,916	3,148	2,536	2,920	1,474
West . . . . .	6,239	858	1,943	1,803	1,436	199
Percent distribution of discharges						
All . . . . .	100.0	23.4	24.8	20.1	22.1	9.7
Sex						
Male . . . . .	100.0	22.5	23.8	20.7	22.5	10.6
Female . . . . .	100.0	24.1	25.4	19.7	21.8	9.1
Age						
Under 15 years . . . . .	100.0	19.8	18.5	26.7	27.1	8.0
15–44 years . . . . .	100.0	22.8	25.4	18.8	22.6	10.4
45–64 years . . . . .	100.0	21.2	23.8	19.9	23.4	11.8
65 years and over . . . . .	100.0	26.1	26.1	19.9	19.8	8.1
Region						
Northeast . . . . .	100.0	22.9	24.0	15.2	25.7	12.2
Midwest . . . . .	100.0	33.7	21.1	18.2	17.2	9.8
South . . . . .	100.0	22.4	24.2	19.5	22.5	11.3
West . . . . .	100.0	13.8	31.2	28.9	23.0	3.2
Number of days of care in thousands						
All . . . . .	164,152	33,883	40,270	33,085	38,525	18,390
Sex						
Male . . . . .	71,076	13,694	16,782	14,615	17,176	8,809
Female . . . . .	93,076	20,189	23,488	18,469	21,349	9,581
Age						
Under 15 years . . . . .	11,531	1,805	2,005	3,180	3,533	1,007
15–44 years . . . . .	40,091	8,799	9,474	7,167	9,580	5,071
45–64 years . . . . .	38,298	7,058	8,851	7,431	9,528	5,430
65 years and over . . . . .	74,233	16,221	19,940	15,306	15,884	6,882
Region						
Northeast . . . . .	39,371	8,491	10,080	5,805	9,976	5,020
Midwest . . . . .	33,277	9,500	7,391	6,358	6,187	3,841
South . . . . .	63,403	12,433	14,805	12,299	15,386	8,480
West . . . . .	28,101	3,459	7,994	8,623	6,976	1,049

**Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by bed size of hospital, according to sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Percent distribution of days of care						
All .....	100.0	20.6	24.5	20.2	23.5	11.2
Sex						
Male .....	100.0	19.3	23.6	20.6	24.2	12.4
Female .....	100.0	21.7	25.2	19.8	22.9	10.3
Age						
Under 15 years .....	100.0	15.7	17.4	27.6	30.6	8.7
15–44 years .....	100.0	21.9	23.6	17.9	23.9	12.6
45–64 years .....	100.0	18.4	23.1	19.4	24.9	14.2
65 years and over .....	100.0	21.9	26.9	20.6	21.4	9.3
Region						
Northeast .....	100.0	21.6	25.6	14.7	25.3	12.8
Midwest .....	100.0	28.5	22.2	19.1	18.6	11.5
South .....	100.0	19.6	23.4	19.4	24.3	13.4
West .....	100.0	12.3	28.4	30.7	24.8	3.7
Average length of stay in days						
All .....	4.9	4.3	4.8	4.9	5.2	5.6
Sex						
Male .....	5.3	4.5	5.3	5.3	5.7	6.2
Female .....	4.6	4.1	4.5	4.6	4.8	5.2
Age						
Under 15 years .....	4.5	3.6	4.3	4.7	5.1	5.0
15–44 years .....	3.7	3.6	3.5	3.5	3.9	4.6
45–64 years .....	5.0	4.3	4.8	4.8	5.3	6.0
65 years and over .....	5.8	4.9	6.0	6.0	6.3	6.7
Region						
Northeast .....	5.6	5.3	6.0	5.5	5.5	5.9
Midwest .....	4.4	3.8	4.7	4.7	4.8	5.2
South .....	4.9	4.3	4.7	4.8	5.3	5.8
West .....	4.5	4.0	4.1	4.8	4.9	5.3

**Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by hospital ownership, according to sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	All	Proprietary	Government	Nonprofit
Number of discharges in thousands				
All	33,727	4,445	4,153	25,129
Sex				
Male	13,389	1,668	1,671	10,050
Female	20,338	2,777	2,482	15,079
Age				
Under 15 years	2,540	270	305	1,964
15–44 years	10,736	1,621	1,539	7,577
45–64 years	7,723	954	1,032	5,738
65 years and over	12,727	1,599	1,277	9,851
Region				
Northeast	6,990	104	306	6,580
Midwest	7,503	233	551	6,719
South	12,994	3,304	2,441	7,250
West	6,239	804	855	4,580
Percent distribution of discharges				
All	100.0	13.2	12.3	74.5
Sex				
Male	100.0	12.5	12.5	75.1
Female	100.0	13.7	12.2	74.1
Age				
Under 15 years	100.0	10.7	12.0	77.4
15–44 years	100.0	15.1	14.3	70.6
45–64 years	100.0	12.4	13.4	74.3
65 years and over	100.0	12.6	10.0	77.4
Region				
Northeast	100.0	1.5	4.4	94.1
Midwest	100.0	3.1	7.3	89.6
South	100.0	25.4	18.8	55.8
West	100.0	12.9	13.7	73.4
Number of days of care in thousands				
All	164,152	22,337	19,933	121,882
Sex				
Male	71,076	9,348	9,200	52,528
Female	93,076	12,989	10,733	69,354
Age				
Under 15 years	11,531	1,352	1,184	8,994
15–44 years	40,091	6,089	6,041	27,961
45–64 years	38,298	4,970	5,601	27,726
65 years and over	74,233	9,926	7,107	57,200
Region				
Northeast	39,371	473	1,855	37,044
Midwest	33,277	1,134	2,486	29,658
South	63,403	17,018	11,720	34,665
West	28,101	3,712	3,873	20,515

**Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay by hospital ownership, according to sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Sex, age, and region	All	Proprietary	Government	Nonprofit
Percent distribution of days of care				
All .....	100.0	13.6	12.1	74.2
Sex				
Male .....	100.0	13.2	12.9	73.9
Female .....	100.0	14.0	11.5	74.5
Age				
Under 15 years .....	100.0	11.7	10.3	78.0
15–44 years .....	100.0	15.2	15.1	69.7
45–64 years .....	100.0	13.0	14.6	72.4
65 years and over .....	100.0	13.4	9.6	77.1
Region				
Northeast .....	100.0	1.2	4.7	94.1
Midwest .....	100.0	3.4	7.5	89.1
South .....	100.0	26.8	18.5	54.7
West .....	100.0	13.2	13.8	73.0
Average length of stay in days				
All .....	4.9	5.0	4.8	4.9
Sex				
Male .....	5.3	5.6	5.5	5.2
Female .....	4.6	4.7	4.3	4.6
Age				
Under 15 years .....	4.5	5.0	3.9	4.6
15–44 years .....	3.7	3.8	3.9	3.7
45–64 years .....	5.0	5.2	5.4	4.8
65 years and over .....	5.8	6.2	5.6	5.8
Region				
Northeast .....	5.6	4.6	6.1	5.6
Midwest .....	4.4	4.9	4.5	4.4
South .....	4.9	5.2	4.8	4.8
West .....	4.5	4.6	4.5	4.5



**Table 7. Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Admission type and source	All ages	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over
Number of discharges in thousands							
All types and sources . . . . .	33,727	13,389	20,338	2,540	10,736	7,723	12,727
Type of admission							
Emergency . . . . .	11,494	5,211	6,283	742	2,822	2,840	5,090
Urgent . . . . .	7,778	2,857	4,922	629	2,894	1,544	2,712
Elective . . . . .	7,010	2,340	4,670	325	2,643	1,737	2,305
Item not available . . . . .	7,445	2,981	4,463	844	2,377	1,602	2,621
Source of admission							
Physician referral <sup>1</sup> . . . . .	12,595	4,160	8,436	894	5,110	2,771	3,820
Emergency department . . . . .	12,198	5,565	6,633	647	2,826	3,005	5,720
Transfer from a hospital . . . . .	902	431	472	109	168	222	403
Transfer from other health facility <sup>2</sup> . . . . .	554	229	325	53	70	88	343
Other sources <sup>3</sup> . . . . .	681	255	426	40	364	148	129
Item not available . . . . .	6,795	2,749	4,046	796	2,198	1,490	2,312
Percent distribution							
All types and sources . . . . .	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Type of admission							
Emergency . . . . .	34.1	38.9	30.9	29.2	26.3	36.8	40.0
Urgent . . . . .	23.1	21.3	24.2	24.8	27.0	20.0	21.3
Elective . . . . .	20.8	17.5	23.0	12.8	24.6	22.5	18.1
Item not available . . . . .	22.1	22.3	21.9	33.3	22.1	20.7	20.6
Source of admission							
Physician referral <sup>1</sup> . . . . .	37.3	31.1	41.5	35.2	47.6	35.9	30.0
Emergency department . . . . .	36.2	41.6	32.6	25.5	26.3	38.9	44.9
Transfer from a hospital . . . . .	2.7	3.2	2.3	4.3	1.6	2.9	3.2
Transfer from other health facility <sup>2</sup> . . . . .	1.6	1.7	1.6	2.1	0.7	1.1	2.7
Other sources <sup>3</sup> . . . . .	2.0	1.9	2.1	1.6	3.4	1.9	1.0
Item not available . . . . .	20.1	20.5	19.9	31.3	20.5	19.3	18.2

<sup>1</sup>Referral from personal physician, clinic physician, or HMO physician.<sup>2</sup>Transfer from skilled nursing facility or other health facility.<sup>3</sup>Includes admissions directed by a court of law or requested by a law enforcement agent and other sources.**Table 8. Number and rate of deaths for discharges from short-stay hospitals, by sex and age: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Age	Number of deaths in thousands			Rate per 100 discharges		
	Both sexes	Male	Female	Both sexes	Male	Female
All ages . . . . .	811	398	414	2.4	3.0	2.0
Under 65 years . . . . .	209	119	91	1.0	1.5	0.7
Under 15 years . . . . .	18	11	7	0.7	0.7	0.6
15-44 years . . . . .	46	25	21	0.4	0.9	0.3
45-64 years . . . . .	145	83	62	1.9	2.2	1.6
65 years and over . . . . .	602	279	323	4.7	5.3	4.4
65-74 years . . . . .	151	82	68	3.2	3.7	2.8
75-84 years . . . . .	258	127	131	4.9	5.7	4.3
85 years and over . . . . .	193	70	123	6.9	7.9	6.4

**Table 9. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Category of first-listed diagnosis and ICD-9-CM code	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 10,000 population <sup>1</sup>	Number in thousands	Rate per 10,000 population <sup>1</sup>	
All conditions <sup>2</sup> . . . . .	33,727	1,174.6	164,152	5,717.1	4.9
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4,446	154.8	20,351	708.8	4.6
Acute myocardial infarction . . . . . 410	818	28.5	4,586	159.7	5.6
Coronary atherosclerosis . . . . . 414.0	1,096	38.2	3,885	135.3	3.5
Other ischemic heart disease . . . . . 411-413,414.1-414.9	211	7.3	546	19.0	2.6
Cardiac dysrhythmias . . . . . 427	788	27.5	2,830	98.6	3.6
Congestive heart failure . . . . . 428.0	970	33.8	5,255	183.0	5.4
Females with deliveries . . . . . V27	3,951	137.6	10,207	355.5	2.6
Psychoses . . . . . 290-299	1,704	59.4	13,666	476.0	8.0
Pneumonia . . . . . 480-486	1,312	45.7	7,539	262.6	5.7
Malignant neoplasms . . . . . 140-208,230-234	1,208	42.1	8,515	296.6	7.1
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	159	5.5	1,381	48.1	8.7
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	160	5.6	1,198	41.7	7.5
Malignant neoplasm of breast . . . . . 174-175,198.81	85	3.0	246	8.6	2.9
Fractures, all sites <sup>3</sup> . . . . . 800-829	995	34.7	5,388	187.7	5.4
Fracture of neck of femur <sup>3</sup> . . . . . 820	315	11.0	2,064	71.9	6.6
Cerebrovascular disease . . . . . 430-438	942	32.8	4,974	173.2	5.3
Diabetes mellitus . . . . . 250	577	20.1	2,893	100.8	5.0
Osteoarthritis and allied disorders . . . . . 715	568	19.8	2,376	82.7	4.2
Chronic bronchitis . . . . . 491	520	18.1	2,749	95.7	5.3
Volume depletion . . . . . 276.5	508	17.7	1,918	66.8	3.8
Asthma . . . . . 493	484	16.8	1,531	53.3	3.2
Benign neoplasms . . . . . 210-229	427	14.9	1,423	49.5	3.3
Cellulitis and abscess . . . . . 681-682	422	14.7	2,188	76.2	5.2
Cholelithiasis . . . . . 574	359	12.5	1,450	50.5	4.0
Intervertebral disc disorder . . . . . 722	353	12.3	992	34.5	2.8
Septicemia . . . . . 038	341	11.9	2,849	99.2	8.3
Noninfectious enteritis and colitis . . . . . 555-558	310	10.8	1,441	50.2	4.7

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>Includes data for diagnostic conditions not shown in table.<sup>3</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 10. Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
	Discharges in thousands			
All conditions <sup>1</sup> . . . . .	12,727	4,642	5,271	2,814
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	2,843	1,048	1,180	615
Acute myocardial infarction . . . . . 410	501	181	200	120
Coronary atherosclerosis . . . . . 414.0	615	307	241	68
Other ischemic heart disease . . . . . 411-413,414.1-414.9	107	46	43	19
Cardiac dysrhythmias . . . . . 427	562	190	244	128
Congestive heart failure . . . . . 428.0	717	214	293	210
Pneumonia . . . . . 480-486	776	217	326	233
Cerebrovascular disease . . . . . 430-438	669	222	295	152
Malignant neoplasms . . . . . 140-208,230-234	643	301	260	81
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	107	43	47	18
Fractures, all sites <sup>2</sup> . . . . . 800-829	524	110	210	204
Fracture of neck of femur <sup>2</sup> . . . . . 820	282	46	110	126
Osteoarthritis and allied disorders . . . . . 715	356	177	154	24
Chronic bronchitis . . . . . 491	343	160	139	43
Volume depletion . . . . . 276.5	249	59	112	78
Psychoses . . . . . 290-299	224	78	94	51
Septicemia . . . . . 038	219	60	96	63
Diabetes mellitus . . . . . 250	193	95	77	21
Diverticula of intestine . . . . . 562	149	48	67	33
Cellulitis and abscess . . . . . 681-682	147	56	58	32
Cholelithiasis . . . . . 574	133	63	47	23
Noninfectious enteritis and colitis . . . . . 555-558	100	35	42	24
	Discharge rate per 10,000 population <sup>3</sup>			
All conditions <sup>1</sup> . . . . .	3,574.9	2,540.2	4,138.9	6,127.6
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	798.7	573.6	926.6	1,339.4
Acute myocardial infarction . . . . . 410	140.8	99.2	156.7	262.0
Coronary atherosclerosis . . . . . 414.0	172.9	167.7	189.1	148.2
Other ischemic heart disease . . . . . 411-413,414.1-414.9	30.0	25.0	33.4	40.6
Cardiac dysrhythmias . . . . . 427	157.8	104.1	191.4	278.6
Congestive heart failure . . . . . 428.0	201.4	117.3	229.8	457.3
Pneumonia . . . . . 480-486	218.0	118.6	256.3	507.1
Cerebrovascular disease . . . . . 430-438	187.9	121.3	231.4	331.9
Malignant neoplasms . . . . . 140-208,230-234	180.5	164.6	204.5	177.4
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	30.2	23.4	36.8	38.6
Fractures, all sites <sup>2</sup> . . . . . 800-829	147.2	60.2	165.1	443.7
Fracture of neck of femur <sup>2</sup> . . . . . 820	79.3	25.1	86.7	274.8
Osteoarthritis and allied disorders . . . . . 715	99.9	97.0	121.0	52.7
Chronic bronchitis . . . . . 491	96.3	87.7	109.2	94.4
Volume depletion . . . . . 276.5	70.0	32.5	88.0	169.3
Psychoses . . . . . 290-299	62.9	42.9	73.8	112.0
Septicemia . . . . . 038	61.4	32.8	75.2	137.2
Diabetes mellitus . . . . . 250	54.1	52.0	60.1	45.8
Diverticula of intestine . . . . . 562	41.7	26.4	52.8	72.1
Cellulitis and abscess . . . . . 681-682	41.2	30.6	45.9	70.1
Cholelithiasis . . . . . 574	37.4	34.5	36.7	50.7
Noninfectious enteritis and colitis . . . . . 555-558	28.2	19.0	32.8	52.0
	Average length of stay in days			
All conditions <sup>1</sup> . . . . .	5.8	5.6	6.0	6.0
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4.9	4.6	5.1	5.1
Acute myocardial infarction . . . . . 410	6.4	6.2	6.5	6.6
Coronary atherosclerosis . . . . . 414.0	3.7	3.6	4.0	3.5
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2.9	3.0	2.6	3.3
Cardiac dysrhythmias . . . . . 427	3.9	3.3	4.1	4.4
Congestive heart failure . . . . . 428.0	5.5	5.6	5.8	5.1
Pneumonia . . . . . 480-486	6.3	6.1	6.3	6.5
Cerebrovascular disease . . . . . 430-438	5.1	4.6	5.5	5.3

See footnotes at end of table.

**Table 10. Number and rate of discharges from short-stay hospitals with average length of stay, for patients 65 years of age and over by age and selected first-listed diagnostic categories: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
	Average length of stay in days			
Malignant neoplasms . . . . . 140-208,230-234	7.5	7.4	7.7	7.1
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	9.4	8.9	9.4	10.5
Fractures, all sites <sup>2</sup> . . . . . 800-829	6.1	6.0	6.2	6.1
Fracture of neck of femur <sup>2</sup> . . . . . 820	6.6	6.5	6.7	6.5
Osteoarthritis and allied disorders . . . . . 715	4.4	4.3	4.5	4.7
Chronic bronchitis . . . . . 491	5.4	5.2	5.6	5.8
Volume depletion . . . . . 276.5	4.9	4.7	4.9	5.0
Psychoses . . . . . 290-299	9.4	10.5	9.4	7.7
Septicemia . . . . . 038	8.3	8.6	8.3	8.1
Diabetes mellitus . . . . . 250	5.9	5.8	6.2	5.1
Diverticula of intestine . . . . . 562	6.1	5.4	6.2	6.9
Cellulitis and abscess . . . . . 681-682	6.1	5.6	6.4	6.6
Cholelithiasis . . . . . 574	5.5	4.4	5.7	8.4
Noninfectious enteritis and colitis . . . . . 555-558	6.3	6.0	6.0	7.4

<sup>1</sup>Includes data for diagnostic conditions not shown in table.<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.<sup>3</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.

**Table 11. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
		Number in thousands			
All conditions . . . . .	33,727	2,540	10,736	7,723	12,727
Infectious and parasitic diseases . . . . . 001-139	877	156	204	185	332
Septicemia . . . . . 038	341	12	34	76	219
Neoplasms . . . . . 140-239	1,682	33	313	611	725
Malignant neoplasms . . . . . 140-208,230-234	1,208	25	121	419	643
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	159	*	5	46	107
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	160	*	*5	55	100
Malignant neoplasm of breast . . . . . 174-175,198.81	85	*	11	36	38
Benign neoplasms . . . . . 210-229	427	*	183	178	61
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	1,619	187	328	454	649
Diabetes mellitus . . . . . 250	577	33	147	205	193
Volume depletion . . . . . 276.5	508	129	53	77	249
Diseases of the blood and blood-forming organs . . . . . 280-289	446	71	106	81	188
Mental disorders . . . . . 290-319	2,464	149	1,422	620	273
Psychoses . . . . . 290-299	1,704	*	957	431	224
Alcohol dependence syndrome . . . . . 303	145	*	87	53	*
Diseases of the nervous system and sense organs . . . . . 320-389	518	81	134	114	189
Diseases of the circulatory system . . . . . 390-459	6,373	31	434	1,871	4,037
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4,446	17	273	1,313	2,843
Acute myocardial infarction . . . . . 410	818	*	57	259	501
Coronary atherosclerosis . . . . . 414.0	1,096	*	48	432	615
Other ischemic heart disease . . . . . 411-413,414.1-414.9	211	*	18	86	107
Cardiac dysrhythmias . . . . . 427	788	*	45	174	562
Congestive heart failure . . . . . 428.0	970	*	32	219	717
Cerebrovascular disease . . . . . 430-438	942	*	40	229	669
Diseases of the respiratory system . . . . . 460-519	3,542	730	382	697	1,732
Acute bronchitis and bronchiolitis . . . . . 466	279	209	8	22	40
Pneumonia . . . . . 480-486	1,312	204	116	216	776
Chronic bronchitis . . . . . 491	520	*	16	161	343
Asthma . . . . . 493	484	187	109	109	80
Diseases of the digestive system . . . . . 520-579	3,320	216	839	955	1,310
Appendicitis . . . . . 540-543	295	70	155	50	21
Noninfectious enteritis and colitis . . . . . 555-558	310	54	89	66	100
Diverticula of intestine . . . . . 562	262	*	36	77	149
Cholelithiasis . . . . . 574	359	*	116	109	133
Diseases of the genitourinary system . . . . . 580-629	1,817	88	580	458	690
Calculus of kidney and ureter . . . . . 592	176	*	81	62	29
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	528	*	524	*	...
Diseases of the skin and subcutaneous tissue . . . . . 680-709	601	*	158	151	199
Cellulitis and abscess . . . . . 681-682	422	41	109	126	147
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	1,736	41	326	598	770
Osteoarthritis and allied disorders . . . . . 715	568	*	20	193	356
Intervertebral disc disorders . . . . . 722	353	*	133	153	67
Congenital anomalies . . . . . 740-759	178	124	31	15	*7
Certain conditions originating in the perinatal period . . . . . 760-779	166	165	*	*	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	283	63	99	68	54
Injury and poisoning . . . . . 800-999	2,697	233	780	621	1,063
Fractures, all sites <sup>2</sup> . . . . . 800-829	995	68	239	165	524
Fracture of neck of femur <sup>2</sup> . . . . . 820	315	*	6	24	282
Poisonings . . . . . 960-989	214	18	120	48	28
Supplementary classifications . . . . . V01-V83	4,880	75	4,073	224	508
Females with deliveries . . . . . V27	3,951	12	3,934	*5	...

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.



**Table 12. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Rate per 10,000 population <sup>1</sup>				
		Under 15 years	15-44 years	45-64 years	65 years and over	
All conditions . . . . .	1,174.6	418.7	864.2	1,158.9	3,574.9	
Infectious and parasitic diseases . . . . . 001-139	30.5	25.8	16.4	27.8	93.1	
Septicemia . . . . . 038	11.9	1.9	2.8	11.5	61.4	
Neoplasms . . . . . 140-239	58.6	5.5	25.2	91.7	203.6	
Malignant neoplasms . . . . . 140-208,230-234	42.1	4.1	9.7	62.9	180.5	
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	5.5	*	0.4	6.8	30.2	
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	5.6	*	*0.4	8.3	28.0	
Malignant neoplasm of breast . . . . . 174-175,198.81	3.0	*	0.9	5.5	10.6	
Benign neoplasms . . . . . 210-229	14.9	*	14.7	26.7	17.0	
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	56.4	30.8	26.4	68.1	182.4	
Diabetes mellitus . . . . . 250	20.1	5.5	11.8	30.7	54.1	
Volume depletion . . . . . 276.5	17.7	21.3	4.2	11.6	70.0	
Diseases of the blood and blood-forming organs . . . . . 280-289	15.5	11.7	8.6	12.2	52.7	
Mental disorders . . . . . 290-319	85.8	24.5	114.5	93.0	76.8	
Psychoses . . . . . 290-299	59.4	*	77.0	64.7	62.9	
Alcohol dependence syndrome . . . . . 303	5.1	*	7.0	8.0	*	
Diseases of the nervous system and sense organs . . . . . 320-389	18.0	13.3	10.8	17.1	53.2	
Diseases of the circulatory system . . . . . 390-459	222.0	5.2	35.0	280.7	1,133.9	
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	154.8	2.7	22.0	197.0	798.7	
Acute myocardial infarction . . . . . 410	28.5	*	4.6	38.9	140.8	
Coronary atherosclerosis . . . . . 414.0	38.2	*	3.9	64.8	172.9	
Other ischemic heart disease . . . . . 411-413,414.1-414.9	7.3	*	1.5	12.8	30.0	
Cardiac dysrhythmias . . . . . 427	27.5	*	3.6	26.2	157.8	
Congestive heart failure . . . . . 428.0	33.8	*	2.6	32.8	201.4	
Cerebrovascular disease . . . . . 430-438	32.8	*	3.2	34.4	187.9	
Diseases of the respiratory system . . . . . 460-519	123.4	120.4	30.7	104.6	486.5	
Acute bronchitis and bronchiolitis . . . . . 466	9.7	34.4	0.7	3.3	11.2	
Pneumonia . . . . . 480-486	45.7	33.6	9.3	32.4	218.0	
Chronic bronchitis . . . . . 491	18.1	*	1.3	24.1	96.3	
Asthma . . . . . 493	16.8	30.8	8.7	16.3	22.5	
Diseases of the digestive system . . . . . 520-579	115.6	35.6	67.6	143.3	367.9	
Appendicitis . . . . . 540-543	10.3	11.5	12.5	7.5	5.8	
Noninfectious enteritis and colitis . . . . . 555-558	10.8	9.0	7.2	9.9	28.2	
Diverticula of intestine . . . . . 562	9.1	*	2.9	11.6	41.7	
Cholelithiasis . . . . . 574	12.5	*	9.3	16.3	37.4	
Diseases of the genitourinary system . . . . . 580-629	63.3	14.6	46.7	68.7	193.9	
Calculus of kidney and ureter . . . . . 592	6.1	*	6.5	9.3	8.2	
Complications of pregnancy, childbirth, and the puerperium <sup>2</sup> . . . . . 630-677	18.4	*	42.2	*	...	
Diseases of the skin and subcutaneous tissue . . . . . 680-709	20.9	*	12.7	22.6	56.0	
Cellulitis and abscess . . . . . 681-682	14.7	6.7	8.8	18.9	41.2	
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	60.5	6.8	26.2	89.8	216.4	
Osteoarthritis and allied disorders . . . . . 715	19.8	*	1.6	29.0	99.9	
Intervertebral disc disorders . . . . . 722	12.3	*	10.7	22.9	18.7	
Congenital anomalies . . . . . 740-759	6.2	20.5	2.5	2.2	*1.9	
Certain conditions originating in the perinatal period . . . . . 760-779	5.8	27.2	*	*	*	
Symptoms, signs, and ill-defined conditions . . . . . 780-799	9.9	10.3	8.0	10.1	15.1	
Injury and poisoning . . . . . 800-999	93.9	38.5	62.8	93.2	298.5	
Fractures, all sites <sup>3</sup> . . . . . 800-829	34.7	11.2	19.2	24.7	147.2	
Fracture of neck of femur <sup>3</sup> . . . . . 820	11.0	*	0.5	3.7	79.3	
Poisonings . . . . . 960-989	7.4	2.9	9.6	7.2	7.9	
Supplementary classifications . . . . . V01-V83	170.0	12.4	327.9	33.6	142.8	
Females with deliveries . . . . . V27	137.6	2.0	316.7	*0.7	...	

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>3</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 13. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions . . . . .	4.9	4.5	3.7	5.0	5.8
Infectious and parasitic diseases . . . . . 001-139	6.4	3.6	5.9	6.7	7.8
Septicemia . . . . . 038	8.3	4.4	9.7	8.4	8.3
Neoplasms . . . . . 140-239	6.1	9.7	4.3	5.3	7.3
Malignant neoplasms . . . . . 140-208,230-234	7.1	12.1	6.6	6.2	7.5
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	8.7	*	8.0	7.2	9.4
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	7.5	*	*5.1	7.3	7.7
Malignant neoplasm of breast . . . . . 174-175,198.81	2.9	*	3.4	2.6	3.0
Benign neoplasms . . . . . 210-229	3.3	2.7	2.7	3.3	5.5
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	4.3	2.9	3.5	4.4	5.0
Diabetes mellitus . . . . . 250	5.0	3.1	4.0	5.2	5.9
Volume depletion . . . . . 276.5	3.8	2.0	2.6	3.9	4.9
Diseases of the blood and blood-forming organs . . . . . 280-289	4.4	3.4	5.1	4.2	4.3
Mental disorders . . . . . 290-319	7.1	9.2	6.5	7.4	8.5
Psychoses . . . . . 290-299	8.0	7.8	7.4	8.8	9.4
Alcohol dependence syndrome . . . . . 303	5.8	*	6.1	5.0	8.2
Diseases of the nervous system and sense organs . . . . . 320-389	5.1	4.3	4.4	5.9	5.5
Diseases of the circulatory system . . . . . 390-459	4.7	6.6	3.9	4.2	5.0
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4.6	5.6	3.7	4.0	4.9
Acute myocardial infarction . . . . . 410	5.6	*	3.9	4.5	6.4
Coronary atherosclerosis . . . . . 414.0	3.5	*	3.0	3.3	3.7
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2.6	*	2.1	2.3	2.9
Cardiac dysrhythmias . . . . . 427	3.6	4.0	2.3	3.0	3.9
Congestive heart failure . . . . . 428.0	5.4	*	4.8	5.1	5.5
Cerebrovascular disease . . . . . 430-438	5.3	*9.6	5.3	5.6	5.1
Diseases of the respiratory system . . . . . 460-519	5.3	3.2	4.4	5.5	6.3
Acute bronchitis and bronchiolitis . . . . . 466	3.4	3.2	3.5	3.4	4.4
Pneumonia . . . . . 480-486	5.7	3.9	4.9	6.0	6.3
Chronic bronchitis . . . . . 491	5.3	*	4.6	5.0	5.4
Asthma . . . . . 493	3.2	2.3	2.9	3.8	4.8
Diseases of the digestive system . . . . . 520-579	4.9	3.1	3.9	4.8	5.9
Appendicitis . . . . . 540-543	3.2	3.3	2.6	3.6	6.4
Noninfectious enteritis and colitis . . . . . 555-558	4.7	2.4	4.1	4.7	6.3
Diverticula of intestine . . . . . 562	5.8	*	5.2	5.4	6.1
Cholelithiasis . . . . . 574	4.0	*	2.9	3.4	5.5
Diseases of the genitourinary system . . . . . 580-629	3.7	3.2	2.7	3.3	4.8
Calculus of kidney and ureter . . . . . 592	2.2	*	2.1	2.2	2.7
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	2.7	*	2.7	*	...
Diseases of the skin and subcutaneous tissue . . . . . 680-709	5.3	3.0	4.4	6.0	6.5
Cellulitis and abscess . . . . . 681-682	5.2	2.9	4.2	5.7	6.1
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	3.9	3.5	3.2	3.5	4.6
Osteoarthritis and allied disorders . . . . . 715	4.2	*	3.4	3.9	4.4
Intervertebral disc disorders . . . . . 722	2.8	*	2.4	2.7	3.8
Congenital anomalies . . . . . 740-759	5.8	6.4	4.1	4.2	*5.7
Certain conditions originating in the perinatal period . . . . . 760-779	11.1	11.2	*	*	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	2.7	2.9	2.0	2.3	4.0
Injury and poisoning . . . . . 800-999	5.3	4.2	4.4	5.7	6.0
Fractures, all sites <sup>2</sup> . . . . . 800-829	5.4	3.1	4.6	5.3	6.1
Fracture of neck of femur <sup>2</sup> . . . . . 820	6.6	*	6.0	7.1	6.6
Poisonings . . . . . 960-989	2.7	2.7	2.2	3.3	3.9
Supplementary classifications . . . . . V01-V83	3.8	5.4	2.7	7.4	10.9
Females with deliveries . . . . . V27	2.6	2.7	2.6	*2.6	...

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 14. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
All conditions . . . . .	33,727	13,389	20,338
Infectious and parasitic diseases . . . . . 001-139	877	417	460
Septicemia . . . . . 038	341	156	186
Neoplasms . . . . . 140-239	1,682	650	1,033
Malignant neoplasms . . . . . 140-208,230-234	1,208	579	629
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	159	76	83
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	160	86	74
Malignant neoplasm of breast . . . . . 174-175,198.81	85	*	85
Benign neoplasms . . . . . 210-229	427	49	378
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	1,619	661	958
Diabetes mellitus . . . . . 250	577	283	294
Volume depletion . . . . . 276.5	508	204	304
Diseases of the blood and blood-forming organs . . . . . 280-289	446	192	254
Mental disorders . . . . . 290-319	2,464	1,236	1,227
Psychoses . . . . . 290-299	1,704	817	887
Alcohol dependence syndrome . . . . . 303	145	101	44
Diseases of the nervous system and sense organs . . . . . 320-389	518	217	301
Diseases of the circulatory system . . . . . 390-459	6,373	3,209	3,164
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4,446	2,319	2,127
Acute myocardial infarction . . . . . 410	818	480	338
Coronary atherosclerosis . . . . . 414.0	1,096	666	429
Other ischemic heart disease . . . . . 411-413,414.1-414.9	211	103	108
Cardiac dysrhythmias . . . . . 427	788	377	411
Congestive heart failure . . . . . 428.0	970	441	529
Cerebrovascular disease . . . . . 430-438	942	432	509
Diseases of the respiratory system . . . . . 460-519	3,542	1,678	1,864
Acute bronchitis and bronchiolitis . . . . . 466	279	152	127
Pneumonia . . . . . 480-486	1,312	618	694
Chronic bronchitis . . . . . 491	520	230	291
Asthma . . . . . 493	484	196	288
Diseases of the digestive system . . . . . 520-579	3,320	1,466	1,854
Appendicitis . . . . . 540-543	295	167	129
Noninfectious enteritis and colitis . . . . . 555-558	310	128	182
Diverticula of intestine . . . . . 562	262	106	157
Cholelithiasis . . . . . 574	359	113	246
Diseases of the genitourinary system . . . . . 580-629	1,817	560	1,257
Calculus of kidney and ureter . . . . . 592	176	99	77
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	528	...	528
Diseases of the skin and subcutaneous tissue . . . . . 680-709	601	299	303
Cellulitis and abscess . . . . . 681-682	422	211	212
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	1,736	747	989
Osteoarthritis and allied disorders . . . . . 715	568	221	347
Intervertebral disc disorders . . . . . 722	353	180	174
Congenital anomalies . . . . . 740-759	178	97	80
Certain conditions originating in the perinatal period . . . . . 760-779	166	93	73
Symptoms, signs, and ill-defined conditions . . . . . 780-799	283	132	151
Injury and poisoning . . . . . 800-999	2,697	1,331	1,366
Fractures, all sites <sup>2</sup> . . . . . 800-829	995	433	562
Fracture of neck of femur <sup>2</sup> . . . . . 820	315	85	230
Poisonings . . . . . 960-989	214	96	118
Supplementary classifications . . . . . V01-V83	4,880	403	4,477
Females with deliveries . . . . . V27	3,951	...	3,951

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 15. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Rate per 10,000 population <sup>1</sup>		
All conditions . . . . .	1,174.6	952.3	1,388.0
Infectious and parasitic diseases . . . . . 001-139	30.5	29.7	31.4
Septicemia . . . . . 038	11.9	11.1	12.7
Neoplasms . . . . . 140-239	58.6	46.2	70.5
Malignant neoplasms . . . . . 140-208,230-234	42.1	41.2	42.9
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	5.5	5.4	5.6
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	5.6	6.1	5.1
Malignant neoplasm of breast . . . . . 174-175,198.81	3.0	*	5.8
Benign neoplasms . . . . . 210-229	14.9	3.5	25.8
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	56.4	47.0	65.4
Diabetes mellitus . . . . . 250	20.1	20.1	20.1
Volume depletion . . . . . 276.5	17.7	14.5	20.7
Diseases of the blood and blood-forming organs . . . . . 280-289	15.5	13.7	17.3
Mental disorders . . . . . 290-319	85.8	87.9	83.8
Psychoses . . . . . 290-299	59.4	58.1	60.5
Alcohol dependence syndrome . . . . . 303	5.1	7.2	3.0
Diseases of the nervous system and sense organs . . . . . 320-389	18.0	15.4	20.6
Diseases of the circulatory system . . . . . 390-459	222.0	228.2	216.0
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	154.8	164.9	145.2
Acute myocardial infarction . . . . . 410	28.5	34.1	23.1
Coronary atherosclerosis . . . . . 414.0	38.2	47.4	29.3
Other ischemic heart disease . . . . . 411-413,414.1-414.9	7.3	7.4	7.3
Cardiac dysrhythmias . . . . . 427	27.5	26.8	28.0
Congestive heart failure . . . . . 428.0	33.8	31.4	36.1
Cerebrovascular disease . . . . . 430-438	32.8	30.7	34.8
Diseases of the respiratory system . . . . . 460-519	123.4	119.3	127.2
Acute bronchitis and bronchiolitis . . . . . 466	9.7	10.8	8.7
Pneumonia . . . . . 480-486	45.7	44.0	47.3
Chronic bronchitis . . . . . 491	18.1	16.3	19.8
Asthma . . . . . 493	16.8	13.9	19.7
Diseases of the digestive system . . . . . 520-579	115.6	104.3	126.5
Appendicitis . . . . . 540-543	10.3	11.9	8.8
Noninfectious enteritis and colitis . . . . . 555-558	10.8	9.1	12.4
Diverticula of intestine . . . . . 562	9.1	7.5	10.7
Cholelithiasis . . . . . 574	12.5	8.0	16.8
Diseases of the genitourinary system . . . . . 580-629	63.3	39.8	85.8
Calculus of kidney and ureter . . . . . 592	6.1	7.0	5.3
Complications of pregnancy, childbirth, and the puerperium <sup>2</sup> . . . . . 630-677	18.4	...	36.0
Diseases of the skin and subcutaneous tissue . . . . . 680-709	20.9	21.2	20.6
Cellulitis and abscess . . . . . 681-682	14.7	15.0	14.4
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	60.5	53.1	67.5
Osteoarthritis and allied disorders . . . . . 715	19.8	15.7	23.7
Intervertebral disc disorders . . . . . 722	12.3	12.8	11.8
Congenital anomalies . . . . . 740-759	6.2	6.9	5.5
Certain conditions originating in the perinatal period . . . . . 760-779	5.8	6.6	5.0
Symptoms, signs, and ill-defined conditions . . . . . 780-799	9.9	9.4	10.3
Injury and poisoning . . . . . 800-999	93.9	94.7	93.2
Fractures, all sites <sup>3</sup> . . . . . 800-829	34.7	30.8	38.4
Fracture of neck of femur <sup>3</sup> . . . . . 820	11.0	6.0	15.7
Poisonings . . . . . 960-989	7.4	6.8	8.0
Supplementary classifications . . . . . V01-V83	170.0	28.7	305.5
Females with deliveries . . . . . V27	137.6	...	269.7

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>3</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 16. Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Average length of stay in days		
All conditions . . . . .	4.9	5.3	4.6
Infectious and parasitic diseases . . . . . 001-139	6.4	6.6	6.1
Septicemia . . . . . 038	8.3	8.8	8.0
Neoplasms . . . . . 140-239	6.1	7.0	5.4
Malignant neoplasms . . . . . 140-208,230-234	7.1	7.3	6.9
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	8.7	8.5	8.9
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	7.5	7.7	7.3
Malignant neoplasm of breast . . . . . 174-175,198.81	2.9	*	2.9
Benign neoplasms . . . . . 210-229	3.3	4.6	3.2
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	4.3	4.5	4.1
Diabetes mellitus . . . . . 250	5.0	5.3	4.7
Volume depletion . . . . . 276.5	3.8	3.8	3.7
Diseases of the blood and blood-forming organs . . . . . 280-289	4.4	4.3	4.4
Mental disorders . . . . . 290-319	7.1	7.2	7.1
Psychoses . . . . . 290-299	8.0	8.1	7.9
Alcohol dependence syndrome . . . . . 303	5.8	5.5	6.3
Diseases of the nervous system and sense organs . . . . . 320-389	5.1	5.6	4.7
Diseases of the circulatory system . . . . . 390-459	4.7	4.6	4.8
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4.6	4.4	4.7
Acute myocardial infarction . . . . . 410	5.6	5.5	5.8
Coronary atherosclerosis . . . . . 414.0	3.5	3.5	3.7
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2.6	2.5	2.6
Cardiac dysrhythmias . . . . . 427	3.6	3.4	3.8
Congestive heart failure . . . . . 428.0	5.4	5.5	5.4
Cerebrovascular disease . . . . . 430-438	5.3	5.4	5.2
Diseases of the respiratory system . . . . . 460-519	5.3	5.2	5.4
Acute bronchitis and bronchiolitis . . . . . 466	3.4	3.3	3.6
Pneumonia . . . . . 480-486	5.7	5.6	5.9
Chronic bronchitis . . . . . 491	5.3	5.1	5.4
Asthma . . . . . 493	3.2	2.7	3.5
Diseases of the digestive system . . . . . 520-579	4.9	4.9	4.8
Appendicitis . . . . . 540-543	3.2	3.3	3.0
Noninfectious enteritis and colitis . . . . . 555-558	4.7	4.6	4.7
Diverticula of intestine . . . . . 562	5.8	6.0	5.6
Cholelithiasis . . . . . 574	4.0	4.7	3.8
Diseases of the genitourinary system . . . . . 580-629	3.7	4.1	3.5
Calculus of kidney and ureter . . . . . 592	2.2	2.1	2.4
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	2.7	...	2.7
Diseases of the skin and subcutaneous tissue . . . . . 680-709	5.3	5.0	5.6
Cellulitis and abscess . . . . . 681-682	5.2	5.1	5.3
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	3.9	3.7	4.1
Osteoarthritis and allied disorders . . . . . 715	4.2	4.1	4.2
Intervertebral disc disorders . . . . . 722	2.8	2.4	3.2
Congenital anomalies . . . . . 740-759	5.8	5.8	5.8
Certain conditions originating in the perinatal period . . . . . 760-779	11.1	10.8	11.6
Symptoms, signs, and ill-defined conditions . . . . . 780-799	2.7	2.5	2.8
Injury and poisoning . . . . . 800-999	5.3	5.4	5.2
Fractures, all sites <sup>2</sup> . . . . . 800-829	5.4	5.4	5.4
Fracture of neck of femur <sup>2</sup> . . . . . 820	6.6	7.0	6.4
Poisonings . . . . . 960-989	2.7	2.9	2.6
Supplementary classifications . . . . . V01-V83	3.8	8.5	3.4
Females with deliveries . . . . . V27	2.6	...	2.6

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.



**Table 17. Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions . . . . .	33,727	6,990	7,503	12,994	6,239
Infectious and parasitic diseases . . . . . 001-139	877	183	190	339	166
Septicemia . . . . . 038	341	74	77	126	64
Neoplasms . . . . . 140-239	1,682	358	353	647	324
Malignant neoplasms . . . . . 140-208,230-234	1,208	268	266	447	226
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	159	34	42	55	27
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	160	38	38	58	26
Malignant neoplasm of breast . . . . . 174-175,198.81	85	17	17	32	19
Benign neoplasms . . . . . 210-229	427	79	77	183	88
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	1,619	350	367	621	281
Diabetes mellitus . . . . . 250	577	128	122	228	99
Volume depletion . . . . . 276.5	508	103	128	198	79
Diseases of the blood and blood-forming organs . . . . . 280-289	446	100	103	184	59
Mental disorders . . . . . 290-319	2,464	681	484	1,036	262
Psychoses . . . . . 290-299	1,704	458	336	*	165
Alcohol dependence syndrome . . . . . 303	145	30	*	*	16
Diseases of the nervous system and sense organs . . . . . 320-389	518	115	124	194	85
Diseases of the circulatory system . . . . . 390-459	6,373	1,426	1,500	2,394	1,054
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4,446	1,025	1,059	1,636	726
Acute myocardial infarction . . . . . 410	818	179	181	297	161
Coronary atherosclerosis . . . . . 414.0	1,096	232	281	412	170
Other ischemic heart disease . . . . . 411-413,414.1-414.9	211	51	46	86	28
Cardiac dysrhythmias . . . . . 427	788	200	191	274	123
Congestive heart failure . . . . . 428.0	970	221	235	372	142
Cerebrovascular disease . . . . . 430-438	942	202	215	351	174
Diseases of the respiratory system . . . . . 460-519	3,542	722	820	1,387	613
Acute bronchitis and bronchiolitis . . . . . 466	279	44	60	111	64
Pneumonia . . . . . 480-486	1,312	260	305	518	228
Chronic bronchitis . . . . . 491	520	111	123	226	60
Asthma . . . . . 493	484	110	104	181	89
Diseases of the digestive system . . . . . 520-579	3,320	680	772	1,222	646
Appendicitis . . . . . 540-543	295	57	65	83	91
Noninfectious enteritis and colitis . . . . . 555-558	310	63	76	119	51
Diverticula of intestine . . . . . 562	262	64	68	93	37
Cholelithiasis . . . . . 574	359	69	82	130	77
Diseases of the genitourinary system . . . . . 580-629	1,817	347	419	722	330
Calculus of kidney and ureter . . . . . 592	176	36	48	67	24
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	528	91	88	247	102
Diseases of the skin and subcutaneous tissue . . . . . 680-709	601	137	127	249	88
Cellulitis and abscess . . . . . 681-682	422	109	95	152	67
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	1,736	368	432	600	336
Osteoarthritis and allied disorders . . . . . 715	568	132	150	186	101
Intervertebral disc disorders . . . . . 722	353	64	79	137	74
Congenital anomalies . . . . . 740-759	178	*	*	55	*
Certain conditions originating in the perinatal period . . . . . 760-779	166	27	*	49	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	283	61	56	102	*
Injury and poisoning . . . . . 800-999	2,697	540	626	976	556
Fractures, all sites <sup>2</sup> . . . . . 800-829	995	203	252	332	209
Fracture of neck of femur <sup>2</sup> . . . . . 820	315	70	81	108	57
Poisonings . . . . . 960-989	214	39	53	80	42
Supplementary classifications . . . . . V01-V83	4,880	750	984	1,972	1,175
Females with deliveries . . . . . V27	3,951	582	736	1,660	974

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 18. Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Rate per 10,000 population <sup>1</sup>		
			Midwest	South	West
All conditions . . . . .	1,174.6	1,290.3	1,153.5	1,266.5	955.4
Infectious and parasitic diseases . . . . . 001-139	30.5	33.8	29.1	33.0	25.3
Septicemia . . . . . 038	11.9	13.6	11.9	12.3	9.9
Neoplasms . . . . . 140-239	58.6	66.1	54.2	63.1	49.7
Malignant neoplasms . . . . . 140-208,230-234	42.1	49.5	41.0	43.6	34.6
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	5.5	6.4	6.5	5.4	4.1
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	5.6	7.1	5.8	5.6	3.9
Malignant neoplasm of breast . . . . . 174-175,198.81	3.0	3.1	2.6	3.1	3.0
Benign neoplasms . . . . . 210-229	14.9	14.6	11.8	17.9	13.5
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	56.4	64.6	56.4	60.5	43.1
Diabetes mellitus . . . . . 250	20.1	23.6	18.8	22.3	15.1
Volume depletion . . . . . 276.5	17.7	19.0	19.7	19.3	12.1
Diseases of the blood and blood-forming organs . . . . . 280-289	15.5	18.5	15.9	17.9	9.0
Mental disorders . . . . . 290-319	85.8	125.7	74.5	101.0	40.2
Psychoses . . . . . 290-299	59.4	84.5	51.7	*	25.2
Alcohol dependence syndrome . . . . . 303	5.1	5.5	*	*	2.5
Diseases of the nervous system and sense organs . . . . . 320-389	18.0	21.3	19.0	19.0	13.0
Diseases of the circulatory system . . . . . 390-459	222.0	263.2	230.5	233.3	161.4
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	154.8	189.2	162.8	159.4	111.2
Acute myocardial infarction . . . . . 410	28.5	33.0	27.8	29.0	24.6
Coronary atherosclerosis . . . . . 414.0	38.2	42.9	43.2	40.2	26.0
Other ischemic heart disease . . . . . 411-413,414.1-414.9	7.3	9.5	7.1	8.4	4.3
Cardiac dysrhythmias . . . . . 427	27.5	37.0	29.4	26.7	18.8
Congestive heart failure . . . . . 428.0	33.8	40.8	36.1	36.3	21.8
Cerebrovascular disease . . . . . 430-438	32.8	37.2	33.1	34.2	26.6
Diseases of the respiratory system . . . . . 460-519	123.4	133.2	126.0	135.2	93.9
Acute bronchitis and bronchiolitis . . . . . 466	9.7	8.1	9.2	10.8	9.9
Pneumonia . . . . . 480-486	45.7	48.0	46.9	50.5	34.9
Chronic bronchitis . . . . . 491	18.1	20.5	19.0	22.0	9.1
Asthma . . . . . 493	16.8	20.2	15.9	17.7	13.7
Diseases of the digestive system . . . . . 520-579	115.6	125.5	118.7	119.1	99.0
Appendicitis . . . . . 540-543	10.3	10.5	10.0	8.1	13.9
Noninfectious enteritis and colitis . . . . . 555-558	10.8	11.7	11.7	11.6	7.9
Diverticula of intestine . . . . . 562	9.1	11.9	10.5	9.1	5.6
Cholelithiasis . . . . . 574	12.5	12.7	12.7	12.7	11.8
Diseases of the genitourinary system . . . . . 580-629	63.3	64.1	64.4	70.3	50.5
Calculus of kidney and ureter . . . . . 592	6.1	6.7	7.4	6.5	3.7
Complications of pregnancy, childbirth, and the puerperium <sup>2</sup> . . . . . 630-677	18.4	16.9	13.5	24.1	15.6
Diseases of the skin and subcutaneous tissue . . . . . 680-709	20.9	25.3	19.5	24.3	13.4
Cellulitis and abscess . . . . . 681-682	14.7	20.0	14.6	14.8	10.3
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	60.5	67.9	66.5	58.5	51.4
Osteoarthritis and allied disorders . . . . . 715	19.8	24.3	23.0	18.1	15.5
Intervertebral disc disorders . . . . . 722	12.3	11.8	12.1	13.4	11.3
Congenital anomalies . . . . . 740-759	6.2	*	*	5.3	*
Certain conditions originating in the perinatal period . . . . . 760-779	5.8	4.9	*	4.8	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	9.9	11.2	8.5	9.9	*
Injury and poisoning . . . . . 800-999	93.9	99.7	96.2	95.1	85.1
Fractures, all sites <sup>3</sup> . . . . . 800-829	34.7	37.5	38.7	32.3	32.0
Fracture of neck of femur <sup>3</sup> . . . . . 820	11.0	12.9	12.4	10.5	8.7
Poisonings . . . . . 960-989	7.4	7.1	8.2	7.8	6.4
Supplementary classifications . . . . . V01-V83	170.0	138.4	151.2	192.2	179.9
Females with deliveries . . . . . V27	137.6	107.4	113.1	161.8	149.1

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>3</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 19. Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Average length of stay in days				
All conditions . . . . .	4.9	5.6	4.4	4.9	4.5
Infectious and parasitic diseases . . . . . 001-139	6.4	7.6	5.2	6.5	6.2
Septicemia . . . . . 038	8.3	10.5	6.7	8.3	8.0
Neoplasms . . . . . 140-239	6.1	6.8	5.9	6.0	5.5
Malignant neoplasms . . . . . 140-208,230-234	7.1	7.8	6.6	7.1	6.6
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	8.7	9.9	7.9	9.4	7.2
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	7.5	7.4	6.6	8.1	7.6
Malignant neoplasm of breast . . . . . 174-175,198.81	2.9	3.1	3.4	2.7	2.6
Benign neoplasms . . . . . 210-229	3.3	3.5	3.5	3.4	2.8
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	4.3	5.0	3.9	4.2	4.1
Diabetes mellitus . . . . . 250	5.0	5.7	4.1	4.9	5.6
Volume depletion . . . . . 276.5	3.8	4.8	3.5	3.6	3.4
Diseases of the blood and blood-forming organs . . . . . 280-289	4.4	4.7	3.5	4.7	4.0
Mental disorders . . . . . 290-319	7.1	9.1	5.8	6.4	7.1
Psychoses . . . . . 290-299	8.0	10.4	6.5	7.1	8.9
Alcohol dependence syndrome . . . . . 303	5.8	6.3	6.3	5.0	*
Diseases of the nervous system and sense organs . . . . . 320-389	5.1	6.2	4.4	5.4	4.0
Diseases of the circulatory system . . . . . 390-459	4.7	5.3	4.2	4.8	4.3
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	4.6	5.2	4.1	4.7	4.1
Acute myocardial infarction . . . . . 410	5.6	6.3	5.2	5.6	5.2
Coronary atherosclerosis . . . . . 414.0	3.5	3.9	3.0	3.9	3.1
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2.6	2.8	2.3	2.6	2.5
Cardiac dysrhythmias . . . . . 427	3.6	4.2	3.3	3.6	3.1
Congestive heart failure . . . . . 428.0	5.4	6.5	4.7	5.5	4.6
Cerebrovascular disease . . . . . 430-438	5.3	5.9	4.9	5.4	4.8
Diseases of the respiratory system . . . . . 460-519	5.3	5.8	4.7	5.5	5.1
Acute bronchitis and bronchiolitis . . . . . 466	3.4	3.9	2.8	3.4	3.7
Pneumonia . . . . . 480-486	5.7	6.2	5.1	6.0	5.6
Chronic bronchitis . . . . . 491	5.3	6.0	4.6	5.4	4.9
Asthma . . . . . 493	3.2	3.1	2.9	3.4	3.0
Diseases of the digestive system . . . . . 520-579	4.9	5.5	4.4	5.0	4.5
Appendicitis . . . . . 540-543	3.2	3.6	3.0	3.2	3.0
Noninfectious enteritis and colitis . . . . . 555-558	4.7	5.2	4.5	4.6	4.3
Diverticula of intestine . . . . . 562	5.8	6.3	5.1	6.2	5.0
Cholelithiasis . . . . . 574	4.0	4.3	3.9	4.0	4.0
Diseases of the genitourinary system . . . . . 580-629	3.7	4.4	3.3	3.8	3.3
Calculus of kidney and ureter . . . . . 592	2.2	2.6	1.7	2.3	2.3
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	2.7	2.9	2.5	2.7	2.4
Diseases of the skin and subcutaneous tissue . . . . . 680-709	5.3	6.0	5.2	5.0	5.0
Cellulitis and abscess . . . . . 681-682	5.2	5.5	4.9	5.4	4.4
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	3.9	4.1	3.8	4.0	3.7
Osteoarthritis and allied disorders . . . . . 715	4.2	4.3	4.1	4.2	4.0
Intervertebral disc disorders . . . . . 722	2.8	2.9	2.8	2.6	3.0
Congenital anomalies . . . . . 740-759	5.8	6.5	7.7	4.8	4.9
Certain conditions originating in the perinatal period . . . . . 760-779	11.1	8.6	10.8	14.1	10.0
Symptoms, signs, and ill-defined conditions . . . . . 780-799	2.7	2.6	2.0	2.5	3.5
Injury and poisoning . . . . . 800-999	5.3	5.7	4.6	5.7	4.9
Fractures, all sites <sup>2</sup> . . . . . 800-829	5.4	6.0	4.9	5.7	5.1
Fracture of neck of femur <sup>2</sup> . . . . . 820	6.6	7.4	6.0	6.3	6.7
Poisonings . . . . . 960-989	2.7	2.7	2.5	3.0	2.6
Supplementary classifications . . . . . V01-V83	3.8	4.0	4.1	3.6	3.7
Females with deliveries . . . . . V27	2.6	2.8	2.5	2.6	2.4

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 20. Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	153,529	6,869	38,740	36,844	71,076
Infectious and parasitic diseases . . . . . 001-139	3,895	442	966	940	1,547
Septicemia . . . . . 038	707	33	85	166	423
Neoplasms . . . . . 140-239	4,362	103	676	1,479	2,104
Malignant neoplasms . . . . . 140-208,230-234	3,276	85	312	1,081	1,798
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	258	*	9	82	167
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	452	*	25	144	280
Malignant neoplasm of breast . . . . . 174-175,198.81	165	*	19	71	75
Benign neoplasms . . . . . 210-229	935	13	344	363	215
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	17,362	591	2,279	5,230	9,262
Diabetes mellitus . . . . . 250	5,117	43	572	1,774	2,728
Volume depletion . . . . . 276.5	2,455	344	341	417	1,352
Diseases of the blood and blood-forming organs . . . . . 280-289	4,603	186	940	1,056	2,422
Mental disorders . . . . . 290-319	11,398	387	4,991	3,207	2,814
Psychoses . . . . . 290-299	3,949	*	1,451	852	1,515
Alcohol dependence syndrome . . . . . 303	754	*	364	307	82
Diseases of the nervous system and sense organs . . . . . 320-389	3,762	409	654	829	1,870
Diseases of the circulatory system . . . . . 390-459	32,356	137	1,871	8,257	22,090
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	18,964	75	865	4,390	13,633
Acute myocardial infarction . . . . . 410	982	*	64	292	626
Coronary atherosclerosis . . . . . 414.0	4,434	*	138	1,242	3,052
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2,269	*	105	727	1,436
Cardiac dysrhythmias . . . . . 427	4,119	27	158	675	3,259
Congestive heart failure . . . . . 428.0	3,381	*	92	641	2,635
Cerebrovascular disease . . . . . 430-438	1,916	*	69	425	1,411
Diseases of the respiratory system . . . . . 460-519	11,181	1,289	1,374	2,505	6,014
Acute bronchitis and bronchiolitis . . . . . 466	417	252	29	52	84
Pneumonia . . . . . 480-486	2,181	271	221	386	1,303
Chronic bronchitis . . . . . 491	1,089	*	41	293	753
Asthma . . . . . 493	1,624	343	449	428	403
Diseases of the digestive system . . . . . 520-579	10,232	510	2,160	2,999	4,564
Appendicitis . . . . . 540-543	335	72	176	58	29
Noninfectious enteritis and colitis . . . . . 555-558	660	128	169	139	224
Diverticula of intestine . . . . . 562	576	*	49	143	384
Cholelithiasis . . . . . 574	636	*	179	186	269
Diseases of the genitourinary system . . . . . 580-629	8,123	180	2,141	2,021	3,781
Calculus of kidney and ureter . . . . . 592	288	*	128	88	68
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	9,450	32	9,402	*16	...
Diseases of the skin and subcutaneous tissue . . . . . 680-709	2,116	176	412	561	968
Cellulitis and abscess . . . . . 681-682	938	59	205	279	395
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	6,230	93	908	1,789	3,440
Osteoarthritis and allied disorders . . . . . 715	1,687	*	56	429	1,202
Intervertebral disc disorders . . . . . 722	563	*	187	234	141
Congenital anomalies . . . . . 740-759	714	405	154	87	68
Certain conditions originating in the perinatal period . . . . . 760-779	472	463	*3	*	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	8,221	700	1,798	2,244	3,479
Injury and poisoning . . . . . 800-999	6,311	459	1,961	1,503	2,388
Fractures, all sites <sup>2</sup> . . . . . 800-829	1,609	97	451	299	762
Fracture of neck of femur <sup>2</sup> . . . . . 820	360	*	9	30	318
Poisonings . . . . . 960-989	377	25	222	87	43
Supplementary classifications . . . . . V01-V83	12,740	308	6,049	2,123	4,261
Females with deliveries . . . . . V27	3,951	12	3,934	*5	...

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 21. Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
All conditions . . . . .	153,529	62,073	91,456
Infectious and parasitic diseases . . . . . 001-139	3,895	1,739	2,155
Septicemia . . . . . 038	707	329	377
Neoplasms . . . . . 140-239	4,362	1,834	2,528
Malignant neoplasms . . . . . 140-208,230-234	3,276	1,618	1,658
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	258	126	132
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	452	235	217
Malignant neoplasm of breast . . . . . 174-175,198.81	165	*	164
Benign neoplasms . . . . . 210-229	935	143	792
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	17,362	7,252	10,110
Diabetes mellitus . . . . . 250	5,117	2,360	2,757
Volume depletion . . . . . 276.5	2,455	1,004	1,451
Diseases of the blood and blood-forming organs . . . . . 280-289	4,603	1,846	2,757
Mental disorders . . . . . 290-319	11,398	5,428	5,970
Psychoses . . . . . 290-299	3,949	1,768	2,182
Alcohol dependence syndrome . . . . . 303	754	543	211
Diseases of the nervous system and sense organs . . . . . 320-389	3,762	1,630	2,132
Diseases of the circulatory system . . . . . 390-459	32,356	15,393	16,962
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	18,964	9,454	9,510
Acute myocardial infarction . . . . . 410	982	563	419
Coronary atherosclerosis . . . . . 414.0	4,434	2,485	1,949
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2,269	1,283	987
Cardiac dysrhythmias . . . . . 427	4,119	1,990	2,129
Congestive heart failure . . . . . 428.0	3,381	1,476	1,905
Cerebrovascular disease . . . . . 430-438	1,916	886	1,030
Diseases of the respiratory system . . . . . 460-519	11,181	5,236	5,945
Acute bronchitis and bronchiolitis . . . . . 466	417	215	203
Pneumonia . . . . . 480-486	2,181	1,044	1,137
Chronic bronchitis . . . . . 491	1,089	491	598
Asthma . . . . . 493	1,624	568	1,057
Diseases of the digestive system . . . . . 520-579	10,232	4,353	5,878
Appendicitis . . . . . 540-543	335	179	156
Noninfectious enteritis and colitis . . . . . 555-558	660	269	390
Diverticula of intestine . . . . . 562	576	221	355
Cholelithiasis . . . . . 574	636	215	421
Diseases of the genitourinary system . . . . . 580-629	8,123	2,827	5,296
Calculus of kidney and ureter . . . . . 592	288	143	145
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	9,450	...	9,450
Diseases of the skin and subcutaneous tissue . . . . . 680-709	2,116	1,022	1,095
Cellulitis and abscess . . . . . 681-682	938	463	475
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	6,230	2,173	4,057
Osteoarthritis and allied disorders . . . . . 715	1,687	566	1,121
Intervertebral disc disorders . . . . . 722	563	275	288
Congenital anomalies . . . . . 740-759	714	368	345
Certain conditions originating in the perinatal period . . . . . 760-779	472	270	202
Symptoms, signs, and ill-defined conditions . . . . . 780-799	8,221	3,649	4,572
Injury and poisoning . . . . . 800-999	6,311	3,226	3,085
Fractures, all sites <sup>2</sup> . . . . . 800-829	1,609	764	845
Fracture of neck of femur <sup>2</sup> . . . . . 820	360	102	258
Poisonings . . . . . 960-989	377	160	217
Supplementary classifications . . . . . V01-V83	12,740	3,824	8,916
Females with deliveries . . . . . V27	3,951	...	3,951

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.



**Table 22. Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	North-east	Midwest	South	West
	Number in thousands				
All conditions . . . . .	153,529	32,417	36,636	56,332	28,144
Infectious and parasitic diseases . . . . . 001-139	3,895	827	829	1,494	746
Septicemia . . . . . 038	707	159	149	262	136
Neoplasms . . . . . 140-239	4,362	1,003	932	1,603	824
Malignant neoplasms . . . . . 140-208,230-234	3,276	779	714	1,185	599
Malignant neoplasm of large intestine and rectum . . . . . 153-154,197.5	258	54	65	93	45
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	452	112	99	163	77
Malignant neoplasm of breast . . . . . 174-175,198.81	165	34	39	61	31
Benign neoplasms . . . . . 210-229	935	188	183	365	199
Endocrine, nutritional and metabolic diseases, and immunity disorders . . . . . 240-279	17,362	3,701	4,256	6,420	2,985
Diabetes mellitus . . . . . 250	5,117	1,130	1,166	2,004	816
Volume depletion . . . . . 276.5	2,455	478	644	916	417
Diseases of the blood and blood-forming organs . . . . . 280-289	4,603	976	1,064	1,742	821
Mental disorders . . . . . 290-319	11,398	2,791	2,719	4,095	1,793
Psychoses . . . . . 290-299	3,949	971	899	1,572	506
Alcohol dependence syndrome . . . . . 303	754	190	*	233	131
Diseases of the nervous system and sense organs . . . . . 320-389	3,762	813	914	1,289	746
Diseases of the circulatory system . . . . . 390-459	32,356	7,403	7,991	11,646	5,316
Heart disease . . . . . 391-392.0,393-398,402,404,410-416,420-429	18,964	4,456	4,769	6,663	3,075
Acute myocardial infarction . . . . . 410	982	220	222	349	191
Coronary atherosclerosis . . . . . 414.0	4,434	1,048	1,145	1,576	666
Other ischemic heart disease . . . . . 411-413,414.1-414.9	2,269	537	581	799	353
Cardiac dysrhythmias . . . . . 427	4,119	983	1,023	1,395	718
Congestive heart failure . . . . . 428.0	3,381	748	909	1,239	485
Cerebrovascular disease . . . . . 430-438	1,916	411	473	687	345
Diseases of the respiratory system . . . . . 460-519	11,181	2,359	2,656	4,159	2,007
Acute bronchitis and bronchiolitis . . . . . 466	417	70	95	165	88
Pneumonia . . . . . 480-486	2,181	433	505	850	393
Chronic bronchitis . . . . . 491	1,089	225	270	450	144
Asthma . . . . . 493	1,624	397	385	539	303
Diseases of the digestive system . . . . . 520-579	10,232	2,145	2,421	3,808	1,859
Appendicitis . . . . . 540-543	335	62	74	100	99
Noninfectious enteritis and colitis . . . . . 555-558	660	135	164	248	113
Diverticula of intestine . . . . . 562	576	137	143	208	87
Cholelithiasis . . . . . 574	636	129	148	229	129
Diseases of the genitourinary system . . . . . 580-629	8,123	1,633	1,906	3,122	1,462
Calculus of kidney and ureter . . . . . 592	288	65	72	106	45
Complications of pregnancy, childbirth, and the puerperium <sup>1</sup> . . . . . 630-677	9,450	1,496	1,822	3,873	2,260
Diseases of the skin and subcutaneous tissue . . . . . 680-709	2,116	482	500	776	358
Cellulitis and abscess . . . . . 681-682	938	221	219	336	162
Diseases of the musculoskeletal system and connective tissue . . . . . 710-739	6,230	1,289	1,654	2,125	1,162
Osteoarthritis and allied disorders . . . . . 715	1,687	364	467	577	280
Intervertebral disc disorders . . . . . 722	563	100	130	216	116
Congenital anomalies . . . . . 740-759	714	182	141	186	*
Certain conditions originating in the perinatal period . . . . . 760-779	472	72	*	149	*
Symptoms, signs, and ill-defined conditions . . . . . 780-799	8,221	1,673	2,231	2,856	1,461
Injury and poisoning . . . . . 800-999	6,311	1,232	1,503	2,255	1,321
Fractures, all sites <sup>2</sup> . . . . . 800-829	1,609	307	390	564	348
Fracture of neck of femur <sup>2</sup> . . . . . 820	360	76	95	124	65
Poisonings . . . . . 960-989	377	64	106	136	70
Supplementary classifications . . . . . V01-V83	12,740	2,342	3,000	4,733	2,665
Females with deliveries . . . . . V27	3,951	582	736	1,660	974

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 23. Number and rate of discharges from short-stay hospitals with a human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years 1990–2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Data are for discharges with at least one of the following *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) codes: 042-044, 279.19, 795.8 (prior to 1995); 042, V08 (beginning in 1995)]

Characteristic	1990	1995	1998	1999	2000	2001	2002
	Number in thousands						
All HIV discharges <sup>1</sup>	146	249	189	180	173	185	189
Sex							
Male	114	183	132	121	115	115	124
Female	32	66	57	59	58	70	65
Age							
Under 25 years	14	17	9	8	7	10	10
25–29 years	20	26	15	13	11	8	11
30–34 years	38	48	30	41	28	28	23
35–39 years	36	59	42	34	35	41	35
40–44 years	17	49	43	34	34	43	42
45 years and over	21	51	51	49	58	56	68
Region							
Northeast	66	95	69	70	68	61	57
Midwest	20	27	23	20	20	18	20
South	37	87	71	67	64	86	81
West	24	41	26	23	21	21	31
	Rate per 100,000 population <sup>2</sup>						
All HIV discharges <sup>1</sup>	58.1	93.9	69.1	65.3	61.6	65.2	65.8
Sex							
Male	93.4	141.6	98.8	89.5	83.9	82.8	88.5
Female	24.7	48.4	40.7	42.1	40.1	48.2	44.1
Age							
Under 25 years	14.8	17.6	9.1	8.1	7.5	10.1	9.5
25–29 years	95.1	132.2	77.0	70.5	56.6	41.3	57.5
30–34 years	172.8	216.2	145.2	208.3	134.9	137.6	113.0
35–39 years	179.0	262.7	185.4	149.1	154.4	182.9	159.7
40–44 years	94.8	241.4	194.4	152.0	152.6	187.4	185.2
45 years and over	27.4	60.2	55.6	52.9	59.8	55.8	66.6
Region							
Northeast	128.8	183.3	133.4	135.0	126.6	112.3	107.9
Midwest	33.6	43.5	36.2	30.7	30.8	27.9	31.7
South	42.1	93.1	73.1	68.4	64.2	84.5	81.7
West	44.1	69.5	42.0	37.0	33.4	32.6	49.9

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>HIV is human immunodeficiency virus.

<sup>2</sup>Rates are based on the U.S. Census Bureau estimates of the civilian population. Populations for 1990–99 were adjusted for the net underenumeration in the 1990 census. Populations for 2000–02 were estimated from the 2000 census. Rates for 2000–01 differ from rates shown in a previous report because they were calculated using revised population estimates.

**Table 24. Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2000–2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. *International Classification of Diseases, 9th Revision, Clinical Modification* code numbers for Barell matrix categories are shown in table III]

Nature of injury	All body regions	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
Number in thousands <sup>1</sup>									
All injuries . . . . .	1,802	159	117	10	83	231	228	688	34
Fracture . . . . .	992	42	33	7	68	95	149	597	*
Dislocation . . . . .	20	...	*	...	4	*	7	9	–
Sprains and strains . . . . .	57	...	–	...	11	6	19	20	*2
Internal . . . . .	203	117	...	3	...	82	...	...	*
Open wound . . . . .	93	...	36	...	...	13	23	21	*
Amputation . . . . .	9	...	...	...	...	...	7	*1	...
Blood vessels . . . . .	7	...	*	...	...	*2	3	*2	*
Contusions and superficial injuries . . . . .	60	...	12	...	...	17	4	19	7
Crush . . . . .	2	...	*	...	...	*	*	*	–
Burns . . . . .	66	...	9	...	...	*	12	15	*
Nerves and spinal cord . . . . .	4	*	*	...	...	*	2	...	*
Unspecified . . . . .	36	...	25	...	...	5	*	*3	*3
System-wide . . . . .	253	...	...	...	...	...	...	...	...
Rate per 10,000 population <sup>2</sup>									
All injuries . . . . .	63.4	5.6	4.1	0.4	2.9	8.1	8.0	24.2	1.2
Fracture . . . . .	34.9	1.5	1.2	0.2	2.4	3.4	5.3	21.0	*
Dislocation . . . . .	0.7	...	*	...	0.1	*	0.2	0.3	–
Sprains and strains . . . . .	2.0	...	–	...	0.4	0.2	0.7	0.7	*0.1
Internal . . . . .	7.1	4.1	...	0.1	...	2.9	...	...	*
Open wound . . . . .	3.3	...	1.2	...	...	0.5	0.8	0.7	*
Amputation . . . . .	0.3	...	...	...	...	...	0.3	*0.0	...
Blood vessels . . . . .	0.3	...	*	...	...	*0.1	0.1	*0.1	*
Contusions and superficial injuries . . . . .	2.1	...	0.4	...	...	0.6	0.1	0.7	0.3
Crush . . . . .	0.1	...	*	...	...	*	*	*	–
Burns . . . . .	2.3	...	0.3	...	...	*	0.4	0.5	*
Nerves and spinal cord . . . . .	0.1	*	*	...	...	*	0.1	...	*
Unspecified . . . . .	1.3	...	0.9	...	...	0.2	*	*0.1	*0.1
System-wide . . . . .	8.9	...	...	...	...	...	...	...	...

... Category not applicable.

– Quantity zero.

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>The annual average number of discharges was computed by adding the number of discharges with a first-listed injury diagnosis during the 3-year period and dividing by 3.<sup>2</sup>Rates were calculated by dividing the annual average number of discharges by the 2001 population. The population is the U.S. Bureau of the Census estimate of the civilian population based on the 2000 census.**Table 25. Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)*]

First-listed diagnosis and ICD–9–CM code	All ages	Under 65 years	65 years and over	All ages	Under 65 years	65 years and over
	Number of deaths in thousands			Rate per 100 discharges		
All conditions <sup>1</sup> . . . . .	811	209	602	2.4	1.0	4.7
Heart disease . . . . . 391–392.0,393–398,402,404,410–416,420–429	157	29	128	3.5	1.8	4.5
Acute myocardial infarction . . . . . 410	67	10	57	8.2	3.1	11.4
Cardiac dysrhythmias . . . . . 427	23	*8	15	2.9	*3.5	2.7
Congestive heart failure . . . . . 428.0	35	*4	31	3.6	*1.4	4.4
Malignant neoplasms . . . . . 140–208,230–234	81	28	53	6.7	4.9	8.3
Malignant neoplasm of trachea, bronchus, and lung . . . . . 162,176.4,197.0,197.3	22	7	15	13.9	11.9	15.1
Pneumonia . . . . . 480–486	71	9	62	5.4	1.7	7.9
Septicemia . . . . . 038	64	15	49	18.7	12.4	22.3
Cerebrovascular disease . . . . . 430–438	61	14	46	6.5	5.3	6.9
Fractures, all sites <sup>2</sup> . . . . . 800–829	22	*7	15	2.2	*1.5	2.8
Volume depletion . . . . . 276.5	15	*	14	3.0	*	5.7
Chronic bronchitis . . . . . 491	13	*	11	2.5	*	3.1

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Includes data for deaths not shown in table.<sup>2</sup>Excludes fractures coded as 733.1, pathologic fracture.

**Table 26. Number of discharges from short-stay hospitals with and without procedures and percent with procedures, by selected characteristics: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants]

Characteristic	All discharges	Discharges without procedures	Discharges with procedure			
			Total	With surgical procedures	Total	With surgical procedures
			Number in thousands		Percent	
All patients . . . . .	33,727	13,249	20,477	15,235	60.7	45.2
Age						
Under 15 years . . . . .	2,540	1,418	1,122	634	44.2	25.0
15–44 years . . . . .	10,736	3,215	7,521	5,947	70.1	55.4
45–64 years . . . . .	7,723	2,940	4,783	3,678	61.9	47.6
65 years and over . . . . .	12,727	5,677	7,051	4,976	55.4	39.1
65–74 years . . . . .	4,642	1,788	2,854	2,130	61.5	45.9
75–84 years . . . . .	5,271	2,371	2,900	2,042	55.0	38.7
85 years and over . . . . .	2,814	1,517	1,297	804	46.1	28.6
Sex						
Male . . . . .	13,389	5,668	7,720	5,491	57.7	41.0
Female . . . . .	20,338	7,581	12,757	9,745	62.7	47.9
Region						
Northeast . . . . .	6,990	2,629	4,361	2,961	62.4	42.4
Midwest . . . . .	7,503	3,213	4,290	3,208	57.2	42.8
South . . . . .	12,994	5,401	7,593	5,896	58.4	45.4
West . . . . .	6,239	2,007	4,233	3,170	67.8	50.8

NOTE: All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.

**Table 27. Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals, by selected procedure categories: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Procedure category and ICD-9-CM code	Number in thousands	Rate per 10,000 population <sup>1</sup>
Surgical procedures <sup>2</sup>	25,255	890.5
Cardiac catheterization . . . . . 37.21-37.23	1,328	46.8
Repair of current obstetric laceration <sup>3</sup> . . . . . 75.5-75.6	1,234	43.5
Removal of coronary artery obstruction and insertion of stent(s) <sup>4</sup> . . . . . 36.0	1,204	42.4
Cesarean section <sup>3</sup> . . . . . 74.0-74.2,74.4,74.99	1,059	37.3
Artificial rupture of membranes <sup>3</sup> . . . . . 73.0	901	31.8
Episiotomy with or without forceps or vacuum extraction <sup>3</sup> . . . . . 72.1,72.21,72.31,72.71,73.6	780	27.5
Hysterectomy . . . . . 68.3-68.7,68.9	669	23.6
Reduction of fracture . . . . . 76.7,79.0-79.3	606	21.4
Open reduction of fracture with internal fixation . . . . . 79.3	414	14.6
Endoscopy of small intestine with biopsy . . . . . 45.14,45.16	585	20.6
Oophorectomy and salpingo-oophorectomy . . . . . 65.3-65.6	533	18.8
Coronary artery bypass graft <sup>4</sup> . . . . . 36.1	515	18.2
Cholecystectomy . . . . . 51.2	436	15.4
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	420	14.8
Total knee replacement . . . . . 81.54	381	13.4
Debridement of wound, infection, or burn . . . . . 86.22,86.28	361	12.7
Lysis of peritoneal adhesions . . . . . 54.5	342	12.0
Bilateral destruction or occlusion of fallopian tubes . . . . . 66.2-66.3	329	11.6
Appendectomy, excluding incidental . . . . . 47.0	329	11.6
Excision or destruction of intervertebral disc . . . . . 80.5	323	11.4
Partial excision of large intestine . . . . . 45.7	263	9.3
Partial excision of bone . . . . . 76.2-76.3,77.6-77.8	218	7.7
Prostatectomy . . . . . 60.2-60.6	195	6.9
Nonsurgical procedures <sup>5</sup>	17,278	601.8
Arteriography and angiocardiology using contrast material . . . . . 88.4-88.5	2,058	71.7
Respiratory therapy . . . . . 93.9,96.7	1,070	37.3
Diagnostic ultrasound . . . . . 88.7	773	26.9
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	703	24.5
Hemodialysis . . . . . 39.95	552	19.2
Insertion of endotracheal tube . . . . . 96.04	477	16.6
Endoscopy of small intestine without biopsy . . . . . 45.11-45.13	447	15.6
Endoscopy of large intestine without biopsy . . . . . 45.21-45.24	364	12.7
Spinal tap . . . . . 03.31	316	11.0
Injection or infusion of cancer chemotherapeutic substance . . . . . 99.25	217	7.6

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.<sup>3</sup>See table 38 for rate per 100 deliveries.<sup>4</sup>See table 29 for comparison of number of procedures with number of discharges with procedures.<sup>5</sup>Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.



**Table 28. Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
Surgical procedures <sup>1</sup>	8,423	3,722	3,465	1,237
Cardiac catheterization . . . . . 37.21-37.23	659	346	262	50
Removal of coronary artery obstruction and insertion of stent(s) <sup>2</sup> . . . . . 36.0	608	329	239	41
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	358	96	159	103
Endoscopy of small intestine with biopsy . . . . . 45.14,45.16	320	111	142	67
Coronary artery bypass graft <sup>2</sup> . . . . . 36.1	279	153	117	*9
Reduction of fracture . . . . . 76.7,79.0-79.3	247	65	98	84
Open reduction of fracture with internal fixation . . . . . 79.3	185	50	71	64
Total knee replacement . . . . . 81.54	241	126	102	12
Cholecystectomy . . . . . 51.2	151	76	53	22
Partial excision of large intestine . . . . . 45.7	145	55	63	27
Debridement of wound, infection, or burn . . . . . 86.22,86.28	140	57	44	39
Prostatectomy . . . . . 60.2-60.6	128	66	49	13
Total hip replacement . . . . . 81.51	114	51	50	13
Nonsurgical procedures <sup>3</sup>	6,863	2,721	2,834	1,307
Arteriography and angiocardiography using contrast material . . . . . 88.4-88.5	1,028	516	424	88
Respiratory therapy . . . . . 93.9,96.7	476	173	207	96
Diagnostic ultrasound . . . . . 88.7	404	151	168	85
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	336	119	140	77
Endoscopy of small intestine without biopsy . . . . . 45.11-45.13	257	90	106	62
Hemodialysis . . . . . 39.95	251	122	101	28
Insertion of endotracheal tube . . . . . 96.04	241	86	104	50
Endoscopy of large intestine without biopsy . . . . . 45.21-45.24	238	79	111	48
		Rate per 10,000 population <sup>4</sup>		
Surgical procedures <sup>1</sup>	2,366.0	2,036.8	2,720.6	2,692.8
Cardiac catheterization . . . . . 37.21-37.23	185.0	189.3	206.1	109.3
Removal of coronary artery obstruction and insertion of stent(s) <sup>2</sup> . . . . . 36.0	170.9	180.0	187.3	88.9
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	100.5	52.5	124.9	223.9
Endoscopy of small intestine with biopsy . . . . . 45.14,45.16	89.8	61.0	111.3	145.0
Coronary artery bypass graft <sup>2</sup> . . . . . 36.1	78.4	83.8	91.6	*19.8
Reduction of fracture . . . . . 76.7,79.0-79.3	69.5	35.8	76.9	182.9
Open reduction of fracture with internal fixation . . . . . 79.3	51.9	27.5	55.6	139.2
Total knee replacement . . . . . 81.54	67.6	69.1	80.2	27.0
Cholecystectomy . . . . . 51.2	42.3	41.6	41.6	47.0
Partial excision of large intestine . . . . . 45.7	40.7	30.2	49.6	58.0
Debridement of wound, infection, or burn . . . . . 86.22,86.28	39.4	31.1	34.6	85.5
Prostatectomy . . . . . 60.2-60.6	35.9	36.0	38.8	27.6
Total hip replacement . . . . . 81.51	32.1	28.0	39.5	28.1
Nonsurgical procedures <sup>3</sup>	1,927.7	1,489.3	2,225.7	2,846.2
Arteriography and angiocardiography using contrast material . . . . . 88.4-88.5	288.7	282.5	332.7	191.7
Respiratory therapy . . . . . 93.9,96.7	133.7	94.6	162.5	209.6
Diagnostic ultrasound . . . . . 88.7	113.4	82.7	131.7	184.6
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	94.3	64.9	110.0	167.5
Endoscopy of small intestine without biopsy . . . . . 45.11-45.13	72.3	49.0	83.4	134.1
Hemodialysis . . . . . 39.95	70.5	66.8	79.0	61.4
Insertion of endotracheal tube . . . . . 96.04	67.6	47.2	81.6	109.8
Endoscopy of large intestine without biopsy . . . . . 45.21-45.24	66.8	43.2	86.9	105.1

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.<sup>2</sup>See table 29 for a comparison of number of procedures with number of discharges with procedures.<sup>3</sup>Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.<sup>4</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.

**Table 29. Number and rate of discharges from short-stay hospitals with a coronary artery procedure and number and rate of procedures, by sex, age, and geographic region: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Sex, age, and region	Removal of coronary artery obstruction and insertion of stent(s) <sup>1</sup>		Coronary artery bypass graft <sup>2</sup>	
	Number in thousands	Rate per 10,000 population <sup>3</sup>	Number in thousands	Rate per 10,000 population <sup>3</sup>
Discharges with procedures				
Total .....	653	22.7	306	10.6
Sex				
Male .....	430	30.6	218	15.5
Female .....	223	15.2	87	6.0
Age				
Under 15 years .....	*	*	*	*
15-44 years .....	42	3.4	12	1.0
45-64 years .....	283	42.4	128	19.2
65 years and over .....	326	91.7	166	46.7
65-74 years .....	177	96.7	90	49.3
75-84 years .....	127	99.9	69	54.4
85 years and over .....	22	48.7	*7	*14.7
Region				
Northeast .....	113	20.9	60	11.0
Midwest .....	176	27.1	68	10.4
South .....	230	22.4	125	12.2
West .....	134	20.5	54	8.2
Procedures				
Total .....	1,204	41.9	515	17.9
Sex				
Male .....	802	57.0	373	26.6
Female .....	402	27.4	142	9.7
Age				
Under 15 years .....	*	*	*	*
15-44 years .....	76	6.1	19	1.5
45-64 years .....	517	77.6	217	32.6
65 years and over .....	608	170.9	279	78.4
65-74 years .....	329	180.0	153	83.8
75-84 years .....	239	187.3	117	91.6
85 years and over .....	41	88.9	*9	*19.8
Region				
Northeast .....	211	38.9	104	19.3
Midwest .....	323	49.7	117	17.9
South .....	416	40.5	204	19.9
West .....	254	38.8	90	13.7

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>ICD-9-CM code 36.0.<sup>2</sup>ICD-9-CM code 36.1.<sup>3</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.

NOTE: The number of procedures exceeds the number of discharges because these procedures are often assigned more than one code for a single operation. See "Methods" section.

**Table 30. Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over	
						Number in thousands
All procedures	42,533	2,084	14,419	10,743	15,286	
Operations on the nervous system	01-05	1,101	215	337	256	293
Spinal tap	03.31	316	142	93	42	39
Operations on the endocrine system	06-07	102	*	30	40	31
Operations on the eye	08-16	86	12	27	21	25
Operations on the ear	18-20	46	19	*	*7	*
Operations on the nose, mouth, and pharynx	21-29	269	66	93	59	52
Operations on the respiratory system	30-34	1,022	57	179	304	482
Bronchoscopy with or without biopsy	33.21-33.24,33.27	251	15	39	69	127
Operations on the cardiovascular system	35-39	6,813	210	681	2,384	3,538
Removal of coronary artery obstruction and insertion of stent(s) <sup>1</sup>	36.0	1,204	*	76	517	608
Coronary artery bypass graft <sup>1</sup>	36.1	515	*	19	217	279
Cardiac catheterization	37.21-37.23	1,328	*	107	555	659
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7-37.8	420	*	*	47	358
Hemodialysis	39.95	552	*	85	210	251
Operations on the hemic and lymphatic system	40-41	354	22	55	130	147
Operations on the digestive system	42-54	5,597	222	1,307	1,607	2,461
Endoscopy of small intestine with or without biopsy	45.11-45.14,45.16	1,032	16	160	279	577
Endoscopy of large intestine with or without biopsy	45.21-45.25	578	*	76	137	358
Partial excision of large intestine	45.7	263	*	35	79	145
Appendectomy, excluding incidental	47.0	329	70	176	57	25
Cholecystectomy	51.2	436	*	142	140	151
Lysis of peritoneal adhesions	54.5	342	*3	143	103	93
Operations on the urinary system	55-59	955	38	232	299	386
Cystoscopy with or without biopsy	57.31-57.33	173	*	40	52	77
Operations on the male genital organs	60-64	262	18	12	79	152
Prostatectomy	60.2-60.6	195	*	*	66	128
Operations on the female genital organs	65-71	2,161	5	1,256	668	231
Oophorectomy and salpingo-oophorectomy	65.3-65.6	533	*	224	249	60
Bilateral destruction or occlusion of fallopian tubes	66.2-66.3	329	*	329	*	*
Hysterectomy	68.3-68.7,68.9	669	*	344	262	63
Obstetrical procedures	72-75	6,646	21	6,617	*8	...
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	780	*	776	*	...
Artificial rupture of membranes	73.0	901	*	899	*	...
Cesarean section	74.0-74.2,74.4,74.99	1,059	*	1,055	*	...
Repair of current obstetric laceration	75.5-75.6	1,234	*	1,228	*	...
Operations on the musculoskeletal system	76-84	3,442	173	811	1,087	1,371
Partial excision of bone	76.2-76.3,77.6-77.8	218	13	67	89	49
Reduction of fracture	76.7,79.0-79.3	606	50	186	122	247
Open reduction of fracture with internal fixation	79.3	414	15	129	86	185
Excision or destruction of intervertebral disc	80.5	323	*	120	147	55
Total hip replacement	81.51	193	*	11	67	114
Total knee replacement	81.54	381	*	10	131	241
Operations on the integumentary system	85-86	1,348	*	388	415	429
Debridement of wound, infection, or burn	86.22,86.28	361	*	88	114	140
Miscellaneous diagnostic and therapeutic procedures	87-99	12,332	890	2,382	3,381	5,679
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	703	42	157	169	336
Arteriography and angiocardiology using contrast material	88.4-88.5	2,058	*	194	823	1,028
Diagnostic ultrasound	88.7	773	*	133	192	404
Respiratory therapy	93.9,96.7	1,070	207	144	243	476
Insertion of endotracheal tube	96.04	477	48	62	126	241
Injection or infusion of cancer chemotherapeutic substance	99.25	217	34	51	72	61

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>See table 29 for comparison of number of procedures with number of discharges with procedures.

**Table 31. Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over	
						Rate per 10,000 population <sup>1</sup>
All procedures	1,481.4	343.7	1,160.7	1,612.0	4,293.7	
Operations on the nervous system	01-05	38.3	35.5	27.2	38.4	82.3
Spinal tap	03.31	11.0	23.4	7.5	6.4	10.9
Operations on the endocrine system	06-07	3.6	*	2.4	6.0	8.6
Operations on the eye	08-16	3.0	2.0	2.2	3.2	7.2
Operations on the ear	18-20	1.6	3.2	*	*1.0	*
Operations on the nose, mouth, and pharynx	21-29	9.4	10.9	7.4	8.8	14.6
Operations on the respiratory system	30-34	35.6	9.4	14.4	45.6	135.4
Bronchoscopy with or without biopsy	33.21-33.24,33.27	8.7	2.5	3.2	10.3	35.8
Operations on the cardiovascular system	35-39	237.3	34.7	54.8	357.6	993.7
Removal of coronary artery obstruction and insertion of stent(s) <sup>2</sup>	36.0	41.9	*	6.1	77.6	170.9
Coronary artery bypass graft <sup>2</sup>	36.1	17.9	*	1.5	32.6	78.4
Cardiac catheterization	37.21-37.23	46.2	*	8.6	83.3	185.0
Insertion, replacement, removal, and revision of pacemaker leads or device	37.7-37.8	14.6	*	*	7.0	100.5
Hemodialysis	39.95	19.2	*	6.8	31.5	70.5
Operations on the hemic and lymphatic system	40-41	12.3	3.6	4.5	19.5	41.2
Operations on the digestive system	42-54	194.9	36.6	105.2	241.1	691.3
Endoscopy of small intestine with or without biopsy	45.11-45.14,45.16	35.9	2.6	12.9	41.8	162.1
Endoscopy of large intestine with or without biopsy	45.21-45.25	20.1	*	6.1	20.6	100.6
Partial excision of large intestine	45.7	9.2	*	2.8	11.9	40.7
Appendectomy, excluding incidental	47.0	11.5	11.5	14.2	8.6	7.2
Cholecystectomy	51.2	15.2	*	11.4	21.1	42.3
Lysis of peritoneal adhesions	54.5	11.9	*0.5	11.5	15.4	26.0
Operations on the urinary system	55-59	33.2	6.3	18.7	44.8	108.4
Cystoscopy with or without biopsy	57.31-57.33	6.0	*	3.2	7.7	21.6
Operations on the male genital organs	60-64	9.1	2.9	1.0	11.9	42.7
Prostatectomy	60.2-60.6	6.8	*	*	9.9	35.9
Operations on the female genital organs	65-71	75.3	0.8	101.1	100.3	65.0
Oophorectomy and salpingo-oophorectomy	65.3-65.6	18.6	*	18.0	37.4	16.8
Bilateral destruction or occlusion of fallopian tubes	66.2-66.3	11.5	*	26.5	*	*
Hysterectomy	68.3-68.7,68.9	23.3	*	27.7	39.4	17.7
Obstetrical procedures	72-75	231.5	3.5	532.6	*1.2	...
Episiotomy with or without forceps or vacuum extraction	72.1,72.21,72.31,72.71,73.6	27.2	*	62.5	*	...
Artificial rupture of membranes	73.0	31.4	*	72.4	*	...
Cesarean section	74.0-74.2,74.4,74.99	36.9	*	84.9	*	...
Repair of current obstetric laceration	75.5-75.6	43.0	*	98.8	*	...
Operations on the musculoskeletal system	76-84	119.9	28.6	65.3	163.0	385.1
Partial excision of bone	76.2-76.3,77.6-77.8	7.6	2.2	5.4	13.4	13.7
Reduction of fracture	76.7,79.0-79.3	21.1	8.3	15.0	18.3	69.5
Open reduction of fracture with internal fixation	79.3	14.4	2.5	10.4	12.8	51.9
Excision or destruction of intervertebral disc	80.5	11.3	*	9.7	22.0	15.4
Total hip replacement	81.51	6.7	*	0.9	10.1	32.1
Total knee replacement	81.54	13.3	*	0.8	19.6	67.6
Operations on the integumentary system	85-86	46.9	*	31.3	62.3	120.5
Debridement of wound, infection, or burn	86.22,86.28	12.6	*	7.1	17.1	39.4
Miscellaneous diagnostic and therapeutic procedures	87-99	429.5	146.7	191.8	507.2	1,595.3
Computerized axial tomography	87.03,87.41,87.71,88.01,88.38	24.5	6.9	12.6	25.3	94.3
Arteriography and angiocardiology using contrast material	88.4-88.5	71.7	*	15.6	123.4	288.7
Diagnostic ultrasound	88.7	26.9	*	10.7	28.8	113.4
Respiratory therapy	93.9,96.7	37.3	34.2	11.6	36.4	133.7
Insertion of endotracheal tube	96.04	16.6	7.9	5.0	18.9	67.6
Injection or infusion of cancer chemotherapeutic substance	99.25	7.6	5.6	4.1	10.9	17.1

\* Figure does not meet standard of reliability or precision.

... Category not applicable.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>See table 29 for comparison of rate of procedures with rate of discharges with procedures.

**Table 32. Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
All procedures . . . . .	42,533	16,834	25,700
Operations on the nervous system . . . . . 01-05	1,101	514	587
Spinal tap . . . . . 03.31	316	161	156
Operations on the endocrine system . . . . . 06-07	102	30	72
Operations on the eye . . . . . 08-16	86	47	39
Operations on the ear . . . . . 18-20	46	24	22
Operations on the nose, mouth, and pharynx . . . . . 21-29	269	159	110
Operations on the respiratory system . . . . . 30-34	1,022	583	439
Bronchoscopy with or without biopsy . . . . . 33.21-33.24,33.27	251	147	104
Operations on the cardiovascular system . . . . . 35-39	6,813	3,967	2,845
Removal of coronary artery obstruction and insertion of stent(s) <sup>1</sup> . . . . . 36.0	1,204	802	402
Coronary artery bypass graft <sup>1</sup> . . . . . 36.1	515	373	142
Cardiac catheterization . . . . . 37.21-37.23	1,328	799	529
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	420	213	207
Hemodialysis . . . . . 39.95	552	288	264
Operations on the hemic and lymphatic system . . . . . 40-41	354	178	175
Operations on the digestive system . . . . . 42-54	5,597	2,350	3,247
Endoscopy of small intestine with or without biopsy . . . . . 45.11-45.14,45.16	1,032	450	582
Endoscopy of large intestine with or without biopsy . . . . . 45.21-45.25	578	235	343
Partial excision of large intestine . . . . . 45.7	263	113	150
Appendectomy, excluding incidental . . . . . 47.0	329	173	156
Cholecystectomy . . . . . 51.2	436	143	293
Lysis of peritoneal adhesions . . . . . 54.5	342	62	280
Operations on the urinary system . . . . . 55-59	955	454	501
Cystoscopy with or without biopsy . . . . . 57.31-57.33	173	86	87
Operations on the male genital organs . . . . . 60-64	262	262	...
Prostatectomy . . . . . 60.2-60.6	195	195	...
Operations on the female genital organs . . . . . 65-71	2,161	...	2,161
Oophorectomy and salpingo-oophorectomy . . . . . 65.3-65.6	533	...	533
Bilateral destruction or occlusion of fallopian tubes . . . . . 66.2-66.3	329	...	329
Hysterectomy . . . . . 68.3-68.7,68.9	669	...	669
Obstetrical procedures . . . . . 72-75	6,646	...	6,646
Episiotomy with or without forceps or vacuum extraction . . . . . 72.1,72.21,72.31,72.71,73.6	780	...	780
Artificial rupture of membranes . . . . . 73.0	901	...	901
Cesarean section . . . . . 74.0-74.2,74.4,74.99	1,059	...	1,059
Repair of current obstetric laceration . . . . . 75.5-75.6	1,234	...	1,234
Operations on the musculoskeletal system . . . . . 76-84	3,442	1,652	1,790
Partial excision of bone . . . . . 76.2-76.3,77.6-77.8	218	112	106
Reduction of fracture . . . . . 76.7,79.0-79.3	606	288	318
Open reduction of fracture with internal fixation . . . . . 79.3	414	186	229
Excision or destruction of intervertebral disc . . . . . 80.5	323	171	152
Total hip replacement . . . . . 81.51	193	81	112
Total knee replacement . . . . . 81.54	381	146	235
Operations on the integumentary system . . . . . 85-86	1,348	616	732
Debridement of wound, infection, or burn . . . . . 86.22,86.28	361	199	161
Miscellaneous diagnostic and therapeutic procedures . . . . . 87-99	12,332	5,996	6,336
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	703	324	378
Arteriography and angiocardiology using contrast material . . . . . 88.4-88.5	2,058	1,142	915
Diagnostic ultrasound . . . . . 88.7	773	347	427
Respiratory therapy . . . . . 93.9,96.7	1,070	546	524
Insertion of endotracheal tube . . . . . 96.04	477	247	229
Injection or infusion of cancer chemotherapeutic substance . . . . . 99.25	217	124	94

... Category not applicable.

<sup>1</sup>See table 29 for comparison of number of procedures with number of discharges with procedures.

**Table 33. Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
	Rate per 10,000 population <sup>1</sup>		
All procedures	1,481.4	1,197.3	1,753.9
Operations on the nervous system . . . . . 01-05	38.3	36.6	40.0
Spinal tap . . . . . 03.31	11.0	11.4	10.6
Operations on the endocrine system . . . . . 06-07	3.6	2.2	4.9
Operations on the eye . . . . . 08-16	3.0	3.4	2.6
Operations on the ear . . . . . 18-20	1.6	1.7	1.5
Operations on the nose, mouth, and pharynx . . . . . 21-29	9.4	11.3	7.5
Operations on the respiratory system . . . . . 30-34	35.6	41.5	29.9
Bronchoscopy with or without biopsy . . . . . 33.21-33.24,33.27	8.7	10.4	7.1
Operations on the cardiovascular system . . . . . 35-39	237.3	282.2	194.2
Removal of coronary artery obstruction and insertion of stent(s) <sup>2</sup> . . . . . 36.0	41.9	57.0	27.4
Coronary artery bypass graft <sup>2</sup> . . . . . 36.1	17.9	26.6	9.7
Cardiac catheterization . . . . . 37.21-37.23	46.2	56.8	36.1
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	14.6	15.1	14.2
Hemodialysis . . . . . 39.95	19.2	20.5	18.1
Operations on the hemic and lymphatic system . . . . . 40-41	12.3	12.7	12.0
Operations on the digestive system . . . . . 42-54	194.9	167.2	221.6
Endoscopy of small intestine with or without biopsy . . . . . 45.11-45.14,45.16	35.9	32.0	39.7
Endoscopy of large intestine with or without biopsy . . . . . 45.21-45.25	20.1	16.7	23.4
Partial excision of large intestine . . . . . 45.7	9.2	8.1	10.2
Appendectomy, excluding incidental . . . . . 47.0	11.5	12.3	10.6
Cholecystectomy . . . . . 51.2	15.2	10.2	20.0
Lysis of peritoneal adhesions . . . . . 54.5	11.9	4.4	19.1
Operations on the urinary system . . . . . 55-59	33.2	32.3	34.2
Cystoscopy with or without biopsy . . . . . 57.31-57.33	6.0	6.1	6.0
Operations on the male genital organs . . . . . 60-64	9.1	18.6	...
Prostatectomy . . . . . 60.2-60.6	6.8	13.9	...
Operations on the female genital organs . . . . . 65-71	75.3	...	147.5
Oophorectomy and salpingo-oophorectomy . . . . . 65.3-65.6	18.6	...	36.4
Bilateral destruction or occlusion of fallopian tubes . . . . . 66.2-66.3	11.5	...	22.5
Hysterectomy . . . . . 68.3-68.7,68.9	23.3	...	45.7
Obstetrical procedures . . . . . 72-75	231.5	...	453.6
Episiotomy with or without forceps or vacuum extraction . . . . . 72.1,72.21,72.31,72.71,73.6	27.2	...	53.2
Artificial rupture of membranes . . . . . 73.0	31.4	...	61.5
Cesarean section . . . . . 74.0-74.2,74.4,74.99	36.9	...	72.2
Repair of current obstetric laceration . . . . . 75.5-75.6	43.0	...	84.2
Operations on the musculoskeletal system . . . . . 76-84	119.9	117.5	122.1
Partial excision of bone . . . . . 76.2-76.3,77.6-77.8	7.6	8.0	7.3
Reduction of fracture . . . . . 76.7,79.0-79.3	21.1	20.5	21.7
Open reduction of fracture with internal fixation . . . . . 79.3	14.4	13.2	15.6
Excision or destruction of intervertebral disc . . . . . 80.5	11.3	12.2	10.4
Total hip replacement . . . . . 81.51	6.7	5.8	7.6
Total knee replacement . . . . . 81.54	13.3	10.4	16.0
Operations on the integumentary system . . . . . 85-86	46.9	43.8	49.9
Debridement of wound, infection, or burn . . . . . 86.22,86.28	12.6	14.2	11.0
Miscellaneous diagnostic and therapeutic procedures . . . . . 87-99	429.5	426.5	432.4
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	24.5	23.1	25.8
Arteriography and angiocardiology using contrast material . . . . . 88.4-88.5	71.7	81.3	62.5
Diagnostic ultrasound . . . . . 88.7	26.9	24.7	29.1
Respiratory therapy . . . . . 93.9,96.7	37.3	38.8	35.8
Insertion of endotracheal tube . . . . . 96.04	16.6	17.6	15.7
Injection or infusion of cancer chemotherapeutic substance . . . . . 99.25	7.6	8.8	6.4

... Category not applicable.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>See table 29 for comparison of rate of procedures with rate of discharges with procedures.



**Table 34. Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures . . . . .	42,533	9,389	9,082	15,394	8,668
Operations on the nervous system . . . . . 01-05	1,101	240	277	345	239
Spinal tap . . . . . 03.31	316	67	72	105	73
Operations on the endocrine system . . . . . 06-07	102	22	13	39	29
Operations on the eye . . . . . 08-16	86	15	17	33	21
Operations on the ear . . . . . 18-20	46	15	6	17	*
Operations on the nose, mouth, and pharynx . . . . . 21-29	269	72	45	91	60
Operations on the respiratory system . . . . . 30-34	1,022	248	208	356	209
Bronchoscopy with or without biopsy . . . . . 33.21-33.24,33.27	251	65	44	95	47
Operations on the cardiovascular system . . . . . 35-39	6,813	1,370	1,463	2,601	1,378
Removal of coronary artery obstruction and insertion of stent(s) <sup>1</sup> . . . . . 36.0	1,204	211	323	416	254
Coronary artery bypass graft <sup>1</sup> . . . . . 36.1	515	104	117	204	90
Cardiac catheterization . . . . . 37.21-37.23	1,328	243	312	541	231
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	420	107	77	158	78
Hemodialysis . . . . . 39.95	552	94	90	242	127
Operations on the hemic and lymphatic system . . . . . 40-41	354	81	66	130	77
Operations on the digestive system . . . . . 42-54	5,597	1,185	1,226	2,111	1,075
Endoscopy of small intestine with or without biopsy . . . . . 45.11-45.14,45.16	1,032	226	235	391	179
Endoscopy of large intestine with or without biopsy . . . . . 45.21-45.25	578	131	130	227	91
Partial excision of large intestine . . . . . 45.7	263	58	64	96	45
Appendectomy, excluding incidental . . . . . 47.0	329	62	73	99	96
Cholecystectomy . . . . . 51.2	436	85	91	170	89
Lysis of peritoneal adhesions . . . . . 54.5	342	66	71	148	57
Operations on the urinary system . . . . . 55-59	955	218	206	364	166
Cystoscopy with or without biopsy . . . . . 57.31-57.33	173	42	41	65	25
Operations on the male genital organs . . . . . 60-64	262	57	58	98	49
Prostatectomy . . . . . 60.2-60.6	195	39	42	74	40
Operations on the female genital organs . . . . . 65-71	2,161	345	432	943	442
Oophorectomy and salpingo-oophorectomy . . . . . 65.3-65.6	533	80	115	232	107
Bilateral destruction or occlusion of fallopian tubes . . . . . 66.2-66.3	329	35	49	180	65
Hysterectomy . . . . . 68.3-68.7,68.9	669	107	153	279	131
Obstetrical procedures . . . . . 72-75	6,646	1,054	1,387	2,569	1,635
Episiotomy with or without forceps or vacuum extraction . . . . . 72.1,72.21,72.31,72.71,73.6	780	119	151	344	167
Artificial rupture of membranes . . . . . 73.0	901	127	249	306	220
Cesarean section . . . . . 74.0-74.2,74.4,74.99	1,059	162	175	489	233
Repair of current obstetric laceration . . . . . 75.5-75.6	1,234	197	252	443	341
Operations on the musculoskeletal system . . . . . 76-84	3,442	704	789	1,234	715
Partial excision of bone . . . . . 76.2-76.3,77.6-77.8	218	44	44	83	46
Reduction of fracture . . . . . 76.7,79.0-79.3	606	130	145	206	124
Open reduction of fracture with internal fixation . . . . . 79.3	414	88	100	145	82
Excision or destruction of intervertebral disc . . . . . 80.5	323	56	62	135	70
Total hip replacement . . . . . 81.51	193	54	48	55	36
Total knee replacement . . . . . 81.54	381	81	104	132	65
Operations on the integumentary system . . . . . 85-86	1,348	291	260	576	221
Debridement of wound, infection, or burn . . . . . 86.22,86.28	361	76	77	147	61
Miscellaneous diagnostic and therapeutic procedures . . . . . 87-99	12,332	3,472	2,629	3,886	2,344
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	703	319	*	137	119
Arteriography and angiocardiology using contrast material . . . . . 88.4-88.5	2,058	391	518	818	331
Diagnostic ultrasound . . . . . 88.7	773	271	137	200	165
Respiratory therapy . . . . . 93.9,96.7	1,070	274	231	311	254
Insertion of endotracheal tube . . . . . 96.04	477	108	99	181	88
Injection or infusion of cancer chemotherapeutic substance . . . . . 99.25	217	53	35	80	49

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>See table 29 for comparison of number of procedures with number of discharges with procedures.

**Table 35. Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures . . . . .	1,481.4	1,733.3	1,396.3	1,500.4	1,327.2
Operations on the nervous system . . . . . 01-05	38.3	44.2	42.6	33.6	36.6
Spinal tap . . . . . 03.31	11.0	12.3	11.1	10.2	11.2
Operations on the endocrine system . . . . . 06-07	3.6	4.1	1.9	3.8	4.4
Operations on the eye . . . . . 08-16	3.0	2.8	2.6	3.3	3.1
Operations on the ear . . . . . 18-20	1.6	2.8	1.0	1.6	*
Operations on the nose, mouth, and pharynx . . . . . 21-29	9.4	13.4	7.0	8.9	9.2
Operations on the respiratory system . . . . . 30-34	35.6	45.8	32.0	34.7	32.0
Bronchoscopy with or without biopsy . . . . . 33.21-33.24,33.27	8.7	12.0	6.8	9.2	7.2
Operations on the cardiovascular system . . . . . 35-39	237.3	253.0	224.9	253.5	211.0
Removal of coronary artery obstruction and insertion of stent(s) <sup>2</sup> . . . . . 36.0	41.9	38.9	49.7	40.5	38.8
Coronary artery bypass graft <sup>2</sup> . . . . . 36.1	17.9	19.3	17.9	19.9	13.7
Cardiac catheterization . . . . . 37.21-37.23	46.2	44.9	48.0	52.8	35.3
Insertion, replacement, removal, and revision of pacemaker leads or device . . . . . 37.7-37.8	14.6	19.7	11.8	15.4	12.0
Hemodialysis . . . . . 39.95	19.2	17.4	13.8	23.6	19.4
Operations on the hemic and lymphatic system . . . . . 40-41	12.3	14.9	10.2	12.7	11.8
Operations on the digestive system . . . . . 42-54	194.9	218.7	188.5	205.8	164.6
Endoscopy of small intestine with or without biopsy . . . . . 45.11-45.14,45.16	35.9	41.7	36.2	38.1	27.5
Endoscopy of large intestine with or without biopsy . . . . . 45.21-45.25	20.1	24.1	19.9	22.1	14.0
Partial excision of large intestine . . . . . 45.7	9.2	10.7	9.9	9.3	7.0
Appendectomy, excluding incidental . . . . . 47.0	11.5	11.4	11.2	9.6	14.7
Cholecystectomy . . . . . 51.2	15.2	15.7	14.0	16.6	13.7
Lysis of peritoneal adhesions . . . . . 54.5	11.9	12.2	10.9	14.5	8.7
Operations on the urinary system . . . . . 55-59	33.2	40.3	31.7	35.5	25.4
Cystoscopy with or without biopsy . . . . . 57.31-57.33	6.0	7.7	6.4	6.3	3.9
Operations on the male genital organs . . . . . 60-64	9.1	10.5	8.9	9.5	7.5
Prostatectomy . . . . . 60.2-60.6	6.8	7.2	6.5	7.2	6.1
Operations on the female genital organs . . . . . 65-71	75.3	63.6	66.4	91.9	67.7
Oophorectomy and salpingo-oophorectomy . . . . . 65.3-65.6	18.6	14.7	17.6	22.6	16.3
Bilateral destruction or occlusion of fallopian tubes . . . . . 66.2-66.3	11.5	6.4	7.6	17.6	9.9
Hysterectomy . . . . . 68.3-68.7,68.9	23.3	19.7	23.6	27.2	20.0
Obstetrical procedures . . . . . 72-75	231.5	194.6	213.2	250.4	250.4
Episiotomy with or without forceps or vacuum extraction . . . . . 72.1,72.21,72.31,72.71,73.6	27.2	21.9	23.2	33.5	25.5
Artificial rupture of membranes . . . . . 73.0	31.4	23.4	38.3	29.8	33.7
Cesarean section . . . . . 74.0-74.2,74.4,74.99	36.9	29.9	27.0	47.6	35.6
Repair of current obstetric laceration . . . . . 75.5-75.6	43.0	36.4	38.8	43.2	52.2
Operations on the musculoskeletal system . . . . . 76-84	119.9	130.0	121.2	120.2	109.5
Partial excision of bone . . . . . 76.2-76.3,77.6-77.8	7.6	8.2	6.8	8.1	7.1
Reduction of fracture . . . . . 76.7,79.0-79.3	21.1	24.1	22.3	20.1	19.0
Open reduction of fracture with internal fixation . . . . . 79.3	14.4	16.2	15.4	14.1	12.5
Excision or destruction of intervertebral disc . . . . . 80.5	11.3	10.4	9.5	13.2	10.7
Total hip replacement . . . . . 81.51	6.7	10.1	7.4	5.4	5.4
Total knee replacement . . . . . 81.54	13.3	14.9	15.9	12.9	9.9
Operations on the integumentary system . . . . . 85-86	46.9	53.6	40.0	56.1	33.8
Debridement of wound, infection, or burn . . . . . 86.22,86.28	12.6	13.9	11.8	14.3	9.4
Miscellaneous diagnostic and therapeutic procedures . . . . . 87-99	429.5	641.0	404.2	378.8	358.9
Computerized axial tomography . . . . . 87.03,87.41,87.71,88.01,88.38	24.5	58.8	*	13.4	18.2
Arteriography and angiocardiology using contrast material . . . . . 88.4-88.5	71.7	72.2	79.6	79.7	50.6
Diagnostic ultrasound . . . . . 88.7	26.9	50.1	21.0	19.5	25.3
Respiratory therapy . . . . . 93.9,96.7	37.3	50.5	35.6	30.3	38.9
Insertion of endotracheal tube . . . . . 96.04	16.6	19.9	15.3	17.7	13.5
Injection or infusion of cancer chemotherapeutic substance . . . . . 99.25	7.6	9.8	5.4	7.8	7.5

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>See table 29 for comparison of number of procedures with number of discharges with procedures.

**Table 36. Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals by age and geographic region, according to type of delivery: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population <sup>1</sup>	Number in thousands	Percent distribution	Rate per 1,000 population <sup>1</sup>	
All deliveries							
10–54 years . . . . .	3,951	100.0	42.7	10,207	100.0	110.2	2.6
10–19 years . . . . .	455	11.5	22.5	1,167	11.4	57.8	2.6
20–24 years . . . . .	997	25.2	101.8	2,432	23.8	248.3	2.4
25–29 years . . . . .	1,026	26.0	110.4	2,574	25.2	276.9	2.5
30–34 years . . . . .	928	23.5	89.5	2,436	23.9	234.9	2.6
35–39 years . . . . .	444	11.2	40.6	1,271	12.4	116.1	2.9
40–54 years . . . . .	100	2.5	3.1	327	3.2	10.2	3.3
All regions . . . . .	3,951	100.0	42.7	10,207	100.0	110.2	2.6
Northeast . . . . .	582	14.7	33.4	1,658	16.2	95.1	2.8
Midwest . . . . .	736	18.6	35.2	1,845	18.1	88.4	2.5
South . . . . .	1,660	42.0	50.1	4,320	42.3	130.5	2.6
West . . . . .	974	24.6	46.1	2,383	23.3	112.7	2.4
Vaginal deliveries <sup>2</sup>							
10–54 years . . . . .	2,893	100.0	31.2	6,278	100.0	67.8	2.2
10–19 years . . . . .	371	12.8	18.4	846	13.5	41.9	2.3
20–24 years . . . . .	774	26.8	79.1	1,632	26.0	166.7	2.1
25–29 years . . . . .	748	25.8	80.4	1,572	25.0	169.2	2.1
30–34 years . . . . .	666	23.0	64.2	1,461	23.3	140.9	2.2
35–39 years . . . . .	277	9.6	25.3	616	9.8	56.3	2.2
40–54 years . . . . .	57	2.0	1.8	150	2.4	4.7	2.6
All regions . . . . .	2,893	100.0	31.2	6,278	100.0	67.8	2.2
Northeast . . . . .	420	14.5	24.1	1,000	15.9	57.4	2.4
Midwest . . . . .	560	19.4	26.8	1,201	19.1	57.5	2.1
South . . . . .	1,172	40.5	35.4	2,588	41.2	78.2	2.2
West . . . . .	741	25.6	35.1	1,488	23.7	70.4	2.0
Cesarean deliveries <sup>3</sup>							
10–54 years . . . . .	1,058	100.0	11.4	3,929	100.0	42.4	3.7
10–19 years . . . . .	84	7.9	4.2	321	8.2	15.9	3.8
20–24 years . . . . .	223	21.1	22.8	800	20.4	81.7	3.6
25–29 years . . . . .	279	26.3	30.0	1,002	25.5	107.8	3.6
30–34 years . . . . .	262	24.7	25.2	975	24.8	94.0	3.7
35–39 years . . . . .	167	15.8	15.3	654	16.7	59.8	3.9
40–54 years . . . . .	43	4.1	1.4	177	4.5	5.5	4.1
All regions . . . . .	1,058	100.0	11.4	3,929	100.0	42.4	3.7
Northeast . . . . .	162	15.3	9.3	658	16.7	37.7	4.1
Midwest . . . . .	175	16.6	8.4	644	16.4	30.8	3.7
South . . . . .	489	46.2	14.7	1,732	44.1	52.3	3.5
West . . . . .	233	22.0	11.0	895	22.8	42.3	3.8

<sup>1</sup>Rates were calculated using U.S. Census Bureau estimates of the civilian population based on the 2000 census.<sup>2</sup>Discharges of females with deliveries except those with cesarean deliveries.<sup>3</sup>Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

**Table 37. Number and percent distribution of females with deliveries discharged from short-stay hospitals by length of stay and type of delivery: United States, selected years 1980–2002**[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Length of stay and type of delivery	1980	1985	1990	1995	2000	2002
Length of stay		Number in thousands				
All deliveries . . . . .	3,762	3,854	4,025	3,766	3,738	3,951
Less than 1 day . . . . .	23	34	37	36	16	*6
1 day . . . . .	216	373	623	1,368	551	526
2 days . . . . .	747	1,037	1,649	1,418	1,863	1,897
3 days . . . . .	1,205	1,140	900	606	863	991
4 days . . . . .	691	586	468	193	292	337
5–7 days . . . . .	726	598	282	100	112	134
8 days or more . . . . .	154	86	67	46	41	61
Vaginal deliveries <sup>1</sup> . . . . .	3,143	2,977	3,080	2,981	2,882	2,893
Less than 1 day . . . . .	23	33	35	34	14	*6
1 day . . . . .	215	371	620	1,362	546	519
2 days . . . . .	743	1,030	1,624	1,271	1,726	1,720
3 days . . . . .	1,185	1,054	618	228	500	535
4 days . . . . .	618	338	106	39	53	60
5–7 days . . . . .	317	125	55	27	25	30
8 days or more . . . . .	42	26	23	21	17	23
Cesarean deliveries <sup>2</sup> . . . . .	619	877	945	785	855	1,058
Less than 1 day . . . . .	*	*	*	*	*	*
1 day . . . . .	*	*	*	*6	*	*
2 days . . . . .	*	*	25	147	137	177
3 days . . . . .	20	86	283	379	363	456
4 days . . . . .	73	249	361	154	239	277
5–7 days . . . . .	409	473	227	73	87	104
8 days or more . . . . .	112	60	44	25	24	38
		Percent distribution				
All deliveries . . . . .	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day . . . . .	0.6	0.9	0.9	1.0	0.4	*0.1
1 day . . . . .	5.7	9.7	15.5	36.3	14.7	13.3
2 days . . . . .	19.9	26.9	41.0	37.7	49.9	48.0
3 days . . . . .	32.0	29.6	22.4	16.1	23.1	25.1
4 days . . . . .	18.4	15.2	11.6	5.1	7.8	8.5
5–7 days . . . . .	19.3	15.5	7.0	2.7	3.0	3.4
8 days or more . . . . .	4.1	2.2	1.7	1.2	1.1	1.5
Vaginal deliveries <sup>1</sup> . . . . .	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day . . . . .	0.7	1.1	1.1	1.1	0.5	*0.2
1 day . . . . .	6.8	12.5	20.1	45.7	19.0	17.9
2 days . . . . .	23.6	34.6	52.7	42.6	59.9	59.5
3 days . . . . .	37.7	35.4	20.1	7.6	17.3	18.5
4 days . . . . .	19.7	11.4	3.4	1.3	1.8	2.1
5–7 days . . . . .	10.1	4.2	1.8	0.9	0.9	1.0
8 days or more . . . . .	1.3	0.9	0.7	0.7	0.6	0.8
Cesarean deliveries <sup>2</sup> . . . . .	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day . . . . .	*	*	*	*	*	*
1 day . . . . .	*	*	*	*0.8	*	*
2 days . . . . .	*	*	2.6	18.7	16.0	16.8
3 days . . . . .	3.2	9.8	29.9	48.3	42.4	43.1
4 days . . . . .	11.8	28.4	38.2	19.6	27.9	26.1
5–7 days . . . . .	66.1	53.9	24.0	9.3	10.2	9.8
8 days or more . . . . .	18.1	6.8	4.7	3.2	2.8	3.6
Type of delivery		Average length of stay in days				
All deliveries . . . . .	3.8	3.3	2.8	2.1	2.5	2.6
Vaginal deliveries <sup>1</sup> . . . . .	3.2	2.7	2.3	1.7	2.1	2.2
Cesarean deliveries <sup>2</sup> . . . . .	6.5	5.2	4.5	3.6	3.7	3.7

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Discharges of females with deliveries except those with cesarean deliveries.<sup>2</sup>Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

**Table 38. Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2002**[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Procedure category and ICD–9–CM code	1980	1985	1990	1995	2000	2002
Number of procedures in thousands						
All obstetrical procedures . . . . . 72–75	3,526	4,239	6,660	6,282	6,145	6,587
Repair of current obstetric laceration . . . . . 75.5–75.6	350	543	789	961	1,134	1,231
Cesarean section . . . . . 74.0–74.2,74.4,74.99	619	877	945	785	855	1,059
Manually assisted delivery . . . . . 73.5	70	193	750	866	898	993
Artificial rupture of membranes . . . . . 73.0	120	315	691	752	833	901
Episiotomy . . . . . 72.1,72.21,72.31,72.71,73.6	2,012	1,820	1,714	1,410	944	780
Medical induction of labor . . . . . 73.4	42	66	225	366	435	504
Vacuum extraction <sup>1</sup> . . . . . 72.7	22	66	190	279	245	234
Forceps delivery <sup>1</sup> . . . . . 72.0–72.4,72.51,72.53,72.6,73.3	555	376	269	176	117	98
Rate per 100 deliveries <sup>2</sup>						
All obstetrical procedures . . . . . 72–75	93.7	110.0	165.5	166.8	164.4	166.7
Cesarean section . . . . . 74.0–74.2,74.4,74.99	16.5	22.7	23.5	20.8	22.9	26.8
Artificial rupture of membranes . . . . . 73.0	3.2	8.2	17.2	20.0	22.3	22.8
Medical induction of labor . . . . . 73.4	1.1	1.7	5.6	9.7	11.7	12.8
Rate per 100 vaginal deliveries <sup>3</sup>						
Repair of current obstetric laceration . . . . . 75.5–75.6	11.1	18.1	25.5	32.0	39.2	42.4
Manually assisted delivery . . . . . 73.5	2.2	6.5	24.3	29.0	31.1	34.2
Episiotomy . . . . . 72.1,72.21,72.31,72.71,73.6	64.0	61.1	55.6	47.2	32.7	26.9
Vacuum extraction <sup>1</sup> . . . . . 72.7	0.7	2.2	6.1	9.2	8.4	7.9
Forceps delivery <sup>1</sup> . . . . . 72.0–72.4,72.51,72.53,72.6,73.3	17.6	12.5	8.6	5.8	4.0	3.3

<sup>1</sup>Forceps delivery and vacuum extraction with episiotomy are also included in the episiotomy category.<sup>2</sup>Procedures per 100 discharges of females with deliveries.<sup>3</sup>Procedures for females with vaginal deliveries per 100 discharges with vaginal deliveries. Vaginal deliveries are all deliveries except those with a cesarean section.**Table 39. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery: United States, selected years 1980–2002**[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Type of delivery	1980	1985	1990	1995	2000	2002
Number in thousands						
Females with deliveries . . . . .	3,762	3,854	4,025	3,766	3,738	3,951
Previous cesarean delivery . . . . .	192	324	426	413	447	499
No previous cesarean delivery . . . . .	3,571	3,530	3,600	3,353	3,290	3,453
Vaginal deliveries . . . . .	3,143	2,977	3,080	2,981	2,882	2,893
Vaginal birth after cesarean delivery (VBAC) . . . . .	*6	21	87	147	123	79
Other vaginal deliveries . . . . .	3,137	2,956	2,993	2,835	2,759	2,814
Cesarean deliveries . . . . .	619	877	945	785	855	1,058
No previous cesarean delivery (primary) . . . . .	434	574	606	519	531	638
Previous cesarean delivery (repeat) . . . . .	185	303	339	266	324	420
Rate per 100 deliveries						
Cesarean deliveries per 100 deliveries . . . . .	16.5	22.7	23.5	20.8	22.9	26.8
Primary cesarean deliveries per 100 deliveries without previous cesarean delivery . . . . .	12.1	16.3	16.8	15.5	16.1	18.5
Repeat cesarean deliveries per 100 cesarean deliveries . . . . .	29.9	34.6	35.9	33.9	37.9	39.7
Vaginal birth after cesarean deliveries (VBAC) per 100 deliveries with previous cesarean delivery . . . . .	*3.4	6.6	20.4	35.5	27.6	15.8

\* Figure does not meet standard of reliability or precision.

NOTE: Previous cesarean deliveries are discharges with ICD–9–CM diagnostic code 654.2. Cesarean deliveries are discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4 or 74.99.

**Table 40. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Cesarean deliveries			
	Total <sup>1</sup>	Primary <sup>2</sup>	Repeat <sup>3</sup>	VBAC <sup>4,5</sup>
Number in thousands				
10–54 years . . . . .	1,058	638	420	79
10–19 years . . . . .	84	73	11	*
20–24 years . . . . .	223	144	79	18
25–29 years . . . . .	279	167	112	20
30–34 years . . . . .	262	139	123	23
35–39 years . . . . .	167	93	74	14
40–54 years . . . . .	43	22	21	*
All regions . . . . .	1,058	638	420	79
Northeast . . . . .	162	98	64	12
Midwest . . . . .	175	102	73	15
South . . . . .	489	304	185	33
West . . . . .	233	134	98	18
Rate per 100 deliveries				
10–54 years . . . . .	26.8	18.5	39.7	15.8
10–19 years . . . . .	18.5	16.6	12.5	*
20–24 years . . . . .	22.4	16.0	35.6	18.1
25–29 years . . . . .	27.2	18.6	40.2	15.0
30–34 years . . . . .	28.2	17.8	46.9	15.8
35–39 years . . . . .	37.7	26.3	44.2	16.1
40–54 years . . . . .	43.2	28.9	49.2	*
All regions . . . . .	26.8	18.5	39.7	15.8
Northeast . . . . .	27.8	19.4	39.4	16.0
Midwest . . . . .	23.8	15.8	41.7	17.4
South . . . . .	29.4	21.1	37.9	15.1
West . . . . .	23.9	15.7	42.3	15.4

\* Figure does not meet standard of reliability or precision.

<sup>1</sup>Discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99. Rate per 100 total deliveries.<sup>2</sup>Cesarean deliveries without ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries without previous cesarean delivery.<sup>3</sup>Cesarean deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 cesarean deliveries.<sup>4</sup>VBAC is vaginal birth after cesarean delivery.<sup>5</sup>Vaginal deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries with previous cesarean delivery.



**Table 41. Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2002**

[Infants born in non-Federal hospitals]

Sex, region, and health status	Discharges		Average length of stay in days
	Number in thousands	Percent distribution	
All newborn infants . . . . .	3,789	100.0	3.2
Sex			
Male . . . . .	1,940	51.2	3.3
Female . . . . .	1,849	48.8	3.2
Region			
Northeast . . . . .	725	19.1	3.4
Midwest . . . . .	764	20.2	2.9
South . . . . .	1,354	35.7	3.4
West . . . . .	946	25.0	3.0
Health status			
Well <sup>1</sup> . . . . .	2,290	60.4	2.0
Male . . . . .	1,125	29.7	2.0
Female . . . . .	1,165	30.7	2.0
Sick <sup>2</sup> . . . . .	1,499	39.6	5.1
Male . . . . .	815	21.5	5.0
Female . . . . .	684	18.1	5.2

<sup>1</sup>Without any illness or risk-related diagnoses.

<sup>2</sup>With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

**Table 42. Number and percent distribution of newborn infants discharged from short-stay hospitals by length of stay, and average length of stay, by health status: United States, selected years 1980-2002**

[Infants born in non-Federal hospitals]

Length of stay and health status	1980	1985	1990	1995	2000	2002
Length of stay						
Number in thousands						
All newborn infants . . . . .	3,824	3,794	3,869	3,631	3,643	3,789
Less than 1 day . . . . .	69	84	79	87	45	34
1 day . . . . .	270	428	693	1,523	722	707
2 days . . . . .	842	1,057	1,624	1,145	1,877	1,927
3 days . . . . .	1,175	1,034	636	486	546	614
4 days . . . . .	618	528	447	134	205	234
5-7 days . . . . .	633	497	232	106	94	107
8 days or more . . . . .	218	167	157	149	153	166
Percent distribution						
All newborn infants . . . . .	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day . . . . .	1.8	2.2	2.0	2.4	1.2	0.9
1 day . . . . .	7.1	11.3	17.9	41.9	19.8	18.7
2 days . . . . .	22.0	27.9	42.0	31.5	51.5	50.9
3 days . . . . .	30.7	27.3	16.4	13.4	15.0	16.2
4 days . . . . .	16.2	13.9	11.6	3.7	5.6	6.2
5-7 days . . . . .	16.5	13.1	6.0	2.9	2.6	2.8
8 days or more . . . . .	5.7	4.4	4.1	4.1	4.2	4.4
Health status						
Average length of stay in days						
All newborn infants . . . . .	4.3	3.8	3.3	2.8	3.2	3.2
Well <sup>1</sup> . . . . .	3.2	2.8	2.3	1.7	2.1	2.0
Sick <sup>2</sup> . . . . .	7.1	5.7	4.8	4.4	5.1	5.1

<sup>1</sup>Without any illness or risk-related diagnoses.

<sup>2</sup>With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

**Table 43. Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2002**[Infants born in non-Federal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Diagnostic category and ICD-9-CM code	Both sexes	Male	Female
Number in thousands			
Sick newborn diagnoses . . . . . 001-999	2,844	1,567	1,277
Hemolytic disease of fetus or newborn, due to isoimmunization, and other perinatal jaundice . . . . . 773-774	623	327	296
Neonatal jaundice associated with preterm delivery . . . . . 774.2	127	66	61
Unspecified fetal and neonatal jaundice . . . . . 774.6	430	228	202
Respiratory distress syndrome and other respiratory conditions of fetus and newborn . . . . . 769-770	374	208	166
Transitory tachypnea of newborn . . . . . 770.6	104	59	45
Disorders relating to short gestation and unspecified low birthweight (prematurity) . . . . . 765	323	166	157
Congenital anomalies . . . . . 740-759	229	132	97
Disorders relating to long gestation and high birthweight . . . . . 766	188	116	72
Endocrine and metabolic disturbances specific to the fetus and newborn . . . . . 775	124	68	56
Birth trauma . . . . . 767	109	59	50
Injuries to scalp . . . . . 767.1	72	44	28
Infections specific to the perinatal period . . . . . 771	98	59	39

**Table 44. Number and percent of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years 1980-2002**[Infants born in non-Federal hospitals with *International Classification of Diseases, 9th Revision, Clinical Modification*, procedure code 64.0]

	1980	1985	1990	1995	2000	2002
Number circumcised in thousands						
United States . . . . .	1,261	1,162	1,169	1,200	1,166	1,166
Northeast . . . . .	253	221	238	249	256	255
Midwest . . . . .	403	331	341	317	330	312
South . . . . .	379	393	396	454	438	434
West . . . . .	226	218	195	180	143	165
Percent circumcised						
United States . . . . .	64.7	59.5	59.0	64.1	62.4	60.1
Northeast . . . . .	67.4	65.2	62.6	68.3	64.6	68.9
Midwest . . . . .	75.9	70.5	76.0	79.8	81.4	81.0
South . . . . .	56.0	56.0	57.1	66.1	63.9	64.0
West . . . . .	61.8	49.0	42.4	42.6	37.3	32.6

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	33,727	13,389	20,338	2,540	10,736	7,723	12,727	6,990	7,503	12,994	6,239
001-139	877	417	460	156	204	185	332	183	190	339	166
005	*5	*	*	*	*	*	*	*	*	*	*
008	140	51	88	35	31	24	50	29	42	48	22
008.4	48	15	33	*	11	10	25	13	17	12	*6
008.45	43	14	29	*	10	9	22	11	15	11	*5
008.6	19	10	9	18	*	*	*	*	*7	*6	*
008.61	18	9	*8	17	*	-	*	*	*6	*5	*
008.8	70	25	45	14	18	14	24	13	16	29	11
009	18	*8	10	*	*8	*	*	*	*	*8	*
009.0	10	*	*7	*	*	*	*	*	*	*	*
011	*6	*	*	*	*	*	*	*	*	*	*
034	12	*	*7	*6	*	*	*	*	*	*	*
034.0	11	*	*6	*	*	*	*	*	*	*	*
038	341	156	186	12	34	76	219	74	77	126	64
038.0	22	11	11	*	*	*	15	*6	*6	*	*6
038.1	44	22	22	*	*6	10	26	10	10	14	10
038.10	*8	*	*	*	*	*	*	*	*	*	*
038.11	24	13	10	*	*	*6	14	*6	*	*8	*
038.19	13	*6	*7	*	*	*	*9	*	*5	*	*
038.2	10	*	*6	*	*	*	*	*	*	*	*
038.4	83	35	48	*	*7	20	55	18	20	26	19
038.40	10	*	*5	-	*	*	*6	*	*	*5	*
038.42	47	17	30	*	*	10	34	11	13	12	11
038.49	20	10	10	*	*	*7	11	*	*	*8	*5
038.8	*9	*	*5	*	*	*	*6	*	*	*	*
038.9	171	78	93	*8	16	35	112	34	35	76	26
041	14	*5	*8	*	*	*	*	*	*	*6	*
042	67	43	24	*	43	22	*	20	*6	31	11
047	44	23	22	21	19	*	*	*7	*8	17	12
047.9	40	21	19	18	17	*	*	*6	*7	16	10
053	19	*9	10	*	*	*	14	*	*	*6	*
053.9	10	*	*6	*	*	*	*7	*	*	*	*
054	*6	*	*	*	*	*	*	*	*	*	*
070	24	14	10	*	*7	13	*	*	*	10	*5
070.5	11	*6	*	*	*	*7	*	*	*	*	*
070.51	*6	*	*	*	*	*	*	*	*	*	*
075	9	*	*5	*	*	*	-	*	*	*	*
079	71	37	34	43	12	*6	10	16	12	31	13
079.9	65	34	32	38	12	*6	10	14	11	28	13
079.99	65	34	31	38	12	*6	10	14	11	28	13
112	15	*7	*8	*	*	*	*6	*	*	*7	*
117	*6	*	*	*	*	*	*	*	*	*	*
135	*7	*	*6	-	*	*	*	*	*	*	*
136	*6	*	*	*	*	*	*	*	*	*	*
140-239	1,682	650	1,033	33	313	611	725	358	353	647	324
150	13	*9	*	-	*	*	*8	*	*	*	*
150.9	*5	*	*	-	*	*	*	-	*	*	*
151	22	15	*7	-	*	*8	11	*	*	10	*
151.0	*6	*	*	-	*	*	*	*	*	*	*
151.2	*6	*	*	-	*	*	*	*	*	*	*
153	113	47	66	*	*	29	80	25	29	39	21
153.0	*6	*	*	-	*	*	*5	*	*	*	*
153.1	*8	*	*6	-	*	*	*5	*	*	*	*
153.2	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
153.3	32	16	16	—	*	11	20	*5	*7	13	*7
153.4	22	9	13	—	*	*	17	*6	*8	*	*
153.6	19	*6	13	—	*	*	16	*	*	*8	*
153.9	12	*	*7	—	*	*	*9	*	*	*	*
154	42	27	15	—	*	15	25	9	13	14	*6
154.0	16	10	*6	—	*	*6	10	*	*6	*6	*
154.1	22	16	*7	—	*	*8	14	*	*7	*7	*
155	15	11	*	*	*	*7	*8	*	*	*6	*
155.0	10	*8	*	*	*	*5	*	*	*	*	*
156	*9	*	*5	—	*	*	*7	*	*	*	*
157	30	16	15	*	*	*8	20	*6	*8	13	*
157.0	15	*8	*6	—	*	*	11	*	*	*6	*
157.9	11	*	*6	—	*	*	*7	*	*	*5	*
161	*8	*7	*	—	*	*	*	*	*	*	*
162	139	77	62	*	*	48	88	33	30	52	23
162.2	*7	*	*	*	*	*	*	*	*	*	*
162.3	51	26	25	—	*	19	31	12	11	17	11
162.5	24	12	12	—	*	*9	15	*6	*6	*7	*
162.8	13	*7	*6	—	*	*	*8	*	*	*	*
162.9	42	27	15	—	*	11	30	*7	*9	21	*
171	*8	*	*	*	*	*	*	*	*	*	*
172	*5	*	*	—	*	*	*	*	*	*	*
173	*6	*	*	*	*	*	*	*	*	*	*
174	85	—	85	—	11	36	38	17	17	32	19
174.4	12	—	12	—	*	*	*5	*	*	*6	*
174.8	28	—	28	—	*	11	13	*9	*	*8	*7
174.9	30	—	30	—	*	14	12	*	*6	14	*7
180	17	—	17	*	*	*8	*	*	*	*7	*5
180.8	*6	—	*6	—	*	*	*	*	*	*	*
180.9	*7	—	*7	*	*	*	*	*	*	*	*
182	29	—	29	—	*	9	17	*7	*7	10	*
182.0	28	—	28	—	*	9	17	*7	*7	10	*
183	23	—	23	*	*	11	9	*	*5	10	*
183.0	23	—	23	*	*	10	9	*	*5	9	*
185	92	92	—	—	*	44	47	19	22	30	21
188	33	24	*8	—	*	*6	26	*7	*7	13	*6
188.8	13	10	*	—	*	*	10	*	*	*	*
188.9	*7	*6	*	—	—	*	*6	*	*	*	*
189	34	23	11	*	*	13	17	*6	*9	17	*
189.0	30	20	10	*	*	11	14	*	*8	14	*
191	34	18	16	*5	*8	11	10	9	*7	10	*8
191.1	*7	*	*	*	*	*	*	*	*	*	*
191.9	*7	*	*	*	*	*	*	*	*	*	*
193	21	*5	16	*	9	*6	*5	*	*	*8	*5
196	12	*6	*6	*	*	*	*7	*	*	*	*
197	97	40	57	*	*6	35	56	24	25	34	15
197.0	20	*8	12	*	*	*7	12	*	*8	*6	*
197.2	21	*7	14	*	*	*6	14	*6	*	*7	*
197.6	18	*6	12	—	*	*7	9	*	*	*7	*
197.7	29	15	15	—	*	10	17	9	*7	10	*
198	89	37	52	*	*9	36	44	20	19	33	17
198.3	38	16	22	*	*	17	18	*9	*8	13	*8
198.5	32	13	18	*	*	12	18	*7	*7	12	*5
198.8	11	*	*7	*	*	*	*	*	*	*	*
198.89	10	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
199	24	13	11	*	*	*8	13	*5	*6	*8	*
199.0	17	9	*8	*	*	*6	*8	*	*	*5	*
199.1	*7	*	*	—	*	*	*	*	*	*	*
200	11	*7	*	*	*	*	*	*	*	*	*
200.0	*9	*5	*	—	*	*	*	*	*	*	*
202	37	21	16	*	*	11	22	*8	*8	14	*8
202.8	32	18	14	*	*	10	18	*6	*8	12	*6
202.80	12	*6	*6	*	*	*	*8	*	*	*	*
202.83	*7	*	*	*	*	*	*	*	*	*	*
202.88	*6	*	*	—	*	*	*	*	*	*	*
203	26	13	13	—	*	10	16	*	*6	14	*
203.0	26	13	13	—	*	9	16	*	*6	13	*
203.00	26	13	13	—	*	9	16	*	*6	13	*
204	25	12	13	*9	*	*	10	*6	*	*7	*8
204.0	13	*6	*6	*9	*	*	*	*	*	*	*
204.00	10	*5	*	*6	*	*	*	*	*	*	*
204.1	12	*6	*6	—	*	*	10	*	*	*	*
204.10	11	*6	*6	—	*	*	9	*	*	*	*
205	22	13	*9	*	*6	*5	*9	*	*	*8	*
205.0	17	10	*7	*	*6	*	*	*	*	*6	*
205.00	16	*9	*7	*	*	*	*	*	*	*6	*
205.1	*5	*	*	*	*	*	*	*	*	*	*
205.10	*5	*	*	*	*	*	*	*	*	*	*
211	46	21	26	*	*6	16	25	12	*7	18	10
211.3	36	17	19	—	*	13	19	9	*6	12	*8
218	256	—	256	—	134	117	*5	41	48	116	50
218.0	28	—	28	—	17	11	*	*8	*6	10	*
218.1	77	—	77	—	38	38	*	15	12	33	16
218.2	28	—	28	—	14	14	*	*	*5	14	*
218.9	123	—	123	—	65	53	*	14	25	60	25
220	44	—	44	*	23	16	*6	*8	9	17	10
225	21	*7	15	*	*	10	*7	*	*	10	*
225.2	15	*	11	*	*	*8	*5	*	*	*7	*
226	9	*	*7	*	*	*	*	*	*	*	*
227	18	*8	10	—	*5	*6	*7	*	*	*7	*6
227.1	9	*	*6	—	*	*	*	*	*	*	*
227.3	*7	*	*	—	*	*	*	*	*	*	*
233	28	*	27	—	14	10	*	*5	*	13	*5
233.0	10	—	10	—	*	*5	*	*	*	*	*
233.1	15	—	15	—	12	*	*	*	*	*8	*
235	14	*7	*7	*	*	*	*7	*	*	*7	*
235.2	9	*	*	*	*	*	*5	*	*	*5	*
236	*6	*	*	*	*	*	*	*	*	*	*
237	*7	*	*	*	*	*	*	*	*	*	*
238	10	*6	*	*	*	*	*6	*	*	*	*
238.7	*8	*	*	*	*	*	*5	*	*	*	*
239	11	*	*7	*	*	*	*5	*	*	*	*
240-279	1,619	661	958	187	328	454	649	350	367	621	281
241	21	*	18	*	*6	*9	*7	*	*	*7	*8
241.0	*9	*	*7	*	*	*	*	*	*	*	*
241.1	10	*	*9	*	*	*5	*	*	*	*	*
242	10	*	*6	*	*	*	*	*	*	*	*
242.0	*5	*	*	*	*	*	*	*	*	*	*
244	17	*	14	*	*	*6	*9	*7	*	*	*
244.9	16	*	13	*	*	*5	*8	*6	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
250 . . . . .	577	283	294	33	147	205	193	128	122	228	99
250.0 . . . . .	158	72	86	12	29	68	49	36	40	62	20
250.00 . . . . .	65	28	37	*	*8	30	25	16	16	26	*7
250.01 . . . . .	23	11	12	*8	*	*5	*5	*	*7	*8	*
250.02 . . . . .	50	21	29	*	*8	24	17	10	11	24	*6
250.03 . . . . .	21	13	*8	*	*9	*8	*	*6	*7	*	*
250.1 . . . . .	112	57	55	17	69	20	*7	20	28	42	21
250.10 . . . . .	14	*7	*7	*	*8	*	*	*	*	*	*
250.11 . . . . .	55	30	25	12	36	*7	*	11	14	19	12
250.12 . . . . .	13	*6	*8	*	*6	*	*	*	*	*5	*
250.13 . . . . .	30	14	16	*	19	*	*	*	*7	14	*
250.2 . . . . .	17	*8	*9	*	*	*6	*7	*6	*	*7	*
250.20 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
250.22 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
250.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
250.4 . . . . .	33	18	15	—	*	15	13	*	*6	13	10
250.40 . . . . .	20	11	9	—	*	11	*8	*	*	*8	*8
250.41 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
250.6 . . . . .	65	29	36	*	17	25	21	15	12	30	*8
250.60 . . . . .	30	12	18	*	*5	11	13	*8	*	13	*
250.61 . . . . .	14	*7	*7	*	*	*5	*	*	*	*7	*
250.62 . . . . .	12	*6	*6	—	*	*6	*	*	*	*6	*
250.63 . . . . .	9	*	*5	*	*	*	*	*	*	*	*
250.7 . . . . .	42	27	15	—	*	16	25	12	*7	15	*8
250.70 . . . . .	27	17	10	—	*	9	18	*6	*	10	*7
250.71 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
250.72 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
250.8 . . . . .	129	62	67	*	15	48	66	32	23	49	25
250.80 . . . . .	83	38	45	*	*	30	48	20	14	30	19
250.81 . . . . .	22	10	12	*	*	10	*7	*	*	11	*
250.82 . . . . .	15	*9	*6	—	*	*5	*7	*	*	*7	*
250.83 . . . . .	9	*	*	*	*	*	*	*	*	*	*
250.9 . . . . .	13	*7	*6	*	*	*	*	*	*	*6	*
250.92 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
251 . . . . .	14	*	10	*	*	*	*8	*	*	*6	*
251.2 . . . . .	11	*	*7	*	*	*	*7	*	*	*5	*
252 . . . . .	*9	*	*7	—	*	*	*	*	*	*	*
252.0 . . . . .	*8	*	*6	—	*	*	*	*	*	*	*
253 . . . . .	16	*6	10	*	*	*	9	*	*	*5	*
253.6 . . . . .	11	*	*8	*	*	*	9	*	*	*	*
255 . . . . .	13	*5	*8	*	*	*	*7	*	*	*6	*
255.4 . . . . .	10	*	*6	*	*	*	*6	*	*	*5	*
263 . . . . .	*9	*5	*	*	*	*	*	*	*	*	*
263.9 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
272 . . . . .	40	23	17	*	*8	23	*9	11	11	13	*
272.0 . . . . .	18	*9	*9	*	*	11	*	*6	*	*7	*
272.4 . . . . .	18	13	*6	—	*	11	*	*	*7	*5	*
274 . . . . .	13	*8	*	—	*	*	*8	*	*	*	*
274.0 . . . . .	10	*7	*	—	*	*	*6	*	*	*	*
275 . . . . .	21	*6	15	*	*	*7	10	*6	*7	*6	*
275.4 . . . . .	20	*5	14	*	*	*7	10	*	*7	*5	*
275.41 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
275.42 . . . . .	11	*	*9	*	*	*	*8	*	*	*	*
276 . . . . .	709	266	443	136	73	130	370	139	174	266	130
276.0 . . . . .	11	*	*8	*	*	*	*9	*	*	*	*
276.1 . . . . .	91	21	69	*	*5	18	65	14	21	33	23
276.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
276.5 . . . . .	508	204	304	129	53	77	249	103	128	198	79

See footnotes at end of table.



**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
276.6	*6	*	*	*	*	*	*	*	*	*	*
276.7	36	15	21	*	*	12	20	*5	*8	10	13
276.8	43	14	29	*	10	14	18	10	9	14	9
276.9	*6	*	*	—	*	*	*	*	*	*	—
277	18	10	*8	*5	10	*	*	*7	*	*5	*
277.0	12	*6	*6	*	*8	*	—	*	*	*	*
277.00	12	*6	*5	*	*8	*	—	*	*	*	*
278	106	22	84	*	59	44	*	22	22	46	16
278.0	102	22	80	*	57	42	*	20	21	45	16
278.00	9	*	*6	*	*	*	*	*	*	*	*
278.01	93	18	74	*	53	38	*	16	19	42	15
280-289	446	192	254	71	106	81	188	100	103	184	59
280	79	21	58	*	10	18	49	18	23	28	10
280.0	34	10	24	*	*	10	20	*9	10	13	*
280.9	43	10	33	*	*6	*7	29	9	13	15	*6
281	*5	*	*	*	*	*	*	*	*	*	*
282	88	41	47	27	52	*7	*	26	16	42	*
282.6	84	40	43	26	51	*6	*	24	15	41	*
282.60	*8	*6	*	*	*	*	*	*	*	*5	*
282.62	73	33	41	20	48	*6	—	23	13	35	*
283	*7	*	*	*	*	*	*	*	*	*	*
284	29	16	13	*	*	9	14	*	*7	13	*
284.8	25	13	12	*	*	*9	12	*	*7	9	*
285	104	41	64	*	10	21	71	23	24	39	18
285.1	13	*	10	*	*	*	*9	*	*6	*	*
285.2	11	*5	*6	—	*	*	*6	*	*	*	*
285.9	78	31	47	*	*8	14	54	18	15	30	15
286	23	10	13	*	*	*	13	*	*6	12	*
286.7	*6	*	*	*	*	*	*	*	*	*	*
286.9	*9	*	*	*	*	*	*5	*	*	*	*
287	31	13	17	*9	*8	*	*9	*8	*	11	*7
287.3	16	*	11	*6	*	*	*	*	*	*6	*
287.5	*8	*	*	*	*	*	*	*	*	*	*
288	59	31	28	14	13	13	19	13	14	25	*7
288.0	43	22	21	10	*8	11	13	9	*8	19	*6
288.8	16	*9	*7	*	*6	*	*6	*	*6	*6	*
289	21	13	*8	11	*	*	*	*	*	*9	*
289.2	*7	*5	*	*5	*	—	*	*	*	*	*
290-319	2,464	1,236	1,227	149	1,422	620	273	681	484	1,036	262
290	35	12	23	—	—	*	33	11	10	11	*
290.2	*8	*	*6	—	—	*	*8	*	*	*	*
290.4	17	*6	11	—	—	*	15	*	*	*6	*
290.40	*8	*	*6	—	—	*	*7	*	*	*	*
291	132	98	34	—	69	55	*8	31	47	22	32
291.0	19	16	*	—	*8	10	*	*	*	*	*6
291.8	107	78	29	—	59	42	*5	25	41	16	25
291.81	104	76	28	—	57	42	*5	25	39	15	25
292	82	49	33	*	52	17	12	36	15	20	11
292.0	45	27	18	*	33	11	*	23	9	*8	*5
292.8	28	17	11	*	14	*	9	11	*	*9	*
292.81	13	*7	*6	*	*	*	9	*	*	*6	*
292.84	14	9	*	*	12	*	—	*7	*	*	*
293	25	10	15	*	*	*7	15	*6	*7	*8	*
293.0	17	*6	11	*	*	*	12	*	*	*6	*
293.8	*6	*	*	*	*	*	*	*	*	*	*
293.83	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
294	46	21	26	*	*	*	41	10	*8	22	*6
294.8	43	18	25	—	*	*	39	*9	*8	20	*6
295	310	171	140	*	199	96	14	102	51	126	32
295.3	96	63	33	*	59	33	*	35	15	39	*8
295.30	33	21	12	—	21	11	*	*	*7	17	*
295.32	22	13	*8	—	12	*8	*	*9	*	11	*
295.34	40	28	12	*	25	13	*	21	*6	10	*
295.6	13	*7	*6	—	*8	*	*	*	*	*6	*
295.64	*8	*	*	—	*	*	*	*	*	*	*
295.7	161	75	86	*	109	44	*7	49	25	68	19
295.70	114	53	61	*	79	28	*5	25	21	54	15
295.74	42	21	22	*	26	14	*	22	*	12	*
295.9	30	18	11	—	15	13	*	13	*6	*9	*
295.90	19	11	*8	—	11	*7	*	*	*6	*7	*
295.94	*8	*	*	—	*	*	*	*7	*	*	*
296	961	395	565	85	574	227	75	224	184	491	62
296.2	233	97	136	27	140	46	21	38	43	137	16
296.20	111	44	66	13	70	20	*8	16	26	62	*6
296.22	31	12	18	*	21	*	*	*	*	24	*
296.23	49	22	27	*7	26	12	*	12	*7	25	*5
296.24	41	18	23	*	21	*9	*7	*8	*7	24	*
296.3	318	124	194	10	183	92	33	85	79	134	20
296.30	69	27	42	*	40	17	11	18	19	29	*
296.32	15	*6	9	*	*9	*	*	*	*	*7	*
296.33	151	53	98	*5	90	44	12	47	37	57	9
296.34	80	37	43	*	42	26	*9	17	18	40	*
296.4	70	33	38	*	45	19	*	21	11	32	*7
296.40	31	14	16	*	21	*7	*	*5	*	18	*
296.43	*6	*	*	*	*5	*	*	*	*	*	*
296.44	32	15	17	*	18	11	*	12	*5	12	*
296.5	97	31	65	*	61	31	*	26	14	50	*6
296.50	45	19	26	*	31	12	*	10	10	22	*
296.53	30	*7	23	*	17	11	*	13	*	16	*
296.54	20	*	15	*	12	*7	*	*	*	11	*
296.6	90	36	54	15	56	15	*	17	11	57	*
296.60	44	16	28	*	31	*8	*	*6	*7	29	*
296.63	21	10	11	*6	12	*	*	*	*	15	*
296.64	24	9	14	*	12	*5	*	*6	*	13	*
296.7	68	29	38	*6	43	13	*6	20	18	24	*5
296.8	58	30	29	18	29	9	*	*6	*	45	*
296.80	39	17	21	16	16	*6	*	*	*	35	*
296.89	18	11	*7	*	12	*	*	*	*	10	*
296.9	25	15	10	*6	16	*	*	10	*	11	*
296.90	22	13	*8	*6	13	*	*	*6	*	11	*
297	*7	*	*	—	*	*	*	*	*	*	*
298	99	55	45	*	54	21	23	34	12	40	13
298.9	98	54	44	*	53	21	23	33	12	40	13
299	*6	*	*	*	*	*	—	*	*	*	*
300	97	34	63	*7	49	28	13	22	26	33	16
300.0	44	17	28	*	18	14	10	9	16	14	*5
300.00	35	13	22	*	14	12	*9	*7	12	11	*
300.01	*6	*	*	*	*	*	*	*	*	*	*
300.1	9	*	*7	*	*5	*	*	*	*	*	*
300.11	*8	*	*6	*	*	*	*	*	*	*	*
300.4	28	10	19	*	17	*6	*	*9	*5	10	*
300.9	10	*	*6	*	*6	*	*	*	*	*	*
301	*8	*	*6	*	*6	*	*	*	*	*	*
303	145	101	44	*	87	53	*5	30	35	64	16
303.0	37	24	13	*	22	14	*	*	15	10	*8

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
303.00	19	11	*9	*	13	*6	*	*	10	*	*
303.01	17	13	*	*	*9	*8	*	*	*	*6	*
303.9	108	77	31	—	65	39	*	25	20	54	*8
303.90	50	35	15	—	29	19	*	*	*8	36	*
303.91	56	41	15	—	35	20	*	22	11	18	*
304	97	61	35	*	77	18	*	25	9	54	*9
304.0	44	26	17	—	35	*8	*	18	*	18	*
304.00	11	*7	*	—	10	*	*	*	*	*9	*
304.01	32	20	12	—	25	*6	*	18	*	9	*
304.2	14	*9	*5	—	11	*	*	*	*	*8	*
304.20	*8	*	*	—	*6	*	*	*	*	*6	*
304.21	*6	*	*	—	*	*	—	*	*	*	*
304.7	*6	*	*	—	*6	*	—	*	*	*	—
304.71	*5	*	*	—	*	*	—	*	*	*	—
304.8	*7	*5	*	—	*6	*	—	*	*	*5	*
304.9	16	12	*	*	12	*	*	*	*	15	*
304.90	15	11	*	*	11	*	*	*	*	14	*
305	99	73	26	*	57	34	*7	14	21	39	24
305.0	49	39	9	*	22	19	*5	*8	11	14	16
305.00	33	26	*7	*	15	12	*	*5	*9	12	*7
305.01	14	13	*	—	*7	*6	*	*	*	*	*9
305.1	21	12	*9	—	11	*9	*	*	*6	10	*
305.2	*6	*6	*	—	*6	*	—	*	*	*	*
305.6	*8	*7	*	—	*5	*	*	*	*	*	*
305.9	10	*5	*5	*	9	*	*	*	*	*7	*
305.90	9	*	*	*	*9	*	*	*	*	*7	*
307	17	*	14	*	9	*	*	*	*	*7	*
309	88	43	45	10	62	14	*	42	17	18	11
309.0	25	13	11	*	18	*	*	13	*	*	*
309.2	10	*6	*	*	*7	*	*	*	*	*	*
309.28	9	*5	*	*	*7	*	*	*	*	*	*
309.4	20	11	*9	*	15	*	*	13	*	*	*
309.8	24	*6	17	*6	14	*	—	*9	*	*8	*
309.81	23	*6	17	*6	13	*	—	*9	*	*8	*
309.9	*6	*	*	—	*5	*	*	*	*	*	*
310	*6	*	*	*	*	*	*	*	*	*	*
311	143	62	81	10	90	33	10	63	19	51	10
312	22	17	*6	*7	13	*	*	12	*	*5	*
312.3	19	14	*	*	12	*	*	10	*	*	*
312.30	12	*8	*	*	*7	*	*	*6	*	*	*
312.34	*6	*5	*	*	*	*	*	*	*	*	*
313	*6	*	*	*	*	—	—	*	*	*	*
313.8	*6	*	*	*	*	—	—	*	*	*	*
313.81	*5	*	*	*	*	—	—	*	*	*	*
314	13	10	*	*8	*6	—	—	*	*	*8	*
314.0	11	9	*	*7	*	—	—	*	*	*5	*
314.01	10	*9	*	*7	*	—	—	*	*	*5	*
319	*5	*	*	*	*	*	*	*	*	*	*
320-389	518	217	301	81	134	114	189	115	124	194	85
320	*8	*	*	*	*	*	*	*	*	*	*
323	*7	*	*	*	*	*	*	*	*	*	*
323.9	*6	*	*	*	*	*	*	*	*	*	*
324	*7	*	*	*	*	*	*	*	*	*	*
331	75	30	45	*	*	*	65	16	19	29	11
331.0	58	20	38	—	*	*	57	13	15	22	*7
331.3	*6	*	*	*	*	*	*	*	*	*	*
331.4	10	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
332 . . . . .	21	14	*7	—	*	*	19	*	*	*7	*
332.0 . . . . .	21	14	*7	—	*	*	19	*	*	*6	*
333 . . . . .	14	*5	*8	*	*	*	*7	*	*	*	*
333.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
337 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
340 . . . . .	24	*	20	*	*8	14	*	*7	*6	*9	*
343 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
344 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
345 . . . . .	53	26	27	17	17	12	*6	15	13	15	9
345.1 . . . . .	10	*	*5	*	*	*	*	*	*	*	*
345.10 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
345.3 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*
345.4 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
345.5 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
345.9 . . . . .	14	*7	*7	*5	*	*	*	*	*	*	*
345.90 . . . . .	10	*	*	*	*	*	*	*	*	*	*
346 . . . . .	53	16	37	*	29	18	*	9	13	22	*8
346.8 . . . . .	*7	*	*6	*	*	*	*	*	*	*	*
346.80 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
346.9 . . . . .	37	11	26	*	20	14	*	*5	9	18	*
346.90 . . . . .	26	*8	18	*	15	*9	*	*	*7	12	*
346.91 . . . . .	11	*	*8	*	*5	*6	*	*	*	*6	*
348 . . . . .	29	11	18	*	*9	*6	10	*6	*8	9	*5
348.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
348.3 . . . . .	10	*	*6	*	*	*	*6	*	*	*	*
349 . . . . .	11	*	*8	*	*6	*	*	*	*	*	*
349.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
351 . . . . .	10	*	*6	*	*	*	*	*	*	*	*
351.0 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
353 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
355 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
356 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
356.9 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
357 . . . . .	*9	*5	*	*	*5	*	*	*	*	*	*
357.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
358 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
358.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
362 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
376 . . . . .	10	*	*	*	*	*	*	*	*	*	*
376.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
376.01 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
380 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
380.1 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
382 . . . . .	18	11	*8	16	*	*	*	*	*	11	*
382.9 . . . . .	17	9	*7	15	*	*	*	*	*	11	*
386 . . . . .	39	15	24	—	*	*7	29	*9	12	14	*
386.1 . . . . .	18	*6	12	—	*	*	13	*	*	*7	*
386.11 . . . . .	13	*5	*8	—	*	*	10	*	*	*5	*
386.3 . . . . .	18	*8	11	—	*	*	14	*	*6	*6	*
386.30 . . . . .	16	*7	*9	—	*	*	13	*	*6	*6	*
390-459 . . . . .	6,373	3,209	3,164	31	434	1,871	4,037	1,426	1,500	2,394	1,054
394 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
396 . . . . .	19	*7	13	—	*	*	16	*	*	*8	*
396.2 . . . . .	*5	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
396.3	*6	*	*	—	*	*	*	*	*	*	*
398	27	9	18	*	*	*	22	*9	*5	10	*
398.9	27	9	18	*	*	*	22	*9	*5	10	*
398.91	27	9	18	*	*	*	22	*9	*5	10	*
401	308	121	187	*	36	127	144	63	68	135	41
401.0	31	10	21	*	*	14	13	*8	*	16	*
401.9	275	110	165	*	32	112	130	54	64	118	39
402	96	35	62	—	*6	28	62	21	17	38	21
402.0	*5	*	*	—	*	*	*	*	*	*	*
402.9	90	32	58	—	*	26	59	20	16	34	20
402.90	14	*	9	—	*	*6	*7	*	*	*6	*
402.91	76	27	49	—	*	20	52	17	14	28	17
403	102	53	49	*	23	38	39	14	16	53	19
403.9	97	50	47	*	21	36	38	13	15	51	18
403.91	93	49	44	*	19	36	36	12	15	47	18
404	29	15	14	—	*	*7	19	*7	*7	9	*6
404.9	26	13	13	—	*	*6	18	*7	*6	*8	*5
404.93	22	11	10	—	*	*	16	*5	*5	*6	*5
410	818	480	338	*	57	259	501	179	181	297	161
410.0	27	18	*9	—	*	11	15	*	*7	11	*
410.01	26	17	*9	—	*	11	14	*	*7	11	*
410.1	85	54	31	—	9	35	41	17	21	28	19
410.11	84	54	30	—	9	35	40	17	20	27	19
410.2	15	9	*6	—	*	*5	*8	*	*	*5	*
410.21	15	9	*6	—	*	*5	*8	*	*	*5	*
410.3	12	*8	*	—	*	*	*6	*	*	*6	*
410.31	12	*8	*	—	*	*	*6	*	*	*6	*
410.4	118	79	39	—	11	57	50	25	21	49	23
410.41	116	78	38	—	11	56	50	25	21	48	23
410.5	9	*7	*	—	*	*	*5	*	*	*	*
410.51	9	*7	*	—	*	*	*5	*	*	*	*
410.7	392	221	171	—	22	106	264	98	90	137	68
410.71	388	219	170	—	22	105	261	97	89	136	67
410.8	16	*8	*8	—	*	*	11	*	*6	*	*
410.81	15	*8	*7	—	*	*	10	*	*6	*	*
410.9	140	73	67	*	*8	32	99	22	30	56	32
410.91	137	72	65	*	*8	31	97	22	30	54	31
411	124	63	61	—	11	54	59	33	24	49	18
411.1	102	53	49	—	9	47	45	26	20	41	14
411.8	21	10	11	—	*	*7	13	*6	*	*8	*
411.89	20	10	10	—	*	*6	13	*6	*	*7	*
412	*8	*	*5	—	*	*	*	*	*	*	*
413	58	23	35	—	*	23	31	13	13	25	*7
413.9	57	22	35	—	*	22	30	13	13	24	*7
414	1,117	681	436	*	49	437	630	237	287	421	172
414.0	1,096	666	429	*	48	432	615	232	281	412	170
414.00	179	98	80	—	*9	66	104	41	49	77	12
414.01	878	538	339	*	39	352	486	183	221	322	151
414.02	30	23	*7	—	*	10	20	*6	9	10	*
414.8	12	10	*	—	*	*	*8	*	*	*	*
414.9	*8	*	*	—	*	*	*6	*	*	*	*
415	101	36	65	*	19	28	54	22	24	38	17
415.1	99	36	63	*	19	27	53	21	24	37	17
415.19	94	35	60	*	18	26	51	20	22	36	16
416	10	*	*7	*	*	*	*	*	*	*	*
420	15	9	*5	*	*5	*6	*	*	*	*	*
420.9	15	9	*5	*	*5	*6	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
420.90	*9	*5	*	—	*	*	*	*	*	*	*
421	18	10	*7	*	*	*6	*7	*	*	*	*
421.0	17	10	*7	*	*	*6	*7	*	*	*	*
423	13	*7	*6	*	*	*	*	*	*	*5	*
423.9	9	*5	*	*	*	*	*	*	*	*	*
424	98	48	50	*	10	23	64	30	22	32	14
424.0	39	15	25	*	*7	13	19	12	9	13	*5
424.1	56	32	24	*	*	10	43	17	12	19	*9
425	36	22	14	*	*7	11	16	10	*9	12	*
425.4	30	19	11	*	*5	*9	14	*7	*7	12	*
426	70	37	33	*	*7	9	52	18	19	19	13
426.0	30	16	14	*	*	*	25	*6	*8	*9	*7
426.1	20	12	*9	*	*	*	18	*7	*	*6	*
426.13	10	*6	*	*	*	*	*9	*	*	*	*
426.4	*5	*	*	—	*	*	*	*	*	*	*
427	788	377	411	*7	45	174	562	200	191	274	123
427.0	30	11	18	*	*	*8	16	*8	*6	11	*
427.1	57	42	15	*	*	20	34	14	13	23	*7
427.3	465	217	248	*	21	97	345	117	119	159	70
427.31	418	186	232	*	18	84	317	109	110	140	59
427.32	47	31	16	*	*	14	28	*8	9	19	10
427.4	*8	*5	*	—	*	*	*	*	*	*	*
427.41	*8	*5	*	—	*	*	*	*	*	*	*
427.5	14	*6	*9	*	*	*6	*8	*	*	*	*
427.6	*9	*	*	*	*	*	*5	*	*	*	*
427.69	*6	*	*	—	*	*	*	*	*	*	*
427.8	199	89	110	*	14	37	145	52	45	70	32
427.81	83	34	49	—	*	*7	74	27	14	32	11
427.89	116	55	61	*	12	29	72	25	31	38	22
427.9	*5	*	*	—	*	*	*	*	*	*	*
428	973	443	530	*	32	219	720	222	235	373	143
428.0	970	441	529	*	32	219	717	221	235	372	142
429	16	*7	*9	*	*	*	*9	*	*	*7	*
430	24	10	14	*	*	10	9	*	*6	*9	*
431	65	28	37	*	*	22	38	16	12	20	18
432	24	16	*7	*	*	*	18	*	*6	*6	*7
432.1	20	13	*6	*	*	*	16	*	*5	*5	*6
433	172	95	77	*	*	43	126	43	37	69	23
433.1	138	77	61	—	*	34	102	36	28	57	18
433.10	120	70	50	—	*	30	89	30	23	52	15
433.11	18	*7	11	—	*	*	13	*6	*	*5	*
433.3	27	15	12	—	*	*7	20	*7	*7	*9	*
433.30	21	12	9	—	—	*6	15	*6	*6	*8	*
433.31	*6	*	*	—	*	*	*	*	*	*	*
434	283	131	153	*	10	68	204	58	70	101	54
434.0	10	*	*6	*	*	*	*7	*	*	*	*
434.01	*8	*	*	*	*	*	*6	*	*	*	*
434.1	36	17	19	—	*	*6	27	11	*6	*8	11
434.10	*6	*	*	—	*	*	*	*	*	*	*
434.11	30	14	16	—	*	*5	22	*9	*5	*7	*8
434.9	238	110	128	*	*7	59	170	46	60	90	42
434.90	*7	*5	*	—	*	*	*5	*	*	*	*
434.91	230	105	126	*	*7	58	165	44	58	87	40
435	198	79	119	*	*8	49	141	47	44	74	33
435.3	10	*	*8	—	—	*	*9	*	*	*	—
435.8	*7	*	*	—	*	*	*	*	*	*	*
435.9	178	71	107	*	*8	45	125	42	37	68	31

See footnotes at end of table.



**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
436	124	53	71	*	*	18	102	19	27	52	26
437	32	10	22	*	*	*9	19	*6	10	11	*
437.1	*7	*	*6	*	*	*	*7	*	*	*	*
437.3	*7	*	*6	*	*	*	*	*	*	*	*
438	20	11	*9	*	*	*6	12	*	*	*7	*
438.2	*6	*	*	*	*	*	*	*	*	*	*
438.20	*5	*	*	*	*	*	*	*	*	*	*
438.8	10	*5	*	*	*	*	*7	*	*	*	*
438.89	*8	*	*	*	*	*	*6	*	*	*	*
440	111	56	55	*	*	32	76	20	30	43	17
440.1	*8	*	*5	*	*	*	*6	*	*	*	*
440.2	94	49	45	*	*	27	64	18	26	36	15
440.20	*9	*	*	—	*	*	*5	*	*	*	*
440.21	25	15	10	—	*	10	14	*	*8	*7	*5
440.22	10	*5	*	*	*	*	*6	*	*	*	*
440.23	16	*6	9	—	*	*	12	*	*	*5	*
440.24	33	16	17	—	*	*6	27	*7	*8	15	*
440.3	*5	*	*	—	*	*	*	*	*	*	*
441	61	44	17	—	*	13	46	15	13	22	11
441.0	*9	*5	*	—	*	*	*	*	*	*	*
441.01	*5	*	*	—	*	*	*	*	*	*	*
441.2	*5	*	*	—	*	*	*	*	*	*	*
441.3	*8	*6	*	—	*	*	*7	*	*	*	*
441.4	38	29	*8	—	—	*6	32	9	*9	14	*5
442	11	*8	*	*	*	*	*7	*	*	*	*
442.3	*6	*5	*	*	*	*	*	*	*	*	*
443	25	12	13	*	*	*8	15	*5	*7	9	*
443.9	22	11	11	—	*	*7	14	*	*6	*9	*
444	33	15	18	—	*	10	19	*8	*8	10	*7
444.2	26	12	13	—	*	*7	15	*6	*6	*7	*6
444.22	22	11	11	—	*	*7	12	*5	*6	*7	*
446	17	*8	*9	*7	*	*	*7	*	*5	*5	*
446.1	*6	*	*	*6	—	—	*	*	*	*	*
446.5	*6	*	*	—	—	*	*5	*	*	*	*
447	14	*6	*8	*	*	*5	*7	*	*	*	*
447.1	*7	*	*	—	*	*	*	*	*	*	*
451	19	*8	11	*	*	*6	*9	*7	*	*6	*
451.0	*5	*	*	—	*	*	*	*	*	*	*
451.1	*8	*	*	—	*	*	*	*	*	*	*
451.19	*6	*	*	—	*	*	*	*	*	*	*
453	135	57	78	*	27	40	68	26	35	57	17
453.8	127	54	73	*	26	38	62	24	34	53	16
453.9	*5	*	*	—	*	*	*	*	*	*	*
454	*7	*	*	—	*	*	*	*	*	*	*
455	29	12	16	—	*6	10	13	*	*6	11	*7
455.2	16	*6	10	—	*	*5	*8	*	*	*6	*
456	*7	*	*	*	*	*	*	*	*	*	*
458	74	33	41	*	*	19	50	14	20	29	12
458.0	35	17	19	*	*	*5	28	*8	10	13	*5
458.9	31	11	19	*	*	10	19	*	*8	14	*
459	25	14	11	*	*	*7	13	*	*5	11	*
459.0	*5	*	*	—	*	*	*	*	*	*	*
459.8	15	*7	*8	*	*	*	10	*	*	*	*
459.81	14	*7	*8	—	*	*	9	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
460-519 . . . . .	3,542	1,678	1,864	730	382	697	1,732	722	820	1,387	613
461 . . . . .	12	*	9	*	*5	*	*	*	*	*	*
461.9 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
462 . . . . .	12	*7	*5	*7	*	*	*	*	*	*	*
463 . . . . .	15	*6	9	*	12	*	—	*	*	*	*
464 . . . . .	20	14	*6	16	*	*	*	*	*5	*	*6
464.4 . . . . .	14	10	*	13	—	*	—	*	*	*	*
465 . . . . .	38	22	16	24	*	*	*7	*7	*9	13	9
465.9 . . . . .	37	21	16	24	*	*	*7	*6	*9	13	*9
466 . . . . .	279	152	127	209	*8	22	40	44	60	111	64
466.0 . . . . .	76	31	45	9	*8	21	38	15	20	32	*9
466.1 . . . . .	203	121	82	199	*	*	*	29	40	79	56
466.11 . . . . .	121	72	50	120	*	*	*	16	23	46	36
466.19 . . . . .	82	50	32	80	*	*	*	13	17	33	20
473 . . . . .	18	10	*8	*5	*6	*	*	*	*	*9	*
473.9 . . . . .	*8	*	*	*	*	*	*	*	*	*5	*
474 . . . . .	15	*9	*6	12	*	*	—	*	*	*5	*5
474.1 . . . . .	11	*8	*	10	*	*	—	*	*	*	*
474.10 . . . . .	9	*7	*	9	*	—	—	*	*	*	*
475 . . . . .	15	*6	*9	*	10	*	*	*	*	*6	*
478 . . . . .	18	11	*7	*6	*	*	*	*5	*	*6	*
478.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
480 . . . . .	39	23	16	33	*	*	*	*	10	14	12
480.1 . . . . .	21	14	*7	19	*	*	*	*	*6	*8	*5
480.9 . . . . .	17	*9	*9	13	*	*	*	*	*	*5	*7
481 . . . . .	39	20	19	*	*6	*7	23	*8	12	11	*8
482 . . . . .	138	67	70	10	11	22	95	26	33	56	23
482.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
482.1 . . . . .	27	15	12	*	*	*	19	*7	*	13	*
482.2 . . . . .	15	*9	*7	*	*	*	11	*	*	*7	*
482.3 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
482.4 . . . . .	39	17	22	*	*	*7	27	*7	*6	18	*8
482.41 . . . . .	37	16	21	*	*	*6	26	*6	*5	18	*7
482.8 . . . . .	25	13	12	*	*	*	20	*	11	*7	*
482.83 . . . . .	20	10	10	*	*	*	17	*	*8	*6	*
482.9 . . . . .	17	*7	10	*	*	*	11	*5	*	*	*
485 . . . . .	21	11	9	*6	*	*	11	*5	*	10	*
486 . . . . .	1,072	495	577	150	96	182	644	217	247	426	183
487 . . . . .	28	12	15	12	*	*6	*7	*5	10	*7	*
487.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
487.1 . . . . .	19	9	9	*8	*	*	*	*	*6	*5	*
490 . . . . .	27	10	17	*	*	*5	14	*	*5	15	*
491 . . . . .	520	230	291	*	16	161	343	111	123	226	60
491.2 . . . . .	518	229	289	*	16	160	342	111	123	225	59
491.20 . . . . .	21	9	12	—	*	*6	14	*6	*	11	*
491.21 . . . . .	497	220	278	*	15	154	327	105	121	214	56
492 . . . . .	16	*8	*8	—	*	*	10	*	*	*6	*
492.8 . . . . .	13	*5	*8	—	*	*	*9	*	*	*	*
493 . . . . .	484	196	288	187	109	109	80	110	104	181	89
493.0 . . . . .	43	28	15	33	*6	*	*	*6	*	*7	27
493.00 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
493.01 . . . . .	*8	*	*	*6	*	*	—	*	*	*	*6
493.02 . . . . .	30	19	11	22	*	*	*	*	*	*5	18
493.2 . . . . .	97	28	69	*	9	44	43	20	24	36	16
493.20 . . . . .	23	*7	16	*	*	11	11	*	*8	9	*
493.21 . . . . .	*7	*	*6	*	*	*	*	*	*	*	*
493.22 . . . . .	66	19	47	*	*7	30	30	14	16	24	12

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
493.9	343	140	203	153	93	60	37	83	76	137	46
493.90	80	29	52	31	17	16	16	16	17	34	13
493.91	69	30	38	41	19	*6	*	13	17	28	11
493.92	194	81	114	81	56	38	19	54	42	76	22
494	9	*	*7	*	*	*	*6	*	*	*	*
494.1	*5	*	*	—	*	*	*	*	*	*	*
496	101	46	55	—	*6	28	67	20	23	39	19
507	189	101	87	*8	*8	24	149	48	40	69	32
507.0	188	101	87	*8	*8	23	149	47	40	69	31
510	16	11	*5	*	*	*	*7	*	*	*	*
510.9	15	10	*	*	*	*	*7	*	*	*	*
511	56	32	24	*	*6	16	33	13	14	18	11
511.0	*7	*	*	*	*	*	*	*	*	*	*
511.8	*7	*	*	—	*	*	*	*	*	*	*
511.9	41	24	17	*	*	10	26	10	11	14	*7
512	40	24	16	*	15	9	14	*6	11	13	9
512.1	*8	*	*5	*	*	*	*	*	*	*	*
512.8	29	19	10	*	13	*5	9	*	*8	9	*7
513	*7	*	*	*	*	*	*	*	*	*	*
513.0	*7	*	*	*	*	*	*	*	*	*	*
514	*8	*	*	*	*	*	*5	*	*	*	*
515	24	13	11	—	*	*	16	*	*7	*9	*
516	11	*5	*6	*	*	*	*6	*	*	*	*
516.3	*5	*	*	—	*	*	*	*	*	*	*
516.8	*6	*	*	*	*	*	*	*	*	*	*
518	219	102	118	11	22	55	132	45	55	84	35
518.0	*9	*	*6	*	*	*	*	*	*	*	*
518.3	*6	*	*	*	*	*	*	*	*	*	*
518.8	193	92	101	*7	17	49	120	42	47	73	32
518.81	146	70	76	*	11	37	94	34	32	57	23
518.82	13	*7	*6	*	*	*	*	*	*	*	*
518.84	21	*9	12	*	*	*6	14	*	*7	*8	*
518.89	11	*5	*6	*	*	*	*5	*	*	*	*
519	23	14	9	*8	*	*	*8	*	*6	11	*
519.1	*8	*5	*	*	*	*	*	*	*	*	*
519.8	10	*5	*	*	*	*	*	*	*	*	*
520-579	3,320	1,466	1,854	216	839	955	1,310	680	772	1,222	646
522	*9	*	*6	*	*5	*	*	*	*	*	*
522.5	*8	*	*6	*	*5	*	*	*	*	*	*
524	15	*	10	*	13	*	*	*	*	*7	*
524.0	*6	*	*	*	*6	*	—	*	*	*	*
527	*7	*	*	*	*	*	*	*	*	*	*
527.2	*6	*	*	*	*	*	*	*	*	*	*
528	*8	*	*	*	*	*	*	*	*	*	*
530	215	102	113	21	49	68	77	46	49	80	40
530.0	*6	*	*	*	*	*	*	*	*	*	*
530.1	37	19	18	*	12	10	14	*6	9	14	*8
530.10	11	*5	*6	*	*	*	*	*	*	*	*
530.11	10	*6	*	*	*	*	*	*	*	*	*
530.19	13	*5	*8	*	*	*	*5	*	*	*6	*
530.2	17	12	*	*	*	*	12	*	*	*6	*
530.3	9	*5	*	*	*	*	*7	*	*	*	*
530.5	*6	*	*	—	*	*	*	*	*	*	*
530.7	13	*8	*	—	*6	*	*	*	*	*	*
530.8	124	51	73	18	24	48	34	30	29	44	21
530.81	122	49	73	17	24	48	33	30	28	43	21

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
531	104	49	55	*	*9	30	65	17	28	36	22
531.0	14	*8	*6	*	*	*	9	*	*	*	*
531.00	13	*7	*6	*	*	*	*9	*	*	*	*
531.4	63	30	33	*	*	19	40	10	17	23	13
531.40	61	29	32	*	*	18	39	10	17	22	13
531.5	*5	*	*	—	*	*	*	*	*	*	*
531.50	*5	*	*	—	*	*	*	*	*	*	*
531.9	17	*7	10	—	*	*	9	*	*	*6	*5
531.90	16	*6	10	—	*	*	*8	*	*	*	*5
532	61	34	28	*	*8	20	33	12	15	21	14
532.0	11	*7	*	*	*	*	*8	*	*	*	*
532.00	11	*7	*	*	*	*	*8	*	*	*	*
532.4	37	20	17	*	*	12	20	*	10	13	10
532.40	37	20	17	*	*	12	20	*	10	13	10
533	13	*	*9	—	*	*	*7	*	*	*7	*
533.4	*6	*	*	—	*	*	*	*	*	*	*
533.40	*6	*	*	—	*	*	*	*	*	*	*
535	147	52	95	*6	34	45	62	33	34	58	21
535.0	27	11	16	*	*	*8	11	*6	*8	10	*
535.00	21	*7	13	*	*	*6	*7	*	*6	*8	*
535.01	*7	*	*	*	*	*	*	*	*	*	*
535.1	22	*7	15	*	*5	*6	11	*9	*	*8	*
535.10	13	*	*9	*	*	*	*5	*5	*	*	*
535.11	10	*	*7	*	*	*	*6	*	*	*	*
535.3	*6	*	*	—	*	*	*	*	*	*	*
535.4	39	16	23	*	*7	14	17	*8	10	15	*6
535.40	20	*6	14	*	*	*8	*7	*	*5	*9	*
535.41	19	10	9	*	*	*6	10	*	*	*6	*
535.5	49	13	35	*	14	13	20	*8	10	22	*8
535.50	30	*7	24	*	11	9	*9	*5	*7	14	*
535.51	18	*7	11	*	*	*	11	*	*	*8	*
536	47	15	31	*	12	11	19	10	*8	19	10
536.2	11	*	*8	*	*	*	*	*	*	*5	*
536.3	*5	*	*	*	*	*	*	*	*	*	*
536.4	15	*6	10	*	*	*	9	*	*	*5	*
536.41	*7	*	*	*	—	*	*	*	*	*	*
536.8	13	*	*9	*	*	*	*	*	*	*	*
537	29	13	16	*	*	*6	19	*6	*	12	*6
537.0	10	*	*6	*	*	*	*	*	*	*	*
537.8	17	*7	10	*	*	*	14	*	*	*7	*
537.83	15	*5	9	—	*	*	13	*	*	*6	*
540	279	159	120	67	144	47	20	55	60	76	88
540.0	61	37	24	17	21	15	*8	9	15	18	19
540.1	32	21	11	11	10	*6	*6	*	*6	10	11
540.9	186	101	85	39	114	26	*7	41	39	48	57
541	*9	*	*	*	*7	*	*	*	*	*6	*
550	39	32	*8	*7	*6	*	22	10	*8	15	*7
550.1	23	17	*5	*	*	*	14	*6	*	9	*
550.10	19	14	*	*	*	*	11	*	*	*8	*
550.9	16	13	*	*	*	*	*8	*	*	*5	*
550.90	11	*9	*	*	*	*	*6	*	*	*	*
552	60	18	41	—	11	22	27	15	14	20	11
552.1	13	*5	*8	—	*	*5	*	*	*	*	*
552.2	39	10	29	—	*6	15	17	10	*9	13	*7
552.20	*8	*	*6	—	*	*	*	*	*	*	*
552.21	28	*7	21	—	*	11	12	*9	*5	9	*
553	86	36	50	*	18	37	30	18	19	35	15
553.1	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
553.2	60	27	33	*	12	27	21	13	14	23	10
553.20	10	*	*6	—	*	*6	*	*	*	*5	*
553.21	49	22	26	—	10	21	18	11	11	17	9
553.3	20	*5	14	*	*6	*6	*7	*	*	*9	*
555	45	19	26	*	25	11	*6	11	13	16	*5
555.0	14	*7	*6	*	*6	*	*	*	*	*	*
555.1	10	*	*6	*	*7	*	*	*	*	*	*
555.9	18	*7	11	*	9	*5	*	*	*	*8	*
556	35	18	17	*	16	9	*9	*7	*7	14	*8
556.6	*5	*	*	*	*	*	*	*	*	*	*
556.9	16	*8	*8	*	*6	*	*5	*	*	*7	*
557	47	14	32	*	*5	11	30	13	12	13	*8
557.0	26	10	17	*	*	*7	16	*8	*7	*6	*
557.9	17	*	13	—	*	*	12	*	*	*6	*
558	183	77	106	50	43	35	56	33	44	76	30
558.9	177	75	102	49	42	32	53	32	42	74	29
560	292	121	172	12	44	76	162	52	81	105	55
560.1	60	27	33	*	*9	14	36	*9	19	25	*8
560.2	*8	*5	*	*	*	*	*	*	*	*	*
560.3	33	15	17	*	*	*	22	*8	*6	14	*
560.39	31	15	16	*	*	*	20	*8	*6	13	*
560.8	86	31	55	*	15	27	43	12	26	30	18
560.81	64	23	42	*	11	20	31	*8	19	22	14
560.89	22	*9	13	*	*	*6	12	*	*7	*7	*
560.9	101	39	62	*	14	30	54	19	27	32	23
562	262	106	157	*	36	77	149	64	68	93	37
562.1	261	105	156	*	36	77	148	64	68	93	37
562.10	25	*9	16	—	*	*7	15	*7	*5	10	*
562.11	159	65	94	*	31	60	68	44	46	50	20
562.12	65	27	38	—	*	*7	58	11	14	28	13
562.13	11	*	*7	—	*	*	*7	*	*	*5	*
564	73	26	47	9	18	18	27	16	18	24	14
564.0	39	16	22	*8	*6	*8	17	10	*9	12	*9
564.00	30	13	18	*7	*	*5	14	*8	*7	10	*5
564.09	*8	*	*	*	*	*	*	*	*	*	*
564.1	19	*	15	*	*8	*	*6	*	*7	*5	*
564.8	*9	*	*5	*	*	*	*	*	*	*	*
564.89	*9	*	*5	*	*	*	*	*	*	*	*
566	22	12	9	*	*9	*8	*	*	*	*7	*6
567	22	12	10	*	*5	9	*7	*5	*	*8	*
567.2	14	*7	*7	*	*	*7	*	*	*	*5	*
567.8	*5	*	*	—	*	*	*	*	*	*	*
568	19	*	16	*	*7	*6	*6	*	*	*9	*
568.0	11	*	9	*	*	*	*	*	*	*7	*
568.8	*8	*	*6	*	*	*	*	*	*	*	*
568.89	*5	*	*	—	*	*	*	*	*	*	*
569	97	41	56	*	12	28	54	20	21	43	12
569.1	*6	*	*	*	*	*	*	*	*	*	*
569.3	20	11	9	*	*	*	12	*	*	10	*
569.4	13	*6	*7	*	*	*	*8	*	*	*	*
569.49	*6	*	*	*	*	*	*	*	*	*	*
569.6	13	*	9	*	*	*5	*6	*	*	*6	*
569.69	*8	*	*5	*	*	*	*	*	*	*	*
569.8	40	18	21	*	*	12	21	*7	11	18	*
569.81	*6	*	*	—	*	*	*	*	*	*	*
569.83	12	*5	*7	*	*	*5	*	*	*	*6	*
569.85	13	*6	*6	—	*	*	10	*	*	*6	*
570	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
571	104	67	37	*	16	59	27	18	20	44	22
571.1	10	*7	*	—	*	*5	*	*	*	*	*
571.2	45	37	*8	—	*7	32	*6	*8	*9	22	*6
571.5	41	20	21	*	*	19	18	*6	*8	17	10
572	35	20	15	*	*	18	12	*8	*6	12	*9
572.2	25	15	10	*	*	13	*9	*5	*	10	*5
573	11	*	*7	*	*	*	*	*	*	*5	*
573.3	*6	*	*	*	*	*	*	*	*	*	*
574	359	113	246	*	116	109	133	69	82	130	77
574.0	119	48	71	*	35	44	39	22	31	42	24
574.00	113	45	68	*	32	43	37	19	30	41	22
574.01	*6	*	*	—	*	*	*	*	*	*	*
574.1	131	33	98	*	45	39	46	27	28	46	31
574.10	129	33	96	*	45	39	45	26	28	45	30
574.2	33	*6	27	*	*9	*9	15	*7	*9	12	*5
574.20	32	*6	26	*	*8	*9	15	*6	*8	12	*
574.3	*5	*	*	—	*	*	*	*	*	*	*
574.4	*8	*	*	—	*	*	*	*	*	*	*
574.5	25	10	15	—	*7	*	14	*5	*	10	*6
574.50	16	*6	10	—	*	*	9	*	*	*5	*
574.51	9	*	*5	—	*	*	*	*	*	*	*
574.6	*6	*	*	—	*	*	*	*	*	*	*
574.7	16	*	13	*	*7	*	*	*	*	*7	*
574.70	11	*	10	*	*6	*	*	*	*	*6	*
574.8	12	*	*7	—	*	*	*	*	*	*5	*
574.80	*9	*	*5	—	*	*	*	*	*	*	*
575	69	24	45	*	24	23	22	14	15	30	11
575.0	22	10	11	*	*7	*6	*9	*6	*5	*8	*
575.1	36	11	26	*	14	11	11	*5	*7	17	*7
575.10	*6	*	*	—	*	*	*	*	*	*	*
575.11	23	*	18	*	11	*6	*5	*	*	13	*
575.12	*7	*	*	—	*	*	*	*	*	*	*
575.8	*8	*	*6	*	*	*	*	*	*	*	*
576	31	14	17	*	*	11	14	*6	*6	9	9
576.1	10	*6	*	*	*	*	*6	*	*	*	*
576.2	11	*	*7	*	*	*5	*	*	*	*	*
576.8	*8	*	*5	*	*	*	*	*	*	*	*
577	272	141	131	*	94	100	75	47	55	115	54
577.0	233	123	110	*	79	84	67	42	48	102	41
577.1	29	13	15	*	12	11	*5	*	*5	11	9
578	175	84	91	*	18	33	120	48	45	55	27
578.0	22	12	10	*	*	*6	10	*5	*	*6	*7
578.1	39	20	19	*	*6	*6	27	14	10	10	*
578.9	114	53	61	*	*8	22	84	28	31	40	16
579	*8	*	*6	*	*	*	*	*	*	*	*
580-629	1,817	560	1,257	88	580	458	690	347	419	722	330
581	*7	*	*	*	*	*	*	*	*	*	*
581.9	*6	*	*	*	*	*	*	*	*	*	*
584	145	72	73	*	*9	38	96	36	31	47	30
584.5	*8	*	*	*	*	*	*5	*	*	*	*
584.9	133	66	67	*	*8	35	89	34	29	44	27
585	21	15	*6	*	*	*5	11	*6	*	*8	*
590	142	25	117	24	53	30	35	20	27	59	36
590.1	74	12	62	10	31	18	15	10	15	31	18
590.10	73	12	61	10	31	18	15	10	15	31	18
590.8	66	13	53	14	21	11	19	10	12	28	17
590.80	66	13	53	14	21	11	19	10	12	28	17

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
591	11	*5	*6	*	*	*	*	*	*	*	*
592	176	99	77	*	81	62	29	36	48	67	24
592.0	57	34	23	*	25	24	*6	12	16	22	*6
592.1	117	65	53	*	55	37	23	24	32	43	18
593	46	17	28	9	10	*8	19	11	10	15	10
593.4	10	*	*7	*	*	*	*	*	*	*	*
593.7	*8	*	*7	*7	*	—	*	*	*	*	*
593.70	*7	*	*5	*6	*	—	*	*	*	*	*
593.8	*6	*	*	*	*	*	*	*	*	*	*
593.9	15	*6	*8	*	*	*	10	*	*	*6	*
594	*8	*7	*	*	*	*	*	*	*	*	*
594.1	*6	*5	*	*	*	*	*	*	*	*	*
595	18	*6	12	*	*	*	11	*5	*	*7	*
595.0	*6	*	*	*	*	*	*	*	*	*	*
595.9	*6	*	*	—	*	*	*	*	*	*	*
596	20	17	*	*	*	*	14	*	*6	*8	*
596.0	10	10	—	*	*	*	*7	*	*	*	*
598	*6	*	*	*	*	*	*	*	*	*	*
599	404	142	262	32	36	62	274	94	90	158	62
599.0	359	115	244	28	32	52	246	84	81	142	52
599.6	*5	*	*	*	—	*	*	*	*	*	*
599.7	33	22	12	*	*	*7	21	*7	*8	11	*7
600	99	99	—	—	*	20	80	17	25	39	19
600.0	84	84	—	—	*	17	67	16	22	30	16
600.9	*8	*8	—	—	—	*	*7	*	*	*5	*
601	10	10	—	—	*	*	*8	*	*	*6	*
601.0	*7	*7	—	—	*	*	*6	*	*	*	*
604	*7	*7	—	*	*	*	*	*	*	*	*
604.9	*6	*6	—	*	*	*	*	*	*	*	*
607	*6	*6	—	*	*	*	*	*	*	*	*
608	13	13	—	*	*	*5	*	*	*	*6	*
611	20	*	19	*	10	*7	*	*6	*	*7	*
611.0	*7	*	*7	*	*	*	*	*	*	*	*
611.1	9	*	*9	*	*6	*	*	*	*	*	*
614	59	—	59	*	46	11	*	10	12	26	11
614.1	*7	—	*7	—	*	*	*	*	*	*	*
614.2	*9	—	*9	*	*6	*	*	*	*	*	*
614.6	18	—	18	*	12	*	*	*	*	9	*
614.9	17	—	17	*	16	*	—	*	*	*	*
616	16	—	16	*	11	*	*	*	*	*7	*
616.0	9	—	9	—	*8	*	*	*	*	*	*
617	86	—	86	*	68	17	*	14	22	38	13
617.0	50	—	50	—	37	12	*	*9	12	22	*6
617.1	15	—	15	—	12	*	—	*	*	*	*
617.3	13	—	13	*	12	*	*	*	*	*8	*
618	153	—	153	—	27	65	61	26	35	63	28
618.0	36	—	36	—	*5	17	14	*5	*8	16	*7
618.1	13	—	13	—	*7	*	*	*	*	*6	*
618.2	22	—	22	—	*	9	*9	*	*6	*8	*
618.3	18	—	18	—	*	*8	*7	*6	*	*5	*
618.4	20	—	20	—	*	*7	*9	*	*5	10	*
618.5	34	—	34	—	*	15	18	*	*8	14	*8
618.8	*6	—	*6	—	*	*	*	*	*	*	*
619	*8	—	*8	—	*	*	*	*	*	*	*
619.1	*6	—	*6	—	*	*	*	*	*	*	*
620	77	—	77	*	53	15	*	12	18	32	15

See footnotes at end of table.



**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
620.0	*5	—	*5	*	*	*	*	*	*	*	*
620.1	12	—	12	*	10	*	—	*	*	*	*
620.2	52	—	52	*	35	10	*	*7	13	22	10
621	14	—	14	—	*	*8	*	*	*	*	*
621.3	*6	—	*6	—	*	*	*	*	*	*	*
622	13	—	13	—	*7	*6	*	*	*	*5	*
622.1	12	—	12	—	*6	*6	*	*	*	*5	*
623	*6	—	*6	*	*	*	*	*	*	*	*
625	80	—	80	*	47	24	*9	11	17	36	16
625.3	21	—	21	—	18	*	—	*	*	10	*
625.6	34	—	34	—	*8	18	*8	*	*8	15	*7
625.9	20	—	20	—	17	*	*	*	*	9	*
626	110	—	110	*	76	33	*	13	28	45	23
626.2	77	—	77	*	55	22	—	*9	21	31	16
626.8	27	—	27	—	18	9	*	*	*7	11	*5
627	13	—	13	—	*	*8	*	*	*	*	*
627.1	11	—	11	—	*	*7	*	*	*	*	*
630-677	528	—	528	*	524	*	—	91	88	247	102
632	12	—	12	*	12	*	—	*	*	*6	*
633	30	—	30	—	30	*	—	*5	*5	11	*9
633.1	24	—	24	—	24	*	—	*	*	10	*6
634	25	—	25	*	25	—	—	*	*	14	*
634.9	18	—	18	*	18	—	—	*	*	10	*
634.91	15	—	15	—	15	—	—	*	*	*8	*
635	*7	—	*7	*	*7	*	—	*	*	*	*
640	*5	—	*5	—	*5	—	—	*	*	*	*
641	16	—	16	*	16	—	—	*	*	*	*
641.1	*8	—	*8	—	*8	—	—	*	*	*	*
641.13	*8	—	*8	—	*8	—	—	*	*	*	*
641.9	*6	—	*6	—	*6	—	—	*	*	*	*
641.93	*6	—	*6	—	*6	—	—	*	*	*	*
642	28	—	28	*	27	*	—	*	*	16	*
642.3	12	—	12	—	12	*	—	*	*	*8	*
642.33	11	—	11	—	11	*	—	*	*	*7	*
642.4	*6	—	*6	*	*6	—	—	*	*	*	*
643	48	—	48	*	48	—	—	*6	*8	23	10
643.0	19	—	19	—	19	—	—	*	*	10	*
643.03	19	—	19	—	19	—	—	*	*	10	*
643.1	20	—	20	*	20	—	—	*	*	*8	*
643.13	20	—	20	*	20	—	—	*	*	*8	*
643.2	*8	—	*8	—	*8	—	—	*	*	*5	*
643.23	*8	—	*8	—	*8	—	—	*	*	*5	*
644	96	—	96	*	96	*	—	20	16	45	15
644.0	86	—	86	*	86	*	—	19	15	39	13
644.03	86	—	86	*	86	*	—	19	15	39	13
644.1	10	—	10	—	10	—	—	*	*	*6	*
644.13	10	—	10	—	10	—	—	*	*	*6	*
646	65	—	65	*	65	—	—	*7	12	32	15
646.2	*6	—	*6	—	*6	—	—	*	*	*	*
646.23	*6	—	*6	—	*6	—	—	*	*	*	*
646.6	38	—	38	*	38	—	—	*	*7	19	*8
646.63	36	—	36	*	36	—	—	*	*6	18	*8
646.8	20	—	20	—	20	—	—	*	*	10	*6
646.83	18	—	18	—	18	—	—	*	*	10	*
647	10	—	10	—	10	—	—	*	*	*6	*
647.8	*6	—	*6	—	*6	—	—	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
647.83	*6	—	*6	—	*6	—	—	*	*	*	*
648	101	—	101	*	100	*	—	20	17	45	18
648.0	10	—	10	—	10	—	—	*	*	*7	*
648.03	10	—	10	—	10	—	—	*	*	*7	*
648.4	11	—	11	—	11	—	—	*	*	*	*
648.43	*8	—	*8	—	*8	—	—	*	*	*	*
648.9	65	—	65	*	65	*	—	13	11	30	10
648.93	63	—	63	*	63	*	—	13	11	30	10
654	13	—	13	—	13	—	—	*	*	*6	*
654.5	*8	—	*8	—	*8	—	—	*	*	*	*
654.53	*8	—	*8	—	*8	—	—	*	*	*	*
658	*6	—	*6	—	*6	—	—	*	*	*	*
659	*6	—	*6	—	*6	*	—	*	*	*	*
666	*5	—	*5	—	*5	*	—	*	*	*	*
670	*6	—	*6	*	*6	—	—	*	*	*	*
670.0	*6	—	*6	*	*6	—	—	*	*	*	*
670.04	*6	—	*6	*	*6	—	—	*	*	*	*
674	13	—	13	—	13	—	—	*	*	*7	*
674.3	*6	—	*6	—	*6	—	—	*	*	*	*
674.34	*6	—	*6	—	*6	—	—	*	*	*	*
680-709	601	299	303	93	158	151	199	137	127	249	88
681	18	11	*7	*	*6	*	*6	*	*6	*6	*
681.0	9	*6	*	*	*	*	*	*	*	*	*
681.00	*8	*	*	*	*	*	*	*	*	*	*
681.1	*9	*	*	*	*	*	*	*	*	*	*
681.10	*9	*	*	*	*	*	*	*	*	*	*
682	404	200	204	38	103	123	140	105	89	145	65
682.0	37	16	21	12	11	*5	9	10	*7	15	*6
682.1	*8	*	*	*	*	*	*	*	*	*	*
682.2	31	14	17	*	*9	12	*8	*	*5	16	*6
682.3	43	24	19	*	17	13	*8	11	10	16	*6
682.4	25	15	10	*	*8	*8	*8	*6	*6	*8	*6
682.5	9	*	*6	*	*	*	*	*	*	*	*
682.6	208	102	106	*6	47	68	87	61	48	68	31
682.7	38	19	19	*5	*7	12	15	10	*8	13	*7
683	*8	*	*5	*7	*	*	*	*	*	*	*
693	*6	*	*	*	*	*	*	*	*	*	*
693.0	*6	*	*	*	*	*	*	*	*	*	*
695	*7	*	*	*	*	*	*	*	*	*	*
695.1	*6	*	*	*	*	*	*	*	*	*	*
701	10	*6	*	*	*	*	*	*	*	*9	*
701.4	*8	*6	*	*	*	*	—	*	—	*8	—
707	76	34	42	*	16	14	45	17	22	26	11
707.0	38	17	20	*	10	*6	22	*8	9	14	*6
707.1	36	15	21	*	*5	*8	22	*9	12	11	*
707.13	*6	*	*	*	*	*	*	*	*	*	*
707.15	10	*	*5	*	*	*	*5	*	*	*	*
707.19	*6	*	*	*	*	*	*	*	*	*	*
709	44	26	18	30	13	*	*	*	*	43	*
709.2	43	25	17	29	13	*	*	*	*	42	*
710-739	1,736	747	989	41	326	598	770	368	432	600	336
710	25	*	21	*	14	*7	*	*	*7	11	*
710.0	20	*	17	*	12	*5	*	*	*7	*8	*
711	18	12	*6	*	*	*	*7	*	*	*6	*
711.0	17	11	*6	*	*	*	*6	*	*	*5	*
711.06	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
714	24	*6	18	*	*	9	10	*5	*8	*8	*
714.0	19	*	15	—	*	*8	*8	*	*7	*6	*
715	568	221	347	—	20	193	356	132	150	186	101
715.1	44	18	27	—	*	18	25	9	*7	17	11
715.15	13	*7	*7	—	*	*	*8	*	*	*	*
715.16	29	11	18	—	*	13	16	*	*	13	*7
715.3	292	117	175	—	9	95	188	68	78	97	49
715.35	81	34	47	—	*	26	52	26	23	19	13
715.36	202	80	122	—	*6	64	131	41	54	73	33
715.9	222	82	140	—	9	76	138	52	60	70	41
715.90	*9	*	*6	—	*	*	*	*	*	*	*
715.91	*7	*	*	—	*	*	*	*	*	*	*
715.95	69	27	41	—	*	23	42	18	17	19	15
715.96	134	49	85	—	*	45	86	29	38	43	23
716	26	11	15	*	*	10	10	*	*7	9	*5
716.1	*5	*	*	—	*	*	*	*	*	*	*
716.9	17	*7	10	*	*	*5	*8	*	*	*7	*
716.96	*7	*	*	*	*	*	*	*	*	*	*
717	9	*	*	*	*7	*	*	*	*	*	*
717.8	*6	*	*	*	*	*	—	*	*	*	*
717.83	*6	*	*	*	*	*	—	*	*	*	*
718	17	*8	*9	*	*6	*	*	*	*	*	*5
718.3	*6	*	*	*	*	*	*	*	*	*	*
719	42	17	25	*	11	*9	21	*9	13	12	*8
719.4	33	13	20	*	9	*7	16	*	11	10	*6
719.41	*8	*	*	*	*	*	*	*	*	*	*
719.45	15	*6	*9	*	*	*	10	*	*	*6	*
721	89	41	48	—	11	35	44	14	21	35	19
721.0	25	13	12	—	*	13	*8	*	*5	10	*7
721.1	19	10	9	—	*	*7	10	*	*	9	*
721.3	38	15	22	—	*	12	22	*5	9	13	9
722	353	180	174	*	133	153	67	64	79	137	74
722.0	55	24	31	—	24	25	*6	*8	10	28	*8
722.1	186	106	80	*	81	74	31	38	41	66	41
722.10	185	105	79	*	80	73	31	38	41	64	41
722.4	12	*6	*6	—	*	*7	*	*	*	*5	*
722.5	53	23	30	—	17	22	14	*8	14	17	14
722.52	53	23	30	—	17	22	14	*8	14	17	14
722.7	22	9	12	—	*	12	*	*	*	10	*
722.71	15	*7	*8	—	*	10	*	*	*	*7	*
722.73	*6	*	*	—	*	*	*	*	*	*	*
722.8	*9	*	*6	—	*	*	*	*	*	*6	*
722.83	*8	*	*6	—	*	*	*	*	*	*6	*
722.9	14	*7	*7	*	*	*7	*	*	*	*	*
722.93	9	*	*	*	*	*	*	*	*	*	*
723	27	16	12	*	*6	14	*6	*	*6	14	*
723.0	9	*7	*	—	*	*6	*	*	*	*	*
723.1	*8	*	*	*	*	*	*	*	*	*	*
723.4	*9	*	*	—	*	*6	*	*	*	*6	*
724	133	60	73	*	21	40	71	33	37	43	19
724.0	74	38	36	—	*	26	46	17	19	25	12
724.02	73	37	36	—	*	26	45	17	19	25	12
724.2	28	10	18	*	11	*7	10	*8	10	*6	*
724.4	9	*	*	—	*	*	*	*	*	*6	*
724.5	15	*	10	*	*	*	*9	*	*	*	*
726	39	21	18	*	*9	17	12	*9	*9	*9	12
726.1	16	*9	*7	—	*	9	*5	*	*	*	*6
726.10	15	*9	*6	—	*	*9	*5	*	*	*	*5

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
726.2	*5	*	*	—	*	*	*	*	*	*	*
726.3	*6	*5	*	—	*	*	*	*	*	*	*
726.33	*6	*5	*	—	*	*	*	*	*	*	*
726.6	*5	*	*	*	*	*	*	*	*	*	*
727	24	10	14	*	*	*7	9	*6	*	9	*5
727.0	*8	*	*	*	*	*	*	*	*	*	*
728	31	17	14	*	*8	*5	16	*	9	12	*
728.8	25	13	12	*	*7	*	14	*	*8	*8	*
728.89	19	11	*8	*	*	*	13	*	*7	*7	*
729	57	23	35	*	17	21	17	11	12	23	11
729.1	15	*6	9	*	*	*8	*	*	*	*8	*
729.5	22	9	12	*	*7	*6	*7	*	*6	*7	*
729.8	13	*6	*8	*	*	*	*6	*	*	*6	*
729.89	10	*	*5	*	*	*	*5	*	*	*	*
730	45	29	16	*	9	15	17	*8	12	16	*9
730.0	10	*8	*	*	*	*	*5	*	*	*	*
730.07	*5	*	*	*	*	*	*	*	*	*	*
730.1	*8	*	*	*	*	*	*	*	*	*	*
730.2	26	16	10	*	*5	10	*9	*	*6	10	*6
730.27	10	*7	*	*	*	*	*5	*	*	*	*
732	*5	*	*	*	*	*	—	*	*	*	*
733	146	43	103	*	22	37	82	33	38	47	28
733.1	62	14	48	*	*	*8	53	16	20	15	12
733.13	44	9	35	*	*	*	40	11	13	11	*9
733.14	*8	*	*6	—	*	*	*	*	*	*	*
733.4	22	*8	14	*	*	10	*8	*	*5	10	*
733.42	18	*8	10	*	*	*9	*5	*	*	*8	*
733.6	20	*	16	*	*7	10	*	*	*	*8	*
733.8	29	13	16	*	*9	*6	13	*6	*6	11	*6
733.81	*7	*	*	*	*	*	*	*	*	*	*
733.82	22	11	12	*	*8	*5	*9	*	*6	9	*
733.9	*6	*	*	*	*	*	*	*	*	*	*
736	*6	*	*	*	*	*	*	*	*	*	*
737	16	*	11	*6	*6	*	*	*5	*	*	*
737.3	14	*	10	*6	*5	*	*	*	*	*	*
737.30	11	*	*8	*	*	*	*	*	*	*	*
738	28	11	17	*	*7	11	9	*7	*6	*7	*9
738.4	22	*9	14	—	*	*9	9	*	*5	*	*8
740-759	178	97	80	124	31	15	*7	54	26	55	43
742	*9	*5	*	*7	*	*	*	*	*	*	*
745	23	10	13	19	*	*	*	10	*	*6	*
745.4	*8	*	*	*7	*	*	*	*	*	*	*
745.5	*6	*	*	*	*	*	*	*	*	*	*
746	12	*6	*5	*6	*	*	*	*	*	*	*
747	16	*9	*7	*9	*	*	*	*6	*	*	*
748	*7	*	*	*7	*	—	—	*	*	*	*
749	13	*8	*6	13	*	*	—	*	*	*7	*
749.0	*7	*	*	*7	*	*	—	*	*	*5	*
749.00	*5	*	*	*	*	—	—	*	*	*	*
749.2	*5	*	*	*5	*	—	—	*	*	*	*
750	17	14	*	16	*	—	*	*	*	*	*7
750.5	13	11	*	13	*	—	—	*	*	*	*5
751	13	*7	*6	9	*	*	*	*	*	*5	*
753	*9	*6	*	*	*	*	*	*	*	*	*
754	11	*6	*5	*8	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
755	*5	*	*	*	*	*	*	*	*	*	*
756	19	*8	10	9	*	*	*	*	*5	*5	*
756.1	10	*	*6	*	*	*	*	*	*	*	*
756.12	*8	*	*	*	*	*	*	*	*	*	*
757	*7	*	*	*	*	—	—	*	*	*7	*
757.3	*7	*	*	*	*	—	—	*	*	*6	*
757.32	*6	*	*	*	*	—	—	—	—	*6	—
759	*5	*	*	*	*	*	*	*	*	*	*
760-779	166	93	73	165	*	—	*	27	35	49	55
765	24	14	10	24	—	—	—	*	*	13	*
765.1	19	11	*9	19	—	—	—	*	*	10	*
765.18	*6	*	*	*6	—	—	—	*	*	*	*
769	17	*8	*9	17	—	—	*	*	*	*	*8
770	32	19	14	32	—	—	*	*8	*7	*7	10
770.6	*8	*	*	*8	—	—	—	*	*	*	*
770.8	16	*9	*7	16	—	—	*	*	*	*	*6
771	14	*8	*6	14	—	—	—	*	*	*	*6
771.8	12	*7	*	12	—	—	—	*	*	*	*
774	38	22	17	38	—	—	—	*5	*8	12	14
774.6	32	18	15	32	—	—	—	*	*7	*9	13
777	*5	*	*	*5	—	—	—	*	*	*	*
778	*8	*	*	*8	—	—	—	*	*	*	*
778.4	*7	*	*	*7	—	—	—	*	*	*	*
779	10	*6	*	10	—	—	—	*	*	*	*
780-799	283	132	151	63	99	68	54	61	56	102	65
780	62	34	28	30	16	*6	10	16	11	23	12
780.2	10	*	*5	*	*	*	*	*	*	*	*
780.3	28	16	12	15	9	*	*	*6	*6	9	*7
780.39	26	15	11	13	9	*	*	*6	*5	*9	*7
780.6	16	*9	*7	13	*	*	*	*	*	*7	*
784	*7	*	*5	*	*	*	*	*	*	*	*
786	77	39	38	*5	26	33	13	16	18	30	14
786.0	*8	*	*5	*	*	*	*	*	*	*	*
786.09	*5	*	*	*	*	—	*	*	*	*	*
786.5	67	36	32	*	24	33	11	14	14	27	12
786.50	31	17	15	*	11	14	*6	*	*8	15	*
786.59	31	17	15	*	11	17	*	*9	*6	*9	*8
787	*6	*	*	*	*	*	*	*	*	*	*
788	*6	*	*	*	*	*	*	*	*	*5	*
789	41	19	22	9	24	*6	*	*7	11	16	*7
789.0	41	19	22	9	24	*6	*	*7	11	16	*7
789.00	12	*6	*7	*	*9	*	*	*	*	*7	*
789.03	9	*	*5	*	*6	*	*	*	*	*	*
789.09	*6	*	*	*	*	*	*	*	*	*	*
799	66	27	39	9	18	15	24	15	12	14	26
799.9	66	27	39	9	17	15	24	15	12	13	26
800-999	2,697	1,331	1,366	233	780	621	1,063	540	626	976	556
800	14	9	*	*8	*	*	*	*	*	*	*
801	30	22	*7	*	14	*7	*	*7	*7	*8	*7
801.0	11	*8	*	*	*	*	*	*	*	*	*
801.1	*6	*	*	*	*	*	*	*	*	*	*
801.2	*7	*6	*	*	*	*	*	*	*	*	*
802	32	24	*7	*	20	*6	*	*7	*6	13	*6
802.2	11	*9	*	*	*7	*	*	*	*	*6	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
802.4	*6	*	*	*	*	*	*	*	*	*	*
802.8	*6	*	*	*	*	*	*	*	*	*	*
805	75	33	42	*	20	17	38	14	23	27	12
805.0	16	11	*5	*	*	*6	*6	*	*	*6	*
805.2	18	*6	12	*	*5	*	*9	*	*7	*5	*
805.4	38	15	23	*	10	*7	21	*5	11	14	*7
806	9	*8	*	*	*6	*	*	*	*	*	*
807	37	20	17	*	*7	10	19	*6	10	10	10
807.0	32	18	15	*	*6	*8	18	*6	9	*8	*9
807.01	*6	*	*	—	*	*	*	*	*	*	*
807.03	*7	*	*	*	*	*	*	*	*	*	*
808	64	19	45	*	14	*5	43	10	21	23	11
808.0	10	*7	*	*	*6	*	*	*	*	*5	*
808.2	44	9	35	*	*5	*	34	*7	15	14	*8
808.8	*6	*	*	—	*	—	*	*	*	*	*
812	64	22	42	15	*7	*8	33	17	14	15	17
812.0	31	*8	24	*	*	*	25	*7	*8	*9	*8
812.00	9	*	*6	*	*	*	*8	*	*	*	*
812.01	*9	*	*8	*	*	*	*7	*	*	*	*
812.09	*8	*	*6	—	*	*	*6	*	*	*	*
812.2	*9	*	*6	*	*	*	*	*	*	*	*
812.21	*7	*	*5	*	*	*	*	*	*	*	*
812.4	21	9	12	14	*	*	*	*7	*	*	*7
812.41	14	*6	*8	11	*	*	*	*	*	*	*5
813	52	28	24	*9	17	10	17	10	10	17	15
813.0	*9	*	*	*	*	*	*	*	*	*	*
813.2	*6	*	*	*	*	*	*	*	*	*	*
813.4	24	11	13	*	*7	*	10	*	*6	*7	*8
813.41	*6	*	*	*	*	*	*	*	*	*	*
813.42	11	*	*7	*	*	*	*6	*	*	*	*
813.44	*7	*	*	*	*	*	*	*	*	*	*
815	*5	*	*	*	*	*	*	*	*	*	*
816	*7	*5	*	*	*	*	*	*	*	*	*
816.1	*5	*	*	*	*	*	*	*	*	*	*
820	315	85	230	*	*6	24	282	70	81	108	57
820.0	93	23	71	*	*	*6	85	22	25	31	16
820.00	*6	*	*5	—	—	*	*6	*	*	*	*
820.02	*8	*	*	*	—	*	*8	*	*	*	*
820.03	*6	*	*6	*	*	*	*	*	*	*	*
820.09	71	16	55	*	*	*	66	17	21	23	11
820.2	149	38	111	*	*	10	136	36	40	50	23
820.20	11	*	10	*	—	*	10	*	*	*	*
820.21	128	32	96	*	*	*8	118	32	35	43	18
820.22	10	*	*6	*	*	*	*8	*	*	*	*
820.8	70	22	48	*	*	*7	61	12	14	26	17
821	65	31	35	12	19	12	23	13	17	22	13
821.0	37	20	17	9	13	*	10	*7	10	14	*6
821.00	*6	*	*	*	*	*	*	*	*	*	*
821.01	30	17	13	*8	12	*	*7	*6	9	10	*5
821.2	23	*7	16	*	*	*6	13	*	*7	*6	*5
821.20	*7	*	*	*	*	*	*	*	*	*	*
821.23	10	*	*8	*	*	*	*6	*	*	*	*
822	21	10	11	*	*5	*6	9	*	*	*7	*6
822.0	20	9	11	*	*	*6	9	*	*	*6	*6
823	63	41	22	*6	31	14	12	15	17	19	12
823.0	24	12	12	*	9	*7	*8	*6	*6	*7	*
823.00	20	10	10	*	*8	*	*6	*	*5	*6	*
823.2	16	13	*	*	9	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
823.22	10	*8	*	*	*5	*	*	*	*	*	*
823.3	*8	*7	*	*	*	*	*	*	*	*	*
823.32	*7	*6	*	*	*	*	*	*	*	*	*
823.8	11	*6	*	*	*5	*	*	*	*	*	*
823.82	*7	*	*	*	*	*	*	*	*	*	*
824	103	48	55	*	43	29	27	19	25	36	23
824.0	*6	*	*	*	*	*	*	*	*	*	*
824.2	*5	*	*	*	*	*	*	*	*	*	*
824.4	26	11	15	*	*9	*8	*9	*6	*7	*7	*6
824.6	25	10	16	*	10	*7	*8	*	*6	10	*7
824.8	28	16	12	*	14	*6	*	*6	*6	9	*6
824.9	*6	*	*	*	*	*	*	*	*	*	*
825	23	14	9	*	14	*6	*	*	*5	*8	*
825.0	10	*8	*	*	*5	*	*	*	*	*	*
825.2	9	*	*6	*	*6	*	*	*	*	*	*
825.25	*7	*	*	*	*	*	*	*	*	*	*
831	*5	*	*	—	*	*	*	*	*	*	*
831.0	*5	*	*	—	*	*	*	*	*	*	*
840	17	*8	9	*	*	*9	*6	*	*	*6	*
840.4	14	*6	*8	—	*	*7	*6	*	*	*	*
844	12	*7	*	—	*5	*	*	*	*	*	*
844.8	*7	*5	*	—	*	*	*	*	*	*	*
847	10	*6	*	*	*	*	*	*	*	*	*
850	42	26	16	*9	16	*7	11	10	*8	11	14
850.0	13	*6	*7	*	*	*	*	*	*	*	*
850.1	16	10	*6	*	*7	*	*5	*	*	*	*5
850.5	*8	*6	*	*	*	*	*	*	*	*	*
850.9	*6	*	*	*	*	*	*	*	*	*	*
851	19	9	10	*	*8	*	*7	*	*	*6	*6
851.8	14	*6	*8	*	*	*	*6	*	*	*	*
852	44	25	19	*	12	*7	22	*7	9	13	15
852.0	11	*7	*	*	*5	*	*	*	*	*	*
852.2	27	14	13	*	*	*5	17	*	*6	*8	10
852.20	*6	*	*	*	*	*	*	*	*	*	*
852.21	12	*5	*7	*	*	*	*8	*	*	*	*
853	*7	*	*	*	*	*	*	*	*	*	*
853.0	*7	*	*	*	*	*	*	*	*	*	*
854	*6	*	*	*	*	*	*	*	*	*	*
854.0	*6	*	*	*	*	*	*	*	*	*	*
860	31	23	*8	*	12	12	*7	*5	*6	11	*9
860.0	17	11	*6	*	*8	*	*	*	*	*6	*6
860.4	*8	*6	*	*	*	*	*	*	*	*	*
861	10	*8	*	*	*6	*	*	*	*	*	*
861.2	*7	*6	*	*	*	*	*	*	*	*	*
861.21	*7	*6	*	*	*	*	*	*	*	*	*
863	*8	*8	*	*	*6	*	*	*	*	*	*
864	10	*7	*	*	*9	*	*	*	*	*	*
864.0	*7	*	*	*	*5	*	*	*	*	*	*
865	14	*8	*5	*	9	*	*	*	*	*6	*
865.0	13	*8	*5	*	*9	*	*	*	*	*6	*
871	*7	*7	*	*	*5	*	*	*	*	*	*
873	23	14	9	*5	*8	*	*7	*	*5	*8	*6
873.0	*5	*	*	*	*	*	*	*	*	*	*
873.4	*9	*6	*	*	*	*	*	*	*	*	*
881	*7	*	*	*	*	*	*	*	*	*	*
882	*7	*5	*	*	*	*	*	*	*	*	*

See footnotes at end of table.



**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
883	*6	*	*	*	*	*	*	*	*	*	*
886	*7	*	*	*	*	*	*	*	*	*	*
886.0	*5	*	*	*	*	*	*	*	*	*	*
891	*8	*	*	*	*	*	*	*	*	*	*
892	*6	*	*	*	*	*	*	*	*	*	*
920	*7	*	*	*	*	*	*	*	*	*	*
922	15	*8	*7	*	*	*	*7	*	*	*	*
922.1	*5	*	*	*	*	*	*	*	*	*	*
922.2	*5	*	*	*	*	*	*	*	*	*	*
924	23	*5	18	*	*	*	18	*7	*7	*7	*
924.0	13	*	10	*	*	*	11	*6	*	*	*
924.01	12	*	*9	*	*	*	11	*5	*	*	*
935	*8	*6	*	*	*	*	*	*	*	*	*
935.1	*8	*5	*	*	*	*	*	*	*	*	*
941	10	*7	*	*	*	*	*	*	*	*5	*
941.2	*8	*6	*	*	*	*	*	*	*	*	*
942	11	*8	*	*5	*	*	*	*	*	*8	*
942.3	*6	*	*	*	*	*	*	*	*	*	—
943	*7	*6	*	*	*	*	*	*	*	*	*
944	*7	*	*	*	*	*	*	*	*	*5	*
945	16	10	*6	*5	*	*	*	*	*	10	*
945.2	*7	*	*	*	*	*	*	*	*	*	*
945.3	9	*5	*	*	*	*	*	*	*	*6	*
946	*6	*6	*	*	*	*	*	*	—	*6	—
949	*8	*	*	*7	*	—	—	—	—	*8	—
949.0	*8	*	*	*7	*	—	—	—	—	*8	—
958	*5	*	*	*	*	*	*	*	*	*	*
959	38	22	16	*8	15	*6	*9	*6	10	14	*8
959.0	28	17	10	*7	10	*	*6	*	*7	10	*6
959.01	26	16	10	*7	9	*	*6	*	*7	9	*6
962	*6	*	*	*	*	*	*	*	*	*	*
962.3	*5	*	*	*	*	*	*	*	*	*	*
965	50	18	32	*	32	9	*	10	13	19	*8
965.0	17	*9	*8	*	10	*	*	*	*	*7	*
965.09	*8	*	*	*	*	*	*	*	*	*	*
965.1	*5	*	*	*	*	*	*	*	*	*	*
965.4	21	*5	16	*	15	*	*	*	*	*9	*
966	10	*	*7	*	*7	*	*	*	*	*	*
966.3	*7	*	*5	*	*6	*	*	*	*	*	*
967	*8	*	*	*	*5	*	*	*	*	*	*
968	10	*7	*	—	*7	*	*	*	*	*	*
968.5	*7	*6	*	—	*	*	*	*	*	*	*
969	67	30	37	*	40	18	*5	10	14	27	16
969.0	21	9	12	*	15	*	*	*	*	9	*
969.4	28	12	17	*	13	11	*	*	*6	11	*7
969.7	*6	*	*	*	*	*	—	*	*	*	*
972	12	*5	*7	*	*	*	*7	*	*	*	*
972.6	*5	*	*	*	*	*	*	*	*	*	*
977	9	*	*8	*	*5	*	*	*	*	*	*
977.9	*7	*	*6	*	*	*	*	*	*	*	*
980	*8	*6	*	*	*5	*	*	*	*	*	*
980.0	*6	*	*	*	*	*	*	*	*	*	*
989	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
992	*7	*	*	*	*	*	*	*	*	*	*
994	*8	*6	*	*	*	*	—	*	*	*	*
995	36	12	23	*	*7	12	12	*7	*7	16	*5
995.1	13	*	*9	*	*	*	*6	*	*	*5	*
995.2	*9	*	*7	*	*	*	*	*	*	*	*
996	473	242	231	31	88	149	206	95	103	184	92
996.0	19	10	*9	*	*	*	13	*6	*	*6	*
996.01	10	*	*6	*	*	*	*8	*	*	*	*
996.04	*5	*	*	—	*	*	*	*	*	*	*
996.1	15	*	10	*	*	*5	*6	*	*	*8	*
996.2	21	12	9	*9	9	*	*	*5	*	*6	*5
996.4	95	38	57	*	11	22	61	20	22	34	19
996.5	13	*5	*8	*	*	*6	*	*	*	*6	*
996.59	*7	*	*	*	*	*	*	*	*	*	*
996.6	147	82	65	11	27	45	63	29	39	56	22
996.61	10	*8	*	—	*	*	*7	*	*	*	*
996.62	77	41	36	*9	16	26	25	15	16	33	12
996.64	9	*7	*	—	*	*	*7	*	*	*	*
996.66	21	11	10	*	*	*	15	*	*9	*7	*
996.67	11	*7	*	*	*	*	*	*	*	*	*
996.69	*8	*	*	*	*	*	*	*	*	*	*
996.7	137	74	63	*5	23	53	56	25	25	53	34
996.72	27	17	10	*	*	12	12	*6	*6	10	*
996.73	47	27	20	*	9	20	17	*7	*5	20	14
996.74	25	14	11	*	*	*9	12	*	*6	10	*6
996.77	11	*	*7	—	*	*	*	*	*	*	*
996.78	14	*6	*8	*	*	*	*5	*	*	*	*
996.8	24	15	*8	*	10	11	*	*	*	14	*
996.81	13	9	*	*	*8	*	*	*	*	*7	*
997	115	51	64	*6	26	40	44	26	27	36	27
997.1	13	9	*	*	*	*	*8	*	*	*	*
997.2	12	*	*9	—	*	*5	*5	*	*	*	*
997.3	10	*5	*	*	*	*	*	*	*	*	*
997.4	44	15	29	*	13	16	13	9	11	12	11
997.5	10	*	*5	—	*	*	*	*	*	*	*
997.6	18	10	*7	*	*	*7	*7	*	*	*8	*
997.62	10	*6	*	*	*	*	*	*	*	*	*
997.69	*8	*	*	*	*	*	*	*	*	*	*
998	233	107	126	10	58	81	84	53	54	80	46
998.1	54	30	24	*	12	14	24	13	14	16	11
998.11	34	23	11	*	*7	*8	15	*8	*9	9	*7
998.12	18	*7	11	*	*	*6	*8	*	*	*	*
998.2	12	*	*7	*	*	*	*	*	*	*	*
998.3	17	*8	*9	*	*	*6	*7	*	*	*7	*
998.5	122	52	70	*	33	44	41	28	27	42	25
998.59	119	51	68	*	32	42	40	27	26	41	24
998.6	*5	*	*	*	*	*	*	*	*	*	*
998.8	22	10	12	*	*6	*9	*5	*6	*	*7	*
998.83	*7	*	*	—	*	*	*	*	*	*	*
998.89	15	*7	*8	*	*	*6	*	*	*	*	*
999	*8	*	*	*	*	*	*	*	*	*	*
V01-V82	4,880	403	4,477	75	4,073	224	508	750	984	1,972	1,175
V10	*5	*	*	*	*	*	*	*	*	*	*
V12	16	*6	11	*	*	*	11	*	*	10	*
V12.5	14	*	10	*	*	*	10	*	*	9	*
V12.59	9	*	*7	*	*	*	*7	*	*	*7	*
V15	12	*7	*	*	*	*5	*	*	*	*	*
V15.8	9	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
V15.81	*6	*	*	*	*	*	*	*	*	*	*
V17	*6	*	*	*	*	*	*	*	*	*	*
V22	*7	-	*7	-	*7	-	-	*	*	*	*
V27	3,951	-	3,951	12	3,934	*	-	582	736	1,660	974
V27.0	3,739	-	3,739	12	3,723	*	-	534	673	1,591	941
V27.1	21	-	21	-	21	*	-	*	*	11	*
V27.2	49	-	49	*	49	*	-	*9	*8	21	11
V27.9	137	-	137	*	137	*	-	36	52	33	16
V30	10	*	*6	10	-	-	-	*	*	*	*
V30.1	10	*	*6	10	-	-	-	*	*	*	*
V43	*5	*	*	-	*	*	*	*	*	*	*
V45	20	*9	11	*	*	10	*	*	*7	*6	*
V45.8	12	*6	*6	*	*	*7	*	*	*	*	*
V45.89	*6	*	*	*	*	*	*	*	*	*	*
V53	11	*7	*	*	*	*	*8	*	*	*	*
V53.3	11	*7	*	*	*	*	*8	*	*	*	*
V53.31	*7	*	*	*	*	*	*6	*	*	*	*
V54	9	*	*8	*	*	*	*	*	*	*	*
V54.8	*6	*	*	*	*	*	*	*	*	*	*
V55	41	20	21	*	*8	17	13	*9	13	13	*6
V55.2	*9	*5	*	*	*	*	*	*	*	*	*
V55.3	28	13	15	*	*	13	9	*	10	9	*
V57	521	202	318	*	32	100	386	80	159	163	118
V57.1	126	50	76	*	*8	17	100	22	19	50	35
V57.8	386	151	235	*	23	82	278	50	139	113	83
V57.89	386	151	235	*	23	82	278	50	139	113	83
V57.9	*7	*	*5	-	-	*	*7	*7	*	*	-
V58	186	99	87	29	41	59	57	47	37	64	39
V58.0	14	*6	*8	*	*	*	*6	*	*	9	*
V58.1	134	78	56	24	34	46	30	35	23	47	29
V58.4	22	10	11	*	*	*	10	*8	*	*	*7
V58.49	20	10	11	*	*	*	10	*8	*	*	*6
V58.8	14	*	9	*	*	*	10	*	*8	*	*
V58.89	12	*	*8	*	*	*	9	*	*8	*	*
V59	*7	*	*	*	*	*	-	*	*	*	*
V70	13	*8	*6	*	*8	*	*	*	*	*	*
V70.7	13	*8	*5	*	*8	*	*	*	*	*	*
V71	*8	*	*	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.

- Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
All codes	164,152	71,076	93,076	11,531	40,091	38,298	74,233	39,371	33,277	63,403	28,101
001-139	5,588	2,761	2,826	560	1,203	1,248	2,576	1,382	976	2,198	1,032
005	*26	*	*	*	*	*	*	*	*	*	*
008	568	208	360	91	101	98	279	130	170	181	88
008.4	311	111	200	*	49	53	202	86	106	79	*41
008.45	272	103	168	*	45	51	171	82	77	77	*36
008.6	52	24	28	40	*	*	*	*	*15	*20	*
008.61	38	21	*17	37	*	—	*	*	*13	*10	*
008.8	186	58	128	30	48	34	74	35	44	79	27
009	58	*23	35	*	*20	*	*	*	*	*22	*
009.0	29	*	*22	*	*	*	*	*	*	*	*
011	*88	*	*	*	*	*	*	*	*	*	*
034	26	*	*15	*13	*	*	*	*	*	*	*
034.0	21	*	*11	*	*	*	*	*	*	*	*
038	2,849	1,364	1,485	52	333	639	1,824	775	518	1,040	517
038.0	268	110	158	*	*	*	162	*116	*54	*	*63
038.1	465	239	226	*	*63	108	286	131	78	145	111
038.10	*86	*	*	*	*	*	*	*	*	*	*
038.11	270	142	128	*	*	*63	181	*89	*	*83	*
038.19	109	*62	*48	*	*	*	*67	*	*35	*	*
038.2	97	*	*45	*	*	*	*	*	*	*	*
038.4	691	322	369	*	*59	172	451	185	140	189	176
038.40	59	*	*35	—	*	*	*42	*	*	*34	*
038.42	366	160	206	*	*	79	266	116	83	88	79
038.49	195	102	93	*	*	*75	98	*	*	*54	*72
038.8	*75	*	*41	*	*	*	*47	*	*	*	*
038.9	1,237	601	637	*26	133	236	843	296	189	613	140
041	91	*46	*44	*	*	*	*	*	*	*49	*
042	578	355	223	*	368	185	*	184	*46	273	76
047	140	67	73	57	65	*	*	*25	*26	59	30
047.9	123	63	60	53	52	*	*	*17	*23	54	28
053	89	*42	47	*	*	*	62	*	*	*39	*
053.9	36	*	*23	*	*	*	*25	*	*	*	*
054	*49	*	*	*	*	*	*	*	*	*	*
070	130	68	62	*	*33	81	*	*	*	53	*36
070.5	61	*27	*	*	*	*43	*	*	*	*	*
070.51	*25	*	*	*	*	*	*	*	*	*	*
075	23	*	*14	*	*	*	—	*	*	*	*
079	184	92	92	102	35	*15	32	41	29	89	26
079.9	169	83	86	89	34	*14	31	38	26	80	25
079.99	168	83	85	89	34	*14	31	37	26	80	25
112	94	*41	*53	*	*	*	*50	*	*	*45	*
117	*74	*	*	*	*	*	*	*	*	*	*
135	*39	*	*30	—	*	*	*	*	*	*	*
136	*58	*	*	*	*	*	*	*	*	*	*
140-239	10,180	4,556	5,624	326	1,357	3,231	5,267	2,425	2,074	3,889	1,793
150	122	*65	*	—	*	*	*85	*	*	*	*
150.9	*37	*	*	—	*	*	*	—	*	*	*
151	291	221	*70	—	*	*133	126	*	*	103	*
151.0	*74	*	*	—	*	*	*	*	*	*	*
151.2	*108	*	*	—	*	*	*	*	*	*	*
153	1,014	423	591	*	*	209	776	247	242	372	153
153.0	*69	*	*	—	*	*	*61	*	*	*	*
153.1	*74	*	*56	—	*	*	*57	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
153.2	*69	*	*	*	*	*	*	*	*	*	*
153.3	262	140	121	—	*	64	183	*51	*56	105	*49
153.4	184	75	109	—	*	*	152	*61	*62	*	*
153.6	204	*81	123	—	*	*	181	*	*	*92	*
153.9	82	*	*58	—	*	*	*54	*	*	*	*
154	332	207	126	—	*	108	212	91	85	118	*38
154.0	121	73	*48	—	*	*42	76	*	*33	*50	*
154.1	173	120	*53	—	*	*58	107	*	*48	*62	*
155	93	60	*	*	*	*37	*53	*	*	*26	*
155.0	64	*44	*	*	*	*23	*	*	*	*	*
156	*70	*	*44	—	*	*	*54	*	*	*	*
157	276	153	123	*	*	*75	185	*47	*62	121	*
157.0	176	*101	*75	—	*	*	121	*	*	*86	*
157.9	62	*	*30	—	*	*	*40	*	*	*26	*
161	*87	*67	*	—	*	*	*	*	*	*	*
162	1,094	608	486	*	*	370	705	257	221	431	185
162.2	*49	*	*	*	*	*	*	*	*	*	*
162.3	473	254	219	—	*	178	290	107	90	172	103
162.5	201	103	98	—	*	*63	134	*54	*47	*64	*
162.8	85	*44	*41	—	*	*	*54	*	*	*	*
162.9	268	169	99	—	*	72	192	*44	*50	140	*
171	*43	*	*	*	*	*	*	*	*	*	*
172	*15	*	*	—	*	*	*	*	*	*	*
173	*26	*	*	*	*	*	*	*	*	*	*
174	246	—	246	—	37	94	115	53	57	86	50
174.4	24	—	24	—	*	*	*9	*	*	*9	*
174.8	76	—	76	—	*	23	45	*32	*	*21	*14
174.9	114	—	114	—	*	46	45	*	*32	45	*28
180	83	—	83	*	*	*25	*	*	*	*36	*17
180.8	*27	—	*27	—	*	*	*	*	*	*	*
180.9	*43	—	*43	*	*	*	*	*	*	*	*
182	129	—	129	—	*	36	82	*36	*27	49	*
182.0	126	—	126	—	*	36	80	*34	*26	49	*
183	129	—	129	*	*	58	55	*	*28	49	*
183.0	128	—	128	*	*	57	54	*	*28	48	*
185	302	302	—	—	*	141	158	57	78	99	68
188	196	150	*46	—	*	*33	162	*43	*45	75	*32
188.8	103	86	*	—	*	*	85	*	*	*	*
188.9	*41	*32	*	—	—	*	*32	*	*	*	*
189	183	118	65	*	*	62	96	*31	*47	92	*
189.0	155	98	57	*	*	49	81	*	*45	75	*
191	261	112	149	*57	*47	67	90	59	*60	78	*65
191.1	*38	*	*	*	*	*	*	*	*	*	*
191.9	*55	*	*	*	*	*	*	*	*	*	*
193	46	*13	34	*	17	*12	*17	*	*	*16	*16
196	60	*26	*34	*	*	*	*38	*	*	*	*
197	732	277	455	*	*29	216	486	201	153	264	115
197.0	100	*48	53	*	*	*32	62	*	*31	*36	*
197.2	163	*57	106	*	*	*48	107	*44	*	*56	*
197.6	194	*34	159	—	*	*48	141	*	*	*81	*
197.7	191	100	91	—	*	57	125	79	*48	50	*
198	570	236	334	*	*36	216	317	157	120	178	115
198.3	239	97	142	*	*	88	138	*68	*51	66	*55
198.5	215	103	112	*	*	79	124	*50	*39	87	*38
198.8	66	*	*43	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
198.89	63	*	*40	*	*	*	*	*	*	*	*
199	212	127	86	*	*	*72	119	*43	*60	*83	*
199.0	161	96	*66	*	*	*47	*94	*	*	*68	*
199.1	*51	*	*	—	*	*	*	*	*	*	*
200	144	*75	*	*	*	*	*	*	*	*	*
200.0	*113	*53	*	—	*	*	*	*	*	*	*
202	328	175	153	*	*	91	194	*62	*55	143	*68
202.8	284	144	140	*	*	85	163	*50	*53	131	*49
202.80	98	*39	*59	*	*	*	*65	*	*	*	*
202.83	*70	*	*	*	*	*	*	*	*	*	*
202.88	*46	*	*	—	*	*	*	*	*	*	*
203	256	99	157	—	*	79	173	*	*75	129	*
203.0	248	99	149	—	*	78	166	*	*75	123	*
203.00	248	99	149	—	*	77	166	*	*75	123	*
204	231	140	91	*77	*	*	63	*72	*	*66	*69
204.0	157	*104	*53	*77	*	*	*	*	*	*	*
204.00	133	*86	*	*67	*	*	*	*	*	*	*
204.1	71	*36	*35	—	*	*	57	*	*	*	*
204.10	68	*36	*33	—	*	*	54	*	*	*	*
205	408	215	*193	*	*123	*89	*99	*	*	*141	*
205.0	358	179	*179	*	*118	*	*	*	*	*108	*
205.00	328	*149	*179	*	*	*	*	*	*	*105	*
205.1	*48	*	*	*	*	*	*	*	*	*	*
205.10	*46	*	*	*	*	*	*	*	*	*	*
211	273	118	156	*	*36	81	156	73	*51	98	51
211.3	201	90	112	—	*	68	119	55	*36	64	*47
218	686	—	686	—	336	333	*17	115	130	327	114
218.0	75	—	75	—	44	31	*	*20	*17	26	*
218.1	208	—	208	—	105	102	*	45	37	85	41
218.2	74	—	74	—	33	41	*	*	*14	39	*
218.9	329	—	329	—	155	159	*	38	63	176	52
220	131	—	131	*	61	49	*20	*30	27	48	27
225	152	*39	113	*	*	64	*69	*	*	75	*
225.2	114	*	90	*	*	*49	*57	*	*	*55	*
226	17	*	*14	*	*	*	*	*	*	*	*
227	52	*25	27	—	*17	*14	*21	*	*	*25	*15
227.1	20	*	*10	—	*	*	*	*	*	*	*
227.3	*28	*	*	—	*	*	*	*	*	*	*
233	78	*	67	—	35	30	*	*12	*	42	*12
233.0	28	—	28	—	*	*11	*	*	*	*	*
233.1	36	—	36	—	26	*	*	*	*	*18	*
235	77	*36	*41	*	*	*	*36	*	*	*35	*
235.2	53	*	*	*	*	*	*27	*	*	*29	*
236	*24	*	*	*	*	*	*	*	*	*	*
237	*24	*	*	*	*	*	*	*	*	*	*
238	68	*38	*	*	*	*	*42	*	*	*	*
238.7	*57	*	*	*	*	*	*39	*	*	*	*
239	49	*	*22	*	*	*	*18	*	*	*	*
240-279	6,906	2,994	3,912	538	1,154	1,979	3,236	1,751	1,416	2,580	1,159
241	41	*	34	*	*9	*14	*19	*	*	*16	*10
241.0	*14	*	*11	*	*	*	*	*	*	*	*
241.1	24	*	*20	*	*	*9	*	*	*	*	*
242	35	*	*22	*	*	*	*	*	*	*	*
242.0	*14	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
244	56	*	44	*	*	*20	*32	*22	*	*	*
244.9	53	*	42	*	*	*18	*31	*20	*	*	*
250	2,893	1,502	1,391	103	589	1,067	1,134	725	500	1,116	552
250.0	477	230	247	32	76	186	182	115	106	204	52
250.00	178	88	90	*	*16	64	94	37	35	89	*18
250.01	54	26	28	*20	*	*10	*12	*	*16	*18	*
250.02	184	77	106	*	*20	91	72	44	37	83	*21
250.03	61	38	*22	*	*28	*21	*	*23	*18	*	*
250.1	448	231	217	51	266	89	*43	81	106	152	109
250.10	54	*30	*24	*	*26	*	*	*	*	*	*
250.11	211	126	85	32	140	*34	*	39	42	62	68
250.12	72	*29	*43	*	*36	*	*	*	*	*20	*
250.13	112	46	66	*	64	*	*	*	*23	54	*
250.2	104	*28	*76	*	*	*46	*47	*33	*	*49	*
250.20	*51	*	*	*	*	*	*	*	*	*	*
250.22	*27	*	*	—	*	*	*	*	*	*	*
250.3	*37	*	*	*	*	*	*	*	*	*	*
250.4	177	108	68	—	*	85	62	*	*25	69	59
250.40	107	63	43	—	*	63	*43	*	*	*38	*43
250.41	*39	*	*	—	*	*	*	*	*	*	*
250.6	361	159	202	*	77	147	124	90	51	181	*39
250.60	158	76	83	*	*20	57	79	*49	*	70	*
250.61	84	*34	*50	*	*	*37	*	*	*	*49	*
250.62	68	*32	*37	—	*	*30	*	*	*	*37	*
250.63	50	*	*33	*	*	*	*	*	*	*	*
250.7	497	318	179	—	*	163	305	148	*73	157	*119
250.70	322	198	123	—	*	89	212	*84	*	87	*102
250.71	*85	*	*	—	*	*	*	*	*	*	*
250.72	*63	*	*	—	*	*	*	*	*	*	*
250.8	735	397	338	*	76	310	348	215	114	258	148
250.80	474	246	228	*	*	200	247	127	73	171	102
250.81	126	73	53	*	*	64	*37	*	*	44	*
250.82	73	*40	*33	—	*	*24	*38	*	*	*30	*
250.83	63	*	*	*	*	*	*	*	*	*	*
250.9	40	*17	*23	*	*	*	*	*	*	*14	*
250.92	*23	*	*	*	*	*	*	*	*	*	*
251	52	*	29	*	*	*	*29	*	*	*27	*
251.2	40	*	*23	*	*	*	*25	*	*	*22	*
252	*25	*	*21	—	*	*	*	*	*	*	*
252.0	*21	*	*17	—	*	*	*	*	*	*	*
253	80	*28	52	*	*	*	53	*	*	*23	*
253.6	62	*	*46	*	*	*	52	*	*	*	*
255	54	*16	*39	*	*	*	*28	*	*	*25	*
255.4	43	*	*30	*	*	*	*26	*	*	*21	*
263	*68	*37	*	*	*	*	*	*	*	*	*
263.9	*61	*	*	*	*	*	*	*	*	*	*
272	75	37	38	*	*20	37	*17	19	19	27	*
272.0	29	*13	*16	*	*	17	*	*11	*	*9	*
272.4	31	21	*10	—	*	17	*	*	*11	*7	*
274	52	*31	*	—	*	*	*37	*	*	*	*
274.0	42	*25	*	—	*	*	*29	*	*	*	*
275	152	*33	119	*	*	*61	74	*25	*42	*73	*
275.4	144	*31	113	*	*	*58	73	*	*40	*70	*
275.41	*29	*	*	*	*	*	*	*	*	*	*
275.42	76	*	*58	*	*	*	*63	*	*	*	*
276	2,656	1,020	1,636	281	187	493	1,696	656	615	958	428
276.0	55	*	*35	*	*	*	*48	*	*	*	*
276.1	336	88	248	*	*11	66	256	68	69	117	83

See footnotes at end of table.



**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
276.2	*23	*	*	*	*	*	*	*	*	*	*
276.5	1,918	785	1,133	263	136	300	1,220	497	450	703	268
276.6	*21	*	*	*	*	*	*	*	*	*	*
276.7	116	40	76	*	*	29	80	*21	*29	33	33
276.8	147	44	103	*	25	59	60	33	31	56	27
276.9	*29	*	*	—	*	*	*	*	*	*	—
277	157	81	*76	*53	81	*	*	*56	*	*35	*
277.0	112	*53	*60	*	*72	*	—	*	*	*	*
277.00	109	*51	*58	*	*72	*	—	*	*	*	*
278	336	64	272	*	175	146	*	65	76	145	50
278.0	323	63	260	*	173	138	*	56	75	143	50
278.00	20	*	*13	*	*	*	*	*	*	*	*
278.01	303	57	247	*	164	127	*	49	69	137	48
280-289	1,942	818	1,124	242	542	343	815	475	366	868	233
280	269	75	193	*	23	50	192	78	72	91	29
280.0	114	39	75	*	*	30	75	*34	27	46	*
280.9	145	29	116	*	*14	*18	110	44	43	43	*15
281	*22	*	*	*	*	*	*	*	*	*	*
282	457	218	239	100	317	*34	*	135	61	242	*
282.6	445	215	230	96	315	*32	*	129	59	238	*
282.60	*26	*23	*	*	*	*	*	*	*	*21	*
282.62	412	188	224	83	297	*31	—	126	53	215	*
283	*43	*	*	*	*	*	*	*	*	*	*
284	145	86	58	*	*	33	79	*	*28	70	*
284.8	130	75	55	*	*	*31	76	*	*28	57	*
285	369	139	231	*	35	76	251	101	74	130	65
285.1	38	*	29	*	*	*	*26	*	*18	*	*
285.2	48	*24	*24	—	*	*	*27	*	*	*	*
285.9	279	103	176	*	*29	49	195	75	47	100	57
286	99	37	61	*	*	*	60	*	*25	55	*
286.7	*33	*	*	*	*	*	*	*	*	*	*
286.9	*37	*	*	*	*	*	*20	*	*	*	*
287	161	45	116	*20	*43	*	*71	*35	*	85	*24
287.3	96	*	83	*12	*	*	*	*	*	*60	*
287.5	*39	*	*	*	*	*	*	*	*	*	*
288	308	139	169	54	67	88	99	63	55	146	*44
288.0	238	112	126	42	*53	68	75	51	*37	122	*28
288.8	67	*25	*42	*	*14	*	*24	*	*18	*23	*
289	70	44	*26	28	*	*	*	*	*	*27	*
289.2	*16	*11	*	*10	*	—	*	*	*	*	*
290-319	17,517	8,852	8,665	1,364	9,284	4,556	2,313	6,177	2,821	6,653	1,866
290	327	108	219	—	—	*	303	95	89	111	*
290.2	*84	*	*64	—	—	*	*83	*	*	*	*
290.4	147	*55	92	—	—	*	130	*	*	*52	*
290.40	*56	*	*41	—	—	*	*50	*	*	*	*
291	673	468	206	—	341	277	*55	150	227	96	201
291.0	131	80	*	—	*44	72	*	*	*	*	*48
291.8	504	352	151	—	289	190	*25	113	197	52	141
291.81	492	348	145	—	281	187	*25	110	190	52	141
292	380	207	172	*	227	80	70	162	64	93	61
292.0	194	100	94	*	132	45	*	91	35	*33	*34
292.8	139	88	51	*	65	*	49	63	*	*40	*
292.81	62	*38	*24	*	*	*	49	*	*	*24	*
292.84	72	47	*	*	58	*	—	*44	*	*	*
293	154	54	99	*	*	*39	92	*35	*44	*44	*
293.0	75	*29	46	*	*	*	54	*	*	*21	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
293.8	*61	*	*	*	*	*	*	*	*	*	*
293.83	*47	*	*	*	*	*	*	*	*	*	*
294	335	192	143	*	*	*	266	106	*47	160	*22
294.8	296	155	141	—	*	*	259	*73	*45	157	*21
295	3,732	1,965	1,767	*	2,366	1,170	185	1,666	466	1,237	363
295.3	1,330	818	512	*	895	380	*	653	151	487	*39
295.30	400	278	122	—	256	139	*	*	*71	237	*
295.32	301	175	*126	—	185	*87	*	*151	*	133	*
295.34	601	348	253	*	431	150	*	405	*68	114	*
295.6	116	*61	*56	—	*64	*	*	*	*	*50	*
295.64	*72	*	*	—	*	*	*	*	*	*	*
295.7	1,774	743	1,031	*	1,172	475	*117	681	211	614	268
295.70	1,201	519	682	*	811	294	*86	333	173	468	227
295.74	513	205	308	*	312	169	*	319	*	121	*
295.9	400	271	128	—	149	243	*	250	*55	*65	*
295.90	184	133	*50	—	102	*77	*	*	*46	*46	*
295.94	*173	*	*	—	*	*	*	*164	*	*	*
296	7,164	3,142	4,022	646	3,666	1,970	882	2,150	1,145	3,209	660
296.2	1,451	610	842	167	765	296	223	317	236	794	104
296.20	613	245	368	83	354	108	*68	117	130	329	*36
296.22	143	65	78	*	96	*	*	*	*	114	*
296.23	303	140	162	*42	159	62	*	76	*29	165	*32
296.24	388	158	230	*	152	*103	*101	*112	*66	184	*
296.3	2,548	1,199	1,349	61	1,114	953	420	775	470	936	367
296.30	472	198	274	*	205	111	142	165	100	191	*
296.32	86	*34	52	*	*44	*	*	*	*	*39	*
296.33	1,014	361	653	*25	553	307	130	397	229	346	41
296.34	960	601	359	*	305	508	*126	183	127	352	*
296.4	646	283	363	*	388	169	*	246	113	226	*62
296.40	289	121	169	*	182	*74	*	*88	*	126	*
296.43	*49	*	*	*	*33	*	*	*	*	*	*
296.44	298	136	162	*	169	81	*	133	*55	82	*
296.5	697	235	463	*	400	252	*	220	97	341	*40
296.50	334	143	191	*	207	104	*	96	66	151	*
296.53	203	*43	159	*	105	88	*	81	*	109	*
296.54	151	*	106	*	84	*55	*	*	*	77	*
296.6	703	265	438	115	418	125	*	168	84	413	*
296.60	328	122	206	*	210	*68	*	*58	*48	209	*
296.63	149	60	89	*34	90	*	*	*	*	86	*
296.64	217	81	136	*	111	*38	*	*56	*	117	*
296.7	492	191	301	*53	258	99	*82	207	96	159	*30
296.8	392	211	181	121	196	66	*	*95	*	264	*
296.80	223	97	127	95	101	*27	*	*	*	198	*
296.89	161	108	*54	*	88	*	*	*	*	65	*
296.9	228	144	84	*86	120	*	*	119	*	75	*
296.90	208	135	*73	*86	104	*	*	*100	*	74	*
297	*52	*	*	—	*	*	*	*	*	*	*
298	794	452	341	*	401	157	220	325	84	293	92
298.9	781	448	334	*	393	156	219	320	83	287	92
299	*56	*	*	*	*	*	—	*	*	*	*
300	436	187	249	*85	209	102	40	175	93	119	49
300.0	147	79	68	*	71	35	23	63	36	41	*7
300.00	101	62	38	*	55	23	*19	*52	20	21	*
300.01	*25	*	*	*	*	*	*	*	*	*	*
300.1	68	*	*24	*	*23	*	*	*	*	*	*
300.11	*31	*	*20	*	*	*	*	*	*	*	*
300.4	124	44	80	*	58	*28	*	*59	*18	38	*
300.9	47	*	*37	*	*21	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
301	*42	*	*34	*	*22	*	*	*	*	*	*
303	837	558	279	*	525	267	*44	189	218	321	108
303.0	133	74	59	*	86	42	*	*	53	27	*35
303.00	75	33	*42	*	58	*15	*	*	34	*	*
303.01	55	41	*	*	*28	*25	*	*	*	*18	*
303.9	704	484	220	—	439	224	*	172	165	294	*73
303.90	297	201	96	—	160	106	*	*	*26	201	*
303.91	390	266	124	—	269	111	*	147	126	91	*
304	492	331	161	*	388	90	*	88	62	297	*45
304.0	154	95	59	—	121	*32	*	63	*	68	*
304.00	44	*26	*	—	35	*	*	*	*	*32	*
304.01	109	69	40	—	85	*23	*	62	*	35	*
304.2	86	*56	*30	—	59	*	*	*	*	*60	*
304.20	*57	*	*	—	*40	*	*	*	*	*48	*
304.21	*29	*	*	—	*	*	—	*	*	*	*
304.7	*31	*	*	—	*30	*	—	*	*	*	—
304.71	*26	*	*	—	*	*	—	*	*	*	—
304.8	*29	*22	*	—	*26	*	—	*	*	*23	*
304.9	83	64	*	*	64	*	*	*	*	80	*
304.90	74	57	*	*	56	*	*	*	*	72	*
305	254	186	69	*	155	76	*20	30	42	108	74
305.0	115	97	18	*	51	46	*16	*16	18	36	45
305.00	62	50	*13	*	28	23	*	*10	*15	28	*10
305.01	48	44	*	—	*20	*21	*	*	*	*	*35
305.1	35	21	*14	—	20	*11	*	*	*9	17	*
305.2	*15	*15	*	—	*15	*	—	*	*	*	*
305.6	*30	*27	*	—	*19	*	*	*	*	*	*
305.9	40	*15	*25	*	36	*	*	*	*	*29	*
305.90	37	*	*	*	*34	*	*	*	*	*27	*
307	104	*	88	*	66	*	*	*	*	*29	*
309	474	245	229	137	266	62	*	267	66	104	37
309.0	92	49	43	*	64	*	*	53	*	*	*
309.2	35	*19	*	*	*24	*	*	*	*	*	*
309.28	32	*19	*	*	*22	*	*	*	*	*	*
309.4	79	42	*37	*	63	*	*	55	*	*	*
309.8	224	*105	119	*120	77	*	—	*131	*	*67	*
309.81	223	*104	119	*120	76	*	—	*131	*	*67	*
309.9	*23	*	*	—	*20	*	*	*	*	*	*
310	*30	*	*	*	*	*	*	*	*	*	*
311	684	306	379	50	430	159	46	366	64	220	34
312	248	191	*57	*144	97	*	*	161	*	*60	*
312.3	216	164	*	*	81	*	*	138	*	*	*
312.30	82	*44	*	*	*40	*	*	*33	*	*	*
312.34	*122	*113	*	*	*	*	*	*	*	*	*
313	*61	*	*	*	*	—	—	*	*	*	*
313.8	*61	*	*	*	*	—	—	*	*	*	*
313.81	*45	*	*	*	*	—	—	*	*	*	*
314	140	116	*	*115	*25	—	—	*	*	*59	*
314.0	124	113	*	*112	*	—	—	*	*	*42	*
314.01	122	*112	*	*112	*	—	—	*	*	*42	*
319	*10	*	*	*	*	*	*	*	*	*	*
320-389	2,651	1,219	1,432	351	592	675	1,033	713	545	1,050	342
320	*63	*	*	*	*	*	*	*	*	*	*
323	*99	*	*	*	*	*	*	*	*	*	*
323.9	*92	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
324	*112	*	*	*	*	*	*	*	*	*	*
331	496	208	287	*	*	*	439	121	110	215	50
331.0	411	153	258	—	*	*	402	106	95	176	*33
331.3	*28	*	*	*	*	*	*	*	*	*	*
331.4	55	*37	*	*	*	*	*	*	*	*	*
332	109	79	*30	—	*	*	95	*	*	*27	*
332.0	108	79	*29	—	*	*	95	*	*	*27	*
333	81	*34	*47	*	*	*	*49	*	*	*	*
333.9	*28	*	*	*	*	*	*	*	*	*	*
337	*23	*	*	*	*	*	*	*	*	*	*
340	97	*	78	*	*33	55	*	*30	*29	*29	*
343	*23	*	*	*	*	*	*	*	*	*	*
344	*37	*	*	*	*	*	*	*	*	*	*
345	367	267	100	63	71	198	*35	201	45	80	40
345.1	41	*	*15	*	*	*	*	*	*	*	*
345.10	*33	*	*	*	*	*	*	*	*	*	*
345.3	207	*176	*31	*	*	*	*	*	*	*	*
345.4	*24	*	*	*	*	*	*	*	*	*	*
345.5	*30	*	*	*	*	*	*	*	*	*	*
345.9	44	*22	*22	*13	*	*	*	*	*	*	*
345.90	27	*	*	*	*	*	*	*	*	*	*
346	182	46	136	*	106	62	*	29	53	71	*30
346.8	*31	*	*27	*	*	*	*	*	*	*	*
346.80	*23	*	*	*	*	*	*	*	*	*	*
346.9	114	25	89	*	64	43	*	*17	32	51	*
346.90	77	*20	57	*	43	*27	*	*	*27	31	*
346.91	37	*	*32	*	*21	*16	*	*	*	*20	*
348	154	66	87	*	*37	*35	64	*30	*42	60	*21
348.1	*27	*	*	*	*	*	*	*	*	*	*
348.3	76	*	*41	*	*	*	*43	*	*	*	*
349	47	*	*33	*	*21	*	*	*	*	*	*
349.0	*18	*	*	*	*	*	*	*	*	*	*
351	28	*	*15	*	*	*	*	*	*	*	*
351.0	26	*	*14	*	*	*	*	*	*	*	*
353	*15	*	*	*	*	*	*	*	*	*	*
355	*26	*	*	—	*	*	*	*	*	*	*
356	*40	*	*	*	*	*	*	*	*	*	*
356.9	*37	*	*	—	*	*	*	*	*	*	*
357	*59	*32	*	*	*34	*	*	*	*	*	*
357.0	*47	*	*	*	*	*	*	*	*	*	*
358	*47	*	*	*	*	*	*	*	*	*	*
358.0	*47	*	*	*	*	*	*	*	*	*	*
362	*19	*	*	*	*	*	*	*	*	*	*
376	29	*	*	*	*	*	*	*	*	*	*
376.0	*24	*	*	*	*	*	*	*	*	*	*
376.01	*24	*	*	*	*	*	*	*	*	*	*
380	*26	*	*16	*	*	*	*	*	*	*	*
380.1	*16	*	*	*	*	*	*	*	*	*	*
382	47	29	*19	40	*	*	*	*	*	33	*
382.9	42	25	*17	36	*	*	*	*	*	30	*
386	90	31	59	—	*	*18	65	*21	27	31	*
386.1	44	*13	31	—	*	*	30	*	*	*18	*
386.11	30	*10	*20	—	*	*	20	*	*	*12	*
386.3	38	*15	23	—	*	*	29	*	*12	*13	*
386.30	33	*14	*19	—	*	*	25	*	*11	*11	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
390-459 . . . . .	29,934	14,829	15,105	207	1,694	7,944	20,089	7,598	6,278	11,508	4,550
394 . . . . .	*53	*	*	*	*	*	*	*	*	*	*
396 . . . . .	156	*49	107	—	*	*	130	*	*	*43	*
396.2 . . . . .	*41	*	*	—	*	*	*	*	*	*	*
396.3 . . . . .	*39	*	*	—	*	*	*	*	*	*	*
398 . . . . .	158	55	103	*	*	*	133	*65	*25	53	*
398.9 . . . . .	158	55	103	*	*	*	133	*65	*25	53	*
398.91 . . . . .	154	54	100	*	*	*	129	*65	*25	49	*
401 . . . . .	778	287	490	*	74	289	411	162	150	377	89
401.0 . . . . .	104	28	75	*	*	57	37	*23	*	63	*
401.9 . . . . .	668	258	410	*	65	231	370	136	135	311	86
402 . . . . .	468	165	303	—	*19	119	330	106	75	187	100
402.0 . . . . .	*32	*	*	—	*	*	*	*	*	*	*
402.9 . . . . .	432	149	284	—	*	106	309	102	72	169	90
402.90 . . . . .	49	*	38	—	*	*14	*32	*	*	*15	*
402.91 . . . . .	384	138	246	—	*	92	277	92	65	154	73
403 . . . . .	595	328	268	*	98	203	262	113	83	273	127
403.9 . . . . .	568	312	256	*	87	195	253	109	77	258	124
403.91 . . . . .	528	283	245	*	83	193	220	105	75	225	124
404 . . . . .	197	110	87	—	*	*42	145	*45	*39	79	*33
404.9 . . . . .	177	99	78	—	*	*35	137	*44	*34	*68	*31
404.93 . . . . .	156	89	67	—	*	*	125	*37	*28	*61	*29
410 . . . . .	4,586	2,622	1,964	*	223	1,162	3,199	1,127	950	1,667	842
410.0 . . . . .	137	90	*47	—	*	52	79	*	*39	55	*
410.01 . . . . .	133	86	*47	—	*	52	75	*	*39	53	*
410.1 . . . . .	471	295	177	—	41	172	258	105	112	166	88
410.11 . . . . .	466	293	174	—	41	171	255	104	109	166	87
410.2 . . . . .	94	56	*38	—	*	*19	*66	*	*	*37	*
410.21 . . . . .	94	56	*38	—	*	*19	*66	*	*	*37	*
410.3 . . . . .	63	*37	*	—	*	*	*39	*	*	*25	*
410.31 . . . . .	62	*36	*	—	*	*	*38	*	*	*24	*
410.4 . . . . .	542	343	199	—	34	243	265	134	85	206	117
410.41 . . . . .	538	340	198	—	34	240	263	133	84	204	117
410.5 . . . . .	47	*27	*	—	*	*	*29	*	*	*	*
410.51 . . . . .	47	*27	*	—	*	*	*29	*	*	*	*
410.7 . . . . .	2,406	1,324	1,082	—	103	489	1,814	678	524	834	369
410.71 . . . . .	2,381	1,314	1,067	—	103	486	1,793	664	522	832	363
410.8 . . . . .	88	*36	*52	—	*	*	71	*	*25	*	*
410.81 . . . . .	78	*36	*42	—	*	*	61	*	*25	*	*
410.9 . . . . .	712	393	320	*	*18	130	563	113	126	314	160
410.91 . . . . .	700	392	308	*	*18	121	559	113	125	303	159
411 . . . . .	304	151	153	—	19	124	161	85	51	125	44
411.1 . . . . .	247	124	123	—	15	107	125	68	42	101	36
411.8 . . . . .	54	26	28	—	*	*16	36	*15	*	*23	*
411.89 . . . . .	53	25	27	—	*	*15	36	*15	*	*22	*
412 . . . . .	*16	*	*9	—	*	*	*	*	*	*	*
413 . . . . .	123	41	82	—	*	38	76	30	32	48	*12
413.9 . . . . .	121	40	82	—	*	37	76	29	32	48	*12
414 . . . . .	3,987	2,377	1,610	*	151	1,479	2,357	936	854	1,650	547
414.0 . . . . .	3,885	2,313	1,572	*	143	1,446	2,295	910	837	1,604	535
414.00 . . . . .	557	313	244	—	*49	183	324	122	114	294	27
414.01 . . . . .	3,182	1,901	1,282	*	93	1,211	1,878	762	694	1,247	480
414.02 . . . . .	115	82	*33	—	*	42	72	*19	24	53	*
414.8 . . . . .	74	58	*	—	*	*	*43	*	*	*	*
414.9 . . . . .	*24	*	*	—	*	*	*17	*	*	*	*
415 . . . . .	637	225	412	*	85	170	381	150	137	261	88
415.1 . . . . .	623	224	399	*	85	167	370	147	136	252	88
415.19 . . . . .	598	216	381	*	81	160	356	136	131	247	84

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
416	59	*	*43	*	*	*	*	*	*	*	*
420	73	40	*32	*	*16	*25	*	*	*	*	*
420.9	73	40	*32	*	*16	*25	*	*	*	*	*
420.90	*41	*24	*	—	*	*	*	*	*	*	*
421	253	118	*134	*	*	*98	*75	*	*	*	*
421.0	247	117	*130	*	*	*96	*72	*	*	*	*
423	65	*40	*25	*	*	*	*	*	*	*18	*
423.9	43	*28	*	*	*	*	*	*	*	*	*
424	625	298	327	*	42	148	434	197	131	221	77
424.0	250	98	152	*	*26	87	137	79	54	89	*27
424.1	365	196	170	*	*	61	293	113	73	130	*49
425	131	82	49	*	*27	36	57	38	*25	51	*
425.4	115	71	43	*	*23	*32	51	*31	*19	49	*
426	263	144	119	*	*13	43	204	83	52	65	64
426.0	123	74	49	*	*	*	90	*29	*27	*30	*37
426.1	92	45	*47	*	*	*	87	*38	*	*22	*
426.13	31	*15	*	*	*	*	*27	*	*	*	*
426.4	*9	*	*	—	*	*	*	*	*	*	*
427	2,830	1,265	1,565	*29	104	523	2,174	834	633	983	381
427.0	96	48	48	*	*	*19	66	*45	*15	25	*
427.1	234	178	56	*	*	71	155	52	47	112	*23
427.3	1,692	740	951	*	49	310	1,322	497	403	582	209
427.31	1,532	631	902	*	40	265	1,227	461	372	526	174
427.32	159	110	50	*	*	45	94	*36	32	56	35
427.4	*34	*21	*	—	*	*	*	*	*	*	*
427.41	*34	*21	*	—	*	*	*	*	*	*	*
427.5	53	*14	*39	*	*	*14	*36	*	*	*	*
427.6	*17	*	*	*	*	*	*11	*	*	*	*
427.69	*13	*	*	—	*	*	*	*	*	*	*
427.8	678	250	428	*	30	90	548	216	147	220	96
427.81	335	105	230	—	*	*25	303	128	51	115	41
427.89	343	145	198	*	22	65	246	88	96	105	55
427.9	*23	*	*	—	*	*	*	*	*	*	*
428	5,268	2,429	2,839	*	154	1,118	3,975	1,443	1,114	2,048	662
428.0	5,255	2,422	2,833	*	154	1,115	3,965	1,437	1,113	2,045	660
429	73	*29	*44	*	*	*	*46	*	*	*36	*
430	276	105	171	*	*	137	104	*	*89	*103	*
431	462	219	243	*	*	167	259	137	85	130	110
432	158	113	*44	*	*	*	115	*	*33	*37	*64
432.1	139	99	*40	*	*	*	109	*	*29	*34	*54
433	622	322	300	*	*	153	458	169	123	267	64
433.1	471	239	232	—	*	110	354	139	80	210	43
433.10	317	171	146	—	*	72	243	78	54	154	32
433.11	154	*68	86	—	*	*	111	*61	*	*56	*
433.3	106	60	45	—	*	*25	79	*27	*31	*35	*
433.30	75	41	34	—	—	*22	52	*17	*21	*29	*
433.31	*31	*	*	—	*	*	*	*	*	*	*
434	1,755	832	923	*	53	432	1,254	456	414	636	249
434.0	60	*	*37	*	*	*	*43	*	*	*	*
434.01	*52	*	*	*	*	*	*38	*	*	*	*
434.1	236	102	134	—	*	*50	175	89	*36	*63	49
434.10	*39	*	*	—	*	*	*	*	*	*	*
434.11	198	82	116	—	*	*39	149	*70	*28	*57	*42
434.9	1,459	707	752	*	*40	371	1,036	352	355	556	195
434.90	*51	*41	*	—	*	*	*43	*	*	*	*
434.91	1,408	666	742	*	*36	368	993	348	347	532	182
435	688	307	380	*	*18	216	454	177	127	294	90

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
435.3	32	*	*25	—	—	*	*29	*	*	*	—
435.8	*22	*	*	—	*	*	*	*	*	*	*
435.9	628	286	342	*	*18	205	406	161	106	275	86
436	705	328	377	*	*	83	597	141	107	312	145
437	151	40	111	*	*	*34	94	*34	40	51	*
437.1	*31	*	*24	*	*	*	*30	*	*	*	*
437.3	*39	*	*34	*	*	*	*	*	*	*	*
438	156	66	*90	*	*	*43	106	*	*	*66	*
438.2	*62	*	*	*	*	*	*	*	*	*	*
438.20	*59	*	*	*	*	*	*	*	*	*	*
438.8	64	*19	*	*	*	*	*57	*	*	*	*
438.89	*59	*	*	*	*	*	*53	*	*	*	*
440	760	381	380	*	*	196	527	168	168	332	92
440.1	*21	*	*12	*	*	*	*16	*	*	*	*
440.2	695	352	343	*	*	178	482	159	155	301	79
440.20	*44	*	*	—	*	*	*28	*	*	*	*
440.21	109	61	48	—	*	42	64	*	*28	*42	*19
440.22	57	*29	*	*	*	*	*34	*	*	*	*
440.23	123	*43	81	—	*	*	98	*	*	*44	*
440.24	352	185	166	—	*	*70	256	*81	*75	175	*
440.3	*28	*	*	—	*	*	*	*	*	*	*
441	419	300	120	—	*	107	299	100	86	149	85
441.0	*64	*35	*	—	*	*	*	*	*	*	*
441.01	*38	*	*	—	*	*	*	*	*	*	*
441.2	*33	*	*	—	*	*	*	*	*	*	*
441.3	*71	*63	*	—	*	*	*43	*	*	*	*
441.4	225	171	*55	—	—	*32	194	50	*53	87	*35
442	70	*53	*	*	*	*	*35	*	*	*	*
442.3	*44	*39	*	*	*	*	*	*	*	*	*
443	132	70	62	*	*	*39	87	*34	*24	56	*
443.9	121	65	55	—	*	*37	83	*	*20	*53	*
444	205	91	115	—	*	66	118	*54	*48	63	*40
444.2	155	74	81	—	*	*52	85	*39	*39	*44	*34
444.22	137	67	70	—	*	*50	68	*35	*36	*41	*
446	85	*41	*44	*29	*	*	*39	*	*25	*33	*
446.1	*27	*	*	*27	—	—	*	*	*	*	*
446.5	*28	*	*	—	—	*	*27	*	*	*	*
447	83	*36	*48	*	*	*40	*40	*	*	*	*
447.1	*36	*	*	—	*	*	*	*	*	*	*
451	86	*36	50	*	*	*25	*38	*32	*	*31	*
451.0	*26	*	*	—	*	*	*	*	*	*	*
451.1	*42	*	*	—	*	*	*	*	*	*	*
451.19	*26	*	*	—	*	*	*	*	*	*	*
453	732	272	460	*	126	206	400	160	178	312	83
453.8	698	262	436	*	121	196	380	149	175	297	77
453.9	*22	*	*	—	*	*	*	*	*	*	*
454	*51	*	*	—	*	*	*	*	*	*	*
455	89	37	52	—	*18	25	46	*	*13	37	*25
455.2	56	*19	36	—	*	*14	*30	*	*	*23	*
456	*39	*	*	*	*	*	*	*	*	*	*
458	285	118	167	*	*	68	204	66	67	115	36
458.0	137	67	70	*	*	*21	112	*37	32	52	*16
458.9	122	35	87	*	*	40	79	*	*26	55	*
459	151	86	65	*	*	*48	80	*	*34	65	*
459.0	*36	*	*	—	*	*	*	*	*	*	*
459.8	90	*45	*45	*	*	*	63	*	*	*	*
459.81	85	*41	*44	—	*	*	58	*	*	*	*

See footnotes at end of table.



**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
460-519 . . . . .	18,830	8,788	10,042	2,357	1,677	3,830	10,967	4,204	3,861	7,641	3,125
461 . . . . .	37	*	29	*	*9	*	*	*	*	*	*
461.9 . . . . .	*16	*	*14	*	*	*	*	*	*	*	*
462 . . . . .	26	*15	*11	*13	*	*	*	*	*	*	*
463 . . . . .	34	*16	18	*	27	*	-	*	*	*	*
464 . . . . .	53	35	*18	46	*	*	*	*	*14	*	*18
464.4 . . . . .	28	17	*	28	-	*	-	*	*	*	*
465 . . . . .	96	57	39	57	*	*	*26	*18	*27	30	20
465.9 . . . . .	94	56	39	56	*	*	*26	*18	*27	29	*20
466 . . . . .	953	495	458	673	*28	75	176	172	168	375	238
466.0 . . . . .	288	110	178	22	*27	71	168	69	67	118	*35
466.1 . . . . .	665	385	280	651	*	*	*	103	101	257	203
466.11 . . . . .	448	256	191	443	*	*	*	64	64	170	150
466.19 . . . . .	217	129	89	208	*	*	*	39	37	87	53
473 . . . . .	49	24	*24	*16	*16	*	*	*	*	*24	*
473.9 . . . . .	*20	*	*	*	*	*	*	*	*	*13	*
474 . . . . .	25	*12	*13	18	*	*	-	*	*	*12	*6
474.1 . . . . .	15	*10	*	14	*	*	-	*	*	*	*
474.10 . . . . .	13	*9	*	13	*	-	-	*	*	*	*
475 . . . . .	29	*11	*19	*	16	*	*	*	*	*12	*
478 . . . . .	73	43	*30	*32	*	*	*	*26	*	*22	*
478.2 . . . . .	*23	*	*	*	*	*	*	*	*	*	*
480 . . . . .	212	134	78	177	*	*	*	*	33	117	51
480.1 . . . . .	139	110	*29	132	*	*	*	*	*17	*99	*18
480.9 . . . . .	68	*21	*47	40	*	*	*	*	*	*14	*33
481 . . . . .	272	120	152	*	*31	*71	155	*65	56	104	*47
482 . . . . .	1,210	571	639	123	110	179	798	199	234	551	226
482.0 . . . . .	*64	*	*	*	*	*	*	*	*	*	*
482.1 . . . . .	315	154	161	*	*	*	194	*58	*	142	*
482.2 . . . . .	88	*50	*38	*	*	*	65	*	*	*40	*
482.3 . . . . .	*74	*	*51	*	*	*	*	*	*	*	*
482.4 . . . . .	403	176	227	*	*	*68	292	*66	*63	196	*78
482.41 . . . . .	384	168	216	*	*	*64	281	*58	*60	191	*75
482.8 . . . . .	177	83	94	*	*	*	121	*	63	*77	*
482.83 . . . . .	136	60	76	*	*	*	97	*	*48	*62	*
482.9 . . . . .	90	*38	52	*	*	*	59	*25	*	*	*
485 . . . . .	101	55	46	*14	*	*	63	*32	*	38	*
486 . . . . .	5,729	2,580	3,150	468	417	1,004	3,841	1,294	1,201	2,304	929
487 . . . . .	90	38	52	35	*	*15	*30	*18	30	*22	*
487.0 . . . . .	*40	*	*	*	*	*	*	*	*	*	*
487.1 . . . . .	46	21	25	*16	*	*	*	*	*19	*12	*
490 . . . . .	88	31	57	*	*	*15	56	*	*17	42	*
491 . . . . .	2,749	1,172	1,577	*	74	801	1,868	664	565	1,228	292
491.2 . . . . .	2,744	1,169	1,575	*	73	799	1,866	663	564	1,225	292
491.20 . . . . .	103	41	62	-	*	*37	63	*29	*	53	*
491.21 . . . . .	2,641	1,128	1,513	*	70	762	1,803	634	557	1,173	277
492 . . . . .	91	*46	*46	-	*	*	57	*	*	*42	*
492.8 . . . . .	64	*25	*39	-	*	*	*42	*	*	*	*
493 . . . . .	1,531	520	1,011	427	313	408	383	345	303	615	268
493.0 . . . . .	104	68	36	77	*17	*	*	*15	*	*15	65
493.00 . . . . .	*12	*	*	*	*	*	*	*	*	*	*
493.01 . . . . .	*24	*	*	*16	*	*	-	*	*	*	*16
493.02 . . . . .	68	43	26	51	*	*	*	*	*	*11	42
493.2 . . . . .	424	106	318	*	32	200	191	94	98	165	68
493.20 . . . . .	76	*23	53	*	*	37	34	*	*23	34	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
493.21	*34	*	*29	*	*	*	*	*	*	*	*
493.22	313	78	235	*	*25	142	146	75	67	116	55
493.9	1,001	346	656	348	264	201	189	236	197	433	135
493.90	248	73	175	72	39	52	85	39	45	127	37
493.91	193	78	115	97	61	*26	*	35	40	81	37
493.92	560	194	366	178	164	123	95	162	112	225	61
494	53	*	*39	*	*	*	*29	*	*	*	*
494.1	*34	*	*	—	*	*	*	*	*	*	*
496	492	235	257	—	*28	113	351	73	99	201	119
507	1,577	851	726	*60	*58	244	1,215	450	290	581	256
507.0	1,570	846	724	*59	*56	242	1,212	448	289	579	253
510	211	144	*67	*	*	*	*85	*	*	*	*
510.9	199	135	*	*	*	*	*82	*	*	*	*
511	315	181	133	*	*29	83	195	76	71	114	53
511.0	*22	*	*	*	*	*	*	*	*	*	*
511.8	*57	*	*	—	*	*	*	*	*	*	*
511.9	227	135	92	*	*	56	148	58	54	82	*33
512	226	146	80	*	94	58	70	*35	64	73	53
512.1	*45	*	*25	*	*	*	*	*	*	*	*
512.8	162	110	52	*	86	*27	45	*	*45	45	*46
513	*88	*	*	*	*	*	*	*	*	*	*
513.0	*88	*	*	*	*	*	*	*	*	*	*
514	*43	*	*	*	*	*	*35	*	*	*	*
515	167	77	90	—	*	*	120	*	*31	*60	*
516	83	*35	*48	*	*	*	*45	*	*	*	*
516.3	*40	*	*	—	*	*	*	*	*	*	*
516.8	*44	*	*	*	*	*	*	*	*	*	*
518	1,960	928	1,031	63	183	469	1,245	451	449	777	282
518.0	*35	*	*25	*	*	*	*	*	*	*	*
518.3	*31	*	*	*	*	*	*	*	*	*	*
518.8	1,837	893	944	*51	171	447	1,168	438	409	731	260
518.81	1,494	750	744	*	126	373	967	376	313	609	195
518.82	64	*27	*37	*	*	*	*	*	*	*	*
518.84	194	*88	105	*	*	*37	146	*	*52	*81	*
518.89	51	*22	*29	*	*	*	*25	*	*	*	*
519	107	68	40	*35	*	*	*38	*	*20	48	*
519.1	*30	*21	*	*	*	*	*	*	*	*	*
519.8	45	*32	*	*	*	*	*	*	*	*	*
520-579	16,159	7,188	8,971	673	3,252	4,560	7,675	3,706	3,409	6,143	2,901
522	*25	*	*18	*	*13	*	*	*	*	*	*
522.5	*23	*	*17	*	*13	*	*	*	*	*	*
524	34	*	22	*	28	*	*	*	*	*14	*
524.0	*14	*	*	*	*13	*	—	*	*	*	*
527	*28	*	*	*	*	*	*	*	*	*	*
527.2	*24	*	*	*	*	*	*	*	*	*	*
528	*33	*	*	*	*	*	*	*	*	*	*
530	696	360	337	74	131	180	311	156	133	297	111
530.0	*42	*	*	*	*	*	*	*	*	*	*
530.1	113	50	63	*	35	32	41	*22	24	47	*20
530.10	30	*13	*16	*	*	*	*	*	*	*	*
530.11	28	*14	*	*	*	*	*	*	*	*	*
530.19	48	*17	*31	*	*	*	*17	*	*	*23	*
530.2	112	92	*	*	*	*	87	*	*	*62	*
530.3	34	*17	*	*	*	*	*24	*	*	*	*
530.5	*25	*	*	—	*	*	*	*	*	*	*
530.7	44	*26	*	—	*16	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
530.8	305	135	170	66	50	95	94	81	59	110	54
530.81	295	129	166	65	49	93	88	78	56	108	53
531	584	284	299	*	*43	141	399	129	128	167	159
531.0	61	*38	*23	*	*	*	46	*	*	*	*
531.00	60	*38	*22	*	*	*	*45	*	*	*	*
531.4	312	163	149	*	*	84	200	42	75	110	85
531.40	297	160	136	*	*	72	196	42	74	96	85
531.5	*44	*	*	—	*	*	*	*	*	*	*
531.50	*42	*	*	—	*	*	*	*	*	*	*
531.9	64	*27	37	—	*	*	41	*	*	*25	*20
531.90	58	*23	34	—	*	*	*34	*	*	*	*20
532	396	184	212	*	*32	158	204	148	78	98	72
532.0	61	*37	*	*	*	*	*50	*	*	*	*
532.00	61	*37	*	*	*	*	*50	*	*	*	*
532.4	184	95	89	*	*	55	115	*	48	63	45
532.40	182	94	87	*	*	55	112	*	46	63	44
533	65	*	*42	—	*	*	*49	*	*	*39	*
533.4	*37	*	*	—	*	*	*	*	*	*	*
533.40	*36	*	*	—	*	*	*	*	*	*	*
535	613	204	409	*16	111	201	286	160	121	244	88
535.0	91	36	55	*	*	*29	45	*26	*21	36	*
535.00	65	*21	44	*	*	*26	*23	*	*11	*30	*
535.01	*26	*	*	*	*	*	*	*	*	*	*
535.1	127	*36	90	*	*27	*48	52	*53	*	*46	*
535.10	70	*	*53	*	*	*	*21	*26	*	*	*
535.11	56	*	*37	*	*	*	*31	*	*	*	*
535.3	*20	*	*	—	*	*	*	*	*	*	*
535.4	173	60	113	*	*19	67	85	*40	37	66	*30
535.40	76	*16	60	*	*	*32	*33	*	*17	*39	*
535.41	96	44	53	*	*	*35	51	*	*	*27	*
535.5	184	48	136	*	43	40	93	*36	37	80	*31
535.50	98	*17	82	*	31	25	*35	*20	*24	44	*
535.51	86	*31	55	*	*	*	58	*	*	*36	*
536	226	64	162	*	41	50	113	51	*34	100	41
536.2	41	*	*31	*	*	*	*	*	*	*21	*
536.3	*38	*	*	*	*	*	*	*	*	*	*
536.4	114	*38	76	*	*	*	75	*	*	*51	*
536.41	*52	*	*	*	—	*	*	*	*	*	*
536.8	30	*	*22	*	*	*	*	*	*	*	*
537	187	76	111	*	*	*34	136	*35	*	90	*36
537.0	85	*	*65	*	*	*	*	*	*	*	*
537.8	90	*51	38	*	*	*	72	*	*	*27	*
537.83	72	*38	34	—	*	*	62	*	*	*23	*
540	896	530	366	225	370	171	130	194	184	253	265
540.0	316	205	111	84	94	77	*60	69	73	86	88
540.1	206	121	85	66	57	*37	*46	*	*36	64	69
540.9	373	204	170	75	219	56	*24	89	74	103	108
541	*18	*	*	*	*11	*	*	*	*	*10	*
550	117	96	*21	*10	*12	*	82	34	*24	45	*15
550.1	75	60	*15	*	*	*	56	*18	*	32	*
550.10	54	42	*	*	*	*	38	*	*	*20	*
550.9	38	32	*	*	*	*	*23	*	*	*11	*
550.90	28	*22	*	*	*	*	*18	*	*	*	*
552	336	94	242	—	43	128	165	93	76	112	55
552.1	58	*15	*43	—	*	*24	*	*	*	*	*
552.2	216	55	161	—	*25	94	97	60	*38	74	*44
552.20	*52	*	*40	—	*	*	*	*	*	*	*
552.21	150	*37	113	—	*	62	69	*51	*18	53	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
553	304	140	164	*	46	124	131	61	60	142	40
553.1	*14	*	*	*	*	*	*	*	*	*	*
553.2	215	109	106	*	30	94	91	40	43	104	29
553.20	38	*	*23	—	*	*19	*	*	*	*24	*
553.21	172	93	79	—	25	73	74	34	33	79	27
553.3	72	*22	50	*	*13	*22	*36	*	*	*30	*
555	236	113	123	*	128	55	*37	53	67	82	*35
555.0	75	*46	*28	*	*28	*	*	*	*	*	*
555.1	64	*	*35	*	*46	*	*	*	*	*	*
555.9	78	*30	48	*	40	*21	*	*	*	*36	*
556	251	141	111	*	107	58	*81	*50	*39	108	*53
556.6	*61	*	*	*	*	*	*	*	*	*	*
556.9	108	*56	*52	*	*37	*	*40	*	*	*45	*
557	408	117	291	*	*34	72	288	111	93	145	*59
557.0	260	87	173	*	*	*43	183	*68	*67	*95	*
557.9	119	*	93	—	*	*	85	*	*	*41	*
558	546	216	329	95	96	126	229	113	144	213	76
558.9	508	201	307	93	94	110	211	108	125	204	72
560	1,848	785	1,063	45	245	403	1,155	393	474	658	323
560.1	349	182	166	*	*48	62	237	*110	83	124	*33
560.2	*68	*43	*	*	*	*	*	*	*	*	*
560.3	151	79	73	*	*	*	120	*59	*30	43	*
560.39	135	72	63	*	*	*	104	*53	*29	38	*
560.8	795	324	471	*	123	208	445	99	233	282	181
560.81	652	267	385	*	100	166	371	*79	199	234	140
560.89	144	*57	87	*	*	*42	75	*	*35	*48	*
560.9	466	150	316	*	51	110	296	96	111	175	84
562	1,516	632	884	*	190	415	911	406	350	577	184
562.1	1,510	630	880	*	189	414	907	405	347	575	184
562.10	137	*54	83	—	*	*31	92	*59	*27	42	*
562.11	926	368	558	*	148	345	432	265	237	315	109
562.12	375	169	206	—	*	*28	345	66	75	173	61
562.13	73	*	*34	—	*	*	*37	*	*	*45	*
564	275	101	175	17	77	81	100	54	72	99	50
564.0	110	45	65	*15	*16	*24	55	31	*27	29	*24
564.00	86	34	52	*12	*	*16	45	*25	*22	24	*16
564.09	*22	*	*	*	*	*	*	*	*	*	*
564.1	76	*	63	*	*28	*	*27	*	*29	*19	*
564.8	*52	*	*23	*	*	*	*	*	*	*	*
564.89	*51	*	*23	*	*	*	*	*	*	*	*
566	73	36	37	*	*22	*27	*	*	*	*21	*20
567	151	73	78	*	*31	74	*42	*40	*	*60	*
567.2	122	*58	*64	*	*	*63	*	*	*	*50	*
567.8	*20	*	*	—	*	*	*	*	*	*	*
568	112	*	88	*	*35	*30	*45	*	*	*60	*
568.0	71	*	57	*	*	*	*	*	*	*47	*
568.8	*40	*	*31	*	*	*	*	*	*	*	*
568.89	*26	*	*	—	*	*	*	*	*	*	*
569	574	200	374	*	50	183	334	120	116	283	55
569.1	*40	*	*	*	*	*	*	*	*	*	*
569.3	65	28	37	*	*	*	50	*	*	27	*
569.4	55	*20	*35	*	*	*	*37	*	*	*	*
569.49	*22	*	*	*	*	*	*	*	*	*	*
569.6	96	*	73	*	*	*48	*41	*	*	*52	*
569.69	*53	*	*41	*	*	*	*	*	*	*	*
569.8	282	121	160	*	*	101	152	*48	67	146	*
569.81	*61	*	*	—	*	*	*	*	*	*	*
569.83	111	*49	*62	*	*	*46	*	*	*	*71	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
569.85	67	*34	*33	—	*	*	56	*	*	*24	*
570	*41	*	*	*	*	*	*	*	*	*	*
571	660	430	231	*	119	356	167	140	108	288	124
571.1	57	*44	*	—	*	*37	*	*	*	*	*
571.2	313	264	*49	—	*70	201	*42	*85	*50	140	*38
571.5	249	102	148	*	*	102	113	*37	*40	124	47
572	214	110	104	*	*	90	90	*52	*30	68	*64
572.2	129	72	58	*	*	68	*49	*28	*	53	*29
573	57	*	*31	*	*	*	*	*	*	*27	*
573.3	*34	*	*	*	*	*	*	*	*	*	*
574	1,450	528	922	*	334	374	736	295	323	523	309
574.0	501	213	288	*	101	152	246	104	123	177	98
574.00	475	200	275	*	93	148	232	95	117	170	94
574.01	*26	*	*	—	*	*	*	*	*	*	*
574.1	439	129	310	*	114	115	209	89	107	168	74
574.10	426	127	300	*	112	112	201	87	105	162	72
574.2	95	*15	80	*	*19	*21	54	*22	*27	35	*11
574.20	90	*13	77	*	*17	*20	52	*21	*25	35	*
574.3	*16	*	*	—	*	*	*	*	*	*	*
574.4	*65	*	*	—	*	*	*	*	*	*	*
574.5	124	55	69	—	*25	*	78	*19	*	41	*46
574.50	84	*41	43	—	*	*	55	*	*	*23	*
574.51	40	*	*26	—	*	*	*	*	*	*	*
574.6	*36	*	*	—	*	*	*	*	*	*	*
574.7	77	*	67	*	*30	*	*	*	*	*35	*
574.70	51	*	47	*	*27	*	*	*	*	*28	*
574.8	71	*	*32	—	*	*	*	*	*	*25	*
574.80	*53	*	*23	—	*	*	*	*	*	*	*
575	303	114	189	*	75	83	143	73	60	120	51
575.0	102	48	54	*	*20	*24	*57	*35	*22	*37	*
575.1	158	54	104	*	47	40	71	*24	*25	71	*39
575.10	*22	*	*	—	*	*	*	*	*	*	*
575.11	88	*	73	*	35	*22	*30	*	*	49	*
575.12	*48	*	*	—	*	*	*	*	*	*	*
575.8	*19	*	*12	*	*	*	*	*	*	*	*
576	168	63	105	*	*	55	89	*44	*27	50	48
576.1	63	*35	*	*	*	*	*37	*	*	*	*
576.2	56	*	*39	*	*	*26	*	*	*	*	*
576.8	*42	*	*31	*	*	*	*	*	*	*	*
577	1,773	919	854	*	591	633	531	309	332	784	349
577.0	1,504	804	700	*	502	499	488	262	289	707	246
577.1	187	71	116	*	62	93	*30	*	*32	51	83
578	770	359	412	*	69	121	572	250	161	262	97
578.0	83	37	46	*	*	*22	38	*19	*	*32	*23
578.1	198	107	91	*	*20	*24	152	101	35	43	*
578.9	489	214	275	*	*30	75	381	131	116	187	55
579	*57	*	*32	*	*	*	*	*	*	*	*
580-629	6,712	2,294	4,418	286	1,567	1,525	3,334	1,522	1,365	2,739	1,086
581	*37	*	*	*	*	*	*	*	*	*	*
581.9	*31	*	*	*	*	*	*	*	*	*	*
584	1,123	542	581	*	*61	309	737	313	194	403	213
584.5	*86	*	*	*	*	*	*47	*	*	*	*
584.9	1,000	496	504	*	*57	269	661	277	170	376	177
585	112	79	*33	*	*	*28	58	*30	*	*51	*
590	514	97	417	90	159	109	156	79	92	201	143
590.1	260	46	214	35	101	64	61	37	48	104	71
590.10	260	46	214	35	101	63	61	37	48	104	71

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
590.8	228	46	182	43	53	42	90	38	38	94	58
590.80	228	46	182	43	53	42	90	38	38	94	58
591	35	*19	*17	*	*	*	*	*	*	*	*
592	388	203	185	*	167	133	79	94	85	152	57
592.0	136	72	64	*	56	59	*19	35	31	53	*17
592.1	249	130	119	*	111	73	61	58	53	97	40
593	179	73	105	26	28	*34	90	46	46	53	35
593.4	38	*	*25	*	*	*	*	*	*	*	*
593.7	*23	*	*18	*21	*	—	*	*	*	*	*
593.70	*18	*	*14	*17	*	—	*	*	*	*	*
593.8	*36	*	*	*	*	*	*	*	*	*	*
593.9	54	*26	*28	*	*	*	43	*	*	*21	*
594	*15	*14	*	*	*	*	*	*	*	*	*
594.1	*11	*10	*	*	*	*	*	*	*	*	*
595	113	*32	81	*	*	*	68	*35	*	*56	*
595.0	*36	*	*	*	*	*	*	*	*	*	*
595.9	*40	*	*	—	*	*	*	*	*	*	*
596	93	79	*	*	*	*	61	*	*26	*42	*
596.0	47	47	—	*	*	*	*33	*	*	*	*
598	*15	*	*	*	*	*	*	*	*	*	*
599	1,952	679	1,273	96	142	253	1,462	515	378	827	232
599.0	1,802	585	1,217	89	133	221	1,359	479	345	770	209
599.6	*19	*	*	*	—	*	*	*	*	*	*
599.7	108	72	36	*	*	*24	75	*29	*27	37	*14
600	251	251	—	—	*	39	212	59	68	86	38
600.0	218	218	—	—	*	35	183	56	59	68	34
600.9	*18	*18	—	—	—	*	*17	*	*	*12	*
601	46	46	—	—	*	*	*39	*	*	*30	*
601.0	*35	*35	—	—	*	*	*32	*	*	*	*
604	*26	*26	—	*	*	*	*	*	*	*	*
604.9	*24	*24	—	*	*	*	*	*	*	*	*
607	*17	*17	—	*	*	*	*	*	*	*	*
608	72	72	—	*	*	*32	*	*	*	*46	*
611	47	*	42	*	14	*18	*	*17	*	*14	*
611.0	*25	*	*24	*	*	*	*	*	*	*	*
611.1	14	*	*10	*	*7	*	*	*	*	*	*
614	212	—	212	*	151	45	*	36	36	100	41
614.1	*28	—	*28	—	*	*	*	*	*	*	*
614.2	*37	—	*37	*	*25	*	*	*	*	*	*
614.6	55	—	55	*	37	*	*	*	*	25	*
614.9	45	—	45	*	43	*	—	*	*	*	*
616	43	—	43	*	26	*	*	*	*	*20	*
616.0	20	—	20	—	*17	*	*	*	*	*	*
617	215	—	215	*	169	44	*	36	54	92	33
617.0	118	—	118	—	88	29	*	*23	31	49	*16
617.1	41	—	41	—	31	*	—	*	*	*	*
617.3	31	—	31	*	26	*	*	*	*	*19	*
618	349	—	349	—	60	132	157	60	76	143	70
618.0	73	—	73	—	*11	32	29	*11	*14	30	*17
618.1	28	—	28	—	*15	*	*	*	*	*14	*
618.2	48	—	48	—	*	18	*22	*	*15	*15	*
618.3	42	—	42	—	*	*15	*20	*16	*	*13	*
618.4	48	—	48	—	*	*16	*21	*	*12	25	*
618.5	83	—	83	—	*	27	52	*	*19	31	*23
618.8	*17	—	*17	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
619	*46	—	*46	—	*	*	*	*	*	*	*
619.1	*33	—	*33	—	*	*	*	*	*	*	*
620	179	—	179	*	122	42	*	30	38	79	33
620.0	*14	—	*14	*	*	*	*	*	*	*	*
620.1	31	—	31	*	25	*	—	*	*	*	*
620.2	115	—	115	*	77	24	*	*16	25	54	20
621	45	—	45	—	*	*24	*	*	*	*	*
621.3	*15	—	*15	—	*	*	*	*	*	*	*
622	32	—	32	—	*19	*13	*	*	*	*18	*
622.1	31	—	31	—	*18	*13	*	*	*	*18	*
623	*22	—	*22	*	*	*	*	*	*	*	*
625	178	—	178	*	109	50	*17	23	38	83	33
625.3	46	—	46	—	42	*	—	*	*	23	*
625.6	70	—	70	—	*18	39	*13	*	*17	32	*15
625.9	48	—	48	—	40	*	*	*	*	25	*
626	248	—	248	*	169	77	*	31	62	102	53
626.2	173	—	173	*	125	48	—	*20	43	71	39
626.8	61	—	61	—	38	23	*	*	*15	26	*10
627	33	—	33	—	*	*17	*	*	*	*	*
627.1	27	—	27	—	*	*16	*	*	*	*	*
630-677	1,411	—	1,411	*	1,404	*	—	268	223	672	248
632	19	—	19	*	19	*	—	*	*	*9	*
633	60	—	60	—	60	*	—	*11	*10	23	*17
633.1	50	—	50	—	49	*	—	*	*	21	*11
634	35	—	35	*	35	—	—	*	*	18	*
634.9	24	—	24	*	24	—	—	*	*	13	*
634.91	19	—	19	—	19	—	—	*	*	*10	*
635	*11	—	*11	*	*11	*	—	*	*	*	*
640	*15	—	*15	—	*15	—	—	*	*	*	*
641	46	—	46	*	46	—	—	*	*	*	*
641.1	*27	—	*27	—	*27	—	—	*	*	*	*
641.13	*27	—	*27	—	*27	—	—	*	*	*	*
641.9	*9	—	*9	—	*9	—	—	*	*	*	*
641.93	*9	—	*9	—	*9	—	—	*	*	*	*
642	63	—	63	*	63	*	—	*	*	37	*
642.3	24	—	24	—	24	*	—	*	*	*16	*
642.33	21	—	21	—	21	*	—	*	*	*14	*
642.4	*21	—	*21	*	*21	—	—	*	*	*	*
643	132	—	132	*	132	—	—	*16	*22	58	36
643.0	48	—	48	—	48	—	—	*	*	23	*
643.03	48	—	48	—	48	—	—	*	*	23	*
643.1	64	—	64	*	64	—	—	*	*	*21	*
643.13	64	—	64	*	64	—	—	*	*	*21	*
643.2	*18	—	*18	—	*18	—	—	*	*	*13	*
643.23	*18	—	*18	—	*18	—	—	*	*	*13	*
644	280	—	280	*	279	*	—	69	44	125	42
644.0	268	—	268	*	267	*	—	67	42	118	40
644.03	268	—	268	*	267	*	—	67	42	118	40
644.1	12	—	12	—	12	—	—	*	*	*7	*
644.13	12	—	12	—	12	—	—	*	*	*7	*
646	171	—	171	*	169	—	—	*17	25	92	37
646.2	*11	—	*11	—	*11	—	—	*	*	*	*
646.23	*11	—	*11	—	*11	—	—	*	*	*	*
646.6	114	—	114	*	113	—	—	*	*17	65	*21
646.63	110	—	110	*	109	—	—	*	*16	63	*21
646.8	39	—	39	—	39	—	—	*	*	17	*13

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
646.83	34	—	34	—	34	—	—	*	*	17	*
647	26	—	26	—	26	—	—	*	*	*15	*
647.8	*16	—	*16	—	*16	—	—	*	*	*	*
647.83	*15	—	*15	—	*15	—	—	*	*	*	*
648	292	—	292	*	292	*	—	58	49	145	41
648.0	33	—	33	—	33	—	—	*	*	*23	*
648.03	33	—	33	—	33	—	—	*	*	*23	*
648.4	60	—	60	—	60	—	—	*	*	*	*
648.43	*50	—	*50	—	*50	—	—	*	*	*	*
648.9	147	—	147	*	147	*	—	34	26	65	22
648.93	141	—	141	*	140	*	—	31	26	63	21
654	59	—	59	—	59	—	—	*	*	*27	*
654.5	*48	—	*48	—	*48	—	—	*	*	*	*
654.53	*48	—	*48	—	*48	—	—	*	*	*	*
658	*29	—	*29	—	*29	—	—	*	*	*	*
659	*8	—	*8	—	*8	*	—	*	*	*	*
666	*12	—	*12	—	*11	*	—	*	*	*	*
670	*15	—	*15	*	*15	—	—	*	*	*	*
670.0	*15	—	*15	*	*15	—	—	*	*	*	*
670.04	*15	—	*15	*	*15	—	—	*	*	*	*
674	49	—	49	—	49	—	—	*	*	*27	*
674.3	*23	—	*23	—	*23	—	—	*	*	*	*
674.34	*23	—	*23	—	*23	—	—	*	*	*	*
680-709	3,177	1,487	1,690	281	692	909	1,296	830	658	1,250	439
681	82	51	*30	*	*26	*	*35	*	*31	*32	*
681.0	36	*25	*	*	*	*	*	*	*	*	*
681.00	*29	*	*	*	*	*	*	*	*	*	*
681.1	*45	*	*	*	*	*	*	*	*	*	*
681.10	*45	*	*	*	*	*	*	*	*	*	*
682	2,106	1,016	1,091	112	432	700	863	592	431	792	291
682.0	137	51	86	32	34	*20	52	50	*20	49	*18
682.1	*39	*	*	*	*	*	*	*	*	*	*
682.2	202	104	98	*	*50	96	*48	*	*27	127	*26
682.3	186	96	90	*	65	63	*47	43	42	85	*16
682.4	89	51	38	*	*22	*26	*36	*21	*16	*36	*16
682.5	57	*	*48	*	*	*	*	*	*	*	*
682.6	1,130	558	572	*21	205	378	526	364	237	372	157
682.7	245	115	130	*12	*33	91	109	74	*50	79	*43
683	*35	*	*25	*27	*	*	*	*	*	*	*
693	*28	*	*	*	*	*	*	*	*	*	*
693.0	*25	*	*	*	*	*	*	*	*	*	*
695	*51	*	*	*	*	*	*	*	*	*	*
695.1	*48	*	*	*	*	*	*	*	*	*	*
701	24	*12	*	*	*	*	*	*	*	*20	*
701.4	*18	*11	*	*	*	*	—	*	—	*18	—
707	631	252	379	*	105	152	363	160	153	217	100
707.0	362	137	225	*	72	*89	199	*78	81	125	*79
707.1	238	102	136	*	*20	*59	155	*73	65	79	*
707.13	*25	*	*	*	*	*	*	*	*	*	*
707.15	59	*	*32	*	*	*	*32	*	*	*	*
707.19	*40	*	*	*	*	*	*	*	*	*	*
709	123	70	53	83	37	*	*	*	*	119	*
709.2	117	66	51	79	37	*	*	*	*	116	*
710-739	6,805	2,777	4,028	143	1,036	2,098	3,528	1,511	1,651	2,392	1,251
710	123	*	98	*	71	*25	*	*	*25	58	*
710.0	85	*	75	*	58	*19	*	*	*21	*43	*

See footnotes at end of table.



**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
711	148	73	*74	*	*	*	*95	*	*	*68	*
711.0	145	72	*73	*	*	*	*93	*	*	*66	*
711.06	*51	*	*	*	*	*	*	*	*	*	*
714	84	*25	59	*	*	30	38	*24	*20	*31	*
714.0	65	*	49	—	*	*26	*29	*	*16	*29	*
715	2,376	905	1,471	—	68	749	1,560	569	620	788	399
715.1	178	69	109	—	*	76	100	41	*32	69	35
715.15	54	*26	*28	—	*	*	*29	*	*	*	*
715.16	118	42	75	—	*	51	66	*	*	53	*24
715.3	1,236	494	742	—	31	368	837	280	337	415	205
715.35	346	137	209	—	*	108	227	113	104	76	53
715.36	868	350	518	—	*20	251	597	165	230	330	143
715.9	920	328	592	—	31	287	602	233	233	296	158
715.90	*29	*	*21	—	*	*	*	*	*	*	*
715.91	*17	*	*	—	*	*	*	*	*	*	*
715.95	291	121	171	—	*	90	185	83	71	83	54
715.96	570	192	379	—	*	176	383	136	148	188	99
716	109	41	67	*	*	40	51	*	*20	56	*16
716.1	*20	*	*	—	*	*	*	*	*	*	*
716.9	69	*27	42	*	*	*20	*39	*	*	*39	*
716.96	*43	*	*	*	*	*	*	*	*	*	*
717	22	*	*	*	*9	*	*	*	*	*	*
717.8	*9	*	*	*	*	*	—	*	*	*	*
717.83	*8	*	*	*	*	*	—	*	*	*	*
718	35	*15	*19	*	*12	*	*	*	*	*	*10
718.3	*11	*	*	*	*	*	*	*	*	*	*
719	138	60	78	*	34	*21	81	*39	35	42	*22
719.4	98	36	62	*	31	*16	51	*	31	37	*14
719.41	*15	*	*	*	*	*	*	*	*	*	*
719.45	53	*20	*33	*	*	*	37	*	*	*24	*
721	325	128	197	—	27	104	194	53	81	131	61
721.0	62	30	32	—	*	31	*24	*	*14	25	*11
721.1	86	38	48	—	*	*22	60	*	*	50	*
721.3	148	47	100	—	*	41	94	*23	38	49	38
722	992	438	554	*	320	417	252	187	225	362	219
722.0	93	38	55	—	39	41	*12	*14	16	46	*16
722.1	493	244	249	*	187	205	101	102	111	171	108
722.10	479	233	246	*	184	194	100	101	110	160	108
722.4	24	*10	*15	—	*	*10	*	*	*	*8	*
722.5	218	85	132	—	61	91	66	*32	59	69	58
722.52	216	85	131	—	61	90	66	*32	58	69	58
722.7	63	21	41	—	*	29	*	*	*	28	*
722.71	35	*16	*19	—	*	19	*	*	*	*17	*
722.73	*22	*	*	—	*	*	*	*	*	*	*
722.8	*36	*	*25	—	*	*	*	*	*	*25	*
722.83	*34	*	*25	—	*	*	*	*	*	*25	*
722.9	59	*27	*32	*	*	*21	*	*	*	*	*
722.93	45	*	*	*	*	*	*	*	*	*	*
723	69	42	27	*	*9	34	*23	*	*16	33	*
723.0	33	*24	*	—	*	*18	*	*	*	*	*
723.1	*16	*	*	*	*	*	*	*	*	*	*
723.4	*16	*	*	—	*	*10	*	*	*	*9	*
724	463	196	267	*	71	122	267	145	113	144	61
724.0	256	118	138	—	*	84	164	62	66	89	38
724.02	250	117	134	—	*	84	158	59	64	89	38
724.2	102	32	70	*	32	*20	48	*43	29	*17	*
724.4	28	*	*	—	*	*	*	*	*	*17	*
724.5	60	*	34	*	*	*	*36	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
726 . . . . .	112	62	50	*	*26	46	36	*24	*29	*26	33
726.1 . . . . .	26	*12	*14	—	*	15	*8	*	*	*	*9
726.10 . . . . .	24	*12	*12	—	*	*14	*8	*	*	*	*9
726.2 . . . . .	*10	*	*	—	*	*	*	*	*	*	*
726.3 . . . . .	*31	*24	*	—	*	*	*	*	*	*	*
726.33 . . . . .	*28	*24	*	—	*	*	*	*	*	*	*
726.6 . . . . .	*28	*	*	*	*	*	*	*	*	*	*
727 . . . . .	66	26	40	*	*	*16	33	*16	*	24	*11
727.0 . . . . .	*26	*	*	*	*	*	*	*	*	*	*
728 . . . . .	235	128	107	*	*49	*39	140	*	94	80	*
728.8 . . . . .	206	110	96	*	*46	*	128	*	*91	*61	*
728.89 . . . . .	139	76	*63	*	*	*	109	*	*66	*46	*
729 . . . . .	181	89	91	*	52	64	58	30	28	81	42
729.1 . . . . .	39	*16	23	*	*	*22	*	*	*	*19	*
729.5 . . . . .	60	28	32	*	*15	*16	*26	*	*15	*23	*
729.8 . . . . .	53	*30	*23	*	*	*	*21	*	*	*33	*
729.89 . . . . .	42	*	*13	*	*	*	*17	*	*	*	*
730 . . . . .	446	257	189	*	93	145	186	*85	86	168	*107
730.0 . . . . .	86	*59	*	*	*	*	*42	*	*	*	*
730.07 . . . . .	*39	*	*	*	*	*	*	*	*	*	*
730.1 . . . . .	*83	*	*	*	*	*	*	*	*	*	*
730.2 . . . . .	276	140	137	*	*51	109	*105	*	*51	117	*65
730.27 . . . . .	95	*48	*	*	*	*	*62	*	*	*	*
732 . . . . .	*12	*	*	*	*	*	—	*	*	*	*
733 . . . . .	596	153	443	*	54	145	384	154	146	200	96
733.1 . . . . .	339	69	270	*	*	*55	274	92	88	98	61
733.13 . . . . .	219	41	178	*	*	*	203	58	55	66	*40
733.14 . . . . .	*61	*	*51	—	*	*	*	*	*	*	*
733.4 . . . . .	92	*33	59	*	*	44	*36	*	*23	37	*
733.42 . . . . .	78	*31	47	*	*	*41	*26	*	*	*33	*
733.6 . . . . .	39	*	32	*	*14	18	*	*	*	*17	*
733.8 . . . . .	85	36	49	*	*17	*19	47	*21	*20	33	*11
733.81 . . . . .	*19	*	*	*	*	*	*	*	*	*	*
733.82 . . . . .	66	29	37	*	*15	*16	*35	*	*18	29	*
733.9 . . . . .	*24	*	*	*	*	*	*	*	*	*	*
736 . . . . .	*22	*	*	*	*	*	*	*	*	*	*
737 . . . . .	99	*	68	*36	*45	*	*	*33	*	*	*
737.3 . . . . .	86	*	60	*36	*38	*	*	*	*	*	*
737.30 . . . . .	67	*	*46	*	*	*	*	*	*	*	*
738 . . . . .	130	46	84	*	*34	45	50	*31	*22	*35	*42
738.4 . . . . .	104	*34	69	—	*	*36	49	*	*21	*	*41
740-759 . . . . .	1,027	559	468	797	130	61	*39	349	203	265	211
742 . . . . .	*35	*18	*	*24	*	*	*	*	*	*	*
745 . . . . .	156	86	70	138	*	*	*	65	*	*41	*
745.4 . . . . .	*42	*	*	*39	*	*	*	*	*	*	*
745.5 . . . . .	*22	*	*	*	*	*	*	*	*	*	*
746 . . . . .	107	*71	*35	*81	*	*	*	*	*	*	*
747 . . . . .	133	*66	*67	*92	*	*	*	*41	*	*	*
748 . . . . .	*66	*	*	*64	*	—	—	*	*	*	*
749 . . . . .	31	*17	*15	31	*	*	—	*	*	*16	*
749.0 . . . . .	*18	*	*	*17	*	*	—	*	*	*12	*
749.00 . . . . .	*11	*	*	*	*	—	—	*	*	*	*
749.2 . . . . .	*12	*	*	*12	*	—	—	*	*	*	*
750 . . . . .	80	63	*	78	*	—	*	*	*	*	*38
750.5 . . . . .	30	26	*	30	*	—	—	*	*	*	*11

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
751	115	*76	*39	87	*	*	*	*	*	*33	*
753	*39	*23	*	*	*	*	*	*	*	*	*
754	34	*19	*15	*23	*	*	*	*	*	*	*
755	*20	*	*	*	*	*	*	*	*	*	*
756	135	*50	85	97	*	*	*	*	*36	*56	*
756.1	43	*	*25	*	*	*	*	*	*	*	*
756.12	*30	*	*	*	*	*	*	*	*	*	*
757	*8	*	*	*	*	—	—	*	*	*7	*
757.3	*7	*	*	*	*	—	—	*	*	*6	*
757.32	*6	*	*	*	*	—	—	—	—	*6	—
759	*20	*	*	*	*	*	*	*	*	*	*
760-779	1,847	1,006	841	1,845	*	—	*	229	378	689	552
765	636	363	272	636	—	—	—	*	*	351	*
765.1	424	219	*206	424	—	—	—	*	*	244	*
765.18	*79	*	*	*79	—	—	—	*	*	*	*
769	528	*206	*322	528	—	—	*	*	*	*	*275
770	251	161	90	251	—	—	*	*62	*70	*55	63
770.6	*46	*	*	*46	—	—	—	*	*	*	*
770.8	116	*83	*33	115	—	—	*	*	*	*	*30
771	90	*52	*38	90	—	—	—	*	*	*	*29
771.8	66	*44	*	66	—	—	—	*	*	*	*
774	80	44	36	80	—	—	—	*10	*13	23	33
774.6	68	35	33	68	—	—	—	*	*11	*18	31
777	*63	*	*	*63	—	—	—	*	*	*	*
778	*18	*	*	*18	—	—	—	*	*	*	*
778.4	*16	*	*	*16	—	—	—	*	*	*	*
779	99	*75	*	99	—	—	—	*	*	*	*
780-799	750	325	426	183	196	155	216	158	112	256	225
780	150	83	68	70	33	*14	33	34	20	58	38
780.2	17	*	*9	*	*	*	*	*	*	*	*
780.3	63	35	28	32	19	*	*	*15	*9	19	*19
780.39	60	33	27	28	19	*	*	*14	*9	*18	*19
780.6	47	*26	*21	35	*	*	*	*	*	*22	*
784	*13	*	*9	*	*	*	*	*	*	*	*
786	119	61	58	*13	37	50	19	27	23	46	23
786.0	*17	*	*9	*	*	*	*	*	*	*	*
786.09	*8	*	*	*	*	—	*	*	*	*	*
786.5	99	53	47	*	32	50	17	24	17	37	21
786.50	50	29	21	*	16	23	*11	*	*9	22	*
786.59	44	21	23	*	14	23	*	*13	*7	*12	*12
787	*17	*	*	*	*	*	*	*	*	*	*
788	*11	*	*	*	*	*	*	*	*	*9	*
789	73	37	36	15	42	*8	*	*14	13	31	*16
789.0	73	37	36	15	42	*8	*	*14	13	31	*16
789.00	21	*10	*11	*	*16	*	*	*	*	*12	*
789.03	17	*	*7	*	*8	*	*	*	*	*	*
789.09	*9	*	*	*	*	*	*	*	*	*	*
799	326	116	211	67	51	68	140	68	45	79	134
799.9	325	115	210	67	50	68	140	68	45	79	133
800-999	14,238	7,177	7,061	971	3,402	3,537	6,328	3,069	2,909	5,519	2,741
800	63	42	*	*26	*	*	*	*	*	*	*
801	209	174	*35	*	107	*48	*	*39	*60	*74	*36
801.0	51	*40	*	*	*	*	*	*	*	*	*
801.1	*68	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
801.2	*54	*46	*	*	*	*	*	*	*	*	*
802	103	74	*29	*	54	*25	*	*24	*19	44	*16
802.2	32	*25	*	*	*18	*	*	*	*	*17	*
802.4	*23	*	*	*	*	*	*	*	*	*	*
802.8	*21	*	*	*	*	*	*	*	*	*	*
805	409	180	229	*	101	87	218	91	99	168	51
805.0	110	71	*39	*	*	*43	*47	*	*	*40	*
805.2	94	*20	74	*	*29	*	*49	*	*22	*37	*
805.4	187	83	105	*	50	*29	107	*35	43	86	*23
806	125	*104	*	*	*75	*	*	*	*	*	*
807	206	126	80	*	*28	62	113	*31	38	73	64
807.0	182	109	73	*	*25	*52	102	*28	37	*62	*55
807.01	*28	*	*	—	*	*	*	*	*	*	*
807.03	*36	*	*	*	*	*	*	*	*	*	*
808	356	118	237	*	83	*41	227	55	99	137	65
808.0	85	*59	*	*	*44	*	*	*	*	*51	*
808.2	216	47	169	*	*25	*	163	*36	68	67	*45
808.8	*34	*	*	—	*	—	*	*	*	*	*
812	223	86	137	18	*26	*29	150	70	44	62	47
812.0	128	*38	90	*	*	*	108	*32	*26	*41	*29
812.00	40	*	*20	*	*	*	*35	*	*	*	*
812.01	*39	*	*33	*	*	*	*36	*	*	*	*
812.09	*34	*	*25	—	*	*	*26	*	*	*	*
812.2	*30	*	*21	*	*	*	*	*	*	*	*
812.21	*23	*	*17	*	*	*	*	*	*	*	*
812.4	49	27	22	17	*	*	*	*23	*	*	*9
812.41	31	*20	*11	12	*	*	*	*	*	*	*7
813	171	96	75	*15	48	38	71	33	31	63	44
813.0	*32	*	*	*	*	*	*	*	*	*	*
813.2	*12	*	*	*	*	*	*	*	*	*	*
813.4	64	32	33	*	*14	*	34	*	*16	*21	*18
813.41	*15	*	*	*	*	*	*	*	*	*	*
813.42	27	*	*19	*	*	*	*16	*	*	*	*
813.44	*20	*	*	*	*	*	*	*	*	*	*
815	*17	*	*	*	*	*	*	*	*	*	*
816	*15	*13	*	*	*	*	*	*	*	*	*
816.1	*11	*	*	*	*	*	*	*	*	*	*
820	2,064	595	1,468	*	*36	172	1,851	516	481	686	379
820.0	599	163	436	*	*	*37	554	159	149	169	121
820.00	*42	*	*35	—	—	*	*41	*	*	*	*
820.02	*55	*	*	*	—	*	*54	*	*	*	*
820.03	*40	*	*38	*	*	*	*	*	*	*	*
820.09	458	120	338	*	*	*	427	131	124	124	79
820.2	993	272	721	*	*	73	900	283	239	331	140
820.20	51	*	48	*	—	*	49	*	*	*	*
820.21	854	237	617	*	*	*59	788	253	209	274	118
820.22	88	*	*57	*	*	*	*62	*	*	*	*
820.8	456	147	309	*	*	*56	395	73	88	179	116
821	449	219	231	57	151	85	156	111	92	179	67
821.0	224	113	111	45	94	*	62	*40	57	103	*23
821.00	*34	*	*	*	*	*	*	*	*	*	*
821.01	190	100	90	*38	86	*	*47	*37	54	81	*17
821.2	157	*48	109	*	*	*45	93	*	*31	*57	*35
821.20	*34	*	*	*	*	*	*	*	*	*	*
821.23	82	*	*59	*	*	*	*53	*	*	*	*
822	80	42	38	*	*15	*18	47	*	*	*29	*20
822.0	78	39	38	*	*	*18	47	*	*	*27	*20

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
823 . . . . .	316	185	131	*13	135	72	96	68	70	103	75
823.0 . . . . .	109	52	57	*	29	*25	*54	*34	*25	*32	*
823.00 . . . . .	86	42	44	*	*26	*	*41	*	*21	*20	*
823.2 . . . . .	55	46	*	*	32	*	*	*	*	*	*
823.22 . . . . .	39	*35	*	*	*23	*	*	*	*	*	*
823.3 . . . . .	*61	*45	*	*	*	*	*	*	*	*	*
823.32 . . . . .	*52	*41	*	*	*	*	*	*	*	*	*
823.8 . . . . .	69	*25	*	*	*22	*	*	*	*	*	*
823.82 . . . . .	*51	*	*	*	*	*	*	*	*	*	*
824 . . . . .	380	153	227	*	125	112	135	76	82	118	105
824.0 . . . . .	*20	*	*	*	*	*	*	*	*	*	*
824.2 . . . . .	*16	*	*	*	*	*	*	*	*	*	*
824.4 . . . . .	89	30	60	*	*24	*25	*39	*25	*19	*17	*27
824.6 . . . . .	79	35	44	*	21	*27	*30	*	*16	22	*29
824.8 . . . . .	90	49	41	*	40	*18	*	*25	*22	25	*18
824.9 . . . . .	*36	*	*	*	*	*	*	*	*	*	*
825 . . . . .	94	55	40	*	49	*29	*	*	*24	*31	*
825.0 . . . . .	31	*23	*	*	*18	*	*	*	*	*	*
825.2 . . . . .	30	*	*18	*	*14	*	*	*	*	*	*
825.25 . . . . .	*19	*	*	*	*	*	*	*	*	*	*
831 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
831.0 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
840 . . . . .	42	*20	22	*	*	*22	*16	*	*	*12	*
840.4 . . . . .	37	*17	*20	—	*	*20	*15	*	*	*	*
844 . . . . .	29	*19	*	—	*11	*	*	*	*	*	*
844.8 . . . . .	*22	*16	*	—	*	*	*	*	*	*	*
847 . . . . .	26	*13	*	*	*	*	*	*	*	*	*
850 . . . . .	103	55	49	*11	38	*15	39	19	*18	24	42
850.0 . . . . .	26	*11	*16	*	*	*	*	*	*	*	*
850.1 . . . . .	38	20	*18	*	*13	*	*16	*	*	*	*15
850.5 . . . . .	*23	*12	*	*	*	*	*	*	*	*	*
850.9 . . . . .	*15	*	*	*	*	*	*	*	*	*	*
851 . . . . .	127	65	62	*	*43	*	*48	*	*	*39	*24
851.8 . . . . .	88	*45	*43	*	*	*	*33	*	*	*	*
852 . . . . .	292	182	111	*	76	*33	141	*28	65	97	102
852.0 . . . . .	60	*36	*	*	*24	*	*	*	*	*	*
852.2 . . . . .	194	111	83	*	*	*25	107	*	*46	*53	75
852.20 . . . . .	*33	*	*	*	*	*	*	*	*	*	*
852.21 . . . . .	78	*36	*42	*	*	*	*56	*	*	*	*
853 . . . . .	*66	*	*	*	*	*	*	*	*	*	*
853.0 . . . . .	*66	*	*	*	*	*	*	*	*	*	*
854 . . . . .	*21	*	*	*	*	*	*	*	*	*	*
854.0 . . . . .	*20	*	*	*	*	*	*	*	*	*	*
860 . . . . .	214	156	*58	*	70	84	*53	*41	*61	65	*47
860.0 . . . . .	78	43	*35	*	*25	*	*	*	*	*32	*22
860.4 . . . . .	*90	*71	*	*	*	*	*	*	*	*	*
861 . . . . .	71	*60	*	*	*30	*	*	*	*	*	*
861.2 . . . . .	*55	*49	*	*	*	*	*	*	*	*	*
861.21 . . . . .	*53	*47	*	*	*	*	*	*	*	*	*
863 . . . . .	*92	*65	*	*	*59	*	*	*	*	*	*
864 . . . . .	74	*63	*	*	*65	*	*	*	*	*	*
864.0 . . . . .	*35	*	*	*	*28	*	*	*	*	*	*
865 . . . . .	112	*92	*19	*	52	*	*	*	*	*68	*
865.0 . . . . .	82	*63	*19	*	*50	*	*	*	*	*41	*
871 . . . . .	*17	*16	*	*	*10	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
873	71	45	25	*8	*34	*	*21	*	*13	*31	*17
873.0	*16	*	*	*	*	*	*	*	*	*	*
873.4	*24	*19	*	*	*	*	*	*	*	*	*
881	*11	*	*	*	*	*	*	*	*	*	*
882	*20	*16	*	*	*	*	*	*	*	*	*
883	*20	*	*	*	*	*	*	*	*	*	*
886	*28	*	*	*	*	*	*	*	*	*	*
886.0	*25	*	*	*	*	*	*	*	*	*	*
891	*31	*	*	*	*	*	*	*	*	*	*
892	*35	*	*	*	*	*	*	*	*	*	*
920	*17	*	*	*	*	*	*	*	*	*	*
922	45	*23	*22	*	*	*	*29	*	*	*	*
922.1	*15	*	*	*	*	*	*	*	*	*	*
922.2	*13	*	*	*	*	*	*	*	*	*	*
924	109	*16	93	*	*	*	85	*37	*19	*35	*
924.0	59	*	47	*	*	*	54	*35	*	*	*
924.01	54	*	*44	*	*	*	52	*33	*	*	*
935	*16	*13	*	*	*	*	*	*	*	*	*
935.1	*15	*12	*	*	*	*	*	*	*	*	*
941	108	*89	*	*	*	*	*	*	*	*84	*
941.2	*24	*12	*	*	*	*	*	*	*	*	*
942	88	*63	*	*33	*	*	*	*	*	*55	*
942.3	*70	*	*	*	*	*	*	*	*	*	—
943	*55	*46	*	*	*	*	*	*	*	*	*
944	*34	*	*	*	*	*	*	*	*	*26	*
945	108	56	*53	*27	*	*	*	*	*	62	*
945.2	*30	*	*	*	*	*	*	*	*	*	*
945.3	70	*36	*	*	*	*	*	*	*	*48	*
946	*105	*104	*	*	*	*	*	*	—	*99	—
949	*91	*	*	*72	*	—	—	—	—	*91	—
949.0	*90	*	*	*72	*	—	—	—	—	*90	—
958	*22	*	*	*	*	*	*	*	*	*	*
959	98	48	51	*11	44	*17	*27	*16	18	49	*16
959.0	73	36	37	*10	35	*	*18	*	*12	39	*10
959.01	69	33	36	*10	31	*	*18	*	*12	36	*10
962	*12	*	*	*	*	*	*	*	*	*	*
962.3	*11	*	*	*	*	*	*	*	*	*	*
965	161	79	83	*	96	37	*	30	31	73	*28
965.0	73	*50	*23	*	45	*	*	*	*	*40	*
965.09	*21	*	*	*	*	*	*	*	*	*	*
965.1	*15	*	*	*	*	*	*	*	*	*	*
965.4	62	*20	41	*	32	*	*	*	*	*22	*
966	29	*	*22	*	*14	*	*	*	*	*	*
966.3	*23	*	*19	*	*12	*	*	*	*	*	*
967	*28	*	*	*	*12	*	*	*	*	*	*
968	20	*14	*	—	*13	*	*	*	*	*	*
968.5	*16	*12	*	—	*	*	*	*	*	*	*
969	170	83	87	*	71	59	*23	22	40	74	34
969.0	55	33	22	*	25	*	*	*	*	32	*
969.4	74	30	44	*	21	28	*	*	*21	28	*14
969.7	*9	*	*	*	*	*	—	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
972	37	*15	*22	*	*	*	*29	*	*	*	*
972.6	*17	*	*	*	*	*	*	*	*	*	*
977	19	*	*17	*	*9	*	*	*	*	*	*
977.9	*15	*	*13	*	*	*	*	*	*	*	*
980	*20	*17	*	*	*12	*	*	*	*	*	*
980.0	*16	*	*	*	*	*	*	*	*	*	*
989	*15	*	*	*	*	*	*	*	*	*	*
992	*12	*	*	*	*	*	*	*	*	*	*
994	*28	*24	*	*	*	*	—	*	*	*	*
995	126	37	89	*	*14	28	54	*25	*22	52	*27
995.1	30	*	*24	*	*	*	*19	*	*	*13	*
995.2	*39	*	*33	*	*	*	*	*	*	*	*
996	2,889	1,480	1,408	196	485	929	1,278	621	585	1,087	596
996.0	80	42	*38	*	*	*	59	*29	*	*25	*
996.01	27	*	*16	*	*	*	*24	*	*	*	*
996.04	*21	*	*	—	*	*	*	*	*	*	*
996.1	84	*	52	*	*	*13	*58	*	*	*36	*
996.2	90	48	42	*36	42	*	*	*22	*	*20	*32
996.4	415	162	253	*	38	86	286	94	104	136	80
996.5	57	*17	*40	*	*	*17	*	*	*	*22	*
996.59	*30	*	*	*	*	*	*	*	*	*	*
996.6	1,237	648	589	101	215	382	539	287	290	507	154
996.61	91	*64	*	—	*	*	*60	*	*	*	*
996.62	657	345	312	*67	121	213	256	143	136	283	95
996.64	42	*28	*	—	*	*	*30	*	*	*	*
996.66	172	77	94	*	*	*	126	*	*57	*74	*
996.67	80	*46	*	*	*	*	*	*	*	*	*
996.69	*80	*	*	*	*	*	*	*	*	*	*
996.7	722	394	328	*18	94	314	296	122	105	225	270
996.72	80	45	35	*	*	32	39	*21	*18	28	*
996.73	322	202	120	*	36	178	108	*22	*26	68	205
996.74	140	79	62	*	*	*45	70	*	*32	59	*16
996.77	41	*	*28	—	*	*	*	*	*	*	*
996.78	48	*28	*20	*	*	*	*25	*	*	*	*
996.8	190	130	*60	*	71	91	*	*	*	106	*
996.81	90	70	*	*	*56	*	*	*	*	*40	*
997	665	316	349	*22	110	272	261	159	132	220	153
997.1	75	53	*	*	*	*	*51	*	*	*	*
997.2	50	*	*41	—	*	*23	*24	*	*	*	*
997.3	41	*22	*	*	*	*	*	*	*	*	*
997.4	270	110	160	*	54	115	87	54	54	93	69
997.5	57	*	*35	—	*	*	*	*	*	*	*
997.6	135	86	*49	*	*	*58	*53	*	*	*67	*
997.62	85	*55	*	*	*	*	*	*	*	*	*
997.69	*49	*	*	*	*	*	*	*	*	*	*
998	1,491	711	779	46	299	570	575	375	282	568	265
998.1	207	109	98	*	30	52	121	74	45	50	38
998.11	109	84	26	*	*11	*26	67	*48	*22	23	*17
998.12	89	*24	65	*	*	*24	*49	*	*	*	*
998.2	112	*	*42	*	*	*	*	*	*	*	*
998.3	91	*37	*54	*	*	*38	*41	*	*	*40	*
998.5	924	429	495	*	176	365	354	236	165	344	179
998.59	907	421	486	*	172	357	349	230	163	338	176
998.6	*59	*	*	*	*	*	*	*	*	*	*
998.8	96	38	58	*	*20	*40	*25	*22	*	*30	*
998.83	*45	*	*	—	*	*	*	*	*	*	*
998.89	50	*19	*31	*	*	*15	*	*	*	*	*
999	*41	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
V01-V82 . . . . .	18,477	3,446	15,031	403	10,910	1,645	5,520	3,005	4,035	7,091	4,347
V10 . . . . .	*15	*	*	*	*	*	*	*	*	*	*
V12 . . . . .	39	*11	28	*	*	*	24	*	*	26	*
V12.5 . . . . .	35	*	27	*	*	*	22	*	*	24	*
V12.59 . . . . .	22	*	*16	*	*	*	*13	*	*	*20	*
V15 . . . . .	26	*14	*	*	*	*8	*	*	*	*	*
V15.8 . . . . .	19	*12	*	*	*	*	*	*	*	*	*
V15.81 . . . . .	*14	*	*	*	*	*	*	*	*	*	*
V17 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
V22 . . . . .	*13	—	*13	—	*13	—	—	*	*	*	*
V27 . . . . .	10,207	—	10,207	33	10,161	*	—	1,658	1,845	4,320	2,383
V27.0 . . . . .	9,418	—	9,418	30	9,377	*	—	1,493	1,634	4,021	2,270
V27.1 . . . . .	44	—	44	—	44	*	—	*	*	24	*
V27.2 . . . . .	215	—	215	*	213	*	—	*34	*42	101	37
V27.9 . . . . .	493	—	493	*	491	*	—	124	163	145	61
V30 . . . . .	89	*	*59	89	—	—	—	*	*	*	*
V30.1 . . . . .	89	*	*59	89	—	—	—	*	*	*	*
V43 . . . . .	*25	*	*	—	*	*	*	*	*	*	*
V45 . . . . .	52	*27	25	*	*	19	*	*	*11	*14	*
V45.8 . . . . .	31	*21	*10	*	*	*10	*	*	*	*	*
V45.89 . . . . .	*10	*	*	*	*	*	*	*	*	*	*
V53 . . . . .	21	*13	*	*	*	*	*16	*	*	*	*
V53.3 . . . . .	20	*13	*	*	*	*	*16	*	*	*	*
V53.31 . . . . .	*13	*	*	*	*	*	*10	*	*	*	*
V54 . . . . .	60	*	*54	*	*	*	*	*	*	*	*
V54.8 . . . . .	*52	*	*	*	*	*	*	*	*	*	*
V55 . . . . .	302	125	177	*	*41	113	124	*47	123	86	*47
V55.2 . . . . .	*46	*24	*	*	*	*	*	*	*	*	*
V55.3 . . . . .	233	84	150	*	*	93	108	*	107	69	*
V57 . . . . .	6,189	2,507	3,682	*	391	1,162	4,561	988	1,773	2,078	1,349
V57.1 . . . . .	1,504	582	923	*	*94	191	1,206	326	224	658	296
V57.8 . . . . .	4,605	1,909	2,695	*	294	967	3,282	599	1,545	1,408	1,053
V57.89 . . . . .	4,605	1,909	2,695	*	294	967	3,282	599	1,545	1,408	1,053
V57.9 . . . . .	*57	*	*46	—	—	*	*56	*54	*	*	—
V58 . . . . .	952	484	468	108	174	274	397	194	200	322	236
V58.0 . . . . .	57	*25	*32	*	*	*	*32	*	*	30	*
V58.1 . . . . .	672	380	292	96	151	233	191	163	99	241	169
V58.4 . . . . .	110	44	66	*	*	*	87	*17	*	*	*49
V58.49 . . . . .	105	40	65	*	*	*	84	*17	*	*	*48
V58.8 . . . . .	101	*	66	*	*	*	81	*	*65	*	*
V58.89 . . . . .	95	*	*65	*	*	*	80	*	*64	*	*
V59 . . . . .	*23	*	*	*	*	*	—	*	*	*	*
V70 . . . . .	28	*19	*9	*	*22	*	*	*	*	*	*
V70.7 . . . . .	27	*19	*8	*	*21	*	*	*	*	*	*
V71 . . . . .	*13	*	*	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in table 45; see "Methods" section.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	153,529	62,073	91,456	6,869	38,740	36,844	71,076	32,417	36,636	56,332	28,144
001-139	3,895	1,739	2,155	442	966	940	1,547	827	829	1,494	746
003	10	*	*6	*	*	*	*	*	*	*5	*
003.0	*6	*	*	*	*	*	*	*	*	*	*
005	*8	*5	*	*	*	*	*	*	*	*	*
005.9	*7	*	*	*	*	*	*	*	*	*	*
008	337	137	200	69	55	58	155	75	87	121	55
008.4	179	69	110	*5	23	38	114	47	47	59	26
008.45	169	66	102	*	21	35	109	45	44	56	24
008.6	33	17	16	31	*	*	*	*	10	11	*8
008.61	31	16	15	30	*	*	*	*	*9	10	*8
008.8	119	48	72	32	29	19	39	23	27	50	20
009	25	11	14	*	10	*6	*7	*7	*	10	*
009.0	15	*6	*9	*	*6	*	*	*	*	*6	*
009.2	*5	*	*	*	*	*	*	*	*	*	*
011	12	*6	*5	*	*	*	*	*	*	*6	*
011.9	*9	*	*	*	*	*	*	*	*	*5	*
031	10	*6	*	*	*	*	*	*	*	*6	*
031.0	*5	*	*	-	*	*	*	*	*	*	*
033	*6	*	*	*5	-	*	*	*	*	*	*
034	30	12	18	16	10	*	*	*	*8	10	*8
034.0	29	12	17	15	10	*	*	*	*7	10	*7
038	707	329	377	33	85	166	423	159	149	262	136
038.0	41	20	22	*	*6	10	22	10	10	11	10
038.1	138	67	71	*7	18	35	77	35	29	46	28
038.10	20	9	11	*	*	*6	9	*	*	10	*
038.11	83	43	40	*	11	22	48	24	17	26	16
038.19	35	15	19	*	*5	*6	21	*7	10	10	*7
038.2	17	*8	*9	*	*	*6	*7	*	*	*	*
038.3	*6	*	*	-	*	*	*	*	*	*	*
038.4	140	61	79	*6	15	33	85	33	32	44	30
038.40	16	*8	*8	*	*	*	*8	*	*	*7	*
038.42	69	25	43	*	*	14	49	18	17	17	17
038.43	11	*6	*5	*	*	*	*7	*	*	*	*
038.49	39	19	20	*	*6	12	19	*8	*9	15	*8
038.8	23	12	11	*	*	*6	12	*7	*	*7	*5
038.9	342	158	184	15	37	74	216	68	67	147	59
040	*5	*	*	*	*	*	*	*	*	*	*
041	1,281	495	786	106	281	270	624	247	297	500	237
041.0	213	83	129	18	60	43	93	43	54	83	32
041.00	25	12	14	*	*	*	16	*7	*6	*8	*
041.01	20	*8	12	*7	*	*	*	*	*	*9	*6
041.02	47	10	37	*	31	*7	*7	*9	13	20	*6
041.04	86	40	46	*	10	21	52	20	22	35	10
041.09	31	11	20	*	*8	*5	13	*	9	12	*5
041.1	303	158	145	23	76	80	124	56	64	126	58
041.10	29	15	14	*	*	*7	14	*7	*	10	*7
041.11	218	115	103	16	58	58	87	40	45	91	43
041.19	56	29	28	*	14	14	23	9	14	25	*8
041.2	*5	*	*	*	*	*	*	*	*	*	*
041.3	89	27	61	*5	12	15	56	15	17	35	21
041.4	340	84	257	34	63	50	193	70	75	124	72
041.5	*8	*5	*	*	*	*	*	*	*	*	*
041.6	44	13	31	*	*5	*	32	*8	11	21	*
041.7	90	42	48	*8	18	21	43	18	20	36	17
041.8	169	75	94	11	37	49	72	31	49	58	31
041.84	16	9	*7	*	*6	*	*6	*	*	*6	*
041.85	52	25	27	*6	*6	15	24	*8	18	18	*8

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
041.86	66	28	38	*	11	21	32	16	14	23	13
041.89	33	12	21	*	13	*8	10	*	11	11	*7
041.9	20	*	16	*	*8	*	*6	*	*	12	*
042	137	93	45	*	86	48	*	39	12	57	29
047	48	24	24	21	22	*	*	*7	11	18	12
047.9	43	22	21	18	20	*	*	*6	9	18	10
052	*7	*	*	*	*	*	*	*	*	*	*
053	61	25	37	*	*	14	41	13	17	18	14
053.1	21	*8	14	—	*	*	16	*	*7	*6	*
053.19	17	*6	11	—	*	*	13	*	*5	*	*
053.7	*5	*	*	*	*	*	*	*	*	*	*
053.9	33	13	20	*	*	*9	20	*8	*8	9	*7
054	59	14	46	*	40	*5	10	11	11	26	11
054.1	16	*	11	*	14	*	*	*	*	*7	*
054.10	10	*	*7	*	*8	*	*	*	*	*	*
054.7	*6	*	*	*	*	*	*	*	*	*	*
054.9	30	*5	25	*	22	*	*	*5	*	13	*7
057	*9	*	*	*6	*	*	*	*	*	*	*
057.9	*6	*	*	*	*	*	*	*	*	*	*
070	353	215	139	*	138	176	36	90	52	140	71
070.1	*7	*	*	*	*	*	*	*	*	*	*
070.3	62	40	22	*	24	30	*8	15	*8	27	12
070.30	30	17	12	*	13	14	*	*9	*	12	*
070.32	32	22	10	*	10	16	*5	*6	*	14	*8
070.4	10	*7	*	—	*	*8	*	*	*	*	*
070.44	*7	*	*	—	*	*5	*	*	*	*	*
070.5	270	163	107	*	109	134	25	71	42	103	54
070.51	128	78	50	*	60	57	10	30	20	54	24
070.54	141	85	56	*	48	77	15	41	21	49	30
075	16	*7	9	*5	10	*	*	*	*6	*6	*
078	19	10	9	*	12	*	*	*	*	10	*
078.1	*9	*	*6	*	*7	*	*	*	*	*5	*
078.5	*6	*	*	*	*	*	*	*	*	*	*
079	175	86	90	95	40	17	23	35	34	70	36
079.6	*8	*	*	*7	*	*	*	*	*	*	*
079.8	11	*	*7	*8	*	*	*	*	*	*8	*
079.89	11	*	*7	*8	*	*	*	*	*	*7	*
079.9	149	74	74	78	33	15	23	32	29	57	31
079.99	146	74	71	77	31	15	23	31	29	56	30
094	10	*6	*	—	*	*	*	*	*	*	*
094.0	*5	*	*	—	*	*	*	*	*	*	*
098	*6	*	*5	*	*	*	*	*	*	*	*
099	*7	*	*5	—	*6	*	*	*	*	*	*
110	56	31	25	*	11	13	30	15	15	20	*6
110.1	28	14	14	*	*	*5	21	9	*7	10	*
110.4	13	*9	*	*	*	*5	*	*	*	*	*
112	293	113	179	35	65	71	121	62	70	105	56
112.0	158	68	91	21	36	39	62	35	42	53	29
112.1	23	—	23	*	10	*7	*	*5	*5	*8	*
112.2	24	*9	16	*	*	*7	13	*5	*6	10	*
112.3	19	*6	13	*7	*	*	*6	*	*	*6	*6
112.4	*8	*	*	*	*	*	*5	*	*	*	*
112.5	13	*	*8	*	*	*	*5	*	*	*5	*
112.8	43	20	23	*	*8	10	23	9	*9	17	*8
112.84	34	16	17	*	*7	*8	19	*7	*7	14	*6
112.89	*8	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
117	24	15	9	*	*8	*7	*6	*	*	10	*7
117.3	*8	*	*	*	*	*	*	*	*	*	*
117.5	*6	*	*	—	*5	*	*	*	*	*	*
117.9	9	*	*	*	*	*	*	*	*	*	*
131	11	*	11	*	*8	*	*	*	*	*7	*
131.0	*8	*	*8	*	*	*	*	*	*	*	*
135	43	11	32	*	13	20	9	10	*8	18	*8
136	22	14	*9	*	12	*5	*	*	*	10	*
136.3	15	11	*	*	10	*	*	*	*	*7	*
136.9	*6	*	*	*	*	*	*	*	*	*	*
138	19	*9	11	—	*	*7	12	*	*	*7	*
140-239	4,362	1,834	2,528	103	676	1,479	2,104	1,003	932	1,603	824
141	*8	*6	*	*	*	*	*	*	*	*	*
150	29	20	9	—	*	*8	18	*7	*6	11	*6
150.9	16	12	*	—	*	*	10	*	*	*9	*
151	34	21	13	—	*	12	18	*8	*	15	*6
151.0	*9	*7	*	—	*	*	*5	*	*	*	*
151.2	*6	*	*	—	*	*	*	*	*	*	*
151.8	*5	*	*	—	*	*	*	*	*	*	*
151.9	*8	*	*	—	*	*	*5	*	*	*	*
152	*5	*	*	—	*	*	*	*	*	*	*
153	178	82	96	*	*6	51	120	38	43	65	32
153.0	*7	*	*	—	*	*	*6	*	*	*	*
153.1	*9	*	*6	—	*	*	*6	*	*	*	*
153.2	*7	*	*	*	*	*	*	*	*	*	*
153.3	38	18	20	—	*	12	25	*6	*7	17	*8
153.4	28	11	18	—	*	*5	22	*7	9	*5	*7
153.6	23	*8	15	—	*	*	19	*6	*	*8	*
153.8	*7	*	*	—	*	*	*	*	*	*	*
153.9	51	28	22	—	*	18	31	*9	14	21	*8
154	68	39	29	—	*	25	41	14	19	23	12
154.0	25	13	12	—	*	9	15	*	*7	9	*
154.1	38	24	14	—	*	14	23	*7	11	12	*8
155	34	25	9	*	*	16	16	10	*5	14	*5
155.0	23	17	*6	*	*	12	10	*	*	12	*
155.1	*8	*6	*	—	*	*	*6	*	*	*	*
156	14	*6	*8	—	*	*	10	*	*	*5	*
156.1	*5	*	*	—	—	*	*	*	*	*	*
157	54	30	24	*	*	17	35	14	15	18	*7
157.0	18	10	*8	—	*	*	13	*	*	*7	*
157.9	28	16	12	—	*	10	17	*6	9	*9	*
158	*8	*	*5	*	*	*	*	*	*	*	*
161	13	10	*	—	*	*	*6	*	*	*5	*
162	342	185	157	*	11	105	226	82	75	129	57
162.2	17	*8	*8	*	*	*5	9	*	*	*6	*
162.3	79	40	39	—	*	25	52	17	16	27	19
162.4	*5	*	*	—	*	*	*	*	*	*	*
162.5	38	18	20	—	*	11	26	10	*8	11	*8
162.8	27	14	13	—	*	*9	17	10	*6	*8	*
162.9	175	102	73	*	*	52	118	39	39	76	21
170	28	17	10	9	13	*	*	12	*	*8	*6
170.7	10	*8	*	*6	*	*	*	*7	*	*	*
170.9	*6	*	*	*	*	*	*	*	*	*	*
171	21	11	11	*	*	*6	*8	*5	*	*8	*
171.3	*9	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
172	10	*6	*	—	*	*	*	*	*	*	*
173	17	*8	*9	*	*	*5	11	*5	*	*6	*
174	161	—	161	*	18	69	74	33	39	59	30
174.1	*5	—	*5	—	*	*	*	*	*	*	*
174.4	15	—	15	—	*	*5	*7	*	*	*6	*
174.8	36	—	36	*	*	16	17	12	*6	10	*8
174.9	92	—	92	*	10	41	41	15	24	36	16
180	29	—	29	*	10	11	*8	*6	*	12	*7
180.8	*8	—	*8	—	*	*	*	*	*	*	*
180.9	17	—	17	*	*6	*	*6	*	*	*8	*
182	41	—	41	—	*	14	23	10	*8	14	*8
182.0	41	—	41	—	*	14	23	10	*8	14	*8
183	50	—	50	*	*7	20	23	10	12	21	*8
183.0	49	—	49	*	*6	19	22	9	12	19	*8
184	*5	—	*5	*	*	*	*	*	*	*	*
185	208	208	—	*	*	58	148	49	46	71	42
186	*7	*7	—	*	*6	*	*	*	*	*	*
186.9	*7	*7	—	*	*6	*	*	*	*	*	*
188	59	43	16	*	*	10	48	15	14	20	10
188.8	18	13	*5	*	*	*	15	*6	*	*5	*
188.9	25	18	*7	*	*	*	20	*6	*6	*8	*
189	55	38	17	*	*	19	30	10	13	24	*7
189.0	48	33	15	*	*	16	26	*8	13	21	*6
191	56	31	25	12	12	17	15	15	12	15	13
191.1	*8	*	*	*	*	*	*	*	*	*	*
191.3	*5	*	*	*	*	*	*	*	*	*	*
191.9	16	10	*6	*	*	*7	*	*	*	*7	*
192	*5	*	*	*	*	*	*	*	*	*	*
193	30	*8	22	*	11	9	*9	*5	*	12	*8
194	*6	*	*	*	*	*	*	*	*	*	*
195	12	*6	*6	*	*	*5	*5	*	*	*6	*
195.0	*6	*	*	—	*	*	*	*	*	*	*
196	171	74	97	*	18	67	85	39	36	63	34
196.0	20	13	*8	*	*	9	*8	*	*	*8	*
196.1	26	13	13	*	*	10	14	*6	*	10	*5
196.2	64	34	30	*	*6	22	36	15	16	22	12
196.3	35	*	33	—	*	15	16	*7	*7	15	*6
196.5	*6	*	*	—	*	*	*	*	*	*	*
196.6	10	*	*5	—	*	*	*5	*	*	*	*
196.9	*7	*	*	*	*	*	*	*	*	*	*
197	451	195	256	*	37	165	245	118	104	153	76
197.0	107	49	59	*	14	38	52	29	25	34	20
197.1	10	*	*6	*	*	*	*	*	*	*	*
197.2	51	19	32	*	*	14	34	13	12	15	11
197.4	9	*	*6	*	*	*	*5	*	*	*	*
197.5	12	*5	*7	*	*	*5	*6	*	*	*5	*
197.6	66	20	46	*	*7	27	33	13	14	24	15
197.7	178	86	92	*	10	65	102	48	45	62	24
197.8	15	*8	*7	*	*	*7	*7	*6	*	*	*
198	414	192	221	*5	33	157	219	101	89	150	74
198.1	10	*5	*	*	*	*	*7	*	*	*	*
198.2	*8	*	*	*	*	*	*5	*	*	*	*
198.3	98	47	50	*	*9	45	43	22	18	37	20
198.4	10	*	*6	*	*	*5	*	*	*	*	*
198.5	188	95	94	*	10	64	112	47	47	64	30
198.7	12	*7	*5	—	*	*6	*6	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
198.8	81	30	52	*	12	30	38	20	15	31	16
198.82	14	*	11	*	*	*7	*	*	*	*5	*
198.89	64	27	37	*	10	21	32	16	12	23	12
199	133	65	68	*	13	41	77	30	28	54	21
199.0	58	27	31	*	*8	18	31	13	10	24	12
199.1	75	38	37	*	*5	23	47	17	19	30	10
200	34	25	*8	*	11	11	10	*7	*9	*8	10
200.0	20	14	*6	*	*	10	*8	*	*6	*6	*
200.00	*7	*5	*	*	*	*	*	*	*	*	*
200.2	*8	*7	*	*	*7	*	*	*	*	*	*6
201	22	12	9	*	10	*5	*6	*6	*7	*6	*
201.9	17	*8	*9	*	*7	*	*	*6	*5	*	*
201.90	12	*6	*6	*	*6	*	*	*	*	*	*
202	128	71	56	*	16	35	75	31	28	42	27
202.1	*6	*	*	*	*	*	*	*	*	*	*
202.8	113	61	52	*	14	31	67	28	27	37	21
202.80	69	33	35	*	*7	17	44	18	18	20	13
202.81	*8	*5	*	*	*	*	*	*	*	*	*
202.82	*6	*	*	*	*	*	*	*	*	*	*
202.83	15	9	*6	*	*	*	*8	*	*	*6	*
202.88	9	*6	*	—	*	*	*	*	*	*	*
203	76	38	38	—	*	29	45	16	17	34	10
203.0	73	36	37	—	*	28	44	15	17	32	10
203.00	73	36	37	—	*	28	44	15	16	32	9
204	109	57	52	25	13	13	58	27	21	32	29
204.0	42	23	20	25	12	*	*	12	*5	12	13
204.00	29	16	13	15	*9	*	*	10	*	*9	*6
204.01	13	*7	*7	10	*	*	—	*	*	*	*7
204.1	67	34	32	—	*	10	56	14	16	20	16
204.10	64	34	30	—	*	9	54	14	15	20	15
205	57	33	24	*6	11	16	24	13	12	19	13
205.0	40	24	16	*	10	14	12	11	*6	14	9
205.00	32	18	14	*	*8	9	12	9	*6	10	*7
205.01	*8	*6	*	*	*	*	*	*	*	*	*
205.1	15	*9	*6	*	*	*	11	*	*	*5	*
205.10	13	*7	*6	*	*	*	9	*	*	*	*
208	*7	*	*	*	*	*	*5	*	*	*	*
210	*5	*	*	*	*	*	*	*	*	*	*
211	198	86	111	*	15	59	124	47	39	72	40
211.1	18	*6	12	—	*	*6	11	*	*	*7	*
211.2	*5	*	*	—	*	*	*	*	*	*	*
211.3	154	69	85	—	*9	44	101	37	31	55	31
211.4	9	*	*5	—	*	*	*6	*	*	*	*
212	*8	*	*6	*	*	*	*	*	*	*	*
213	*6	*	*	*	*	*	*	*	*	*	*
214	17	10	*6	*	*	*5	*8	*	*	*	*
216	13	*	10	*	*7	*	*	*	*	*	*
216.5	*7	*	*7	*	*	*	*	*	*	*	*
218	494	—	494	—	250	219	25	89	97	210	98
218.0	48	—	48	—	26	21	*	11	11	16	10
218.1	138	—	138	—	64	68	*6	29	25	59	26
218.2	75	—	75	—	34	40	*	15	13	33	15
218.9	233	—	233	—	126	91	16	34	49	103	48
219	*7	—	*7	—	*	*	*	*	*	*	*
220	74	—	74	*	35	30	*9	13	16	25	20

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
221 . . . . .	*5	—	*5	—	*	*	*	*	*	*	*
225 . . . . .	34	10	24	*	*	12	17	*8	*7	14	*
225.2 . . . . .	24	*6	18	*	*	*9	14	*6	*6	10	*
226 . . . . .	12	*	10	*	*	*	*	*	*	*	*
227 . . . . .	26	11	15	*	*7	*8	11	*	*	10	*9
227.0 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
227.1 . . . . .	11	*	*7	—	*	*	*6	*	*	*	*
227.3 . . . . .	10	*5	*	—	*	*	*	*	*	*	*
228 . . . . .	18	*6	12	*	*	*6	*	*	*	*5	*6
228.0 . . . . .	16	*5	11	*	*	*5	*	*	*	*	*6
228.04 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
230 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
233 . . . . .	39	*	37	—	16	14	*8	*8	*6	17	*7
233.0 . . . . .	16	—	16	—	*	*8	*	*	*	*	*
233.1 . . . . .	18	—	18	—	12	*	*	*	*	10	*
235 . . . . .	24	12	12	*	*	*6	15	*	*7	10	*
235.2 . . . . .	15	*8	*7	*	*	*	10	*	*	*7	*
236 . . . . .	*9	*	*7	*	*	*	*	*	*	*	*
237 . . . . .	16	11	*6	*	*	*5	*	*	*	*	*6
237.7 . . . . .	9	*7	*	*	*	*	*	*	*	*	*
237.70 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
238 . . . . .	76	38	39	*	*	16	55	22	18	26	10
238.4 . . . . .	17	9	*8	*	*	*7	9	*6	*	*6	*
238.7 . . . . .	52	26	27	*	*	*8	42	15	15	16	*7
239 . . . . .	26	12	14	*	*6	*	14	*6	*6	10	*
239.6 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
240-279 . . . . .	17,362	7,252	10,110	591	2,279	5,230	9,262	3,701	4,256	6,420	2,985
240 . . . . .	15	*	11	*	*6	*	*6	*	*6	*	*
240.9 . . . . .	14	*	11	*	*	*	*6	*	*6	*	*
241 . . . . .	42	*6	36	*	*9	14	20	*9	*	18	11
241.0 . . . . .	17	*	14	*	*5	*	*9	*	*	*6	*5
241.1 . . . . .	21	*	18	*	*	*8	*9	*	*	10	*
242 . . . . .	85	22	64	*	20	23	42	22	18	33	12
242.0 . . . . .	22	*6	15	*	9	*6	*6	*5	*	*8	*
242.00 . . . . .	20	*6	15	*	*8	*6	*6	*	*	*7	*
242.9 . . . . .	55	12	43	*	9	15	31	15	11	23	*6
242.90 . . . . .	52	12	40	*	*9	15	29	14	10	23	*6
244 . . . . .	1,529	313	1,217	*7	173	369	980	376	409	454	290
244.0 . . . . .	40	*8	32	*	*	10	25	*8	10	12	11
244.1 . . . . .	*8	*	*7	—	*	*	*	*	*	*	*
244.8 . . . . .	*5	*	*	*	*	*	*	*	*	*	—
244.9 . . . . .	1,472	299	1,173	*6	164	354	948	364	395	437	277
245 . . . . .	16	*	16	*	*6	*7	*	*	*	*	*
245.2 . . . . .	12	*	12	—	*	*5	*	*	*	*	*
246 . . . . .	14	*	11	—	*	*	*6	*	*	*6	*
246.9 . . . . .	11	*	*9	—	*	*	*	*	*	*	*
250 . . . . .	5,117	2,360	2,757	43	572	1,774	2,728	1,130	1,166	2,004	816
250.0 . . . . .	3,728	1,671	2,056	19	329	1,247	2,132	837	869	1,472	550
250.00 . . . . .	3,019	1,373	1,646	*	221	998	1,796	683	703	1,158	475
250.01 . . . . .	375	161	214	12	58	126	180	93	82	162	37
250.02 . . . . .	273	109	164	*	30	102	141	46	66	128	32
250.03 . . . . .	60	28	32	*	21	22	15	15	17	23	*5
250.1 . . . . .	146	74	72	17	87	28	14	25	33	56	33
250.10 . . . . .	21	10	11	*	10	*8	*	*	*	*7	*6
250.11 . . . . .	66	37	30	13	44	*8	*	13	15	24	14
250.12 . . . . .	19	*8	10	*	*8	*6	*	*	*	*7	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
250.13	40	19	21	*	25	*6	*	*5	*9	17	9
250.2	25	12	12	*	*	10	11	*9	*	*9	*
250.20	9	*	*5	*	*	*6	*	*	*	*	*
250.22	11	*6	*	—	*	*	*6	*	*	*	*
250.3	*8	*	*5	*	*	*	*	*	*	*	*
250.4	306	161	145	—	33	131	142	63	59	112	72
250.40	192	103	89	—	*7	81	103	42	34	64	53
250.41	71	36	35	—	19	32	19	13	13	32	13
250.42	30	15	15	—	*	12	17	*5	*9	12	*
250.43	13	*7	*6	—	*	*6	*	*	*	*5	*
250.5	139	60	79	—	21	65	53	26	29	57	28
250.50	73	33	41	—	*5	33	35	15	14	26	18
250.51	41	16	25	—	11	20	11	*7	10	20	*
250.52	15	*8	*7	—	*	*8	*5	*	*	*8	*
250.53	10	*	*6	—	*	*	*	*	*	*	*
250.6	375	181	194	*	57	145	170	80	90	146	58
250.60	214	102	112	*	17	75	121	45	54	76	38
250.61	84	42	42	*	22	34	26	19	17	37	11
250.62	49	24	25	—	*7	26	16	9	13	22	*5
250.63	28	13	15	*	11	11	*6	*7	*7	11	*
250.7	95	60	35	—	*	36	55	23	20	34	18
250.70	60	37	22	—	*	22	37	12	13	20	15
250.71	20	11	*8	—	*	9	9	*5	*	9	*
250.72	11	*8	*	—	*	*	*6	*	*	*	*
250.8	247	116	130	*	25	90	131	61	52	90	44
250.80	151	67	84	*	*8	51	93	36	31	53	31
250.81	44	22	22	*	*8	20	15	12	*8	19	*
250.82	34	18	16	—	*5	13	16	*8	*8	14	*
250.83	17	9	*8	*	*	*5	*8	*	*	*	*
250.9	48	21	27	*	11	18	18	*6	10	24	9
250.90	13	*7	*6	*	*	*6	*	*	*	*8	*
250.92	23	10	14	*	*5	*8	9	*	*6	10	*
250.93	10	*	*7	*	*	*	*	*	*	*5	*
251	75	29	45	9	15	20	30	21	18	24	12
251.1	*6	*	*	*	*	*	*	*	*	*	*
251.2	31	13	18	*6	*	*7	13	*6	*6	12	*6
251.8	33	13	21	*	*8	10	15	12	9	*8	*
252	39	12	27	*	*8	12	18	*7	*9	13	10
252.0	25	*9	16	*	*7	10	*9	*6	*6	9	*
252.1	13	*	10	*	*	*	*9	*	*	*	*5
253	82	37	45	*6	11	20	46	19	22	27	15
253.2	14	10	*	*	*	*	*6	*	*	*6	*
253.5	10	*	*6	*	*	*	*	*	*	*	*
253.6	49	18	30	*	*	*8	37	12	15	12	10
255	72	30	42	*	14	21	33	18	15	27	12
255.0	*8	*	*7	*	*	*	*	*	*	*	*
255.4	47	21	26	*	*8	14	23	13	10	18	*5
255.9	*7	*	*	*	*	*	*	*	*	*	*
256	16	—	16	*	13	*	*	*	*	*7	*
256.4	11	—	11	—	10	*	—	*	*	*	*
260	11	*	*8	*	*	*	*9	*	*	*	*
261	26	12	14	*	*	*7	16	*5	*	11	*5
262	*6	*	*	*	*	*	*	*	*	*	*
263	379	171	208	*8	37	76	258	76	99	153	51
263.8	23	11	12	*	*	*	16	*	*7	10	*
263.9	347	155	193	*5	32	71	238	69	90	140	49
266	40	16	24	*	*	*5	32	11	*7	15	*7
266.2	37	14	23	*	*	*	31	10	*7	14	*6

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
269	12	*6	*6	*	*	*	*6	*	*	*5	*
269.9	*6	*	*	—	*	*	*	*	*	*	*
270	16	11	*	*	*	*8	*	*7	*	*	*
271	19	*6	12	*	*	*	10	*	*5	*	*5
271.3	17	*6	11	*	*	*	*9	*	*	*	*
272	2,731	1,413	1,318	*	198	1,042	1,487	663	675	933	460
272.0	1,115	556	559	*	82	441	591	351	263	331	170
272.1	43	24	18	*	13	19	10	*	10	20	*8
272.2	13	*6	*6	*	*	*	*8	*	*	*	*
272.4	1,509	797	712	*	95	558	855	290	387	561	271
272.5	*6	*	*	*	*	*	*	*	*	*	*
272.8	14	10	*	*	*	*6	*6	*	*	*6	*
272.9	26	15	11	—	*	11	14	*8	*9	*6	*
273	55	23	32	*	*8	16	27	12	*9	25	9
273.1	*7	*	*	—	*	*	*6	*	*	*	*
273.8	41	16	25	*	*7	11	18	*7	*6	20	*7
274	232	161	71	—	20	63	149	52	64	76	42
274.0	49	36	13	—	*5	13	31	10	13	16	11
274.9	178	121	57	—	15	48	115	40	48	59	30
275	258	95	163	*7	51	76	123	45	76	97	39
275.0	11	*8	*	*	*	*	*	*	*	*	*
275.2	96	33	63	*	24	28	44	18	32	41	*6
275.3	33	14	19	*	*9	*9	13	*	9	12	*7
275.4	116	40	77	*	15	36	61	20	32	40	25
275.41	54	18	35	*	11	19	21	9	14	18	12
275.42	52	16	36	*	*	16	33	*9	14	19	10
275.49	10	*	*	*	*	*	*7	*	*	*	*
276	5,296	2,095	3,200	448	714	1,160	2,974	979	1,368	2,014	935
276.0	123	54	69	*7	*9	20	86	27	31	44	21
276.1	848	325	523	19	73	213	543	138	235	319	156
276.2	269	124	146	40	32	68	130	38	51	116	64
276.3	51	23	27	*	*9	11	27	*7	12	20	11
276.4	29	13	16	*	*	*7	17	*	*6	10	*9
276.5	2,455	1,004	1,451	344	341	417	1,352	478	644	916	417
276.6	90	42	48	*	12	35	41	10	22	40	17
276.7	400	184	216	*7	35	104	254	83	98	142	78
276.8	943	299	644	20	191	258	474	185	233	373	153
276.9	88	28	60	*	*9	27	49	10	37	32	9
277	53	28	24	14	24	*7	*8	14	13	14	11
277.0	26	13	13	*9	16	*	*	*8	*7	*7	*
277.00	26	13	13	*8	16	*	*	*8	*7	*7	*
277.3	*5	*	*	*	*	*	*	*	*	*	*
277.4	*7	*5	*	*	*	*	*	*	*	*	*
278	1,086	374	712	17	352	482	234	202	245	421	217
278.0	1,066	369	697	17	345	474	230	198	241	413	214
278.00	673	246	428	14	197	293	169	126	153	247	147
278.01	393	123	270	*	147	181	61	71	88	166	68
278.1	11	*	10	—	*	*6	*	*	*	*	*
278.8	9	*	*	*	*	*	*	*	*	*	*
279	24	10	14	*6	*6	*7	*	*	*5	10	*
279.0	13	*7	*6	*	*	*	*	*	*	*	*
280-289	4,603	1,846	2,757	186	940	1,056	2,422	976	1,064	1,742	821
280	747	255	491	14	134	172	427	158	181	276	131
280.0	420	155	265	*	70	101	246	88	102	159	72
280.8	9	*	*6	*	*	*	*	*	*	*	*
280.9	317	98	219	10	62	69	176	69	78	112	58

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
281	85	37	48	*	*7	17	60	19	26	26	14
281.0	26	11	15	—	*	*	21	*8	*8	*8	*
281.1	12	*	*7	—	*	*	*8	*	*	*	*
281.9	44	20	24	*	*5	10	30	*8	13	13	10
282	149	62	87	41	81	19	*7	46	28	65	11
282.4	15	*	11	*	*6	*	*	*6	*	*	*
282.5	21	*	17	*	12	*	*	*5	*	*8	*
282.6	106	50	56	33	62	10	*	32	20	50	*
282.60	17	11	*6	*6	*8	*	*	*	*	9	*
282.61	*8	*	*	*5	*	*	*	*	*	*	*
282.62	80	35	46	22	52	*7	*	25	14	38	*
283	29	15	14	*	*	*6	15	*5	*7	10	*7
283.0	14	*7	*6	*	*	*	*8	*	*	*	*
283.9	11	*7	*	*	*	*	*6	*	*	*	*
284	183	97	86	12	29	60	82	45	36	65	37
284.8	169	90	79	11	26	55	77	40	35	58	36
284.9	13	*6	*7	*	*	*	*5	*	*	*6	*
285	2,374	854	1,520	39	475	503	1,358	478	564	900	432
285.1	616	221	394	*6	103	130	376	109	197	194	116
285.2	248	114	133	*	36	65	144	51	55	96	45
285.21	80	41	39	*	13	27	39	13	15	35	17
285.22	47	26	21	*	*	14	29	12	13	18	*
285.29	120	47	73	*	19	24	76	27	27	43	24
285.8	37	13	24	*	10	*	21	*5	*8	12	12
285.9	1,472	505	967	29	325	302	816	313	304	597	258
286	218	108	110	*9	39	64	106	50	46	85	37
286.0	*6	*6	*	*	*	*	*	*	*	*	*
286.3	16	*8	*9	*	*	*6	*7	*6	*	*5	*
286.4	*9	*	*7	*	*	*	*	*	*	*	*
286.5	12	*	*7	—	*	*	*7	*	*	*6	*
286.6	20	10	10	*	*	*7	*9	*	*	*7	*
286.7	41	23	18	*	*	11	25	11	*7	17	*6
286.9	112	54	57	*	21	34	54	21	26	44	21
287	389	214	176	20	67	109	194	87	77	152	74
287.0	*5	*	*	*	*	*	*	*	*	*	*
287.3	36	11	25	*8	*9	*6	13	10	*6	13	*8
287.4	56	34	23	*	11	19	26	14	14	19	10
287.5	288	164	124	*8	45	81	154	60	55	119	54
288	320	151	169	31	77	77	135	67	72	125	56
288.0	153	78	75	20	33	37	63	36	30	59	28
288.3	*6	*	*	*	*	*	*	*	*	*	*
288.8	157	68	88	10	42	36	68	28	40	62	26
289	109	54	55	16	27	30	36	22	27	38	22
289.0	10	*6	*	—	*	*5	*	*	*	*	*
289.2	10	*6	*	*6	*	*	*	*	*	*	*
289.3	*8	*	*	*5	*	*	*	*	*	*	*
289.4	*6	*	*	*	*	*	*	*	*	*	*
289.5	10	*	*6	*	*	*	*	*	*	*	*
289.59	*8	*	*	*	*	*	*	*	*	*	*
289.8	35	16	20	*	9	11	15	*6	*8	14	*8
289.9	25	*9	16	*	*	*	13	*6	*6	9	*
290-319	11,398	5,428	5,970	387	4,991	3,207	2,814	2,791	2,719	4,095	1,793
290	234	84	150	—	*	*9	225	74	55	78	28
290.0	106	30	76	—	*	*	105	38	23	32	12
290.2	19	*6	13	—	—	*	19	*7	*6	*5	*
290.20	9	*	*6	—	—	*	9	*	*	*	*
290.21	10	*	*7	—	—	*	10	*	*	*	—
290.3	14	*6	*8	—	—	*	14	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
290.4	88	39	49	—	*	*5	83	22	20	34	12
290.40	68	31	37	—	*	*	63	16	13	29	10
290.41	*7	*	*	—	—	*	*6	*	*	*	*
290.42	*7	*	*	—	—	*	*7	*	*	*	*
290.43	*7	*	*	—	*	*	*6	*	*	*	*
291	306	232	73	*	142	123	40	73	100	74	59
291.0	58	48	9	—	21	27	9	12	12	22	11
291.1	*5	*	*	—	*	*	*	*	*	*	*
291.2	16	12	*	*	*	*8	*7	*	*6	*	*
291.3	*5	*	*	—	*	*	*	*	*	*	*
291.8	219	162	58	*	114	84	21	55	76	46	42
291.81	199	148	50	*	99	78	21	53	60	44	41
291.89	21	13	*7	*	15	*5	*	*	16	*	*
292	214	116	98	*	106	49	58	67	52	65	31
292.0	87	51	36	*	59	22	*6	37	18	22	10
292.1	14	*8	*6	*	*6	*	*6	*	*	*	*
292.12	11	*5	*5	*	*	*	*6	*	*	*	*
292.8	102	52	50	*	36	21	45	26	28	33	15
292.81	61	28	33	*	*6	13	41	11	17	22	10
292.84	35	21	14	*	26	*7	*	13	10	9	*
292.89	*5	*	*	—	*	*	*	*	*	*	*
292.9	11	*6	*5	—	*6	*	*	*	*	*6	*
293	121	56	66	*	13	25	83	28	44	31	18
293.0	82	38	44	*	*	16	64	19	27	21	14
293.8	27	12	14	*	9	*7	10	*5	14	*6	*
293.83	19	9	9	*	*7	*5	*6	*	9	*	*
293.9	11	*	*7	*	*	*	9	*	*	*	*
294	813	285	528	*	11	32	769	189	203	293	128
294.1	322	112	210	*	*	10	310	82	84	106	51
294.10	263	90	172	*	*	*7	253	70	64	85	44
294.11	60	22	38	*	*	*	56	12	20	21	*7
294.8	462	160	302	*	*	16	441	97	116	179	70
294.9	26	11	15	*	*	*5	17	9	*	*8	*7
295	529	274	255	*5	274	178	72	150	103	206	70
295.3	133	85	48	*	70	49	14	44	23	51	16
295.30	58	35	23	—	28	23	*7	11	13	25	10
295.32	31	19	12	—	15	11	*	11	*	15	*
295.34	42	29	13	*	26	14	*	21	*6	10	*
295.4	*5	*	*	—	*	*	—	*	*	*	*
295.6	27	14	13	—	10	11	*6	*	*8	12	*
295.60	*7	*	*	—	*	*	*	*	*	*	*
295.62	10	*	*6	—	*	*	*	*	*	*6	*
295.64	*8	*	*	—	*	*	*	*	*	*	*
295.7	234	110	125	*	141	67	20	60	46	97	31
295.70	180	84	96	*	109	50	16	34	41	80	25
295.72	*7	*	*	—	*	*	*	*	*	*	*
295.74	46	23	24	*	28	15	*	23	*	14	*6
295.9	121	57	63	*	43	46	31	37	23	42	18
295.90	105	47	58	*	36	40	29	27	22	38	18
295.92	*7	*	*	—	*	*	*	*	*	*	*
295.94	*8	*	*	—	*	*	*	*7	*	*	*
296	1,476	595	881	108	822	390	157	324	293	727	133
296.2	355	144	211	31	194	77	52	62	69	193	31
296.20	194	77	117	17	107	42	28	34	46	97	17
296.22	36	14	22	*	24	*6	*	*	*	27	*
296.23	70	29	41	*7	34	17	11	15	*9	41	*5
296.24	48	21	27	*	25	10	*8	10	*8	28	*
296.3	394	153	241	13	219	117	45	102	95	171	26
296.30	99	39	60	*	52	29	16	26	27	40	*
296.32	22	*7	15	*	13	*7	*	*	*5	9	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
296.33	178	65	113	*7	105	52	14	52	41	75	11
296.34	89	40	49	*	46	28	12	18	21	44	*6
296.4	85	37	48	*	52	23	*6	22	12	43	*8
296.40	37	16	21	*	24	*9	*	*6	*5	22	*
296.43	*7	*	*	*	*5	*	*	*	*	*	*
296.44	38	17	21	*	23	12	*	12	*6	18	*
296.5	132	44	88	*	82	40	*6	33	19	69	11
296.50	68	26	43	*	45	18	*	14	14	34	*7
296.53	34	9	25	*	20	12	*	14	*	18	*
296.54	26	*8	17	*	16	*8	*	*	*	16	*
296.6	117	46	71	17	71	25	*	19	14	73	11
296.60	58	20	38	*	37	15	*	*8	*9	36	*6
296.63	27	10	17	*8	15	*	*	*6	*	20	*
296.64	30	15	15	*	17	*6	*	*6	*	16	*
296.7	241	94	147	11	119	76	36	60	63	83	35
296.8	93	45	48	19	51	20	*	11	10	67	*5
296.80	63	27	36	16	29	15	*	*	*6	50	*
296.89	27	16	11	*	20	*	*	*5	*	16	*
296.9	56	31	25	10	32	10	*	15	10	27	*
296.90	48	26	23	10	26	9	*	10	9	26	*
296.99	*7	*	*	*	*6	*	*	*	*	*	*
297	28	9	19	—	*6	*5	17	*8	*7	*9	*
297.1	10	*	*7	—	*	*	*6	*	*	*	*
297.9	17	*	12	—	*	*	11	*6	*	*5	*
298	207	102	105	*	70	40	94	52	40	85	30
298.9	202	100	102	*	68	39	92	51	38	83	30
299	20	14	*6	11	*7	*	*	*6	*	*6	*
299.0	12	*9	*	*6	*	*	*	*	*	*	*
299.00	12	*9	*	*6	*	*	*	*	*	*	*
299.8	*6	*	*	*	*	*	—	*	*	*	*
299.80	*6	*	*	*	*	*	—	*	*	*	*
300	1,090	341	749	23	369	349	348	251	276	382	180
300.0	595	176	419	*7	163	189	235	137	159	202	97
300.00	491	142	349	*5	114	157	214	115	133	160	82
300.01	75	24	51	*	37	23	14	16	18	31	11
300.02	29	10	19	*	12	9	*7	*6	*8	11	*
300.1	28	*5	22	*	17	*9	*	*6	*5	11	*
300.11	14	*	10	*	*6	*6	*	*	*	*6	*
300.14	*6	*	*6	—	*	*	*	*	*	*	*
300.2	24	*7	17	*	16	*6	*	*5	*5	*7	*6
300.21	13	*	10	—	*9	*	*	*	*	*	*
300.23	*5	*	*	*	*	*	*	*	*	*	*
300.3	34	16	18	*	21	10	*	10	*6	12	*6
300.4	311	90	221	*7	90	107	106	78	71	116	46
300.8	*6	*	*	*	*	*	*	*	*	*	*
300.9	89	43	46	*	58	25	*	14	25	32	19
301	341	144	197	*	239	89	9	138	97	69	37
301.2	*6	*	*	—	*	*	*	*	*	*	*
301.5	*8	*	*6	*	*	*	*	*	*	*	*
301.50	*6	*	*6	*	*	*	*	*	*	*	*
301.6	22	*7	15	*	13	*9	*	*	10	*5	*5
301.7	31	25	*6	*	26	*	*	*8	13	*	*6
301.8	136	36	100	*	97	34	*	51	50	24	11
301.81	11	*8	*	*	*6	*	*	*	*	*	*
301.83	104	15	89	*	76	25	*	36	39	20	*8
301.89	15	9	*6	*	12	*	*	9	*	*	*
301.9	126	65	61	*	87	34	*	73	16	27	10
302	*7	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
303	754	543	211	*	364	307	82	190	201	233	131
303.0	140	103	38	*	71	55	13	18	54	31	38
303.00	60	42	18	*	32	21	*6	*	21	15	21
303.01	77	59	18	*	37	33	*7	14	31	16	16
303.9	614	441	173	*	292	252	69	171	147	202	93
303.90	269	191	78	*	119	109	41	44	57	126	42
303.91	274	200	73	*	140	112	21	108	64	65	37
303.92	23	17	*6	—	15	*6	*	*7	*9	*	*5
303.93	49	32	16	—	18	26	*	13	17	9	*9
304	521	304	217	*	386	118	15	165	100	187	69
304.0	176	98	79	*	120	52	*	72	25	46	33
304.00	54	25	28	*	36	16	*	*8	12	24	10
304.01	116	69	47	—	80	33	*	61	12	22	21
304.03	*6	*	*	—	*	*	—	*	*	*	*
304.1	28	12	16	*	17	*7	*	*6	*	12	*
304.10	14	*6	*8	*	*9	*	*	*	*	*8	*
304.11	13	*5	*7	—	*8	*	*	*	*	*	*
304.2	95	58	36	*	73	21	*	32	15	40	*8
304.20	46	29	17	*	35	10	*	*7	*8	28	*
304.21	41	26	16	*	32	9	*	21	*	11	*
304.3	68	44	24	*	60	*7	*	14	32	17	*
304.30	24	15	9	*	21	*	—	*	*9	*9	*
304.31	36	25	12	*	32	*	*	10	17	*8	*
304.4	16	*9	*7	—	14	*	*	*	*	*	*7
304.40	9	*	*	—	*8	*	*	*	*	*	*5
304.41	*6	*	*	—	*5	*	*	*	*	*	*
304.6	11	*6	*5	—	*6	*	*	*	*	*	*
304.61	*6	*	*	—	*	*	*	*	*	*	*
304.7	32	18	14	*	23	*7	*	17	*	*9	*
304.70	*6	*	*	*	*	*	*	*	*	*	*
304.71	25	15	10	*	17	*5	*	15	*	*6	*
304.8	36	24	12	*	30	*5	—	11	*6	16	*
304.80	23	14	9	*	19	*	—	*6	*	14	*
304.81	10	*8	*	—	9	*	—	*	*	*	*
304.9	59	35	24	*	42	14	*	*9	*	40	*7
304.90	47	29	18	*	34	12	*	*	*	36	*5
304.91	*7	*	*	—	*6	*	*	*	*	*	*
305	2,794	1,534	1,259	16	1,404	1,013	361	538	654	1,038	565
305.0	467	327	140	*	239	167	57	117	98	163	89
305.00	314	220	95	*	164	105	41	63	78	116	57
305.01	98	73	25	*	49	39	10	29	13	33	23
305.02	26	15	11	*	15	10	*	17	*	*	*
305.03	29	20	9	*	11	13	*	*7	*	11	*7
305.1	1,707	846	862	*	697	709	298	261	441	661	345
305.2	157	96	61	*5	131	21	*	49	33	49	27
305.20	101	57	44	*	84	13	*	20	28	37	16
305.21	34	23	11	*	27	*6	*	15	*	*7	*8
305.22	14	10	*	*	12	*	—	12	*	*	*
305.23	*8	*6	*	*	*7	*	—	*	*	*	*
305.3	*6	*	*	*	*5	*	—	*	*	*	*
305.4	17	*8	*9	—	13	*	*	*6	*	*5	*
305.40	11	*	*7	—	*8	*	*	*	*	*	*
305.41	*5	*	*	—	*	*	—	*	*	*	*
305.5	56	36	20	*	36	19	*	17	11	16	12
305.50	36	21	15	*	22	13	*	*8	9	12	*7
305.51	15	11	*	—	11	*	*	*7	*	*	*
305.6	199	118	80	*	140	56	*	54	40	66	39
305.60	135	75	60	*	94	39	*	26	35	51	25
305.61	41	29	13	—	28	13	*	16	*	12	11
305.62	13	10	*	—	11	*	—	10	*	*	*
305.63	*8	*	*	—	*7	*	—	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
305.7	35	17	18	*	27	*8	*	*	*	10	23
305.70	24	12	11	*	18	*6	—	*	*	*7	14
305.71	*6	*	*	—	*5	*	*	*	*	*	*
305.9	147	80	67	*	114	26	*	33	25	64	25
305.90	114	61	53	*	92	17	*	18	23	55	19
305.91	12	*7	*	*	*9	*	*	*6	*	*	*
305.93	16	*8	*8	*	10	*5	*	*6	*	*	*
306	9	*	*7	*	*	*	*	*	*	*	*
307	67	19	48	*5	35	12	14	18	16	20	13
307.1	10	*	9	*	*8	*	*	*5	*	*	*
307.2	*6	*	*	*	*	*	*	*	*	*	*
307.5	15	*	14	*	12	*	*	*5	*	*	*
307.50	*6	*	*6	*	*	*	*	*	*	*	*
307.51	*6	*	*6	—	*6	*	—	*	*	*	*
307.8	13	*	10	*	*8	*	*	*	*	*	*
307.81	11	*	*8	*	*6	*	*	*	*	*	*
307.9	18	*6	12	*	*	*	11	*	*	*6	*
308	22	*9	13	*	11	*7	*	*	*5	11	*
308.3	*8	*	*	*	*	*	*	*	*	*	*
308.9	*9	*	*6	*	*	*	*	*	*	*	*
309	293	114	179	23	179	63	28	102	77	69	45
309.0	67	30	37	*	40	12	12	21	19	15	12
309.2	31	12	19	*	16	*7	*8	*8	12	*8	*
309.28	26	11	16	*	14	*5	*6	*6	11	*6	*
309.4	25	13	13	*	18	*	*	15	*	*	*
309.8	141	44	97	15	87	36	*	51	36	34	20
309.81	136	42	93	14	84	35	*	50	36	30	19
309.89	*5	*	*	*	*	*	*	*	*	*	*
309.9	20	10	9	*	14	*	*	*	*	*	*
310	49	21	28	*	*7	11	29	10	12	18	*9
310.1	13	*7	*6	*	*	*	*	*	*	*	*5
310.2	11	*5	*5	*	*	*	*	*	*	*	*
310.9	24	*8	16	*	*	*	22	*5	*	12	*
311	1,094	376	718	22	360	330	381	284	274	361	175
312	72	44	28	20	43	*6	*	27	19	18	*8
312.3	46	28	18	9	30	*	*	21	12	*9	*
312.30	25	16	*9	*6	17	*	*	12	*8	*	*
312.34	12	9	*	*	*8	*	*	*6	*	*	*
312.8	*9	*	*	*	*	*	*	*	*	*	*
312.89	*6	*	*	*	*	*	*	*	*	*	*
312.9	16	12	*	*7	*7	*	*	*	*6	*	*
313	43	24	19	28	15	*	—	9	12	16	*6
313.8	41	22	19	27	14	*	—	*8	12	16	*5
313.81	37	20	17	24	13	—	—	*7	11	14	*
314	94	65	29	52	40	*	*	21	22	42	10
314.0	91	64	26	51	37	*	*	21	22	38	10
314.00	21	16	*6	10	11	*	*	*	*6	*8	*
314.01	69	49	20	41	26	*	—	16	16	30	*7
315	50	30	20	31	16	*	*	11	14	11	14
315.2	*8	*5	*	*	*	*	*	*	*	*	*
315.3	*5	*	*	*	*	—	—	*	*	*	*
315.8	13	*7	*6	12	*	*	*	*	*	*	*
315.9	19	10	*9	*9	*8	*	*	*	*5	*	*5
317	31	18	13	*	17	9	*	14	*7	*8	*
318	29	15	13	*	13	10	*	14	*6	*7	*
318.0	*6	*	*	*	*	*	*	*	*	*	*
318.1	15	10	*5	*	*5	*6	*	*8	*	*	*
318.2	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
319 . . . . .	89	51	38	17	34	23	15	24	23	27	14
320-389 . . . . .	3,762	1,630	2,132	409	654	829	1,870	813	914	1,289	746
320 . . . . .	14	*7	*7	*6	*	*	*	*	*	*6	*
321 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
322 . . . . .	12	*7	*	*	*	*	*	*	*	*	*
322.9 . . . . .	10	*5	*	*	*	*	*	*	*	*	*
323 . . . . .	14	*7	*7	*	*	*	*	*	*	*6	*
323.9 . . . . .	9	*	*	*	*	*	*	*	*	*	*
324 . . . . .	14	*8	*7	*	*	*7	*	*	*	*7	*
324.0 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
324.1 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
331 . . . . .	477	177	301	21	16	21	419	112	121	169	75
331.0 . . . . .	385	129	256	—	*	*8	376	93	95	143	54
331.3 . . . . .	14	9	*	*	*	*	11	*	*	*	*
331.4 . . . . .	63	30	33	19	13	10	21	14	15	17	17
331.9 . . . . .	13	*8	*	*	*	*	*9	*	*6	*	*
332 . . . . .	256	134	123	*	*	21	233	68	58	83	46
332.0 . . . . .	255	133	122	*	*	21	232	68	58	83	46
333 . . . . .	87	35	52	*	14	20	51	22	28	22	16
333.0 . . . . .	*6	*	*	*	*	*	*5	*	*	*	*
333.1 . . . . .	26	12	14	*	*	*	19	*6	9	*7	*
333.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
333.8 . . . . .	*9	*	*5	—	*	*	*6	*	*	*	*
333.82 . . . . .	*8	*	*	—	*	*	*5	*	*	*	*
333.9 . . . . .	31	10	20	*	*6	9	15	*8	11	*7	*5
333.90 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
333.99 . . . . .	22	*7	15	*	*	*7	12	*5	*8	*5	*
334 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
335 . . . . .	11	*5	*6	*	*	*	*6	*	*	*	*
335.2 . . . . .	9	*	*5	—	*	*	*6	*	*	*	*
335.20 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
336 . . . . .	18	11	*7	*	*	*6	*8	*	*	*7	*
336.3 . . . . .	*8	*5	*	—	*	*	*	*	*	*	*
337 . . . . .	88	39	49	*	29	29	27	20	15	39	14
337.1 . . . . .	63	29	34	*	20	20	22	13	10	31	9
337.2 . . . . .	14	*	11	*	*8	*6	*	*5	*	*	*
337.22 . . . . .	*6	*	*	*	*	*	—	*	*	*	*
337.9 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
340 . . . . .	99	23	75	*	22	52	25	26	29	27	17
342 . . . . .	205	91	114	*6	14	45	140	29	56	63	57
342.0 . . . . .	10	*	*6	—	*	*	*8	*	*	*	*
342.00 . . . . .	*7	*	*5	—	*	*	*7	*	*	*	*
342.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
342.9 . . . . .	188	83	105	*5	13	42	128	27	50	58	53
342.90 . . . . .	152	67	85	*	11	34	103	22	40	49	42
342.91 . . . . .	18	*8	10	*	*	*	15	*	*	*5	*7
342.92 . . . . .	18	*8	10	*	*	*5	11	*	*7	*	*
343 . . . . .	83	41	42	42	27	9	*	22	20	22	19
343.2 . . . . .	12	*	*7	*8	*	*	*	*	*	*	*
343.9 . . . . .	61	32	30	30	20	*8	*	16	15	17	13
344 . . . . .	125	81	44	*6	49	38	32	23	28	52	22
344.0 . . . . .	54	39	15	*	23	16	12	10	13	24	*7
344.00 . . . . .	47	34	13	*	20	13	11	*9	11	22	*
344.1 . . . . .	41	30	11	*	20	13	*8	*8	*9	16	*8
344.6 . . . . .	9	*	*	*	*	*	*	*	*	*	*
344.60 . . . . .	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
344.9 . . . . .	12	*5	*7	*	*	*	*	*	*	*	*
345 . . . . .	111	47	64	32	39	25	16	27	28	33	23
345.1 . . . . .	15	*6	*9	*	*7	*	*	*	*	*	*
345.10 . . . . .	13	*6	*8	*	*7	*	*	*	*	*	*
345.3 . . . . .	20	9	11	*7	*	*	*	*	*	*8	*
345.4 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
345.5 . . . . .	12	*6	*6	*	*	*	*	*6	*	*	*
345.50 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
345.9 . . . . .	45	18	27	12	17	10	*6	*8	15	14	*8
345.90 . . . . .	40	16	24	10	15	10	*6	*6	14	13	*7
346 . . . . .	213	51	163	*6	110	81	17	46	49	77	42
346.0 . . . . .	*8	*	*6	—	*	*	*	*	*	*	*
346.00 . . . . .	*7	*	*6	—	*	*	*	*	*	*	*
346.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
346.8 . . . . .	12	*	11	*	*7	*	*	*	*	*	*
346.80 . . . . .	11	*	9	*	*6	*	*	*	*	*	*
346.9 . . . . .	184	44	140	*	96	70	13	38	43	70	32
346.90 . . . . .	171	41	130	*	90	63	13	37	41	62	30
346.91 . . . . .	13	*	10	*	*6	*7	*	*	*	*7	*
347 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
348 . . . . .	172	86	86	23	35	47	66	30	40	61	41
348.0 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
348.1 . . . . .	56	31	25	*6	*8	18	23	11	12	20	14
348.2 . . . . .	*7	*	*6	*	*6	*	*	*	*	*	*
348.3 . . . . .	59	27	32	*6	*	16	32	10	13	22	13
348.4 . . . . .	10	*	*	*	*	*	*	*	*	*	*
348.5 . . . . .	*9	*5	*	*	*	*	*	*	*	*	*
348.8 . . . . .	21	11	10	*	*9	*5	*	*	*	*7	*6
349 . . . . .	54	14	40	*	32	10	10	*8	12	21	13
349.0 . . . . .	31	*	27	*	26	*	*	*5	*7	13	*6
349.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
349.8 . . . . .	12	*	*8	*	*	*	*6	*	*	*	*
349.82 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
350 . . . . .	20	*7	13	*	*	*	14	*	*6	*7	*
350.1 . . . . .	19	*6	13	*	*	*	14	*	*6	*7	*
351 . . . . .	30	11	19	*	*9	*9	11	10	*6	10	*
351.0 . . . . .	28	10	18	*	*8	*8	11	10	*5	*9	*
353 . . . . .	11	*6	*5	*	*5	*	*	*	*	*	*
353.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
354 . . . . .	34	12	23	*	*8	14	12	*8	*8	9	*9
354.0 . . . . .	22	*6	16	—	*6	*8	*8	*	*6	*6	*6
354.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
355 . . . . .	54	26	28	*	10	17	27	11	12	19	11
355.8 . . . . .	24	11	14	*	*	*9	12	*	*7	*7	*6
355.9 . . . . .	18	*9	10	*	*	*6	9	*	*	*8	*
356 . . . . .	98	43	55	*	12	26	59	15	18	40	25
356.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
356.9 . . . . .	86	39	47	*	*9	23	53	11	16	37	22
357 . . . . .	247	129	119	*	35	89	122	48	68	90	42
357.0 . . . . .	13	*7	*6	*	*6	*	*	*	*	*	*
357.2 . . . . .	219	111	108	*	27	81	112	44	58	81	36
358 . . . . .	26	12	14	*	*	*6	14	*8	*7	*7	*
358.0 . . . . .	24	11	13	*	*	*6	14	*8	*7	*6	*
359 . . . . .	27	14	12	*	*7	*8	*8	*8	*5	*8	*5
359.1 . . . . .	12	*8	*	*	*	*	*	*	*	*	*
361 . . . . .	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
362	175	71	104	*	17	55	99	37	36	66	37
362.0	100	42	57	—	14	46	40	20	18	42	20
362.01	98	42	56	—	14	46	39	20	17	41	19
362.3	12	*7	*5	—	*	*	*9	*	*	*	*
362.34	*9	*5	*	—	*	*	*6	*	*	*	*
362.5	49	14	35	*	*	*	47	11	13	13	11
362.50	48	14	35	—	*	*	46	11	13	13	11
365	182	65	117	*	*	28	151	45	49	53	35
365.9	176	62	114	*	*	27	147	44	47	52	34
366	22	*7	15	*	*	*	17	*	*7	*7	*
366.9	18	*6	13	*	*	*	14	*	*6	*7	*
368	58	27	31	*	15	15	26	13	15	19	12
368.1	10	*	*6	*	*	*	*6	*	*	*	*
368.16	*7	*	*	*	*	*	*	*	*	*	*
368.2	14	*6	*9	*	*	*6	*	*	*	*	*
368.4	18	10	*9	—	*	*	11	*	*5	*5	*
368.46	14	*8	*6	—	*	*	*8	*	*	*	*
368.8	10	*	*6	*	*	*	*	*	*	*	*
369	99	43	55	*	14	24	58	23	21	36	18
369.0	32	11	21	*	*	*7	21	*8	*7	11	*6
369.00	31	11	20	*	*	*7	20	*8	*7	10	*6
369.4	31	13	18	*	*	*7	19	*8	*5	11	*6
369.6	23	12	11	*	*7	*6	10	*	*6	10	*
369.60	22	12	10	*	*6	*5	10	*	*6	10	*
369.9	*5	*	*	*	*	*	*	*	*	*	*
370	*5	*	*	*	*	*	*	*	*	*	*
372	50	27	23	15	10	*7	18	9	11	20	10
372.0	13	10	*	*5	*	*	*6	*	*	*	*
372.00	11	*9	*	*	*	*	*5	*	*	*	*
372.3	29	13	16	*9	*6	*5	9	*6	*	13	*6
372.30	27	12	15	*8	*	*	9	*5	*	11	*5
372.7	*5	*	*	*	*	*	*	*	*	*	*
372.72	*5	*	*	*	*	*	*	*	*	*	*
373	10	*	*7	*	*	*	*	*	*	*	*
373.1	*7	*	*	*	*	*	*	*	*	*	*
373.13	*6	*	*	*	*	*	*	*	*	*	*
374	*7	*	*	*	*	*	*	*	*	*	*
374.3	*5	*	*	*	*	*	*	*	*	*	*
376	17	*8	10	*6	*6	*	*	*	*	*5	*
376.0	11	*5	*6	*	*	*	*	*	*	*	*
376.01	10	*	*6	*	*	*	*	*	*	*	*
377	14	*6	*8	*	*	*	*	*	*	*	*
378	22	*8	14	*	*6	*8	*5	*9	*5	*	*
378.5	12	*	*7	*	*	*	*	*5	*	*	*
378.51	*5	*	*	*	*	*	*	*	*	*	*
379	15	*6	*9	*	*	*	*	*	*	*	*5
380	35	13	21	*5	11	*8	10	*8	*9	11	*7
380.1	20	*7	13	*	*7	*	*	*	*	*8	*
380.10	19	*7	12	*	*7	*	*	*	*	*7	*
380.4	9	*	*	*	*	*	*	*	*	*	*
381	21	10	12	14	*	*	*	*	*5	*7	*
381.0	*5	*	*	*	*	*	*	*	*	*	*
381.4	*8	*	*	*	*	*	*	*	*	*	*
382	170	95	75	145	10	*8	*7	26	41	70	34
382.0	*5	*	*	*	*	*	—	*	*	*	*
382.9	162	90	72	139	9	*7	*7	24	37	68	33

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
383	9	*	*6	*	*	*	*	*	*	*	*
385	*8	*	*	*	*	*	*	*	*	*	*
386	75	25	50	*	*7	16	51	18	23	25	10
386.0	15	*5	10	—	*	*6	*8	*	*	*	*
386.00	14	*	10	—	*	*5	*7	*	*	*	*
386.1	32	10	22	—	*	*6	23	*7	10	12	*
386.11	21	*7	15	—	*	*	16	*5	*5	*8	*
386.12	*6	*	*	—	*	*	*	*	*	*	*
386.3	24	*9	16	*	*	*	18	*6	*8	*8	*
386.30	22	*9	13	*	*	*	17	*	*7	*8	*
388	19	9	10	*	*6	*	*7	*	*5	*5	*
388.6	*5	*	*	*	*	*	*	*	*	*	*
388.7	*6	*	*	*	*	*	*	*	*	*	*
388.70	*5	*	*	*	*	*	*	*	*	*	*
389	86	38	47	*7	12	13	54	22	22	22	20
389.1	*8	*	*	*	*	*	*5	*	*	*	*
389.10	*6	*	*	*	*	*	*	*	*	*	*
389.9	68	31	37	*5	*8	10	45	17	19	17	14
390-459	32,356	15,393	16,962	137	1,871	8,257	22,090	7,403	7,991	11,646	5,316
394	30	*5	25	*	*	*8	20	*	*7	10	10
394.0	12	*	12	*	*	*	9	*	*	*	*
394.1	*6	*	*	*	*	*	*	*	*	*	*
394.2	*7	*	*6	—	*	*	*5	*	*	*	*
396	172	62	110	*	*	18	152	45	38	67	22
396.0	*5	*	*	—	*	*	*	*	*	*	*
396.2	42	15	27	—	*	*	40	15	10	13	*
396.3	87	35	52	*	*	11	74	21	16	39	11
396.8	20	*7	13	*	*	*	17	*	*5	*6	*
396.9	16	*	12	—	*	*	15	*	*	*7	*
397	156	56	100	*	*7	28	119	43	32	57	24
397.0	154	55	99	*	*7	28	117	42	32	56	24
398	77	25	52	*	*	10	64	23	15	25	13
398.9	77	25	52	*	*	10	64	23	15	25	13
398.90	*7	*	*5	*	*	*	*	*	*	*	*
398.91	70	23	46	*	*	*8	59	20	15	23	12
401	8,006	3,336	4,671	18	587	2,447	4,955	1,808	1,923	2,980	1,296
401.0	78	29	49	*	*8	29	40	16	12	43	*6
401.1	105	45	60	*	*7	31	66	65	*	34	*5
401.9	7,824	3,262	4,562	17	571	2,387	4,848	1,727	1,909	2,902	1,285
402	378	135	244	*	10	76	291	99	75	138	66
402.0	10	*	*6	—	*	*	*6	*	*	*6	*
402.01	*6	*	*	—	*	*	*	*	*	*	*
402.9	364	130	235	*	*9	73	282	97	72	131	64
402.90	188	71	117	*	*	42	142	52	37	68	31
402.91	176	58	118	*	*	31	140	45	35	62	34
403	898	454	444	*6	110	305	476	169	188	375	166
403.0	17	*9	*8	*	*5	*6	*	*	*	10	*
403.01	15	*8	*8	*	*5	*6	*	*	*	10	*
403.1	*8	*	*	*	*	*	*	*	*	*	—
403.11	*7	*	*	—	*	*	*	*	*	*	—
403.9	874	441	432	*6	104	296	468	163	186	362	163
403.90	47	23	24	*	*5	11	31	9	14	19	*6
403.91	826	418	408	*5	99	285	437	154	172	344	158
404	82	43	39	*	*6	18	58	20	20	26	17
404.0	*6	*	*	—	*	*	*	*	*	*	*
404.9	75	39	36	*	*5	15	55	18	18	23	15
404.91	*6	*	*	—	*	*	*5	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
404.92	16	*8	*8	—	*	*	10	*	*	*7	*
404.93	50	27	23	*	*	10	37	13	11	14	12
405	11	*	*7	*	*	*	*5	*	*	*	*
405.9	10	*	*7	*	*	*	*	*	*	*	*
405.91	*8	*	*6	*	*	*	*	*	*	*	*
410	982	563	419	*	64	292	626	220	222	349	191
410.0	28	18	10	—	*	11	15	*	*8	11	*
410.01	26	17	9	—	*	11	14	*	*7	11	*
410.1	100	62	38	—	10	38	52	20	25	31	24
410.11	92	57	35	—	9	36	47	18	23	29	23
410.12	*6	*	*	—	*	*	*	*	*	*	*
410.2	16	10	*6	—	*	*5	*8	*	*	*5	*
410.21	15	9	*6	—	*	*5	*8	*	*	*5	*
410.3	13	*8	*	—	*	*	*7	*	*	*6	*
410.31	12	*8	*	—	*	*	*6	*	*	*6	*
410.4	131	85	45	—	11	61	58	28	24	54	25
410.41	123	81	42	—	11	57	55	26	23	50	23
410.42	*8	*	*	—	*	*	*	*	*	*	*
410.5	11	*8	*	—	*	*	*7	*	*	*	*
410.51	11	*8	*	—	*	*	*7	*	*	*	*
410.6	*5	*	*	—	*	*	*	*	*	*	*
410.7	475	261	214	—	25	121	328	124	110	160	81
410.71	445	246	199	—	23	115	306	115	104	152	74
410.72	28	15	14	—	*	*5	21	*9	*6	*7	*7
410.8	21	10	10	—	*	*	14	*	*7	*	*
410.81	19	10	*9	—	*	*	13	*	*6	*	*
410.9	183	96	87	*	10	39	133	29	41	72	41
410.91	165	89	76	*	10	35	121	27	36	64	38
410.92	15	*6	*9	—	*	*	10	*	*5	*7	*
411	734	407	327	—	37	277	420	183	164	272	115
411.0	*5	*	*	—	*	*	*	*	*	*	*
411.1	673	378	295	—	33	260	380	170	149	252	103
411.8	56	26	30	—	*	16	37	13	14	18	11
411.81	*5	*	*	—	*	*	*	*	*	*	*
411.89	50	24	26	—	*	14	33	12	10	17	11
412	755	446	309	*	38	228	489	172	206	245	132
413	479	242	237	—	20	151	308	111	131	174	63
413.9	474	239	235	—	19	149	306	111	130	171	62
414	4,736	2,673	2,063	*	148	1,313	3,272	1,119	1,225	1,684	708
414.0	4,434	2,485	1,949	*	138	1,242	3,052	1,048	1,145	1,576	666
414.00	1,275	718	556	—	26	298	951	342	353	443	136
414.01	3,019	1,677	1,342	*	110	902	2,004	675	757	1,080	507
414.02	96	67	29	—	*	33	62	20	27	31	19
414.04	*8	*	*	—	*	*	*5	*	*	*	*
414.05	35	19	16	—	*	*6	29	*7	*6	20	*
414.1	*6	*	*	—	*	*	*	*	*	*	*
414.8	239	156	82	—	*6	54	178	56	64	81	37
414.9	56	28	28	—	*	14	39	13	15	25	*
415	163	66	97	*	28	45	90	39	38	58	28
415.0	*6	*	*	—	—	*	*	*	*	*	*
415.1	157	63	94	*	28	43	85	36	37	57	27
415.11	13	*	9	*	*	*	*7	*	*	*	*
415.19	144	60	84	*	26	39	79	32	33	54	25
416	261	103	158	*	24	69	164	61	61	88	50
416.0	41	16	25	*	*	9	25	16	*9	12	*
416.8	147	56	91	*	17	36	93	29	35	52	32
416.9	73	31	42	*	*	24	46	16	18	24	15
420	31	19	12	*	*9	13	*9	*8	*8	*8	*7
420.9	28	18	10	*	*8	12	*8	*7	*8	*7	*6

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
420.90	18	12	*7	*	*	*8	*6	*	*6	*5	*
420.91	*7	*	*	*	*	*	*	*	*	*	*
421	35	22	13	*	*6	10	18	9	*6	11	*8
421.0	32	20	12	*	*6	9	16	*9	*6	9	*8
423	44	23	21	*	*8	12	21	11	*8	14	11
423.9	32	17	16	*	*6	10	15	10	*5	11	*6
424	1,011	391	620	*5	100	184	721	264	239	337	171
424.0	581	190	390	*	84	122	373	151	142	195	93
424.1	365	179	187	*	10	48	306	99	82	119	66
424.2	25	*7	18	*	*	*	17	*5	*5	*9	*6
424.3	*8	*	*	*	*	*	*	*	*	*	*
424.9	32	11	21	*	*	*6	23	*7	*9	10	*6
424.90	32	11	21	*	*	*6	23	*7	*9	10	*6
425	643	360	283	*6	58	198	381	134	158	240	111
425.1	17	*6	11	*	*	*	11	*	*	*6	*
425.4	575	326	249	*	49	170	350	120	138	219	98
425.5	17	14	*	—	*	11	*	*	*5	*	*
425.8	22	9	12	*	*	*8	11	*	*5	10	*5
425.9	*8	*	*6	*	*	*	*	*	*	*	*
426	436	221	216	*	24	74	336	100	115	131	90
426.0	68	37	30	*	*	9	56	14	19	19	15
426.1	121	65	56	*	*6	21	94	29	32	37	23
426.10	10	*6	*	*	*	*	*8	*6	*	*	*
426.11	65	33	32	*	*	14	48	11	21	21	13
426.12	9	*	*	—	*	*	*8	*	*	*	*
426.13	36	22	15	*	*	*	30	10	*8	11	*8
426.2	13	*7	*6	—	*	*	10	*	*	*7	*
426.3	97	33	64	*	*	17	78	25	22	28	21
426.4	70	42	28	*	*	12	53	17	17	20	16
426.5	33	21	12	—	*	*	29	*6	11	*9	*7
426.52	11	*8	*	—	—	*	10	*	*	*	*
426.53	15	*8	*7	—	*	*	13	*	*	*	*
426.6	*7	*	*	—	*	*	*6	*	*	*	*
426.7	13	*	*8	*	*	*5	*	*	*	*	*
426.8	*7	*	*	*	*	*	*	*	*	*	*
426.89	*6	*	*	*	*	*	*	*	*	*	*
426.9	*8	*5	*	*	*	*	*6	*	*	*	*
427	4,119	1,990	2,129	27	158	675	3,259	983	1,023	1,395	718
427.0	105	36	68	*	9	20	73	24	23	39	19
427.1	251	170	81	*	16	72	162	57	61	95	38
427.3	2,723	1,298	1,425	*	50	365	2,304	684	685	895	459
427.31	2,515	1,178	1,337	*	41	314	2,158	644	637	821	412
427.32	208	120	88	*	9	51	146	40	48	74	47
427.4	45	30	14	*	*	20	22	11	*8	15	11
427.41	44	30	14	*	*	20	22	11	*8	14	11
427.5	104	52	51	*	*9	30	63	20	23	34	27
427.6	97	49	48	*	*7	18	72	18	25	32	22
427.61	25	12	14	*	*	*	21	*6	*7	*5	*8
427.69	70	37	33	*	*	15	50	11	18	27	14
427.8	735	332	403	17	61	136	521	156	181	263	135
427.81	197	86	111	*	*	15	177	55	40	74	28
427.89	538	246	292	16	57	121	344	101	141	189	107
427.9	58	22	36	*	*	13	40	13	17	21	*7
428	3,401	1,485	1,916	14	93	643	2,651	752	912	1,248	490
428.0	3,381	1,476	1,905	13	92	641	2,635	748	909	1,239	485
428.1	*9	*	*	*	*	*	*7	*	*	*	*
428.9	11	*5	*6	*	*	*	*9	*	*	*6	*
429	232	113	118	*	18	44	165	53	66	86	27
429.2	89	47	42	*	*	14	73	25	19	37	*9

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
429.3	97	44	53	*	13	19	64	16	36	34	11
429.4	9	*	*	*	*	*	*	*	*	*	*
429.7	*5	*	*	*	*	*	*	*	*	*	*
429.9	22	9	13	*	*	*	15	*5	*7	*7	*
430	29	11	18	*	*5	12	12	*6	*7	10	*6
431	78	34	44	*	*6	26	45	19	14	24	20
432	37	21	16	*	*	*	28	*6	*9	12	10
432.1	31	18	14	*	*	*	25	*5	*7	11	*8
433	315	169	146	*	*	71	240	78	75	120	42
433.1	240	132	109	*	*	53	185	60	57	92	32
433.10	216	121	95	—	*	47	168	53	51	84	28
433.11	24	11	13	*	*	*6	17	*7	*6	*7	*
433.2	*5	*	*	*	*	*	*	*	*	*	*
433.3	62	32	31	*	*	14	48	17	15	22	*8
433.30	54	26	28	*	*	13	41	15	12	19	*7
433.31	*8	*6	*	—	*	*	*7	*	*	*	*
434	363	162	201	*	13	83	264	71	91	131	70
434.0	13	*	*8	*	*	*	9	*	*	*	*
434.01	10	*	*6	*	*	*	*8	*	*	*	*
434.1	48	22	26	*	*	10	34	13	*8	12	15
434.10	*7	*	*	—	*	*	*5	*	*	*	*
434.11	41	19	22	*	*	*9	29	10	*8	10	12
434.9	302	136	166	*	9	70	221	57	78	114	53
434.90	10	*6	*	—	*	*	*6	*	*	*	*
434.91	293	129	163	*	*8	68	215	54	76	111	52
435	262	107	155	*	11	61	190	60	58	99	44
435.3	14	*	10	—	*	*	13	*	*	*	*
435.8	10	*	*6	—	*	*	*7	*	*	*	*
435.9	230	95	135	*	10	53	167	54	49	88	39
436	167	72	95	*	*5	24	138	29	38	70	31
437	164	69	95	*	*7	28	128	32	46	62	24
437.0	71	31	39	—	*	*8	63	16	23	23	*8
437.1	16	*	11	*	*	*	13	*	*5	*5	*
437.2	13	*5	*7	*	*	*	*8	*	*	*5	*
437.3	17	*6	11	*	*	*8	*8	*	*	*5	*
437.9	38	18	20	*	*	*	32	*	*9	19	*6
438	500	240	260	*	15	116	366	110	134	159	97
438.0	*9	*	*	—	*	*	*5	*	*	*	*
438.1	82	43	40	*	*	14	66	16	25	24	17
438.10	*5	*	*	*	*	*	*	*	*	*	*
438.11	54	29	25	*	*	*9	44	11	19	15	9
438.12	12	*6	*6	—	*	*	10	*	*	*	*
438.19	12	*	*7	—	*	*	*9	*	*	*	*
438.2	225	102	123	*	*8	60	156	60	59	67	40
438.20	199	89	110	*	*7	53	138	53	53	62	32
438.21	11	*6	*5	*	*	*	*6	*	*	*	*
438.22	14	*7	*7	—	*	*	12	*	*	*	*5
438.5	*6	*	*	—	*	*	*	*	*	*	*
438.8	152	77	75	*	*	32	116	23	40	55	33
438.82	38	22	16	—	*	*6	33	*6	*6	16	11
438.89	113	54	58	*	*	26	82	17	35	39	22
438.9	24	10	14	*	*	*	19	*6	*6	9	*
440	380	198	182	*	*8	93	278	76	108	130	66
440.0	24	13	11	—	*	*6	17	*6	*7	*6	*5
440.1	42	14	27	*	*	*8	32	*8	13	14	*6
440.2	224	125	99	*	*	59	160	40	62	78	44
440.20	60	34	26	*	*	11	48	*7	17	22	14
440.21	42	24	17	—	*	17	23	*7	13	13	*8

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
440.22	12	*6	*6	*	*	*	*7	*	*	*	*
440.23	34	16	18	—	*	*7	25	*7	9	11	*7
440.24	71	40	31	—	*	17	53	17	18	27	10
440.29	*5	*	*	*	*	*	*	*	*	*	*
440.3	*8	*	*	—	*	*	*	*	*	*	*
440.8	*9	*	*	—	*	*	*	*	*	*	*
440.9	74	38	36	*	*	12	61	20	22	26	*6
441	166	115	51	—	*	25	138	40	41	53	31
441.0	17	10	*7	—	*	*6	9	*	*	*6	*5
441.01	*8	*6	*	—	*	*	*	*	*	*	*
441.02	*5	*	*	—	*	*	*	*	*	*	*
441.2	17	10	*7	—	*	*	11	*	*	*	*6
441.3	10	*7	*	—	*	*	*9	*	*	*	*
441.4	114	84	30	—	*	12	102	28	31	39	16
442	32	25	*8	*	*	*8	22	*8	*7	9	*8
442.2	15	13	*	*	*	*	11	*	*	*	*
442.3	11	*8	*	*	*	*	*8	*	*	*	*
443	464	231	233	*	14	116	334	100	122	189	54
443.0	17	*	14	*	*	*7	*5	*6	*	*	*
443.8	41	25	16	*	*	15	25	10	9	13	9
443.81	27	19	*9	—	*	12	15	*7	*5	*9	*6
443.89	14	*7	*7	*	*	*	11	*	*	*	*
443.9	404	202	202	—	*7	94	303	84	108	171	42
444	83	44	40	*	*8	26	49	16	19	29	20
444.0	*8	*	*	—	*	*	*	*	*	*	*
444.2	55	29	27	*	*5	16	34	11	13	19	13
444.21	*7	*	*	—	*	*	*6	*	*	*	*
444.22	48	25	23	*	*	15	29	9	12	17	9
444.8	17	10	*7	*	*	*6	*8	*	*	*6	*
444.81	10	*7	*	*	*	*	*	*	*	*	*
444.89	*7	*	*	*	*	*	*	*	*	*	*
446	45	18	27	*7	*	*9	25	9	11	14	11
446.1	*7	*	*	*7	—	—	*	*	*	*	*
446.4	*7	*	*	—	*	*	*	*	*	*	*
446.5	19	*5	14	—	—	*	18	*	*6	*5	*
447	60	28	32	*	*	15	39	14	15	23	*8
447.1	28	14	13	—	*	*7	19	*7	*7	11	*
447.6	13	*	*8	*	*	*	*6	*	*	*7	*
447.9	*7	*	*	—	*	*	*6	*	*	*	*
448	*7	*	*	*	*	*	*	*	*	*	*
448.0	*6	*	*	*	*	*	*	*	*	*	*
451	61	23	37	*	13	15	31	13	12	25	11
451.0	11	*	*7	—	*	*	*7	*	*	*	*
451.1	14	*6	*8	*	*	*	*9	*	*	*6	*
451.11	*6	*	*	—	*	*	*	*	*	*	*
451.19	*8	*	*	*	*	*	*5	*	*	*	*
451.2	9	*	*7	*	*	*	*6	*	*	*5	*
451.8	25	11	14	*	*8	*7	9	*	*	10	*6
451.82	*7	*	*	—	*	*	*	*	*	*	*
451.84	13	*6	*7	*	*	*	*	*	*	*7	*
453	341	143	198	*	51	105	184	77	82	128	54
453.2	*5	*	*	*	*	*	*	*	*	*	*
453.8	305	129	176	*	48	96	160	69	74	113	48
453.9	26	9	17	*	*	*6	18	*5	*7	11	*
454	31	11	20	—	*	9	20	*8	10	*8	*5
454.0	*6	*	*	—	*	*	*	*	*	*	*
454.2	*7	*	*	—	*	*	*	*	*	*	*
454.9	16	*5	11	—	*	*	11	*	*6	*5	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
455	168	67	101	*	37	51	80	38	29	68	33
455.0	71	30	40	—	11	21	38	19	10	29	13
455.2	27	10	16	*	*	10	12	*	*6	10	*6
455.3	16	*5	10	*	*	*5	*6	*	*	*	*5
455.5	11	*	*8	—	*	*	*	*	*	*6	*
455.6	27	10	17	—	*9	*5	13	*7	*	11	*
455.8	10	*	*5	—	*	*	*	*	*	*	*
456	71	40	31	*	12	36	23	16	13	23	19
456.0	*5	*	*	*	*	*	*	*	*	*	*
456.1	*9	*	*	*	*	*5	*	*	*	*	*
456.2	47	27	20	*	*8	23	16	*8	10	15	14
456.20	29	18	11	*	*6	12	11	*5	*6	10	*8
456.21	18	*8	10	*	*	11	*5	*	*	*	*6
456.8	*9	*6	*	*	*	*	*	*	*	*	*
457	33	16	17	*	*8	14	9	*8	*8	10	*7
457.1	20	9	10	*	*	*8	*7	*5	*	*6	*
457.2	*5	*	*	*	*	*	*	*	*	*	*
457.8	*5	*	*	*	*	*	*	*	*	*	*
458	446	214	232	*8	50	114	273	91	112	162	80
458.0	112	53	59	*	11	21	79	23	29	41	19
458.2	39	20	19	*	*	12	22	11	10	10	*9
458.8	29	15	14	*	*	*9	16	*	9	11	*5
458.9	265	126	139	*6	30	72	156	54	64	100	47
459	164	79	85	*	17	47	99	38	41	61	24
459.0	18	10	*9	*	*	*6	*9	*	*	*7	*
459.2	14	*7	*7	*	*	*5	*7	*	*	*7	*
459.8	121	56	65	*	11	34	75	31	34	40	16
459.81	113	51	61	*	10	32	70	27	32	38	15
459.89	*8	*	*	*	*	*	*5	*	*	*	*
459.9	10	*6	*	—	*	*	*7	*	*	*6	*
460-519	11,181	5,236	5,945	1,289	1,374	2,505	6,014	2,359	2,656	4,159	2,007
460	*5	*	*	*	*	*	*	*	*	*	*
461	57	17	40	10	19	11	17	11	17	22	*7
461.0	13	*	10	*	*	*	*	*	*	*	*
461.2	*6	*	*5	*	*	*	*	*	*	*	*
461.9	33	12	21	*6	10	*6	10	*	10	15	*
462	55	26	29	18	21	*8	9	14	13	20	*9
463	24	10	14	*7	16	*	*	*	*7	*9	*
464	37	20	17	25	*	*	*	*8	9	12	*8
464.0	*7	*	*	*	*	*	*	*	*	*	*
464.00	*7	*	*	*	*	*	*	*	*	*	*
464.1	*7	*	*	*6	*	*	*	*	*	*	*
464.10	*7	*	*	*6	*	*	*	*	*	*	*
464.4	19	11	*8	19	*	*	—	*	*	*7	*
465	155	72	83	65	33	23	35	29	36	56	34
465.9	152	71	82	63	32	22	34	28	36	54	34
466	417	215	203	252	29	52	84	70	95	165	88
466.0	171	67	105	13	29	49	80	35	44	70	22
466.1	246	148	98	239	*	*	*	34	51	95	66
466.11	144	87	57	142	*	*	*	17	30	55	42
466.19	102	61	41	97	*	*	*	17	21	40	23
470	10	*8	*	*	*	*	*	*	*	*	*
472	10	*	*7	*	*	*	*	*	*	*	*
472.0	10	*	*7	*	*	*	*	*	*	*	*
473	173	78	94	31	51	43	47	32	38	81	22
473.0	22	10	11	*	*7	*	*9	*	*	13	*
473.2	10	*5	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
473.8	16	*7	10	*	*	*6	*	*	*	*6	*
473.9	118	54	64	22	34	30	32	21	28	54	15
474	28	17	11	19	*6	*	—	*7	*5	*9	*7
474.0	*8	*	*	*	*	*	—	*	*	*	*
474.1	19	13	*6	15	*	*	—	*5	*	*6	*
474.10	13	9	*	12	*	*	—	*	*	*	*
475	19	*8	10	*5	11	*	*	*	*5	*7	*
477	70	27	43	*6	16	19	29	*6	22	24	16
477.9	68	26	42	*6	15	19	28	*6	22	24	16
478	96	55	41	19	23	28	25	17	20	36	23
478.0	*6	*	*	—	*	*	*	*	*	*	*
478.1	23	12	10	*7	*8	*	*	*	*	*9	*6
478.2	30	19	10	*6	*7	12	*5	*	*8	12	*5
478.29	21	14	*8	*	*	10	*	*	*7	*8	*
478.3	19	10	9	*	*	*6	11	*	*	*6	*6
478.30	*9	*	*	*	*	*	*6	*	*	*	*
478.31	*5	*	*	*	*	*	*	*	*	*	*
478.7	11	*6	*	*	*	*	*	*	*	*	*
478.74	*6	*	*	*	*	*	*	*	*	*	*
480	45	26	19	36	*	*	*	*	11	15	14
480.1	23	15	*8	21	*	*	*	*	*7	*9	*6
480.9	21	10	11	14	*	*	*	*	*	*6	*8
481	58	28	30	*	10	13	32	13	16	17	11
482	239	127	112	16	23	41	159	48	50	99	42
482.0	19	12	*6	*	*	*	12	*	*	10	*
482.1	43	25	18	*	*	*8	27	11	*8	18	*7
482.2	22	11	11	*	*	*	15	*	*6	*8	*
482.3	18	11	*6	*	*	*	11	*	*	10	*
482.30	*6	*	*	*	*	*	*	*	*	*	*
482.31	*7	*5	*	*	*	—	*5	*	*	*6	*
482.4	75	37	38	*	*8	12	53	17	12	32	14
482.41	70	34	36	*	*7	11	50	14	11	31	13
482.8	38	21	18	*	*	*6	28	*6	14	14	*
482.82	*5	*	*	*	*	*	*	*	*	*	*
482.83	28	14	13	*	*	*	20	*	10	10	*
482.9	25	10	15	*	*	*	14	*7	*6	*6	*6
483	*8	*	*	*	*	*	*	*	*	*	*
483.0	*6	*	*	*	*	*	*	*	*	*	*
485	36	18	18	11	*	*7	16	*7	*	21	*
486	1,791	837	953	199	180	322	1,089	359	421	694	317
487	47	21	27	19	*6	*9	14	*8	18	14	*8
487.0	*8	*	*	*	*	*	*	*	*	*	*
487.1	35	16	19	15	*5	*7	*9	*7	11	11	*6
490	110	45	66	*5	26	22	57	17	21	56	17
491	1,089	491	598	*	41	293	753	225	270	450	144
491.2	1,042	473	569	*	30	278	734	219	252	435	136
491.20	114	52	62	—	*	33	76	27	23	43	21
491.21	928	421	507	*	25	245	658	192	229	392	115
491.9	37	12	24	*	*9	11	16	*	14	13	*5
492	197	105	92	*	13	52	130	49	49	69	30
492.0	17	14	*	—	*7	*	*6	*	*	*6	*
492.8	179	91	88	*	*6	48	125	47	44	63	26
493	1,624	568	1,057	343	449	428	403	397	385	539	303
493.0	78	44	34	54	13	*9	*	12	*7	14	45
493.00	34	16	17	21	*6	*	*	*5	*	*6	19
493.01	*9	*5	*	*7	*	*	—	*	*	*	*6
493.02	36	22	13	27	*6	*	*	*6	*	*6	21

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
493.2	315	106	209	*	26	112	176	66	89	103	56
493.20	196	67	128	*	14	66	116	40	58	64	34
493.21	11	*	*8	*	*	*6	*	*	*	*	*
493.22	108	36	73	*	10	41	57	22	30	35	21
493.9	1,230	417	812	288	410	307	224	319	288	421	202
493.90	884	277	607	133	321	244	186	235	211	280	157
493.91	80	36	45	47	21	*8	*	15	18	35	12
493.92	266	105	161	108	68	55	34	68	59	106	33
494	41	18	23	*	*	*7	29	*7	12	17	*
494.0	30	15	15	*	*	*5	22	*	10	13	*
494.1	10	*	*8	*	*	*	*7	*	*	*	*
496	2,021	1,029	992	*	58	457	1,505	441	511	735	335
500	*9	*8	*	—	*	*	*8	*6	*	*	*
501	22	22	*	—	—	*	18	*6	*	10	*
507	350	195	155	15	23	56	255	92	71	121	66
507.0	347	193	154	14	23	56	254	91	71	119	65
508	*6	*	*	*	*	*	*	*	*	*	*
510	33	24	*9	*	*6	11	12	*8	*7	9	9
510.0	*5	*	*	*	*	*	*	*	*	*	*
510.9	27	19	*8	*	*5	*8	11	*7	*6	*8	*6
511	432	203	229	14	51	103	264	94	103	149	86
511.0	22	11	11	*	*5	*8	*8	*	*	10	*
511.8	29	16	13	*	*5	9	13	*7	*6	*8	*8
511.9	378	175	203	11	40	85	242	83	92	131	73
512	106	58	48	*8	26	26	47	18	26	38	24
512.0	*7	*	*	*	*	*	*	*	*	*	*
512.1	48	23	26	*	*6	12	26	10	10	17	11
512.8	51	31	20	*	17	12	17	*7	14	18	12
513	15	10	*6	*	*	*7	*	*	*	*7	*
513.0	15	10	*6	*	*	*7	*	*	*	*6	*
514	39	13	25	*	*	9	23	*6	12	15	*6
515	170	76	93	*	10	36	124	32	50	62	25
516	28	14	14	*	*	*8	16	*7	*8	*8	*5
516.3	16	*6	9	—	*	*5	9	*	*	*6	*
516.8	11	*7	*	*	*	*	*7	*	*	*	*
517	18	*6	12	*	*5	*8	*	*	*	*7	*
517.8	17	*6	11	*	*5	*8	*	*	*	*7	*
518	1,391	675	716	122	164	361	744	282	302	511	296
518.0	352	172	180	27	57	95	173	62	75	131	84
518.1	*8	*	*	*	*	*	*	*	*	*	*
518.3	19	*9	10	*	*	*	11	*	*	*8	*
518.4	21	*6	15	*	*	*5	11	*	*	*8	*5
518.5	111	61	50	*6	17	31	57	24	29	35	23
518.8	879	422	456	85	82	223	489	188	188	327	176
518.81	584	279	306	20	56	158	350	134	116	231	103
518.82	121	68	54	55	11	16	39	18	20	32	51
518.83	48	24	24	*	*	17	24	*9	20	16	*
518.84	44	16	28	*	*	14	28	*6	12	20	*5
518.89	81	36	45	*6	*9	17	49	20	20	26	14
519	82	42	41	16	15	16	35	15	21	34	12
519.0	9	*	*	*	*	*	*	*	*	*	*
519.1	33	16	17	*6	*6	*6	14	*6	10	12	*5
519.4	*7	*	*	*	*	*	*	*	*	*	*
519.8	28	14	13	*6	*5	*	12	*	*7	14	*
520-579	10,232	4,353	5,878	510	2,160	2,999	4,564	2,145	2,421	3,808	1,859
521	21	14	*7	*	9	*	*	*	*	9	*

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
521.0	20	14	*6	*	*8	*	*	*	*	9	*
521.00	15	11	*	*	*7	*	*	*	*	*7	*
522	24	*8	16	*	13	*	*	*5	*5	9	*
522.5	20	*6	14	*	11	*	*	*	*	*8	*
523	13	*7	*6	*	*6	*	*	*	*	*	*
523.1	*5	*	*	*	*	*	*	*	*	*	*
524	31	11	20	*	25	*	*	*	*	16	*7
524.0	13	*6	*7	*	12	*	—	*	*	*7	*
524.6	10	*	*9	*	*6	*	*	*	*	*	*
525	9	*6	*	*	*	*	*	*	*	*	*
526	*9	*	*6	*	*	*	*	*	*	*	*
527	17	*6	11	*	*	*	*8	*	*	*6	*
527.2	11	*	*8	*	*	*	*5	*	*	*	*
528	40	18	21	9	11	10	9	9	9	13	*8
528.0	18	*9	9	*	*	*	*	*5	*	*	*
528.3	*7	*	*	*	*	*	*	*	*	*	*
528.9	*9	*	*	*	*	*	*	*	*	*	*
530	2,043	837	1,206	101	339	648	956	451	457	789	346
530.0	10	*	*6	*	*	*	*	*	*	*5	*
530.1	216	105	112	*	47	65	100	52	53	75	36
530.10	91	42	48	*	20	29	40	27	20	30	13
530.11	63	31	32	*	14	20	27	16	14	23	11
530.12	*8	*	*	*	*	*	*	*	*	*	*
530.19	55	27	28	*	11	15	29	*9	17	20	9
530.2	67	43	24	*	13	21	33	15	14	24	14
530.3	51	24	27	*	*	*7	39	10	12	21	*8
530.5	21	*8	14	*	*	*9	10	*	*	*7	*
530.6	*7	*	*	—	*	*	*6	*	*	*	*
530.7	24	15	*8	*	*9	*	9	*	*6	10	*
530.8	1,642	633	1,010	92	259	538	752	360	365	643	275
530.81	1,615	620	996	92	256	531	737	355	358	633	269
530.82	14	*7	*7	*	*	*	*8	*	*	*6	*
530.89	12	*5	*6	*	*	*	*7	*	*	*	*
531	166	77	89	*	14	45	107	33	44	55	35
531.0	18	10	*8	*	*	*5	13	*	*6	*6	*
531.00	18	10	*8	*	*	*5	12	*	*5	*6	*
531.4	77	37	41	*	*	22	50	14	21	27	15
531.40	76	36	39	*	*	21	50	14	21	26	15
531.5	*7	*	*	*	*	*	*	*	*	*	*
531.50	*7	*	*	*	*	*	*	*	*	*	*
531.9	53	22	31	*	*7	13	33	10	12	18	12
531.90	51	21	30	*	*7	13	32	10	12	17	12
532	101	52	49	*	14	30	56	19	27	34	21
532.0	13	*8	*	*	*	*	*9	*	*	*	*
532.00	13	*8	*	*	*	*	*9	*	*	*	*
532.4	50	26	23	*	*5	16	29	*6	13	18	12
532.40	49	26	23	*	*5	15	28	*6	13	17	12
532.9	24	11	12	*	*6	*7	11	*6	*7	*7	*
532.90	23	11	12	*	*6	*6	10	*6	*6	*7	*
533	129	62	67	*	20	37	72	30	32	47	21
533.4	11	*	*6	—	*	*	*8	*	*	*	*
533.40	10	*	*6	—	*	*	*8	*	*	*	*
533.7	*6	*	*	—	*	*	*	*	*	*	*
533.70	*6	*	*	—	*	*	*	*	*	*	*
533.9	111	53	58	*	17	33	61	24	26	41	19
533.90	110	53	57	*	17	33	60	24	26	40	19
534	*7	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
535	606	244	362	13	133	180	280	135	133	241	97
535.0	61	25	36	*7	11	18	25	12	16	27	*6
535.00	47	17	30	*6	10	14	17	10	13	20	*
535.01	14	*8	*6	*	*	*	*8	*	*	*7	*
535.1	93	34	59	*	15	27	50	29	15	30	19
535.10	74	26	48	*	14	22	38	24	11	25	13
535.11	20	*8	12	*	*	*6	12	*	*	*5	*5
535.3	17	12	*	*	9	*6	*	*	*	*	*
535.30	13	9	*	*	*7	*	*	*	*	*	*
535.4	129	47	81	*	24	41	62	26	30	52	21
535.40	93	30	63	*	20	31	40	18	22	40	13
535.41	36	18	18	*	*	10	22	*8	*8	12	*8
535.5	248	98	151	*	60	70	114	52	52	109	36
535.50	208	81	127	*	55	60	91	44	44	91	29
535.51	40	17	23	*	*6	10	24	*8	*8	18	*7
535.6	57	27	30	*	14	17	27	12	16	18	11
535.60	50	24	26	*	13	14	23	11	13	16	10
535.61	*7	*	*	*	*	*	*	*	*	*	*
536	260	98	161	19	67	78	95	46	66	112	36
536.2	36	11	25	*9	12	*8	*8	*6	*8	17	*5
536.3	96	36	59	*	33	33	28	18	13	52	12
536.4	41	17	24	*6	*	*9	24	10	*7	17	*6
536.41	15	*6	10	*	*	*	10	*	*	*8	*
536.42	16	*7	*9	*	*	*	9	*	*	*6	*
536.49	*9	*	*5	*	*	*	*	*	*	*	*
536.8	79	31	48	*	20	26	31	9	37	22	10
536.9	*7	*	*	*	*	*	*	*	*	*	*
537	75	29	45	*	*8	18	45	16	17	27	15
537.0	24	10	14	*	*	*7	12	*	*	10	*6
537.3	*7	*	*5	*	*	*	*	*	*	*	*
537.8	37	15	22	*	*	*8	25	*9	10	12	*6
537.83	20	*7	13	—	*	*	18	*	*	*8	*
537.89	11	*5	*6	*	*	*	*	*	*	*	*
540	295	163	132	68	153	49	25	57	64	83	91
540.0	65	38	27	17	23	15	10	10	15	20	21
540.1	34	21	13	11	10	*7	*6	*	*7	11	11
540.9	196	103	93	40	120	27	*9	42	42	53	59
541	15	*7	*8	*	9	*	*	*	*	*9	*
542	*6	*	*	*	*	*	*	*	*	*	*
543	19	*6	13	*	10	*	*	*	*5	*6	*
543.9	16	*5	11	*	9	*	*	*	*5	*	*
550	77	62	15	9	10	15	43	18	19	27	12
550.1	26	20	*6	*	*	*	15	*6	*	11	*
550.10	22	16	*5	*	*	*	13	*5	*	*9	*
550.9	50	41	*9	*5	*6	12	27	12	14	16	*8
550.90	35	28	*8	*	*5	*9	19	*8	11	11	*5
550.91	*6	*6	—	*	*	*	*	*	*	*	*
550.92	*8	*7	*	*	*	*	*	*	*	*	*
552	89	28	61	—	14	31	44	21	22	29	17
552.1	24	11	14	—	*	*9	10	*5	*6	*8	*5
552.2	50	12	38	—	*7	20	23	12	12	17	10
552.20	11	*	*9	—	*	*	*6	*	*	*	*
552.21	34	*8	26	—	*5	14	15	10	*7	12	*6
552.29	*5	*	*	—	*	*	*	*	*	*	*
552.3	*6	*	*	—	*	*	*5	*	*	*	*
553	524	193	331	*7	85	161	270	109	129	206	79
553.1	54	25	29	*	13	20	18	*9	16	18	11
553.2	102	43	59	*	18	44	39	23	21	37	21
553.20	27	12	15	*	*	12	11	*	*5	13	*6

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
553.21	70	29	41	*	13	30	27	18	15	23	14
553.3	362	122	240	*	53	95	211	75	91	149	47
555	105	44	61	*	51	32	19	23	29	36	17
555.0	19	10	9	*	*9	*5	*	*5	*	*6	*
555.1	14	*5	9	*	9	*	*	*	*5	*	*
555.2	*5	*	*	*	*	*	*	*	*	*	*
555.9	67	28	39	*	31	22	13	14	16	26	11
556	64	35	29	*	26	18	19	15	15	20	15
556.6	*7	*5	*	*	*	*	*	*	*	*	*
556.9	40	23	17	*	14	11	14	9	10	11	10
557	79	26	53	*	*8	19	50	21	20	25	12
557.0	44	16	29	*	*6	12	25	12	11	14	*7
557.1	*6	*	*	*	*	*	*	*	*	*	*
557.9	29	*9	20	*	*	*5	21	*7	*8	10	*
558	411	164	246	121	84	70	136	76	100	167	68
558.1	*6	*	*	—	*	*	*	*	*	*	*
558.9	402	162	240	120	83	66	132	75	97	164	66
560	709	303	405	28	119	183	379	125	193	252	138
560.0	*5	*	*	*	*	*	*	*	*	*	*
560.1	319	145	175	13	62	86	158	50	87	118	65
560.2	17	*8	*9	*	*	*	10	*	*6	*5	*
560.3	81	34	47	*5	*8	9	58	18	16	35	12
560.39	77	32	44	*5	*8	*9	55	17	16	32	11
560.8	133	51	82	*	22	39	70	23	41	41	27
560.81	83	31	52	*	14	24	43	12	24	28	19
560.89	50	20	29	*	*8	14	27	11	17	13	*9
560.9	153	62	91	*	23	46	81	28	42	51	32
562	576	221	355	*	49	143	384	137	143	208	87
562.0	11	*6	*6	—	*	*	*9	*	*	*	*
562.00	*8	*	*	—	*	*	*6	*	*	*	*
562.1	565	215	349	*	48	141	375	134	141	204	86
562.10	241	85	156	*	11	49	181	63	55	88	35
562.11	221	88	133	*	34	76	110	54	65	72	30
562.12	87	36	52	*	*	12	75	14	18	38	17
562.13	15	*6	*9	—	*	*	9	*	*	*7	*
564	560	205	356	36	116	142	267	116	161	187	97
564.0	398	168	230	32	65	86	215	80	112	139	67
564.00	356	151	205	27	57	73	199	71	104	125	55
564.09	41	17	24	*6	*7	12	15	*8	*7	14	11
564.1	119	20	98	*	37	41	40	25	37	33	23
564.7	*6	*	*	*	*	*	*	*	*	*	*
564.8	21	9	12	*	*6	*6	*6	*6	*6	*6	*
564.81	*5	*	*	*	*	*	*	*	*	*	*
564.89	16	*6	10	*	*	*5	*6	*	*	*5	*
565	14	10	*	*	*	*5	*	*	*	*	*
565.0	*5	*	*	*	*	*	*	*	*	*	*
565.1	9	*8	*	*	*	*	*	*	*	*	*
566	30	18	12	*	12	10	*6	*6	*7	*9	*9
567	108	55	54	*	24	46	35	20	22	43	23
567.2	72	38	34	*	18	31	21	12	16	30	14
567.8	16	9	*7	*	*	*6	*6	*	*	*	*
567.9	20	*7	13	*	*	9	*8	*	*	*9	*
568	190	45	144	*	67	59	60	36	39	79	35
568.0	158	38	119	*	54	50	52	31	30	69	27
568.8	31	*7	24	*	13	9	*8	*	*9	10	*8
568.81	16	*	14	*	*8	*	*	*	*	*6	*
568.89	13	*	10	*	*	*6	*	*	*5	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
569	272	117	155	*6	43	76	147	63	65	109	35
569.0	12	*5	*7	*	*	*5	*6	*	*	*6	*
569.1	*7	*	*6	*	*	*	*5	*	*	*	*
569.3	57	28	29	*	10	14	32	13	11	22	10
569.4	30	15	16	*	*6	*9	15	*8	*9	9	*
569.41	*8	*5	*	*	*	*	*	*	*	*	*
569.49	18	*8	10	*	*	*	10	*6	*7	*	*
569.5	22	11	11	*	*	*7	11	*7	*6	*6	*
569.6	26	*9	17	*	*	10	13	*	*7	13	*
569.62	*6	*	*	*	*	*	*	*	*	*	*
569.69	17	*6	12	*	*	*6	9	*	*	*8	*
569.8	109	45	65	*	17	28	61	24	27	47	11
569.81	13	*6	*7	*	*	*6	*	*	*	*5	*
569.82	*8	*	*6	*	*	*	*	*	*	*	*
569.83	27	12	15	*	*	11	11	*5	*6	13	*
569.84	*8	*	*	*	*	*	*6	*	*	*	*
569.85	23	9	14	—	*	*	19	*6	*5	11	*
569.89	30	12	19	*	*7	*	19	*7	*7	12	*
570	25	13	12	*	*7	*9	*8	*5	*	*8	*7
571	421	262	159	*	78	213	127	93	87	150	92
571.1	44	35	10	—	19	20	*	10	10	11	13
571.2	142	109	34	*	25	91	27	34	28	51	29
571.3	22	17	*5	—	*7	10	*5	*	*8	*5	*8
571.4	11	*	*7	*	*	*	*	*	*	*5	*
571.49	*8	*	*5	*	*	*	*	*	*	*	*
571.5	158	81	77	*	13	72	72	36	28	61	33
571.6	*8	*	*7	*	*	*	*	*	*	*	*
571.8	27	9	17	*	9	11	*6	*6	*6	11	*
572	140	85	55	*	20	70	47	31	29	50	30
572.0	*8	*	*	*	*	*	*	*	*	*	*
572.2	62	40	23	*	*8	33	21	14	13	23	13
572.3	41	23	18	*	*7	20	14	*7	10	13	11
572.4	*9	*5	*	*	*	*	*	*	*	*	*
572.8	19	12	*8	*	*	10	*6	*6	*	*7	*
573	84	37	47	*	17	28	35	18	19	33	14
573.3	33	16	17	*	11	9	11	9	*5	13	*6
573.8	27	11	16	*	*	10	13	*	*8	*8	*5
573.9	16	*6	10	*	*	*7	*8	*	*	*7	*
574	636	215	421	*	179	186	269	129	148	229	129
574.0	145	58	87	*	41	49	55	26	39	53	27
574.00	139	55	83	*	38	48	52	23	37	52	26
574.01	*6	*	*	—	*	*	*	*	*	*	*
574.1	246	74	172	*	75	81	89	51	53	91	51
574.10	239	71	168	*	73	79	86	49	52	89	49
574.11	*7	*	*	—	*	*	*	*	*	*	*
574.2	137	48	89	*	32	34	70	31	34	46	26
574.20	135	46	88	*	31	34	69	29	34	46	26
574.3	*7	*	*5	—	*	*	*	*	*	*	*
574.31	*5	*	*	—	*	*	*	*	*	*	*
574.4	11	*	*6	—	*	*	*	*	*	*	*
574.40	*6	*	*	—	*	*	*	*	*	*	*
574.41	*5	*	*	—	*	*	*	*	*	*	*
574.5	42	15	27	—	*8	*9	26	9	*7	16	10
574.50	26	10	16	—	*	*6	17	*6	*	*9	*8
574.51	16	*5	11	—	*	*	*9	*	*	*7	*
574.6	*7	*	*	—	*	*	*	*	*	*	*
574.7	19	*	17	*	*8	*	*8	*	*5	*8	*
574.70	14	*	13	*	*7	*	*	*	*	*6	*
574.71	*5	*	*	*	*	*	*	*	*	*	*
574.8	13	*	*8	—	*	*	*6	*	*	*5	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
574.80	10	*	*6	—	*	*	*	*	*	*	*
574.9	*8	*	*	—	*	*	*	*	*	*	*
575	139	47	92	*	53	41	44	25	29	58	27
575.0	29	14	15	*	*8	*7	13	*8	*6	11	*
575.1	66	21	46	*	28	19	19	10	14	30	13
575.10	12	*	10	*	*5	*	*	*	*	*	*
575.11	42	12	30	*	19	12	11	*5	10	21	*6
575.12	12	*5	*6	—	*	*	*	*	*	*	*
575.3	*7	*	*5	—	*	*	*	*	*	*	*
575.8	24	*7	18	*	9	11	*	*	*	11	*5
576	122	50	71	*	18	34	66	23	26	41	31
576.1	43	23	21	*	*	10	29	9	11	11	12
576.2	31	11	20	*	*	11	15	*6	*5	12	*8
576.8	43	15	28	*	10	13	18	*7	*9	17	10
577	460	227	233	*6	159	170	125	82	93	196	89
577.0	333	163	170	*	110	118	100	60	67	146	60
577.1	86	45	41	*	35	38	12	13	18	35	21
577.2	19	10	*9	*	*8	*6	*	*	*	*7	*
577.8	13	*5	*8	*	*	*	*	*	*	*	*
577.9	9	*	*6	—	*	*	*5	*	*	*	*
578	453	211	243	14	54	95	290	114	117	146	77
578.0	61	33	27	*5	15	13	27	17	11	17	16
578.1	120	57	63	*	14	29	72	33	30	40	18
578.9	272	121	152	*	25	53	191	64	76	89	43
579	48	19	29	14	*8	11	14	*7	14	15	12
579.0	*6	*	*	*	*	*	*	*	*	*	*
579.3	22	*8	14	*8	*	*5	*7	*	*7	*6	*6
579.8	13	*6	*8	*5	*	*	*	*	*	*5	*
579.9	*5	*	*	*	*	*	*	*	*	*	*
580-629	8,123	2,827	5,296	180	2,141	2,021	3,781	1,633	1,906	3,122	1,462
580	*9	*	*	*	*	*	*	*	*	*	*
580.9	*5	*	*	*	*	*	*	*	*	*	*
581	39	21	18	*	*8	14	13	*8	*7	17	*8
581.8	17	10	*7	*	*	*6	*8	*	*	*8	*
581.81	17	10	*7	—	*	*6	*8	*	*	*8	*
581.9	18	*8	10	*	*	*8	*	*	*	*8	*
582	28	15	13	*	*8	*5	14	*9	*6	10	*
582.8	*7	*	*	*	*	*	*	*	*	*	*
582.81	*6	*	*	*	*	*	*	*	*	*	*
582.9	18	9	*8	—	*	*	12	*6	*5	*5	*
583	139	74	65	*	20	55	63	29	30	45	35
583.8	125	67	59	*	18	50	57	27	27	40	32
583.81	119	63	56	*	16	47	55	26	25	39	29
583.89	*7	*	*	*	*	*	*	*	*	*	*
583.9	10	*6	*	*	*	*	*	*	*	*	*
584	664	350	314	10	49	176	428	160	151	237	116
584.5	68	40	29	*	*6	25	36	19	11	20	19
584.8	10	*	*6	*	*	*	*	*	*	*	*
584.9	585	306	279	*6	41	150	388	138	138	215	94
585	266	157	108	*6	27	71	161	56	61	92	56
586	40	22	18	*	*	10	28	*9	10	16	*6
588	19	*7	12	*	*	*6	*7	*6	*	*7	*
588.8	14	*	*9	*	*	*	*6	*	*	*5	*
590	232	41	192	29	101	42	60	33	47	95	57
590.0	*6	*	*6	*	*	*	*	*	*	*	*
590.00	*6	*	*6	*	*	*	*	*	*	*	*
590.1	108	19	89	11	49	21	26	13	23	44	26

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
590.10	108	19	89	11	49	21	26	13	23	44	26
590.8	115	20	95	16	48	19	32	18	22	47	28
590.80	114	19	95	16	48	19	30	18	22	47	27
591	194	83	112	*9	53	52	81	48	47	62	38
592	288	143	145	*	128	88	68	65	72	106	45
592.0	126	60	66	*	53	40	31	32	30	48	16
592.1	159	83	76	*	73	47	37	32	41	57	29
593	692	358	334	21	52	129	490	143	187	253	109
593.2	15	10	*	—	*	*	*8	*	*	*5	*
593.3	12	*	*8	*	*	*	*	*	*	*	*
593.4	27	10	17	*	*	*8	13	*5	*7	11	*
593.5	9	*	*6	*	*	*	*	*	*	*	*
593.7	16	*5	11	14	*	*	*	*5	*	*	*
593.70	12	*	*8	11	*	*	*	*	*	*	*
593.8	18	*8	10	*	*	*6	*7	*	*	*6	*
593.81	*7	*	*	—	*	*	*	*	*	*	*
593.89	11	*	*7	*	*	*	*	*	*	*	*
593.9	594	318	276	*	32	107	452	117	163	223	91
594	21	18	*	*	*	*6	13	*	*	*9	*
594.1	19	16	*	*	*	*	13	*	*	*7	*
595	61	22	40	*	10	13	38	14	16	23	*9
595.0	13	*	9	*	*	*	*8	*	*	*	*
595.1	*8	*	*8	*	*	*	*	*	*	*	*
595.2	*7	*	*	—	*	*	*6	*	*	*	*
595.8	*8	*	*	—	*	*	*6	*	*	*	*
595.82	*6	*	*	—	—	*	*	*	*	*	*
595.9	24	11	13	*	*	*	15	*	*8	9	*
596	199	135	64	*7	24	50	118	41	58	63	37
596.0	66	64	*	*	*	12	53	13	21	21	11
596.1	*9	*6	*	*	*	*	*6	*	*	*	*
596.3	*7	*	*	*	*	*	*	*	*	*	*
596.4	10	*5	*	—	*	*	*8	*	*	*	*
596.5	74	38	36	*	17	25	29	16	25	21	12
596.51	*5	*	*	*	*	*	*	*	*	*	*
596.54	62	34	28	*	15	21	22	13	20	18	11
596.59	*6	*	*	*	*	*	*	*	*	*	*
596.7	*5	*	*	—	*	*	*	*	*	*	*
596.8	23	11	13	*	*	*6	11	*	*	*8	*7
597	*6	*	*	—	*	*	*	*	*	*	*
597.8	*5	*	*	—	*	*	*	*	*	*	*
598	30	25	*6	*	*6	*8	17	*7	*8	13	*
598.9	25	20	*5	*	*	*6	14	*5	*6	11	*
599	2,179	751	1,428	62	270	361	1,487	470	523	842	345
599.0	1,836	522	1,314	50	221	295	1,270	396	442	709	289
599.6	42	35	*8	*	*	*8	29	11	*7	20	*
599.7	282	188	94	*8	43	52	178	60	69	106	47
599.8	14	*	11	*	*	*5	*5	*	*	*5	*
599.81	*5	—	*5	—	*	*	*	*	*	*	*
600	449	449	—	*	*	79	368	110	102	146	91
600.0	403	403	—	*	*	68	334	103	93	126	81
600.1	*8	*8	—	—	—	*	*	*	*	*	*
600.9	33	33	—	—	*	*6	26	*	*6	16	*7
601	50	50	—	—	*	12	35	11	13	19	*7
601.0	16	16	—	—	*	*	11	*	*	*7	*
601.1	19	19	—	—	*	*	14	*5	*5	*6	*
601.9	13	13	—	—	*	*	*9	*	*	*	*
602	*6	*6	—	—	*	*	*6	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
603	9	9	—	*	*	*	*	*	*	*	*
603.9	*7	*7	—	*	*	*	*	*	*	*	*
604	15	15	—	*	*	*6	*	*	*	*5	*
604.9	14	14	—	*	*	*6	*	*	*	*	*
604.90	10	10	—	*	*	*	*	*	*	*	*
605	10	10	—	*	*	*	*	*	*	*	*
607	25	25	—	*	*6	10	*7	*	*	10	*6
607.8	15	15	—	*	*	*6	*	*	*	*6	*
607.84	*8	*8	—	—	*	*	*	*	*	*	*
607.89	*5	*5	—	*	*	*	*	*	*	*	*
608	29	29	—	*	*8	11	*6	*6	*6	13	*
608.4	11	11	—	*	*	*	*	*	*	*	*
608.8	11	11	—	*	*	*	*	*	*	*	*
608.86	*6	*6	—	*	*	*	*	*	*	*	*
610	20	*	19	—	*5	10	*	*5	*5	*6	*
610.1	16	*	16	—	*	*7	*	*	*	*5	*
611	51	*	46	*	21	18	12	12	12	17	11
611.0	13	*	13	*	*5	*	*	*	*	*5	*
611.1	15	*	12	*	*8	*5	*	*	*	*6	*
611.7	14	*	13	—	*	*	*6	*	*5	*	*
611.72	10	*	10	—	*	*	*	*	*	*	*
611.8	*7	*	*6	—	*	*	*	*	*	*	*
614	277	—	277	*	189	72	15	55	52	121	49
614.1	30	—	30	*	17	11	*	*7	*7	13	*
614.2	16	—	16	*	12	*	*	*	*	*7	*
614.3	*6	—	*6	*	*	*	*	*	*	*	*
614.4	10	—	10	—	*6	*	*	*	*	*6	*
614.6	183	—	183	*	122	52	*8	35	32	82	33
614.9	26	—	26	*	25	*	—	*6	*5	10	*5
615	27	—	27	*	25	*	*	*	*	16	*
615.0	*6	—	*6	*	*6	*	*	*	*	*	*
615.9	17	—	17	—	16	*	*	*	*	11	*
616	162	—	162	*	89	58	14	31	32	71	28
616.0	122	—	122	—	66	49	*8	24	25	53	21
616.1	28	—	28	*	16	*7	*	*	*6	14	*
616.10	28	—	28	*	16	*7	*	*	*5	14	*
617	337	—	337	*	220	106	10	53	76	139	69
617.0	198	—	198	—	110	78	9	35	45	81	38
617.1	46	—	46	—	35	11	*	*6	*9	18	13
617.2	11	—	11	—	*8	*	—	*	*	*	*
617.3	50	—	50	*	41	*9	*	*5	12	23	9
617.8	*6	—	*6	—	*5	*	*	*	*	*	*
617.9	21	—	21	—	19	*	—	*	*6	*8	*
618	254	—	254	—	58	108	88	41	55	107	51
618.0	64	—	64	—	12	31	22	*9	14	27	14
618.1	23	—	23	—	13	*8	*	*	*6	12	*
618.2	28	—	28	—	*7	11	9	*6	*7	10	*
618.3	19	—	19	—	*	*8	*8	*6	*	*6	*
618.4	28	—	28	—	*9	9	10	*5	*7	13	*
618.5	41	—	41	—	*	19	19	*	9	19	9
618.6	35	—	35	—	*7	14	14	*7	*6	14	*8
618.8	15	—	15	—	*	*7	*	*	*	*5	*5
619	15	—	15	—	*6	*	*6	*	*	*8	*
619.1	12	—	12	—	*	*	*5	*	*	*5	*
620	357	—	357	*6	223	112	17	57	82	147	71
620.0	63	—	63	*	43	18	*	*8	14	31	10
620.1	71	—	71	*	50	20	—	12	17	29	13

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
620.2	154	—	154	*	97	43	11	25	35	61	33
620.3	*7	—	*7	—	*	*5	*	*	*	*	*
620.5	9	—	9	*	*6	*	*	*	*	*	*
620.8	51	—	51	*	25	23	*	*9	13	17	12
621	89	—	89	—	36	37	16	18	22	36	14
621.0	37	—	37	—	11	19	*7	*7	*8	14	*7
621.2	10	—	10	—	*7	*	*	*	*	*	*
621.3	19	—	19	—	*7	*9	*	*	*	*7	*
621.8	13	—	13	—	*	*	*	*	*	*6	*
622	37	—	37	*	18	16	*	*5	12	13	*8
622.1	22	—	22	*	13	*8	*	*	*8	*9	*
622.7	9	—	9	—	*	*6	*	*	*	*	*
623	27	—	27	*	13	*8	*6	*6	*5	*9	*7
623.8	22	—	22	*	10	*7	*	*	*	*7	*6
624	*7	—	*7	*	*	*	*	*	*	*	*
625	367	—	367	*	215	110	42	43	85	168	71
625.0	31	—	31	—	27	*	*	*	*5	17	*6
625.3	101	—	101	*	82	19	*	*7	21	51	23
625.4	*6	—	*6	—	*	*	—	*	*	*	*
625.6	148	—	148	—	40	69	39	21	36	61	29
625.8	*7	—	*7	—	*	*	*	*	*	*	*
625.9	69	—	69	*	55	13	*	9	19	31	9
626	324	—	324	*	210	112	*	45	76	137	66
626.2	240	—	240	*	157	82	*	31	56	101	50
626.6	*7	—	*7	—	*	*	*	*	*	*	*
626.8	57	—	57	*	36	21	*	*9	13	26	10
626.9	11	—	11	*	*7	*	*	*	*	*5	*
627	60	—	60	—	*8	32	21	*7	15	23	15
627.0	*6	—	*6	—	*	*	—	*	*	*	*
627.1	30	—	30	—	*	17	12	*	*8	11	*7
627.2	12	—	12	—	*	*7	*	*	*	*5	*
627.9	*5	—	*5	—	*	*	*	*	*	*	*
628	*6	—	*6	*	*6	*	—	*	*	*	*
630-677	9,450	—	9,450	32	9,402	16	—	1,496	1,822	3,873	2,260
632	14	—	14	*	14	*	—	*	*	*7	*
633	31	—	31	—	31	*	—	*5	*5	11	9
633.1	25	—	25	—	25	*	—	*	*	10	*6
634	27	—	27	*	27	—	—	*	*	15	*
634.9	18	—	18	*	18	—	—	*	*	10	*
634.91	15	—	15	—	15	—	—	*	*	*8	*
635	*9	—	*9	*	*9	*	—	*	*	*	*
635.9	*6	—	*6	*	*5	*	—	*	*	*	*
639	*8	—	*8	*	*8	—	—	*	*	*	*
640	*9	—	*9	—	*9	—	—	*	*	*6	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*6	—	*6	—	*6	—	—	*	*	*	*
641	95	—	95	*	95	*	—	23	17	32	23
641.0	9	—	9	—	9	*	—	*	*	*	*
641.01	*8	—	*8	—	*8	—	—	*	*	*	*
641.1	22	—	22	—	21	*	—	*7	*	*7	*
641.11	13	—	13	—	13	*	—	*	*	*5	*
641.13	*9	—	*9	—	*9	—	—	*	*	*	*
641.2	42	—	42	*	42	—	—	11	*6	14	10
641.21	40	—	40	—	40	—	—	10	*6	14	10
641.8	*6	—	*6	—	*6	—	—	*	*	*	*
641.9	14	—	14	—	14	*	—	*	*	*	*
641.91	*7	—	*7	—	*7	*	—	*	*	*	*

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
641.93	*7	—	*7	—	*7	—	—	*	*	*	*
642	338	—	338	*	336	*	—	55	56	156	71
642.0	44	—	44	—	44	*	—	9	*5	23	*7
642.01	35	—	35	—	34	*	—	*7	*	17	*6
642.03	9	—	9	—	9	*	—	*	*	*6	*
642.3	125	—	125	*	124	*	—	16	21	60	27
642.31	109	—	109	*	109	*	—	14	20	50	25
642.33	13	—	13	—	13	*	—	*	*	*9	*
642.4	90	—	90	*	89	*	—	17	16	38	19
642.41	82	—	82	*	81	*	—	15	14	35	18
642.5	36	—	36	*	36	*	—	*6	*6	16	*9
642.51	33	—	33	*	32	*	—	*5	*6	13	*8
642.7	13	—	13	—	13	*	—	*	*	*7	*
642.71	11	—	11	—	11	—	—	*	*	*5	*
642.9	21	—	21	*	21	*	—	*	*	*8	*
642.91	15	—	15	*	15	—	—	*	*	*5	*
643	59	—	59	*	59	—	—	*8	11	27	13
643.0	23	—	23	—	23	—	—	*	*	12	*5
643.03	22	—	22	—	22	—	—	*	*	11	*
643.1	22	—	22	*	22	—	—	*	*	*8	*
643.13	22	—	22	*	22	—	—	*	*	*8	*
643.2	11	—	11	—	11	—	—	*	*	*7	*
643.23	10	—	10	—	10	—	—	*	*	*6	*
644	395	—	395	*	393	*	—	73	61	183	78
644.0	100	—	100	*	100	*	—	22	18	44	16
644.03	100	—	100	*	100	*	—	22	18	44	16
644.1	10	—	10	—	10	—	—	*	*	*6	*
644.13	10	—	10	—	10	—	—	*	*	*6	*
644.2	284	—	284	*	283	*	—	49	42	133	60
644.21	284	—	284	*	282	*	—	49	42	133	60
645	329	—	329	*	327	*	—	55	69	115	91
645.1	321	—	321	*	319	*	—	53	67	113	88
645.11	317	—	317	*	315	*	—	51	66	111	88
645.2	*9	—	*9	—	*9	—	—	*	*	*	*
645.21	*9	—	*9	—	*9	—	—	*	*	*	*
646	244	—	244	*	243	*	—	33	49	110	51
646.1	18	—	18	—	18	—	—	*	*	*7	*5
646.11	16	—	16	—	16	—	—	*	*	*6	*
646.2	16	—	16	*	16	—	—	*	*5	*6	*
646.21	*6	—	*6	—	*6	—	—	*	*	*	*
646.23	10	—	10	*	10	—	—	*	*	*	*
646.6	136	—	136	*	136	*	—	20	25	69	22
646.61	65	—	65	*	64	*	—	12	12	33	*8
646.62	*6	—	*6	—	*6	—	—	*	*	*	*
646.63	62	—	62	*	62	—	—	*7	12	31	12
646.8	60	—	60	—	60	—	—	*6	13	22	19
646.81	32	—	32	—	32	—	—	*	*7	9	12
646.83	23	—	23	—	23	—	—	*	*	11	*5
647	97	—	97	*	97	—	—	11	15	53	19
647.6	52	—	52	*	52	—	—	*5	*7	29	11
647.61	42	—	42	*	42	—	—	*	*6	23	9
647.63	*8	—	*8	—	*8	—	—	*	*	*	*
647.8	39	—	39	—	39	—	—	*	*7	21	*7
647.81	22	—	22	—	22	—	—	*	*	11	*
647.83	15	—	15	—	15	—	—	*	*	*9	*
648	1,428	—	1,428	*	1,424	*	—	242	294	585	308
648.0	42	—	42	*	42	*	—	*7	*5	24	*6
648.01	25	—	25	*	25	*	—	*6	*	12	*
648.03	17	—	17	—	17	*	—	*	*	11	*
648.1	53	—	53	—	53	*	—	13	12	18	9

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
648.11	46	—	46	—	46	*	—	11	11	15	*8
648.13	*6	—	*6	—	*6	—	—	*	*	*	*
648.2	277	—	277	*	276	*	—	46	50	125	57
648.21	167	—	167	*	166	—	—	25	29	82	31
648.22	78	—	78	*	78	*	—	15	14	28	20
648.23	23	—	23	*	23	—	—	*	*	10	*
648.24	9	—	9	—	9	—	—	*	*	*	*
648.3	11	—	11	—	11	—	—	*	*	*	*
648.31	*7	—	*7	—	*7	—	—	*	*	*	*
648.4	176	—	176	*	176	*	—	27	48	60	41
648.41	140	—	140	*	139	*	—	22	39	46	32
648.43	29	—	29	—	29	—	—	*5	*7	10	*7
648.44	*5	—	*5	—	*5	—	—	*	*	*	*
648.6	39	—	39	—	39	—	—	11	*8	14	*7
648.61	29	—	29	—	29	—	—	*9	*6	10	*
648.63	*7	—	*7	—	*7	—	—	*	*	*	*
648.7	*9	—	*9	—	*9	—	—	*	*	*	*
648.8	155	—	155	*	154	*	—	35	31	52	37
648.81	139	—	139	*	139	*	—	29	28	48	34
648.83	15	—	15	—	15	—	—	*6	*	*	*
648.9	662	—	662	*	660	*	—	98	136	282	146
648.91	536	—	536	*	534	*	—	72	112	230	122
648.92	13	—	13	—	13	—	—	*	*	*5	*
648.93	110	—	110	*	109	*	—	24	20	45	21
650	411	—	411	—	410	*	—	56	58	195	103
651	61	—	61	*	60	*	—	14	10	24	14
651.0	55	—	55	*	54	*	—	13	9	21	12
651.01	37	—	37	*	37	*	—	*7	*6	15	*9
651.03	18	—	18	*	18	—	—	*6	*	*5	*
652	299	—	299	*	297	*	—	48	58	131	62
652.1	*8	—	*8	—	*8	—	—	*	*	*	*
652.11	*6	—	*6	—	*6	—	—	*	*	*	*
652.2	133	—	133	*	133	—	—	20	27	60	27
652.21	130	—	130	*	130	—	—	19	26	58	26
652.3	29	—	29	*	29	*	—	*	*	15	*7
652.31	29	—	29	*	28	*	—	*	*	14	*7
652.4	*7	—	*7	—	*7	—	—	*	*	*	*
652.41	*7	—	*7	—	*7	—	—	*	*	*	*
652.5	43	—	43	*	42	*	—	*6	10	20	*8
652.51	43	—	43	*	42	*	—	*6	10	20	*8
652.6	24	—	24	—	24	*	—	*	*	11	*
652.61	23	—	23	—	23	*	—	*	*	11	*
652.8	48	—	48	*	48	—	—	*9	9	18	12
652.81	48	—	48	*	48	—	—	*9	9	18	12
653	103	—	103	*	103	—	—	13	15	47	29
653.4	86	—	86	*	86	—	—	10	11	40	25
653.41	86	—	86	*	86	—	—	10	11	40	25
653.5	12	—	12	—	12	—	—	*	*	*6	*
653.51	12	—	12	—	12	—	—	*	*	*6	*
654	630	—	630	*	628	*	—	102	110	277	141
654.0	12	—	12	—	12	—	—	*	*	*	*
654.01	11	—	11	—	11	—	—	*	*	*	*
654.1	27	—	27	—	27	*	—	*6	*	11	*6
654.11	25	—	25	—	25	*	—	*	*	10	*6
654.2	514	—	514	*	512	*	—	78	91	225	120
654.21	499	—	499	*	498	*	—	76	89	218	116
654.23	15	—	15	—	14	*	—	*	*	*7	*
654.4	23	—	23	*	23	—	—	*	*	*9	*6
654.41	19	—	19	*	19	—	—	*	*	*7	*5
654.5	31	—	31	—	31	*	—	*6	*	18	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
654.51	18	—	18	—	18	*	—	*	*	10	*
654.53	13	—	13	—	13	—	—	*	*	*7	*
654.6	13	—	13	*	13	—	—	*	*	*6	*
654.61	10	—	10	*	10	—	—	*	*	*5	*
654.9	*7	—	*7	—	*7	—	—	*	*	*	*
654.91	*6	—	*6	—	*6	—	—	*	*	*	*
655	48	—	48	*	47	*	—	*	*7	23	13
655.7	18	—	18	*	17	—	—	*	*	9	*
655.71	15	—	15	*	14	—	—	*	*	*7	*
655.8	23	—	23	—	22	*	—	*	*	11	*6
655.81	18	—	18	—	18	*	—	*	*	10	*
656	378	—	378	*	376	*	—	56	83	163	76
656.1	76	—	76	*	76	*	—	11	17	38	11
656.11	72	—	72	*	72	*	—	10	17	36	11
656.3	38	—	38	*	38	—	—	*5	10	17	*7
656.31	38	—	38	*	38	—	—	*5	10	17	*7
656.4	21	—	21	—	21	—	—	*	*	11	*5
656.41	21	—	21	—	21	—	—	*	*	11	*5
656.5	64	—	64	*	63	*	—	11	9	31	12
656.51	62	—	62	*	61	*	—	10	*9	31	12
656.6	90	—	90	*	90	*	—	11	20	37	22
656.61	90	—	90	*	89	*	—	11	20	37	22
656.7	32	—	32	—	32	—	—	*6	17	*6	*
656.71	30	—	30	—	30	—	—	*	17	*5	*
656.8	53	—	53	*	53	—	—	10	*7	21	15
656.81	53	—	53	*	53	—	—	10	*7	21	15
657	23	—	23	—	23	*	—	*	*	10	*
657.0	23	—	23	—	23	*	—	*	*	10	*
657.01	22	—	22	—	22	*	—	*	*	10	*
658	361	—	361	*	359	*	—	67	61	143	89
658.0	101	—	101	*	100	*	—	18	14	43	26
658.01	98	—	98	*	96	*	—	17	13	42	25
658.1	133	—	133	*	133	*	—	27	24	52	29
658.11	130	—	130	*	129	*	—	26	23	51	29
658.2	59	—	59	*	59	*	—	11	12	23	13
658.21	55	—	55	*	55	*	—	11	12	20	12
658.4	64	—	64	*	64	—	—	10	9	25	20
658.41	63	—	63	*	63	—	—	10	9	24	20
659	870	—	870	*6	859	*	—	132	165	358	216
659.1	45	—	45	*	44	*	—	*7	10	19	*8
659.11	36	—	36	*	36	*	—	*6	*8	15	*7
659.13	*8	—	*8	—	*8	*	—	*	*	*	*
659.2	48	—	48	*	48	*	—	*	*8	22	14
659.21	47	—	47	*	47	*	—	*	*7	22	14
659.4	20	—	20	—	20	*	—	*	*	*7	*8
659.41	20	—	20	—	20	*	—	*	*	*7	*8
659.5	41	—	41	—	40	*	—	*8	*9	12	12
659.51	40	—	40	—	39	*	—	*8	*9	11	12
659.6	219	—	219	*	216	*	—	35	34	86	65
659.61	207	—	207	*	205	*	—	33	32	80	62
659.63	12	—	12	—	11	*	—	*	*	*6	*
659.7	437	—	437	*	435	*	—	68	90	187	92
659.71	434	—	434	*	432	*	—	67	89	186	91
659.8	54	—	54	*	49	*	—	*6	11	22	15
659.81	53	—	53	*	48	—	—	*6	11	21	15
660	209	—	209	*	207	—	—	26	44	74	65
660.0	40	—	40	*	39	—	—	*6	*7	15	11
660.01	40	—	40	*	39	—	—	*6	*7	15	11
660.1	42	—	42	—	42	—	—	*	*6	13	19
660.11	42	—	42	—	42	—	—	*	*6	13	19

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
660.2	*6	—	*6	—	*6	—	—	*	*	*	*
660.21	*6	—	*6	—	*6	—	—	*	*	*	*
660.3	45	—	45	*	45	—	—	*5	*9	19	13
660.31	45	—	45	*	45	—	—	*5	*9	19	13
660.4	61	—	61	—	61	—	—	*7	19	18	17
660.41	61	—	61	—	61	—	—	*7	19	18	17
660.6	*6	—	*6	—	*6	—	—	*	*	*	*
660.61	*6	—	*6	—	*6	—	—	*	*	*	*
660.7	*8	—	*8	*	*7	—	—	*	*	*	*
660.71	*8	—	*8	*	*7	—	—	*	*	*	*
661	343	—	343	*	341	*	—	56	65	135	87
661.0	80	—	80	*	79	*	—	14	15	34	17
661.01	79	—	79	*	78	*	—	13	15	34	17
661.1	80	—	80	*	79	*	—	15	12	30	23
661.11	80	—	80	*	79	*	—	15	12	30	23
661.2	83	—	83	*	83	—	—	14	16	37	17
661.21	83	—	83	*	82	—	—	14	15	37	17
661.3	89	—	89	*	89	*	—	10	21	28	29
661.31	89	—	89	*	89	*	—	10	21	28	29
661.9	*7	—	*7	—	*7	—	—	*	*	*	*
661.91	*6	—	*6	—	*6	—	—	*	*	*	*
662	39	—	39	—	39	—	—	*7	11	11	9
662.0	9	—	9	—	9	—	—	*	*	*	*
662.01	*9	—	*9	—	*9	—	—	*	*	*	*
662.2	25	—	25	—	25	—	—	*	*7	*7	*6
662.21	25	—	25	—	25	—	—	*	*7	*7	*6
663	786	—	786	*	785	*	—	115	168	302	201
663.0	11	—	11	—	11	—	—	*	*	*5	*
663.01	10	—	10	—	10	—	—	*	*	*	*
663.1	130	—	130	*	130	*	—	14	38	42	36
663.11	129	—	129	*	129	*	—	14	38	41	36
663.2	41	—	41	*	41	*	—	*6	*9	14	13
663.21	41	—	41	*	41	*	—	*6	*9	14	12
663.3	563	—	563	*	563	*	—	86	112	228	138
663.31	563	—	563	*	563	*	—	86	112	228	138
663.4	16	—	16	*	16	—	—	*	*	*6	*5
663.41	16	—	16	*	16	—	—	*	*	*6	*5
663.8	21	—	21	*	21	—	—	*	*	*6	*7
663.81	21	—	21	*	21	—	—	*	*	*6	*7
664	1,221	—	1,221	*5	1,215	*	—	192	262	431	336
664.0	539	—	539	*	538	*	—	85	115	197	142
664.01	538	—	538	*	537	*	—	85	115	197	142
664.1	507	—	507	*	504	*	—	82	114	165	145
664.11	506	—	506	*	503	*	—	82	114	165	145
664.2	105	—	105	*	103	—	—	16	20	41	27
664.21	105	—	105	*	103	—	—	16	20	41	27
664.3	37	—	37	—	37	—	—	*	*7	13	12
664.31	37	—	37	—	37	—	—	*	*7	13	12
664.4	14	—	14	—	14	—	—	*	*	*7	*
664.41	14	—	14	—	14	—	—	*	*	*7	*
664.8	*9	—	*9	—	*9	—	—	*	*	*	*
664.81	*9	—	*9	—	*9	—	—	*	*	*	*
664.9	*7	—	*7	—	*7	—	—	*	*	*	*
664.91	*7	—	*7	—	*7	—	—	*	*	*	*
665	168	—	168	*	168	*	—	20	44	67	36
665.3	*9	—	*9	—	*9	—	—	*	*	*	*
665.31	*8	—	*8	—	*8	—	—	*	*	*	*
665.4	46	—	46	*	46	—	—	*6	10	21	*9
665.41	46	—	46	*	46	—	—	*6	10	21	*9
665.5	101	—	101	*	101	*	—	12	32	35	22

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
665.51	101	—	101	*	101	*	—	12	32	35	22
666	113	—	113	*	112	*	—	15	23	42	33
666.0	14	—	14	—	14	*	—	*	*	*	*
666.02	13	—	13	—	13	*	—	*	*	*	*
666.1	84	—	84	*	83	*	—	11	18	30	25
666.12	82	—	82	*	82	*	—	11	18	29	24
666.2	11	—	11	—	11	*	—	*	*	*6	*
666.22	*7	—	*7	—	*7	—	—	*	*	*	*
667	21	—	21	—	21	—	—	*6	*	*6	*5
667.0	14	—	14	—	14	—	—	*	*	*	*
667.02	14	—	14	—	14	—	—	*	*	*	*
667.1	*7	—	*7	—	*7	—	—	*	*	*	*
667.12	*7	—	*7	—	*7	—	—	*	*	*	*
668	18	—	18	—	18	—	—	*	*5	*8	*
668.8	17	—	17	—	17	—	—	*	*5	*7	*
668.81	*6	—	*6	—	*6	—	—	*	*	*	*
668.82	10	—	10	—	10	—	—	*	*	*	*
669	127	—	127	*	125	—	—	17	20	57	32
669.4	*5	—	*5	—	*5	—	—	*	*	*	*
669.5	59	—	59	*	59	—	—	*6	9	33	10
669.51	59	—	59	*	59	—	—	*6	9	33	10
669.7	*7	—	*7	*	*7	—	—	*	*	*	*
669.71	*7	—	*7	*	*7	—	—	*	*	*	*
669.8	48	—	48	*	47	—	—	*8	*8	16	16
669.81	45	—	45	*	45	—	—	*8	*8	16	14
670	26	—	26	*	26	*	—	*	*	15	*
670.0	26	—	26	*	26	*	—	*	*	15	*
670.02	19	—	19	*	18	—	—	*	*	10	*
670.04	*8	—	*8	*	*8	*	—	*	*	*	*
671	25	—	25	—	25	—	—	*	*	10	*6
671.8	13	—	13	—	13	—	—	*	*	*6	*
671.81	*8	—	*8	—	*8	—	—	*	*	*	*
672	17	—	17	*	16	—	—	*	*	*5	*6
672.0	17	—	17	*	16	—	—	*	*	*5	*6
672.02	16	—	16	*	16	—	—	*	*	*	*
674	42	—	42	*	42	—	—	*6	*6	21	9
674.3	16	—	16	—	16	—	—	*	*	*9	*
674.32	*9	—	*9	—	*9	—	—	*	*	*	*
674.34	*8	—	*8	—	*8	—	—	*	*	*	*
674.8	18	—	18	—	18	—	—	*	*	10	*
674.82	12	—	12	—	12	—	—	*	*	*7	*
674.84	*6	—	*6	—	*6	—	—	*	*	*	*
675	*7	—	*7	—	*7	—	—	*	*	*	*
675.2	*5	—	*5	—	*5	—	—	*	*	*	*
676	13	—	13	—	13	—	—	*	*	9	*
676.2	*7	—	*7	—	*7	—	—	*	*	*	*
676.22	*6	—	*6	—	*6	—	—	*	*	*	*
680-709	2,116	1,022	1,095	176	412	561	968	482	500	776	358
680	*7	*	*	*	*	*	*	*	*	*	*
681	57	34	23	*	14	18	21	12	16	19	10
681.0	19	13	*7	*	9	*	*	*	*5	*5	*
681.00	15	9	*5	*	*6	*	*	*	*	*	*
681.1	37	21	16	*	*	13	17	*7	11	14	*
681.10	34	19	15	*	*	12	15	*6	10	13	*
682	881	428	452	55	190	261	374	209	203	316	153
682.0	52	24	28	14	16	*9	14	13	11	19	9
682.1	14	*7	*7	*	*	*5	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
682.2	112	39	74	*9	33	37	34	21	18	55	19
682.3	90	48	41	*7	29	28	25	21	20	32	17
682.4	39	22	17	*	12	12	12	*9	*8	13	9
682.5	18	*6	12	*	*7	*	*	*	*	*7	*
682.6	421	206	215	10	72	121	218	108	109	136	68
682.7	121	72	49	*7	15	42	57	28	26	43	23
682.8	*6	*	*	*	*	*	*	*	*	*	*
682.9	*8	*	*6	*	*	*	*	*	*	*	*
683	10	*	*6	*8	*	*	*	*	*	*	*
684	*7	*	*	*	*	*	*	*	*	*	*
685	*7	*	*	*	*	*	*	*	*	*	*
686	17	*8	*9	*	*5	*	*5	*	*	*6	*
686.9	11	*5	*6	*	*	*	*	*	*	*	*
690	9	*6	*	*	*	*	*	*	*	*	*
690.1	*8	*6	*	*	*	*	*	*	*	*	*
690.10	*6	*	*	*	*	*	*	*	*	*	*
691	25	14	11	22	*	*	*	*	*7	*7	*8
691.0	22	12	9	21	*	*	*	*	*	*6	*8
692	52	25	27	15	13	10	15	13	13	14	13
692.9	45	22	22	13	11	*7	14	11	11	11	11
693	51	15	36	*6	12	13	21	14	12	14	11
693.0	50	14	36	*	12	13	21	14	12	13	11
694	*5	*	*	*	*	*	*	*	*	*	*
695	34	13	21	*5	*7	10	12	*7	*6	14	*6
695.1	9	*6	*	*	*	*	*	*	*	*	*
695.3	*8	*	*	—	*	*	*	*	*	*	*
695.8	*5	*	*	*	*	*	*	*	*	*	*
695.89	*5	*	*	*	*	*	*	*	*	*	*
696	39	23	16	*	11	16	11	13	11	*9	*7
696.0	*7	*	*	*	*	*	*	*	*	*	*
696.1	31	20	11	*	*9	13	*9	10	*9	*6	*6
698	39	9	29	*	20	*7	11	*6	12	15	*6
698.9	29	*6	23	*	16	*5	*8	*	*9	12	*
701	26	11	15	*5	*8	10	*	*	*	15	*
701.4	9	*7	*	*	*	*	*	*	*	*9	*
702	*9	*	*6	—	*	*	*	*	*	*	*
703	10	*	*6	*	*	*	*	*	*	*	*
703.8	*5	*	*	*	*	*	*	*	*	*	*
704	12	*	*7	*	*6	*	*	*	*	*7	*
704.0	*6	*	*	*	*	*	*	*	*	*	*
705	*7	*	*	*	*	*	*	*	*	*	*
705.8	*6	*	*	*	*	*	*	*	*	*	*
706	20	10	10	*	9	*	*5	*5	*5	*6	*
706.1	*9	*	*	*	*6	*	*	*	*	*	*
706.2	*9	*	*	*	*	*	*	*	*	*	*
707	687	342	346	*	66	168	449	164	177	240	107
707.0	328	155	174	*	30	55	241	76	77	128	48
707.1	340	179	161	*	32	109	199	83	95	104	57
707.10	34	14	20	*	*	9	23	10	*8	10	*6
707.11	*7	*	*	—	*	*	*	*	*	*	*
707.12	28	10	17	*	*	*7	18	*8	*9	*	*7
707.13	26	12	14	*	*	*	17	*5	10	*6	*5
707.14	52	28	25	*	*	15	34	16	13	17	*7
707.15	149	89	60	*	15	58	75	32	43	50	24
707.19	45	22	22	*	*	13	27	10	12	16	*6
707.8	17	*7	10	*	*	*	*8	*	*	*7	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
708	22	9	13	*	*6	*7	*	*	*5	*9	*
708.0	12	*	*8	*	*	*	*	*	*	*6	*
708.9	*8	*	*	*	*	*	*	*	*	*	*
709	78	38	39	35	22	*8	12	*7	*9	55	*7
709.0	*5	*	*	*	*	*	*	*	*	*	*
709.2	53	28	25	31	17	*	*	*	*	47	*
709.8	9	*	*	*	*	*	*	*	*	*	*
709.9	*7	*	*	*	*	*	*	*	*	*	*
710-739	6,230	2,173	4,057	93	908	1,789	3,440	1,289	1,654	2,125	1,162
710	189	30	159	*	60	77	51	35	36	85	32
710.0	135	19	116	*	53	57	24	23	27	63	22
710.1	19	*	16	*	*	*9	*7	*	*	*8	*
710.2	16	*	13	*	*	*	10	*	*	*5	*
710.3	*5	*	*	*	*	*	*	*	*	*	*
710.4	11	*	*7	—	*	*	*	*	*	*	*
711	38	25	13	*	*7	11	15	*	10	15	*8
711.0	33	22	11	*	*6	10	14	*	*9	13	*7
711.06	14	10	*	*	*	*6	*5	*	*	*5	*
712	*7	*	*	—	*	*	*5	*	*	*	*
712.3	*5	*	*	—	—	*	*	*	*	*	*
713	13	*6	*7	*	*	*8	*	*	*	*8	*
713.5	10	*	*6	—	*	*6	*	*	*	*7	*
714	252	60	192	*	31	65	152	55	73	88	36
714.0	234	56	178	*	26	62	145	51	69	81	33
714.3	9	*	*8	*	*	*	*	*	*	*	*
714.30	9	*	*8	*	*	*	*	*	*	*	*
715	1,687	566	1,121	*	56	429	1,202	364	467	577	280
715.0	39	12	28	—	*	*	33	15	*9	11	*
715.00	*8	*	*	—	—	*	*7	*	*	*	*
715.09	31	*9	22	—	*	*	26	13	*8	*7	*
715.1	62	22	40	—	*	24	36	12	10	25	15
715.15	19	*9	10	—	*	*6	12	*	*	*5	*
715.16	37	13	25	—	*	16	20	*7	*6	16	*9
715.3	401	156	245	—	13	124	264	89	112	128	71
715.30	*6	*	*	—	*	*	*	*	*	*	*
715.31	18	*7	11	—	*	*	15	*	*	*	*5
715.35	106	44	62	—	*	31	71	30	31	26	18
715.36	255	95	160	—	*8	81	166	53	71	88	43
715.37	*8	*	*	—	*	*	*	*	*	*	*
715.8	66	21	46	—	*	14	49	10	27	23	*6
715.89	66	21	45	—	*	14	49	10	27	23	*6
715.9	1,114	354	759	*	36	259	818	236	307	389	182
715.90	749	226	524	*	21	151	577	157	203	275	114
715.91	20	9	11	—	*	*6	13	*	*6	*8	*
715.95	101	39	62	*	*6	28	66	23	27	28	23
715.96	208	71	136	—	*5	66	137	42	64	67	35
715.97	*7	*	*	—	*	*	*	*	*	*	*
715.98	25	*6	19	—	*	*5	19	*7	*	*8	*7
716	277	100	176	*	27	96	152	44	82	107	43
716.1	9	*	*	*	*	*	*	*	*	*	*
716.5	15	*	10	*	*	*	*9	*	*6	*	*
716.59	10	*	*7	*	*	*	*7	*	*	*	*
716.9	249	89	161	*	24	85	139	37	73	100	39
716.90	176	61	116	*	13	58	105	26	46	76	29
716.91	12	*6	*6	—	*	*5	*5	*	*	*	*
716.94	*5	*	*	—	*	*	*	*	*	*	*
716.95	*7	*	*	*	*	*	*	*	*	*	*
716.96	24	9	15	*	*	*8	12	*	*9	*9	*
716.98	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
716.99	14	*	9	*	*	*6	*5	*	*7	*	*
717	33	18	14	*	16	10	*5	*6	*	*8	14
717.4	*5	*	*	—	*	*	*	*	*	*	*
717.7	*8	*	*	*	*	*	*	*	*	*	*
717.8	9	*	*5	*	*6	*	*	*	*	*	*5
717.83	*7	*	*	*	*	*	—	*	*	*	*
718	48	24	24	*6	13	14	15	10	10	16	12
718.3	9	*	*6	*	*	*	*	*	*	*	*
718.4	16	*9	*7	*	*	*	*7	*	*	*8	*
718.46	*6	*	*	*	*	*	*	*	*	*	*
718.5	*8	*	*	*	*	*	*	*	*	*	*
718.56	*5	*	*	—	*	*	*	*	*	*	*
718.8	*6	*	*	*	*	*	*	*	*	*	*
719	234	92	142	*6	49	57	122	47	75	60	52
719.0	20	10	10	*	*	*5	10	*	*	*6	*6
719.06	16	*9	*7	*	*	*	*9	*	*	*	*5
719.4	147	54	93	*	39	36	69	20	44	49	35
719.40	*6	*	*	—	*	*	*	*	*	*	*
719.41	43	18	26	*	13	12	18	*6	15	13	10
719.45	48	16	32	*	*5	*8	33	*5	16	16	11
719.46	24	*7	17	*	*6	*7	11	*	*	9	*7
719.47	*5	*	*	*	*	*	*	*	*	*	*
719.49	10	*	*6	*	*6	*	*	*	*	*	*
719.5	*6	*	*	—	*	*	*	*	*	*	*
719.7	51	18	32	*	*	12	37	18	25	*	*5
719.70	21	*9	12	—	*	*	16	*8	11	*	*
719.75	13	*	*9	*	*	*	10	*	*7	*	*
719.76	14	*	11	*	*	*6	*8	*5	*6	*	*
720	*9	*6	*	*	*	*	*	*	*	*	*
720.0	*6	*	*	—	*	*	*	*	*	*	*
721	282	115	167	*	31	88	163	47	83	94	58
721.0	72	31	40	—	9	30	32	10	17	25	19
721.1	27	16	11	—	*	9	15	*5	*6	13	*
721.2	*6	*	*	—	*	*	*	*	*	*	*
721.3	114	46	68	*	11	31	73	19	36	34	25
721.9	53	16	37	—	*5	12	36	10	20	16	*7
721.90	53	16	37	—	*5	11	36	10	19	16	*7
722	563	275	288	*	187	234	141	100	130	216	116
722.0	66	29	37	—	28	31	*8	11	13	32	11
722.1	227	127	100	*	96	91	39	47	55	78	48
722.10	223	124	99	*	94	90	39	46	53	76	48
722.2	*6	*	*	—	*	*	*	*	*	*	*
722.4	37	17	21	—	*8	18	11	*	*8	14	10
722.5	119	53	66	—	31	44	44	15	29	44	30
722.52	116	52	64	—	31	43	42	15	28	43	29
722.6	20	*5	15	—	*	*7	9	*	*6	*9	*
722.7	28	13	15	—	*6	15	*7	*5	*5	12	*
722.71	19	10	9	—	*	11	*	*	*	*8	*
722.73	*7	*	*	—	*	*	*	*	*	*	*
722.8	17	*7	11	—	*	*7	*5	*	*	10	*
722.83	15	*6	9	—	*	*6	*	*	*	9	*
722.9	41	22	19	*	*8	17	16	9	*9	14	9
722.91	*7	*	*	—	*	*	*	*	*	*	*
722.93	30	16	14	*	*6	12	12	*7	*7	10	*6
723	84	43	41	*	21	35	25	14	23	35	13
723.0	17	12	*5	—	*	*8	*8	*	*	*9	*
723.1	39	17	22	*	14	13	10	*	14	14	*7
723.4	22	11	11	—	*	12	*6	*	*	11	*

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
724 . . . . .	511	207	304	*	120	165	223	113	146	156	96
724.0 . . . . .	139	58	80	—	*5	40	93	38	40	40	21
724.00 . . . . .	26	*7	20	—	*	*	23	*9	*9	*	*
724.02 . . . . .	110	51	60	—	*	37	69	29	30	35	16
724.2 . . . . .	161	66	95	*	54	53	53	31	45	49	36
724.3 . . . . .	14	*	10	*	*	*	*7	*5	*	*	*
724.4 . . . . .	32	13	19	*	*8	12	12	*8	*8	14	*
724.5 . . . . .	146	57	89	*	44	50	50	27	46	43	30
724.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
725 . . . . .	43	*9	34	*	*	*	41	*8	15	12	*8
726 . . . . .	109	52	57	*	16	46	46	23	30	26	29
726.0 . . . . .	*7	*	*5	—	*	*	*	*	*	*	*
726.1 . . . . .	36	17	19	—	*	17	17	*9	*9	*8	10
726.10 . . . . .	33	16	17	—	*	15	15	*9	*7	*8	*8
726.2 . . . . .	21	*8	12	—	*	11	*7	*	*6	*	*7
726.3 . . . . .	16	13	*	*	*	*7	*	*	*7	*	*
726.33 . . . . .	16	13	*	*	*	*7	*	*	*7	*	*
726.5 . . . . .	14	*	12	—	*	*	10	*	*	*6	*
726.6 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
727 . . . . .	70	26	44	*5	16	21	28	16	13	26	15
727.0 . . . . .	28	13	15	*	*9	*8	10	*6	*	12	*6
727.00 . . . . .	15	*6	9	*	*	*	*7	*	*	*7	*
727.05 . . . . .	*7	*5	*	—	*	*	*	*	*	*	*
727.1 . . . . .	*5	*	*5	*	*	*	*	*	*	*	*
727.4 . . . . .	11	*	*9	*	*	*	*6	*	*	*5	*
727.40 . . . . .	9	*	*7	*	*	*	*6	*	*	*	*
727.6 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
727.61 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
727.8 . . . . .	10	*	*5	*	*	*	*	*	*	*	*
728 . . . . .	143	76	67	*6	33	37	67	28	32	55	28
728.2 . . . . .	*9	*	*6	—	*	*	*7	*5	*	*	*
728.3 . . . . .	13	*6	*7	*	*	*	10	*	*	12	*
728.8 . . . . .	94	55	40	*	27	25	39	17	23	34	21
728.84 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
728.85 . . . . .	15	*7	*8	*	*	*6	*	*	*	*6	*
728.86 . . . . .	12	*7	*6	*	*	*	*	*	*	*	*5
728.89 . . . . .	62	39	23	*	18	12	31	12	16	23	11
728.9 . . . . .	16	*7	9	*	*	*	10	*	*	*6	*
729 . . . . .	345	110	235	9	82	127	126	55	94	124	72
729.1 . . . . .	144	23	121	*	40	75	28	27	34	56	27
729.2 . . . . .	10	*6	*	—	*	*6	*	*	*	*	*
729.3 . . . . .	*6	*	*5	*	*	*	*	*	*	*	*
729.5 . . . . .	77	30	46	*	21	22	30	12	22	24	19
729.8 . . . . .	101	45	56	*	17	19	63	12	32	33	24
729.81 . . . . .	17	*8	*9	*	*7	*	*6	*	*	*6	*
729.82 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
729.89 . . . . .	77	35	42	*	*8	15	53	10	26	25	18
730 . . . . .	123	77	46	*7	22	45	49	30	29	42	23
730.0 . . . . .	33	22	11	*	*5	12	13	*9	11	*8	*6
730.07 . . . . .	22	14	*8	*	*	*8	9	*6	*7	*5	*
730.1 . . . . .	25	15	10	*	*	9	10	*8	*	*8	*
730.17 . . . . .	13	*8	*	—	*	*7	*	*	*	*	*
730.2 . . . . .	64	40	25	*	11	23	26	13	13	26	13
730.25 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
730.26 . . . . .	*9	*6	*	*	*	*	*	*	*	*	*
730.27 . . . . .	29	18	11	*	*	12	15	*6	*6	10	*7
730.28 . . . . .	12	*8	*	*	*	*	*5	*	*	*	*
731 . . . . .	34	21	14	—	*	16	16	*7	9	14	*
731.0 . . . . .	*9	*5	*	—	—	*	*8	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
731.8	25	15	10	—	*	15	*7	*	*7	11	*
732	9	*6	*	*	*	*	*	*	*	*	*
733	889	149	740	9	52	140	688	210	225	285	169
733.0	555	53	502	*	11	61	483	138	144	173	101
733.00	535	50	486	*	10	58	467	132	140	168	95
733.01	*7	*	*7	—	*	*	*6	*	*	*	*
733.09	14	*	10	—	*	*	10	*	*	*	*
733.1	148	35	112	*	*	22	122	39	36	42	31
733.10	*5	*	*	—	—	*	*	*	*	*	*
733.11	*6	*	*	*	*	*	*	*	*	*	*
733.13	110	24	86	*	*	13	97	30	26	33	22
733.14	11	*	*9	*	*	*	*6	*	*	*	*
733.15	*6	*	*	*	*	*	*	*	*	*	*
733.19	*7	*	*6	—	*	*	*5	*	*	*	*
733.2	*6	*	*	*	*	*	*	*	*	*	*
733.4	40	16	24	*	*7	15	17	*7	10	17	*6
733.42	29	14	16	*	*6	12	11	*6	*7	13	*
733.6	34	10	23	*	10	14	*9	*6	*7	13	*8
733.8	46	19	28	*	13	10	21	9	9	18	9
733.81	11	*	*7	*	*	*	*6	*	*	*	*
733.82	36	15	21	*	11	*8	15	*6	*8	15	*6
733.9	60	14	46	*	*8	15	35	10	18	20	13
733.90	46	10	36	*	*	11	29	*8	14	14	10
733.99	*9	*	*6	*	*	*	*	*	*	*	*
735	20	*	18	*	*	*6	9	*	10	*	*
735.0	*7	*	*7	*	*	*	*	*	*	*	*
735.4	*6	*	*	—	*	*	*	*	*	*	*
736	43	24	19	*	*9	12	17	9	10	15	*9
736.4	*8	*	*	*	*	*	*5	*	*	*	*
736.42	*6	*	*	*	*	*	*	*	*	*	*
736.7	22	12	9	*	*7	*6	*7	*	*	*7	*5
736.79	17	10	*7	*	*5	*5	*6	*	*	*6	*
736.8	*7	*	*	*	*	*	*	*	*	*	*
737	86	21	65	10	21	15	40	20	22	29	16
737.1	17	*6	11	*	*	*	10	*	*	*8	*
737.10	14	*	9	*	*	*	9	*	*	*7	*
737.3	59	14	46	10	16	12	21	14	15	18	12
737.30	51	12	39	*7	14	10	19	12	14	16	9
737.4	10	*	*9	*	*	*	*9	*	*	*	*
737.43	*5	*	*	*	*	*	*5	*	*	*	*
738	65	20	44	*	13	22	26	14	12	18	21
738.1	*6	*	*	*	*	*	*	*	*	*	*
738.19	*6	*	*	*	*	*	*	*	*	*	*
738.4	51	16	35	—	*9	18	25	11	11	13	17
739	24	10	14	*	11	*	*7	18	*	*	—
739.2	*7	*	*	*	*	*	*	*6	*	*	—
739.3	10	*	*8	—	*8	*	*	10	*	*	—
740-759	714	368	345	405	154	87	68	182	141	186	205
741	27	14	13	*8	15	*	*	*7	*7	*8	*6
741.0	12	*6	*6	*6	*6	*	*	*	*	*	*
741.00	*8	*	*	*	*	*	*	*	*	*	*
741.9	14	*8	*7	*	10	*	*	*	*	*	*
741.90	11	*5	*6	*	*8	*	*	*	*	*	*
742	41	20	20	31	*6	*	*	10	*7	9	13
742.1	10	*5	*	*9	*	*	*	*	*	*	*
742.3	9	*5	*	*7	*	*	*	*	*	*	*
742.4	*5	*	*	*	*	*	*	*	*	*	*
742.5	*7	*	*	*6	*	*	*	*	*	*	*
742.59	*7	*	*	*6	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
743	*8	*	*	*6	*	*	—	*	*	*	*
744	*6	*	*	*	*	*	*	*	*	*	*
745	103	48	55	73	14	*8	*9	31	15	23	34
745.1	*8	*	*	*7	*	—	—	*	*	*	*
745.11	*5	*	*	*	*	—	—	*	*	*	*
745.2	12	*6	*6	*8	*	*	*	*	*	*	*
745.4	31	12	19	24	*	*	*	*7	*7	*7	10
745.5	43	20	23	27	*	*5	*7	13	*	11	14
746	53	29	24	28	12	*8	*5	18	*9	14	12
746.0	*6	*	*	*	*	*	*	*	*	*	*
746.4	10	*7	*	*	*	*	*	*	*	*	*
746.8	21	9	11	11	*	*	*	*6	*5	*7	*
746.89	11	*5	*6	*6	*	*	*	*	*	*	*
747	66	36	30	44	9	*	9	16	11	14	25
747.0	20	10	10	19	*	—	*	*	*	*	9
747.1	10	*7	*	*7	*	*	*	*	*	*	*
747.10	*7	*	*	*	*	*	*	*	*	*	*
747.3	10	*6	*	*9	*	*	*	*	*	*	*
747.6	10	*5	*5	*	*	*	*	*	*	*	*
747.61	*6	*	*	*	*	*	*	*	*	*	*
747.8	10	*5	*5	*	*	*	*	*	*	*	*
748	24	16	*8	23	*	—	*	*	*	*	11
748.3	17	12	*	17	—	—	*	*	*	*	*8
749	19	12	*7	18	*	*	—	*	*	*9	*
749.0	10	*	*5	*9	*	*	—	*	*	*6	*
749.00	*7	*	*	*7	*	—	—	*	*	*	*
749.2	*7	*6	*	*7	*	—	—	*	*	*	*
750	25	18	*7	18	*	*	*	*	*5	*7	*8
750.3	*6	*	*	*	*	*	*	*	*	*	*
750.5	13	11	*	13	*	—	—	*	*	*	*5
751	32	16	16	17	*6	*	*	*7	9	9	*7
751.0	*6	*	*	*	*	*	*	*	*	*	*
751.2	*6	*	*	*6	—	—	—	*	*	*	*
751.5	*7	*	*	*	*	*	*	*	*	*	*
752	37	13	24	14	16	*	*	*7	*7	10	13
752.1	10	—	10	*	*6	*	*	*	*	*	*
752.11	9	—	9	*	*6	*	*	*	*	*	*
752.3	*8	—	*8	—	*7	*	*	*	*	*	*
752.5	*5	*5	—	*	*	—	*	*	*	*	*
752.51	*5	*5	—	*	*	—	*	*	*	*	*
752.6	*6	*6	—	*	*	*	*	*	*	*	*
753	70	36	34	19	18	20	13	18	14	18	20
753.0	11	*	*6	*	*	*	*	*	*	*	*
753.1	35	20	15	*	*6	15	9	*8	*7	9	11
753.10	*7	*5	*	*	*	*	*	*	*	*	*
753.12	14	*7	*7	*	*	*9	*	*	*	*	*7
753.2	9	*6	*	*6	*	*	*	*	*	*	*
753.3	*6	*	*	*	*	*	*	*	*	*	*
754	27	14	13	19	*5	*	*	*6	*	*8	*9
754.2	*7	*	*	*	*	*	—	*	*	*	*
754.7	*6	*	*	*	*	*	*	*	*	*	*
755	17	*8	*8	*8	*	*	*	*5	*	*	*6
755.6	10	*	*6	*	*	*	*	*	*	*	*
755.63	*6	*	*	*	*	*	*	*	*	*	*
756	63	31	33	25	12	16	11	13	15	19	16
756.0	*9	*	*	*8	*	*	—	*	*	*	*
756.1	29	14	15	*	*7	10	*8	*6	9	*8	*6
756.12	18	*8	10	*	*	*8	*6	*	*7	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
756.7	*7	*	*	*5	*	*	*	*	*	*	*
756.71	*6	*	*	*	*	*	*	*	*	*	*
756.8	*6	*	*	*	*	*	*	*	*	*	*
757	12	*	*7	*5	*	*	*	*	*	*8	*
757.3	*9	*	*	*5	*	*	*	*	*	*7	*
757.32	*7	*	*	*	*	*	—	—	*	*6	—
758	47	26	21	23	15	*7	*	12	14	10	10
758.0	35	21	14	18	10	*6	*	*9	12	*8	*6
759	37	18	19	20	10	*	*	13	11	*8	*6
759.8	24	12	12	15	*6	*	*	11	*5	*	*
759.89	19	*9	10	15	*	*	*	*8	*	*	*
760-779	472	270	202	463	*	*	*	72	97	149	154
765	65	33	32	65	*	—	*	*9	13	25	18
765.0	9	*	*	9	*	—	—	*	*	*	*
765.1	56	29	28	56	*	—	*	*8	11	22	15
765.10	*8	*	*	*8	—	—	—	*	*	*5	*
765.15	*5	*	*	*5	—	—	*	*	*	*	*
765.16	*6	*	*	*5	—	—	*	*	*	*	*
765.17	*7	*	*	*7	—	—	—	*	*	*	*
765.18	12	*7	*6	12	—	—	—	*	*	*	*
765.19	12	*6	*6	12	—	—	—	*	*	*	*5
767	10	*6	*	9	*	—	—	*	*	*	*
769	30	16	14	29	*	—	*	*	*7	9	11
770	91	54	37	88	*	*	*	19	20	28	25
770.6	13	*8	*6	13	*	—	—	*	*	*	*5
770.7	20	14	*6	20	*	*	*	*6	*	*6	*
770.8	43	23	20	41	*	*	*	*7	10	13	13
771	36	21	15	35	*	*	*	*7	*6	9	13
771.8	28	17	11	27	*	*	*	*	*5	*8	10
772	10	*6	*	9	*	*	—	*	*	*	*
772.1	*5	*	*	*5	*	*	—	*	*	*	*
773	9	*6	*	9	*	—	—	*	*	*	*
774	83	50	33	83	*	—	—	10	16	26	31
774.2	33	20	13	33	—	—	—	*	*6	11	13
774.6	45	27	18	44	*	—	—	*6	9	13	17
775	37	20	16	36	*	*	*	*6	*6	11	14
775.4	*6	*	*	*6	—	*	—	*	*	*	*
775.5	15	9	*6	15	*	*	—	*	*	*	*8
775.6	*8	*	*	*7	*	—	—	*	*	*	*
776	17	10	*7	16	*	*	*	*	*	*8	*
776.6	10	*6	*	10	*	—	*	*	*	*5	*
777	10	*5	*	10	—	—	—	*	*	*	*
778	15	*8	*7	14	*	*	—	*	*	*	*
778.4	*8	*	*	*8	*	—	—	*	*	*	*
779	40	23	17	39	*	—	*	*5	10	10	14
779.0	*7	*	*	*7	*	—	—	*	*	*	*
779.3	27	16	11	27	*	—	—	*	*7	*6	11
780-799	8,221	3,649	4,572	700	1,798	2,244	3,479	1,673	2,231	2,856	1,461
780	2,464	1,150	1,314	239	523	643	1,059	564	628	848	425
780.0	125	54	71	*6	28	29	62	22	31	40	32
780.01	20	*8	12	*	*	*	12	*	*	*5	*7
780.02	*8	*	*	*	*	*	*5	*	*	*	*
780.09	93	39	54	*	22	22	44	17	24	28	24
780.1	18	*9	9	*	*	*	12	*	*5	*8	*
780.2	432	189	243	*	49	94	285	124	93	152	63
780.3	823	411	412	99	231	223	270	204	207	278	134

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
780.31	20	13	*7	20	*	*	*	*5	*	*7	*5
780.39	803	399	404	79	231	222	270	199	204	271	129
780.4	144	48	97	*	23	34	87	32	44	43	26
780.5	275	153	122	14	77	118	65	64	73	84	53
780.50	*5	*	*	*	*	*	*	*	*	*	*
780.52	30	10	20	*	*9	*8	14	*6	*8	10	*7
780.57	229	135	93	14	62	104	49	56	61	68	43
780.6	344	162	181	104	74	72	94	61	75	140	68
780.7	216	89	127	*6	23	49	139	37	73	70	36
780.79	211	88	123	*6	21	46	138	35	73	68	35
780.8	*5	*	*	*	*	*	*	*	*	*	*
780.9	82	33	49	*	14	19	44	15	24	32	10
781	244	99	145	9	21	48	166	66	69	70	40
781.0	23	*9	14	*	*5	*6	*9	*6	*5	*7	*
781.2	156	60	96	*	*8	28	119	45	42	47	23
781.3	49	23	26	*	*	11	31	10	16	13	10
781.9	10	*	*7	*	*	*	*	*	*	*	*
781.99	10	*	*7	*	*	*	*	*	*	*	*
782	218	100	118	27	47	59	86	34	68	70	46
782.0	54	29	25	*	20	18	14	*7	18	17	12
782.1	45	21	24	12	12	9	11	*8	15	13	*8
782.3	82	30	53	*	9	24	48	13	26	26	17
782.4	23	13	10	*	*	*	9	*	*5	9	*6
782.7	*8	*	*	*	*	*	*	*	*	*	*
783	196	88	108	65	25	26	80	42	49	65	41
783.0	39	15	23	*	*6	*7	22	*5	10	17	*6
783.2	48	23	25	*	10	11	22	*7	16	16	9
783.21	47	23	24	*	9	11	22	*7	16	16	*8
783.3	18	*9	*9	14	*	*	*	*	*	*6	*5
783.4	47	26	21	42	*	*	*	16	10	12	*8
783.40	24	15	9	20	*	*	*	*8	*	*6	*7
783.41	21	10	11	20	*	*	*	*8	*5	*6	*
783.7	36	13	23	*	*	*	31	*8	*6	11	11
784	374	152	222	23	101	101	149	65	113	124	72
784.0	185	62	122	11	83	53	37	34	49	70	32
784.2	12	*6	*6	*	*	*	*	*	*	*	*
784.3	61	27	34	*	*	15	42	*9	22	17	14
784.5	58	27	31	*	*	16	37	*8	20	17	14
784.7	42	21	21	*	*	10	24	10	12	12	*7
785	357	165	192	22	73	100	162	74	101	123	58
785.0	80	34	47	*5	25	22	29	13	27	28	13
785.1	40	13	27	*	*6	15	19	10	10	14	*6
785.2	36	13	23	*8	10	*5	12	*5	12	14	*
785.4	67	38	29	*	*7	19	41	17	12	27	12
785.5	87	44	43	*	11	26	47	22	29	22	14
785.50	*8	*	*	*	*	*	*	*	*	*	*
785.51	25	13	12	*	*	*8	14	*	9	*6	*
785.59	54	27	27	*	*6	16	29	15	18	13	*8
785.6	41	22	20	*	14	12	12	*7	*8	16	10
785.9	*5	*	*	*	*	*	*	*	*	*	*
786	1,434	641	793	75	278	537	543	288	356	549	241
786.0	208	91	117	54	37	45	73	37	61	74	36
786.03	20	*9	11	13	*	*	*	*	*6	*6	*
786.05	60	23	37	*	13	19	26	10	17	23	10
786.06	*7	*	*	*	*	*	*	*	*	*	*
786.07	10	*	*	*	*	*	*	*	*	*	*
786.09	104	49	55	31	17	19	36	19	32	34	19
786.1	*8	*	*	*6	*	*	*	*	*	*	*
786.2	35	13	22	*7	*7	*8	13	*6	*9	13	*8

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
786.3	49	30	19	*	11	13	24	12	12	18	*7
786.5	1,090	475	614	*	218	460	409	223	265	422	180
786.50	451	205	246	*	91	183	175	83	108	186	74
786.51	32	13	18	—	*	13	14	*	*8	12	*7
786.52	57	25	32	*	18	19	20	9	18	21	9
786.59	549	232	317	*	104	245	200	127	130	202	91
786.6	34	19	15	*	*	*9	21	*6	*8	14	*7
786.8	*6	*6	*	—	*	*	*	*	*	*	*
787	808	305	504	110	193	191	314	133	233	279	163
787.0	430	142	288	70	123	110	127	67	124	150	89
787.01	253	72	181	17	81	76	79	40	65	100	48
787.02	90	28	62	*5	27	25	33	14	29	24	23
787.03	87	43	44	48	15	*8	16	12	30	26	19
787.1	*8	*	*6	—	*	*	*	*	*	*	*
787.2	122	61	61	*	14	21	83	24	36	40	21
787.3	11	*	*8	*	*	*	*	*	*	*6	*
787.6	14	*7	*7	*	*	*	*	*	*	*	*
787.9	223	89	133	30	47	54	92	37	64	77	44
787.91	221	88	133	30	46	53	91	37	63	77	44
788	449	248	201	11	69	98	271	102	146	134	67
788.0	21	12	*9	*	11	*6	*	*	*6	10	*
788.1	14	*7	*7	*	*5	*	*	*	*5	*	*
788.2	251	170	81	*	34	51	163	66	76	71	38
788.20	211	140	71	*	30	43	136	53	65	61	33
788.29	35	26	9	*	*	*7	24	12	9	10	*
788.3	109	34	75	*	13	27	65	20	40	31	17
788.30	73	25	48	*	*8	15	47	16	29	20	*8
788.31	*9	*	*	—	*	*	*6	*	*	*	*
788.33	13	*	13	—	*	*6	*6	*	*6	*	*
788.39	10	*	*7	*	*	*	*	*	*	*	*
788.4	17	*8	*8	*	*	*	*8	*	*	*	*
788.41	11	*	*7	*	*	*	*6	*	*	*	*
788.5	*6	*	*	*	*	*	*	*	*	*	*
788.9	26	12	14	—	*	*	21	*	10	*8	*
789	621	249	372	45	214	186	177	117	159	237	109
789.0	445	157	288	35	179	112	119	76	119	172	78
789.00	171	69	102	19	62	40	50	25	40	75	32
789.01	45	11	34	*	22	13	*8	*8	11	17	10
789.02	11	*6	*	*	*	*5	*	*	*5	*	*
789.03	55	17	38	*5	34	*8	*8	11	13	20	12
789.04	20	*	16	*	9	*	*6	*	*6	*7	*
789.05	*5	*	*	*	*	*	*	*	*	*	*
789.06	52	19	33	*	15	16	18	9	15	20	*8
789.07	19	*7	12	*	*6	*	*7	*5	*	*8	*
789.09	66	20	46	*	27	19	18	12	25	22	*6
789.1	10	*	*7	*	*	*	*	*	*	*5	*
789.2	18	11	*6	*	*8	*	*	*	*	*	*6
789.3	18	*	13	*	*	*6	*7	*	*5	*9	*
789.30	*9	*	*7	*	*	*	*	*	*	*	*
789.39	*5	*	*	*	*	*	*	*	*	*	*
789.5	129	72	57	*	18	62	45	33	29	46	21
790	386	178	208	28	74	107	178	84	107	120	74
790.2	29	*8	20	*	12	10	*7	*6	*7	*8	*7
790.4	17	*7	10	*	*6	*6	*	*	*5	*	*
790.5	36	16	20	*	10	10	14	10	11	*8	*7
790.6	130	56	73	*8	17	37	68	26	38	48	18
790.7	111	59	52	16	19	31	46	23	27	36	24
790.9	58	28	30	*	*8	11	38	13	17	15	12
790.92	37	12	24	*	*	*7	25	10	11	*9	*7
790.93	11	11	—	—	—	*	*8	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
790.99	*8	*	*	*	*	*	*	*	*	*	*
791	26	10	16	*	*9	*7	*7	*	*8	11	*
791.0	13	*	*8	*	*6	*	*	*	*	*6	*
791.9	9	*	*6	*	*	*	*	*	*	*	*
792	64	12	52	*	40	*5	18	9	24	20	12
792.1	26	12	14	*	*	*	17	*6	*9	*8	*
792.3	38	—	38	*	38	*	—	*	15	11	*8
793	25	10	15	*	*	*6	14	*	10	*9	*
793.1	*8	*	*	*	*	*	*	*	*	*	*
794	138	68	70	*6	34	44	53	22	37	58	22
794.3	52	27	24	*	9	18	24	*7	20	17	*8
794.31	37	17	19	*	*7	14	14	*6	11	13	*6
794.39	12	*8	*	—	*	*	*8	*	*7	*	*
794.8	76	35	41	*5	22	23	26	13	14	37	13
795	34	10	23	*	19	*8	*6	14	*7	*7	*6
795.5	19	*8	11	*	11	*	*	*8	*	*	*5
795.7	10	*	*8	*	*	*	*	*5	*	*	*
795.79	9	*	*8	*	*	*	*	*5	*	*	*
796	64	25	39	*	27	17	18	*7	20	25	12
796.0	10	*5	*5	*	*	*	*	*	*	*5	*
796.2	43	16	27	*	21	12	*8	*	13	17	9
797	11	*	*7	*	*	—	11	*	9	*	*
799	305	134	172	30	45	63	167	45	89	105	67
799.0	108	50	58	19	16	26	47	16	29	43	19
799.1	10	*	*	*	*	*	*6	*	*	*	*
799.3	79	29	50	*	*	13	63	*	39	27	*8
799.4	40	23	18	*	*6	*8	26	*6	*6	17	11
799.9	66	27	39	9	17	15	24	15	12	13	26
800-999	6,311	3,226	3,085	459	1,961	1,503	2,388	1,232	1,503	2,255	1,321
800	19	14	*5	10	*6	*	*	*	*	*5	*7
800.0	*6	*	*	*	*	*	*	*	*	*	*
800.2	*5	*	*	*	*	*	*	*	*	*	*
801	47	35	12	*7	22	11	*7	9	12	13	13
801.0	19	14	*5	*	*6	*	*	*	*6	*5	*
801.01	*7	*6	*	*	*	*	*	*	*	*	*
801.1	*8	*6	*	*	*6	*	*	*	*	*	*
801.2	12	*9	*	*	*6	*	*	*	*	*	*
802	94	66	28	*6	52	21	15	17	18	44	16
802.0	17	10	*8	*	*7	*	*5	*	*	*7	*
802.2	21	17	*	*	14	*	*	*	*	12	*
802.4	18	14	*	*	11	*5	*	*	*	*8	*
802.6	12	*7	*	*	*5	*	*	*	*	*6	*
802.8	15	12	*	*	*8	*	*	*	*	*5	*
803	*7	*	*	*	*	*	*	*	*	*	*
805	165	77	88	*	53	35	75	29	39	61	35
805.0	34	22	13	*	11	13	10	*6	*8	11	9
805.02	11	*6	*	*	*	*	*5	*	*	*	*
805.07	*8	*7	*	*	*	*	*	*	*	*	*
805.2	41	17	24	*	14	*7	20	9	10	16	*6
805.4	69	29	40	*	18	12	37	11	19	27	12
805.6	15	*6	9	*	*8	*	*5	*	*	*6	*6
806	13	11	*	*	*8	*	*	*	*	*	*5
806.0	*7	*6	*	*	*	*	*	*	*	*	*
807	125	73	52	*	32	36	55	22	26	40	36
807.0	114	66	48	*	30	32	50	21	25	36	33
807.00	*9	*	*	*	*	*	*	*	*	*	*
807.01	25	14	11	*	*7	*8	10	*	*6	*8	*6

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
807.02	18	10	*8	*	*	*	10	*	*	*5	*7
807.03	16	*9	*7	*	*5	*	*6	*	*	*5	*
807.04	10	*6	*	—	*	*	*6	*	*	*	*
807.05	*6	*	*	—	*	*	*	*	*	*	*
807.09	25	15	10	*	*7	*8	9	*	*6	*8	*8
807.2	*6	*	*	—	*	*	*	*	*	*	*
808	117	43	74	*	30	14	71	16	34	42	25
808.0	24	16	*8	*	11	*	*8	*	*6	10	*6
808.2	70	17	52	*	11	*7	50	10	23	22	15
808.4	11	*	*7	*	*5	*	*	*	*	*	*
808.41	*6	*	*	*	*	*	*	*	*	*	*
808.8	11	*	*7	—	*	*	*8	*	*	*	*
810	25	14	11	*	10	*	9	*6	*6	*9	*
810.0	25	14	11	*	10	*	9	*6	*6	*9	*
810.00	13	*8	*6	*	*5	*	*	*	*	*6	*
810.02	*7	*	*	*	*	*	*	*	*	*	*
811	12	*8	*	*	*	*	*	*	*	*	*
811.0	12	*7	*	*	*	*	*	*	*	*	*
812	96	34	62	18	12	13	54	25	23	27	21
812.0	53	14	38	*	*	*8	41	13	13	17	10
812.00	13	*5	*8	*	*	*	11	*	*	*5	*
812.01	19	*	15	*	*	*	16	*5	*5	*6	*
812.03	*6	*	*	—	*	*	*	*	*	*	*
812.09	13	*	9	*	*	*	9	*	*	*	*
812.2	15	*6	9	*	*	*	*6	*	*	*6	*
812.21	11	*	*7	*	*	*	*	*	*	*	*
812.4	24	11	13	15	*	*	*6	*7	*6	*	*7
812.41	16	*8	*8	11	*	*	*	*5	*	*	*5
813	100	50	49	12	32	20	35	16	20	36	28
813.0	17	*8	*9	*	*	*	*6	*	*	*6	*
813.01	*5	*	*	*	*	*	*	*	*	*	*
813.05	*6	*	*	*	*	*	*	*	*	*	*
813.2	10	*6	*	*	*6	*	*	*	*	*	*
813.4	54	24	30	*	16	10	24	*8	12	19	15
813.41	11	*	*6	*	*	*	*	*	*	*	*
813.42	26	10	16	*	*7	*	14	*	*5	*8	*8
813.43	*6	*	*	*	*	*	*	—	*	*	*
813.44	12	*6	*6	*	*	*	*	*	*	*	*
814	*6	*	*	*	*	*	*	*	*	*	*
814.0	*6	*	*	*	*	*	*	*	*	*	*
815	14	10	*	*	*6	*	*	*	*	*7	*
815.0	13	*9	*	*	*	*	*	*	*	*6	*
816	18	12	*6	*	10	*	*	*	*	*7	*5
816.0	*8	*	*	*	*	*	*	*	*	*	*
816.1	10	*8	*	*	*7	*	*	*	*	*	*
816.11	*5	*	*	*	*	*	*	*	*	*	*
820	360	102	258	*	9	30	318	76	95	124	65
820.0	101	25	76	*	*	*7	91	23	27	34	17
820.00	*7	*	*6	—	—	*	*7	*	*	*	*
820.02	*9	*	*	*	—	*	*8	*	*	*	*
820.03	*7	*	*6	*	*	*	*5	*	*	*	*
820.09	77	18	59	*	*	*5	70	17	23	26	11
820.2	168	45	123	*	*	13	150	40	48	55	26
820.20	13	*	11	*	*	*	12	*	*	*	*
820.21	139	37	102	*	*	10	126	34	39	46	20
820.22	16	*6	9	*	*	*	12	*	*5	*5	*
820.8	87	29	58	*	*	*9	76	14	19	33	22

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
821	81	38	43	14	24	14	30	17	21	27	16
821.0	43	23	20	10	16	*5	13	*9	11	16	*7
821.00	10	*	*6	*	*	*	*	*	*	*	*
821.01	34	19	15	*8	14	*	*8	*7	10	11	*6
821.2	30	10	20	*	*	*7	17	*6	*9	*9	*7
821.20	10	*	*6	*	*	*	*	*	*	*	*
821.21	*5	*	*	*	*	*	*	*	*	*	*
821.23	11	*	*9	*	*	*	*7	*	*	*	*
822	28	14	14	*	*8	*7	14	*5	*6	10	*8
822.0	27	13	14	*	*7	*6	14	*	*5	9	*7
823	89	55	34	*7	42	20	20	21	24	28	17
823.0	38	20	18	*	13	11	12	*8	10	13	*7
823.00	28	13	15	*	11	*8	*8	*6	*7	9	*5
823.01	*6	*	*	*	*	*	*	*	*	*	*
823.2	22	17	*5	*	12	*	*	*	*6	*6	*5
823.20	*6	*	*	*	*	*	*	*	*	*	*
823.22	12	9	*	*	*6	*	*	*	*	*	*
823.3	10	*8	*	*	*6	*	*	*	*	*	*
823.32	*8	*6	*	*	*	*	*	*	*	*	*
823.8	14	*8	*7	*	*6	*	*	*	*	*5	*
823.82	10	*	*6	*	*	*	*	*	*	*	*
824	130	61	69	*5	53	37	34	23	34	46	26
824.0	10	*6	*	*	*	*	*	*	*	*	*
824.2	12	*	*7	*	*	*	*	*	*	*	*
824.4	30	12	18	*	10	*9	10	*7	*7	9	*6
824.6	26	10	16	*	10	*8	*8	*	*6	11	*7
824.8	38	22	16	*	19	10	*6	*8	12	12	*7
824.9	*7	*	*	*	*	*	*	*	*	*	*
825	47	27	20	*	25	13	*6	9	11	18	*9
825.0	13	10	*	*	*7	*	*	*	*	*	*
825.2	25	11	14	*	14	*	*	*	*5	10	*5
825.25	15	*6	9	*	*8	*	*	*	*	*	*
825.3	*6	*	*	*	*	*	*	*	*	*	*
826	*9	*6	*	*	*	*	*	*	*	*	*
826.0	*6	*	*	*	*	*	*	*	*	*	*
831	12	9	*	*	*8	*	*	*	*	*	*
831.0	12	9	*	*	*8	*	*	*	*	*	*
835	*6	*	*	*	*	*	*	*	*	*	*
835.0	*6	*	*	*	*	*	*	*	*	*	*
836	16	*7	10	*	*8	*	*	*	*	*7	*
836.0	*7	*	*	*	*	*	*	*	*	*	*
836.1	*5	*	*	—	*	*	*	*	*	*	*
839	*7	*	*	*	*	*	*	*	*	*	*
840	34	15	20	*	*	15	15	*6	11	11	*6
840.4	23	9	13	—	*	10	12	*	*7	*7	*
843	*6	*	*	*	*	*	*	*	*	*	*
844	27	15	12	*	14	*7	*6	*5	*8	*9	*
844.2	*7	*	*	*	*6	*	*	*	*	*	*
844.8	10	*7	*	—	*	*	*	*	*	*	*
844.9	*7	*	*	—	*	*	*	*	*	*	*
845	14	*7	*7	*	*7	*	*	*	*	*	*
845.0	13	*7	*6	*	*6	*	*	*	*	*	*
845.00	*8	*	*	*	*	*	*	*	*	*	*
846	*5	*	*	—	*	*	*	*	*	*	*
847	37	16	21	*	15	10	10	*5	*7	14	10
847.0	20	*9	11	*	*9	*6	*	*	*	*8	*6
847.2	11	*	*6	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
848	*7	*	*	—	*	*	*	*	*	*	*
850	69	42	26	11	29	11	18	18	13	18	20
850.0	18	10	*8	*5	*7	*	*	*5	*	*	*5
850.1	22	14	*8	*	10	*	*6	*5	*	*6	*6
850.5	15	11	*	*	*7	*	*	*	*	*	*6
850.9	13	*7	*6	*	*6	*	*	*	*	*	*
851	25	12	13	*	10	*	*9	*	*5	*8	*7
851.8	17	*8	9	*	*5	*	*7	*	*	*5	*5
852	59	33	26	*	16	*8	30	10	12	19	19
852.0	17	9	*7	*	*7	*	*6	*	*	*5	*
852.06	*5	*	*	*	*	*	*	*	*	*	*
852.2	36	19	17	*	*6	*6	22	*6	*7	11	12
852.20	10	*	*5	*	*	*	*7	*	*	*	*
852.21	15	*7	*8	*	*	*	*9	*	*	*	*6
853	9	*	*5	*	*	*	*	*	*	*	*
853.0	9	*	*5	*	*	*	*	*	*	*	*
854	12	*7	*	*	*7	*	*	*	*	*	*
854.0	12	*7	*	*	*7	*	*	*	*	*	*
860	70	48	21	*	30	20	18	12	12	27	18
860.0	37	25	13	*	17	9	10	*6	*6	14	12
860.2	11	*6	*	*	*	*	*	*	*	*	*
860.4	15	11	*	*	*	*6	*	*	*	*6	*
861	39	29	10	*	22	*9	*6	*7	9	13	10
861.2	32	25	*7	*	19	*7	*	*6	*7	11	*8
861.21	29	22	*7	*	17	*6	*	*5	*7	*9	*8
862	14	11	*	*	*5	*	*6	*	*	*	*7
862.2	*8	*6	*	*	*	*	*5	*	*	*	*5
862.29	*7	*6	*	*	*	*	*5	*	*	*	*5
863	18	15	*	*	13	*	*	*	*6	*6	*
864	21	13	*8	*	14	*	*	*	*	*8	*5
864.0	16	*8	*8	*	*9	*	*	*	*	*5	*
864.05	*7	*	*	*	*	*	*	*	*	*	*
864.1	*5	*	*	*	*	*	—	*	*	*	*
865	25	16	*9	*	16	*5	*	*	*	12	*5
865.0	23	14	*9	*	14	*	*	*	*	11	*
865.00	*6	*	*	*	*	*	*	*	*	*	*
866	12	*9	*	*	*8	*	*	*	*	*5	*
866.0	11	*8	*	*	*7	*	*	*	*	*	*
866.01	*6	*	*	*	*	*	*	*	*	*	*
867	22	16	*5	*	*	*	13	*	*6	9	*
867.0	19	15	*	*	*	*	12	*	*6	*8	*
868	10	*6	*	*	*8	*	*	*	*	*5	*
868.0	*6	*	*	*	*	*	*	*	*	*	*
870	14	10	*	*	*8	*	*	*	*	*	*
870.8	*6	*	*	*	*	*	*	*	*	*	*
871	9	*8	*	*	*6	*	*	*	*	*5	*
872	9	*6	*	*	*6	*	*	*	*	*	*
872.0	*6	*5	*	*	*	*	*	*	*	*	*
873	175	106	69	18	75	29	53	35	45	57	38
873.0	47	24	23	*	16	10	20	*9	11	19	*8
873.2	*8	*5	*	*	*	*	*	*	*	*	*
873.20	*7	*	*	*	*	*	*	*	*	*	*
873.4	88	58	30	*7	38	15	28	19	25	26	19
873.41	*6	*	*	*	*	*	*	*	*	*	*
873.42	48	31	17	*	17	*8	20	11	12	13	11
873.43	15	10	*5	*	*7	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
873.44	*7	*	*	*	*	*	*	*	*	*	*
873.49	*8	*6	*	*	*	*	*	*	*	*	*
873.5	*8	*	*	*	*	*	—	*	*	*	*
873.6	16	11	*	*7	*7	*	*	*	*5	*	*
873.63	*6	*	*	*	*	*	*	*	*	*	*
873.64	*6	*	*	*	*	*	*	*	*	*	*
874	*6	*	*	*	*	*	*	*	*	*	*
875	*8	*8	*	*	*7	*	*	*	*	*	*
875.0	*7	*7	*	*	*6	*	*	*	*	*	*
879	15	11	*	*	*8	*	*	*	*	*6	*
879.8	*6	*	*	*	*	*	*	*	*	*	*
880	10	*8	*	*	*7	*	*	*	*	*7	*
880.0	*7	*	*	*	*	*	*	*	*	*5	*
881	50	30	21	*	27	*7	12	*6	16	15	13
881.0	38	21	17	*	20	*5	10	*	13	12	10
881.00	16	*9	*7	*	11	*	*	*	*5	*6	*
881.01	12	*8	*	*	*	*	*6	*	*	*	*
881.02	11	*	*6	*	*8	*	*	*	*	*	*
881.1	*6	*	*	*	*	*	*	*	*	*	*
881.2	*6	*	*	*	*	*	*	*	*	*	*
882	25	16	9	*	14	*6	*	*	*	12	*
882.0	12	*6	*7	*	*8	*	*	*	*	*6	*
882.1	*8	*5	*	*	*	*	*	*	*	*	*
882.2	*6	*5	*	*	*	*	*	—	*	*	*
883	28	18	10	*	15	*7	*	*6	*7	10	*6
883.0	11	*5	*6	*	*	*	*	*	*	*	*
883.1	*7	*	*	*	*	*	*	*	*	*	*
883.2	10	*8	*	*	*7	*	*	*	*	*	*
886	*7	*5	*	*	*	*	*	*	*	*	*
886.0	*6	*	*	*	*	*	*	*	*	*	*
890	12	*7	*5	*	*6	*	*	*	*	*	*
890.0	*8	*	*	*	*	*	*	*	*	*	*
891	34	21	12	*	18	*5	*9	*8	*6	12	*7
891.0	23	14	*9	*	14	*	*	*6	*	*8	*
891.1	10	*6	*	*	*	*	*	*	*	*	*
892	14	*8	*6	*	*	*	*	*	*	*6	*
892.1	*9	*5	*	*	*	*	*	*	*	*	*
902	*5	*	*	*	*	*	*	*	*	*	*
903	12	10	*	*	*8	*	*	*	*	*	*
903.2	*5	*	*	*	*	*	*	*	*	*	*
905	60	26	35	*	16	17	24	12	18	19	12
905.0	*8	*	*	*	*	*	*	*	*	*	*
905.1	*8	*	*	*	*	*	*	*	*	*	*
905.2	*8	*	*5	*	*	*	*	*	*	*	*
905.3	16	*	13	—	*	*	14	*	*7	*	*
905.4	18	*8	9	*	*6	*7	*	*	*	*6	*
906	26	19	*8	12	9	*	*	*	*	20	*
906.9	14	10	*	10	*	—	—	—	—	14	—
907	44	31	12	*	23	14	*6	*9	14	15	*7
907.0	20	15	*5	*	9	*6	*	*5	*7	*	*
907.2	21	15	*6	*	13	*7	*	*	*6	10	*
908	16	11	*	*	*8	*	*	*	*	*	*6
908.9	12	9	*	*	*6	*	*	*	*	*	*
909	14	*8	*7	*	*	*	*6	*	*	*	*
909.2	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
910 . . . . .	40	26	15	*6	17	*7	9	*6	13	11	10
910.0 . . . . .	36	23	13	*6	15	*6	*9	*5	12	10	*9
911 . . . . .	16	*9	*7	*	10	*	*	*	*	*6	*
911.0 . . . . .	11	*7	*	*	*7	*	*	*	*	*	*
912 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
912.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
913 . . . . .	30	16	14	*	15	*	10	*	11	*8	*8
913.0 . . . . .	25	15	11	*	12	*	10	*	*9	*7	*8
914 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
916 . . . . .	28	16	12	*	11	*6	*6	*6	*6	*9	*7
916.0 . . . . .	18	12	*6	*	*7	*	*	*	*	*	*
917 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
919 . . . . .	24	15	9	*	12	*	*5	*	9	*6	*5
919.0 . . . . .	20	11	*9	*	11	*	*	*	*7	*	*
920 . . . . .	67	29	38	*7	16	11	33	11	16	23	17
921 . . . . .	22	11	11	*	*8	*	*6	*	*6	*7	*
921.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
921.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
921.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
922 . . . . .	67	28	38	*	19	14	32	10	17	22	18
922.1 . . . . .	31	12	19	*	*8	*8	14	*	*7	12	*8
922.2 . . . . .	16	10	*6	*	*7	*	*6	*	*	*	*7
922.3 . . . . .	16	*5	11	*	*	*	10	*	*5	*	*
922.31 . . . . .	11	*	*7	*	*	*	*7	*	*	*	*
922.32 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
923 . . . . .	47	17	30	*	17	*8	21	*8	16	12	10
923.0 . . . . .	22	*9	14	*	9	*	*8	*5	*5	*7	*5
923.00 . . . . .	17	*6	11	*	*7	*	*6	*	*	*5	*
923.1 . . . . .	13	*	*8	*	*	*	*7	*	*6	*	*
923.11 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
923.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
923.20 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
924 . . . . .	99	31	69	*	30	15	50	18	28	31	22
924.0 . . . . .	29	*7	23	*	*	*	22	*8	*9	*8	*
924.00 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
924.01 . . . . .	23	*	18	*	*	*	18	*7	*7	*6	*
924.1 . . . . .	29	*7	22	*	9	*5	14	*	10	10	*6
924.10 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
924.11 . . . . .	21	*	17	*	*7	*	10	*	*7	*7	*6
924.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
924.8 . . . . .	27	11	17	*	14	*	*8	*5	*8	*7	*7
933 . . . . .	13	*7	*6	*	*	*	*7	*	*	*6	*
933.1 . . . . .	12	*6	*6	*	*	*	*6	*	*	*5	*
934 . . . . .	13	*7	*5	*	*	*	*6	*	*	*	*
934.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
935 . . . . .	14	*8	*6	*	*	*	*6	*	*	*	*
935.1 . . . . .	10	*7	*	*	*	*	*	*	*	*	*
941 . . . . .	22	17	*	10	*7	*	*	*6	*5	*9	*
941.2 . . . . .	15	11	*	*6	*6	*	*	*	*	*7	*
941.29 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*
942 . . . . .	23	16	*6	11	*6	*	*	*	*	14	*
942.2 . . . . .	13	11	*	*8	*	*	*	*	*	*7	*
942.3 . . . . .	*8	*	*	*	*	*	*	*	*	*6	*
943 . . . . .	19	13	*7	*	10	*	*	*	*5	9	*
943.2 . . . . .	12	*8	*	*	*6	*	*	*	*	*6	—
943.21 . . . . .	*6	*	*	*	*	*	*	*	*	*	—

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
943.3	*6	*	*	*	*	*	*	*	*	*	*
944	14	*8	*6	*	*5	*	*	*	*	*7	*
944.2	*7	*	*	*	*	*	*	*	*	*	*
945	26	18	*8	*8	*7	*7	*	*	*7	12	*
945.2	12	9	*	*6	*	*	*	*	*	*5	*
945.3	11	*7	*	*	*	*	*	*	*	*6	*
946	*7	*7	*	*	*	*	*	*	—	*7	—
948	43	31	12	12	16	*9	*6	*	*7	27	*
948.0	29	20	10	*8	11	*5	*	*	*6	17	*
948.00	29	20	10	*8	11	*5	*	*	*6	17	*
948.1	*7	*6	*	*	*	*	*	*	*	*6	*
948.10	*6	*6	*	*	*	*	*	*	*	*	—
949	9	*	*	*7	*	*	—	*	—	9	—
949.0	*8	*	*	*7	*	—	—	—	—	*8	—
952	*7	*5	*	*	*	*	*	*	*	*	*
955	14	13	*	*	10	*	*	*	*	*6	*
958	41	28	13	*	17	*8	13	10	10	15	*6
958.3	11	*6	*	*	*	*	*	*	*	*	*
958.4	*7	*	*	*	*	*	*	*	*	*	*
958.7	*8	*6	*	*	*	*	*	*	*	*	*
958.8	*8	*6	*	*	*	*	*	*	*	*	*
959	82	46	36	13	31	14	23	15	23	27	17
959.0	50	30	19	10	19	*8	13	*8	14	16	11
959.01	45	27	18	*9	17	*6	13	*8	13	14	11
959.1	10	*5	*	*	*5	*	*	*	*	*	*
959.7	*6	*	*	*	*	*	*	*	*	*	*
962	10	*	*6	*	*	*	*	*	*	*	*
962.3	*8	*	*	*	*	*	*	*	*	*	*
963	*8	*	*6	*	*6	*	*	*	*	*	*
963.0	*7	*	*	*	*5	*	*	*	*	*	*
965	86	29	57	*6	58	15	*7	17	26	29	15
965.0	27	12	15	*	16	*8	*	*5	*6	11	*5
965.00	*6	*	*	—	*	*	*	*	*	*	*
965.01	*5	*	*	—	*	*	—	*	*	*	*
965.09	13	*5	*8	*	*7	*	*	*	*	*6	*
965.1	*7	*	*5	*	*6	*	*	*	*	*	*
965.4	31	*9	23	*	22	*	*	*6	9	11	*5
965.6	12	*	*8	*	10	*	—	*	*5	*	*
965.61	11	*	*8	*	10	*	—	*	*5	*	*
965.8	*5	*	*	*	*	*	*	*	*	*	*
966	16	*6	10	*	12	*	*	*	*6	*6	*
966.3	12	*	*9	*	10	*	*	*	*5	*	*
967	17	*6	11	*	9	*5	*	*	*	*9	*
967.8	*7	*	*5	*	*	*	*	*	*	*	*
968	16	11	*	*	10	*	*	*	*	*5	*
968.5	10	*8	*	*	*7	*	*	*	*	*	*
969	111	46	66	*5	72	28	*6	17	30	40	25
969.0	35	13	22	*	25	*7	*	*6	10	12	*8
969.3	10	*	*7	*	*7	*	*	*	*	*	*
969.4	43	17	26	*	23	15	*	*7	11	16	10
969.7	*8	*6	*	*	*6	*	—	*	*	*	*
969.8	*5	*	*	*	*	*	*	*	*	*	*
970	*6	*	*	*	*	*	—	*	*	*	*
971	*5	*	*	*	*	*	*	*	*	*	*
972	19	*9	11	*	*	*	10	*	*	*6	*6
972.1	*6	*	*	—	*	*	*6	*	*	*	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
972.6	*8	*	*	*	*	*	*	*	*	*	*
975	*7	*	*	*	*	*	*	*	*	*	*
977	15	*	12	*	*8	*	*	*	*	*6	*
977.9	11	*	*9	*	*6	*	*	*	*	*	*
980	21	13	*8	*	14	*6	*	*	*6	*7	*
980.0	17	10	*7	*	11	*5	*	*	*5	*6	*
986	*5	*	*	*	*	*	*	*	*	*	*
987	*8	*	*	*	*	*	*	*	*	*	*
989	10	*7	*	*	*	*	*	*	*	*5	*
989.5	*6	*	*	*	*	*	*	*	*	*	*
991	12	*8	*	*	*	*	*6	*	*	*	*
991.6	11	*7	*	*	*	*	*6	*	*	*	*
992	9	*7	*	*	*	*	*	*	*	*	*
994	11	*7	*	*	*	*	*	*	*	*	*
995	77	29	48	10	17	21	29	17	15	35	10
995.0	*6	*	*	*	*	*	*	*	*	*	*
995.1	16	*6	10	*	*	*5	*7	*	*	*7	*
995.2	30	10	20	*	*	*8	17	*5	*6	15	*
995.3	*9	*	*	*	*	*	*	*	*	*	*
995.5	*8	*	*	*6	*	*	*	*	*	*	*
995.8	*5	*	*	*	*	*	*	*	*	*	*
996	792	405	387	48	143	252	349	163	171	304	155
996.0	33	17	16	*	*	*	25	10	*8	11	*
996.01	18	*7	10	*	*	*	15	*	*	*8	*
996.02	*6	*	*	*	*	*	*	*	*	*	*
996.04	*7	*	*	—	*	*	*	*	*	*	*
996.1	30	11	19	*	*6	10	12	*	*5	16	*
996.2	25	13	12	10	11	*	*	*6	*5	*8	*6
996.3	9	*6	*	*	*	*	*	*	*	*	*
996.39	*6	*	*	*	*	*	*	*	*	*	*
996.4	118	45	72	*	14	27	74	25	27	41	25
996.5	27	13	15	*	*7	10	*8	*	*	10	*8
996.59	18	*8	9	*	*	*6	*7	*	*	*6	*6
996.6	237	128	109	18	44	74	101	52	58	83	43
996.61	12	9	*	—	*	*	*8	*	*	*	*
996.62	137	71	66	16	29	46	46	31	27	53	26
996.64	17	10	*7	—	*	*	13	*	*7	*5	*
996.66	28	13	15	*	*	*6	20	*5	10	*9	*
996.67	14	10	*	*	*	*	*	*	*	*	*
996.68	*6	*	*	*	*	*	*	*	*	*	*
996.69	14	*7	*7	*	*	*6	*5	*	*	*	*
996.7	258	139	119	*9	37	94	119	49	54	100	55
996.72	57	36	22	*	*	24	28	14	13	22	*9
996.73	69	36	33	*	12	31	26	13	9	31	17
996.74	65	33	32	*	*9	21	33	9	19	23	14
996.76	16	12	*	*	*	*	12	*	*	*8	*
996.77	13	*	*8	—	*	*5	*5	*	*	*	*
996.78	22	10	12	*	*6	*6	*8	*	*	*7	*6
996.79	*8	*	*6	*	*	*	*	*	*	*	*
996.8	54	33	21	*	20	27	*	10	*7	31	*5
996.81	35	21	14	*	15	16	*	*	*6	19	*
996.82	*9	*	*	*	*	*7	—	*	*	*6	*
997	589	286	303	16	100	178	295	135	145	181	128
997.0	31	13	17	*	*	10	16	*7	*9	9	*6
997.02	13	*6	*7	*	*	*	10	*	*	*	*
997.09	14	*6	*8	*	*	*6	*	*	*	*	*
997.1	127	76	51	*	*8	33	84	35	31	32	29
997.2	21	*8	13	*	*	*8	10	*	*6	*6	*5

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
997.3	112	51	61	*	20	33	53	27	29	34	23
997.4	182	75	107	*5	44	62	70	36	45	57	45
997.5	72	36	36	*	14	17	41	17	16	26	13
997.6	33	20	13	*	*	12	16	*7	*7	13	*5
997.62	20	13	*7	*	*	*8	11	*	*5	*7	*
997.69	12	*6	*5	*	*	*	*	*	*	*6	*
997.9	*9	*5	*	*	*	*	*	*	*	*	*
997.99	*7	*	*	*	*	*	*	*	*	*	*
998	720	321	400	23	150	234	314	158	159	252	151
998.1	219	111	108	*6	40	63	110	52	50	70	46
998.11	109	62	47	*6	20	26	57	25	27	32	25
998.12	99	45	55	*	16	33	50	26	22	33	18
998.13	10	*	*6	*	*	*	*	*	*	*	*
998.2	101	36	65	*	24	32	43	22	21	33	25
998.3	52	25	27	*	11	19	21	10	13	21	*9
998.5	205	92	113	*8	44	71	81	46	45	74	40
998.59	200	89	111	*8	44	68	80	44	44	73	39
998.6	14	*6	*8	*	*	*7	*	*	*	*7	*
998.8	118	48	70	*	24	40	51	24	25	43	27
998.83	18	*7	11	*	*	*8	*7	*	*	*6	*
998.89	96	40	56	*	21	29	42	19	19	36	22
998.9	*5	*	*	—	*	*	*	*	*	*	*
999	52	24	27	*	15	15	18	*9	12	18	12
999.2	20	9	11	*	*5	*7	*7	*	*	*7	*
999.3	11	*7	*	*	*	*	*	*	*	*	*
999.8	11	*6	*6	*	*	*	*5	*	*	*	*
999.9	*8	*	*5	*	*	*	*	*	*	*	*
V01-V82	12,740	3,824	8,916	308	6,049	2,123	4,261	2,342	3,000	4,733	2,665
V01	*7	*	*	*	*	*	*	*	*	*	*
V02	442	15	426	*	424	11	*	62	82	199	98
V02.5	415	*	413	*	410	*	*	55	78	189	92
V02.51	404	*	404	*	403	*	—	54	75	187	87
V02.59	*7	*	*5	*	*	*	*	*	*	*	*
V02.6	25	13	12	*	13	9	*	*6	*	*9	*6
V02.61	*7	*	*	*	*	*	*	*	*	*	*
V02.62	16	9	*6	—	*8	*6	*	*	*	*5	*
V04	11	*	10	*	10	*	*	*	*	*	*
V04.3	*9	—	*9	—	*9	—	—	*	*	*	*
V05	10	*6	*	9	*	*	—	*	*	*5	*
V05.3	*7	*	*	*7	*	*	—	*	*	*	*
V06	11	*	*8	*	*8	*	*	*	*	*	*5
V06.4	*7	—	*7	—	*7	—	—	*	—	*	*5
V07	51	*	48	*	43	*	*	9	11	15	16
V07.2	35	*	34	*	34	*	—	*7	*8	*9	12
V07.3	*6	*	*5	—	*	*	*	*	*	*	*
V07.39	*6	*	*5	—	*	*	*	*	*	*	*
V07.4	*6	—	*6	—	*	*	*	*	*	*	*
V08	52	32	20	*	34	16	*	17	*8	24	*
V09	92	48	44	*	23	21	44	18	22	42	11
V09.0	86	45	41	*	23	19	40	16	20	39	10
V10	991	413	578	*5	66	248	672	248	273	302	168
V10.0	208	99	110	*	*7	48	153	56	61	63	28
V10.01	*6	*	*	—	*	*	*	*	*	*	*
V10.02	*9	*	*	—	*	*	*6	*	*	*	*
V10.03	*8	*6	*	—	*	*	*5	*	*	*	*
V10.04	11	*6	*	—	*	*	*8	*	*	*	*
V10.05	141	61	80	*	*	26	110	36	43	41	21
V10.06	21	11	10	*	*	*7	13	*7	*	*8	*

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V10.09	*8	*	*	—	*	*	*	*	*	*	*
V10.1	69	33	36	*	*	19	49	17	22	21	*8
V10.11	68	33	36	*	*	18	49	17	22	21	*8
V10.2	16	12	*	*	*	*6	10	*7	*	*	*
V10.21	13	10	*	—	*	*	*9	*6	*	*	*
V10.3	247	*	244	*	16	75	157	61	59	83	45
V10.4	237	155	82	*	17	41	178	55	73	66	43
V10.41	27	—	27	—	9	10	*8	*6	*8	9	*
V10.42	26	—	26	—	*	*7	19	*6	*9	*6	*6
V10.43	24	—	24	*	*	*6	13	*	9	*	*6
V10.46	148	148	—	*	*	16	131	37	42	43	26
V10.5	81	49	32	*	*	14	63	18	20	27	15
V10.51	42	31	11	*	*	*6	35	10	12	12	*8
V10.52	34	16	19	*	*	*7	24	*8	*8	13	*6
V10.7	22	9	13	*	*	*8	11	*5	*5	*7	*
V10.72	*5	*	*	—	*	*	*	*	*	*	*
V10.79	17	*8	10	*	*	*6	10	*	*	*	*
V10.8	105	51	54	*	17	36	49	27	27	30	21
V10.82	30	16	14	—	*5	12	14	10	*6	*8	*7
V10.83	33	17	16	—	*	12	21	*6	11	11	*6
V10.85	*6	*	*	*	*	*	*	*	*	*	*
V10.87	16	*5	10	*	*	*6	*5	*	*	*6	*
V10.89	10	*	*6	*	*	*	*	*	*	*	*
V11	35	19	16	—	11	11	13	*	16	*8	*7
V11.3	18	14	*	—	*	*6	*8	*	*8	*	*
V11.8	*7	*	*	—	*	*	*	*	*	*	*
V12	828	360	468	10	81	201	536	166	233	288	141
V12.0	41	17	23	*	14	11	14	*8	*6	17	9
V12.01	12	*7	*6	—	*	*6	*	*	*	*	*
V12.02	10	*	*7	*	*	*	*7	*	*	*	*
V12.09	18	*8	10	*	11	*	*	*	*	*9	*
V12.2	13	*	10	*	*	*6	*6	*	*6	*	*
V12.3	*5	*	*	*	*	*	*	*	*	*	—
V12.4	12	*7	*6	*	*	*	*7	*	*	*	*
V12.49	*8	*	*	*	*	*	*	*	*	*	*
V12.5	595	248	347	*	44	137	410	129	164	207	94
V12.50	22	*8	14	*	*	*	19	*	*9	*8	*
V12.51	159	58	101	*	28	49	82	31	47	48	32
V12.52	*6	*	*	—	*	*	*	*	*	*	*
V12.59	408	179	228	*	15	85	304	93	106	150	58
V12.6	18	*7	11	*	*	*	10	*	*	*8	*
V12.7	142	74	68	*	16	40	85	20	47	44	32
V12.71	87	46	41	*	10	27	50	*9	29	24	25
V12.72	27	16	11	*	*	*6	19	*	9	11	*
V12.79	25	11	14	*	*	*7	14	*7	*7	*8	*
V13	86	42	45	11	27	18	30	15	27	27	17
V13.0	55	29	26	*	17	15	22	10	19	18	*8
V13.01	36	20	16	*	14	13	9	*7	11	12	*6
V13.09	18	*8	10	*	*	*	12	*	*7	*5	*
V13.2	*7	*	*6	*	*	*	*	*	*	*	*
V13.5	*5	*	*	—	*	*	*	*	*	*	*
V13.7	9	*5	*	*8	*	—	*	—	*	*	*
V14	90	28	62	*	26	32	29	17	18	32	23
V14.0	44	16	29	*	14	13	15	*9	10	15	10
V14.1	*5	*	*	*	*	*	*	*	*	*	*
V14.2	13	*	11	—	*	*5	*	*	*	*	*
V14.5	10	*	*8	*	*	*	*	*	*	*	*
V14.8	*9	*	*5	*	*	*	*	*	*	*	*
V15	847	468	379	34	197	281	335	129	208	288	221
V15.0	17	*7	10	*	*5	*	*	*	*	*	*5

See footnotes at end of table.



**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V15.1	30	15	15	18	*	*	*	11	*	*6	*8
V15.3	26	13	12	*	*	9	13	*5	*7	*7	*6
V15.4	24	*8	16	*6	14	*	*	*	*	16	*
V15.41	22	*7	15	*5	13	*	*	*	*	15	*
V15.5	30	18	12	*	10	9	11	*	12	10	*5
V15.8	716	404	312	*6	159	248	303	104	178	243	191
V15.81	212	107	105	*	95	69	43	32	59	80	40
V15.82	482	286	196	*	60	174	247	67	115	150	150
V15.84	*6	*	*	—	*	*	*	*	*	*	*
V15.89	15	*6	9	*	*	*	*8	*	*	*8	*
V16	47	14	33	*	15	23	10	*8	11	16	13
V16.0	17	*9	*8	*	*	*7	*5	*	*	*	*
V16.3	14	*	13	*	*	*9	*	*	*	*5	*
V16.4	*7	*	*6	—	*	*	*	*	*	*	*
V17	148	86	62	*	41	71	32	25	33	56	34
V17.3	108	65	43	*	25	55	29	21	25	41	21
V17.4	28	15	13	*	11	14	*	*	*	11	9
V18	18	*9	9	*	9	*6	*	*	*	*6	*7
V18.0	16	*8	*8	*	*8	*5	*	*	*	*5	*7
V21	*5	*	*	*	*	*	—	*	*	*	*
V21.3	*5	*	*	*	*	*	—	*	*	*	*
V22	15	—	15	—	15	—	—	*	*	*	*
V22.2	11	—	11	—	11	—	—	*	*	*	*
V23	120	—	120	*	118	*	—	12	18	63	27
V23.2	*7	—	*7	—	*7	—	—	*	*	*	*
V23.4	10	—	10	—	10	—	—	*	*	*5	*
V23.7	83	—	83	*	81	—	—	*7	12	43	21
V23.8	12	—	12	*	12	—	—	*	*	*8	*
V23.89	10	—	10	*	10	—	—	*	*	*7	*
V24	*5	—	*5	—	*5	—	—	*	*	*	*
V25	313	*	313	—	313	*	—	33	44	179	58
V25.0	10	—	10	—	10	—	—	—	*	9	*
V25.02	*6	—	*6	—	*6	—	—	—	*	*5	*
V25.2	302	*	302	—	302	*	—	33	43	168	58
V26	*7	*	*7	—	*	*	*	*	*	*	*
V26.5	*6	*	*6	—	*	*	*	*	*	*	*
V26.51	*6	—	*6	—	*	*	*	*	*	*	*
V27	3,951	—	3,951	12	3,934	*	—	582	736	1,660	974
V27.0	3,739	—	3,739	12	3,723	*	—	534	673	1,591	941
V27.1	21	—	21	—	21	*	—	*	*	11	*
V27.2	49	—	49	*	49	*	—	*9	*8	21	11
V27.9	137	—	137	*	137	*	—	36	52	33	16
V28	*8	—	*8	—	*8	—	—	*6	*	—	—
V28.6	*6	—	*6	—	*6	—	—	*6	—	—	—
V29	13	*8	*	13	—	—	—	*	*	*8	*
V29.0	11	*7	*	11	—	—	—	*	*	*6	*
V30	10	*	*6	10	—	—	—	*	*	*	*
V30.1	10	*	*6	10	—	—	—	*	*	*	*
V42	129	73	56	10	32	51	36	33	25	47	24
V42.0	56	32	24	*	21	23	*8	11	12	22	11
V42.1	11	*7	*	*	*	*7	*	*	*	*6	*
V42.2	23	13	9	—	*	*	19	*8	*	*	*6
V42.7	18	11	*7	*	*	10	*	*	*	*8	*
V42.8	15	*7	*8	*	*6	*5	*	*5	*	*6	*
V42.81	*7	*	*	*	*	*	*	*	*	*	*
V43	486	183	303	*	23	109	353	118	149	136	83
V43.3	133	61	72	*	12	33	88	41	38	35	18

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V43.6 . . . . .	346	118	228	*	11	74	261	76	108	99	63
V43.61 . . . . .	*7	*	*6	—	*	*	*	*	*	*	*
V43.64 . . . . .	154	56	98	*	*6	31	117	34	44	41	35
V43.65 . . . . .	181	59	122	*	*	39	137	38	61	55	26
V44 . . . . .	184	95	90	43	33	33	76	52	42	61	30
V44.0 . . . . .	34	20	14	11	*5	*7	10	12	*8	9	*5
V44.1 . . . . .	74	39	35	28	*7	*8	30	22	14	23	15
V44.2 . . . . .	24	11	14	*	*9	*7	*8	*	*8	*8	*
V44.3 . . . . .	39	18	21	*	*8	*7	23	9	10	16	*
V44.6 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V45 . . . . .	1,801	1,022	778	27	158	505	1,110	425	514	592	269
V45.0 . . . . .	279	146	133	*	9	32	237	80	79	92	28
V45.01 . . . . .	244	120	124	*	*7	21	214	70	67	81	26
V45.02 . . . . .	32	25	*7	*	*	10	21	10	10	10	*
V45.1 . . . . .	76	41	35	*	16	29	30	15	17	33	11
V45.2 . . . . .	31	14	16	12	*9	*	*8	*8	*7	*9	*7
V45.3 . . . . .	32	10	22	*	11	16	*6	*	11	10	*7
V45.4 . . . . .	43	18	25	*	11	17	15	*6	15	12	11
V45.6 . . . . .	*9	*	*5	*	*	*	*6	*	*	*	*
V45.61 . . . . .	*7	*	*	—	*	*	*6	*	*	*	*
V45.7 . . . . .	130	38	92	*	24	46	58	22	39	38	31
V45.71 . . . . .	28	*	28	—	*	10	15	*	*7	*7	10
V45.72 . . . . .	19	11	*8	*	*	*7	*8	*	*8	*	*
V45.73 . . . . .	29	12	17	*	*	*9	15	*	*7	12	*5
V45.76 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
V45.77 . . . . .	23	*	22	—	*	10	*8	*	9	*5	*
V45.79 . . . . .	19	*7	12	*	*8	*6	*	*	*	*5	*6
V45.8 . . . . .	1,201	752	449	11	76	361	752	288	344	396	172
V45.81 . . . . .	634	430	204	*	11	173	450	166	169	214	84
V45.82 . . . . .	382	242	140	—	18	138	225	85	108	126	63
V45.89 . . . . .	184	80	104	11	47	50	77	37	67	56	25
V46 . . . . .	24	14	10	*	*7	*	9	*6	*6	*8	*
V46.1 . . . . .	20	13	*7	*	*6	*	*6	*5	*	*8	*
V49 . . . . .	79	48	31	*	*9	28	41	13	20	29	16
V49.6 . . . . .	*6	*6	*	*	*	*	*	*	*	*	*
V49.7 . . . . .	65	42	23	*	*7	25	33	11	17	24	14
V49.72 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
V49.75 . . . . .	34	22	12	—	*	11	20	*	10	10	*9
V49.76 . . . . .	17	10	*7	*	*	*8	*8	*	*	*8	*
V49.8 . . . . .	*7	*	*7	—	*	*	*	*	*	*	*
V49.81 . . . . .	*6	—	*6	—	—	*	*	*	*	*	*
V50 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V53 . . . . .	17	10	*7	*	*	*	12	*	*	*5	*5
V53.3 . . . . .	14	9	*5	*	*	*	11	*	*	*	*
V53.31 . . . . .	11	*6	*	*	*	*	*9	*	*	*	*
V54 . . . . .	74	23	51	*	*8	13	51	10	18	26	20
V54.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V54.8 . . . . .	58	18	40	*	*6	*8	43	*6	17	20	15
V54.9 . . . . .	10	*	*7	*	*	*	*6	*	*	*	*
V55 . . . . .	51	25	26	*6	9	19	18	10	16	17	*8
V55.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V55.2 . . . . .	9	*5	*	*	*	*	*	*	*	*	*
V55.3 . . . . .	30	14	16	*	*	13	11	*5	11	10	*
V56 . . . . .	101	52	49	*	21	40	37	12	*6	53	29
V56.0 . . . . .	97	50	47	*	20	39	36	12	*6	52	28
V57 . . . . .	626	249	377	9	48	119	450	104	178	202	142
V57.1 . . . . .	161	63	98	*	10	25	124	30	26	60	45
V57.2 . . . . .	46	20	27	*	*9	*7	30	9	10	16	11

See footnotes at end of table.

**Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V57.21	46	20	27	*	*9	*7	30	9	10	16	11
V57.8	408	163	245	*5	29	87	287	58	140	124	85
V57.89	407	163	244	*5	29	87	287	58	140	124	85
V57.9	*8	*	*6	—	—	*	*7	*8	*	*	—
V58	508	253	255	34	77	137	259	104	134	161	108
V58.0	33	17	16	*	*5	11	15	*5	*	18	*6
V58.1	161	93	68	26	43	55	38	38	25	57	40
V58.4	30	14	17	*	*5	*6	15	*9	*	*6	11
V58.49	29	13	16	*	*	*5	15	*9	*	*6	11
V58.6	265	122	143	*	23	62	179	49	93	75	48
V58.61	222	104	119	*	15	48	160	47	70	61	44
V58.69	42	17	24	*	*8	14	19	*	22	13	*
V58.8	16	*6	10	*	*	*	10	*	*9	*	*
V58.89	13	*	*8	*	*	*	9	*	*8	*	*
V59	*7	*	*	*	*	*	—	*	*	*	*
V59.4	*5	*	*	*	*	*	—	*	*	*	—
V60	53	31	23	*	29	12	12	10	13	15	15
V60.0	26	19	*7	*	18	*7	*	*	*6	*7	10
V60.2	10	*	*7	*	*6	*	*	*	*	*	*
V60.3	*6	*	*	—	*	*	*5	*	*	*	*
V61	59	23	36	10	38	9	*	*5	24	18	11
V61.0	13	*5	*8	*	*8	*	*	*	*5	*	*
V61.1	14	*8	*6	—	13	*	*	*	*9	*	*
V61.10	14	*8	*6	—	12	*	*	*	*9	*	*
V61.2	10	*	*6	*5	*	*	—	*	*	*	*
V61.20	*8	*	*	*	*	—	—	*	*	*	*
V61.5	*6	—	*6	—	*6	*	—	*	*	*	*
V61.8	*7	*	*	*	*	*	*	*	*	*	*
V62	68	36	33	*	47	13	*	*6	27	23	13
V62.0	10	*7	*	—	*7	*	—	*	*	*	*
V62.5	*7	*6	*	*	*6	*	—	*	*	*	*
V62.8	40	19	22	*	28	*9	*	*5	20	12	*
V62.81	23	11	12	*	17	*5	*	*	15	*6	*
V62.89	14	*7	*7	*	10	*	*	*	*	*	*
V63	*5	*	*	*	*	*	*	*	*	*	*
V64	114	50	64	*	46	37	28	24	27	39	24
V64.1	22	11	11	*	*5	*7	*9	*5	*5	*7	*5
V64.2	16	*8	*8	—	*7	*6	*	*	*	*8	*
V64.3	17	*7	9	*	*7	*5	*	*	*	*8	*
V64.4	58	23	35	*	27	18	13	11	16	17	14
V65	33	13	19	*7	18	*6	*	*	*	22	*
V65.2	*8	*	*	*	*6	*	*	*	*	*	*
V65.3	20	*8	12	*7	*8	*	*	*	*	15	*
V66	36	20	16	*	*	*9	26	*7	11	12	*6
V66.7	32	17	14	*	*	*7	23	*6	10	12	*
V70	18	10	*8	*	10	*	*	*6	*	*6	*
V70.7	17	10	*7	*	9	*	*	*6	*	*6	*
V71	11	*5	*5	*	*	*	*	*	*	*	*
V72	10	*6	*	*	*	*	*	*6	*	*	*
V72.8	*7	*	*	—	*	*	*	*5	*	*	*

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
All codes . . . . .	42,533	16,834	25,700	2,084	14,419	10,743	15,286	9,389	9,082	15,394	8,668
01-05 . . . . .	1,101	514	587	215	337	256	293	240	277	345	239
01 . . . . .	131	64	67	17	34	38	42	25	29	39	39
01.0 . . . . .	10	*	*6	*	*	*	*	*	*	*	*
01.02 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
01.1 . . . . .	19	10	*9	*	*7	*6	*	*	*5	*	*
01.13 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
01.18 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
01.2 . . . . .	21	12	10	*	*6	*	*6	*	*	*7	*8
01.24 . . . . .	13	*7	*6	*	*	*	*	*	*	*	*5
01.3 . . . . .	32	17	15	*	*	*9	18	*6	*8	*7	11
01.31 . . . . .	21	14	*7	*	*	*	15	*	*6	*	*7
01.39 . . . . .	11	*	*8	*	*	*	*	*	*	*	*
01.5 . . . . .	48	20	27	*5	12	17	13	*9	*8	18	13
01.51 . . . . .	10	*	*7	*	*	*	*	*	*	*	*
01.59 . . . . .	36	17	19	*	10	12	*9	*7	*6	12	11
02 . . . . .	117	64	53	31	35	26	25	25	25	35	32
02.0 . . . . .	23	12	11	*8	*6	*6	*	*	*	*8	*6
02.02 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
02.06 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
02.1 . . . . .	13	*5	*8	*	*	*	*	*	*	*	*
02.12 . . . . .	12	*	*8	*	*	*	*	*	*	*	*
02.2 . . . . .	24	13	11	*	*8	*8	*	*	*5	*8	*7
02.3 . . . . .	16	*8	*9	*5	*	*	*	*	*	*5	*5
02.34 . . . . .	13	*6	*7	*	*	*	*	*	*	*	*
02.4 . . . . .	23	13	10	11	*7	*	*	*6	*7	*	*6
02.42 . . . . .	19	11	*8	10	*5	*	*	*	*5	*	*5
02.9 . . . . .	17	13	*	*	*5	*	*7	*5	*	*	*
02.94 . . . . .	*9	*7	*	*	*	*	*	*	*	*	*
03 . . . . .	779	354	424	163	245	166	205	170	210	248	151
03.0 . . . . .	181	95	87	*	24	66	90	43	37	66	35
03.02 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
03.09 . . . . .	176	92	83	*	21	64	88	42	36	62	35
03.3 . . . . .	318	161	157	142	94	43	40	67	72	106	73
03.31 . . . . .	316	161	156	142	93	42	39	67	72	105	73
03.4 . . . . .	15	*9	*6	*	*	*	*	*	*	*6	*
03.5 . . . . .	32	16	16	*	*9	*8	10	*7	*6	*9	10
03.53 . . . . .	15	*7	*7	*	*6	*	*6	*	*	*	*5
03.59 . . . . .	16	*8	*8	*	*	*6	*	*	*	*	*
03.6 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
03.9 . . . . .	217	66	152	*6	110	43	59	47	89	56	26
03.90 . . . . .	76	26	50	*	31	21	20	18	25	20	13
03.91 . . . . .	75	15	60	*	49	10	16	12	46	13	*
03.92 . . . . .	39	17	22	*	*7	*9	21	13	13	10	*
03.95 . . . . .	18	*	15	*	16	*	*	*	*	*7	*
04 . . . . .	69	29	40	*	21	25	19	18	13	22	16
04.0 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
04.07 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
04.3 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
04.4 . . . . .	16	*6	10	*	*6	*	*	*	*	*6	*
04.43 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
04.8 . . . . .	29	10	19	*	*	12	12	11	*7	*6	*5
04.81 . . . . .	26	10	17	*	*	11	11	11	*	*6	*
06-07 . . . . .	102	30	72	*	30	40	31	22	13	39	29
06 . . . . .	86	22	65	*	25	34	27	18	10	33	25
06.1 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
06.2 . . . . .	24	*	20	*	*6	10	*7	*	*	*8	*8
06.3 . . . . .	12	*	*8	—	*6	*	*	*	*	*	*
06.39 . . . . .	10	*	*7	—	*	*	*	*	*	*	*
06.4 . . . . .	19	*	15	*	*7	*8	*	*	*	*8	*5

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
06.8 . . . . .	17	*	12	—	*	*6	*7	*	*	*8	*
06.89 . . . . .	14	*	10	—	*	*	*7	*	*	*7	*
07 . . . . .	16	*9	*7	*	*5	*6	*	*	*	*5	*
07.6 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
08-16 . . . . .	86	47	39	12	27	21	25	15	17	33	21
08 . . . . .	35	20	15	*	13	*8	10	*7	*7	10	11
08.8 . . . . .	23	14	*9	*	10	*6	*6	*	*5	*7	*6
08.81 . . . . .	19	12	*7	*	*8	*	*6	*	*5	*	*6
11 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
12 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
13 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
14 . . . . .	21	*8	13	*	*	*6	*7	*	*	9	*
14.7 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
16 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
18-20 . . . . .	46	24	22	19	11	*7	*9	15	*6	17	*8
18 . . . . .	16	10	*6	*	*9	*	*	*6	*	*6	*
18.4 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
20 . . . . .	26	13	12	16	*	*	*5	*8	*	10	*
20.0 . . . . .	17	10	*8	13	*	*	*	*5	*	*7	*
20.01 . . . . .	15	*9	*6	13	*	*	*	*5	*	*6	*
21-29 . . . . .	269	159	110	66	93	59	52	72	45	91	60
21 . . . . .	61	37	24	*	21	14	21	17	10	19	15
21.0 . . . . .	22	12	10	*	*	*5	14	*7	*	*5	*5
21.01 . . . . .	11	*6	*	*	*	*	*8	*	*	*	*
21.03 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
21.6 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
21.69 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
21.7 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
21.8 . . . . .	18	12	*7	*	*9	*	*	*6	*	*6	*
21.81 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
21.88 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
22 . . . . .	41	22	20	*7	17	11	*8	9	*	16	12
22.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
22.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
22.4 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
22.6 . . . . .	20	11	*9	*	*8	*	*	*	*	*8	*7
22.63 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*
23 . . . . .	21	11	10	*	9	*	*	*5	*	10	*
23.0 . . . . .	10	*5	*	*	*	*	*	*	*	*	*
23.09 . . . . .	10	*5	*	*	*	*	*	*	*	*	*
23.1 . . . . .	10	*5	*	*	*5	*	*	*	*	*	*
23.19 . . . . .	9	*5	*	*	*	*	*	*	*	*	*
24 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
25 . . . . .	14	9	*	*	*	*	*	*	*	*	*
26 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*
26.3 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
27 . . . . .	49	32	17	20	16	*9	*5	14	*8	17	9
27.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
27.5 . . . . .	16	11	*5	*6	*6	*	*	*	*	*5	*
27.51 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
27.6 . . . . .	19	13	*6	12	*	*	*	*6	*	*7	*
27.62 . . . . .	9	*5	*	*9	*	*	—	*	*	*5	*
27.69 . . . . .	*8	*7	*	*	*	*	*	*	*	*	*
28 . . . . .	47	28	19	24	15	*5	*	11	*9	16	11
28.0 . . . . .	10	*	*5	*	*6	*	*	*	*	*	*
28.2 . . . . .	10	*7	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
28.3 . . . . .	19	11	*8	16	*	*	—	*	*	*6	*6
29 . . . . .	18	11	*7	*	*5	*7	*	*8	*	*	*
29.4 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
30-34 . . . . .	1,022	583	439	57	179	304	482	248	208	356	209
30 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
31 . . . . .	143	85	58	18	25	45	55	42	28	52	21
31.1 . . . . .	66	41	25	*	12	23	28	18	13	28	*8
31.2 . . . . .	18	11	*8	*	*	*	*9	*	*	*8	*
31.29 . . . . .	18	11	*8	*	*	*	*9	*	*	*8	*
31.4 . . . . .	46	28	18	10	*8	12	16	17	10	11	*7
31.42 . . . . .	39	24	16	10	*8	9	13	16	*8	*9	*7
31.7 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
32 . . . . .	83	41	42	*	13	34	33	20	18	24	20
32.2 . . . . .	40	20	20	*	10	15	14	10	*8	12	10
32.29 . . . . .	37	19	18	*	*9	14	14	9	*7	11	*9
32.3 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
32.4 . . . . .	31	14	17	*	*	14	14	*8	*7	9	*7
33 . . . . .	314	182	133	17	50	92	156	77	60	119	59
33.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
33.2 . . . . .	298	173	125	16	45	87	151	74	55	112	57
33.21 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
33.22 . . . . .	52	32	21	*	*8	17	23	16	10	16	11
33.23 . . . . .	37	22	15	*7	*7	*8	16	9	*7	15	*6
33.24 . . . . .	111	64	46	*	19	29	60	27	18	45	20
33.26 . . . . .	27	15	12	*	*	*8	17	*6	*5	10	*6
33.27 . . . . .	42	25	18	*	*	11	26	11	*7	16	*9
33.28 . . . . .	20	12	*9	*	*	9	*7	*	*6	*8	*
34 . . . . .	474	271	203	19	89	131	235	108	99	158	109
34.0 . . . . .	174	106	69	12	45	49	69	38	36	63	38
34.03 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
34.04 . . . . .	129	78	51	*8	33	36	52	31	26	49	23
34.09 . . . . .	33	20	13	*	10	10	12	*	*8	10	12
34.2 . . . . .	49	29	21	*	10	15	24	14	9	12	15
34.21 . . . . .	16	11	*5	*	*	*	*7	*	*	*	*
34.22 . . . . .	11	*5	*6	*	*	*	*6	*	*	*	*
34.24 . . . . .	13	9	*	—	*	*	*7	*	*	*	*
34.5 . . . . .	22	17	*	*	*5	*8	*8	*	*	*7	*7
34.51 . . . . .	20	15	*	*	*	*7	*6	*	*	*5	*7
34.6 . . . . .	*8	*6	*	—	*5	*	*	*	*	*	*
34.7 . . . . .	*7	*6	*	*	*	*	*	*	*	*	*
34.8 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
34.9 . . . . .	201	100	100	*	19	52	127	46	46	67	41
34.91 . . . . .	177	88	89	*	16	46	113	40	41	58	37
34.92 . . . . .	23	11	11	*	*	*6	14	*5	*	*9	*
35-39 . . . . .	6,813	3,967	2,845	210	681	2,384	3,538	1,370	1,463	2,601	1,378
35 . . . . .	136	71	65	29	14	29	64	45	23	40	28
35.1 . . . . .	15	*7	*8	*	*	*	*7	*5	*	*	*
35.12 . . . . .	10	*5	*5	*	*	*	*6	*	*	*	*
35.2 . . . . .	78	42	36	*	9	19	49	20	15	25	18
35.21 . . . . .	24	13	11	*	*	*	21	*9	*	*6	*6
35.22 . . . . .	32	19	13	*	*	11	17	*6	*7	12	*7
35.23 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
35.24 . . . . .	14	*6	*8	*	*	*6	*6	*	*	*	*
35.3 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
35.33 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
35.6 . . . . .	*9	*	*	*6	*	*	*	*	*	*	*
35.62 . . . . .	*5	*	*	*	*	—	*	*	*	*	*
35.7 . . . . .	10	*	*5	*7	*	*	*	*	*	*	*
35.71 . . . . .	*8	*	*	*5	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
35.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
36 . . . . .	1,724	1,179	545	*	95	737	889	316	441	622	345
36.0 . . . . .	1,204	802	402	*	76	517	608	211	323	416	254
36.01 . . . . .	544	358	186	*	37	238	268	98	154	186	105
36.02 . . . . .	16	10	*6	—	*	*6	*9	*	*	*6	*
36.05 . . . . .	97	66	31	—	*	39	54	12	28	31	25
36.06 . . . . .	537	363	174	*	34	228	273	95	138	186	118
36.09 . . . . .	*6	*	*	—	*	*	*	*	*	*	—
36.1 . . . . .	515	373	142	—	19	217	279	104	117	204	90
36.11 . . . . .	53	33	20	—	*	21	28	13	13	19	*8
36.12 . . . . .	100	66	35	—	*	43	55	22	23	37	18
36.13 . . . . .	94	72	22	—	*	36	56	18	21	38	17
36.14 . . . . .	40	35	*5	—	*	19	20	*	*8	20	*8
36.15 . . . . .	207	150	57	—	*5	90	112	44	48	79	35
36.16 . . . . .	12	10	*	—	*	*	*	*	*	*6	*
37 . . . . .	2,064	1,218	845	16	158	709	1,181	430	461	803	370
37.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
37.1 . . . . .	14	*8	*6	*	*	*	*6	*	*	*5	*
37.12 . . . . .	13	*8	*5	*	*	*	*5	*	*	*	*
37.2 . . . . .	1,463	884	579	10	123	597	732	281	342	585	255
37.21 . . . . .	35	17	18	*	*	14	15	*5	*6	18	*5
37.22 . . . . .	1,122	687	435	*	90	488	542	194	268	461	199
37.23 . . . . .	171	95	76	*5	13	52	101	44	39	63	26
37.26 . . . . .	97	64	34	*	9	29	58	30	23	30	14
37.27 . . . . .	22	12	10	*	*	*7	10	*	*	*8	*5
37.29 . . . . .	9	*5	*	—	*	*	*	*	*	*	*
37.3 . . . . .	39	20	19	*	*8	12	17	10	*9	12	*8
37.34 . . . . .	31	17	14	*	*6	10	15	*9	*7	10	*6
37.6 . . . . .	44	31	14	—	*	21	21	10	10	14	10
37.61 . . . . .	42	29	13	—	*	20	19	10	9	13	10
37.7 . . . . .	221	112	109	*	*8	25	186	56	41	80	43
37.71 . . . . .	29	15	14	*	*	*	26	*8	*8	*9	*
37.72 . . . . .	136	66	70	*	*	15	117	34	24	55	24
37.75 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
37.76 . . . . .	*8	*	*	*	*	*	*6	*	*	*	*
37.78 . . . . .	30	17	13	*	*	*	22	*7	*5	*7	10
37.8 . . . . .	199	101	99	*	*	21	172	50	36	78	36
37.80 . . . . .	*7	*	*	—	*	*	*6	*	*	*	*
37.81 . . . . .	14	*8	*6	—	*	*	12	*	*	*6	*
37.82 . . . . .	14	*6	*8	*	*	*	12	*5	*	*	*
37.83 . . . . .	141	70	71	*	*	15	122	36	22	57	25
37.87 . . . . .	13	*8	*	*	*	*	11	*	*	*	*
37.9 . . . . .	73	57	17	*	*	25	44	17	18	26	12
37.94 . . . . .	56	45	11	*	*	21	31	13	14	20	*9
37.99 . . . . .	*7	*	*	*	*	*	*5	*	*	*	*
38 . . . . .	1,604	780	824	125	256	455	769	334	289	630	351
38.0 . . . . .	33	17	16	*	*	*8	20	*6	*7	12	*8
38.03 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
38.08 . . . . .	18	10	*8	—	*	*	13	*	*	*7	*
38.1 . . . . .	158	93	65	—	*	40	116	38	34	64	23
38.12 . . . . .	134	79	56	—	*	33	101	33	29	55	17
38.16 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
38.18 . . . . .	15	*8	*6	—	*	*	*9	*	*	*6	*
38.2 . . . . .	10	*	*6	—	*	*	9	*	*	*	*
38.21 . . . . .	9	*	*6	—	*	*	*8	*	*	*	*
38.4 . . . . .	44	32	12	*	*	9	31	11	*9	14	10
38.44 . . . . .	25	20	*6	*	*	*	21	*8	*5	*7	*5
38.45 . . . . .	11	*7	*	*	*	*	*	*	*	*	*
38.6 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
38.7 . . . . .	59	25	34	—	*7	16	36	18	10	25	*5
38.8 . . . . .	26	16	10	*9	*8	*	*	*7	*	10	*7

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
38.85	10	*	*5	*8	*	*	—	*	*	*	*
38.86	*5	*	*	—	*	*	*	*	*	*	*
38.9	1,259	584	675	112	230	373	545	248	222	498	290
38.91	93	52	42	19	12	23	39	16	20	37	20
38.92	15	*8	*7	14	—	*	*	*	*	9	*
38.93	948	427	522	73	188	280	408	195	173	361	220
38.94	9	*5	*	*	*	*	*	*	*	*	*
38.95	162	80	82	*	25	63	72	24	26	72	41
38.98	26	9	16	*	*	*	20	*7	*	17	*
38.99	*5	*	*	*	*	*	*	*	*	*	*
39	1,286	719	567	38	159	453	635	245	249	507	285
39.0	*5	*	*	*	*	*	*	*	*	*	*
39.2	138	75	64	*	*9	44	83	28	24	56	30
39.25	21	12	9	—	*	*7	14	*	*6	*8	*
39.27	37	18	18	*	*6	12	19	*6	*	19	*9
39.29	73	40	33	*	*	22	48	17	15	27	14
39.3	23	14	*9	*	*8	*	9	*	*	*6	*9
39.31	16	11	*5	*	*6	*	*6	*	*	*5	*5
39.32	*6	*	*	*	*	*	*	*	*	*	*
39.4	78	42	36	*	11	31	34	13	13	32	19
39.42	22	13	9	—	*5	12	*5	*	*	10	*5
39.49	50	26	23	*	*5	17	26	9	9	19	12
39.5	135	73	62	*	16	41	73	29	33	49	24
39.50	97	52	45	*	*9	30	56	20	25	37	16
39.51	10	*	*7	—	*	*	*	*	*	*	*
39.52	*5	*	*	*	*	*	*	*	*	*	*
39.59	9	*8	*	*	*	*	*	*	*	*	*
39.6	270	178	92	18	20	97	135	58	61	91	60
39.61	249	165	84	16	17	90	125	56	59	85	49
39.62	*6	*	*	—	*	*	*	*	*	*	*5
39.63	*6	*	*	*	*	*	*	*	*	*	*
39.64	*6	*	*	*	*	*	*	*	*	*	*
39.7	18	14	*	*	*	*	13	*	*	*7	*
39.71	12	10	*	—	*	*	11	*	*	*5	*
39.79	*7	*	*	*	*	*	*	*	*	*	*
39.9	614	318	296	9	92	228	285	107	104	264	139
39.90	44	22	21	*	*	13	28	9	10	17	*7
39.93	*6	*	*	*	*	*	*	*	*	*	*
39.95	552	288	264	*7	85	210	251	94	90	242	127
39.98	*7	*	*	*	*	*	*	*	*	*	*
40-41	354	178	175	22	55	130	147	81	66	130	77
40	203	103	100	*5	22	89	87	46	41	71	45
40.1	57	24	33	*	*8	23	23	14	9	24	11
40.11	55	24	31	*	*8	22	22	13	*9	23	10
40.2	53	22	30	*	*5	19	27	*9	12	19	13
40.23	14	*	13	—	*	*6	*7	*	*	*	*
40.24	*7	*	*	*	*	*	*	*	*	*	*
40.29	29	15	13	*	*	10	14	*5	*8	10	*6
40.3	64	37	28	*	*6	31	28	16	12	19	17
40.4	12	*7	*	*	*	*6	*	*	*	*	*
40.41	9	*6	*	*	*	*5	*	*	*	*	*
40.5	15	11	*	*	*	9	*	*	*	*6	*
40.59	*7	*5	*	—	*	*	*	*	*	*	*
41	150	75	75	16	33	41	60	35	25	58	32
41.0	10	*6	*	*	*	*	*	*	*	*	*
41.04	*6	*	*	*	*	*	*	*	*	*	*
41.3	116	59	57	14	20	29	53	27	19	45	24
41.31	116	59	57	14	20	29	53	27	19	45	24
41.5	21	*9	12	*	*7	*7	*6	*	*	*8	*

See footnotes at end of table.



**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
42-54 . . . . .	5,597	2,350	3,247	222	1,307	1,607	2,461	1,185	1,226	2,111	1,075
42 . . . . .	156	87	69	11	20	41	84	46	31	47	32
42.2 . . . . .	72	39	33	*7	9	20	36	29	12	19	11
42.23 . . . . .	67	37	30	*6	*9	19	33	27	12	18	10
42.3 . . . . .	31	21	11	*	*	11	15	*5	*7	11	*8
42.33 . . . . .	31	21	10	*	*	11	15	*5	*7	10	*8
42.9 . . . . .	38	18	20	*	*	*6	28	*8	9	15	*6
42.92 . . . . .	37	18	19	*	*	*5	28	*8	9	15	*6
43 . . . . .	235	116	119	27	21	46	141	52	46	92	44
43.1 . . . . .	174	78	97	11	14	28	120	41	37	71	26
43.11 . . . . .	142	63	79	*6	*8	20	108	33	30	59	20
43.19 . . . . .	32	14	18	*5	*6	*8	12	*8	*6	13	*5
43.3 . . . . .	16	14	*	15	*	*	*	*	*	*	*7
43.4 . . . . .	18	*9	*9	*	*	*	11	*	*	*6	*
43.41 . . . . .	12	*5	*7	-	*	*	9	*	*	*	*
43.7 . . . . .	15	*8	*7	-	*	*8	*	*	*	*7	*
44 . . . . .	279	106	174	*8	90	93	89	57	52	114	56
44.0 . . . . .	*7	*	*	-	*	*	*	*	*	*	*
44.1 . . . . .	28	*9	19	*	*7	*5	15	*	*6	16	*
44.13 . . . . .	15	*	11	*	*5	*	*8	*	*	*7	*
44.14 . . . . .	11	*	*8	*	*	*	*8	*	*	*9	*
44.2 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
44.29 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
44.3 . . . . .	92	20	72	*	49	39	*	18	19	40	15
44.31 . . . . .	65	12	53	-	39	26	*	*8	15	29	13
44.39 . . . . .	26	*8	18	*	10	12	*	10	*	10	*
44.4 . . . . .	81	43	38	-	*6	21	54	17	14	28	23
44.42 . . . . .	*8	*	*	-	*	*	*	*	*	*	*
44.43 . . . . .	66	35	32	-	*	16	46	12	12	23	19
44.6 . . . . .	55	23	31	*7	20	20	*8	14	10	19	12
44.66 . . . . .	41	18	24	*5	14	15	*7	*9	*9	15	9
44.69 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
45 . . . . .	2,179	926	1,252	34	310	577	1,258	478	491	839	371
45.0 . . . . .	10	*	*6	*	*	*	*6	*	*	*	*
45.02 . . . . .	*6	*	*	-	*	*	*	*	*	*	*
45.1 . . . . .	1,035	451	584	16	161	280	578	227	235	392	180
45.13 . . . . .	444	205	238	*	65	119	256	98	109	156	81
45.14 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
45.16 . . . . .	580	240	339	12	94	156	318	126	125	231	97
45.2 . . . . .	581	237	344	*7	77	138	359	132	130	227	92
45.23 . . . . .	314	131	183	*	30	75	208	67	69	131	47
45.24 . . . . .	45	16	29	*	*7	12	27	*6	11	23	*6
45.25 . . . . .	214	85	129	*6	38	50	120	56	49	72	37
45.3 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
45.33 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
45.4 . . . . .	133	59	74	-	*7	35	91	30	28	51	25
45.41 . . . . .	*6	*	*	-	*	*	*	*	*	*	*
45.42 . . . . .	105	48	57	-	*	28	72	24	24	37	20
45.43 . . . . .	21	*7	14	-	*	*6	14	*	*	10	*
45.6 . . . . .	67	26	41	*	13	20	31	13	14	30	10
45.62 . . . . .	63	25	39	*	12	19	30	13	12	29	9
45.7 . . . . .	263	113	150	*	35	79	145	58	64	96	45
45.72 . . . . .	13	*5	*8	*	*	*	*	*	*	*	*
45.73 . . . . .	90	39	52	*	10	23	57	19	22	30	19
45.74 . . . . .	11	*	*6	*	*	*	*6	*	*	*	*
45.75 . . . . .	27	12	16	*	*	*8	17	*7	*7	10	*
45.76 . . . . .	95	43	52	*	15	37	43	23	25	33	14
45.79 . . . . .	27	10	16	*	*	*6	16	*	*6	13	*
45.8 . . . . .	11	*6	*	*	*	*	*	*	*	*6	*
45.9 . . . . .	69	26	43	*	10	16	41	13	14	29	13

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
45.91	15	*	10	*	*	*	*7	*	*	*6	*
45.93	26	10	15	*	*	*	16	*	*6	11	*
45.94	23	*9	14	*	*	*	16	*6	*	10	*
46	217	105	113	9	43	69	96	43	58	80	36
46.0	16	*7	9	*	*	*	9	*	*5	*	*
46.01	*6	*	*	*	*	*	*	*	*	*	*
46.03	9	*	*6	*	*	*	*6	*	*	*	*
46.1	50	25	25	*	*8	14	27	11	13	18	*7
46.10	28	12	16	*	*	*9	16	*	*7	11	*
46.11	15	10	*5	*	*	*	*6	*	*	*5	*
46.13	*6	*	*	*	*	*	*	*	*	*	*
46.2	15	*7	*8	*	*	*	*7	*	*	*6	*
46.20	*8	*	*	*	*	*	*	*	*	*	*
46.3	25	15	10	*	*	*8	12	*5	*6	*8	*6
46.32	*7	*	*	*	*	*	*	*	*	*	*
46.39	18	11	*7	*	*	*5	*9	*	*	*5	*
46.4	11	*	*9	*	*	*6	*	*	*	*	*
46.41	*6	*	*	*	*	*	*	*	*	*	*
46.5	40	20	20	*	*8	18	11	*8	13	13	*7
46.51	11	*6	*5	*	*	*	*	*	*	*	*
46.52	28	13	15	*	*	13	9	*5	10	9	*
46.7	37	16	21	*	13	*9	14	*7	*8	16	*6
46.73	16	*6	9	*	*6	*	*7	*	*	*9	*
46.75	10	*5	*	*	*	*	*	*	*	*	*
46.79	*6	*	*	*	*	*	*	*	*	*	*
46.8	17	9	*8	*	*	*	*8	*	*6	*6	*
46.85	10	*5	*	*	*	*	*7	*	*	*	*
47	389	193	197	74	202	76	37	71	92	121	106
47.0	329	173	156	70	176	57	25	62	73	99	96
47.01	117	55	63	25	68	18	*6	17	27	31	42
47.09	212	118	93	45	108	39	20	45	46	67	54
47.1	58	19	40	*	24	19	12	*9	19	22	9
47.19	54	18	36	*	21	18	11	*8	18	20	*8
48	139	73	66	*8	26	41	65	34	36	47	23
48.2	53	26	27	*	11	14	23	14	13	16	9
48.23	21	*8	13	*	*	*7	10	*5	*5	*8	*
48.24	30	17	13	*	*7	*7	12	*8	*8	*7	*7
48.3	21	10	10	*	*	*6	13	*	*7	*7	*
48.36	15	*7	*8	*	*	*	*9	*	*	*5	*
48.5	10	*7	*	—	*	*	*6	*	*	*	*
48.6	31	16	15	*	*	12	17	*8	*6	13	*
48.63	19	10	*9	—	*	*8	10	*6	*	*6	*
48.69	9	*	*5	*	*	*	*	*	*	*5	*
48.8	12	*8	*	*	*6	*	*	*	*	*	*
48.81	11	*7	*	*	*6	*	*	*	*	*	*
49	49	25	24	*	15	19	12	9	10	16	14
49.0	10	*6	*	*	*	*	*	*	*	*	*
49.01	*7	*	*	*	*	*	*	*	*	*	*
49.4	18	*8	10	*	*	9	*	*	*	*7	*
49.46	12	*6	*6	—	*	*6	*	*	*	*	*
49.7	*7	*	*	*	*	*	*	*	*	*	*
50	117	63	53	*7	24	43	43	27	22	44	23
50.1	87	45	43	*5	18	29	34	21	18	30	18
50.11	64	33	31	*	15	21	24	14	9	25	15
50.12	23	12	12	*	*	*8	10	*7	9	*	*
50.2	11	*7	*	*	*	*7	*	*	*	*5	*
50.22	*5	*	*	*	*	*	*	*	*	*	*
50.29	*5	*	*	—	*	*	*	*	*	*	*
50.9	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
51	717	260	458	*	207	215	292	143	141	269	164
51.0	*5	*	*	*	*	*	*	*	*	*	*
51.1	74	33	41	*	14	23	36	14	14	28	19
51.10	60	26	34	*	13	19	28	9	10	25	16
51.14	*7	*	*	—	*	*	*5	*	*	*	*
51.2	436	143	293	*	142	140	151	85	91	170	89
51.22	112	49	63	*	21	37	53	26	27	37	22
51.23	323	93	230	*	121	103	97	60	64	133	66
51.3	*7	*	*	*	*	*	*	*	*	*	*
51.4	*8	*	*6	—	*	*	*7	*	*	*	*
51.41	*5	*	*	—	*	*	*	*	*	*	*
51.5	*6	*	*	—	*	*	*	*	*	*	*
51.51	*5	*	*	—	*	*	*	*	*	*	*
51.8	162	61	101	*	44	42	76	31	26	60	45
51.85	75	28	47	—	19	21	35	16	13	27	20
51.87	37	16	21	*	9	10	18	*8	*5	12	12
51.88	44	15	29	*	13	10	21	*7	*7	18	12
51.9	13	*8	*	*	*	*	*9	*	*	*	*
51.98	11	*7	*	*	*	*	*8	*	*	*	*
52	35	17	18	*	10	12	12	*6	*8	14	*8
52.1	*7	*	*	—	*	*	*	*	*	*	*
52.5	*5	*	*	*	*	*	*	*	*	*	*
52.7	*6	*	*	—	*	*	*	*	*	*	*
52.9	10	*	*6	*	*	*	*	*	*	*	*
52.93	*6	*	*	—	*	*	*	*	*	*	*
53	251	117	134	10	47	92	100	57	60	88	47
53.0	43	35	*7	*	*6	*8	25	11	10	15	*7
53.00	*6	*	*	*	*	*	*	*	*	*	*
53.01	*5	*	*	*	*	*	*	*	*	*	*
53.02	*6	*	*	*	*	*	*	*	*	*	*
53.03	*9	*8	*	—	*	*	*6	*	*	*	*
53.04	*9	*8	*	—	*	*	*6	*	*	*	*
53.05	*8	*7	*	—	*	*	*	*	*	*	*
53.1	*7	*6	*	*	*	*	*	*	*	*	*
53.4	58	24	34	*	14	23	21	11	16	20	11
53.41	10	*	*5	*	*	*5	*	*	*	*	*
53.49	49	19	29	*	13	17	18	10	13	17	*8
53.5	43	13	30	*	11	19	12	*7	11	16	*8
53.51	25	*8	18	*	*8	*9	*8	*5	*6	*8	*5
53.59	17	*6	12	*	*	10	*	*	*	*8	*
53.6	80	32	49	*	12	36	32	21	18	27	14
53.61	66	27	39	—	10	29	27	18	14	22	11
53.69	15	*	10	*	*	*8	*	*	*	*	*
53.7	*8	*	*5	*	*	*	*	*	*	*	*
54	833	263	571	28	292	282	232	161	178	342	152
54.0	18	*7	10	*	*	*8	*5	*	*	*8	*
54.1	50	20	31	*	18	15	14	11	11	18	10
54.11	25	12	13	*	9	*7	*7	*7	*6	*8	*
54.12	12	*	*8	*	*	*	*	*	*	*	*
54.19	13	*	10	*	*	*	*	*	*	*5	*
54.2	130	37	93	*	54	42	32	24	26	57	23
54.21	72	18	54	*	38	22	11	13	13	33	13
54.23	23	*6	17	*	*5	9	*8	*	*6	*8	*5
54.24	14	*6	*9	*	*	*	*7	*	*	*7	*
54.25	19	*7	12	*	*	*7	*6	*	*	*9	*
54.3	14	*	10	*	*	*	*5	*	*	*	*
54.4	66	17	49	*	23	20	20	12	14	24	15
54.5	342	62	280	*	143	103	93	66	71	148	57
54.51	52	*9	43	*	22	15	14	12	10	20	10
54.59	290	53	237	*	121	87	79	54	61	129	47
54.6	15	*8	*7	*	*8	*	*	*	*	*8	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
54.61	*7	*	*	*	*	*	*	*	*	*	*
54.7	*6	*	*	*	*	*	*	*	*	*	*
54.9	192	104	88	14	37	86	56	35	43	73	41
54.91	140	78	62	*	25	68	44	27	35	55	24
54.93	*7	*	*	*	*	*	*	*	*	*	*
54.95	13	*6	*7	*5	*	*	*	*	*	*	*
54.98	24	12	11	*	*	10	*6	*	*5	9	*5
55-59	955	454	501	38	232	299	386	218	206	364	166
55	133	77	56	*9	42	48	35	34	22	59	18
55.0	33	20	14	*	9	14	9	*8	*6	14	*
55.03	19	10	*9	*	*	*8	*7	*	*	*8	*
55.04	*8	*	*	*	*	*	*	*	*	*	*
55.2	23	13	10	*	9	*7	*	*6	*	10	*
55.23	21	12	9	*	*9	*6	*	*	*	10	*
55.4	*6	*	*	*	*	*	*	*	*	*	*
55.5	40	22	18	*	12	14	12	*8	*6	20	*5
55.51	37	20	17	*	10	13	11	*7	*6	19	*5
55.6	15	11	*	*	*7	*7	*	*	*	*7	*
55.69	15	11	*	*	*7	*7	*	*	*	*7	*
55.8	*6	*	*	*	*	*	*	*	*	*	*
55.87	*6	*	*	*	*	*	*	*	*	*	*
56	98	51	47	13	28	32	25	25	20	37	16
56.0	45	26	19	*	17	17	10	*8	11	20	*5
56.3	22	11	11	*	*8	*8	*6	*	*	10	*
56.31	21	10	11	*	*8	*8	*5	*	*	10	*
56.5	*7	*	*	—	*	*	*	*	*	*	*
56.51	*6	*	*	—	*	*	*	*	*	*	*
56.7	16	*7	*9	10	*	*	*	*8	*	*	*
56.74	15	*7	*8	10	*	*	*	*8	*	*	*
57	398	219	179	10	67	98	223	99	90	134	75
57.0	29	25	*	—	*	*6	21	*6	*7	10	*6
57.1	33	18	16	*	*5	11	17	*6	*7	15	*6
57.17	15	*9	*6	*	*	*	*9	*	*	*9	*
57.18	13	*7	*7	*	*	*5	*6	*	*	*	*
57.3	174	87	87	*	40	52	77	42	41	65	26
57.32	165	81	84	*	39	50	72	38	40	63	24
57.33	*8	*5	*	*	*	*	*5	*	*	*	*
57.4	42	33	9	—	*	*8	33	11	11	14	*6
57.49	42	33	9	—	*	*8	33	11	11	14	*6
57.7	*7	*5	*	—	*	*	*	*	*	*	*
57.71	*6	*5	*	—	*	*	*	*	*	*	*
57.8	21	*6	16	*	*8	*6	*5	*	*	9	*5
57.81	14	*	11	*	*6	*	*	*	*	*7	*
57.9	85	44	42	*	10	12	62	28	17	16	24
57.94	74	33	41	*	9	10	54	27	14	11	22
58	43	34	*9	*	*6	11	23	9	11	17	*6
58.5	*6	*5	*	*	*	*	*	*	*	*	*
58.6	24	19	*	*	*	*5	14	*	*7	11	*
59	283	72	211	*	89	111	80	52	64	117	50
59.0	*6	*	*6	*	*	*	*	*	*	*	*
59.3	*6	—	*6	—	*	*	*	*	*	*	*
59.4	*7	*	*6	*	*	*	*	*	*	*	*
59.5	43	*	43	—	15	22	*7	*	11	18	10
59.7	67	*	67	—	14	32	22	9	15	28	15
59.79	66	*	66	—	13	31	21	*9	15	28	15
59.8	143	68	75	*	54	45	42	33	31	57	22
59.9	*6	*	*	*	*	*	*	*	*	*	*
60-64	262	262	—	18	12	79	152	57	58	98	49
60	206	206	—	—	*	67	138	43	45	77	41

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
60.1	*8	*8	—	—	—	*	*7	*	*	*	*
60.11	*8	*8	—	—	—	*	*7	*	*	*	*
60.2	129	129	—	—	*	23	106	26	28	54	21
60.29	126	126	—	—	*	22	104	26	27	52	21
60.5	57	57	—	—	*	41	16	10	13	18	16
61	12	12	—	*	*5	*	*	*	*	*6	*
61.0	*6	*6	—	*	*	*	*	*	*	*	*
62	10	10	—	*	*	*	*	*	*	*	*
63	*6	*6	—	*	*	*	*	*	*	*	*
64	27	27	—	13	*	*6	*6	*6	*7	11	*
64.0	16	16	—	12	*	*	*	*	*	*6	*
64.9	*8	*8	—	*	*	*	*	*	*	*	*
65-71	2,161	—	2,161	*5	1,256	668	231	345	432	943	442
65	630	—	630	*	303	264	62	99	133	267	131
65.2	49	—	49	*	42	*5	*	9	*8	19	13
65.25	10	—	10	*	*9	*	*	*	*	*	*
65.29	39	—	39	*	33	*	*	*7	*6	15	10
65.3	23	—	23	*	17	*	*	*	*5	*9	*
65.31	*6	—	*6	—	*6	*	—	*	*	*	*
65.39	17	—	17	*	11	*	*	*	*	*6	*
65.4	102	—	102	*	66	29	*7	16	21	48	17
65.41	13	—	13	*	*8	*	*	*	*	*	*
65.49	89	—	89	*	57	25	*6	15	17	44	14
65.5	11	—	11	—	*	*	*	*	*	*	*
65.51	*8	—	*8	—	*	*	*	*	*	*	*
65.6	397	—	397	—	138	213	47	57	87	172	81
65.61	347	—	347	—	118	187	41	52	74	148	72
65.62	13	—	13	—	*6	*	*	*	*	*	*
65.63	36	—	36	—	13	19	*	*	*8	18	*6
65.8	29	—	29	*	21	*7	*	*	*7	11	*7
65.81	*5	—	*5	—	*	*	—	*	*	*	*
65.89	24	—	24	*	18	*6	*	*	*6	9	*
65.9	*8	—	*8	*	*7	*	*	*	*	*	*
65.91	*7	—	*7	*	*6	*	*	*	*	*	*
66	393	—	393	*	387	*	*	46	61	205	80
66.0	*6	—	*6	—	*5	*	*	*	*	*	*
66.2	*7	—	*7	—	*7	—	—	*	*	*5	*
66.3	322	—	322	—	322	*	—	34	49	175	64
66.32	217	—	217	—	217	*	—	23	28	119	47
66.39	105	—	105	—	105	*	—	11	21	56	17
66.4	11	—	11	*	10	*	*	*	*	*	*
66.5	*8	—	*8	—	*6	*	*	*	*	*	*
66.51	*7	—	*7	—	*5	*	*	*	*	*	*
66.6	30	—	30	*	28	*	*	*5	*7	11	*7
66.61	*6	—	*6	*	*6	*	—	*	*	*	*
66.62	19	—	19	—	19	*	—	*	*	*7	*
67	17	—	17	—	11	*	*	*	*	10	*
67.5	*7	—	*7	—	*7	—	—	*	*	*	*
67.59	*7	—	*7	—	*7	—	—	*	*	*	*
68	731	—	731	*	390	274	68	120	162	305	144
68.1	15	—	15	*	*9	*	*	*	*	*5	*
68.12	*8	—	*8	—	*	*	*	*	*	*	*
68.2	45	—	45	—	37	*7	*	*9	*7	21	*9
68.29	41	—	41	—	34	*7	*	*8	*6	20	*8
68.3	32	—	32	—	19	12	*	*8	*6	12	*7
68.4	424	—	424	—	214	176	35	68	96	184	77
68.5	207	—	207	—	109	71	26	30	50	82	45
68.51	68	—	68	—	38	26	*	*7	18	31	11
68.59	139	—	139	—	71	45	23	23	32	50	34

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
69	105	—	105	*	86	14	*5	20	17	46	22
69.0	55	—	55	—	42	9	*	*8	*8	25	14
69.02	29	—	29	—	29	*	—	*	*	14	*9
69.09	26	—	26	—	13	9	*	*	*5	10	*5
69.1	*6	—	*6	*	*	*	*	*	*	*	*
69.19	*6	—	*6	*	*	*	*	*	*	*	*
69.2	*8	—	*8	—	*	*	*	*	*	*	*
69.22	*6	—	*6	—	*	*	*	*	*	*	*
69.5	23	—	23	—	23	*	*	*7	*	9	*
69.52	16	—	16	—	16	—	—	*	*	*8	*
69.9	12	—	12	—	11	*	*	*	*	*6	*
69.96	10	—	10	—	10	*	—	*	*	*	*
70	249	—	249	*	62	94	90	46	51	100	51
70.2	*6	—	*6	*	*	*	*	*	*	*	*
70.3	12	—	12	—	10	*	*	*	*	*5	*
70.32	*7	—	*7	—	*7	*	*	*	*	*	*
70.5	136	—	136	*	24	60	52	22	30	54	29
70.50	68	—	68	—	11	28	29	13	15	26	14
70.51	39	—	39	—	*7	17	15	*6	*9	18	*6
70.52	29	—	29	*	*6	15	*8	*	*6	11	*9
70.7	40	—	40	*	9	14	17	9	*7	15	10
70.77	27	—	27	—	*	10	13	*6	*	9	*7
70.79	*6	—	*6	*	*	*	*	*	*	*	*
70.9	44	—	44	*	12	16	16	10	*8	18	*8
70.92	43	—	43	—	11	16	15	10	*8	17	*8
71	35	—	35	*	17	13	*	10	*6	9	11
71.3	*9	—	*9	*	*5	*	*	*	*	*	*
71.7	12	—	12	*	*	*	*	*	*	*	*
71.79	*8	—	*8	*	*	*	*	*	*	*	*
72-75	6,646	—	6,646	21	6,617	*8	—	1,054	1,387	2,569	1,635
72	337	—	337	*	334	—	—	39	62	149	87
72.0	29	—	29	—	29	—	—	*	*5	14	*7
72.1	56	—	56	*	56	—	—	*6	*9	34	*7
72.2	*5	—	*5	—	*5	—	—	—	*	*	*
72.4	*5	—	*5	—	*5	—	—	*	*	*	*
72.7	234	—	234	*	232	—	—	29	46	89	71
72.71	125	—	125	*	124	—	—	18	22	48	37
72.79	110	—	110	*	108	—	—	11	24	41	33
73	3,016	—	3,016	*7	3,005	*	—	450	647	1,210	709
73.0	901	—	901	*	899	*	—	127	249	306	220
73.01	157	—	157	*	157	*	—	14	49	58	36
73.09	744	—	744	*	742	*	—	113	200	248	184
73.1	*9	—	*9	—	*9	—	—	*	*	*	*
73.4	514	—	514	*	512	*	—	71	106	196	140
73.5	993	—	993	*	988	*	—	154	167	446	225
73.51	*8	—	*8	—	*8	—	—	*	*	*	*
73.59	984	—	984	*	980	*	—	153	165	443	223
73.6	593	—	593	*	591	*	—	94	120	257	122
73.9	*5	—	*5	—	*5	—	—	*	*	*	*
74	1,060	—	1,060	*	1,057	*	—	162	176	489	234
74.0	13	—	13	*	13	—	—	*	*	*7	*
74.1	1,035	—	1,035	*	1,032	*	—	160	170	477	228
74.9	*7	—	*7	*	*7	—	—	—	*	*	*
74.99	*7	—	*7	*	*7	—	—	—	*	*	*
75	2,232	—	2,232	*9	2,221	*	—	403	502	721	606
75.1	15	—	15	*	15	—	—	*	*	*7	*
75.3	910	—	910	*	905	*	—	191	235	239	246
75.32	183	—	183	*	181	*	—	34	64	44	41
75.34	614	—	614	*	611	*	—	141	139	165	169
75.35	72	—	72	*	71	*	—	11	26	16	19

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
75.37	39	—	39	—	39	*	—	*	*6	14	15
75.4	41	—	41	*	41	—	—	*6	*8	15	12
75.5	10	—	10	—	10	—	—	*	*	*	*
75.51	*8	—	*8	—	*8	—	—	*	*	*	*
75.6	1,224	—	1,224	*5	1,218	*	—	196	251	439	339
75.61	37	—	37	*	37	—	—	*	*6	16	10
75.62	120	—	120	*	118	—	—	19	24	43	34
75.69	1,067	—	1,067	*	1,063	*	—	173	221	379	294
75.7	24	—	24	—	24	*	—	*	*	14	*
76-84	3,442	1,652	1,790	173	811	1,087	1,371	704	789	1,234	715
76	73	47	26	*7	46	15	*	12	10	34	17
76.6	19	*8	11	*	16	*	*	*	*	*9	*5
76.65	*6	*	*	*	*5	*	—	*	*	*	*
76.7	31	26	*	*	19	*9	*	*	*5	14	*7
76.75	*6	*	*	*	*	*	*	*	*	*	*
76.76	11	9	*	*	*7	*	*	*	*	*7	*
76.9	12	*9	*	*	*7	*	*	*	*	*6	*
77	293	144	148	26	83	103	80	59	63	109	62
77.2	*8	*6	*	*	*	*	*	*	*	*	*
77.3	13	*7	*6	*6	*	*	*	*5	*	*	*
77.4	35	16	19	*	*6	*6	22	*8	*7	14	*7
77.45	*5	*	*	*	*	*	*	*	*	*	*
77.49	20	9	11	*	*	*	13	*	*	9	*
77.5	13	*	12	*	*	*	*7	*	*6	*	*
77.6	50	30	19	*5	12	18	15	12	10	18	10
77.65	*7	*	*	*	*	*	*	*	*	*	*
77.67	*7	*	*	*	*	*	*	*	*	*	*
77.68	*9	*5	*	*	*	*	*	*	*	*	*
77.69	13	*7	*7	*	*	*6	*	*	*	*6	*
77.7	123	59	64	*6	46	50	21	24	24	49	27
77.79	113	56	57	*	42	47	19	22	22	45	25
77.8	40	20	20	*	*8	19	12	*7	10	15	*8
77.81	15	*6	*9	—	*	10	*	*	*	*6	*
77.88	*6	*	*	*	*	*	*	*	*	*	*
77.89	9	*5	*	*	*	*	*	*	*	*	*
77.9	*6	*	*	*	*	*	*	*	*	*	*
78	236	106	130	27	55	72	82	50	45	88	53
78.0	35	15	20	*	11	11	11	*6	*6	14	*9
78.03	*5	*	*	*	*	*	*	*	*	*	*
78.05	9	*	*7	*	*	*	*5	*	*	*	*
78.07	12	*7	*6	*	*	*	*	*	*	*5	*
78.1	36	20	15	*	12	12	*8	*8	*9	13	*6
78.13	*7	*	*5	*	*	*	*	*	*	*	*
78.17	15	10	*	*	*6	*	*	*	*	*5	*
78.4	24	9	15	*	*	*	14	*	*	10	*
78.49	16	*6	10	*	*	*	13	*	*	*5	*
78.5	58	23	35	10	11	11	26	13	11	18	16
78.55	35	10	25	*6	*	*5	21	9	*5	12	9
78.57	*8	*6	*	*	*	*	*	*	*	*	*
78.59	*6	*	*	*	*	*	*	*	*	*	*
78.6	72	33	40	*5	17	30	20	15	13	28	16
78.65	21	*8	13	*	*	*	11	*6	*	*5	*
78.67	16	*9	*7	*	*	*8	*	*	*	*6	*
78.69	23	11	12	*	*7	13	*	*	*	12	*
79	667	319	348	60	199	137	271	145	162	225	136
79.0	74	38	36	18	19	13	24	17	20	21	16
79.01	10	*	*7	*	*	*	*6	*	*	*	*
79.02	22	10	12	*	*5	*	*8	*5	*5	*7	*
79.05	*9	*	*	*	*	*	*	*	*	*	*
79.06	25	14	11	*	*7	*	*8	*6	*7	*7	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
79.1 . . . . .	80	35	45	13	17	13	37	21	18	23	19
79.11 . . . . .	11	*5	*6	*8	*	*	*	*	*	*	*
79.12 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
79.15 . . . . .	43	14	29	*	*6	*5	29	12	12	11	*8
79.16 . . . . .	11	*8	*	*	*6	*	*	*	*	*	*
79.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
79.3 . . . . .	414	186	229	15	129	86	185	88	100	145	82
79.31 . . . . .	29	11	18	*	*7	*7	11	*9	*6	*8	*7
79.32 . . . . .	35	22	13	*	15	*8	*8	*7	*8	10	10
79.34 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
79.35 . . . . .	175	57	118	*	18	21	133	41	46	62	26
79.36 . . . . .	137	69	67	*	62	42	30	24	34	49	29
79.37 . . . . .	16	10	*7	*	12	*	*	*	*	*5	*
79.39 . . . . .	10	*7	*	*	*8	*	*	*	*	*	*
79.6 . . . . .	44	31	13	*5	21	13	*5	*8	10	18	*9
79.62 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
79.66 . . . . .	17	11	*6	*	*8	*	*	*	*	*7	*
79.7 . . . . .	36	18	18	*	*7	*8	18	*9	10	10	*7
79.71 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
79.75 . . . . .	23	11	12	*	*	*5	13	*6	*7	*7	*
79.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
80 . . . . .	481	259	222	*8	161	196	116	87	94	194	106
80.0 . . . . .	10	*6	*	—	*	*	*7	*	*	*	*
80.1 . . . . .	24	16	*8	*	*8	*8	*8	*	*	10	*5
80.15 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
80.16 . . . . .	9	*6	*	*	*	*	*	*	*	*	*
80.2 . . . . .	33	18	15	*	*7	11	15	*5	*6	16	*7
80.21 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
80.26 . . . . .	25	14	11	*	*5	*6	14	*	*	13	*
80.4 . . . . .	15	*5	9	*	*	*	*6	*	*	*	*7
80.46 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
80.5 . . . . .	323	171	152	*	120	147	55	56	62	135	70
80.51 . . . . .	319	169	150	*	118	145	54	56	61	133	69
80.6 . . . . .	15	9	*6	—	*6	*6	*	*	*	*	*
80.7 . . . . .	15	*7	*8	*	*	*5	*7	*	*	*7	*
80.76 . . . . .	12	*	*7	*	*	*	*6	*	*	*6	*
80.8 . . . . .	38	22	16	*	12	13	13	*7	10	13	*8
80.81 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
80.85 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
80.86 . . . . .	14	*8	*6	*	*5	*	*	*	*	*	*
81 . . . . .	1,277	538	740	16	182	422	657	266	324	434	253
81.0 . . . . .	296	146	151	*9	98	132	58	50	64	116	66
81.02 . . . . .	111	58	54	—	34	59	19	17	21	52	20
81.03 . . . . .	14	*8	*6	*	*	*7	*	*	*	*7	*
81.04 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
81.05 . . . . .	18	*9	*9	*	10	*	*	*	*	*5	*
81.06 . . . . .	22	9	13	*	12	*9	*	*	*5	*8	*
81.07 . . . . .	14	*7	*7	—	*	*5	*5	*	*6	*	*
81.08 . . . . .	109	52	57	*	32	48	27	16	24	38	31
81.1 . . . . .	21	*8	13	*	*6	*8	*7	*	*8	*6	*
81.11 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
81.3 . . . . .	18	*7	11	*	*7	*8	*	*	*	*8	*
81.38 . . . . .	10	*	*6	*	*	*	*	*	*	*5	*
81.4 . . . . .	32	15	18	*	19	*7	*	*7	*	12	*8
81.45 . . . . .	12	*6	*7	*	10	*	—	*	*	*	*
81.47 . . . . .	12	*5	*7	*	*7	*	*	*	*	*	*
81.5 . . . . .	762	290	472	*	29	226	506	170	201	251	140
81.51 . . . . .	193	81	112	—	11	67	114	54	48	55	36
81.52 . . . . .	109	29	80	*	*	10	97	18	30	35	26
81.53 . . . . .	43	19	23	*	*5	9	28	10	10	14	*9
81.54 . . . . .	381	146	235	—	10	131	241	81	104	132	65

See footnotes at end of table.



**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
81.55	35	14	21	—	*	*7	26	*6	10	14	*5
81.8	43	16	27	*	*5	16	22	*7	15	11	11
81.80	*7	*	*	—	*	*	*	*	*	*	*
81.81	16	*	12	—	*	*	12	*	*7	*	*
81.83	17	*8	9	*	*	9	*	*	*	*	*7
81.9	97	53	45	*	15	23	55	26	26	28	18
81.91	59	34	25	*	*7	16	32	15	15	18	12
81.92	29	13	17	—	*	*	22	*9	10	*7	*
82	29	21	*8	*	18	*7	*	*	*	15	*
82.0	*7	*	*	*	*	*	*	*	*	*	*
82.4	15	13	*	*	10	*	*	*	*	10	*
82.45	*8	*8	*	*	*	*	*	*	*	*6	*
83	220	115	106	25	52	74	69	44	51	71	54
83.0	24	15	*9	*	*9	*8	*6	*5	*6	*7	*6
83.09	15	*9	*6	*	*5	*5	*	*	*	*	*
83.1	21	12	*9	*7	*6	*	*	*	*	*6	*6
83.13	*6	*	*	*	*	*	*	*	*	*	*
83.14	10	*7	*	*	*	*	*	*	*	*	*
83.2	15	*7	*8	*	*	*	*5	*	*	*6	*
83.21	15	*7	*8	*	*	*	*5	*	*	*6	*
83.3	16	*6	10	*	*	*6	*7	*	*	*8	*
83.39	14	*	9	*	*	*5	*6	*	*	*7	*
83.4	28	14	14	*	*7	9	10	*5	*7	*9	*7
83.45	17	*9	*8	*	*	*7	*	*	*	*6	*
83.5	*7	*	*	*	*	*	*	*	*	*	*
83.6	61	33	29	*	10	27	22	11	15	19	17
83.63	40	20	21	—	*	22	17	*8	11	10	11
83.64	10	*6	*	*	*	*	*	*	*	*	*
83.65	9	*6	*	*	*	*	*	*	*	*	*
83.7	*8	*	*	*	*	*	*	*	*	*	*
83.8	32	17	15	*8	*8	*8	*7	*8	*6	9	9
83.82	*7	*	*	*	*	*	*	*	*	*	*
83.85	12	*6	*5	*7	*	*	*	*	*	*	*
83.88	*7	*	*	*	*	*	*	*	*	*	*
83.9	*9	*	*	*	*	*	*6	*	*	*	*
84	166	103	63	*	17	60	87	38	35	64	29
84.0	11	*8	*	*	*	*	*	*	*	*	*
84.01	*9	*6	*	*	*	*	*	*	*	*	*
84.1	134	81	53	*	9	46	79	30	28	52	25
84.11	49	32	18	*	*	22	24	14	11	16	*8
84.12	12	*9	*	*	*	*	*7	*	*	*	*
84.15	40	22	18	—	*	14	23	*6	9	16	9
84.17	30	17	13	*	*	*6	23	*7	*5	13	*
84.3	19	13	*6	*	*	9	*6	*	*	*8	*
85-86	1,348	616	732	115	388	415	429	291	260	576	221
85	183	*	181	*	41	89	53	44	33	70	36
85.1	13	—	13	*	*	*5	*6	*	*	*6	*
85.11	*5	—	*5	—	*	*	*	*	*	*	*
85.12	*6	—	*6	*	*	*	*	*	*	*	*
85.2	18	*	18	*	*	*8	*7	*	*	*6	*
85.21	11	*	11	*	*	*5	*	*	*	*	*
85.23	*5	—	*5	—	*	*	*	*	*	*	*
85.3	13	*	12	*	*8	*	*	*	*	*	*
85.32	10	*	10	*	*7	*	*	*	*	*	*
85.4	83	*	82	—	12	36	35	16	17	31	18
85.41	21	*	21	—	*	11	*8	*	*	*9	*
85.43	51	*	51	—	*8	21	23	10	12	20	9
85.5	*7	—	*7	—	*	*	*	*	*	*	*
85.7	10	—	10	—	*	*7	*	*	*	*	*
85.8	11	*	10	—	*	*6	*	*	*	*5	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
85.89	*6	—	*6	—	*	*	*	*	*	*	*
85.9	20	*	19	—	*5	13	*	*6	*	*7	*
85.95	12	*	12	—	*	*8	*	*	*	*5	*
86	1,164	614	551	115	347	326	376	247	227	506	185
86.0	293	150	144	30	86	94	83	65	57	119	52
86.01	12	*6	*6	*	*	*	*	*	*	*	*
86.04	137	73	65	12	50	44	32	26	29	53	29
86.05	40	19	20	*	11	13	12	10	*7	18	*5
86.06	*7	*	*	*	*	*	*	*	*	*	*
86.07	71	34	37	*7	14	21	29	18	12	31	9
86.09	23	12	11	*	*	9	*8	*6	*	*8	*
86.1	23	12	11	*	*	*6	*8	*8	*	*8	*
86.11	23	12	11	*	*	*6	*8	*8	*	*8	*
86.2	406	217	189	19	94	121	172	94	91	158	64
86.22	287	159	128	13	72	92	110	62	60	120	45
86.23	*6	*	*	*	*	*	*	*	*	*	*
86.27	36	13	23	*	*	*5	29	15	12	*7	*
86.28	74	41	33	*6	16	22	30	14	17	26	16
86.3	65	24	40	*7	25	16	17	11	12	32	10
86.4	12	*5	*7	*	*	*	*7	*	*	*	*
86.5	163	95	68	*6	73	33	51	34	40	54	34
86.59	163	95	68	*6	73	33	51	34	40	54	34
86.6	124	77	47	39	33	28	24	13	10	92	*8
86.66	*9	*6	*	*	*	*	*	*	*	*8	—
86.69	96	59	38	34	27	19	17	10	*8	70	*8
86.7	37	22	15	*	13	10	10	*8	*6	19	*
86.74	24	14	10	*	*8	*7	*6	*	*	13	*
86.8	38	10	28	*	16	14	*	10	*6	17	*5
86.83	18	*	17	*	9	*8	*	*6	*	*6	*
86.84	*9	*	*	*	*	*	*	*	*	*7	*
86.89	*7	*	*	*	*	*	*	*	*	*	*
87-99	12,332	5,996	6,336	890	2,382	3,381	5,679	3,472	2,629	3,886	2,344
87	854	376	478	51	185	216	401	253	156	264	181
87.0	313	148	165	20	58	64	171	141	59	59	53
87.03	310	146	164	20	56	64	171	140	59	58	53
87.2	20	*8	12	*	*8	*	*6	*	*8	*8	*
87.21	*9	*	*6	*	*	*	*	*	*	*	*
87.4	213	96	116	13	39	49	112	53	30	75	55
87.41	111	48	62	*	19	31	56	49	16	24	21
87.44	97	46	51	*8	18	17	53	*	13	49	33
87.5	136	41	94	*	39	43	52	23	24	57	32
87.51	*7	*	*	*	*	*	*	*	*	*	*
87.53	120	35	85	*	38	39	43	20	21	51	29
87.54	*7	*	*	*	*	*	*	*	*	*	*
87.6	37	17	20	*6	*7	10	15	*7	*	11	14
87.61	*8	*6	*	*	*	*	*	*	*	*	*
87.62	16	*6	10	*	*	*	*	*	*	*5	*5
87.63	*6	*	*	*	*	*	*	*	*	*	*
87.7	132	64	67	10	34	44	43	27	29	53	22
87.73	20	9	11	*	*6	9	*5	*	*	11	*
87.74	84	39	45	*	23	30	28	20	20	33	11
87.76	*7	*	*	*	*	*	*	*	*	*	*
87.79	*7	*	*	*	*	*	*	*	*	*	*
88	3,428	1,765	1,663	103	464	1,186	1,675	896	773	1,164	595
88.0	208	95	113	10	58	58	82	94	39	41	35
88.01	207	94	112	10	57	57	82	93	39	40	34
88.1	19	11	*8	*	*	*6	*6	*	*	12	*
88.19	14	*9	*6	*	*	*	*	*	*	*8	*
88.2	18	*5	13	*	*5	*	*9	*	*	*	*9
88.26	*7	*	*6	—	*	*	*	—	*	*	*
88.3	79	37	42	*8	24	19	27	36	15	15	12

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
88.38	73	34	38	*7	23	16	27	35	14	14	10
88.4	456	237	219	*7	48	146	255	79	121	171	84
88.41	123	55	68	*	18	44	60	27	31	47	18
88.42	131	75	56	*	*7	40	82	18	40	50	22
88.43	21	*9	12	*	*	*5	9	*	*	*5	10
88.44	11	*7	*	*	*	*	*5	*	*	*	*
88.45	33	14	18	*	*	9	21	*	*7	17	*6
88.47	28	15	13	*	*	10	15	*	*7	12	*6
88.48	85	50	35	*	*8	26	51	18	24	29	14
88.49	22	11	10	*	*5	*6	10	*	*6	*6	*7
88.5	1,602	906	696	*6	146	677	773	312	397	647	246
88.51	13	*7	*6	*	*	*	*6	*	*	*6	*
88.53	632	349	283	*	60	278	291	124	153	268	87
88.54	22	12	10	*	*	9	10	*8	*	*7	*
88.55	17	11	*6	-	*	*8	*6	*5	*	*6	*
88.56	854	490	364	*	76	349	429	152	222	333	147
88.57	57	34	24	*	*	27	26	19	11	24	*
88.6	22	10	12	*	*	9	*8	*	*5	9	*
88.67	11	*	*7	*	*	*	*	*	*	*	*
88.7	773	347	427	45	133	192	404	271	137	200	165
88.71	49	22	27	*	*	11	29	18	*8	12	11
88.72	472	224	248	20	53	124	274	158	80	141	92
88.73	*7	*5	*	*	*	*	*	*	*	*	*
88.74	17	*7	9	*	*6	*	*	*	*	*5	*
88.75	37	19	17	*7	*9	*	17	14	*	*8	11
88.76	88	35	53	11	26	21	31	36	14	11	27
88.77	69	30	39	*	*9	23	37	30	19	10	10
88.78	21	-	21	*	21	-	-	*6	*	*	*6
88.79	15	*	10	*	*	*	*6	*6	*	*6	*
88.9	251	117	134	24	40	76	111	98	51	64	37
88.91	150	65	85	16	23	49	62	63	30	35	22
88.93	42	24	19	*	*9	13	17	14	10	11	*8
88.94	13	*8	*	*	*	*	*6	*7	*	*	*
88.97	41	18	23	*	*5	*9	25	13	*8	15	*
89	746	361	385	51	115	200	379	271	127	185	163
89.0	62	32	30	*	*9	18	33	22	-	*8	32
89.04	*6	*	*	*	*	*	*	*6	-	*	-
89.05	14	*7	*8	-	*	*	9	14	-	-	*
89.06	15	*8	*7	-	*	*5	*9	-	-	*	15
89.07	*6	*	*	*	-	*	*	-	-	-	*6
89.08	*8	*	*	*	*	*	*	*	-	*6	-
89.09	12	*7	*	-	*	*	*6	*	-	-	11
89.1	86	41	45	25	21	14	26	36	*9	28	13
89.14	55	26	29	14	11	9	21	22	*	17	11
89.15	*6	*	*	*	*	*	*	*	*	*	*
89.19	21	10	12	*8	*8	*	*	11	*	*8	*
89.2	17	*8	9	-	*	*	*8	*7	*	*6	*
89.22	*8	*	*	-	*	*	*6	*5	*	*	*
89.3	26	15	11	10	*5	*	*6	11	*7	*	*
89.37	*6	*	*	*	*	*	*	*	*	*	*
89.38	*7	*	*	*	*	*	*	*	*	*	*
89.39	9	*	*	*5	*	*	*	*	*	*	*
89.4	143	73	70	*	18	54	71	49	16	32	46
89.41	48	28	20	*	*6	21	20	19	*	9	15
89.44	91	42	49	*	12	32	48	29	11	21	31
89.5	284	130	154	*8	42	63	171	107	61	76	39
89.50	*7	*	*5	*	*	*	*6	*	*	*	*
89.52	115	51	64	*	21	27	63	18	15	58	23
89.54	151	71	79	*	19	33	95	83	40	14	13
89.59	11	*5	*6	*	*	*	*7	*	*	*	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
89.6 . . . . .	129	63	65	*6	16	42	65	40	32	31	27
89.61 . . . . .	10	*	*5	*	*	*	*	*	*	*	*
89.62 . . . . .	10	*	*5	*	*	*	*5	*	*	*	*
89.64 . . . . .	44	23	20	*	*	13	26	10	14	12	*8
89.65 . . . . .	43	19	24	*	*7	13	22	15	*7	11	10
89.68 . . . . .	17	*9	*8	*	*	*7	*7	*5	*6	*	*
90 . . . . .	18	*7	11	*	*	*	*8	*	*7	*7	*
90.5 . . . . .	*8	*	*5	*	*	*	*	*	*	*6	*
90.59 . . . . .	*6	*	*	*	*	*	*	*	—	*	*
90.8 . . . . .	*8	*	*	—	*	*	*	*	*7	*	—
90.81 . . . . .	*7	*	*	—	*	*	*	*	*7	*	—
91 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
92 . . . . .	257	129	128	*7	34	84	132	94	30	71	62
92.0 . . . . .	96	47	49	*	11	32	50	34	10	19	32
92.02 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
92.04 . . . . .	11	*6	*	*	*	*	*6	*	*	*	*
92.05 . . . . .	71	34	37	*	*6	26	40	24	*8	15	25
92.1 . . . . .	58	24	34	*	10	15	31	31	*	11	12
92.14 . . . . .	22	12	10	*	*	*5	14	12	*	*	*5
92.15 . . . . .	30	10	20	—	*7	*9	14	16	*	*5	*6
92.2 . . . . .	96	56	41	*	12	33	49	26	15	39	16
92.24 . . . . .	17	13	*	*	*	*	11	*6	*	*6	*
92.27 . . . . .	28	17	11	—	*	10	15	*7	*6	*8	*7
92.29 . . . . .	45	23	21	*	*5	18	20	10	*8	22	*
92.3 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
92.32 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
93 . . . . .	1,415	641	774	166	176	282	792	366	301	429	319
93.0 . . . . .	34	17	17	*	*6	10	17	19	*8	*	*
93.01 . . . . .	30	15	15	*	*6	*9	15	18	*8	*	*
93.1 . . . . .	24	10	13	*	*	*6	17	17	*	*	*
93.11 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
93.12 . . . . .	12	*6	*6	*	*	*	*9	*9	—	*	—
93.2 . . . . .	32	15	17	*	*	12	16	22	*	*5	*
93.22 . . . . .	22	10	12	*	*	*6	14	18	*	*	*
93.3 . . . . .	317	130	186	*6	22	58	231	57	69	119	72
93.38 . . . . .	22	11	10	*	*	*7	14	*	*	11	*8
93.39 . . . . .	288	115	173	*5	20	49	214	51	67	106	64
93.4 . . . . .	12	*8	*	*	*	*	*	*	*	*	*
93.46 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
93.5 . . . . .	103	54	48	20	28	18	37	26	21	39	17
93.53 . . . . .	19	10	*9	*5	*	*	*7	*6	*	*	*5
93.54 . . . . .	24	11	13	*	*8	*	10	*	*6	*7	*7
93.57 . . . . .	24	16	*8	10	*8	*	*	*	*	16	*
93.59 . . . . .	25	10	15	*	*	*8	13	*8	*5	*8	*
93.7 . . . . .	38	20	18	*	*	*8	27	*5	14	*9	11
93.75 . . . . .	38	20	18	*	*	*8	27	*5	14	*8	10
93.8 . . . . .	430	184	245	11	57	91	271	90	84	166	90
93.81 . . . . .	46	23	23	*	16	11	17	*6	12	21	*8
93.83 . . . . .	202	90	111	*	24	46	127	47	59	63	33
93.89 . . . . .	177	69	108	*	14	33	127	37	12	78	49
93.9 . . . . .	422	200	223	125	51	76	171	128	94	80	121
93.90 . . . . .	90	46	44	13	*9	24	44	30	27	16	17
93.94 . . . . .	181	83	98	70	23	26	62	58	41	28	53
93.96 . . . . .	131	62	69	37	13	22	59	32	23	30	47
93.99 . . . . .	14	*7	*7	*	*	*	*	*5	*	*	*
94 . . . . .	726	430	297	42	453	189	42	344	169	153	60
94.0 . . . . .	10	*6	*	*	*	*	*	*	—	*9	—
94.2 . . . . .	152	74	78	11	91	33	17	109	19	22	*
94.22 . . . . .	*5	*	*	*	*	*	*	*	*	*	—
94.23 . . . . .	34	18	16	*	24	*6	*	32	—	*	—

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
94.25	82	42	41	*6	54	18	*	63	11	*7	*
94.27	29	11	18	—	9	*9	11	*8	*7	13	*
94.3	96	49	46	13	59	18	*6	54	28	13	*
94.37	*7	*	*	*	*	—	—	*7	—	—	—
94.38	39	19	20	*	27	9	*	39	—	*	—
94.39	45	23	21	*6	27	*9	*	*	27	13	*
94.4	109	55	54	13	69	22	*5	52	26	31	—
94.42	14	*7	*7	*	*7	*	*	10	*	*	—
94.44	85	40	45	*8	53	18	*	33	24	28	—
94.6	355	243	112	*	227	114	13	125	96	76	58
94.61	12	*6	*6	*	*9	*	*	*	*6	*	*
94.62	169	122	47	—	89	69	10	52	61	30	25
94.63	16	9	*7	—	9	*6	*	*	*6	*	*5
94.64	*8	*5	*	*	*7	*	*	*	*	*	*
94.65	76	46	30	*	57	17	*	39	*8	20	9
94.66	*7	*	*	—	*	*	*	*	*	*	*
94.67	*9	*6	*	*	*8	*	*	—	*	*	*
94.68	53	41	12	—	39	14	*	28	*	*9	11
94.69	*7	*	*	*	*	*	*	*	*	*	*
95	11	*7	*	10	*	*	*	*	*	*6	*
95.4	9	*5	*	*9	*	*	*	*	*	*	*
95.47	*5	*	*	*5	—	—	—	*	*	*	—
96	1,537	751	786	170	302	365	700	374	339	512	312
96.0	567	288	280	51	76	154	286	132	124	201	111
96.04	477	247	229	48	62	126	241	108	99	181	88
96.05	*6	*	*	*	*	*	*	*	*	*	*
96.07	78	33	45	*	13	24	38	19	22	15	21
96.2	10	*5	*	*	*	*	*	*	*	*	*
96.3	24	10	15	*	11	*	*7	*	*9	*7	*
96.33	15	*	10	*	10	*	*	*	*7	*	*
96.38	*6	*	*	*	*	*	*	*	*	*	*
96.4	100	*	96	*	96	*	*	10	24	32	33
96.49	96	*	95	*	95	*	*	9	23	31	32
96.5	60	36	25	*8	15	17	20	15	11	16	17
96.52	*5	*	*	*	*	*	*	*	*	*	*
96.56	33	21	12	*5	*6	*8	13	*6	*6	10	10
96.59	18	9	*9	*	*8	*5	*	*5	*	*6	*
96.6	126	63	63	22	9	20	75	64	31	21	11
96.7	648	346	301	83	93	167	305	146	138	231	133
96.70	21	14	*7	*	*	*6	11	*7	*	14	*
96.71	402	210	193	51	59	102	190	85	95	138	83
96.72	224	123	101	31	30	59	104	54	42	79	49
97	105	48	58	13	22	28	42	27	19	37	23
97.0	34	16	18	*	*	*7	20	10	*7	11	*7
97.02	23	10	13	*	*	*	13	*7	*	*7	*
97.05	*8	*	*	*	*	*	*6	*	*	*	*
97.2	11	*5	*6	*	*	*	*	*	*	*	*
97.23	*8	*	*	*	*	*	*	*	*	*	*
97.3	*5	*	*	*	*	*	*	*	*	*	*
97.4	10	*	*7	*	*	*	*	*	*	*6	*
97.49	*9	*	*6	*	*	*	*	*	*	*5	*
97.5	11	*5	*6	*	*	*	*6	*	*	*	*
97.55	*5	*	*	*	*	*	*	*	*	*	*
97.6	14	*6	*9	*	*	*	*6	*	*	*6	*
97.62	12	*	*8	*	*	*	*	*	*	*5	*
97.8	17	*8	*8	*	*5	*6	*	*	*	*6	*
97.82	*6	*	*	*	*	*	*	*	*	*	*
98	26	16	10	*7	*6	*	*8	*6	*	10	*5
98.0	12	*7	*5	*	*	*	*	*	*	*5	*
98.02	10	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

**Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2002—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
98.5 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
98.51 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
99 . . . . .	3,201	1,463	1,738	266	618	821	1,496	839	701	1,044	618
99.0 . . . . .	1,486	646	840	56	180	380	870	290	302	585	308
99.00 . . . . .	20	9	10	*	*	*6	12	*6	*	*6	*6
99.02 . . . . .	62	25	37	*	*5	26	31	11	13	18	20
99.03 . . . . .	25	*8	17	*	*5	*7	12	*6	*6	12	*
99.04 . . . . .	1,187	502	685	40	143	287	717	228	241	478	240
99.05 . . . . .	64	33	31	*8	10	19	27	11	11	28	13
99.07 . . . . .	113	61	53	*	12	33	65	26	24	38	25
99.09 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
99.1 . . . . .	375	147	227	42	129	83	120	111	74	104	87
99.10 . . . . .	48	29	19	*	*5	18	24	9	*8	22	*9
99.11 . . . . .	75	*	75	*	74	*	*	14	16	27	19
99.14 . . . . .	11	*7	*	*	*	*	*	*	*	*	*
99.15 . . . . .	134	61	73	26	24	31	53	40	30	32	33
99.17 . . . . .	29	12	17	*	*6	11	9	15	*6	*	*
99.18 . . . . .	36	18	18	*8	*7	*9	13	15	*	*8	13
99.19 . . . . .	41	21	20	*	9	13	19	14	11	*7	9
99.2 . . . . .	1,056	522	534	104	267	306	379	371	261	270	153
99.20 . . . . .	172	120	53	—	13	77	82	36	36	59	42
99.21 . . . . .	262	115	147	41	89	59	73	99	80	57	26
99.23 . . . . .	95	39	56	*5	18	26	47	34	28	23	10
99.25 . . . . .	217	124	94	34	51	72	61	53	35	80	49
99.28 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
99.29 . . . . .	298	121	177	23	93	70	112	148	76	47	26
99.4 . . . . .	15	*	14	*	15	*	—	*	*	*	*6
99.47 . . . . .	*8	—	*8	—	*8	—	—	*	*	*	*
99.48 . . . . .	*6	—	*6	—	*6	—	—	*	*	*	*
99.5 . . . . .	23	12	11	9	*	*5	*	*6	10	*6	*
99.52 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
99.55 . . . . .	11	*6	*5	*8	*	*	*	*	*	*5	*
99.6 . . . . .	164	91	73	*	15	37	109	42	35	47	39
99.60 . . . . .	58	31	27	*	*6	13	38	14	12	17	15
99.61 . . . . .	40	23	16	*	*	9	25	11	*7	10	11
99.62 . . . . .	60	33	27	*	*	14	41	15	15	17	12
99.69 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
99.7 . . . . .	17	*7	10	*	*6	*5	*	*6	*	*6	*
99.71 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
99.8 . . . . .	52	30	23	50	*	*	*	*6	*8	17	22
99.82 . . . . .	*6	*	*	*6	—	—	—	*	*	*	*
99.83 . . . . .	44	26	18	44	*	*	*	*5	*5	15	19
99.9 . . . . .	11	*6	*	*	*	*	*6	*	*	*5	—
99.99 . . . . .	11	*6	*	*	*	*	*6	*	*	*5	—

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

# Appendix

## Technical Notes

### Abstract Form

Figure I shows the medical abstract form used for the 2002 NHDS. The hospital number, date of birth, and patient ZIP Code are confidential information and are not released to the public.

### Population Estimates

Population estimates used to compute rates are shown in table I. These estimates are for the civilian population of the United States, including institutionalized persons, as of July 1, 2002. The data are postcensal estimates based on the 2000 census and are from unpublished tabulations provided by the U.S. Census Bureau.

### Reliability of Estimates

The statistics presented in this report are based on a sample survey and may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from the NHDS were calculated with SUDAAN software, which takes into account the complex sample design. A description of the software and the approach it uses has been published (21).

The chances are about 32 in 100 that an estimate from the sample would differ from a complete census by more than the standard error. The chances are 5 in 100 that the difference would be more than twice the standard error, and about 1 in 100 that the difference would be more than 2.5 times as large as the standard error. The relative standard error (RSE) expresses the standard error as a percent of the estimate and can be multiplied by the estimate to obtain the standard error.

To derive approximate standard errors for the estimates in this report,

**Table I. Civilian population by age, geographic region, and sex: United States, 2002**

Age and geographic region	Both sexes	Male	Female
All ages			
Total . . . . .	287,123	140,598	146,525
Region:			
Northeast . . . . .	54,170	26,200	27,970
Midwest . . . . .	65,044	31,882	33,162
South . . . . .	102,600	50,024	52,576
West . . . . .	65,309	32,492	32,816
Under 15 years			
Total . . . . .	60,646	31,038	29,609
Under 1 year . . . . .	4,034	2,064	1,970
1–4 years . . . . .	15,575	7,962	7,614
5–14 years . . . . .	41,037	21,013	20,025
Region:			
Northeast . . . . .	10,676	5,464	5,212
Midwest . . . . .	13,602	6,964	6,638
South . . . . .	21,664	11,080	10,584
West . . . . .	14,704	7,530	7,174
15–44 years			
Total . . . . .	124,230	62,363	61,866
15–24 years . . . . .	40,059	20,379	19,680
25–34 years . . . . .	39,512	19,847	19,666
35–44 years . . . . .	44,659	22,138	22,521
Region:			
Northeast . . . . .	23,051	11,468	11,584
Midwest . . . . .	27,952	14,064	13,888
South . . . . .	44,276	22,082	22,194
West . . . . .	28,951	14,749	14,201
45–64 years			
Total . . . . .	66,645	32,425	34,220
45–54 years . . . . .	40,045	19,642	20,402
55–64 years . . . . .	26,600	12,783	13,817
Region:			
Northeast . . . . .	13,018	6,270	6,747
Midwest . . . . .	15,192	7,449	7,742
South . . . . .	23,921	11,570	12,351
West . . . . .	14,514	7,135	7,379
65 years and over			
Total . . . . .	35,602	14,772	20,830
65–74 years . . . . .	18,274	8,301	9,973
75–84 years . . . . .	12,735	5,081	7,654
85 years and over . . . . .	4,593	1,390	3,203
Region:			
Northeast . . . . .	7,425	2,998	4,427
Midwest . . . . .	8,298	3,405	4,893
South . . . . .	12,739	5,292	7,447
West . . . . .	7,139	3,077	4,062

NOTE: Unpublished estimates from the U.S. Census Bureau based on the 2000 census.

first calculate the approximate RSE of the estimate using the parameters shown in table II. Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate  $X$ , may be estimated from the formula:

$$RSE(X) = \sqrt{a + b/X}$$

where  $a$  and  $b$  are defined in table II.

When multiplied by 100,  $RSE(X)$  is expressed as a percent of  $X$ .

The approximate RSE for a percent,  $100p$ , in which the numerator is a subclass of the denominator, may be calculated directly using the formula:

$$RSE(p) = \sqrt{b(1-p) / (pX)}$$

where  $100p$  is the percent of interest,  $X$



OMB No. 0920-0212; Approval Expires 08/31/2005

**Notice** – All information which would permit identification of an individual or an establishment will be held confidential, will be used only by persons engaged in and for the purposes of the survey, and will not be disclosed or released to other persons or used for any other purpose. Public reporting burden of this collection of information is estimated to average 4 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road, MS D-24, Atlanta, GA 30333, ATTN: PRA (0920-0212)

FORM **HDS-1**  
(9-27-2003)

U.S. DEPARTMENT OF COMMERCE  
Economics and Statistics Administration  
U.S. CENSUS BUREAU  
ACTING AS COLLECTING AGENT FOR  
DEPARTMENT OF HEALTH AND HUMAN SERVICES  
CENTERS FOR DISEASE CONTROL AND PREVENTION  
NATIONAL CENTER FOR HEALTH STATISTICS

## MEDICAL ABSTRACT – NATIONAL HOSPITAL DISCHARGE SURVEY

### A. PATIENT IDENTIFICATION

<b>1.</b> Hospital number . . . . .	<input type="text"/>	<b>4.</b> Date of admission . . . . .	Month <input type="text"/> <input type="text"/> - Day <input type="text"/> <input type="text"/> - Year <input type="text"/> <input type="text"/> <input type="text"/>
<b>2.</b> HDS number . . . . .	<input type="text"/>	<b>5.</b> Date of discharge . . . . .	Month <input type="text"/> <input type="text"/> - Day <input type="text"/> <input type="text"/> - Year <input type="text"/> <input type="text"/> <input type="text"/>
<b>3.</b> (Item deleted)		<b>6.</b> Residence ZIP Code . . . . .	<input type="text"/>

### B. PATIENT CHARACTERISTICS

<b>7.</b> Date of birth	Month <input type="text"/> <input type="text"/> - Day <input type="text"/> <input type="text"/> - Year <input type="text"/> <input type="text"/> <input type="text"/>	<b>11.</b> Race – Mark <i>all that apply</i>
<b>8.</b> Age – Complete only if date of birth not given	Units <input type="text"/> <input type="text"/> <input type="text"/> { 1 <input type="checkbox"/> Years 2 <input type="checkbox"/> Months 3 <input type="checkbox"/> Days	
<b>9.</b> Sex – Mark (X) one	1 <input type="checkbox"/> Male    2 <input type="checkbox"/> Female    3 <input type="checkbox"/> Not stated	4 <input type="checkbox"/> Black or African American
<b>10.</b> Ethnicity – Mark (X) one	1 <input type="checkbox"/> Hispanic or Latino    2 <input type="checkbox"/> Not Hispanic or Latino    3 <input type="checkbox"/> Not stated	5 <input type="checkbox"/> American Indian or Alaska Native
		6 <input type="checkbox"/> Other – Specify <input type="text"/>
		7 <input type="checkbox"/> Not stated
		<b>12.</b> Marital status – Mark (X) one
		1 <input type="checkbox"/> Married    2 <input type="checkbox"/> Single    3 <input type="checkbox"/> Widowed    4 <input type="checkbox"/> Divorced    5 <input type="checkbox"/> Separated    6 <input type="checkbox"/> Not stated

### C. ADMINISTRATIVE INFORMATION

<b>13.</b> Type of Admission – Mark (X) one	1 <input type="checkbox"/> Emergency    2 <input type="checkbox"/> Urgent    3 <input type="checkbox"/> Elective    4 <input type="checkbox"/> Newborn    5 <input type="checkbox"/> Items not available/unknown	<b>16.</b> Expected source(s) of payment	Principal	Other additional sources	
<b>14.</b> Source of Admission – Mark (X) one	1 <input type="checkbox"/> Physician referral    2 <input type="checkbox"/> Clinical referral    3 <input type="checkbox"/> HMO referral    4 <input type="checkbox"/> Transfer from a hospital    5 <input type="checkbox"/> Transfer from SNF    6 <input type="checkbox"/> Transfer from other health facility    7 <input type="checkbox"/> Emergency room    8 <input type="checkbox"/> Court/Law enforcement    9 <input type="checkbox"/> Other – Specify <input type="text"/> 10 <input type="checkbox"/> Item not available		Mark one only	Mark all that apply	
<b>15.</b> Status/Disposition of patient – Mark (X) appropriate box(es)	Status 1 <input type="checkbox"/> Alive		1. Worker's compensation . . . . .	<input type="checkbox"/>	<input type="checkbox"/>
	Disposition a. <input type="checkbox"/> Routine discharge/discharged home b. <input type="checkbox"/> Left against medical advice c. <input type="checkbox"/> Discharged, transferred to another short-term hospital d. <input type="checkbox"/> Discharged, transferred to long-term care institution e. <input type="checkbox"/> Other disposition/not stated 2 <input type="checkbox"/> Died 3 <input type="checkbox"/> Status not stated		2. Medicare . . . . .	<input type="checkbox"/>	<input type="checkbox"/>
		3. Medicaid . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		4. Other government payments . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		5. Blue Cross/Blue Shield . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		6. HMO/PPO . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		7. Other private or commercial insurance . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		8. Self pay . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		9. No charge . . . . .	<input type="checkbox"/>	<input type="checkbox"/>	
		10. Other – Specify <input type="text"/>			
		<input type="checkbox"/> No source of payment indicated	<input type="checkbox"/>	<input type="checkbox"/>	

(Over)

Figure I. Medical abstract form for the National Hospital Discharge Survey, 2002



<b>D. MEDICAL INFORMATION</b>							
<b>17. Final Diagnoses (including E-code diagnoses) (Enter ICD-9-CM codes as well as narrative if available)</b>							
Principal: _____							
Other/additional: _____							
<b>18. Surgical and Diagnostic Procedures (Enter ICD-9-CM codes as well as narrative if available)</b>					Date of procedure(s)		
					Month	Day	Year
Principal: _____							
Other/additional: _____							
<input type="checkbox"/> NONE							
Completed by					Date		

Figure I. Medical abstract form for the National Hospital Discharge Survey, 2002—Con.

**Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics, 2002**

Characteristic	Discharges or first-listed diagnoses		Days of care		All-listed diagnoses		All-listed procedures	
	a	b	a	b	a	b	a	b
Total	0.00158	379.947	0.00269	1124.056	0.00164	395.929	0.00242	333.960
Sex								
Male	0.00174	346.105	0.00324	1430.558	0.00179	377.598	0.00296	332.448
Female	0.00157	384.179	0.00917	1881.266	0.00165	405.835	0.00243	331.119
Age								
Under 15 years	0.02110	241.964	0.03148	585.935	0.02265	265.264	0.03370	224.177
15–44 years	0.00194	333.180	0.00335	928.419	0.00202	326.273	0.00266	313.905
45–64 years	0.00167	328.374	0.00296	1238.451	0.00178	311.460	0.00293	278.111
65 years and over	0.00181	351.120	0.00325	1753.537	0.00180	403.465	0.00314	280.950
Region								
Northeast	0.00632	181.435	0.01100	492.047	0.01270	248.829	0.01206	184.247
Midwest	0.01073	270.823	0.01411	757.568	0.01347	210.823	0.02523	320.157
South	0.00272	315.316	0.00452	1125.453	0.00381	312.859	0.00449	286.033
West	0.00450	406.684	0.00828	1336.183	0.00538	498.904	0.00686	408.739
Principal expected source of payment								
Medicare	0.00189	359.897	0.00345	1707.735	0.00194	400.414	0.00356	250.559
Medicaid	0.00718	320.939	0.01069	923.765	0.00604	331.594	0.00965	237.653
Worker's compensation and other government payment	0.00993	325.814	0.01616	1272.519	0.01227	325.761	0.01167	325.796
HMO/PPO <sup>1</sup>	0.00498	242.924	0.00702	623.505	0.00431	284.866	0.00612	255.188
Blue Cross/Blue Shield and other private insurance	0.00394	306.593	0.00615	904.019	0.00446	347.031	0.00593	295.779
Self pay	0.00313	304.686	0.00706	932.033	0.00426	296.254	0.00721	220.949
No charge and other	0.04132	185.401	0.04812	661.567	0.03535	203.770	0.04466	201.837

<sup>1</sup>Health maintenance organization and preferred provider organization.

NOTE: The relative standard error (RSE) can be determined from the equation  $RSE(X) = \sqrt{a + b/X}$ .

is the denominator of the percent, and  $b$  is the parameter  $b$  in the formula for approximating the  $RSE(X)$ . The values for  $b$  are given in table II. When multiplied by 100,  $RSE(p)$  is expressed as a percent of  $p$ . This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (22,23).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained by the method described above using figures from table II) if the denominator is an estimate produced by the U.S. Census Bureau for the total population or one or more of the age-sex groups of the total population.

The approximate RSE of a ratio ( $X/Y$ ) in which the numerator ( $X$ ) and the denominator ( $Y$ ) are both estimated from the same survey, but the numerator is not a subclass of the denominator, is calculated using the formula:

$$RSE(X/Y) = \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

When multiplied by 100, the  $RSE(X/Y)$  is expressed as a percent of the ratio  $X/Y$ . This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (22,23).

## Definitions of Selected Terms

**Admission source**—The source of admission variable was added to the NHDS beginning in 2001. It was modeled after the variable contained in the National Uniform Bill (UB-92), the standard billing form used by Medicare for reimbursement of hospital inpatient care. The categories for admission source shown in this report are as follows:

**Physician referral**—The patient was admitted to the hospital upon the recommendation of his/her personal

physician, the hospital's clinic physician, or a health maintenance organization physician.

**Emergency department**—The patient was admitted to the hospital upon the recommendation of the hospital's emergency department physician. Emergency department and emergency room are used interchangeably in this report.

**Transfer from a hospital**—The patient was admitted to the hospital as a transfer from an acute care hospital where he/she was an inpatient.

**Transfer from other health facility**—The patient was admitted to the hospital as a transfer from a skilled nursing facility or other health facility where he/she was an inpatient.

**Other sources**—The patient was admitted to the hospital upon the direction of a court of law, on the request of a law enforcement

agency representative, or for other reasons not categorized above.

*Item not available*—Information about the source of admission to the hospital was not available for the patient.

*Admission type*—Like admission source, the admission type variable was added to the NHDS in 2001 and is modeled after the variable in the UB-92. Four admission types are shown in this report:

*Emergency*—The patient required immediate medical intervention as a result of severe life threatening or potentially disabling conditions. Patients in this group are generally admitted through the emergency room.

*Urgent*—The patient required immediate care and treatment of a physical or mental disorder. Patients in this group are usually admitted to the first available and suitable accommodation.

*Elective*—The patient's condition permitted adequate time to schedule the availability of suitable accommodation.

*Item not available*—Information about the admission type for the patient was not available.

*Age*—Patient's age at the birthday prior to admission to the hospital.

*Average length of stay*—The mean length of stay for discharges. It is calculated by dividing the number of days of care by the number of discharges.

*Barell injury diagnosis matrix*—A two-dimensional array that includes all the *International Classification of Diseases, 9th revision, Clinical Modification* (ICD-9-CM) codes describing injury (17). The code numbers from the matrix are shown in [table III](#). This matrix was developed to provide a standard format for describing the site and type of injuries for use in epidemiological and clinical analyses of injury data.

*Bed size of hospital*—The number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed

for use) for patients, not including bassinets for newborn infants. In this report, the classification of hospitals by bed size is based on the number of beds listed in the SMG Hospital Market Database (14).

*Days of care*—The total number of days a patient spent in the hospital. A stay of less than 1 day (patient admitted as an inpatient and discharged on the same day) is counted as one day. For patients admitted and discharged on different days, the number of days of care is computed by counting all the days from (and including) the day of admission to (but not including) the day of discharge.

*Death rate*—Calculated by dividing the number of deaths for a category by the total number of discharges in that category and multiplying by 100. Discharges whose status is unknown (1.0 percent) are assumed to be discharged alive.

*Diagnosis*—A disease or injury (or other factor that influences a patient's health status or causes contact with health services) that is listed on the medical record of the patient at discharge. The diagnosis established after study to be chiefly responsible for the admission of the patient to the hospital is the principal diagnosis. The first-listed diagnosis is the one specified as the principal diagnosis on the face sheet or discharge summary of the medical record, or if the principal diagnosis is not specified, the first one listed on the face sheet or discharge summary of the medical record. It is usually the main cause of the hospitalization. The number of first-listed diagnoses is the same as the number of discharges. Secondary diagnoses are all the diagnoses listed except the first-listed one and generally refer to conditions that did not cause the hospitalization, but may complicate the treatment of the first-listed diagnosis. All-listed diagnoses include first-listed diagnoses and secondary diagnoses and are included for users who want to know the total number of times conditions have been reported for hospitalized patients. A minimum of one and a maximum of seven diagnoses are coded per discharge in the NHDS. The HIV data ([table 23](#)) are for discharges

with an HIV diagnosis listed in any position among the coded diagnoses.

*Discharge*—A completed inpatient hospitalization. A hospitalization may be completed by death or by releasing the patient to the customary place of residence, a nursing home, another hospital, or other locations.

*Hospital*—Hospitals with an average length of stay for all inpatients of less than 30 days (short stay) or those whose specialty is general (medical or surgical) or children's general. Federal hospitals, hospital units of institutions, and hospitals with fewer than six beds staffed for patient use are excluded.

*Ischemic heart disease*—Pathological condition in which there is a lack of oxygen supply to the heart (ICD-9-CM codes 410–414). Also called coronary artery disease, ischemic heart disease includes conditions such as acute myocardial infarction, angina, and coronary atherosclerosis. Ischemic heart disease does not include heart conditions such as cardiac dysrhythmia (abnormal heart rhythm) or congestive heart failure.

*Newborn infant*—Inpatients admitted to a hospital by birth. Sick newborn infants are those that had at least one of the ICD-9-CM (1) diagnostic codes 001–999.

*Ownership of hospital*—The type of ownership is determined by the organization that controls and operates the hospital. Hospitals are grouped as follows:

*Proprietary*—Hospitals operated by individuals, partnerships, or corporations for profit.

*Government*—Hospitals operated by State or local governments.

*Nonprofit*—Hospitals operated by a nonprofit organization or church.

*Principal expected source of payment*—The primary expected method of payment for the hospitalization. This is classified as reported on the patient's medical record. Types of payment are as follows:

*Medicare*—The health insurance program for the aged and disabled administered by the Centers for Medicare and Medicaid Services

**Table III. Code numbers for Barel injury diagnosis matrix categories in table 24**

[Code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification*]

Type of injury	All body sites	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
All types of injury . . .	800–908, 909.0–909.2, 909.4, 909.9, 910–994, 995.50–995.59, 995.80–995.85	800–801, 803, 804, 850–854, 950.1–950.3, 995.55	802, 807.5–807.6, 830, 848.0–848.2, 870–874, 900, 910, 918, 920–921, 925, 940–941, 947.0, 950.0, 950.9, 951, 953.0, 954.0, 957.0, 959.0	806, 952	805, 839.0–839.5, 847.0–847.4	807.0–807.4, 808–809, 839.6–839.7, 846, 847.9, 848.3–848.5, 860–868, 875–878, 879.0–879.7, 901, 902.0–902.5, 902.81–902.82, 911, 922, 926, 942, 947.3–947.4, 953.1–953.3, 953.5, 954.1–954.9, 959.1	810–818, 831–834, 840–842, 880–887, 903, 912–915, 923, 927, 943–944, 953.4, 955, 959.2–959.5	820–827, 835–838, 843–845, 890–897, 904.0–904.8, 916, 917, 924.0–924.5, 928, 945, 959.6–959.7	819, 828–829, 839.8–839.9, 848.8–848.9, 869, 879.8–879.9, 902.87, 902.89, 902.9, 904.9, 919, 924.8–924.9, 929, 946, 947.1–947.2, 947.8–947.9, 948, 949, 953.8–953.9, 956, 957.1–957.9, 959.8–959.9
Fracture . . . . .	800–829	800–801, 803, 804	802, 807.5–807.6	806	805	807.0–807.4, 808, 809	810–818	820–827	819, 828–829
Dislocation . . . . .	830–839	None	830	None	839.0–839.5	839.6–839.7	831–834	835–838	839.8–839.9
Sprains and strains . .	840–848	None	848.0–848.2	None	847.0–847.4	846, 847.9, 848.3–848.5	840–842	843–845	848.8–848.9
Internal . . . . .	850–854, 860–869, 952, 995.55	850–854, 995.55	None	952	None	860–868	None	None	869
Open wound . . . . .	870–884, 890–894	None	870–874	None	None	875–878, 879.0–879.7	880–884	890–894	879.8–879.9
Amputation . . . . .	885–887, 895–897	None	None	None	None	None	885–887	895–897	None
Blood vessels . . . . .	900–904	None	900	None	None	901, 902.0–902.5, 902.81–902.82	903	904.0–904.8	902.87, 902.89, 902.9, 904.9
Contusions and superficial injuries . . .	910–924	None	910, 918, 920–921	None	None	911, 922	912–915, 923	916–917, 924.0–924.5	919, 924.8–924.9
Crush . . . . .	925–929	None	925	None	None	926	927	928	929
Burns . . . . .	940–949	None	940–941, 947.0	None	None	942, 947.3–947.4	943–944	945	946, 947.1–947.2, 947.8–947.9, 948–949
Nerves and spinal cord . . . . .	950–951, 953–957	950.1–950.3	950.0, 950.9, 951, 953.0, 954.0, 957.0	None	None	953.1–953.3, 953.5, 954.1–954.9	953.4, 955	None	953.8–953.9, 956, 957.1–957.9
Unspecified . . . . .	959	None	959.0	None	None	959.1	959.2–959.5	959.6–959.7	959.8–959.9
System-wide . . . . .	905–908, 909.0–909.2, 909.4, 909.9, 930–939, 958, 960–994, 995.50–995.54, 995.59, 995.80–995.85								

SOURCE: <http://www.cdc.gov/nchs/about/otheract/ice/barellmatrix.htm>.

(formerly the Health Care Financing Administration). A person 65 years of age and over is included in this category only if Medicare is reported as the principal expected source of payment.

**Medicaid**—A jointly funded Federal-State health insurance program providing medical care to those unable to afford it. If Medicaid and HMO/PPO are reported, the discharge is included in the category listed first (i.e., the principal expected source of payment).

**Worker’s compensation**—A State or municipal disability insurance or industrial accident insurance.

**Other government**—This includes payments under the Title V Program, the State-funded Maternal and Child Health Program that provides coverage for children up to 21 years of age with specific chronic medical conditions (e.g., Crippled Children’s Programs), CHAMPUS (Civilian Health and Medical Program of the Uniformed Services), or TRICARE (the Defense Department’s health care program for service families), casualty insurance paid by the State, vocational rehabilitation, Federal or State research grant (medical

research), or legal hold (prisoner in medical detention).

**HMO/PPO**—Any health maintenance organization (HMO) or preferred provider organization (PPO) sponsored by consumers, insurers, organizations, physicians, or hospitals. Medicare and Medicaid discharges are included in this category when HMO/PPO is reported as their principal expected source of payment.

**Blue Cross/Blue Shield and other private**—A private insurance plan not specified as an HMO/PPO. This includes Blue Cross/Blue Shield plans, medical coverage provided by life insurance companies, casualty insurance companies, health insurance companies, and independent plans such as employer/union sponsored plans and/or self-funded plans (partial or total).

**Self-pay**—The majority of the costs for the hospitalization were expected to be paid by the patient, spouse, family, or next of kin.

**No charge**—Patients admitted with the understanding that the medical services would be free. This includes hospital-sponsored special research or teaching patients.

**Other**—Includes any nonprofit source of payment, such as church welfare, United Way (United Appeal), or Shriner’s Hospitals for Children.

**Procedure**—Surgical procedures (such as appendectomy), diagnostic procedures (such as spinal tap), and therapeutic treatments (such as injection or infusion of cancer chemotherapeutic substance) reported on a patient’s medical record. Data are for all-listed procedures, which include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded per discharge in the NHDS. The procedures classified as nonsurgical in this report are listed in [table IV](#) by ICD-9-CM code numbers. All other procedures were considered surgical. This classification was begun in 1979 for the presentation of NHDS data and still remains an important distinction. However, the development of minimally invasive and noninvasive procedures has resulted in less difference between surgical and nonsurgical procedures with regard to operative or anesthetic risk and requirements for highly trained personnel or special equipment.

**Region**—Hospitals are classified by location in one of the four geographic regions of the United States that

**Table IV. Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD-9-CM category, 2002**

[Procedure categories and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Nonsurgical codes
Operations on the nervous system . . . . .	01-05 01.18-01.19,03.31,03.39,04.19,05.19
Operations on the endocrine system . . . . .	06-07 06.19, 07.19
Operations on the eye . . . . .	08-16 08.19,09.19,09.41-09.49,10.29,11.29, 12.29,14.19,15.09,16.21,16.29
Operations on the ear . . . . .	18-20 18.01,18.11,18.19,20.31,20.39
Operations on the nose, mouth, and pharynx . . . . .	21-29 21.00-21.02,21.21,21.29,22.19,24.19, 25.09,26.19,27.29,28.19,29.11,29.19
Operations on the respiratory system . . . . .	30-34 31.41-31.42,31.48-31.49,33.21-33.23, 33.29,34.21-34.22,34.28-34.29
Operations on the cardiovascular system . . . . .	35-39 37.26-37.28,37.29,38.29,39.95
Operations on the hemic and lymphatic system . . . . .	40-41 40.19,41.38-41.39
Operations on the digestive system . . . . .	42-54 42.22-42.23,42.29,44.11-44.13,44.19,44.32,45.11-45.13,45.19,45.21-45.24, 45.28-45.29, 48.21-48.23,48.29,49.21, 49.29,50.19,51.10-51.11,51.19,52.19,54.21,54.29
Operations on the urinary system . . . . .	55-59 55.21-55.22,55.29,56.31,56.35,56.39, 57.31-57.32,57.39,57.94-57.95, 58.21-58.22,58.29,59.29
Operations on the male genital organs . . . . .	60-64 60.18-60.19,61.19,62.19,63.09,64.19, 64.94
Operations on the female genital organs . . . . .	65-71 65.19,66.19,67.19,68.11,68.19,69.92,70.21-70.22,70.29,71.19
Obstetrical procedures . . . . .	72-75 73.4-73.59,73.91-73.92,75.31-75.32,75.34-75.35,75.38,75.94
Operations on the musculoskeletal system . . . . .	76-84 76.19,78.80-78.89,80.20-80.29,81.98, 83.29,84.41-84.43,84.45-84.47
Operations on the integumentary system . . . . .	85-86 85.19,86.19,86.92
Miscellaneous diagnostic and therapeutic procedures . . . . .	87-99 87.01-99.99

correspond to those used by the U.S. Census Bureau.

<i>Region</i>	<i>States included</i>
Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania
Midwest	Michigan, Ohio, Illinois, Indiana, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
South	Delaware, Maryland, District of Columbia, Virginia, West Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Mississippi, Arkansas, Louisiana, Oklahoma, and Texas
West	Montana, Idaho, Wyoming, Colorado, New Mexico, Arizona, Utah, Nevada, Washington, Oregon, California, Hawaii, and Alaska

data using code changes introduced in the October 1, 2002, addendum, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

Tables V and VI show the new ICD-9-CM codes from the 1986–2001 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned before development of the new code. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1).

## Changes in ICD-9-CM Codes

Although the ICD-9-CM has been used for coding NHDS data since 1979, the classification system undergoes annual updating. Assignment of new diagnostic and procedure codes, fourth- and fifth-digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Centers for Medicare and Medicaid Services (formerly Health Care Financing Administration). Addenda to the ICD-9-CM have become effective October 1 of each year since 1986, except in 1999. No addendum was released in 1999 because of concerns about instituting coding changes just before the millennium crossover.

The data in this report were coded using the addenda for October 1, 1986–2001. When hospitals reported



**Table V. Changes in diagnosis codes**

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81	1995	005.8	204.90	1991	204.9
005.89	1995	005.8	204.91	1991	V10.61
007.4	1997	007.8	205.00	1991	205.0
007.5	2000	007.8	205.01	1991	V10.62
008.00-008.09	1992	008.0	205.10	1991	205.1
008.43-008.47	1992	008.49	205.11	1991	V10.62
008.61-008.69	1992	008.6	205.20	1991	205.2
031.2	1997	031.8	205.21	1991	V10.62
038.10-038.11	1997	038.1	205.30	1991	205.3
038.19	1997	038.1	205.31	1991	V10.62
041.00-041.03	1992	041.0	205.80	1991	205.8
041.04	1992	041.0	205.81	1991	V10.62
041.04 (code title restated)	1997	041.04	205.90	1991	205.9
041.05	1992	041.0	205.91	1991	V10.62
041.09	1992	041.0	206.00	1991	206.0
041.10-041.19	1992	041.1	206.01	1991	V10.63
041.81-041.85	1992	041.8	206.10	1991	206.1
041.86	1995	041.84	206.11	1991	V10.63
041.89	1992	041.8	206.20	1991	206.2
042	1994	042.0-042.2, 042.9, 043.0-043.3, 043.9, 044.0, 044.9 (Codes deleted)	206.21	1991	V10.63
042.0-042.9	1986	279.19	206.80	1991	206.8
043.0-043.9	1986	279.19	206.81	1991	V10.63
044.0-044.9	1986	279.19	206.90	1991	206.9
070.20-070.21	1991	070.2	206.91	1991	V10.63
070.22	1994	070.20	207.00	1991	207.0
070.23	1994	070.21	207.01	1991	V10.69
070.30-070.31	1991	070.3	207.10	1991	207.1
070.32	1994	070.30	207.11	1991	V10.69
070.33	1994	070.31	207.20	1991	207.2
070.41-070.43	1991	070.4	207.21	1991	V10.69
070.44	1994	070.41	207.80	1991	207.8
070.49	1991	070.4	207.81	1991	V10.69
070.51-070.53	1991	070.5	208.00	1991	208.0
070.54	1994	070.51	208.01	1991	V10.60
070.59	1991	070.5	208.10	1991	208.1
077.98-077.99	1993	077.9	208.11	1991	V10.60
078.10-078.11,078.19	1993	078.1	208.20	1991	208.2
078.88	1993	078.89	208.21	1991	V10.60
079.4	1993	079.8	208.80	1991	208.8
079.50-079.53,079.59	1993	079.8	208.81	1991	V10.60
079.6	1996	079.89	208.90	1991	208.9
079.81	1995	079.89	208.91	1991	V10.60
079.88-079.89	1993	079.8	237.70-237.72	1990	237.7
079.98-079.99	1993	079.9	250.02	1993	250.90
082.40-082.41, 082.49	2000	082.8	250.03	1993	250.91
088.81	1989	088.8	250.12	1993	250.10
088.82	1993	088.89	250.13	1993	250.11
088.89	1989	088.8	250.22	1993	250.20
099.40-099.49	1992	099.4	250.23	1993	250.21
099.50-099.59	1992	078.89	250.32	1993	250.30
112.84-112.85	1992	112.89	250.33	1993	250.31
114.4-114.5	1993	114.3	250.42	1993	250.40
176.0-176.9	1991	173.0-173.9	250.43	1993	250.41
203.00	1991	203.0	250.52	1993	250.50
203.01	1991	V10.79	250.53	1993	250.51
203.10	1991	203.1	250.62	1993	250.60
203.11	1991	V10.79	250.63	1993	250.61
203.80	1991	203.8	250.72	1993	250.70
203.81	1991	V10.79	250.73	1993	250.71
204.00	1991	204.0	250.82	1993	250.80
204.01	1991	V10.61	250.83	1993	250.81
204.10	1991	204.1	250.92	1993	250.90
204.11	1991	V10.61	250.93	1993	250.91
204.20	1991	204.2	256.31	2001	256.3
204.21	1991	V10.61	275.40-275.42	1997	275.4
204.80	1991	204.8	275.49	1997	275.4
204.81	1991	V10.61	277.7	2001	277.8
			278.00-278.01	1995	278.0
			283.10-283.11,283.19	1993	283.1

Table V. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
285.21–285.22,285.29 . . . . .	2000	285.8	433.20–433.21 . . . . .	1993	433.2
291.81 . . . . .	1996	291.8	433.30–433.31 . . . . .	1993	433.3
291.89 . . . . .	1996	291.8	433.80–433.81 . . . . .	1993	433.8
293.84 . . . . .	1996	293.89	433.90–433.91 . . . . .	1993	433.9
294.10–294.11 . . . . .	2000	294.1	434.00–434.01 . . . . .	1993	434.0
300.82 . . . . .	1996	300.81	434.10–434.11 . . . . .	1993	434.1
305.1 . . . . .	1994	305.10, 305.11, 305.12, 305.13 (Codes deleted)	434.90–434.91 . . . . .	1993	434.9
312.81–312.82,312.89 . . . . .	1994	312.8	435.3 . . . . .	1995	435.0 & 435.1
315.32 . . . . .	1996	315.39	437.7 . . . . .	1992	780.9
320.81–320.89 . . . . .	1992	320.8	438.0 . . . . .	1997	294.9 & 438
333.92–333.93 . . . . .	1994	333.99	438.10 . . . . .	1997	784.5 & 438
337.20–337.22,337.29 . . . . .	1993	337.9	438.11 . . . . .	1997	784.3 & 438
337.3 . . . . .	1998	337.9	438.12 . . . . .	1997	784.4 & 438
342.00–342.02 . . . . .	1994	342.0	438.19 . . . . .	1997	784.5 & 438
342.10–342.12 . . . . .	1994	342.1	438.20 . . . . .	1997	342.90 & 438
342.80–342.82 . . . . .	1994	342.9	438.21 . . . . .	1997	342.91 & 438
342.90–342.92 . . . . .	1994	342.9	438.22 . . . . .	1997	342.92 & 438
344.00–344.04,344.09 . . . . .	1994	344.0	438.30 . . . . .	1997	344.40 & 438
344.30–344.32 . . . . .	1994	344.3	438.31 . . . . .	1997	344.41 & 438
344.40–344.42 . . . . .	1994	344.4	438.32 . . . . .	1997	344.42 & 438
344.81,344.89 . . . . .	1993	344.8	438.40 . . . . .	1997	344.30 & 438
345.00–345.01 . . . . .	1989	345.0	438.41 . . . . .	1997	344.31 & 438
345.10–345.11 . . . . .	1989	345.1	438.42 . . . . .	1997	344.32 & 438
345.40–345.41 . . . . .	1989	345.4	438.50–438.52 . . . . .	1997	344.89 & 438
345.50–345.51 . . . . .	1989	345.5	438.53 . . . . .	1998	438.50
345.60–345.61 . . . . .	1989	345.6	438.81 . . . . .	1997	784.69 & 438
345.70–345.71 . . . . .	1989	345.7	438.82 . . . . .	1997	787.2 & 438
345.80–345.81 . . . . .	1989	345.8	438.89 . . . . .	1997	438
345.90–345.91 . . . . .	1989	345.9	438.9 . . . . .	1997	438
346.00–346.01 . . . . .	1992	346.0	440.20–440.22 . . . . .	1992	440.2
346.10–346.11 . . . . .	1992	346.1	440.23 . . . . .	1993	440.20 & 707.1 or 707.8 or 707.9
346.20–346.21 . . . . .	1992	346.2	440.24 . . . . .	1993	440.20 & 785.4
346.80–346.81 . . . . .	1992	346.8	440.29 . . . . .	1993	440.2
346.90–346.91 . . . . .	1992	346.9	440.30–440.32 . . . . .	1994	996.1
355.71 . . . . .	1993	354.4	441.00–441.03 . . . . .	1994	441.0
355.79 . . . . .	1993	355.7	441.6 . . . . .	1993	441.1 & 441.3
371.82 . . . . .	1992	371.89	441.7 . . . . .	1993	441.2 & 441.4
372.81,372.89 . . . . .	2000	372.8	446.20–446.21,446.29 . . . . .	1990	446.2
374.87 . . . . .	1990	374.89	451.82–451.84 . . . . .	1993	451.89
403.00–403.01 . . . . .	1989	403.0	458.2 . . . . .	1995	997.9 & 458.9
403.10–403.11 . . . . .	1989	403.1	458.8 . . . . .	1997	458.9
403.90–403.91 . . . . .	1989	403.9	464.00–464.01 . . . . .	2001	464.0
404.00–404.03 . . . . .	1989	404.0	464.50–464.51 . . . . .	2001	464.0
404.10–404.13 . . . . .	1989	404.1	466.11 . . . . .	1996	466.1
404.90–404.93 . . . . .	1989	404.9	466.19 . . . . .	1996	466.1
410.00–410.02 . . . . .	1989	410.0	474.0 (Code title restated) . . . . .	1997	474.0
410.10–410.12 . . . . .	1989	410.1	474.00–474.02 . . . . .	1997	474.0
410.20–410.22 . . . . .	1989	410.2	477.1 . . . . .	2000	477.8
410.30–410.32 . . . . .	1989	410.3	482.30–482.39 . . . . .	1992	482.3
410.40–410.42 . . . . .	1989	410.4	482.40 . . . . .	1998	482.4
410.50–410.52 . . . . .	1989	410.5	482.41 . . . . .	1998	482.4
410.60–410.62 . . . . .	1989	410.6	482.49 . . . . .	1998	482.4
410.70–410.72 . . . . .	1989	410.7	482.81–482.83 . . . . .	1992	482.8
410.80–410.82 . . . . .	1989	410.8	482.84 . . . . .	1997	482.83
410.90–410.92 . . . . .	1989	410.9	482.89 . . . . .	1992	482.8
411.81 . . . . .	1989	410.9	483.0 . . . . .	1992	483
411.89 . . . . .	1989	411.8	483.1 . . . . .	1996	078.88 & 484.8
414.00–414.01 . . . . .	1994	414.0	483.8 . . . . .	1992	483
414.02–414.03 . . . . .	1994	996.03	491.20–491.21 . . . . .	1991	491.2
414.04–414.05 . . . . .	1996	414.00	493.20 . . . . .	1989	493.90
415.11 . . . . .	1995	997.3 & 415.1	493.21 . . . . .	1989	493.91
415.19 . . . . .	1995	415.1	493.02 . . . . .	2000	493.00
429.71 . . . . .	1989	410.0–410.9	493.12 . . . . .	2000	493.10
429.79 . . . . .	1989	410.0–410.9	493.22 . . . . .	2000	493.20
433.00–433.01 . . . . .	1993	433.0	493.92 . . . . .	2000	493.90
433.10–433.11 . . . . .	1993	433.1	494.0–494.1 . . . . .	2000	494
			512.1 . . . . .	1994	997.3



**Table V. Changes in diagnosis codes—Con.**

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
518.6 . . . . .	1997	518.89	651.30–651.31,651.33 . . . . .	1989	651.00–651.01, 651.03
518.81 . . . . .	1987	799.1	651.40–651.41,651.43 . . . . .	1989	651.10–651.11, 651.13
518.82–518.89 . . . . .	1987	518.8	651.50–651.51,651.53 . . . . .	1989	651.20–651.21, 651.23
518.83 . . . . .	1998	518.81	651.60–651.61,651.63 . . . . .	1989	651.80–651.81, 651.83
518.84 . . . . .	1998	518.81	654.20–654.21,654.23 . . . . .	1990	654.2,654.9
519.00–519.02, 519.09 . . . . .	1998	519.0	654.90–654.94 . . . . .	1990	654.2,654.9
521.00–521.09 . . . . .	2001	521.0	655.70&655.71 . . . . .	1997	655.8
524.00–524.09 . . . . .	1992	524.0	655.73 . . . . .	1997	655.8
524.10–524.19 . . . . .	1992	524.1	657.0 . . . . .	1991	657
524.60–524.69 . . . . .	1991	524.6	659.60,659.61,659.63 . . . . .	1992	659.80–659.81, 659.83
524.70–524.79 . . . . .	1992	524.8	659.70,659.71,659.73 . . . . .	1998	656.30, 656.31, 656.33
525.10–525.19 . . . . .	2001	525.1	665.10,665.11 . . . . .	1992	665.10,665.11,665.12, 665.14 <sup>1</sup>
530.10–530.11, 530.19 . . . . .	1993	530.1	670.0 . . . . .	1991	670
530.12 . . . . .	2001	530.10	672.0 . . . . .	1991	672
530.81 . . . . .	1993	530.1	677 . . . . .	1994	None
530.82–530.84, 530.89 . . . . .	1993	530.8	686.00 & 686.01 . . . . .	1997	686.0
535.00–535.01 . . . . .	1991	535.0	686.09 . . . . .	1997	686.0
535.10–535.11 . . . . .	1991	535.1	690.10 . . . . .	1995	690
535.20–535.21 . . . . .	1991	535.2	690.11 . . . . .	1995	691.8 & 704.8
535.30–535.31 . . . . .	1991	535.3	690.12 . . . . .	1995	691.8
535.40–535.41 . . . . .	1991	535.4	690.18 . . . . .	1995	690
535.50–535.51 . . . . .	1991	535.5	690.8 . . . . .	1995	690
535.60–535.61 . . . . .	1991	535.6	692.72–692.74 . . . . .	1992	692.79
536.3 . . . . .	1994	536.8	692.75 . . . . .	2000	692.79
536.40–536.42,536.49 . . . . .	1998	997.4	692.76–692.77 . . . . .	2001	692.71
537.82 . . . . .	1990	537.89	692.82–692.83 . . . . .	1992	692.89
537.83 . . . . .	1991	537.82	702.0–702.8 . . . . .	1991	702
558.3 . . . . .	2000	558.9	702.11,702.19 . . . . .	1994	702.1
556.0–556.6 . . . . .	1994	556	704.02 . . . . .	1993	704.09
556.8–556.9 . . . . .	1994	556	707.10–707.15, 707.19 . . . . .	2000	707.1
562.02 . . . . .	1991	562.00	709.00–709.01,709.09 . . . . .	1994	709.0
562.03 . . . . .	1991	562.01	710.5 . . . . .	1992	288.3,729.1
562.12 . . . . .	1991	562.10	718.70–718.79 . . . . .	2001	718.80–718.89
562.13 . . . . .	1991	562.11	727.83 . . . . .	2000	727.89
564.00–564.09 . . . . .	2001	564.0	728.86 . . . . .	1995	729.4
564.81 . . . . .	1998	564.8	733.10–733.16,733.19 . . . . .	1993	733.1
564.89 . . . . .	1998	564.8	733.93 . . . . .	2001	733.16
569.60–569.61 . . . . .	1995	569.6	733.94 . . . . .	2001	733.19
569.62 . . . . .	1998	569.69	733.95 . . . . .	2001	733.19
569.69 . . . . .	1995	569.6	738.10–738.19 . . . . .	1992	738.1
569.84 . . . . .	1990	557.1	747.60–747.64,747.69 . . . . .	1993	747.6
569.85 . . . . .	1991	569.84	747.82 . . . . .	1993	747.89
574.60 . . . . .	1996	574.00 & 574.30	752.51–752.52 . . . . .	1996	752.5
574.61 . . . . .	1996	574.01 & 574.31	752.61–752.63 . . . . .	1996	752.6
574.70 . . . . .	1996	574.10 & 574.40	752.64–752.65 . . . . .	1996	752.8
574.71 . . . . .	1996	574.11 & 574.41	752.69 . . . . .	1996	752.8
574.80 . . . . .	1996	574.00 & 574.10 574.30 & 574.40	753.10–753.17,753.19 . . . . .	1990	753.1
574.81 . . . . .	1996	574.01 & 574.11 574.31 & 574.41	753.20–753.23 . . . . .	1996	753.2
574.90 . . . . .	1996	574.20 & 574.50	753.29 . . . . .	1996	753.2
574.91 . . . . .	1996	574.21 & 574.51	756.70–756.71 . . . . .	1997	756.7
575.10–575.11 . . . . .	1996	575.1	756.79 . . . . .	1997	756.7
575.12 . . . . .	1996	575.0 & 575.1	758.81 . . . . .	1996	758.8
593.70–593.73 . . . . .	1994	593.7	758.89 . . . . .	1996	758.9
596.51–596.53 . . . . .	1992	596.5	759.81–759.82 . . . . .	1989	759.8
596.54 . . . . .	1992	344.61	759.83 . . . . .	1994	759.89
596.55–596.59 . . . . .	1992	596.5	759.89 . . . . .	1989	759.8
599.81–599.89 . . . . .	1992	599.8	760.75 . . . . .	1991	760.79
600.0–600.3, 600.9 . . . . .	2000	600	760.76 . . . . .	1994	760.79
602.3 . . . . .	2001	602.8	763.81–763.83,763.89 . . . . .	1998	763.8
608.82 . . . . .	2001	608.83	764.00–764.09 . . . . .	1988	764.0
608.87 . . . . .	2001	608.89	764.10–764.19 . . . . .	1988	764.1
645.0 . . . . .	1991	645	764.20–764.29 . . . . .	1988	764.2
645.10–645.11, 645.13 . . . . .	2000	645.00–645.01, 645.03			
645.20–645.21, 645.23 . . . . .	2000	645.00–645.01, 645.03			

See footnote at end of table.

Table V. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
764.90–764.99	1988	764.9	996.04	1994	996.09
765.00–765.09	1988	765.0	996.51–996.59	1987	996.5
765.10–765.19	1988	765.1	996.55	1998	996.52
772.10–772.14	2001	772.1	996.56	1998	996.59
779.7	2001	772.1	996.60–996.69	1989	996.6
780.01–780.02	1992	780.0	996.68	1998	996.69
780.03	1993	780.01	996.70–996.79	1989	996.7
780.09	1992	780.0	996.80–996.84	1987	996.8
780.31	1997	780.3	996.85	1990	996.8
780.39	1997	780.3	996.86	1987	996.8
780.57	1992	780.51,780.53	996.87	2000	996.89
780.71	1998	780.7	996.89	1987	996.8
780.79	1998	780.7	997.00–997.01	1995	997.0
781.8	1994	781.9	997.02	1995	997.9 & 430–434, 436
781.91–781.92	2000	781.9			
781.99	2000	781.9	997.09	1995	997.0
783.21	2000	783.2	997.71	2001	997.4
783.22	2000	783.4	997.72	2001	997.5
783.40–783.43	2000	783.4	997.79	2001	997.2
783.7	2000	783.4	997.91	1995	997.9
786.03	1998	786.09	997.99	1995	997.9
786.04	1998	786.09	998.11–998.12	1996	998.1
786.05	1998	786.09	998.13	1996	998.89
786.06	1998	786.09	998.51	1996	998.5
786.07	1998	786.09	998.59	1996	998.5
787.01–787.03	1994	787.0	998.81–998.82,998.89	1994	998.8
787.91	1995	558.9	998.83	1996	998.89
787.99	1995	787.9	V02.51	1998	V02.5
788.20–788.21, 788.29	1993	788.2	V02.52	1998	V02.5
788.30–788.39	1992	788.3	V02.59	1998	V02.5
788.41–788.43	1993	788.4	V02.60–V02.62	1997	V02.6
788.61–788.62, 788.69	1993	788.6	V02.69	1997	V02.6
789.00–789.07, 789.09	1994	789.0	V03.81–V03.82,V03.89	1994	V03.8
789.30–789.37, 789.39	1994	789.3	V05.3–V05.4	1993	V05.8
789.40–789.47, 789.49	1994	789.4	V06.5–V06.6	1994	V06.8
789.60–789.67, 789.69	1994	789.6	V07.31,V07.39	1994	V07.3
790.01, 790.09	2000	790.0	V07.4	1992	V07.8
790.91	1993	790.9	V08	1994	044.9,795.8 (Codes deleted)
790.92	1993	286.9	V09.0–V09.91	1993	None
790.93, 790.99	1993	790.9	V10.48	1998	V10.49
790.94	1997	790.99	V10.53	2001	V10.59
792.5	2000	792.9	V12.00–V12.03,V12.09	1994	V12.0
793.80–793.81, 793.89	2001	793.8	V12.40–V12.41	1997	V12.4
795.71	1994	795.8 (Code deleted)	V12.49	1997	V12.4
			V12.50–V12.52	1995	V12.5
795.79	1994	795.7	V12.59	1995	V12.5
795.8	1986	795.7	V12.70–V12.72,V12.79	1994	V12.7
796.5	1997	796.9	V13.00–V13.01,V13.09	1994	V13.0
840.7	2001	840.8	V13.61	1998	V13.6
864.05	1992	864.09	V13.69	1998	V13.6
864.15	1992	864.19	V15.01–V15.09	2000	V15.0
909.5	1994	909.9	V15.41–V15.42	1996	V15.4
922.31–922.33	1996	922.3	V15.49	1996	V15.4
925.1–925.2	1993	925	V15.82	1994	305.13 (Codes deleted)
959.0 (Code title restated)	1997	959.0	V15.84–V15.86	1995	V15.89
959.01	1997	854.00	V16.40–V16.43	1997	V16.4
959.09	1997	959.0	V16.49	1997	V16.4
965.61	1998	965.6	V16.51	1998	V16.5
965.69	1998	965.6	V16.59	1998	V16.5
989.81–989.84	1995	989.8	V18.61	1998	V18.6
989.89	1995	989.8	V18.69	1998	V18.6
995.50–995.55	1996	995.5	V21.30–V21.35	2000	None
995.59	1996	995.5	V23.7	1989	V23.8
995.60–995.69	1993	995.0	V23.81	1998	V23.8
995.7	2000	None	V23.82	1998	V23.8
995.80	1996	995.81	V23.83	1998	V23.8
995.81 (Code title restated)	1996	995.81			
995.82–995.85	1996	995.81			
995.86	1998	995.89			

Table V. Changes in diagnosis codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
V23.84 . . . . .	1998	V23.8	V56.31–V56.32 . . . . .	2000	V56.8
V23.89 . . . . .	1998	V23.8	V57.21–V57.22 . . . . .	1994	V57.2
V25.43 . . . . .	1992	V25.49	V58.41,V58.49 . . . . .	1994	V58.4
V25.5 . . . . .	1992	V25.8	V58.61 . . . . .	1995	V67.51
V26.21–V26.22,V26.29 . . . . .	2000	V26.2	V58.62 . . . . .	1998	V58.69
V26.51 . . . . .	1998	None	V58.69 . . . . .	1995	V67.51
V26.52 . . . . .	1998	None	V58.81,V58.89 . . . . .	1994	V58.8
V28.6 . . . . .	1997	V28.8	V58.82 . . . . .	1995	V58.89
V29.0–V29.1 . . . . .	1992	V71.8	V58.83 . . . . .	2000	V58.89
V29.2 . . . . .	1994	V29.8	V59.01–V59.02 . . . . .	1995	V59.0
V29.3 . . . . .	1998	V29.8	V59.09 . . . . .	1995	V59.0
V29.8 . . . . .	1992	V71.8	V59.6 . . . . .	1995	V59.8
V29.9 . . . . .	1992	V71.9	V61.10–V61.12 . . . . .	1996	V61.1
V30.00–V30.01 . . . . .	1989	V30.0	V61.22 . . . . .	1996	V61.21
V31.00–V31.01 . . . . .	1989	V31.0	V62.83 . . . . .	1996	V65.49
V32.00–V32.01 . . . . .	1989	V32.0	V64.4 . . . . .	1997	None
V33.00–V33.01 . . . . .	1989	V33.0	V65.40–V65.45,V65.49 . . . . .	1994	V65.4
V34.00–V34.01 . . . . .	1989	V34.0	V66.7 . . . . .	1996	None
V35.00–V35.01 . . . . .	1989	V35.0	V67.00–V67.01,V67.09 . . . . .	2000	V67.0
V36.00–V36.01 . . . . .	1989	V36.0	V69.0–V69.3 . . . . .	1994	None
V37.00–V37.01 . . . . .	1989	V37.0	V69.8–V69.9 . . . . .	1994	None
V39.00–V39.01 . . . . .	1989	V39.0	V71.81 . . . . .	2000	V71.8
V42.81–V82.83 . . . . .	1997	V42.8	V71.89 . . . . .	2000	V71.8
V42.84 . . . . .	2000	V42.89	V72.81–V72.85 . . . . .	1993	V72.8
V42.89 . . . . .	1997	V42.8	V73.88–V73.89 . . . . .	1993	V73.8
V43.60–V43.66,V43.69 . . . . .	1994	V43.6	V73.98–V73.99 . . . . .	1993	V73.9
V43.81–V43.82 . . . . .	1995	V43.8	V76.10–V76.12 . . . . .	1997	V76.1
V43.83 . . . . .	1998	V43.89	V76.19 . . . . .	1997	V76.1
V43.89 . . . . .	1995	V43.8	V76.44 . . . . .	1998	V76.49
V44.50 . . . . .	1998	V44.5	V76.45 . . . . .	1998	V76.49
V44.51 . . . . .	1998	V44.5	V76.46–V76.47 . . . . .	2000	V76.49
V44.52 . . . . .	1998	V44.5	V76.50–V76.52 . . . . .	2000	V76.49
V44.59 . . . . .	1998	V44.5	V76.81, V76.89 . . . . .	2000	V76.8
V45.00 . . . . .	1994	V45.89	V77.91, V77.99 . . . . .	2000	V77.9
V45.01 . . . . .	1994	V45.0	V82.81, V82.89 . . . . .	2000	V82.8
V45.02,V45.09 . . . . .	1994	V45.89	V83.01–V83.02 . . . . .	2001	None
V45.51 . . . . .	1994	V45.5	E854.8 . . . . .	1995	E858.8
V45.52,V45.59 . . . . .	1994	V45.89	E869.4 . . . . .	1994	E869.8
V45.61 . . . . .	1997	V45.6	E880.1 . . . . .	1995	E884.9
V45.63 . . . . .	1997	V45.6	E884.3–E884.4 . . . . .	1995	E884.2
V45.71 . . . . .	1997	611.8	E884.5–E884.6 . . . . .	1995	E884.9
V45.72 . . . . .	1997	569.89	E885.1–E885.4,E885.9 . . . . .	2000	E885
V45.73 . . . . .	1997	593.89	E888.0 . . . . .	2001	E920
V45.74 . . . . .	2000	593.89,596.8	E888.1 . . . . .	2001	E888
V45.75 . . . . .	2000	V45.89	E888.8–E888.9 . . . . .	2001	E888
V45.76 . . . . .	2000	518.89	E906.5 . . . . .	1995	E906.3
V45.77 . . . . .	2000	602.8,607.89, 608.89,620.8, 621.8,622.8	E908.0–E908.4 . . . . .	1995	E908
V45.78 . . . . .	2000	360.89	E908.8–E908.9 . . . . .	1995	E908
V45.79 . . . . .	2000	255.8,289.59, 388.8,569.49, 577.8,V45.89	E909.0–E909.4 . . . . .	1995	E909
V45.82 . . . . .	1994	V45.89	E909.8–E909.9 . . . . .	1995	E909
V45.83 . . . . .	1995	V45.89	E917.3 . . . . .	2001	E917.9
V45.84 . . . . .	2001	None	E917.4 . . . . .	2001	E917.9
V49.60–V49.67 . . . . .	1994	V49.5	E917.5 . . . . .	2001	E886.0
V49.70–V49.77 . . . . .	1994	V49.5	E917.6 . . . . .	2001	E917.1
V49.81 . . . . .	2000	None	E917.7 . . . . .	2001	E888
V49.82 . . . . .	2001	None	E917.8 . . . . .	2001	E888
V49.89 . . . . .	2000	V49.8	E920.5 . . . . .	1995	E920.4
V50.41–V50.42,V50.49 . . . . .	1994	V50.8	E922.4 . . . . .	1997	E917.9
V53.01–V53.02 . . . . .	1997	V53.0	E924.2 . . . . .	1995	E924.0
V53.09 . . . . .	1997	V53.0	E928.3 . . . . .	2000	E928.8
V53.31 . . . . .	1994	V53.3	E955.6 . . . . .	1997	E955.9
V53.32,V53.39 . . . . .	1994	V53.9	E967.2 . . . . .	1996	E967.0
V56.1 . . . . .	1995	V58.89	E967.3 . . . . .	1996	None
V56.1 (Code title restated) . . . . .	1998	V56.1	E967.4–E967.8 . . . . .	1996	E967.1
V56.2 . . . . .	1998	V56.1	E968.7 . . . . .	2000	E968.8
			E968.5 . . . . .	1995	E968.8
			E968.6 . . . . .	1997	E968.8
			E985.6 . . . . .	1997	E985.4

<sup>1</sup>The title for the subcategory 665.1 has been changed, making the fifth-digit subclassifications 665.12 and 665.14 invalid.

Table VI. Changes in procedure codes

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
02.96 . . . . .	1992	89.19	37.83 (Device only) . . . . .	1987	37.73–37.77 (Leads/Device)
03.90 . . . . .	1987	03.99 (Insertion of catheter)	37.85–37.87 . . . . .	1987	37.85
05.25 . . . . .	1995	39.7	37.89 . . . . .	1987	37.86 + 37.89
11.75 . . . . .	1989	11.79	37.94–37.98 . . . . .	1986	37.99
11.76 . . . . .	1989	11.62	38.22 . . . . .	1986	38.29
20.96–20.98 . . . . .	1986	20.95	38.44 (Abdominal aorta only) . . . . .	1986	38.44 (entire aorta)
22.12 . . . . .	1988	22.11	38.45 (Thoracic aorta added) . . . . .	1986	38.44–38.45
26.12 . . . . .	1988	26.11	38.95 . . . . .	1989	38.93
29.31 . . . . .	1991	83.02	39.28 . . . . .	1991	39.29
29.32 . . . . .	1991	29.3	39.50 . . . . .	1995	39.59
29.33 . . . . .	1991	29.3	39.65 . . . . .	1988	39.61
29.39 . . . . .	1991	29.3	39.66 . . . . .	1990	39.65
31.45 . . . . .	1988	31.43–31.44	39.71 . . . . .	2000	39.52
31.95 . . . . .	1989	31.75	39.79 . . . . .	2000	39.52
32.01 . . . . .	1989	32.0	39.90 . . . . .	1996	39.50
32.09 . . . . .	1989	32.0	41.00–41.03 . . . . .	1988	41.0
32.22 . . . . .	1995	32.29,32.9	41.04 . . . . .	1994	99.79
32.28 . . . . .	1989	32.29	41.05 . . . . .	1997	None
33.27 . . . . .	1987	33.22+33.27	41.06 . . . . .	1997	None
33.28 . . . . .	1987	33.27	41.07 . . . . .	2000	41.04
33.29 . . . . .	1987	33.28–33.29	41.08 . . . . .	2000	41.05
33.50 . . . . .	1995	33.5	41.09 . . . . .	2000	41.01
33.51 . . . . .	1995	33.5	42.25 . . . . .	1988	42.24
33.52 . . . . .	1995	33.5	42.33 . . . . .	1989	42.32,42.39
33.6 . . . . .	1990	33.5+37.5	42.33 . . . . .	1990	42.91
34.05 . . . . .	1994	34.99	43.11 . . . . .	1989	43.1
35.84 . . . . .	1988	35.82	43.19 . . . . .	1989	43.1,43.2
35.96 . . . . .	1986	35.03	43.41 . . . . .	1989	43.41,43.49
36.00–36.03 . . . . .	1986	36.0	44.21 . . . . .	1986	44.2
36.04 . . . . .	1986	39.97	44.22 . . . . .	1986	44.99
36.05 . . . . .	1987	36.01	44.29 . . . . .	1986	44.2
36.05 . . . . .	1986	36.01 <sup>1</sup> ,36.02	44.32 . . . . .	2001	44.39
36.06 . . . . .	1995	36.01,36.02,36.03,36.05	44.43 . . . . .	1989	43.49,45.32
36.09 . . . . .	1986	36.0	44.44 . . . . .	1989	38.86
36.09 . . . . .	1991	36.00 (Code deleted)	44.49 . . . . .	1989	43.0
36.17 . . . . .	1996	36.19	44.93–44.94 . . . . .	1986	44.99
36.31 . . . . .	1998	36.3	45.16 . . . . .	1988	45.14 (45.15 before 1987)
36.32 . . . . .	1998	36.3	45.30 . . . . .	1989	45.31,45.32
36.39 . . . . .	1998	36.3	45.42 . . . . .	1988	45.41
37.26–37.27 . . . . .	1988	37.29	45.43 . . . . .	1989	45.49
37.28 . . . . .	2001	88.72	45.75 (Hartmann resection added) . . . . .	1988	48.66 (Code deleted)
37.34 . . . . .	1988	37.33	45.95 . . . . .	1987	45.93
37.35 . . . . .	1997	37.33	46.13 . . . . .	1992	46.12 (Code deleted)
37.65 . . . . .	1995	37.62	46.32 . . . . .	1989	46.39
37.66 . . . . .	1995	37.62	46.85 . . . . .	1989	46.99
37.67 . . . . .	1998	37.4	46.97 . . . . .	2000	46.99
37.70 (Leads only) . . . . .	1987	37.70 (Leads/Device)	47.01 . . . . .	1996	47.0
37.71–37.72 (Leads only) . . . . .	1987	37.74 (Leads/Device)	47.09 . . . . .	1996	47.0
37.73 (Leads only) . . . . .	1987	37.73 (Leads/Device)	47.11 . . . . .	1996	47.1
37.74 (Leads only) . . . . .	1987	37.76 (Leads/Device)	47.19 . . . . .	1996	47.1
37.75 (Leads only) . . . . .	1987	37.89 (Leads/Device)	48.36 . . . . .	1995	45.42
37.76 (Leads only) . . . . .	1987	37.81 (Leads/Device)	49.31 . . . . .	1989	49.3
37.77 (Leads only) . . . . .	1987	37.83–37.84 (Leads/Device)	49.39 . . . . .	1989	49.3
37.78 . . . . .	1987	37.71–37.72	51.10 . . . . .	1989	51.97
37.79 . . . . .	1987	86.09	51.11 . . . . .	1989	51.11,51.97
37.80–37.87 . . . . .	1992	89.49 (Code deleted. This procedure is included in the code for pacemaker insertion/replacement.)	51.14 . . . . .	1989	51.12
37.80 (Device only) . . . . .	1987	37.73–37.77 (Leads/Device)	51.15 . . . . .	1989	51.97
37.81 (Device only) . . . . .	1987	37.73–37.77 (Leads/Device)	51.21 . . . . .	1996	51.22,51.23
37.82 (Device only) . . . . .	1987	37.73–37.77 (Leads/Device)	51.22 . . . . .	1991	51.21 (Code deleted), 51.22
			51.23 . . . . .	1991	51.22
			51.24 . . . . .	1996	51.22,51.23
			51.64 . . . . .	1989	51.69
			51.84–51.88 . . . . .	1989	51.97
			51.97 . . . . .	1986	52.91,51.99, or 51.82

See footnotes at end of table.

Table VI. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
51.98	1986	51.99	75.38	2001	75.34
52.13	1989	51.97,52.91	77.56	1989	77.89,78.49,81.18
52.14	1989	52.11	77.57	1989	77.89,80.48,81.18,83.85
52.21	1989	52.2	77.58	1989	77.59,81.18
52.22	1989	52.2	78.10	1991	78.40
52.84	1996	99.29	78.11	1991	78.41
52.85	1996	99.29	78.12	1991	78.42
52.86	1996	99.29	78.13	1991	78.43
52.93	1989	52.93 + 52.91	78.14	1991	78.44
52.94	1989	52.09	78.15	1991	78.45
52.97	1989	52.91	78.16	1991	78.46
52.98	1989	52.91	78.17	1991	78.47
52.99	1989	52.93,52.94,52.99	78.18	1991	78.48
54.24	1987	54.23	78.19	1991	78.49
54.25	1993	54.98	78.20	1991	78.10,78.20,78.30
54.51	1996	54.5	78.21	1991	78.11,78.31
54.59	1996	54.5	78.22	1991	78.12,78.22,78.32
55.03–55.04	1986	55.02	78.23	1991	78.13,78.23,78.33
56.33–56.34	1987	56.33	78.24	1991	78.14,78.34
56.35	1987	45.12	78.25	1991	78.15,78.25,78.35
57.17–57.18	1989	57.21	78.27	1991	78.17,78.27,78.37
57.22	1989	57.22,57.82	78.28	1991	78.18,78.38
58.31	1990	58.3	78.29	1991	78.11,78.16,78.19,78.29, 78.39
58.39	1990	58.3	78.39	1991	78.31
58.93	1986	57.99	78.90 <sup>2</sup>	1987	78.40
59.03	1996	59.02	78.91 <sup>2</sup>	1987	78.41
59.12	1996	59.11	78.92 <sup>2</sup>	1987	78.42
59.72	1995	59.79	78.93 <sup>2</sup>	1987	78.43
59.96	1986	59.95	78.94 <sup>2</sup>	1987	78.44
60.21	1995	60.2	78.95 <sup>2</sup>	1987	78.45
60.29	1995	60.2	78.96 <sup>2</sup>	1987	78.46
60.95	1991	60.99	78.97 <sup>2</sup>	1987	78.47
60.96	2000	60.29	78.98 <sup>2</sup>	1987	78.48
60.97	2000	60.29	78.99 <sup>2</sup>	1987	78.49
64.97	1986	64.95	80.50–80.59	1986	80.5
65.01	1996	65.0	81.03	1989	81.02
65.09	1996	65.0	81.04–81.05	1989	81.03,81.04,81.05
65.13	1996	65.12	81.06–81.07	1989	81.06,81.07
65.14	1996	65.19	81.08	1989	81.06,81.07,81.08
65.23	1996	65.21	81.09	1989	81.08
65.24	1996	65.22	81.30	2001	81.09
65.25	1996	65.29	81.31	2001	81.09
65.31	1996	65.3	81.32	2001	81.09
65.39	1996	65.3	81.33	2001	81.09
65.41	1996	65.4	81.34	2001	81.09
65.49	1996	65.4	81.35	2001	81.09
65.53	1996	65.51	81.36	2001	81.09
65.54	1996	65.52	81.37	2001	81.09
65.63	1996	65.61	81.38	2001	81.09
65.64	1996	65.62	81.39	2001	81.09
65.74	1996	65.71	81.40	1989	81.69
65.75	1996	65.72	81.51	1989	81.51,81.59
65.76	1996	65.73	81.52	1989	81.61,81.62,81.63,81.64
65.81	1996	65.8	81.53	1989	81.51,81.59,81.61,81.62, 81.63,81.64
65.89	1996	65.8	81.54–81.55	1989	81.41
66.01	1992	66.0	81.56	1989	81.48
66.02	1992	66.73	81.57	1989	81.31,81.39
67.51	2001	67.5	81.59	1989	81.39
67.59	2001	67.5	81.72	1989	81.79
68.15	1987	68.14	81.73–81.74	1989	81.86
68.16	1987	68.13	81.75	1989	81.87
68.23	1996	68.29	81.79	1989	81.79,81.87
68.51	1996	68.5	81.80	1989	81.81
68.59	1996	68.5	81.97	1992	81.59
68.9	1992	68.4	85.95	1987	85.99
74.3	1992	69.11 (Code deleted)	85.96	1987	85.99
75.37	1998	99.29			

See footnotes at end of table.

Table VI. Changes in procedure codes—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
86.06 . . . . .	1987	86.09	92.33 . . . . .	1998	92.3
86.07 . . . . .	1990	86.09	92.39 . . . . .	1998	92.3
86.27 . . . . .	1986	86.22–86.23	93.90 . . . . .	1988	93.92
86.28 . . . . .	1988	86.22	94.61–94.69 . . . . .	1989	94.25
86.67 . . . . .	1998	86.65	96.29 . . . . .	1998	96.39
86.93 . . . . .	1987	86.89	96.6 . . . . .	1986	96.35
88.90 . . . . .	1986	88.39	96.70 . . . . .	1991	93.92 (Code deleted)
88.91 . . . . .	1986	89.15	96.71 . . . . .	1991	93.92 (Code deleted)
88.92 . . . . .	1986	89.39	96.72 . . . . .	1991	93.92 (Code deleted)
88.93 . . . . .	1986	89.15	97.05 . . . . .	1989	51.97
			97.44 . . . . .	2001	37.64
88.94 . . . . .	1986	89.39	98.51–98.52 . . . . .	1989	59.96 (Code deleted)
88.95 . . . . .	1986	89.29	98.59 . . . . .	1989	59.96 (Code deleted)
88.97 . . . . .	1989	88.99	99.00 . . . . .	1995	99.02
88.98 . . . . .	1989	88.90	99.10 . . . . .	1998	99.29
88.99 . . . . .	1986	89.39	99.15 . . . . .	1986	99.29
89.10 . . . . .	1989	89.15	99.20 . . . . .	1998	99.29
89.17–89.18 . . . . .	1988	89.15	99.28 . . . . .	1994	99.25
89.19 . . . . .	1989	89.15	99.71–99.79 <sup>3</sup> . . . . .	1988	99.07
89.50 . . . . .	1991	89.54	99.75 . . . . .	2000	99.29
92.3 . . . . .	1995	01.59,04.07,07.63,07.68	99.85 . . . . .	1987	93.35
92.30 . . . . .	1998	92.3	99.86 . . . . .	1987	93.39
92.31 . . . . .	1998	92.3	99.88 . . . . .	1988	99.83
92.32 . . . . .	1998	92.3			

<sup>1</sup>Before October 1986 contents of current code 36.05 would have been assigned to 36.0.<sup>2</sup>Codes 78.90–78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90–78.99 were reassigned to codes 78.40–78.49.<sup>3</sup>Codes 99.71–99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71–99.79 were reclassified as "Therapeutic apheresis."

**U.S. DEPARTMENT OF  
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention  
National Center for Health Statistics  
3311 Toledo Road  
Hyattsville, Maryland 20782

---

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

MEDIA MAIL  
POSTAGE & FEES PAID  
CDC/NCHS  
PERMIT NO. G-284