Lis	t of Chartbook Figures		16.	Limitation of activity caused by selected chronic health conditions among older adults, by age:
Po	pulation	17	17.	United States, 2004–2005
1.	Total population and older population: United States, 1950–2050	17		45 years of age and over, by age and percent of poverty level: United States, 2005 49
2.	Foreign-born population, by citizenship: United States, 1970–2004.	19	Мо	rtality
3. 4.	Population in selected race and Hispanic origin groups, by age: United States, 1980–2006 Poverty by age: United States, 1966–2005	21 23	18. 19.	Life expectancy at birth and at 65 years of age, by race and sex: United States, 1970–2004 51 Infant, neonatal, and postneonatal mortality rates:
5.	Low income by age, race, and Hispanic origin: United States, 2005	25	20.	United States, 1950–2004
He	alth Care Expenditures		20.	ages: United States, 1950–2004
6.	Personal health care expenditures, by source of		Spe	ecial Feature: Access to Health Care
	funds and type of expenditures: United States, 2005	27	21.	
7.	National expenditures for mental health services, by source of funds: United States, 1986–2003	29		did not receive needed health-related services in the past 12 months because they could not afford them, by age and type of service: United States,
8.	National expenditures for substance abuse treatment, by source of funds: United States,			2005
	1986–2003	31	22.	Patient care physicians per 10,000 population, by county: United States, 2004 61
Health Risk Factors			23.	Obstetricians or gynecologists per 10,000 females age 15 years and over, by county: United States,
9.	Cigarette smoking among men, women, high school students, and mothers during pregnancy: United States, 1965–2005	33	24.	2004
10.	Blood cotinine levels among children 4–17 years of age, by percent of poverty level: United States, 1988–1994 and 2001–2004	35	25.	1988, 1996, and 2006
11.	Alcohol-related emergency department (ED) visit rates among persons 14–28 years of age, by age and sex: United States, 2002–2004	37	26.	origin: United States, 1988, 1996, and 2004 67 No usual source of care among adults 45–64 years of age, by selected diagnosed chronic
12.	Weekly restaurant meal consumption among people 1 year of age and over, by age: United States,		27.	conditions and race and Hispanic origin: United States, 2004–2005 69 Delayed medical care in the past 12 months due
13.	1999–2004	3941	21.	to lack of transportation among adults 18 years of age and over, by sex, percent of poverty level,
Morbidity and Limitation of Activity			28.	and age: United States, 2004–2005
14.	Limitation of activity caused by selected chronic health conditions among children, by age:	40	29.	United States, 1984–2005
15.	United States, 2004–2005	43		length of time uninsured and selected characteristics: United States, 2005
	age: United States, 2004–2005	45		

Health, United States, 2007 xiii

30.	The uninsured population under 65 years of age, by selected characteristics: United States, 2005	77
31.	Persons under 65 years of age who spent more than 10% of after-tax family income on out-of-pocket medical expenditures, by percent of poverty level: United States, 1996 and 2004	79
32.	Adults 20–64 years of age with undiagnosed high cholesterol or elevated blood pressure, by health insurance status and age: United States,	
	1999–2004	81
33.	Persons under 65 years of age who did not get needed medical care in the past year due to cost, by duration of health insurance coverage and percent	+
	of poverty level: United States, 2005	ι 83
34.	No dental visit in the past year among persons with natural teeth, by age and percent of poverty level:	
	United States, 2005	85
35.	Adults 50 years of age and over ever having a colorectal scope procedure, by selected characteristics: United States, annual average	
	2000, 2003, and 2005	87
36.	Adults 18 years of age and over reporting antidepressant drug use in the past month by sex and race and Hispanic origin: United States,	
	1988–1994 and 1999–2002	89

xiv Health, United States, 2007