

THE NEW YORK FERTILITY CENTER FLUSHING, NEW YORK

Comparison of success rates across clinics may not be meaningful. Patient medical characteristics and treatment approaches vary (see pages 11–20).

| 2014 ART CYCLE PROFILE | | | | Data verified by Tony Tsai, MD | | | | |
|---|------|-----------|-----|--------------------------------|-----|----------------|-----|--|
| Type of ART and Procedural Factors ^a | | | | Patient Diagnosis ^b | | | | |
| IVF | 100% | With ICSI | 88% | Tubal factor | 10% | Uterine factor | 8% | Multiple Factors: Female factors only 0% Female & male factors <1% |
| Unstimulated | 2% | PGD/PGS | 5% | Ovulatory dysfunction | 13% | Male factor | 8% | |
| Used gestational carrier | 0% | | | Diminished ovarian reserve | <1% | Other factor | 7% | |
| | | | | Endometriosis | 4% | Unknown factor | 49% | |

| 2014 ART SUCCESS RATES ^c | | Total number of cycles ^d : 251 (includes 0 cycle[s] using frozen eggs) | | | | | |
|---|--|---|-------|-------|-----------------------------------|--------|--------|
| Type of Cycle | | Age of Woman | | | | | |
| | | <35 | 35–37 | 38–40 | 41–42 | 43–44 | >44 |
| Fresh Embryos from Nondonor Eggs | | | | | | | |
| Number of cycles | | 80 | 34 | 37 | 24 | 16 | 28 |
| Percentage of cancellations before retrieval (%) | | 6.3 | 14.7 | 10.8 | 4.2 | 1 / 16 | 14.3 |
| Average number of embryos transferred | | 3.2 | 3.3 | 3.3 | 3.0 | 3.1 | 2.2 |
| Percentage of embryos transferred resulting in implantation (%) | | 16.0 | 6.7 | 11.5 | 3.6 | 2.9 | 0 / 18 |
| Percentage of elective single embryo transfers (eSET) (%) | | 0.0 | 0.0 | 0.0 | 0 / 13 | 0 / 9 | 0 / 6 |
| Outcomes per Cycle | | | | | | | |
| Percentage of cycles resulting in term, normal weight & singleton live births ^e (%) | | 6.3 | 2.9 | 8.1 | 0.0 | 0 / 16 | 0.0 |
| Percentage of cycles resulting in singleton live births (%) | | 11.3 | 5.9 | 8.1 | 4.2 | 0 / 16 | 0.0 |
| Percentage of cycles resulting in twin live births (%) | | 3.8 | 2.9 | 2.7 | 0.0 | 0 / 16 | 0.0 |
| Percentage of cycles resulting in live births (%) | | 15.0 | 8.8 | 10.8 | 4.2 | 0 / 16 | 0.0 |
| Percentage of cycles resulting in pregnancies (%) | | 36.3 | 17.6 | 21.6 | 12.5 | 1 / 16 | 3.6 |
| Outcomes per Transfer | | | | | | | |
| Number of transfers | | 65 | 26 | 26 | 19 | 11 | 12 |
| Percentage of transfers resulting in term, normal weight & singleton live births ^e (%) | | 7.7 | 3.8 | 11.5 | 0 / 19 | 0 / 11 | 0 / 12 |
| Percentage of transfers resulting in singleton live births (%) | | 13.8 | 7.7 | 11.5 | 1 / 19 | 0 / 11 | 0 / 12 |
| Percentage of transfers resulting in twin live births (%) | | 4.6 | 3.8 | 3.8 | 0 / 19 | 0 / 11 | 0 / 12 |
| Percentage of transfers resulting in live births (%) | | 18.5 | 11.5 | 15.4 | 1 / 19 | 0 / 11 | 0 / 12 |
| Percentage of transfers resulting in pregnancies (%) | | 44.6 | 23.1 | 30.8 | 3 / 19 | 1 / 11 | 1 / 12 |
| Frozen Embryos from Nondonor Eggs | | | | | | | |
| Number of cycles | | 12 | 8 | 0 | 1 | 3 | 2 |
| Number of transfers | | 11 | 8 | 0 | 1 | 3 | 2 |
| Estimated average number of transfers per retrieval | | 1.6 | 1.1 | | | 0.8 | 0.7 |
| Average number of embryos transferred | | 3.2 | 3.5 | | 1.0 | 3.0 | 1.0 |
| Percentage of embryos transferred resulting in implantation (%) | | 3.1 | 3.6 | | 0 / 1 | 0 / 9 | 1 / 2 |
| Percentage of transfers resulting in term, normal weight & singleton live births ^e (%) | | 0 / 11 | 0 / 8 | | 0 / 1 | 0 / 3 | 0 / 2 |
| Percentage of transfers resulting in singleton live births (%) | | 0 / 11 | 0 / 8 | | 0 / 1 | 0 / 3 | 1 / 2 |
| Percentage of transfers resulting in twin live births (%) | | 0 / 11 | 0 / 8 | | 0 / 1 | 0 / 3 | 0 / 2 |
| Percentage of transfers resulting in live births (%) | | 0 / 11 | 0 / 8 | | 0 / 1 | 0 / 3 | 1 / 2 |
| Percentage of transfers resulting in pregnancies (%) | | 2 / 11 | 1 / 8 | | 0 / 1 | 0 / 3 | 1 / 2 |
| Number of Egg/Embryo Banking Cycles | | | | | | | |
| | | 0 | 0 | 0 | 0 | 0 | 0 |
| Donor Eggs | | Fresh Embryos^f | | | Frozen Embryos^f | | |
| Number of cycles | | 0 | | | 6 | | |
| Number of transfers | | 0 | | | 6 | | |
| Average number of embryos transferred | | | | | 3.3 | | |
| Percentage of embryos transferred resulting in implantation (%) | | | | | 0.0 | | |
| Percentage of transfers resulting in term, normal weight & singleton live births ^e (%) | | | | | 0 / 6 | | |
| Percentage of transfers resulting in singleton live births (%) | | | | | 0 / 6 | | |
| Percentage of transfers resulting in twin live births (%) | | | | | 0 / 6 | | |
| Percentage of transfers resulting in live births (%) | | | | | 0 / 6 | | |
| Percentage of transfers resulting in pregnancies (%) | | | | | 0 / 6 | | |

| CURRENT SERVICES & PROFILE | | Current Name: The New York Fertility Center | | | | | |
|----------------------------|-----|---|-----|---------------|-----|-------------------------------|-----|
| Donor eggs? | Yes | Gestational carriers? | No | Single women? | Yes | Verified lab accreditation? | Yes |
| Donor embryos? | No | Embryo cryopreservation? | Yes | SART member? | No | (See Appendix C for details.) | |

^a Reflects features of fresh nondonor cycles. If IVF is less than 100%, the remaining cycles are GIFT, ZIFT, or a combination of these procedures with IVF.
^b Total patient diagnosis percentages may be greater than 100% because more than one diagnosis can be reported for each ART cycle.
^c Fractions are used for rates with denominators less than 20. Multiple-infant births (for example, twins) with at least one live infant are counted as one live birth.
^d Total cycle number includes those using frozen eggs. It excludes 0 cycle(s) evaluating new procedures. Both cycle types are excluded from ART success rates.
^e In this report, births are defined as term if at least 37 full weeks gestation and normal birth weight if at least 2,500 grams (approximately 5 pounds, 8 ounces).
^f All ages are reported together because previous data show that patient age does not materially affect success with donor eggs.