Table ALC-2a. Age-adjusted percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Both sexes | 100.0 | 34.8 (0.27) | 13.1 (0.14) | 31.6 (0.20) | 15.5 (0.17) | 5.1 (0.09) |
| Age | 100.0 | 34.0 (0.21) | 13.1 (0.14) | 31.0 (0.20) | 15.5 (0.17) | 3.1 (0.03) |
| 18–24 years | 100.0 | 41.0 (0.74) | 10.2 (0.42) | 30.6 (0.57) | 13.0 (0.45) | 5.2 (0.28) |
| 25–44 years | 100.0 | 26.5 (0.32) | 13.4 (0.21) | 37.7 (0.30) | 17.3 (0.25) | 5.1 (0.14) |
| 45–64 years | 100.0 | 33.4 (0.37) | 14.4 (0.22) | 30.4 (0.32) | 16.0 (0.25) | 5.7 (0.16) |
| 65–74 years | 100.0 | 44.7 (0.62) | 13.5 (0.34) | 23.4 (0.45) | 13.6 (0.39) | 4.9 (0.22) |
| 75 years and over | 100.0 | 59.3 (0.66) | 11.2 (0.36) | 16.5 (0.44) | 10.6 (0.39) | 2.4 (0.18) |
| Race | | | | | | |
| One race ² | 100.0 | 34.8 (0.27) | 13.1 (0.14) | 31.6 (0.20) | 15.5 (0.17) | 5.1 (0.09) |
| White | 100.0 | 31.8 (0.30) | 13.0 (0.16) | 32.8 (0.23) | 16.9 (0.19) | 5.5 (0.10) |
| Black or African American | 100.0 | 46.4 (0.52) | 13.9 (0.32) | 26.4 (0.42) | 10.0 (0.30) | 3.3 (0.17) |
| American Indian or Alaska Native | 100.0 | 46.1 (2.03) | 11.6 (1.15) | 26.2 (1.72) | 10.0 (1.02) | 6.2 (1.01) |
| Asian | 100.0 | 51.3 (0.72) | 12.7 (0.46) | 26.1 (0.62) | 8.1 (0.39) | 1.8 (0.20) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 46.3 (4.83) | *11.8 (3.64) | 27.7 (4.18) | 8.8 (2.40) | *5.5 (1.74) |
| Two or more races ³ | 100.0 | 36.8 (1.32) | 14.3 (0.97) | 29.3 (1.22) | 13.8 (0.93) | 5.7 (0.55) |
| Black or African American, white | 100.0 | 32.7 (3.34) | 18.0 (2.77) | 32.0 (3.34) | 9.9 (1.57) | 7.4 (1.67) |
| American Indian or Alaska Native, white | 100.0 | 35.7 (1.97) | 16.4 (1.51) | 28.1 (1.95) | 14.0 (1.60) | 5.8 (0.88) |
| Hispanic or Latino origin ⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 44.8 (0.47) | 13.3 (0.29) | 27.8 (0.41) | 11.1 (0.27) | 2.9 (0.14) |
| Mexican or Mexican American | 100.0 | 44.6 (0.58) | 13.4 (0.37) | 27.2 (0.50) | 11.4 (0.34) | 3.4 (0.19) |
| Not Hispanic or Latino | 100.0 | 33.0 (0.29) | 13.0 (0.15) | 32.3 (0.22) | 16.2 (0.18) | 5.5 (0.10) |
| White, single race | 100.0 | 29.0 (0.33) | 12.9 (0.19) | 33.9 (0.26) | 18.1 (0.22) | 6.1 (0.12) |
| Black or African American, single race | 100.0 | 46.5 (0.54) | 13.9 (0.33) | 26.2 (0.43) | 10.0 (0.31) | 3.3 (0.17) |
| American Indian or Alaska Native, single race | 100.0 | 48.9 (2.52) | 9.3 (1.31) | 23.4 (2.13) | 11.7 (1.53) | 6.7 (1.36) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 52.3 (0.52) | 12.6 (0.33) | 20.3 (0.40) | 10.4 (0.32) | 4.4 (0.22) |
| GED or equivalent ⁶ | 100.0 | 41.7 (1.04) | 14.8 (0.75) | 24.1 (0.89) | 12.9 (0.77) | 6.5 (0.53) |
| High school graduate | 100.0 | 38.0 (0.43) | 14.3 (0.28) | 28.9 (0.41) | 13.5 (0.30) | 5.3 (0.19) |
| Some college, no degree | 100.0 | 30.8 (0.46) | 15.9 (0.34) | 32.5 (0.44) | 15.4 (0.34) | 5.3 (0.20) |
| Associate's degree | 100.0 | 30.1 (0.52) | 16.0 (0.39) | 33.8 (0.50) | 15.2 (0.41) | 4.9 (0.28) |
| Bachelor's degree | 100.0 | 24.0 (0.40) | 11.6 (0.28) | 38.0 (0.42) | 20.7 (0.36) | 5.7 (0.20) |
| Master's, professional, or doctoral degree | 100.0 | 23.7 (0.52) | 11.2 (0.37) | 39.5 (0.57) | 21.5 (0.50) | 4.1 (0.22) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 51.6 (0.54) | 12.9 (0.31) | 21.9 (0.37) | 8.9 (0.28) | 4.7 (0.20) |
| 100% to less than 200% of the poverty threshold | 100.0 | 45.3 (0.48) | 14.7 (0.31) | 25.3 (0.36) | 10.3 (0.26) | 4.3 (0.17) |
| 200% to less than 400% of the poverty threshold | 100.0 | 34.9 (0.39) | 14.0 (0.25) | 31.5 (0.33) | 14.7 (0.26) | 5.0 (0.15) |
| 400% of the poverty threshold or greater | 100.0 | 23.2 (0.34) | 11.5 (0.21) | 38.4 (0.33) | 21.2 (0.31) | 5.7 (0.15) |

Table ALC-2a. Age-adjusted percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Science characteristic | and over | Nonamiker | difficol | Light dilliker | annicon | dillikoi |
| Both sexes (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 32.5 (0.34) | 14.1 (0.20) | 33.7 (0.28) | 15.6 (0.22) | 4.1 (0.11) |
| Widowed | 100.0 | 42.7 (2.05) | 14.2 (1.38) | 30.3 (2.21) | 10.1 (1.38) | 2.7 (0.39) |
| Divorced or separated | 100.0 | 32.5 (0.47) | 14.4 (0.35) | 31.9 (0.49) | 15.0 (0.38) | 6.1 (0.25) |
| Never married | 100.0 | 41.0 (0.50) | 11.8 (0.30) | 27.8 (0.39) | 13.9 (0.30) | 5.6 (0.19) |
| Living with a partner | 100.0 | 25.7 (0.70) | 12.6 (0.57) | 33.6 (0.74) | 20.0 (0.62) | 8.1 (0.41) |
| Region | | | | | | |
| Northeast | 100.0 | 31.5 (0.58) | 12.0 (0.34) | 34.9 (0.49) | 16.1 (0.40) | 5.5 (0.23) |
| Midwest | 100.0 | 30.9 (0.55) | 14.2 (0.33) | 33.7 (0.44) | 16.4 (0.35) | 4.8 (0.16) |
| South | 100.0 | 39.2 (0.47) | 13.0 (0.21) | 28.7 (0.32) | 14.1 (0.27) | 5.0 (0.15) |
| West | 100.0 | 34.2 (0.53) | 12.9 (0.25) | 31.5 (0.37) | 16.0 (0.33) | 5.3 (0.18) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 33.1 (0.30) | 12.9 (0.20) | 32.8 (0.25) | 16.2 (0.22) | 5.0 (0.11) |
| Small MSA | 100.0 | 34.5 (0.53) | 12.9 (0.24) | 31.7 (0.39) | 15.7 (0.31) | 5.3 (0.16) |
| Not in MSA | 100.0 | 41.0 (0.88) | 14.5 (0.34) | 27.0 (0.59) | 12.6 (0.41) | 4.9 (0.24) |

Table ALC-2a. Age-adjusted percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Men | 100.0 | 29.5 (0.31) | 9.9 (0.17) | 32.5 (0.26) | 22.6 (0.25) | 5.5 (0.13) |
| Age | | | | | | |
| 18–24 years | 100.0 | 37.6 (0.97) | 8.2 (0.51) | 30.0 (0.80) | 18.8 (0.72) | 5.4 (0.38) |
| 25–44 years | 100.0 | 21.2 (0.38) | 9.4 (0.28) | 38.3 (0.43) | 25.5 (0.39) | 5.6 (0.20) |
| 45–64 years | 100.0 | 28.7 (0.46) | 11.0 (0.27) | 31.3 (0.43) | 22.9 (0.39) | 6.1 (0.26) |
| 65–74 years | 100.0 | 38.4 (0.83) | 10.4 (0.46) | 25.4 (0.66) | 20.5 (0.65) | 5.4 (0.35) |
| 75 years and over | 100.0 | 50.0 (1.03) | 10.6 (0.61) | 20.4 (0.76) | 16.5 (0.72) | 2.5 (0.30) |
| Race | | | | | | |
| One race ² | 100.0 | 29.5 (0.31) | 9.9 (0.17) | 32.5 (0.26) | 22.7 (0.26) | 5.5 (0.13) |
| White | 100.0 | 27.3 (0.33) | 9.5 (0.19) | 33.0 (0.30) | 24.3 (0.29) | 5.9 (0.15) |
| Black or African American | 100.0 | 40.0 (0.78) | 10.8 (0.44) | 29.0 (0.68) | 16.2 (0.54) | 4.0 (0.26) |
| American Indian or Alaska Native | 100.0 | 38.0 (2.38) | 10.3 (1.47) | 26.9 (2.24) | 16.0 (1.98) | 8.7 (1.60) |
| Asian | 100.0 | 40.4 (0.99) | 12.5 (0.70) | 32.8 (0.95) | 12.5 (0.69) | 1.9 (0.34) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 39.1 (4.70) | * | 28.8 (5.35) | 15.6 (4.25) | * |
| Two or more races ³ | 100.0 | 33.8 (1.85) | 10.1 (1.46) | 29.3 (1.91) | 20.8 (1.66) | 6.0 (0.81) |
| Black or African American, white | 100.0 | 26.0 (5.09) | 12.8 (3.54) | 33.4 (4.81) | 16.5 (3.10) | *11.3 (4.52) |
| American Indian or Alaska Native, white | 100.0 | 34.8 (2.99) | 12.2 (2.41) | 25.3 (2.99) | 20.8 (2.76) | 6.9 (1.41) |
| Hispanic or Latino origin ⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 34.7 (0.60) | 10.2 (0.38) | 32.8 (0.60) | 18.1 (0.50) | 4.2 (0.24) |
| Mexican or Mexican American | 100.0 | 33.6 (0.77) | 10.1 (0.50) | 32.5 (0.73) | 18.9 (0.65) | 4.9 (0.33) |
| Not Hispanic or Latino | 100.0 | 28.6 (0.34) | 9.8 (0.19) | 32.4 (0.30) | 23.5 (0.28) | 5.7 (0.14) |
| White, single race | 100.0 | 25.6 (0.37) | 9.3 (0.22) | 33.1 (0.34) | 25.7 (0.33) | 6.2 (0.17) |
| Black or African American, single race | 100.0 | 40.3 (0.81) | 10.8 (0.45) | 28.8 (0.70) | 16.2 (0.55) | 3.9 (0.27) |
| American Indian or Alaska Native, single race | 100.0 | 40.5 (3.11) | 9.4 (1.99) | 21.5 (3.18) | 19.4 (3.14) | 9.3 (1.99) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 40.6 (0.69) | 10.2 (0.45) | 26.0 (0.61) | 17.0 (0.55) | 6.1 (0.37) |
| GED or equivalent ⁶ | 100.0 | 35.8 (1.51) | 11.3 (1.03) | 26.2 (1.32) | 18.4 (1.22) | 8.3 (0.85) |
| High school graduate | 100.0 | 31.7 (0.56) | 10.7 (0.35) | 30.8 (0.62) | 20.1 (0.50) | 6.7 (0.29) |
| Some college, no degree | 100.0 | 25.3 (0.62) | 12.0 (0.45) | 33.4 (0.65) | 23.5 (0.59) | 5.8 (0.32) |
| Associate's degree | 100.0 | 25.3 (0.74) | 11.2 (0.53) | 35.1 (0.83) | 22.9 (0.73) | 5.5 (0.51) |
| Bachelor's degree | 100.0 | 20.8 (0.52) | 8.3 (0.36) | 37.4 (0.61) | 28.9 (0.57) | 4.7 (0.28) |
| Master's, professional, or doctoral degree | 100.0 | 21.8 (0.69) | 8.4 (0.46) | 38.1 (0.77) | 28.9 (0.77) | 2.8 (0.26) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 44.2 (0.78) | 10.0 (0.41) | 24.5 (0.59) | 15.0 (0.51) | 6.3 (0.34) |
| 100% to less than 200% of the poverty threshold | 100.0 | 38.8 (0.65) | 11.4 (0.41) | 27.6 (0.57) | 16.6 (0.46) | 5.6 (0.29) |
| 200% to less than 400% of the poverty threshold | 100.0 | 30.3 (0.51) | 10.6 (0.32) | 31.9 (0.47) | 21.6 (0.42) | 5.6 (0.22) |
| 400% of the poverty threshold or greater | 100.0 | 20.4 (0.42) | 8.5 (0.27) | 37.6 (0.46) | 28.5 (0.46) | 5.1 (0.21) |

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| | 4114 6761 | - Trondinino | - Carrinton | Light difficol | dillino. | dillitor |
| Men (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 27.1 (0.38) | 10.8 (0.26) | 35.7 (0.37) | 22.4 (0.35) | 4.0 (0.16) |
| Widowed | 100.0 | 29.7 (3.17) | 15.5 (3.77) | 31.5 (4.16) | 19.7 (3.29) | 3.5 (0.74) |
| Divorced or separated | 100.0 | 26.8 (0.70) | 9.5 (0.45) | 30.7 (0.76) | 24.5 (0.75) | 8.5 (0.45) |
| Never married | 100.0 | 37.2 (0.69) | 9.2 (0.41) | 27.8 (0.58) | 19.6 (0.48) | 6.2 (0.30) |
| Living with a partner | 100.0 | 21.4 (0.90) | 8.3 (0.63) | 32.6 (1.02) | 28.8 (1.02) | 8.9 (0.57) |
| Region | | | | | | |
| Northeast | 100.0 | 27.3 (0.80) | 9.6 (0.44) | 35.1 (0.69) | 23.0 (0.62) | 5.1 (0.30) |
| Midwest | 100.0 | 26.9 (0.62) | 10.2 (0.39) | 33.6 (0.57) | 24.1 (0.52) | 5.3 (0.25) |
| South | 100.0 | 33.0 (0.54) | 9.9 (0.25) | 30.3 (0.43) | 20.9 (0.43) | 5.9 (0.23) |
| West | 100.0 | 28.6 (0.56) | 9.8 (0.33) | 32.8 (0.49) | 23.4 (0.50) | 5.3 (0.25) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 28.0 (0.37) | 9.9 (0.24) | 33.8 (0.36) | 23.3 (0.34) | 5.1 (0.17) |
| Small MSA | 100.0 | 29.2 (0.58) | 9.4 (0.29) | 32.2 (0.46) | 23.2 (0.48) | 6.0 (0.23) |
| Not in MSA | 100.0 | 35.1 (0.93) | 11.2 (0.43) | 28.4 (0.71) | 19.3 (0.67) | 6.0 (0.37) |

Table ALC-2a. Age-adjusted percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Colotted Ghardelensite | and over | TTOTIGHTING | dillikoi | Light diffici | dillikoi | dillikoi |
| Women | 100.0 | 39.6 (0.33) | 16.1 (0.20) | 30.8 (0.26) | 8.7 (0.15) | 4.7 (0.11) |
| Age | | | | | | |
| 18–24 years | 100.0 | 44.5 (0.96) | 12.1 (0.61) | 31.1 (0.79) | 7.2 (0.41) | 5.0 (0.37) |
| 25–44 years | 100.0 | 31.6 (0.43) | 17.3 (0.31) | 37.1 (0.40) | 9.3 (0.24) | 4.7 (0.18) |
| 45–64 years | 100.0 | 37.8 (0.46) | 17.7 (0.33) | 29.6 (0.40) | 9.6 (0.26) | 5.3 (0.19) |
| 65–74 years | 100.0 | 50.1 (0.73) | 16.2 (0.50) | 21.6 (0.57) | 7.6 (0.37) | 4.5 (0.27) |
| 75 years and over | 100.0 | 65.6 (0.75) | 11.7 (0.45) | 13.8 (0.49) | 6.5 (0.36) | 2.4 (0.21) |
| Race | | | | | | |
| One race ² | 100.0 | 39.7 (0.34) | 16.1 (0.20) | 30.8 (0.26) | 8.8 (0.15) | 4.7 (0.11) |
| White | 100.0 | 36.1 (0.39) | 16.3 (0.24) | 32.6 (0.31) | 9.7 (0.18) | 5.2 (0.13) |
| Black or African American | 100.0 | 51.5 (0.64) | 16.5 (0.44) | 24.3 (0.52) | 4.9 (0.25) | 2.8 (0.19) |
| American Indian or Alaska Native | 100.0 | 53.3 (2.64) | 12.9 (1.74) | 25.6 (2.29) | 4.4 (1.05) | 3.7 (0.85) |
| Asian | 100.0 | 60.8 (0.96) | 12.9 (0.64) | 20.2 (0.78) | 4.3 (0.39) | 1.8 (0.25) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 57.7 (5.72) | *10.4 (3.31) | 24.0 (5.07) | * | *5.5 (1.98) |
| Two or more races ³ | 100.0 | 39.5 (1.77) | 18.0 (1.34) | 29.4 (1.57) | 7.6 (0.93) | 5.5 (0.77) |
| Black or African American, white | 100.0 | 36.7 (4.31) | 19.8 (3.64) | 30.3 (3.97) | *6.3 (1.93) | 7.0 (1.97) |
| American Indian or Alaska Native, white | 100.0 | 36.7 (2.49) | 20.5 (1.93) | 30.5 (2.48) | 7.5 (1.50) | 4.8 (1.18) |
| Hispanic or Latino origin⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 54.7 (0.61) | 16.5 (0.44) | 23.0 (0.49) | 4.1 (0.22) | 1.7 (0.13) |
| Mexican or Mexican American | 100.0 | 55.7 (0.78) | 16.8 (0.57) | 21.8 (0.62) | 3.8 (0.27) | 1.8 (0.18) |
| Not Hispanic or Latino | 100.0 | 37.0 (0.36) | 16.1 (0.22) | 32.2 (0.28) | 9.5 (0.17) | 5.2 (0.13) |
| White, single race | 100.0 | 32.0 (0.43) | 16.2 (0.27) | 34.8 (0.35) | 10.9 (0.21) | 6.0 (0.16) |
| Black or African American, single race | 100.0 | 51.4 (0.66) | 16.6 (0.45) | 24.2 (0.54) | 4.9 (0.26) | 2.8 (0.20) |
| American Indian or Alaska Native, single race | 100.0 | 55.8 (3.21) | 9.2 (1.78) | 25.2 (2.82) | 5.3 (1.52) | 4.5 (1.22) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 64.0 (0.69) | 15.1 (0.51) | 14.6 (0.50) | 3.6 (0.26) | 2.8 (0.24) |
| GED or equivalent ⁶ | 100.0 | 48.8 (1.45) | 18.9 (1.15) | 21.8 (1.17) | 6.2 (0.79) | 4.3 (0.63) |
| High school graduate | 100.0 | 44.5 (0.59) | 18.1 (0.46) | 27.0 (0.53) | 6.5 (0.28) | 3.9 (0.22) |
| Some college, no degree | 100.0 | 35.5 (0.61) | 19.2 (0.46) | 31.9 (0.58) | 8.5 (0.33) | 4.9 (0.25) |
| Associate's degree | 100.0 | 34.0 (0.69) | 19.9 (0.60) | 32.7 (0.65) | 8.9 (0.40) | 4.4 (0.29) |
| Bachelor's degree | 100.0 | 27.0 (0.54) | 14.7 (0.43) | 38.5 (0.57) | 13.1 (0.39) | 6.7 (0.30) |
| Master's, professional, or doctoral degree | 100.0 | 25.7 (0.68) | 13.9 (0.56) | 40.7 (0.79) | 14.4 (0.57) | 5.3 (0.34) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 56.9 (0.61) | 15.0 (0.41) | 20.2 (0.44) | 4.5 (0.24) | 3.4 (0.22) |
| 100% to less than 200% of the poverty threshold | 100.0 | 50.7 (0.63) | 17.6 (0.44) | 23.5 (0.48) | 5.0 (0.24) | 3.2 (0.18) |
| 200% to less than 400% of the poverty threshold | 100.0 | 39.3 (0.54) | 17.3 (0.38) | 31.2 (0.47) | 7.9 (0.26) | 4.4 (0.20) |
| 400% of the poverty threshold or greater | 100.0 | 26.1 (0.47) | 14.7 (0.33) | 39.3 (0.46) | 13.5 (0.32) | 6.4 (0.23) |
| | | , , | , , | | | , , |

Table ALC-2a. Age-adjusted percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| | | | | | | |
| Women (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 38.0 (0.43) | 17.3 (0.29) | 31.7 (0.37) | 8.8 (0.21) | 4.2 (0.14) |
| Widowed | 100.0 | 46.1 (2.38) | 14.0 (1.42) | 29.9 (2.54) | 7.5 (1.51) | 2.5 (0.45) |
| Divorced or separated | 100.0 | 36.5 (0.63) | 17.8 (0.49) | 32.7 (0.65) | 8.5 (0.37) | 4.4 (0.25) |
| Never married | 100.0 | 44.8 (0.69) | 14.5 (0.45) | 28.0 (0.52) | 7.8 (0.32) | 4.9 (0.24) |
| Living with a partner | 100.0 | 29.9 (1.08) | 16.8 (0.88) | 34.7 (1.04) | 11.3 (0.66) | 7.3 (0.60) |
| Region | | | | | | |
| Northeast | 100.0 | 35.4 (0.71) | 14.3 (0.45) | 34.8 (0.63) | 9.7 (0.39) | 5.9 (0.34) |
| Midwest | 100.0 | 34.7 (0.72) | 18.1 (0.47) | 33.8 (0.60) | 9.1 (0.32) | 4.3 (0.21) |
| South | 100.0 | 44.8 (0.56) | 15.9 (0.32) | 27.3 (0.40) | 7.9 (0.24) | 4.1 (0.17) |
| West | 100.0 | 39.5 (0.67) | 16.0 (0.37) | 30.3 (0.51) | 8.9 (0.31) | 5.3 (0.23) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 37.8 (0.39) | 15.8 (0.28) | 31.9 (0.33) | 9.5 (0.21) | 5.0 (0.16) |
| Small MSA | 100.0 | 39.3 (0.67) | 16.2 (0.34) | 31.3 (0.51) | 8.6 (0.28) | 4.6 (0.20) |
| Not in MSA | 100.0 | 46.5 (1.03) | 17.5 (0.52) | 25.8 (0.71) | 6.4 (0.30) | 3.8 (0.24) |

^{*} Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and estimates were weighted using the Sample Adult weight. Unknowns for the columns were not included in the denominators when calculating percentages. Percentages may not add to totals due to rounding. "Both sexes," "men," and "women" include persons of other races and unknown race and ethnicity, unknown education, and unknown marital status. Unless otherwise specified, estimates are age-adjusted using the projected 2000 U.S. population as the standard population and using three age groups: 18–44, 45–64, and 65 and over. Estimates for age groups are not age-adjusted. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from http://www.cdc.gov/nchs/nhis/SHS/tables.htm). Suggested citation: Villarroel MA, Clarke TC, Schoenborn CA. Tables of adult health behaviors, alcohol use: National Health Interview Survey, 2011–2014. 2016. Available from: http://www.cdc.gov/nchs/nhis/SHS/tables.htm.

¹Derived from several source questions: "In any one year, have you had at least 12 drinks of any type of alcoholic beverage?", "In your entire life, have you had at least 12 drinks of any type of alcoholic beverage?", "In the past year, how often did you drink any type of alcoholic beverage?" and "In the past year, on those days that you drank alcoholic beverages, on the average, how many drinks did you have?" A nondrinker had zero drinks in the past year (includes former drinkers and lifetime abstainers). An infrequent drinker had at least 12 drinks in his or her lifetime and 1–11 drinks in the past year. A light drinker had 3 drinks or less per week, on average, in the past year. A moderate drinker had more than 3 and up to and including 7 drinks per week for women, on average, in the past year. A heavier drinker had more than 14 drinks per week for women, on average, in the past year.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

⁵Shown only for adults aged 25 and over. Estimates are age adjusted to the projected 2000 U.S. population as the standard population using three age groups: 25–44, 45–64, and 65 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷Based on family income and family size using the U.S. Census Bureau poverty thresholds for 2010, 2011, 2012, and 2013.

⁸MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Both sexes | 235,845 | 81,421 | 30,557 | 72,145 | 35,695 | 11,805 |
| Age | | | | | | |
| 18–24 years | 30,153 | 12,109 | 3,004 | 9,026 | 3,848 | 1,530 |
| 25–44 years | 81,276 | 21,168 | 10,700 | 30,123 | 13,811 | 4,073 |
| 45–64 years | 81,945 | 26,880 | 11,628 | 24,473 | 12,902 | 4,599 |
| 65–74 years | 24,278 | 10,667 | 3,217 | 5,575 | 3,239 | 1,168 |
| 75 years and over | 18,194 | 10,596 | 2,008 | 2,949 | 1,895 | 434 |
| Race | | | | | | |
| One race ² | 232,068 | 80,139 | 30,048 | 71,000 | 35,166 | 11,581 |
| White | 188,610 | 60,173 | 24,255 | 59,607 | 31,090 | 10,255 |
| Black or African American | 28,367 | 12,602 | 3,903 | 7,416 | 2,822 | 950 |
| American Indian or Alaska Native | 1,954 | 853 | 226 | 507 | 193 | 119 |
| Asian | 12,745 | 6,342 | 1,621 | 3,358 | 1,024 | 233 |
| Native Hawaiian or Other Pacific Islander | 393 | 169 | *44 | 113 | *37 | *23 |
| Two or more races ³ | 3,776 | 1,282 | 508 | 1,145 | 529 | 224 |
| Black or African American, white | 753 | 223 | 97 | 253 | 102 | 58 |
| American Indian or Alaska Native, white | 1,557 | 545 | 258 | 421 | 216 | 89 |
| Hispanic or Latino origin⁴ and race | | | | | | |
| Hispanic or Latino | 34,996 | 14,738 | 4,579 | 10,035 | 3,940 | 1,053 |
| Mexican or Mexican American | 21,611 | 8,963 | 2,831 | 6,142 | 2,540 | 750 |
| Not Hispanic or Latino | 200,849 | 66,683 | 25,978 | 62,110 | 31,756 | 10,751 |
| White, single race | 156,720 | 46,632 | 20,102 | 50,570 | 27,468 | 9,316 |
| Black or African American, single race | 27,201 | 12,139 | 3,745 | 7,035 | 2,711 | 918 |
| American Indian or Alaska Native, single race | 1,230 | 584 | 111 | 276 | 135 | 79 |
| Education ⁵ | | | | | | |
| Less than high school diploma | 27,998 | 15,111 | 3,365 | 5,177 | 2,637 | 1,152 |
| GED or equivalent ⁶ | 6,018 | 2,419 | 885 | 1,433 | 758 | 383 |
| High school graduate | 46,650 | 18,261 | 6,529 | 12,582 | 5,942 | 2,412 |
| Some college, no degree | 35,745 | 10,977 | 5,570 | 11,265 | 5,419 | 1,872 |
| Associate's degree | 23,771 | 6,959 | 3,764 | 7,898 | 3,610 | 1,171 |
| Bachelor's degree | 40,524 | 9,357 | 4,678 | 15,406 | 8,256 | 2,285 |
| Master's, professional, or doctoral degree | 23,877 | 5,637 | 2,674 | 9,194 | 5,107 | 965 |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 32,666 | 15,903 | 4,138 | 7,328 | 2,962 | 1,526 |
| 100% to less than 200% of the poverty threshold | 45,025 | 20,495 | 6,445 | 10,926 | 4,459 | 1,856 |
| 200% to less than 400% of the poverty threshold | 70,656 | 24,932 | 9,730 | 21,346 | 10,052 | 3,419 |
| 400% of the poverty threshold or greater | 87,498 | 20,090 | 10,244 | 32,545 | 18,223 | 5,004 |

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|--|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Both Sexes (continued) Marital status | | | | | | |
| Married | 124,885 | 41,004 | 17,279 | 40,128 | 19,430 | 5,310 |
| Widowed | 14,165 | 8,138 | 1,749 | 2,489 | 1,085 | 418 |
| Divorced or separated | 26,937 | 9,138 | 3,916 | 7,747 | 3,873 | 1,659 |
| Never married | 52,245 | 19,175 | 5,396 | 15,590 | 7,857 | 3,012 |
| Living with a partner | 17,226 | 3,826 | 2,168 | 6,088 | 3,419 | 1,390 |
| Region | | | | | | |
| Northeast | 41,864 | 13,053 | 5,014 | 14,070 | 6,680 | 2,218 |
| Midwest | 53,927 | 16,559 | 7,600 | 17,639 | 8,656 | 2,530 |
| South | 86,250 | 33,662 | 11,045 | 23,849 | 11,876 | 4,229 |
| West | 53,805 | 18,147 | 6,898 | 16,587 | 8,483 | 2,828 |
| Place of residence ⁸ | | | | | | |
| Large MSA | 126,734 | 41,066 | 16,174 | 40,679 | 20,171 | 6,320 |
| Small MSA | 72,525 | 24,889 | 9,286 | 22,245 | 11,134 | 3,745 |
| Not in MSA | 36,586 | 15,466 | 5,097 | 9,221 | 4,391 | 1,740 |

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Men | 113,760 | 32,977 | 11,109 | 35,991 | 25,187 | 6,145 |
| Age | | | | | | |
| 18–24 years | 15,174 | 5,562 | 1,216 | 4,448 | 2,781 | 795 |
| 25–44 years | 40,089 | 8,326 | 3,682 | 15,046 | 10,030 | 2,181 |
| 45–64 years | 39,742 | 11,171 | 4,289 | 12,197 | 8,898 | 2,386 |
| 65–74 years | 11,317 | 4,259 | 1,148 | 2,812 | 2,271 | 596 |
| 75 years and over | 7,437 | 3,658 | 773 | 1,488 | 1,207 | 186 |
| Race | | | | | | |
| One race ² | 111,961 | 32,423 | 10,935 | 35,457 | 24,813 | 6,034 |
| White | 92,063 | 24,873 | 8,719 | 29,491 | 21,828 | 5,317 |
| Black or African American | 12,796 | 4,844 | 1,355 | 3,673 | 2,067 | 512 |
| American Indian or Alaska Native | 938 | 336 | 96 | 246 | 149 | 80 |
| Asian | 5,969 | 2,314 | 740 | 1,982 | 737 | 111 |
| Native Hawaiian or Other Pacific Islander | 195 | 55 | * | 65 | *33 | * |
| Two or more races ³ | 1,799 | 554 | 174 | 535 | 374 | 111 |
| Black or African American, white | 304 | 68 | 35 | 102 | 64 | 21 |
| American Indian or Alaska Native, white | 757 | 256 | 95 | 186 | 158 | 51 |
| Hispanic or Latino origin⁴ and race | | | | | | |
| Hispanic or Latino | 17,639 | 5,649 | 1,745 | 5,888 | 3,208 | 755 |
| Mexican or Mexican American | 11,070 | 3,373 | 1,073 | 3,705 | 2,114 | 550 |
| Not Hispanic or Latino | 96,121 | 27,328 | 9,364 | 30,103 | 21,979 | 5,390 |
| White, single race | 75,967 | 19,692 | 7,143 | 24,158 | 18,864 | 4,638 |
| Black or African American, single race | 12,261 | 4,687 | 1,291 | 3,475 | 1,982 | 491 |
| American Indian or Alaska Native, single race | 549 | 215 | 50 | 115 | 102 | 48 |
| Education ⁵ | | | | | | |
| Less than high school diploma | 13,693 | 5,688 | 1,377 | 3,330 | 2,166 | 794 |
| GED or equivalent ⁶ | 3,268 | 1,115 | 360 | 845 | 591 | 265 |
| High school graduate | 22,609 | 7,138 | 2,372 | 6,667 | 4,404 | 1,499 |
| Some college, no degree | 16,464 | 4,100 | 1,963 | 5,341 | 3,803 | 941 |
| Associate's degree | 10,637 | 2,590 | 1,169 | 3,679 | 2,422 | 597 |
| Bachelor's degree | 19,491 | 3,923 | 1,608 | 7,231 | 5,508 | 895 |
| Master's, professional, or doctoral degree | 11,883 | 2,632 | 1,001 | 4,360 | 3,399 | 334 |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 13,870 | 5,682 | 1,345 | 3,442 | 2,113 | 874 |
| 100% to less than 200% of the poverty threshold | 20,335 | 7,734 | 2,260 | 5,473 | 3,287 | 1,111 |
| 200% to less than 400% of the poverty threshold | 34,669 | 10,497 | 3,629 | 10,729 | 7,309 | 1,885 |
| 400% of the poverty threshold or greater | 44,886 | 9,063 | 3,875 | 16,348 | 12,478 | 2,275 |

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Men (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 63,247 | 17,767 | 6,799 | 21,215 | 13,871 | 2,590 |
| Widowed | 3,037 | 1,414 | 299 | 635 | 483 | 131 |
| Divorced or separated | 11,148 | 3,101 | 1,084 | 3,123 | 2,582 | 967 |
| Never married | 27,722 | 9,142 | 2,259 | 8,085 | 5,781 | 1,701 |
| Living with a partner | 8,467 | 1,515 | 654 | 2,901 | 2,450 | 748 |
| Region | | | | | | |
| Northeast | 20,132 | 5,393 | 1,897 | 6,826 | 4,559 | 994 |
| Midwest | 26,300 | 6,914 | 2,650 | 8,640 | 6,195 | 1,371 |
| South | 40,974 | 13,305 | 4,006 | 12,021 | 8,357 | 2,397 |
| West | 26,354 | 7,366 | 2,556 | 8,505 | 6,075 | 1,381 |
| Place of residence ⁸ | | | | | | |
| Large MSA | 61,372 | 16,629 | 5,950 | 20,384 | 14,041 | 3,082 |
| Small MSA | 34,977 | 10,046 | 3,265 | 10,948 | 7,942 | 2,057 |
| Not in MSA | 17,411 | 6,302 | 1,894 | 4,660 | 3,204 | 1,006 |

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Women | 122,085 | 48,444 | 19,448 | 36,154 | 10,509 | 5,660 |
| Age | ,000 | , | .0,0 | 33,131 | . 0,000 | 0,000 |
| 18–24 years | 14,979 | 6,547 | 1,787 | 4,578 | 1,066 | 734 |
| 25–44 years | 41,187 | 12,841 | 7,018 | 15,077 | 3,781 | 1,892 |
| 45–64 years | 42,203 | 15,710 | 7,339 | 12,276 | 4,004 | 2,213 |
| 65–74 years | 12,960 | 6,408 | 2,068 | 2,763 | 968 | 572 |
| 75 years and over | 10,757 | 6,938 | 1,235 | 1,461 | 688 | 249 |
| Race | | | | | | |
| One race ² | 120,107 | 47,716 | 19,113 | 35,543 | 10,353 | 5,547 |
| White | 96,547 | 35,300 | 15,536 | 30,117 | 9,262 | 4,938 |
| Black or African American | 15,570 | 7,758 | 2,547 | 3,743 | 755 | 438 |
| American Indian or Alaska Native | 1,015 | 517 | 129 | 260 | 44 | 39 |
| Asian | 6,776 | 4,028 | 881 | 1,376 | 287 | 122 |
| Native Hawaiian or Other Pacific Islander | 198 | 114 | *20 | 48 | * | *10 |
| Two or more races ³ | 1,978 | 728 | 334 | 611 | 155 | 113 |
| Black or African American, white | 449 | 155 | 62 | 152 | *38 | 38 |
| American Indian or Alaska Native, white | 800 | 289 | 163 | 235 | 58 | 38 |
| Hispanic or Latino origin ⁴ and race | | | | | | |
| Hispanic or Latino | 17,357 | 9,090 | 2,834 | 4,147 | 732 | 299 |
| Mexican or Mexican American | 10,541 | 5,590 | 1,757 | 2,437 | 426 | 201 |
| Not Hispanic or Latino | 104,728 | 39,354 | 16,614 | 32,007 | 9,777 | 5,361 |
| White, single race | 80,752 | 26,941 | 12,959 | 26,413 | 8,603 | 4,678 |
| Black or African American, single race | 14,940 | 7,452 | 2,453 | 3,560 | 729 | 427 |
| American Indian or Alaska Native, single race | 681 | 369 | 61 | 162 | *33 | 31 |
| Education ⁵ | | | | | | |
| Less than high school diploma | 14,305 | 9,423 | 1,988 | 1,847 | 471 | 358 |
| GED or equivalent ⁶ | 2,750 | 1,304 | 525 | 588 | 167 | 118 |
| High school graduate | 24,041 | 11,124 | 4,157 | 5,916 | 1,537 | 913 |
| Some college, no degree | 19,281 | 6,877 | 3,607 | 5,924 | 1,616 | 930 |
| Associate's degree | 13,134 | 4,369 | 2,595 | 4,219 | 1,188 | 574 |
| Bachelor's degree | 21,032 | 5,434 | 3,070 | 8,176 | 2,748 | 1,390 |
| Master's, professional, or doctoral degree | 11,994 | 3,005 | 1,673 | 4,834 | 1,708 | 631 |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 18,796 | 10,221 | 2,793 | 3,886 | 849 | 652 |
| 100% to less than 200% of the poverty threshold | 24,690 | 12,762 | 4,184 | 5,453 | 1,172 | 745 |
| 200% to less than 400% of the poverty threshold | 35,988 | 14,435 | 6,101 | 10,618 | 2,743 | 1,533 |
| 400% of the poverty threshold or greater | 42,612 | 11,027 | 6,369 | 16,197 | 5,745 | 2,729 |

Table ALC-2b. Frequency distribution (number in thousands) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Women (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 61,638 | 23,238 | 10,480 | 18,913 | 5,559 | 2,720 |
| Widowed | 11,128 | 6,724 | 1,450 | 1,854 | 602 | 287 |
| Divorced or separated | 15,788 | 6,037 | 2,833 | 4,624 | 1,292 | 692 |
| Never married | 24,523 | 10,032 | 3,138 | 7,505 | 2,076 | 1,311 |
| iving with a partner | 8,759 | 2,310 | 1,514 | 3,187 | 969 | 641 |
| Region | | | | | | |
| Northeast | 21,732 | 7,660 | 3,117 | 7,244 | 2,121 | 1,224 |
| Midwest | 27,626 | 9,645 | 4,951 | 8,999 | 2,460 | 1,158 |
| South | 45,276 | 20,358 | 7,039 | 11,828 | 3,519 | 1,832 |
| Vest | 27,451 | 10,781 | 4,342 | 8,083 | 2,408 | 1,446 |
| Place of residence ⁸ | | | | | | |
| _arge MSA | 65,362 | 24,437 | 10,224 | 20,295 | 6,129 | 3,238 |
| Small MSA | 37,548 | 14,843 | 6,021 | 11,297 | 3,192 | 1,689 |
| Not in MSA | 19,175 | 9,164 | 3,203 | 4,562 | 1,187 | 733 |

^{*} Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and estimates were weighted using the Sample Adult weight. Unknowns for the columns were not included in the frequencies, but they were included in the "All adults aged 18 and over" column. "Both sexes," "men," and "women" include persons of other races and unknown race and ethnicity, unknown education, and unknown marital status. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from

http://www.cdc.gov/nchs/nhis/SHS/tables.htm). Suggested citation: Villarroel MA, Clarke TC, Schoenborn CA. Tables of adult health behaviors, alcohol use: National Health Interview Survey, 2011–2014. 2016. Available from: http://www.cdc.gov/nchs/nhis/SHS/tables.htm.

SOURCE: NCHS, National Health Interview Survey, 2011-2014.

¹Derived from several source questions: "In any one year, have you had at least 12 drinks of any type of alcoholic beverage?", "In your entire life, have you had at least 12 drinks of any type of alcoholic beverage?", "In the past year, how often did you drink any type of alcoholic beverage?" and "In the past year, on those days that you drank alcoholic beverages, on the average, how many drinks did you have?" A nondrinker had zero drinks in the past year (includes former drinkers and lifetime abstainers). An infrequent drinker had at least 12 drinks in his or her lifetime and 1–11 drinks in the past year. A light drinker had 3 drinks or less per week, on average, in the past year. A moderate drinker had more than 3 and up to and including 14 drinks per week for men, and more than 3 and up to and including 7 drinks per week for women, on average, in the past year. A heavier drinker had more than 14 drinks per week for women, on average, in the past year.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, recardless of race.

⁵Shown only for adults aged 25 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷Based on family income and family size using the U.S. Census Bureau poverty thresholds for 2010, 2011, 2012, and 2013.

⁸MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Both sexes | 100.0 | 35.2 (0.27) | 13.2 (0.14) | 31.1 (0.20) | 15.4 (0.16) | 5.1 (0.09) |
| Age | 100.0 | 33.2 (0.21) | 13.2 (0.14) | 31.1 (0.20) | 13.4 (0.10) | 5.1 (0.09) |
| 18–24 years | 100.0 | 41.0 (0.74) | 10.2 (0.42) | 30.6 (0.57) | 13.0 (0.45) | 5.2 (0.28) |
| 25–44 years | 100.0 | 26.5 (0.32) | 13.4 (0.21) | 37.7 (0.30) | 17.3 (0.25) | 5.1 (0.14) |
| 45–64 years | 100.0 | 33.4 (0.37) | 14.4 (0.22) | 30.4 (0.32) | 16.0 (0.25) | 5.7 (0.16) |
| 65–74 years | 100.0 | 44.7 (0.62) | 13.5 (0.34) | 23.4 (0.45) | 13.6 (0.39) | 4.9 (0.22) |
| 75 years and over | 100.0 | 59.3 (0.66) | 11.2 (0.36) | 16.5 (0.44) | 10.6 (0.39) | 2.4 (0.18) |
| Race | | | | | | |
| One race ² | 100.0 | 35.2 (0.27) | 13.2 (0.14) | 31.1 (0.20) | 15.4 (0.17) | 5.1 (0.09) |
| White | 100.0 | 32.5 (0.30) | 13.1 (0.16) | 32.2 (0.23) | 16.8 (0.19) | 5.5 (0.10) |
| Black or African American | 100.0 | 45.5 (0.54) | 14.1 (0.32) | 26.8 (0.43) | 10.2 (0.31) | 3.4 (0.17) |
| American Indian or Alaska Native | 100.0 | 44.9 (2.18) | 11.9 (1.18) | 26.7 (1.83) | 10.2 (1.06) | 6.3 (1.02) |
| Asian | 100.0 | 50.4 (0.75) | 12.9 (0.47) | 26.7 (0.65) | 8.1 (0.39) | 1.9 (0.21) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 43.8 (4.86) | *11.4 (3.64) | 29.2 (4.39) | 9.7 (2.65) | *6.0 (2.01) |
| Two or more races ³ | 100.0 | 34.8 (1.30) | 13.8 (0.91) | 31.1 (1.26) | 14.3 (0.95) | 6.1 (0.58) |
| Black or African American, white | 100.0 | 30.4 (2.92) | 13.2 (1.79) | 34.5 (3.13) | 13.9 (2.14) | 7.9 (1.62) |
| American Indian or Alaska Native, white | 100.0 | 35.7 (1.96) | 16.9 (1.64) | 27.5 (1.90) | 14.1 (1.59) | 5.8 (0.88) |
| Hispanic or Latino origin⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 42.9 (0.48) | 13.3 (0.29) | 29.2 (0.42) | 11.5 (0.28) | 3.1 (0.14) |
| Mexican or Mexican American | 100.0 | 42.2 (0.60) | 13.3 (0.35) | 28.9 (0.53) | 12.0 (0.35) | 3.5 (0.20) |
| Not Hispanic or Latino | 100.0 | 33.8 (0.30) | 13.2 (0.15) | 31.5 (0.22) | 16.1 (0.18) | 5.4 (0.10) |
| White, single race | 100.0 | 30.3 (0.33) | 13.0 (0.18) | 32.8 (0.25) | 17.8 (0.21) | 6.0 (0.12) |
| Black or African American, single race | 100.0 | 45.7 (0.56) | 14.1 (0.33) | 26.5 (0.44) | 10.2 (0.31) | 3.5 (0.18) |
| American Indian or Alaska Native, single race | 100.0 | 49.2 (2.73) | 9.4 (1.30) | 23.3 (2.21) | 11.4 (1.52) | 6.7 (1.30) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 55.1 (0.52) | 12.3 (0.31) | 18.9 (0.37) | 9.6 (0.29) | 4.2 (0.21) |
| GED or equivalent ⁶ | 100.0 | 41.2 (1.09) | 15.1 (0.76) | 24.4 (0.92) | 12.9 (0.78) | 6.5 (0.53) |
| High school graduate | 100.0 | 39.9 (0.43) | 14.3 (0.27) | 27.5 (0.39) | 13.0 (0.29) | 5.3 (0.18) |
| Some college, no degree | 100.0 | 31.3 (0.47) | 15.9 (0.33) | 32.1 (0.44) | 15.4 (0.34) | 5.3 (0.20) |
| Associate's degree | 100.0 | 29.7 (0.53) | 16.1 (0.39) | 33.8 (0.51) | 15.4 (0.42) | 5.0 (0.31) |
| Bachelor's degree | 100.0 | 23.4 (0.39) | 11.7 (0.29) | 38.5 (0.42) | 20.6 (0.36) | 5.7 (0.20) |
| Master's, professional, or doctoral degree | 100.0 | 23.9 (0.52) | 11.3 (0.38) | 39.0 (0.56) | 21.7 (0.51) | 4.1 (0.22) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 49.9 (0.60) | 13.0 (0.31) | 23.0 (0.41) | 9.3 (0.31) | 4.8 (0.21) |
| 100% to less than 200% of the poverty threshold | 100.0 | 46.4 (0.49) | 14.6 (0.30) | 24.7 (0.36) | 10.1 (0.26) | 4.2 (0.17) |
| 200% to less than 400% of the poverty threshold | 100.0 | 35.9 (0.40) | 14.0 (0.25) | 30.7 (0.32) | 14.5 (0.26) | 4.9 (0.15) |
| 400% of the poverty threshold or greater | 100.0 | 23.3 (0.32) | 11.9 (0.21) | 37.8 (0.31) | 21.2 (0.29) | 5.8 (0.15) |

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Both sexes (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 33.3 (0.34) | 14.0 (0.20) | 32.6 (0.27) | 15.8 (0.22) | 4.3 (0.12) |
| Widowed | 100.0 | 58.6 (0.62) | 12.6 (0.39) | 17.9 (0.45) | 7.8 (0.31) | 3.0 (0.20) |
| Divorced or separated | 100.0 | 34.7 (0.44) | 14.9 (0.30) | 29.4 (0.42) | 14.7 (0.31) | 6.3 (0.22) |
| Never married | 100.0 | 37.6 (0.50) | 10.6 (0.25) | 30.6 (0.38) | 15.4 (0.32) | 5.9 (0.19) |
| Living with a partner | 100.0 | 22.7 (0.60) | 12.8 (0.56) | 36.0 (0.71) | 20.2 (0.55) | 8.2 (0.38) |
| Region | | | | | | |
| Northeast | 100.0 | 31.8 (0.58) | 12.2 (0.34) | 34.3 (0.48) | 16.3 (0.41) | 5.4 (0.22) |
| Midwest | 100.0 | 31.3 (0.56) | 14.3 (0.32) | 33.3 (0.44) | 16.3 (0.35) | 4.8 (0.16) |
| South | 100.0 | 39.8 (0.48) | 13.0 (0.20) | 28.2 (0.32) | 14.0 (0.27) | 5.0 (0.15) |
| West | 100.0 | 34.3 (0.53) | 13.0 (0.25) | 31.3 (0.37) | 16.0 (0.33) | 5.3 (0.18) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 33.0 (0.30) | 13.0 (0.20) | 32.7 (0.25) | 16.2 (0.22) | 5.1 (0.11) |
| Small MSA | 100.0 | 34.9 (0.54) | 13.0 (0.24) | 31.2 (0.39) | 15.6 (0.31) | 5.3 (0.16) |
| Not in MSA | 100.0 | 43.1 (0.91) | 14.2 (0.31) | 25.7 (0.55) | 12.2 (0.41) | 4.8 (0.24) |

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Men | 100.0 | 29.6 (0.31) | 10.0 (0.16) | 32.3 (0.26) | 22.6 (0.25) | 5.5 (0.13) |
| Age | 100.0 | 20.0 (0.01) | 10.0 (0.10) | 02.0 (0.20) | 22.0 (0.20) | 0.0 (0.10) |
| 18–24 years | 100.0 | 37.6 (0.97) | 8.2 (0.51) | 30.0 (0.80) | 18.8 (0.72) | 5.4 (0.38) |
| 25–44 years | 100.0 | 21.2 (0.38) | 9.4 (0.28) | 38.3 (0.43) | 25.5 (0.39) | 5.6 (0.20) |
| 45–64 years | 100.0 | 28.7 (0.46) | 11.0 (0.27) | 31.3 (0.43) | 22.9 (0.39) | 6.1 (0.26) |
| 65–74 years | 100.0 | 38.4 (0.83) | 10.4 (0.46) | 25.4 (0.66) | 20.5 (0.65) | 5.4 (0.35) |
| 75 years and over | 100.0 | 50.0 (1.03) | 10.6 (0.61) | 20.4 (0.76) | 16.5 (0.72) | 2.5 (0.30) |
| Race | | | | | | |
| One race ² | 100.0 | 29.6 (0.31) | 10.0 (0.16) | 32.3 (0.26) | 22.6 (0.25) | 5.5 (0.13) |
| White | 100.0 | 27.6 (0.34) | 9.7 (0.19) | 32.7 (0.29) | 24.2 (0.29) | 5.9 (0.15) |
| Black or African American | 100.0 | 38.9 (0.79) | 10.9 (0.45) | 29.5 (0.70) | 16.6 (0.55) | 4.1 (0.28) |
| American Indian or Alaska Native | 100.0 | 37.0 (2.41) | 10.6 (1.54) | 27.1 (2.28) | 16.4 (2.05) | 8.9 (1.66) |
| Asian | 100.0 | 39.3 (1.01) | 12.6 (0.71) | 33.7 (0.99) | 12.5 (0.70) | 1.9 (0.34) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 29.2 (6.16) | * | 34.2 (6.59) | 17.3 (4.98) | * |
| Two or more races ³ | 100.0 | 31.7 (1.83) | 10.0 (1.43) | 30.6 (1.92) | 21.4 (1.67) | 6.4 (0.85) |
| Black or African American, white | 100.0 | 23.5 (4.08) | 12.2 (2.99) | 35.1 (4.52) | 22.1 (3.55) | 7.1 (1.98) |
| American Indian or Alaska Native, white | 100.0 | 34.4 (2.94) | 12.7 (2.71) | 24.9 (2.89) | 21.2 (2.77) | 6.8 (1.39) |
| Hispanic or Latino origin ⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 32.8 (0.60) | 10.1 (0.37) | 34.1 (0.61) | 18.6 (0.50) | 4.4 (0.24) |
| Mexican or Mexican American | 100.0 | 31.2 (0.74) | 9.9 (0.47) | 34.3 (0.76) | 19.5 (0.64) | 5.1 (0.35) |
| Not Hispanic or Latino | 100.0 | 29.0 (0.35) | 9.9 (0.18) | 32.0 (0.29) | 23.3 (0.28) | 5.7 (0.15) |
| White, single race | 100.0 | 26.4 (0.38) | 9.6 (0.21) | 32.4 (0.33) | 25.3 (0.32) | 6.2 (0.18) |
| Black or African American, single race | 100.0 | 39.3 (0.82) | 10.8 (0.46) | 29.1 (0.72) | 16.6 (0.57) | 4.1 (0.29) |
| American Indian or Alaska Native, single race | 100.0 | 40.4 (3.23) | 9.5 (2.03) | 21.7 (3.18) | 19.3 (3.16) | 9.1 (1.95) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 42.6 (0.71) | 10.3 (0.44) | 24.9 (0.59) | 16.2 (0.53) | 5.9 (0.35) |
| GED or equivalent ⁶ | 100.0 | 35.1 (1.56) | 11.4 (1.03) | 26.6 (1.38) | 18.6 (1.23) | 8.3 (0.85) |
| High school graduate | 100.0 | 32.3 (0.57) | 10.7 (0.35) | 30.2 (0.61) | 19.9 (0.48) | 6.8 (0.29) |
| Some college, no degree | 100.0 | 25.4 (0.63) | 12.2 (0.45) | 33.1 (0.64) | 23.6 (0.60) | 5.8 (0.32) |
| Associate's degree | 100.0 | 24.8 (0.75) | 11.2 (0.52) | 35.2 (0.83) | 23.2 (0.74) | 5.7 (0.58) |
| Bachelor's degree | 100.0 | 20.5 (0.51) | 8.4 (0.37) | 37.7 (0.62) | 28.7 (0.57) | 4.7 (0.28) |
| Master's, professional, or doctoral degree | 100.0 | 22.4 (0.69) | 8.5 (0.46) | 37.2 (0.76) | 29.0 (0.76) | 2.8 (0.27) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 42.2 (0.83) | 10.0 (0.41) | 25.6 (0.62) | 15.7 (0.55) | 6.5 (0.35) |
| 100% to less than 200% of the poverty threshold | 100.0 | 38.9 (0.67) | 11.4 (0.40) | 27.6 (0.58) | 16.5 (0.46) | 5.6 (0.29) |
| 200% to less than 400% of the poverty threshold | 100.0 | 30.8 (0.53) | 10.7 (0.31) | 31.5 (0.47) | 21.5 (0.42) | 5.5 (0.22) |
| 400% of the poverty threshold or greater | 100.0 | 20.6 (0.41) | 8.8 (0.27) | 37.1 (0.44) | 28.3 (0.43) | 5.2 (0.22) |

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---------------------------------|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Men (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 28.5 (0.38) | 10.9 (0.24) | 34.1 (0.36) | 22.3 (0.33) | 4.2 (0.17) |
| Widowed | 100.0 | 47.7 (1.24) | 10.1 (0.73) | 21.4 (1.03) | 16.3 (0.86) | 4.4 (0.49) |
| Divorced or separated | 100.0 | 28.6 (0.63) | 10.0 (0.40) | 28.8 (0.63) | 23.8 (0.61) | 8.9 (0.40) |
| Never married | 100.0 | 33.9 (0.63) | 8.4 (0.32) | 30.0 (0.54) | 21.4 (0.47) | 6.3 (0.27) |
| Living with a partner | 100.0 | 18.3 (0.75) | 7.9 (0.52) | 35.1 (1.05) | 29.6 (0.95) | 9.0 (0.55) |
| Region | | | | | | |
| Northeast | 100.0 | 27.4 (0.77) | 9.6 (0.43) | 34.7 (0.67) | 23.2 (0.62) | 5.1 (0.30) |
| Midwest | 100.0 | 26.8 (0.63) | 10.3 (0.38) | 33.5 (0.57) | 24.0 (0.51) | 5.3 (0.25) |
| South | 100.0 | 33.2 (0.56) | 10.0 (0.24) | 30.0 (0.42) | 20.8 (0.43) | 6.0 (0.24) |
| West | 100.0 | 28.5 (0.55) | 9.9 (0.33) | 32.9 (0.49) | 23.5 (0.50) | 5.3 (0.25) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 27.7 (0.37) | 9.9 (0.24) | 33.9 (0.36) | 23.4 (0.33) | 5.1 (0.18) |
| Small MSA | 100.0 | 29.3 (0.59) | 9.5 (0.29) | 32.0 (0.46) | 23.2 (0.48) | 6.0 (0.23) |
| Not in MSA | 100.0 | 36.9 (0.94) | 11.1 (0.39) | 27.3 (0.67) | 18.8 (0.65) | 5.9 (0.35) |

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011-2014

| Selected characteristic | All adults aged 18 and over | Nondrinker ¹ | Infrequent drinker ¹ | Light drinker ¹ | Moderate drinker ¹ | Heavier drinker ¹ |
|---|-----------------------------------|-------------------------|------------------------------------|----------------------------|----------------------------------|---------------------------------|
| Women | 100.0 | 40.3 (0.33) | 16.2 (0.19) | 30.1 (0.25) | 8.7 (0.15) | 4.7 (0.11) |
| Age | 100.0 | 40.3 (0.33) | 10.2 (0.19) | 30.1 (0.23) | 0.7 (0.13) | 4.7 (0.11) |
| 18–24 years | 100.0 | 44.5 (0.96) | 12.1 (0.61) | 31.1 (0.79) | 7.2 (0.41) | 5.0 (0.37) |
| 25–44 years | 100.0 | 31.6 (0.43) | 17.3 (0.31) | 37.1 (0.40) | 9.3 (0.24) | 4.7 (0.18) |
| 45–64 years | 100.0 | 37.8 (0.46) | 17.7 (0.33) | 29.6 (0.40) | 9.6 (0.26) | 5.3 (0.19) |
| 65–74 years | 100.0 | 50.1 (0.73) | 16.2 (0.50) | 21.6 (0.57) | 7.6 (0.37) | 4.5 (0.27) |
| 75 years and over | 100.0 | 65.6 (0.75) | 11.7 (0.45) | 13.8 (0.49) | 6.5 (0.36) | 2.4 (0.21) |
| Race | | | | | | |
| One race ² | 100.0 | 40.3 (0.34) | 16.2 (0.20) | 30.1 (0.25) | 8.8 (0.15) | 4.7 (0.11) |
| White | 100.0 | 37.1 (0.39) | 16.3 (0.23) | 31.7 (0.30) | 9.7 (0.18) | 5.2 (0.13) |
| Black or African American | 100.0 | 50.9 (0.66) | 16.7 (0.45) | 24.6 (0.53) | 5.0 (0.25) | 2.9 (0.20) |
| American Indian or Alaska Native | 100.0 | 52.2 (2.91) | 13.1 (1.78) | 26.3 (2.51) | 4.5 (1.06) | 3.9 (0.89) |
| Asian | 100.0 | 60.2 (1.01) | 13.2 (0.65) | 20.6 (0.80) | 4.3 (0.39) | 1.8 (0.26) |
| Native Hawaiian or Other Pacific Islander | 100.0 | 57.8 (5.59) | *10.1 (3.23) | 24.5 (5.09) | * | *5.3 (2.16) |
| Two or more races ³ | 100.0 | 37.5 (1.78) | 17.2 (1.25) | 31.5 (1.66) | 8.0 (0.97) | 5.8 (0.82) |
| Black or African American, white | 100.0 | 34.9 (3.93) | 13.9 (2.26) | 34.1 (4.18) | 8.6 (2.51) | 8.5 (2.35) |
| American Indian or Alaska Native, white | 100.0 | 36.9 (2.54) | 20.9 (2.01) | 30.0 (2.51) | 7.4 (1.46) | 4.8 (1.19) |
| Hispanic or Latino origin ⁴ and race | | | | | | |
| Hispanic or Latino | 100.0 | 53.2 (0.63) | 16.6 (0.44) | 24.2 (0.51) | 4.3 (0.23) | 1.7 (0.14) |
| Mexican or Mexican American | 100.0 | 53.7 (0.80) | 16.9 (0.56) | 23.4 (0.64) | 4.1 (0.29) | 1.9 (0.19) |
| Not Hispanic or Latino | 100.0 | 38.2 (0.36) | 16.1 (0.21) | 31.0 (0.27) | 9.5 (0.17) | 5.2 (0.12) |
| White, single race | 100.0 | 33.8 (0.42) | 16.3 (0.25) | 33.2 (0.33) | 10.8 (0.21) | 5.9 (0.15) |
| Black or African American, single race | 100.0 | 51.0 (0.68) | 16.8 (0.46) | 24.3 (0.54) | 5.0 (0.26) | 2.9 (0.21) |
| American Indian or Alaska Native, single race | 100.0 | 56.3 (3.44) | 9.3 (1.82) | 24.6 (2.98) | 5.0 (1.41) | 4.7 (1.22) |
| Education ⁵ | | | | | | |
| Less than high school diploma | 100.0 | 66.9 (0.64) | 14.1 (0.46) | 13.1 (0.44) | 3.3 (0.23) | 2.5 (0.21) |
| GED or equivalent ⁶ | 100.0 | 48.3 (1.50) | 19.4 (1.20) | 21.8 (1.20) | 6.2 (0.80) | 4.4 (0.64) |
| High school graduate | 100.0 | 47.0 (0.57) | 17.6 (0.41) | 25.0 (0.49) | 6.5 (0.26) | 3.9 (0.21) |
| Some college, no degree | 100.0 | 36.3 (0.62) | 19.0 (0.46) | 31.3 (0.58) | 8.5 (0.33) | 4.9 (0.24) |
| Associate's degree | 100.0 | 33.7 (0.69) | 20.0 (0.59) | 32.6 (0.65) | 9.2 (0.42) | 4.4 (0.29) |
| Bachelor's degree | 100.0 | 26.1 (0.53) | 14.7 (0.43) | 39.3 (0.58) | 13.2 (0.40) | 6.7 (0.30) |
| Master's, professional, or doctoral degree | 100.0 | 25.4 (0.67) | 14.1 (0.56) | 40.8 (0.78) | 14.4 (0.57) | 5.3 (0.34) |
| Poverty status ⁷ | | | | | | |
| Less than 100% of the poverty threshold | 100.0 | 55.5 (0.67) | 15.2 (0.41) | 21.1 (0.49) | 4.6 (0.26) | 3.5 (0.23) |
| 100% to less than 200% of the poverty threshold | 100.0 | 52.5 (0.62) | 17.2 (0.42) | 22.4 (0.46) | 4.8 (0.22) | 3.1 (0.17) |
| 200% to less than 400% of the poverty threshold | 100.0 | 40.7 (0.51) | 17.2 (0.37) | 30.0 (0.43) | 7.7 (0.26) | 4.3 (0.19) |
| 400% of the poverty threshold or greater | 100.0 | 26.2 (0.44) | 15.1 (0.32) | 38.5 (0.43) | 13.7 (0.31) | 6.5 (0.21) |

Table ALC-2c. Crude percent distribution (with standard errors) of current alcohol drinking status among adults aged 18 and over, by sex and selected characteristics: United States, annualized, 2011–2014

| | All adults aged 18 | 1 | Infrequent | 1 | Moderate | Heavier |
|---------------------------------|-----------------------|-------------------------|----------------------|----------------------------|----------------------|----------------------|
| Selected characteristic | and over | Nondrinker ¹ | drinker ¹ | Light drinker ¹ | drinker ¹ | drinker ¹ |
| Women (continued) | | | | | | |
| Marital status | | | | | | |
| Married | 100.0 | 38.2 (0.44) | 17.2 (0.29) | 31.1 (0.36) | 9.1 (0.22) | 4.5 (0.15) |
| Widowed | 100.0 | 61.6 (0.69) | 13.3 (0.44) | 17.0 (0.49) | 5.5 (0.30) | 2.6 (0.20) |
| Divorced or separated | 100.0 | 39.0 (0.57) | 18.3 (0.44) | 29.9 (0.55) | 8.3 (0.31) | 4.5 (0.22) |
| Never married | 100.0 | 41.7 (0.67) | 13.0 (0.38) | 31.2 (0.53) | 8.6 (0.33) | 5.4 (0.26) |
| Living with a partner | 100.0 | 26.8 (0.90) | 17.6 (0.86) | 37.0 (0.91) | 11.2 (0.59) | 7.4 (0.52) |
| Region | | | | | | |
| Northeast | 100.0 | 35.9 (0.71) | 14.6 (0.45) | 33.9 (0.60) | 9.9 (0.40) | 5.7 (0.32) |
| Midwest | 100.0 | 35.4 (0.72) | 18.2 (0.46) | 33.1 (0.58) | 9.0 (0.31) | 4.3 (0.20) |
| South | 100.0 | 45.7 (0.56) | 15.8 (0.32) | 26.5 (0.39) | 7.9 (0.24) | 4.1 (0.16) |
| West | 100.0 | 39.8 (0.67) | 16.0 (0.36) | 29.9 (0.51) | 8.9 (0.30) | 5.3 (0.23) |
| Place of residence ⁸ | | | | | | |
| Large MSA | 100.0 | 38.0 (0.39) | 15.9 (0.28) | 31.6 (0.33) | 9.5 (0.22) | 5.0 (0.16) |
| Small MSA | 100.0 | 40.1 (0.65) | 16.3 (0.34) | 30.5 (0.50) | 8.6 (0.27) | 4.6 (0.20) |
| Not in MSA | 100.0 | 48.6 (1.04) | 17.0 (0.48) | 24.2 (0.65) | 6.3 (0.29) | 3.9 (0.23) |

^{*} Estimates are considered unreliable. Data preceded by an asterisk have a relative standard error (RSE) greater than 30% and less than or equal to 50% and should be used with caution. Data not shown have an RSE greater than 50%.

¹Derived from several source questions: "In any one year, have you had at least 12 drinks of any type of alcoholic beverage?", "In your entire life, have you had at least 12 drinks of any type of alcoholic beverage?", "In the past year, how often did you drink any type of alcoholic beverage?" and "In the past year, on those days that you drank alcoholic beverages, on the average, how many drinks did you have?" A nondrinker had zero drinks in the past year (includes former drinkers and lifetime abstainers). An infrequent drinker had at least 12 drinks in his or her lifetime and 1–11 drinks in the past year. A light drinker had 3 drinks or less per week, on average, in the past year. A moderate drinker had more than 3 and up to and including 14 drinks per week for men, and more than 3 and up to and including 7 drinks per week for women, on average, in the past year. A heavier drinker had more than 14 drinks per week for men, and more than 7 drinks per week for women, on average, in the past year.

²Refers to persons who indicated only a single race group, including those of Hispanic or Latino origin.

³Refers to persons who indicated more than one race group, including those of Hispanic or Latino origin. Only two combinations of multiple race groups are shown due to small sample sizes for other combinations.

⁴Refers to persons who are of Hispanic or Latino origin and may be of any race or combination of races. "Not Hispanic or Latino" refers to persons who are not of Hispanic or Latino origin, regardless of race.

⁵Shown only for adults aged 25 and over.

⁶GED is General Educational Development high school equivalency diploma.

⁷Based on family income and family size using the U.S. Census Bureau poverty thresholds for 2010, 2011, 2012, and 2013.

⁸MSA is metropolitan statistical area. Large MSAs have a population of 1 million or more; small MSAs have a population of less than 1 million. "Not in MSA" consists of persons not living in a metropolitan statistical area.

NOTES: Estimates are based on household interviews of a sample of the civilian noninstitutionalized population. This table is based on data from the Sample Adult file and estimates were weighted using the Sample Adult weight. Unknowns for the columns were not included in the denominators when calculating percentages. Percentages may not add to totals due to rounding. "Both sexes," "men," and "women" include persons of other races and unknown race and ethnicity, unknown education, and unknown marital status. For more information on the data source, methods, and definitions used for this table, refer to Technical Notes for Summary Health Statistics Tables: National Health Interview Survey (available from http://www.cdc.gov/nchs/nhis/SHS/tables.htm). Suggested citation: Villarroel MA, Clarke TC, Schoenborn CA. Tables of adult health behaviors, alcohol use: National Health Interview Survey, 2011–2014. 2016. Available from: http://www.cdc.gov/nchs/nhis/SHS/tables.htm.

SOURCE: NCHS, National Health Interview Survey, 2011–2014.