

Advance Data No. 350 • March 9, 2005

Table 2-Supplement. Number of women 15–44 years of age who have ever had sexual intercourse and percent who have ever used the specified contraceptive method, by new race and Hispanic origin classification: United States, 2002

Method	Hispanic origin and race		
	Hispanic or Latina	Not Hispanic or Latina	
		White, single race	Black or African American, single race
		Number in thousands	
All women . . . . .	7,887	34,999	7,403
		Percent (with standard errors) who have ever used specified method	
Any method . . . . .	96.5 (0.7)	98.8 (0.2)	97.3 (0.6)
Female sterilization . . . . .	24.1 (1.8)	19.2 (1.0)	26.9 (1.6)
Male sterilization . . . . .	4.4 (0.5)	16.7(1.0)	5.0 (0.9)
Pill . . . . .	68.5 (1.8)	87.0 (0.8)	79.8 (1.5)
Implant . . . . .	4.0 (0.6)	1.4 (0.2)	3.3 (0.7)
1-Month Injectable (Lunelle) . . . . .	3.1 (0.4)	0.5 (0.1)	1.0 (0.4)
3-Month Injectable (Depo-Provera) . . . . .	24.3 (1.8)	13.6 (0.9)	23.3 (1.5)
Emergency contraception . . . . .	3.8 (0.5)	4.2 (0.4)	4.1 (0.8)
Contraceptive patch . . . . .	1.1 (0.3)	0.7 (0.2)	1.4 (0.4)
Today™ Sponge . . . . .	2.7 (0.4)	8.8 (0.6)	6.3 (0.8)
Intrauterine device (IUD) . . . . .	10.0 (0.9)	4.7 (0.5)	5.4 (0.8)
Diaphragm . . . . .	3.4 (0.6)	10.4 (0.7)	6.9 (0.9)
Condom . . . . .	78.2 (1.7)	92.2 (0.7)	91.8 (1.2)
Female condom . . . . .	1.4 (0.3)	1.2 (0.2)	5.4 (0.7)
Periodic abstinence — calendar rhythm . . . . .	16.2 (1.1)	16.6 (0.8)	13.7 (1.4)
Periodic abstinence — natural family planning . . . . .	4.7 (0.7)	3.6 (0.4)	1.7 (0.4)
Withdrawal . . . . .	47.2 (1.9)	59.7 (1.4)	50.5 (1.8)
Foam alone . . . . .	8.2 (0.7)	12.5 (0.6)	15.5 (1.3)
Jelly/cream alone . . . . .	3.4 (0.5)	7.7 (0.5)	9.6 (0.8)
Suppository/insert . . . . .	4.5 (0.5)	7.9 (0.6)	8.8 (0.8)
Other methods . . . . .	1.3 (0.3)	0.8 (0.2)	1.3 (0.5)

## Advance Data No. 350 • March 9, 2005

**Table 3-Supplement. Number of women 15–44 years of age who have ever had premarital sexual intercourse and percent who used the specified contraceptive method at first intercourse, by new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	Used any method	Percent			
			Pill	Condom	Withdrawal	All other methods
All women <sup>1</sup> . . . . .	48,649	63.0	17.8	45.0	7.7	3.4
Hispanic origin and race						
Hispanic or Latina . . . . .	6,373	46.2	9.5	33.1	6.1	4.4
Not Hispanic or Latina:						
White, single race . . . . .	31,946	67.5	18.6	48.1	8.7	3.2
Black or African American, single race . . . . .	7,251	60.1	22.5	43.8	4.5	2.9

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

NOTE: This table refers to first sexual intercourse after menarche.

**Advance Data No. 350 • March 22, 2005**

**Table 7-Supplement. Number of women 15–44 years of age and percent distribution by current contraceptive status and specific method, according to new race and Hispanic origin classification: United States, 2002**

Contraceptive status and method	All Women <sup>1</sup>	Hispanic origin and race		
		Hispanic or Latina	Not Hispanic or Latina	
			White, single race	Black or African American, single race
		Number in thousands		
All women . . . . .	61,561	9,107	39,498	8,250
		Percent distribution		
Total . . . . .	100.0	100.0	100.0	100.0
Using contraception (contraceptors) . . . . .	61.9	59.0	64.6	57.6
Female sterilization . . . . .	16.7	19.9	15.4	22.6
Male sterilization . . . . .	5.7	2.6	7.6	1.3
Pill . . . . .	18.9	13.0	22.2	13.1
Implant, Lunelle or Patch . . . . .	0.8	1.8	0.5	0.6
3-month Injectable (Depo-Provera) . . . . .	3.3	4.3	2.7	5.4
Intrauterine device (IUD) . . . . .	1.3	3.2	1.0	0.8
Diaphragm . . . . .	0.2	–	0.2	0.1
Condom . . . . .	11.1	10.9	10.7	11.4
Periodic abstinence — calendar rhythm . . . . .	0.7	0.6	0.8	0.3
Periodic abstinence — natural family planning . . . . .	0.2	0.3	0.3	0.1
Withdrawal . . . . .	2.5	2.2	2.6	1.5
Other methods <sup>2</sup> . . . . .	0.6	0.3	0.7	0.5
Not using contraception . . . . .	38.1	41.0	35.4	42.4
Surgically sterile-female (noncontraceptive) . . . . .	1.5	0.9	1.6	1.6
Nonsurgically sterile-female or male . . . . .	1.6	1.7	1.7	1.4
Pregnant or postpartum . . . . .	5.3	6.9	4.7	5.7
Seeking pregnancy . . . . .	4.2	5.2	4.0	4.3
Other nonuse:				
Never had intercourse or no sex in 3 months before interview . . . . .	18.1	18.7	17.0	19.0
Had intercourse in 3 months before interview . . . . .	7.4	7.7	6.5	10.4
All other nonusers <sup>3</sup> . . . . .	0.0	–	0.1	0.1

0.0=Quantity greater than zero but less than 0.05.

– Quantity zero.

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

<sup>2</sup>Includes Today Sponge, Cervical Cap, Female Condom, and other methods not shown separately.

<sup>3</sup>Includes male sterility of unknown origin and other small groups, not shown separately.

NOTE: Percents may not add to 100 due to rounding.

## Advance Data No. 350 • March 35, 2005

**Table 9-Supplement. Number of women 15–44 years of age, number at risk of unintended pregnancy, and percent of women currently using a method of contraception, by new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	Percent currently using a method	Number at risk of unintended pregnancy	Percent currently using a method
All women <sup>1</sup> . . . . .	61,561	61.9	42,683	89.3
Hispanic origin and race				
Hispanic or Latina . . . . .	9,107	59.0	6,075	88.4
Not Hispanic or Latina:				
White, single race . . . . .	39,498	64.6	28,079	90.9
Black or African American, single race . . . . .	8,250	57.6	5,611	84.7

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

NOTE: "At risk of unintended pregnancy" is defined as codes 1–21 and 42 on CONSTAT1, the recode for current contraceptive status. These codes represent all current contraceptors plus women who have had sex in the last 3 months but are not current contraceptors.

**Advance Data No. 350 • March 9, 2005**

**Table 10-Supplement. Number of women 15–44 years of age who are currently using a method of contraception and percent distribution by method, according to new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	Using any method	Sterilization		Pill	Condom	3-month Injectable	Other methods
			Female	Male				
Percent distribution								
All women <sup>1</sup> . . . . .	38,109	100.0	27.0	9.2	30.6	18.0	5.3	9.9
Hispanic origin and race								
Hispanic or Latina . . . . .	5,371	100.0	33.8	4.4	22.0	18.5	7.3	14.1
Not Hispanic or Latina:								
White, single race . . . . .	25,513	100.0	23.9	11.7	34.4	16.6	4.2	9.3
Black or African American, single race . . . . .	4,754	100.0	39.2	2.3	22.7	19.8	9.4	6.6

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

## Advance Data No. 350 • March 35, 2005

**Table 12-Supplement. Number of women 15–44 years of age, percent currently using contraception, and percent who used each of the specified contraceptive methods in the month of interview, according to new race and Hispanic origin classification: United States, 2002**

Contraceptive status and method	All women <sup>1</sup>	Hispanic origin and race		
		Hispanic or Latina	Not Hispanic or Latina	
			White, single race	Black or African American, single race
		Number in thousands		
All women . . . . .	61,561	9,107	39,498	8,250
		Percent		
Currently using contraception . . . . .	61.9	59.0	64.6	57.6
Female sterilization . . . . .	16.7	19.9	15.4	22.6
Male sterilization . . . . .	6.3	2.8	8.3	1.5
Pill . . . . .	19.2	13.0	22.5	13.3
Implant, Lunelle, or Patch . . . . .	0.8	1.8	0.5	0.6
3-month Injectable (Depo-Provera) . . . . .	3.3	4.3	2.7	5.4
Intrauterine device (IUD) . . . . .	1.3	3.2	1.1	0.8
Diaphragm . . . . .	0.4	–	0.6	0.1
Condom . . . . .	14.7	14.2	14.0	17.0
Periodic abstinence—calendar . . . . .	1.3	0.8	1.5	0.3
Periodic abstinence—temperature . . . . .	0.2	0.3	0.3	0.1
Withdrawal . . . . .	5.4	3.7	6.2	2.8
Other methods . . . . .	1.0	0.7	1.1	1.1

– Quantity zero.

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

NOTE: Percents may not add to the total who were using contraception because more than one method could have been used in the month of interview. Up to four methods were coded in recodes CONSTAT1–4.

## Advance Data No. 350 • March 35, 2005

**Table 14-Supplement. Number of women 15–44 years of age who are using contraception in the month of interview and percent distribution by specified method or method combination, according to new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	Total	Condom only	Condom and pill	Condom and withdrawal	Condom and other methods	All other users	Percent using condom at all
Percent distribution								
All women <sup>1</sup> . . . . .	38,109	100.0	14.3	4.1	2.7	2.7	76.2	23.8
Hispanic origin and race								
Hispanic or Latina . . . . .	5,370	100.0	15.5	2.4	2.0	4.2	75.9	24.1
Not Hispanic or Latina:								
White, single race . . . . .	25,513	100.0	12.7	4.3	3.0	1.7	78.3	21.7
Black or African American, single race . . . . .	4,754	100.0	17.7	4.6	1.0	6.2	70.5	29.6

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

NOTE: Percents may not add to 100 due to rounding.

Advance Data No. 350 • March 22, 2005

Table 15-Supplement. Number of women 15–44 years of age, percent who received at least one family planning service from a medical care provider in the 12 months prior to interview, and percent who received specified family planning services, by new Hispanic origin and race classification: United States, 2002

Characteristic	Services received in past 12 months						
	Number in thousands	At least one family planning service	Birth control method	Birth control counseling	Birth control checkup or test	Sterilization counseling	Sterilization operation
All women <sup>1</sup> . . . . .	61,561	41.7	33.9	18.6	23.6	4.4	1.9
Hispanic origin and race							
Hispanic or Latina . . . . .	9,107	39.7	28.9	22.6	20.6	7.0	2.3
Not Hispanic or Latina:							
White, single race . . . . .	39,498	43.1	36.4	17.3	25.4	3.6	1.7
Black or African American, single race . . . . .	8,250	39.4	30.6	20.6	21.8	5.1	2.1

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.



## Advance Data No. 350 • March 35, 2005

**Table 16-Supplement. Number of women 15–44 years of age and percent who received the specified medical services from a medical care provider in the 12 months prior to interview, by new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	At least one medical service	Pregnancy test	Pap smear	Pelvic exam	Counseling/ test/treatment for STD
		Percent				
All women <sup>1</sup> . . . . .	61,561	69.1	19.7	64.4	59.7	12.6
Hispanic origin and race						
Hispanic or Latina . . . . .	9,107	63.4	24.3	57.1	48.4	12.5
Not Hispanic or Latina:						
White, single race . . . . .	39,498	70.1	17.3	66.1	63.2	11.8
Black or African American, single race . . . . .	8,250	74.4	23.7	69.1	58.1	15.7

<sup>1</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

**Advance Data No. 350 • March 35, 2005**

**Table 17-Supplement. Number of women 15–44 years of age, percent who received at least one family planning or medical service in the 12 months prior to interview, and percent who used the specified type of provider, by new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	At least one family planning or medical service	Type of provider			
			Clinic		Private doctor or HMO <sup>1</sup>	Other <sup>2</sup>
			Total	Title X		
Percent						
All women <sup>2</sup> . . . . .	61,561	72.7	21.9	8.4	56.0	2.0
Hispanic origin and race						
Hispanic or Latina . . . . .	9,107	68.6	34.5	13.5	37.7	2.0
Not Hispanic or Latina:						
White, single race . . . . .	39,498	73.3	16.7	6.5	61.9	1.8
Black or African American, single race . . . . .	8,250	77.1	30.6	14.3	52.6	1.9

<sup>1</sup>Other sources" may include hospitals, school or university health centers, military health facilities, and other sources.

<sup>2</sup>Includes women of other, unknown, or multiple race and origin groups, not shown separately.

**Advance Data No. 350 • March 35, 2005**

**Table 18-Supplement. Number of women 15–44 years of age, number who received at least one family planning or medical service from the specified type of provider in the 12 months prior to interview, and percent distribution by new Hispanic origin and race classification: United States, 2002**

Characteristic	Number in thousands	At least one family planning or medical service	Type of provider			
			Clinic		Private doctor or HMO <sup>1</sup>	Other <sup>2</sup>
			Total	Title X		
Number in thousands						
All women <sup>1</sup> . . . . .	61,561	44,747	13,453	5,393	34,444	1,214
Percent distribution						
Total. . . . .	61,561	100.0	100.0	100.0	100.0	100.0
Hispanic or Latina . . . . .	9,107	14.0	23.4	22.7	10.0	15.0
Not Hispanic or Latina:						
White, single race . . . . .	39,498	64.7	49.1	47.8	71.0	58.9
Black or African American, single race . . . . .	8,250	14.2	18.7	21.9	12.6	13.2

<sup>1</sup> Includes women of other, unknown, or multiple race and origin groups, not shown separately.

Note: Percents may not add to 100 due to rounding.