| List of Chartbook Figures Population | | | Morbidity and Activity Limitation | | | |
|---------------------------------------|---|----------|--|--|----|--|
| | | 15. | Three or more chronic conditions among adults 45 years of age and over, by age and percent of poverty level: United States, 2004 | 43 | | |
| 1. 2. | Total population and older population: United States, 1950–2050 | 17 | 16. | Selected chronic health conditions causing limitation of activity among children, by age: United States, | | |
| 3. | United States, 1990 and 2000 Population in selected race and Hispanic origin | 19 | 17. | 2003–2004 | 45 | |
| 4. 5. | groups, by age: United States, 1980–2005 Poverty by age: United States, 1966–2004 Low income by age, race, and Hispanic origin: | 21 23 | 18. | United States, 2003–2004 | 47 | |
| | United States, 2004 | 25 | | limitation of activity among older adults, by age: United States, 2003–2004 | 49 | |
| Health Insurance and Expenditures | | | Hea | alth Care Utilization | | |
| 6. | Health insurance coverage at the time of interview among persons under 65 years of age: United States, 1984–2004 | 27 | 19. | Dental visit in the past year among children, by age, race and Hispanic origin: United States, 2004 | 51 | |
| 7. | Uninsured for at least part of the 12 months prior to interview among persons under 65 years of | 2, | 20. | Influenza vaccination during the past year among adults 50 years of age and over, by race and Hispanic origin: United States, 2003–2004 | 53 | |
| | age, by length of time uninsured and selected characteristics: United States, 2004 | 29 | 21. | Emergency department visits for falls, by sex and age: United States, 2002–2004 | 55 | |
| 8. | National health expenditures as a percentage of Gross Domestic Product: United States, 1960–2004 | 30 | 22. | Visits to physician offices and hospital outpatient departments, by sex and age: United States, | | |
| 9. | Personal health care expenditures, by source of funds and type of expenditures: United States, | 30 | 23. | 1997–2004 | 57 | |
| | 2004 | 31 | | 18–64 years of age with obesity, by sex and age: United States, 1999–2004 | 59 | |
| Hea | Ith Risk Factors | | | | | |
| 10. | Cigarette smoking among men, women, high | | Мо | rtality | | |
| 11. | school students, and mothers during pregnancy: United States, 1965–2005 | 33 | 24. 25. | Life expectancy at birth and at 65 years of age, by sex: United States, 1900–2003 | 61 | |
| | high school students, by sex, grade level, and year: United States, 1991, 1993, 2003, and 2005 | 35 | | United States, 1950–2003 | 63 | |
| 12. | Regular leisure-time physical activity among adults 18 years of age and over, by percent of | 33 | 26. 27. | Infant mortality rates, by detailed race and Hispanic origin of mother: United States, 2001–2003 Death rates for leading causes of death for all ages: | 65 | |
| | poverty level, race and Hispanic origin: | 37 | 21. | United States, 1950–2003 | 67 | |
| 13. | United States, 2004 Overweight and obesity, by age: United States, | 31 | | | | |
| 14. | 1960–2004 | 39 | | | | |
| | 1971–2002 | 41 | | | | |

Health, United States, 2006 xiii

Special Feature: Pain

| 28. | Adults 20 years of age and over reporting pain in the | Э |
|-----|--|----|
| | month prior to interview, by selected characteristics: | |
| | United States, 1999–2002 | 69 |
| 29. | Duration of pain among adults reporting pain, by | |
| | age: United States, 1999–2002 | 71 |
| 30. | Adults reporting low back pain, migraine, neck, and | |
| | face pain in the 3 months prior to interview, by age: | |
| | United States, 2004 | 73 |
| 31. | Adults 18 years of age and over reporting low back | |
| | pain in the 3 months prior to interview, by selected | |
| | characteristics: United States, 2004 | 75 |
| 32. | Adults 18 years of age and over reporting joint | |
| | pain in the 30 days prior to interview, by | |
| | severity level and selected characteristics: | |
| 00 | United States, 2003 | 77 |
| 33. | Narcotic analgesic drug visits to the emergency department among visits with a severe pain level | |
| | recorded, by age, sex, and race: United States, | |
| | 2003–2004 | 79 |
| 34. | Adults 18 years of age and over reporting narcotic | 73 |
| 04. | drug use in the month prior to interview, by sex | |
| | and age: United States, 1988–1994 and | |
| | 1999–2002 | 81 |
| 35. | Hospital discharges for knee and nonfracture hip | |
| | replacement surgery among adults 45 years of | |
| | age and over, by sex and age: United States, | |
| | 1992–1993 and 2003–2004 | 83 |
| 36. | Adults reporting ambulatory medical care use for | |
| | headaches and associated ambulatory medical | |
| | care expenses, by age and sex: United States, | |
| | 2002–2003 | 85 |
| 37. | Health status measures among adults 18 years of | |
| | age and over with and without low back pain, by | |
| | age: United States, 2004 | 87 |

xiv Health, United States, 2006