

User Guide to the 2019 Fetal Death Public Use File



2019 Fetal Death Data Set

User's Guide

ACKNOWLEDGMENTS

Preparation of this user's guide was coordinated by Elizabeth C.W. Gregory and Claudia P. Valenzuela of the Division of Vital Statistics (DVS) under the general direction of Joyce A. Martin, Lead Statistician, Reproductive Statistics Branch (RSB). The vital statistics computer file on which it is based was prepared by DVS staff.

The Division of Vital Statistics Director, Steven Schwartz, manages the Vital Statistics Cooperative Program, through which the vital registration offices of all states, the District of Columbia, New York City, Puerto Rico, Virgin Islands, Guam, American Samoa, and the Commonwealth of the Northern Mariana Islands, provided the data to the National Center for Health Statistics (NCHS). This Division, and its Data Acquisition, Classification, and Evaluation Branch (DACEB), and Information Technology Branch (ITB), processed computer edits, designed and programmed the tabulations, reviewed the data, prepared documentation for this publication, and were responsible for receipt and processing of the basic data file.

NCHS acknowledges the essential role of the vital registration offices of all states and territories in maintaining the system through which vital statistics data are obtained and their cooperation in providing the information on which this publication is based.

Questions on the documentation or substantive questions concerning the data should be directed to the Reproductive Statistics Branch, Division of Vital Statistics, NCHS, 3311 Toledo Road, Hyattsville, MD 20782-2003 (1-800) 232-4636.

Department of Health and Human Services
Centers for Disease Control and Prevention
National Center for Health Statistics
Division of Vital Statistics

2019 Fetal Death Data Set

Contents

Introduction

Control count of records

List of data elements and locations

Record layout and definition of item and codes

Documentation tables

Figure 1. U.S. Standard Report of fetal Death

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2019

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2019

External links

Fetal and Perinatal Mortality: United States, 2013

http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64_08.pdf

Cause of fetal death: Data from the fetal death report, 2014

https://www.cdc.gov/nchs/data/nvsr/nvsr65/nvsr65_07.pdf

Cause-of-death Data From the Fetal Death File, 2015–2017

<https://www.cdc.gov/nchs/data/nvsr/nvsr69/nvsr69-04-508.pdf>

Births: Final Data for 2019 (Forthcoming)

User Guide to the 2019 Period Linked Birth/Infant Death Public Use File (Forthcoming)

Introduction

This User Guide is for use with the 2019 fetal death public use data. The 2019 fetal death micro-data file may be downloaded at:

http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm [[1]. The micro-data fetal death file for the U.S. does not include geographic detail (e.g., state or county of birth). The territories file contains selected geographic detail. Selected fetal death data for 2005-2018, including some geographic data and cause of death data, are available in CDC WONDER (<http://wonder.cdc.gov>), an interactive online data access tool. Selected fetal death data for 2019 are expected to be available in WONDER in early 2021.

Definition of fetal death

“Fetal death” means death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy and which is not an induced termination of pregnancy. The death is indicated by the fact that after such expulsion or extraction, the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. Heartbeats are to be distinguished from transient cardiac contractions; respirations are to be distinguished from fleeting respiratory efforts or gasps.

This definition [2] has been adopted by the Centers for Disease Control and Prevention’s National Center for Health Statistics (NCHS) as the nationally recommended standard and is based on the definition published by the World Health Organization in 1950 and revised in 1988. The term “fetal death” is defined on an all-inclusive basis to end confusion arising from the use of such terms as stillbirth, spontaneous abortion, and miscarriage. All U.S. states and registration areas have definitions similar to the standard definition, except for Puerto Rico and Wisconsin,

which have no formal definition [3]. Fetal deaths do not include induced terminations of pregnancy.

Reporting requirements for fetal death data

Reporting requirements for fetal deaths vary by state, and these differences have implications for comparisons of fetal and perinatal mortality rates by state [2]. Table A shows the period of gestation at which fetal death reporting is required for each reporting area for 2019. The majority of states require reporting of fetal deaths at 20 weeks of gestation or more, or a minimum of 350 grams birthweight (roughly equivalent to 20 weeks), or some combination of the two. Six states (Colorado, Georgia, Hawaii, New York, Rhode Island, and Virginia), American Samoa, and the U.S. Virgin Islands require reporting of fetal deaths at all periods of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), two states require reporting beginning at 12 weeks of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), and one state requires reporting beginning at 16 weeks of gestation. At the other end of the spectrum, one state requires reporting of fetal deaths with birthweights of 500 grams or more (roughly equivalent to 22 weeks of gestation).

Classification of fetal deaths at 20 weeks of gestation or greater

To maximize the comparability of data by year and by state while including as much meaningful data as possible, most tables published by NCHS are based on fetal deaths occurring at gestations of 20 weeks or more. These tabulations also include fetal deaths for which gestation is not stated for those states requiring reporting at 20 weeks of gestation or more only. Fetal deaths of not-stated gestation are excluded for states requiring reporting of all products of conception, except for those with a stated birthweight of 350 grams or more. In 2019, this rule

was applied to the following reporting areas: Arkansas, Colorado, Georgia, Hawaii, New York (including New York City), Oklahoma, Pennsylvania, Rhode Island, and Virginia (American Samoa and the Virgin Islands did not report data for 2019) (see Table A). The summation of these rules results in the creation of the tabulation flag (position 10), which, when invoked, produces the number of records with a stated or presumed period of gestation of 20 weeks or more.

Classification of fetal deaths by occurrence and residence

In tabulations by place of residence, fetal deaths occurring within the United States to U.S. citizens and to residents who are not citizens are allocated to the usual place of residence of the mother in the United States, as reported on the report of fetal death. Fetal deaths to U.S. residents occurring outside this country are not included in tabulations by place of residence or place of occurrence. Accordingly, the total count of fetal deaths for the United States by place of residence and by place of occurrence will not be identical. Fetal deaths to nonresidents of the United States are included in data by place of occurrence but excluded from data by place of residence, as previously indicated. See Tables 2 and 3, respectively, for the number of births by residence and occurrence for the 50 states and the District of Columbia for 2019.

Geographic classification: The geographic code structure for the 2019 fetal death file is given in the NCHS manual, “Vital Records Geographic Classification, 2014,” and in the country, county, and place geographic code files [4]. The geographic code structure on the 2019 file is based on results of the 2010 Census of Population.

Standard Report of Fetal Death

The U.S. Standard Report of Fetal Death, issued by the U.S. Department of Health and Human Services, has served for many years as the principal means for attaining uniformity in the

content of the documents used to collect information on fetal deaths in the United States. The U.S. Standard Report of Fetal Death has historically been revised every 10-15 years. Most state reports conform closely in content to the standard report, but are modified to the extent required by the particular state's needs or by special provisions of the state's vital statistics law.

The 2003 revision: In 2003, a revised U.S. Standard Report of Fetal Death was adopted (Figure 1). For more information on the 2003 standard report and details regarding the certificate revision and links to the documents referenced below, see the NCHS website of the 2003 report revision at http://www.cdc.gov/nchs/nvss/vital_certificate_revisions.htm. The 2003 report of fetal death replaces the previous 1989 U.S. Standard Report of Fetal Death [5,6]. Implementation of the 2003 U.S. Standard Report of Fetal Death (revised) by the states and independent reporting areas was phased in from 2003 to 2018. All states, the District of Columbia, Guam, the Northern Marianas, and Puerto Rico had implemented the revised report of fetal death as of January 1, 2018. American Samoa and the Virgin Islands continue to use the 1989 (unrevised) revision of the U.S. Standard Report of Fetal Death (see User Guide to the 2017 Fetal Death Public Use File [7] for a detailed implementation schedule).

The 2003 Revision of the U.S. Standard Report of Fetal Death introduced substantial changes to data content and quality. Many key data items are common between revisions; however, a number of items were substantively modified. The 2003 revision also includes many new items never before collected on the Standard Report [5,6]. For details on data items comparable between revisions see the User Guide to the 2013 Fetal Death Public Use File [8].

A key aspect of the 2003 revision of the U.S. Standard Report of Fetal Death was the reengineering of the data collection and transmission system to improve data quality, speed of data collection and transmission, and to enhance standardization of data [5,9]. To encourage

collection of data from the best sources, two worksheets were developed: the “Patient’s Worksheet” (available at: <https://www.cdc.gov/nchs/data/dvs/patientwkstfetaldeath.pdf>) [10] and the “Facility Worksheet” (available at: <https://www.cdc.gov/nchs/data/dvs/FacilityFetal04.pdf>) [11]. For the Patient’s Worksheet, data are directly obtained from the mother and include items such as race, Hispanic origin, and educational attainment. For the Facility Worksheet, data are obtained directly from the medical records of the mother for items such as date of first prenatal care visit, pregnancy risk factors, and method of delivery. To assist hospital staff in completing the Facility Worksheet, a comprehensive instruction manual was developed: Guide to Completing the Facility Worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 Revision) (“Guide to the Facility Worksheet”; available at: <https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf>) [12]. Detailed definitions and instructions for data items that are collected from the Facility Worksheet are in the “Guide to the Facility Worksheet”.

Additionally, a review of 2003-based report of fetal death revision items in 2014 by a collaborative effort among representatives from several vital statistics jurisdictions: The National Association for Public Health Statistics and Information Systems (NAPHSIS), and NCHS, resulted in the decision to drop a number of items from the national report of fetal death data file for reasons of poor data quality. For more information on this effort and for a full list of items that were dropped, see:

https://www.cdc.gov/nchs/nvss/deleted_items_from_birth_fetal_death_files.htm.

The first ever eLearning training, “Applying Best Practices for Reporting Medical and Health Information on Birth Certificates,” on completing the medical and health information for the birth certificate and report of fetal death was launched in October 2016. The training

emphasizes the importance and uses of birth certificate and report of fetal death data and best practices for collecting specific medical and health items. The audience for the training includes birth information specialists, physicians, nurses, and hospital administrators. Continuing education credits for nurses, physicians, and non-clinical staff are also available. The training is internet-based and approximately 45 minutes in length. It is available at:

www.cdc.gov/nchs/training/BirthCertificateElearning.

Fetal death data files

Micro-data files: Fetal death micro-data files for data years 1982-2019 may be downloaded at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm. The general rules used to classify characteristics of fetal deaths are presented in several NCHS manuals [4, 9, 13]. These instructions are for states to use to collect and code the data items; they do not include NCHS edit recodes.

The 2003-2006 fetal death micro-data files include data items common to both the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death, as well as items exclusive to the 2003 revision. For 2007-2013, in an effort to improve the timeliness of fetal death data, the fetal death files only included items that were comparable across the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death. From 2014-2017, the fetal mortality files again included data comparable between the two revisions, as well as items exclusive to the 2003 revision. Beginning with 2018, the fetal death file only includes data exclusive to the 2003 revision.

The U.S. micro-data fetal death files for 2005 through 2019 do not include geographic detail (e.g., mother's state of residence). Tabulations of fetal death data by state-level geographic detail and many other demographic and health items are available in the CDC WONDER web

application (<http://wonder.cdc.gov/>) for data years 2005-2018. Additionally, certain geographic level data may also be available upon request: See “NCHS Data Release and Access Policy for Micro-data and Compressed Vital Statistics Files”, available at:

http://www.cdc.gov/nchs/nvss/dvs_data_release.htm.

The territories file includes data on fetal deaths occurring in Guam, the Northern Marianas, and Puerto Rico (American Samoa and the Virgin Islands did not report data for 2019). It includes limited geographical data such as information identifying individual territories and counties (or their equivalent) with populations of 100,000 or more by place of occurrence and residence.

Race of Mother

The 2003 revision of the U.S. Standard Report of Fetal Death allows the reporting of the five race categories either alone (i.e., single-race) or in combination (i.e., more than one race or multiple races) for the mother [6], in accordance with the revised standards issued by the Office of Management and Budget (OMB) in 1997 [14]. The five categories for race specified in the revised standards are: American Indian or Alaska Native (AIAN), Asian, Black or African American, Native Hawaiian or Other Pacific Islander (NHOPI), and White. Information on this change is presented elsewhere [15-17]. Starting in 2018, all states and the District of Columbia, in addition to Puerto Rico, Guam, and Northern Marianas, reported race according to the 1997 revised OMB standards.

Data from American Samoa and the Virgin Islands still follow the 1977 OMB standards, which required a minimum set of four single-race categories [American Indian or Alaska Native (AIAN), Asian or Pacific Islander (API, with Asian combined with Pacific Islander), black, and white] and to allow respondents to select only one race category [18].

Prior to 2018, in order to provide uniformity and comparability of the data before all or most of the data were available in the new multiple-race format, it was necessary to “bridge” the responses of those for whom more than one race was reported (multiple race) to one, single race. Beginning in 2018, data by race are shown according to the 1997 OMB standards. For a comparison of 2018 data by race between the 1977 and 1997 OMB standards, see Table B in the 2018 User Guide (19).

Hispanic Origin of Mother

Hispanic origin and race Hispanic origin: Hispanic origin and race are reported separately on the birth certificate (Figure 1). It is recommended that this information be reported directly by the mother via the Mother’s Worksheet [10]. For 1989 through 2018, data on the public use file and in NCHS reports for specified Hispanic groups are shown in most cases for five specified Hispanic groups: Mexican, Puerto Rican, Cuban, Central and South American, and “other and unknown Hispanic.” Starting with 2019, data are presented for the additional Hispanic group, Dominican (see items MHISPX in file position 143). This subgroup was previously included in “other and unknown Hispanic.” In tabulations of fetal death data by race and Hispanic origin, data for persons of Hispanic origin are not further classified by race because the vast majority of Hispanic women are reported as white. In tabulations of fetal death data by race only, data for persons of Hispanic origin are included in the data for each race group according to the mother’s reported race. In tabulations that include Hispanic origin, data for non-Hispanic persons are classified according to the race of the mother.

Cause of Death Data

A cause-of-fetal death item has been included on the report of fetal death since 1939 because it was considered critical information. However, the data were not released on public–

use files or published until 2014 due to resource constraints and quality concerns. National data on cause of death are now released annually and are available for 2014-2019.

A major redesign of the cause of death (COD) section of the U.S. Standard Report of Fetal Death was undertaken with the 2003 revision. The goals of the redesign were to be consistent with instructions in the World Health Organization's (WHO) International Classification of Diseases (ICD) [20], and to thereby improve the quality and specificity of COD information, provide additional reporting guidance, and retain the flexibility to report any cause of death [21].

As with other deaths, the intent is for an attending physician or medical examiner or coroner to report cause of death [20]. The cause-of-fetal-death item requests a medical opinion from this person on the conditions and diseases resulting in or contributing to death, but also asks the medical certifier to report one cause separately (item 18a) from all other causes (item 18b) reported on the fetal death report. The certifier may form this medical opinion upon various medical tests, investigations, and examinations as is the case with other deaths. However, the term "initiating cause" used to refer to the one cause reported separately is unique to fetal deaths because of differences in the format of the cause item and how the initiating cause is determined for fetal deaths compared with the "underlying cause" term used with other deaths [21].

NCHS codes cause of fetal death reported by the certifier using the ICD-10 classification [20]. Coding is accomplished through a combination of automated and manual processes following the guidelines laid out in Instruction manual part 2k, Instructions for the Automated Classification of the Initiating and Multiple Causes of Fetal Deaths [22]. Literal text stated on the fetal death report is assigned ICD-10 codes, and a single cause of death, the initiating cause of death, is selected from the conditions entered by the medical certifier in the cause-of-death

section of the fetal death report. One section of the report (18a) is for the medical certifier to state the single condition that he or she considers the cause that initiated or triggered problems that resulted in the fetus dying at this point in time, so this is anticipated to be the initiating cause of death. If more than one cause or condition is entered by the medical certifier, the initiating cause is determined by the placement of the condition on the report, provisions of the ICD, and associated selection rule and modifications. A second section of the report (18b) is for the medical certifier to state any other conditions or causes that he or she felt played a role in causing the fetal death. Since more medical information may be reported on the report than is directly reflected in the initiating cause of death, this additional information is captured in multiple cause-of-fetal-death data.

The fetal death data file also includes a cause of death data item for 124 Selected Causes of Fetal Death (fetal cause list) (tables 7A and 7B). The reporting area for these tables includes areas that used the 2003 revision format and where less than 50 percent of an area's cause data were attributed to the Fetal death of unspecified cause (unspecified cause) (P95).

Incomplete national reporting: The use of reporting flags

The 2019 national fetal death file includes items that are not reported by all vital statistics jurisdictions. Reporting flags were developed to help the user more readily identify the correct number of fetal deaths for items with less than full national reporting. Reporting flags are available for many items in the file, and positions for reporting flags are noted along with each data item in the file layout.

Translating "blanks"

In the 2019 fetal death file, for data items which are not reported by all states, fetal deaths to residents of a state reporting the item that occurred in a state not reporting the item, are

represented by “blanks.” Blanks should be treated as “unknowns” for tabulations. In sum, the correct use of reporting flags and translation of blanks will result in an accurate tally of fetal deaths for items with incomplete national reporting. For an example of SAS code that may be used to incorporate the correct use of reporting flags and the translation of blanks, see below.

Example of SAS code using reporting flags (and translating blanks)

The example below is for the receipt of WIC item, for which data for 48 states are included for 2019 (New York City does not report this item and the District of Columbia, Hawaii, and Rhode Island reported this item for 2019 but had unknown levels of 50% or higher and were flagged as not reporting).

Sample SAS program

```
01    DATA work;
02        INFILE 'c:fet16us.dat' LRECL=3050;
03        INPUT
04            oe_tabflg    10
05            wic    $229
06            restatus 131
07            f_wic    $377;
08
09        /*Include if 20 weeks gestation or more*/
10        IF oe_tabflg EQ 2;
11        /*Exclude foreign residents*/
12        IF restatus NE 4;
13        /*Select reporting area*/
14        IF f_wic=1;
15        /*Convert blanks to unknown*/
```

```
16           IF wic = ' ' THEN wic = 'U';
17
18   PROC FREQ;
19           TABLE wic;
20   RUN;
```

In this example, “oe_tabflg” is used to select fetal deaths of 20 weeks gestation or more. Also, “restatus” is used to exclude fetal deaths to foreign residents (this is standard practice for all NCHS tabulations). As well in this example, blanks are represented by character values SAS code = (‘ ’). However, for some items in the file, e.g., number of previous cesarean deliveries, blanks are represented by numeric values for which the SAS code is (‘.’).

Quality of Data

Percentage of unknown responses

Information on the completeness of reporting of fetal death (with a stated or presumed period of gestation of 20 weeks or more) data is shown in Table B, which presents a listing of items and the percentage of records that were not stated for each reporting area: all 50 states, the District of Columbia, New York City, Puerto Rico, Guam, American Samoa, the Northern Marianas, and the Virgin Islands. In general, percentages of unknown responses are considerably higher for fetal deaths than for live births, and among fetal deaths the percentage unknown is higher for fetal deaths that occur earlier in the gestational period. In fetal mortality tables, unknown responses are shown in frequencies tables but are excluded from the computation of percent distributions and fetal and perinatal mortality rates. Thus, rates for

variables with a substantial percentage of unknown responses (such as birthweight) may understate the “true” rates of fetal mortality for that characteristic.

As high levels of missing data are a data quality concern, beginning with the 2017 data year, jurisdictions with a 50% or greater unknown level for a particular item are flagged as not reporting that item.

State Specific Issues

The items below had unknown levels of 50% or higher and were flagged as not reporting for the following jurisdictions in 2019:

- Father’s date of birth (all components) - Alabama, Arkansas, the District of Columbia, Georgia, New York (including New York City), Tennessee, Virginia, Wisconsin, and Guam
- WIC- District of Columbia, Hawaii, Rhode Island
- Attendant-Indiana
- Estimated time of fetal death-District of Columbia, Hawaii, Kansas, Nevada, New York, Tennessee, Virginia, Guam, Northern Marianas, and Puerto Rico
- Mother’s education-Rhode Island
- Mother’s pre-pregnancy weight-Hawaii
- Date of last live birth-Washington
- Cause of death- California, Georgia, Hawaii, Michigan, Mississippi, North Carolina, North Dakota, Vermont, New York City, Guam, and Northern Marianas

Computation of rates

Fetal mortality rates are computed as the number of fetal deaths at 20 weeks of gestation or more per 1,000 live births and fetal deaths at 20 weeks or more. Perinatal mortality rates are

computed in a similar fashion, as shown below. The denominators for all fetal and perinatal mortality rates are live births plus fetal deaths in the specified gestational age group, thus representing the population at risk of the event [2].

$$\text{Fetal mortality rate} = \frac{\text{Fetal deaths at 20 weeks of gestation or more}}{\text{Live births and fetal deaths at 20 weeks or more}} \times 1,000$$

Perinatal mortality rate, definition I =

$$\frac{\text{Fetal deaths at 28 weeks or more and infant deaths under 7 days}}{\text{Live births and fetal deaths at 28 weeks or more}} \times 1,000$$

Perinatal mortality rate, definition II =

$$\frac{\text{Fetal deaths at 20 weeks or more and infant deaths under 28 days}}{\text{Live births and fetal deaths at 20 weeks or more}} \times 1,000$$

In each case, the fetal deaths included in the denominator of each rate mirror the fetal deaths included in the numerator. A previous NCHS report [23] contains information on the historical development of various perinatal measures. An asterisk (*) is shown in place of any rate based on fewer than 20 fetal or perinatal deaths in the numerator.

Random variation in fetal and perinatal mortality

The number of fetal deaths, perinatal deaths, and live births reported for an area represent complete counts of such events. As such, they are not subject to sampling error, although they are subject to nonsampling error in the registration process. However, when the figures are used for analytic purposes, such as the comparison of rates over time, for different areas, or among

different subgroups, the number of events that actually occurred may be considered as one of a large series of possible results that could have arisen under the same circumstances [24]. As a result, numbers of births, fetal deaths, perinatal deaths, and fetal and perinatal mortality rates are subject to random variation. The probable range of values may be estimated from the actual figures according to certain statistical assumptions.

In general, distributions of vital events may be assumed to follow the normal distribution. When the number of events is large, the relative standard error (RSE) is usually small. When the number of events is small (perhaps less than 100) and the probability of such an event is small, considerable caution must be used in interpreting the data. Such infrequent events may be assumed to follow a Poisson probability distribution. Estimates of RSEs and 95% confidence intervals are shown below. In the formulas, D is the number of fetal or perinatal deaths and B is the number of live births plus fetal deaths used as the denominator in computing fetal and perinatal mortality rates.

The formulas for the two RSEs are as follows:

$$\text{RSE}(D)=100*\sqrt{\frac{1}{D}} \text{ and}$$

$$\text{RSE}(B)=100*\sqrt{\frac{1}{B}} .$$

For example, let us say that for group A the number of fetal deaths was 238, whereas the number of live births plus fetal deaths in the denominator was 32,650 yielding a fetal mortality rate of 7.29 fetal deaths per 1,000 live births and fetal deaths.

$$\text{The RSE of the deaths} = 100*\sqrt{\frac{1}{238}} = 6.48,$$

whereas the RSE for the births plus fetal deaths in the denominator is

$$\text{RSE} = 100 * \sqrt{\frac{1}{32,650}} = 0.55.$$

The formula for the RSE of the fetal mortality rate is:

$$\text{RSE} = 100 * \sqrt{\frac{1}{D} + \frac{1}{B}}$$

Thus the RSE for the example above is:

$$= 100 * \sqrt{\frac{1}{238} + \frac{1}{32,650}} = 6.51.$$

Normal distribution

When the number of events is greater than 100, the normal distribution is used to estimate the 95 percent confidence intervals of a rate, R_I , as follows:

$$\text{Lower: } R_I - 1.96 * R_I * \frac{\text{RSE}(R_I)}{100}$$

$$\text{Upper: } R_I + 1.96 * R_I * \frac{\text{RSE}(R_I)}{100}$$

Thus, for Group A:

$$\text{Lower: } 7.29 - (1.96 * 7.29 * \frac{6.51}{100}) = 6.36$$

$$\text{Upper: } 7.29 + (1.96 * 7.29 * \frac{6.51}{100}) = 8.22$$

Thus the chances are 95 out of 100 that the true fetal or perinatal mortality rate for Group A lies somewhere in the 6.36–8.22 interval.

Poisson distribution

When the number of events in the numerator is less than 100 the confidence interval for the rate, R_I , can be estimated based on the Poisson distribution using the values in Table III (2).

Lower: $R_I * L(.95, D_{adj})$

Upper: $R_I * U(.95, D_{adj})$

where D_{adj} is the adjusted number of fetal or perinatal deaths (rounded to the nearest integer) used to take into account the RSE of the number of deaths in the numerator and the number of live births plus fetal deaths in the denominator, and is computed as follows:

$$D_{adj} = \frac{D * B}{D + B}$$

$L(.95, D_{adj})$ and $U(.95, D_{adj})$ refer to the values in Table III corresponding to the value of D_{adj} .

For example, let us say that for Group B the number of fetal deaths was 73, and the number of live births plus fetal deaths in the denominator was 11,422, and the fetal mortality rate was 6.39.

$$D_{adj} = \frac{73 * 11,422}{73 + 11,422} = 73$$

Therefore the 95 percent confidence interval (using the formula in Table III for 1\N99 infant deaths) =

$$\text{Lower: } 6.39 * 0.78384 = 5.01$$

$$\text{Upper: } 6.39 * 1.25735 = 8.03$$

Comparison of two fetal or perinatal mortality rates

If either of the two rates to be compared is based on less than 100 deaths, compute the confidence intervals for both rates and check to see if they overlap. If so, the difference is not statistically significant at the 95 percent level. If they do not overlap, the difference is statistically significant. If both of the two rates (R_1 and R_2) to be compared are based on 100 or more deaths, the following z-test should be used to define a significance test statistic:

$$z = \frac{R_1 - R_2}{\sqrt{R_1^2 \left(\frac{RSE(R_1)}{100} \right)^2 + R_2^2 \left(\frac{RSE(R_2)}{100} \right)^2}}$$

If $|z| \geq 1.96$, then the difference is statistically significant at the 0.05 level and if $|z| < 1.96$, the difference is not significant.

References

1. National Center for Health Statistics. Fetal Death 2019. Public use file. Hyattsville, Maryland: National Center for Health Statistics. Annual internet product. 2021. Available at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.
2. MacDorman MF, Gregory ECW. Fetal and perinatal mortality: United States, 2013. National vital statistics reports; vol 64 no 8. Hyattsville, MD: National Center for Health Statistics. 2015.
3. Kowaleski J. State definitions and reporting requirements for live births, fetal deaths, and induced terminations of pregnancy (1997 revision). Hyattsville, MD: National Center for Health Statistics. 1997. Available from: <http://www.cdc.gov/nchs/data/misc/itop97.pdf>.
4. Division of Vital Statistics. Instruction manual part 8: Geographic classification, 2014. Hyattsville, Maryland: National Center for Health Statistics. Available at: https://www.cdc.gov/nchs/data/dvs/IMP8_2014.pdf.
5. National Center for Health Statistics. Report of the Panel to Evaluate the U.S. Standard Certificates. Hyattsville, Maryland: National Center for Health Statistics. 2000
6. National Center for Health Statistics. 2003 revision of the U.S. Standard Report of Fetal Death. 2003.
7. National Center for Health Statistics. User guide to the 2017 fetal death public use file. Hyattsville, MD. Available from: https://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm.
8. National Center for Health Statistics. User Guide to the 2013 Fetal death Public Use File. Hyattsville, Maryland: National Center for Health Statistics. Annual product 2015. Available at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.

9. National Center for Health Statistics. Fetal death edit specifications for the 2003 Revision of the U.S. Standard Certificate of Birth. 2012.

https://www.cdc.gov/nchs/data/dvs/fetal_death_edit_specifications.pdf.

10. National Center for Health Statistics. Patient's Worksheet for Report of Fetal Death.

2019. Available at: <https://www.cdc.gov/nchs/data/dvs/fetal-death-mother-worksheet-english-2019-508.pdf>.

11. National Center for Health Statistics. Facility Worksheet for Report of Fetal Death. 2019.

Available at: <https://www.cdc.gov/nchs/data/dvs/fetal-death-facility-worksheet-2019-508.pdf>.

12. National Center for Health Statistics. Guide to completing the facility worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 revision). 2019 update. Hyattsville, Maryland: National Center for Health Statistics.

<https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf>.

13. Division of Vital Statistics. Computer edits fetal death data --2003 Revised Certificate. Hyattsville, Maryland: National Center for Health Statistics.

14. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. Fed Regist 62FR58781–58790. October 30, 1997.

15. Ingram DD, Parker JD, Schenker N, et al. United States Census 2000 population with bridged race categories. National Center for Health Statistics. Vital Health Stat 2(135). 2003.

16. Johnson D. Coding and editing multiple race. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP. Portland, Oregon. June 6–10, 2004.

17. Weed JA. NCHS procedures for multiple-race and Hispanic origin data: Collection, coding, editing, and transmitting. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP. Portland, Oregon. June 6–10, 2004.

18. Office of Management and Budget. Race and ethnic standards for federal statistics and administrative reporting. Statistical Policy Directive 15. May 12, 1977.
19. National Center for Health Statistics. User guide to the 2018 fetal death public use file. Hyattsville, MD. Available from: https://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm.
20. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. Geneva, Switzerland. 2009.
21. Hoyert DL, Gregory ECW. Cause-of-death data from the fetal death file, 2015–2017. National Vital Statistics Reports; vol 69 no 4. Hyattsville, MD: National Center for Health Statistics. 2020.
22. National Center for Health Statistics. Instruction manual, part 2k: Instructions for the automated classification of the initiating and multiple causes of fetal deaths, 2014. Hyattsville, MD. 2014.
23. Hoyert DL. Perinatal mortality in the United States, 1985–91. National Center for Health Statistics. Vital Health Stat 20(26). 1995. Available from: http://www.cdc.gov/nchs/data/series/sr_20/sr20_026.pdf.
24. Brillinger DR. The natural variability of vital rates and associated statistics. Biometrics 42(4):693–734. 1986.

Control Count of Records

File Characteristics: Fetal Public Use File 2019

All files:

Record format: Blocked, Fixed, Format
Code scheme: Numeric/Alphabetic/Blank
Record length: 3150

Record counts

U.S. file record count:

Counts of all fetal deaths:

Record count	46,007
By occurrence	46,007
By residence	45,876
To foreign residents	131

Counts of 20 weeks or more:

Record count	21,556
By occurrence	21,556
By residence	21,478
To foreign residents	78

Counts of less than 20 weeks:

Record count	24,451
By occurrence	24,451
By residence	24,398
To foreign residents	53

U.S. territories file record count:

Counts of all fetal deaths:

Record count	247
By occurrence	247
By residence	246
To foreign residents	1

Counts of 20 weeks or more:

Record count	247
By occurrence	247
By residence	246
To foreign residents	1

Counts of less than 20 weeks:

Record count	0
By occurrence	0
By residence	0
To foreign residents	0

2019 LIST OF DATA ELEMENTS AND LOCATIONS

<u>Data Items</u>	<u>Locations</u>
1. General	
a) Tabulation flags (gestational age)	10
b) Data year	11-14
c) Resident status	131
2. Prenatal Care	
a) Month began	202-203
3. Fetus	
a) Sex	316
b) Plurality	301
c) Weight at delivery	349-355
d) Gestational age	340-344(NCHS Standard item), 331-335
e) Month/year of delivery	11-16
f) Time of delivery	21-24
g) Day of week of delivery	25
4. Mother	
a) Age	86-90
b) Race	132-137
c) Education	145
d) Hispanic origin	139-144
5. Pregnancy History	
a) Live birth order	187
b) Birth interval	197-201
6. Father	
a) Age	177-180
7. Other Items	
a) Attendant at delivery	283
b) Place of delivery	34, 52
c) WIC receipt	229
8. Medical and Health Data	
a) Method of delivery	274-277, 280
b) Medical risk factors	257-267
c) Other risk factors	
i. Tobacco	230-238
ii. Mother's pre-pregnancy body mass index (BMI)	245-249
d) Maternal morbidity	281-282
9. Residence reporting flags	372-440

10. Cause of Death

- | | |
|--|-----------|
| a.) Initiating cause or condition code | 2603-2607 |
| b.) Initiating cause or condition recode | 2643-2645 |
| c.) Initiating cause or condition flag | 2651 |

**2019 Fetal Death
Public Use Record Layout**

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
1-6	6	FILLER					
7	1	VERSION	Version		A,S	A S	State used the 2003 version of the US Standard Report of Fetal Death (Revised) State used the 1989 version of the US Standard Report of Fetal Death (Unrevised)
8-9	2	FILLER					
10	1	OE_TABFLG	Obstetric Estimate Tabulation Flag (Use with NCHS Standard item)		A,S	1 2	Under 20 weeks (exclude) 20 weeks or more (include)
11-14	4	DOD_YY	Delivery Year		A,S	2014	Year of delivery
15-16	2	DOD_MM	Delivery Month		A,S	01 02 03 04 05 06 07 08 09 10 11 12	January February March April May June July August September October November December
17-20	4	FILLER					
21-24	4	DOD_TT	Delivery Time	c409	A	0000-2359 9999	Time of Birth Not Stated
25	1	DOD_WK	Weekday		A,S	1 2 3	Sunday Monday Tuesday

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						4	Wednesday
						5	Thursday
						6	Friday
						7	Saturday
26-27	2	OSTATE	Territory of Occurrence <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A,S		
			<u>Territories</u>			AS	American Samoa
						GU	Guam
						MP	Northern Marianas
						PR	Puerto Rico
						VI	Virgin Islands
28-29	2	FILLER					
30-32	3	OCNTYFIPS	Occurrence FIPS County <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file) Data are suppressed if population < 100,000</i>		A,S		
						000	No county level geography
						001-998	County of Occurrence (See Geographic Table)
						999	County of less than 100,000
33	1	OCNTYPOP	Occurrence Territory County Population <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A,S		
						0	County of 1,000,000 or more
						1	County of 500,000 to 1,000,000
						2	County of 250,000 to 500,000
						3	County of 100,000 to 250,000

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						9	County less than 100,000
34	1	BFACIL	Delivery Place (Revised)	409	A	1 2 3 4 5 6 7 9	Hospital Freestanding Birth Center Home (intended) Home (not intended) Home (unknown if intended) Clinic/Doctor's Office Other Unknown
35-51	17	FILLER					
52	1	BFACIL3	Delivery Place Recode		A,S	1 2 3	In Hospital Not in Hospital Unknown/not stated
53-83	31	FILLER					
84	1	MAGE_IMPFLG	Mother's Age Imputed		A,S	Blank 1	Age not imputed Age imputed
85	1	MAGE_REPFLG	Reported Age of Mother Flag		A,S	Blank 1	Reported age not used Reported age used
86-87	2	MAGER	Mother's Single Year of Age	432	A,S	12 13 14 15 16	12 years and under 13 years 14 years 15 years 16 years

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						17	17 years
						18	18 years
						19	19 years
						20	20 years
						21	21 years
						22	22 years
						23	23 years
						24	24 years
						25	25 years
						26	26 years
						27	27 years
						28	28 years
						29	29 years
						30	30 years
						31	31 years
						32	32 years
						33	33 years
						34	34 years
						35	35 years
						36	36 years
						37	37 years
						38	38 years
						39	39 years
						40	40 years
						41	41 years
						42	42 years
						43	43 years
						44	44 years
						45	45 years
						46	46 years
						47	47 years
						48	48 years
						49	49 years
						50	50 years and over
88-89	2	MAGER14	Mother's Age Recode 14	432	A,S	01	Under 15 Years
						03	15 years
						04	16 years
						05	17 years
						06	18 years
						07	19 years

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						08	20-24 years
						09	25-29 years
						10	30-34 years
						11	35-39 years
						12	40-44 years
						13	45-49 years
						14	50-54 years
90	1	MAGER9	Mother's Age Recode 9	432	A,S	1	Under 15 years
						2	15-19 years
						3	20-24 years
						4	25-29 years
						5	30-34 years
						6	35-39 years
						7	40-44 years
						8	45-49 years
						9	50-54 years
91-92	2	MBCNTRY	Mother's Birth Country c409 <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A	AA-ZZ	See Geographic Documentation
93-96	4	FILLER					
97	1	MBSTATE_REC	Mother's Birth State Recode c409		A	1	Native born (50 U.S. States)
						2	Foreign born (includes territories)
						3	Unknown or Not Stated
98-101	4	FILLER					
102-103	2	MRSTATEPSTL	Mother's Residence Postal State <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A,S		
			<u>United States</u>			AK	Alaska
						AL	Alabama
						AR	Arkansas

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						AZ	Arizona
						CA	California
						CO	Colorado
						CT	Connecticut
						DE	Delaware
						DC	District of Columbia
						FL	Florida
						GA	Georgia
						HI	Hawaii
						ID	Idaho
						IL	Illinois
						IN	Indiana
						IA	Iowa
						KS	Kansas
						KY	Kentucky
						LA	Louisiana
						MA	Massachusetts
						MD	Maryland
						ME	Maine
						MI	Michigan
						MN	Minnesota
						MO	Missouri
						MS	Mississippi
						MT	Montana
						NC	North Carolina
						ND	North Dakota
						NE	Nebraska
						NH	New Hampshire
						NJ	New Jersey
						NM	New Mexico
						NV	Nevada
						NY	New York
						OH	Ohio
						OK	Oklahoma
						OR	Oregon
						PA	Pennsylvania
						RI	Rhode Island
						SC	South Carolina
						SD	South Dakota
						TN	Tennessee
						TX	Texas

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						UT	Utah
						VA	Virginia
						VT	Vermont
						WA	Washington
						WI	Wisconsin
						WV	West Virginia
						WY	Wyoming
			<u>Territories</u>			AS	American Samoa
						GU	Guam
						MP	Northern Marianas
						PR	Puerto Rico
						VI	Virgin Islands
			<u>Canadian Provinces</u>			AB	Alberta
						BC	British Columbia
						MB	Manitoba
						NB	New Brunswick
						NL	Newfoundland and Labrador
						NT	Northwest Territories
						NS	Nova Scotia
						NU	Nunavut
						ON	Ontario
						PE	Prince Edward Island
						QC	Quebec
						SK	Saskatchewan
						YT	Yukon Territory
			<u>Foreign</u>			XX	Foreign (Rest of world)
						ZZ	Not Classifiable
104-106	3	MRCNTYFIPS	Mother's FIPS County of Residence <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file) Data are suppressed if population < 100,000</i>	A,s		000	No county level geography
						001-998	County of residence (See Geographic Table)

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						999	County of less than 100,000
107-111	5	FILLER					
112	1	RCNTY_POP	Population of Residence County <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A,S	0 1 2 3 9 Z	County of 1,000,000 or more County of 500,000 to 1,000,000 County of 250,000 to 500,000 County of 100,000 to 250,000 County less than 100,000 Foreign resident
113-129	17	FILLER					
130	1	RECTYPE	Record Type <i>(This item is available in the territory file only, geographic codes are not available in the U.S. file)</i>		A,S	1 2	RESIDENT: State and county of occurrence and residence are the same. NONRESIDENT: State and county of occurrence and residence are different.
131	1	RESTATUS	Residence Status <u>United States</u>		A,S	1 2 3 4 1	RESIDENT: State and county of occurrence and residence are the same. INTRASTATE NONRESIDENT: State of occurrence and residence are the same but county is different. INTERSTATE NONRESIDENT: State of occurrence and residence are different but both are one of the 50 US states or District of Columbia. FOREIGN RESIDENT: The state of residence is not one of the 50 US states or District of Columbia. <u>Territories</u> RESIDENT: State and county of occurrence and residence

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
							are the same. (Unique to Guam, all US residents are considered residents of Guam and thus are assigned 1.)
						2	INTRATERRITORY NONRESIDENT: Territory of occurrence and residence are the same but county is different.
						3	INTERTERRITORY RESIDENT: Territory of occurrence and residence are different but both are US Territories.
						4	FOREIGN RESIDENT: The residence is not a US Territory.
132-133	2	MRACE31	Mother's Race Recode 31	425	A		
						01	White (alone)
						02	Black (alone)
						03	AIAN (American Indian or Alaskan Native) (alone)
						04	Asian (alone)
						05	NHOPI (Native Hawaiian or Other Pacific Islander) (alone)
						06	Black And White
						07	Black and AIAN
						08	Black and Asian
						09	Black and NHOPI
						10	AIAN and White
						11	AIAN and Asian
						12	AIAN and NHOPI
						13	Asian and White
						14	Asian and NHOPI
						15	NHOPI and White
						16	Black, AIAN, and White
						17	Black, AIAN, and Asian
						18	Black, AIAN, and NHOPI
						19	Black, Asian, and White
						20	Black, Asian, and NHOPI
						21	Black, NHOPI, and White
						22	AIAN, Asian, and White
						23	AIAN, NHOPI, and White
						24	AIAN, Asian, and NHOPI
						25	Asian, NHOPI, and White
						26	Black, AIAN, Asian, and White
						27	Black, AIAN, Asian, and NHOPI
						28	Black, AIAN, NHOPI, and White
						29	Black, Asian, NHOPI, and White
						30	AIAN, Asian, NHOPI, and White
						31	Black, AIAN, Asian, NHOPI, and White

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
134	1	MRACE6	Mother's Race Recode 6	425	A	1 2 3 4 5 6	White (alone) Black (alone) AIAN (alone) Asian (alone) NHOPI (alone) More than one race
135-136	2	MRACE15	Mother's Race Recode 15	425	A	01 02 03 04 05 06 07 08 09 10 11 12 13 14 15	White (alone) Black (alone) AIAN (alone) Asian Indian (alone) Chinese (alone) Filipino (alone) Japanese (alone) Korean (alone) Vietnamese (alone) Other Asian (alone) Hawaiian (alone) Guamanian (alone) Samoan (alone) Other Pacific Islander (alone) More than one race
137	1	MRACEREC	Mother's Bridged Race Recode <u>United States and non-Puerto Rican Territories</u>	439		A,S 1 2 3 4	White Black American Indian or Alaskan Native Asian or Pacific Islander
138	1	MRACEIMP	Mother's Race Imputed		A,S	Blank 1 2	Mother's race not imputed Unknown race imputed All other races, formerly coded 09, imputed.
139-141	3	FILLER					

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
142	1	UMHISP	Mother's Hispanic Origin	424	A,S	0 1 2 3 4 5 9	Non-Hispanic Mexican Puerto Rican Cuban Central or South American Other and Unknown Hispanic Origin unknown or not stated
143	1	MHISPX	Mother's Hispanic Origin			0 1 2 3 4 5 6 9	Non-Hispanic Mexican Puerto Rican Cuban Central or South American Dominican Other and Unknown Hispanic Origin unknown or not stated
144	1	SR_MRACEHISP	Mother's Race/Hispanic Origin	c425	A	1 2 3 4 5 6 7 8	Non-Hispanic White (alone) Non-Hispanic Black (alone) Non-Hispanic AIAN (alone) Non-Hispanic Asian (alone) Non-Hispanic NHOPI (alone) Non-Hispanic more than one race Hispanic Origin unknown or not stated
145	1	MEDUC	Mother's Educ-Revised	372	A	1 2 3 4 5 6 7 8	8 th grade or less 9 th through 12 th grade with no diploma High school graduate or GED completed Some college credit, but not a degree Associate degree (AA, AS) Bachelor's degree (BA, AB, BS) Master's degree (MA, MS, MEng, Med, MSW, MBA) Doctorate (PhD, EdD) or Professional degree (MD, DDS, DVM, LLB, JD)

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						9	Unknown
146-171	26	FILLER					
172	1	FAGERPT_FLG	Father's Reported Age Used		A,S	Blank 1	Father's reported age not used Father's reported age used
173-176	4	FILLER					
177-178	2	FAGECOMB	Father's Combined Age	426	A,S	09-98 99	Father's combined age in years Unknown or not stated
179-180	2	FAGEREC11	Father's Age Recode 11	426	A,S	01 02 03 04 05 06 07 08 09 10 11	Under 15 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years 55-98 years Not stated
181-182	2	PRIORLIVE	Prior Births Now Living	440	A,S	00-30 99	Number of children still living from previous live births. Unknown or not stated
183-184	2	PRIORDEAD	Prior Births Now Dead	440	A,S	00-30 99	Number of children dead from previous live births. Unknown or not stated
185-186	2	FILLER					

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
187	1	LBO_REC	Live Birth Order Recode	440	A,S	0-7 8 9	Number of live birth order. 8 or more live births Unknown or not stated
188-196	9	FILLER					
197-199	3	ILLB_R	Interval Since Last Live Birth Recode	427	A	000-003 004-300 888 999	Plural delivery Months since last live birth Not applicable Unknown or not stated
200-201	2	ILLB_R11	Interval Since Last Live Birth Recode 11	427	A	00 01 02 03 04 05 06 07 08 88 99	Zero to 3 months (plural delivery) 4 to 11 months 12 to 17 months 18 to 23 months 24 to 35 months 36 to 47 months 48 to 59 months 60 to 71 months 72 months and over Not applicable Unknown or not stated
202-203	2	PRECARE	Month Prenatal Care Began (Revised)	403	A	00 01-10 99	No prenatal care Month prenatal care began Unknown or not stated
204	1	PRECARE_REC	Month Prenatal Care Began Recode (Revised)	403	A	1 2 3 4 5	1 st to 3 rd month 4 th to 6 th month 7 th to final month No prenatal care Unknown or not stated
205-228	24	FILLER					

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
229	1	WIC	WIC	377	A	Y N U	Yes No Unknown or not stated
230-231	2	CIG_0	Cigarettes Before Pregnancy	405	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
232-233	2	CIG_1	Cigarettes 1st Trimester	406	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
234-235	2	CIG_2	Cigarettes 2nd Trimester	407	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
236-237	2	CIG_3	Cigarettes 3rd Trimester	408	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
238	1	CIG_REC	Cigarette Recode (Revised)	374	A	Y N U	Yes No Unknown or not stated
239-242	4	FILLER					
243-244	2	M_Ht_In	Mother's Height in Inches	375	A	30-78 99	Height in inches Unknown or not stated
245-248	4	BMI	BMI (prepregnancy)	376	A	13.0-69.9 99.9	Body Mass Index Unknown or not stated

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
249	1	BMI_R	Body Mass Index Recode (prepregnancy)	376	A	1 2 3 4 5 6 9	Underweight <18.5 Normal 18.5-24.9 Overweight 25.0-29.9 Obesity I 30.0-34.9 Obesity II 35.0-39.9 Extreme Obesity III ≥ 40.0 Unknown or not stated
250-252	3	FILLER					
253-255	3	PWgt_R	Prepregnancy Weight Recode	376	A	075-375 999	Weight in pounds Unknown or not stated
256	1	FILLER					
257-262	6	<u>Risk Factors (Revised)</u>			A		
			The checkbox items indented below follow this code structure:			Y N U	Yes No Unknown or not stated
257	1	RF_DIAB	Prepregnancy Diabetes	378			
258	1	RF_GEST	Gestational Diabetes	379			
259	1	RF_PHYP	Prepregnancy Hypertension	380			
260	1	RF_GHYP	Gestational Hypertension	381			
261	1	RF_ECLAM	Hypertension Eclampsia	382			
262	1	RF_INFTR	Infertility Treatment	383			
263	1	RF_FEDRG	Fertility Enhancing Drugs	384	A	Y N X U	Yes No Not applicable Unknown or not stated
264	1	RF_ARTEC	Asst. Reproductive Technology	385	A	Y N X U	Yes No Not applicable Unknown

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
265	1	RF_CESAR	Previous Cesareans	386	A	Y N U	Yes No Unknown or not stated
266-267	2	RF_CESARN	Previous Cesareans Num	387	A	00 01-30 99	None Number of previous cesareans Unknown or not stated
268-273	6	FILLER					
274-277	6	<u>Method of Delivery (Revised)</u>					
274	1	ME_PRES	Fetal Presentation	388	A	1 2 3 9	Cephalic Breech Other Unknown or not stated
275	1	ME_ROUT	Route & Method of Delivery	389	A	1 2 3 4 9	Spontaneous Forceps Vacuum Cesarean Unknown or not stated
276	1	ME_TRIAL	Trial of Labor Attempted	390	A	Y N X U	Yes No Not applicable Unknown or not stated
277	1	RDMETH_REC	Delivery Method Recode (Revised)	c386	A	1 2 3 4 5 6 9	Vaginal (excludes vaginal after previous C section) Vaginal after previous c-section Primary C-section Repeat C-section Vaginal (unknown if previous c-section) (2003 Standard only) C-section (unknown if previous c-section) (2003 Standard only) Not stated

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
278-279	2	FILLER					
280	1	DMETH_REC	Delivery Method Recode	410	A,S	1 2 9	Vaginal C-section Not stated
281-282	2		<u>Maternal Morbidity</u> The checkbox items indented below follow this structure:		A	Y N U	Yes No Unknown or not stated
281	1	MM_RUPT	Ruptured Uterus	391			
282	1	MM_ICU	Admit to Intensive Care	392			
283	1	ATTEND	Attendant	433	A,S	1 2 3 4 5 9	Doctor of Medicine (MD) Doctor of Osteopathy (DO) Certified Nurse Midwife (CNM) Other Midwife Other Unknown or not stated
284-300	17	FILLER					
301	1	DPLURAL	Plurality Recode		A,S	1 2 3 4	Single Twin Triplet Quadruplet or higher
302	1	FILLER					
303	1	IMP_PLUR	Plurality Imputed		A,S	Blank 1	Plurality is not imputed Plurality is imputed

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
304-315	12	FILLER					
316	1	SEX	Sex of Infant		A,S	M F	Male Female
317	1	IMP_SEX	Sex Imputed Only where gestation is stated or presumed to be 20 weeks or more		A,S	Blank 1	Infant Sex not Imputed Infant Sex is Imputed
		<u>Gestation</u>					
318-319	2	DLMP_MM	Last Normal Menses Month	429	A,S	01 02 03 04 05 06 07 08 09 10 11 12 99	January February March April May June July August September October November December Unknown or not stated
320-321	2	FILLER					
322-325	4	DLMP_YY	Last Normal Menses Year	431	A,S	nnnn 9999	Year of last normal menses Unknown or not stated
326-328	3	FILLER					
329	1	GEST_IMP	Gestation Imputed Flag		A,S	Blank 1	Gestation is not imputed Gestation is imputed
330	1	OBGEST_FLG	Obstetric Estimate of Gestation Used Flag		A,S	Blank 1	Obstetric Estimate is not used Obstetric Estimate is used

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
331-332	2	COMBGEST	Gestation – Detail in Weeks	404	A,S	02-47 99	2 nd through 47 th week of Gestation Unknown
333-334	2	GESTREC12	Gestation Recode 12	404	A,S	01 02 03 04 05 06 07 08 09 10 11 12	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks 28-31 weeks 32-33 weeks 34-36 weeks 37-38 weeks 39-40 weeks 41 weeks 42 weeks and over Unknown
335	1	GESTREC5	Gestation Recode 5	404	A,S	1 2 3 4 5	Under 20 weeks 20-23 weeks 24-27 weeks 28 weeks and over Unknown
336-337	2	OEGest_Unedt	Obstetric Gestation 373	A,S		0-98 02-47 99	Range of obstetric estimate of gestation - revised Range of clinical estimate of gestation - unrevised Unknown or not stated
338	1	FILLER					
339	1	COMBGEST_USED	Combined Gestation Used Flag		A,S	blank 1	Combined gestation not used Combined gestation used
340-341	2	OEGest_Comb	Obstetric Estimate Edited (NCHS Standard item)	373	A,S	02-47 99	Weeks of gestation Not stated
342-343	2	OEGest_R12	Obstetric Estimate Recode12 (NCHS Standard item)	373	A,S	01 02 03 04	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						05	28-31 weeks
						06	32-33 weeks
						07	34-36 weeks
						08	37-38 weeks
						09	39-40 weeks
						10	41 weeks
						11	42 weeks and over
						12	Unknown
344	1	OEGest_R5	Obstetric Estimate Recode 5 (NCHS Standard item)	373	A,S	1	Under 20 weeks
						2	20-23 weeks
						3	24-27 weeks
						4	28 weeks and over
						5	Unknown
345-348	4	FILLER					
349-352	4	DBWT	Birth Weight – Detail in Grams		A,S	0001-8165	Number of grams
						9999	Not stated birth weight
353-354	2	BWTR14	Birth Weight Recode 14		A,S	01	0249 grams or less
						02	0250 – 0349 grams
						03	0350 – 0499 grams
						04	0500 – 0999 grams
						05	1000 – 1499 grams
						06	1500 – 1999 grams
						07	2000 – 2499 grams
						08	2500 – 2999 grams
						09	3000 – 3499 grams
						10	3500 – 3999 grams
						11	4000 – 4499 grams
						12	4500 – 4999 grams
						13	5000 – 8165 grams
						14	Unknown or Not Stated

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition	
355	1	BWTR4	Birth Weight Recode 4		A,S	1 2 3 4	1499 grams or less 1500 – 2499 grams 2500 - 8165 grams Unknown or not stated	
356	1	FILLER						
357	1	ESTOFD	Estimated Time of Fetal Death	435	A	N L A U	At assessment, no labor At assessment, labor Labor, no assessment Unknown	
358	1	AUTOPSY	Was Autopsy Performed?	436	A	Y N P	Yes No Planned	
359	1	HISTOPF	Was Histological Placental Exam Performed?	437	A	Y N P	Yes No Planned	
360	1	AUTOPF	Was Autop/Hist results used in cause?	438	A	Y N X	Yes No Not applicable	
361-371	11	FILLER						
372-438		<u>Flag File for Reporting Flags</u>						
		The reporting flags indented below follow this coding structure:					0 1	Not reporting Reporting
	372	F_MEDUC	Education of Mother		A			
	373	F_CLINEST	Obstetric Estimate of Gestation		A,S			
	374	F_TOBACO	Tobacco Use		A			
	375	F_M_HT	Mother's Height		A			
	376	F_PWGT	Prepregnancy Weight		A			
	377	F_WIC	WIC		A			
	378	F_RF_PDIAB	Prepregnancy Diabetes		A			

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
379	1	F_RF_GDIAB	Gestational Diabetes		A		
380	1	F_RF_PHYPER	Prepregnancy Hypertension		A		
381	1	F_RF_GHYPER	Gestational Hypertension		A		
382	1	F_RF_ECLAMP	Hypertension Eclampsia		A		
383	1	F_RF_INFT	Infertility Treatment		A		
384	1	F_RF_INFT_DRG	Fertility Enhance Drug		A		
385	1	F_RF_INFT_ART	Reproductive Technology		A		
386	1	F_RF_CESAR	Previous Cesarean		A		
387	1	F_RF_NCESAR	Number of Previous Cesareans		A		
388	1	F_MD_PRESENT	Fetal Presentation		A		
389	1	F_MD_ROUTE	Final Route and Method of Delivery		A		
390	1	F_MD_TRIAL	Trial of Labor Attempted		A		
391	1	F_MM_RUPTUR	Ruptured Uterus		A		
392	1	F_MM_ICU	Admission to Intensive Care		A		
393-402	10	FILLER					
403	1	F_MPCB	Month Prenatal Care Began		A		
404	1	F_CMBGST	Combined Gestation		A,S		
405	1	F_CIGS_0	Cigarettes Before Pregnancy		A		
406	1	F_CIGS_1	Cigarettes 1 st Trimester		A		
407	1	F_CIGS_2	Cigarettes 2 nd Trimester		A		
408	1	F_CIGS_3	Cigarettes 3 rd Trimester		A		
409	1	F_FACILITY	Facility		A		
410	1	DelMethRecF	Delivery Method Recode Flag		A,S		
411-423	13	FILLER					
424	1	F_MORIGIN	Mother's Hispanic origin		A,S		
425	1	F_MRACE_R	Mother's Race (Revised)		A		
426	1	F_FAGE_u	Father's Age		A,S		
427	1	F_DLLB_MM	Last Live Birth Month (Revised)		A		
428	1	FILLER					
429	1	F_DLMP_MM	Last normal menses month		A,S		
430	1	FILLER					
431	1	F_DLMP_YY	Last normal menses year		A,S		
432	1	F_MAGE	Mother's Age		A,S		
433	1	F_ATTENDANT	Attendant		A,S		
434	1	FILLER					
435	1	F_ESTOFD	Estimated Time of Fetal Death (Revised)		A		
436	1	F_AUTOPSY	Autopsy (Revised)		A		
437	1	F_HISTOPF	Histological Placental Exam (Revised)		A		
438	1	F_AUTOPF	Autopsy/Histological Exam Used in Cause		A		
439	1	F_MRACEREC	Mother's Bridged Race Recode		A,S		
440	1	F_PLB	Number of previous live births		A,S		

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
441-2602	2164	FILLER					
2603-2642		40	<u>Coded Causes & Conditions</u> The fields indented below follow this structure:			A ICD10 Blank	Code Coded Value None reported
	2603-2607	5	IICOD	Initiating Cause or Condition Code	2651		
	2608-2642	35	FILLER				
2643-2645	3	IC_124_Fetal	Initiating Fetal Recode 124	2651		A	001-124
2646-2650	5	FILLER					
2651			<u>Flag File for Reporting Flags</u> The reporting flags indented below follow this coding structure:			0 1	Not reporting Reporting
	2651	1	F_ICOD Init Cause or condition	A			
2652-3150	499	FILLER					
3150		END OF RECORD					

* For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Table 1. Number of fetal deaths and fetal mortality rates: United States, 2014-2019

	Fetal deaths			Fetal mortality rate\1		
	Total\2	20-27 weeks\3	28 weeks or more\3	Total\2	20-27 weeks\3	28 weeks or more\3
Obstetric estimate of gestation\4						
2019	21,478	11,216	10,262	5.70	2.98	2.73
2018	22,459	11,844	10,615	5.89	3.11	2.79
2017	22,827	11,861	10,966	5.89	3.07	2.84
2016	23,880	12,486	11,394	6.02	3.15	2.88
2015	23,776	12,407	11,369	5.94	3.11	2.85
2014	23,980	12,652	11,328	5.98	3.16	2.83
LMP-based gestation\5						
2014	23,999	11,948	12,051	5.98	2.99	3.01

\1Rate is number of fetal deaths in specified group per 1,000 live births and fetal deaths.

\2Fetal deaths with stated or presumed period of gestation of 20 weeks or more.

\3Not stated gestational age proportionally distributed.

\4Beginning with the 2014 data year, the obstetric estimate of gestation at delivery replaces the measure based on the date of last normal menses as the standard for measuring gestational age.

\5Date of last normal menses (LMP) was used for measuring gestational

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 2. Fetal Deaths by Period of Gestation: United States, Each State and Territory, 2019

[Nonresidents of the United States are excluded]

State/territory of residence	Fetal Deaths with Stated or Presumed Gestation of 20 weeks or More			
	Total	20-27 Weeks	28 Weeks and Over	Not stated
United States\1	21,478	11,159	10,209	110
Alabama	526	280	246	-
Alaska	55	30	24	1
Arizona	497	246	251	-
Arkansas	306	173	133	-
California	2,171	1,083	1,069	19
Colorado	350	178	172	-
Connecticut	133	72	61	-
Delaware	59	21	38	-
District of Columbia	86	46	40	-
Florida	1,538	809	690	39
Georgia	976	519	453	4
Hawaii	85	57	28	-
Idaho	112	51	60	1
Illinois	829	459	370	-
Indiana	489	258	230	1
Iowa	196	94	101	1
Kansas	196	106	90	-
Kentucky	306	158	148	-
Louisiana	290	163	124	3
Maine	65	32	31	2
Maryland	462	257	205	-
Massachusetts	298	138	160	-
Michigan	652	329	321	2
Minnesota	359	179	180	-
Mississippi	348	181	166	1
Missouri	402	204	197	1
Montana	54	18	36	-
Nebraska	125	64	61	-
Nevada	230	128	102	-
New Hampshire	55	21	34	-
New Jersey	694	423	269	2
New Mexico	69	27	41	1
New York	591	286	301	4
New York City	740	457	283	-
North Carolina	784	413	371	-
North Dakota	66	35	31	-
Ohio	850	436	412	2
Oklahoma	282	135	145	2
Oregon	189	92	94	3
Pennsylvania	802	406	394	2

Rhode Island	49	19	30	-
South Carolina	345	144	201	-
South Dakota	69	40	29	-
Tennessee	498	258	238	2
Texas	1,504	774	724	6
Utah	246	121	125	-
Vermont	20	4	16	-
Virginia	485	249	233	3
Washington	484	271	206	7
West Virginia	79	36	43	-
Wisconsin	342	162	179	1
Wyoming	40	17	23	-
American Samoa	---	---	---	---
Guam	34	12	22	-
Northern Marianas	8	2	6	-
Puerto Rico	204	124	80	-
Virgin Islands	---	---	---	---

- Quantity zero.

--- Data unavailable.

1 Excludes data for the territories.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 3. Fetal Deaths by Period of Gestation and State/Territory of Occurrence: United States, 2019

State / territory	All Fetal Deaths			
	Total	Under 20 Weeks	20 Weeks and Over	Not stated
United States\1	46,007	23,017	21,438	1,552
Alabama	509	-	509	-
Alaska	53	2	50	1
Arizona	489	2	487	-
Arkansas	628	315	311	2
California	2,190	-	2,167	23
Colorado	376	1	375	-
Connecticut	141	2	139	-
Delaware	61	3	58	-
District of Columbia	127	1	126	-
Florida	1,573	4	1,527	42
Georgia	8,320	6,760	981	579
Hawaii	803	605	92	106
Idaho	133	29	104	-
Illinois	794	-	794	-
Indiana	493	-	492	1
Iowa	192	-	191	1
Kansas	196	-	196	-
Kentucky	297	4	293	-
Louisiana	328	45	280	3
Maine	63	-	61	2
Maryland	439	-	439	-
Massachusetts	297	1	296	-
Michigan	646	5	639	2
Minnesota	388	27	361	-
Mississippi	349	-	348	1
Missouri	433	12	420	1
Montana	51	-	51	-
Nebraska	133	-	133	-
Nevada	233	2	231	-
New Hampshire	55	1	54	-
New Jersey	686	19	665	2
New Mexico	61	3	57	1
New York	4,917	3,873	535	509
New York City	7,515	6,711	794	10
North Carolina	789	1	788	-
North Dakota	73	-	73	-
Ohio	861	-	859	2
Oklahoma	268	-	263	5
Oregon	229	37	191	1

Pennsylvania	1,215	415	798	2
Rhode Island	557	471	52	34
South Carolina	329	2	327	-
South Dakota	83	3	80	-
Tennessee	541	5	533	3
Texas	1,744	218	1,520	6
Utah	257	-	257	-
Vermont	24	-	24	-
Virginia	4,104	3,424	478	202
Washington	511	10	491	10
West Virginia	88	3	85	-
Wisconsin	334	1	332	1
Wyoming	31	-	31	-
American Samoa	---	---	---	---
Guam	35	-	35	-
Northern Marianas	8	-	8	-
Puerto Rico	204	-	204	-
Virgin Islands	---	---	---	---

- Quantity zero.

--- Data unavailable.

1 Excludes data for the territories.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 4. Number and percentage of fetal deaths by race and Hispanic origin of mother: United States, 2019

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Race	Number		Percentage	
	Total	Non-Hispanic	Total\1	Non-Hispanic
All races\2	21,478	16,464	100.0	100.0
One race	21,041	16,109	98.0	97.8
White	13,420	9,067	62.5	55.1
Black	6,198	5,766	28.9	35.0
American Indian and Alaska Native (AIAN)	249	214	1.2	1.3
Asian	1,036	963	4.8	5.8
Native Hawaiian and Other Pacific Islander (NHOPI)	138	99	0.6	0.6
More than one race	475	379	2.2	2.3
Two races	399	331	1.9	2.0
Black and White	196	163	0.9	1.0
Black and AIAN	21	19	0.1	0.1
Black and Asian	12	12	0.1	0.1
Black and NHOPI	10	8	0.0	0.0
AIAN and White	65	53	0.3	0.3
AIAN and Asian	5	3	0.0	0.0
AIAN and NHOPI	1	-	0.0	-
Asian and White	62	53	0.3	0.3
Asian and NHOPI	17	15	0.1	0.1
NHOPI and White	10	5	0.0	0.0
Three races	35	21	0.2	0.1
Black, AIAN and White	21	12	0.1	0.1
Black AIAN and Asian	-	-	-	-
Black, AIAN and NHOPI	-	-	-	-
Black, Asian and White	4	1	0.0	0.0
Black, Asian and NHOPI	1	1	0.0	0.0
Black, NHOPI, and White	1	1	0.0	0.0
AIAN, Asian and White	1	1	0.0	0.0
AIAN, NHOPI and White	-	-	-	-
AIAN, Asian and NHOPI	-	-	-	-
Asian, NHOPI and White	7	5	0.0	0.0
Four races	3	3	0.0	0.0
Black, AIAN, Asian and White	1	1	0.0	0.0
Black, AIAN, Asian, and NHOPI	-	-	-	-
Black, AIAN, NHOPI and White	-	-	-	-
Black, Asian, NHOPI and White	1	1	0.0	0.0
AIAN, Asian, NHOPI and White	1	1	0.0	0.0
Five races	-	-	-	-
Black, AIAN, Asian, NHOPI and White	-	-	-	-

- Quantity zero.

0.0 Quantity more than zero but less than 0.5.

\1 Includes births to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, non-Hispanic, multiple-race women, and births with origin not stated.

\2 Race and Hispanic origin are reported separately on birth certificates. Persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards.

NOTE: Race categories are consistent with the 1997 Office of Management and Budget standards.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 5A. Number and percentage of fetal deaths by selected demographic and health characteristics: United States, 2019

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

	Number	Percent
Day of week		
Total	21,478	100
Sunday	2,319	10.8
Monday	2,591	12.1
Tuesday	3,357	15.6
Wednesday	3,472	16.2
Thursday	3,362	15.7
Friday	3,433	16
Saturday	2,944	13.7
Month of delivery		
Total	21,478	100
January	1,836	8.5
February	1,609	7.5
March	1,844	8.6
April	1,798	8.4
May	1,982	9.2
June	1,845	8.6
July	1,826	8.5
August	1,866	8.7
September	1,771	8.2
October	1,814	8.4
November	1,653	7.7
December	1,634	7.6
Place of delivery		
Total	21,478	100
Hospital\1	20,923	97.4
Freestanding birthing center	29	0.1
Clinic or doctor's office	73	0.3
Residence	305	1.4
Other	109	0.5
Not specified	39	0.2
Age of mother		
Total	21,478	100
<15	17	0.1
15-19	1,190	5.5
20-24	4,112	19.1
25-29	5,750	26.8
30-34	5,577	26
35-39	3,638	16.9
40-44	1,099	5.1
45-49	88	0.4
50-54	7	0

Race and Hispanic origin of mother		
Total\2	21,478	100
Non-Hispanic, single race\3		
Non-Hispanic white	9,067	42.2
Non-Hispanic black	5,766	26.8
Non-Hispanic American Indian or Alaska Native	214	1
Non-Hispanic Asian	963	4.5
Non-Hispanic Native Hawaiian or Other Pacific Islander	99	0.5
Non-Hispanic multiple race	355	1.7
Hispanic\4	4,264	19.9
Unknown	750	3.5
Hispanic origin of mother		
Total	21,478	100
Non-Hispanic		
Mexican	2,427	11.3
Puerto Rican	369	1.7
Cuban	112	0.5
Central or South American	683	3.2
Dominican	155	0.7
Otherand unknown Hispanic	518	2.4
Unknown	750	3.5
Education of mother\5		
Total	21,429	100
8th grade or less	732	3.4
9th-12th grade with no diploma	2,056	9.6
High school graduate\6	6,487	30.3
Some college credit, but no degree	3,734	17.4
Associate's degree\7	1,474	6.9
Bachelor's degree\8	2,900	13.5
Master's degree\9	1,138	5.3
Doctorate or professional degree\10	346	1.6
Unknown	2,562	12
Age of father\11		
Total	16,928	100
<15	5	0
15-19	335	2
20-24	1,899	11.2
25-29	3,195	18.9
30-34	3,569	21.1
35-39	2,755	16.3
40-44	1,288	7.6
45-49	511	3
50-54	166	1
55-98	99	0.6
Unknown	3,106	18.3
Trimester prenatal care began		
Total	21,478	100
1st trimester	13,609	63.4

2nd trimester	3,163	14.7
3rd trimester	356	1.7
No care	1,721	8
Unknown	2,629	12.2
Mother received WIC food for herself during this pregnancy\12		
Total	20,518	100
Yes	4,690	22.9
No	12,740	62.1
Not stated	3,088	15.1
Tobacco use during pregnancy		
Total	21,478	100
Yes	1,909	8.9
No	17,832	83
Unknown	1,737	8.1
Maternal pre-pregnancy body mass index\13		
Total	21,393	100
Underweight (BMI less than 18.5)	504	2.4
Normal weight (BMI of 18.5 to 24.9)	6,050	28.3
Overweight (BMI of 25.0 to 29.9)	4,961	23.2
Obese (BMI of 30.0 and over)	7,203	33.7
Not stated\3	2,675	12.5
Method of delivery		
Total	21,478	100
Vaginal (excludes vaginal after previous C-section)	14,967	69.7
Vaginal after previous C-section	1,657	7.7
Primary C-section	1,905	8.9
Repeat C-section	1,263	5.9
Vaginal (unknown if previous c-section)	621	2.9
C-section (unknown if previous c-section)	83	0.4
Not stated	982	4.6
Attendant at delivery\14		
Total	20,989	100
Doctor of medicine	17,840	85
Doctor of osteopathy	1,754	8.4
Certified nurse midwife	672	3.2
Other midwife	36	0.2
Other	536	2.6
Unspecified	151	0.7
Plurality		
Total	21,478	100
Singleton	19,937	92.8
Twin	1,455	6.8
Triplet	78	0.4
Quadruplet or higher	8	0
Sex		

Total	21,478	100
Male	11,268	52.5
Female	10,210	47.5
Obstetric estimate of gestation		
Total	21,478	100
20-23 weeks	7,892	36.7
24-27 weeks	3,267	15.2
28-31 weeks	2,625	12.2
32-33 weeks	1,506	7
34-36 weeks	2,636	12.3
37-38 weeks	2,002	9.3
39-40 weeks	1,277	5.9
41 weeks	123	0.6
42 weeks and over	40	0.2
Unknown	110	0.5
Birthweight		
Total	21,478	100
<500 grams	7,308	34
0500 - 0999 grams	3,729	17.4
1000 - 1499 gram	1,920	8.9
1500 - 1999 gram	1,736	8.1
2000 - 2499 gram	1,641	7.6
2500 - 2999 gram	1,581	7.4
3000 - 3499 gram	1,191	5.5
3500 - 3999 gram	560	2.6
4000 - 4499 gram	200	0.9
4500 - 4999 gram	91	0.4
5000 - 8165 gram	45	0.2
Unknown	1,476	6.9

0.0 Quantity more than zero but less than 0.05.

- Quantity zero.

\1Includes fetal deaths occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and births with origin not stated.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death.

\4Includes all persons of Hispanic origin of any race.

\5Education of mother is not reported by Rhode Island for 2019.

\6Includes General Educational Development (GED).

\7Includes Associate in Arts and Associate in Science.

\8Includes Bachelor in Arts and Bachelor in Science.

\9Includes Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\10Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

\11Alabama, Arkansas, the District of Columbia, Georgia, New York (including New York City), Tennessee, Virginia, and Wisconsin reported father's age for 2019 but had an unknown level of 50% or higher and were flagged as not reporting.

\12Mother received WIC food for herself during this pregnancy is not reported by New York City. The District of Columbia, Hawaii, and Rhode Island reported WIC for 2019 but had an unknown level of 50% or higher and were flagged as not reporting.

\13Hawaii reported maternal pre-pregnancy body mass index for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

\14Indiana reported attendant at delivery for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 5B. Number and percentage of fetal deaths by selected demographic and health characteristics: Guam, Northern Marianas, and Puerto Rico, 2019

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

	Number	Percent
Day of week		
Total	246	100.0
Sunday	23	9.3
Monday	22	8.9
Tuesday	37	15.0
Wednesday	37	15.0
Thursday	46	18.7
Friday	49	19.9
Saturday	32	13.0
Month of delivery		
Total	246	100.0
January	19	7.7
February	19	7.7
March	20	8.1
April	25	10.2
May	30	12.2
June	23	9.3
July	21	8.5
August	20	8.1
September	19	7.7
October	15	6.1
November	22	8.9
December	13	5.3
Place of delivery		
Total	246	100.0
Hospital\1	243	98.8
Freestanding birthing center	-	-
Clinic or doctor's office	-	-
Residence	1	0.4
Other	1	0.4
Not specified	1	0.4
Age of mother		
Total	246	100.0
<15	1	0.4
15-19	27	11.0
20-24	56	22.8
25-29	72	29.3
30-34	41	16.7
35-39	29	11.8
40-44	19	7.7
45-49	1	0.4
50-54	-	-

Race and Hispanic origin of mother		
Total\2	246	100.0
Non-Hispanic, single race\3		
Non-Hispanic white	6	2.4
Non-Hispanic black	-	-
Non-Hispanic American Indian or Alaska I	-	-
Non-Hispanic Asian	7	2.8
Non-Hispanic Native Hawaiian or Other P:	32	13.0
Non-Hispanic multiple race	1	0.4
Hispanic\4	198	80.5
Unknown	2	0.8
Hispanic origin of mother		
Total	246	100.0
Non-Hispanic		
Mexican	-	-
Puerto Rican	193	78.5
Cuban	1	0.4
Central or South American	-	-
Dominican	4	1.6
Otherand unknown Hispanic	-	-
Unknown	2	0.8
Education of mother		
Total	246	100.0
8th grade or less	7	2.8
9th-12th grade with no diploma	21	8.5
High school graduate\5	75	30.5
Some college credit, but no degree	38	15.4
Associate's degree\6	37	15.0
Bachelor's degree\7	49	19.9
Master's degree\8	13	5.3
Doctorate or professional degree\9	3	1.2
Unknown	3	1.2
Age of father\10		
Total	212	100.0
<15	-	-
15-19	10	4.7
20-24	28	13.2
25-29	44	20.8
30-34	18	8.5
35-39	21	9.9
40-44	12	5.7
45-49	8	3.8
50-54	1	0.5
55-98	-	-
Unknown	70	33.0
Trimester prenatal care began		
Total	246	100.0

1st trimester	165	67.1
2nd trimester	36	14.6
3rd trimester	10	4.1
No care	26	10.6
Unknown	9	3.7

Mother received WIC food for herself during this pregnancy

Total	246	100.0
Yes	157	63.8
No	68	27.6
Not stated	21	8.5

Tobacco use during pregnancy

Total	246	100.0
Yes	8	3.3
No	230	93.5
Unknown	8	3.3

Maternal pre-pregnancy body mass index

Total	246	100.0
Underweight (BMI less than 18.5)	12	4.9
Normal weight (BMI of 18.5 to 24.9)	80	32.5
Overweight (BMI of 25.0 to 29.9)	60	24.4
Obese (BMI of 30.0 and over)	80	32.5
Not stated\3	14	5.7

Method of delivery

Total	246	100.0
Vaginal (excludes vaginal after previous C-section)	174	70.7
Vaginal after previous C-section	19	7.7
Primary C-section	28	11.4
Repeat C-section	17	6.9
Vaginal (unknown if previous c-section)	3	1.2
C-section (unknown if previous c-section)	3	1.2
Not stated	2	0.8

Attendant at delivery

Total	246	100.0
Doctor of medicine	234	95.1
Doctor of osteopathy	4	1.6
Certified nurse midwife	5	2.0
Other midwife	1	0.4
Other	1	0.4
Unspecified	1	0.4

Plurality

Total	246	100.0
Singleton	237	96.3
Twin	9	3.7
Triplet	-	-
Quadruplet or higher	-	-

Sex		
Total	246	100.0
Male	129	52.4
Female	117	47.6
Obstetric estimate of gestation		
Total	246	100.0
20-23 weeks	100	40.7
24-27 weeks	38	15.4
28-31 weeks	32	13.0
32-33 weeks	11	4.5
34-36 weeks	26	10.6
37-38 weeks	26	10.6
39-40 weeks	13	5.3
41 weeks	-	-
42 weeks and over	-	-
Unknown	-	-
Birthweight		
Total	246	100.0
<500 grams	84	34.1
0500 - 0999 grams	56	22.8
1000 - 1499 gram	19	7.7
1500 - 1999 gram	21	8.5
2000 - 2499 gram	21	8.5
2500 - 2999 gram	15	6.1
3000 - 3499 gram	10	4.1
3500 - 3999 gram	4	1.6
4000 - 4499 gram	5	2.0
4500 - 4999 gram	1	0.4
5000 - 8165 gram	-	-
Unknown	10	4.1

0.0 Quantity more than zero but less than 0.05.

- Quantity zero.

\1Includes births occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and births with origin not stated.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death.

\4Includes all persons of Hispanic origin of any race.

\5Includes General Educational Development (GED).

\6Includes Associate in Arts and Associate in Science.

\7Includes Bachelor in Arts and Bachelor in Science.

\8Includes Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\9Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

10\Guam reported father's age for 2019 but had an unknown level of 50% or higher and were flagged as not reporting.

NOTE: American Samoa and the Virgin Islands did not report data for 2019

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 6A. Number and percentage of fetal deaths by selected health characteristics: United States, 2019

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Risk factor	Total	Number reported	Unknown	Per 1,000
Diabetes				
Prepregnancy (Diagnosis prior to this pregnancy)	21,478	781	1,089	38.3
Gestational (Diagnosis in this pregnancy)	21,478	936	1,089	45.9
Hypertension				
Prepregnancy (Chronic)	21,478	1,233	1,089	60.5
Gestational (PIH, preeclampsia)	21,478	1,446	1,089	70.9
Eclampsia	21,478	154	1,089	7.6
Pregnancy resulted from infertility treatment	21,478	609	1,089	29.9
Mother had a previous cesarean delivery\1	21,478	2,980	1,089	14.6
Maternal morbidity\2				
Ruptured uterus	21,478	134	827	648.9
Admission to intensive care unit	21,478	478	827	2,314.7

\1 Mother had a previous cesarean delivery per 100 fetal deaths.

\2 Maternal morbidities per 100,000 fetal deaths.

NOTE: PIH is pregnancy-induced hypertension.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 6B. Number and percentage of fetal deaths by selected health characteristics: Guam, Northern Marianas, and Puerto Rico, 2019

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Risk factor	Total	Number reported	Unknown	Per 1,000
Diabetes				
Pregpregnancy (Diagnosis prior to this pregnancy)	246	8	6	33.3
Gestational (Diagnosis in this pregnancy)	246	10	6	41.7
Hypertension				
Pregpregnancy (Chronic)	246	18	6	75.0
Gestational (PIH, preeclampsia)	246	14	6	58.3
Eclampsia	246	2	6	8.3
Pregnancy resulted from infertility treatment	246	3	6	12.5
Mother had a previous cesarean delivery\1	246	36	6	15.0
Maternal morbidity\2				
Ruptured uterus	246	4	12	1,709.4
Admissioin to intensive care unit	246	3	12	1,282.1

\1 Mother had a previous cesarean delivery per 100 fetal deaths.

\2 Maternal morbidities per 100,000 fetal deaths.

NOTE: PIH is pregnancy-induced hypertension.

NOTE: American Samoa and the Virgin Islands did not report data for 2019

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 7A. Deaths according to 124 selected causes of fetal death: 42 states and the District of Columbia, 2019
 [Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Cause of death (based on ICD-10)	Number
All causes	15,636
Certain infectious and parasitic diseases	-
Congenital syphilis	-
Human immunodeficiency virus (HIV) disease	-
Other viral diseases	-
Other and unspecified infectious and parasitic diseases	-
Malignant neoplasms	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior	88
Anemias	8
Endocrine, nutritional and metabolic diseases	2
Short stature, not elsewhere classified	-
Cystic fibrosis	-
Other endocrine, nutritional and metabolic diseases	2
Meningitis	4
Other diseases of nervous system and sense organs	-
Umbilical hernia	-
Other hernia	-
Other and unspecified diseases of digestive system	-
All other diseases, excluding perinatal conditions, congenital anomalies, and symptoms, signs and ill-defined conditions	12
Certain conditions originating in the perinatal period	13,889
Fetus affected by maternal conditions that may be unrelated to present pregnancy	1,614
Fetus affected by maternal hypertensive disorders	927
Fetus affected by maternal renal and urinary tract diseases	27
Fetus affected by maternal infectious and parasitic diseases	113
Fetus affected by other maternal circulatory and respiratory diseases	64
Fetus affected by maternal nutritional disorders	4
Fetus affected by maternal injury	56
Fetus affected by surgical procedure on mother	6
Fetus affected by other medical procedures and maternal conditions	393
Fetus affected by unspecified maternal condition	24
Fetus affected by maternal complications of pregnancy	1,960
Fetus affected by incompetent cervix	264
Fetus affected by premature rupture of membranes	1,301
Fetus affected by oligohydramnios	87
Fetus affected by polyhydramnios	48
Fetus affected by ectopic pregnancy	5
Fetus affected by multiple pregnancy	193
Fetus affected by maternal death	14
Fetus affected by malpresentation before labor	7
Fetus affected by other and unspecified maternal complications of pregnancy	41
Fetus affected by complications of placenta, cord and membranes	3,981
Fetus affected by placenta previa	27
Fetus affected by other forms of placental separation and hemorrhage	1,314
Fetus affected by other and unspecified morphological and functional abnormalities of placenta	694
Fetus affected by placental transfusion syndromes	111
Fetus affected by prolapsed cord	108
Fetus affected by other compression of umbilical cord	902
Fetus affected by other and unspecified conditions of umbilical cord	362
Fetus affected by chorioamnionitis	447
Fetus affected by other and unspecified abnormalities of membranes	16
Fetus affected by other complications of labor and delivery	114
Fetus affected by breech delivery and extraction	9
Fetus affected by other malpresentation, malposition and disproportion during labor and delivery	7
Fetus affected by forceps delivery	-
Fetus affected by delivery by vacuum extractor (ventouse)	-
Fetus affected by cesarean delivery	-
Fetus affected by precipitate delivery	1
Fetus affected by abnormal uterine contractions	2
Fetus affected by other and unspecified complications of labor and delivery	95
Fetus affected by noxious influences transmitted via placenta	162
Slow fetal growth and fetal malnutrition	173
Disorders related to short gestation and low birth weight, not elsewhere classified	301
Extremely low birth weight or extreme immaturity	169
Other low birth weight and preterm	132
Disorders related to long gestation and high birth weight	5
Exceptionally large size and other heavy for gestational age fetus	2
Post-term, not heavy for gestational age fetus	3
Birth trauma	1
Intracranial laceration and hemorrhage due to birth injury and other injuries to central nervous system	1
Other birth trauma	-
Intrauterine hypoxia and birth asphyxia	12
Intrauterine hypoxia first noted before onset of labor	-
Intrauterine hypoxia first noted during labor and delivery	-
Birth asphyxia, unspecified	11
Other respiratory condition originating in the perinatal period	1
Congenital pneumonia	6
Aspiration syndromes	-
Interstitial emphysema and related conditions originating in the perinatal period	3
Atelectasis	-
Other respiratory system disorders	-
Infections specific to the perinatal period	12
Congenital rubella syndrome	-
Congenital cytomegalovirus infection	2
Congenital herpes (herpes simplex) infection	1
Congenital viral hepatitis	-
Bacterial sepsis	-
Congenital tuberculosis	-
Congenital toxoplasmosis	1
Other infections specific to the perinatal period	8
Fetal hemorrhage	21
Hemolytic disease of fetus	17
Rh isoimmunization of fetus	9
ABO isoimmunization of fetus	1
Other hemolytic disease of fetus	7
Hydrops fetalis due to hemolytic disease	-
Perinatal jaundice	2
Hematological disorders	3
Transient endocrine and metabolic disorders specific to fetus	381
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus	376
Other transient endocrine and metabolic disorders specific to fetus	5
Digestive system disorders of fetus	2
Other conditions originating in the perinatal period	5,122
Hydrops fetalis not due to hemolytic disease	194
Fetal death of unspecified cause	4,899
Withdrawal symptoms from maternal use of drugs of addiction	-
Termination of pregnancy	-
Complications of intrauterine procedures, not elsewhere classified	-
All other specified conditions originating in the perinatal period	29
Condition originating in the perinatal period, unspecified	-
Congenital malformations, deformations and chromosomal abnormalities	1,613
Congenital malformations of nervous system	297
Anencephaly and similar malformations	152
Encephalocele	8
Microcephaly	6
Congenital hydrocephalus	47
Reduction deformities of brain	41
Other congenital malformations of brain	16
Spina bifida	15
Other congenital malformations of spinal cord and nervous system	12
Congenital malformations of eye, ear, face and neck	5
Congenital malformations of heart	195

Other congenital malformations of circulatory system.....	Q25-Q28	57
Congenital malformations of lung.....	Q33	9
Other congenital malformations of respiratory system.....	Q30-Q32,Q34	13
Congenital malformations of digestive system.....	Q35-Q45	34
Congenital malformations of genital organs.....	Q50-Q56	14
Congenital malformations of urinary system.....	Q60-Q64	120
Renal agenesis and other reduction defects of kidney.....	Q60	61
Cystic kidney disease.....	Q61	30
Other congenital malformations of urinary system.....	Q62-Q64	29
Congenital malformations and deformations of musculoskeletal system, limbs and integument.....	Q65-Q85	203
Other congenital malformations.....	Q86-Q89	115
Conjoined twins.....	Q89.4	2
Multiple congenital malformations, not elsewhere classified.....	Q89.7	45
All other congenital malformations.....	Q86-Q87,Q89.0-Q89.3,Q89.8-Q89.9	68
Chromosomal abnormalities, not elsewhere classified.....	Q90-Q99	551
Down's syndrome.....	Q90	120
Edward's syndrome.....	Q91.0-Q91.3	199
Patau's syndrome.....	Q91.4-Q91.7	55
Other chromosomal abnormalities, not elsewhere classified.....	Q92-Q99	177
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified.....	R00-R99	7
External causes of mortality.....	*U01.V01-Y84	13
Accidents (unintentional injuries).....	V01-X59	13
Assault (homicide).....	*U01.X85-Y09	-
Complications of medical and surgical care.....	Y40-Y84	-
Other external causes.....	Y10-Y36	-

- Quantity zero.

NOTE: Excludes data for jurisdictions (Georgia, Hawaii, Michigan, Mississippi, New York City, North Carolina, North Dakota, and Vermont) for which the cause of death was unspecified (P95) for 50% or more of records. Also excludes data for California which did not report cause of death based on the 2003 revision of the U.S. Standard Report of Fetal Death.

Source: National Center for Health Statistics, National Vital Statistics System.

Table 7B. Deaths according to 124 selected causes of fetal death: Puerto Rico, 2019
 [Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Cause of death (based on ICD-10)	Number
All causes ¹	204
Certain infectious and parasitic diseases.....A00-B99	-
Congenital syphilis.....A50	-
Human immunodeficiency virus (HIV) disease.....B20-B24	-
Other viral diseases.....A80-B19, B25-B34	-
Other and unspecified infectious and parasitic diseases.....A00-A49, A51-A79, B35-B99	-
Malignant neoplasms, excluding perinatal conditions: congenital anomalies, and symptoms, signs and ill-defined conditions.....C00-C97	-
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior.....D00-D48	-
Anemias.....D50-D64	-
Endocrine, nutritional and metabolic diseases.....E00-E88	-
Short stature, not elsewhere classified.....E34.3	-
Cystic fibrosis.....E84	-
Other endocrine, nutritional and metabolic diseases.....E00-E32, E34.0-E34.2, E34.4-E34.9, E40-E83, E85-E88	-
Meningitis.....G00, G03	-
Other diseases of nervous system and sense organs.....G04-H93	-
Umbilical hernia.....K42	-
Other hernia.....K40-K41, K43-K46	-
Other and unspecified diseases of digestive system.....K00-K38, K50-K92	-
All other diseases, excluding perinatal conditions: congenital anomalies, and symptoms, signs and ill-defined conditions.....D65-D89, I00-J98, L00-N98, U04	170
Certain conditions originating in the perinatal period.....P00-P96	28
Fetus affected by maternal conditions that may be unrelated to present pregnancy.....P00	17
Fetus affected by maternal hypertensive disorders.....P00.0	1
Fetus affected by maternal renal and urinary tract diseases.....P00.1	2
Fetus affected by maternal infectious and parasitic diseases.....P00.2	1
Fetus affected by other maternal circulatory and respiratory diseases.....P00.3	-
Fetus affected by maternal nutritional disorders.....P00.4	-
Fetus affected by maternal injury.....P00.5	-
Fetus affected by surgical procedure on mother.....P00.6	-
Fetus affected by other medical procedures and maternal conditions.....P00.7-P00.8	7
Fetus affected by unspecified maternal condition.....P00.9	46
Fetus affected by maternal complications of pregnancy.....P01	18
Fetus affected by incompetent cervix.....P01.0	19
Fetus affected by premature rupture of membranes.....P01.1	4
Fetus affected by oligohydramnios.....P01.2	-
Fetus affected by polyhydramnios.....P01.3	-
Fetus affected by ectopic pregnancy.....P01.4	-
Fetus affected by multiple pregnancy.....P01.5	4
Fetus affected by maternal death.....P01.6	-
Fetus affected by malpresentation before labor.....P01.7	1
Fetus affected by other and unspecified maternal complications of pregnancy.....P01.8-P01.9	48
Fetus affected by complications of placenta, cord and membranes.....P02	2
Fetus affected by placenta previa.....P02.0	14
Fetus affected by other forms of placental separation and hemorrhage.....P02.1	14
Fetus affected by other and unspecified morphological and functional abnormalities of placenta.....P02.2	-
Fetus affected by placental transfusion syndromes.....P02.3	-
Fetus affected by prolapsed cord.....P02.4	-
Fetus affected by other compression of umbilical cord.....P02.5	11
Fetus affected by other and unspecified conditions of umbilical cord.....P02.6	5
Fetus affected by chorioamnionitis.....P02.7	2
Fetus affected by other and unspecified abnormalities of membranes.....P02.8-P02.9	-
Fetus affected by other complications of labor and delivery.....P03	1
Fetus affected by breech delivery and extraction.....P03.0	-
Fetus affected by other malpresentation, malposition and disproportion during labor and delivery.....P03.1	-
Fetus affected by forceps delivery.....P03.2	-
Fetus affected by delivery by vacuum extractor (ventouse).....P03.3	-
Fetus affected by cesarean delivery.....P03.4	-
Fetus affected by precipitate delivery.....P03.5	-
Fetus affected by abnormal uterine contractions.....P03.6	-
Fetus affected by other and unspecified complications of labor and delivery.....P03.8-P03.9	1
Fetus affected by noxious influences transmitted via placenta.....P04	2
Slow fetal growth and fetal malnutrition.....P05	12
Disorders related to short gestation and low birth weight, not elsewhere classified.....P07	5
Extremely low birth weight or extreme immaturity.....P07.0, P07.2	7
Other low birth weight and preterm.....P07.1, P07.3	-
Disorders related to long gestation and high birth weight.....P08	-
Exceptionally large size and other heavy for gestational age fetus.....P08.0-P08.1	-
Post-term, not heavy for gestational age fetus.....P08.2	-
Birth trauma.....P10-P15	-
Intracranial laceration and hemorrhage due to birth injury and other injuries to central nervous system.....P10-P11	-
Other birth trauma.....P12-P15	-
Intrauterine hypoxia and birth asphyxia.....P20-P21	-
Intrauterine hypoxia first noted before onset of labor.....P20.0	-
Intrauterine hypoxia first noted during labor and delivery.....P20.1	-
Intrauterine hypoxia, unspecified.....P20.9	-
Birth asphyxia.....P21	-
Other respiratory conditions originating in the perinatal period.....P22.8-P22.9, P23-P28	-
Congenital pneumonia.....P23	-
Aspiration syndromes.....P24	-
Interstitial emphysema and related conditions originating in the perinatal period.....P25	-
Atelectasis.....P28.0-P28.1	-
Other respiratory system disorders.....P22.8-P22.9, P26-P27, P28.2-P28.9	-
Infections specific to the perinatal period.....P35-P39	-
Congenital rubella syndrome.....P35.0	-
Congenital cytomegalovirus infection.....P35.1	-
Congenital herpesviral (herpes simplex) infection.....P35.2	-
Congenital viral hepatitis.....P35.3	-
Bacterial sepsis.....P36	-
Congenital tuberculosis.....P37.0	-
Congenital toxoplasmosis.....P37.1	-
Other infections specific to the perinatal period.....P35.8-P35.9, P37.2-P37.9, P38-P39	-
Fetal hemorrhage.....P50-P54	-
Hemolytic disease of fetus.....P55-P56	-
Rh isoimmunization of fetus.....P55.0	-
ABO isoimmunization of fetus.....P55.1	-
Other hemolytic disease of fetus.....P55.8-P55.9	-
Hydrops fetalis due to hemolytic disease.....P56	-
Perinatal jaundice.....P57-P59	-
Hematological disorders.....P60-P61	-
Transitory endocrine and metabolic disorders specific to fetus.....P70-P74	8
Syndrome of infant of a diabetic mother and neonatal diabetes mellitus.....P70.0-P70.2	8
Other transitory endocrine and metabolic disorders specific to fetus.....P70.3-P70.9, P71-P74	-
Digestive system disorders of fetus.....P76-P78	-
Other conditions originating in the perinatal period.....P29, P80-P96	25
Hydrops fetalis not due to hemolytic disease.....P83.2	6
Fetal death of unspecified cause.....P95	16
Withdrawal symptoms from maternal use of drugs of addiction.....P96.1	-
Termination of pregnancy.....P96.4	-
Complications of intrauterine procedures, not elsewhere classified.....P96.5	-
All other specified conditions originating in the perinatal period.....P29, P83.0-P83.1, P83.3-P83.9, P91, P94, P96.0, P96.3, P96.8	3
Condition originating in the perinatal period, unspecified.....P96.9	-
Congenital malformations, deformations and chromosomal abnormalities.....Q00-Q99	33
Congenital malformations of nervous system.....Q00-Q07	7
Anencephaly and similar malformations.....Q00	2
Encephalocele.....Q01	1
Microcephaly.....Q02	-
Congenital hydrocephalus.....Q03	-
Reduction deformities of brain.....Q04.0-Q04.3	2
Other congenital malformations of brain.....Q04.4-Q04.9	1
Spina bifida.....Q05	-

Other congenital malformations of spinal cord and nervous system.....	Q06-Q07	1
Congenital malformations of eye, ear, face and neck.....	Q10-Q18	1
Congenital malformations of heart.....	Q20-Q24	3
Other congenital malformations of circulatory system.....	Q25-Q28	1
Congenital malformations of lung.....	Q33	-
Other congenital malformations of respiratory system.....	Q30-Q32,Q34	-
Congenital malformations of digestive system.....	Q35-Q45	1
Congenital malformations of genital organs.....	Q50-Q56	-
Congenital malformations of urinary system.....	Q60-Q64	2
Renal agenesis and other reduction defects of kidney.....	Q60	-
Cystic kidney disease.....	Q61	2
Other congenital malformations of urinary system.....	Q62-Q64	-
Congenital malformations and deformations of musculoskeletal system, limbs and integument.....	Q65-Q85	7
Other congenital malformations.....	Q86-Q89	3
Conjoined twins.....	Q89.4	-
Multiple congenital malformations, not elsewhere classified.....	Q89.7	1
All other congenital malformations.....	Q86-Q87,Q89.0-Q89.3,Q89.8-Q89.9	2
Chromosomal abnormalities, not elsewhere classified.....	Q90-Q99	8
Down's syndrome.....	Q90	1
Edward's syndrome.....	Q91.0-Q91.3	5
Patau's syndrome.....	Q91.4-Q91.7	1
Other chromosomal abnormalities, not elsewhere classified.....	Q92-Q99	1
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified.....	R00-R99	-
External causes of mortality.....	*U01,V01-Y84	-
Accidents (unintentional injuries).....	V01-X59	-
Assault (homicide).....	*U01,X85-Y09	-
Complications of medical and surgical care.....	Y40-Y84	-
Other external causes.....	Y10-Y36	-

- Quantity zero.

NOTE: Excludes data for jurisdictions (Guam and Northern Marianas) for which the cause of death was unspecified (P95) for 50% or more of records.

Source: National Center for Health Statistics, National Vital Statistics System.

MOTHER	1. NAME OF FETUS (optional-at the discretion of the parents)		2. TIME OF DELIVERY (24hr)	3. SEX (M/F/Unk)	4. DATE OF DELIVERY (Mo/Day/Yr)		
	5a. CITY, TOWN, OR LOCATION OF DELIVERY	7. PLACE WHERE DELIVERY OCCURRED (Check one)		8. FACILITY NAME (If not institution, give street and number)			
	5b. ZIP CODE OF DELIVERY	<input type="checkbox"/> Hospital <input type="checkbox"/> Freestanding birthing center					
	6. COUNTY OF DELIVERY	<input type="checkbox"/> Home Delivery: Planned to deliver at home? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Clinic/Doctor's office <input type="checkbox"/> Other (Specify) _____					
	10a. MOTHER'S CURRENT LEGAL NAME (First, Middle, Last, Suffix)				10b. DATE OF BIRTH (Mo/Day/Yr)		
	10c. MOTHER'S NAME PRIOR TO FIRST MARRIAGE (First, Middle, Last, Suffix)			10d. BIRTHPLACE (State, Territory, or Foreign Country)			
	11a. RESIDENCE OF MOTHER-STATE	11b. COUNTY		11c. CITY, TOWN, OR LOCATION			
	11d. STREET AND NUMBER		11e. APT. NO.	11f. ZIP CODE	11g. INSIDE CITY LIMITS? <input type="checkbox"/> Yes <input type="checkbox"/> No		
	12a. FATHER'S CURRENT LEGAL NAME (First, Middle, Last, Suffix)		12b. DATE OF BIRTH (Mo/Day/Yr)	12c. BIRTHPLACE (State, Territory, or Foreign Country)			
	13. METHOD OF DISPOSITION: <input type="checkbox"/> Burial <input type="checkbox"/> Cremation <input type="checkbox"/> Hospital Disposition <input type="checkbox"/> Donation <input type="checkbox"/> Removal from State <input type="checkbox"/> Other (Specify) _____						
ATTENDANT AND REGISTRATION INFORMATION		14. ATTENDANT'S NAME, TITLE, AND NPI NAME: _____ NPI: _____ TITLE: <input type="checkbox"/> MD <input type="checkbox"/> DO <input type="checkbox"/> CNM/CM <input type="checkbox"/> OTHER MIDWIFE <input type="checkbox"/> OTHER (Specify) _____	15. NAME AND TITLE OF PERSON COMPLETING REPORT Name _____ Title _____	16. DATE REPORT COMPLETED ____/____/____ MM DD YYYY	17. DATE RECEIVED BY REGISTRAR ____/____/____ MM DD YYYY		
		18. CAUSE/CONDITIONS CONTRIBUTING TO FETAL DEATH					
CAUSE OF FETAL DEATH		18a. INITIATING CAUSE/CONDITION (AMONG THE CHOICES BELOW, PLEASE SELECT THE <u>ONE</u> WHICH MOST LIKELY BEGAN THE SEQUENCE OF EVENTS RESULTING IN THE DEATH OF THE FETUS) Maternal Conditions/Diseases (Specify) _____ Complications of Placenta, Cord, or Membranes <input type="checkbox"/> Rupture of membranes prior to onset of labor <input type="checkbox"/> Abruptio placenta <input type="checkbox"/> Placental insufficiency <input type="checkbox"/> Prolapsed cord <input type="checkbox"/> Chorioamnionitis <input type="checkbox"/> Other (Specify) _____ Other Obstetrical or Pregnancy Complications (Specify) _____ Fetal Anomaly (Specify) _____ Fetal Injury (Specify) _____ Fetal Infection (Specify) _____ Other Fetal Conditions/Disorders (Specify) _____ <input type="checkbox"/> Unknown		18b. OTHER SIGNIFICANT CAUSES OR CONDITIONS (SELECT OR SPECIFY ALL OTHER CONDITIONS CONTRIBUTING TO DEATH IN ITEM 18b) Maternal Conditions/Diseases (Specify) _____ Complications of Placenta, Cord, or Membranes <input type="checkbox"/> Rupture of membranes prior to onset of labor <input type="checkbox"/> Abruptio placenta <input type="checkbox"/> Placental insufficiency <input type="checkbox"/> Prolapsed cord <input type="checkbox"/> Chorioamnionitis <input type="checkbox"/> Other (Specify) _____ Other Obstetrical or Pregnancy Complications (Specify) _____ Fetal Anomaly (Specify) _____ Fetal Injury (Specify) _____ Fetal Infection (Specify) _____ Other Fetal Conditions/Disorders (Specify) _____ <input type="checkbox"/> Unknown			
		18c. WEIGHT OF FETUS (grams preferred, specify unit) _____ <input type="checkbox"/> grams <input type="checkbox"/> lb/oz		18e. ESTIMATED TIME OF FETAL DEATH <input type="checkbox"/> Dead at time of first assessment, no labor ongoing <input type="checkbox"/> Dead at time of first assessment, labor ongoing <input type="checkbox"/> Died during labor, after first assessment <input type="checkbox"/> Unknown time of fetal death		18f. WAS AN AUTOPSY PERFORMED? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Planned	
		18d. OBSTETRIC ESTIMATE OF GESTATION AT DELIVERY _____ (completed weeks)				18g. WAS A HISTOLOGICAL PLACENTAL EXAMINATION PERFORMED? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Planned	
						18h. WERE AUTOPSY OR HISTOLOGICAL PLACENTAL EXAMINATION RESULTS USED IN DETERMINING THE CAUSE OF FETAL DEATH? <input type="checkbox"/> Yes <input type="checkbox"/> No	

Mother's Name _____
 Mother's Medical Record No. _____

MOTHER

19. MOTHER'S EDUCATION (Check the box that best describes the highest degree or level of school completed at the time of delivery)		20. MOTHER OF HISPANIC ORIGIN? (Check the box that best describes whether the mother is Spanish/Hispanic/Latina. Check the "No" box if mother is not Spanish/Hispanic/Latina)		21. MOTHER'S RACE (Check one or more races to indicate what the mother considers herself to be)																					
<input type="checkbox"/> 8th grade or less <input type="checkbox"/> 9th - 12th grade, no diploma <input type="checkbox"/> High school graduate or GED completed <input type="checkbox"/> Some college credit but no degree <input type="checkbox"/> Associate degree (e.g., AA, AS) <input type="checkbox"/> Bachelor's degree (e.g., BA, AB, BS) <input type="checkbox"/> Master's degree (e.g., MA, MS, MEng, MEd, MSW, MBA) <input type="checkbox"/> Doctorate (e.g., PhD, EdD) or Professional degree (e.g., MD, DDS, DVM, LLB, JD)		<input type="checkbox"/> No, not Spanish/Hispanic/Latina <input type="checkbox"/> Yes, Mexican, Mexican American, Chicana <input type="checkbox"/> Yes, Puerto Rican <input type="checkbox"/> Yes, Cuban <input type="checkbox"/> Yes, other Spanish/Hispanic/Latina (Specify) _____		<input type="checkbox"/> White <input type="checkbox"/> Black or African American <input type="checkbox"/> American Indian or Alaska Native (Name of the enrolled or principal tribe) _____ <input type="checkbox"/> Asian Indian <input type="checkbox"/> Chinese <input type="checkbox"/> Filipino <input type="checkbox"/> Japanese <input type="checkbox"/> Korean <input type="checkbox"/> Vietnamese <input type="checkbox"/> Other Asian (Specify) _____ <input type="checkbox"/> Native Hawaiian <input type="checkbox"/> Guamanian or Chamorro <input type="checkbox"/> Samoan <input type="checkbox"/> Other Pacific Islander (Specify) _____ <input type="checkbox"/> Other (Specify) _____																					
22. MOTHER MARRIED? (At delivery, conception, or anytime between) <input type="checkbox"/> Yes <input type="checkbox"/> No		23a. DATE OF FIRST PRENATAL CARE VISIT _____ / _____ / _____ <input type="checkbox"/> No Prenatal Care MM / DD / YYYY		23b. DATE OF LAST PRENATAL CARE VISIT _____ / _____ / _____ MM / DD / YYYY																					
24. TOTAL NUMBER OF PRENATAL VISITS FOR THIS PREGNANCY _____ (If none, enter "0".)		25. MOTHER'S HEIGHT _____ (feet/inches)		26. MOTHER'S PREPREGNANCY WEIGHT _____ (pounds)																					
27. MOTHER'S WEIGHT AT DELIVERY _____ (pounds)		28. DID MOTHER GET WIC FOOD FOR HERSELF DURING THIS PREGNANCY? <input type="checkbox"/> Yes <input type="checkbox"/> No		29. NUMBER OF PREVIOUS LIVE BIRTHS																					
29a. Now Living Number _____ <input type="checkbox"/> None		29b. Now Dead Number _____ <input type="checkbox"/> None		30. NUMBER OF OTHER PREGNANCY OUTCOMES (spontaneous or induced losses or ectopic pregnancies) 30a. Other Outcomes Number (Do not include this fetus) _____ <input type="checkbox"/> None																					
29c. DATE OF LAST LIVE BIRTH _____ / _____ MM / YYYY		30b. DATE OF LAST OTHER PREGNANCY OUTCOME _____ / _____ MM / YYYY		31. CIGARETTE SMOKING BEFORE AND DURING PREGNANCY For each time period, enter either the number of cigarettes or the number of packs of cigarettes smoked. IF NONE, ENTER "0". Average number of cigarettes or packs of cigarettes smoked per day. <table border="0"> <tr> <td></td> <td># of cigarettes</td> <td>OR</td> <td># of packs</td> </tr> <tr> <td>Three Months Before Pregnancy</td> <td>_____</td> <td>OR</td> <td>_____</td> </tr> <tr> <td>First Three Months of Pregnancy</td> <td>_____</td> <td>OR</td> <td>_____</td> </tr> <tr> <td>Second Three Months of Pregnancy</td> <td>_____</td> <td>OR</td> <td>_____</td> </tr> <tr> <td>Third Trimester of Pregnancy</td> <td>_____</td> <td>OR</td> <td>_____</td> </tr> </table>			# of cigarettes	OR	# of packs	Three Months Before Pregnancy	_____	OR	_____	First Three Months of Pregnancy	_____	OR	_____	Second Three Months of Pregnancy	_____	OR	_____	Third Trimester of Pregnancy	_____	OR	_____
	# of cigarettes	OR	# of packs																						
Three Months Before Pregnancy	_____	OR	_____																						
First Three Months of Pregnancy	_____	OR	_____																						
Second Three Months of Pregnancy	_____	OR	_____																						
Third Trimester of Pregnancy	_____	OR	_____																						
32. DATE LAST NORMAL MENSES BEGAN _____ / _____ / _____ MM / DD / YYYY		33. PLURALITY - Single, Twin, Triplet, etc. (Specify) _____		34. IF NOT SINGLE BIRTH - Born First, Second, Third, etc. (Specify) _____																					
35. MOTHER TRANSFERRED FOR MATERNAL MEDICAL OR FETAL INDICATIONS FOR DELIVERY? <input type="checkbox"/> Yes <input type="checkbox"/> No IF YES, ENTER NAME OF FACILITY MOTHER TRANSFERRED FROM: _____																									

MEDICAL AND HEALTH INFORMATION

36. RISK FACTORS IN THIS PREGNANCY (Check all that apply): Diabetes <input type="checkbox"/> Prepregnancy (Diagnosis prior to this pregnancy) <input type="checkbox"/> Gestational (Diagnosis in this pregnancy) Hypertension <input type="checkbox"/> Prepregnancy (Chronic) <input type="checkbox"/> Gestational (PIH, preeclampsia) <input type="checkbox"/> Eclampsia <input type="checkbox"/> Previous preterm birth <input type="checkbox"/> Other previous poor pregnancy outcome (Includes perinatal death, small-for-gestational age/intrauterine growth restricted birth) <input type="checkbox"/> Pregnancy resulted from infertility treatment-If yes, check all that apply: <input type="checkbox"/> Fertility-enhancing drugs, Artificial insemination or Intrauterine insemination <input type="checkbox"/> Assisted reproductive technology (e.g., in vitro fertilization (IVF), gamete intrafallopian transfer (GIFT)) <input type="checkbox"/> Mother had a previous cesarean delivery If yes, how many _____ <input type="checkbox"/> None of the above		37. INFECTIONS PRESENT AND/OR TREATED DURING THIS PREGNANCY (Check all that apply) <input type="checkbox"/> Gonorrhea <input type="checkbox"/> Syphilis <input type="checkbox"/> Chlamydia <input type="checkbox"/> Listeria <input type="checkbox"/> Group B Streptococcus <input type="checkbox"/> Cytomegalovirus <input type="checkbox"/> Parvovirus <input type="checkbox"/> Toxoplasmosis <input type="checkbox"/> None of the above <input type="checkbox"/> Other (Specify) _____	
38. METHOD OF DELIVERY A. Was delivery with forceps attempted but unsuccessful? <input type="checkbox"/> Yes <input type="checkbox"/> No B. Was delivery with vacuum extraction attempted but unsuccessful? <input type="checkbox"/> Yes <input type="checkbox"/> No C. Fetal presentation at delivery <input type="checkbox"/> Cephalic <input type="checkbox"/> Breech <input type="checkbox"/> Other D. Final route and method of delivery (Check one) <input type="checkbox"/> Vaginal/Spontaneous <input type="checkbox"/> Vaginal/Forceps <input type="checkbox"/> Vaginal/Vacuum <input type="checkbox"/> Cesarean If cesarean, was a trial of labor attempted? <input type="checkbox"/> Yes <input type="checkbox"/> No E. Hysterotomy/Hysterectomy <input type="checkbox"/> Yes <input type="checkbox"/> No		39. MATERNAL MORBIDITY (Check all that apply) (Complications associated with labor and delivery) <input type="checkbox"/> Maternal transfusion <input type="checkbox"/> Third or fourth degree perineal laceration <input type="checkbox"/> Ruptured uterus <input type="checkbox"/> Unplanned hysterectomy <input type="checkbox"/> Admission to intensive care unit <input type="checkbox"/> Unplanned operating room procedure following delivery <input type="checkbox"/> None of the above	
40. CONGENITAL ANOMALIES OF THE FETUS (Check all that apply) <input type="checkbox"/> Anencephaly <input type="checkbox"/> Meningocele/Spina bifida <input type="checkbox"/> Cyanotic congenital heart disease <input type="checkbox"/> Congenital diaphragmatic hernia <input type="checkbox"/> Omphalocele <input type="checkbox"/> Gastroschisis <input type="checkbox"/> Limb reduction defect (excluding congenital amputation and dwarfing syndromes) <input type="checkbox"/> Cleft Lip with or without Cleft Palate <input type="checkbox"/> Cleft Palate alone <input type="checkbox"/> Down Syndrome <input type="checkbox"/> Karyotype confirmed <input type="checkbox"/> Karyotype pending <input type="checkbox"/> Suspected chromosomal disorder <input type="checkbox"/> Karyotype confirmed <input type="checkbox"/> Karyotype pending <input type="checkbox"/> Hypospadias <input type="checkbox"/> None of the anomalies listed above			

NOTE: This recommended standard fetal death report is the result of an extensive evaluation process. Information on the process and resulting recommendations as well as plans for future activities is available on the Internet at: http://www.cdc.gov/nchs/vital_certs_rev.htm.

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2019

Area	All periods of gestation	12 weeks	16 weeks	20 weeks	20 weeks or 350 grams	20 weeks or 400 grams	20 weeks or 500 grams	350 grams	500grams
Alabama				X					
Alaska				X					
Arizona					X				
Arkansas		X							
California				X					
Colorado	X ¹								
Connecticut				X					
Delaware								X ²	
District of Columbia							X		
Florida				X					
Georgia	X								
Hawaii	X								
Idaho					X				
Illinois				X					
Indiana				X					
Iowa				X					
Kansas				X					
Kentucky					X				
Louisiana					X				
Maine				X					
Maryland				X ¹					
Massachusetts					X				
Michigan						X			
Minnesota				X					
Mississippi					X				
Missouri					X				
Montana								X ²	
Nebraska				X					
Nevada				X					
New Hampshire					X				
New Jersey				X					
New Mexico				X ⁴					
New York	X								
New York (excluding New York City)	X								
New York City	X								
North Carolina				X					
North Dakota				X					
Ohio				X					
Oklahoma		X ⁵							
Oregon				X					
Pennsylvania			X						
Rhode Island	X								
South Carolina					X				
South Dakota									X
Tennessee					X				
Texas				X					
Utah				X					
Vermont				X ⁶					
Virginia	X								
Washington				X					
West Virginia				X					
Wisconsin					X				
Wyoming				X					
American Samoa	X								
Guam					X				
Northern Marianas								X ²	
Puerto Rico					X				
Virgin Islands	X								

1 Although State law requires the reporting of fetal deaths of all periods of gestation, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

2 If weight is unknown, 20 completed weeks of gestation or more.

3 If gestational age is unknown, weight of 500 grams or more.

4 If gestational age is unknown, weight of 350 grams or more.

5 Although State law requires the reporting of fetal deaths at 12 weeks of gestation or more, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

6 If gestational age is unknown, weight of 400 grams or more, 15 ounces or more.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2019

[By place of residence]

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Reporting areas	Total Fetal deaths	Time of birth	Mother's birthplace	Education of mother ²	Father's age ³	Mother's Hispanic origin
Total of reporting areas ¹	21,478	1.2	4.7	12.0	18.3	3.5
Alabama	526	-	0.8	3.0	---	-
Alaska	55	1.8	7.3	18.2	32.7	10.9
Arizona	497	2.0	1.4	2.6	12.5	2.0
Arkansas	306	-	0.7	12.4	---	0.3
California	2,171	1.4	4.1	11.4	10.9	4.1
Colorado	350	-	14.0	22.3	10.0	0.6
Connecticut	133	4.5	0.8	18.0	9.0	3.8
Delaware	59	-	3.4	6.8	23.7	1.7
District of Columbia	86	-	27.9	19.8	---	-
Florida	1,538	2.5	7.5	10.0	15.7	5.6
Georgia	976	-	1.0	19.2	---	4.0
Hawaii	85	16.5	24.7	44.7	38.8	35.3
Idaho	112	-	0.9	-	12.5	-
Illinois	829	1.4	4.0	10.9	12.9	0.4
Indiana	489	-	0.6	3.1	10.0	-
Iowa	196	0.5	3.1	17.9	13.3	1.5
Kansas	196	0.5	1.0	8.2	10.7	5.6
Kentucky	306	-	1.3	3.9	39.9	0.7
Louisiana	290	-	1.7	6.9	25.5	-
Maine	65	9.2	12.3	16.9	26.2	7.7
Maryland	462	0.9	6.5	14.1	15.2	3.2
Massachusetts	298	0.7	13.4	32.6	29.9	19.1
Michigan	652	0.5	6.6	11.3	19.6	2.0
Minnesota	359	0.8	0.3	4.2	10.0	1.1
Mississippi	348	0.3	3.4	8.3	43.7	0.3
Missouri	402	0.7	4.7	6.2	17.4	2.2
Montana	54	-	-	1.9	5.6	-
Nebraska	125	-	0.8	39.2	7.2	-
Nevada	230	0.4	1.3	4.8	17.8	3.0
New Hampshire	55	-	1.8	3.6	16.4	5.5
New Jersey	694	0.3	0.6	4.3	46.0	0.9
New Mexico	69	-	5.8	4.3	23.2	4.3
New York (excluding New York City)	591	6.9	1.5	39.4	---	17.6
New York City	740	0.3	8.1	40.5	---	10.4
North Carolina	784	-	1.0	3.4	39.4	0.8
North Dakota	66	-	9.1	6.1	18.2	6.1
Ohio	850	0.4	2.0	2.4	12.0	0.9
Oklahoma	282	0.4	3.2	3.2	13.5	2.5
Oregon	189	-	1.6	3.7	10.6	1.6
Pennsylvania	802	2.6	7.5	11.5	15.5	0.1
Rhode Island	49	-	20.4	---	30.6	14.3
South Carolina	345	0.3	9.3	8.7	31.6	1.7
South Dakota	69	-	-	1.4	5.8	-
Tennessee	498	0.8	11.2	14.9	---	0.8
Texas	1,504	0.1	5.1	9.0	14.9	2.6
Utah	246	-	2.8	4.5	8.1	-
Vermont	20	-	-	5.0	10.0	-
Virginia	485	4.3	1.6	16.5	---	3.7
Washington	484	2.5	17.4	14.9	16.5	10.3
West Virginia	79	2.5	1.3	7.6	19.0	1.3
Wisconsin	342	0.3	0.9	9.1	---	0.9
Wyoming	40	-	2.5	7.5	17.5	-
American Samoa	---	---	---	---	---	---
Guam	34	-	-	-	---	-
Northern Marianas	8	25.0	12.5	25.0	25.0	-
Puerto Rico	204	-	-	0.5	33.3	1.0
Virgin Islands	---	---	---	---	---	---

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2019
[By place of residence]

Reporting areas	Place of delivery	Attendant at delivery\4	Month prenatal care began	Mother's height	Mother's pre-pregnancy weight\5	Did mother get WIC food for herself during this pregnancy\6	Live-birth order
Total of reporting areas ¹	0.2	0.7	12.2	5.9	11.3	15.1	5.2
Alabama	-	-	2.9	2.5	8.7	4.6	0.8
Alaska	-	3.6	12.7	7.3	16.4	16.4	7.3
Arizona	0.2	1.8	3.4	3.2	6.0	8.0	2.4
Arkansas	0.3	-	7.2	1.3	17.6	13.7	2.0
California	0.5	-	13.4	8.2	15.0	20.8	6.7
Colorado	0.9	2.9	35.4	21.7	25.1	36.9	11.7
Connecticut	-	-	9.0	3.0	5.3	20.3	2.3
Delaware	-	1.7	6.8	1.7	6.8	3.4	3.4
District of Columbia	-	1.2	26.7	7.0	12.8	---	1.2
Florida	-	0.1	12.5	6.9	11.2	15.9	5.0
Georgia	0.1	0.1	20.9	4.4	12.2	24.4	11.0
Hawaii	1.2	-	47.1	41.2	---	---	32.9
Idaho	-	-	7.1	-	0.9	1.8	0.9
Illinois	0.7	2.5	12.9	3.4	12.8	3.7	5.1
Indiana	-	---	6.1	4.5	3.7	43.6	2.2
Iowa	-	-	5.1	3.1	8.7	31.6	1.0
Kansas	-	0.5	5.1	1.5	3.6	14.3	0.5
Kentucky	-	1.6	4.9	0.3	3.3	6.2	0.3
Louisiana	-	0.3	5.5	2.8	7.6	17.2	0.3
Maine	-	-	18.5	12.3	16.9	18.5	27.7
Maryland	-	-	13.6	2.8	7.6	21.0	0.4
Massachusetts	-	-	4.7	2.3	13.8	44.0	1.0
Michigan	-	0.3	12.0	3.2	13.0	15.6	5.8
Minnesota	-	-	1.1	2.5	5.3	3.1	0.6
Mississippi	-	0.3	7.8	1.4	4.0	8.3	0.6
Missouri	-	-	6.2	1.0	5.7	12.4	2.0
Montana	-	1.9	1.9	1.9	5.6	1.9	1.9
Nebraska	-	0.8	9.6	5.6	12.0	12.0	2.4
Nevada	-	-	10.9	0.4	10.9	14.3	0.9
New Hampshire	-	-	3.6	3.6	5.5	9.1	-
New Jersey	-	0.1	4.8	1.2	3.2	7.5	1.2
New Mexico	-	-	2.9	1.4	1.4	10.1	-
New York (excluding New York City)	0.2	-	38.1	24.7	32.1	48.6	31.5
New York City	-	-	36.4	6.2	17.4	---	9.7
North Carolina	-	1.0	1.8	1.3	5.1	4.2	0.4
North Dakota	-	1.5	22.7	1.5	12.1	10.6	9.1
Ohio	0.6	0.1	4.8	3.1	8.2	3.6	0.2
Oklahoma	-	-	7.1	4.3	7.1	7.1	6.7
Oregon	-	0.5	7.4	2.6	4.8	4.2	2.1
Pennsylvania	-	-	11.6	12.5	15.2	22.8	13.5
Rhode Island	-	-	30.6	36.7	36.7	---	22.4
South Carolina	-	-	2.9	2.6	8.1	14.5	0.6
South Dakota	-	-	5.8	-	4.3	1.4	-
Tennessee	-	0.8	12.9	4.4	8.4	15.9	1.0
Texas	0.2	0.9	6.3	4.9	7.5	0.4	1.7
Utah	-	-	4.5	2.0	6.5	-	0.4
Vermont	-	-	-	-	-	5.0	-
Virginia	0.8	1.2	29.7	11.8	22.9	14.6	0.2
Washington	0.4	11.2	27.3	17.4	24.4	22.1	17.8
West Virginia	-	-	3.8	2.5	8.9	11.4	-
Wisconsin	-	-	1.8	2.0	5.3	9.4	0.3
Wyoming	-	7.5	22.5	7.5	7.5	15.0	5.0
American Samoa	---	---	---	---	---	---	---
Guam	-	-	11.8	26.5	29.4	29.4	8.8
Northern Marianas	-	-	-	-	-	-	-
Puerto Rico	0.5	0.5	2.5	1.5	1.0	5.4	1.5
Virgin Islands	---	---	---	---	---	---	---

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2019
[By place of residence]

Reporting areas	Birth interval\7	Cigarette smoking before and during pregnancy	Risk Factors in this Pregnancy	Method of Delivery	
				Fetal presentation	Final route and method of delivery
Total of reporting areas ¹	11.5	8.1	5.1	10.6	4.6
Alabama	3.4	3.2	1.0	2.9	0.4
Alaska	16.4	20.0	12.7	7.3	3.6
Arizona	3.6	5.0	3.6	6.8	1.2
Arkansas	7.8	12.4	-	-	-
California	11.9	11.5	13.1	32.9	10.1
Colorado	21.7	46.6	-	10.0	9.7
Connecticut	9.8	4.5	3.8	9.0	6.8
Delaware	10.2	-	-	1.7	1.7
District of Columbia	37.2	15.1	-	2.3	-
Florida	14.9	4.5	5.8	17.4	9.2
Georgia	17.6	12.5	12.0	16.5	7.2
Hawaii	43.5	27.1	9.4	49.4	44.7
Idaho	7.1	-	3.6	-	-
Illinois	11.6	6.3	0.7	4.2	1.6
Indiana	9.0	4.9	9.2	22.9	0.8
Iowa	6.1	2.6	0.5	0.5	0.5
Kansas	4.1	12.8	-	0.5	0.5
Kentucky	3.9	3.6	1.6	2.9	0.7
Louisiana	4.8	2.1	-	0.3	0.3
Maine	23.1	12.3	-	7.7	4.6
Maryland	8.2	1.3	0.4	1.3	0.2
Massachusetts	6.0	3.7	-	1.0	2.0
Michigan	11.0	10.3	4.9	7.2	0.8
Minnesota	3.6	2.2	0.6	3.3	0.8
Mississippi	1.7	1.7	2.3	2.3	0.6
Missouri	9.0	7.2	-	3.0	0.7
Montana	3.7	1.9	-	1.9	1.9
Nebraska	9.6	6.4	0.8	0.8	0.8
Nevada	4.3	7.0	0.4	0.4	0.4
New Hampshire	-	3.6	-	-	1.8
New Jersey	6.5	6.6	1.0	1.6	2.9
New Mexico	4.3	4.3	-	2.9	-
New York (excluding New York City)	34.5	22.8	3.9	24.4	19.3
New York City	28.6	13.5	19.3	18.8	8.6
North Carolina	3.3	0.9	0.6	0.8	0.5
North Dakota	15.2	1.5	1.5	6.1	-
Ohio	12.7	0.2	-	8.8	3.4
Oklahoma	14.2	3.9	3.5	-	-
Oregon	5.8	3.2	1.1	2.1	1.6
Pennsylvania	10.1	21.9	7.4	11.2	7.2
Rhode Island	34.7	30.6	12.2	12.2	8.2
South Carolina	12.2	1.7	3.8	4.3	2.9
South Dakota	1.4	1.4	-	-	-
Tennessee	7.6	0.6	-	1.2	0.2
Texas	10.8	4.7	3.7	0.3	1.3
Utah	2.0	1.6	-	-	-
Vermont	-	-	-	-	-
Virginia	12.8	6.4	-	20.6	-
Washington	---	16.7	24.4	22.1	15.9
West Virginia	7.6	1.3	1.3	6.3	2.5
Wisconsin	2.6	2.3	1.8	5.6	0.6
Wyoming	37.5	17.5	-	10.0	5.0
American Samoa	---	---	---	---	---
Guam	8.8	23.5	11.8	38.2	-
Northern Marianas	-	-	-	12.5	25.0
Puerto Rico	5.4	-	1.0	2.9	-
Virgin Islands	---	---	---	---	---

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2019
 [By place of residence]

Reporting areas	Maternal morbidity	Weight of fetus	Obstetric estimate of gestation	Cause of death ⁸
Total of reporting areas ¹	3.9	6.9	0.5	31.3
Alabama	0.2	1.5	-	36.5
Alaska	9.1	7.3	1.8	36.4
Arizona	6.0	0.8	-	48.5
Arkansas	-	3.6	-	41.5
California	20.2	5.6	0.9	---
Colorado	-	16.9	-	39.7
Connecticut	2.3	12.8	-	21.8
Delaware	-	1.7	-	39.0
District of Columbia	-	-	-	22.1
Florida	6.4	9.5	2.5	19.6
Georgia	6.5	15.4	0.4	---
Hawaii	11.8	41.2	-	---
Idaho	-	2.7	0.9	25.0
Illinois	0.6	2.3	-	26.7
Indiana	-	4.5	0.2	30.9
Iowa	-	6.1	0.5	25.5
Kansas	-	1.0	-	24.5
Kentucky	0.7	1.0	-	46.1
Louisiana	-	8.6	1.0	25.9
Maine	-	24.6	3.1	23.1
Maryland	0.2	3.9	-	29.0
Massachusetts	-	5.7	-	24.5
Michigan	6.4	4.3	0.3	---
Minnesota	0.3	4.2	-	37.3
Mississippi	3.7	1.4	0.3	---
Missouri	-	2.7	0.2	39.3
Montana	-	3.7	-	31.5
Nebraska	-	4.8	-	20.0
Nevada	0.4	2.2	-	43.0
New Hampshire	-	-	-	32.7
New Jersey	0.6	1.6	0.3	28.8
New Mexico	-	5.8	1.4	43.5
New York (excluding New York City)	-	21.2	0.7	45.9
New York City	-	11.8	-	---
North Carolina	0.4	5.5	-	---
North Dakota	-	1.5	-	---
Ohio	0.5	2.8	0.2	29.8
Oklahoma	3.2	3.9	0.7	23.8
Oregon	0.5	3.7	1.6	40.7
Pennsylvania	5.9	10.6	0.2	28.1
Rhode Island	14.3	4.1	-	34.7
South Carolina	3.5	1.2	-	39.1
South Dakota	1.4	-	-	26.1
Tennessee	-	3.0	0.4	45.0
Texas	1.5	3.2	0.4	22.3
Utah	-	11.8	-	30.9
Vermont	-	5.0	-	---
Virginia	-	13.4	0.6	42.1
Washington	-	27.5	1.4	20.0
West Virginia	1.3	-	-	20.3
Wisconsin	0.3	3.5	0.3	48.2
Wyoming	-	7.5	-	27.5
American Samoa	---	---	---	---
Guam	11.8	-	-	---
Northern Marianas	12.5	25.0	-	---
Puerto Rico	3.4	3.9	-	7.8
Virgin Islands	---	---	---	---

0.0 Quantity more than zero but less than 0.05.

---Data not available.

- Quantity zero.

1\Excludes data for American Samoa, Guam, the Commonwealth of the Northern Marianas, Puerto Rico, and the Virgin Islands.

2\rhode island reported education of mother for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

3\Alabama, Arkansas, the District of Columbia, Georgia, New York (including New York City), Tennessee, Virginia, Wisconsin and Guam reported father's age for 2019 but had an unknown level of 50% or higher and were flagged as not reporting.

4\Indiana reported attendant at delivery for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

5\Hawaii reported mother's pre-pregnancy weight for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

6\New York City does not report WIC and the District of Columbia, Hawaii, and Rhode Island reported this item for 2019 but had unknown levels of 50% or higher and were flagged as not reporting

7\Washington reported birth interval for 2019 but had an unknown level of 50% or higher and was flagged as not reporting.

8\Georgia, Hawaii, Michigan, Mississippi, North Carolina, North Dakota, Vermont, New York City, Guam, and the Northern Marianas reported cause of death for 2019 but had an unknown level of 50% or higher and were flagged as not reporting. California did not report cause of death based on the 2003 revision of the U.S. Standard Report of Fetal Death and was also flagged as not reporting.