User Guide to the 2022 Fetal Death Public Use File



2022 Fetal Death Data Set

User's Guide

ACKNOWLEDGMENTS

Preparation of this user's guide was coordinated by Elizabeth C.W. Gregory and Claudia P. Valenzuela of the Division of Vital Statistics (DVS) under the general direction of Joyce A. Martin, Lead Statistician, Reproductive Statistics Branch (RSB). The vital statistics computer file on which it is based was prepared by DVS staff.

The Division of Vital Statistics Acting Director, Paul Sutton, manages the Vital Statistics Cooperative Program, through which the vital registration offices of all states, the District of Columbia, New York City, Puerto Rico, Virgin Islands, Guam, American Samoa, and the Commonwealth of the Northern Mariana Islands, provided the data to the National Center for Health Statistics (NCHS). This Division, and its Data Acquisition, Classification, and Evaluation Branch (DACEB), and Information Technology Branch (ITB), processed computer edits, designed and programmed the tabulations, reviewed the data, prepared documentation for this publication, and were responsible for receipt and processing of the basic data file.

NCHS acknowledges the essential role of the vital registration offices of all states and territories in maintaining the system through which vital statistics data are obtained and their cooperation in providing the information on which this publication is based.

Questions on the documentation or substantive questions concerning the data should be directed to the Reproductive Statistics Branch, Division of Vital Statistics, NCHS, 3311 Toledo Road, Hyattsville, MD 20782-2003 (1-800) 232-4636.

Department of Health and Human Services Centers for Disease Control and Prevention National Center for Health Statistics Division of Vital Statistics

2022 Fetal Death Data Set

Contents

Introduction

Control count of records

List of data elements and locations

Record layout and definition of item and codes

Documentation tables

Figure 1. U.S. Standard Report of Fetal Death

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2022

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2022

External links

Fetal Mortality in the United States: 2020-2021 and 2021-Provisional 2022 https://www.cdc.gov/nchs/data/vsrr/vsrr032.pdf

Fetal Mortality: United States, 2021 https://www.cdc.gov/nchs/data/nvsr/nvsr72/nvsr72-08.pdf

Perinatal Mortality in the United States, 2020–2021 https://www.cdc.gov/nchs/data/databriefs/db489.pdf

Cause-of-Death Data from the Fetal Death File, 2018–2020 https://www.cdc.gov/nchs/data/nvsr/nvsr71/nvsr71-07.pdf

Births: Final Data for 2022 https://www.cdc.gov/nchs/data/nvsr/nvsr73/nvsr73-02.pdf

Infant Mortality in the United States: Provisional Data From the 2022 Period Linked Birth/Infant Death File https://wcms-wp.cdc.gov/nchs/data/vsrr/vsrr033.pdf

User Guide to the 2022 Period Linked Birth/Infant Death Public Use File (Forthcoming)

Introduction

This User Guide is for use with the 2022 fetal death public use data. The 2022 fetal death micro-data file may be downloaded at:

http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm [1]. The micro-data fetal death file for the U.S. does not include geographic detail (e.g., state or county of birth). The territories file contains selected geographic detail. Selected fetal death data for 2005-2021, including some geographic data and cause of death data, are available in CDC WONDER

(<u>http://wonder.cdc.gov</u>), an interactive online data access tool. Selected fetal death data for 2022 are expected to be available in WONDER in early 2024.

For 2022, fetal death data are included on the file for all 50 states, the District Columbia, Guam and Puerto Rico. American Samoa did not report data for 2022 and data for the Northern Marianas and the U.S. Virgin Islands are excluded from the file due to the small number of events reported (6 and 14, respectively).

Definition of fetal death

"Fetal death" means death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy and which is not an induced termination of pregnancy. The death is indicated by the fact that after such expulsion or extraction, the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. Heartbeats are to be distinguished from transient cardiac contractions; respirations are to be distinguished from fleeting respiratory efforts or gasps.

This definition [2] has been adopted by the Centers for Disease Control and Prevention's National Center for Health Statistics (NCHS) as the nationally recommended standard and is based on the definition published by the World Health Organization in 1950 and revised in 1988. The term "fetal death" is defined on an all-inclusive basis to end confusion arising from the use of such terms as stillbirth, spontaneous abortion, and miscarriage. All U.S. states and registration areas have definitions similar to the standard definition, except for Puerto Rico and Wisconsin, which have no formal definition [3]. Fetal deaths do not include induced terminations of pregnancy.

Reporting requirements for fetal death data

Reporting requirements for fetal deaths vary by state, and these differences can have implications for comparisons of fetal and perinatal mortality rates by state [2]. Table A shows the period of gestation at which fetal death reporting is required for each reporting area for 2022. All jurisdictions require reporting of fetal deaths of at least 20 weeks of gestation or more, or a minimum of 350 grams birthweight (roughly equivalent to 20 weeks), or some combination of the two, but some require reporting at earlier gestational ages. Six states (Colorado, Georgia, Hawaii, New York, Rhode Island, and Virginia), American Samoa, and the U.S. Virgin Islands require reporting of fetal deaths at all periods of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), two states require reporting beginning at 12 weeks of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), and one state requires reporting beginning at 16 weeks of gestation.

User Guides for the Fetal Death Public Use files for 2007-2020 (Table A) indicated that South Dakota reported fetal deaths with birthweights of 500 grams or more (approximately 22 weeks of gestation or more). Over this period and to the present, however, South Dakota has reported fetal deaths at 20 weeks of gestation or more. Note that numbers of fetal deaths in the public use fetal death file reflect the correct reporting period of 20 weeks of gestation or more.

Classification of fetal deaths at 20 weeks of gestation or greater

To maximize the comparability of data by year and by state while including as much meaningful data as possible, most tables published by NCHS are based on fetal deaths occurring at gestations of 20 weeks or more. These tabulations also include fetal deaths for which gestation is not stated for those states requiring reporting at 20 weeks of gestation or more only. Fetal deaths of not-stated gestation are excluded for states requiring reporting of all products of conception, except for those with a stated birthweight of 350 grams or more. In 2022, this rule was applied to the following reporting areas: Arkansas, Colorado, Georgia, Hawaii, New York (including New York City), Oklahoma, Pennsylvania, Rhode Island, and Virginia. American Samoa and the Virgin Islands also requiring reporting of all products of conception when data are available (see Table A). The summation of these rules results in the creation of the tabulation flag (position 10), which, when invoked, produces the number of records with a stated or presumed period of gestation of 20 weeks or more.

Data for fetal deaths occurring before 20 weeks of gestation should be used with caution, if at all, as they represent a limited, subnational reporting area and are not evaluated for data quality due to limited resources.

Classification of fetal deaths by occurrence and residence

In tabulations by place of residence, fetal deaths occurring within the United States to U.S. citizens and to residents who are not citizens are allocated to the usual place of residence of the mother in the United States, as reported on the report of fetal death. Fetal deaths to U.S. residents occurring outside this country are not included in tabulations by place of residence or place of occurrence. Accordingly, the total count of fetal deaths for the United States by place of residence of residence and by place of occurrence will not be identical. Fetal deaths to nonresidents of the

United States are included in data by place of occurrence but excluded from data by place of residence, as previously indicated. See Tables 2 and 3, respectively, for the number of fetal deaths by residence and occurrence for the 50 states and the District of Columbia for 2022.

Geographic classification: The geographic code structure for the 2022 fetal death file is given in the NCHS manual, "Vital Records Geographic Classification, 2014," and in the country, county, and place geographic code files [4]. The geographic code structure on the 2022 file is based on results of the 2010 Census of Population.

Standard Report of Fetal Death

The U.S. Standard Report of Fetal Death, issued by the U.S. Department of Health and Human Services, has served for many years as the principal means for attaining uniformity in the content of the documents used to collect information on fetal deaths in the United States. The U.S. Standard Report of Fetal Death has historically been revised every 10-15 years. Most state reports conform closely in content to the standard report but are modified to the extent required by the particular state's needs or by special provisions of the state's vital statistics law.

The 2003 revision: In 2003, a revised U.S. Standard Report of Fetal Death was adopted (Figure 1). For more information on the 2003 standard report and details regarding the certificate revision and links to the documents referenced below, see the NCHS website of the 2003 report revision at http://www.cdc.gov/nchs/nvss/vital_certificate_revisions.htm. The 2003 report of fetal death replaces the previous 1989 U.S. Standard Report of Fetal Death [5,6]. Implementation of the 2003 U.S. Standard Report of Fetal Death (revised) by the states and independent reporting areas was phased in from 2003 to 2018. All states, the District of Columbia, Guam, the Northern Marianas, Puerto Rico, and the Virgin Islands reported data based on the 2003 U.S. Standard Report of fetal Death as of 2018.

The 2003 Revision of the U.S. Standard Report of Fetal Death introduced substantial changes to data content and quality. Many key data items are common between revisions; however, a number of items were substantively modified. The 2003 revision also includes many new items never before collected on the Standard Report [5,6]. For details on data items comparable between revisions see the User Guide to the 2013 Fetal Death Public Use File [7].

A key aspect of the 2003 revision of the U.S. Standard Report of Fetal Death was the reengineering of the data collection and transmission system to improve data quality, speed of data collection and transmission, and to enhance standardization of data [5,8]. To encourage collection of data from the best sources, two worksheets were developed: the "Patient's Worksheet" (available at: <u>https://www.cdc.gov/nchs/data/dvs/patientwkstfetaldth.pdf</u>) [9] and the "Facility Worksheet" (available at: <u>https://www.cdc.gov/nchs/data/dvs/FacilityFetal04.pdf</u>) [10]. For the Patient's Worksheet, data are directly obtained from the mother and include items such as race, Hispanic origin, and educational attainment. For the Facility Worksheet, data are obtained directly from the medical records of the mother for items such as date of first prenatal care visit, pregnancy risk factors, and method of delivery. To assist hospital staff in completing the Facility Worksheet, a comprehensive instruction manual was developed: Guide to Completing the Facility Worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 Revision) ("Guide to the Facility Worksheet"; available at:

<u>https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf</u>) [11]. Detailed definitions and instructions for data items that are collected from the Facility Worksheet are in the "Guide to the Facility Worksheet".

Additionally, a review of 2003-based report of fetal death revision items in 2014 by a collaborative effort among representatives from several vital statistics jurisdictions: The National

Association for Public Health Statistics and Information Systems (NAPHSIS), and NCHS, resulted in the decision to drop a number of items from the national report of fetal death data file for reasons of poor data quality. For more information on this effort and for a full list of items that were dropped, see:

https://www.cdc.gov/nchs/nvss/deleted items from birth fetal death files.htm.

The first ever eLearning training, "Applying Best Practices for Reporting Medical and Health Information on Birth Certificates," on completing the medical and health information for the birth certificate and report of fetal death was launched in October 2016 and has recently been updated. The training emphasizes the importance and uses of birth certificate and report of fetal death data and best practices for collecting specific medical and health items, including cause of fetal death. The audience for the training includes birth information specialists, physicians, nurses, and hospital administrators. Continuing education credits for nurses, physicians, and nonclinical staff are also available. The training is internet-based and approximately 45 minutes in length. It is available at: www.cdc.gov/nchs/training/BirthCertificateElearning.

Fetal death data files

Micro-data files: Fetal death micro-data files for data years 1982-2022 may be downloaded at: <u>http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm</u>. The general rules used to classify characteristics of fetal deaths are presented in several NCHS manuals [4, 8, 12]. These instructions are for states to use to collect and code the data items; they do not include NCHS edit recodes.

The 2003-2006 fetal death micro-data files include data items common to both the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death, as well as items exclusive to the 2003 revision. For 2007-2013, in an effort to improve the timeliness of fetal death data, the fetal death files only included items that were comparable across the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death. From 2014-2017, the fetal mortality files again included data comparable between the two revisions, as well as well as items exclusive to the 2003 revision. Beginning with 2018, the fetal death file only includes data exclusive to the 2003 revision.

The U.S. micro-data fetal death files for 2005 through 2022 do not include geographic detail (e.g., mother's state of residence). Tabulations of fetal death data by state-level geographic detail and many other demographic and health items are available in the CDC WONDER web application (<u>http://wonder.cdc.gov/</u>) for data years 2005-2021. Additionally, certain geographic level data may also be available upon request: See "NCHS Data Release and Access Policy for Micro-data and Compressed Vital Statistics Files", available at:

http://www.cdc.gov/nchs/nvss/dvs_data_release.htm.

The territories file includes data on fetal deaths occurring in Guam and Puerto Rico (American Samoa did not report data for 2022 and data for the Northern Marianas and the Virgin Islands were excluded due to small numbers). It includes limited geographical data such as information identifying individual territories and counties (or their equivalent) with populations of 100,000 or more by place of occurrence and residence.

Race of Mother

The 2003 revision of the U.S. Standard Report of Fetal Death allows the reporting of the five race categories either alone (i.e., single-race) or in combination (i.e., more than one race or multiple races) for the mother [6], in accordance with the revised standards issued by the Office of Management and Budget (OMB) in 1997 [13]. The five categories for race specified in the revised standards are: American Indian or Alaska Native (AIAN), Asian, Black or African

American, Native Hawaiian or Other Pacific Islander (NHOPI), and White. Information on this change is presented elsewhere [14-16]. Starting in 2018, all states and the District of Columbia, in addition to Puerto Rico, Guam, the Northern Marianas, and the Virgin Islands, began reporting race according to the 1997 revised OMB standards.

Prior to 2018, to provide uniformity and comparability of the data before all or most of the data were available in the new multiple-race format, it was necessary to "bridge" the responses of those for whom more than one race was reported (multiple race) to one, single race. Beginning in 2018, data by race are shown according to the 1997 OMB standards. Bridged-race data are no longer available on the public use file beginning with the 2020 data year.

Hispanic Origin of Mother

Hispanic origin and race: Hispanic origin and race are reported separately on the birth certificate (Figure 1). It is recommended that this information be reported directly by the mother via the Patient's Worksheet [9]. For 1989 through 2018, data on the public use file and in NCHS reports for specified Hispanic groups are shown in most cases for five specified Hispanic groups: Mexican, Puerto Rican, Cuban, Central and South American, and "other and unknown Hispanic." Starting with 2019, data are presented for the additional Hispanic group, Dominican (see item MHISPX in file position 143). This subgroup was previously included in "other and unknown Hispanic." In tabulations of fetal death data by race and Hispanic origin, data for persons of Hispanic origin are not further classified by race because the vast majority of Hispanic women are reported as white. In tabulations of fetal death data by race only, data for persons of Hispanic origin are included in the data for each race group according to the mother's reported race. In tabulations that include Hispanic origin, data for non-Hispanic persons are classified according to the race of the mother. Beginning in 2022, records for which mother's detailed Hispanic origin is coded to "Latin American" will be recoded to "Other and Unknown Hispanic origin" instead of "Central and South American" for the mother's Hispanic origin recodes at positions 142 and 143 in the file layout.

Cause of Death Data

A cause-of-fetal death item has been included on the report of fetal death since 1939 because it was considered critical information. However, the data were not released on public–use files or published until 2014 due to resource constraints and quality concerns. National data on cause of death are now released annually and are available for 2014-2022.

A major redesign of the cause of death (COD) section of the U.S. Standard Report of Fetal Death was undertaken with the 2003 revision. The goals of the redesign were to be consistent with instructions in the World Health Organization's (WHO) International Classification of Diseases (ICD) [17], and to thereby improve the quality and specificity of COD information, provide additional reporting guidance, and retain the flexibility to report any cause of death [18].

As with other deaths, the intent is for an attending physician or medical examiner or coroner to report cause of death [17]. The cause-of-fetal-death item requests a medical opinion from this person on the conditions and diseases resulting in or contributing to death, but also asks the medical certifier to report one cause separately (item 18a) from all other causes (item 18b) reported on the fetal death report. The certifier may form this medical opinion upon various medical tests, investigations, and examinations as is the case with other deaths. However, the term "initiating cause" used to refer to the one cause reported separately is unique to fetal deaths because of differences in the format of the cause item and how the initiating cause is determined for fetal deaths compared with the "underlying cause" term used with other deaths [18].

NCHS codes cause of fetal death reported by the certifier using the ICD-10 classification [17]. Coding is accomplished through a combination of automated and manual processes following the guidelines laid out in Instruction manual part 2k, Instructions for the Automated Classification of the Initiating and Multiple Causes of Fetal Deaths [19]. Literal text stated on the fetal death report is assigned ICD-10 codes, and a single cause of death, the initiating cause of death, is selected from the conditions entered by the medical certifier in the cause-of-death section of the fetal death report. One section of the report (18a) is for the medical certifier to state the single condition that he or she considers the cause that initiated or triggered problems that resulted in the fetus dying at this point in time, so this is anticipated to be the initiating cause of death. If more than one cause or condition is entered by the medical certifier, the initiating cause is determined by the placement of the condition on the report, provisions of the ICD, and associated selection rule and modifications. A second section of the report (18b) is for the medical certifier to state any other conditions or causes that he or she felt played a role in causing the fetal death. Since more medical information may be reported on the report than is directly reflected in the initiating cause of death, this additional information is captured in multiple cause-of-fetal-death data.

The fetal death data file also includes a cause of death data item for 124 Selected Causes of Fetal Death (fetal cause list) (tables 7A and 7B). The reporting area for these tables includes areas that used the 2003 revision format and where less than 50 percent of an area's cause data were attributed to the Fetal death of unspecified cause (unspecified cause) (P95).

Incomplete national reporting: The use of reporting flags

The 2022 national fetal death file includes items that are not reported by some vital statistics jurisdictions or are reported but have an unknown level of 50% or higher and are flagged as not reporting. Reporting flags were developed to help the user more readily identify the correct number of fetal deaths for items with less than full national reporting. Reporting flags are available for many items in the file, and positions for reporting flags are noted along with each data item in the file layout.

Translating "blanks"

In the 2022 fetal death file, for data items which are not reported by all states, fetal deaths to residents of a state reporting the item that occurred in a state not reporting the item, are represented by "blanks." Blanks should be treated as "unknowns" for tabulations. In sum, the correct use of reporting flags and translation of blanks will result in an accurate tally of fetal deaths for items with incomplete national reporting. For an example of SAS code that may be used to incorporate the correct use of reporting flags and the translation of blanks, see below.

Example of SAS code using reporting flags (and translating blanks)

The example below is for the receipt of WIC item, for which data for 47 states and the District of Columbia are included for 2022 (New York City does not report this item and Hawaii, Indiana, and Rhode Island reported this item for 2022 but had unknown levels of 50% or higher and were flagged as not reporting).

Sample SAS program

01	DATA work;
02	INFILE 'c:fet22us.dat' LRECL=3150;
03	INPUT
04	oe_tabflg 10
05	wic \$229

06	restatus 131
07	f_wic \$377;
08	
09	/*Include if 20 weeks gestation or more*/
10	IF oe_tabflg EQ 2;
11	/*Exclude foreign residents*/
12	IF restatus NE 4;
13	/*Select reporting area*/
14	IF f_wic=1;
15	/*Convert blanks to unknown*/
16	IF wic =' ' THEN wic ='U';
17	
18	PROC FREQ;
19	TABLE wic;
20	RUN;

In this example, "oe_tabflg" is used to select fetal deaths of 20 weeks gestation or more. Also, "restatus" is used to exclude fetal deaths to foreign residents (this is standard practice for all NCHS tabulations). As well in this example, blanks are represented by character values SAS code = (``). However, for some items in the file, e.g., number of previous cesarean deliveries, blanks are represented by numeric values for which the SAS code is (`.').

Quality of Data

State-Specific Data Quality Issues

These state-specific data quality issues are of concern due to potential underreporting and/or inaccurate reporting for 2022.

- <u>Previous live births now dead</u> Florida-Some records erroneously reported the current fetal death as a previous live birth now dead. Data for this item for this state should be used with caution.
- <u>Attendant</u> New Jersey-No records have been reported as having a certified nurse midwife as the attendant since mid-2021. Data for this item for this state should be used with caution.
- <u>Plurality</u> Hawaii-The number of multiple deliveries may be underreported due to the high percentage of records reported as unknown plurality; records with unknown plurality are imputed to a singleton delivery. Data for this item for this state should be used with caution.

Percentage of unknown responses

Information on the completeness of reporting of fetal death (with a stated or presumed period of gestation of 20 weeks or more) data is shown in Table B, which presents a listing of items and the percentage of records that were not stated for each reporting area: all 50 states, the District of Columbia, New York City, Puerto Rico, Guam, American Samoa, the Northern Marianas, and the Virgin Islands. In general, percentages of unknown responses are considerably higher for fetal deaths than for live births, and among fetal deaths the percentage unknown is higher for fetal deaths that occur earlier in the gestational period. In fetal mortality tables, unknown responses are shown in frequencies tables but are excluded from the computation of percent distributions and fetal and perinatal mortality rates. Thus, rates for variables with a substantial percentage of unknown responses (such as birthweight) may understate the "true" rates of fetal mortality for that characteristic.

As high levels of missing data are a data quality concern, beginning with the 2017 data year, jurisdictions with a 50% or greater unknown level for a particular item are flagged as not reporting that item.

State Specific Issues

The items below had unknown levels of 50% or higher for the jurisdiction(s) noted and are flagged as not reporting for those jurisdictions in 2022:

- Mother's education-Hawaii and Rhode Island
- <u>Father's age</u>- Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New York, Rhode Island, Virginia, and Guam
- Previous live births dead-Hawaii
- Live birth order-Hawaii
- <u>Date of last live birth</u>-Hawaii
- Date of first prenatal care-Hawaii and Rhode Island
- <u>WIC</u>-Hawaii, Indiana, Rhode Island, and Guam
- Mother's pre-pregnancy weight-Hawaii
- Mother's pre-pregnancy BMI-Hawaii and Guam
- All smoking items-Rhode Island and Guam
- <u>Fetal presentation</u>-Hawaii
- Route and method of delivery-Hawaii
- <u>Trial of labor attempted</u>-Hawaii
- Date of last normal menses-Colorado and Hawaii
- <u>Estimated time of fetal death</u>-District of Columbia, Hawaii, Kansas, Nevada, New York, Oklahoma, Virginia, and Guam

 <u>Cause of death</u>- Arizona, California, Georgia, Hawaii, Mississippi, New York City, North Dakota, Vermont, and Guam

Change in items reported by New York state

Due to reinterpretation of state law, as of September 2022, New York state no longer reports data on the following items to NCHS:

- Day of delivery
- Mother's date of birth
- City of residence
- Inside city limits
- Father's date of birth
- Match number

In order to help data users generate correct counts, additional reporting flags have been added where appropriate.

Development of new imputation for day of delivery for New York state

For all jurisdictions, when a record has an unknown value for the day of delivery, the day is imputed to 15. Because the date of delivery is used to create several derived data items, such as month prenatal care began, and the day of delivery is missing for all records that occurred in New York State after September 2022, a more robust imputation for day of delivery was developed for fetal deaths that occurred in New York state in order to create a more natural distribution. That is, for any record with an unknown day of delivery with a complete date of last normal menses and a valid value for the obstetric estimate of gestational age, a date of delivery is calculated. If the month of this calculated date matches the month of delivery reported on the record or the calculated date falls within 4 days of the start of the reported month, then the calculated day of delivery is used. Otherwise, the day of delivery is imputed to 15.

Computation of rates

Fetal mortality rates are computed as the number of fetal deaths at 20 weeks of gestation or more per 1,000 live births and fetal deaths at 20 weeks or more. Perinatal mortality rates are computed in a similar fashion, as shown below. The denominators for all fetal and perinatal mortality rates are live births plus fetal deaths in the specified gestational age group, thus representing the population at risk of the event [2].

Fetal mortality rate = Fetal deaths at 20 weeks of gestation or more x 1,000

Live births and fetal deaths at 20 weeks or more

Perinatal mortality rate, definition I =

<u>Fetal deaths at 28 weeks or more and infant deaths under 7 days x 1,000</u> Live births and fetal deaths at 28 weeks or more

Perinatal mortality rate, definition II =

<u>Fetal deaths at 20 weeks or more and infant deaths under 28 days</u> x 1,000 Live births and fetal deaths at 20 weeks or more

In each case, the fetal deaths included in the denominator of each rate mirror the fetal deaths included in the numerator. A previous NCHS report [20] contains information on the historical development of various perinatal measures. An asterisk (*) is shown in place of any rate based on fewer than 20 fetal or perinatal deaths in the numerator.

Random variation in fetal and perinatal mortality

The number of fetal deaths, perinatal deaths, and live births reported for an area represent complete counts of such events. As such, they are not subject to sampling error, although they are subject to nonsampling error in the registration process. However, when the figures are used for analytic purposes, such as the comparison of rates over time, for different areas, or among different subgroups, the number of events that actually occurred may be considered as one of a large series of possible results that could have arisen under the same circumstances [21]. As a result, numbers of births, fetal deaths, perinatal deaths, and fetal and perinatal mortality rates are subject to random variation. The probable range of values may be estimated from the actual figures according to certain statistical assumptions.

In general, distributions of vital events may be assumed to follow the normal distribution. When the number of events is large, the relative standard error (RSE) is usually small. When the number of events is small (perhaps less than 100) and the probability of such an event is small, considerable caution must be used in interpreting the data. Such infrequent events may be assumed to follow a Poisson probability distribution. Estimates of RSEs and 95% confidence intervals are shown below. In the formulas, D is the number of fetal or perinatal deaths and B is the number of live births plus fetal deaths used as the denominator in computing fetal and perinatal mortality rates.

The formulas for the two RSEs are as follows:

RSE(D)=100*
$$\sqrt{\frac{1}{D}}$$
 and
RSE(B)=100* $\sqrt{\frac{1}{B}}$.

For example, let us say that for group A the number of fetal deaths was 238,

whereas the number of live births plus fetal deaths in the denominator was 32,650 yielding a fetal mortality rate of 7.29 fetal deaths per 1,000 live births and fetal deaths.

The RSE of the deaths =
$$100*\sqrt{\frac{1}{238}} = 6.48$$
,

whereas the RSE for the births plus fetal deaths in the denominator is

$$RSE=100^* \sqrt{\frac{1}{32,650}} = 0.55.$$

The formula for the RSE of the fetal mortality rate is:

$$RSE=100*\sqrt{\frac{1}{D}+\frac{1}{B}}$$

Thus the RSE for the example above is:

$$= 100^* \sqrt{\frac{1}{238} + \frac{1}{32,650}} = 6.51.$$

Normal distribution

When the number of events is greater than 100, the normal distribution is used to estimate the 95 percent confidence intervals of a rate, R_{I} , as follows:

Lower:
$$R_1 - 1.96 * R_1 * \frac{\text{RSE}(\text{R1})}{100}$$

Upper:
$$R_{l}$$
+ 1.96 * R_{l} * $\frac{\text{RSE}(\text{R1})}{100}$

Thus, for Group A:

Lower:
$$7.29 - (1.96 * 7.29 * \frac{6.51}{100}) = 6.36$$

Upper: 7.29 + (1.96 * 7.29 * $\frac{6.51}{100}$) = 8.22

Thus the chances are 95 out of 100 that the true fetal or perinatal mortality rate for Group A lies somewhere in the 6.36–8.22 interval.

Poisson distribution

When the number of events in the numerator is less than 100 the confidence interval for the rate, R_{l} , can be estimated based on the Poisson distribution using the values in Table III (2).

Lower: R_1 *L(.95, D_{adj})

Upper: R_1 *U(.95, D_{adj})

where D_{adj} is the adjusted number of fetal or perinatal deaths (rounded to the nearest integer) used to take into account the RSE of the number of deaths in the numerator and the number of live births plus fetal deaths in the denominator, and is computed as follows:

$$\mathbf{D}_{\mathrm{adj}} = \frac{D * B}{D + B}$$

 $L(.95, D_{adj})$ and $U(.95, D_{adj})$ refer to the values in Table III corresponding to the value of D_{adj} .

For example, let us say that for Group B the number of fetal deaths was 73, and the number of live births plus fetal deaths in the denominator was 11,422, and the fetal mortality rate was 6.39.

$$D_{adj} = \frac{73*11,422}{73+11,422} = 73$$

Therefore the 95 percent confidence interval (using the formula in Table III for 1\N99 infant deaths) =

Lower:
$$6.39*0.78384 = 5.01$$

Upper:
$$6.39*1.25735 = 8.03$$

Comparison of two fetal or perinatal mortality rates

If either of the two rates to be compared is based on less than 100 deaths, compute the confidence intervals for both rates and check to see if they overlap. If so, the difference is not statistically significant at the 95 percent level. If they do not overlap, the difference is statistically significant. If both of the two rates (R_1 and R_2) to be compared are based on 100 or more deaths, the following z-test should be used to define a significance test statistic:

$$z = \frac{\mathbf{R}_1 - \mathbf{R}_2}{\sqrt{\mathbf{R}_1^2 \left(\frac{RSE(\mathbf{R}_1)}{100}\right)^2 + \mathbf{R}_2^2 \left(\frac{RSE(\mathbf{R}_2)}{100}\right)^2}}$$

If $|z| \ge 1.96$, then the difference is statistically significant at the 0.05 level and if |z| < 1.96, the difference is not significant.

References

 National Center for Health Statistics. Fetal Death 2022. Public use file. Hyattsville, Maryland: National Center for Health Statistics. Annual internet product. 2024. Available at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.

 MacDorman MF, Gregory ECW. Fetal and perinatal mortality: United States, 2013.
 National vital statistics reports; vol 64 no 8. Hyattsville, MD: National Center for Health Statistics. 2015. 3. Kowaleski J. State definitions and reporting requirements for live births, fetal deaths, and induced terminations of pregnancy (1997 revision). Hyattsville, MD: National Center for Health Statistics. 1997. Available from: <u>http://www.cdc.gov/nchs/data/misc/itop97.pdf</u>.

4. Division of Vital Statistics. Instruction manual part 8: Geographic classification, 2014.Hyattsville, Maryland: National Center for Health Statistics. Available at:

https://www.cdc.gov/nchs/data/dvs/IMP8_2014.pdf.

5. National Center for Health Statistics. Report of the Panel to Evaluate the U.S. Standard Certificates. Hyattsville, Maryland: National Center for Health Statistics. 2000

 National Center for Health Statistics. 2003 revision of the U.S. Standard Report of Fetal Death. 2003.

7. National Center for Health Statistics. User Guide to the 2013 Fetal death Public Use File. Hyattsville, Maryland: National Center for Health Statistics. Annual product 2015. Available at: <u>http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm</u>.

8. National Center for Health Statistics. Fetal death edit specifications for the 2003 Revision of the U.S. Standard Certificate of Birth. 2012.

https://www.cdc.gov/nchs/data/dvs/fetal_death_edit_specifications.pdf.

9. National Center for Health Statistics. Patient's Worksheet for Report of Fetal Death.

2019. Available at: https://www.cdc.gov/nchs/data/dvs/fetal-death-mother-worksheet-english-2019-508.pdf.

10. National Center for Health Statistics. Facility Worksheet for Report of Fetal Death. 2019. Available at: <u>https://www.cdc.gov/nchs/data/dvs/fetal-death-facility-worksheet-2019-508.pdf</u>.

11. National Center for Health Statistics. Guide to completing the facility worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 revision). 2019 update. Hyattsville,

Maryland: National Center for Health Statistics.

https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf.

Division of Vital Statistics. Computer edits fetal death data --2003 Revised Certificate.
 Hyattsville, Maryland: National Center for Health Statistics.

13. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. Fed Regist 62FR58781–58790. October 30, 1997.

14. Ingram DD, Parker JD, Schenker N, et al. United States Census 2000 population with bridged race categories. National Center for Health Statistics. Vital Health Stat 2(135). 2003.

15. Johnson D. Coding and editing multiple race. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP. Portland, Oregon. June 6–10, 2004.

Weed JA. NCHS procedures for multiple-race and Hispanic origin data: Collection,
 coding, editing, and transmitting. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP.
 Portland, Oregon. June 6–10, 2004.

17. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. Geneva, Switzerland. 2009.

 Hoyert DL, Gregory ECW. Cause-of-death data from the fetal death file, 2015–2017.
 National Vital Statistics Reports; vol 69 no 4. Hyattsville, MD: National Center for Health Statistics. 2020.

National Center for Health Statistics. Instruction manual, part 2k: Instructions for the automated classification of the initiating and multiple causes of fetal deaths, 2014. Hyattsville, MD. 2014.

20. Hoyert DL. Perinatal mortality in the United States, 1985–91. National Center for Health Statistics. Vital Health Stat 20(26). 1995. Available from:

http://www.cdc.gov/nchs/data/series/sr_20/sr20_026.pdf.

Brillinger DR. The natural variability of vital rates and associated statistics. Biometrics
 42(4):693–734. 1986.

Control Count of Records

File Characteristics: Fetal Public Use File 2022

All files:

Record format:	Blocked, Fixed, Format
Code scheme:	Numeric/Alphabetic/Blank
Record length:	3150

Record counts

U.S. file record count:

Counts of a	ll fetal deaths:
-------------	------------------

Record count	40,113
By occurrence	40,113
By residence	39,918
To foreign residents	195

Counts of 20 weeks or more:

Record count	20,368
By occurrence	20,368
By residence	20,202
To foreign residents	166

Counts of less than 20 weeks:

Record count	19,745
By occurrence	19,745
By residence	19,716
To foreign residents	29

U.S. territories file record count:

Counts of all fetal deaths:

Record count	259
By occurrence	259
By residence	254
To foreign residents	5

Counts of 20 weeks or more:

Record count	257
By occurrence	257
By residence	252
To foreign residents	5

Counts of less than 20 weeks:

Record count	2
By occurrence	2
By residence	2
To foreign residents	-

2022 LIST OF DATA ELEMENTS AND LOCATIONS

Data Items	Locations
 General Tabulation flags (gestational age) Data year Resident status 	10 11-14 131
2. Prenatal Carea) Month began	202-203
 3. Fetus a) Sex b) Plurality c) Weight at delivery d) Gestational age e) Month/year of delivery f) Time of delivery 	316 301 349-355 340-344(NCHS Standard item), 331-335 11-16 21-24
 g) Day of week of delivery 4. Mother a) Age b) Race c) Education d) Hispanic origin 	25 86-90 132-136 145 142-144
5. Pregnancy Historya) Live birth orderb) Birth interval	187 197-201
6. Father a) Age	177-180
 7. Other Items a) Attendant at delivery b) Place of delivery c) WIC receipt 	283 34, 52 229
 8. Medical and Health Data a) Method of delivery b) Medical risk factors c) Other risk factors i. Tobacco ii. Mother's pre-pregnancy body mass index (BMI) d) Maternal morbidity 	274-277, 280 257-267 230-238 245-249 281-282
9. Residence reporting flags	372-445

10. Cause of Death

a.) Initiating cause or condition code	2603-2607
b.) Initiating cause or condition recode	2643-2645
c.) Initiating cause or condition flag	2651

2022 Fetal Death Public Use Record Layout

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
1-6 7	6 1	FILLER VERSION	Version		A,S	A S	State used the 2003 version of the US Standard Report of Fetal Death (Revised) State used the 1989 version of the US Standard Report of Fetal Death (Unrevised)
8-9	2	FILLER					
10	1	OE_TABFLG	Obstetric Estimate Tabu (Use with NCHS Standard		A,S	1 2	Under 20 weeks (exclude) 20 weeks or more (include)
11-14	4	DOD_YY	Delivery Year		A,S	2014	Year of delivery
15-16	2	DOD_MM	Delivery Month		A,S	01 02 03 04 05 06 07 08 09 10 11 12	January February March April May June July August September October November December
17-20	4	FILLER					
21-24	4	DOD_TT	Delivery Time	c409	А	0000-2 9999	2359 Time of Birth Not Stated
25 * For A S	2003 V 1989 V	Version of US Stat Version of US Stat	Weekday the following codes apply: ndard Birth Certificate ndard Birth Certificate pdard Birth Certificate but		A,S	1 2 3	Sunday Monday Tuesday

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	L	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
							4 5 6 7	Wednesday Thursday Friday Saturday
26-27		2	OSTATE	Territory of Occurrence (This item is available in the geographic codes are not a		A,S île)		
				<u>Territories</u>			AS GU MP PR VI	American Samoa Guam Northern Marianas Puerto Rico Virgin Islands
28-29		2	FILLER					
30-32		3	OCNTYFIPS	Occurrence FIPS County (This item is available in the geographic codes are not a Data are suppressed if popu	vailable in the U.S. f	A,S îile)	000 001-998 999	No county level geography County of Occurrence (See Geographic Table) County of less than 100,000
33		1	OCNTYPOP	Occurrence Territory Cou (This item is available in the geographic codes are not a	e territory file only,	A,S îile)		
							0 1 2 3	County of 1,000,000 or more County of 500,000 to 1,000,000 County of 250,000 to 500,000 County of 100,000 to 250,000
	* For th A S s	2003 V 1989 V	Version of US Star Version of US Star	the following codes apply: adard Birth Certificate adard Birth Certificate adard Birth Certificate but t	his element confor	ms to the	: 2003 stat	ndards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition	
						9	County less than 100,000	
34	1	BFACIL	Delivery Place (Revised)	409	Α	1 2 3 4 5 6 7 9	Hospital Freestanding Birth Center Home (intended) Home (not intended) Home (unknown if intended) Clinic/Doctor's Office Other Unknown	
35-51	17	FILLER						
52	1	BFACIL3	Delivery Place Recode		A,S	1 2 3	In Hospital Not in Hospital Unknown/not stated	
53-83	31	FILLER						
84	1	MAGE_IMPFLG	Mother's Age Imputed		A,S	Blank 1	Age not imputed Age imputed	
85	1	MAGE_REPFLG	Reported Age of Mother I	Flag	A,S	Blank 1	Reported age not used Reported age used	
86-87	2	MAGER	Mother's Single Year of A	.ge 432	A,S	12 13 14 15 16	12 years and under 13 years 14 years 15 years 16 years	
 * For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate S 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards 								

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						17	17 years
						18	18 years
						19	19 years
						20	20 years
						21	21 years
						22	22 years
						23	23 years
						24	24 years
						25 26	25 years
						26 27	26 years
						27 28	27 years
						28 29	28 years 29 years
						30	30 years
						31	31 years
						32	32 years
						33	33 years
						34	34 years
						35	35 years
						36	36 years
						37	37 years
						38	38 years
						39	39 years
						40	40 years
						41	41 years
						42	42 years
						43	43 years
						44	44 years
						45	45 years
						46 47	46 years
						47	47 years
						48	48 years
						49 50	49 years 50 years and over
						30	50 years and over
88-89	2	MAGER14	Mother's Age Recode 14	432	A,S	01	Under 15 Years
						03	15 years
						04	16 years
						05	17 years
						06	18 years
						07	19 years
	or the Versi	on column (Vers) the following codes apply:				
А			andard Birth Certificate				
S	1989	Version of US St	andard Birth Certificate				

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition	
						08 09 10 11 12 13 14	20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years	
90	1	MAGER9	Mother's Age Recode 9	432	A,S	1 2 3 4 5 6 7 8 9	Under 15 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years	
91-92	2	MBCNTRY	Mother's Birth Country (This item is available in the geographic codes are not av			A AA	-ZZ See Geographic Documentation	
93-96	4	FILLER						
97	1	MBSTATE_REC	Mother's Birth State Recod	e c409	А	1 2 3	Native born (50 U.S. States) Foreign born (includes territories) Unknown or Not Stated	
98-101	4	FILLER						
102-103	2	MRSTATEPSTL	Mother's Residence Postal (This item is available in the geographic codes are not av	e territory file only,	A,S file)			
			United States			AK AL AR	Alaska Alabama Arkansas	
 * For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate S 1989 Version of US Standard Birth Certificate s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards 								

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						AZ	Arizona
						CA	California
						CO	Colorado
						CT	Connecticut
						DE DC	Delaware District of Columbia
						DC FL	Florida
						FL GA	Georgia
						HI	Hawaii
						ID	Idaho
						IL	Illinois
						IN	Indiana
						IA	Iowa
						KS	Kansas
						KY	Kentucky
						LA	Louisiana
						MA	Massachusetts
						MD	Maryland
						ME	Maine
						MI	Michigan
						MN	Minnesota
						MO	Missouri
						MS	Mississippi
						MT	Montana
						NC	North Carolina
						ND	North Dakota
						NE	Nebraska
						NH	New Hampshire
						NJ NM	New Jersey New Mexico
						NV	Nevada
						NY	New York
						OH	Ohio
						OK	Oklahoma
						OR	Oregon
						PA	Pennsylvania
						RI	Rhode Island
						SC	South Carolina
						SD	South Dakota
						TN	Tennessee
						TX	Texas

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

- S
- 1989 Version of US Standard Birth Certificate 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position		Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
							UT VA VT WA WI WV WY	Utah Virginia Vermont Washington Wisconsin West Virginia Wyoming
				<u>Territories</u>			AS GU MP PR VI	American Samoa Guam Northern Marianas Puerto Rico Virgin Islands
				<u>Canadian Province</u>	e <u>s.</u>		AB BC MB NL NT NS NU ON PE QC SK YT	Alberta British Columbia Manitoba New Brunswick Newfoundland and Labrador Northwest Territories Nova Scotia Nunavut Ontario Prince Edward Island Quebec Saskatchewan Yukon Territory
				<u>Foreign</u>			XX ZZ	Foreign (Rest of world) Not Classifiable
				Mother's FIPS County of 1 (This item is available in the geographic codes are not av Data are suppressed if popu he following codes apply: dard Birth Certificate	e territory file only, vailable in the U.S. fil	le)	000 001-998	No county level geography County of residence (See Geographic Table)
	A S S	1989 V	ersion of US Stan	dard Birth Certificate dard Birth Certificate dard Birth Certificate but th	nis element conforr 8	ns to the	2003 star	ndards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						999	County of less than 100,000
107-111	5	FILLER					
112	1		Population of Residence (ilable in the territory file only, s are not available in the U.S.		A,S	0 1 2 3 9 Z	County of 1,000,000 or more County of 500,000 to 1,000,000 County of 250,000 to 500,000 County of 100,000 to 250,000 County less than 100,000 Foreign resident
113-129	17	FILLER					
130	1	RECTYPE	Record Type (This item is available in th geographic codes are not a		A,S le)	1	
						1 2	RESIDENT: State and county of occurrence and residence are the same. NONRESIDENT: State and county of occurrence and residence are different.
131	1	RESTATUS	Residence Status <u>United States</u>		A,S	1 2 3 4	RESIDENT: State and county of occurrence and residence are the same. INTRASTATE NONRESIDENT: State of occurrence and residence are the same but county is different. INTERSTATE NONRESIDENT: State of occurrence and residence are different but both are one of the 50 US states or District of Columbia. FOREIGN RESIDENT: The state of residence is not one of the 50 US states or District of Columbia.
* For t A S s	2003 V 1989 V	Version of US Star Version of US Star	<u>Territories</u> the following codes apply: ndard Birth Certificate ndard Birth Certificate ndard Birth Certificate but t	this element conform	ns to the	1 2003 star	RESIDENT: State and county of occurrence and residence

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						2 3 4	are the same. (Unique to Guam, all US residents are considered residents of Guam and thus are assigned 1.) INTRATERRITORY NONRESIDENT: Territory of occurrence and residence are the same but county is different. INTERTERRITORY RESIDENT: Territory of occurrence and residence are different but both are US Territories. FOREIGN RESIDENT: The residence is not a US Territory.
132-133	2	MRACE31	Mother's Race Recode 31	425	А	01	
						01	White (alone)
						02	Black (alone)
						03 04	AIAN (American Indian or Alaskan Native) (alone) Asian (alone)
						04	NHOPI (Native Hawaiian or Other Pacific Islander) (alone)
						05	Black And White
						07	Black and AIAN
						08	Black and Asian
						09	Black and NHOPI
						10	AIAN and White
						11	AIAN and Asian
						12	AIAN and NHOPI
						13	Asian and White
						14	Asian and NHOPI
						15	NHOPI and White
						16	Black, AIAN, and White
						17	Black, AIAN, and Asian
						18	Black, AIAN, and NHOPI
						19	Black, Asian, and White
						20	Black, Asian, and NHOPI
						21	Black, NHOPI, and White
						22	AIAN, Asian, and White
						23	AIAN, NHOPI, and White
						24	AIAN, Asian, and NHOPI
						25	Asian, NHOPI, and White
						26	Black, AIAN, Asian, and White
						27	Black, AIAN, Asian, and NHOPI
						28	Black, AIAN, NHOPI, and White
						29 20	Black, Asian, NHOPI, and White
						30 21	AIAN, Asian, NHOPI, and White
						31	Black, AIAN, Asian, NHOPI, and White
* For t A			the following codes apply: ndard Birth Certificate				

- S
- 1989 Version of US Standard Birth Certificate 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
134	1	MRACE6	Mother's Race Recode 6	425	А	1 2 3 4 5 6	White (alone) Black (alone) AIAN (alone) Asian (alone) NHOPI (alone) More than one race
135-136	2	MRACE15	Mother's Race Recode 15	425	Α	01 02 03 04 05 06 07 08 09 10 11 12 13 14 15	White (alone) Black (alone) AIAN (alone) Asian Indian (alone) Chinese (alone) Filipino (alone) Japanese (alone) Korean (alone) Vietnamese (alone) Other Asian (alone) Hawaiian (alone) Guamanian (alone) Samoan (alone) Other Pacific Islander (alone) More than one race
137	1	FILLER					
138	1	MRACEIMP	Mother's Race Imputed		A,S	Blank 1 2	Mother's race not imputed Unknown race imputed All other races, formerly coded 09, imputed.
139-141	3	FILLER					
142 * For t A S s	2003 V 1989 V	Version of US Sta Version of US Sta	Mother's Hispanic Origin the following codes apply: ndard Birth Certificate ndard Birth Certificate ndard Birth Certificate but th	424 nis element conf	A,S Forms to the	0 e 2003 sta	Non-Hispanic ndards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						1 2 3 4 5 9	Mexican Puerto Rican Cuban Central or South American Other and Unknown Hispanic Origin unknown or not stated
143	1	MHISPX	Mother's Hispanic Ori	gin		0 1 2 3 4 5 6 9	Non-Hispanic Mexican Puerto Rican Cuban Central or South American Dominican Other and Unknown Hispanic Origin unknown or not stated
144	1	SR_MRACEHIS	P Mother's Race/Hispanic C	Drigin c425	Α	1 2 3 4 5 6 7 8	Non-Hispanic White (alone) Non-Hispanic Black (alone) Non-Hispanic AIAN (alone) Non-Hispanic Asian (alone) Non-Hispanic NHOPI (alone) Non-Hispanic more than one race Hispanic Origin unknown or not stated
145	1	MEDUC	Mother's Educ-Revised	372	Α	1 2 3 4 5 6 7 8 9	8 th grade or less 9 th through 12 th grade with no diploma High school graduate or GED completed Some college credit, but not a degree Associate degree (AA, AS) Bachelor's degree (BA, AB, BS) Master's degree (MA, MS, MEng, Med, MSW, MBA) Doctorate (PhD, EdD) or Professional degree (MD, DDS, DVM, LLB, JD) Unknown

146-171 26 FILLER

* For the Version column (Vers) the following codes apply:

- 2003 Version of US Standard Birth Certificate А
- S
- 1989 Version of US Standard Birth Certificate 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
172	1	FAGERPT_FLG	Father's Reported Age Us	ed	A,S	Blank 1	Father's reported age not used Father's reported age used
173-176	4	FILLER					
177-178	2	FAGECOMB	Father's Combined Age	426 A,	S	09-98 99	Father's combined age in years Unknown or not stated
179-180	2	FAGEREC11	Father's Age Recode 11	426	A,S	01 02 03 04 05 06 07 08 09 10 11	Under 15 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years 55-98 years Not stated
181-182	2	PRIORLIVE	Prior Births Now Living	440	A,S	00-30 99	Number of children still living from previous live births. Unknown or not stated
183-184	2	PRIORDEAD	Prior Births Now Dead	441	A,S	00-30 99	Number of children dead from previous live births. Unknown or not stated
185-186	2	FILLER					
187	1	LBO_REC	Live Birth Order Recode	441	A,S	0-7 8 9	Number of live birth order. 8 or more live births Unknown or not stated
* For t A		Version of US Stan	he following codes apply: dard Birth Certificate				

S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
188-196	9	FILLER					
197-199	3	ILLB_R	Interval Since Last Live B	irth Recode 427	A	000-003 004-300 888 999	Plural delivery Months since last live birth Not applicable Unknown or not stated
200-201	2	ILLB_R11	Interval Since Last Live B	irth Recode 11 427	Α	00 01 02 03 04 05 06 07 08 88 99	Zero to 3 months (plural delivery) 4 to 11 months 12 to 17 months 18 to 23 months 24 to 35 months 36 to 47 months 48 to 59 months 60 to 71 months 72 months and over Not applicable Unknown or not stated
202-203	2	PRECARE	Month Prenatal Care Beg	an (Revised) 403	А	00 01-10 99	No prenatal care Month prenatal care began Unknown or not stated
204	1	PRECARE_REC	Month Prenatal Care Beg	an Recode (Revised) 403) A	1 2 3 4 5	1 st to 3 rd month 4 th to 6 th month 7 th to final month No prenatal care Unknown or not stated
205-228	24	FILLER					
229 * For t A S	2003	Version of US Stand	WIC ne following codes apply: dard Birth Certificate dard Birth Certificate	377	А	Y N	Yes No
S	1989 \	Version of US Stand	dard Birth Certificate but th	nis element conform	ns to the	2003 star	ndards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						U	Unknown or not stated
230-231	2	CIG_0	Cigarettes Before Pregnan	cy 405	Α	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
232-233	2	CIG_1	Cigarettes 1 st Trimester	406	А	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
234-235	2	CIG_2	Cigarettes 2 nd Trimester	407	Α	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
236-237	2	CIG_3	Cigarettes 3 rd Trimester	408	Α	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
238	1	CIG_REC	Cigarette Recode (Revised) 374	А	Y N U	Yes No Unknown or not stated
239-242	4	FILLER					
243-244	2	M_Ht_In	Mother's Height in Inches	375	А	30-78 99	Height in inches Unknown or not stated
245-248	4	BMI	BMI (prepregnancy)	445	А	13.0-69.9 99.9	9 Body Mass Index Unknown or not stated
249 * For	1 the Versi	BMI_R	Body Mass Index Recode (prepregnancy)	445	А	1 2 3 4	Underweight <18.5 Normal 18.5-24.9 Overweight 25.0-29.9 Obesity I 30.0-34.9

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						5 6 9	Obesity II 35.0-39.9 Extreme Obesity III ≥ 40.0 Unknown or not stated
250-252	3	FILLER					
253-255	3	PWgt_R	Prepregnancy Weight Re	code 376	A	075-375 999	Weight in pounds Unknown or not stated
256	1	FILLER					
257-262	6	Risk Factors (Re The cho	e <mark>vised)</mark> eckbox items indented below t	follow this code stru	A cture:		Y Yes N No U Unknown or not stated
257 258 259 260 261 262	1 1 1 1 1	RF_DIAB RF_GEST RF_PHYP RF_GHYP RF_ECLAM RF_INFTR	Prepregnancy Diabetes Gestational Diabetes Prepregnancy Hypertensi Gestational Hypertension Hypertension Eclampsia Infertility Treatment				C CIKIOWI OF INCESTATE
263	1	RF_FEDRG	Fertility Enhancing Drug	s 384	Α	Y N X U	Yes No Not applicable Unknown or not stated
264	1	RF_ARTEC	Asst. Reproductive Techn	ology 385	A	Y N X U	Yes No Not applicable Unknown
265	1	RF_CESAR	Previous Cesareans	386	A	Y N U	Yes No Unknown or not stated

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

Position	Len	Field	Description	Reporting Flag Positi		Vers*	Values	Definition
266-267	2	RF_CESARN	Previous Cesareans Num	387		A	00 01-30 99	None Number of previous cesareans Unknown or not stated
268-273	6	FILLER						
274-277 274	6 1	<u>Method of Deliv</u> ME_PRES	<u>erv (Revised)</u> Fetal Presentation	388		A	1 2 3 9	Cephalic Breech Other Unknown or not stated
275	1	ME_ROUT	Route & Method of Delive	ry :	389	Α	1 2 3 4 9	Spontaneous Forceps Vacuum Cesarean Unknown or not stated
276	1	ME_TRIAL	Trial of Labor Attempted	390	0	А	Y N X U	Yes No Not applicable Unknown or not stated
277	1	RDMETH_REC	Delivery Method Recode (389	Α	1 2 3 4 5 6 9	Vaginal (excludes vaginal after previous C section) Vaginal after previous c-section Primary C-section Repeat C-section Vaginal (unknown if previous c-section) (2003 Standard only) C-section (unknown if previous c-section) (2003 Standard only) Not stated

278-279 2 FILLER

* For the Version column (Vers) the following codes apply:

- A 2003 Version of US Standard Birth Certificate
- S 1989 Version of US Standard Birth Certificate
- s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
280	1	DMETH_REC	Delivery Method Recode	410	A,S	1 2 9	Vaginal C-section Not stated
281-282	2	<u>Materr</u> The che	nal Morbidity eckbox items indented below f	ollow this structure:	А	Y N U	Yes No Unknown or not stated
281 282	1 1	MM_RUPT MM_ICU	Ruptured Uterus Admit to Intensive Care	391 392			
283	1	ATTEND	Attendant	433	A,S	1 2 3 4 5 9	Doctor of Medicine (MD) Doctor of Osteopathy (DO) Certified Nurse Midwife (CNM) Other Midwife Other Unknown or not stated
284-300	17	FILLER					
301	1	DPLURAL	Plurality Recode		A,S	1 2 3 4	Single Twin Triplet Quadruplet or higher
302	1	FILLER					
303	1	IMP_PLUR	Plurality Imputed		A,S	Blank 1	Plurality is not imputed Plurality is imputed
304-315	12	FILLER					

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

- S
- 1989 Version of US Standard Birth Certificate 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards s

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
316	1	SEX	Sex of Infant		A,S	M F	Male Female
317	1	IMP_SEX	Sex Imputed Only where gestation is sta to be 20 weeks or more	ted or presumed	A,S	Blank 1	Infant Sex not Imputed Infant Sex is Imputed
		Gestation					
318-319	2	DLMP_MM	Last Normal Menses Mon	nth 429	A,S	01 02 03 04 05 06 07 08 09 10 11 12 99	January February March April May June July August September October November December Unknown or not stated
320-321	2	FILLER					
322-325	4	DLMP_YY	Last Normal Menses Year	r 431	A,S	nnnn 9999	Year of last normal menses Unknown or not stated
326-328	3	FILLER					
329	1	GEST_IMP	Gestation Imputed Flag		A,S	Blank 1	Gestation is not imputed Gestation is imputed
330	1	OBGEST_FLG	Obstetric Estimate of Ges	station Used Flag	A,S	Blank 1	Obstetric Estimate is not used Obstetric Estimate is used
331-332	2	COMBGEST	Gestation – Detail in Wee	ks 404	A,S	02-47 99	2 nd through 47 th week of Gestation Unknown

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

- S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
333-334	2	GESTREC12	Gestation Recode 12	404	A,S	01 02 03 04 05 06 07 08 09 10 11 12	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks 28-31 weeks 32-33 weeks 34-36 weeks 37-38 weeks 39-40 weeks 41 weeks 42 weeks and over Unknown
335	1	GESTREC5	Gestation Recode 5	404	A,S	1 2 3 4 5	Under 20 weeks 20-23 weeks 24-27 weeks 28 weeks and over Unknown
336-337	2	OEGest_Unedt	Obstetric Gestation 373	A,S		0-98 02-47 99	Range of obstetric estimate of gestation - revised Range of clinical estimate of gestation - unrevised Unknown or not stated
338	1	FILLER					
339	1	COMBGEST_US	GED Combined Gesta	ation Used Flag	A,S	blank 1	Combined gestation not used Combined gestation used
340-341	2	OEGest_Comb	Obstetric Estimate Edited (NCHS Standard item)	I 373 A,S		02-47 99	Weeks of gestation Not stated
342-343	2	OEGest_R12	Obstetric Estimate Recod (NCHS Standard item)	e12 373 A,S		01 02 03 04 05 06 07 08	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks 28-31 weeks 32-33 weeks 34-36 weeks 37-38 weeks
* For t A			the following codes apply: adard Birth Certificate				
S			idard Birth Certificate				
S	1989 \	Version of US Star	ndard Birth Certificate but t	this element confo	rms to the	e 2003 sta	ndards

20

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
344	1	OEGest_R5	Obstetric Estimate Recod (NCHS Standard item)	e 5 373 A,S		09 10 11 12 1 2 3 4 5	 39-40 weeks 41 weeks 42 weeks and over Unknown Under 20 weeks 20-23 weeks 24-27 weeks 28 weeks and over Unknown
345-348	4	FILLER					
349-352	4	DBWT	Birth Weight – Detail in C	Grams 442	A,S	0001-81 9999	65 Number of grams Not stated birth weight
353-354	2	BWTR14	Birth Weight Recode 14	442	A,S	01 02 03 04 05 06 07 08 09 10 11 12 13 14	0249 grams or less 0250 – 0349 grams 0350 – 0499 grams 0500 – 0999 grams 1000 – 1499 grams 1500 – 1999 grams 2000 – 2499 grams 2500 – 2999 grams 3000 – 3499 grams 3500 – 3999 grams 4000 – 4499 grams 4500 – 4999 grams 5000 – 8165 grams Unknown or Not Stated
355 * For t		BWTR4	Birth Weight Recode 4 the following codes apply:	442	A,S	1 2	1499 grams or less 1500 – 2499 grams

2003 Version of US Standard Birth Certificate А

S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						3 4	2500 - 8165 grams Unknown or not stated
356	1	FILLER					
357	1	ESTOFD	Estimated Time of Fetal D	eath 435	А	N L A U	At assessment, no labor At assessment, labor Labor, no assessment Unknown
358	1	AUTOPSY	Was Autopsy Performed?	436	А	Y N P	Yes No Planned
359	1	HISTOPF	Was Histological Placenta	l Exam Performed?	437 A	Y N P	Yes No Planned
360	1	AUTOPF	Was Autop/Hist results us	ed in cause? 438	А	Y N X	Yes No Not applicable
361-371	11	FILLER					
372-438 372 373 374 375 376 377 378 379 380 381 382 * For th	1 1 1 1 1 1 1 1 1 1 1 1 1	F_MEDUC F_CLINEST F_TOBACO F_M_HT F_PWGT F_WIC F_RF_PDIAB F_RF_PDIAB F_RF_GDIAB F_RF_PHYPER F_RF_GHYPER F_RF_ECLAMP	orting Flags s indented below follow this c Education of Mother Obstetric Estimate of Gestat Tobacco Use Mother's Height Prepregnancy Weight WIC Prepregnancy Diabetes Gestational Diabetes Prepregnancy Hypertension Gestational Hypertension Hypertension Eclampsia he following codes apply:	ion	A A,S A A A A A A A A	01	Not reporting Reporting
A S s	1989 V	version of US Stand	dard Birth Certificate dard Birth Certificate dard Birth Certificate but tl	nis element conform 22	ns to the	2003 star	ndards

Position	Len
Position	L

Flag	Position
------	----------

383	1	F_RF_INFT Infe	ertility Treatment		А
384	1	F_RF_INFT_DRG Fert			A
385	1	F_RF_INFT_ART Rep	productive Technology		A
386	1		vious Cesarean		A
387	1		nber of Previous Cesareans		A
388	1	F_MD_PRESENT Feta	al Presentation	1	4
389	1	F_MD_ROUTE Fina	al Route and Method of De	livery	A
390	1	F_MD_TRIAL Tria	al of Labor Attempted		A
391	1	F_MM_RUPTUR Rupt	tured Uterus	A	A
392	1	F_MM_ICU Adm	nission to Intensive Care	I	4
393-402	10	FILLER			
403	1	F_MPCB Mor	nth Prenatal Care Began		А
404	1	F_CMBGST Con	nbined Gestation		A,S
405	1	F_CIGS_0 Cig	garettes Before Pregnancy		А
406	1		garettes 1 st Trimester		А
407	1		garettes 2 nd Trimester		А
408	1	F_CIGS_3 Cig	garettes 3 rd Trimester		А
409	1	F_FACILITY Fac	cility		А
410	1	DelMethRecF Deli	ivery Method Recode Flag		A,S
411-423	13	FILLER			
424	1	F_MORIGIN Mot	ther's Hispanic origin		A,S
425	1	F_MRACE_R Mo	other's Race (Revised)		Α
426	1	F FAGE u Fath	her's Age		A,S
427	1	F_DLLB_MM Last I	Live Birth Month (Revised)		А
428	1	FILLER			
429	1	F_DLMP_MM Last r	normal menses month		A,S
430	1	FILLER			
431	1	F DLMP YY Last n	normal menses year		A,S
432	1	F_MAGE Mother's A	ge		A,S
433	1	F_ATTENDANT Atte	endant		A,S
434	1	FILLER			
435	1	F_ESTOFD Estim	mated Time of Fetal Death	(Revised)	Α
436	1	F AUTOPSY Auto	opsy (Revised)		А
437	1	F_HISTOPF Hist	tological Placental Exam (R	levised)	Α
438	1	F AUTOPF Auto	opsy/Histological Exam Us	ed in Cause	А
439	1	F_MRACEREC Mot	ther's Bridged Race Recode	2	A,S
440	1	F_PLBL Nu	umber of previous live birth	ns living	
441	1	F_PLBD Nu	mber of previous live births	s dead	A
442	1	F_DBWT Birt	thweight	A,S	
443-444	2	FILLER	-		
445	1	F_BMI Bod	ly mass index	A	

* For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

- S 1989 Version of US Standard Birth Certificate
- 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

A,S A,S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition

446-2602 2164 FILLER 2603-2642 40 Coded Causes & Conditions А The fields indented below follow this structure: ICD10 Code Coded Value Blank None reported Initiating Cause or Condition Code 2603-2607 5 IICOD 2651 2608-2642 35 FILLER 2643-2645 3 IC 124 Fetal Initiating Fetal Recode 124 2651 А 001-124 2646-2650 5 FILLER 2651 Flag File for Reporting Flags The reporting flags indented below follow this coding structure: Not reporting 0 Reporting 1 F_ICOD Init Cause or condition 2651 1 А 2652-3150 499 FILLER

3150 END OF RECORD

* For the Version column (Vers) the following codes apply:

- A 2003 Version of US Standard Birth Certificate
- S 1989 Version of US Standard Birth Certificate
- s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

_		Fetal de	eaths	Fetal m	nortality	rate\1
	Total\2	20-27 28 weeks weeks\3 or more\3		Total\2	20-27 weeks\3	28 weeks or more\3
_		Obstet	ric estimat	e of gestati	on\4	
2022	20,202	10,246	9,956	5.48	2.79	2.71
2021	21,105	10,824	10,281	5.73	2.95	2.80
2020	20,854	10,764	10,090	5.74	2.97	2.78
2019	21,478	11,216	10,262	5.70	2.98	2.73
2018	22,459	11,844	10,615	5.89	3.11	2.79
2017	22 , 827	11,861	10,966	5.89	3.07	2.84
2016	23,880	12,486	11,394	6.02	3.15	2.88
2015	23 , 776	12,407	11,369	5.94	3.11	2.85
2014	23,980	12,652	11,328	5.98	3.16	2.83
			LMP-based g	estation\5		
2014 -	23,999	11,948	12,051	5.98	2.99	3.01
	is number and fetal		eaths in spe	ecified group	per 1,00	0 live
	deaths wi or more.	th stated c	or presumed	period of ge	station o	f 20
\3Not s	stated gest	ational age	e proportion	nally distrib	outed.	

Table 1. Number of fetal deaths and fetal mortality rates: United States, 2014-2022

\4Beginning with the 2014 data year, the obstetric estimate of gestation at delivery replaces the measure based on the date of last normal menses as the standard for measuring gestational age. \5Date of last normal menses (LMP) was used for measuring gestational age.

Table 2. Fetal Deaths by Period of Gestation: United States, Each State and Territory, 2022

Fetal Deaths with Stated or Presumed State/territory of Gestation of 20 weeks or More residence 20-27 28 Weeks Total Not stated Weeks and Over United States \1 20,202 10,173 9,886 Alabama _ Alaska _ Arizona Arkansas California 2,115 1,103 Colorado _ Connecticut _ Delaware District of Columbia _ Florida 1,548 Georgia 1,000 Hawaii Idaho _ Illinois Indiana Iowa _ Kansas Kentucky _ Louisiana _ Maine _ Maryland _ Massachusetts _ Michigan Minnesota _ Mississippi Missouri Montana _ Nebraska Nevada _ New Hampshire _ New Jersey _ New Mexico _ New York New York City _ North Carolina North Dakota _ Ohio _ Oklahoma Oregon Pennsylvania _ Rhode Island _ South Carolina

[Nonresidents of the United States are excluded]

South Dakota	54	24	30	-
Tennessee	538	263	273	2
Texas	1,613	753	853	7
Utah	252	124	128	-
Vermont	21	7	14	-
Virginia	497	257	233	7
Washington	421	213	208	-
West Virginia	70	26	44	-
Wisconsin	320	171	147	2
Wyoming	36	16	20	-
American Samoa				
Guam	34	12	21	1
Northern Marianas				
Puerto Rico	218	144	73	1
Virgin Islands				

NOTE: American Samoa did not report data for 2022. Data for the Northern Marianas and the Virgin Islands were excluded due to small numbers. - Quantity zero. --- Data unavailable. \lExcludes data for the territories. SOURCE: National Center for Health Statistics, National Vital Statistics System.

_		All Fetal		
State / territory	Total	Under 20 Weeks	20 Weeks and Over	Not stated
United States\1	40,113	18,424	20,218	1,471
Alabama	410	_	410	-
Alaska	53	2	51	-
Arizona	497	6	487	4
Arkansas	484	230	253	1
California	2,144	1	2,119	24
Colorado	325	4	321	-
Connecticut	180	-	180	-
Delaware	57	-	56	1
District of Columbia	99	2	97	-
Florida	1 , 570	3	1,504	63
Georgia	8,214	6,477	1,006	731
Hawaii	474	286	96	92
Idaho	114	30	84	-
Illinois	733	-	732	1
Indiana	446	1	441	4
Iowa	160	1	159	-
Kansas	215	-	212	3
Kentucky	274	-	274	-
Louisiana	349	58	291	-
Maine	57	-	57	-
Maryland	351	_	351	-
Massachusetts	285	-	285	-
Michigan	611	7	603	1
Minnesota	360	-	360	-
Mississippi	316	6	307	3
Missouri	380	7	369	4
Montana	52	-	52	-
Nebraska	107	-	106	1
Nevada	238	4	234	-
New Hampshire	54	-	54	-
New Jersey	487	-	487	-
New Mexico	68	5	63	-
New York	4,228	3,372	509	347
New York City	5,058	4,351	701	6
North Carolina	708	2	700	6
North Dakota	53	-	53	-
Ohio	742	-	742	-
Oklahoma	287	1	283	3
Oregon	235	41	191	3
Pennsylvania	1,035	369	663	3
Rhode Island	361	233	60	68
South Carolina	263	1	262	-
South Dakota	64	2	62	-
Tennessee	587	3	582	2
Texas	1,948	247	1,694	7

Table 3. Fetal Deaths by Period of Gestation and State/Territory of Occurrence: United States, 2022

Utah	258	-	258	-
Vermont	18	-	18	-
Virginia	3,250	2,665	494	91
Washington	433	-	433	-
West Virginia	84	6	78	-
Wisconsin	301	1	298	2
Wyoming	36	-	36	-
American Samoa				
Guam	34	-	33	1
Northern Marianas				
Puerto Rico	225	2	222	1
Virgin Islands				

NOTE: American Samoa did not report data for 2022. Data for the Northern Marianas and the Virgin Islands were excluded due to small numbers.

- Quantity zero.

--- Data unavailable.

\1Excludes data for the territories.

Table 4. Number and percentage of fetal deaths by race and Hispanic origin of mother: United States, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Race		umber	Per	centage
	Total\1	Non-Hispanic	Total\1	Non-Hispanic
All races\2	20,202	14,926	100.0	100.0
One race	19 , 777	14,580	97.9	97.7
White	12,714	8,280	62.9	55.5
Black	5,764	5,194	28.5	34.8
American Indian and Alaska Native (AIAN)	224	187	1.1	1.3
Asian	925	813	4.6	5.4
Native Hawaiian and Other Pacific Islander (NHOPI)	150	106	0.7	0.7
More than one race	425	346	2.1	2.3
Two races	377	311	1.9	2.1
Black and White	197	157	1.0	1.1
Black and AIAN	15	14	0.1	0.1
Black and Asian	11	10	0.1	0.1
Black and NHOPI	3	3	0.0	0.0
AIAN and White	51	45	0.3	0.3
AIAN and Asian	2	1	0.0	0.0
AIAN and NHOPI	2	2	0.0	0.0
Asian and White	73	63	0.4	0.4
Asian and NHOPI	12	9	0.1	0.1
NHOPI and White	11	7	0.1	0.0
Three races	46	35	0.2	0.2
Black, AIAN and White	16	13	0.1	0.1
Black AIAN and Asian	-	-	-	-
Black, AIAN and NHOPI	1	1	0.0	0.0
Black, Asian and White	8	8	0.0	0.1
Black, Asian and NHOPI	1	1	0.0	0.0
Black, NHOPI, and White	1	1	0.0	0.0
AIAN, Asian and White	-	-	-	-
AIAN, NHOPI and White	-	-	-	-
AIAN, Asian and NHOPI	-	-	-	-
Asian, NHOPI and White	19	11	0.1	0.1
Four races	2	-	0.0	-
Black, AIAN, Asian and White	-	-	-	-
Black, AIAN, Asian, and NHOPI	-	-	-	-
Black, AIAN, NHOPI and White	1	-	0.0	-
Black, Asian, NHOPI and White	-	-	_	-
AIAN, Asian, NHOPI and White	1	-	0.0	-
Five races	-	-	-	-
Black, AIAN, Asian, NHOPI and White	-	-	_	-

- Quantity zero.

0.0 Quantity more than zero but less than 0.05.

\1 Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

\2 Race and Hispanic origin are reported separately on reports of fetal death. Persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 5A. Number and percentage of fetal deaths by selected demographic and health characteristics: United States, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

more.]	Number	Percent
Day of week		
Total	20,202	100.0
Sunday	2,138	10.6
Monday	2,361	11.7
Tuesday	3,064	15.2
Wednesday	3,182	15.8
Thursday	3,265	16.2
Friday	3,336	16.5
Saturday	2,856	14.1
Month of delivery		
Total	20,202	100.0
January	1,829	9.1
February	1,521	7.5
March	1,711	8.5
April	1,659	8.2
Мау	1,702	8.4
June	1,692	8.4
July	1,762	8.7
August	1,672	8.3
September	1,666	8.2
October	1,645	8.1
November	1,655	8.2
December	1,688	8.4
Place of delivery		
Total	20,202	100.0
Hospital\1	19 , 617	97.1
Freestanding birthing center	15	0.1
Clinic or doctor's office	42	0.2
Residence	357	1.8
Other	137	0.7
Not specified	34	0.2
Age of mother		
Total	20,202	100.0
<15	16	0.1
15-19	991	4.9
20-24	3,631	18.0
25-29	5,071	25.1
30-34	5,634	27.9
35-39	3,613	17.9
40-44	1,138	5.6
45-49	96	0.5
50-54	12	0.1
Race and Hispanic origin of mother		
Total\2	20,202	100.0
Non-Hispanic, single race\3		
Non-Hispanic white	8,280	41.0
Non-Hispanic black	5,194	25.7
Non-Hispanic American Indian or Alaska Native	187	0.9
Non-Hispanic Asian	813	4.0
Non-Hispanic Native Hawaiian or Other Pacific Islander	106	0.5
Non-Hispanic multiple race	346	1.7
Hispanic\4	4,359	21.6
Unknown	917	4.5

Hispanic origin of mother		
Total	20,202	100.0
Non-Hispanic	14,926	73.9
Mexican	2,400	11.9
Puerto Rican	323	1.6
Cuban	102	0.5
Central or South American	779	3.9
Dominican	142 613	0.7
Other and unknown Hispanic Unknown	917	3.0 4.5
UIKIIOWII	917	4.5
Education of mother\5		
Total	20,053	100.0
8th grade or less	780	3.9
9th-12th grade with no diploma	1,696	8.5
High school graduate\6	6,121	30.5
Some college credit, but no degree	3,123	15.6
Associate's degree\7	1,300	6.5
Bachelor's degree\8 Master's degree\9	2,746 1,173	13.7 5.8
Doctorate or professional degree\10	333	1.7
Unknown	2,781	13.9
	_,	2019
Age of father\11		
Total	17,223	100.0
<15	4	0.0
15-19	320	1.9
20-24	1,664	9.7
25-29	2,932	17.0
30-34 35-39	3,734	21.7
40-44	2,723 1,421	15.8 8.3
45-49	483	2.8
50-54	177	1.0
55-98	119	0.7
Unknown	3,646	21.2
Trimester prenatal care began\12		100 0
Total 1st trimester	20,053 12,391	100.0 61.8
2nd trimester	2,896	14.4
3rd trimester	339	1.7
No care	1,945	9.7
Unknown	2,482	12.4
Mother received WIC food for herself during this pregnancy\13		
Total	18,980	100.0
Yes	3,676	19.4
No	12,527	66.0
Not stated	2,777	14.6
Tobacco use during pregnancy\14		
Total	20,150	100.0
Yes	1,318	6.5
No	17,126	85.0
Unknown	1,706	8.5
Maternal pre-pregnancy body mass index\15		
Total	20,105	100.0
Underweight (BMI less than 18.5)	465	2.3
Normal weight (BMI of 18.5 to 24.9) Overweight (BMI of 25.0 to 29.9)	5,303 4,581	26.4 22.8
Obese (BMI of 30.0 and over)	4,581 7,118	22.8 35.4
Not stated	2,638	13.1
	2,000	⊥ V • ⊥

Method of delivery\16 Total	20,105	100.0
Vaginal (excludes vaginal after previous C-section)	13,608	67.7
Vaginal after previous C-section	1,602	8.0
Primary C-section	1,959	9.7
Repeat C-section	1,255	6.2
Vaginal (unknown if previous c-section)	617	3.1
C-section (unknown if previous c-section)	124	0.6
Not stated	940	4.7
Attendant at delivery		
Total	20,202	100.0
Doctor of medicine	16,865	83.5
Doctor of osteopathy	1,804	8.9
Certified nurse midwife	720	3.6
Other midwife	44	0.2
Other Users si final	567	2.8
Unspecified	202	1.0
Plurality		100.0
Total	20,202 18,805	100.0 93.1
Singleton Twin	1,326	93.1 6.6
Triplet	£, 520	0.3
Quadruplet or higher	6	0.0
gaaalapice of higher	0	0.0
Sex		100.0
Total Male	20,202 10,519	100.0 52.1
Female	9,683	47.9
	5,000	11.5
Obstetric estimate of gestation		
Total	20,202	100.0
20-23 weeks	7,087	35.1
24-27 weeks	3,086	15.3
28-31 weeks	2,487	12.3
32-33 weeks	1,425	7.1
34-36 weeks	2,613	12.9
37-38 weeks	1,873	9.3
39-40 weeks	1,275	6.3
41 weeks 42 weeks and over	159 54	0.8 0.3
Unknown	143	0.3
	110	0.7
Birthweight		
Total	20,202	100.0
<500 grams	6,554	32.4
0500 - 0999 grams	3,321	16.4
1000 - 1499 gram	1,915	9.5
1500 - 1999 gram	1,698	8.4
2000 - 2499 gram 2500 - 2999 gram	1,595	7.9
2500 - 2999 gram 3000 - 3499 gram	1,553 1,141	7.7 5.6
3500 - 3999 gram	573	2.8
4000 - 4499 gram	207	1.0
4500 - 4999 gram	92	0.5
5000 - 8165 gram	52	0.3
Unknown	1,494	7.4
0.0 Quantity more than zero but less than 0.05.	•	

0.0 Quantity more than zero but less than 0.05.

- Quantity zero.

 $\label{eq:lincludes}$ fetal deaths occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death. \4Includes all persons of Hispanic origin of any race.

5Hawaii and Rhode Island reported education of mother for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

\6Includes General Educational Development (GED).

\7Includes Associate in Arts and Associate in Science.

 \Bachelor in Arts and Bachelor in Science.

 $\9Includes$ Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\10Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

\11Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New York, Rhode Island, and Virginia reported father's age for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

12Hawaii and Rhode Island reported date of first prenatal care for 2022 but had unknown levels of 50% or higher and were flagged as not reporting.

\13Mother received WIC food for herself durng this pregnancy is not reported by New York City. Hawaii, Indiana, and Rhode Island reported WIC for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

\14Rhode Island reported tobacco use during pregnancy for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

15Hawaii reported maternal pre-pregnancy body mass index for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

 $\16Hawaii$ reported method of delivery for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

Table 5B. Number and percentage of fetal deaths by selected demographic and health characteristics: Guam and Puerto Rico, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

more.]	Number	Percent
Day of week		
Total	252	100.0
Sunday	27	10.7
Monday	27	10.7
Tuesday	43	17.1
Wednesday	49	19.4
Thursday	37	14.7
Friday	43	17.1
Saturday	26	10.3
Month of delivery		
Total	252	100.0
January	29	11.5
February	19	7.5
March	16	6.3
April	13	5.2
Мау	26	10.3
June	30	11.9
July	21	8.3
August	19	7.5
September	17	6.7
October	21	8.3
November	24	9.5
December	17	6.7
Place of delivery		
Total	252	100.0
Hospital\1	246	97.6
Freestanding birthing center	-	-
Clinic or doctor's office	-	-
Residence	3	1.2
Other	2	0.8
Not specified	1	0.4
Age of mother		
Total	252	100.0
<15	-	-
15-19	17	6.7
20-24	55	21.8
25-29	78	31.0
30-34	59	23.4
35-39	28	11.1
40-44	14	5.6
45-49 50-54	1	0.4
Race and Hispanic origin of mother Total\2	252	100.0
Non-Hispanic, single race\3		
Non-Hispanic white	3	1.2
Non-Hispanic black	1	0.4
Non-Hispanic American Indian or Alaska Native	-	_
Non-Hispanic Asian	5	2.0
Non-Hispanic Native Hawaiian or Other Pacific Islander	27	10.7
Non-Hispanic multiple race	2	0.8
Hispanic\4	214	84.9
Unknown	-	_

Hispanic origin of mother		
Total	252	100.0
Non-Hispanic	38	15.1
Mexican	1	0.4
Puerto Rican	207	82.1
Cuban	-	-
Central or South American	1	0.4
Dominican	5	2.0
Other and unknown Hispanic Unknown	-	_
Education of mother		
Total	252	100.0
8th grade or less	4	1.6
9th-12th grade with no diploma	23	9.1
High school graduate\5	81	32.1
Some college credit, but no degree	27	10.7
Associate's degree\6	36	14.3
Bachelor's degree\7	62	24.6 6.0
Master's degree\8 Doctorate or professional degree\9	15 2	0.8
Unknown	2	0.8
	2	0.0
Age of father\10		
Total	218	100.0
<15	-	-
15-19	8	3.7
20-24	33	15.1
25-29	39	17.9
30-34	33	15.1
35-39 40-44	29 17	13.3 7.8
45-49	4	1.8
50-54	3	1.4
55-98	-	_
Unknown	52	23.9
Trimester prenatal care began		
Total	252	100.0
1st trimester 2nd trimester	185 36	73.4 14.3
3rd trimester	1	0.4
No care	15	6.0
Unknown	15	6.0
Mother received WIC food for herself during this pregnancy $\setminus 11$		
Total	218	100.0
Yes	152	69.7
No	57	26.1
Not stated	9	4.1
Tobacco use during pregnancy\12		
Total	218	100.0
Yes	1	0.5
No	216	99.1
Unknown	1	0.5
Maternal pre-pregnancy body mass index\13	010	100.0
Total Underweight (PMI less than 18 5)	218 6	100.0 2.8
Underweight (BMI less than 18.5) Normal weight (BMI of 18.5 to 24.9)	61	2.8
Overweight (BMI of 25.0 to 29.9)	63	28.0
Obese (BMI of 30.0 and over)	82	37.6
Not stated	6	2.8

Decision of certivery 120 100.0 Vaginal first previous C-section 130 75.4 Vaginal first previous C-section 24 05.5 Frinary C-section 26 10.3 Vaginal first previous C-section 7 2.9 Vaginal (unknown if previous c-section) - - C-section (unknown if previous c-section) 4 0.4 Not stated 4 1.6 Attendant at delivery 5 2.0 Doctor of cateopathy 5 2.0 Octor of cateopathy 5 2.0 Octor of cateopathy 5 2.0 Other miduife - - Other miduife - - Unspecified 13 5.2 Divin 13 5.2 Striptet - - Outary 13 5.2 Outary 13 5.2 Outary - - Outary - - Outary 12			
Taginal rescludes vaginal after previous C-section 180 75.4 Vaginal fract previous C-section 26 10.3 Repeat C-section 7 2.8 Vaginal fract previous c-section) - - C-section (unknown if previous c-section) 1 0.4 Attendant at delivery - - Total 263 100.0 Doctor of selection 5 2.0 Obstrated - - Other - - Other - - Obstrated 252 100.0 Singlaten 233 94.8 Wain 13 5.2 Obstrate - - Obstrate - - Outher - - Obstrate 252 100.0 Singlaten 253 100.0 Son - - Obstrate - - Obstrate 252 100.0 Ide	Method of delivery	252	100 0
Yajial ifer previous C-section 24 9.5 Primary C-section 7 2.8 Vaginal (unknown if previous c-section) 1 0.4 Not stated 4 1.6 Attendant at delivery 252 100.0 Doctor of calcione 233 96.4 Doctor of calcione 239 96.4 Doctor of calcione 232 100.0 Cartified nurse siduife - - Other - - Unapacified 13 5.2 Plurality - - Triplet - - Cuadruptet or higher - - Costoric cestinate of gestation 122 100.0 Dotal 222 100.0 Dotal 222 100.0 Cartified nures </td <td></td> <td>-</td> <td></td>		-	
Primary C-section 26 10.3 Repeat C-section - - C-section (unknown if previous c-section) 1 0.4 Not stated 4 1.6 Attendant at delivery - - Total 223 100.0 Doctor of nedicine 243 96.4 Doctor of costeopathy 5 2.0 Chars mixedife - - Chars mixedife - - Obspecified 4 1.6 Plorality - - Triplet - - Quadpulet or higher - - Sax - - Costertic estimate of gestation - - Obstertic estimate of gestation - - Costertic est			
Repeat C-section 7 2.8 Vaginal (unknown if previous c-section) 1 0.4 Not stated 4 1.6 Attendant at delivery 252 100.0 Doctor of sedicine 243 9.4 Dector of sedicine 24 1.6 Timitifie - - - Other mitwife - - - Output 13 5.2 100.0 Singleton 13 5.2 100.0 Nale 128 50.8 12 Nale 128 50.8 12 Total 25 9.9 12.4 4.3			
vaginal (unknown if previous c-section) - - C-section (unknown if previous c-section) 1 0.4 Not stated 4 1.6 Attendant at delivery - - Total 252 100.0 Doctor of medicine 243 96.4 Doctor of seteopathy 5 2.0 Certified untre midwife - - Other midwife - - Unspecified 4 1.6 Plurality - - Triplet - - Cadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Female 122 100.0 20-23 weeks 122 100.0 22-33 weeks 12 10.0 23-33 weeks 20 0.7 31-4 22 0.7 22-3 weeks 11 4.4 32-3 weeks	-		
c-section (unknown if previous c-section) 1 0.4 Not stated 4 1.6 Attendant at delivery 252 100.0 Doctor of medicine 243 96.4 Doctor of satespathy 5 2.0 Certified nurse midwife - - Other midwife - - Plurality - - Cotal 252 100.0 Singleton 239 94.8 Twin 13 5.2 Potal 252 100.0 Male 128 5.8 Perale 22 10.0 Obstetric estimate of gestation		_	-
Not. stated 4 1.6 Attendant at delivery 252 100.0 Doctor of nedicine 243 96.4 Doctor of soteopathy 5 2.0 Certified nurse midwife - - Other midwife - - Unspecified 4 1.6 Plarality - - Total 252 100.0 Singleton 239 94.8 Twin 13 5.2 Triplet - - Oudruplet or higher - - Sex - - Total 252 100.0 Male 128 53.8 Penale 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 212 48.4 21-23 weeks 22 8.7 21-33 weeks 20 7.9 21-40 4.4 4.4		1	0.4
Total 252 100.0 Doctor of medicine 243 96.4 Doctor of osteopathy 3 2.0 Certified nurse midwife - - Other - - Other - - Unspecified 4 1.6 Plurality - - Total 252 100.0 Singleton 233 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Female 122 40.2 Obsterric estimate of gestation - - Total 252 100.0 24-27 weeks 34 13.5 24-27 weeks 24 49.2 33 1.2 44.4 436 weeks 2 8.7 340	Not stated	4	1.6
Doctor of madicine 243 96.4 Doctor of osteopathy 5 2.0 Other midwife - - Unspecified 4 1.6 Plurality 232 100.0 Singleton 233 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex 7 - Total 252 100.0 Male 233 94.8 Twin 13 5.2 Total 252 100.0 Male 128 5.8 Female 124 49.2 Obstetric estimate of gestation 7 - Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 25 5.9 32-33 weeks 20 7.9 32-40 weeks 3 1.2 41 weeks 3 1.2	Attendant at delivery		
Doctor of osteopathy 5 2.0 Cartified nurse midwife - - Other - - Other - - Unspecified 4 1.6 Plurality 239 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Pemale 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 34 13.5 24-27 weeks 34 13.5 23-31 weeks 22 8.7 23-33 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 3 1.2 42 weeks and over 104 4.4 1000 - 1499 gram 18 7.1		-	
Certified nusse midwife - - Other - - Unspecified 4 1.6 Plurality 252 100.0 Total 239 94.8 Twin 13 5.2 Quadruplet or higher - - Sex - - Total 252 100.0 Male 13 5.2 Total 252 100.0 Male - - Total 252 100.0 Male 128 55.8 Female 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 12 48.4 24-31 weeks 25 9.9 32-33 weeks 20 7.9 33-40 weeks 12 4.8 41 weeks 10.4 4 42 weeks and over 10.3 40.9 000 -		-	
Other midwife - - Other - - Other - - Objectified 252 100.0 Singleton 239 94.8 Tvin 13 5.22 Triplet - - Quadruplet or higher - - Sex - - Total 252 100.0 Male - - Total 252 100.0 Male 128 50.8 Female 122 40.4 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 12 48.4 24-21 weeks 25 9.9 32-33 weeks 11 4.4 34 13.12 4.8 24-30 weeks 20 7.9 37-38 weeks 20 7.9 34-40 weeks 3 1.2 41 weeks		5	2.0
Other - - - Unspecified 4 1.6 Plurality 239 94.8 Twin 1.3 5.2 Quadruplet or higher - - Quadruplet or higher - - Sex Total 252 100.0 Male 252 100.0 Male 252 100.0 Male 252 100.0 Sex Total 252 100.0 Male 252 100.0 10 Cobstetric estimate of gestation Total 252 100.0 Cotal 252 100.0 10 44.3 20-23 weeks 34 13.5 2 10.2 22-33 weeks 21 24.8 3 1.2 23-34 weeks 20 7.9 3-40 4.4 34-36 41 weeks 1 1.4 4.4 34-36 2.2 8.7 100 or 1499 gram 1.03		-	-
Unspecified 4 1.6 Plurality 252 100.0 Singleton 239 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Female 124 50.8 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 12 48.4 24-27 weeks 24 48.4 14.4 24-36 weeks 12 48.4 24-27 weeks 20 7.9 34-40 weeks 3 1.2 34-36 weeks 1 0.4 0.4 0.4 0.4 41 weeks 3 1.2 4.8 1.0 4.4		-	-
Plurality 252 100.0 Singleton 239 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 23 44.4 24-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 10 0.4 Unknown 2 0.8 Birthweight - -			_ 1 C
Total 252 100.0 Singleton 239 94.8 Triplat - - Quadruplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 Vala 252 100.0 Vala 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 34 13.5 22-33 weeks 20 7.9 32-33 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 0500 - 0999 grams 103 40.9 0500 - 0999 grams 13 7.1 1000 - 1499 gram 16 6.	Unspecified	4	1.0
Singleton 239 94.8 Twin 13 5.2 Triplet - - Quadruplet or higher - - Sex 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 12 4.8 41 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 18 7.1 1500 - 1999 gram 16 6.3 2000 - 2499 gram 12 4.8 <td></td> <td></td> <td>100 0</td>			100 0
Twin 13 5.2 Triplet - - Quaduplet or higher - - Sex - - Total 252 100.0 Male 128 50.8 Pemale 124 49.2 Obstetric estimate of gestation - - Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 12 4.8 7000 - 1499 gram 16 6.3 2000 - 2499 gram 103 40.9 2000 - 2499 gram 12 4.8 2000 - 2499 gram 16 6.3 2000 - 2499 gram <td></td> <td></td> <td></td>			
Triplet - - Quadruplet or higher - - Sex 70tal 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation 252 100.0 70tal 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Uhknown 2 0.8 Birthweight 25 100.0 7000 - 1499 gram 18 7.1 1000 - 1499 gram 18 7.1 1000 - 1499 gram 12 4.8 2000 - 2499 gram 12 4.8 2000 - 2499 gram 12 4.8 2000 - 2499 gram 12 4.8	-		
Quadruplet or higher - - Sex 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation 252 100.0 Total 252 100.0 Obstetric estimate of gestation 252 100.0 Total 252 100.0 20-23 weeks 122 48.4 24-37 weeks 25 9.9 22-33 weeks 21 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 1 0.4 Manom 2 8.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 252 100.0 5000 - 0999 grams 59 23.4 1000 - 1499 gram 18 7.1 5000 - 2999 gram 12 4.8 2000 - 2499 gram 12 4.8 2000 - 2499 gram 12 4.8 <td></td> <td>-</td> <td>J.2 _</td>		-	J.2 _
Sex Total 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation 122 48.4 Total 252 100.0 20-23 weeks 34 13.5 24-27 weeks 34 13.5 25-31 weeks 25 9.9 32-33 weeks 22 8.7 32-33 weeks 22 8.7 34-36 weeks 20 7.9 34-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.0 500 grams 59 23.4 1000 - 1499 gram 18 7.1 1000 - 1499 gram 18 7.1 1000 - 1499 gram 12 4.8 2000 - 2499 gram 12 4.8 2000 - 2499 gram 12 4.8 2000 - 1499 gram 16 6.3 2000 - 24	-	_	_
Total 252 100.0 Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation 252 100.0 Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 43-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 3 1.2 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 7500 grams 103 40.9 0500 - 1999 gram 16 6.3 2000 - 2499 gram 12 4.8 2100 - 1499 gram 16 6.3 2000 - 2499 gram 12 4.8 2100 - 2499 gram 12 4.8 2500 - 2999 gram 12 4.8			
Male 128 50.8 Female 124 49.2 Obstetric estimate of gestation 252 100.0 C0-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 3 1.2 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 7 7.1 Total 252 100.0 <500 grams		252	100 0
Female 124 49.2 Obstetric estimate of gestation 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 21 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 3 1.2 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 C500 grams 59 23.4 1000 - 1499 gram 18 7.1 1500 - 1999 gram 16 6.3 200 - 2499 gram 12 4.8 200 - 2499 gram 17 6.7 3000 - 3499 gram 16 6.3 200 - 2499 gram 10 4.0 200 - 3499 gram 10 4.0 200 - 3499 gram 3 1.2 300 - 3499			
Obstatric estimate of gestation 252 100.0 Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 21 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 10.3 Birthweight 252 100.0 500 - 0999 grams 59 23.4 1000 - 1499 gram 18 7.1 1500 - 1999 gram 16 6.3 2000 - 2499 gram 17 6.7 3000 - 3499 gram 17 6.7 3000 - 3499 gram 9 3.6 3500 - 3999 gram 10 4.0 4000 - 4499 gram 3 1.2 4500 - 4999 gram 3 1.2 <td< td=""><td></td><td></td><td></td></td<>			
Total 252 100.0 20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 21 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 103 40.9 0500 - 0999 grams 59 23.4 1000 - 1499 gram 16 6.3 2000 - 2499 gram 12 4.8 2500 - 2999 gram 12 4.8 2500 - 2999 gram 17 6.7 3000 - 3499 gram 9 3.6 3500 - 3999 gram 10 4.0 400 - 4499 gram 3 1.2 4500 - 4999 gram 3 1.2 4500 - 8165 gram 1 0.4	remare	121	49.2
20-23 weeks 122 48.4 24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 77-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams	Obstetric estimate of gestation		
24-27 weeks 34 13.5 28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams	Total	252	100.0
28-31 weeks 25 9.9 32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 20 7.9 39-40 weeks 3 1.2 41 weeks 3 1.2 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 7500 grams 103 40.9 0500 - 0999 grams 13 7.1 1500 - 1999 gram 16 6.3 2000 - 2499 gram 12 4.8 2500 - 2999 gram 12 4.8 2500 - 2999 gram 12 4.8 2500 - 2999 gram 10 4.0 3000 - 3499 gram 9 3.6 3500 - 3999 gram 3 1.2 4500 - 4999 gram 3 1.2 4500 - 4999 gram 3 1.2 4500 - 4999 gram 3 1.2 4500 - 8165 gram 1 0.4	20-23 weeks	122	48.4
32-33 weeks 11 4.4 34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams			
34-36 weeks 22 8.7 37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams			
37-38 weeks 20 7.9 39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 2 0.8 Total 252 100.0 <500 grams			
39-40 weeks 12 4.8 41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight Total 252 100.0 <500 grams			
41 weeks 3 1.2 42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams			
42 weeks and over 1 0.4 Unknown 2 0.8 Birthweight 252 100.0 <500 grams			
Unknown 2 0.8 Birthweight 700 000 Total 252 100.0 <500 grams			
BirthweightTotal252100.0<500 grams			
Total252100.0<500 grams	UIKIIOWII	Σ	0.0
<500 grams			
0500 - 0999 grams5923.41000 - 1499 gram187.11500 - 1999 gram166.32000 - 2499 gram124.82500 - 2999 gram176.73000 - 3499 gram93.63500 - 3999 gram104.04000 - 4499 gram31.24500 - 4999 gram10.45000 - 8165 gram10.4			
1000 - 1499 gram187.11500 - 1999 gram166.32000 - 2499 gram124.82500 - 2999 gram176.73000 - 3499 gram93.63500 - 3999 gram104.04000 - 4499 gram31.24500 - 4999 gram10.45000 - 8165 gram10.4			
1500 - 1999 gram166.32000 - 2499 gram124.82500 - 2999 gram176.73000 - 3499 gram93.63500 - 3999 gram104.04000 - 4499 gram31.24500 - 4999 gram10.45000 - 8165 gram10.4			
2000 - 2499 gram124.82500 - 2999 gram176.73000 - 3499 gram93.63500 - 3999 gram104.04000 - 4499 gram31.24500 - 4999 gram10.45000 - 8165 gram10.4			
2500 - 2999 gram176.73000 - 3499 gram93.63500 - 3999 gram104.04000 - 4499 gram31.24500 - 4999 gram10.45000 - 8165 gram10.4			
3000 - 3499 gram 9 3.6 3500 - 3999 gram 10 4.0 4000 - 4499 gram 3 1.2 4500 - 4999 gram 1 0.4 5000 - 8165 gram 1 0.4	-		
3500 - 3999 gram 10 4.0 4000 - 4499 gram 3 1.2 4500 - 4999 gram 1 0.4 5000 - 8165 gram 1 0.4	-		
4000 - 4499 gram 3 1.2 4500 - 4999 gram 1 0.4 5000 - 8165 gram 1 0.4	-		
4500 - 4999 gram10.45000 - 8165 gram10.4	-		
5000 - 8165 gram 1 0.4	-		
-			
	-	3	

0.0 Quantity more than zero but less than 0.05.

- Quantity zero.

 $\label{eq:lincludes}$ fetal deaths occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death. \4Includes all persons of Hispanic origin of any race.

 $\5$ Includes General Educational Development (GED).

 $\ \$ Associate in Arts and Associate in Science.

\7Includes Bachelor in Arts and Bachelor in Science.

\8Includes Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\9Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

<code>\10Guam reported father's age for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.</code>

 $\11Guam$ reported mother received WIC food for herself durng this pregnancy for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

 $\12Guam$ reported to bacco use during pregnancy for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

 $\13Guam$ reported maternal pre-pregnancy body mass index for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

Table 6A. Number and percentage of fetal deaths by selected health charcteristics: United States, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Risk factor	Total	Number reported	Unknown	Per 1,000
Diabetes		10001000		-,
Prepregnancy (Diagnosis prior to this pregnancy)	20,202	871	1,084	45.6
Gestational (Diagnosis in this pregnancy)	20,202	959	1,084	50.2
Hypertension				
Prepregnancy (Chronic)	20,202	1,334	1,084	69.8
Gestational (PIH, preeclampsia)	20,202	1 , 598	1,084	83.6
Eclampsia	20,202	141	1,084	7.4
Pregnancy resulted from infertility treatment	20,202	567	1,084	29.7
Mother had a previous cesarean delivery\1	20,202	2,916	1,084	15.3
Maternal morbidity\2				
Ruptured uterus	20,202	145	1,217	763.8
Admissioin to intensive care unit	20,202	465	1,217	2,449.3

 $\1$ Mother had a previous cesarean delivery per 100 fetal deaths.

 $\2$ Maternal morbidities per 100,000 fetal deaths.

NOTE: PIH is pregnancy-induced hypertension.

Table 6B. Number and percentage of fetal deaths by selected health charcteristics: Guam and Puerto Rico, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Risk factor		Number reported	Unknown	Per 1,000
Diabetes				
Prepregnancy (Diagnosis prior to this pregnancy)	252	17	3	68.3
Gestational (Diagnosis in this pregnancy)	252	12	3	48.2
Hypertension				
Prepregnancy (Chronic)	252	14	3	56.2
Gestational (PIH, preeclampsia)	252	13	3	52.2
Eclampsia	252	3	3	12.0
Pregnancy resulted from infertility treatment	252	5	3	20.1
Mother had a previous cesarean delivery\1		31	3	12.4
Maternal morbidity\2				
Ruptured uterus	252	1	13	418.4
Admission to intensive care unit	252	-	13	_

- Quantity zero.

 $\1$ Mother had a previous cesarean delivery per 100 fetal deaths.

\2 Maternal morbidities per 100,000 fetal deaths.

NOTE: PIH is pregnancy-induced hypertension.

Table 7A. Deaths according to 124 selected causes of fetal death: 43 states and the District of Columbia, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.] Cause of death (based on ICD-10)	Number
All causes ¹	15,476
Certain infectious and parasitic diseases	5
Human immunodeficiency virus (HIV) disease	-
Other viral diseases	3
Other and unspecified infectious and parasitic diseasesA00-A49,A51-A79,B35-B99 Malignant neoplasms	- 0
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behaviorDO	0-D48 85
AnemiasD50-D64	5
Endocrine, nutritional and metabolic diseases	2
Cystic fibrosis	2
Other endocrine, nutritional and metabolic diseases	-
Meningitis	-
Other diseases of nervous system and sense organsG04-H93 Umbilical herniaK42	4
Other hernia	-
Other and unspecified diseases of digestive system	1
All other diseases, excluding perinatal conditions, congenital anomalies, and symptoms, signs and ill-defined conditions	L00-N98,U0 7 13,845
Fetus affected by maternal conditions that may be unrelated to present pregnancyP00	1,740
Fetus affected by maternal hypertensive disordersP00.0	908
Fetus affected by maternal renal and urinary tract diseases	19 267
Fetus affected by maternal infectious and parasitic diseasesP00.2 Fetus affected by other maternal circulatory and respiratory diseasesP00.3	69
Fetus affected by maternal nutritional disordersP00.4	2
Fetus affected by maternal injury	51
Fetus affected by surgical procedure on mother	7 417
Fetus affected by unspecified maternal conditionP00.9	/ ± ד -
Fetus affected by maternal complications of pregnancyP01	1,804
Fetus affected by incompetent cervixof membranesP01.0 Fetus affected by premature rupture of membranesP01.1	259 1 , 180
Fetus affected by oligohydramnios	65
Fetus affected by polyhydramniosP01.3	37
Fetus affected by ectopic pregnancy	2 206
Fetus affected by multiple pregnancy	19
Fetus affected by malpresentation before laborP01.7	6
Fetus affected by other and unspecified maternal complications of pregnancyP01.8-P01	
Fetus affected by complications of placenta, cord and membranesP02 Fetus affected by placenta previa	3,743 35
Fetus affected by other forms of placental separation and hemorrhageP02.1	1,351
Fetus affected by other and unspecified morphological and functional abnormalities of placenta	
Fetus affected by placental transfusion syndromesP02.3 Fetus affected by prolapsed cord	112
Fetus affected by other compression of umbilical cordP02.5	838
Fetus affected by other and unspecified conditions of umbilical cordP02.6	244
Fetus affected by chorioamnionitis Fetus affected by other and unspecified abnormalities of membranesP02.8-P02.9	389
Fetus affected by other complications of labor and delivery	136
Fetus affected by breech delivery and extractionP03.0	6
Fetus affected by other malpresentation, malposition and disproportion during labor and delivery Fetus affected by forceps delivery	203.1 9
Fetus affected by delivery by vacuum extractor (ventouse)P03.3	-
Fetus affected by cesarean delivery	4
Fetus affected by precipitate delivery	- 5
Fetus affected by other and unspecified complications of labor and deliveryP03.8-P0	3.9 112
Fetus affected by noxious influences transmitted via placentaP04	141
Slow fetal growth and fetal malnutrition	P07 242
Extremely low birth weight or extreme immaturity	
Other low birth weight and pretermP07.1,P07.3	99
Disorders related to long gestation and high birth weight	2
Exceptionally large size and other heavy for gestational age fetusP08.0-P08.1 Post-term, not heavy for gestational age fetus	2
Birth traumaP10-P15	1
Intracranial laceration and hemorrhage due to birth injury and other injuries to central nervous system	- 11
Other birth traumaP12-P15 Intrauterine hypoxia and birth asphyxia	-
Intrauterine hypoxia first noted before onset of labor	-
Intrauterine hypoxia first noted during labor and deliveryP20.1	-
Intrauterine hypoxia, unspecified Birth asphyxiaP21	
Birth aspnyxiaP21 Other respiratory conditions originating in the perinatal periodP22.8-P22.9,P23-P28	
Congenital pneumoniaP23	
Aspiration syndromes	
Interstitial emphysema and related conditions originating in the perinatal period	
Other respiratory system disorders	
Infections specific to the perinatal periodP35-P39	6
Congenital rubella syndrome infection	-
Congenital herpesviral (herpes simplex) infection	-
Congenital viral hepatitisP35.3	
Bacterial sepsis	
Congenital tuberculosis Congenital toxoplasmosis	
Other infections specific to the perinatal periodP35.8-P35.9,P37.2-P37.9,P38-P39	
Fetal hemorrhageP50-P54	13
Hemolytic disease of fetus	
ABO isoimmunization of fetus	
Other hemolytic disease of fetus	
Hydrops fetalis due to hemolytic disease	
Perinatal jaundice	
Perinatal jaundice	
Hematological disorders Transitory endocrine and metabolic disorders specific to fetus	2 451
Hematological disorders Transitory endocrine and metabolic disorders specific to fetusP70-P74 Syndrome of infant of a diabetic mother and neonatal diabetes mellitusP70.0-P70.2	-
Hematological disorders Transitory endocrine and metabolic disorders specific to fetus	-
Hematological disordersP60-P61 Transitory endocrine and metabolic disorders specific to fetusP70-P74 Syndrome of infant of a diabetic mother and neonatal diabetes mellitusP70.0-P70.2 Other transitory endocrine and metabolic disorders specific to fetus	2 451 451 - 1 5,345

Fetal death of unspecified cause	5,142
Withdrawal symptoms from maternal use of drugs of addiction	-
Termination of pregnancy	-
Complications of intrauterine procedures, not elsewhere classified	-
All other specified conditions originating in the perinatal period	16
Condition originating in the perinatal period, unspecifiedP96.9	_
Congenital malformations, deformations and chromosomal abnormalities	1,502
Congenital malformations of nervous system	276
Anencephaly and similar malformations	128
EncephaloceleQ01	10
MicrocephalyQ02	_
Congenital hydrocephalusQ03	52
Reduction deformities of brain	34
Other congenital malformations of brain	14
Spina bifida	16
· Other congenital malformations of spinal cord and nervous systemQ06-Q07	22
Congenital malformations of eye, ear, face and neckQ10-Q18	3
Congenital malformations of heartQ20-Q24	201
Other congenital malformations of circulatory systemQ25-Q28	47
Congenital malformations of lung	8
Other congenital malformations of respiratory system	10
Congenital malformations of digestive system	24
Congenital malformations of genital organs	8
Congenital malformations of urinary system	88
Renal agenesis and other reduction defects of kidney	44
Cystic kidney disease	18
Other congenital malformations of urinary systemQ62-Q64	26
Congenital malformations and deformations of musculoskeletal system, limbs and integument	162
Other congenital malformations	92
Conjoined twins	-
Multiple congenital malformations, not elsewhere classified	33
All other congenital malformations	59
Chromosomal abnormalities, not elsewhere classified	583
Down's syndrome	112
Edward's syndrome	218
Patau's syndrome	57
Other chromosomal abnormalities, not elsewhere classified	196
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classifiedR00-R9	7
External causes of mortality*U01,V01-Y84	11
Accidents (unintentional injuries)	11
Assault (homicide)*U01,X85-Y09	_
Complications of medical and surgical care	_
Other external causesY10-Y36	-
- Quantity zero	

- Quantity zero.

NOTE: Excludes data for jurisdictions (Arizona, Georgia, Hawaii, Mississippi, New York City, North Dakota, and Vermont) for which the cause of death was unspecified (P95) for 50% or more of records. Also excludes data for California which did not report cause of death based on the 2003 revision of the U.S. Standard Report of Fetal Death. Source: National Center for Health Statistics, National Vital Statistics System.

Table 7B. Deaths according to 124 selected causes of fetal death: Puerto Rico, 2022

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

All causes ¹		218
Certain infectious and parasitic diseases		
Congenital syphilis Human immunodeficiency virus (HIV) disease		-
Other viral diseases		
Other and unspecified infectious and parasitic diseasesA00	0-A49,A51-A79,B35-B99	
Malignant neoplasms		
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unkr Anemias	nown behaviorD00-D48	-
Endocrine, nutritional and metabolic diseases		
Short stature, not elsewhere classified	E34.3	
Cystic fibrosis		
Other endocrine, nutritional and metabolic diseases	E00-E32,E34.0-E34.2,E34.4-E34.9,E40-E83,E85-E88	-
Other diseases of nervous system and sense organs		
Jmbilical hernia	K42	
Other hernia		
Other and unspecified diseases of digestive system	K00-K38,K50-K92 ies, and symptoms, signs and ill-defined conditions	-
Certain conditions originating in the perinatal period		18
Fetus affected by maternal conditions that may be unrelated to pre	esent pregnancyP00	32
Fetus affected by maternal hypertensive disorders		14
Fetus affected by maternal renal and urinary tract diseases Fetus affected by maternal infectious and parasitic diseases		4
Fetus affected by maternal infectious and parasitic diseases.		
Fetus affected by maternal nutritional disorders		
Fetus affected by maternal injury		
Fetus affected by surgical procedure on mother Fetus affected by other medical procedures and maternal condit		-
Fetus affected by unspecified maternal condition		ـ ــ ,
Fetus affected by maternal complications of pregnancy	P01	52
Fetus affected by incompetent cervix		23
Fetus affected by premature rupture of membranes Fetus affected by oligohydramnios		23
Fetus affected by polyhydramnios		
Fetus affected by ectopic pregnancy	P01.4	
Fetus affected by multiple pregnancy		
Fetus affected by maternal death		-
	s of pregnancyP01.8-P01.9	
Fetus affected by complications of placenta, cord and membranes		55
Fetus affected by placenta previa		4
Fetus affected by other forms of placental separation and hemo Fetus affected by other and unspecified morphological and func	orrnagePUZ.1 ctional abnormalities of placentaP02.2	1:
Fetus affected by placental transfusion syndromes		
Fetus affected by prolapsed cord		
Fetus affected by other compression of umbilical cord Fetus affected by other and unspecified conditions of umbilica		11
Fetus affected by chorioamnionitis		
Fetus affected by other and unspecified abnormalities of membr	ranesP02.8-P02.9	
Fetus affected by other complications of labor and delivery		4
Fetus affected by breech delivery and extraction	P03.0 roportion during labor and deliveryP03.1	-
Fetus affected by forceps delivery		
Fetus affected by delivery by vacuum extractor (ventouse) \ldots .		
Fetus affected by cesarean delivery		
Fetus affected by precipitate delivery Fetus affected by abnormal uterine contractions		-
	r and delivery	
Fetus affected by noxious influences transmitted via placenta		2
Slow fetal growth and fetal malnutrition	P05 ere classifiedP07	1
Extremely low birth weight or extreme immaturity		± .
Other low birth weight and preterm		11
Disorders related to long gestation and high birth weight		
Exceptionally large size and other heavy for gestational age fetus Post-term, not heavy for gestational age fetus		-
Birth trauma		,
	her injuries to central nervous systemP10-P11	
Other birth trauma Intrauterine hypoxia and birth asphyxia		-
Intrauterine hypoxia and birth asphyxia Intrauterine hypoxia first noted before onset of labor		4
Intrauterine hypoxia first noted during labor and delivery	P20.1	
Intrauterine hypoxia, unspecified		
Birth asphyxiaBirth asphyxia		-
Congenital pneumonia		
Aspiration syndromes		
	perinatal period	
Atelectasis		-
infections specific to the perinatal period	P35-P39	,
Congenital rubella syndrome		
Congenital cytomegalovirus infection		
Congenital viral hepatitis		-
Bacterial sepsis	P36	
Congenital tuberculosis		
Congenital toxoplasmosis		
Other infections specific to the perinatal periodP35.8-P35.9,		
Other infections specific to the perinatal periodP35.8-P35.9, Fetal hemorrhage	P50-P54	
Other infections specific to the perinatal periodP35.8-P35.9, Fetal hemorrhage Hemolytic disease of fetus Rh isoimmunization of fetus	P50-P54 P55-P56 P55.0	
Other infections specific to the perinatal periodP35.8-P35.9, Fetal hemorrhage Hemolytic disease of fetus Rh isoimmunization of fetus ABO isoimmunization of fetus	P50-P54 P55-P56 P55.0 P55.1	-
Other infections specific to the perinatal periodP35.8-P35.9, Fetal hemorrhage Hemolytic disease of fetus Rh isoimmunization of fetus ABO isoimmunization of fetus Other hemolytic disease of fetus	P50-P54 P55-P56 P55.0 P55.1 P55.8-P55.9	-
Other infections specific to the perinatal periodP35.8-P35.9, Fetal hemorrhage Hemolytic disease of fetus Rh isoimmunization of fetus ABO isoimmunization of fetus.	P50-P54 P55-P56 P55.0 P55.1 P55.8-P55.9 P56	

Transitory endocrine and metabolic disorders specific to fetus	0
Syndrome of infant of a diabetic mother and neonatal diabetes mellitusP70.0-P70.2	0
Other transitory endocrine and metabolic disorders specific to fetus	0
	-
Digestive system disorders of fetus	-
Other conditions originating in the perinatal periodP29,P80-P96	18
Hydrops fetalis not due to hemolytic diseaseP83.2	1
Fetal death of unspecified causeP95	15
Withdrawal symptoms from maternal use of drugs of addictionP96.1	-
Termination of pregnancyP96.4	-
Complications of intrauterine procedures, not elsewhere classifiedP96.5	-
All other specified conditions originating in the perinatal period	2
Condition originating in the perinatal period, unspecifiedP96.9	-
Congenital malformations, deformations and chromosomal abnormalitiesQ00-Q99	28
Congenital malformations of nervous systemsystem	10
Anencephaly and similar malformations	1
EncephaloceleQ01	-
MicrocephalyQ02	-
Congenital hydrocephalusQ03	1
Reduction deformities of brain	4
Other congenital malformations of brain	_
Spina bifida	3
Other congenital malformations of spinal cord and nervous systemQ06-Q07	1
Congenital malformations of eye, ear, face and neckQ10-Q18	_
Congenital malformations of heart	1
Other congenital malformations of circulatory system	_
Congenital malformations of lung	_
Other congenital malformations of respiratory system	_
Congenital malformations of digestive systemQ35-Q45	1
Congenital malformations of genital organs	_
Congenital malformations of urinary system	4
Renal agenesis and other reduction defects of kidney	_
Cystic kidney disease	4
Other congenital malformations of urinary system	_
Congenital malformations and deformations of musculoskeletal system, limbs and integument	4
Other congenital malformations	2
Conjoined twins	_
Multiple congenital malformations, not elsewhere classified	1
All other congenital malformations	1
Chromosomal abnormalities, not elsewhere classified	1 6
Down's syndrome	2
Edward's syndrome	2
	5
Patau's syndrome	-
Other chromosomal abnormalities, not elsewhere classified	Ţ
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	-
External causes of mortality*U01,V01-Y84	-
Accidents (unintentional injuries)	-
Assault (homicide)	-
Complications of medical and surgical care	-
Other external causes	

- Quantity zero.

NOTE: Excludes data for Guam, for which the cause of death was unspecified (P95) for 50% or more of records. American Samoa did not report data for 2022 and data for the Northern Marianas and the Virgin Islands were excluded due to small numbers.

LOCAL FILE NO.	US STANDA	ARD R	REPORT OF	FET	AL DEAT	Ή			STATE FIL	LE NUMBER:	
MOTHER	1. NAME OF FETUS (optional-at the discretion of the pare				ME OF DELIV	ERY 3.	SEX	(M/F/Unk)	4. DATE	OF DELIV ERY (M	lo/Day/Yr)
	5a. CITY, TOWN, OR LOCATION OF DELIVERY 7	PLAC	E WHERE DELIN	/FRY OC	,	lhr) leck one)	8		NAME (If no	ot institution, give str	eet and
		□ Hospita						number)	ti un∠ (n no	st montation, give ou	
			anding birthing ce	enter							
			Delivery: Planned		er at home? □	Yes 🗆 N					
	8. COUNT OF DELIVERT		Doctor's office				c ĉ). Facility II	D. (NPI)		
			(Specify)								
	10a. MOTHER'S CURRENT LEGAL NAME (First, Middle,						-	10b DATE	OF BIRTH	(Mo/Day/Yr)	
		Luot, Ou						IOD. DATE	OF BIRT	(Mo/Day/11)	
	10c. MOTHER'S NAME PRIOR TO FIRST MARRIAGE (Fi	irst, Middl	le, Last, Suffix)					10d. BIRTH	IPLACE (S	tate, Territory, or Fo	reign Country)
	11a. RESIDENCE OF MOTHER-STATE 11b. Co	OUNTY					11c.	CITY, TOWN	I, OR LOC	ATION	
	11d. STREET AND NUMBER				11e. APT. N	0.	11f.	ZIP CODE		11g. INSIDE C	ITY LIMITS?
										□ Yes	□ No
FATHER	12a. FATHER'S CURRENT LEGAL NAME (First, Middle,	Last, Suff	fix) 12	b. DAT	E OF BIRTH (M	/lo/Day/Yr)		12c. BIRTHF	LACE (Sta	te, Territory, or Fore	ign Country)
DISPOSITION	13. METHOD OF DISPOSITION:		Deseties	D	l fra en Obata	- 0	. (0				
ATTENDANT	Burial Cremation Hospital Disposition ATTENDANT'S NAME, TITLE, AND NPI		. NAME AND TIT	LE OF F	PERSON	D Other			LETED 1	7. DATE RECEIV	ED BY
AND	NAME:		COMPLETING	REPOR	RΤ.					REGISTRAR	
REGISTRATION	NPI:	Na	ame				/	///	YY	/	/
INFORMATION		ті	tle			IVI	IVI	זי עע	ŤŤ		TTTT
	OTHER (Specify)										
	18. CAUSE/C			ΟΝΤΙ		G TO	FF		ΔΤΗ		
CAUSE	18a. INITIATING CAUSE/CONDITION				8b. OTHER S					IS	
OF	(AMONG THE CHOICES BELOW, PLEASE SELECT TH LIKELY BEGAN THE SEQUENCE OF EVENTS RESULT THE FETUS)	HE <u>ONE</u> TING IN	WHICH MOST THE DEATH OF		SELECT OR S N ITEM 18b)	PECIFY A	ALL C	OTHER COND	ITIONS CO	ONTRIBUTING TO	DEATH
FETAL DEATH	Maternal Conditions/Diseases (Specify)			N	laternal Condit	ions/Disea	ases	(Specify)			
DEATH	Complications of Placenta, Cord, or Membranes				complications of	f Placenta	Cor	rd or Membra	nes		
1	Rupture of membranes prior to onset of la	abor		Ũ				mbranes prior		labor	
	Abruptio placenta				Abruptio placenta						
	 Placental insufficiency 				 Placental insufficiency 						
	 Prolapsed cord 				 Prolapsed cord 						
	Chorioamnionitis				Chorioamnionitis						
i i	Other Specify)			Other Specify)							
	Other Obstetrical or Pregnancy Complications (Specify)			c	other Obstetrica	al or Pregr	nancy	Complication	s (Specify)	·	
Mother's Name Mother's Medical Record No.	Fetal Anomaly (Specify)			 _ F	etal Anomaly (Specify) _					
dica	Fetal Injury (Specify)			- - F	- etal Injury (Sp	ecify)					
Na Na	Fetal Infection (Specify)			F	etal Infection	(Specify)					
Mother's Name Mother's Medic	Other Fetal Conditions/Disorders (Specify)			_ 0	Other Fetal Co	nditions/Di	sorde	ers (Specify) _			
Mot Mot											
	Unknown Unknown Sc. WEIGHT OF FETUS (grams preferred, specify unit)		18e. ESTIMATE		Unknown OF FETAL DE	ATH				PSY PERFORMED)?
			Dead at time	of first a	ssessment, no	labor ong	oing			No De Planned	
	□ grams □ lb/oz		Dead at time	of first a	ssessment, lat	oor ongoin	g	18g. WAS EXAI	A HISTOL	OGICAL PLACEN PERFORMED?	IAL
			Died during la	abor, afte	or, after first assessment			Yes No Planned Ish. WERE AUTOPSY OR HISTOLOGICAL			
	18d. OBSTETRIC ESTIMATE OF GESTATION AT DELIV		 Unknown tim 	e of fetal	death			PLAC IN DI	CENTAL EX	XAMINATION RES	ULTS USED

REV. 11/2003

MOTHER	 19. MOTHER'S EDUCATION (Check the box that best describes the highest degree or level of school completed at the time of delivery) 8th grade or less 9th - 12th grade, no diploma High school graduate or GED completed 	the box that best describes wh	MOTHER'S RACE (Check one or more races to indicate what the mother considers herself to be) White Black or African American American Indian or Alaska Native (Name of the enrolled or principal tribe)Asian Indian Chinese Filipino Japanese				
	 Some college credit but no degree Associate degree (e.g., AA, AS) Bachelor's degree (e.g., BA, AB, BS) Master's degree (e.g., MA, MS, MEng, MEd, MSW, MBA) Doctorate (e.g., PhD, EdD) or Professional degree (e.g., MD, DDS, DVM, LLB, JD) MOTHER MARRIED? 	, 		Samoan Other Pacific Islander (Specify) Other (Specify)			
		M M D D YYYY	enatal Care CARE VI 	D D YYYY IVERY 28. DID MOTHER DURING THI	VISITS FOR THIS PREGNANCY (If none, enter "0".) R GET WIC FOOD FOR HERSELF IS PREGNANCY? □ Yes □ No		
	LIVE BIRTHS 29a. Now Living 29b. Now Dead 30a	NUMBER OF OTHER PREGNANCY OUTCOMES (spontaneous or induced losses or ectopic pregnancies) Da. Other Outcomes	 CIGARETTE SMOKING For each time period, en cigarettes smoked. IF N Average number of cigarett 	nter either the number of ci NONE, ENTER "0". tes or packs of cigarettes s	igarettes or the number of packs of		
	None None	□ None 30b. DATE OF LAST OTHER	Three Months Before Prec First Three Months of Pre Second Three Months of F Third Trimester of Pregnar 32. DATE LAST NORMAL	gnancy gnancy Pregnancy ncy 33. PLURALITY - Single	OR OR OR OR e, 34. IF NOT SINGLE BIRTH-		
	MM YYYY 35. MOTHER TRANSFERRED FOR MAT		MENSES BEGAN	Twin, Triplet, etc. (Specify) Yes □ No	Born First, Second, Third, etc. (Specify)		
MEDICAL AND HEALTH INFORMATION	36. RISK FACTORS IN THIS PREGNANCY Diabetes Prepregnancy (Diagnosis prior to th Gestational (Diagnosis in this preg Hypertension Prepregnancy (Chronic) Gestational (PIH, preeclampsia) Eclampsia Previous preterm birth Other previous poor pregnancy outcome growth restricted birth) Pregnancy resulted from infertility treatr Fertility-enhancing drugs, Artificial ir Intrauterine insemination Assisted reproductive technology (e. Mother had a previous cesarean deliver If yes, how many None of the above	his pregnancy) egnancy) ne (Includes perinatal death, small-for-ges tment-If yes, check all that apply: insemination or e.g., in vitro fertilization (IVF), gamete intra ery	tational age/intrauterine afallopian transfer (GIFT))	 THIS PREGNANCY ((Gonorrhea Syphilis Chlamydia Listeria Group B Streptococci Cytomegalovirus Parvovirus Toxoplasmosis None of the above Other (Specify) 	205		
	 38. METHOD OF DELIVERY A. Was delivery with forceps attempted but Yes No B. Was delivery with vacuum extraction att unsuccessful? Yes No C. Fetal presentation at delivery Cephalic Breech Other D. Final route and method of delivery (Che Vaginal/Spontaneous Vaginal/Spontaneous Vaginal/Vacuum Cesarean If cesarean, was a trial of labor attem Yes No E. Hysterotomy/Hysterectomy 	eck one) (Complications associations descent of the above associations of the above associations of the above association of the above association of the above association as	gree perineal laceration ectomy nsive care unit ting room procedure following	(Check Anencephaly Meningomyelocele Cyanotic congenita Congenital diaphra Omphalocele Gastroschisis Limb reduction def amputation and dw Cleft Lip with or wi Cleft Palate alone	al heart disease agmatic hernia fect (excluding congenital warfing syndromes) ithout Cleft Palate nfirmed nding iosomal disorder nfirmed nding		

REV. 11/2003

NOTE: This recommended standard fetal death report is the result of an extensive evaluation process. Information on the process and resulting recommendations as well as plans for future activities is available on the Internet at: http://www.cdc.gov/nchs/vital_certs_rev.htm.

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2022

States, 2022		1	1			1	1	r
	All				20 weeks	20 weeks	20 weeks	
Area	periods of	12 weeks	16 weeks	20 weeks	or 350	or 400	or 500	350 grams
Alea	gestation	IZ WEEKS	IO WEEKS	20 WEEKS	grams	grams	grams	550 grains
	gestation				grams	grams	grams	
Alabama				Х				
Alaska				Х				
Arizona					Х			
Arkansas		Х						
California		Δ		37				
	1			Х				
Colorado	X ¹							
Connecticut				Х				
Delaware								x ²
District of Columbia							Х	
Florida				Х				
Georgia	Х							
Hawaii	X							
	^							
Idaho					Х			
Illinois				Х				
Indiana	ļ			Х				
Iowa	ļ			Х				
Kansas				Х				
Kentucky					Х			
Louisiana					Х			
Maine				Х				
Maryland				X ³				
				Λ				
Massachusetts					Х			
Michigan						Х		
Minnesota				Х				
Mississippi					Х			
Missouri					Х			
Montana								X ²
Nebraska				Х				
Nevada				X				
New Hampshire				21	Х			
				Х	А			
New Jersey								
New Mexico				X ³				
New York	Х							
New York (excluding New York City)	Х							
New York City	Х							
North Carolina				Х				
North Dakota				Х				
Ohio				Х				
	1	X ⁴						
Oklahoma		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~						
Oregon				Х				
Pennsylvania	l		Х	ļ	ļ			
Rhode Island	Х							
South Carolina					Х			
South Dakota				Х				
Tennessee					Х			
Texas				Х				
Utah				X				
Vermont				x ⁵				
	17			^				
Virginia	Х							
Washington				X				
West Virginia				Х				
Wisconsin					Х			
Wyoming				Х				
American Samoa	Х							
Guam	1				Х			
	1							X ²
Northern Marianas	+				X			Δ
Puerto Rico				ļ	A			
Virgin Islands	Х							
1 Although Ctote low we wince the work								of 20 modes

1 Although State law requires the reporting of fetal deaths of all periods of gestation, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

2 If weight is unknown, 20 completed weeks of gestation or more.

3 If gestational age is unknown, weight of 350 grams or more.

4 Although State law requires the reporting of fetal deaths at 12 weeks of gestation or more, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

5 If gestational age is unknown, weight of 400 grams or more, 15 ounces or more.

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2022

[By place of residence]

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Reporting areas	Total Fetal deaths	Time of birth	Mother's birthplace	Education of mother ²	Father's age ³	Mother's Hispanic origin
Total of reporting areas ¹	20,202	1.7	5.6	14.2	26.7	4.5
Alabama	426	-	-	5.4	64.1	-
Alaska	54	3.7	3.7	14.8	35.2	7.4
Arizona	492	1.8	0.6	4.1	13.8	1.8
Arkansas	265	1.5	3.0	11.3	50.9	1.1
California	2,115	1.3	5.0	13.0	12.0	6.0
Colorado	, 305	-	16.7	34.1	13.1	1.0
Connecticut	175	4.0	1.7	26.3	9.1	4.6
Delaware	56	_	1.8	14.3	32.1	1.8
District of Columbia	65	1.5	46.2	40.0	60.0	1.5
Florida	1,548	2.8	9.0	8.1	16.5	6.5
Georgia	1,000	1.1	0.2	23.1	59.1	1.7
Hawaii	97	25.8	38.1	55.7	56.7	47.4
Idaho	91	1.1	- 50.1	2.2	13.2	1.1
Illinois	737	2.6	8.8	19.9	16.7	0.3
Indiana	439	2.0	3.4	7.3	27.8	1.4
Iowa	163	-	-	17.8	7.4	-
Kansas	205	1.5	1.0	6.3	11.7	1.5
Kentucky	283	-	1.1	4.9	46.3	-
Louisiana	300	-	0.7	4.3	24.7	-
Maine	63	1.6	6.3	7.9	20.6	-
Maryland	366	1.9	12.6	20.8	19.9	
Massachusetts	284	0.7	20.1	39.8	34.2	
Michigan	614	0.2	3.3	6.4	18.4	2.3
Minnesota	351	0.9	0.9	3.4	10.5	2.0
Mississippi	320	-	1.6	7.2	42.5	-
Missouri	358	0.6	2.0	5.6	17.9	2.2
Montana	54	1.9	-	-	13.0	-
Nebraska	110	1.8	0.9	40.0	7.3	4.5
Nevada	235	-	2.1	6.8	11.1	-
New Hampshire	56	1.8	7.1	8.9	23.2	3.6
New Jersey	498	1.6	1.2	7.6	8.4	5.8
New Mexico	69	1.4	-	2.9	44.9	1.4
New York (excluding New York City)	577	8.7	1.9	30.7	58.1	18.0
New York City	634	3.8	11.4	26.5	45.7	10.4
North Carolina	696	1.1	0.4	1.9	42.0	-
North Dakota	47	_	2.1	2.1	17.0	-
Ohio	741	0.1	3.4	2.8	11.3	1.6
Oklahoma	293	0.7	5.5	6.1	19.1	5.1
Oregon	195	2.1	3.6	9.7	16.4	5.1
Pennsylvania	663	5.1	3.6	15.8	14.5	0.2
Rhode Island	52	3.8	-	63.5	75.0	9.6
South Carolina	288		0.3	7.3	24.7	2.4
South Dakota	54	_	1.9	7.4	3.7	1.9
		1 5	1.9		47.4	
Tennessee	538	1.5		19.7 24.8	47.4 24.6	4.1 7.3
Texas	1,613	0.1	12.3			1.3
Utah	252	-	2.0	9.5	8.3	-
Vermont	21	-	4.8	14.3	23.8	4.8
Virginia	497	3.2	1.6	18.3	56.1	5.6
Washington	421	2.4	13.3	9.7	12.8	5.5
West Virginia	70	1.4	-	-	8.6	-
Wisconsin	320	1.3	1.9	8.4	45.3	0.3
Wyoming	36	-	-	8.3	11.1	2.8

American Samoa						
Guam	34	-	2.9	2.9	73.5	-
Northern Marianas						
Puerto Rico	218	0.9	0.9	0.5	23.9	-
Virgin Islands						

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2022

[By place of residence]

Reporting areas	Place of delivery	Attendant at delivery	Month prenatal care began\4	Mother's height	Mother's pre- pregnancy weight\5	Did mother get WIC food for herself during this pregnancy\6	Live-birth order\7
Total of reporting areas ¹	0.2	1.0	12.7	6.3	12.2	16.0	5.5
Alabama	_	_	2.8	0.9	5.9	3.8	_
Alaska	1.9	_	9.3	7.4	14.8		7.4
Arizona	0.2	1.2	3.9	3.0	6.1		3.3
Arkansas	-		4.2	2.3	14.7		2.6
California	0.7	-	14.1	8.5	15.5		7.1
Colorado	_	2.6	46.9	22.6	34.1		21.3
Connecticut	_	1.1	7.4	2.9	6.9		2.3
Delaware	_	10.7	7.1	5.4	10.7		3.6
District of Columbia	_		18.5	6.2	32.3		-
Florida	-	0.1	14.5	9.5	14.1		7.0
Georgia	0.1	0.2	18.9	6.7	14.5		8.2
Hawaii	-	1.0	64.9	41.2	64.9		54.6
Idaho	_	-	3.3	3.3	3.3		5.5
Illinois	0.1	0.5	16.0	4.2	17.2		0.9
Indiana	-	11.8	10.0	4.2	9.6		8.0
Indiana Iowa	-	-	3.1	4.3	9.6 3.7		8.0
			3.1 4.4	2.4			
Kansas	-	-0.4	4.4	2.4	7.3 1.8		- 0.4
Kentucky							
Louisiana	-	-	3.7	0.7	6.0		0.3
Maine	-	-	7.9	3.2	12.7		-
Maryland	-	0.5	16.4	4.1	7.4		0.8
Massachusetts	-	-	3.9	2.8	14.8		-
Michigan	-	0.3	5.4	1.0	4.9		2.3
Minnesota	-	0.9	2.0	2.0	9.7		2.0
Mississippi	-	0.6	7.8	1.3	5.0		-
Missouri	0.3	-	6.7	2.0	6.1		1.7
Montana	-	-	1.9	1.9	3.7		-
Nebraska	0.9	0.9	10.0	10.0	11.8		5.5
Nevada	-	0.4	5.5	5.5	6.8		0.9
New Hampshire	-	-	3.6	-	1.8		-
New Jersey	-	-	11.6	2.8	8.4		1.8
New Mexico	-	-	4.3	1.4	14.5		-
New York (excluding New York City)	0.3	-	36.9	21.7	27.7	44.7	12.8
New York City	-	-	29.5	4.9	14.8		11.8
North Carolina	-	1.6	2.6	1.3	5.6	4.0	-
North Dakota	-	-	4.3	-	4.3	10.6	-
Ohio	-	0.5	4.0	3.0	6.3	6.2	0.8
Oklahoma	-	-	7.5	7.2	15.4	10.9	28.7
Oregon	-	-	4.6	5.6	6.2	8.7	6.2
Pennsylvania	-	-	15.1	9.2	16.6	25.9	6.2
Rhode Island	-	1.9	51.9	11.5	19.2	88.5	19.2
South Carolina	-	2.1	5.6	2.4	3.8		3.1
South Dakota	-	-	3.7	-	3.7		7.4
Tennessee	-	-	9.5	3.9	8.4		0.9
Texas	0.7	4.6	14.9	8.1	7.7		4.3
Utah	_	0.4	3.2	1.6	6.0		0.4
Vermont	_	-	14.3		4.8		-
Virginia	0.2	0.4	19.5	16.1	34.0		1.0
Washington		1.7	17.6	10.1	18.5		30.6
West Virginia	_	2.9	5.7	2.9	4.3		1.4
Wisconsin	_	2.9	4.1	1.6	5.3		0.3
Wyoming	-	-	4.1 19.4	±.0 _		2.8	
American Samoa							
Guam	-	-	29.4	44.1	47.1	50.0	8.8
Northern Marianas							
Puerto Rico	0.5	1.8	2.3	2.3	0.9	4.1	5.0
Virgin Islands							

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2022

[By place of residence]

				Method of	Delivery
Reporting areas	Birth interval\8	Cigarette smoking before and during pregnancy\9	Risk Factors in this Pregnancy	Fetal presentation\10	Final route and method of delivery\11
Total of reporting areas ¹	11.8	8.6	5.4	10.7	4.9
Alabama	3.8	1.9	0.9	1.9	0.9
Alaska	14.8	24.1	5.6	3.7	-
Arizona	4.9		3.5	5.7	1.6
Arkansas	9.8		0.4	0.8	0.4
California	13.9		10.8	34.1	10.2
Colorado	33.1		-	15.4	14.4
Connecticut	12.6		2.3	5.7	1.7
Delaware	8.9		3.6	1.8	3.6
District of Columbia	21.5		1.5	3.1	_
Florida	14.0		7.6	17.8	8.7
Georgia	13.9		12.8	15.9	3.3
Hawaii	56.7		17.5	53.6	52.6
Idaho	9.9		2.2	3.3	3.3
Illinois	5.2		0.1	8.7	6.1
Indiana	13.7		12.1	26.7	14.1
Iowa	3.1		-	1.2	1.2
Kansas	2.0		-	0.5	
Kentucky	3.2		2.8	4.6	1.4
Louisiana	5.3		2.0		
Maine	3.2		_	7.9	-
Maryland	4.1		0.3	0.5	_
Maryranu Massachusetts	6.7		0.5	0.7	1.8
Massachusetts Michigan	11.4		3.9	6.8	0.7
Minnesota	4.3		1.4	4.0	1.7
Minnesota Mississippi	4.3		5.3	4.0	0.3
Mississippi Missouri	8.1		5.5	5.9	1.1
Missouri Montana	11.1		_	1.9	1.9
Nebraska	14.5		3.6	1.9	1.9
Neblaska Nevada	5.5		5.0	0.9	0.9
Nevada New Hampshire	5.5		-	0.9	0.9
_	6.8		1.8	6.8	1.0
New Jersey New Mexico	2.9		1.0	0.0	1.0
	2.9		4.2	27.0	22.9
New York (excluding New York City) New York City	25.4		4.2	20.7	10.1
North Carolina	23.4		19.4	0.1	0.3
North Dakota	2.3		2.1	14.9	
Ohio	14.3		0.3	2.0	-1.8
Ohloma	14.5		7.5	2.0	1.0
	9.7			- 1 E	
Oregon			-	1.5	1.0
Pennsylvania	6.6		6.3	11.3	9.0
Rhode Island	23.1		9.6	11.5	7.7
South Carolina	10.8		5.9	2.4	1.4
South Dakota	7.4		1.9	1.9	1.9
Tennessee	10.8		0.4	0.2	- 1 ~
Texas	9.7 6.0		8.2	0.2	1.3
Utah			-	-	-
Vermont	9.5		4.8	4.8	-
Virginia	18.5		-	0.8	-
Washington	35.6		14.3	19.7	10.5
West Virginia	5.7		1.4	7.1	2.9
Wisconsin	1.6		0.9	4.7	0.6
Wyoming	30.6	13.9	-	-	-

American Samoa					
Guam	20.6	50.0	-	44.1	2.9
Northern Marianas					
Puerto Rico	9.2	0.5	1.4	2.8	1.4
Virgin Islands					

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2022 [By place of residence]

Reporting areas	Maternal morbidity	Weight of fetus	Obstetric estimate of gestation	Cause of death\12	
Total of reporting areas 1	6.0	7.4	0.7	37.2	
Alabama	0.7	4.7	-	33.3	
Alaska	1.9	13.0	_	35.2	
Arizona	3.3	1.8	0.8	50.6	
Arkansas	0.8	8.3	0.4	44.9	
California	19.0	5.9	0.9		
Colorado	-	24.6	-	41.6	
Connecticut	-	11.4	-	23.4	
Delaware	3.6	-	1.8	32.1	
District of Columbia	-	4.6	-	15.4	
Florida	9.1	13.5	3.9	17.8	
Georgia	7.9	11.8	0.4	67.6	
Hawaii	18.6	49.5	4.1	73.2	
Idaho	-	3.3	-	26.4	
Illinois	0.3	9.4	0.1	19.5	
Indiana	7.1	10.3	0.9	32.6	
Iowa	_	2.5	-	33.7	
Kansas	_	1.5	1.0	23.9	
Kentucky	1.4	2.1	-	47.0	
Louisiana	_	4.3	_	22.3	
Maine	_	14.3	_	20.6	
Maryland	1.1	3.6	_	32.8	
Massachusetts	0.4	5.6	_	29.2	
Michigan	20.4	2.0	0.2	51.5	
Minnesota	0.6	4.6	-	36.5	
Mississippi	5.0	2.2	0.9	57.2	
Missouri	-	4.2	1.1	32.1	
Montana	_	1.9		27.8	
Nebraska	0.9	10.0	0.9	22.7	
Nevada	0.4	2.6	-	42.6	
New Hampshire	-	-	_	21.4	
New Jersey	4.8	4.2	_	45.2	
New Mexico		2.9	_	26.1	
New York (excluding New York City)	4.5	21.3	0.5	48.5	
New York City	19.9	12.3		57.9	
North Carolina		5.6	0.9	49.3	
North Dakota	_	2.1	-	49.3 51.1	
Ohio	2.8	4.0	_	33.3	
Ohio Oklahoma	6.1	4.0	0.7	29.7	
	0	4.4	1.5	43.6	
Oregon	7.5	1.5	1.5	24.7	
Pennsylvania Rhode Island		15.4	-	40.4	
South Carolina	11.5 7.6		0.3		
		1.4		43.4	
South Dakota	-	7.4	-	31.5	
Tennessee	0.2	1.3	0.4	43.1	
Texas	4.0	0.8	0.4	25.5	
Utah	-	9.5	-	29.4	
Vermont	4.8	4.8	-	57.1	
Virginia	0.4	6.4	1.4	46.7	
Washington	-	14.7	-	25.4	
West Virginia	7.1	4.3	-	40.0	
Wisconsin	0.3	5.3	0.6	43.8	
Wyoming	-	2.8	-	33.3	

American Samoa				
Guam	5.9	-	2.9	82.4
Northern Marianas				
Puerto Rico	5.0	1.4	0.5	6.9
Virgin Islands				

0.0 Quantity more than zero but less than 0.05.

---Data not available.

- Quantity zero.

American Samoa

NOTE: American Samoa did not report data for 2022. Data for the Commonwealth of Northern Marianas and the Virgin Islands were excluded due to small numbers. Reporting areas with levels of unknown data for an item of 50% and higher (based on counts by residence) are flagged in the file as not reporting.

1\Excludes data for American Samoa, Guam, the Commonwealth of the Northern Marianas, Puerto Rico, and the Virgin Islands.

\2Hawaii and Rhode Island reported education of mother for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

\3Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New York, Rhode Island, Virginia, and Guam reported father's age for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

\4Hawaii and Rhode Island reported date of first prenatal care for 2022 but had unknown levels of 50% or higher and were flagged as not reporting.

\5Hawaii reported mother's pre-pregnancy weight for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

\6New York City does not report WIC and Hawaii, Indiana, Rhode Island, and Guam reported this item for 2022 but had unknown levels of 50% or higher and were flagged as not reporting

\7Hawaii reported live birth order for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

\8Hawaii reported birth interval for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

\9Rhode Island and Guam reported cigarette smoking before and during pregnancy for 2022 but had an unknown level of 50% or higher and were flagged as not reporting.

\10Hawaii reported fetal presentation for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

\11Hawaii reported final route and method of delivery for 2022 but had an unknown level of 50% or higher and was flagged as not reporting.

\12Arizona, Georgia, Hawaii, Mississippi, North Dakota, New York City, Vermont, and Guam reported cause of death for 2022 but had an unknown level of 50% or higher and were flagged as not reporting. California did not report cause of death based on the 2003 revision of the U.S. Standard Report of Fetal Death and was also flagged as not reporting.