

List of Chartbook Figures

Population

1. Total population, population 65 years and over and 75 years and over: United States, 1950–2050	20
2. Percent of population in 4 age groups: United States, 1950, 2000, and 2050	21
3. Percent of population in selected race and Hispanic origin groups by age: United States, 1980–2000	23
4. Poverty rates by age: United States, 1966–2002	24
5. Low income population by age, race, and Hispanic origin: United States, 2002	25

Health Insurance

6. Health insurance coverage among persons under 65 years of age: United States, 1984–2002	26
7. No health insurance coverage among persons under 65 years of age by selected characteristics: United States, 2002	27

Preventive Health Care

8. Early prenatal care by race and Hispanic origin of mother: United States, 1980–2002	28
9. Early prenatal care by detailed race and Hispanic origin of mother: United States, 2002	29
10. Influenza and pneumococcal vaccination among adults 65 years of age and over: United States, 1989–2002	30
11. Influenza and pneumococcal vaccination among adults 65 years of age and over by race and Hispanic origin: United States, 2000–2002	31

Health Risk Factors

12. Cigarette smoking among men, women, high school students, and mothers during pregnancy: United States, 1965–2003	32
13. Current cigarette smoking among high school students by sex, frequency, and grade level: United States, 2003	33
14. High school students not engaging in recommended amounts of physical activity (neither moderate nor vigorous) by grade and sex: United States, 2003	34
15. Adults not engaging in leisure-time physical activity by age and sex: United States, 1998–2002	34
16. Overweight and obesity by age: United States, 1960–2002	36
17. Obesity among adults 20–74 years of age by sex, race, and Hispanic origin: United States, 1999–2002	37

Limitation of Activity

18. Selected chronic health conditions causing limitation of activity among children by age: United States, 2001–02	39
19. Limitation of activity caused by 1 or more chronic health conditions among working-age adults by selected characteristics: United States, 2000–2002	40
20. Selected chronic health conditions causing limitation of activity among working-age adults by age: United States, 2000–2002	41
21. Limitation of activities of daily living among Medicare beneficiaries 65 years of age and over: United States, 1992–2002	43

Mortality

22.	Life expectancy at birth and at 65 years of age by sex: United States, 1901–2001	45
23.	Infant, neonatal, and postneonatal mortality rates: United States, 1950–2002	46
24.	Infant mortality rates by detailed race and Hispanic origin of mother: United States, 1999–2001	47
25.	Death rates for leading causes of death for all ages: United States, 1950–2002	49

Special Feature: Drugs

26.	Percent of persons reporting prescription drug use in the past month by age: United States, 1988–94 and 1999–2000	51
27.	Percent of physician office and hospital outpatient department visits with 5 or more drugs prescribed, ordered, or provided by age: United States, 1995–2002	53
28.	Percent of asthma visits with quick-relief and long-term control drugs prescribed, ordered, or provided: United States, 1995–2002	55
29.	Percent of asthma visits with selected asthma drugs prescribed, ordered, or provided: United States, 1995–2002	57
30.	Percent of adults 18 years of age and over reporting antidepressant drug use in the past month by sex and age: United States, 1988–94 and 1999–2000.	58
31.	Percent of adults 18 years of age and over reporting antidepressant drug use in the past month by race and ethnicity: United States, 1988–94 and 1999–2000.	59
32.	Selective serotonin reuptake inhibitor (SSRI) antidepressant drug visits among adults 18 years of age and over by sex: United States, 1995–2002	61
33.	Stimulant drug visits among children 5–17 years of age by sex: United States, 1994–2002	63
34.	Antidepressant drug visits among children 5–17 years of age by sex: United States, 1994–2002.	63
35.	Cholesterol-lowering statin drug visits among adults 45 years of age and over by sex and age: United States, 1995–2002	65
36.	Percent of nonsteroidal anti-inflammatory drug (NSAID) visits with selective COX-2 NSAIDs prescribed, ordered, or provided among adults 18 years of age and over by age: United States, 1999–2002	67