

Table 11. Death rates for 113 selected causes and Enterocolitis due to *Clostridium difficile*, by age: United States, 2007

[Rates per 100,000 population in specified group. Populations used for computing death rates are postcensal estimates based on the 2000 census, estimated as of July 1, 2007; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All causes	803.6	684.5	28.6	15.3	79.9	104.9	184.4	420.9	877.7	2,011.3	5,011.6	12,946.5
Salmonella infections (A01-A02)	0.0	*	*	*	*	*	*	*	*	*	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07-A09)	2.2	*	*	*	*	*	0.1	0.3	1.0	4.7	18.8	52.0
Tuberculosis (A16-A19)	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.2	1.6
Respiratory tuberculosis (A16)	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.9	1.3
Other tuberculosis (A17-A19)	0.0	*	*	*	*	*	*	0.0	0.1	0.1	0.2	*
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	0.0	*	*	*	*	*	*	*
Septicemia (A40-A41)	11.5	6.6	0.5	0.2	0.4	0.7	2.1	5.5	12.9	32.8	79.9	174.4
Syphilis (A50-A53)	0.0	*	*	*	*	*	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15-B19)	2.5	*	*	*	*	0.2	1.1	6.4	7.4	4.5	4.4	2.9
Human immunodeficiency virus (HIV) disease (B20-B24)	3.7	*	*	*	0.4	2.7	8.3	9.5	5.3	2.3	0.8	*
Malaria (B50-B54)	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)	1.9	3.6	0.4	0.1	0.2	0.3	0.5	1.2	2.8	5.7	11.2	20.0
Malignant neoplasms (C00-C97)	186.6	1.7	2.2	2.4	3.9	8.5	30.8	114.3	315.4	715.5	1,256.3	1,590.2
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	2.7	*	*	*	*	0.1	0.6	2.6	6.2	9.9	13.4	17.1
Malignant neoplasm of esophagus (C15)	4.5	*	*	*	*	0.1	0.6	3.3	10.3	19.3	25.9	25.0
Malignant neoplasm of stomach (C16)	3.8	*	*	*	*	0.3	1.1	2.6	5.7	13.2	25.0	35.4
Malignant neoplasms of colon, rectum and anus (C18-C21)	17.8	*	*	*	0.1	0.7	3.0	10.9	27.7	60.1	118.4	200.8
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	5.7	*	*	0.1	0.1	0.2	0.9	5.7	12.8	20.1	32.8	32.0
Malignant neoplasm of pancreas (C25)	11.3	*	*	*	*	0.1	1.2	6.4	19.9	44.8	79.2	94.6
Malignant neoplasm of larynx (C32)	1.2	*	*	*	*	*	0.1	1.0	2.8	5.1	6.7	6.0
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	52.6	*	*	*	0.1	0.3	4.3	28.4	95.4	248.8	371.3	299.8
Malignant melanoma of skin (C43)	2.8	*	*	*	0.1	0.4	1.1	2.7	5.3	9.0	15.6	19.9
Malignant neoplasm of breast (C50)	13.6	*	*	*	*	0.8	5.1	13.7	26.8	42.3	69.7	116.6
Malignant neoplasm of cervix uteri (C53)	1.3	*	*	*	*	0.5	1.5	2.1	2.7	3.2	3.9	4.7
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	2.5	*	*	*	*	0.1	0.4	1.3	4.8	10.4	15.0	20.1
Malignant neoplasm of ovary (C56)	4.8	*	*	*	0.1	0.2	0.8	3.5	9.2	18.7	30.3	37.6

Table 11. Death rates for 113 selected causes and Enterocolitis due to *Clostridium difficile*, by age: United States, 2007

[Rates per 100,000 population in specified group. Populations used for computing death rates are postcensal estimates based on the 2000 census, estimated as of July 1, 2007; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Malignant neoplasm of prostate (C61)	9.6	*	*	*	*	*	0.0	1.0	6.9	29.5	86.4	170.5
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.2	*	*	0.1	0.1	0.1	0.6	2.8	7.9	16.3	27.2	33.0
Malignant neoplasm of bladder (C67)	4.6	*	*	*	*	*	0.2	1.3	4.8	14.6	38.5	68.6
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	4.4	*	0.7	0.8	0.5	0.9	2.0	4.4	9.1	15.5	20.0	16.1
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	18.2	0.6	0.7	0.9	1.5	1.9	3.4	8.2	23.5	63.2	137.3	185.5
Hodgkin's disease (C81)	0.4	*	*	*	0.2	0.3	0.3	0.3	0.5	1.1	2.2	2.4
Non-Hodgkin's lymphoma (C82-C85)	6.8	*	*	0.1	0.3	0.5	1.2	3.2	8.9	23.1	52.7	72.1
Leukemia (C91-C95)	7.2	0.5	0.6	0.8	1.0	1.1	1.5	3.1	8.6	23.8	52.7	76.7
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.7	*	*	*	*	*	0.4	1.6	5.5	15.1	29.6	34.1
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	*	*	*	*	*	*	*	*	*	*	*
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	21.0	0.5	0.7	0.6	1.3	1.7	4.0	12.4	33.4	71.6	139.7	206.7
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.7	1.4	0.4	0.2	0.2	0.3	0.8	1.6	4.3	12.4	36.2	76.4
Anemias (D50-D64)	1.6	*	*	0.1	0.2	0.3	0.4	0.6	1.0	2.9	9.4	36.1
Diabetes mellitus (E10-E14)	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.6	78.1	162.7	276.2
Nutritional deficiencies (E40-E64)	0.9	*	*	*	*	*	0.1	0.2	0.6	1.7	6.2	24.4
Malnutrition (E40-E46)	0.9	*	*	*	*	*	0.1	0.2	0.6	1.6	5.8	22.4
Other nutritional deficiencies (E50-E64)	0.1	*	*	*	*	*	*	*	*	*	0.4	2.0
Meningitis (G00,G03)	0.2	1.9	*	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.7	0.7
Parkinson's disease (G20-G21)	6.7	*	*	*	*	*	*	0.1	1.2	11.9	71.9	143.5
Alzheimer's disease (G30)	24.7	*	*	*	*	*	*	0.2	2.2	20.6	176.7	849.1
Major cardiovascular diseases (I00-I78)	267.3	13.4	1.4	0.8	3.2	9.7	34.4	105.5	247.0	597.5	1,758.7	5,679.0
Diseases of heart (I00-I09,I11,I13,I20-I51)	204.3	10.0	1.1	0.6	2.6	7.9	27.4	85.3	200.3	462.9	1,315.0	4,267.7
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.1	*	*	*	*	0.1	0.2	0.4	1.0	2.7	8.1	17.9
Hypertensive heart disease (I11)	10.2	*	*	*	0.1	0.8	3.2	8.2	13.7	20.7	48.6	192.3
Hypertensive heart and renal disease (I13)	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.0	6.1	22.3
Ischemic heart diseases (I20-I25)	134.7	0.6	*	0.1	0.4	2.6	14.4	55.6	141.1	325.7	891.9	2,705.3
Acute myocardial infarction (I21-I22)	44.1	*	*	*	0.1	1.0	5.6	21.6	54.5	121.1	288.9	756.7
Other acute ischemic heart diseases (I24)	1.4	*	*	*	*	*	0.3	0.9	2.1	3.8	7.8	20.8
Other forms of chronic ischemic heart disease (I20,I25)	89.3	*	*	*	0.2	1.6	8.6	33.2	84.5	200.7	595.1	1,927.9
Atherosclerotic cardiovascular disease, so described (I25.0)	19.6	*	*	*	*	0.5	3.2	13.5	30.7	51.1	109.2	314.7

Table 11. Death rates for 113 selected causes and Enterocolitis due to *Clostridium difficile*, by age: United States, 2007

[Rates per 100,000 population in specified group. Populations used for computing death rates are postcensal estimates based on the 2000 census, estimated as of July 1, 2007; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	69.7	*	*	*	0.2	1.0	5.4	19.7	53.8	149.7	485.9	1,613.2
Other heart diseases (I26-I51)	57.3	9.3	1.0	0.5	2.1	4.4	9.5	20.7	43.5	111.8	360.4	1,329.8
Acute and subacute endocarditis (I33)	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.2	2.4	2.4
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.3	0.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.1	2.1
Heart failure (I50)	18.8	0.5	*	*	0.1	0.2	0.7	2.4	8.4	29.7	122.4	554.4
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	37.8	8.2	0.8	0.5	1.8	4.0	8.3	17.5	34.1	80.3	234.6	770.9
Essential hypertension and hypertensive renal disease (I10,I12,I15)	7.9	*	*	*	0.1	0.2	0.9	2.8	6.5	16.2	49.5	191.1
Cerebrovascular diseases (I60-I69)	45.1	3.1	0.3	0.2	0.5	1.2	4.9	14.6	32.1	93.0	322.3	1,015.5
Atherosclerosis (I70)	2.7	*	*	*	*	*	0.1	0.3	1.1	4.3	17.6	83.3
Other diseases of circulatory system (I71-I78)	7.3	*	*	*	0.2	0.3	1.1	2.5	7.0	21.0	54.3	121.4
Aortic aneurysm and dissection (I71)	4.3	*	*	*	0.1	0.2	0.8	1.7	4.5	13.5	33.4	59.5
Other diseases of arteries, arterioles and capillaries (I72-I78)	3.0	*	*	*	0.1	0.1	0.3	0.8	2.5	7.5	20.9	62.0
Other disorders of circulatory system (I80-I99)	1.4	1.2	*	*	0.1	0.3	0.7	1.3	1.8	2.9	6.8	17.4
Influenza and pneumonia (J09-J18) ³	17.5	5.2	0.7	0.3	0.4	0.8	1.8	4.4	9.6	28.7	114.1	463.2
Influenza (J09-J11) ³	0.1	*	*	0.1	*	*	*	*	0.1	0.2	0.6	2.5
Pneumonia (J12-J18)	17.3	4.9	0.5	0.2	0.4	0.8	1.8	4.3	9.5	28.5	113.5	460.7
Other acute lower respiratory infections (J20-J22,U04) ⁴	0.1	1.1	*	*	*	*	*	*	*	0.1	0.3	1.5
Acute bronchitis and bronchiolitis (J20-J21)	0.1	1.1	*	*	*	*	*	*	*	0.1	0.2	1.1
Other and unspecified acute lower respiratory infections (J22,U04) ^{4,5}	0.0	*	*	*	*	*	*	*	*	*	*	0.4
Chronic lower respiratory diseases (J40-J47)	42.4	1.0	0.3	0.3	0.4	0.6	1.8	9.5	39.1	148.1	368.9	596.1
Bronchitis, chronic and unspecified (J40-J42)	0.2	0.6	*	*	*	*	*	0.1	0.2	0.5	1.4	4.8
Emphysema (J43)	4.2	*	*	*	*	*	0.1	1.1	4.9	17.0	37.1	45.5
Asthma (J45-J46)	1.1	*	0.2	0.3	0.3	0.5	0.7	1.2	1.4	2.1	4.4	12.0
Other chronic lower respiratory diseases (J44,J47)	36.8	*	*	*	*	0.1	0.9	7.1	32.6	128.5	326.0	533.8
Pneumoconioses and chemical effects (J60-J66,J68)	0.3	*	*	*	*	*	*	0.0	0.2	0.9	2.9	4.9
Pneumonitis due to solids and liquids (J69)	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	8.9	39.8	153.3
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	9.5	7.5	0.6	0.2	0.4	0.5	1.1	3.1	9.3	28.7	73.2	138.9
Peptic ulcer (K25-K28)	1.0	*	*	*	*	0.0	0.2	0.7	1.2	2.5	6.4	17.3
Diseases of appendix (K35-K38)	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	1.5
Hernia (K40-K46)	0.6	0.8	*	*	*	*	0.1	0.2	0.5	1.3	3.6	11.3
Chronic liver disease and cirrhosis (K70,K73-K74)	9.7	*	*	*	0.1	0.9	6.0	18.7	24.5	26.7	28.4	19.8
Alcoholic liver disease (K70)	4.8	*	*	*	*	0.7	4.2	11.7	13.2	10.1	5.7	2.1
Other chronic liver disease and cirrhosis (K73-K74)	4.9	*	*	*	*	0.2	1.7	7.0	11.3	16.6	22.6	17.7
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.5	7.8	23.3

Table 11. Death rates for 113 selected causes and Enterocolitis due to *Clostridium difficile*, by age: United States, 2007

[Rates per 100,000 population in specified group. Populations used for computing death rates are postcensal estimates based on the 2000 census, estimated as of July 1, 2007; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15.4	3.4	0.1	0.1	0.2	0.6	1.7	5.1	13.6	40.1	113.0	290.6
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	0.9
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	1.0	*	*	*	*	*	0.1	0.3	0.6	2.0	7.1	22.3
Renal failure (N17-N19)	14.3	3.2	*	*	0.2	0.6	1.6	4.8	12.9	37.9	105.3	267.3
Other disorders of kidney (N25,N27)	0.0	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	3.5
Hyperplasia of prostate (N40)	0.2	*	*	*	*	*	*	*	*	0.2	1.1	5.1
Inflammatory diseases of female pelvic organs (N70-N76)	0.0	*	*	*	*	*	*	*	*	0.1	0.2	0.4
Pregnancy, childbirth and the puerperium (O00-O99)	0.3	*	0.4	0.8	0.5	0.1	*	*	*	*
Pregnancy with abortive outcome (O00-O07)	0.0	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.2	*	0.4	0.8	0.5	0.1	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)	4.8	339.8	0.4	0.1	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.5	135.9	3.3	0.9	0.9	1.0	1.2	1.6	2.1	1.8	3.1	4.8
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	11.1	85.0	1.4	0.3	1.4	2.2	3.3	5.0	6.8	12.9	44.8	251.4
All other diseases (Residual)	79.0	31.4	3.6	2.2	4.6	7.8	16.4	37.6	70.0	149.0	469.6	1,701.9
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	41.0	30.2	9.6	5.5	37.4	36.9	39.2	46.3	37.3	45.2	105.5	286.7
Transport accidents (V01-V99,Y85)	15.5	3.0	3.5	3.4	25.7	18.4	15.8	16.4	14.8	16.5	22.9	24.1
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)	14.6	2.9	3.3	3.2	24.9	17.5	14.8	14.9	13.3	15.2	21.8	23.2
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)	0.4	*	*	0.1	0.4	0.3	0.4	0.5	0.4	0.4	0.5	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.6	*	*	0.1	0.4	0.6	0.6	1.0	1.1	0.9	0.6	0.6
Nontransport accidents (W00-X59,Y86)	25.5	27.2	6.1	2.0	11.7	18.5	23.4	29.9	22.5	28.7	82.6	262.6
Falls (W00-W19)	7.5	0.6	0.2	0.1	0.5	0.8	1.4	3.0	5.3	13.4	50.3	166.7
Accidental discharge of firearms (W32-W34)	0.2	*	*	0.1	0.4	0.2	0.2	0.2	0.2	0.2	0.2	*
Accidental drowning and submersion (W65-W74)	1.1	1.3	2.8	0.6	1.5	0.9	1.0	1.1	1.0	1.0	1.4	1.6

Table 11. Death rates for 113 selected causes and Enterocolitis due to *Clostridium difficile*, by age: United States, 2007

[Rates per 100,000 population in specified group. Populations used for computing death rates are postcensal estimates based on the 2000 census, estimated as of July 1, 2007; see "Technical Notes." The asterisks (*) preceding the cause-of-death codes indicate that they are not part of the *International Classification of Diseases, Tenth Revision (ICD-10)*, Second Edition; see "Technical Notes"]

Cause of death (based on ICD-10, 2004)	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)	1.1	0.9	1.2	0.5	0.5	0.5	0.7	1.1	1.5	2.2	3.3	4.9
Accidental poisoning and exposure to noxious substances (X40-X49)	9.9	*	0.2	0.2	7.4	14.0	17.6	20.5	9.5	3.1	2.7	3.5
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64, W75-W99,X10-X39,X50-X59,Y86)	5.7	23.9	1.6	0.6	1.4	2.0	2.6	4.0	5.0	8.9	24.5	85.8
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11.5	0.5	9.7	13.0	15.6	17.7	15.5	12.6	16.3	15.6
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	5.8	0.1	4.5	5.7	6.7	8.0	8.5	8.8	12.2	11.0
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	5.7	0.3	5.3	7.3	8.9	9.7	7.0	3.8	4.1	4.6
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6.1	8.3	2.4	0.9	13.1	11.7	7.1	4.9	3.0	2.1	2.1	1.5
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	4.2	*	0.3	0.5	11.0	9.2	4.7	2.6	1.4	1.0	0.7	0.4
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)	1.9	7.9	2.1	0.4	2.1	2.5	2.3	2.2	1.6	1.2	1.4	1.0
Legal intervention (Y35,Y89.0)	0.1	*	*	*	0.2	0.3	0.2	0.1	*	*	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.8	2.1	0.3	0.1	1.4	2.1	2.9	3.4	1.8	0.9	1.0	1.3
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	*	*	*	0.2	0.1	0.1	0.1	0.1	*	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.7	2.1	0.3	0.1	1.2	2.0	2.8	3.3	1.7	0.9	1.0	1.3
Operations of war and their sequelae (Y36,Y89.1)	0.0	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40-Y84,Y88)	0.9	0.5	0.1	0.1	0.1	0.2	0.3	0.6	1.2	2.5	5.5	7.9
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁶	2.1	*	*	*	*	*	0.1	0.2	1.0	4.5	18.0	48.7

0.0 Quantity more than zero but less than 0.05.

* Figure does not meet standards of reliability or precision; see "Technical Notes."

... Category not applicable.

¹Figures for age not stated included in "All ages" but not distributed among age groups.

²Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see "Technical Notes."

³New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category in 2007. No deaths occurred from this cause in 2007.

⁴New ICD-10 code U04 (Severe acute respiratory syndrome [SARS]) was added to the category in 2007. No deaths occurred from this cause in 2007.

⁵Cause-of-death title was changed in 2007 to reflect the addition of SARS (ICD-10 code U04).

⁶Included in "Certain other intestinal infections (A04,A07-A09)" shown above. Beginning with data year 2006, Enterocolitis due to *Clostridium difficile* (A04.7) is shown separately at the bottom of tables showing 113 selected causes and is included in the list of rankable causes, see "Technical Notes."

NOTE: Complete confirmation of deaths from selected causes of death, considered to be of public health concern, were not provided by the following states--Connecticut, Florida, Indiana, Kentucky, Maryland, North Carolina, Oklahoma, Pennsylvania, Rhode Island, South Carolina, Washington, and West Virginia; see "Technical Notes."