User Guide to the 2021 Fetal Death Public Use File



2021 Fetal Death Data Set

User's Guide

ACKNOWLEDGMENTS

Preparation of this user's guide was coordinated by Elizabeth C.W. Gregory and Claudia P. Valenzuela of the Division of Vital Statistics (DVS) under the general direction of Joyce A. Martin, Lead Statistician, Reproductive Statistics Branch (RSB). The vital statistics computer file on which it is based was prepared by DVS staff.

The Division of Vital Statistics Director, Steven Schwartz, manages the Vital Statistics Cooperative Program, through which the vital registration offices of all states, the District of Columbia, New York City, Puerto Rico, Virgin Islands, Guam, American Samoa, and the Commonwealth of the Northern Mariana Islands, provided the data to the National Center for Health Statistics (NCHS). This Division, and its Data Acquisition, Classification, and Evaluation Branch (DACEB), and Information Technology Branch (ITB), processed computer edits, designed and programmed the tabulations, reviewed the data, prepared documentation for this publication, and were responsible for receipt and processing of the basic data file.

NCHS acknowledges the essential role of the vital registration offices of all states and territories in maintaining the system through which vital statistics data are obtained and their cooperation in providing the information on which this publication is based.

Questions on the documentation or substantive questions concerning the data should be directed to the Reproductive Statistics Branch, Division of Vital Statistics, NCHS, 3311 Toledo Road, Hyattsville, MD 20782-2003 (1-800) 232-4636.

Department of Health and Human Services Centers for Disease Control and Prevention National Center for Health Statistics Division of Vital Statistics

2021 Fetal Death Data Set

Contents

Introduction

Control count of records

List of data elements and locations

Record layout and definition of item and codes

Documentation tables

Figure 1. U.S. Standard Report of Fetal Death

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2021

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2021

External links

Fetal Mortality in the United States: 2019-2020 and 2020-Provisional 2021 https://www.cdc.gov/nchs/data/vsrr/vsrr026.pdf

Fetal Mortality: United States, 2020

https://www.cdc.gov/nchs/data/nvsr/nvsr71/nvsr71-04.pdf

Cause-of-Death Data from the Fetal Death File, 2018–2020 https://www.cdc.gov/nchs/data/nvsr/nvsr71/nvsr71-07.pdf

Births: Final Data for 2021

https://www.cdc.gov/nchs/data/nvsr/nvsr72/nvsr72-01.pdf

User Guide to the 2021 Period Linked Birth/Infant Death Public Use File (Forthcoming)

Introduction

This User Guide is for use with the 2021 fetal death public use data. The 2021 fetal death micro-data file may be downloaded at:

http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm [1]. The micro-data fetal death file for the U.S. does not include geographic detail (e.g., state or county of birth). The territories file contains selected geographic detail. Selected fetal death data for 2005-2020, including some geographic data and cause of death data, are available in CDC WONDER (http://wonder.cdc.gov), an interactive online data access tool. Selected fetal death data for 2021 are expected to be available in WONDER in early 2023.

For 2021, fetal death data are included on the file for all 50 states, the District Columbia, Guam and Puerto Rico. American Samoa did not report data for 2021 and data for the Northern Marianas and the U.S. Virgin Islands are excluded from the file due to the small number of events reported (4 and 15, respectively).

Definition of fetal death

"Fetal death" means death prior to the complete expulsion or extraction from its mother of a product of human conception, irrespective of the duration of pregnancy and which is not an induced termination of pregnancy. The death is indicated by the fact that after such expulsion or extraction, the fetus does not breathe or show any other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. Heartbeats are to be distinguished from transient cardiac contractions; respirations are to be distinguished from fleeting respiratory efforts or gasps.

This definition [2] has been adopted by the Centers for Disease Control and Prevention's National Center for Health Statistics (NCHS) as the nationally recommended standard and is

based on the definition published by the World Health Organization in 1950 and revised in 1988. The term "fetal death" is defined on an all-inclusive basis to end confusion arising from the use of such terms as stillbirth, spontaneous abortion, and miscarriage. All U.S. states and registration areas have definitions similar to the standard definition, except for Puerto Rico and Wisconsin, which have no formal definition [3]. Fetal deaths do not include induced terminations of pregnancy.

Reporting requirements for fetal death data

Reporting requirements for fetal deaths vary by state, and these differences have implications for comparisons of fetal and perinatal mortality rates by state [2]. Table A shows the period of gestation at which fetal death reporting is required for each reporting area for 2021. The majority of states require reporting of fetal deaths at 20 weeks of gestation or more, or a minimum of 350 grams birthweight (roughly equivalent to 20 weeks), or some combination of the two. Six states (Colorado, Georgia, Hawaii, New York, Rhode Island, and Virginia), American Samoa, and the U.S. Virgin Islands require reporting of fetal deaths at all periods of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), two states require reporting beginning at 12 weeks of gestation (although one of these does not send data for fetal deaths at less than 20 weeks to NCHS), and one state requires reporting beginning at 16 weeks of gestation.

User Guides for the Fetal Death Public Use files for 2007-2020 (Table A) indicated that South Dakota reported fetal deaths with birthweights of 500 grams or more (approximately 22 weeks of gestation or more). Over this period and to the present, however, South Dakota has reported fetal deaths at 20 weeks of gestation or more. Note that numbers of fetal deaths in the public use fetal death file reflect the correct reporting period of 20 weeks of gestation or more.

Classification of fetal deaths at 20 weeks of gestation or greater

To maximize the comparability of data by year and by state while including as much meaningful data as possible, most tables published by NCHS are based on fetal deaths occurring at gestations of 20 weeks or more. These tabulations also include fetal deaths for which gestation is not stated for those states requiring reporting at 20 weeks of gestation or more only. Fetal deaths of not-stated gestation are excluded for states requiring reporting of all products of conception, except for those with a stated birthweight of 350 grams or more. In 2021, this rule was applied to the following reporting areas: Arkansas, Colorado, Georgia, Hawaii, New York (including New York City), Oklahoma, Pennsylvania, Rhode Island, and Virginia. American Samoa and the Virgin Islands also requiring reporting of all products of conception when data are available (see Table A). The summation of these rules results in the creation of the tabulation flag (position 10), which, when invoked, produces the number of records with a stated or presumed period of gestation of 20 weeks or more.

Classification of fetal deaths by occurrence and residence

In tabulations by place of residence, fetal deaths occurring within the United States to U.S. citizens and to residents who are not citizens are allocated to the usual place of residence of the mother in the United States, as reported on the report of fetal death. Fetal deaths to U.S. residents occurring outside this country are not included in tabulations by place of residence or place of occurrence. Accordingly, the total count of fetal deaths for the United States by place of residence and by place of occurrence will not be identical. Fetal deaths to nonresidents of the United States are included in data by place of occurrence but excluded from data by place of residence, as previously indicated. See Tables 2 and 3, respectively, for the number of fetal deaths by residence and occurrence for the 50 states and the District of Columbia for 2021.

Geographic classification: The geographic code structure for the 2021 fetal death file is given in the NCHS manual, "Vital Records Geographic Classification, 2014," and in the country, county, and place geographic code files [4]. The geographic code structure on the 2021 file is based on results of the 2010 Census of Population.

Standard Report of Fetal Death

The U.S. Standard Report of Fetal Death, issued by the U.S. Department of Health and Human Services, has served for many years as the principal means for attaining uniformity in the content of the documents used to collect information on fetal deaths in the United States. The U.S. Standard Report of Fetal Death has historically been revised every 10-15 years. Most state reports conform closely in content to the standard report but are modified to the extent required by the particular state's needs or by special provisions of the state's vital statistics law.

The 2003 revision: In 2003, a revised U.S. Standard Report of Fetal Death was adopted (Figure 1). For more information on the 2003 standard report and details regarding the certificate revision and links to the documents referenced below, see the NCHS website of the 2003 report revision at http://www.cdc.gov/nchs/nvss/vital_certificate_revisions.htm. The 2003 report of fetal death replaces the previous 1989 U.S. Standard Report of Fetal Death [5,6]. Implementation of the 2003 U.S. Standard Report of Fetal Death (revised) by the states and independent reporting areas was phased in from 2003 to 2018. All states, the District of Columbia, Guam, the Northern Marianas, Puerto Rico, and the Virgin Islands reported data based on the 2003 U.S. Standard Report of fetal Death as of 2018.

The 2003 Revision of the U.S. Standard Report of Fetal Death introduced substantial changes to data content and quality. Many key data items are common between revisions; however, a number of items were substantively modified. The 2003 revision also includes many

new items never before collected on the Standard Report [5,6]. For details on data items comparable between revisions see the User Guide to the 2013 Fetal Death Public Use File [7].

A key aspect of the 2003 revision of the U.S. Standard Report of Fetal Death was the reengineering of the data collection and transmission system to improve data quality, speed of data collection and transmission, and to enhance standardization of data [5,8]. To encourage collection of data from the best sources, two worksheets were developed: the "Patient's Worksheet" (available at: https://www.cdc.gov/nchs/data/dvs/patientwkstfetaldth.pdf) [9] and the "Facility Worksheet" (available at: https://www.cdc.gov/nchs/data/dvs/FacilityFetal04.pdf) [10]. For the Patient's Worksheet, data are directly obtained from the mother and include items such as race, Hispanic origin, and educational attainment. For the Facility Worksheet, data are obtained directly from the medical records of the mother for items such as date of first prenatal care visit, pregnancy risk factors, and method of delivery. To assist hospital staff in completing the Facility Worksheet, a comprehensive instruction manual was developed: Guide to Completing the Facility Worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 Revision) ("Guide to the Facility Worksheet"; available at: https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf) [11]. Detailed definitions and instructions for data items that are collected from the Facility Worksheet are in the "Guide to the Facility Worksheet".

Additionally, a review of 2003-based report of fetal death revision items in 2014 by a collaborative effort among representatives from several vital statistics jurisdictions: The National Association for Public Health Statistics and Information Systems (NAPHSIS), and NCHS, resulted in the decision to drop a number of items from the national report of fetal death data file for reasons of poor data quality. For more information on this effort and for a full list of items

that were dropped, see:

https://www.cdc.gov/nchs/nvss/deleted items from birth fetal death files.htm.

The first ever eLearning training, "Applying Best Practices for Reporting Medical and Health Information on Birth Certificates," on completing the medical and health information for the birth certificate and report of fetal death was launched in October 2016 and has recently been updated. The training emphasizes the importance and uses of birth certificate and report of fetal death data and best practices for collecting specific medical and health items, including cause of fetal death. The audience for the training includes birth information specialists, physicians, nurses, and hospital administrators. Continuing education credits for nurses, physicians, and non-clinical staff are also available. The training is internet-based and approximately 45 minutes in length. It is available at: www.cdc.gov/nchs/training/BirthCertificateElearning.

Fetal death data files

Micro-data files: Fetal death micro-data files for data years 1982-2021 may be downloaded at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm. The general rules used to classify characteristics of fetal deaths are presented in several NCHS manuals [4, 8, 12]. These instructions are for states to use to collect and code the data items; they do not include NCHS edit recodes.

The 2003-2006 fetal death micro-data files include data items common to both the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death, as well as items exclusive to the 2003 revision. For 2007-2013, in an effort to improve the timeliness of fetal death data, the fetal death files only included items that were comparable across the 1989 and 2003 revisions of the U.S. Standard Report of Fetal Death. From 2014-2017, the fetal mortality files again included data comparable between the two revisions, as well as well as items exclusive to the 2003

revision. Beginning with 2018, the fetal death file only includes data exclusive to the 2003 revision.

The U.S. micro-data fetal death files for 2005 through 2021 do not include geographic detail (e.g., mother's state of residence). Tabulations of fetal death data by state-level geographic detail and many other demographic and health items are available in the CDC WONDER web application (http://wonder.cdc.gov/) for data years 2005-2020. Additionally, certain geographic level data may also be available upon request: See "NCHS Data Release and Access Policy for Micro-data and Compressed Vital Statistics Files", available at:

http://www.cdc.gov/nchs/nvss/dvs data release.htm.

The territories file includes data on fetal deaths occurring in Guam and Puerto Rico (American Samoa did not report data for 2021 and data for the Northern Marianas and the Virgin Islands were excluded due to small numbers). It includes limited geographical data such as information identifying individual territories and counties (or their equivalent) with populations of 100,000 or more by place of occurrence and residence.

Race of Mother

The 2003 revision of the U.S. Standard Report of Fetal Death allows the reporting of the five race categories either alone (i.e., single-race) or in combination (i.e., more than one race or multiple races) for the mother [6], in accordance with the revised standards issued by the Office of Management and Budget (OMB) in 1997 [13]. The five categories for race specified in the revised standards are: American Indian or Alaska Native (AIAN), Asian, Black or African American, Native Hawaiian or Other Pacific Islander (NHOPI), and White. Information on this change is presented elsewhere [14-16]. Starting in 2018, all states and the District of Columbia,

in addition to Puerto Rico, Guam, the Northern Marianas, and the Virgin Islands, began reporting race according to the 1997 revised OMB standards.

Prior to 2018, to provide uniformity and comparability of the data before all or most of the data were available in the new multiple-race format, it was necessary to "bridge" the responses of those for whom more than one race was reported (multiple race) to one, single race. Beginning in 2018, data by race are shown according to the 1997 OMB standards. Bridged-race data are no longer available on the public use file beginning with the 2020 data year.

Hispanic Origin of Mother

Hispanic origin and race: Hispanic origin and race are reported separately on the birth certificate (Figure 1). It is recommended that this information be reported directly by the mother via the Patient's Worksheet [9]. For 1989 through 2018, data on the public use file and in NCHS reports for specified Hispanic groups are shown in most cases for five specified Hispanic groups: Mexican, Puerto Rican, Cuban, Central and South American, and "other and unknown Hispanic." Starting with 2019, data are presented for the additional Hispanic group, Dominican (see item MHISPX in file position 143). This subgroup was previously included in "other and unknown Hispanic." In tabulations of fetal death data by race and Hispanic origin, data for persons of Hispanic origin are not further classified by race because the vast majority of Hispanic women are reported as white. In tabulations of fetal death data by race only, data for persons of Hispanic origin are included in the data for each race group according to the mother's reported race. In tabulations that include Hispanic origin, data for non-Hispanic persons are classified according to the race of the mother.

Cause of Death Data

A cause-of-fetal death item has been included on the report of fetal death since 1939 because it was considered critical information. However, the data were not released on public—use files or published until 2014 due to resource constraints and quality concerns. National data on cause of death are now released annually and are available for 2014-2021.

A major redesign of the cause of death (COD) section of the U.S. Standard Report of Fetal Death was undertaken with the 2003 revision. The goals of the redesign were to be consistent with instructions in the World Health Organization's (WHO) International Classification of Diseases (ICD) [17], and to thereby improve the quality and specificity of COD information, provide additional reporting guidance, and retain the flexibility to report any cause of death [18].

As with other deaths, the intent is for an attending physician or medical examiner or coroner to report cause of death [17]. The cause-of-fetal-death item requests a medical opinion from this person on the conditions and diseases resulting in or contributing to death, but also asks the medical certifier to report one cause separately (item 18a) from all other causes (item 18b) reported on the fetal death report. The certifier may form this medical opinion upon various medical tests, investigations, and examinations as is the case with other deaths. However, the term "initiating cause" used to refer to the one cause reported separately is unique to fetal deaths because of differences in the format of the cause item and how the initiating cause is determined for fetal deaths compared with the "underlying cause" term used with other deaths [18].

NCHS codes cause of fetal death reported by the certifier using the ICD-10 classification [17]. Coding is accomplished through a combination of automated and manual processes following the guidelines laid out in Instruction manual part 2k, Instructions for the Automated Classification of the Initiating and Multiple Causes of Fetal Deaths [19]. Literal text stated on the

fetal death report is assigned ICD-10 codes, and a single cause of death, the initiating cause of death, is selected from the conditions entered by the medical certifier in the cause-of-death section of the fetal death report. One section of the report (18a) is for the medical certifier to state the single condition that he or she considers the cause that initiated or triggered problems that resulted in the fetus dying at this point in time, so this is anticipated to be the initiating cause of death. If more than one cause or condition is entered by the medical certifier, the initiating cause is determined by the placement of the condition on the report, provisions of the ICD, and associated selection rule and modifications. A second section of the report (18b) is for the medical certifier to state any other conditions or causes that he or she felt played a role in causing the fetal death. Since more medical information may be reported on the report than is directly reflected in the initiating cause of death, this additional information is captured in multiple cause-of-fetal-death data.

The fetal death data file also includes a cause of death data item for 124 Selected Causes of Fetal Death (fetal cause list) (tables 7A and 7B). The reporting area for these tables includes areas that used the 2003 revision format and where less than 50 percent of an area's cause data were attributed to the Fetal death of unspecified cause (unspecified cause) (P95).

Incomplete national reporting: The use of reporting flags

The 2021 national fetal death file includes items that are not reported by some vital statistics jurisdictions or are reported but have an unknown level of 50% or higher and are flagged as not reporting. Reporting flags were developed to help the user more readily identify the correct number of fetal deaths for items with less than full national reporting. Reporting flags are available for many items in the file, and positions for reporting flags are noted along with each data item in the file layout.

Translating "blanks"

In the 2021 fetal death file, for data items which are not reported by all states, fetal deaths to residents of a state reporting the item that occurred in a state not reporting the item, are represented by "blanks." Blanks should be treated as "unknowns" for tabulations. In sum, the correct use of reporting flags and translation of blanks will result in an accurate tally of fetal deaths for items with incomplete national reporting. For an example of SAS code that may be used to incorporate the correct use of reporting flags and the translation of blanks, see below.

Example of SAS code using reporting flags (and translating blanks)

The example below is for the receipt of WIC item, for which data for 48 states are included for 2021 (New York City does not report this item and the District of Columbia, Hawaii, and Rhode Island reported this item for 2021 but had unknown levels of 50% or higher and were flagged as not reporting).

Sample SAS program

```
01
       DATA work;
02
              INFILE 'c:fet21us.dat' LRECL=3150;
03
              INPUT
04
                      oe tabflg
                                  10
05
                     wic $229
06
                     restatus 131
07
                     f wic $377;
80
09
              /*Include if 20 weeks gestation or more*/
10
              IF oe tabflg EQ 2;
11
              /*Exclude foreign residents*/
```

```
12
             IF restatus NE 4;
13
             /*Select reporting area*/
14
             IF f wic=1;
15
             /*Convert blanks to unknown*/
16
             IF wic =' 'THEN wic ='U';
17
18
       PROC FREQ;
19
             TABLE wic;
20
       RUN;
```

In this example, "oe_tabflg" is used to select fetal deaths of 20 weeks gestation or more. Also, "restatus" is used to exclude fetal deaths to foreign residents (this is standard practice for all NCHS tabulations). As well in this example, blanks are represented by character values SAS code = (''). However, for some items in the file, e.g., number of previous cesarean deliveries, blanks are represented by numeric values for which the SAS code is ('.').

Quality of Data

State-Specific Data Quality Issues

These state-specific data quality issues are of concern due to potential underreporting and/or inaccurate reporting for 2021.

<u>Date of prenatal care visit</u> – Virginia-No records have been reported as not applicable

 (i.e., receiving no prenatal care) since May 2018. Data for this item for this state should be used with caution.

- <u>Trial of labor attempted</u> Virginia-For 2021, all records reported as cesarean deliveries
 were erroneously reported as "yes" for trial of labor attempted. As a result, this item for
 this state has been flagged as not reporting and should not be used.
- Attendant New Jersey-No records have been reported as having a certified nurse midwife as the attendant since mid-2021. Data for this item for this state should be used with caution.
- <u>Plurality</u> District of Columbia and Hawaii-The number of multiple deliveries may be
 underreported for these jurisdictions due to the high percentage of records reported as
 unknown plurality; records with unknown plurality are imputed to a singleton delivery.
 Data for this item for these reporting areas should be used with caution.

Percentage of unknown responses

Information on the completeness of reporting of fetal death (with a stated or presumed period of gestation of 20 weeks or more) data is shown in Table B, which presents a listing of items and the percentage of records that were not stated for each reporting area: all 50 states, the District of Columbia, New York City, Puerto Rico, Guam, American Samoa, the Northern Marianas, and the Virgin Islands. In general, percentages of unknown responses are considerably higher for fetal deaths than for live births, and among fetal deaths the percentage unknown is higher for fetal deaths that occur earlier in the gestational period. In fetal mortality tables, unknown responses are shown in frequencies tables but are excluded from the computation of percent distributions and fetal and perinatal mortality rates. Thus, rates for variables with a substantial percentage of unknown responses (such as birthweight) may understate the "true" rates of fetal mortality for that characteristic.

As high levels of missing data are a data quality concern, beginning with the 2017 data year, jurisdictions with a 50% or greater unknown level for a particular item are flagged as not reporting that item.

State Specific Issues

The items below had unknown levels of 50% or higher for the jurisdiction(s) noted and are flagged as not reporting for those jurisdictions in 2021:

- <u>Inside city limits</u>-Rhode Island
- Mother's education-Hawaii and Rhode Island
- <u>Father's age-</u> Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New Mexico,
 New York (including New York City), Virginia, Wisconsin, and Guam
- Previous live births dead-Hawaii
- Live birth order-Hawaii
- <u>Date of last live birth</u>-Hawaii and Washington
- <u>Date of first prenatal care</u>-Hawaii and Rhode Island
- WIC- District of Columbia, Hawaii, and Rhode Island
- Mother's height-Hawaii
- Mother's pre-pregnancy weight-Hawaii
- Mother's pre-pregnancy BMI-Hawaii
- Number of previous cesarean deliveries-Washington
- Fetal presentation-Hawaii
- Route and method of delivery-Hawaii
- Trial of labor attempted-Hawaii and Virginia
- <u>Date of last normal menses</u>-Colorado and Hawaii

- Birthweight-Hawaii
- Estimated time of fetal death-District of Columbia, Georgia, Hawaii, Kansas, Missouri,
 Mississippi, Nevada, New York, Oklahoma, Tennessee, Rhode Island, Virginia, Guam,
 and Puerto Rico
- <u>Cause of death</u>- Arizona, California, Georgia, Hawaii, Mississippi, New York (including New York City), North Dakota, Vermont, Wisconsin, and Guam

Computation of rates

Fetal mortality rates are computed as the number of fetal deaths at 20 weeks of gestation or more per 1,000 live births and fetal deaths at 20 weeks or more. Perinatal mortality rates are computed in a similar fashion, as shown below. The denominators for all fetal and perinatal mortality rates are live births plus fetal deaths in the specified gestational age group, thus representing the population at risk of the event [2].

Fetal mortality rate = Fetal deaths at 20 weeks of gestation or more x 1,000

Live births and fetal deaths at 20 weeks or more

Perinatal mortality rate, definition I =

Fetal deaths at 28 weeks or more and infant deaths under 7 days x 1,000

Live births and fetal deaths at 28 weeks or more

Perinatal mortality rate, definition II =

Fetal deaths at 20 weeks or more and infant deaths under 28 days x 1,000

Live births and fetal deaths at 20 weeks or more

In each case, the fetal deaths included in the denominator of each rate mirror the fetal deaths included in the numerator. A previous NCHS report [20] contains information on the historical development of various perinatal measures. An asterisk (*) is shown in place of any rate based on fewer than 20 fetal or perinatal deaths in the numerator.

Random variation in fetal and perinatal mortality

The number of fetal deaths, perinatal deaths, and live births reported for an area represent complete counts of such events. As such, they are not subject to sampling error, although they are subject to nonsampling error in the registration process. However, when the figures are used for analytic purposes, such as the comparison of rates over time, for different areas, or among different subgroups, the number of events that actually occurred may be considered as one of a large series of possible results that could have arisen under the same circumstances [21]. As a result, numbers of births, fetal deaths, perinatal deaths, and fetal and perinatal mortality rates are subject to random variation. The probable range of values may be estimated from the actual figures according to certain statistical assumptions.

In general, distributions of vital events may be assumed to follow the normal distribution. When the number of events is large, the relative standard error (RSE) is usually small. When the number of events is small (perhaps less than 100) and the probability of such an event is small, considerable caution must be used in interpreting the data. Such infrequent events may be assumed to follow a Poisson probability distribution. Estimates of RSEs and 95% confidence intervals are shown below. In the formulas, D is the number of fetal or perinatal deaths and B is the number of live births plus fetal deaths used as the denominator in computing fetal and perinatal mortality rates.

The formulas for the two RSEs are as follows:

RSE(D)=
$$100*\sqrt{\frac{1}{D}}$$
 and

RSE(B)=100*
$$\sqrt{\frac{1}{B}}$$
.

For example, let us say that for group A the number of fetal deaths was 238, whereas the number of live births plus fetal deaths in the denominator was 32,650 yielding a fetal mortality rate of 7.29 fetal deaths per 1,000 live births and fetal deaths.

The RSE of the deaths =
$$100*\sqrt{\frac{1}{238}} = 6.48$$
,

whereas the RSE for the births plus fetal deaths in the denominator is

$$RSE=100*\sqrt{\frac{1}{32,650}}=0.55.$$

The formula for the RSE of the fetal mortality rate is:

RSE=100*
$$\sqrt{\frac{1}{D} + \frac{1}{B}}$$

Thus the RSE for the example above is:

$$= 100*\sqrt{\frac{1}{238} + \frac{1}{32,650}} = 6.51.$$

Normal distribution

When the number of events is greater than 100, the normal distribution is used to estimate the 95 percent confidence intervals of a rate, R_{I_i} as follows:

Lower:
$$R_I - 1.96 * R_I * \frac{RSE(R1)}{100}$$

Upper:
$$R_I + 1.96 * R_I * \frac{RSE(R1)}{100}$$

Thus, for Group A:

Lower:
$$7.29 - (1.96 * 7.29 * \frac{6.51}{100}) = 6.36$$

Upper:
$$7.29 + (1.96 * 7.29 * \frac{6.51}{100}) = 8.22$$

Thus the chances are 95 out of 100 that the true fetal or perinatal mortality rate for Group A lies somewhere in the 6.36–8.22 interval.

Poisson distribution

When the number of events in the numerator is less than 100 the confidence interval for the rate, R_{I_i} can be estimated based on the Poisson distribution using the values in Table III (2).

Lower:
$$R_I$$
*L(.95, D_{adj})

Upper:
$$R_I$$
*U(.95, D_{adj})

where D_{adj} is the adjusted number of fetal or perinatal deaths (rounded to the nearest integer) used to take into account the RSE of the number of deaths in the numerator and the number of live births plus fetal deaths in the denominator, and is computed as follows:

$$D_{adj} = \frac{D * B}{D + B}$$

 $L(.95,\,D_{adj}) \mbox{ and } U(.95,\,D_{adj}) \mbox{ refer to the values in Table III corresponding to the value of} \\ D_{adj}.$

For example, let us say that for Group B the number of fetal deaths was 73, and the number of live births plus fetal deaths in the denominator was 11,422, and the fetal mortality rate was 6.39.

$$D_{adj} = \frac{73 * 11,422}{73 + 11,422} = 73$$

Therefore the 95 percent confidence interval (using the formula in Table III for 1\N99 infant deaths) =

Lower: 6.39*0.78384 = 5.01

Upper: 6.39*1.25735 = 8.03

Comparison of two fetal or perinatal mortality rates

If either of the two rates to be compared is based on less than 100 deaths, compute the confidence intervals for both rates and check to see if they overlap. If so, the difference is not statistically significant at the 95 percent level. If they do not overlap, the difference is statistically significant. If both of the two rates (R_1 and R_2) to be compared are based on 100 or more deaths, the following z-test should be used to define a significance test statistic:

$$z = \frac{R_I - R_2}{\sqrt{R_I^2 \left(\frac{RSE(R_I)}{100}\right)^2 + R_2^2 \left(\frac{RSE(R_2)}{100}\right)^2}}$$

If $|z| \ge 1.96$, then the difference is statistically significant at the 0.05 level and if |z| < 1.96, the difference is not significant.

References

- 1. National Center for Health Statistics. Fetal Death 2021. Public use file. Hyattsville, Maryland: National Center for Health Statistics. Annual internet product. 2023. Available at: http://www.cdc.gov/nchs/data access/VitalStatsOnline.htm.
- 2. MacDorman MF, Gregory ECW. Fetal and perinatal mortality: United States, 2013. National vital statistics reports; vol 64 no 8. Hyattsville, MD: National Center for Health Statistics. 2015.
- 3. Kowaleski J. State definitions and reporting requirements for live births, fetal deaths, and induced terminations of pregnancy (1997 revision). Hyattsville, MD: National Center for Health Statistics. 1997. Available from: http://www.cdc.gov/nchs/data/misc/itop97.pdf.
- 4. Division of Vital Statistics. Instruction manual part 8: Geographic classification, 2014. Hyattsville, Maryland: National Center for Health Statistics. Available at: https://www.cdc.gov/nchs/data/dvs/IMP8 2014.pdf.
- 5. National Center for Health Statistics. Report of the Panel to Evaluate the U.S. Standard Certificates. Hyattsville, Maryland: National Center for Health Statistics. 2000
- 6. National Center for Health Statistics. 2003 revision of the U.S. Standard Report of Fetal Death. 2003.
- 7. National Center for Health Statistics. User Guide to the 2013 Fetal death Public Use File. Hyattsville, Maryland: National Center for Health Statistics. Annual product 2015. Available at: http://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.
- 8. National Center for Health Statistics. Fetal death edit specifications for the 2003 Revision of the U.S. Standard Certificate of Birth. 2012.

https://www.cdc.gov/nchs/data/dvs/fetal death edit specifications.pdf.

- 9. National Center for Health Statistics. Patient's Worksheet for Report of Fetal Death.

 2019. Available at: https://www.cdc.gov/nchs/data/dvs/fetal-death-mother-worksheet-english-2019-508.pdf.
- 10. National Center for Health Statistics. Facility Worksheet for Report of Fetal Death. 2019. Available at: https://www.cdc.gov/nchs/data/dvs/fetal-death-facility-worksheet-2019-508.pdf.
- 11. National Center for Health Statistics. Guide to completing the facility worksheets for the Certificate of Live Birth and Report of Fetal Death (2003 revision). 2019 update. Hyattsville, Maryland: National Center for Health Statistics.

https://www.cdc.gov/nchs/data/dvs/GuidetoCompleteFacilityWks.pdf.

- 12. Division of Vital Statistics. Computer edits fetal death data --2003 Revised Certificate. Hyattsville, Maryland: National Center for Health Statistics.
- 13. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. Fed Regist 62FR58781–58790. October 30, 1997.
- 14. Ingram DD, Parker JD, Schenker N, et al. United States Census 2000 population with bridged race categories. National Center for Health Statistics. Vital Health Stat 2(135). 2003.
- 15. Johnson D. Coding and editing multiple race. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP. Portland, Oregon. June 6–10, 2004.
- 16. Weed JA. NCHS procedures for multiple-race and Hispanic origin data: Collection, coding, editing, and transmitting. Presented at the 2004 Joint Meeting of NAPHSIS and VSCP. Portland, Oregon. June 6–10, 2004.
- 17. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. Geneva, Switzerland. 2009.

- 18. Hoyert DL, Gregory ECW. Cause-of-death data from the fetal death file, 2015–2017. National Vital Statistics Reports; vol 69 no 4. Hyattsville, MD: National Center for Health Statistics. 2020.
- 19. National Center for Health Statistics. Instruction manual, part 2k: Instructions for the automated classification of the initiating and multiple causes of fetal deaths, 2014. Hyattsville, MD. 2014.
- 20. Hoyert DL. Perinatal mortality in the United States, 1985–91. National Center for Health Statistics. Vital Health Stat 20(26). 1995. Available from:

 http://www.cdc.gov/nchs/data/series/sr-20/sr20-026.pdf.
- 21. Brillinger DR. The natural variability of vital rates and associated statistics. Biometrics 42(4):693–734. 1986.

Control Count of Records

File Characteristics: Fetal Public Use File 2021

All files:

Record format: Blocked, Fixed, Format Code scheme: Numeric/Alphabetic/Blank

Record length: 3150

Record counts

<u>U.S. file record count:</u>

Counts of all fetal deaths:

Record count	42,428
By occurrence	42,428
By residence	42,329
To foreign residents	99

Counts of 20 weeks or more:

Record count	21,194
By occurrence	21,194
By residence	21,105
To foreign residents	89

Counts of less than 20 weeks:

Record count	21,234
By occurrence	21,234
By residence	21,224
To foreign residents	10

U.S. territories file record count:

Counts of all fetal deaths:

Record count	275
By occurrence	275
By residence	274
To foreign residents	1

Counts of 20 weeks or more:

Record count	274
By occurrence	274
By residence	273
To foreign residents	1

Counts of less than 20 weeks:

Record count	1
By occurrence	1
By residence	1
To foreign residents	-

2021 LIST OF DATA ELEMENTS AND LOCATIONS

<u>Da</u>	ta Items	Locations
1.	General a) Tabulation flags (gestational age) b) Data year c) Resident status	10 11-14 131
2.	Prenatal Care a) Month began	202-203
3.	Fetus a) Sex b) Plurality c) Weight at delivery d) Gestational age e) Month/year of delivery	316 301 349-355 340-344(NCHS Standard item), 331-335 11-16
	f) Time of delivery g) Day of week of delivery	21-24 25
4.	Mother a) Age b) Race c) Education d) Hispanic origin	86-90 132-136 145 139-144
5.	Pregnancy History a) Live birth order b) Birth interval	187 197-201
6.	Father a) Age	177-180
7.	Other Items a) Attendant at delivery b) Place of delivery c) WIC receipt	283 34, 52 229
8.	Medical and Health Data a) Method of delivery b) Medical risk factors c) Other risk factors i. Tobacco ii. Mother's pre-pregnancy body mass index (BMI) d) Maternal morbidity	274-277, 280 257-267 230-238 245-249 281-282
9.	Residence reporting flags	372-442

10. Cause of Death

a.) Initiating cause or condition code
b.) Initiating cause or condition recode
c.) Initiating cause or condition flag
2603-2607
2643-2645
2651

2021 Fetal Death Public Use Record Layout

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
1-6	6	FILLER					
7	1	VERSION	Version		A,S	A S	State used the 2003 version of the US Standard Report of Fetal Death (Revised) State used the 1989 version of the US Standard Report of Fetal Death (Unrevised)
8-9	2	FILLER					
10	1	OE_TABFLG	Obstetric Estimate Tabul (Use with NCHS Standard		A,S	1 2	Under 20 weeks (exclude) 20 weeks or more (include)
11-14	4	DOD_YY	Delivery Year		A,S	2014	Year of delivery
15-16	2	DOD_MM	Delivery Month		A,S	01 02 03 04 05 06 07 08 09 10 11	January February March April May June July August September October November December
17-20	4	FILLER					
21-24	4	DOD_TT	Delivery Time	c409	A	0000-2 9999	2359 Time of Birth Not Stated
25	1	DOD_WK	Weekday		A,S	1 2 3	Sunday Monday Tuesday

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						4 5 6 7	Wednesday Thursday Friday Saturday
26-27	2	OSTATE	Territory of Occurrence (This item is available in the geographic codes are not av		A,S ìle)		
			<u>Territories</u>			AS GU MP PR VI	American Samoa Guam Northern Marianas Puerto Rico Virgin Islands
28-29	2	FILLER					
30-32	3	OCNTYFIPS	Occurrence FIPS County (This item is available in the geographic codes are not av Data are suppressed if popu	vailable in the U.S. f	A,S île)	000 001-998 999	No county level geography County of Occurrence (See Geographic Table) County of less than 100,000
33	1	OCNTYPOP	Occurrence Territory Cou (This item is available in the geographic codes are not av	territory file only,	A,S ìle)		
J.						0 1 2 3	County of 1,000,000 or more County of 500,000 to 1,000,000 County of 250,000 to 500,000 County of 100,000 to 250,000

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						9	County less than 100,000
34	1	BFACIL	Delivery Place (Revised)	409	A	1 2 3 4 5 6 7 9	Hospital Freestanding Birth Center Home (intended) Home (not intended) Home (unknown if intended) Clinic/Doctor's Office Other Unknown
35-51	17	FILLER					
52	1	BFACIL3	Delivery Place Recode		A,S	1 2 3	In Hospital Not in Hospital Unknown/not stated
53-83	31	FILLER					
84	1	MAGE_IMPFLG	Mother's Age Imputed		A,S	Blank 1	Age not imputed Age imputed
85	1	MAGE_REPFLG	Reported Age of Mother F	lag	A,S	Blank 1	Reported age not used Reported age used
86-87	2	MAGER	Mother's Single Year of A	ge 432	A,S	12 13 14 15 16	12 years and under 13 years 14 years 15 years 16 years

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						17	17 years
						18	18 years
						19	19 years
						20	20 years
						21	21 years
						22	22 years
						23	23 years
						24	24 years
						25	25 years
						26	26 years
						27	27 years
						28	28 years
						29	29 years
						30	30 years
						31	31 years
						32	32 years
						33	33 years
						34	34 years
						35	35 years
						36	36 years
						37	37 years
						38	38 years
						39	39 years
						40	40 years
						41	41 years
						42	42 years
						43	43 years
						44	44 years
						45	45 years
						46	46 years
						47	47 years
						48	48 years
						49	49 years
						50	50 years and over
88-89	2	MAGER14	Mother's Age Recode 14	432	A,S	01	Under 15 Years
			_			03	15 years
						04	16 years
						05	17 years
						06	18 years
						07	19 years
* F 4	T 7 .	1 (37) (1	C 11 ' 1 1				

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						08 09 10 11 12 13	20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years
90	1	MAGER9	Mother's Age Recode 9	432	A,S	1 2 3 4 5 6 7 8	Under 15 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years
91-92	2	MBCNTRY	Mother's Birth Country (This item is available in the geographic codes are not ava			A AA	-ZZ See Geographic Documentation
93-96	4	FILLER					
97	1	MBSTATE_REC	Mother's Birth State Recode	c409	A	1 2 3	Native born (50 U.S. States) Foreign born (includes territories) Unknown or Not Stated
98-101	4	FILLER					
102-103	2	MRSTATEPSTL	Mother's Residence Postal (This item is available in the geographic codes are not ava	territory file only,	A,S		
			<u>United States</u>			AK AL AR	Alaska Alabama Arkansas

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting	Vers*	Values	Definition
				Flag Position			

ΑZ Arizona CACalifornia CO Colorado CTConnecticut DE Delaware DC District of Columbia FL Florida GA Georgia Hawaii HI ID Idaho ILIllinois IN Indiana IΑ Iowa KS Kansas KY Kentucky LA Louisiana Massachusetts MA MD Maryland Maine ME MI Michigan Minnesota MN MO Missouri MS Mississippi MT Montana NC North Carolina ND North Dakota NE Nebraska NH New Hampshire NJ New Jersey NM New Mexico NVNevada NYNew York ОН Ohio Oklahoma OK OR Oregon Pennsylvania PA RI Rhode Island SC South Carolina SD South Dakota TNTennessee TXTexas

^{*} For the Version column (Vers) the following codes apply:

A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

s 1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						LIT	II.
						UT VA	Utah Virginia
						VT	Vermont
						WA	Washington
						WI	Wisconsin
						WV	West Virginia
						WY	Wyoming
			<u>Territories</u>			AS	American Samoa
			remones			GU	Guam
						MP	Northern Marianas
						PR	Puerto Rico
						VI	Virgin Islands
			Canadian Province	es_		AB	Alberta
						BC	British Columbia
						MB	Manitoba
						NB	New Brunswick
						NL NT	Newfoundland and Labrador Northwest Territories
						NS NS	Nova Scotia
						NU	Nunavut
						ON	Ontario
						PE	Prince Edward Island
						QC	Quebec
						SK	Saskatchewan
						YT	Yukon Territory
			<u>Foreign</u>				
						XX	Foreign (Rest of world)
						ZZ	Not Classifiable
104-106	3	MRCNTYFIPS	Mother's FIPS County of I (This item is available in the geographic codes are not av	territory file only,	le)		
			Data are suppressed if popu		,	000 001-998	No county level geography County of residence (See Geographic Table)
* For t	he Versic	on column (Vers) t	he following codes apply:				

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

¹⁹⁸⁹ Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						999	County of less than 100,000
107-111	5	FILLER					
112	1		Population of Residence C lable in the territory file only, are not available in the U.S. f	•	A,S	0 1 2 2	County of 1,000,000 or more County of 500,000 to 1,000,000 County of 250,000 to 500,000
						3 9 Z	County of 100,000 to 250,000 County less than 100,000 Foreign resident
113-129	17	FILLER					
130	1	RECTYPE	Record Type (This item is available in the geographic codes are not av		A,S		
			geographic codes are not av	vanable in the O.S. Ju	<i>e)</i>	1	RESIDENT: State and county of occurrence and residence
						2	are the same. NONRESIDENT: State and county of occurrence and residence are different.
131	1	RESTATUS	Residence Status <u>United States</u>		A,S	1	RESIDENT: State and county of occurrence and residence
						2	are the same. INTRASTATE NONRESIDENT: State of occurrence and
						3	residence are the same but county is different. INTERSTATE NONRESIDENT: State of occurrence and
						4	residence are different but both are one of the 50 US states or District of Columbia. FOREIGN RESIDENT: The state of residence is not one of the 50 US states or District of Columbia.
ቃ ፑ ብ	3 7	1. (1)	<u>Territories</u>			1	RESIDENT: State and county of occurrence and residence

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						2 3 4	are the same. (Unique to Guam, all US residents are considered residents of Guam and thus are assigned 1.) INTRATERRITORY NONRESIDENT: Territory of occurrence and residence are the same but county is different. INTERTERRITORY RESIDENT: Territory of occurrence and residence are different but both are US Territories. FOREIGN RESIDENT: The residence is not a US Territory.
132-133	2	MRACE31	Mother's Race Recode 31	425	A	01 02 03 04 05 06 07 08 09 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31	White (alone) Black (alone) AIAN (American Indian or Alaskan Native) (alone) Asian (alone) NHOPI (Native Hawaiian or Other Pacific Islander) (alone) Black And White Black and AIAN Black and Asian Black and NHOPI AIAN and White AIAN and White AIAN and NHOPI Asian and White Asian and NHOPI NHOPI and White Black, AIAN, and White Black, AIAN, and White Black, Asian, and White AIAN, NHOPI, and White AIAN, NHOPI, and White AIAN, Asian, and NHOPI Black, AIAN, Asian, and White Black, AIAN, Asian, NHOPI, and White Black, Asian, NHOPI, and White Black, Asian, NHOPI, and White Black, AIAN, Asian, NHOPI, and White

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
134	1	MRACE6	Mother's Race Recode 6	425	A	1 2 3 4 5 6	White (alone) Black (alone) AIAN (alone) Asian (alone) NHOPI (alone) More than one race
135-136	2	MRACE15	Mother's Race Recode 15	425	A	01 02 03 04 05 06 07 08 09 10 11 12 13 14	White (alone) Black (alone) AIAN (alone) Asian Indian (alone) Chinese (alone) Filipino (alone) Japanese (alone) Korean (alone) Vietnamese (alone) Other Asian (alone) Hawaiian (alone) Guamanian (alone) Samoan (alone) Other Pacific Islander (alone) More than one race
137	1	FILLER					
138	1	MRACEIMP	Mother's Race Imputed		A,S	Blank 1 2	Mother's race not imputed Unknown race imputed All other races, formerly coded 09, imputed.
139-141	3	FILLER					
142	1	UMHISP	Mother's Hispanic Origin	424	A,S	0	Non-Hispanic

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Report Flag Po		Vers*	Values	Definition
							1 2 3 4 5 9	Mexican Puerto Rican Cuban Central or South American Other and Unknown Hispanic Origin unknown or not stated
143	1	MHISPX	Mother's Hispanic Orig	gin			0 1 2 3 4 5 6	Non-Hispanic Mexican Puerto Rican Cuban Central or South American Dominican Other and Unknown Hispanic Origin unknown or not stated
144	1	SR_MRACEHIS	P Mother's Race/Hispanic O	Origin	c425	A	1 2 3 4 5 6 7 8	Non-Hispanic White (alone) Non-Hispanic Black (alone) Non-Hispanic AIAN (alone) Non-Hispanic Asian (alone) Non-Hispanic NHOPI (alone) Non-Hispanic more than one race Hispanic Origin unknown or not stated
145	1	MEDUC	Mother's Educ-Revised	37	2	A	1 2 3 4 5 6 7 8	8 th grade or less 9 th through 12 th grade with no diploma High school graduate or GED completed Some college credit, but not a degree Associate degree (AA, AS) Bachelor's degree (BA, AB, BS) Master's degree (MA, MS, MEng, Med, MSW, MBA) Doctorate (PhD, EdD) or Professional degree (MD, DDS, DVM, LLB, JD) Unknown

146-171 26 FILLER

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
172	1	FAGERPT_FLG	Father's Reported Age Use	ed	A,S	Blank 1	Father's reported age not used Father's reported age used
173-176	4	FILLER					
177-178	2	FAGECOMB	Father's Combined Age	426 A	.,S	09-98 99	Father's combined age in years Unknown or not stated
179-180	2	FAGEREC11	Father's Age Recode 11	426	A,S	01 02 03 04 05 06 07 08 09 10	Under 15 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years 50-54 years 50-98 years Not stated
181-182	2	PRIORLIVE	Prior Births Now Living	440	A,S	00-30 99	Number of children still living from previous live births. Unknown or not stated
183-184	2	PRIORDEAD	Prior Births Now Dead	441	A,S	00-30 99	Number of children dead from previous live births. Unknown or not stated
185-186	2	FILLER					
187	1	LBO_REC	Live Birth Order Recode	441	A,S	0-7 8 9	Number of live birth order. 8 or more live births Unknown or not stated

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
188-196	9	FILLER					
197-199	3	ILLB_R	Interval Since Last Live B	irth Recode 427	A		Plural delivery Months since last live birth Not applicable Unknown or not stated
200-201	2	ILLB_R11	Interval Since Last Live B	irth Recode 11 427	A	00 01 02 03 04 05 06 07 08 88 99	Zero to 3 months (plural delivery) 4 to 11 months 12 to 17 months 18 to 23 months 24 to 35 months 36 to 47 months 48 to 59 months 60 to 71 months 72 months and over Not applicable Unknown or not stated
202-203	2	PRECARE	Month Prenatal Care Bega	an (Revised) 403	A	00 01-10 99	No prenatal care Month prenatal care began Unknown or not stated
204	1	PRECARE_REC	Month Prenatal Care Bega	an Recode (Revised 403	I) A	1 2 3 4 5	1st to 3rd month 4th to 6th month 7th to final month No prenatal care Unknown or not stated
205-228	24	FILLER					
229	1	WIC	WIC	377	A	Y N	Yes No

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						U	Unknown or not stated
230-231	2	CIG_0	Cigarettes Before Pregnand	ey 405	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
232-233	2	CIG_1	Cigarettes 1st Trimester	406	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
234-235	2	CIG_2	Cigarettes 2 nd Trimester	407	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
236-237	2	CIG_3	Cigarettes 3 rd Trimester	408	A	00-97 98 99	Number of cigarettes daily 98 or more cigarettes daily Unknown or not stated
238	1	CIG_REC	Cigarette Recode (Revised)	374	A	Y N U	Yes No Unknown or not stated
239-242	4	FILLER					
243-244	2	M_Ht_In	Mother's Height in Inches	375	A	30-78 99	Height in inches Unknown or not stated
245-248	4	BMI	BMI (prepregnancy)	376	A	13.0-69.9 99.9	Body Mass Index Unknown or not stated
249	1	BMI_R	Body Mass Index Recode (prepregnancy)	376	A	1 2 3 4	Underweight <18.5 Normal 18.5-24.9 Overweight 25.0-29.9 Obesity I 30.0-34.9

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						5 6 9	Obesity II 35.0-39.9 Extreme Obesity III ≥ 40.0 Unknown or not stated
250-252	3	FILLER					
253-255	3	PWgt_R	Prepregnancy Weight Rec	ode 376	A	075-375 999	Weight in pounds Unknown or not stated
256	1	FILLER					
257-262	6	Risk Factors (Re	vised) ckbox items indented below for	ollow this code struc	A ture:		Y Yes N No U Unknown or not stated
257 258 259 260 261 262	1 1 1 1 1	RF_DIAB RF_GEST RF_PHYP RF_GHYP RF_ECLAM RF_INFTR	Prepregnancy Diabetes Gestational Diabetes Prepregnancy Hypertension Gestational Hypertension Hypertension Eclampsia Infertility Treatment	378 379 on380 381 382 383			Cliniowii oi not stated
263	1	RF_FEDRG	Fertility Enhancing Drugs	384	A	Y N X U	Yes No Not applicable Unknown or not stated
264	1	RF_ARTEC	Asst. Reproductive Techno	ology 385	A	Y N X U	Yes No Not applicable Unknown
265	1	RF_CESAR	Previous Cesareans	386	A	Y N U	Yes No Unknown or not stated

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Positi		Vers*	Values	Definition
266-267	2	RF_CESARN	Previous Cesareans Num	387		A	00 01-30 99	None Number of previous cesareans Unknown or not stated
268-273	6	FILLER						
274-277 274	6 1	Method of Deliv ME_PRES	ery (Revised) Fetal Presentation	388		A	1 2 3 9	Cephalic Breech Other Unknown or not stated
275	1	ME_ROUT	Route & Method of Delive	ry	389	A	1 2 3 4 9	Spontaneous Forceps Vacuum Cesarean Unknown or not stated
276	1	ME_TRIAL	Trial of Labor Attempted	390	0	A	Y N X U	Yes No Not applicable Unknown or not stated
277	1	RDMETH_REC	Delivery Method Recode (389	A	1 2 3 4 5 6 9	Vaginal (excludes vaginal after previous C section) Vaginal after previous c-section Primary C-section Repeat C-section Vaginal (unknown if previous c-section) (2003 Standard only) C-section (unknown if previous c-section) (2003 Standard only) Not stated

²⁷⁸⁻²⁷⁹ 2 FILLER

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

¹⁹⁸⁹ Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
280	1	DMETH_REC	Delivery Method Recode	410	A,S	1 2 9	Vaginal C-section Not stated
281-282	2	Matern The che	nal Morbidity ockbox items indented below for	ollow this structure:	A	Y N U	Yes No Unknown or not stated
281 282	1 1	MM_RUPT MM_ICU	Ruptured Uterus Admit to Intensive Care	391 392			
283	1	ATTEND	Attendant	433	A,S	1 2 3 4 5 9	Doctor of Medicine (MD) Doctor of Osteopathy (DO) Certified Nurse Midwife (CNM) Other Midwife Other Unknown or not stated
284-300	17	FILLER					
301	1	DPLURAL	Plurality Recode		A,S	1 2 3 4	Single Twin Triplet Quadruplet or higher
302	1	FILLER					
303	1	IMP_PLUR	Plurality Imputed		A,S	Blank 1	Plurality is not imputed Plurality is imputed
304-315	12	FILLER					

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
316	1	SEX	Sex of Infant		A,S	M F	Male Female
317	1	IMP_SEX	Sex Imputed Only where gestation is state to be 20 weeks or more	ed or presumed	A,S	Blank 1	Infant Sex not Imputed Infant Sex is Imputed
		Gestation					
318-319	2	DLMP_MM	Last Normal Menses Mont	h 429	A,S	01 02 03 04 05 06 07 08 09 10 11 12 99	January February March April May June July August September October November December Unknown or not stated
320-321	2	FILLER					
322-325	4	DLMP_YY	Last Normal Menses Year	431	A,S	nnnn 9999	Year of last normal menses Unknown or not stated
326-328	3	FILLER					
329	1	GEST_IMP	Gestation Imputed Flag		A,S	Blank 1	Gestation is not imputed Gestation is imputed
330	1	OBGEST_FLG	Obstetric Estimate of Gest	ation Used Flag	A,S	Blank 1	Obstetric Estimate is not used Obstetric Estimate is used
331-332	2	COMBGEST	Gestation – Detail in Week	s 404	A,S	02-47 99	2 nd through 47 th week of Gestation Unknown

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
333-334	2	GESTREC12	Gestation Recode 12	404	A,S	01 02 03 04 05 06 07 08 09 10 11	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks 28-31 weeks 32-33 weeks 34-36 weeks 37-38 weeks 39-40 weeks 41 weeks 42 weeks and over Unknown
335	1	GESTREC5	Gestation Recode 5	404	A,S	1 2 3 4 5	Under 20 weeks 20-23 weeks 24-27 weeks 28 weeks and over Unknown
336-337	2	OEGest_Unedt	Obstetric Gestation 373	A,S		0-98 02-47 99	Range of obstetric estimate of gestation - revised Range of clinical estimate of gestation - unrevised Unknown or not stated
338	1	FILLER					
339	1	COMBGEST_US	SED Combined Gesta	ntion Used Flag	A,S	blank 1	Combined gestation not used Combined gestation used
340-341	2	OEGest_Comb	Obstetric Estimate Edited (NCHS Standard item)	1 373 A,S		02-47 99	Weeks of gestation Not stated
342-343	2	OEGest_R12	Obstetric Estimate Recod (NCHS Standard item)	e12 373 A,S		01 02 03 04 05 06 07 08	Under 16 weeks 16-19 weeks 20-23 weeks 24-27 weeks 28-31 weeks 32-33 weeks 34-36 weeks 37-38 weeks

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description Reporting Flag Position		Vers*	Values	Definition
						09 10 11 12	39-40 weeks 41 weeks 42 weeks and over Unknown
344	1	OEGest_R5	Obstetric Estimate Recode (NCHS Standard item)	5 373 A,S		1	Under 20 weeks
			(FOLIS Standard Rein)			2 3 4 5	20-23 weeks 24-27 weeks 28 weeks and over Unknown
345-348	4	FILLER					
349-352	4	DBWT	Birth Weight – Detail in G	rams 442	A,S	0001-81 9999	65 Number of grams Not stated birth weight
353-354	2	BWTR14	Birth Weight Recode 14	442	A,S	01 02 03 04 05 06 07 08 09 10 11 12 13	0249 grams or less 0250 – 0349 grams 0350 – 0499 grams 0500 – 0999 grams 1000 – 1499 grams 1500 – 1999 grams 2000 – 2499 grams 2500 – 2999 grams 3000 – 3499 grams 3500 – 3999 grams 4000 – 4499 grams 4500 – 4999 grams 5000 – 8165 grams Unknown or Not Stated
355	1	BWTR4	Birth Weight Recode 4	442	A,S	1 2	1499 grams or less 1500 – 2499 grams

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S

¹⁹⁸⁹ Version of US Standard Birth Certificate
1989 Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition
						3 4	2500 - 8165 grams Unknown or not stated
356	1	FILLER					
357	1	ESTOFD	Estimated Time of Fetal D	eath 435	A	N L A U	At assessment, no labor At assessment, labor Labor, no assessment Unknown
358	1	AUTOPSY	Was Autopsy Performed?	436	A	Y N P	Yes No Planned
359	1	HISTOPF	Was Histological Placental	Exam Performed?	437 A	Y N P	Yes No Planned
360	1	AUTOPF	Was Autop/Hist results use	ed in cause? 438	A	Y N X	Yes No Not applicable
361-371	11	FILLER					
372-438		Flag File for Reporting flag	orting Flags s indented below follow this c	oding structure:		0 1	Not reporting Reporting
A	2003 V	ersion of US Stand	Education of Mother Obstetric Estimate of Gestat Tobacco Use Mother's Height Prepregnancy Weight WIC Prepregnancy Diabetes Gestational Diabetes Prepregnancy Hypertension Gestational Hypertension Hypertension Eclampsia ne following codes apply:		A A,S A A A A A A A A		
S s			lard Birth Certificate lard Birth Certificate but th	is element conform	ns to the	2003 star	ndards

			Flag Position	
383	1	F_RF_INFT In	nfertility Treatment	A
384	1	F_RF_INFT_DRG Fe		A
385	1		eproductive Technology	A
386	1		revious Cesarean	A
387	1		umber of Previous Cesareans	A
388	1	F MD PRESENT F	Fetal Presentation	A
389	1	F_MD_ROUTE F	Final Route and Method of Delivery	A
390	1		rial of Labor Attempted	A
391	1	F MM RUPTUR Ru		A
392	1	F MM ICU Ad	dmission to Intensive Care	A
393-402	10	FILLER		
403	1	F MPCB M	Ionth Prenatal Care Began	A
404	1	F CMBGST Co	ombined Gestation	A,S
405	1	F CIGS 0	Cigarettes Before Pregnancy	A
406	1	F CIGS 1	Cigarettes 1 st Trimester	A
407	1	F_CIGS_2	Cigarettes 2 nd Trimester	A
408	1	F_CIGS_3	Cigarettes 3 rd Trimester	A
409	1	F_FACILITY F	Facility	A
410	1	DelMethRecF De	elivery Method Recode Flag	A,S
411-423	13	FILLER		
424	1	F_MORIGIN M	Iother's Hispanic origin	A,S
425	1	F_MRACE_R N	Mother's Race (Revised)	A
426	1	F_FAGE_u Fa	ather's Age	A,S
427	1	F_DLLB_MM Las	st Live Birth Month (Revised)	A
428	1	FILLER		
429	1	F_DLMP_MM Las	st normal menses month	A,S
430	1	FILLER		
431	1	F_DLMP_YY Last	t normal menses year	A,S
432	1	F_MAGE Mother's		A,S
433	1	F_ATTENDANT At	ttendant	A,S
434	1	FILLER		
435	1		stimated Time of Fetal Death (Revised)	A
436	1		utopsy (Revised)	A
437	1		istological Placental Exam (Revised)	A
438	1		utopsy/Histological Exam Used in Cause	A
439	1		Iother's Bridged Race Recode	A,S
440	1		Number of previous live births living	A,S
441	1		Number of previous live births dead	A,S
442	1	F_DBWT B	Birthweight A,S	

Description

Reporting

Position

Len

Field

Vers* Values Definition

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

¹⁹⁸⁹ Version of US Standard Birth Certificate but this element conforms to the 2003 standards S

Position	Len	Field	Description	Reporting Flag Position	Vers*	Values	Definition	
443-2602 2164	FII	LLER						
2603-2642		40 The field	Coded Causes & Conditions s indented below follow this structu	ure:		A ICD10 Blank	Code Coded Value None reported	
2603-26 2608-26		5 35	IICOD Initiating Cause or Condition Code 265 FILLER		2651		•	
2643-2645	3	IC_124_	Fetal Initiating Fetal Recode	124 2651		A	001-124	
2646-2650 5	FILL	ER						
2651			e for Reporting Flags rting flags indented below follow the	his coding structure:		0 1	Not reporting Reporting	
2651	1	F_ICOD	Init Cause or condition	A				
2652-3150 499	F	ILLER						

3150 END OF RECORD

^{*} For the Version column (Vers) the following codes apply: A 2003 Version of US Standard Birth Certificate

S 1989 Version of US Standard Birth Certificate

¹⁹⁸⁹ Version of US Standard Birth Certificate but this element conforms to the 2003 standards

Table 1. Number of fetal deaths and fetal mortality rates: United States, 2014-2021

	Fetal deaths			Fetal m	nortality	rate\1
_	Total\2	20-27	28 weeks	Total\2	20-27	28 weeks
_	10041 (2	weeks\3	or more\3	10041 (2	weeks\3	or more\3
_						
_		Obste	etric estimat	e of gestati	on\4	
2021	21,105	10,824	10,281	5.73	2.95	2.80
2020	20,854	10,764	10,090	5.74	2.97	2.78
2019	21,478	11,216	10,262	5.70	2.98	2.73
2018	22,459	11,844	10,615	5.89	3.11	2.79
2017	22,827	11,861	10,966	5.89	3.07	2.84
2016	23,880	12,486	11,394	6.02	3.15	2.88
2015	23 , 776	12,407	11,369	5.94	3.11	2.85
2014	23,980	12 , 652	11,328	5.98	3.16	2.83
			LMP-based g	estation\5		
2014	23 , 999	11,948	12,051	5.98	2.99	3.01

 \label{lambda} IRate is number of fetal deaths in specified group per 1,000 live births and fetal deaths.

 $\2$ Fetal deaths with stated or presumed period of gestation of 20 weeks or more.

\3Not stated gestational age proportionally distributed.

 $\$ 4Beginning with the 2014 data year, the obstetric estimate of gestation at delivery replaces the measure based on the date of last normal menses as the standard for measuring gestational age. $\$ 5Date of last normal menses (LMP) was used for measuring gestational age.

Table 2. Fetal Deaths by Period of Gestation: United States, Each State and Territory, 2021

[Nonresidents of the United States are excluded]

State/territory of	Fetal Deaths with Stated or Presumed Gestation of 20 weeks or More					
residence	Total	20-27 Weeks	28 Weeks and Over	Not stated		
United States\1	21,105	10,747	10,208	15		
Alabama	510	249	259	:		
Alaska	57	24	33			
Arizona	485	237	248			
Arkansas	304	148	154	:		
California	2,243	1,124	1,088	3		
Colorado	312	164	148			
Connecticut	162	98	64			
Delaware	54	33	20			
District of Columbia	65	29	36			
Florida	1,594	815	732	4		
Georgia	1,016	565	443			
Hawaii	109	62	43			
Idaho	112	60	52			
Illinois	795	462	331			
Indiana	426	244	181			
Iowa	164	74	90			
Kansas	195	78	116			
Kentucky	329	159	169			
Louisiana	302	139	163			
Maine	67	32	33			
Maryland	457	255	202			
Massachusetts						
Michigan	290	131	159			
Minnesota	601	316	284			
Mississippi	349	179	169			
Missouri	355	197	157			
Montana	388	174	211			
	39	14	25			
Nebraska	99	51	48			
Nevada	280	147	133			
New Hampshire	73	28	45			
New Jersey	612	358	253			
New Mexico	84	31	53			
New York	654	316	335			
New York City	631	384	246			
North Carolina	652	300	346			
North Dakota	52	26	26			
Ohio	759	403	356			
Oklahoma	269	132	135			
Oregon	187	72	114			
Pennsylvania	777	375	401			
Rhode Island	56	29	26			
South Carolina	323	121	202			
South Dakota	76	47	29			
Tennessee	537	263	272			
Texas	1,457	701	750			
Utah	309	179	130			
Vermont	27	15	12			
Virginia	493	239	251			
Washington	451	260	178	1		
West Virginia	93	32	60			
Wisconsin	321	146	174			
Wyoming	53	30	23			
American Samoa						
Guam	31	11	20			
Northern Marianas						
Puerto Rico	242	144	94			
Virgin Islands						

NOTE: American Samoa did not report data for 2021. Data for the Northern Marianas and the Virgin Islands were excluded due to small numbers.

⁻ Quantity zero.

⁻⁻⁻ Data unavailable.

Table 3. Fetal Deaths by Period of Gestation and State/Territory of Occurrence: United States, 2021

	A	ll Fetal 1	Deaths	
State / territory	Total	Under 20 Weeks	20 Weeks and Over	Not stated
United States\1	42,428	19,921	21,040	1,46
Alabama	500	_	498	
Alaska	59	3	56	
Arizona	485	2	483	
Arkansas	532	237		
California	2,272	1	2,236	3
Colorado	330	_	329	
Connecticut	166	_	166	
Delaware	60	2	57	
District of Columbia	95	_	95	
Florida	1,628	3	1,577	4
Georgia	8,504	6 , 795		68
Hawaii	476	270	106	10
Idaho	129	27	102	
Illinois	781	1	779	
Indiana	428	1	425	
Iowa	161	1	160	
Kansas	188	_	186	
Kentucky	319	1	317	
Louisiana	336	37	299	
Maine	63	_	61	
Maryland	426	_	426	
Massachusetts	294	1	293	
Michigan	595	4	590	
Minnesota	337	_	337	
Mississippi	362	9	352	
Missouri	422	3	415	
Montana	39	-	39	
Nebraska	100	-	100	
Nevada	284	2	282	
New Hampshire	69	1	68	
New Jersey	601	11	589	
New Mexico	79	3	76	
New York	4,626	3,654	580	39
New York City	5,592	4,882	704	
North Carolina	659	-	653	
North Dakota	72	-	72	
Ohio	762	-	762	
Oklahoma	254	-	253	
Oregon	254	57	196	
Pennsylvania	1,163	392	771	
Rhode Island	509	408	63	3
South Carolina	301	4	297	
South Dakota	85	4	81	
Tennessee	579	6	570	
Texas	1,703	203	1,494	
Utah	322	-	322	
Vermont	31	-	31	
Virginia	3,469	2,873	490	10
Washington	469	14	442	1
West Virginia	103	8	93	
Wisconsin Wyoming	314 41	1	312 41	
	41	_	41	
American Samoa				
Guam Northern Marianas	32	1	31 	
Puerto Rico	243	_	239	
Virgin Islands	245		255	

NOTE: American Samoa did not report data for 2021. Data for the Northern Marianas and the Virgin Islands were excluded due to small numbers.

⁻ Quantity zero.

⁻⁻⁻ Data unavailable.

^{\1}Excludes data for the territories.

Table 4. Number and percentage of fetal deaths by race and Hispanic origin of mother: United States, 2021

Race -	Num	ber	Perce	ntage
Nace	Total	Non-Hispanic	Total\1	Non-Hispanic
All races\2	21,105	15 , 877	100.0	100.0
One race	20,672	15,507	97.9	97.7
White	13,639	9,196	64.6	57.9
Black	5,684	5,173	26.9	32.6
American Indian and Alaska Native (AIAN)	235	197	1.1	1.2
Asian	960	846	4.5	5.3
Native Hawaiian and Other Pacific Islander (NHOPI)	154	95	0.7	0.6
More than one race	433	370	2.1	2.3
Two races	399	341	1.9	2.1
Black and White	200	174	0.9	1.1
Black and AIAN	15	13	0.1	0.1
Black and Asian	11	8	0.1	0.1
Black and NHOPI	4	4	0.0	0.0
AIAN and White	75	62	0.4	0.4
AIAN and Asian	2	2	0.0	0.0
AIAN and NHOPI	1	1	0.0	0.0
Asian and White	65	59	0.3	0.4
Asian and NHOPI	7	5	0.0	0.0
NHOPI and White	19	13	0.1	0.1
Three races	31	26	0.1	0.2
Black, AIAN and White	10	8	0.0	0.1
Black AIAN and Asian	_	_	_	-
Black, AIAN and NHOPI	_	_	_	-
Black, Asian and White	8	7	0.0	0.0
Black, Asian and NHOPI	_	_	-	-
Black, NHOPI, and White	3	3	0.0	0.0
AIAN, Asian and White	_	_	_	-
AIAN, NHOPI and White	_	_	_	-
AIAN, Asian and NHOPI	_	_	_	_
Asian, NHOPI and White	10	8	0.0	0.1
Four races	3	3	0.0	0.0
Black, AIAN, Asian and White	3	3	0.0	0.0
Black, AIAN, Asian, and NHOPI	_	_	_	-
Black, AIAN, NHOPI and White	_	_	_	_
Black, Asian, NHOPI and White	_	_	_	-
AIAN, Asian, NHOPI and White	_	_	_	-
Five races	_	_	_	-
Black, AIAN, Asian, NHOPI and White	_	_	_	-

⁻ Quantity zero.

^{0.0} Quantity more than zero but less than 0.05.

^{\1} Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

^{\2} Race and Hispanic origin are reported separately on reports of fetal death. Persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards.

SOURCE: National Center for Health Statistics, National Vital Statistics System.

Table 5A. Number and percentage of fetal deaths by selected demographic and health characteristics: United States, 2021

more.]	Number	Percent
Day of week		
Total	21,105	
Sunday	2,287	
Monday	2,540	
Tuesday	3,238	
Wednesday	3,374	
Thursday	3,372	16.0
Friday	3,409	16.2
Saturday	2,885	13.7
Month of delivery		
Total	21,105	100.0
January	1,746	8.3
February	1,555	7.4
March	1,675	7.9
April	1,757	
May	1,726	8.2
June	1,790	8.5
July	1,747	
August	1,925	9.1
September	2,005	9.5
October	1,735	
November	1,674	
December	1,770	8.4
Place of delivery		
Total	21,105	100.0
Hospital\1	20,479	
Freestanding birthing center	26	0.1
Clinic or doctor's office	48	0.2
Residence	396	
Other	112	0.5
Not specified	44	0.2
Age of mother		
Total	21,105	100.0
<15	25	
15-19	1,065	
20-24	3,802	
25–29	5,532	
30-34	5 , 747	
35–39	3,741	
40-44	1,085	
45–49	95	
50-54	13	
Race and Hispanic origin of mother		
Total\2	21,105	100.0
Non-Hispanic, single race\3	21,100	100.0
Non-Hispanic white	9,196	43.6
Non-Hispanic black	5,173	
Non-Hispanic American Indian or Alaska Native	197	
Non-Hispanic Asian	846	
Non-Hispanic Asian Non-Hispanic Native Hawaiian or Other Pacific Islander	95	
Non-Hispanic multiple race	370	
Hispanic\4	4,290	
Unknown	4 , 290 938	4.4
OHAHOWH	938	4.4

Hispanic origin of mother Total	21,105	100.0
Non-Hispanic	15 , 877	75.2
Mexican	2 , 325	11.0
Puerto Rican	363	1.7
Cuban	118	0.6
Central or South American	825	3.9
Dominican	147	
Other and unknown Hispanic	512	2.4
Unknown	938	4.4
Education of mother\5		
Total	20,940	100.0
8th grade or less	717	3.4
9th-12th grade with no diploma	1,806	8.6
High school graduate\6	6,363	30.4
Some college credit, but no degree	3,267	15.6
Associate's degree\7	1,433	6.8
Bachelor's degree\8	2,910	13.9
Master's degree\9	1,225	5.9
Doctorate or professional degree\10	303	1.4
Unknown	2,916	13.9
Age of father\11	16 010	100 0
Total	16,918	100.0
<15	1	0.0
15-19	338	2.0
20-24 25-29	1,748 3,004	10.3 17.8
30-34	3,611	21.3
35-39	2,861	16.9
40-44	1,302	7.7
45-49	466	2.8
50-54	177	1.0
55-98	107	0.6
Unknown	3,303	19.5
	·	
Trimester prenatal care began\12		
Total	20,940	100.0
1st trimester	13,238	63.2
2nd trimester	2,889	13.8
3rd trimester	359	1.7
No care	1,718	8.2
Unknown	2,736	13.1
Mother received WIC food for herself during this pregnancy\13		
Total	20,244	100.0
Yes	3,684	18.2
No	13,309	
Not stated	3 , 251	16.1
Mohagaa waa during magnanara		
Tobacco use during pregnancy	01 105	100 0
Total Yes	21,105 1,632	100.0
No	17,776	84.2
NO Unknown	1,697	84.2
OHAHOWH	1, 091	0.0
Maternal pre-pregnancy body mass index\14		
Total	20,996	100.0
Underweight (BMI less than 18.5)	485	2.3
Normal weight (BMI of 18.5 to 24.9)	5 , 821	27.7
Overweight (BMI of 25.0 to 29.9)	4,852	23.1
Obese (BMI of 30.0 and over)	7,270	34.6
Not stated	2,568	12.2
	•	

Method of delivery\15		
Total	20,996	100.0
Vaginal (excludes vaginal after previous C-section)	14,282	68.0
Vaginal after previous C-section	1,665	7.9
Primary C-section	1,950	
Repeat C-section	1,295	
Vaginal (unknown if previous c-section)	662	3.2
C-section (unknown if previous c-section)	116	0.6
Not stated	1,026	4.9
Not Stated	1,020	4.9
Attendant at delivery		
Total	21,105	100.0
Doctor of medicine	17,564	83.2
Doctor of osteopathy	1,861	
Certified nurse midwife	703	
Other midwife	51	0.2
Other	566	2.7
Unspecified	360	1.7
Plurality		
Total	21,105	100.0
Singleton	19,591	92.8
Twin	1,430	6.8
Triplet	78	0.4
Quadruplet or higher	6	0.0
Sex		
Total	21,105	100.0
Male	10,992	
Female	10,113	47.9
Obstatnia agtimata of gostation		
Obstetric estimate of gestation Total	21,105	100.0
20-23 weeks	7,469	35.4
24-27 weeks	3,278	15.5
	2,756	
28-31 weeks 32-33 weeks	1,481	13.1 7.0
34-36 weeks	2,618	12.4
37-38 weeks	1,957	9.3
39-40 weeks	1,216	5.8
41 weeks	130	0.6
42 weeks and over	50	0.2
Unknown	150	0.7
	100	• • •
Birthweight\16		
Total	20,996	100.0
<500 grams	6,896	32.8
0500 - 0999 grams	3,671	17.5
1000 - 1499 gram	1,960	9.3
1500 - 1999 gram	1,770	8.4
2000 - 2499 gram	1,647	7.8
2500 - 2999 gram	1,468	7.0
3000 - 3499 gram	1,215	5.8
3500 - 3999 gram	566	2.7
4000 - 4499 gram	204	1.0
4500 - 4999 gram	83	0.4
5000 - 8165 gram	45	0.2
Unknown	1,471	7.0
0.0 Quantity more than zero but less than 0.05		

^{0.0} Quantity more than zero but less than 0.05.

 $\label{lincludes}$ fetal deaths occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

⁻ Quantity zero.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death. \4Includes all persons of Hispanic origin of any race.

 $\$ 5Hawaii and Rhode Island reported education of mother for 2021 but had an unknown level of 50% or higher and were flagged as not reporting.

\6Includes General Educational Development (GED).

\7Includes Associate in Arts and Associate in Science.

\8Includes Bachelor in Arts and Bachelor in Science.

\9Includes Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\10Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

\11Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New Mexico, New York (including New York City), Virginia, and Wisconsin reported father's age for 2021 but had an unknown level of 50% or higher and were flagged as not reporting.

 $\12$ Hawaii and Rhode Island reported date of first prenatal care for 2021 but had unknown levels of 50% or higher and were flagged as not reporting.

\13Mother received WIC food for herself durng this pregnancy is not reported by New York City. The District of Columbia, Hawaii, and Rhode Island reported WIC for 2021 but had an unknown level of 50% or higher and were flagged as not reporting.

 $\14$ Hawaii reported maternal pre-pregnancy body mass index for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

 $\15$ Hawaii reported method of delivery for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\16Hawaii reported birthweight for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

Table 5B. Number and percentage of fetal deaths by selected demographic and health characteristics: Guam and Puerto Rico, 2021

or more.	Number	Percent
Day of week	0.7.5	100.0
Total	273	
Sunday	24	
Monday	35	
Tuesday	43	
Wednesday	50	
Thursday	41	
Friday	39	
Saturday	41	15.0
Month of delivery		
Total	273	
January	33	
February	11	
March	36	
April	24	
May	19	
June	19	
July	21	
August	24	
September	19	
October	28	
November	26	
December	13	4.8
Place of delivery		
Total	273	
Hospital\1	264	
Freestanding birthing center	1	0.4
Clinic or doctor's office	- -	
Residence	2	
Other	6	2.2
Not specified	-	_
Age of mother		
Total	273	100.0
<15	-	
15-19	21	
20-24	60	
25–29	73	
30-34	64	
35–39	43	
40-44	12	
45-49	-	
50-54	-	_
Race and Hispanic origin of mother		
Total\2	273	100.0
Non-Hispanic, single race\3		
Non-Hispanic white	g	
Non-Hispanic black	2	
Non-Hispanic American Indian or Alaska Native	-	
Non-Hispanic Asian	2	
Non-Hispanic Native Hawaiian or Other Pacific Islander	25	
Non-Hispanic multiple race	2	
Hispanic\4	233	85.3
Unknown	-	_

Wisnessia suinin of method		
Hispanic origin of mother Total	273	100.0
Non-Hispanic	40	14.7
Mexican	_	_
Puerto Rican	227	83.2
Cuban	_	_
Central or South American	3	1.1
Dominican	3	1.1
Other and unknown Hispanic	-	-
Unknown	_	_
Education of mother		
Total	273	100.0
8th grade or less	6	2.2
9th-12th grade with no diploma	25	9.2
High school graduate\5	86	31.5
Some college credit, but no degree	27	9.9
Associate's degree\6	44	16.1
Bachelor's degree\7 Master's degree\8	54 21	19.8 7.7
Doctorate or professional degree\9	5	1.8
Unknown	5	1.8
	-	_,,
Age of father\10		
Total	242	100.0
<15	-	-
15-19	3	1.2
20-24 25-29	41 43	16.9 17.8
30-34	43 27	11.2
35–39	33	13.6
40-44	26	10.7
45-49	7	2.9
50-54	-	_
55-98	2	0.8
Unknown	60	24.8
Trimester prenatal care began		
Total	273	100.0
1st trimester	187	68.5
2nd trimester	42	15.4
3rd trimester	10	3.7
No care	21	7.7
Unknown	13	4.8
Mother received WIC food for herself during this pregnancy		
Total	273	100.0
Yes	165	60.4
No	86	31.5
Not stated	22	8.1
Tobacco use during pregnancy		
Total	273	100.0
Yes No	3 257	1.1 94.1
Unknown	13	4.8
	10	1.0
Maternal pre-pregnancy body mass index		
Total	273	100.0
Underweight (BMI less than 18.5)	18	6.6
Normal weight (BMI of 18.5 to 24.9)	75	27.5
Overweight (BMI of 25.0 to 29.9)	61	22.3
Obese (BMI of 30.0 and over) Not stated\3	103 16	37.7 5.9
	Τ.0	J • J

Method of delivery		
Total	273	100.0
Vaginal (excludes vaginal after previous C-section)	190	69.6
Vaginal after previous C-section	21	7.7
Primary C-section	31	11.4
Repeat C-section	13	4.8
Vaginal (unknown if previous c-section)	10	3.7
C-section (unknown if previous c-section)	3	1.1
Not stated	5	1.8
Not Stated	J	1.0
Attendant at delivery		
Total	273	100.0
Doctor of medicine	261	95.6
Doctor of osteopathy	5	1.8
Certified nurse midwife	2	0.7
Other midwife	-	_
Other	_	_
Unspecified	5	1.8
Plurality		
Total	273	100.0
Singleton	258	94.5
Twin	12	4.4
Triplet	3	1.1
Quadruplet or higher	_	_
Corr		
Sex Total	273	100.0
Male	144	52.7
Female	129	47.3
remare	129	47.5
Obstetric estimate of gestation		
Total	273	100.0
20-23 weeks	110	40.3
24-27 weeks	45	16.5
28-31 weeks	31	11.4
32-33 weeks	24	8.8
34-36 weeks	16	5.9
37-38 weeks	24	8.8
39-40 weeks	18	6.6
41 weeks	1	0.4
42 weeks and over	-	-
Unknown	4	1.5
Birthweight		
Total	273	100.0
<500 grams	79	28.9
0500 - 0999 grams	75	27.5
1000 - 1499 gram	29	10.6
1500 - 1999 gram	23	8.4
2000 - 2499 gram	13	4.8
2500 - 2999 gram	19	7.0
3000 - 3499 gram	11 7	4.0
3500 - 3999 gram 4000 - 4499 gram	•	2.6 1.8
4500 - 4499 gram 4500 - 4999 gram	5 -	Τ.δ
4300 - 4999 gram 5000 - 8165 gram	_	_
Unknown	12	4.4
0.0 Quantity more than zero but less than 0.05	14	4.4

^{0.0} Quantity more than zero but less than 0.05.

 $\label{lincludes}$ fetal deaths occurring en route to or on arrival at hospital.

\2Includes fetal deaths to race and origin groups not shown separately, such as Hispanic, single-race white, Hispanic, single-race black, and non-Hispanic, multiple-race women, and fetal deaths with origin not stated.

⁻ Quantity zero.

\3Race and Hispanic origin are reported separately on reports of fetal death; persons of Hispanic origin may be of any race. In this table, non-Hispanic women are classified by race. Race categories are consistent with the 1997 Office of Management and Budget standards. Single-race is defined as only one race reported on the report of fetal death.

\4Includes all persons of Hispanic origin of any race.

\5Includes General Educational Development (GED). \6Includes Associate in Arts and Associate in Science.

 $\Time 1$ 7Includes Bachelor in Arts and Bachelor in Science.

 \S Includes Master in Arts, Master in Science, Master of Engineering, Master of Education, Master of Social Work, and Master of Business Administration.

\9Includes Doctor of Philosophy, Doctor of Education, Doctor of Medicine, Doctor of Dental Surgery, Doctor of Veterinary Medicine, Doctor of Laws, and Juris Doctor.

 $\10\$ Guam reported father's age for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

Table 6A. Number and percentage of fetal deaths by selected health characteristics: United States, 2021

Risk factor	Total	Number reported	Unknown	Per 1,000
Diabetes				
Prepregnancy (Diagnosis prior to this pregnancy)	21,105	816	1,151	40.9
Gestational (Diagnosis in this pregnancy)	21,105	1,013	1,151	50.8
Hypertension				
Prepregnancy (Chronic)	21,105	1,324	1,151	66.4
Gestational (PIH, preeclampsia)	21,105	1,481	1,151	74.2
Eclampsia	21,105	129	1,151	6.5
Pregnancy resulted from infertility treatment	21,105	595	1,151	29.8
Mother had a previous cesarean delivery\1	21,105	3,032	1,151	15.2
Maternal morbidity\2				
Ruptured uterus	21,105	144	1,152	721.7
Admissioin to intensive care unit	21,105	477	1,152	2,390.6

 $^{\1}$ Mother had a previous cesarean delivery per 100 fetal deaths.

NOTE: PIH is pregnancy-induced hypertension.

^{\2} Maternal morbidities per 100,000 fetal deaths.

Table 6B. Number and percentage of fetal deaths by selected health charcteristics: Guam and Puerto Rico, 2021

Risk factor	Total	Number reported	Unknown	Per 1,000
Diabetes				
Prepregnancy (Diagnosis prior to this pregnancy)	273	14	13	53.8
Gestational (Diagnosis in this pregnancy)	273	10	13	38.5
Hypertension				
Prepregnancy (Chronic)	273	18	13	69.2
Gestational (PIH, preeclampsia)	273	16	13	61.5
Eclampsia	273	-	13	-
Pregnancy resulted from infertility treatment	273	1	13	3.8
Mother had a previous cesarean delivery\1	273	35	13	13.5
Maternal morbidity\2				
Ruptured uterus	273	5	19	1,968.5
Admissioin to intensive care unit	273	5	19	1,968.5

⁻ Quantity zero.

NOTE: PIH is pregnancy-induced hypertension.

 $^{\1}$ Mother had a previous cesarean delivery per 100 fetal deaths.

^{\2} Maternal morbidities per 100,000 fetal deaths.

Table 7A. Deaths according to 124 selected causes of fetal death: 41 states and the District of Columbia, 2021

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.] Cause of death (based on ICD-10) Number All causes¹..... 15,212 Human immunodeficiency virus (HIV) disease......B20-B24 Other and unspecified infectious and parasitic diseases........A00-A49,A51-A79,B35-B99 Umbilical hernia......K42 Other hernia.....K40-K41,K43-K46 Other and unspecified diseases of digestive system......K00-K38,K50-K92 13,559 Fetus affected by maternal conditions that may be unrelated to present pregnancy...P00 1,915 825 26 479 Fetus affected by other maternal circulatory and respiratory diseases............P00.3 Fetus affected by maternal injury......P00.5 Fetus affected by other medical procedures and maternal conditions......P00.7-P00.8 1,804 263 Fetus affected by incompetent cervix......P01.0 1,130 Fetus affected by oligohydramnios......P01.2 Fetus affected by polyhydramnios......P01.3 42 Fetus affected by ectopic pregnancy......P01.4 239 Fetus affected by maternal death......P01.6 Fetus affected by malpresentation before labor......P01.7 36 3,688 Fetus affected by placenta previa......P02.0 30 Fetus affected by other forms of placental separation and hemorrhage............P02.1 1,332 688 89 86 Fetus affected by prolapsed cord......P02.4 786 278 Fetus affected by other and unspecified conditions of umbilical cord............P02.6 390 Fetus affected by other and unspecified abnormalities of membranes......P02.8-P02.9 138 Fetus affected by forceps delivery......P03.2 107 185 179 221 Disorders related to short gestation and low birth weight, not elsewhere classified......P07 Extremely low birth weight or extreme immaturity......P07.0,P07.2 123 Other low birth weight and preterm......P07.1,P07.3 98 Exceptionally large size and other heavy for gestational age fetus......P08.0-P08.1 Other birth trauma......P12-P15 Intrauterine hypoxia and birth asphyxia......P20-P21 Birth asphyxia......P21 Other respiratory conditions originating in the perinatal period....P22.8-P22.9,P23-P28 Congenital pneumonia......P23 Infections specific to the perinatal period......P35-P39 Congenital cytomegalovirus infection......P35.1 Congenital viral hepatitis......P35.3 Bacterial sepsis......P36 Other infections specific to the perinatal period.....P35.8-P35.9,P37.2-P37.9,P38-P39 Hematological disorders......P60-P61 432 432 Syndrome of infant of a diabetic mother and neonatal diabetes mellitus....P70.0-P70.2 Digestive system disorders of fetus......P76-P78 4,945 156 Fetal death of unspecified cause......P95 4,765 Termination of pregnancy......P96.4 24 1,503 295 Anencephaly and similar malformations......Q00 124 17 Microcephaly......Q02 Congenital hydrocephalus......Q03 Reduction deformities of brain......Q04.0-Q04.3 Other congenital malformations of brain......Q04.4-Q04.9 Spina bifida......Q05 Other congenital malformations of spinal cord and nervous system.....Q06-Q07 18 Congenital malformations of heart......Q20-Q24 188 47 10 25 105 58 17 30 Other congenital malformations of urinary system......Q62-Q64 151 Other congenital malformations......Q86-Q89 121 Conjoined twins......Q89.4 56 58 Chromosomal abnormalities, not elsewhere classified.......Q90-Q99 545 113 Edward's syndrome......Q91.0-Q91.3 189 53 190 External causes of mortality......*U01,V01-Y84 11 Assault (homicide)......*U01,X85-Y09 - Quantity zero.

NOTE: Excludes data for jurisdictions (Arizona, Georgia, Hawaii, Mississippi, New York (including New York City), North Dakota, Vermont, and Wisconsin) for which the cause of death was unspecified (P95) for 50% or more of records. Also excludes data for California which did not report cause of death based on the 2003 revision of the U.S. Standard Report of

Table 7B. Deaths according to 124 selected causes of fetal death: Puerto Rico, 2021

causes ¹ ain infectious and parasitic diseases	A00-B99	
Human immunodeficiency virus (HIV) disease	B20-B24	
Other and unspecified infectious and parasitic diseases	A00-A49,A51-A79,B35-B99	
	nknown behavior	
crine, nutritional and metabolic diseases	E34.3	
Cystic fibrosis Other endocrine, nutritional and metabolic diseases	E00-E32	E34.0-E34.2,E34.4-E34.9,E40-E83,E85-E88
ngitisr diseases of nervous system and sense organs		
lical herniar herniar and unspecified diseases of digestive system	K40-K41,K43-K46	
	alies, and symptoms, signs and il	-defined conditions
Fetus affected by maternal conditions that may be unrelated to proceed to be affected by maternal hypertensive disorders	present pregnancyP00	
Fetus affected by maternal renal and urinary tract diseases Fetus affected by maternal infectious and parasitic diseases	P00.1	
Fetus affected by other maternal circulatory and respiratory Fetus affected by maternal nutritional disorders	y diseasesP00.3	
Fetus affected by maternal injury Fetus affected by surgical procedure on mother	P00.5	
Fetus affected by other medical procedures and maternal conference fetus affected by unspecified maternal condition		
Fetus affected by maternal complications of pregnancy Fetus affected by incompetent cervix		
Fetus affected by premature rupture of membranes Fetus affected by oligohydramnios		
Fetus affected by polyhydramnios Fetus affected by ectopic pregnancy		
Fetus affected by multiple pregnancy Fetus affected by maternal death		
	ons of pregnancy	p01.8-p01.9
Fetus affected by complications of placenta, cord and membranes Fetus affected by placenta previa	P02.0	
Fetus affected by other forms of placental separation and he Fetus affected by other and unspecified morphological and further fetus affected by placental transfusion syndromes	unctional abnormalities of placent	a
Fetus affected by prolapsed cord	P02.4	
Fetus affected by other and unspecified conditions of umbil: Fetus affected by chorioamnionitis	ical cord	
Fetus affected by other and unspecified abnormalities of mer Fetus affected by other complications of labor and delivery	P03	
Fetus affected by breech delivery and extraction Fetus affected by other malpresentation, malposition and dispetus affected by forceps delivery	sproportion during labor and deliv	rery
Fetus affected by delivery by vacuum extractor (ventouse) Fetus affected by cesarean delivery	P03.3	
Fetus affected by precipitate delivery Fetus affected by abnormal uterine contractions	P03.5	
s affected by noxious influences transmitted via placenta	P04	p03.8-p03.9
fetal growth and fetal malnutritionrders related to short gestation and low birth weight, not elsew Extremely low birth weight or extreme immaturity	where classified	p07
Other low birth weight and preterm		
Exceptionally large size and other heavy for gestational age fer Post-term, not heavy for gestational age fetus	tusP08.0-P08.1	
	other injuries to central nervous	systemP10-P11
ther birth trauma	P20-P21	
Intrauterine hypoxia first noted before onset of labor Intrauterine hypoxia first noted during labor and delivery Intrauterine hypoxia, unspecified	P20.1	
Birth asphyxiar respiratory conditions originating in the perinatal period	P21	
Congenital pneumonia	P23	
Atelectasis	P28.0-P28.1	P25
Other respiratory system disorders	P35-P39	
Congenital cytomegalovirus infection	P35.1	
Congenital viral hepatitis	P36	
Congenital tuberculosis	P37.1	
hemorrhage	P50-P54	
Rh isoimmunization of fetus	p55.0	
Other hemolytic disease of fetus	P56	
natal jaundice tological disorders sitory endocrine and metabolic disorders specific to fetus	P60-P61	
sitory endocrine and metabolic disorders specific to fetus Syndrome of infant of a diabetic mother and neonatal diabetes me Other transitory endocrine and metabolic disorders specific to	ellitusP70.0-P70.2	
stive system disorders of fetus r conditions originating in the perinatal period	P76-P78	
Hydrops fetalis not due to hemolytic disease Fetal death of unspecified cause	P83.2	
Withdrawal symptoms from maternal use of drugs of addiction Termination of pregnancy	P96.4	
Complications of intrauterine procedures, not elsewhere classificable of the perinatal perination origination originating in the perinatal period, unspecified	iod	29, P83.0-P83.1, P83.3-P83.9, P91, P94, P96.0, P96.3, P96.8
condition originating in the perinatal period, unspecified enital malformations, deformations and chromosomal abnormalities Congenital malformations of nervous system	sQ00-Q99	
Anencephaly and similar malformations Encephalocele	Q00	
Microcephaly Congenital hydrocephalus	Q03	
Reduction deformities of brain	Q04.4-Q04.9	
Spina bifida Other congenital malformations of spinal cord and nervous systems. Congenital malformations of eye, ear, face and neck	ystemQ06-Q07	
Congenital malformations of eye, ear, race and neck Other congenital malformations of circulatory system		
Congenital malformations of lung	Q33 Q30-Q32,Q34	
Congenital malformations of digestive system	Q50-Q56	
Congenital malformations of urinary system Renal agenesis and other reduction defects of kidney Cystic kidney disease	Q60	
Cystic kidney disease Other congenital malformations of urinary system ongenital malformations and deformations of musculoskeletal system	Q62-Q64	Q65-Q85
ther congenital malformations	Q86-Q89	
Multiple congenital malformations, not elsewhere classified All other congenital malformations		
Chromosomal abnormalities, not elsewhere classified Down's syndrome	Q90-Q99	
Edward's syndrome	Q91.4-Q91.7	
Other chromosomal abnormalities, not elsewhere classified toms, signs and abnormal clinical and laboratory findings, not ernal causes of mortality	elsewhere classified	
Accidents (unintentional injuries)	v01-x59	
Assault (homicide)		

NOTE: Excludes data for Guam, for which the cause of death was unspecified (P95) for 50% or more of records. American Samoa did not report data for 2021 and data for the Northern Marianas and the Virgin Islands were excluded due to small numbers.

Source: National Center for Health Statistics, National Vital Statistics System.

MOTHER	MOTHER'S EDUCATION (Check the box that best describes the highest degree or level of school completed at the time of delivery)	the box mother	ER OF HISPANIC OR x that best describes w r is Spanish/Hispanic/L ox if mother is not Spa	hether the atina. Check the	е	what the	he mother cor	nsiders he	e or more races to indica rself to be)	te
	□ 8th grade or less	□ No, no	ot Spanish/Hispanic/La	tina		□ Ame	rican Indian or	Alaska Na		
	□ 9th - 12th grade, no diploma	□ Yes, M	Mexican, Mexican Ame	rican, Chicana			ne of the enroll n Indian	ea or princ	cipai tribe)	
	☐ High school graduate or GED	□ Yes, F	Puerto Rican			□ Chine				
	completed	□ Yes, 0	Cuban			☐ Filipin☐ Japar				
	☐ Some college credit but no degree	□ Yes, o	other Spanish/Hispanic	/Latina		□ Korea				
	☐ Associate degree (e.g., AA, AS)	(Specify	')			□ Vietna	amese Asian (Specif	(1)		
	□ Bachelor's degree (e.g., BA, AB, BS)						Hawaiian	y)		
	 Master's degree (e.g., MA, MS, MEng, MEd, MSW, MBA) 						nanian or Char	norro		
	□ Doctorate (e.g., PhD, EdD) or Professional degree (e.g., MD, DDS, DVM, LLB, JD)					□ Other		er (Specify	y)	
	, ,	DATE OF FIR	RST PRENATAL CARE	VISIT			PRENATAL		TAL NUMBER OF PREI	
	(At delivery, conception, or anytime between) □ Yes □ No	1	/	renatal Care	CARI	E VISIT	,		SITS FOR THIS PREGN (If none, enter	
	M		YYYY	MOTHER'S WE	M M		YYYY 28 DID MOT		WIC FOOD FOR HERS	
	(feet/inches)	(pounds)			ounds)		-		EGNANCY? Yes	
	LIVE BIRTHS	OUTCOMES (s	THER PREGNANCY pontaneous or induced	For each	time period	, enter eith			NANCY tes or the number of pac	cks of
		losses or ectopi Other Outcom		cigarettes	s smoked.	IF NONE,	ENTER "0".			
		mber (Do not in	nclude this fetus)	Average nu	ımber of ciga	arettes or p	acks of cigare	ttes smoke of cigarett		
		•		Three Mon	ths Before F Months of	Pregnancy			OR OR	
	□ None □ None □	None		Second Th	ree Months	of Pregnar			OR	
	29c. DATE OF LAST LIVE BIRTH 30I	b. DATE OF L		32. DATE L		AL 33. F	PLURALITY - S		OR 34. IF NOT SINGLE BI	
	MM YYYY		CY OUTCOME /	/_	ES BEGAN		win, Triplet, e	ic.	Born First, Second, Th	nird, etc.
		MM	YYYY	MM DD		` '	cify)		(Specify)	
	35. MOTHER TRANSFERRED FOR MATE IF YES, ENTER NAME OF FACILITY I			TIONS FOR DE	LIVERY?	Yes	No No			
MEDICAL	36. RISK FACTORS IN THIS PREGNANCY	(Check all that	apply):						ND/OR TREATED DUR	RING
	Diabetes					TH	IIS PREGNAN	CY (Chec	k all that apply)	
AND	 Prepregnancy (Diagnosis prior to this Gestational (Diagnosis in this pregress) 	s pregnancy) nancy)				□ Go	onorrhea			
HEALTH	Hypertension					□ Sy	/philis			
NFORMATION	□ Prepregnancy (Chronic) □ Gestational (PIH, preeclampsia)					□ Cl	nlamydia			
	□ Eclampsia					□ Lis	steria			
	□ Previous preterm birth					□ Gi	roup B Strepto	coccus		
	 Other previous poor pregnancy outcome growth restricted birth) 	(Includes perina	atal death, small-for-ge	stational age/in	trauterine	□ Cy	ytomegalovirus	3		
	,	ent-If yes shor	ok all that apply:			□ Pa	arvovirus			
	□ Pregnancy resulted from infertility treatm	•	л ан шасарру.			п То	oxoplasmosis			
	 Fertility-enhancing drugs, Artificial ins Intrauterine insemination 	semination or				□ No	one of the abo	ve		
	□ Assisted reproductive technology (e.g	g., in vitro fertiliz	zation (IVF), gamete int	rafallopian trans	sfer (GIFT))	□ Of	ther (Specify)_			
	Mother had a previous cesarean delivery If yes, how many	/								
	□ None of the above									
	38. METHOD OF DELIVERY		39. MATERNAL MOI	RBIDITY (Check	k all that app	oly) 40.	CONGENITA	L ANOMA	LIES OF THE FETUS	
	A. Was delivery with forceps attempted but to	unsuccessful?	(Complications asso	ociated with labo	or and delive	ery)	(CI	neck all tha	at apply)	
	□ Yes □ No		 Maternal transfu 	sion			Anencephaly Meningomyel	ocele/Spir	na bifida	
	B. Was delivery with vacuum extraction atte unsuccessful?	empted but	□ Third or fourth d	egree perineal la	aceration		Cyanotic cong Congenital di			
	□ Yes □ No		 Ruptured uterus 				Omphalocele Gastroschisis			
	C. Fetal presentation at delivery		 Unplanned hyste 	erectomy			Limb reductio	n defect (e	excluding congenital g syndromes)	
	□ Cephalic		□ Admission to inte	ensive care unit			Cleft Lip with	or without		
	□ Breech □ Other		□ Unplanned open	ating room proc	edure follow		Cleft Palate a	me	ad.	
	D. Final route and method of delivery (Chec	k one)	delivery				□ Karyotyp	e pending		
	□ Vaginal/Spontaneous □ Vaginal/Forceps		□ None of the abo	ve			Suspected cl	e confirme	ed	
	□ Vaginal/Vacuum □ Cesarean	-410					 Karyotyp Hypospadias 			
	If cesarean, was a trial of labor attemp □ Yes □ No	piea?					None of the	anomalies	listed above	

E. Hysterotomy/Hysterectomy

□ Yes □ No

Table A. Period of gestation and weight minimums at which fetal death reporting is required, by reporting areas: United States, 2021

States, 2021								
Area	All periods of gestation	12 weeks	16 weeks	20 weeks	20 weeks or 350 grams	20 weeks or 400 grams	20 weeks or 500 grams	350 grams
								<u> </u>
Alabama				Х				<u> </u>
Alaska				Х				
Arizona					X			
Arkansas		X						
California				Х				
Colorado	X^1							
Connecticut				X				
Delaware								² X
District of Columbia							X	
Florida				X				
Georgia	X							
Hawaii	Х							
Idaho					X			
Illinois				X				
Indiana				X				
Iowa				X				
Kansas				X				
Kentucky					X			
Louisiana					X			
Maine				Х				
Maryland				X ³				
Massachusetts				71	Х			
Michigan					77	Х		
Minnesota				Х		27		
Mississippi				77	Х			
Missouri					X			
					A			X ²
Montana								X
Nebraska				X				
Nevada				X	37			
New Hampshire				V	X			-
New Jersey				X				-
New Mexico				X ⁴				
New York	X							
New York (excluding New York City)	X							
New York City	Х							
North Carolina				X				
North Dakota				Х				
Ohio		5		Х				<u> </u>
Oklahoma		X ⁵						
Oregon				Х				ļ
Pennsylvania			Х					
Rhode Island	X							ļ
South Carolina					X			ļ
South Dakota				Х				
Tennessee					X			
Texas				X				ļ
Utah				X				
Vermont	<u> </u>			X ⁶				<u> </u>
Virginia	Х							
Washington				Х				
West Virginia				Х				
Wisconsin					X			
Wyoming				X				
American Samoa	Х							
Guam					X			
Northern Marianas								X ²
Puerto Rico					X	 		<u> </u>
Virgin Islands	Х					†		
g		<u> </u>	ļ.	!	<u> </u>	<u> </u>	I	

¹ Although State law requires the reporting of fetal deaths of all periods of gestation, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

² If weight is unknown, 20 completed weeks of gestation or more.

³ If gestational age is unknown, weight of 500 grams or more.

⁴ If gestational age is unknown, weight of 350 grams or more.

⁵ Although State law requires the reporting of fetal deaths at 12 weeks of gestation or more, only data for fetal deaths of 20 weeks of gestation or more are provided to NCHS.

⁶ If gestational age is unknown, weight of 400 grams or more, 15 ounces or more.

Table B. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2021

[By place of residence]

[Fetal deaths include only those with stated or presumed period of gestation of 20 weeks or more.]

Reporting areas	Total Fetal deaths	Time of birth	Mother's birthplace	Education of mother ²	Father's age ³	Mother's Hispanic origin
Total of reporting areas ¹	21,105	1.9	5.1	13.9	19.5	4.4
Alabama	510	0.2	0.2	3.5	62.4	_
Alaska	57	1.8	7.0	10.5	24.6	8.8
Arizona	485	1.4	0.4	3.1	11.1	1.2
Arkansas	304	1.6	3.6	11.8	52.3	1.0
California	2,243	1.4	4.1	12.1	11.7	5.6
Colorado	312	_	17.0	26.3	11.2	
Connecticut	162	9.3	2.5	26.5	14.2	
Delaware	54	_	1.9	14.8	24.1	_
District of Columbia	65	9.2	13.8	46.2	63.1	_
Florida	1,594	2.7	7.8	9.1	15.4	7.0
Georgia	1,016	0.7	0.4	22.8	59.6	1.7
Hawaii	109	36.7	32.1	56.0	52.3	47.7
Idaho	112	1.8	3.6	3.6	13.4	0.9
Illinois	795	3.3	7.5	16.7	17.1	0.4
						0.4
Indiana	426	_	1.6	2.8	8.7	- 0 0
Iowa	164	-	- 1 5	17.1	14.0	0.6
Kansas	195	-	1.5	5.6	7.7	5.1
Kentucky	329	_	0.3	5.2	41.3	0.3
Louisiana	302	_	1.3	9.6	23.8	0.3
Maine	67	4.5	-	7.5	9.0	-
Maryland	457	0.7	7.0	19.0	18.6	7.9
Massachusetts	290	1.4	25.2	31.7	31.7	16.9
Michigan	601	0.7	3.3	5.7	20.5	2.2
Minnesota	349	1.1	3.7	5.4	9.5	2.9
Mississippi	355	0.8	2.5	9.3	45.9	0.6
Missouri	388	0.3	1.8	4.6	16.0	1.8
Montana	39	_	_	_	10.3	_
Nebraska	99	1.0	1.0	28.3	8.1	_
Nevada	280	_	2.5	5.7	12.5	_
New Hampshire	73	2.7	4.1	11.0	21.9	11.0
New Jersey	612	1.6	1.5	8.3	29.6	3.3
New Mexico	84	1.2	11.9	7.1	45.2	4.8
New York (excluding New York City)	654	8.7	1.7	35.9	57.8	19.1
New York City	631	4.1	8.6	27.6	46.8	11.3
North Carolina	652	_	1.8	3.4	38.0	0.3
North Dakota	52	_	11.5	9.6	19.2	
Ohio	759	0.5	1.7	3.0	11.7	0.8
Oklahoma	269	0.4	3.7	8.2	16.4	5.9
	187	0.4	3.2		13.4	2.7
Oregon				3.7		
Pennsylvania	777	3.2	3.1	10.7	12.6	-
Rhode Island	56	5.4	23.2	83.9	37.5	8.9
South Carolina	323	-	5.9	6.5	25.7	0.9
South Dakota	76	-	2.6	6.6	10.5	5.3
Tennessee	537	1.1	10.8	16.6	44.5	1.3
Texas	1,457	-	9.3	29.2	29.2	7.8
Utah	309	1.3	1.0	8.4	7.8	-
Vermont	27	3.7	3.7	7.4	7.4	-
Virginia	493	3.9	1.2	29.8	62.5	4.5
Washington	451	4.9	20.2	15.3	16.2	11.8
West Virginia	93	-	-	3.2	15.1	-
Wisconsin	321	0.9	0.3	10.9	49.8	1.6
Wyoming	53	-	-	7.5	20.8	3.8
American Samoa						
Guam	31	-	3.2	12.9	77.4	-
Northern Marianas						
Puerto Rico	242	0.4	-	0.4	24.8	-
Virgin Islands						

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2021
[By place of residence]

Reporting areas	Place of delivery	Attendant at delivery	Month prenatal care began\4	Mother's height\5	Mother's pre- pregnancy weight\6	Did mother get WIC food for herself during this pregnancy\7	Live-birth order\8
Total of reporting areas ¹	0.2	1.7	13.1	5.9	11.1	16.1	5.1
Alabama	_	_	2.5	2.2	3.5	3.9	-
Alaska	_	-	8.8	5.3	12.3	15.8	5.3
Arizona	_	1.4	3.1	3.7	6.0	5.2	3.9
Arkansas	-	-	5.6	2.6	12.8	4.3	2.0
California	1.1	0.0	14.8	7.5	15.9	21.2	7.0
Colorado	-	2.2	46.8	19.2	27.6	38.8	17.9
Connecticut	_	0.6	12.3	3.7	8.0	22.8	3.1
Delaware	-	9.3	7.4	3.7	7.4	38.9	1.9
District of Columbia	-	-	16.9	18.5	41.5	47.7	3.1
Florida	-	0.1	13.3	8.8	12.9	18.1	6.7
Georgia	0.1	_	17.2	3.0	9.4	22.9	6.5
Hawaii	1.8	_	68.8	50.5	63.3	69.7	58.7
Idaho	_	-	4.5	2.7	4.5	6.3	3.6
Illinois	0.5	1.1	12.1	4.0	14.6	2.8	2.3
Indiana	_	44.4	13.6	6.6	8.7		1.9
Iowa	_	_	1.2	_	5.5		0.6
Kansas	_	_	3.1	2.6	5.1		_
Kentucky	_	0.3	4.3	_	3.6		_
Louisiana	_	_	7.0	1.0	7.6		0.3
Maine	_	1.5	19.4	4.5	23.9		3.0
Maryland	_	-	18.4	4.6	9.0		2.0
Massachusetts	_	_	5.2	3.8	12.1		2.8
Michigan	0.2	_	5.5	1.0	5.7		3.0
Minnesota	-	2.0	2.6	4.3	7.4		0.6
Mississippi	_	0.8	9.9	1.7	4.5		1.1
Missouri	_	-	10.6	1.0	3.6		1.0
Montana	_	_	2.6	2.6	5.1		-
Nebraska	_	_	2.0	4.0	8.1		_
Nevada	_	_	5.0	4.6	5.7	2.9	0.7
New Hampshire	_	2.7	8.2	2.7	4.1		2.7
New Jersey	_	0.2	10.9	2.6	9.6		0.5
New Mexico		0.2	10.7	7.1	11.9		2.4
New York (excluding New York City)	0.2	_	35.5	21.1	30.4		15.3
New York City	0.2	_	32.0	4.1	11.4		9.7
North Carolina	_	0.9	2.8	0.8	3.2		0.5
North Dakota		1.9	11.5	-	3.8		0.5
Ohio	_	0.3	4.6	3.2	7.6		0.3
Oklahoma	_	0.3	7.8	5.6	8.6		22.3
	_	0.4	3.2	1.1	1.6		2.1
Oregon	_		18.0	10.3	14.8		11.8
Pennsylvania Rhode Island	_	-	71.4	26.8	37.5		25.0
South Carolina	0.3	0.9	2.5	1.5	7.1		1.9
South Dakota	0.3	0.9	3.9	1.5	6.6		1.9
	_	-	9.1	3.7	5.2		0.9
Tennessee	_ 0						
Texas	0.5	3.0	12.0 5.5	6.8 1.0	5.9 6.1		4.3
Utah Vermont	_	1.0		1.0	3.7		_
	_		- 26 7		29.6		4 2
Virginia	-	0.8	36.7	15.8			4.3
Washington	0.4	11.3	29.0	16.4	26.6		27.3
West Virginia	_	-	4.3	1.1	1.1		2.2
Wisconsin Wyoming	-	18.9	3.7 28.3	1.2 5.7	3.7 9.4		0.6 7.5
American Samoa							
Guam	-	-	12.9	19.4	25.8	35.5	9.7
Northern Marianas							
Puerto Rico	_	2.1	3.7	2.1	2.1	4.5	4.5
Virgin Islands							

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2021
[By place of residence]

			'	Method of Delivery		
Reporting areas	Birth interval\9	Cigarette smoking before and during pregnancy	Risk Factors in this Pregnancy	Fetal presentation\10	Final route and method of delivery\11	
Total of reporting areas 1	11.5		5.5	11.5	4.9	
Alabama	2.4	2.2	0.8	1.6	0.4	
Alaska	10.5	19.3	7.0	7.0	5.3	
Arizona	4.1	4.3	3.5	7.2	2.1	
Arkansas	11.5	2.6	0.7	0.3	-	
California	12.7	11.9	11.3	33.2	9.4	
Colorado	32.4	30.1	0.3	19.9	16.0	
Connecticut	13.6	4.3	1.2	8.6	2.5	
Delaware	11.1	5.6	1.9	1.9		
District of Columbia	35.4	13.8	1.5	10.8	1.5	
Florida	18.3		8.0	20.6	9.1	
Georgia	12.2		10.4	15.0	3.4	
Hawaii	55.0		15.6	61.5	58.7	
	8.0					
Idaho			1.8	5.4	0.9	
Illinois	14.0	2.8	0.3	9.4	6.7	
Indiana	9.6		12.9	21.8	0.9	
Iowa	1.8	1.2	-	3.0	-	
Kansas	1.0		-	-	-	
Kentucky	4.6		2.1	3.3	0.6	
Louisiana	8.3	3.0	0.3	0.3	0.3	
Maine	7.5	-	-	6.0	4.5	
Maryland	8.3	3.1	1.1	1.5	0.2	
Massachusetts	7.6	3.4	-	1.4	3.4	
Michigan	12.0	4.5	5.2	6.5	0.8	
Minnesota	2.6	3.4	0.9	5.2	1.1	
Mississippi	4.5	0.8	2.8	2.5	0.3	
Missouri	5.4	5.2	_	3.9	0.5	
Montana	_	2.6	_	2.6	-	
Nebraska	6.1	4.0	_	_	_	
Nevada	5.4	1.1	0.4	2.5	1.1	
New Hampshire	2.7	4.1	2.7	1.4	-	
New Jersey	6.2	4.9	0.8	6.4	2.9	
New Mexico	11.9	2.4	2.4	3.6	1.2	
New York (excluding New York City)	28.4	22.2	2.9	31.2	29.2	
New York City	24.1	10.8	17.9	17.7	8.9	
North Carolina	3.1	0.5	0.3	0.5	0.5	
North Dakota	3.8	1.9	-	3.8	0.0	
Ohio	12.3	0.1		5.7	1.8	
			0.1		1.0	
Oklahoma	16.0	7.1	5.2	0.4	-	
Oregon	3.2	1.6	-	- 12.0	-	
Pennsylvania	14.8	20.3	6.3	13.0	9.0	
Rhode Island	37.5	37.5	16.1	28.6	14.3	
South Carolina	10.2		3.1	4.0	1.2	
South Dakota	7.9		1.3	-	-	
Tennessee	9.5		0.2	0.4	0.2	
Texas	9.3		7.1	0.2	1.4	
Utah	3.2	0.6	-	-	-	
Vermont	-	_	-	3.7	3.7	
Virginia	16.4	14.0	0.2	18.9	0.2	
Washington	58.5	21.3	35.7	23.3	17.1	
West Virginia	8.6	8.6	4.3	3.2	-	
Wisconsin	1.6	4.7	0.3	4.4	2.2	
Wyoming	22.6		-	3.8	3.8	
American Samoa						
Guam	6.5	35.5	25.8	38.7	-	
Northern Marianas						
Puerto Rico	11.6	0.8	2.1	2.9	2.1	
Virgin Islands						

See footnotes at end of table.

Table E. Percent of fetal death records on which specified items were not stated: United States and each state and territory, New York City, and the District of Columbia, 2021

[By place of residence]

Reporting areas	Maternal morbidity	Weight of fetus\12	Obstetric estimate of gestation	Cause of death\13
Total of reporting areas ¹	5.5	7.0	0.7	31.
Alabama	0.8	4.5	0.4	34.
Alaska	5.3	3.5	_	36.
Arizona	3.9	1.0	_	52.
Arkansas	0.7	8.2	0.7	43.
California	18.5	6.9	1.4	
Colorado	0.3	21.5	1.1	35.
Connecticut	1.9	4.3	_	22.
			1 0	
Delaware	5.6	_	1.9	24.
District of Columbia	1.5	1.5	-	26.
Florida	8.7	10.4	2.9	17.
Georgia	5.9	7.9	0.8	68.
Hawaii	11.9	56.9	3.7	77.
Idaho	0.9	6.3	-	25.
Illinois	0.8	8.4	0.3	21.
Indiana	_	6.1	0.2	31.
Iowa	_	4.9	_	31.
Kansas	_	1.5	0.5	28.
Kentucky	1.2	0.6	0.3	41.
Louisiana	0.3	9.6	-	22.
Maine	-	14.9	3.0	13.
Maryland	-	2.2	_	34.
Massachusetts	-	7.6	-	23.
Michigan	24.1	2.3	0.2	49.
Minnesota	0.3	3.4	0.3	38.
Mississippi	3.7	3.4	0.3	55.
Missouri	-	2.8	0.8	36.
Montana	-	2.6	-	43.
Nebraska	_	6.1	_	21.
Nevada	0.4	2.9	_	46.
New Hampshire	4.1	5.5	_	19.
New Jersey	3.1	0.7	0.2	34.
New Mexico	-	6.0	-	33.
	2.4	25.2	0.5	51.
New York (excluding New York City)				
New York City	19.7	11.4	0.2	64.
North Carolina	0.2	4.4	0.9	42.
North Dakota	_	1.9	-	67.
Ohio	0.7	4.5	-	31.
Oklahoma	4.5	4.1	0.7	23.
Oregon	-	3.2	0.5	41.
Pennsylvania	7.7	12.5	0.1	25.
Rhode Island	8.9	3.6	1.8	39.
South Carolina	2.8	1.9	_	41.
South Dakota	1.3	2.6	_	25.
Tennessee	0.2	2.0	0.4	46.
Texas	4.0	0.9	0.4	22.
Jtah	0.3	16.8	-	34.
/ermont	-	7.4	-	63.
/irginia	-	8.3	0.6	47.
Washington	-	24.4	2.9	23.
West Virginia	2.2	3.2	1.1	30.
Visconsin	0.6	4.7	0.3	50.
Wyoming	-	15.1	-	32.
American Samoa				
Guam Northern Marianas	32.3	-		93.
Puerto Rico	3.7	5.0	1.7	7.
Virgin Islands				

^{0.0} Quantity more than zero but less than 0.05.

NOTE: American Samoa did not report data for 2021. Data for the Commonwealth of Northern Marianas and the Virgin Islands were excluded due to small numbers. Reporting areas with levels of unknown data for an item of 50% and higher (based on counts by occurrence and prior to the implementation of edits) are not included in the U.S. totals and are flagged in the file as not reporting.

1\Excludes data for American Samoa, Guam, the Commonwealth of the Northern Marianas, Puerto Rico, and the Virgin Islands.

\2Hawaii and Rhode Island reported education of mother for 2021 but had an unknown level of 50% or higher and were flagged as not reporting.

\3Alabama, Arkansas, the District of Columbia, Georgia, Hawaii, New Mexico, New York (including New York City), Virginia, Wisconsin, and Guam reported father's age for 2021 but had an unknown level of 50% or higher and were flagged as not reporting.

\4Hawaii and Rhode Island reported date of first prenatal care for 2021 but had unknown levels of 50% or higher and were flagged as not reporting.

 $\$ Thawaii reported mother's height for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\6Hawaii reported mother's pre-pregnancy weight for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\7New York City does not report WIC and the District of Columbia, Hawaii, and Rhode Island reported this item for 2021 but had unknown levels of 50% or higher and were flagged as not reporting

\8Hawaii reported live birth order for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\9Hawaii and Washington reported birth interval for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\11Hawaii reported final route and method of delivery for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

 $\10$ Hawaii reported fetal presentation for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\12Hawaii reported weight of fetus for 2021 but had an unknown level of 50% or higher and was flagged as not reporting.

\13Arizona, Georgia, Hawaii, Mississippi, North Dakota, New York (including New York City), Vermont, Wisconsin and Guam reported cause of death for 2021 but had an unknown level of 50% or higher and were flagged as not reporting. California did not report cause of death based on the 2003 revision of the U.S. Standard Report of Fetal Death and was also flagged as not reporting.

⁻⁻⁻Data not available.

⁻ Quantity zero.