Healthy People 2010 Operational Definition

25-1. Reduce the proportion of adolescents and young adults with *Chlamydia trachomatis* infections.

25-1c. Males aged 15 to 24 years attending STD clinics.

National Data Source STD Surveillance System (STDSS), CDC,

NCHHSTP.

State Data Source State and local Health Department STD Control

Programs.

Healthy People 2000

Objective

Adapted from 19.2 (Sexually Transmitted Diseases).

Changes since the

2000 Publication

None.

Measure Percent.

Baseline (Year) 15.7 (1997)

Target 3.0

Target-Setting Method Better than the best racial/ethnic subgroup.

For a discussion of target-setting methods, see Part

A, section 4.

Numerator Number of positive tests among men aged 15 to 24

years who attended STD clinics in the past 12

months.

Denominator Number tests administered (unsatisfactory tests,

indeterminate or inconclusive results, or inadequate specimens are excluded) among men aged 15 to 24 years who attended STD clinics in the past 12

years who attended 5 LD clinics in the past

months.

Population Targeted U.S. civilian population.

Questions Used To Obtain the National

Baseline Data

Not applicable.

Expected Periodicity Annual.

Comments A case definition for *Chlamydia trachomatis* is

provided by CDC.1

Data are collected from Regional Infertility Prevention Program laboratory reports.

This measure tracks only tests that are specific for *Chlamydia trachomatis* infection. The number of positive cases is based on test results from persons routinely screened. Routine screening is defined by each clinic. In some clinics, all patients are universally screened. In others, routine screening is done selectively based on clinical findings or behavioral risk factors.

A minimum of 500 valid test results by gender and subpopulation is recommended to derive reliable estimates for this measure. Some clinics may not collect all U.S. Census-defined race and/or ethnicity data categories.

Objective 25-1 is a modification of Healthy People 2000 objective 19.2, which tracked total cases of Chlamydia in women under age 25 years who attended family planning clinics. Objective 25-1 tracks the same measure: 25-1a focuses on women aged 15 to 24 years who attended family planning clinics and 25-1b on women aged 15 to 25 years who attended STD clinics; 25-1c expands the measure to focus on men aged 15 to 24 years who attended STD clinics, while 25-1d is a new subobjective which addresses women aged 24 years and younger who are enrolled in a National Job Training Program.

For some measures, data do not meet the criteria for statistical reliability, data quality, or confidentiality and have been suppressed. Information on suppression of data for the major Healthy People 2010 data systems has been published in a *Healthy People Statistical Note*. ²

See Appendix A for focus area contact information.

References

- 1. Centers for Disease Control and Prevention (CDC). Case definitions for infectious conditions under public health surveillance. *Morbidity and Mortality Weekly Report* 46 (RR-10): 1997.
- 2. Klein, R.J.; Proctor, S.E.; Boudreault, M.A.; Turczyn, K.M. Healthy People 2010 Criteria for Data Suppression. *Statistical Notes* No. 24. Hyattsville, MD: National Center for Health Statistics. 2002.

Operational Definition: Objective 25-1c