

Cohort Mortality and Survivorship: United States Death-Registration States, 1900-1968

An analysis of mortality rates by age, color, and sex of selected generations of 5-year birth cohorts born 1896-1900 through 1926-1930. Compares cohort and period life table survivorship (l_x) by single years of age, color, and sex for selected 5-year cohorts born 1899-1903 through 1928-1932. Based on death and population data for the death-registration States of the United States each year from 1900 to 1968.

DHEW Publication No. (HSM) 73-1400

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service

Health Services and Mental Health Administration
National Center for Health Statistics
Rockville, Md. November 1972



Vital and Health Statistics-Series,3-No. 16

NATIONAL CENTER FOR HEALTH STATISTICS

THEODORE D. WOOLSEY, *Director*
EDWARD B. PERRIN, Ph.D., *Deputy Director*
PHILIP S. LAWRENCE, Sc.D., *Associate Director*
OSWALD K. SAGEN, Ph.D., *Assistant Director for Health Statistics Development*
WALT R. SIMMONS, M.A., *Assistant Director for Research and Scientific Development*
JOHN J. HANLON, M.D., *Medical Advisor*
JAMES E. KELLY, D.D.S., *Dental Advisor*
EDWARD E. MINTY, *Executive Officer*
ALICE HAYWOOD, *Information Officer*

OFFICE OF HEALTH STATISTICS ANALYSIS

IWAO M. MORIYAMA, Ph.D., *Director*
DEAN E. KRUEGER, *Deputy Director*

Vital and Health Statistics-Series 3-No. 16

DHEW Publication No. (HSM) 73-1400

Library of Congress Catalog Card Number 71-179928

CONTENTS

	Page
Introduction -----	1
Earlier Studies-----	1
Data and Methodology-----	3
Qualifications of Data-----	4
Cohort Mortality-----	5
Cohort Survivorship-----	11
Survivorship of Birth Cohorts-----	11
Cohort and Period Survivorship-----	11
Discussion -----	17
References -----	18
List of Detailed Tables-----	19
Appendix I. Relationship Between Cohort and Period Mortality-----	32
Appendix II. Production of the Single-Year Data-----	33
Appendix III. Construction of the Cohort and Period Survivorship Tables----	34
Published Tables-----	34
Method of Estimating Death Rates at Age 0-1-----	34
Deaths at Other Ages and Survivors-----	35

SYMBOLS

Data not available-----	---
Category not applicable-----	...
Quantity zero-----	-
Quantity more than 0 but less than 0.05----	0.0
Figure does not meet standards of reliability or precision-----	*

COHORT MORTALITY AND SURVIVORSHIP: UNITED STATES DEATH-REGISTRATION STATES, 1900-1968

Iwao M. Moriyama, Ph.D., and Susan O. Gustavus, Ph.D.^a

INTRODUCTION

The official death statistics are derived from the mortality experience of a population for a particular time period, usually a calendar year. They represent a slice or a cross section of the mortality surface taken across the time axis, and are known as period mortality data. Another way of looking at death statistics is *along* the diagonal of the age and time axes rather than *across* the time axis. These longitudinal sections of the mortality surface show the mortality experience of cohorts of individuals from birth through the successive ages over their lifetimes (see appendix I). These cohort or generation mortality data are representations of what actually happens in life; nevertheless, data are seldom expressed in this way because a relatively long series of age-specific mortality statistics is needed to do so.

Following the same group of people over the lifetime of the cohort presents quite a different mortality and survival picture from that provided by the official annual mortality statistics. This is because of changes, usually improvements, in mortality rates during the life of the cohort.

^aIwao M. Moriyama, Director, Office of Health Statistics Analysis, National Center for Health Statistics, is presently on leave, serving as Chief, Statistics Department, Atomic Bomb Casualty Commission, Hiroshima, Japan. Susan O. Gustavus is Assistant Professor, Department of Sociology, The University of Utah, Salt Lake City.

This study presents the mortality and survivorship experience of four cohorts born a decade apart and subjected to the force of mortality in the United States during the period 1900 to 1968. These data are unique in that death rates and survivorship rates are derived for each calendar year.

The effects of the 1918 influenza pandemic may be seen in the experience of two of the four cohorts. The effects of other influenza epidemics of lesser proportions are also apparent in the generation curves.

The effects of World War II and of the Korean War are evident in cohort mortality curves of white males. Males of Negro and other races did not experience nearly the same increase in mortality during World War II, nor did the rate peak up to the same extent as the rate for white males in the Korean War.

Because of the decline in mortality over the years, the differences in cohort and period survivorship indicate that past period life tables have not represented the real-life mortality experience of a birth cohort. However, because the rate of decline in the mortality rates is slowing down, future period life tables should become better predictors of mortality in a cohort than were past period life tables.

Earlier Studies

The analysis of mortality patterns by the use of generation data is not new. While past efforts

in examining mortality data by the cohort method have been limited by lack of suitable data, Kermack, McKendrick, and McKinlay¹ studied mortality data for cohorts at 10-year intervals from 1755 to 1925 in Great Britain and Sweden. No projections were attempted; consequently, data for most of the cohorts were incomplete. Their study showed that mortality patterns were fairly constant in each cohort, that is, that the most important factor in later mortality was the experience of the cohort before age 15 years, with each cohort having fairly similar mortality rates after that age. While later findings do not agree with this conclusion, this study was significant in that it was concerned with the mortality experience of cohorts of individuals instead of the total population at a fixed point in time.

Case² presented a review of cohort analysis and a detailed explanation of the logic of the technique including examples with data for England and Wales from 1851 to 1951. He compared the cohort and period approaches to mortality and commented on the "generation effect" and how it could be examined with cohort mortality data. The generation effect is based on the hypothesis that early mortality experience affects, or even determines, later mortality. This may occur in a variety of ways. For example, Pearson³ and others felt that an effect of reducing infant mortality rates would be to raise mortality rates at later ages because such a lowering at the early ages would keep the "weak" alive and prevent natural selection from operating. However, this has not been borne out by later experience—or perhaps, the rapidity with which the death rate has declined may have obscured the effects, if any, of postponing deaths of presumably impaired lives. Case advanced the notion that the existing concepts of the laws of mortality were inadequate and could lead to improper inferences on the nature-nurture complex of problems because environment and therapeutic measures constantly change. He favored the use of the cohort analysis as a narrative or historical technique, and proposed "a synthesis of knowledge derived from social history, medical history, and cohort analysis to be made to interpret the narrative."

By far, the most frequent use of the cohort approach has been to examine mortality from specific diseases. Of these, Frost's⁴ study of

tuberculosis is a classic. He demonstrated that the actual pattern of mortality was not what was expected from previous findings on age-specific death rates for tuberculosis at one point in time. The latter approach showed that the greatest risk of death from tuberculosis was in the older ages, whereas the cohort data made it clear that the groups experiencing the highest risks at later ages had already passed through periods of even higher risks at the younger ages. Picken⁵ confirmed Frost's results after applying Frost's methods to data for England and Wales for the same time period. However, Spicer⁶ found from his analysis that the generation hypothesis gave a good description of mortality from respiratory tuberculosis until about 1930. After this period, the hypothesis no longer agreed satisfactorily with the facts.

The cohort method has been applied in studies of cancer mortality by Korteweg,⁷ Stocks,^{8,9} Haenszel and Shimkin,¹⁰ Cutler and Loveland,¹¹ and others. These studies showed that successive cohorts experienced increased mortality for some sites and decreased mortality for other sites. In the Cutler and Loveland study, the cohort mortality rates for lung cancer were projected to estimate incidence rates. Deaths from other diseases have also been examined using the cohort approach.¹² In general, studies of cohort mortality data relating to specific diseases are much more meaningful than those encompassing all causes of death. The data for all causes of death are composites of the exposure to various diseases, and the cohort patterns of different diseases are averaged out. However, they are useful summaries of the total mortality experience of the cohorts as they are exposed to the actual force of mortality at various stages of life.

While life tables have often been used to determine death and survivorship rates, or years of life remaining at each age, they have generally been constructed for one point in time. Here again, lack of data has required analyses to rest on the assumption that the age-specific mortality rates for a particular year will prevail through the entire lifetime of the population presented in the life table. Dublin and Spiegelman¹³ showed the weakness of such an assumption by demonstrating from life tables

for the period 1871 to 1931 that there was a much greater "saving of life" during this period than would have been anticipated if the 1871 death rates applied in later years.

More recently, Spiegelman¹⁴ used available vital statistics to create data on cohorts at 10-year intervals from 1900 to 1960. This study and others had several limitations. First, none of them was able to observe a cohort at shorter intervals than 5-year periods, and most used 10-year periods. Second, lacking data to complete their cohorts at young or old ages, incomplete cohorts had to be dealt with, or projections made on the basis of a number of assumptions. Alternatively, the analysis had to be limited to one period in each cohort's life, say, after age 45.

Data and Methodology

It is the purpose of this report to present mortality rates for four cohorts whose central years of birth are 1901, 1910, 1920, and 1930. The survivorship of these cohorts is also examined using both cohort and period mortality data in order to see how much difference exists in these two approaches, and the possible implications of this discrepancy.

The data in this report were produced from estimates of the population in 5-year age groups from birth to age 84 years in the period 1900 to 1968, inclusive. The number of deaths by age, sex, and color was obtained from the official vital statistics to which the war deaths were added from data made available by the Department of Defense. In this respect, the material is different from the conventional U.S. mortality statistics which include only deaths registered in the United States.

Data on both population and deaths refer to the expanding death-registration States for the years prior to 1933 and to the United States for subsequent years.^b

Population and death data by single years of age were interpolated from the 5-year age groups by applying Beers' formula (see appendix II).

^bFor information on the death-registration States see the technical appendix of *Vital Statistics of the United States*, Vol. II—Mortality, Part A.

To prepare cohort or generation data, the statistics by age were combined in two ways. First, to produce *cohort mortality* tables, the data by single years of age for single-year birth cohorts were combined into 5-year birth cohorts to show mortality rates for 5-year age groups of cohorts for each calendar year. For example, the death rate for the cohort born 1896-1900 was computed for the year 1910 at which time the cohort was 10-14 years old, or age 12.5 on the average (refer to the column of x 's in table A). A death rate at ages 11-15 was then computed for the same cohort in 1911 when the cohort was a year older, or 13.5 years old on the average (refer to the column of y 's in table A). In this way, the cohort was followed through each calendar year until 1968.

Second, to produce *cohort survivorship* in detailed tables 1-8, the data were combined another way into 5-year birth cohorts to show mortality rates by single years of age for a succession of years of death. Thus, the death rate for a specified age represents the mortality experience of five birth cohorts at that age over a period of 5 calendar years. For example, the mortality rate for age 10 for the 1899-1903 birth cohort is based on population and deaths in the year 1909 for the 1899 component of the 5-year cohort, population and deaths in the year 1910 for the 1900 component, etc. (refer to the diagonal of o 's in table A). The sum of the deaths in the five cohorts was divided by the sum of the five cohort populations to obtain a mortality rate for a single year of age. The life table death rate, q_x , was calculated from these mortality rates for each single year of age. Beginning with a population of 100,000 (radix), the q_x values were applied to the surviving life table populations to obtain the number dying at each age. The number surviving to each successive age was obtained by subtraction.

To produce the *period survivorship* in detailed tables 1-8, deaths and populations by single years of age were averaged over a period of 5 years. Death rates for the 5-year period were then computed from the average numbers of deaths and population. For example, the period survivorship table for 1910 is based on the average death rate by age for the period 1908 to 1912. An exception is the 1901 period

Table A. Example of data used in calculating cohort mortality rates for single years of time and cohort survivorship for single years of age

Year of birth	Year of death				
	1909	1910	1911	1912	1913
1896-----		<i>x</i>	<i>y</i>		
1897-----		<i>x</i>	<i>y</i>		
1898-----		<i>x</i>	<i>y</i>		
1899-----	<i>o</i>	<i>x</i>	<i>y</i>		
1900-----		<i>xo</i>	<i>y</i>		
1901-----			<i>o</i>		
1902-----				<i>o</i>	
1903-----					<i>o</i>

- x* Basis of the mortality rate of the cohort born 1896-1900 for the year 1910 (ages 10-14) at average age 12.5.
y Basis of the mortality rate of the cohort born 1896-1900 for the year 1911 (ages 11-15) at average age 13.5.
o Basis for the mortality rate from which survivorship from age 10 to age 11 was calculated for the cohort born 1899-1903.

survivorship table, which is based on averages for the 4 years 1900-1903. Survivorship in these tables was then computed in the same manner as in the cohort tables. (See appendix III.)

Because the original data on which the interpolation procedure was performed contained population estimates and deaths only for the period 1900 to 1968, the time periods which could be selected for examination were limited. Thus, the earliest 5-year birth cohort chosen for the study of survivorship was the cohort of 1899-1903. Since death rates for the years after 1968 were not included in this study, cohort mortality and survivorship tables are incomplete after this date. Consequently, the birth cohort of 1899-1903 can be followed only until it reaches age 70, the birth cohort of 1908-1912 until it reaches age 61, the birth cohort of 1918-1922 until it reaches 51 years, and the birth cohort of 1928-1932 only until it reaches age 41.

In spite of this time limitation, these data are unique in that they allow birth cohorts to be followed each year in time. Past uses of cohort techniques have been largely limited to looking

at mortality experience of cohorts at 5- or 10-year intervals. By the use of the interpolation procedure, however, it is now possible to see changes in mortality in each calendar year. Such single-year data show variations in mortality that are not apparent from the 5-year estimates.

It would have been possible to examine 1-year birth cohorts, say the 1900 birth cohort, instead of grouping the data into 5-year birth cohorts. The specificity provided by a cohort of births occurring in a single year is a desirable feature. However, 5-year birth cohorts were used to smooth out irregularities that may have arisen in the data as a result of the interpolation procedure.

Qualifications of Data

In this study the populations referred to as cohorts are not cohorts in the true sense of the word. Technically, one would start with a cohort of births and observe the deaths each year until the cohort becomes extinct. In this study the cohort is, loosely speaking, the population at specified ages at a particular time period. The

mortality data were derived from death statistics for the death-registration States, which was an expanding group that did not include all the States in the United States until 1933. Thus, the mortality data do not specifically relate to the original cohort as it aged over the years. However, the observed death rates may be taken as approximations of the true mortality rates of the cohort as it passed through the various ages.

To reduce variability in the rates and to minimize the effect of heaping in the terminal digits of 0 and 5 in the statements of age, the data were grouped into cohorts born over a 5-year period. However, these groupings produced damping effects inherent in the averaging process.

In order to derive survivorship tables, it was necessary to have death rates at specified ages. These were obtained by averaging the death rates for each specific age experienced by the cohort. For a 5-year cohort this meant the averaging of death rates over 5 calendar years. Although this is an acceptable procedure for computing survivorship data, it produces an undesirable effect in the analysis of cohort mortality. Because the death rates at any age are averaged over 5 calendar years, it is not possible to see the correspondence between an event and the exact time at which it occurred. For example, when the death rates over time are averaged, the effect of the influenza pandemic of 1918 appears at a peak in 1920 for the birth cohort of 1899-1903. In order to avoid this kind of distortion along the time axis, cohort mortality rates were computed on a basis different from that used for deriving the survivorship tables, as described above (page 3). This difference needs to be kept in mind in the interpretation of the cohort mortality and survival data.

Another problem is that it is not now possible to produce cohort mortality data for a complete generation. Because the mortality series for the United States is relatively short, the curves will be truncated until sufficient data are available so that a cohort may be followed to extinction.

These qualifications are not unique to this series of data. All cohort material based on data for the death-registration States has the same limitations. All cohort data that are combined in 5- or 10-year age groups are also subject to distortions arising from averaging age-specific rates for the span of the age group.

COHORT MORTALITY

The characteristic pattern of both cohort and period mortality rates is the high death rate at the two extremes of the age scale. The mortality risk is extremely high at birth, declines to a minimum in childhood (age about 10-12 years), and rises again. The highest level is reached in the older ages, but this is not always apparent from the data which do not carry the cohort to the end of its lifetime.

A number of unusual peaks in mortality may be observed in the cohort data, especially for white males (see figure 1). The effects of the 1918 influenza pandemic may be seen in the experience of two cohorts. The highest peak occurred in the 1896-1900 cohort in 1918 when the average age of the group was 20 years. A smaller peak in mortality occurred for the 1906-1910 cohort at the average age of 10; this cohort was not hit nearly as hard by the influenza epidemic as the older cohort. The effects of other respiratory disease epidemics of lesser proportions such as those that occurred in 1929, 1936, and 1937 are also apparent in the cohort curves at the following ages:

Cohort	Year of epidemic	
	1929	1936 and 1937
1896-1900-----	31	38 and 39
1906-1910-----	21	28 and 29
1916-1920-----		18 and 19




Figure 1. Mortality rates for cohorts born 1896-1900, 1906-1910, 1916-1920, and 1926-1930, by sex, color, and age: death-registration States, 1900-1968.




Figure 1. Mortality rates for cohorts born 1896-1900, 1906-1910, 1916-1920, and 1926-1930, by sex, color, and age: death-registration States, 1900-1968—Con.




Figure 1. Mortality rates for cohorts born 1896-1900, 1906-1910, 1916-1920, and 1926-1930, by sex, color, and age: death-registration States, 1900-1968—Con.




Figure 1. Mortality rates for cohorts born 1896-1900, 1906-1910, 1916-1920, and 1926-1930, by sex, color, and age: death-registration States, 1900-1968—Con.

The effect of World War II may be seen in the rates for cohorts of white males born after 1900. The greatest impact was on the cohort born between 1916 and 1920. This group was 24-28 years of age in 1944 when the peak of mortality was experienced in World War II. Lesser peaks appear in the curves for other cohorts. The death rate for the 1926-1930 cohort rose to a maximum in 1945 when the members were 15-19 years of age. If this peak resulted from World War II, only the older members of this cohort were presumably involved. This cohort was further exposed to war risks in 1951 during the Korean conflict. It would also appear from these data that even the 1906-1910 cohort was affected by World War II. A small upswing in mortality can be seen in the rates for the cohort when the individuals in the group were 34-39 years of age in 1944 and 1945.

The mortality rates for cohorts of males of Negro and other races show a little different picture from that of white males, in addition to their generally higher levels. The effects of the influenza pandemic of 1918 are evident in the 1896-1900 cohort and to a lesser degree in the 1906-1910 cohort. Also, there are a number of minor peaks representing the effects of influenza epidemics over the years. The consequences of World War II mortality are not as apparent in the rates for Negro and other races compared with the peak in mortality for white males. The effect of the Korean conflict is seen in the mortality experience of the 1926-1930 cohort of males of Negro and other races, but here again the rate did not peak up to quite the same extent as the rate for white males.

The mortality experience of females is similar to that of males except for the absence of war casualties and the lower level of mortality particularly at the older ages. Prominent are the peaks at ages 20 and 10 years for the 1896-1900 and 1906-1910 cohorts, respectively, resulting from the influenza pandemic of 1918. Lesser peaks from other influenza epidemics are also evident.

There is a greater similarity in the configuration of the cohort death rates between the sexes than between color groups. For Negroes and other races the range of the death rates is much greater and the base of the curve is much narrower than for whites. The death rates for races other than white exhibit greater variability because of the

smaller frequencies of deaths. Also, the respiratory disease epidemics produced much greater upswings in the death rates for this group.

The improved mortality experience of the various cohorts is evidenced by the nest of curves for each color-sex group where most death rates for each succeeding cohort are lower than the rates for the previous cohort. At the older ages, there appears to be a convergence of death rates with those of the previous cohort. The point of convergence seems to be occurring earlier and earlier with each succeeding cohort. This suggests that the upturns in the death rates for the succeeding cohorts are occurring at younger ages. The narrowing of the base of the generation mortality curves appears significant. More is said about this phenomenon in the Discussion section below.

In addition to beginning at younger ages, the upturning death rates appear to be following a steeper rate of increase into the older ages for succeeding cohorts. As a consequence, there are points of crossover where the death rates of some cohorts begin to exceed those of the preceding cohort. Crossover points for the 1926-1930 cohort occur near age 35 for males of both color groups and near age 40 for white females. For females of other races, crossover has not yet occurred but appears imminent from the trend line for the 1926-1930 cohort (see figure 1D). All cohorts of males of other races shown in figure 1C demonstrate the crossover phenomenon. However, in comparison with the preceding cohort, only the 1926-1930 cohort of males of other races has demonstrated substantially higher mortality persisting over a number of years. Their comparatively high death rates between ages 35 and 40 during the 1960's are consistent with the rising death rates in these ages in recent years.¹⁵

The minimum level of mortality was reached in the childhood ages (between 6 and 16 years) in the various cohorts. There does not seem to be any great change in the age of occurrence of minimum mortality over the years. Also, no pattern of differentials by sex is discernible. However, there is a difference in age of minimum mortality for the two color groups. In general, the lowest point of the death rate among whites is at an age several years higher than that for Negro and other races. This age spread by color in the cohort mortality curves results

from the low mortality among whites in the later years of childhood.

COHORT SURVIVORSHIP

The survivorship data for the various selected birth cohorts are given in detailed tables 1-8. As stated previously, these data were computed on a slightly different basis than the cohort mortality data already discussed. For the purpose of generation survivorship computations, the mortality rates over a period of 5 calendar years were averaged to obtain stability in the computed death rates. The same end was achieved in a different manner in computing cohort mortality rates, that is, the rates were averaged over the ages represented in the cohort for any particular year. In this way, the effect of events in a specific year is not obscured by the averaging process. Although the two sets of data are not precisely comparable, they are more suitable for the two purposes of this analysis than if they were computed in the same way.

In the cohort survivorship data the following abbreviations are made for the convenience of discussion:

Birth cohort of:	Cohort of births occurring in:
1901	1899-1903
1910	1908-1912
1920	1918-1922
1930	1928-1932

Survivorship of Birth Cohorts

The survivorship of birth cohorts of different color and sex groups is presented in figure 2. The most favorable survivorship pattern is that of white females, and the least favorable is that of males other than white. Of the white females in the 1901 birth cohort, more than 55 percent survived to age 70 years. Of males of other races born in the same period, less than 20 percent lived to age 70 years.

The effect of improvement in mortality over the years is evident in the survivorship curves for the different birth cohorts. It would appear that the reductions in mortality for the white population have been relatively uniform in time,

whereas the decrease in mortality (or increase in survivorship) for the population other than white was particularly large between the 1901 and 1910 cohorts. It is possible that some of this change is only apparent and resulted from the large increase in the Negro population due to the growth of the death-registration area.

Cohort and Period Survivorship

The generation tables represent more or less what happens in real life as compared with the usual period life tables. Because of the decrease in mortality with attendant increase in life expectancy over time, the number surviving to each successive age is generally higher in the generation or cohort table than in the period life table. This is illustrated by figure 3 which shows the survivorship of the cohort of white males born 1899-1903 in comparison with the survivors as computed from the age-specific mortality rates for the period 1900 to 1903.

The relative differences between the numbers of survivors to successive selected ages in the period and cohort tables are shown in table B. It may be seen from these data that the percentage differences in the numbers of survivors between the generation and period mortality tables increase with age. Also, the differences in survivorship based on these two types of tables vary with time and with the population group involved.

As table B reveals, the largest discrepancies between cohort and period survivorship occur in the 1920 comparisons. This is because of the peculiarity in the mortality data for that period. The 5-year average centered on 1920 includes data for 1918 and 1919, the years of the influenza pandemic which took the largest death toll in the history of U.S. vital statistics. This was followed by a year of unduly low mortality so that basing the period data only on the mortality experience for 1920 would still give an atypical comparison. However, this would not have been nearly as misleading as the inclusion of data for the epidemic years. Therefore, it would be well not to attach much significance to the differences in survivorship in the 1920 generation and period tables.




Figure 2. Survivorship (l_x) of birth cohorts of 1901, 1910, 1920, and 1930, by sex, color, and age: death-registration States, 1900-1968.




Figure 3. Survivorship (L_x) of white males in birth cohort of 1901 as compared with corresponding period survivorship, by single years of age: death-registration States, 1900-1968.

White males.—As may be seen from figure 4A (as well as in table B), the difference between generation and period survivorship of white males is less than 5 percent for the age groups under 30 years. From that point, the difference increases rather sharply. Although the 1930 period table came the closest to the actual mortality experience of white males, the variations in the different periods under comparison were relatively small.

The cohort and period survivorship difference at age 20 in the 1901 table was small because of the influenza epidemic. The birth cohort of 1901 would have been age 20 in 1919-1923. Thus, the number surviving to this age in the cohort table was smaller than would have been expected had there not been an influenza epidemic.

A similar change in the cohort and period survivorship difference is seen in the 1910 comparison for white males. At age 15 and earlier, this cohort experienced higher mortality rates than would have been expected without the outbreak of influenza. Thus, the number surviving to age 15 was closer to the period mortality survivors of 1910 than might have been expected.

A significant dip can be observed at age 25 years in the difference between the cohort and period survivorship of the 1920 table. This resulted from the rise in mortality of white males aged 15-25 years during World War II.

White females.—As in the comparison for white males, the differences in generation and period survivorship for white females are relatively small for the age group under 30 years (see figure 4B). These differences increase rapidly in the older ages. At age 70 years, the difference is near 60 percent, the highest relative difference between generation and period survivorship of any color-sex group.

By age 35 years, the 1930 period table was the best predictor of the actual number of white female survivors from the cohort, but it was only slightly better than the table for 1910. As was true for white males, the variations in difference between periods were relatively small if the 1920 period comparison is excluded.

The events that affected the birth cohorts of white females, making survivorship differences less than expected, were the same that affected the survivorship of white males in the 1901 and 1910 cohorts. A decrease in survivorship differences is apparent at age 20 for the 1901 birth cohort, and before age 15 for the 1910 birth cohort.

Males of other races.—There appears to be an increase in the survivorship differences between the generation and period data for males of other races at age 15 for the 1901 birth cohort and at age 20 for the 1910 birth cohort (figure 4C). The reason for these changes is not clear.

Table B. Percent difference¹ between cohort and period survivorship (l_x), by sex, color, and age: 1901, 1910, 1920, and 1930

Age in years	1901				1910			
	Male		Female		Male		Female	
	White	All other	White	All other	White	All other	White	All other
5-----	0.4	-0.1	0.4	0.2	0.3	1.6	0.3	1.5
10-----	0.7	0.3	0.8	0.8	0.4	2.3	0.4	2.1
15-----	1.0	1.4	1.1	2.3	0.5	3.2	0.6	3.5
20-----	1.1	1.7	1.3	2.6	1.0	4.7	1.1	5.4
25-----	2.3	2.4	2.4	2.4	2.1	5.9	2.1	6.8
30-----	4.1	3.0	4.2	2.4	3.6	7.4	3.6	8.4
31-----	4.6	3.2	4.6	2.6	4.0	7.8	4.0	8.9
32-----	5.0	3.4	5.0	2.8	4.4	8.3	4.4	9.4
33-----	5.5	3.6	5.5	3.2	4.8	8.8	4.8	10.0
34-----	5.9	3.8	6.0	3.6	5.2	9.3	5.2	10.6
35-----	6.4	4.0	6.5	3.9	5.6	10.1	5.7	11.3
40-----	9.1	4.8	9.2	5.1	8.3	14.7	8.3	15.5
45-----	12.3	6.7	12.5	8.2	11.5	19.8	11.2	21.2
50-----	15.6	10.0	16.4	12.4	14.8	25.3	14.8	28.4
55-----	18.8	13.9	21.9	17.3	17.8	31.5	19.4	37.8
60-----	22.9	22.0	29.9	26.2	21.2	37.4	26.2	51.2
65-----	27.2	28.4	41.5	36.0	---	---	---	---
70-----	32.7	30.9	58.9	45.9	---	---	---	---

¹Percent difference is the difference between cohort l_x and period l_x as a percent of the period l_x ; based on data in tables 1-8.

Table B. Percent difference¹ between cohort and period survivorship (l_x), by sex, color, and age: 1901, 1910, 1920, and 1930—Con.

Age in years	1920				1930			
	Male		Female		Male		Female	
	White	All other	White	All other	White	All other	White	All other
5-----	0.7	1.1	0.8	1.1	0.2	0.2	0.2	0.2
10-----	1.2	1.6	1.3	1.8	0.5	0.6	0.4	0.6
15-----	1.7	2.3	1.8	2.9	0.8	1.1	0.7	1.2
20-----	2.8	4.5	2.9	5.3	1.2	2.8	1.4	3.6
25-----	3.3	8.4	5.2	9.6	1.7	5.8	2.6	7.5
30-----	5.3	12.7	8.2	14.6	2.8	9.8	4.0	12.0
31-----	5.9	13.7	8.9	15.8	3.1	10.7	4.3	12.9
32-----	6.6	14.7	9.5	16.9	3.4	11.6	4.6	13.8
33-----	7.3	15.8	10.2	18.2	3.6	12.5	4.9	14.8
34-----	7.9	16.9	10.9	19.5	3.9	13.5	5.2	15.8
35-----	8.6	18.1	11.5	20.8	4.2	14.4	5.5	16.8
40-----	11.8	23.8	14.7	27.5	5.8	19.3	7.3	22.6
45-----	14.5	28.5	17.8	34.4	---	---	---	---
50-----	17.0	32.0	21.3	42.9	---	---	---	---
55-----	---	---	---	---	---	---	---	---
60-----	---	---	---	---	---	---	---	---
65-----	---	---	---	---	---	---	---	---
70-----	---	---	---	---	---	---	---	---

¹Percent difference is the difference between cohort l_x and period l_x as a percent of the period l_x ; based on data in tables 1-8.




Figure 4. Percent difference between cohort and period survivorship (l_x) of birth cohorts of 1901, 1910, 1920, and 1930, by sex, color, and age: death-registration States, 1900-1968.

As compared with the pattern for white males, the spread in the experience of various birth cohorts of males of other races is large. Also, it would appear that the period table that came closest to the actual mortality experience of a birth cohort was that for 1901 for males of other races. The 1930 period table turned out to be a poor third.

Because mortality rates have decreased over time, it was expected that the cohort tables would always show greater numbers of survivors at each age than would the period tables. The one exception to this, indicated by the negative difference in table B, is the difference in the male survivors of other races to age 5 in the 1901 tables. This difference from the expected pattern is a result of higher death rates in the 1901 cohort table than in the 1901 period table until age 3. The death rates affecting males of other races aged 0-3 years in the years 1902, 1903, and 1904 were higher than the corresponding death rates in 1901. While the difference is not great, it is a deviation from the general pattern.

Females of other races.—The general pattern of survivorship differences between the generation and period tables for females of other races resembles that for males of the corresponding color group. However, the differences in survivorship are uniformly greater for females of this group. Also, as may be seen in figure 4D, there is an unusual plateau in the survivorship differences between ages 15 and 30 years for the 1901 birth cohort. This plateau covers the years 1914-1933, but the reasons for the relatively high cohort death rates are not known.

DISCUSSION

The annual mortality statistics have been valuable in following the course of mortality over the years, but such period mortality data do not represent the real-life situation in which a population cohort goes through life being subjected to changing forces of mortality. By generating mortality data on an annual basis for various birth cohorts, it is also possible to see the effects of specific events, such as respiratory disease epidemics and wars, on mortality of specific population groups. Thus, a new dimension (longitudinal) is added to mortality statistics.

The big disadvantage of generation or cohort mortality statistics is that a large body of statistics is needed. The mortality series for the United States is now sufficiently long so that it would be worth while examining the longitudinal experience of various cohorts. As was done in Sweden, it would be desirable to tabulate annually mortality statistics by single years of age. These statistics could then be grafted to this report's data which were derived by an interpolation procedure from mortality statistics by 5-year age groups.

Cohort data by causes of death should provide more insight into mortality from various diseases and their determinants. Because statistics on all causes of death are a weighted average of death rates for the different component diseases, it would be expected that the data in this report would show only the grossest changes in mortality. This turned out to be the case. The influenza epidemic of 1918 and some of the lesser epidemics of other years, as well as the effects of World War II and the Korean War on the male population, appear to be reflected by the cohort data.

Of special interest is the pattern of cohort mortality data which consisted of a nest of U-shaped curves. In these curves, the base of the U's of the cohort mortality curves for whites is much broader than that for all other races. Also, with the improvement in mortality experience for the succeeding cohorts, there is a continuous narrowing of the base. This same phenomenon may be seen in the cohort mortality curves for Sweden presented by Bolander.¹⁶ This seems contrary to expectations. With decreasing mortality, one would expect a broadening of the base. In fact, if all people were constructed like Longfellow's one-hoss shay, the shape of the curve would approach the mirror image of an L which would depict a zero mortality from birth until the appointed age when the death rate would be 100 percent.

The narrowing of the base of the generation mortality curves with improvement in mortality suggests that over the years the decline in mortality has taken place primarily at the younger ages. If this tendency should continue, the point will be reached before too long where large changes in generation mortality will become severely limited.

The examination of differences in cohort and period survivorship for four time periods by color and sex has shown that, generally speaking, past period life tables have not represented the actual mortality experience of a birth cohort. This is largely true because mortality rates have decreased over time so that each birth cohort is exposed to more favorable mortality rates throughout its lifetime than those prevailing at the time of its birth. Thus, to the extent that mortality rates improved, the period survivorship tables gave values that were too low. However, mortality at the older ages is no longer declining very much in the successive cohorts. In the last few years, the mortality rates after age 35 of later born

cohorts have risen above those of earlier born cohorts, demonstrating a crossover effect. Also, the age range in which substantial improvements in mortality are possible is narrowing. At the moment, this age range is from birth to about age 30 years. Unless major breakthroughs are achieved, further declines in mortality will be small compared with past improvements. From this it follows that future period life tables should become better predictors of the mortality experience of a cohort than past period life tables have been. This should be more true of whites than of races other than white and more true for females than for males.

REFERENCES

- ¹Kermack, W. O. McKendrick, A. G., and McKinlay, P. L.: Death rates in Great Britain and Sweden. *Lancet* 1:698-703, Mar. 31, 1934.
- ²Case, R. A. M.: Cohort analysis of mortality rates as an historical or narrative technique. *Brit. J. Prev. & Social Med.* 10:159-171, 1956.
- ³Pearson, K.: The intensity of natural selection in man. *Proc. Roy. Soc. London s. B* 85:469-476, 1912.
- ⁴Frost, W. H.: Age selection of mortality from tuberculosis in successive decades. *Am. J. Hyg.* 30, No. 3, Section A: 91-96, Nov. 1939.
- ⁵Picken, R. M. F.: Age selection of mortality from tuberculosis in successive decades. *Pub. Health, London* 53:145-148, Apr. 1940.
- ⁶Spicer, C. C.: The generation method of analysis applied to mortality from respiratory tuberculosis. *J. Hyg.* 52:361-368, 1954.
- ⁷Korteweg, R.: The age curve in lung cancer. *Brit. J. Cancer* 5:21-27, 1951.
- ⁸Stocks, P.: Studies of cancer death rates at different ages in England and Wales in 1921 to 1950: Uterus, breast and lung. *Brit. J. Cancer* 7:283-302, 1953.
- ⁹Stocks, P.: A study of the age curve for cancer of the stomach in connection with a theory of the cancer producing mechanism. *Brit. J. Cancer* 7:407-417, 1953.
- ¹⁰Haenszel, W., and Shimkin, M.: Smoking patterns and epidemiology of lung cancer in the United States: Are they compatible? *J. Nat. Cancer Inst.* 16(6): 1417-1441, 1956.
- ¹¹Cutler, S. J., and Loveland, D. B.: The risk of developing lung cancer and its relationship to smoking. *J. Nat. Cancer Inst.* 15 (1):201-211, 1954.
- ¹²Yates, P. O.: A change in the pattern of cerebrovascular disease. *Lancet*, pp. 65-69, Jan. 11, 1964.
- ¹³Dublin, L. I., and Spiegelman, M.: Current versus generation life tables. *Human Biol.* 13 (4):439-458, 1941.
- ¹⁴Spiegelman, M.: Segmented generation mortality. *Demography* 6(2):117-123, 1969.
- ¹⁵National Center for Health Statistics: Mortality in persons 15-44 years of age, United States, 1960 and 1968. *Monthly Vital Statistics Report*, Vol. 20, No. 9, Supplement. DHEW Pub. No. (HSM) 72-1128. Washington. U.S. Government Printing Office, Dec. 8, 1971.
- ¹⁶Bolander, A. M.: A study of cohort mortality in the past hundred years. Three studies on generation mortality of Sweden. Presented at the International Conference of the International Union for the Scientific Study of Population, London, Sept. 1969.
- ¹⁷Spiegelman, M.: *Introduction to Demography*, revised ed. Cambridge. Harvard University Press, 1968. p. 153.
- ¹⁸Beers, H. S.: Six term formulas for routine actuarial interpolation. *Record Am. Inst. Actuaries*, pp. 245-260, Nov. 1944; Modified interpolation formulas that minimize fourth differences. *Record Am. Inst. Actuaries*, pp. 14-61, June 1945.
- ¹⁹U.S. Bureau of the Census: U.S. Life Tables 1890, 1901, 1910, and 1901-1910. Washington. U.S. Government Printing Office, 1921.



LIST OF DETAILED TABLES

		Page
Table 1.	Cohort survivorship (l_x) from birth to age 70 of white males and females born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968-----	20
2.	Cohort survivorship (l_x) from birth to age 61 of white males and females born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968-----	22
3.	Cohort survivorship (l_x) from birth to age 51 of white males and females born 1918-1922 compared with corresponding period survivorship: death-registration States, 1900-1968-----	24
4.	Cohort survivorship (l_x) from birth to age 41 of white males and females born 1928-1932 compared with corresponding period survivorship: death registration States, 1900-1968-----	25
5.	Cohort survivorship (l_x) from birth to age 70 of males and females, other than white, born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968-----	26
6.	Cohort survivorship (l_x) from birth to age 61 of males and females, other than white, born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968-----	28
7.	Cohort survivorship (l_x) from birth to age 51 of males and females, other than white, born 1918-1922 compared with corresponding period survivorship: death-registration States, 1900-1968-----	30
8.	Cohort survivorship (l_x) from birth to age 41 of males and females, other than white, born 1928-1932 compared with corresponding period survivorship: death-registration States, 1900-1968-----	31

Table 1. Cohort survivorship (l_x) from birth to age 70 of white males and females born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	86,686	86,686	88,939	88,939
2-----	83,716	83,661	86,164	86,124
3-----	82,503	82,356	84,995	84,858
4-----	81,765	81,533	84,268	84,055
5-----	81,242	80,941	83,755	83,450
6-----	80,822	80,460	83,354	82,972
7-----	80,489	80,063	83,035	82,581
8-----	80,216	79,734	82,773	82,260
9-----	79,994	79,458	82,560	81,993
10-----	79,808	79,221	82,383	81,765
11-----	79,640	79,009	82,225	81,561
12-----	79,480	78,808	82,075	81,367
13-----	79,321	78,606	81,929	81,170
14-----	79,152	78,390	81,775	80,957
15-----	78,964	78,151	81,603	80,718
16-----	78,704	77,880	81,366	80,446
17-----	78,396	77,572	81,092	80,136
18-----	78,038	77,228	80,777	79,790
19-----	77,654	76,851	80,432	79,415
20-----	77,259	76,446	80,062	79,015
21-----	76,955	76,012	79,745	78,592
22-----	76,661	75,550	79,433	78,145
23-----	76,373	75,066	79,129	77,677
24-----	76,083	74,567	78,823	77,192
25-----	75,795	74,058	78,520	76,691
26-----	75,508	73,541	78,221	76,175
27-----	75,216	73,016	77,923	75,645
28-----	74,924	72,483	77,632	75,103
29-----	74,633	71,941	77,350	74,553
30-----	74,343	71,388	77,072	73,997
31-----	74,056	70,824	76,801	73,435
32-----	73,769	70,250	76,537	72,868
33-----	73,476	69,665	76,264	72,296
34-----	73,174	69,068	76,008	71,718
35-----	72,856	68,458	75,737	71,136
36-----	72,527	67,835	75,465	70,550
37-----	72,192	67,200	75,191	69,960
38-----	71,854	66,552	74,920	69,364
39-----	71,515	65,890	74,658	68,759
40-----	71,169	65,214	74,399	68,143
41-----	70,807	64,523	74,136	67,514
42-----	70,427	63,816	73,867	66,871
43-----	70,026	63,093	73,590	66,215
44-----	69,606	62,355	73,305	65,549
45-----	69,164	61,601	73,007	64,875
46-----	68,699	60,829	72,699	64,193
47-----	68,204	60,037	72,378	63,499
48-----	67,673	59,221	72,042	62,785
49-----	67,098	58,380	71,689	62,040

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 1. Cohort survivorship (l_x) from birth to age 70 of white males and females born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968—Con.

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
50-----	66,478	57,512	71,318	61,256
51-----	65,812	56,618	70,926	60,432
52-----	65,097	55,696	70,510	59,564
53-----	64,327	54,742	70,070	58,656
54-----	63,505	53,747	69,610	57,706
55-----	62,629	52,702	69,126	56,713
56-----	61,699	51,600	68,620	55,674
57-----	60,707	50,438	68,084	54,583
58-----	59,645	49,216	67,512	53,438
59-----	58,514	47,940	66,902	52,237
60-----	57,306	46,618	66,246	50,982
61-----	56,016	45,255	65,536	49,675
62-----	54,647	43,849	64,772	48,317
63-----	53,199	42,395	63,956	46,906
64-----	51,658	40,887	63,077	45,438
65-----	50,029	39,317	62,144	43,909
66-----	48,316	37,684	61,117	42,317
67-----	46,530	35,994	60,007	40,663
68-----	44,677	34,253	58,809	38,946
69-----	42,757	32,471	57,518	37,166
70-----	40,723	30,684	56,114	35,325

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 2. Cohort survivorship (l_x) from birth to age 61 of white males and females born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	89,712	89,712	91,510	91,510
2-----	87,452	87,390	89,410	89,349
3-----	86,512	86,372	88,527	88,403
4-----	85,959	85,733	88,009	87,792
5-----	85,555	85,267	87,634	87,348
6-----	85,211	84,880	87,323	86,988
7-----	84,897	84,562	87,038	86,690
8-----	84,632	84,298	86,802	86,441
9-----	84,398	84,075	86,596	86,230
10-----	84,188	83,880	86,410	86,045
11-----	83,997	83,700	86,240	85,875
12-----	83,829	83,525	86,100	85,710
13-----	83,665	83,344	85,964	85,540
14-----	83,499	83,149	85,825	85,358
15-----	83,322	82,933	85,674	85,157
16-----	83,129	82,690	85,506	84,932
17-----	82,916	82,416	85,320	84,680
18-----	82,684	82,111	85,116	84,401
19-----	82,440	81,776	84,900	84,097
20-----	82,189	81,414	84,676	83,770
21-----	81,933	81,024	84,443	83,419
22-----	81,677	80,607	84,207	83,045
23-----	81,421	80,171	83,974	82,652
24-----	81,168	79,724	83,744	82,245
25-----	80,917	79,273	83,516	81,829
26-----	80,666	78,819	83,290	81,404
27-----	80,422	78,361	83,069	80,969
28-----	80,185	77,897	82,857	80,526
29-----	79,954	77,422	82,655	80,074
30-----	79,727	76,934	82,471	79,615
31-----	79,494	76,431	82,290	79,149
32-----	79,250	75,914	82,110	78,675
33-----	78,976	75,380	81,930	78,192
34-----	78,681	74,827	81,749	77,697
35-----	78,390	74,255	81,566	77,189
36-----	78,103	73,662	81,379	76,667
37-----	77,821	73,049	81,192	76,132
38-----	77,548	72,417	81,005	75,586
39-----	77,278	71,769	80,819	75,032
40-----	76,992	71,107	80,630	74,472
41-----	76,688	70,429	80,434	73,906
42-----	76,361	69,734	80,228	73,331
43-----	76,011	69,019	80,011	72,745
44-----	75,636	68,281	79,783	72,142
45-----	75,265	67,516	79,543	71,519
46-----	74,829	66,722	79,289	70,873
47-----	74,361	65,915	79,019	70,203
48-----	73,850	65,061	78,730	69,506
49-----	73,285	64,178	78,417	68,778

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 2. Cohort survivorship (l_x) from birth to age 61 of white males and females born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968—Con.

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
50-----	72,661	63,266	78,077	68,016
51-----	71,971	62,325	77,708	67,217
52-----	71,216	61,353	77,309	66,380
53-----	70,401	60,346	76,879	65,502
54-----	69,524	59,296	76,418	64,579
55-----	68,571	58,192	75,924	63,604
56-----	67,539	57,024	75,389	62,568
57-----	66,438	55,786	74,812	61,462
58-----	65,243	54,474	74,216	60,282
59-----	63,968	53,095	73,554	59,032
60-----	62,605	51,660	72,846	57,719
61-----	61,130	50,176	72,075	56,349

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 3. Cohort survivorship (l_x) from birth to age 51 of white males and females born 1918-1922 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	92,322	92,322	93,898	93,898
2-----	90,823	90,576	92,533	92,288
3-----	90,149	89,750	91,926	91,514
4-----	89,737	89,196	91,549	90,979
5-----	89,431	88,769	91,269	90,573
6-----	89,174	88,406	91,038	90,233
7-----	88,953	88,092	90,849	89,951
8-----	88,760	87,819	90,692	89,713
9-----	88,593	87,580	90,561	89,507
10-----	88,445	87,366	90,448	89,321
11-----	88,311	87,168	90,346	89,145
12-----	88,185	86,975	90,250	88,970
13-----	88,055	86,776	90,153	88,787
14-----	87,920	86,559	90,051	88,589
15-----	87,773	86,312	89,941	88,369
16-----	87,610	86,028	89,821	88,120
17-----	87,434	85,702	89,692	87,836
18-----	87,250	85,336	89,558	87,514
19-----	87,064	84,934	89,422	87,153
20-----	86,877	84,502	89,288	86,753
21-----	86,663	84,038	89,158	86,313
22-----	86,370	83,540	89,027	85,835
23-----	85,862	83,013	88,895	85,326
24-----	85,192	82,466	88,765	84,795
25-----	84,591	81,905	88,639	84,250
26-----	84,100	81,335	88,520	83,694
27-----	83,720	80,757	88,406	83,128
28-----	83,477	80,170	88,301	82,556
29-----	83,321	79,573	88,198	81,980
30-----	83,163	78,963	88,097	81,401
31-----	83,001	78,341	87,998	80,821
32-----	82,837	77,708	87,897	80,241
33-----	82,671	77,070	87,795	79,663
34-----	82,504	76,433	87,690	79,089
35-----	82,334	75,802	87,582	78,523
36-----	82,156	75,179	87,468	77,962
37-----	81,966	74,562	87,347	77,407
38-----	81,761	73,948	87,217	76,856
39-----	81,537	73,334	87,077	76,308
40-----	81,291	72,716	86,925	75,760
41-----	81,022	72,092	86,760	75,212
42-----	80,727	71,460	86,579	74,661
43-----	80,400	70,816	86,376	74,102
44-----	80,040	70,155	86,159	73,528
45-----	79,566	69,473	85,919	72,932
46-----	79,127	68,767	85,655	72,312
47-----	78,643	68,035	85,365	71,665
48-----	78,109	67,278	85,049	70,989
49-----	77,519	66,497	84,707	70,284
50-----	76,872	65,692	84,337	69,550
51-----	76,157	64,862	83,933	68,784

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 4. Cohort survivorship (l_x) from birth to age 41 of white males and females born 1928-1932 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	White males		White females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	93,886	93,886	95,145	95,145
2-----	92,967	92,900	94,324	94,257
3-----	92,531	92,404	93,937	93,815
4-----	92,244	92,068	93,680	93,511
5-----	92,026	91,803	93,483	93,278
6-----	91,828	91,564	93,311	93,064
7-----	91,662	91,358	93,173	92,888
8-----	91,525	91,178	93,063	92,742
9-----	91,410	91,018	92,975	92,617
10-----	91,312	90,872	92,902	92,506
11-----	91,228	90,735	92,839	92,402
12-----	91,144	90,599	92,780	92,298
13-----	91,047	90,458	92,720	92,189
14-----	90,931	90,305	92,658	92,070
15-----	90,812	90,135	92,593	91,937
16-----	90,694	89,943	92,524	91,786
17-----	90,562	89,728	92,453	91,615
18-----	90,409	89,492	92,381	91,423
19-----	90,255	89,240	92,308	91,211
20-----	90,081	88,976	92,237	90,981
21-----	89,883	88,699	92,167	90,731
22-----	89,655	88,408	92,100	90,461
23-----	89,417	88,107	92,034	90,176
24-----	89,200	87,799	91,969	89,883
25-----	89,018	87,488	91,904	89,586
26-----	88,859	87,174	91,840	89,287
27-----	88,719	86,857	91,776	88,985
28-----	88,586	86,535	91,710	88,680
29-----	88,454	86,207	91,640	88,369
30-----	88,320	85,872	91,566	88,051
31-----	88,182	85,529	91,488	87,727
32-----	88,038	85,176	91,406	87,396
33-----	87,886	84,810	91,317	87,056
34-----	87,723	84,426	91,220	86,705
35-----	87,546	84,020	91,116	86,341
36-----	87,354	83,593	91,001	85,963
37-----	87,143	83,143	90,877	85,571
38-----	86,912	82,669	90,743	85,165
39-----	86,658	82,171	90,596	84,745
40-----	86,380	81,648	90,433	84,310
41-----	86,074	81,098	90,253	83,859

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 5. Cohort survivorship (l_x) from birth to age 70 of males and females, other than white, born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	75,511	75,511	78,985	78,985
2-----	69,158	69,186	73,079	73,131
3-----	66,696	66,815	70,581	70,482
4-----	65,406	65,552	69,255	69,126
5-----	64,593	64,682	68,269	68,113
6-----	63,916	63,976	67,556	67,367
7-----	63,403	63,423	66,992	66,751
8-----	63,006	62,975	66,539	66,228
9-----	62,687	62,599	66,168	65,764
10-----	62,420	62,253	65,846	65,320
11-----	62,197	61,919	65,556	64,882
12-----	61,975	61,559	65,278	64,420
13-----	61,734	61,170	64,970	63,917
14-----	61,472	60,742	64,630	63,367
15-----	61,158	60,280	64,222	62,775
16-----	60,693	59,761	63,661	62,132
17-----	60,141	59,193	63,012	61,446
18-----	59,533	58,576	62,316	60,737
19-----	58,889	57,928	61,599	60,033
20-----	58,241	57,263	60,892	59,351
21-----	57,659	56,581	60,238	58,688
22-----	57,042	55,895	59,570	58,040
23-----	56,404	55,201	58,891	57,404
24-----	55,748	54,507	58,207	56,775
25-----	55,095	53,811	57,525	56,150
26-----	54,433	53,120	56,839	55,527
27-----	53,761	52,431	56,153	54,907
28-----	53,111	51,744	55,498	54,290
29-----	52,492	51,064	54,883	53,669
30-----	51,890	50,388	54,297	53,038
31-----	51,300	49,721	53,737	52,388
32-----	50,718	49,059	53,196	51,726
33-----	50,154	48,402	52,677	51,048
34-----	49,580	47,749	52,167	50,374
35-----	48,974	47,097	51,634	49,714
36-----	48,339	46,446	51,066	49,073
37-----	47,695	45,794	50,483	48,439
38-----	47,070	45,138	49,906	47,802
39-----	46,477	44,476	49,353	47,148
40-----	45,893	43,800	48,809	46,449
41-----	45,301	43,106	48,241	45,701
42-----	44,690	42,395	47,632	44,905
43-----	44,065	41,666	47,007	44,064
44-----	43,447	40,912	46,396	43,206
45-----	42,831	40,144	45,807	42,351
46-----	42,209	39,347	45,226	41,494
47-----	41,569	38,528	44,627	40,641
48-----	40,914	37,678	44,016	39,782
49-----	40,237	36,808	43,392	38,908

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 5. Cohort survivorship (l_x) from birth to age 70 of males and females, other than white, born 1899-1903 compared with corresponding period survivorship: death-registration States, 1900-1968—Con.

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
50-----	39,519	35,916	42,743	38,022
51-----	38,755	35,008	42,053	37,120
52-----	37,950	34,087	41,321	36,209
53-----	37,130	33,147	40,576	35,287
54-----	36,322	32,181	39,850	34,338
55-----	35,508	31,173	39,124	33,358
56-----	34,681	30,111	38,386	32,348
57-----	33,821	28,990	37,618	31,282
58-----	32,940	27,820	36,833	30,175
59-----	32,046	26,650	36,048	29,049
60-----	31,115	25,497	35,228	27,910
61-----	30,114	24,363	34,335	26,786
62-----	29,051	23,254	33,366	25,667
63-----	27,962	22,169	32,368	24,554
64-----	26,838	21,080	31,367	23,439
65-----	25,664	19,980	30,349	22,320
66-----	24,396	18,879	29,260	21,196
67-----	23,057	17,768	28,078	20,072
68-----	21,683	16,665	26,842	18,942
69-----	20,331	15,567	25,613	17,822
70-----	18,948	14,478	24,377	16,706

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 6. Cohort survivorship (l_x) from birth to age 61 of males and females, other than white, born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	88,429	88,429	90,114	90,114
2-----	83,895	83,433	85,925	85,494
3-----	82,184	81,403	84,268	83,489
4-----	81,278	80,227	83,349	82,379
5-----	80,725	79,448	82,766	81,554
6-----	80,216	78,777	82,285	80,935
7-----	79,756	78,254	81,804	80,400
8-----	79,406	77,820	81,416	79,932
9-----	79,130	77,453	81,096	79,514
10-----	78,891	77,123	80,821	79,127
11-----	78,670	76,804	80,574	78,747
12-----	78,463	76,474	80,354	78,348
13-----	78,237	76,112	80,118	77,901
14-----	77,976	75,700	79,835	77,382
15-----	77,667	75,226	79,477	76,771
16-----	77,297	74,682	79,027	76,060
17-----	76,863	74,059	78,492	75,257
18-----	76,377	73,372	77,889	74,395
19-----	75,840	72,642	77,247	73,525
20-----	75,255	71,889	76,588	72,678
21-----	74,622	71,117	75,919	71,858
22-----	73,961	70,329	75,264	71,056
23-----	73,292	69,533	74,623	70,271
24-----	72,634	68,734	73,999	69,498
25-----	71,977	67,936	73,387	68,733
26-----	71,303	67,140	72,776	67,977
27-----	70,616	66,346	72,164	67,232
28-----	69,944	65,553	71,566	66,491
29-----	69,299	64,758	70,996	65,748
30-----	68,689	63,958	70,460	64,996
31-----	68,096	63,154	69,944	64,230
32-----	67,501	62,343	69,425	63,448
33-----	66,911	61,519	68,897	62,649
34-----	66,336	60,673	68,381	61,838
35-----	65,846	59,804	67,890	60,975
36-----	65,333	58,911	67,420	60,152
37-----	64,830	57,997	66,963	59,323
38-----	64,320	57,067	66,506	58,483
39-----	63,804	56,125	66,042	57,624
40-----	63,269	55,174	65,560	56,740
41-----	62,703	54,212	65,052	55,823
42-----	62,109	53,238	64,521	54,870
43-----	61,491	52,251	63,977	53,885
44-----	60,859	51,254	63,436	52,879
45-----	60,213	50,250	62,889	51,863
46-----	59,540	49,241	62,322	50,841
47-----	58,828	48,225	61,728	49,810
48-----	58,069	47,193	61,105	48,762
49-----	57,271	46,129	60,456	47,686

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 6. Cohort survivorship (l_x) from birth to age 61 of males and females, other than white, born 1908-1912 compared with corresponding period survivorship: death-registration States, 1900-1968—Con.

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
50-----	56,433	45,027	59,784	46,573
51-----	55,533	43,880	59,076	45,423
52-----	54,549	42,697	58,319	44,241
53-----	53,505	41,485	57,511	43,030
54-----	52,427	40,255	56,693	41,794
55-----	51,301	39,011	55,861	40,532
56-----	50,110	37,748	55,007	39,239
57-----	48,837	36,459	54,118	37,905
58-----	47,492	35,144	53,170	36,543
59-----	46,094	33,823	52,174	35,172
60-----	44,666	32,507	51,132	33,815
61-----	43,137	31,211	50,020	32,483

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 7. Cohort survivorship (l_x) from birth to age 51 of males and females, other than white, born 1918-1922 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	89,935	89,935	91,650	91,650
2-----	87,240	86,848	89,227	88,844
3-----	86,217	85,586	88,272	87,651
4-----	85,670	84,875	87,710	86,901
5-----	85,279	84,375	87,328	86,365
6-----	84,925	83,883	86,960	85,869
7-----	84,648	83,503	86,676	85,474
8-----	84,422	83,201	86,459	85,151
9-----	84,236	82,954	86,290	84,874
10-----	84,072	82,723	86,150	84,618
11-----	83,912	82,491	86,018	84,360
12-----	83,739	82,238	85,872	84,075
13-----	83,545	81,945	85,690	83,741
14-----	83,322	81,596	85,472	83,337
15-----	83,066	81,176	85,208	82,844
16-----	82,768	80,672	84,887	82,250
17-----	82,425	80,076	84,507	81,551
18-----	82,044	79,384	84,082	80,758
19-----	81,635	78,601	83,628	79,893
20-----	81,210	77,736	83,174	78,976
21-----	80,762	76,785	82,727	78,012
22-----	80,264	75,756	82,283	77,004
23-----	79,740	74,687	81,841	75,974
24-----	79,210	73,627	81,413	74,947
25-----	78,718	72,610	81,014	73,938
26-----	78,276	71,651	80,644	72,957
27-----	77,877	70,741	80,300	72,002
28-----	77,499	69,862	79,972	71,066
29-----	77,133	68,986	79,651	70,136
30-----	76,760	68,092	79,335	69,201
31-----	76,377	67,173	79,024	68,258
32-----	75,987	66,235	78,715	67,310
33-----	75,597	65,281	78,404	66,355
34-----	75,219	64,325	78,096	65,368
35-----	74,839	63,376	77,783	64,405
36-----	74,455	62,435	77,460	63,440
37-----	74,065	61,500	77,120	62,473
38-----	73,648	60,572	76,760	61,508
39-----	73,196	59,651	76,380	60,546
40-----	72,705	58,735	75,981	59,589
41-----	72,171	57,824	75,556	58,637
42-----	71,586	56,915	75,096	57,687
43-----	70,951	56,002	74,610	56,731
44-----	70,274	55,077	74,103	55,757
45-----	69,544	54,134	73,573	54,757
46-----	68,759	53,178	73,021	53,727
47-----	67,901	52,207	72,433	52,664
48-----	66,980	51,222	71,804	51,569
49-----	65,999	50,222	71,136	50,444
50-----	64,962	49,206	70,441	49,293
51-----	63,842	48,170	69,702	48,115

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

Table 8. Cohort survivorship (l_x) from birth to age 41 of males and females, other than white, born 1928-1932 compared with corresponding period survivorship: death-registration States, 1900-1968

Age x in years	All other males		All other females	
	Cohort l_x	Period l_x	Cohort l_x	Period l_x
0-----	100,000	100,000	100,000	100,000
1-----	90,888	90,888	92,552	92,552
2-----	89,024	88,950	90,923	90,859
3-----	88,261	88,133	90,239	90,115
4-----	87,827	87,666	89,836	89,674
5-----	87,534	87,323	89,557	89,334
6-----	87,280	86,988	89,311	89,009
7-----	87,087	86,732	89,127	88,760
8-----	86,933	86,529	88,990	88,568
9-----	86,803	86,357	88,882	88,414
10-----	86,687	86,197	88,791	88,279
11-----	86,573	86,033	88,708	88,143
12-----	86,456	85,849	88,620	87,985
13-----	86,325	85,633	88,514	87,782
14-----	86,175	85,375	88,384	87,514
15-----	86,005	85,067	88,232	87,167
16-----	85,814	84,703	88,059	86,732
17-----	85,613	84,277	87,872	86,209
18-----	85,394	83,786	87,674	85,610
19-----	85,154	83,228	87,466	84,955
20-----	84,882	82,604	87,256	84,258
21-----	84,582	81,909	87,052	83,522
22-----	84,254	81,148	86,857	82,749
23-----	83,916	80,339	86,672	81,949
24-----	83,585	79,507	86,491	81,136
25-----	83,272	78,671	86,315	80,320
26-----	82,964	77,837	86,137	79,505
27-----	82,664	77,003	85,956	78,692
28-----	82,365	76,165	85,768	77,879
29-----	82,060	75,317	85,569	77,063
30-----	81,746	74,453	85,358	76,241
31-----	81,425	73,573	85,135	75,413
32-----	81,092	72,678	84,900	74,578
33-----	80,739	71,763	84,651	73,729
34-----	80,355	70,821	84,381	72,859
35-----	79,928	69,850	84,085	71,962
36-----	79,456	68,650	83,765	71,038
37-----	78,909	67,822	83,422	70,089
38-----	78,347	66,771	83,057	69,117
39-----	77,741	65,700	82,672	68,121
40-----	77,082	64,610	82,242	67,099
41-----	76,354	63,500	81,783	66,047

NOTE: l_x refers to the number of survivors, of 100,000 born alive, at the beginning of age x.

APPENDIX I

RELATIONSHIP BETWEEN COHORT AND PERIOD MORTALITY

There are various ways in which age-specific mortality rates may be viewed. Following Spiegelman's¹⁷ presentations, the schematic diagram shown below depicts mortality rates observed over a period of years in three ways—period mortality or mortality rates for a calendar year, time trend of mortality by age, and the generation mortality or mortality for a cohort of individuals born in a particular calendar year.


If $q_{x,t}$ denotes the mortality rate at age x in calendar year t , then:

(1) The vertical lines represent the case where t is constant and x alone varies. In this case the mortality rates by age are for a calendar year. These are the period mortality rates.

(2) The horizontal lines represent the case where x is constant and t alone varies. In this case the observed time trend for age x is shown over a series of calendar years.

(3) The diagonal lines represent the case where both x and t jointly advance by the same unit interval of time, such that $t - x = \theta$ is a constant which defines the year of birth. In this case, the mortality rates are those for a generation traced from birth. Strictly speaking, deaths at age x as of the last birthday during the calendar year t will occur among births in calendar year $\theta - 1$ as well as in calendar year θ . Likewise, there will be deaths at age x in calendar year $t+1$ among births in year θ . To simplify the description, it is assumed that deaths are concentrated at the mid-age and at the middle of the calendar year.

NOTE: The list of references follows the text.



APPENDIX II

PRODUCTION OF THE SINGLE-YEAR DATA

The original data for the survivorship tables presented in this report were of two types: estimates of the population in 5-year age groups from age 0-4 to age 80-84 from 1900 through 1968, and age-specific deaths for the death-registration States during that same time period. In order to obtain deaths and population by single years of age, an interpolation procedure was used.

Interpolation as a generating procedure allows division of grouped data into smaller units. For example, interpolation is often used to produce single-year estimates of population or of deaths by smoothing a 5-year estimate into five single-year estimates. This is done by applying a set of constant multipliers to the 5-year data.

The interpolation formulas used to produce the data in this report are those derived by Beers.¹⁸ These formulas were chosen in preference to osculatory formulas since they are based on fifth differences and are more suitable for smoothing deaths and population, which may have unusual distributions.

Because such a procedure can produce some irregularities, the interpolated data were then recombined into 5-year age groups or birth cohorts in order to minimize any irregularities so produced. Interpolating the data in this way allows examination of the yearly mortality rates of a birth cohort, instead of a view of the cohort only at 5-year intervals.

NOTE: The list of references follows the text.

— ○ ○ ○ —

APPENDIX III

CONSTRUCTION OF THE COHORT AND PERIOD SURVIVORSHIP TABLES (DETAILED TABLES 1-8)

Published Tables

For the years 1901, 1910, 1920, and 1930, published period life tables were available. However, various problems of comparability arose which led to the construction of period life tables from the same set of interpolated data as that used for the cohort tables.

First, the age-specific mortality rates used in the published period life tables covered 3-year periods. The cohort tables used age-specific mortality rates covering 5-year periods. Second, the published period life tables covered only the death-registration States in the 3 years surrounding these dates. The 1910 period life table available in published sources was computed for the death-registration States of 1900. Thus, by using the same set of interpolated data for both sets of tables, the cohort and period tables in this report at least begin using the same death-registration States. However, it was not possible to eliminate the problem of new States coming into the death-registration area during the time covered by the cohort tables. For races other than white there is some reason to believe that this may have produced irregularities. As new States were added with different types and numbers of people of other races (for example, with predominately rural or urban populations), the mortality rates may show fluctuation that would not be expected on the basis of age alone.

Finally, the published period tables were available only for Negroes, while the cohort tables were available only for all races other than white. The error that might have been introduced by assuming these were the same, while probably not large, nevertheless was a factor in the decision to construct both period and cohort tables from the same set of data. This is not to say that the period tables constructed from the interpolated data are any more correct than the published period tables for these dates. Such a procedure only makes the period tables more comparable to the cohort tables.

Method of Estimating Death Rates at Age 0-1

Death rates of infants aged 0-1 year in the population are generally computed by dividing the number of deaths of children of that age by the number of births in that year. The number of births are used in this calculation instead of the number of children aged 0-1 enumerated in the population since there is generally an undercount of infants in census data.

However, in constructing survivorship tables for the years included in this report, special problems arose in making an estimate of the births occurring in these years. The death-registration area and the birth-registration area were not the same prior to 1933. Consequently, even though some scant data were available to estimate births during these years, these data did not cover the same States as the death data available from the interpolated set.

The method of estimating births for these years was adapted from Glover.¹⁹ In this procedure, population and death estimates for older ages were used to construct estimates of births at an earlier time. For example, births in 1910 should be equal to the sum of the population aged 3-4 in 1913, added to the deaths of children aged 2-3 in 1912, deaths of children aged 1-2 in 1911, and deaths of children aged 0-1 in 1910. In symbols this procedure appears as follows:

$$B^{1910} = P_{3-4}^{1913} + D_{2-3}^{1912} + D_{1-2}^{1911} + D_{0-1}^{1910}$$

where B = births, P = population, and D = deaths.

Each of these estimates of population and deaths was adjusted by separation factors so as to include only those persons who were part of the birth cohort being estimated. This procedure was necessary since children aged 0-1 in 1910, for example, may not have been born in 1910. A child born in September of 1909 would still be age 0-1 in 1910. Likewise, children born in September of 1910 would still be age 0-1 in 1911.

NOTE: The list of references follows the text.

Separation factors were used to attempt to separate out those who were actually part of the cohort being estimated by considering infant mortality rates during the years in question. In order to calculate such separation factors, deaths by month of age were necessary. Monthly mortality data show what proportion of the infant deaths during a given year were of children actually born in that year versus what proportion of those deaths were of children born in the previous year.

The table shows the separation factors used for each year by color, sex, and age. The 1900 and 1910 estimates for whites were available from published sources. The 1930 white and other than white estimates were also available. The 1920 estimates for whites and people of other races were calculated to be congruent with the other sets. For the years 1900 and 1910 estimates for races other than white were similarly calculated.

One additional adjustment to these birth estimates was necessary in order to construct estimates of deaths of infants aged 0-1. This adjustment was made in order to allow for a changing death-registration area during the time periods in question. In the above estimate of births in 1910, for example, Kentucky and Missouri were admitted to the death-registration area in 1911, and Virginia was admitted in 1913. Consequently, the estimate of deaths for those 0-1 year of age in 1910 did not include those children in Kentucky and Missouri who died at that age and were part of the 1911 birth cohort. The population estimate of those aged 3-4 in 1913 included the children in Virginia at that age, but none of the death estimates in 1912, 1911, or 1910 included the children dying in this State before reaching age 3.

To compensate for this underestimation, data were obtained from the annual *Vital Statistics* volumes on mortality by color, sex, and single years of age under 5 in those States which entered the registration area during one of the birth estimation periods. In the above estimate, for example, the deaths of those children aged 2-3, 1-2, and 0-1 in Virginia in 1913—the first year for which such data were available for Virginia—were added to the birth estimate. These figures are only an approximation since they are not actually for the year in question. They do constitute, however, a needed, and perhaps not grossly inaccurate, adjustment.

Separation factors used in estimates, by color, sex, and age: percent dying from cohort born in previous year

Year and age	White		All other	
	Male	Female	Male	Female
1900 and 1910:				
0-----	.28	.29	.31	.32
1-----	.41	.41	.41	.41
2-----	.47	.47	.47	.47
3-----	.48	.48	.48	.48
4-----	.48	.48	.48	.48
1920:				
0-----	.23	.24	.26	.27
1-----	.41	.41	.41	.41
2-----	.47	.47	.47	.47
3-----	.48	.48	.48	.48
4-----	.48	.48	.48	.48
1930:				
0-----	.19	.20	.21	.22
1-----	.41	.41	.41	.41
2-----	.47	.47	.47	.47
3-----	.48	.48	.48	.48
4-----	.48	.48	.48	.48

SOURCES: 1900 and 1910 estimates for whites from M. Spiegelman, *Introduction to Demography*, Chicago, The Society of Actuaries, 1955, p. 75. 1930 estimates from *U.S. Life Tables and Actuarial Tables 1939-41* (1947), p. 118. 1920 estimates for whites and all others and 1900 and 1910 estimates for all others calculated to be congruent with other sets.

Deaths at Other Ages and Survivors

As noted above, the number of deaths in each year were combined in different ways to produce the cohort and period death rates for survivorship tables. In order to produce the cohort tables, the data were combined by 5-year birth cohorts (as shown earlier in table A) to produce mortality rates by single years of age for a succession of birth dates. For a given succession of birth dates (cohort), the particular single year of age determines the calendar years of data to be combined

in computing the cohort's average age-specific death rate. The 1908-1912 cohort, for example, was 50 years old in 1958-1962, so the deaths at age 50 in this latter period of years were combined in computing their average death rate at age 50. Their death rate at age 51 was based on deaths occurring in 1959-1963, at age 52 in 1960-1964, and at other ages in like fashion. To produce the period tables, the average single-year age-specific mortality rates were computed from the same 5-year period of data, irrespective of age, with the central year the same as the central year of birth of the cohorts being examined. For example, the period rates for 1908-1912, like the 1908-1912 cohort rates, have a central year of 1910. But unlike the 1908-1912 cohort death rate at age 50, the corresponding period rate is based on death and population at age 50 in 1908-1912. The period rate at age 51 and all other ages is based on 1908-1912 data. The period rates are thus based on the average age-specific death rate prevailing at one 5-year period in time.

The death rates, m_x , produced by the combination procedures were converted into life table death rates, q_x . This procedure was necessary since the death rate m_x was calculated for those alive at the midpoint of the age interval. For life table purposes, the q_x death rate shows the death rate for those alive at the beginning of the age interval. This conversion was accomplished by use of the approximation formula:

$$q_x = \frac{m_x}{1 + 1/2 m_x}$$

After the death rates were computed, the number dying at each year of age was obtained by multiplying the number alive at the beginning of that age interval (l_x) by these death rates (q_x). The result is the number dying during that year of age (d_x). Beginning with 100,000 alive at age 0, the number surviving to each successive age was then computed by subtraction (l_x).



VITAL AND HEALTH STATISTICS PUBLICATION SERIES

Formerly Public Health Service Publication No. 1000

- Series 1. Programs and collection procedures.*—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions, data collection methods used, definitions, and other material necessary for understanding the data.
- Series 2. Data evaluation and methods research.*—Studies of new statistical methodology including: experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, contributions to statistical theory.
- Series 3. Analytical studies.*—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and committee reports.*—Final reports of major committees concerned with vital and health statistics, and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data from the Health Interview Survey.*—Statistics on illness, accidental injuries, disability, use of hospital, medical, dental, and other services, and other health-related topics, based on data collected in a continuing national household interview survey.
- Series 11. Data from the Health Examination Survey.*—Data from direct examination, testing, and measurement of national samples of the civilian, noninstitutional population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics; and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data from the Institutional Population Surveys.*—Statistics relating to the health characteristics of persons in institutions, and their medical, nursing, and personal care received, based on national samples of establishments providing these services and samples of the residents or patients.
- Series 13. Data from the Hospital Discharge Survey.*—Statistics relating to discharged patients in short-stay hospitals, based on a sample of patient records in a national sample of hospitals.
- Series 14. Data on health resources: manpower and facilities.*—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on mortality.*—Various statistics on mortality other than as included in regular annual or monthly reports—special analyses by cause of death, age, and other demographic variables, also geographic and time series analyses.
- Series 21. Data on natality, marriage, and divorce.*—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports—special analyses by demographic variables, also geographic and time series analyses, studies of fertility.
- Series 22. Data from the National Natality and Mortality Surveys.*—Statistics on characteristics of births and deaths not available from the vital records, based on sample surveys stemming from these records, including such topics as mortality by socioeconomic class, hospital experience in the last year of life, medical care during pregnancy, health insurance coverage, etc.

For a list of titles of reports published in these series, write to:

Office of Information
National Center for Health Statistics
Public Health Service, HSMHA
Rockville, Md. 20852

DHEW Publication No. (HSM) 73-14 00
Series 3 - No. 16

**U.S. DEPARTMENT OF
HEALTH, EDUCATION, AND WELFARE**

Public Health Service
HEALTH SERVICES AND MENTAL HEALTH ADMINISTRATION
5600 Fishers Lane
Rockville, Maryland 20852

OFFICIAL BUSINESS
Penalty for Private Use, \$300

POSTAGE AND FEES PAID
U.S. DEPARTMENT OF HEW

HEW 396



THIRD CLASS
BLK. RT.