

Series 10

No. 202



Vital and Health Statistics

From the CENTERS FOR DISEASE CONTROL AND PREVENTION / National Center for Health Statistics

Injury and Poisoning Episodes and Conditions: National Health Interview Survey, 1997

July 2000



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics



Copyright Information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested Citation

Warner M, Barnes PM, and Fingerhut LA. Injury and poisoning episodes and conditions; National Health Interview Survey, 1997. *Vital Health Stat* 10(202). 2000.

Library of Congress-in-Publication Data

Injury and poisoning episodes and conditions: National Health Interview Survey, 1997.

p.; cm.—(Vital and health statistics. Series 10, Data from the National Health Survey ; no. 202) (DHHS publication ; no. (PHS) 2000-1530) "June 2000."

Includes bibliographical references.

ISBN 0-8406-0563-3

1. Wounds and injuries—United States—Statistics. 2. Poisoning—United States—Statistics. 3. United States—Statistics, Medical. I. National Center for Health Statistics (U.S.) II. Series. III. Series: DHHS publication ; no. (PHS) 2000-1530.

[DNLN: 1. National Health Interview Survey (U.S.) 2. Wounds and Injuries—United States—Statistics. 3. Health Surveys—United States.

4. Poisoning—United States—Statistics. W2 A N148VJ no. 202 2000] RA407.3.A346 no. 202

[RA645.T73]

362.1'0973'021 s—dc21

[617.1'00973021]

00-041852

For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.

Vital and Health Statistics

Injury and Poisoning Episodes and Conditions: National Health Interview Survey, 1997

Series 10:
Data From the National Health
Interview Survey
No. 202

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
July 2000
DHHS Publication No. (PHS) 2000-1530

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for International Statistics*

Jennifer H. Madans, Ph.D., *Associate Director for Science*

Lester R. Curtin, Ph.D., *Acting Associate Director for Research and Methodology*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Analysis, Epidemiology, and Health Promotion*

P. Douglas Williams, *Acting Associate Director for Data Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Vital and Health Statistics Systems*

Douglas L. Zinn, *Acting Associate Director for Management*

Charles J. Rothwell, *Associate Director for Data Processing and Services*

Office of Analysis, Epidemiology, and Health Promotion

Jennifer H. Madans, Ph.D., *Acting Associate Director for Analysis, Epidemiology, and Health Promotion*

Lois A. Fingerhut, M.A., *Special Assistant for Injury Epidemiology*

Division of Health Interview Statistics

Jane F. Gentleman, Ph.D., *Director*

Susan S. Jack, M.S., *Acting Chief, Illness and Disability Statistics Branch*

Cooperation of the U.S. Bureau of the Census

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the U.S. Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

Contents

Abstract	1
Highlights.....	1
Introduction	2
Methods	2
Source of Data	2
Capturing Episodes of Injury and Poisoning	2
Analysis Fields	4
Sample Size and Response Rates	4
Data Files	4
Units of Analysis, Annual Estimates, and Rates	4
Variance Estimation and Significance Testing	5
Age-adjusted Rates	5
Results	6
Age, Sex, and Race/Ethnicity	6
Leading External Causes of Injury	6
Injury Conditions	8
Place of Injury	8
Activity at Time of Injury	9
Hospitalization	9
School- and Work-loss Days	9
Limitations After Injury	10
Discussion	10
Strengths of the Redesign	10
Limitations and Lessons Learned	11
Comparability with Prior Years	12
Conclusion.....	13
References	13

Appendixes

I. NHIS Questionnaire, Family Core: Section II—Injuries	26
II. Matrix for External Cause of Injury Mortality and Morbidity Data	33
III. Injury Morbidity Matrix	34
Codes for Body Region of Injury	34
Codes for Nature of Injury	37
IV. Technical Notes	38

Detailed Tables

1. Number, percent distribution, standard error, and annual rate of injury and poisoning episodes by selected demographic characteristics: United States, 1997	16
---	----

2.	Annual rate and standard error of leading external causes of injury and poisoning episodes by sex and age: United States, 1997	18
3.	Number, percent, and standard error of fall-related episodes and percent by type of fall and sex: United States, 1997 . .	19
4.	Number, percent, and standard error of transportation-related episodes by selected characteristics and sex: United States, 1997	19
5.	Number of poisoning episodes, number and percent of poisoning episodes involving a call to a poison control center, and standard error by sex and age: United States, 1997	20
6.	Annual rate and standard error of conditions by the nature of the injury and sex and age: United States, 1997	21
7.	Annual rate and standard error of injury conditions by body region injured and sex and age: United States, 1997	22
8.	Number, percent, and standard error of injury episodes by place of occurrence, activity engaged in, and sex: United States, 1997	23
9.	Number, percent, and standard error of injury and poisoning episodes involving hospitalization by sex and age: United States, 1997	24
10.	Number, annual rates, and standard error of injury episodes resulting in time lost from work or school: United States, 1997	24
11.	Number, percent, and standard error of injury episodes by whether persons required help with daily routine activities or personal care and age and sex: United States, 1997	25

List of Figures

1.	Injury and poisoning episode rates by age: United States, 1997	5
2.	Injury and poisoning episode rates by age and sex: United States, 1997	6
3.	Injury and poisoning episode rates by age, race, and ethnicity: United States, 1997	6
4.	Age-adjusted external cause of injury and poisoning episode rates by mechanism of injury and sex: United States, 1997	7
5.	Injury episode rates for falls by age and sex: United States, 1997	7
6.	Injury episode rates for being struck by or against an object or person by age and sex: United States, 1997	8
7.	Injury episode rates for transportation-related injuries by age and sex: United States, 1997	8
8.	Injury episode rates for overexertion by age and sex: United States, 1997	9
9.	Injury episode rates for cutting and piercing injuries by age and sex: United States, 1997	9
10.	Poisoning episode rates by age and sex: United States, 1997	10
11.	Age-adjusted injury condition rates by nature of injury and sex: United States, 1997	10
12.	Injury condition rates for sprains and strains by age and sex: United States, 1997	11
13.	Injury condition rates for open wounds by age and sex: United States, 1997	11
14.	Injury condition rates for fractures by age and sex: United States, 1997	12
15.	Injury condition rates for contusions by age and sex: United States, 1997	12
16.	Age-adjusted injury condition rates by body region and sex: United States, 1997	13
17.	Upper extremity injury condition rates by age and sex: United States, 1997	13
18.	Lower extremity injury condition rates by age and sex: United States, 1997	14
19.	Percent of injury episodes by place of injury and sex: United States, 1997	14
20.	Percent of injury episodes by activity at the time of injury and sex: United States, 1997	15

Objective

This report provides a descriptive overview of the first year of data from the injury section of the redesigned National Health Interview Survey. It documents the Survey's design methodologies and presents detailed national estimates of nonfatal injury and poisoning episodes for 1997.

Methods

Data for the U.S. civilian noninstitutionalized population were collected using Computer Assisted Personal Interview (CAPI). The data on all medically attended injuries and poisonings occurring to any family member during the 3-month period prior to the interview were obtained from an adult member of the family.

Results

In 1997, 34.4 million medically-attended episodes of injury and poisoning were reported, resulting in an age-adjusted rate of 128.9 episodes per 1,000 persons. Injury episodes resulted in 40.9 million injury conditions for a rate of 153.7 conditions per 1,000 persons. Falls were the leading external cause of injury, followed by episodes resulting from being struck by or against a person or an object, transportation, overexertion, cutting and piercing instruments, and poisoning. Sprains and strains were the most frequently reported injury condition followed by open wounds, fractures, and contusions. Upper extremity and lower extremity injuries were the leading body regions for these conditions. Leisure activities and paid work were most often reported as the activities the person was engaged in when the injury episode occurred, and the home was the most likely place for the injury to have occurred.

Conclusion

The redesigned NHIS is a useful source of information about medically-attended nonfatal injuries and poisonings. A single year now provides enough data to produce stable national estimates on details of injury and poisoning episodes.

Keywords: *National Health Interview Survey • injury episodes • poisoning episodes • injury conditions*

Injury and Poisoning Episodes and Conditions: National Health Interview Survey, 1997

by Margaret Warner, Ph.D., Office of Analysis, Epidemiology, and Health Promotion; Patricia M. Barnes, M.A., Division of Health Interview Statistics; and Lois A. Fingerhut, M.A., Office of Analysis, Epidemiology, and Health Promotion

Highlights

- In 1997, there were 34.4 million medically-attended episodes of injury and poisoning reported among the U.S. civilian noninstitutionalized population at a rate of 129 episodes of injury and poisoning per 1,000 persons (similar for crude and adjusted rates). The injury and poisoning episodes resulted in 40.9 million conditions, at a rate of 154 conditions per 1,000 persons (similar for crude and adjusted rates).
- The age-adjusted injury and poisoning episode rate for males was 21% higher than the rate for females.
- The age-adjusted injury and poisoning episode rate was higher for non-Hispanic white persons than for either non-Hispanic black or Hispanic persons.
- In 1997, *falls* were the leading external cause of injury; 11.3 million episodes of falls were reported at an age-adjusted rate of 43.1 per 1,000 persons. *Falls* were followed by episodes resulting from *being struck by or against a person or an object, transportation, overexertion, cutting and piercing instruments, and poisoning*.
- In 1997, *sprains and strains* were the leading injury conditions reported followed by *open wounds, fractures and contusions*; the respective age-adjusted rates were 38.5, 29.0, 23.7, and 19.0 per 1,000 persons.
- The *home* was the most frequently reported place of injury, with 24% of injuries occurring *inside the home*, and another 18% *outside the home*. Females were located *inside the home* at the time of the injuries almost twice as often as males.
- *Leisure activities* and *paid work* were most often reported as the activities the person was engaged in when the injury episode occurred, accounting for 22% and 19%, respectively, of all episodes of injury. The percent of people engaged in *leisure activities* were similar for males and females, while males were engaged in *paid work* almost twice as often as females.
- Respondents reported that the injured was hospitalized in 2.5 million episodes or about 7% of the injury and poisoning episodes.
- After the injury, about 7% of the injured needed help in handling their *daily routine*, such as household chores and shopping, and about 6% needed help attending to their personal care such as eating and bathing. The percent of persons with limitations in both *personal care* and *daily routine* increased with age.

We would like to thank Nathaniel Schenker and Van Parsons of the Office of Research Methodology for their help with the technical aspects of this analysis and Jennifer Madans and Jane Gentleman for their constructive comments. This report was edited by Klaudia M. Cox, and typeset by Jacqueline M. Davis of the Publications Branch, Division of Data Services.

Introduction

The National Health Interview Survey (NHIS) has been used to make estimates of the frequency and rate of nonfatal injuries and poisonings in the civilian noninstitutionalized population since its inception in 1957. Prior to 1997, the injury and poisoning statistics extracted from the NHIS were limited because there was very little detail about the circumstances of the injuries and poisonings and because the annual sample sizes were, in general, too small to make reliable estimates with a single year of data (1). However, because the NHIS was potentially such a rich source of data about nonfatal injuries, and because there were no other comparable national data, the number of questions related to injuries and poisonings was expanded when the questionnaire was redesigned in 1997 in order to answer basic epidemiological questions about the injuries and the circumstances of the traumatic event causing the injuries.

National estimates of nonfatal injury rates in the United States are available from several data systems operated by the National Center for Health Statistics (NCHS). This includes the National Hospital Ambulatory Medical Care Survey and the National Ambulatory Medical Care Survey, which capture data from ambulatory care settings (2–4), and the National Hospital Discharge Survey, which captures data from inpatient settings (5,6). Data from these sample surveys are based on medical records, rather than personal interviews. The ambulatory care surveys obtain data on both the external cause and the diagnosis of injury-related visits to emergency departments, outpatient clinics, and physicians' offices. The inpatient hospital survey obtains detailed diagnostic information on injuries, but relatively poor information on the external causes. None of the surveys have detailed information on the circumstances of the injury or on other relevant variables, such as place or activity at the time of the injury.

The National Electronic Injury Surveillance System (NEISS) from the

U.S. Consumer Product Safety Commission is another national source of nonfatal injury data (7,8). The estimates from NEISS are of consumer product-related injuries treated in a sample of emergency departments and are based on information abstracted from medical records. Consumer product injuries are those resulting from products such as toys, appliances, and lawn mowers. It also includes injuries where any kind of sports equipment is mentioned (8). Certain products, such as firearms and motor vehicles, are specifically excluded from the NEISS system. However, work is currently underway to expand the scope of NEISS to include all injuries treated in emergency departments (9).

The NHIS injury estimates for 1997 are not directly comparable to estimates of injury conditions or episodes from prior years of the NHIS. When the NHIS questionnaire was redesigned in 1997, extensive changes were made in the way injury and poisoning data were collected. Differences between 1997 and previous years include changes to: a) the questions that screen for injuries and poisonings; b) the inclusion criteria; c) the recall period; d) the severity threshold; and e) the phrasing, placement, and number of questions. The changes were introduced to increase the level of detail available on injury episodes and to enable the user to make more reliable annual estimates by increasing the sample size.

This report presents detailed national estimates of nonfatal injury and poisoning episodes and injury conditions from the first year of the redesigned NHIS. In addition, detail is provided on the design, methodologies, and rationale used in making these estimates and a discussion of the strengths and weaknesses of the new design.

Methods

Source of Data

The NHIS, conducted by the National Center for Health Statistics, collects demographic

and health data on a nationally representative sample of the civilian noninstitutionalized population residing in the United States. The NHIS has conducted face-to-face interviews continuously since its inception in 1957. In 1997, the NHIS was extensively redesigned, including the introduction of computer-assisted personal interviewing (CAPI).

The new Basic Module will remain largely unchanged from year to year and functions in a similar manner to the previous core questionnaire. The Basic Module contains three components: the Family Core, the Sample Adult Core, and the Sample Child Core (10).

The Family Core component collects information on all members of the family residing in the home at the time of the interview and will serve as a sampling frame for additional integrated national surveys. Information collected in the Family Core includes household composition and sociodemographic characteristics, tracking information, information for matches to administrative databases, and basic indicators of health status and utilization of health care services (10). The Injury Section, which contains questions pertaining to both injuries and poisonings, is in the Family Core; thus the questions on injuries and poisonings will be asked annually about every person in the family.

Capturing Episodes of Injury and Poisoning

Data are collected from an adult member (18 years or older) of the household about medically-attended injuries and medically-attended poisonings occurring in the previous 3-month period to any member of the family residing in the home at the time of the interview. If other adult members of the family are present during the interview, they can respond for themselves. The Injury Section of the Family Core of the NHIS survey questionnaire is included as [appendix I](#).

The focus of the redesigned injury section is a medically-attended injury episode or poisoning episode rather than an acute condition resulting from the

episode, as was the case prior to 1997 (1). An injury episode refers to the traumatic event in which the person was injured one or more times from an external cause (e.g., fall downstairs, motor vehicle traffic crash). An injury condition is the acute condition or the physical harm caused by the traumatic event (e.g., a fracture or a sprain). An injury episode can result in multiple conditions to the same person. If two people from the same household are injured seriously enough to require medical attention in the same episode (e.g., a car crash), the episode will be counted for each person injured.

Poisoning episodes include ingestion or contact with harmful substances and overdoses or wrong use of any drug or medication. The instructions to the respondent specifically exclude illness such as food poisoning and poison ivy. Food poisoning is an intestinal infectious disease, and poison ivy is classified as a dermatological condition (11). The *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD–9–CM) was relied upon for the definitions of injury and poisoning conditions and for the classification of external causes of the injury conditions used in the questionnaire and during the editing process (11).

A medically-attended injury or poisoning is one for which a health care professional was contacted either in person or by telephone for advice or treatment. Calls to poison control centers are considered contact with a health care professional and are included in this definition of medical attendance.

The questions that screen for the injuries and poisonings (appendix I) are:

“During the past three-months, were you or anyone in your family injured seriously enough that you got medical advice or treatment?” and

“During the past three-months, were you or anyone in your family poisoned seriously enough that you got medical advice or treatment?”

The two important changes in the screening questions in the 1997 redesign to highlight are that the recall period was increased from 2 weeks to 3 months and that the severity threshold was changed from events resulting in at

least half a day of restricted activity and/or medical attendance to only events resulting in medical advice or treatment.

The recall period was increased to capture more events on which to base national estimates. Increasing the recall period allows more episodes to be captured from the same number of people interviewed. With a longer recall period, the respondents may be more likely to forget events that happened further in the past (that is, memory decay), and the events more frequently forgotten are likely to be less severe events (12). Therefore, to counterbalance the increase in the recall period, the severity threshold was increased to include only episodes resulting in a visit or advice from a medical professional, as events of this severity may be better remembered.

During the editing process, injury episodes were limited to those including at least one health condition classified as an injury as defined by ICD–9–CM nature-of-injury codes 800–959 and 990–999 (11,13). Using CAPI to conduct the interview facilitated the collection of verbatim text information on how the injury happened, the body part injured, and the type of injury. This information, along with responses to questions about specific types of injury episodes, place of occurrence, and activity was used to assign ICD–9–CM diagnostic and external cause codes for all injury episodes.

For poisoning episodes in 1997, the cause of the poisoning was recorded in a precategorized list of causes rather than as a verbatim response. Therefore there are no external cause codes (E-codes) for the poisoning episodes. However, there was an option for the respondent to choose the open-ended category *“something else.”* If that response was chosen, the interviewer entered the respondent’s verbatim description of the cause. During the editing, the descriptions were reviewed and recategorized. Episodes involving illnesses such as food poisoning or poison ivy were sometimes reported and recorded, despite specific instruction to the interviewer not to include them. Therefore, episodes of this type were excluded during the editing process. In addition, the tables in this report

showing poisoning episodes do not include the 47 cases (unweighted) coded as *“Allergic/adverse reaction to medical or other substance”* or *“Something else—NOT poisoning.”* For additional information, refer to the NHIS Survey Description that is on the NHIS Web site (10).

While editing the 1997 injury data, it was discovered that respondents and interviewers sometimes had a difficult time differentiating between injuries and poisonings even with the available *“help screens.”* Poisonings or toxic effects within ICD–9–CM, (i.e., diagnosis codes 960–989 and external cause codes E850.0–E869.9, E950.0–E952.9, E962.0–E962.9, E980.0–E982.9, and E972) were sometimes reported following the screening question focusing on injuries, and thus the respondents answered the questions pertinent to injuries. When these injury episodes were assigned ICD–9–CM codes, the coders gave these episodes *“flags”* to indicate they were poisonings. In addition, the question on cause of poisoning contained a response category *“a venomous animal or plant.”* Coders also flagged injury episodes that appeared to fit into this cause of poisoning category. All episodes flagged as poisonings in the injury file were moved to the poisoning file. After editing was completed, it was discovered that seven episodes coded as E905.0–E905.9 (venomous animals and plants as the cause of poisoning and toxic reactions) were not given flags to indicate they were poisonings and, therefore, were not moved from the injury file to the poisoning file.

Respondents were not asked about the health conditions resulting from poisoning episodes and, therefore, it is not possible to count the number of poisoning conditions. However, for this report, it was assumed that there was only a single condition per poisoning episode to estimate the total number of injury and poisoning conditions. This assumption may result in an underestimate of the number of poisoning conditions. Beginning with data year 2000, there will be one set of questions for injury and poisoning episodes. This will allow ICD external cause codes and diagnostic codes to be

assigned for both injury and poisoning episodes. From that point, the inclusion criteria will rely entirely on the ICD classification codes.

Analysis Fields

The injury-specific questions on the NHIS elicit information on the external cause of the injury episode, the injury diagnoses, what the person was doing at the time of the injury episode, place of occurrence, and whether the injury episode caused any limitation of activity. Edited verbatim response to the question on how the injury occurred (up to 336 characters) and the four responses to the question on the body part injured (up to 34 characters each) and four responses to the question on the kind of injury (up to 44 characters each) are available in the verbatim text file. The descriptions were edited to protect the injured person's confidentiality. Grammatical and/or spelling errors were not corrected. Poisoning-specific questions include the external cause of the poisoning and whether a poison control center was contacted.

Questions asked about both injuries and poisonings include the date of the episode, whether the person was hospitalized, and whether the person missed any days from work or school. The month and year of the date of episode are available for analysis. During the editing process, an elapsed time (i.e., recall period) variable was derived from the date of the episode and the date of the interview.

The NHIS collects detailed information on the four most recent injury episodes and the four most recent poisoning episodes. Up to four ICD-9-CM injury diagnosis codes and three E-codes were assigned for each injury episode and are available for analysis in the data files. Tables and figures include the first-listed external cause and include all conditions reported (11).

The external cause code categories in [table 2](#) are based on the mechanism axis of the recommended framework for presenting injury data with the exception of the *transportation* category (13) ([appendix II](#)). The *transportation* category includes the external cause codes for most modes of transportation

(i.e., “Motor vehicle traffic;” “Pedal cyclist, other;” “Pedestrian, other;” and “Transport, other).” These codes were selected because they correspond with several additional questions asked only for transportation episodes. The external causes are categorized by mechanism of injury regardless of intent.

There are no questions in the injury section that specifically address the question of intentionality because when the questionnaire was tested in the NCHS Cognitive Questionnaire Lab during the early phases of its design, it was determined that the setting of the interview was not conducive to identifying assaults and intentionally self-inflicted injuries.

The ICD-9-CM diagnosis codes used to categorize the injury conditions in [tables 3](#) and [4](#) are based on a matrix of nature of injury by body region injured developed by Mackenzie and Champion (14). This matrix cross classifies the ICD-9-CM diagnosis codes 800-994 by nature of injury (e.g., fracture, sprain/strain) and by body region injured (e.g., upper extremity, abdomen). The diagnosis codes used are shown in [appendix III](#).

Selection of the age groups used in this report—under 12 years, 12-21 years, 22-44 years, 45-64 years, 65-79 years, and 80 years and over—were based on a graphical analysis of the 1997 injury episode rate by single year of age. The groups were determined by compromise between the observed pattern of the data and standard age groups often used by NCHS ([figure 1](#)). The age groups will be reconsidered when more data are available.

Data by race and ethnicity are shown for non-Hispanic white, non-Hispanic black and for Hispanic persons, but not for persons of other races, because sample sizes were not large enough to make reliable estimates.

Sample Size and Response Rates

The interviewed sample in 1997 consisted of 39,832 households, which yielded 103,477 persons in 40,623 families. The total noninterview rate was approximately 8.2%: 5.0% due to

respondent refusal and the remainder primarily due to failure to locate an eligible respondent at home after repeated calls or unacceptable partial interviews (10).

Injury and poisoning information was reported for 2,949 people having 3,114 episodes of injury or poisoning. These episodes resulted in 3,712 conditions including 3,533 injury conditions and 179 poisoning conditions (assuming one poisoning condition per poisoning episode). Ninety-five percent of the people had only one episode reported during the 3-month period.

Data Files

Data on injuries are in three public use data files—the Person file, Injury Episode file, and the Injury Verbatim file. Data on poisonings are found in two files—the Person file and the Poison Episode file. Each of these files can be linked so that information about the injury or the poisoning can be combined with any of the other variables from the NHIS.

Units of Analysis, Annual Estimates, and Rates

Weighted data are used to estimate the number of injury and poisoning episodes, the injury and poisoning conditions, and the size of the population. The Final Annual Weight available on the data tape is based on information from the sample design and sampling ratio and is adjusted for nonresponse and post-stratified based on census totals for sex, age, and race/ethnicity.

The injury and poisonings episodes are collected with a 3-month recall. To annualize the estimate, each 3-month estimate should be multiplied by 4. Estimates shown in the figures and tables are based on the number of episodes and conditions occurring in 1 year. Rates are calculated either as the annual number of injury and poisoning episodes per 1,000 population or as the annual number of injury conditions per 1,000 population.

It is not possible to estimate the number of people injured or poisoned




Figure 1. Injury and poisoning episode rates by age: United States, 1997

annually using the NHIS. Although the number of persons who were injured during the 3-month recall period is known, this number cannot be assumed to be uniform over a 12-month period. For example, if it is known that 100 people responded that they were injured during the 3-month recall period, one cannot assume that 400 different people were injured in a 12-month period because some people may be injured multiple times and some may be injured once. On the other hand, it is appropriate to estimate the number of injuries over the 12-month period (by multiplying the 3-month estimate by 4) because that figure is the same whether or not individuals had multiple injuries.

Variance Estimation and Significance Testing

Variance estimates were produced to indicate the reliability of point estimates. The NHIS data are obtained through a complex sample design involving stratification, clustering, and multistage sampling. The variance estimates were calculated using the SUDAAN software package and incorporated the design

information available on the public use data tapes. The Taylor series linearization method was chosen for variance estimation in SUDAAN. Taylor series linearization is used to approximate the functions of linear statistics. SUDAAN then computes the design-specific variance for the linearized values (15).

Standard errors are shown for all rates and percents in tables. Rates or percents with relative standard errors greater than 30% are considered unreliable and are indicated with an asterisk. The relative standard errors are calculated as follows:

Relative standard error = $(SE/EST)100$
 where SE is the standard error of the estimate and EST is the estimate.

Data points in the figures are shown with 95% confidence intervals (i.e., error bars) that were calculated as follows:

$PE \pm 1.96(SE)$
 where PE is the point estimate, SE is the standard error of the estimate, and 1.96 is the reliability coefficient corresponding to a confidence level of 95%.

While statistical tests of differences among the point estimates were not used

in this report, confidence intervals are given to graphically display the stability of the individual point estimates. The objective of the analysis is to identify patterns or trends in the data by investigating variation in specific injury rates or proportions by age, race, ethnicity, and sex as displayed in the figures. Variability in the rates is summarized by the standard errors and confidence intervals provided. Key results are highlighted in the text when large differences in rates were observed. See [appendix IV](#) for more specific information on issues of statistical testing.

Age-Adjusted Rates

Data were age-adjusted by the direct method to the 2000 projected population. The age groups used to adjust are the same ones used in the age-specific analyses in this report: under 12 years, 12–21 years, 22–44 years, 45–64 years, 65–79 years, and 80 years and over. Age adjustment is used to adjust for differences in the age distribution of populations being compared. Age adjustment is often done

when trends are being analyzed or when populations being compared are known to have different age distributions. The NCHS standard for age adjustment was recently changed from the 1940 standard to a 2000 standard (16). Age-adjusted and crude rates of injury and poisoning in 1997 are similar.

Results

In 1997, there were 34.4 million episodes of injury and poisoning reported among the U.S. civilian noninstitutionalized population at a rate of 129 episodes of injury and poisoning per 1,000 persons (similar for crude and adjusted rates) (table 1). The injury and poisoning episodes resulted in 40.9 million conditions, at a rate of 154 conditions per 1,000 persons (similar for crude and adjusted rates).

Age, Sex, and Race/Ethnicity

The age-adjusted injury and poisoning episode rate for males was 21% higher than the rate for females (140.5 and 115.7 per 1,000 persons, respectively). In general, age-specific episode rates for males were higher than for females among persons younger than age 45 years, but were lower at the older ages (figure 2). The age-adjusted injury and poisoning episode rate was higher for non-Hispanic white persons than for either non-Hispanic black or Hispanic persons (142.8 and 109.1 and 89.3 per 1,000 persons, respectively) (table 1). Rates for persons under 22 years of age were generally higher for non-Hispanic white persons than for others. For persons 22–64 years, it appears that Hispanic persons have lower episode rates than other persons (figure 3).

Leading External Causes of Injury

In 1997, *falls* were the leading external cause of injury; 11.3 million episodes of *falls* were reported at an age-adjusted rate of 43.1 per 1,000 persons (table 2). *Falls* were followed




Figure 2. Injury and poisoning episode rates by age and sex: United States, 1997

by episodes resulting from *being struck by or against a person or an object, transportation, overexertion, cutting and piercing instruments, and poisoning*. The external cause category *transportation* includes most modes of transportation (i.e., motor vehicles, pedal cyclist, pedestrian, and other transportation).

Falls is the only category of external causes for which the

age-adjusted rate for females exceeded that for males (47.2 versus 37.6 per 1,000 persons, respectively) (figure 4). The higher episode rate for *falls* among females results from the higher rates for persons older than 45 years and especially for those 65 years and older where the rate for females was more than twice the rate for males (figure 5).




Figure 3. Injury and poisoning episode rates by age, race, and ethnicity: United States, 1997




Figure 4. Age-adjusted external cause of injury and poisoning episode rates by mechanism of injury and sex: United States, 1997

The fall was on level ground in 35–40% of the episodes. In 25% of the falls, the type of fall did not fit into one of the predefined categories of falls (table 3).

Sex differences in the rate for being struck by or against an object or person are largest at ages 12–21 years; the rate for males at that age was 3.5 times the rate for females (figure 6). The rates for

males 22–44 years old were considerably lower than for males 12–21 years old, but were still more than twice the rates for females at ages 22–44 years.

The transportation-related injury episode rates were higher for persons 12–21 and 22–44 years old than for younger or older persons (figure 7). Although transportation-related episodes

include most modes of travel, 75% of these episodes could be classified as motor vehicle traffic-related (E810–E819). About 58% of transportation-related episodes involved drivers of vehicles, 27% involved passengers, 10% involved bicycles, and 3% involved pedestrians (table 4). Of those injured as drivers or passengers, nearly 70% were in passenger cars, 13% were in light trucks, and 6% were on motorcycles. Over three quarters (78%) of those injured as drivers or passengers in passenger cars, light trucks, or large trucks, were reported to be restrained at the time of the incident by wearing a seatbelt or by being buckled into a safety seat. Only about 40% of those injured while riding a bicycle, motorcycle, an all-terrain vehicle, or a snowmobile were reported to be wearing a helmet (table 4).

The rates of reported overexertion episodes were higher among persons ages 12–64 years than among the younger or older age groups (figure 8). Although the rates were much lower, patterns for injuries associated with cutting and piercing instruments were similar to those for being struck by something (figures 6 and 9).

Poisoning episode rates were higher for children under 12 years than for others with no other readily observable patterns by age or by sex (figure 10). For about 45% of all reported poisoning episodes, a call was made to a poison control center—about 877,000 calls in 1997 (table 5). This number is less than half the number of reported calls made as reported by the American Association of Poison Control Centers (17). However, there are differences in the reporting methods between the two surveillance systems. NHIS estimates are based on reporting by the patient or other family member with a 3-month recall and the other is based on direct reporting by the poison control centers. Nonetheless, both data systems show higher proportions of calls involving a young child compared with older persons. Based on NHIS data, about 3 out of 4 poisonings involving a child less than 6 years old results in a call to a poison control center regarding the incident.




Figure 5. Injury episode rates for falls by age and sex: United States, 1997




Figure 6. Injury episode rates for being struck by or against an object or person by age and sex: United States, 1997

Injury Conditions

In 1997, *sprains and strains* were the injury condition most frequently reported followed by *open wounds* and *fractures* and *contusions*. The age-adjusted rates were 38.5, 29.0, 23.7, and 19.0 per 1,000 persons, respectively (table 6). For each of these conditions,

with the exception of *open wounds*, there were no rate differences by sex (figure 11). The age-adjusted *open wound* rate for males was about twice the rates for females.

The condition rate for *sprains and strains* were lower for children under 12 years than for others. Rates generally declined with age, starting at ages 12




Figure 7. Injury episode rates for transportation-related injuries by age and sex: United States, 1997

and older (figure 12). Among elderly females, the rate was about twice that for elderly males.

Sex differences in *open wound* rates at ages 12–21 years were far higher than for almost any other condition (table 6). The rate for males was close to 6 times that for females. At ages under 12 years and 22–44 years, the rates for males were about twice those for females (figure 13).

The age distribution for *fracture* rates was bimodal with peaks at 12–21 years and at 65 years and over (figure 14). Finally, rates for *contusions* were higher for persons 65 years and over and for females, in particular, than for younger persons (figure 15).

In addition to the nature of the injury, the body region injured can also describe the condition. *Upper extremity* and *lower extremity* injuries were the most often reported sites of the conditions (with age-adjusted rates of 43.2 and 40.3 per 1,000 persons, respectively) followed by injuries to the *spine and back*, *face*, and *skull and brain* (table 7). Both *upper* and *lower extremity* injury condition rates were higher than those for other regions of the body (figure 16).

Upper extremity injury rates were higher for males 12–44 years old than for males of other ages or for females of any age (figure 17). The rate for *lower extremity* injuries for males 12–21 years old was 1.5 times higher than for females of the same age, but at ages 45 years and older, *lower extremity* injuries were more likely among females than males (figure 18). For females 65 and over, the rate was 3 times that for their male counterparts. The rate of *skull and brain* injuries among males under 12 years old was 3 times that for females, and for males 22–44 years, the rate was closer to 2 times that for females (table 7).

Place of Injury

The *home* was the most frequently reported place of injury with 24% of injuries occurring *inside the home*, and another 18% *outside the home* (table 8). Females were located *inside the home* at the time of the injuries almost twice as often as males (32% vs. 18%). Injuries




Figure 8. Injury episode rates for overexertion by age and sex: United States, 1997

occurring while on the *street and highway* accounted for a further 13% of the episodes with no difference between males and females (figure 19). Injuries occurring at *sport facilities, industrial and construction areas, schools, and trade/service areas* each accounted for 6–7% of the injuries. Males were reported as injured at *sports facilities*

more often than females (10% vs. 4%). Male-female differences were greatest at *industrial/construction sites*, with 11% of male injury episodes occurring at these sites compared with 2% of their female counterparts. Eight percent of male injuries, compared with 5% of female injuries, occurred at *schools*, a location where equal exposure is expected.




Figure 9. Injury episode rates for cutting and piercing instruments by age and sex: United States, 1997

Activity at Time of Injury

Leisure activities and paid work were most often reported as the activities the person was engaged in when the injury episode occurred, accounting for 22% and 19% respectively of all episodes of injury (table 8). The percent of persons engaged in *leisure activities* were similar for males and females (figure 20). Males were engaged in *paid work* at the time of injury almost twice as often as females (24% vs. 14%). Injuries resulting during *sport-related activities* were reported for 14% of the episodes, more than twice as often by males than by females (18% vs 8%). *Working around the house or the yard* was reported for about 10% of the injury episodes, somewhat more often for females than for males. For 8% of the episodes, *driving* was the reported activity and for 3%, *attending school* was reported.

Hospitalization

Respondents reported that the person was hospitalized in 2.5 million injury and poisoning episodes, or about 7% of the episodes (table 9). For those hospitalized, the number of days in a hospital ranged from 1–70 days. However, 62% of those hospitalized had a 1–3 day stay and 90% were in the hospital for 2 weeks or less. The percent of persons hospitalized increased with age with 4% of those under 22 years hospitalized after the episode compared with 15% of those ages 65 and over.

School- and Work-Loss Days

There were 3.3 million injury and poisoning episodes resulting in time lost from school. About 85% of the episodes with school loss occurred among 5–24 year olds at a rate of 36 episodes per 1,000 persons (table 10). For those ages 5–24 years, about 22% of the episodes with time lost from school resulted in less than 1 day lost, 65% resulted in 1–5 days lost, and 13% resulted in 6 days or more lost.

There were 10.1 million injury and poisoning episodes resulting in time lost from work. About 99% of the episodes with work loss occurred among 14–75 year olds at a rate of 51 episodes per




Figure 10. Poisoning episode rates by age and sex: United States, 1997

1,000 persons. For those ages 14–75 years, about 17% of the episodes with time lost from work resulted in less than 1 day lost, 46% in 1–5 days lost, and 37% in 6 days or more lost.

Limitations After injury

After the injury, about 7% of the injured people needed help in handling

daily routines such as household chores and shopping, and about 6% of injured people needed help attending to *personal care* such as eating and bathing (table 11). The percent of persons with limitations in both *personal care* and *daily routine* increased with age. At ages 65 years and over, 21% of the injured persons needed help with *daily routine* compared with 2% at ages 5–21 years.




Figure 11. Age-adjusted injury condition rates by nature of injury and sex: United States, 1997

At these same ages, 19% needed help with *personal care* compared with 4% at the younger ages. A higher percent of females were limited after injuries than males were; 10% of females were limited in their *daily routine* versus 5% of males.

Discussion

This report provides national estimates of injury and poisoning episodes and conditions for the first year of the redesigned National Health Interview Survey. The first year of any ongoing survey is a time of learning for everyone, from the interviewers and data processors to the analysts. Analyses are for a single year, so it is possible that some of the findings presented will change in subsequent years as more data are accumulated. The following discussion presents the strengths and limitations of the redesign and some methodological issues for consideration.

Strengths of the Redesign

With only 1 year of data, reliable national estimates can be made for many age, sex, and cause-specific rates of injury and poisoning episodes and conditions. This is the result of increasing the recall period to 3 months because it increases the likelihood that a person will have had an injury or poisoning. Therefore, although the same numbers of people are interviewed, there are more episodes on which to make the national estimates.

An acknowledged limitation of the NHIS prior to 1997 was the limited information on the circumstances of the injury (18). In the redesigned survey, there are multiple questions on the circumstances of the injury episode, including cause, place, activity, and outcome of the injury. In addition, for certain cause-specific injuries, there are additional questions that address issues related to that cause. Work-related and sports-related injury episode rates can be estimated on the national level from the activity and place fields. Detail on transportation-related episodes is




Figure 12. Injury condition rates for sprains and strains by age and sex: United States, 1997

available and will make the data compatible with the Tenth Revision of the ICD (19). For example, for persons injured in transportation-related incidents, questions on the type of vehicle, if the person was the driver or passenger and on the type of vehicle if the person was a pedestrian struck by it.

A narrative description of how the injury occurred, the part of the body injured, and the nature of the injury are available for analysis. These narratives are the verbatim responses to the interviewer’s queries with minimal editing for confidentiality. The injury data were ICD coded for NCHS




Figure 13. Injury condition rates for open wounds by age and sex: United States, 1997

purposes, but researchers can code them according to other cause or diagnostic classification schemes. In addition, it is possible to search for text strings of words of interest (20, 21, 22). The analyst should be cautioned, however, against making estimates based on small numbers of observations.

Limitations and Lessons Learned

In 1997, there are separate sets of questions for injuries and poisonings, although some questions are common to both categories. In general, there is more detail for injuries than for poisonings. For example, the narrative field is available for injury episodes only. Not having all the questions asked of all respondents limits the completeness of the analysis. In addition, respondents did not always understand the subtle distinctions that separate an injury from a poisoning, so occasionally injuries were reported as poisonings and poisonings as injuries. These problems were addressed in the data editing process, but editing was limited by missing responses when injuries and poisonings were inappropriately reported. The confusion between an injury and a poisoning is understandable because there is often disagreement among injury researchers about appropriate categorization. For example, are venomous bites or stings poisonings? The ICD-9-CM does not include them in the section of codes for “accidental poisonings” although the text descriptions of the appropriate ICD-9-CM code includes the words “as the cause of poisoning or toxic reaction.” Because of these difficulties, the injury and poisoning questions were modified for the year 2000 survey so there is one set of questions for all persons injured or poisoned.

In 1997, respondents were asked if anyone in the family had any injuries. However, there was variability in what respondents considered to be an injury. For example, while repetitive motion injuries or carpal tunnel syndrome were reported in response to the screening question, these are not injuries according to ICD-9-CM and were not included in




Figure 14. Injury condition rates for fractures by age and sex: United States, 1997

this data set. It is important to remember that the NHIS is not based on physician diagnoses, but on descriptions of injuries provided by the respondents. Thus, estimates should be made only on broad injury categories rather than specific diagnoses.

Respondents were prompted to choose an activity to best fit what they were doing at the time of injury. A

review of the verbatim responses, however, indicated that some of the “other” responses fit into one of the predefined categories such as “driving” and “leisure” activities. Some of the categories in the activity were also difficult for the respondent to interpret. For example, if the injured person was doing housework at the time of the injury, two categories were suitable:

“Working around the house or yard” or “Unpaid work (including housework, shopping, volunteer work).” As a result, new guidelines were given to interviewers to select the first category if the response involved “housework” and the response categories are being evaluated for future years.

Comparability With Prior Years

Estimates of all injury conditions for 1997 are not directly comparable to estimates from prior years of the NHIS. As discussed in the Methods section of this report, differences between 1997 and previous years include changes to the screening questions; the inclusion criteria; the recall period; the severity threshold; and the phrasing, placement, and number of questions. The following discussion is a brief introduction to comparisons between the redesign and previous years. A more detailed analysis will be provided in the near future.

An evaluation of changes in the 1995 and 1997 injury and poisoning condition rates requires that data from the two surveys be reanalyzed. To make the 1995 data more comparable to the 1997 data, injuries that were not medically attended and adverse effects and complications of medical care (ICD-9-CM 995.0-995.4, 995.6, 995.86, 995.89-999 and E870-E878, and E930-E949) were excluded from the 1995 estimates. Comparisons were made with 1995 because only a partial year of data is available for 1996.

In the redesigned 1997 survey, where respondents were asked directly about injury conditions, there were only four reported conditions (unweighted) that were coded to “adverse effects” or “complications of surgical and medical care” (ICD-9-CM 995.0-995.4, 995.6, 995.86, and 995.89-999), accounting for 0.1% of all injury conditions. In contrast, in 1995, 13% of all acute injury conditions (about 7.5 million conditions) in the NHIS were adverse effects or complications. This finding mirrors the many discussions among injury researchers and prevention specialists, including those participating in the International Collaborative Effort




Figure 15. Injury condition rates for contusions by age and sex: United States, 1997




Figure 16. Age-adjusted injury condition rates by body region and sex: United States, 1997

on Injury Statistics (1996) and in the injury data committee of the Injury Section of the American Public Health Association (1996), in which it was agreed that these conditions and their comparable external causes (E870–E878 and E930–E949) were not injuries that could be prevented in the same manner as injuries resulting, for example, from

motor vehicle crashes or falls (13, 23). Thus, they have been excluded from some recent counts of injuries and injury deaths published by the National Center for Health Statistics (24, 25).

When limited to the medically attended injuries and excluding the adverse effects and complications of surgical and medical care, the 1995 rate

of 19.0 injury and poisoning conditions per 100 persons is 25% greater than the 1997 rate of 15.4 injury and poisoning conditions per 100 persons. However, the 1995 estimate does not address the increased recall period. Analysis of recall bias was complicated by missing values in the date of injury field, but will be addressed in a forthcoming report.

Preliminary analyses comparing specific injury conditions between 1995 and 1997 show similar numbers of fractures and sprains and strains, but differences in the numbers of “superficial injuries” as classified by the ICD (appendix III). This suggests that the recall of less severe injuries may be playing a role in the decrease in estimated number of injuries. However, the estimates were made more stable with the 3-month recall because of the greater number of observations on which the estimates are based.

Conclusion

The redesigned NHIS is a useful source of information about medically-attended nonfatal injuries and poisonings occurring in the United States. A single year of data provides enough data to produce stable national estimates on details of injury and poisoning episodes. With each successive year of data, the amount of information about the causes and diagnoses of nonfatal injuries will increase and thus so should the ability to expand and enhance prevention efforts.

For complete documentation and data tapes, see: <http://www.cdc.gov/nchs/nhis.htm>




Figure 17. Upper extremity injury condition rates by age and sex: United States, 1997




Figure 18. Lower extremity injury condition rates by age and sex: United States, 1997




Figure 19. Percent of injury episodes by place of injury and sex: United States, 1997




Figure 20. Percent of injury episodes by activity at the time of injury and sex: United States, 1997

References

- Benson V, Marano MA. Current estimates from the National Health Interview Survey, 1995. National Center for Health Statistics. *Vital Health Stat* 10(199). 1998.
- Burt CW, Fingerhut LA. Injury visits to hospital emergency departments: United States, 1992–95. National Center for Health Statistics. *Vital Health Stat* 13(131). 1998.
- Burt CW. Injury-related visits to hospital emergency departments: United States, 1992. Advance data from vital and health statistics; no 261. Hyattsville, Maryland: National Center for Health Statistics. 1995.
- Woodwell, DA. National Ambulatory Medical Care Survey: 1997 Summary. Advance Data: no 305. Hyattsville, Maryland: National Center for Health Statistics. 1999.
- Hall MJ, Owings MF. Hospitalizations for injury and poisonings in the United States, 1991. Advance data from vital and health statistics; no 252. Hyattsville, Maryland: National Center for Health Statistics. 1994.
- Lawrence, L, Hall, MJ. 1997 Summary: National Hospital Discharge Survey. Advance data from vital and health statistics; no 308. Hyattsville, Maryland: National Center for Health Statistics. 1999.
- US Consumer Product Safety Commission, Consumer Product-Related Statistics Page, <http://www.cpsc.gov/library/data.html>.
- US Consumer Product Safety Commission, National Electronic Injury Surveillance System Coding Manual, Washington, DC, 1997.
- Quinlan KP, Thompson MP, Annett JL, et al. Expanding the National Electronic Injury Surveillance System to Monitor All Nonfatal Injuries Treated in US Hospital Emergency Departments. *Annals of Emergency Medicine*. 1999; 34: 637–645.
- 1997 National Health Interview Survey (NHIS) Public Use Data Release, NHIS Survey Description, Division of Health Interview Statistics, National Center for Health Statistics, Hyattsville, MD, Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, February 2000 (<http://www.cdc.gov/nchs/nhis.htm#1997NHIS>.)
- Public Health Service and Health Care Financing Administration. International Classification of Diseases, 9th Revision, Clinical Modification. Washington: Public Health Service. 1991.
- Harel Y, Overpeck MD, Jones DH, et al. The effects of recall on estimating annual nonfatal injury rates for children and adolescents. *American Journal of Public Health*. 1994; 84: 599–605.
- McLoughlin E, Annett JL, Fingerhut L, et al. Recommended framework for presenting injury mortality data. *MMWR*. Centers for Disease Control and Prevention. 1997; 46 (no. RR-14):1–32.
- Mackenzie EJ, Champion H. Injury diagnosis morbidity matrix. In: Proceedings of the International Collaborative Effort on Injury Statistics, Vol 3. Hyattsville, MD: National Center for Health Statistics. 2000.
- Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual, Release 7.0, 1st edition, Research Triangle Institute, Research Triangle Park, NC. 1996.
- Anderson RN, Rosenberg HM. Age Standardization of Death Rates: Implementation of the Year 2000 Standard. National Center for Health Statistics. National Vital Statistics Reports. 47(3). 1998.
- Litovitz TL, Klein-Schwartz W, Dyer KS, et al. 1997 Annual Report of the American Association of Poison Control Centers Toxic Exposure Surveillance System. *American Journal of Emergency Medicine*, Vol 16 (5). 1998: 443–497.
- Baker SP, O'Neil B, Ginsburg MJ, Li G. The Injury Factbook. 2nd Edition. New York: Oxford University Press. 1992.
- World Health Organization. International Statistical Classification of Diseases and Related Health Problems, 10th Revision. Geneva: World Health Organization. 1992.
- Jenkins EL, Hard DL. Implications for the use of E codes of the International Classification of Diseases and narrative data in identifying tractor-related deaths in agriculture, United States, 1980–1986. *Scandinavian Journal of Work, Environment, and Health* 1992; 18 Supp. 2: 49–50.
- Langley JD. Experiences using New Zealand's hospital based surveillance system for injury prevention research. *Methods of Information in Medicine* 1994; 34: 340–344.
- Sorock GS, Smith GS, Reeve GR, et al. Three perspectives on work-related injury surveillance systems. *American Journal of Industrial Medicine* 1997; 32: 116–128.
- Proceedings of the International Collaborative Effort on Injury Statistics Volume 1. National Center for Health Statistics, Hyattsville, MD. DHHS publication number (PHS) 95–1252. 1995.
- Fingerhut LA, Warner M. Injury Chartbook. Health, United States, 1996–97. Hyattsville, MD: National Center for Health Statistics. 1997.
- Hoyert DL, Kochanek KD, Murphy SL. Deaths: Final Data for 1997. National Vital Statistics Reports; Vol 47 no 19. Hyattsville, Maryland: National Center for Health Statistics. 1999.

Table 1. Number, percent distribution, standard error, and annual rate of injury and poisoning episodes by selected demographic characteristics: United States, 1997

Selected demographic characteristic	Number of episodes in thousands	Percent distribution	Standard error (percent)	Rate per 1,000 population	Standard error (rate)
Sex and age					
Both sexes:					
All	34,383	100.0	...	129.0	2.6
All, age-adjusted	128.9	2.6
Under 12 years	5,384	15.7	0.7	112.2	5.9
12–21 years	6,542	19.0	0.9	171.4	8.4
22–44 years	12,766	37.1	1.1	136.4	4.5
45–64 years	5,470	15.9	0.8	99.7	5.1
65 years and over	4,221	12.3	0.7	131.9	7.0
65–79 years	2,850	8.3	0.6	114.9	7.7
80 years and over	1,371	4.0	0.4	190.5	19.4
Male:					
All	18,544	53.9	1.1	142.4	4.0
All, age-adjusted	140.5	4.1
Under 12 years	3,063	8.9	0.6	125.1	8.2
12–21 years	4,394	12.8	0.7	225.3	13.3
22–44 years	7,406	21.5	0.9	160.3	7.1
45–64 years	2,483	7.2	0.6	93.5	7.3
65 years and over	1,197	3.5	0.4	88.9	9.8
65–79 years	800	2.3	0.3	73.3	9.4
80 years and over	397	1.2	0.3	155.3	34.0
Female:					
All	15,840	46.1	1.1	116.1	3.4
All, age-adjusted	115.7	3.4
Under 12 years	2,320	6.7	0.5	98.7	7.7
12–21 years	2,149	6.2	0.6	115.1	11.1
22–44 years	5,360	15.6	0.8	113.1	5.8
45–64 years	2,988	8.7	0.5	105.5	6.9
65 years and over	3,023	8.8	0.6	163.1	10.4
65–79 years	2,050	6.0	0.5	147.5	11.7
80 years and over	973	2.8	0.3	209.9	24.3
Race/ethnicity and age					
White non-Hispanic:					
All	27,299	79.4	0.8	141.8	3.3
All, age-adjusted	142.8	3.3
Under 12 years	4,096	11.9	0.7	133.4	8.2
12–21 years	5,329	15.5	0.8	208.6	11.4
22–44 years	9,747	28.3	1.0	147.3	5.7
45–64 years	4,381	12.7	0.7	101.9	5.9
65 years and over	3,745	10.9	0.6	138.2	8.0
65–79 years	2,489	7.2	0.5	119.6	9.0
80 years and over	1,256	3.7	0.4	199.8	21.5
Black non-Hispanic:					
All	3,513	10.2	0.6	107.9	6.6
All, age-adjusted	109.1	6.7
Under 12 years	579	1.7	0.2	78.8	10.0
12–21 years	613	1.8	0.2	109.7	15.0
22–44 years	1,482	4.3	0.4	128.5	11.1
45–64 years	608	1.8	0.3	110.5	15.4
65 years and over	231	0.7	0.1	89.8	19.7
65–79 years	141	0.4	0.1	69.3	18.3
80 years and over	*90	*0.3	0.1	*168.1	54.2

See footnotes at end of table.

Table 1. Number, percent distribution, standard error, and annual rate of injury and poisoning episodes by selected demographic characteristics: United States, 1997—Con.

Selected demographic characteristic	Number of episodes in thousands	Percent distribution	Standard error (percent)	Rate per 1,000 population	Standard error (rate)
Hispanic:					
All	2,658	7.7	0.5	88.8	5.2
All, age-adjusted	89.3	5.3
Age group:					
Under 12 years	629	1.8	0.2	82.8	10.4
12–21 years	483	1.4	0.2	93.2	12.0
22–44 years	1,063	3.1	0.3	93.7	8.6
45–64 years	276	0.8	0.1	66.4	9.1
65 years and over	207	0.6	0.1	126.1	27.3
65–79 years	182	0.5	0.1	134.2	30.5
80 years and over	*25	0.1	0.0	*87.8	52.1
Geographic region					
Northeast	6,732	19.6	0.9	128.4	5.4
Midwest	8,723	25.4	0.9	132.7	5.5
South	12,214	35.5	1.0	128.3	4.2
West	6,715	19.5	0.9	126.0	6.2

. . . Category not applicable.

* Figure does not meet standard of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

Table 2. Annual rate and standard error of leading external causes of injury and poisoning episodes by sex and age: United States, 1997

Sex and age	Fall		Struck by or against a person or an object		Transportation ¹		Overexertion		Cutting-piercing instruments		Poisoning	
	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error
Both sexes												
All.	42.4	1.5	19.6	1.0	16.7	1.0	13.9	0.9	10.0	0.7	7.3	0.6
All, age adjusted.	43.1	1.5	19.4	1.0	16.5	1.0	13.9	0.8	9.9	0.7	7.2	0.6
Under 12 years	45.7	3.7	17.3	2.2	8.6	1.4	*2.4	1.0	8.0	1.3	13.0	1.8
12–21 years	45.5	4.6	49.1	4.4	23.8	3.0	12.3	2.1	15.4	2.2	5.6	1.4
22–44 years	30.6	2.0	18.9	1.7	22.9	1.9	20.3	1.8	13.4	1.4	6.7	1.0
45–64 years	32.1	2.9	11.5	1.5	11.2	1.6	17.6	1.9	5.5	1.1	5.5	1.3
65 years and over.	86.2	5.8	4.0	1.2	11.5	2.2	7.8	1.9	4.4	1.3	5.4	1.6
Male												
All.	36.9	2.1	27.6	1.7	19.3	1.4	15.1	1.3	13.3	1.2	6.9	0.9
All, age adjusted.	37.6	2.1	26.6	1.6	18.9	1.4	14.9	1.2	12.9	1.1	6.7	0.9
Under 12 years	47.7	5.7	21.9	3.5	13.8	2.5	*3.0	1.7	10.7	2.2	11.3	2.1
12–21 years	49.1	6.1	75.3	7.5	28.2	4.5	17.1	3.5	25.7	4.3	*4.8	1.7
22–44 years	29.9	3.1	27.0	2.9	26.1	2.7	22.5	2.6	15.7	2.1	7.6	1.6
45–64 years	25.0	3.4	11.3	2.2	9.8	2.1	17.5	2.7	6.6	1.7	*5.2	1.7
65 years and over.	47.2	7.6	*3.4	1.5	12.3	3.4	*4.4	1.9	*4.8	2.0	*2.9	1.8
Female												
All.	47.7	2.4	11.9	1.0	14.1	1.2	12.7	1.1	6.9	0.8	7.7	0.8
All, age adjusted.	47.2	2.4	12.1	1.0	14.1	1.2	12.7	1.1	6.8	0.8	7.7	0.8
Under 12 years	43.7	5.0	12.5	2.4	*3.2	1.3	*1.7	1.0	*5.1	1.6	14.7	2.9
12–21 years	41.7	7.3	21.7	4.4	19.2	3.6	*7.4	2.3	*4.6	1.7	*6.4	2.3
22–44 years	31.3	3.0	11.0	1.7	19.7	2.3	18.1	2.0	11.2	1.8	5.9	1.2
45–64 years	38.7	4.5	11.6	2.2	12.6	2.2	17.8	2.9	*4.5	1.4	5.8	1.6
65 years and over.	114.5	8.8	*4.4	1.6	11.0	2.9	10.2	2.9	*4.2	1.6	*7.2	2.4

* Figure does not meet standard of reliability or precision.

¹Transportation includes the categories "Motor vehicle traffic;" "Pedal cycle, other;" "Pedestrian, other;" and "Transport, other" (appendix II).

NOTE: External cause of injury was derived from the verbatim responses to questions FIJ.050 to FIJ.220 of the questionnaire (appendix I). See appendix II for a list of the ICD–9–CM codes in each cause category.

Table 3. Number, percent, and standard error of fall-related episodes by type of fall and sex: United States, 1997

	Both sexes			Male			Female		
	Number in thousands	Percent of all fall episodes	Standard error	Number in thousands	Percent of all fall episodes	Standard error	Number in thousands	Percent of all fall episodes	Standard error
All fall episodes	11,306	100.0	...	5,015	100.0	...	6,291	100.0	...
Type of fall									
All types of falls mentioned ¹	12,285	108.7	...	5,493	109.5	...	6,792	108.0	...
Floor/level ground	4,158	36.8	1.7	1,729	34.5	2.7	2,430	38.6	2.2
Stairs/step.	1,296	11.5	1.0	428	8.5	1.4	869	13.8	1.5
Curb/sidewalk	1,162	10.3	1.0	415	8.3	1.4	747	11.9	1.4
Furniture.	807	7.1	0.9	294	5.9	1.3	513	8.2	1.2
Playground equipment	493	4.4	0.7	246	4.9	1.1	247	3.9	0.8
Ladder/scaffolding.	447	4.0	0.7	327	6.5	1.3	*121	*1.9	0.6
Hole/opening	382	3.4	0.6	172	3.4	0.9	210	3.3	0.9
Other (specified) ²	610	5.4	0.8	386	7.7	1.5	224	3.6	0.8
Other.	2,793	24.7	1.4	1,439	28.7	2.1	1,354	21.5	1.8
Refused/don't know.	*135	*1.2	0.5	*57	*1.1	0.7	*78	*1.2	0.6

... Category not applicable.

* Figure does not meet standard of reliability or precision.

¹"All types of falls mentioned" is greater than the total number of fall episodes because respondents could indicate up to two types of falls.

²"Other (specified)" type of fall includes escalator, building, tree, toilet, bathtub, and pool.

NOTES: Only when fall was selected as the cause of the injury were respondents asked how the injured person fell. Therefore, the total number of fall episodes as shown in this table will not match the total number of fall episodes based on the ICD-9-CM external cause codes. Percents are based on unrounded numbers.

Table 4. Number, percent, and standard error of transportation-related episodes by selected characteristics and sex: United States, 1997

Selected characteristic	Both sexes			Male			Female		
	Number of episodes in thousands	Percent	Standard error	Number of episodes in thousands	Percent	Standard error	Number of episodes in thousands	Percent	Standard error
Injured person ¹									
Driver	2,530	58.0	2.2	1,414	57.2	3.2	1,116	59.0	3.6
Passenger	1,195	27.4	2.2	552	22.3	2.9	643	34.0	3.5
Bicycle.	439	10.1	1.5	422	17.1	2.6	*17	*0.9	0.7
Pedestrian	123	2.8	0.8	*45	*1.8	0.7	*78	*4.1	1.5
Refused/don't know.	*76	*1.8	0.7	*39	*1.6	0.9	*37	*2.0	1.0
Type of vehicle ²									
Passenger car	2,559	68.7	2.9	1,174	59.7	4.1	1,385	78.8	3.3
Light truck.	494	13.3	2.2	249	12.7	2.6	245	13.9	3.0
Motorcycle	231	6.2	1.5	211	10.7	2.7	20	1.1	0.8
Large truck	*146	*3.9	1.2	*134	*6.8	2.0	*13	*0.7	0.7
Other (specified) ³	192	5.2	1.3	130	6.6	1.9	*62	*3.5	1.6
Other.	*102	*2.8	0.9	*69	*3.5	1.4	*34	*1.9	1.0
Refused/don't know.	—	—	—
Seat belt/child restraint ⁴									
Yes.	2,506	78.3	2.7	1,154	74.1	3.9	1,353	82.4	3.4
No	663	20.7	2.7	387	24.8	3.9	276	16.8	3.3
Refused/don't know.	*30	*1.0	0.6	*17	*1.1	0.8	*14	*0.8	0.6
Helmet use ⁵									
Yes.	303	40.7	7.1	286	41.2	7.4	*17	*33.4	21.4
No	432	58.0	7.2	399	57.4	7.5	*33	*66.7	21.4
Refused/don't know.	*10	*1.4	1.4	*10	*1.5	1.5	—

* Figure does not meet standard of reliability or precision.

— Quantity zero.

... Category not applicable.

¹Only persons providing information about injury episodes caused by a vehicle as transportation were asked if the person involved was injured as a driver, passenger, bicycle rider, or pedestrian.

²Only persons providing information about injury episodes in which a person was injured as a driver or passenger of a vehicle were asked about the type of vehicle involved.

³Other types of vehicle included vehicles such as buses, all terrain vehicles, farm equipment, airplanes, and boats.

⁴Only persons providing information about injury episodes in which a person was injured as a driver or a passenger in a passenger car, light truck, or large truck were asked if the injured person was wearing a seat belt or buckled in a safety seat.

⁵Only persons providing information about injury episodes in which a person was injured while riding a bicycle or motorcycle or all-terrain vehicle or ski/snowmobile were asked if the injured person was wearing a helmet.

Table 5. Number of poisoning episodes, number and percent of poisoning episodes involving a call to a poison control center, and standard error by sex and age: United States, 1997

Sex and age	Poisonings	Poison control center called		
	Number of episodes in thousands	Number of episodes in thousands	Percent	Standard error
Both sexes				
All ages	1,945	877	45.1	4.3
Under 6 years	551	415	75.5	6.0
6 years and over.	1,394	461	33.1	4.9
Male				
All ages	898	346	38.6	6.7
Under 6 years	212	155	72.9	9.9
6 years and over.	685	*191	*27.9	7.4
Female				
All ages	1,047	531	50.7	5.5
Under 6 years	338	261	77.1	7.5
6 years and over.	709	270	38.1	6.4

* Figure does not meet standard of reliability or precision.

Table 6. Annual rate and standard error of conditions by the nature of the injury and sex and age : United States, 1997

Sex and age	All conditions ¹		Sprains/strains		Open wounds		Fractures		Contusions	
	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error
Both sexes										
All	153.6	3.4	38.6	1.5	29.3	1.3	23.4	1.0	18.7	1.1
All, age adjusted.	153.7	3.4	38.5	1.5	29.0	1.3	23.7	1.0	19.0	1.2
Under 12 years	118.9	6.4	8.7	1.5	41.9	3.5	18.5	2.1	12.3	1.9
12–21 years	197.3	10.7	58.4	4.8	40.5	4.4	31.1	3.3	20.7	3.3
22–44 years	167.5	6.3	52.0	3.0	29.0	2.3	19.7	1.7	17.2	2.0
45–64 years	120.7	7.0	36.7	3.1	14.1	2.0	19.5	2.3	14.6	2.3
65 years and over.	169.1	9.8	24.0	3.0	24.1	3.2	39.2	4.3	37.2	4.9
Male										
All	168.5	5.3	39.9	2.2	40.3	2.2	25.1	1.6	16.4	1.5
All, age adjusted.	167.0	5.4	39.3	2.2	39.2	2.1	25.5	1.7	16.7	1.6
Under 12 years	134.8	9.3	11.1	2.4	54.2	5.1	15.6	2.7	13.7	2.8
12–21 years	260.4	17.0	67.9	7.2	67.9	8.3	43.2	5.5	24.8	5.0
22–44 years	192.8	9.2	53.9	4.3	38.3	3.6	25.3	3.0	15.2	2.3
45–64 years	115.2	10.4	33.6	4.3	20.3	3.5	17.9	3.6	12.5	3.1
65 years and over.	118.7	14.6	16.1	3.5	21.1	4.4	29.3	5.9	21.4	5.6
Female										
All	139.3	4.5	37.3	2.1	18.8	1.3	21.8	1.4	20.9	1.7
All, age adjusted.	138.5	4.5	37.4	2.1	18.6	1.3	21.7	1.4	20.6	1.7
Under 12 years	102.4	8.1	6.1	1.7	29.0	4.4	21.6	3.1	10.8	2.5
12–21 years	131.4	13.9	48.5	6.9	11.9	3.1	18.5	3.8	16.5	3.8
22–44 years	142.8	8.5	50.1	4.2	19.9	2.8	14.2	2.1	19.3	3.0
45–64 years	125.8	9.0	39.5	4.7	8.3	1.8	21.0	3.1	16.5	3.4
65 years and over.	205.7	14.5	29.7	4.5	26.3	4.3	46.4	6.0	48.8	7.4

* Figure does not meet standard of reliability or precision.

¹Poisoning episodes are assumed to have a single condition resulting from the episode.

NOTES: There may be more than one condition per episode. See appendix III for ICD–9–CM diagnosis codes in category.

Table 7. Annual rate and standard error of injury conditions by body region injured and sex and age: United States, 1997

Sex and age	Upper extremity		Lower extremity		Spine/back		Face		Skull/brain	
	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error	Rate per 1,000 population	Standard error
Both sexes										
All.	43.2	1.6	40.1	1.4	19.8	1.1	15.3	0.8	8.7	0.6
All, age adjusted.	43.2	1.6	40.3	1.4	19.8	1.1	15.1	0.8	8.6	0.6
Under 12 years	27.4	2.6	21.7	2.3	2.8	0.8	30.2	3.0	14.4	1.9
12–21 years	68.7	5.0	60.9	5.0	17.0	2.5	16.6	2.7	12.9	2.7
22–44 years	46.7	2.8	40.3	2.3	31.6	2.5	12.0	1.4	6.4	1.0
45–64 years	30.3	3.0	38.2	3.2	19.7	2.2	5.7	1.1	3.9	0.9
65 years and over.	48.2	4.3	45.4	4.1	14.0	2.5	17.2	2.8	9.7	1.9
Male										
All.	51.2	2.3	38.8	2.2	20.0	1.5	18.9	1.3	10.8	1.0
All, age adjusted.	51.0	2.3	38.4	2.2	20.0	1.6	18.4	1.3	10.5	1.0
Under 12 years	26.9	3.5	25.1	3.9	*4.2	1.4	34.1	4.1	21.4	3.2
12–21 years	95.9	8.1	73.0	7.9	18.4	3.5	27.9	5.0	16.5	3.3
22–44 years	58.8	4.2	40.8	3.7	32.7	3.3	15.9	2.4	8.1	1.5
45–64 years	32.6	4.6	31.8	4.2	17.6	2.8	7.5	2.0	*3.9	1.3
65 years and over.	41.0	6.5	21.1	4.1	*11.9	3.9	*11.1	4.1	*6.3	2.4
Female										
All.	35.6	1.9	41.3	1.9	19.6	1.5	11.8	1.0	6.6	0.9
All, age adjusted.	35.4	1.9	41.2	1.9	19.5	1.5	11.6	1.0	6.6	0.9
Under 12 years	27.8	3.7	18.2	2.8	*1.3	0.8	26.1	4.1	7.1	1.9
12–21 years	40.3	6.2	48.2	6.0	15.6	3.7	*4.7	1.6	*9.0	4.3
22–44 years	35.0	3.4	39.7	3.4	30.6	3.2	8.2	1.4	4.7	1.2
45–64 years	28.3	3.5	44.3	4.3	21.7	3.2	4.1	1.2	*4.0	1.3
65 years and over.	53.5	6.2	63.1	6.6	15.5	3.4	21.6	3.8	12.2	2.7

* Figure does not meet standard of reliability or precision.

NOTES: There may be more than one body part injured per episode. See appendix III for ICD–9–CM diagnosis codes in category.

Table 8. Number, percent, and standard error of injury episodes by place of occurrence, activity engaged in, and sex: United States, 1997

Activity and place	Both sexes			Male			Female		
	Number of episodes in thousands	Percent	Standard error	Number of episodes in thousands	Percent	Standard error	Number of episodes in thousands	Percent	Standard error
All injury episodes	32,438	100.0	...	17,646	100.0	...	14,792	100.0	...
Place									
All places mentioned ¹	32,900	101.4	...	17,911	101.5	...	14,989	101.3	...
Home (inside)	7,832	24.1	0.9	3,098	17.6	1.1	4,734	32.0	1.5
Home (outside)	5,760	17.8	0.7	3,074	17.4	1.0	2,686	18.2	1.2
Street/highway	4,220	13.0	0.7	2,293	13.0	0.9	1,927	13.0	1.0
Sport facility	2,369	7.3	0.5	1,716	9.7	0.8	653	4.4	0.6
Industrial/construction	2,191	6.8	0.5	1,866	10.6	0.8	325	2.2	0.4
School	1,991	6.1	0.5	1,315	7.5	0.7	676	4.6	0.6
Trade/service	1,986	6.1	0.5	1,095	6.2	0.7	892	6.0	0.7
Park/recreation area	1,289	4.0	0.4	866	4.9	0.6	423	2.9	0.5
Other public building	920	2.8	0.3	447	2.5	0.4	473	3.2	0.5
Other (specified) ²	2,414	7.4	0.5	999	5.7	0.6	1,415	9.6	1.0
Other	1,603	4.9	0.4	981	5.6	0.6	622	4.2	0.6
Refused/don't know	324	1.0	0.2	161	0.9	0.3	163	1.1	0.3
Activity									
All activities mentioned ¹	33,060	101.9	...	18,069	102.4	...	14,991	101.3	...
Leisure	7,169	22.1	0.9	3,876	22.0	1.1	3,293	22.3	1.3
Working paid job	6,266	19.3	0.8	4,195	23.8	1.1	2,071	14.0	1.1
Sports	4,458	13.7	0.7	3,233	18.3	1.1	1,225	8.3	0.9
Working (house/yard)	3,363	10.4	0.6	1,521	8.6	0.8	1,842	12.5	1.0
Driving	2,432	7.5	0.5	1,301	7.4	0.7	1,131	7.7	0.7
Attending school	915	2.8	0.3	557	3.2	0.5	358	2.4	0.5
Unpaid work	960	3.0	0.3	420	2.4	0.4	540	3.7	0.5
Other (specified) ³	1,611	5.0	0.4	564	3.2	0.5	1,047	7.1	0.7
Other	5,542	17.1	0.8	2,242	12.7	0.9	3,301	22.3	1.3
Refused/don't know	343	1.1	0.2	160	0.9	0.3	183	1.2	0.3

... Category not applicable.

¹All places mentioned" and "All activities mentioned" are greater than the total number of injury episodes because respondents could indicate up to two places or activities.

²Other (specified)" place includes child care center or preschool, residential institution, health care facility, parking lot, farm, river, lake, stream, ocean, swimming pool, and mine or quarry.

³Other (specified)" activity includes sleeping, resting, eating, drinking, cooking, and being cared for.

NOTES: Percents are calculated based on the sex-specific total number of injury episodes and are also based on unrounded numbers.

Table 9. Number, percent, and standard error of injury and poisoning episodes involving hospitalization by sex and age: United States, 1997

Sex and age	Hospitalized		
	Number of episodes in thousands	Percent hospitalized	Standard error
Both sexes			
All ages	2,466	7.2	0.5
Under 22 years.	506	4.2	0.7
22–64 years.	1,335	7.3	0.7
65 years and over	625	14.8	2.1
Male			
All ages	1,395	7.5	0.7
Under 22 years.	350	4.7	0.8
22–64 years.	837	8.5	1.1
65 years and over	209	17.4	4.1
Female			
All ages	1,071	6.8	0.8
Under 22 years.	156	3.5	1.0
22–64 years.	498	6.0	0.9
65 years and over	417	13.8	2.3

Table 10. Number, annual rate, and standard error of injury and poisoning episodes resulting in time lost from work or school: United States, 1997

Time lost	Number of episodes in thousands	Rate per 1,000 population	Standard error
Time lost from school¹			
Any time lost	2,777	36.2	2.5
Less than 1 day	616	8.0	1.1
1–5 days	1,807	23.6	2.2
6 days or more	354	4.6	0.8
Time lost from work²			
Any time lost	10,054	50.6	1.8
Less than 1 day	1,669	8.4	0.7
1–5 days	4,635	23.3	1.2
6 days or more	3,750	18.9	1.2

¹For persons ages 5–24 years.²For persons 14–75 years.

Table 11. Number, percent, and standard error of injury episodes by whether persons required help with daily routine activities or personal care and sex and age: United States, 1997

Sex and age	Needs help with daily routine activities ¹			Needs help with personal care ²		
	Number of episodes in thousands	Percent	Standard error	Number of episodes in thousands	Percent	Standard error
Both sexes						
All injury episodes, ages 5 years and over	2,232	7.3	0.5	1,785	5.8	0.5
5–21 years	217	2.3	0.6	361	3.9	0.7
22–64 years.	1,158	6.7	0.7	673	3.9	0.5
65 years and over	856	21.2	2.3	751	18.5	2.3
Male						
All injury episodes, ages 5 years and over.	780	4.7	0.6	732	4.4	0.6
5–21 years	*136	*2.2	0.7	200	3.3	0.8
22–64 years.	397	4.2	0.8	278	3.0	0.6
65 years and over	247	21.3	4.8	254	21.9	4.7
Female						
All injury episodes, ages 5 years and over	1,452	10.4	0.9	1,052	7.5	0.8
5–21 years	*81	*2.5	1.0	161	5.0	1.3
22–64 years.	761	9.6	1.2	394	5.0	0.9
65 years and over	609	21.1	2.6	496	17.2	2.5

* Figure does not meet standard of reliability or precision.

¹Daily routine activities includes such things as household chores, doing necessary business, shopping, or getting around for other purposes.²Personal care needs includes such things as eating, bathing, dressing, or getting around the house.

NOTE: The limitation questions were not asked for persons less than 5 years old.

Appendix I

NHIS questionnaire, Family Core: Section II—Injuries

Section II—INJURIES

Injuries are a major health problem. In order to develop new ways to help prevent both accidental and intentional injuries, we need to know more about them. In this next set of questions, I will ask about injuries that happened in the past 3 months; Note here that we are only interested in injuries that required medical advice or treatment.

FIJ.010 DURING THE PAST THREE MONTHS, that is since {91 days before today date}, {were/was} {you/anyone in the family} injured seriously enough that {you/they} got medical advice or treatment?

>FINJ3M< (1) Yes (FIJ.020) (7) Refused (FIJ.300)
 (2) No (FIJ.300) (9) DK (FIJ.300)

FIJ.020 Who was this? (Anyone else?)

>PINJ3MR< [] [] []
 [] [] []

FIJ.030 How many different times in the past three months {were/was} {you/subject's name} injured seriously enough to seek medical advice?

>IJNO3M< Times Injured (01–94): _____

FIJ.040 [If FIJ.030 equals 1, ask:]

When did {subject's name} injury happen?

>IJDATE_M< MONTH: _____
 >IJDATE_D< DAY: _____
 >IJDATE_Y< YEAR: _____

[If FIJ.030 greater than 1, ask:]

Now I'm going to ask a few question about {subject's name} most recent injury. When did that injury happen?

>IJDATE_M< MONTH: _____
 >IJDATE_D< DAY: _____
 >IJDATE_Y< YEAR: _____

[If FIJ.030 equals 2 or more, ask:]

We just talked about {subject's name} injury on {recent injury date}. When did {subject's name} injury BEFORE THAT happen?

>IJDATE_M< MONTH: _____
 >IJDATE_D< DAY: _____
 >IJDATE_Y< YEAR: _____

[FIJ.051 to FIJ.295 are asked for each injury episode]

FIJ.050 At the time of the injury, what part(s) of {subject's name} body was hurt? What kind of injury was it? Anything else?

BODY PART

KIND OF INJURY

>IJBODY1<	_____	>IJKIND1<	_____
>IJBODY2<	_____	>IJKIND2<	_____
>IJBODY3<	_____	>IJKIND3<	_____
>IJBODY4<	_____	>IJKIND4<	_____

FIJ.070 How did {subject's name} injury(s) happen? Please describe fully the circumstances or events leading to the injury(s), and any object, substance, or other person involved.

FR: ENTER THE VERBATIM RESPONSE, PROBING FOR AS MUCH DETAIL AS POSSIBLE, INCLUDING SPECIFICALLY WHAT THE INJURED PERSON WAS DOING AT THE TIME AND ALL CIRCUMSTANCES SURROUNDING THE EVENT. RECORD ALL VOLUNTEERED INFORMATION.

>IJHOW1< _____
 >IJHOW2< _____
 >IJHOW3< _____
 >IJHOW4< _____

FIJ.080

FR: ENTER THE FIRST APPROPRIATE BOX WHICH DESCRIBES THE CAUSE OF THE PERSON'S INJURY FROM THE LIST BELOW.

- >CAUS<
- | | |
|---|-----------------------|
| (1) Vehicle as transportation, including Motor Vehicle/
bicycle/motorcycle/pedestrian/train/boat/airplane(FIJ.090) | (5) Fall (FIJ.170) |
| (2) Gun/being shot (FIJ.190) | (6) Other (FIJ.200) |
| (3) Fire/burn/scald related (FIJ.150) | (7) Refused (FIJ.200) |
| (4) Near drowning/water in lungs (FIJ.160) | (9) DK (FIJ.200) |

FR: THE NEXT SET OF QUESTIONS ARE ASKED TO VERIFY DETAILS OF THE CIRCUMSTANCES SURROUNDING THE INJURY(S). IF YOU ALREADY KNOW THE ANSWER BECAUSE OF THE VERBATIM RESPONSE FOR HOW THE INJURY(S) OCCURRED, VERIFY THE ANSWER WITH THE RESPONDENT. OTHERWISE, ASK THE QUESTION.

FIJ.090 {Were/Was} {you/subject's name} injured as the driver of a vehicle, a passenger in a vehicle, a bicycle rider, or as a pedestrian?

- >MVWHO<
- | | |
|--------------------------------------|--------------------------|
| (1) Driver of a vehicle (FIJ.100) | (4) Pedestrian (FIJ.140) |
| (2) Passenger of a vehicle (FIJ.100) | (7) Refused (FIJ.200) |
| (3) Bicycle rider (FIJ.130) | (9) DK (FIJ.200) |

FIJ.100 What type of vehicle {were/was} {you/subject's name} in?

- >MVTYP<
- | | |
|--|---|
| (01) Passenger car (FIJ.120) | (07) Farm equipment (tractor) (FIJ.200) |
| (02) Light truck (including pickups,
vans and utility vehicles) (FIJ.120) | (08) Airplane (FIJ.200) |
| (03) Bus (FIJ.200) | (09) Boat (FIJ.200) |
| (04) Large truck (FIJ.120) | (10) Train (FIJ.200) |
| (05) Motorcycles (including mopeds,
minibikes) (FIJ.130) | (11) Other (FIJ.200) |
| (06) All terrain vehicle or ski/snow-mobile | (97) Refused (FIJ.200) |
| | (99) DK (FIJ.200) |

FIJ.120 **[If AGE is greater than or equal to 5, ask:]**

{Were/Was} {you/subject's name} wearing a safety belt at the time of the accident?

[Else, ask:]

{Were/Was} {you/subject's name} buckled in a car safety seat at the time of the accident?

- >SBELT<
- | | |
|---------|-------------|
| (1) Yes | (7) Refused |
| (2) No | (9) DK |

(Goto FIJ.200)

(FIJ.130) {Were/Was} {you/subject's name} wearing a helmet at the time of the accident?

- >HELMT<
- | | |
|---------|-------------|
| (1) Yes | (7) Refused |
| (2) No | (9) DK |

(Goto FIJ.200)

FIJ.140 What type of vehicle {were/was} {you/subject's name} struck by?

- >MVHIT<
- | | |
|--|---|
| (01) Passenger car | (07) Farm equipment (tractor) |
| (02) Light truck (including pickups,
vans and utility vehicles) | (08) Bicycle |
| (03) Bus | (09) Train |
| (04) Large truck | (10) Boat (includes all on)
water vehicles |
| (05) Motorcycle (including mopeds
and minibikes) | (11) Other |
| (06) All terrain vehicle or ski or
snow-mobile | (97) Refused |
| | (99) DK |

(Goto FIJ.200)

FIJ.150 What was it that burned/scalded {you/subject's name}?

FR: IF RESPONSE IS FIRE OR SMOKE ASK:

What caused the fire/smoke?

- >BURN<
- | | |
|---|----------------------|
| (01) Cigarette, cigar, pipe | (07) Other explosive |
| (02) Cooking unit | (08) Water or steam |
| (03) Heater | (09) Food |
| (04) Wiring | (10) Chemicals |
| (05) Motor vehicle battery caps,
radiator caps | (11) Other |
| (06) Fireworks | (97) Refused |
| | (99) DK |

(Goto FIJ.200)

FIJ.160 What body of water was involved?

- >WATER<
- | | |
|---------------------|------------------|
| (1) Bathtub | (5) River, creek |
| (2) Swimming pool | (6) Other |
| (3) Lake, pond | (7) Refused |
| (4) Bay, ocean, sea | (9) DK |

(Goto FIJ.200)

FIJ.170 How did {you/subject's name} fall? Anything else?

FR: HAND CARD F4. RECORD UP TO 2 RESPONSES. ENTER 'N' FOR NO MORE.

On or down or from:

- >FALL<
- | | |
|------------------------------|---|
| (1) Escalator | (7) Building or other structure |
| (2) Stairs or steps | (8) Chair, bed, sofa or other furniture |
| (3) Floor/level ground | (9) Tree |
| (4) Curb, including sidewalk | (10) Toilet, commode |
| (5) Ladder or scaffolding | (11) Bathtub, shower |
| (6) Playground equipment | |

Into:

- | | |
|----------------------------|--------------|
| (12) Swimming pool | (97) Refused |
| (13) Hole or other opening | (99) DK |
| (14) Other | |
| [] | [] |

FIJ.180 What caused {you/subject's name} to fall? Was it due to:

- >FWHY<
- | | |
|--|-----------------------|
| (1) Slipping, tripping or stumbling | (5) Or something else |
| (2) Jumping or diving | (7) Refused |
| (3) Collision with/pushing, shoving
by another person | (9) DK |
| (4) Loss of balance/dizziness/becoming faint/seizure | |

(Goto FIJ.200)

FIJ.190 What kind of gun was it?

- <GUNTP>
- | | |
|---------------------------------------|-------------|
| (1) Firearm (handgun, shotgun, rifle) | (4) Other |
| (2) BB or pellet gun | (7) Refused |
| (3) Dart gun | (9) DK |

FIJ.200 What {were/was} {you/subject's name} doing when the injury(s) happened?

FR: HAND CARD F5. RECORD UP TO 2 RESPONSES. ENTER 'N' FOR NO MORE.

- >WHAT<
- | | |
|--|--|
| (1) Driving | (7) Leisure activity (excluding sports) |
| (2) Working at paid job | (8) Sleeping, resting, eating, drinking |
| (3) Working around the house or yard | (9) Cooking |
| (4) Attending school | (10) Being cared for (hands on care from other person) |
| (5) Unpaid work (incl. housework, shopping, volunteer work) | (11) Other |
| (6) Sports (organized team or individual such as running, biking, skating) | (97) Refused sport |
| [] | (99) DK |
| [] | [] |

FIJ.220 Where {were/was} {you/subject's name} when the injury(s) happened?

FR: HAND CARD F6. RECORD UP TO 2 RESPONSES. ENTER 'N' FOR NO MORE.

- >WHERE<
- | | |
|---|---|
| (1) Home (inside) | (11) Farm |
| (2) Home (outside) | (12) Park/recreation area (fields, bike or jog path), |
| (3) School (not residential) | (13) River/lake/stream/ocean |
| (4) Child care center or Preschool | (14) Swimming pool |
| (5) Residential institution (excl. hosp.) | (15) Industrial or construction area |
| (6) Health care facility (incl. hospital) | (16) Mine/quarry |
| (7) Street/highway | (17) Other public building |
| (8) Parking lot | (18) Other |
| (9) Sport facility, ath. field or playground | (97) Refused |
| (10) Trade and service areas (restaurant, store, bank, gas station) | (99) DK |
| [] | [] |

(Goto FIJ.250)

FIJ.240 {Were/Was} {you/subject's name} hospitalized for at least one night as a result of this injury/these injuries?

- >IHOSP<
- | | |
|-------------------|-----------------------|
| (1) Yes (FIJ.250) | (7) Refused (FIJ.260) |
| (2) No (FIJ.260) | (9) DK (FIJ.260) |

FIJ.250 How many nights {were/was} {you/subject's name} in the hospital?

FR: IF "STILL IN HOSPITAL," ASK HOW MANY NIGHTS UP TO TODAY.

- >IHNO<
- | | |
|---------------------|--------------|
| (01-94) 1-94 nights | (97) Refused |
| (95) 95+ nights | (99) DK |

Check item FIJCII: If AGE is greater than 13 then go to FIJ.260; Else
 If AGE is greater than 4 and less than 14 then go to FIJ.270; Else
 If AGE is less than 5 then return to FIJ.040 for next injury event or next person.
 If there are no more persons and no more injuries-events, go to FIJ.300.

FIJ.260 As a result of this injury/these injuries, how much work did {you/subject's name} miss?

FR: HAND CARD F7.

- >WKLS<
- | | |
|----------------------|--|
| (0) None | (6) Not employed at the time of the injury |
| (1) Less than 1 day | (7) Refused |
| (2) 1 to 5 days | (9) DK |
| (3) Six or more days | |

FIJ.270 As a result of this injury/these injuries, how much school did {you/subject's name} miss?

FR: HAND CARD F8.

- >SCLS< (0) None (6) Not in school at the time of the injury
 (1) Less than 1 day (7) Refused
 (2) One to five days (9) DK
 (3) Six or more days

FIJ.280 As a result of this injury/these injuries {do/does} {you/subject's name} now need the help of other persons with {your/his/her} personal care needs, such as eating, bathing, dressing or getting around this home?

- >IJADL< (1) Yes (FIJ.285) (7) Refused (FIJ.290)
 (2) No (FIJ.290) (9) DK (FIJ.290)

FIJ.285 Do you expect {you/subject's name} will need this help for a total of 6 months or longer?

- >LIMTM< (1) Yes (7) Refused
 (2) No (9) DK

FIJ.290 As a result of this injury/these injuries {do/does} {you/subject's name} now need the help of other persons in handling routine needs such as everyday household chores, doing necessary business, shopping or getting around for other purposes?

- >IJIAD< (1) Yes (FIJ.295) (7) Refused (FIJ.040/FIJ.300)
 (2) No (FIJ.040/FIJ.300) (9) DK (FIJ.040/FIJ.300)

FIJ.295 Do you expect {you/subject's name} will need this help for a total of 6 months or longer?

- >HLIMIT< (1) Yes (FIJ.040/FIJ.300) (7) Refused (FIJ.040/FIJ.300)
 (2) No (FIJ.040/FIJ.300) (9) DK (FIJ.040/FIJ.300)

FIJ.300 The next questions are about POISONING, which includes coming into contact with harmful substances, and overdose or wrong use of any drug or medication. Do not include any illnesses such as poison ivy or food poisoning.

FR: HAND CALENDAR CARD.

DURING THE PAST THREE MONTHS, that is since {91 days before today's date}, did {you/anyone in the family} have a poisoning that caused someone to seek medical advice or treatment, including calls to a poison control center?

- >FPOIS3M< (1) Yes (FIJ.310) (7) Refused (FAU.010)
 (2) No (FAU.010) (9) DK (FAU.010)

FIJ.310 Who was this? (Anyone else?)

- >PPOIS3MR< [] [] []
 [] [] []

FIJ.320 How many different times in the PAST THREE MONTHS {were/was} {you/subject's name} poisoned?

- (01-94) 1-94 times (97) Refused
 (95) 95+ times (99) DK

FIJ.330 **[If FIJ.320 equals 1, ask:]**

When did {subject's name} poisoning happen?

- >POIDTEM< MONTH: _____
 >POIDTED< DAY: _____
 >POIDTEY< YEAR: _____

[If FIJ.320 is greater than 1, ask:]

Now I'm going to ask a few question about {subject's name} most recent poisoning. When did that happen?

>POIDTEM< MONTH: _____
 >POIDTED< DAY: _____
 >POIDTEY< YEAR: _____

[IF FIJ.320 is greater than or equal to 2, ask:]

We just talked about {subject's name} poisoning on {recent poisoning}. When did {subject's name} poisoning BEFORE THAT happen?

>POIDTEM< MONTH: _____
 >POIDTED< DAY: _____
 >POIDTEY< YEAR: _____

[FIJ.340 to FIJ.410 are repeated for each poisoning episode.]

FIJ.340 Did {you/subject's name} poisoning result from:

- >POITPR2< (1) a drug or medical substance used mistakenly or in overdose (FIJ.360)
 (2) a harmful or toxic solid or liquid substance (FIJ.360)
 (3) inhaling gases or vapors (FIJ.360)
 (4) eating a poisonous plant or other substance mistaken for food (FIJ.360)
 (5) a venomous animal or plant (FIJ.360)
 (6) something else (FIJ.350)
 (7) Refused (FIJ.360)
 (9) DK (FIJ.360)

FIJ.350

FR: ENTER THE VERBATIM RESPONSE.

>PSPEC_1< _____
 >PSPEC_2< _____
 >PSPEC_3< _____
 >PSPEC_4< _____

FIJ.360 Did you or did someone else call a poison control center for advice in treating {subject's name} poisoning?

- >POICC< (1) Yes (7) Refused
 (2) No (9) DK

FIJ.370 {Were/Was} {you/subject's name} hospitalized for at least one night as a result of this poisoning?

- >PHOSP< (1) Yes (FIJ.380) (7) Refused (FIJ.390)
 (2) No (FIJ.390) (9) DK (FIJ.390)

FIJ.380 How many nights {were/was} {you/subject's name} in the hospital?

FR: IF "STILL IN HOSPITAL," ASK HOW MANY NIGHTS UP TO TODAY.

- >PHNO< (01-94) 1-94 nights (97) Refused
 (95) 95+ nights (99) DK

Check item FIJCCI2: If AGE greater than 13 then go to FIJ.400; Else
 If AGE greater than 4 and less than 14 then go to FIJ.410; Else
 If AGE less than 5 then return to FIJ.330 for the next poisoning event or the next person.
 If there are no more persons and no more poisoning events, go to FAU.010.

FIJ.400 As a result of this poisoning, how much work did {you/subject's name} miss?

FR: HAND CARD F7.

- >PWKLS< (0) None (6) Not employed at the time of poisoning
 (1) Less than 1 day (7) Refused
 (2) One to five days (9) DK
 (3) Six or more days

FIJ.410 As a result of this poisoning, how many days of school did {you/subject's name} miss?

FR: HAND CARD F8.

- >PSCLS<
- | | |
|----------------------|--|
| (0) None | (6) Not in school at the time of poisoning |
| (1) Less than 1 day | (7) Refused |
| (2) One to five days | (9) DK |
| (3) Six or more days | |

(Goto next section—Health Care Access and Utilization)

Appendix II

Matrix for External Cause of Injury Mortality and Morbidity Data

Mechanism or cause	Manner or intent				
	Unintentional	Suicide/ self-inflicted	Homicide/ Assault	Undetermined	Other
Cut/pierce	E920.0–.9	E956	E966	E986	E974
Drowning/submersion	E830.0–.9, 832.0–.9, 910.0–.9	E954	E964	E984	...
Fall	E880.0–886.9, 888	E957.0–.9	E968.1	E987.0–.9	...
Fire/hot object or substance	E890.0–899, 924.0–.9	E958.1,.2,.7	E961, 968.0, .3	E988.1,.2,.7	...
Fire/flame	E890.0–899	E958.1	E968.0	E988.1	...
Hot object/scald	E924.0–.9	E958.2,.7	E961, 968.3	E988.2,.7	...
Firearm	E922.0–.9	E955.0–.4	E965.0–.4	E985.0–.4	E970
Machinery	E919.0–.9
Motor vehicle traffic	E810–819 (.0–.9)	E958.5	E968.5	E988.5	...
Occupant	E810–819 (.0,.1)
Motorcyclist	E810–819 (.2,.3)
Pedal cyclist	E810–819 (.6)
Pedestrian	E810–819 (.7)
Unspecified	E810–819 (.9)
Pedal cyclist, other	E800–807 (.3), E820–825 (.6), E826 (1,.9), 827–829 (.1)
Pedestrian, other	E800–807 (.2), 820–825 (.7), E826–829 (.0)
Transport, other	E800–807 (.0–.1,.8–.9), E820–825 (.0–.5,.8–.9), E826 (.2–.8), 827–829, (.2–.9) E831.0–.9, 833.0–845.9	E958.6	...	E988.6	...
Natural/environmental factors	E900.0–909, 928.0–.2	E958.3	...	E988.3	...
Bites and stings	E905.0–.6,.9, 906.0–.4, .5, .9
Overexertion	E927
Poisoning	E850.0–869.9	E950.0–952.9	E962.0–.9	E980.0–982.9	E972
Struck by, against	E916–917.9	...	E960.0, 968.2	...	E973, 975
Suffocation	E911–913.9	E953.0–.9	E963	E983.0–.9	...
Other specified and classifiable	E846–848, 914–915, 918, E921.0–.9, 923.0–.9, E925.0–926.9, 929.0–.5	E955.5,.9, 958.0,.4	E960.1, 965.5–.9, E967.0–.9, 968.4	E985.5, 988.0,.4	E971, 978 E990–994, 996, E997.0–.2
Other specified, NEC ¹	E928.8, 929.8	E958.8, 959	E968.8, 969	E988.8, 989	E995, 997.8, 977, E998–999
Unspecified	E887, 928.9, 929.9	E958.9	E968.9	E988.9	E976, 997.9
All injury ²	E800–869, 880–929	E950–959	E960–969	E980–989	E970–978 E990–999

¹Not elsewhere classifiable.

²Excludes fatal and nonfatal events caused by adverse events (E-codes E870–E879 and E930–E949).

NOTE: E968.5 and E906.5 are the only codes that are singled out that are in ICD–9–CM but not in ICD-9. All of the other codes that are in CM only are folded into larger groupings in the matrix.

Appendix III

Injury Morbidity Matrix

Codes for Body Region of Injury

Please note: the following list can be used if ICD is coded to the 4th digit; if only 3-digit codes are available follow instructions next to **.

1.	Skull and Brain: excl. face ^(1,2) (incl. scalp)	800–801, 803–804 850–854 873.0–873.1 873.8–873.9 951	** Code 873 under Other ** Code 873 under Other
2.	Face	802 830 848.0–848.1 870–872 873.2–873.7 910 918 920–921 925.1 940 947.0 950	** Code 848 under Other ** Code 873 under Other ** Code 925 under Other ** Code 947 under Other
	Head ^(1,2) (Skull & Brain & Face)	800–804 850–854 870–73 830 848.0–848.1 910 918 920–921 925.1 940 947.0 950–951	** Code 848 under Other ** Code 925 under Other ** Code 947 under Other
3.	Neck ^(2,3,5,8)	807.5–807.6 848.2 874 900 925.2	**Code 807 under Thorax **Code 848 under Other **Code 925 under Other **Code 947 under Other
4.	Thorax ^(4,5)	947.1 807.0–807.4 848.3–848.4 860–862 875 879.0–879.1 901 922.0–922.1 947.2	**Code 807 under Thorax **Code 848 under Other **Code 879 under Other **Code 922 under Other **Code 947 under Other
5.	Abdomen, pelvic contents, genital organs	863–868 878 879.2–879.5 902 922.2 922.4 926.0 947.3–947.4	**Code 879 under Other **Code 922 under Other **Code 922 under Other **Code 926 under Other **Code 947 under Other

See footnotes at end of table.

Please note: the following list can be used if ICD is coded to the 4th digit; if only 3-digit codes are available follow instructions next to **.

6.	Spine and Back ^(6,7,8)	805 806 876–877 922.3 839.0–839.5 847 952–953	**Code 922 under Other **Code 839 under Other
7.	Upper Extremity	810–818 831–834 840–42 880–887 903 912–915 923 927 943–944 955 959.2–959.5	**Code 959 under Other
8.	Lower Extremity and Bony Pelvis ⁽⁶⁾	808 821–27 835–838 843–845 846 848.5 890–897 904 916–917 924 928 945 956 959.6–959.7	**Code 848 under Other **Code 959 under Other
17.	(Neck of femur fracture)⁽⁹⁾	820	
9.	Other and Ill-Defined Body Region	809 819 828 829 839.6–839.9 848.8–848.9 869 879.6–879.9 911 919 922.8–922.9 926.1 926.8–926.9 929 941–42 946 947.8–947.9 948–949 954 957 959.0–959.1 959.8–959.9	** Code 839 under Other ** Code 848 under Other **Code 879 under Other **Code 922 under Other **Code 926 under Other **Code 926 under Other **Code 947 under Other **Code 959 under Other **Code 959 under Other
10.	Foreign Bodies	930–939	
11.	Poisonings	960–979	

See footnotes at end of table.

Please note: the following list can be used if ICD is coded to the 4th digit; if only 3-digit codes are available follow instructions next to **.

12.	Toxic Effects	980–989
13.	Other and Unspec Effects of External Causes	990–995
	Effects of reduced temperature	991
	Effects of heat and light	992
	Drowning	994.1
	Asphyxiation and strangulation	994.7
	Electrocution	994.8
	All other Effects of External Causes	990, 993, 994.0, 994.2–994.6, 994.9, 995
14.	Late Effects	905–909
15.	Early Traumatic Complications	958
16.	Complications of Surgical and Medical Care	996–999
18.	No Injury	No diagnosis codes above 799

¹Include 804 under **Head** (instead of **Multiple Body Regions**) even though it reads: Multiple fractures involving skull or face with other bones: assume that principal fracture is to the skull or face.

²Code all injuries to blood vessels of Head or Neck (900) under **Neck**; it is not easy to distinguish whether blood vessel is part of head or neck based only on third or fourth digit of ICD

³Injuries to trachea (typically categorized at 4th or 5th digit which is not available for mortality data) is classified under **Neck** (instead of **Thorax**)

⁴Injuries to the trunk unless otherwise specified are coded under **Other** since these injuries could be to the region of the thorax, abdomen or back

⁵Fx to larynx and trachea (807.5–807.6) are coded under **Neck** unless 3rd digit code only, then code under **thorax** and assume injury (fx) is more likely to be to ribs and /or sternum.

⁶Injuries to sacrum and coccyx are coded under **Spine** as they are typically only distinguishable from other injuries to the spine at the 4th or 5th digits.

⁷Injuries to buttock region (eg. 877) are coded under **Spine and Back**

⁸Injuries classified under Neck include only those injuries to the front of the neck or soft tissue; injuries to the neck portion of the spine are classified under **Spine and Back**

⁹Neck of femur fractures have been classified separately.

Codes for Nature of Injury

1.	Fractures ^(1,2)	800–805; 807–829
2.	Dislocations	830–839
3.	Sprains and Strains	840–848
4.	Crushing Injury	925–929
5.	Amputation of Limbs	885–887; 895–897
6.	Injury to Internal Organs ^(2,3,4,5) incl. CNS injuries	860–869 850–854 952–953 806
7.	Nerves ⁽⁴⁾	950–951; 954–957
8.	Blood Vessels	900–904
9.	Open Wounds ^(3,5)	870–884, 888–894
10.	Superficial Injuries	910–919
11.	Contusions	920–924
12.	Burns	940–949
13.	Effects of Foreign Bodies	930–939
14.	Other Injury- (other and unspecified)	959
	Multiple sites	959.8
	All other sites	959.0–959.7
	Unspecified sites	959.9
15.	Poisonings	960–979
16.	Toxic Effects	980–989
17.	Other and Unspec. Effects of External Causes	990–995
	Effects of reduced temperature	991
	Effects of heat and light	992
	Drowning	994.1
	Asphyxiation and strangulation	994.7
	Electrocution	994.8
	All other Effects of External Causes	990, 993, 994.0, 994.2–994.6, 994.9, 995
18.	Late Effects of Injuries etc.	905–909
19.	Early Complications of trauma	958
20.	Complications of Surgical and Medical Care	996–999
21.	No Injury	No diagnosis codes above 799

¹**Fractures** include skull fractures with intracranial injury; **HOWEVER**, if data are coded to the fourth digit; include the following codes (i.e. intracranial injuries with skull fx) under Injury to Internal Organs:

800.1– 800.4	801.1–801.4
800.6–800.9	801.6–801.9
803.1–803.4	804.1–804.4
803.6–803.9	804.6–804.9

²**Fractures** exclude spine fxs with SCI; they are classified under **Injuries to Internal Organs**; (3) Injuries to Internal Organs include CNS injuries (injuries to the brain and spinal cord); they also include injuries to larynx, trachea, pharynx and thyroid; they do NOT include injuries to internal structures of the eye, ear, and nose (these are included under Open Wounds);

⁴Injuries to Nerves exclude injuries to nerve roots to spine and spinal plexus (953)—these are included under Injury to Internal Organs;

⁵**Open Wounds** includes injuries to the larynx, trachea, pharynx and thyroid; **HOWEVER**, if data are only coded to the fourth digit, include codes 874.0–874.5 (i.e. injuries to larynx, trachea, pharynx and thyroid) under **Injury to Internal Organs**.

⁶The United States Multiple Cause of Death does not include 4th digit classification for intracranial injuries with skull fx (800–804) or injuries to larynx, trachea, pharynx and thyroid (874.0–874.5).

Appendix IV

Technical Notes

The Methods section of this report mentioned that tests of significance for differences between rates or percents would not be conducted. However, for individual rates and percents, confidence intervals and standard errors are provided to indicate the stability of the point estimates. The confidence intervals are displayed in the figures in a way that graphically portrays the sizes of differences in the point estimates relative to their individual stabilities. Examination of the amount of overlap between intervals is not equivalent to standard significance testing for differences.

If tests of significance for differences were to be performed, there would be several technical considerations. Suppose that it were desired to test whether two population percents, say p_1 and p_2 , are different. The standard error of $\hat{p}_1 - \hat{p}_2$ can be written as

$$SE_{12} = \sqrt{SE_1^2 + SE_2^2 - 2 \text{Cov}(\hat{p}_1, \hat{p}_2)}, \quad (1)$$

where SE_1 and SE_2 are the standard errors of \hat{p}_1 and \hat{p}_2 , respectively, “Cov” denotes covariance, and “hats” are used to label estimates. If an estimate \hat{SE}_{12} of equation (1) were computed, either via separate estimation of the components on the right-hand side of equation (1) or via another technique, then an approximate significance test at the 5% level for whether p_1 and p_2 are different could be performed by checking whether the confidence interval

$$\hat{p}_1 - \hat{p}_2 \pm 1.96 \hat{SE}_{12} \quad (2)$$

contains 0.

A nonzero value for $\text{Cov}(\hat{p}_1, \hat{p}_2)$ in equation (1) could occur as a result of the design of the National Health Interview Survey. For example, point estimates for any two groups might be correlated due to the clustering of subjects by household or primary sampling unit. A nonzero covariance could also occur if the groups being

compared have subjects in common. This is possible, for example, when external causes of injury are being compared because a person could have more than one episode and each episode could have a different external cause.

If the covariance between \hat{p}_1 and \hat{p}_2 were assumed to be equal to 0, then by equations (1) and (2), a test of significance for whether p_1 and p_2 are different could be performed by checking whether the interval

$$\hat{p}_1 - \hat{p}_2 \pm 1.96 \sqrt{\hat{SE}_1^2 + \hat{SE}_2^2} \quad (3)$$

contains 0.

In addition to the above considerations, the issue of adjusting for multiple comparisons arises when more than one test of significance is to be carried out.

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *Monthly Vital Statistics Report* (MVSR). These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the MVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Data Dissemination Branch
National Center for Health Statistics
Centers for Disease Control and Prevention
6525 Belcrest Road, Room 1064
Hyattsville, MD 20782-2003
(301) 458-4636
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs/

**DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
6525 Belcrest Road
Hyattsville, Maryland 20782-2003

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

STANDARD MAIL (A)
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284