

**Data Brief 57: Health Insurance Affects Diagnosis and Control of Hypercholesterolemia and Hypertension Among Adults Aged 20–64: United States, 2005–2008**

Data table for Figure 1. Screening for high cholesterol in the last 5 years among adults aged 20-64, by insurance coverage and age: United States, 2005-2008

Age	Private insurance	Public insurance	No insurance
	Percent (standard error)		
Total	72.2 (1.1)	67.6 (1.7)	37.1 (1.0)
20-39 years	55.8 (1.7)	46.5 (3.0)	26.8 (1.1)
40-64 years	83.4 (1.1)	83.0 (1.5)	52.2 (2.1)

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 2005-2008.

**Data Brief 57: Health Insurance Affects Diagnosis and Control of Hypercholesterolemia and Hypertension Among Adults Aged 20–64: United States, 2005–2008**

Data table for Figure 2. Uncontrolled hypercholesterolemia among adults aged 20-64 with hypercholesterolemia, by insurance coverage and age: United States, 2005-2008

Age	Private insurance	Public insurance	No insurance
Percent (standard error)			
All			
Total	61.6 (2.2)	57.7 (3.6)	81.5 (2.4)
20-39 years	85.7 (3.1)	87.3 (4.2)	93.8 (2.8)
40-64 years	55.7 (2.0)	50.8 (3.7)	75.5 (3.2)
Diagnosed and uncontrolled			
Total	34.4 (2.1)	32.3 (2.7)	33.5 (2.8)
20-39 years	36.5 (4.7)	23.5 (5.1)	28.4 (4.6)
40-64 years	33.9 (2.3)	34.4 (3.2)	36.1 (3.4)
Undiagnosed			
Total	27.1 (1.4)	25.4 (3.4)	48.0 (2.9)
20-39 years	49.2 (5.3)	63.8 (6.4)	65.4 (4.8)
40-64 years	21.8 (1.7)	16.4 (3.0)	39.4 (3.6)

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 2005-2008.

**Data Brief 57: Health Insurance Affects Diagnosis and Control of Hypercholesterolemia and Hypertension Among Adults Aged 20–64: United States, 2005–2008**

Data table for Figure 3. Uncontrolled hypertension among adults aged 20-64 with hypertension, by insurance coverage and age: United States, 2005-2008

Age	Private insurance	Public insurance	No insurance
Percent (standard error)			
All			
Total	51.8 (1.7)	44.6 (3.1)	71.3 (3.2)
20-39 years	64.6 (4.7)	50.8 (7.2)	83.0 (5.7)
40-64 years	49.8 (1.7)	43.6 (2.8)	67.9 (3.5)
Diagnosed and uncontrolled			
Total	31.0 (1.9)	28.3 (2.3)	30.8 (2.9)
20-39 years	29.0 (3.9)	24.6 (6.5)	19.1 (5.0)
40-64 years	31.3 (2.0)	28.9 (2.8)	34.2 (3.5)
Undiagnosed			
Total	20.8 (1.7)	16.3 (2.6)	40.5 (3.4)
20-39 years	35.6 (5.3)	26.1 (7.0)	63.9 (6.5)
40-64 years	18.5 (1.5)	14.7 (2.7)	33.6 (3.7)

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 2005-2008.

**Data Brief 57: Health Insurance Affects Diagnosis and Control of Hypercholesterolemia and Hypertension Among Adults Aged 20–64: United States, 2005–2008**

Data table for Figure 4. Health insurance coverage among adults aged 20-64 with hypercholesterolemia or hypertension, by age: United States, 2005-2008

Age	Private insurance	Public insurance	No insurance
	Percent (standard error)		
Total	68.3 (1.6)	14.2 (0.8)	17.5 (1.3)
20-39 years	61.2 (3.1)	13.6 (1.8)	25.2 (2.8)
40-64 years	70.2 (1.8)	14.4 (0.8)	15.4 (1.5)

SOURCE: CDC/NCHS, National Health and Nutrition Examination Survey, 2005-2008.