

**Services and Protection Objective: Tuberculosis identification**

PHS Agency Assignment: National Immunization Program; National Center for Infectious Diseases

Rev.- 10/22/99

**20.17 for actively identifying cases of tuberculosis and latent infection in populations at high risk for tuberculosis.**

Tuberculosis identification programs	1992-93 Baseline	2000 Target
Local health department	80%	90%

*Note: Local health department refers to any local component of the public health system, defined as an administrative and service unit of local or State government concerned with health and carrying some responsibility for the health of a jurisdiction smaller than a State.*

Data Source: National Profile of Local Health Departments, National Association of County and City Health Officials.

**Services and Protection Objective: Preventive therapy for tuberculosis**

PHS Agency Assignment: National Immunization Program; National Center for Infectious Diseases

Rev.- 10/22/99

**20.18 Increase to at least 85 percent the proportion of people found to have tuberculosis infection who completed courses of preventive therapy.**

Percent completing preventive therapy	Baseline Year	Baseline	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	2000 Target
Tuberculosis-infected person	1987	66.3% <sup>a</sup>	67.7% <sup>b</sup>	61.0% <sup>c</sup>	63.0% <sup>d</sup>	64.9% <sup>e</sup>	66.3% <sup>f</sup>	65.3% <sup>g</sup>	65%	65.40%	65.20%	62.20%	85%

- <sup>a</sup>Reports from 89 reporting areas from which 64% of tuberculosis cases were reported.
- <sup>b</sup>Reports from 89 reporting areas from which 62% of tuberculosis cases were reported.
- <sup>c</sup>Reports from 91 reporting areas from which 77% of tuberculosis cases were reported.
- <sup>d</sup>Reports from 93 reporting areas from which 71% of tuberculosis cases were reported.
- <sup>e</sup>Reports from 88 reporting areas from which 70% of tuberculosis cases were reported.
- <sup>f</sup>Reports from 101 reporting areas from which 78% of tuberculosis cases were reported.
- <sup>g</sup>Reports from 108 reporting areas from which 78% of tuberculosis cases were reported.

**Data Source:** Tuberculosis Program Management Report Data on Completion of Preventive Therapy CDC, NCHSTP.

**Services and Protection Objective: Laboratory capability for influenza diagnosis**

**PHS Agency Assignment: National Immunization Program; National Center for Infectious Diseases**

**Rev.- 10/22/99**

**20.19 Increase to at least 85 percent the proportion of tertiary care hospital laboratories and to at least 50 percent the proportion of secondary care hospital and health maintenance organization laboratories possessing technologies for rapid viral diagnosis of influenza.**

Rapid viral diagnosis of influenza	Baseline		1995	1997	1999 <sup>a</sup>	2000
	Year	Baseline				Target
Tertiary care hospital laboratories	1993	52%	57%	58%	64%	85%
Secondary care hospitals	1993	45%	46%	46%	51%	50%
Health maintenance organizations	1993	<sup>b</sup> 68%	56%	68%	67%	50%

- <sup>a</sup> 1999 Data are provisional.
- <sup>b</sup> Baseline data have been revised.

**Data Sources:** Survey of Laboratories using Rapid Viral Diagnosis of Influenza, CDC, NCID, Viral and Rickettsial Disease Division.