

SAFER • HEALTHIER • PEOPLE™



Monitoring the

Nation's Health

Vital and Health Statistics

October 2006

Series 13, Number 162

National Hospital Discharge Survey: 2004 Annual Summary With Detailed Diagnosis and Procedure Data



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Copyright information

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

Suggested citation

Kozak LJ, DeFrances CJ, Hall MJ. National Hospital Discharge Survey: 2004 annual summary with detailed diagnosis and procedure data. National Center for Health Statistics. Vital Health Stat 13(162). 2006.

Library of Congress Catalog Card Number 97-10617

For sale by the U.S. Government Printing Office
Superintendent of Documents
Mail Stop: SSOP
Washington, DC 20402-9328
Printed on acid-free paper.

Vital and Health Statistics

Series 13, Number 162

National Hospital Discharge Survey: 2004 Annual Summary With Detailed Diagnosis and Procedure Data

Data From the National Health Care
Survey

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Centers for Disease Control and Prevention
National Center for Health Statistics

Hyattsville, Maryland
October 2006
DHHS Publication No. (PHS) 2006-1733

National Center for Health Statistics

Edward J. Sondik, Ph.D., *Director*

Jennifer H. Madans, Ph.D., *Acting Co-Deputy Director*

Michael H. Sadagursky, *Acting Co-Deputy Director*

Jennifer H. Madans, Ph.D., *Associate Director for Science*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Michael H. Sadagursky, *Associate Director for Management and Operations*

Lawrence H. Cox, Ph.D., *Associate Director for Research and Methodology*

Margot A. Palmer, *Director for Information Technology*

Margot A. Palmer, *Acting Director for Information Services*

Linda T. Bilheimer, Ph.D., *Associate Director for Analysis, Epidemiology, and Health Promotion*

Charles J. Rothwell, M.S., *Director for Vital Statistics*

Jane E. Sisk, Ph.D., *Director for Health Care Statistics*

Jane F. Gentleman, Ph.D., *Director for Health Interview Statistics*

Clifford L. Johnson, *Director for Health and Nutrition Examination Surveys*

Division of Health Care Statistics

Jane E. Sisk, Ph.D., *Director*

Robin Remsburg, Dr.N., *Deputy Director*

Irma Arispe, Ph.D., *Associate Director for Science*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

Charles Adams, *Chief, Technical Services Branch*

Contents

Abstract	1
Highlights.....	1
Patient and Hospital Characteristics	1
Diagnoses.....	1
Procedures	2
Deliveries and Newborn Infants	2
Introduction	3
Methods	4
Data Source	4
Estimation	5
Use of Tables.....	5
References	6
Appendix	193
Technical Notes	193

Text Figures

1. Percent distribution of discharges and days of care by age: United States, 2004.....	2
2. Percent of discharges transferred to long-term care institutions by age: United States, 2004	2
3. Rate of discharges in major diagnostic categories by sex: United States, 2004	3
4. Number of discharges with human immunodeficiency virus diagnoses by age: United States, 1995, 2000, and 2004	3
5. Rate of cardiac catheterizations by sex, age, and geographic region: United States, 2004.....	4
6. Average length of stay for women with deliveries by type of delivery: United States, 1980–2004	4

Detailed Tables

Tables on Patient and Hospital Characteristics

1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2004	8
2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2004	9
3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2004	10
4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2004	12
5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2004	15
6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, according to sex, age, and geographic region: United States, 2004	17
7. Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2004.....	19
8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2004	20

Diagnosis Tables

9.	Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2004	22
10.	Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2004	23
11.	Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004	25
12.	Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004	26
13.	Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004..	27
14.	Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004	28
15.	Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004	29
16.	Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004..	30
17.	Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004 ..	31
18.	Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004	32
19.	Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004.....	33
20.	Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2004	34
21.	Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2004	35
22.	Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2004.....	36
23.	Number and rate of discharges from short-stay hospitals with a human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years 1990–2004.....	37
24.	Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2002–2004.....	38
25.	Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2004.....	38

Procedure Tables

26.	Number of discharges from short-stay hospitals with and without procedures and percentage with procedures, by selected characteristics: United States, 2004.....	39
27.	Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals by selected procedure categories: United States, 2004.....	40
28.	Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2004	41
29.	Number and rate of discharges from short-stay hospitals with a coronary artery bypass graft and number and rate of procedures, by sex, age, and geographic region: United States, 2004.....	42
30.	Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2004	43
31.	Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2004.....	44
32.	Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2004.....	45
33.	Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2004.....	46
34.	Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2004.....	47
35.	Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2004.....	48

Tables on Deliveries and Newborn Infants

36.	Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2004 ..	49
37.	Number and percent distribution of females with deliveries discharged from short-stay hospitals, by length of stay and type of delivery: United States, selected years 1980–2004.....	50
38.	Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2004.....	51
39.	Number and rate of females with deliveries discharged from short-stay hospitals, by type of delivery: United States, selected years 1980–2004.....	51

40.	Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 2004.	52
41.	Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2004	53
42.	Number and percent distribution of newborn infants discharged from short-stay hospitals, by length of stay, and average length of stay, by health status: United States, selected years 1980–2004	53
43.	Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2004	54
44.	Number and percentage of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years 1980–2004.	54

Diagnosis, Days of Care, and Procedure Tables by ICD–9–CM Code

45.	Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004.	55
46.	Number of days of care for discharges from short-stay hospitals, by ICD–9–CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004	83
47.	Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004	111
48.	Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004.	173

Appendix Tables

I.	Civilian population by age, geographic region, and sex: United States, 2004	193
II.	Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics, 2004	196
III.	Code numbers for Barell injury diagnosis matrix categories in Table 24	198
IV.	Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD–9–CM category, 2004.	199
V.	Changes in ICD–9–CM diagnosis codes, 1986–2003.	201
VI.	Changes in ICD–9–CM in procedure codes, 1986–2003	207

Appendix Figure

I.	Medical abstract form for the National Hospital Discharge Survey, 2004	194
----	-----------------------------------------------------------------------------	-----

Objectives

This report presents 2004 national estimates and selected trend data on the use of nonfederal short-stay hospitals in the United States. Estimates are provided by selected patient and hospital characteristics, diagnoses, and surgical and nonsurgical procedures performed. Estimates of diagnoses and procedures are presented according to International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM) codes.

Methods

The estimates are based on data collected through the National Hospital Discharge Survey (NHDS). The survey has been conducted annually since 1965. In 2004, data were collected for approximately 371,000 discharges. Of the 476 eligible nonfederal short-stay hospitals in the sample, 439 (92 percent) responded to the survey.

Results

An estimated 34.9 million inpatients were discharged from nonfederal short-stay hospitals in 2004. They used 167.9 million days of care and had an average length of stay of 4.8 days. Hospital use by age ranged from 4.3 million days of care for patients 5–14 years of age to 31.8 million days of care for 75–84 year olds. Almost a third of patients 85 years and over were discharged from hospitals to long-term care institutions. Diseases of the circulatory system was the leading diagnostic category for males. Childbirth was the leading category for females, followed by circulatory diseases. The proportion of HIV discharges who were 40 years of age and over increased from 40 percent in 1995 to 67 percent in 2004. The rate of cardiac catheterizations was higher for males than for females and higher for patients 65–74 and 75–84 years of age than for older or younger groups. The average length of stay for both vaginal and cesarean deliveries decreased from 1980 through 1995 but stays for vaginal deliveries increased 24 percent during the period from 1995 to 2004.

Keywords: *hospitalization • inpatient • diagnoses • procedures*

National Hospital Discharge Survey: 2004 Annual Summary with Detailed Diagnosis and Procedure Data

by *Lola Jean Kozak, Ph.D.; Carol Jean DeFrances, Ph.D.; and Margaret Jean Hall, Ph.D., Division of Health Care Statistics*

Highlights

Patient and Hospital Characteristics

- An estimated 34.9 million inpatients were discharged from nonfederal short-stay hospitals in 2004. They used an estimated 167.9 million days of care and were hospitalized for an average of 4.8 days.
- Hospital use ranged from almost a million discharges (3 percent) and 4.3 million days of care (3 percent) for patients 5–14 years of age to 5.5 million discharges (16 percent) and 31.8 million days of care (19 percent) for patients 75–84 years of age (Figure 1).
- The average length of a hospital stay ranged from 4.4 days in the Midwest to 5.4 days in the Northeast.
- Discharges 15–44 years of age made up 58 percent of discharges in the self-pay category but only 31 percent of total discharges. Hospitals in the South accounted for 54 percent of self-pay discharges, compared with 38 percent of total discharges.
- Average length of stay increases with hospital size. In 2004 it ranged from 4.2 days in hospitals with less than 100 beds to 5.6 days in hospitals with 500 beds or more.
- Only about 2 percent of patients under 65 years of age, but nearly

20 percent of patients 65 years of age and over and almost a third of those 85 years of age and over were discharged to long-term care institutions following their hospitalizations (Figure 2).

Diagnoses

- Figure 3 shows leading diagnostic categories, identified by ranking ICD-9-CM Chapters 1–17 and childbirth. For males, the discharge rate for circulatory diseases was significantly higher than the rate for any other diagnostic category. For females the discharge rate for childbirth was highest and the rate for circulatory diseases was second highest.
- Males and females had similar discharge rates for cardiovascular diseases, respiratory diseases, and injury and poisoning, but females had a higher rate than males for digestive diseases (Figure 3).
- As shown in Figure 4, the number of discharges for patients with HIV diagnoses decreased from 249,000 in 1995 to 173,000 in 2000, corresponding to the introduction of antiretroviral therapy. The number did not change significantly from 2000–2004. HIV discharges for patients aged 40 and over increased from 100,000 in 1995 (40 percent of all HIV discharges) to 136,000 (67 percent) in 2004.




Figure 1. Percent distribution of discharges and days of care by age: United States, 2004




Figure 2. Percent of discharges transferred to long-term care institutions by age: United States, 2004

- The average annual number of discharges with a first listed diagnosis of injury was 1,903,000 in 2002–04. Using the Barell injury diagnosis matrix, fracture (1,029,000) was the most common type of injury reported and lower

extremity, including the hip, was the most common site of fracture (598,000).

Procedures

- In 2004, procedures were performed on 62 percent of discharges:

46 percent had surgical procedures and another 16 percent had only one or more nonsurgical diagnostic or therapeutic procedures.

- Females had 7.0 million obstetrical procedures, including 1.2 million repairs of current obstetric laceration and 1.2 million cesarean sections.
- Almost 7 million inpatient operations were performed on the cardiovascular system in 2004, including 1.3 million cardiac catheterizations; 5.8 million operations were performed on the digestive system, including 1.1 million endoscopies of the small intestine.
- The rate of cardiac catheterizations was higher for males than for females, and higher for patients 65–74 and 75–84 years of age than for older or younger age groups. Rates were not significantly different across geographic regions (Figure 5).

Deliveries and Newborn Infants

- Approximately 4.1 million women were hospitalized for delivery in 2004, and their average stay was 2.6 days.
- Cesarean sections per 100 deliveries rose from 20.8 in 1995 to 29.9 in 2004, and the rate of primary cesareans per 100 deliveries without a previous cesarean increased from 15.5 to 20.6 during this period.
- The rate of vaginal birth after cesarean delivery (VBAC) dropped 69 percent, from 35.5 in 1995 to 10.9 in 2004.
- The average length of stay for both vaginal and cesarean deliveries decreased from 1980 to 1995, but stays for vaginal deliveries increased 24 percent during the period from 1995 to 2004 (Figure 6).
- In 2004, 43 percent of newborn infants had at least one illness or risk-related diagnosis. Perinatal jaundice (714,000), respiratory conditions (391,000), disorders related to prematurity (301,000), and congenital anomalies (275,000) were common diagnoses.




Figure 3. Rate of discharges in major diagnostic categories by sex: United States, 2004




Figure 4. Number of discharges with human immunodeficiency virus diagnoses by age: United States, 1995, 2000, and 2004

- The proportion of male newborn infants circumcised during the birth hospitalization was 57 percent nationally, but ranged from 32 percent in the West to 80 percent in the Midwest.

Introduction

This report presents data from the 2004 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the National Center for Health Statistics

(NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report provides a summary of the 2004 data, including estimates of diagnoses and procedures by the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM) codes (1).

NHDS is the principal source for national data on the characteristics of patients discharged from nonfederal short-stay hospitals. Data from the NHDS are used to evaluate the health status of the population, plan programs to improve health status, study trends in morbidity, and carry out research activities in the health field (2-6). The U.S. Department of Health and Human Services uses NHDS data to develop and monitor goals for the Healthy People 2010 Health Objectives. Other Centers for Disease Control and Prevention centers, including the National Center for Infectious Diseases, the National Center for Injury Prevention and Control, and the National Center for Chronic Disease Prevention and Health Promotion, track a variety of conditions, including human immunodeficiency virus (HIV), injuries, and chronic diseases with NHDS data. The National Institutes of Health makes extensive use of NHDS data on heart disease and cancer. The Veterans Administration and Department of Defense compare NHDS data with statistics on the inpatient care provided in their hospitals (7,8).

NHDS produces estimates of utilization only by inpatients in nonfederal short-stay hospitals. Data on ambulatory medical care services are collected in other NCHS surveys. The National Ambulatory Medical Care Survey obtains information on visits to physicians' offices (9). The National Hospital Ambulatory Medical Care Survey collects data on visits to hospital outpatient and emergency departments (10,11). Information about long-term care is collected in the National Nursing Home Survey (12) and the National Home and Hospice Care Survey (13). Data on ambulatory surgery performed in hospitals and freestanding surgery centers were collected in the National Survey of Ambulatory Surgery from




Figure 5. Rate of cardiac catheterizations by sex, age, and geographic region: United States, 2004




Figure 6. Average length of stay for women with deliveries by type of delivery: United States, 1980-2004

1994-96 and are being collected again for 2006 (14).

Methods

Data Source

NHDS data are collected from a sample of inpatient records acquired

from a national probability sample of hospitals. Because persons with multiple discharges during the year can be sampled more than once, the NHDS produces estimates for discharges, not persons. Only general hospitals, children's general hospitals, or hospitals with an average length of stay of fewer than 30 days for all patients are eligible for inclusion in the survey. Federal, military, and Department of Veterans

Affairs hospitals, as well as hospital units of institutions (such as prison hospitals), and hospitals with fewer than six beds staffed for patient use are excluded.

From 1988 until 2003, the hospital sampling frame for the NHDS was constructed from the SMG Hospital Market Database (15). Beginning in 2003, the sampling frame was constructed from products of Verispan, L.L.C., specifically their "Healthcare Market Index, updated May 15, 2003" and their "Hospital Market Profiling Solution, Second Quarter 2003." These two products were formerly known as the SMG Hospital Market Database. The sampling frame and sample are updated every 3 years to include hospitals that opened or changed their eligibility status since the previous update (16).

Since 1988, the NHDS has used a modified three-stage sample design. Units selected at the first stage consist of either hospitals or geographic areas, such as counties, groups of counties, or metropolitan statistical areas in the 50 states and the District of Columbia. Within sampled geographic areas, additional hospitals are selected. At the last stage, discharges are selected within sampled hospitals using systematic random sampling.

For 2004, the sample consisted of 501 hospitals, of which 25 were out of scope because they had gone out of business or were otherwise ineligible to be in the NHDS universe. Of the 476 in-scope (eligible) hospitals, 439 (92 percent) responded to the survey. Data were collected for approximately 371,000 discharges from the 439 responding hospitals.

Two data collection procedures are used for the survey. One is a manual system in which sample selection and transcription of information from medical records are performed by hospital staff or by staff of the U.S. Census Bureau on behalf of NCHS. Completed forms are sent to NCHS for medical coding, data entry, editing, and estimation.

For the other data collection procedure, NCHS purchases electronic files containing medical record data from commercial organizations, state data systems, hospitals, or hospital

associations. Discharges from these files are systematically sampled.

Approximately 44 percent of respondent hospitals provided data through this automated system for the 2004 survey.

The NHDS medical abstract form (Figure I) and the automated data contain items that relate to the personal characteristics of the patient. These include birth date (converted to age), sex, race, ethnicity, marital status, ZIP code, and expected sources of payment. Administrative items such as admission and discharge dates, admission type and source, and discharge status are also included. The medical information includes final medical diagnoses, procedures performed, and dates of surgery. Medical data are coded according to the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) (1). The date of birth and ZIP code are confidential items not available to the public, although they can be used, with privacy safeguards, through the NCHS Research Data Center.

For the manual data collection in the NHDS, an ongoing quality control program is undertaken for the medical coding and data entry from abstracts to electronic form. Approximately 10 percent of abstracts are independently recoded, with discrepancies resolved by a chief coder. The overall error rate for records manually coded for the 2004 survey year was 0.1 percent for demographic and medical (ICD–9–CM) coding and data entry.

Estimation

Because they are sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components:

- (1) inflation by reciprocals of the probabilities of sample selection,
- (2) adjustment for nonresponse, and
- (3) population weighting ratio adjustments.

These three components of the final weight are described in more detail in

another report (16). Information about the standard errors of statistics for the 2004 NHDS is found in Table II.

Use of Tables

Sociodemographic information about patients and utilization estimates by hospital characteristics are shown in Tables 1–8. Diagnostic data are included in Tables 9–25, and information about procedures is found in Tables 26–35. Tables 36–44 present data on women with deliveries and newborn infants. Estimates of diagnoses, days of care by diagnosis, and procedures are shown by ICD–9–CM code in Tables 45–48. Figures 1–6 illustrate highlights from the survey data. Definitions of terms used in these tables and figures are presented in the appendix.

Diagnostic and procedure data are presented by chapter of the ICD–9–CM with frequently occurring broad and specific subcategories within chapters also shown (see Tables 11–22 and 30–35). The diagnostic categories shown in Table 9 are the same as the subcategories within chapters shown in Tables 11–18. Categories shown in Table 10 are the subcategories in Tables 11–18 with estimates of 100,000 or more for patients 65 years of age and over. The leading categories shown in Figure 3 were identified by ranking the ICD–9–CM Chapters 1–17 and childbirth (code V27).

The procedure categories shown in Table 27 are the same as the subcategories in Tables 30–35, except that they are divided into surgical and nonsurgical components. The categories in Table 28 are the subcategories in Tables 30–35 for patients 65 years of age and over, divided into surgical and nonsurgical components with estimates of 100,000 or more.

In 2002, the ICD–9–CM Coordination and Maintenance Committee created Chapter 00, “Procedures Not Elsewhere Classified” to provide additional codes for procedures. Beginning with the 2003 NHDS, this chapter is included in the miscellaneous diagnostic and therapeutic procedures and new technologies category shown in procedure Tables 30–35.

Diagnoses assigned ICD–9–CM codes E800–E999 (supplementary classification of external causes of injury and poisoning) are not included in this report. However, detailed information about injury diagnoses and E-codes has been published (17).

Injury data are presented in this report using the Barell injury diagnosis matrix (Table 24). This is a two-dimensional array of ICD–9–CM codes (excluding external cause codes) that groups injuries by body site and type of injury. The codes for the matrix are shown in Table III. The matrix provides a standard format for reporting injury data that improves the usefulness and comparability of epidemiological and clinical analyses (18). Data categorized using the Barell matrix are included in this report to provide a national benchmark for States and localities.

Table 29 compares the number of coronary artery bypass grafts (CABG) with the number of inpatients who receive these grafts. More than one code is often assigned to a patient to indicate the number and type of bypasses performed during a single operation. This results in a much larger number of procedures than of inpatients who have the procedures.

Beginning with the 2003 NHDS data, estimates for the procedure of “Insertion of coronary artery stent(s)” are shown separately from estimates for “Balloon angioplasty of coronary artery or coronary atherectomy” (Tables 27, 28 and 30–35). Previously these were combined in the category “Removal of coronary artery obstruction and insertion of stent(s).”

Tables 8 and 25 show rates of death per 100 hospital discharges. These are different from mortality rates published regularly by NCHS in *National Vital Statistics Reports* (NVSR) (19), which are ratios of deaths to the population. Mortality rates published in the NVSR include deaths that occur in hospitals and other settings and are for the underlying cause of death, which may differ from the first-listed diagnosis of a patient who dies in the hospital.

Data for newborn infants are included only in Tables 41–44. Because estimates of newborn infants are based on a sample and because they do not

include out-of-hospital births, these estimates may not agree with data on births published in *National Vital Statistics Reports* (20), which are based on birth certificate data.

Tables 45–47 provide information by three-digit, four-digit, and five-digit ICD–9–CM diagnostic codes. Table 48 shows two-, three-, and four-digit procedure codes. Only ICD–9–CM code numbers appear in Tables 45–48. Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD–9–CM (1). A specific diagnosis or procedure code appears in Tables 45, 47, and 48 if the estimate of the total for the code is 5,000 or more. The codes that appear in Table 46 for days of care are identical to those in Table 45 for first-listed diagnoses.

Estimates in all tables are rounded to the nearest thousand. Totals may include data for categories that are not shown individually in the tables. For these reasons, figures within tables do not always add to the totals. Rates, percents, and average lengths of stay are calculated from unrounded figures and may not precisely agree with measures calculated from rounded data.

Because of low reliability, estimates in Tables 1–44 with relative standard errors of more than 30 percent or that are based on fewer than 30 records are not presented; only an asterisk (*) appears in the tables. Estimates based on 30–59 records are presented but are preceded by an asterisk to indicate that they also have low reliability.

Due to the large number of estimates in Tables 45–48, more generalized criteria for reliability were used. Only an asterisk is shown in Tables 45, 47, and 48 if the estimate of the diagnosis or procedure is less than 5,000, and only an asterisk is shown in Table 46 for estimates of days of care derived from fewer than 5,000 estimated discharges. These estimates generally have a relative standard error of more than 30 percent or a sample size of less than 30. Estimates of diagnoses and procedures of less than 9,000 are preceded by an asterisk because they are generally based on fewer than 60 records. Days of care estimates in table 46 that are derived from fewer

than 9,000 estimated discharges are also preceded by asterisks.

The population estimates used to compute rates are shown in appendix Table I. These are postcensal estimates as of July 1, 2004, based on the 2000 census.

A weighted least squares regression method (21) was used to test the significance of the trends shown in Figures 4 and 6. Differences mentioned among NHDS estimates were tested using the two-sided *t*-test. Terms that express differences such as higher, lower, largest, smallest, leading, increased, or decreased, were only used when the differences were statistically significant at the 0.05 level.

References

- International Classification of Diseases, 9th Revision, Clinical Modification, 6th edition. U.S. Department of Health and Human Services, National Center for Health Statistics, Health Care Financing Administration. 2003. Available from: <http://www.cdc.gov/nchs/icd9.htm#ICD-9-CM>.
- Gupta RS, Carrion-Carire V, Weiss KB. The widening black/white gap in asthma hospitalizations and mortality. *J Allergy Clin Immunol* 117:351–8. 2006.
- Tomashek KM, Crouse CJ, Iyasu S, et al. A comparison of morbidity rates attributable to conditions originating in the perinatal period among newborns discharged from United States hospitals, 1989–90 and 1999–2000. *Paediatr Perinat Epidemiol* 20:24–34. 2006.
- Fry AM, Shay DK, Holman RC, et al. Trends in hospitalizations for pneumonia among persons aged 65 years or older in the United States, 1988–2002. *JAMA* 294:2712–9. 2005.
- Shinogle JA, Owings MF, Kozak LJ. Gastric bypass as treatment for obesity: trends, characteristics, and complications. *Obes Res* 13:2202–9. 2005.
- Curns AT, Holman RC, Sejvar JJ, et al. Infectious disease hospitalizations among older adults in the United States from 1990 through 2002. *Arch Intern Med* 165:2514–20. 2005.
- Rosenthal GE, Kaboli PJ, Barnett MJ. Differences in length of stay in Veterans Health Administration and other United States hospitals: Is the gap closing? *Med Care* 41(8):882–94. 2003.
- Jackson JL, Cheng EY, Jones DL, Meyer G. Comparison of discharge diagnoses and inpatient procedures between military and civilian health care systems. *Mil Med* 164(10): 701–4. 1999.
- Hing E, Cherry DK, Woodwell DA. National Ambulatory Medical Care Survey: 2004 summary. Advance data from vital and health statistics; no 374. Hyattsville, Maryland: National Center for Health Statistics. 2006. Available from: <http://www.cdc.gov/nchs/data/ad/ad374.pdf>.
- Middleton K, Hing E. National Hospital Ambulatory Medical Care Survey: 2004 outpatient department summary. Advance data from vital and health statistics; no 373. Hyattsville, Maryland: National Center for Health Statistics. 2006. Available from: <http://www.cdc.gov/nchs/data/ad/ad373.pdf>.
- McCaig LF, Nawar EW. National Hospital Ambulatory Medical Care Survey: 2004 emergency department summary. Advance data from vital and health statistics; no 372. Hyattsville, Maryland: National Center for Health Statistics. 2006. Available from: <http://www.cdc.gov/nchs/data/ad/ad372.pdf>.
- Jones A. The National Nursing Home Survey: 1999 summary. National Center for Health Statistics. *Vital Health Stat* 13(152). 2002. Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_152.pdf.
- Haupt BJ. Characteristics of hospice care discharges and their length of service: United States, 2000. National Center for Health Statistics. *Vital Health Stat* 13(154). 2003. Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_154.pdf.
- Owings MF, Kozak LJ. Ambulatory and inpatient procedures in the United States, 1996. National Center for Health Statistics. *Vital Health Stat* 13(139). 1998. Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_139.pdf.
- SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. April 1987, April 1991, April 1994, April 1997, April 2000; Verispan, L.L.C. Healthcare Market Index, updated May 15, 2003, and Hospital Market Profiling Solution, Second Quarter, 2003.

16. Dennison CF, Pokras R. Design and operation of the National Hospital Discharge Survey: 1988 redesign. National Center for Health Statistics. Vital Health Stat 1 (39). 2000. Available from: http://www.cdc.gov/nchs/data/series/sr_01/sr01_039.pdf.
17. Heinen M, Hall MJ, Boudreault MA, Fingerhut LA. National trends in injury hospitalizations, 1979–2001. Hyattsville, MD: National Center for Health Statistics. 2005. Available from: <http://www.cdc.gov/nchs/data/injury/InjuryChartbook79-01.pdf>.
18. Barell V, Aharonson-Daniel L, Fingerhut LA, et al. An introduction to the Barell body region by nature of injury diagnosis matrix. Injury Prevention 8:91–6. 2002.
19. Sutton PD, Munson ML. Births, marriages, divorces, and deaths: Provisional data for September 2005. National vital statistics reports; 54 no 15. Hyattsville, MD: National Center for Health Statistics. 2006. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_15.pdf.
20. Hamilton BE, Martin JA, Ventura SJ, et al. Births: Preliminary data for 2004. National vital statistics reports vol 54 no 8. Hyattsville, MD: National Center for Health Statistics. 2005. Available from: http://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_08.pdf.
21. Gillum BS, Graves EJ, Kozak LJ. Trends in hospital utilization: United States, 1988–92. National Center for Health Statistics. Vital Health Stat 13(124). 1996. Available from: http://www.cdc.gov/nchs/data/series/sr_13/sr13_124.pdf.
22. Research Triangle Institute. SUDAAN Language Manual, Release 9.0. Research Triangle Park, NC: Research Triangle Institute. 2004.
23. Hansen MH, Hurwitz WN, Madow WG. Sample survey methods and theory, vol 1. New York: John Wiley & Sons. Wiley Classics Library Edition. 1993.
24. Cochran WG. Sampling techniques, 3rd ed. New York: John Wiley & Sons. 1977.

Table 1. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by age: United States, selected years, 1970–2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Age	1970	1980	1985	1990	1995	2000	2004
Number of discharges in thousands							
All ages	29,127	37,832	35,056	30,788	30,722	31,706	34,864
Under 15 years	3,873	3,672	2,972	2,412	2,405	2,383	2,571
15–44 years	12,664	15,635	13,966	11,799	10,593	9,969	10,800
45–64 years	6,693	8,660	7,610	6,244	6,168	6,958	8,323
65 years and over	5,897	9,864	10,508	10,333	11,556	12,396	13,171
65–74 years	3,163	4,943	5,011	4,689	4,832	4,678	4,786
75–84 years	2,099	3,611	3,969	3,949	4,590	5,119	5,538
85 years and over	635	1,310	1,528	1,694	2,134	2,599	2,846
Rate of discharges per 1,000 population ¹							
All ages	144.3	167.7	148.4	122.3	115.7	112.8	119.2
Under 15 years	66.8	71.6	57.7	43.1	40.4	39.5	42.3
15–44 years	154.6	150.1	125.0	99.3	87.8	80.9	86.6
45–64 years	159.6	194.8	170.8	135.5	118.5	111.4	117.8
65 years and over	293.3	383.7	369.8	334.1	347.7	353.4	362.9
65–74 years	253.2	315.8	297.3	261.6	260.0	254.6	259.2
75–84 years	339.5	463.9	446.4	395.7	415.6	412.4	427.0
85 years and over	443.7	576.4	572.9	560.6	592.5	605.1	585.7
Number of days of care in thousands							
All ages	226,445	274,508	226,217	197,422	164,627	155,857	167,881
Under 15 years	18,189	16,191	13,554	11,655	10,715	10,734	11,652
15–44 years	71,605	81,951	67,397	54,062	40,825	36,593	40,079
45–64 years	62,419	71,008	53,541	42,153	34,207	34,443	41,798
65 years and over	74,232	105,358	91,726	89,552	78,880	74,086	74,352
65–74 years	37,873	49,253	41,090	37,422	31,310	26,438	25,944
75–84 years	27,633	40,355	36,024	35,926	31,974	31,525	31,831
85 years and over	8,726	15,750	14,612	16,204	15,597	16,123	16,576
Rate of days of care per 1,000 population ¹							
All ages	1,121.6	1,216.7	957.7	784.0	620.2	554.6	574.1
Under 15 years	313.9	315.7	263.0	208.4	179.9	178.0	191.6
15–44 years	874.0	786.8	603.3	454.9	338.4	297.1	321.5
45–64 years	1,488.8	1,596.9	1,201.6	914.4	657.5	551.5	591.6
65 years and over	3,691.9	4,098.4	3,228.0	2,895.6	2,373.7	2,111.9	2,048.6
65–74 years	3,031.6	3,147.0	2,437.3	2,087.8	1,684.7	1,439.0	1,405.2
75–84 years	4,469.2	5,183.7	4,052.2	3,599.5	2,894.7	2,539.8	2,454.1
85 years and over	6,102.1	6,932.1	5,478.7	5,361.7	4,330.6	3,753.8	3,411.0
Average length of stay in days							
All ages	7.8	7.3	6.5	6.4	5.4	4.9	4.8
Under 15 years	4.7	4.4	4.6	4.8	4.5	4.5	4.5
15–44 years	5.7	5.2	4.8	4.6	3.9	3.7	3.7
45–64 years	9.3	8.2	7.0	6.8	5.5	5.0	5.0
65 years and over	12.6	10.7	8.7	8.7	6.8	6.0	5.6
65–74 years	12.0	10.0	8.2	8.0	6.5	5.7	5.4
75–84 years	13.2	11.2	9.1	9.1	7.0	6.2	5.7
85 years and over	13.7	12.0	9.6	9.6	7.3	6.2	5.8

¹Rates were calculated using the U.S. Census Bureau estimates of the civilian population. Rates for 1990 and 1995 were based on population estimates adjusted for the net underenumeration in the 1990 census. Rates for 2000 and 2004 were calculated using 2000-based postcensal estimates. Rates for 2000 differ from the ones shown in earlier reports because they were calculated using revised population estimates.

Table 2. Number, percent distribution, and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex and age: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex and age	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	
Both sexes							
All ages	34,864	100.0	119.2	167,881	100.0	574.1	4.8
Under 15 years	2,571	7.4	42.3	11,652	6.9	191.6	4.5
Under 1 year	842	2.4	206.5	5,070	3.0	1,243.5	6.0
1–4 years	734	2.1	45.9	2,266	1.3	141.7	3.1
5–14 years	995	2.9	24.4	4,316	2.6	105.9	4.3
15–44 years	10,800	31.0	86.6	40,079	23.9	321.5	3.7
15–19 years	1,175	3.4	56.9	4,465	2.7	216.3	3.8
20–24 years	1,887	5.4	91.8	6,133	3.7	298.5	3.3
25–34 years	4,046	11.6	102.2	13,572	8.1	342.7	3.4
35–44 years	3,692	10.6	84.2	15,910	9.5	362.7	4.3
45–64 years	8,323	23.9	117.8	41,798	24.9	591.6	5.0
45–54 years	4,146	11.9	99.7	20,420	12.2	491.1	4.9
55–64 years	4,176	12.0	143.6	21,377	12.7	735.2	5.1
65 years and over	13,171	37.8	362.9	74,352	44.3	2,048.6	5.6
65–74 years	4,786	13.7	259.2	25,944	15.5	1,405.2	5.4
75–84 years	5,538	15.9	427.0	31,831	19.0	2,454.1	5.7
85 years and over	2,846	8.2	585.7	16,576	9.9	3,411.0	5.8
Male							
All ages	13,844	100.0	96.5	72,449	100.0	504.9	5.2
Under 15 years	1,432	10.3	46.0	6,488	9.0	208.5	4.5
Under 1 year	490	3.5	235.1	2,879	4.0	1,380.3	5.9
1–4 years	399	2.9	48.9	1,224	1.7	149.7	3.1
5–14 years	542	3.9	26.0	2,386	3.3	114.4	4.4
15–44 years	2,827	20.4	45.0	13,793	19.0	219.7	4.9
15–19 years	331	2.4	31.3	1,689	2.3	159.9	5.1
20–24 years	345	2.5	33.0	1,653	2.3	158.2	4.8
25–34 years	784	5.7	39.3	3,803	5.2	190.4	4.9
35–44 years	1,367	9.9	62.7	6,648	9.2	304.8	4.9
45–64 years	4,073	29.4	118.4	21,075	29.1	612.4	5.2
45–54 years	2,005	14.5	98.2	10,416	14.4	510.2	5.2
55–64 years	2,068	14.9	147.7	10,659	14.7	761.5	5.2
65 years and over	5,512	39.8	363.8	31,093	42.9	2,051.8	5.6
65–74 years	2,263	16.3	268.5	12,155	16.8	1,442.3	5.4
75–84 years	2,369	17.1	454.1	13,807	19.1	2,645.8	5.8
85 years and over	880	6.4	583.8	5,131	7.1	3,402.6	5.8
Female							
All ages	21,020	100.0	141.1	95,432	100.0	640.8	4.5
Under 15 years	1,139	5.4	38.4	5,164	5.4	173.9	4.5
Under 1 year	352	1.7	176.6	2,191	2.3	1,100.1	6.2
1–4 years	335	1.6	42.8	1,042	1.1	133.3	3.1
5–14 years	453	2.2	22.8	1,931	2.0	97.1	4.3
15–44 years	7,973	37.9	128.9	26,286	27.5	425.0	3.3
15–19 years	844	4.0	83.8	2,775	2.9	275.4	3.3
20–24 years	1,541	7.3	152.7	4,480	4.7	443.6	2.9
25–34 years	3,262	15.5	166.2	9,770	10.2	497.6	3.0
35–44 years	2,325	11.1	105.5	9,262	9.7	420.1	4.0
45–64 years	4,249	20.2	117.3	20,723	21.7	571.8	4.9
45–54 years	2,141	10.2	101.2	10,004	10.5	472.8	4.7
55–64 years	2,108	10.0	139.8	10,718	11.2	710.8	5.1
65 years and over	7,658	36.4	362.3	43,259	45.3	2,046.3	5.6
65–74 years	2,523	12.0	251.4	13,789	14.4	1,374.0	5.5
75–84 years	3,169	15.1	408.7	18,025	18.9	2,325.0	5.7
85 years and over	1,966	9.4	586.5	11,446	12.0	3,414.8	5.8

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
Both sexes					
All ages:					
United States	34,864	119.2	167,881	574.1	4.8
Northeast	7,388	135.5	39,876	731.5	5.4
Midwest	7,680	117.0	33,668	512.9	4.4
South	13,239	125.8	64,896	616.7	4.9
West	6,557	97.8	29,441	439.2	4.5
Under 15 years:					
United States	2,571	42.3	11,652	191.6	4.5
Northeast	495	46.8	2,369	223.8	4.8
Midwest	*	*	*	*	3.8
South	977	44.2	4,603	208.3	4.7
West	*	*	*	*	4.6
15–44 years:					
United States	10,800	86.6	40,079	321.5	3.7
Northeast	2,049	89.9	8,979	394.1	4.4
Midwest	2,157	77.6	7,423	267.0	3.4
South	4,381	97.8	16,197	361.6	3.7
West	2,213	75.6	7,481	255.6	3.4
45–64 years:					
United States	8,323	117.8	41,798	591.6	5.0
Northeast	1,765	128.6	9,505	692.4	5.4
Midwest	1,732	107.8	7,711	479.8	4.5
South	3,350	132.6	17,406	688.8	5.2
West	1,476	94.7	7,175	460.4	4.9
65 years and over:					
United States	13,171	362.9	74,352	2,048.6	5.6
Northeast	3,079	414.9	19,023	2,563.4	6.2
Midwest	3,323	396.4	16,735	1,996.3	5.0
South	4,531	346.9	26,691	2,043.5	5.9
West	2,238	301.3	11,903	1,602.3	5.3
Male					
All ages:					
United States	13,844	96.5	72,449	504.9	5.2
Northeast	3,097	117.2	17,637	667.5	5.7
Midwest	3,058	94.9	14,227	441.4	4.7
South	5,146	100.0	27,979	543.9	5.4
West	2,543	76.2	12,605	377.6	5.0
Under 15 years:					
United States	1,432	46.0	6,488	208.5	4.5
Northeast	277	51.2	1,331	245.8	4.8
Midwest	*	*	*	*	3.8
South	536	47.5	2,539	224.7	4.7
West	*	*	*	*	4.6
15–44 years:					
United States	2,827	45.0	13,793	219.7	4.9
Northeast	671	59.0	3,652	321.0	5.4
Midwest	569	40.5	2,511	178.9	4.4
South	1,101	49.1	5,443	242.7	4.9
West	485	32.5	2,187	146.3	4.5
45–64 years:					
United States	4,073	118.4	21,075	612.4	5.2
Northeast	869	131.2	4,768	719.9	5.5
Midwest	841	106.7	3,750	475.5	4.5
South	1,627	133.0	8,800	719.2	5.4
West	736	95.9	3,757	489.8	5.1

See footnotes at end of table.

Table 3. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 1,000 population ¹	Number in thousands	Rate per 1,000 population ¹	
65 years and over:					
United States	5,512	363.8	31,093	2,051.8	5.6
Northeast	1,280	425.4	7,886	2,621.3	6.2
Midwest	1,384	400.0	6,970	2,014.8	5.0
South	1,881	343.0	11,197	2,041.7	6.0
West	968	302.3	5,040	1,574.0	5.2
Female					
All ages:					
United States	21,020	141.1	95,432	640.8	4.5
Northeast	4,290	152.7	22,240	791.7	5.2
Midwest	4,622	138.4	19,440	581.9	4.2
South	8,093	150.5	36,917	686.5	4.6
West	4,014	119.3	16,835	500.2	4.2
Under 15 years:					
United States	1,139	38.4	5,164	173.9	4.5
Northeast	218	42.1	1,038	200.8	4.8
Midwest	*	*	*	*	3.9
South	441	40.8	2,063	191.1	4.7
West	*	*	*	*	4.6
15–44 years:					
United States	7,973	128.9	26,286	425.0	3.3
Northeast	1,377	120.8	5,327	467.1	3.9
Midwest	1,589	115.4	4,912	356.7	3.1
South	3,279	146.6	10,754	480.8	3.3
West	1,727	120.7	5,294	369.8	3.1
45–64 years:					
United States	4,249	117.3	20,723	571.8	4.9
Northeast	896	126.1	4,737	666.8	5.3
Midwest	890	108.8	3,961	483.9	4.4
South	1,723	132.2	8,606	660.2	5.0
West	740	93.5	3,418	431.8	4.6
65 years and over:					
United States	7,658	362.3	43,259	2,046.3	5.6
Northeast	1,799	407.7	11,138	2,523.9	6.2
Midwest	1,939	393.8	9,765	1,983.3	5.0
South	2,649	349.6	15,494	2,044.7	5.8
West	1,271	300.6	6,863	1,623.8	5.4

* Figure does not meet standard of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
United States		Number of discharges in thousands						
All	34,864	13,751	5,734	668	5,253	6,405	1,551	900
Male	13,844	5,834	1,849	376	1,846	2,507	766	388
Female	21,020	7,917	3,885	292	3,407	3,898	785	512
Under 15 years	2,571	13	1,118	50	459	650	67	*
15-44 years	10,800	591	3,199	294	2,406	2,774	894	396
45-64 years	8,323	1,673	1,179	260	1,788	2,456	529	257
65 years and over	13,171	11,474	239	65	599	525	62	65
Northeast								
All	7,388	3,067	992	85	1,085	1,517	240	*
Male	3,097	1,283	412	58	402	626	123	*
Female	4,290	1,785	580	27	682	891	117	*
Under 15 years	495	*	135	*	102	*	20	*
15-44 years	2,049	133	534	32	453	615	133	44
45-64 years	1,765	334	276	35	377	566	73	26
65 years and over	3,079	2,597	48	*	153	163	14	*
Midwest								
All	7,680	3,518	996	102	1,083	1,537	242	93
Male	3,058	1,473	321	59	382	614	124	*
Female	4,622	2,044	676	42	701	923	118	53
Under 15 years	*	*	153	*	*	*	*	*
15-44 years	2,157	111	598	*	503	680	143	45
45-64 years	1,732	340	217	38	418	578	79	*
65 years and over	3,323	3,066	28	*	84	81	*	*6
South								
All	13,239	5,015	2,346	318	1,748	2,466	838	357
Male	5,146	2,137	676	174	586	936	403	*
Female	8,093	2,878	1,669	144	1,161	1,530	435	192
Under 15 years	977	*	473	16	129	207	27	*
15-44 years	4,381	266	1,334	143	844	1,099	479	145
45-64 years	3,350	748	456	130	579	1,001	303	87
65 years and over	4,531	3,995	82	29	196	159	30	21
West								
All	6,557	2,151	1,400	164	1,337	885	232	*
Male	2,543	941	440	85	475	331	116	*
Female	4,014	1,210	960	79	863	554	116	*
Under 15 years	*	*	*	*	*	*	8	*
15-44 years	2,213	81	732	81	606	381	139	*
45-64 years	1,476	251	229	57	414	311	74	*
65 years and over	2,238	1,816	81	*	167	121	*	*
United States		Number of days of care in thousands						
All	167,881	79,061	26,979	3,094	20,181	25,489	6,301	3,825
Male	72,449	33,705	10,912	1,959	8,079	11,168	3,347	1,764
Female	95,432	45,356	16,068	1,136	12,102	14,321	2,954	2,062
Under 15 years	11,652	51	5,659	267	1,881	2,675	248	*
15-44 years	40,079	3,751	12,236	1,272	7,655	9,508	3,117	1,534
45-64 years	41,798	10,267	7,625	1,226	7,582	10,348	2,587	1,160
65 years and over	74,352	64,992	1,458	330	3,063	2,958	349	372

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
Northeast		Number of days of care in thousands						
All	39,876	19,263	6,132	388	4,448	6,463	1,035	*
Male	17,637	8,134	2,771	275	1,874	2,859	594	*
Female	22,240	11,129	3,361	113	2,574	3,604	441	*
Under 15 years	2,369	*	797	*	351	*	*	*
15–44 years	8,979	989	2,856	129	1,527	2,282	516	185
45–64 years	9,505	2,099	2,141	158	1,717	2,428	378	112
65 years and over	19,023	16,169	338	*	853	964	67	*
Midwest								
All	33,668	17,944	4,184	456	3,841	5,547	822	437
Male	14,227	7,512	1,621	*	1,539	2,434	443	202
Female	19,440	10,432	2,563	*	2,302	3,113	379	234
Under 15 years	*	*	*	*	*	*	*	*
15–44 years	7,423	610	2,177	*	1,524	2,180	409	202
45–64 years	7,711	1,949	1,149	*	1,606	2,277	298	*
65 years and over	16,735	15,383	141	*	*	394	*	*
South								
All	64,896	30,173	10,408	1,555	7,045	9,953	3,546	1,500
Male	27,979	13,083	3,906	999	2,734	4,284	1,827	778
Female	36,917	17,091	6,502	556	4,311	5,669	1,719	722
Under 15 years	4,603	*	2,343	*	*	934	91	*
15–44 years	16,197	1,750	4,754	670	2,735	3,793	1,714	518
45–64 years	17,406	4,808	2,761	654	2,614	4,276	1,569	463
65 years and over	26,691	23,587	551	153	1,004	949	172	*
West								
All	29,441	11,681	6,255	696	4,848	3,526	899	*
Male	12,605	4,976	2,613	391	1,933	1,591	483	*
Female	16,835	6,704	3,642	305	2,915	1,935	415	*
Under 15 years	*	*	*	*	*	*	27	*
15–44 years	7,481	402	2,450	288	1,870	1,253	479	*
45–64 years	7,175	1,411	1,575	262	1,645	1,367	342	*
65 years and over	11,903	9,853	428	*	715	651	*	*
United States		Average length of stay						
All	4.8	5.7	4.7	4.6	3.8	4.0	4.1	4.3
Male	5.2	5.8	5.9	5.2	4.4	4.5	4.4	4.6
Female	4.5	5.7	4.1	3.9	3.6	3.7	3.8	4.0
Under 15 years	4.5	3.9	5.1	5.4	4.1	4.1	3.7	4.2
15–44 years	3.7	6.3	3.8	4.3	3.2	3.4	3.5	3.9
45–64 years	5.0	6.1	6.5	4.7	4.2	4.2	4.9	4.5
65 years and over	5.6	5.7	6.1	5.1	5.1	5.6	5.6	5.7
Northeast								
All	5.4	6.3	6.2	4.6	4.1	4.3	4.3	5.0
Male	5.7	6.3	6.7	4.7	4.7	4.6	4.8	5.4
Female	5.2	6.2	5.8	4.2	3.8	4.0	3.8	4.5
Under 15 years	4.8	*	5.9	6.6	3.4	4.6	3.8	5.7
15–44 years	4.4	7.4	5.3	4.0	3.4	3.7	3.9	4.2
45–64 years	5.4	6.3	7.8	4.5	4.6	4.3	5.2	4.3
65 years and over	6.2	6.2	7.1	5.2	5.6	5.9	4.7	6.5

See footnotes at end of table.

Table 4. Number of discharges from short-stay hospitals and of days of care, with average length of stay, by principal expected source of payment, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Region, sex, and age	All principal expected sources of payment ¹	Medicare	Medicaid	Worker's compensation and other government payments	HMO or PPO ²	Blue Cross Blue Shield and other private insurance	Self-pay	Other payments and no charge
Midwest				Average length of stay				
All	4.4	5.1	4.2	4.5	3.5	3.6	3.4	4.7
Male	4.7	5.1	5.1	4.9	4.0	4.0	3.6	5.0
Female	4.2	5.1	3.8	3.8	3.3	3.4	3.2	4.5
Under 15 years	3.8	*	4.7	3.8	2.9	3.5	4.7	5.0
15–44 years	3.4	5.5	3.6	4.9	3.0	3.2	2.9	4.5
45–64 years	4.5	5.7	5.3	4.0	3.8	3.9	3.8	4.9
65 years and over	5.0	5.0	5.1	*5.2	5.8	4.9	7.4	*4.6
South								
All	4.9	6.0	4.4	4.9	4.0	4.0	4.2	4.2
Male	5.4	6.1	5.8	5.8	4.7	4.6	4.5	4.7
Female	4.6	5.9	3.9	3.8	3.7	3.7	3.9	3.8
Under 15 years	4.7	*4.1	5.0	5.0	5.4	4.5	3.4	3.7
15–44 years	3.7	6.6	3.6	4.7	3.2	3.5	3.6	3.6
45–64 years	5.2	6.4	6.1	5.0	4.5	4.3	5.2	5.3
65 years and over	5.9	5.9	6.7	5.2	5.1	5.9	5.8	6.5
West								
All	4.5	5.4	4.5	4.3	3.6	4.0	3.9	3.9
Male	5.0	5.3	5.9	4.6	4.1	4.8	4.2	3.7
Female	4.2	5.5	3.8	3.9	3.4	3.5	3.6	4.0
Under 15 years	4.6	*	5.0	*	4.1	3.5	3.4	2.7
15–44 years	3.4	5.0	3.3	3.5	3.1	3.3	3.4	3.9
45–64 years	4.9	5.6	6.9	4.6	4.0	4.4	4.6	3.8
65 years and over	5.3	5.4	5.3	4.5	4.3	5.4	*	5.0

* Figure does not meet standard of reliability or precision.

¹Includes discharges for which no expected source of payment was provided.²HMO is defined as health maintenance organization and PPO is preferred provider organization.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Number of discharges in thousands						
All	34,864	8,200	8,754	6,931	6,908	4,071
Sex						
Male	13,844	3,158	3,371	2,780	2,798	1,737
Female	21,020	5,042	5,383	4,152	4,110	2,333
Age						
Under 15 years	2,571	395	620	*	632	220
15–44 years	10,800	2,222	2,863	2,070	2,318	1,328
45–64 years	8,323	1,801	1,999	1,597	1,736	1,189
65 years and over	13,171	3,783	3,271	2,560	2,222	1,334
Region						
Northeast	7,388	1,765	1,677	1,232	1,564	1,150
Midwest	7,680	3,058	1,290	*	*	813
South	13,239	2,661	3,727	2,823	2,128	1,899
West	6,557	*	2,060	1,798	1,775	*
Percent distribution of discharges						
All	100.0	23.5	25.1	19.9	19.8	11.7
Sex						
Male	100.0	22.8	24.3	20.1	20.2	12.5
Female	100.0	24.0	25.6	19.8	19.6	11.1
Age						
Under 15 years	100.0	15.3	24.1	27.4	24.6	8.6
15–44 years	100.0	20.6	26.5	19.2	21.5	12.3
45–64 years	100.0	21.6	24.0	19.2	20.9	14.3
65 years and over	100.0	28.7	24.8	19.4	16.9	10.1
Region						
Northeast	100.0	23.9	22.7	16.7	21.2	15.6
Midwest	100.0	39.8	16.8	*	*	10.6
South	100.0	20.1	28.2	21.3	16.1	14.3
West	100.0	*	31.4	27.4	27.1	*
Number of days of care in thousands						
All	167,881	34,596	42,186	33,211	35,203	22,684
Sex						
Male	72,449	13,972	17,977	14,367	15,598	10,536
Female	95,432	20,624	24,209	18,845	19,605	12,149
Age						
Under 15 years	11,652	1,166	*	*	3,130	1,141
15–44 years	40,079	7,338	10,192	7,359	9,254	5,936
45–64 years	41,798	7,931	9,653	7,854	9,609	6,751
65 years and over	74,352	18,161	19,431	14,693	13,210	8,856
Region						
Northeast	39,876	8,583	9,443	6,701	8,586	6,564
Midwest	33,668	11,747	5,837	*	*	4,263
South	64,896	11,249	18,115	13,756	11,050	10,727
West	29,441	*	8,792	7,995	8,506	*

See footnotes at end of table.

Table 5. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by bed size of hospital, according to sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All bed sizes	6–99 beds	100–199 beds	200–299 beds	300–499 beds	500 beds or more
Percent distribution of days of care						
All	100.0	20.6	25.1	19.8	21.0	13.5
Sex						
Male	100.0	19.3	24.8	19.8	21.5	14.5
Female	100.0	21.6	25.4	19.7	20.5	12.7
Age						
Under 15 years	100.0	10.0	25.0	*	26.9	9.8
15–44 years	100.0	18.3	25.4	18.4	23.1	14.8
45–64 years	100.0	19.0	23.1	18.8	23.0	16.2
65 years and over	100.0	24.4	26.1	19.8	17.8	11.9
Region						
Northeast	100.0	21.5	23.7	16.8	21.5	16.5
Midwest	100.0	34.9	17.3	*	*	12.7
South	100.0	17.3	27.9	21.2	17.0	16.5
West	100.0	*	29.9	27.2	28.9	*
Average length of stay in days						
All	4.8	4.2	4.8	4.8	5.1	5.6
Sex						
Male	5.2	4.4	5.3	5.2	5.6	6.1
Female	4.5	4.1	4.5	4.5	4.8	5.2
Age						
Under 15 years	4.5	3.0	4.7	4.7	5.0	5.2
15–44 years	3.7	3.3	3.6	3.6	4.0	4.5
45–64 years	5.0	4.4	4.8	4.9	5.5	5.7
65 years and over	5.6	4.8	5.9	5.7	5.9	6.6
Region						
Northeast	5.4	4.9	5.6	5.4	5.5	5.7
Midwest	4.4	3.8	4.5	4.4	4.9	5.2
South	4.9	4.2	4.9	4.9	5.2	5.6
West	4.5	4.2	4.3	4.4	4.8	5.4

* Figure does not meet standard of reliability or precision.

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, according to sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All	Proprietary	Government	Nonprofit
Number of discharges in thousands				
All	34,864	4,442	4,072	26,351
Sex				
Male	13,844	1,617	1,679	10,548
Female	21,020	2,825	2,393	15,802
Age				
Under 15 years	2,571	273	237	2,061
15–44 years	10,800	1,588	1,414	7,798
45–64 years	8,323	1,012	1,110	6,201
65 years and over	13,171	1,568	1,312	10,291
Region				
Northeast	7,388	*	*	6,989
Midwest	7,680	*	*	6,794
South	13,239	3,327	2,306	7,606
West	6,557	*	*	4,962
Percent distribution of discharges				
All	100.0	12.7	11.7	75.6
Sex				
Male	100.0	11.7	12.1	76.2
Female	100.0	13.4	11.4	75.2
Age				
Under 15 years	100.0	10.6	9.2	80.2
15–44 years	100.0	14.7	13.1	72.2
45–64 years	100.0	12.2	13.3	74.5
65 years and over	100.0	11.9	10.0	78.1
Region				
Northeast	100.0	*	*	94.6
Midwest	100.0	*	*	88.5
South	100.0	25.1	17.4	57.5
West	100.0	*	*	75.7
Number of days of care in thousands				
All	167,881	21,978	19,851	126,052
Sex				
Male	72,449	9,078	8,890	54,481
Female	95,432	12,900	10,961	71,571
Age				
Under 15 years	11,652	1,100	947	9,605
15–44 years	40,079	5,740	5,778	28,561
45–64 years	41,798	5,505	6,098	30,195
65 years and over	74,352	9,633	7,027	57,691
Region				
Northeast	39,876	*	*	37,556
Midwest	33,668	*	*	29,499
South	64,896	17,017	10,979	36,900
West	29,441	*	*	22,097

See footnotes at end of table.

Table 6. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by hospital ownership, according to sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All	Proprietary	Government	Nonprofit
		Percent distribution of days of care		
All	100.0	13.1	11.8	75.1
Sex				
Male	100.0	12.5	12.3	75.2
Female	100.0	13.5	11.5	75.0
Age				
Under 15 years	100.0	9.4	8.1	82.4
15–44 years	100.0	14.3	14.4	71.3
45–64 years	100.0	13.2	14.6	72.2
65 years and over	100.0	13.0	9.5	77.6
Region				
Northeast	100.0	*	*	94.2
Midwest	100.0	*	*	87.6
South	100.0	26.2	16.9	56.9
West	100.0	*	*	75.1
		Average length of stay in days		
All	4.8	4.9	4.9	4.8
Sex				
Male	5.2	5.6	5.3	5.2
Female	4.5	4.6	4.6	4.5
Age				
Under 15 years	4.5	4.0	4.0	4.7
15–44 years	3.7	3.6	4.1	3.7
45–64 years	5.0	5.4	5.5	4.9
65 years and over	5.6	6.1	5.4	5.6
Region				
Northeast	5.4	4.4	6.1	5.4
Midwest	4.4	5.0	4.7	4.3
South	4.9	5.1	4.8	4.9
West	4.5	4.4	4.9	4.5

* Figure does not meet standard of reliability or precision.

Table 7. Number and percent distribution of discharges by type and source of admission to short-stay hospitals, according to sex and age: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Admission type and source	All ages	Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over
Number of discharges in thousands							
All types and sources	34,864	13,844	21,020	2,571	10,800	8,323	13,171
Type of admission							
Emergency	13,398	6,135	7,263	931	3,096	3,449	5,921
Urgent	8,954	3,216	5,738	780	3,189	1,795	3,190
Elective	8,379	2,882	5,497	530	3,084	2,080	2,685
Item not available	4,133	1,611	2,522	330	1,430	999	1,374
Source of admission							
Physician referral ¹	13,639	4,479	9,160	906	5,458	3,036	4,238
Emergency department	14,195	6,455	7,739	927	3,194	3,692	6,382
Transfer from a hospital	1,115	557	558	154	197	276	488
Transfer from other health facility ²	548	248	300	16	62	107	362
Other sources ³	698	269	429	*	338	162	135
Item not available	4,669	1,836	2,834	504	1,550	1,050	1,565
Percent distribution							
All types and sources	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Type of admission							
Emergency	38.4	44.3	34.6	36.2	28.7	41.4	45.0
Urgent	25.7	23.2	27.3	30.3	29.5	21.6	24.2
Elective	24.0	20.8	26.2	20.6	28.6	25.0	20.4
Item not available	11.9	11.6	12.0	12.8	13.2	12.0	10.4
Source of admission							
Physician referral ¹	39.1	32.4	43.6	35.2	50.5	36.5	32.2
Emergency department	40.7	46.6	36.8	36.1	29.6	44.4	48.5
Transfer from a hospital	3.2	4.0	2.7	6.0	1.8	3.3	3.7
Transfer from other health facility ²	1.6	1.8	1.4	0.6	0.6	1.3	2.8
Other sources ³	2.0	1.9	2.0	2.5	3.1	1.9	1.0
Item not available	13.4	13.3	13.5	19.6	14.4	12.6	11.9

* Figure does not meet standard of reliability or precision.

¹Referral from personal physician, clinic physician, or health maintenance organization physician.²Transfer from skilled nursing facility or other health facility.³Includes admissions directed by a court of law or requested by a law enforcement agent and other sources.

Table 8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All dispositions	Discharged alive					Died	Status not stated
		Routine or discharged home	Transferred to					
			Other short-term hospital	Long-term care institution	Other			
Number of discharges in thousands								
All	34,864	26,715	1,555	3,104	2,288	802	401	
Sex								
Male	13,844	10,379	719	1,162	1,001	408	175	
Female	21,020	16,336	836	1,942	1,287	394	226	
Age								
Under 65 years	21,694	19,013	705	518	1,024	212	223	
Under 15 years	2,571	2,409	48	12	72	16	14	
15–44 years	10,800	9,873	239	111	436	40	101	
45–64 years	8,323	6,731	418	395	516	156	107	
65 years and over	13,171	7,702	850	2,586	1,264	590	179	
65–74 years	4,786	3,348	297	528	395	148	69	
75–84 years	5,538	3,179	372	1,130	532	249	77	
85 years and over	2,846	1,175	181	928	338	192	33	
Region								
Northeast	7,388	5,318	299	866	656	174	*	
Midwest	7,680	5,715	318	830	553	203	*	
South	13,239	10,360	603	973	783	278	241	
West	6,557	5,322	334	435	296	147	24	
Percent distribution of discharges								
All	100.0	76.6	4.5	8.9	6.6	2.3	1.2	
Sex								
Male	100.0	75.0	5.2	8.4	7.2	2.9	1.3	
Female	100.0	77.7	4.0	9.2	6.1	1.9	1.1	
Age								
Under 65 years	100.0	87.6	3.2	2.4	4.7	1.0	1.0	
Under 15 years	100.0	93.7	1.9	0.5	2.8	0.6	0.5	
15–44 years	100.0	91.4	2.2	1.0	4.0	0.4	0.9	
45–64 years	100.0	80.9	5.0	4.7	6.2	1.9	1.3	
65 years and over	100.0	58.5	6.5	19.6	9.6	4.5	1.4	
65–74 years	100.0	70.0	6.2	11.0	8.3	3.1	1.4	
75–84 years	100.0	57.4	6.7	20.4	9.6	4.5	1.4	
85 years and over	100.0	41.3	6.4	32.6	11.9	6.8	1.1	
Region								
Northeast	100.0	72.0	4.1	11.7	8.9	2.4	*	
Midwest	100.0	74.4	4.1	10.8	7.2	2.6	*	
South	100.0	78.3	4.6	7.4	5.9	2.1	1.8	
West	100.0	81.2	5.1	6.6	4.5	2.2	*	
Number of days of care in thousands								
All	167,881	109,294	9,195	25,321	14,516	6,707	2,847	
Sex								
Male	72,449	46,156	4,465	10,454	6,663	3,383	1,328	
Female	95,432	63,137	4,731	14,867	7,854	3,324	1,519	
Age								
Under 65 years	93,529	73,833	4,144	5,475	6,411	2,179	1,488	
Under 15 years	11,652	10,217	335	*	*	*	*	
15–44 years	40,079	34,089	1,370	1,273	2,303	428	617	
45–64 years	41,798	29,528	2,439	4,028	3,560	1,490	753	
65 years and over	74,352	35,461	5,052	19,846	8,105	4,529	1,360	
65–74 years	25,944	15,083	1,710	4,801	2,563	1,279	509	
75–84 years	31,831	14,711	2,219	8,702	3,483	2,116	602	
85 years and over	16,576	5,667	1,123	6,344	2,060	1,135	249	

See footnotes at end of table.

Table 8. Number and percent distribution of discharges from short-stay hospitals and of days of care, with average length of stay, by disposition of patient, according to sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Sex, age, and region	All dispositions	Discharged alive					Died	Status not stated
		Routine or discharged home	Transferred to					
			Other short-term hospital	Long-term care institution	Other			
Region		Number of days of care in thousands						
Northeast	39,876	24,120	1,792	7,399	4,276	1,756	*	
Midwest	33,668	21,165	1,349	5,885	3,558	1,194	*	
South	64,896	43,501	3,817	8,459	4,958	2,503	1,659	
West	29,441	20,508	2,238	3,577	1,725	1,254	*	
All		100.0	65.1	5.5	15.1	8.6	4.0	1.7
Sex		Percent distribution of days of care						
Male	100.0	63.7	6.2	14.4	9.2	4.7	1.8	
Female	100.0	66.2	5.0	15.6	8.2	3.5	1.6	
Age		Average length of stay						
Under 65 years	100.0	78.9	4.4	5.9	6.9	2.3	1.6	
Under 15 years	100.0	87.7	2.9	*	4.7	2.2	*	
15–44 years	100.0	85.1	3.4	3.2	5.7	1.1	1.5	
45–64 years	100.0	70.6	5.8	9.6	8.5	3.6	1.8	
65 years and over	100.0	47.7	6.8	26.7	10.9	6.1	1.8	
65–74 years	100.0	58.1	6.6	18.5	9.9	4.9	2.0	
75–84 years	100.0	46.2	7.0	27.3	10.9	6.6	1.9	
85 years and over	100.0	34.2	6.8	38.3	12.4	6.8	1.5	
Region		Average length of stay						
Northeast	100.0	60.5	4.5	18.6	10.7	4.4	*	
Midwest	100.0	62.9	4.0	17.5	10.6	3.5	*	
South	100.0	67.0	5.9	13.0	7.6	3.9	2.6	
West	100.0	69.7	7.6	12.2	5.9	4.3	*	
All		4.8	4.1	5.9	8.2	6.3	8.4	7.1
Sex		Average length of stay						
Male	5.2	4.4	6.2	9.0	6.7	8.3	7.6	
Female	4.5	3.9	5.7	7.7	6.1	8.4	6.7	
Age		Average length of stay						
Under 65 years	4.3	3.9	5.9	10.6	6.3	10.3	6.7	
Under 15 years	4.5	4.2	7.0	14.5	7.6	15.9	*	
15–44 years	3.7	3.5	5.7	11.5	5.3	10.8	6.1	
45–64 years	5.0	4.4	5.8	10.2	6.9	9.5	7.0	
65 years and over	5.6	4.6	5.9	7.7	6.4	7.7	7.6	
65–74 years	5.4	4.5	5.8	9.1	6.5	8.6	7.4	
75–84 years	5.7	4.6	6.0	7.7	6.6	8.5	7.8	
85 years and over	5.8	4.8	6.2	6.8	6.1	5.9	7.6	
Region		Average length of stay						
Northeast	5.4	4.5	6.0	8.5	6.5	10.1	7.1	
Midwest	4.4	3.7	4.2	7.1	6.4	5.9	8.4	
South	4.9	4.2	6.3	8.7	6.3	9.0	6.9	
West	4.5	3.9	6.7	8.2	5.8	8.5	5.9	

* Figure does not meet standard of reliability or precision.

Table 9. Number and rate of discharges from short-stay hospitals and of days of care, with average length of stay, by selected first-listed diagnostic categories: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM)]

Category of first-listed diagnosis and ICD–9–CM code	Discharges		Days of care		Average length of stay in days
	Number in thousands	Rate per 10,000 population ¹	Number in thousands	Rate per 10,000 population ¹	
All conditions ²	34,864	1,192.3	167,881	5,741.2	4.8
Heart disease	4,375	149.6	20,220	691.5	4.6
Acute myocardial infarction	732	25.0	4,218	144.3	5.8
Coronary atherosclerosis	1,065	36.4	3,743	128.0	3.5
Other ischemic heart disease	184	6.3	491	16.8	2.7
Cardiac dysrhythmias	762	26.1	2,822	96.5	3.7
Congestive heart failure	1,092	37.3	5,840	199.7	5.4
Females with deliveries	4,136	141.4	10,791	369.0	2.6
Psychoses	1,601	54.8	12,643	432.4	7.9
Schizophrenic disorders	331	11.3	3,557	121.7	10.8
Major depressive disorder	463	15.8	3,148	107.6	6.8
Pneumonia	1,329	45.5	7,339	251.0	5.5
Malignant neoplasms	1,205	41.2	7,927	271.1	6.6
Malignant neoplasm of large intestine and rectum	161	5.5	1,385	47.4	8.6
Malignant neoplasm of trachea, bronchus, and lung	164	5.6	1,135	38.8	6.9
Fractures, all sites ³	1,039	35.5	5,814	198.8	5.6
Fracture of neck of femur ³	329	11.3	2,107	72.1	6.4
Cerebrovascular disease	906	31.0	4,719	161.4	5.2
Certain complications of surgical and medical care	871	29.8	5,381	184.0	6.2
Osteoarthritis and allied disorders	696	23.8	2,697	92.2	3.9
Diabetes mellitus	599	20.5	2,825	96.6	4.7
Cellulitis and abscess	562	19.2	2,685	91.8	4.8
Volume depletion	518	17.7	1,699	58.1	3.3
Asthma	497	17.0	1,599	54.7	3.2
Chronic bronchitis	486	16.6	2,413	82.5	5.0
Urinary tract infection	429	14.7	2,081	71.2	4.9
Septicemia	410	14.0	3,438	117.6	8.4
Benign neoplasms	380	13.0	1,228	42.0	3.2
Benign neoplasm of uterus	216	7.4	536	18.3	2.5
Intervertebral disc disorder	369	12.6	1,022	35.0	2.8
Cholelithiasis	354	12.1	1,426	48.8	4.0
Essential hypertension	341	11.7	787	26.9	2.3
Anemias	316	10.8	1,542	52.7	4.9
Noninfectious enteritis and colitis	315	10.8	1,481	50.6	4.7
Appendicitis	313	10.7	982	33.6	3.1
Diverticula of intestine	312	10.7	1,567	53.6	5.0
Intestinal obstruction	305	10.4	1,881	64.3	6.2
Acute bronchitis and bronchiolitis	265	9.1	868	29.7	3.3
Poisonings	256	8.8	794	27.1	3.1
Acute pancreatitis	246	8.4	1,399	47.8	5.7
Calculus of kidney and ureter	171	5.9	403	13.8	2.4

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²Includes data for diagnostic conditions not shown in table.³Excludes fractures coded as 733.1, pathologic fracture.

Table 10. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2004[Discharges of inpatients from nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM)]

Category of first-listed diagnosis and ICD–9–CM code	65 years and over	65–74 years	75–84 years	85 years and over
		Discharges in thousands		
All conditions ¹	13,171	4,786	5,538	2,846
Heart disease 391–392.0,393–398,402,404,410–416,420–429	2,787	975	1,209	603
Acute myocardial infarction 410	460	149	185	125
Coronary atherosclerosis 414.0	576	288	237	51
Cardiac dysrhythmias 427	526	175	234	118
Congestive heart failure 428.0,428.2–428.4	816	220	367	229
Pneumonia 480–486	800	228	346	226
Cerebrovascular disease 430–438	637	197	277	163
Malignant neoplasms 140–208,230–234	625	304	244	77
Malignant neoplasm of large intestine and rectum 153–154,197.5	103	45	40	17
Fractures, all sites ² 800–829	534	123	213	197
Fracture of neck of femur ² 820	289	48	116	125
Osteoarthritis and allied disorders 715	427	225	175	27
Certain complications of surgical and medical care 996–999	327	155	127	44
Chronic bronchitis 491	323	139	133	50
Urinary tract infection 599.0	302	69	124	109
Septicemia 038	285	81	124	80
Volume depletion 276.5	245	56	110	79
Psychoses 290–299	220	91	82	47
Diabetes mellitus 250	211	97	88	26
Diverticula of intestine 562	177	67	74	36
Intestinal obstruction 560	174	60	79	36
Cellulitis and abscess 681–682	174	64	74	35
Essential hypertension 401	152	61	63	28
Anemias 280–285	136	48	55	33
Cholelithiasis 574	127	57	50	20
Asthma 493	104	51	35	18
		Discharge rate per 10,000 population ³		
All conditions ¹	3,628.9	2,592.3	4,269.7	5,856.7
Heart disease 391–392.0,393–398,402,404,410–416,420–429	767.9	528.2	931.8	1,241.6
Acute myocardial infarction 410	126.6	80.9	143.0	256.5
Coronary atherosclerosis 414.0	158.6	155.7	182.4	105.9
Cardiac dysrhythmias 427	145.0	94.7	180.1	242.6
Congestive heart failure 428.0,428.2–428.4	225.0	119.4	282.7	472.0
Pneumonia 480–486	220.4	123.4	267.0	464.9
Cerebrovascular disease 430–438	175.6	106.9	213.5	335.4
Malignant neoplasms 140–208,230–234	172.2	164.7	187.8	159.0
Malignant neoplasm of large intestine and rectum 153–154,197.5	28.3	24.6	31.2	34.6
Fractures, all sites ² 800–829	147.0	66.8	164.1	406.3
Fracture of neck of femur ² 820	79.6	26.1	89.1	257.4
Osteoarthritis and allied disorders 715	117.7	121.8	135.1	55.8
Certain complications of surgical and medical care 996–999	90.1	84.1	98.3	90.7
Chronic bronchitis 491	88.9	75.4	102.8	103.1
Urinary tract infection 599.0	83.2	37.5	95.7	223.6
Septicemia 038	78.5	43.8	95.9	163.9
Volume depletion 276.5	67.6	30.5	84.7	162.9
Psychoses 290–299	60.7	49.5	63.1	96.9
Diabetes mellitus 250	58.2	52.6	67.6	54.5
Diverticula of intestine 562	48.8	36.4	56.7	74.8
Intestinal obstruction 560	48.0	32.4	60.7	73.6
Cellulitis and abscess 681–682	47.9	34.8	57.1	72.9
Essential hypertension 401	41.9	33.0	48.4	58.5
Anemias 280–285	37.5	26.0	42.5	67.6
Cholelithiasis 574	35.0	31.1	38.4	40.9
Asthma 493	28.7	27.6	27.2	36.6

See footnotes at end of table.

Table 10. Number and rate of discharges from short-stay hospitals, with average length of stay, for patients 65 years of age and over, by age and selected first-listed diagnostic categories: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)*]

Category of first-listed diagnosis and ICD–9–CM code	65 years and over	65–74 years	75–84 years	85 years and over
	Average length of stay in days			
All conditions ¹	5.6	5.4	5.7	5.8
Heart disease 391–392.0,393–398,402,404,410–416,420–429	4.9	4.5	5.0	5.3
Acute myocardial infarction 410	6.2	5.4	6.7	6.3
Coronary atherosclerosis 414.0	3.8	3.7	4.1	3.2
Cardiac dysrhythmias 427	3.9	3.4	3.9	4.5
Congestive heart failure 428.0,428.2–428.4	5.4	5.4	5.3	5.4
Pneumonia 480–486	6.1	6.0	6.2	6.2
Cerebrovascular disease 430–438	5.1	4.7	5.2	5.3
Malignant neoplasms 140–208,230–234	6.9	7.0	6.8	6.5
Malignant neoplasm of large intestine and rectum 153–154,197.5	9.4	9.2	9.4	10.0
Fractures, all sites ² 800–829	6.1	6.2	6.2	5.9
Fracture of neck of femur ² 820	6.5	6.8	6.5	6.3
Osteoarthritis and allied disorders 715	4.0	3.9	4.1	4.0
Certain complications of surgical and medical care 996–999	6.6	5.9	7.5	6.3
Chronic bronchitis 491	5.1	5.0	5.0	5.9
Urinary tract infection 599.0	5.2	4.5	5.1	5.7
Septicemia 038	8.1	8.9	8.1	7.5
Volume depletion 276.5	4.0	3.8	4.1	4.0
Psychoses 290–299	9.5	10.9	8.6	8.2
Diabetes mellitus 250	5.3	5.5	5.1	5.5
Diverticula of intestine 562	5.4	5.5	5.5	4.8
Intestinal obstruction 560	6.2	6.2	6.4	5.8
Cellulitis and abscess 681–682	5.9	5.6	6.3	5.6
Essential hypertension 401	2.4	2.0	2.6	2.8
Anemias 280–285	3.9	3.7	4.0	4.1
Cholelithiasis 574	5.1	4.4	5.4	6.2
Asthma 493	4.8	4.8	4.9	4.8

¹Includes data for diagnostic conditions not shown in table.²Excludes fractures coded as 733.1, pathologic fracture.³Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.

Table 11. Number of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Number in thousands				
All conditions	34,864	2,571	10,800	8,323	13,171
Infectious and parasitic diseases 001-139	949	155	172	212	409
Septicemia 038	410	7	28	90	285
Neoplasms 140-239	1,640	38	286	606	709
Malignant neoplasms 140-208,230-234	1,205	28	125	428	625
Malignant neoplasm of large intestine and rectum 153-154,197.5	161	*	10	48	103
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	*	12	54	97
Benign neoplasms 210-229	380	8	150	164	58
Benign neoplasm of uterus 218-219	216	*	109	104	*
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,753	189	386	484	693
Diabetes mellitus 250	599	31	148	208	211
Volume depletion 276.5	518	127	64	82	245
Diseases of the blood and blood-forming organs 280-289	449	58	109	98	184
Anemias 280-285	316	27	89	63	136
Mental disorders 290-319	2,311	127	1,253	662	268
Psychoses 290-299	1,601	*	845	463	220
Schizophrenic disorders 295	331	*	177	123	28
Major depressive disorder 296.2-296.3	463	*	268	123	48
Diseases of the nervous system and sense organs 320-389	553	80	133	132	208
Diseases of the circulatory system 390-459	6,363	31	448	1,881	4,003
Essential hypertension 401	341	*	49	140	152
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,375	19	265	1,304	2,787
Acute myocardial infarction 410	732	*	42	230	460
Coronary atherosclerosis 414.0	1,065	*	48	440	576
Other ischemic heart disease 411-413,414.1-414.9	184	*	15	76	92
Cardiac dysrhythmias 427	762	5	56	175	526
Congestive heart failure 428.0,428.2-428.4	1,092	*3	38	233	816
Cerebrovascular disease 430-438	906	*	48	218	637
Diseases of the respiratory system 460-519	3,550	687	346	736	1,780
Acute bronchitis and bronchiolitis 466	265	200	16	15	34
Pneumonia 480-486	1,329	182	104	244	800
Chronic bronchitis 491	486	*	11	150	323
Asthma 493	497	190	91	112	104
Diseases of the digestive system 520-579	3,580	248	859	1,070	1,404
Appendicitis 540-543	313	82	157	54	21
Noninfectious enteritis and colitis 555-558	315	56	94	70	96
Intestinal obstruction 560	305	14	32	85	174
Diverticula of intestine 562	312	*	42	93	177
Cholelithiasis 574	354	*2	122	103	127
Acute pancreatitis 577.0	246	*4	76	94	72
Diseases of the genitourinary system 580-629	1,941	84	580	496	782
Calculus of kidney and ureter 592	171	*	73	68	29
Urinary tract infection 599.0	429	26	35	66	302
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	523	*	520	*	...
Diseases of the skin and subcutaneous tissue 680-709	764	*	214	210	234
Cellulitis and abscess 681-682	562	54	165	170	174
Diseases of the musculoskeletal system and connective tissue 710-739	1,920	38	311	681	889
Osteoarthritis and allied disorders 715	696	*	23	246	427
Intervertebral disc disorders 722	369	*	130	153	85
Congenital anomalies 740-759	206	143	32	22	9
Certain conditions originating in the perinatal period 760-779	203	203	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	248	59	79	64	47
Injury and poisoning 800-999	2,845	240	828	709	1,069
Fractures, all sites ² 800-829	1,039	73	241	191	534
Fracture of neck of femur ² 820	329	*	*7	31	289
Poisonings 960-989	256	19	150	62	25
Certain complications of surgical and medical care 996-999	871	45	181	318	327
Supplementary classifications V01-V83	5,068	82	4,244	259	483
Females with deliveries V27	4,136	*5	4,120	10	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 12. Rate of discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	1,192.3	422.7	866.5	1,177.9	3,628.9
Infectious and parasitic diseases 001-139	32.4	25.6	13.8	30.1	112.6
Septicemia 038	14.0	1.2	2.2	12.7	78.5
Neoplasms 140-239	56.1	6.2	23.0	85.8	195.5
Malignant neoplasms 140-208,230-234	41.2	4.6	10.0	60.5	172.2
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.5	*	0.8	6.9	28.3
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.6	*	1.0	7.6	26.8
Benign neoplasms 210-229	13.0	1.3	12.1	23.2	15.9
Benign neoplasm of uterus 218-219	7.4	*	8.7	14.7	*
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	59.9	31.1	31.0	68.5	190.9
Diabetes mellitus 250	20.5	5.0	11.9	29.5	58.2
Volume depletion 276.5	17.7	20.8	5.1	11.6	67.6
Diseases of the blood and blood-forming organs 280-289	15.3	9.6	8.7	13.9	50.6
Anemias 280-285	10.8	4.5	7.2	8.9	37.5
Mental disorders 290-319	79.0	21.0	100.6	93.7	73.9
Psychoses 290-299	54.8	*	67.8	65.5	60.7
Schizophrenic disorders 295	11.3	*	14.2	17.5	7.6
Major depressive disorder 296.2-296.3	15.8	*	21.5	17.5	13.3
Diseases of the nervous system and sense organs 320-389	18.9	13.2	10.7	18.7	57.3
Diseases of the circulatory system 390-459	217.6	5.1	35.9	266.3	1,103.0
Essential hypertension 401	11.7	*	3.9	19.8	41.9
Heart disease 391-392.0,393-398,402,404,410-416,420-429	149.6	3.1	21.3	184.6	767.9
Acute myocardial infarction 410	25.0	*	3.4	32.5	126.6
Coronary atherosclerosis 414.0	36.4	*	3.9	62.3	158.6
Other ischemic heart disease 411-413,414.1-414.9	6.3	*	1.2	10.8	25.2
Cardiac dysrhythmias 427	26.1	0.8	4.5	24.8	145.0
Congestive heart failure 428.0,428.2-428.4	37.3	*0.6	3.1	33.0	225.0
Cerebrovascular disease 430-438	31.0	*	3.8	30.8	175.6
Diseases of the respiratory system 460-519	121.4	113.0	27.7	104.2	490.6
Acute bronchitis and bronchiolitis 466	9.1	32.8	1.3	2.2	9.4
Pneumonia 480-486	45.5	29.8	8.3	34.5	220.4
Chronic bronchitis 491	16.6	*	0.9	21.2	88.9
Asthma 493	17.0	31.2	7.3	15.9	28.7
Diseases of the digestive system 520-579	122.4	40.8	68.9	151.4	386.9
Appendicitis 540-543	10.7	13.5	12.6	7.7	5.7
Noninfectious enteritis and colitis 555-558	10.8	9.2	7.5	9.9	26.4
Intestinal obstruction 560	10.4	2.3	2.6	12.0	48.0
Diverticula of intestine 562	10.7	*	3.3	13.1	48.8
Cholelithiasis 574	12.1	*0.4	9.8	14.6	35.0
Acute pancreatitis 577.0	8.4	*0.7	6.1	13.3	19.7
Diseases of the genitourinary system 580-629	66.4	13.8	46.5	70.2	215.4
Calculus of kidney and ureter 592	5.9	*	5.8	9.6	7.9
Urinary tract infection 599.0	14.7	4.3	2.8	9.3	83.2
Complications of pregnancy, childbirth, and the puerperium ² 630-677	17.9	*	41.7	*	...
Diseases of the skin and subcutaneous tissue 680-709	26.1	*	17.2	29.8	64.4
Cellulitis and abscess 681-682	19.2	8.8	13.2	24.1	47.9
Diseases of the musculoskeletal system and connective tissue 710-739	65.7	6.3	25.0	96.4	245.1
Osteoarthritis and allied disorders 715	23.8	*	1.8	34.8	117.7
Intervertebral disc disorders 722	12.6	*	10.4	21.7	23.4
Congenital anomalies 740-759	7.0	23.4	2.6	3.1	2.5
Certain conditions originating in the perinatal period 760-779	7.0	33.3	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	8.5	9.7	6.3	9.0	12.8
Injury and poisoning 800-999	97.3	39.5	66.4	100.3	294.4
Fractures, all sites ³ 800-829	35.5	12.0	19.3	27.0	147.0
Fracture of neck of femur ³ 820	11.3	*	*0.6	4.4	79.6
Poisonings 960-989	8.8	3.2	12.0	8.8	6.9
Certain complications of surgical and medical care 996-999	29.8	7.5	14.5	45.0	90.1
Supplementary classifications V01-V83	173.3	13.4	340.5	36.6	133.1
Females with deliveries V27	141.4	*0.9	330.5	1.5	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."³Excludes fractures coded as 733.1, pathologic fracture.

Table 13. Average length of stay for discharges from short-stay hospitals, by age and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
	Average length of stay in days				
All conditions	4.8	4.5	3.7	5.0	5.6
Infectious and parasitic diseases 001-139	6.6	2.9	6.0	7.8	7.6
Septicemia 038	8.4	5.1	9.5	9.1	8.1
Neoplasms 140-239	5.7	7.3	4.1	5.4	6.6
Malignant neoplasms 140-208,230-234	6.6	8.7	5.7	6.3	6.9
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	*	6.2	7.3	9.4
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.9	*	7.8	6.9	6.9
Benign neoplasms 210-229	3.2	3.3	2.7	3.1	5.0
Benign neoplasm of uterus 218-219	2.5	*	2.5	2.4	*3.0
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.0	3.0	3.4	4.0	4.5
Diabetes mellitus 250	4.7	2.6	4.0	4.9	5.3
Volume depletion 276.5	3.3	2.1	2.9	3.2	4.0
Diseases of the blood and blood-forming organs 280-289	4.7	4.0	6.0	4.8	4.1
Anemias 280-285	4.9	4.3	6.5	4.9	3.9
Mental disorders 290-319	7.1	10.3	6.4	7.2	8.7
Psychoses 290-299	7.9	8.7	7.1	8.4	9.5
Schizophrenic disorders 295	10.8	*	10.2	11.3	12.2
Major depressive disorder 296.2-296.3	6.8	5.8	5.8	7.5	11.1
Diseases of the nervous system and sense organs 320-389	5.3	4.6	4.2	4.8	6.5
Diseases of the circulatory system 390-459	4.7	6.9	4.1	4.3	4.9
Essential hypertension 401	2.3	*	2.3	2.2	2.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.6	8.1	4.0	4.1	4.9
Acute myocardial infarction 410	5.8	*	3.4	5.4	6.2
Coronary atherosclerosis 414.0	3.5	*	2.8	3.2	3.8
Other ischemic heart disease 411-413,414.1-414.9	2.7	*	2.1	2.2	2.7
Cardiac dysrhythmias 427	3.7	4.9	*	3.1	3.9
Congestive heart failure 428.0,428.2-428.4	5.4	*11.0	4.5	5.3	5.4
Cerebrovascular disease 430-438	5.2	*8.5	5.5	5.5	5.1
Diseases of the respiratory system 460-519	5.2	2.9	4.5	5.4	6.1
Acute bronchitis and bronchiolitis 466	3.3	3.1	3.0	4.6	3.8
Pneumonia 480-486	5.5	3.2	4.7	5.6	6.1
Chronic bronchitis 491	5.0	*	5.5	4.6	5.1
Asthma 493	3.2	2.2	2.9	3.7	4.8
Diseases of the digestive system 520-579	4.7	3.6	3.8	4.8	5.4
Appendicitis 540-543	3.1	3.7	2.4	3.7	4.8
Noninfectious enteritis and colitis 555-558	4.7	2.5	3.9	6.0	5.9
Intestinal obstruction 560	6.2	5.8	5.2	6.5	6.2
Diverticula of intestine 562	5.0	*	4.7	4.5	5.4
Cholelithiasis 574	4.0	*3.2	3.2	3.7	5.1
Acute pancreatitis 577.0	5.7	*6.0	5.6	5.3	6.3
Diseases of the genitourinary system 580-629	3.8	3.1	2.8	3.5	4.9
Calculus of kidney and ureter 592	2.4	*	2.2	2.2	3.1
Urinary tract infection 599.0	4.9	3.1	4.2	4.4	5.2
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.7	*	2.7	*	...
Diseases of the skin and subcutaneous tissue 680-709	5.1	2.6	4.3	5.5	6.5
Cellulitis and abscess 681-682	4.8	2.9	4.0	4.9	5.9
Diseases of the musculoskeletal system and connective tissue 710-739	3.9	4.3	3.5	3.5	4.2
Osteoarthritis and allied disorders 715	3.9	*	3.6	3.6	4.0
Intervertebral disc disorders 722	2.8	*	2.4	2.6	3.7
Congenital anomalies 740-759	6.8	7.2	3.5	*	5.5
Certain conditions originating in the perinatal period 760-779	10.7	10.7	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	2.7	2.0	2.3	2.5	4.4
Injury and poisoning 800-999	5.5	3.6	4.6	5.9	6.2
Fractures, all sites ² 800-829	5.6	2.4	5.1	6.1	6.1
Fracture of neck of femur ² 820	6.4	*	*5.8	6.4	6.5
Poisonings 960-989	3.1	1.8	2.7	3.3	*
Certain complications of surgical and medical care 996-999	6.2	5.8	5.0	6.5	6.6
Supplementary classifications V01-V83	3.7	4.9	2.7	7.4	10.2
Females with deliveries V27	2.6	*2.7	2.6	3.3	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 14. Number of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
All conditions	34,864	13,844	21,020
Infectious and parasitic diseases 001-139	949	441	507
Septicemia 038	410	188	222
Neoplasms 140-239	1,640	655	985
Malignant neoplasms 140-208,230-234	1,205	576	629
Malignant neoplasm of large intestine and rectum 153-154,197.5	161	82	79
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	89	75
Benign neoplasms 210-229	380	51	329
Benign neoplasm of uterus 218-219	216	...	216
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,753	723	1,030
Diabetes mellitus 250	599	299	300
Volume depletion 276.5	518	216	302
Diseases of the blood and blood-forming organs 280-289	449	175	274
Anemias 280-285	316	115	201
Mental disorders 290-319	2,311	1,176	1,135
Psychoses 290-299	1,601	781	820
Schizophrenic disorders 295	331	179	151
Major depressive disorder 296.2-296.3	463	190	273
Diseases of the nervous system and sense organs 320-389	553	238	316
Diseases of the circulatory system 390-459	6,363	3,227	3,136
Essential hypertension 401	341	136	206
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,375	2,306	2,069
Acute myocardial infarction 410	732	423	309
Coronary atherosclerosis 414.0	1,065	662	403
Other ischemic heart disease 411-413,414.1-414.9	184	95	89
Cardiac dysrhythmias 427	762	364	398
Congestive heart failure 428.0,428.2-428.4	1,092	520	571
Cerebrovascular disease 430-438	906	416	490
Diseases of the respiratory system 460-519	3,550	1,712	1,838
Acute bronchitis and bronchiolitis 466	265	142	123
Pneumonia 480-486	1,329	637	692
Chronic bronchitis 491	486	225	261
Asthma 493	497	207	290
Diseases of the digestive system 520-579	3,580	1,580	2,001
Appendicitis 540-543	313	182	131
Noninfectious enteritis and colitis 555-558	315	123	192
Intestinal obstruction 560	305	128	177
Diverticula of intestine 562	312	132	180
Cholelithiasis 574	354	112	242
Acute pancreatitis 577.0	246	117	129
Diseases of the genitourinary system 580-629	1,941	583	1,358
Calculus of kidney and ureter 592	171	88	83
Urinary tract infection 599.0	429	125	304
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	523	...	523
Diseases of the skin and subcutaneous tissue 680-709	764	395	368
Cellulitis and abscess 681-682	562	296	267
Diseases of the musculoskeletal system and connective tissue 710-739	1,920	805	1,115
Osteoarthritis and allied disorders 715	696	259	436
Intervertebral disc disorders 722	369	178	191
Congenital anomalies 740-759	206	112	94
Certain conditions originating in the perinatal period 760-779	203	112	91
Symptoms, signs, and ill-defined conditions 780-799	248	115	133
Injury and poisoning 800-999	2,845	1,392	1,452
Fractures, all sites ² 800-829	1,039	459	580
Fracture of neck of femur ² 820	329	93	236
Poisonings 960-989	256	107	149
Certain complications of surgical and medical care 996-999	871	407	464
Supplementary classifications V01-V83	5,068	403	4,665
Females with deliveries V27	4,136	...	4,136

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 15. Rate of discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Rate per 10,000 population ¹	
		Male	Female
All conditions	1,192.3	964.9	1,411.3
Infectious and parasitic diseases 001-139	32.4	30.7	34.1
Septicemia 038	14.0	13.1	14.9
Neoplasms 140-239	56.1	45.6	66.1
Malignant neoplasms 140-208,230-234	41.2	40.1	42.3
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.5	5.7	5.3
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.6	6.2	5.0
Benign neoplasms 210-229	13.0	3.5	22.1
Benign neoplasm of uterus 218-219	7.4	...	14.5
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	59.9	50.4	69.1
Diabetes mellitus 250	20.5	20.8	20.1
Volume depletion 276.5	17.7	15.0	20.3
Diseases of the blood and blood-forming organs 280-289	15.3	12.2	18.4
Anemias 280-285	10.8	8.0	13.5
Mental disorders 290-319	79.0	82.0	76.2
Psychoses 290-299	54.8	54.4	55.1
Schizophrenic disorders 295	11.3	12.5	10.2
Major depressive disorder 296.2-296.3	15.8	13.2	18.3
Diseases of the nervous system and sense organs 320-389	18.9	16.6	21.2
Diseases of the circulatory system 390-459	217.6	224.9	210.6
Essential hypertension 401	11.7	9.4	13.8
Heart disease 391-392.0,393-398,402,404,410-416,420-429	149.6	160.7	138.9
Acute myocardial infarction 410	25.0	29.5	20.8
Coronary atherosclerosis 414.0	36.4	46.1	27.1
Other ischemic heart disease 411-413,414.1-414.9	6.3	6.6	6.0
Cardiac dysrhythmias 427	26.1	25.4	26.7
Congestive heart failure 428.0,428.2-428.4	37.3	36.3	38.4
Cerebrovascular disease 430-438	31.0	29.0	32.9
Diseases of the respiratory system 460-519	121.4	119.3	123.4
Acute bronchitis and bronchiolitis 466	9.1	9.9	8.3
Pneumonia 480-486	45.5	44.4	46.5
Chronic bronchitis 491	16.6	15.7	17.5
Asthma 493	17.0	14.5	19.4
Diseases of the digestive system 520-579	122.4	110.1	134.3
Appendicitis 540-543	10.7	12.7	8.8
Noninfectious enteritis and colitis 555-558	10.8	8.6	12.9
Intestinal obstruction 560	10.4	8.9	11.9
Diverticula of intestine 562	10.7	9.2	12.1
Cholelithiasis 574	12.1	7.8	16.2
Acute pancreatitis 577.0	8.4	8.1	8.7
Diseases of the genitourinary system 580-629	66.4	40.6	91.2
Calculus of kidney and ureter 592	5.9	6.1	5.6
Urinary tract infection 599.0	14.7	8.7	20.4
Complications of pregnancy, childbirth, and the puerperium ² 630-677	17.9	...	35.1
Diseases of the skin and subcutaneous tissue 680-709	26.1	27.5	24.7
Cellulitis and abscess 681-682	19.2	20.6	17.9
Diseases of the musculoskeletal system and connective tissue 710-739	65.7	56.1	74.9
Osteoarthritis and allied disorders 715	23.8	18.1	29.3
Intervertebral disc disorders 722	12.6	12.4	12.8
Congenital anomalies 740-759	7.0	7.8	6.3
Certain conditions originating in the perinatal period 760-779	7.0	7.8	6.1
Symptoms, signs, and ill-defined conditions 780-799	8.5	8.0	8.9
Injury and poisoning 800-999	97.3	97.0	97.5
Fractures, all sites ³ 800-829	35.5	32.0	38.9
Fracture of neck of femur ³ 820	11.3	6.5	15.8
Poisonings 960-989	8.8	7.5	10.0
Certain complications of surgical and medical care 996-999	29.8	28.4	31.2
Supplementary classifications V01-V83	173.3	28.1	313.2
Females with deliveries V27	141.4	...	277.7

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."³Excludes fractures coded as 733.1, pathologic fracture.

Table 16. Average length of stay for discharges from short-stay hospitals, by sex and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	Both sexes	Average length of stay in days	
		Male	Female
All conditions	4.8	5.2	4.5
Infectious and parasitic diseases 001-139	6.6	6.9	6.4
Septicemia 038	8.4	8.8	8.1
Neoplasms 140-239	5.7	6.6	5.2
Malignant neoplasms 140-208,230-234	6.6	6.8	6.4
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	8.2	9.0
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.9	6.8	7.1
Benign neoplasms 210-229	3.2	5.5	2.9
Benign neoplasm of uterus 218-219	2.5	...	2.5
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.0	4.1	3.9
Diabetes mellitus 250	4.7	4.8	4.6
Volume depletion 276.5	3.3	3.1	3.4
Diseases of the blood and blood-forming organs 280-289	4.7	4.6	4.8
Anemias 280-285	4.9	4.7	5.0
Mental disorders 290-319	7.1	7.0	7.2
Psychoses 290-299	7.9	7.9	7.9
Schizophrenic disorders 295	10.8	10.3	11.3
Major depressive disorder 296.2-296.3	6.8	6.8	6.8
Diseases of the nervous system and sense organs 320-389	5.3	5.7	5.0
Diseases of the circulatory system 390-459	4.7	4.5	4.9
Essential hypertension 401	2.3	2.2	2.4
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.6	4.4	4.8
Acute myocardial infarction 410	5.8	5.7	5.8
Coronary atherosclerosis 414.0	3.5	3.4	3.7
Other ischemic heart disease 411-413,414.1-414.9	2.7	2.4	3.0
Cardiac dysrhythmias 427	3.7	3.3	4.1
Congestive heart failure 428.0,428.2-428.4	5.4	5.2	5.5
Cerebrovascular disease 430-438	5.2	5.0	5.4
Diseases of the respiratory system 460-519	5.2	5.2	5.2
Acute bronchitis and bronchiolitis 466	3.3	3.1	3.4
Pneumonia 480-486	5.5	5.3	5.7
Chronic bronchitis 491	5.0	4.9	5.0
Asthma 493	3.2	2.8	3.5
Diseases of the digestive system 520-579	4.7	4.7	4.7
Appendicitis 540-543	3.1	3.2	3.1
Noninfectious enteritis and colitis 555-558	4.7	4.8	4.7
Intestinal obstruction 560	6.2	5.7	6.5
Diverticula of intestine 562	5.0	4.9	5.1
Cholelithiasis 574	4.0	4.6	3.8
Acute pancreatitis 577.0	5.7	5.8	5.6
Diseases of the genitourinary system 580-629	3.8	4.3	3.6
Calculus of kidney and ureter 592	2.4	2.1	2.6
Urinary tract infection 599.0	4.9	4.9	4.8
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.7	...	2.7
Diseases of the skin and subcutaneous tissue 680-709	5.1	5.0	5.2
Cellulitis and abscess 681-682	4.8	4.5	5.0
Diseases of the musculoskeletal system and connective tissue 710-739	3.9	3.7	4.0
Osteoarthritis and allied disorders 715	3.9	3.8	3.9
Intervertebral disc disorders 722	2.8	2.5	3.1
Congenital anomalies 740-759	6.8	6.7	6.9
Certain conditions originating in the perinatal period 760-779	10.7	10.9	10.4
Symptoms, signs, and ill-defined conditions 780-799	2.7	2.7	2.6
Injury and poisoning 800-999	5.5	5.5	5.4
Fractures, all sites ² 800-829	5.6	5.7	5.5
Fracture of neck of femur ² 820	6.4	6.6	6.3
Poisonings 960-989	3.1	3.1	3.1
Certain complications of surgical and medical care 996-999	6.2	6.3	6.1
Supplementary classifications V01-V83	3.7	8.4	3.3
Females with deliveries V27	2.6	...	2.6

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 17. Number of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
	Number in thousands				
All conditions	34,864	7,388	7,680	13,239	6,557
Infectious and parasitic diseases 001-139	949	196	204	354	195
Septicemia 038	410	80	102	138	90
Neoplasms 140-239	1,640	373	343	620	303
Malignant neoplasms 140-208,230-234	1,205	285	259	444	217
Malignant neoplasm of large intestine and rectum 153-154,197.5	161	40	40	60	22
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	164	34	42	65	23
Benign neoplasms 210-229	380	77	72	156	75
Benign neoplasm of uterus 218-219	216	37	41	97	41
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	1,753	399	382	671	300
Diabetes mellitus 250	599	144	108	237	109
Volume depletion 276.5	518	101	137	203	76
Diseases of the blood and blood-forming organs 280-289	449	111	94	178	66
Anemias 280-285	316	81	68	126	41
Mental disorders 290-319	2,311	707	477	861	267
Psychoses 290-299	1,601	456	356	627	163
Schizophrenic disorders 295	331	*	52	145	31
Major depressive disorder 296.2-296.3	463	117	115	202	30
Diseases of the nervous system and sense organs 320-389	553	130	118	203	102
Diseases of the circulatory system 390-459	6,363	1,441	1,471	2,397	1,055
Essential hypertension 401	341	74	69	145	53
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4,375	1,010	1,029	1,620	716
Acute myocardial infarction 410	732	169	172	259	132
Coronary atherosclerosis 414.0	1,065	231	237	421	175
Other ischemic heart disease 411-413,414.1-414.9	184	44	39	71	30
Cardiac dysrhythmias 427	762	194	180	270	118
Congestive heart failure 428.0,428.2-428.4	1,092	247	280	409	155
Cerebrovascular disease 430-438	906	192	211	346	157
Diseases of the respiratory system 460-519	3,550	769	860	1,306	615
Acute bronchitis and bronchiolitis 466	265	42	53	104	67
Pneumonia 480-486	1,329	271	380	454	223
Chronic bronchitis 491	486	117	109	202	57
Asthma 493	497	130	110	166	92
Diseases of the digestive system 520-579	3,580	756	809	1,305	710
Appendicitis 540-543	313	68	64	94	88
Noninfectious enteritis and colitis 555-558	315	73	77	114	51
Intestinal obstruction 560	305	65	73	105	62
Diverticula of intestine 562	312	80	75	109	48
Cholelithiasis 574	354	73	71	123	87
Acute pancreatitis 577.0	246	44	55	100	46
Diseases of the genitourinary system 580-629	1,941	361	457	776	347
Calculus of kidney and ureter 592	171	34	46	65	25
Urinary tract infection 599.0	429	85	102	170	72
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	523	87	90	243	103
Diseases of the skin and subcutaneous tissue 680-709	764	150	151	351	111
Cellulitis and abscess 681-682	562	120	123	234	85
Diseases of the musculoskeletal system and connective tissue 710-739	1,920	403	479	663	375
Osteoarthritis and allied disorders 715	696	159	190	221	126
Intervertebral disc disorders 722	369	62	89	140	77
Congenital anomalies 740-759	206	53	*	72	*
Certain conditions originating in the perinatal period 760-779	203	26	*	72	*
Symptoms, signs, and ill-defined conditions 780-799	248	57	37	109	46
Injury and poisoning 800-999	2,845	591	612	1,056	586
Fractures, all sites ² 800-829	1,039	214	235	375	215
Fracture of neck of femur ² 820	329	68	71	126	64
Poisonings 960-989	256	52	54	95	55
Certain complications of surgical and medical care 996-999	871	187	175	327	182
Supplementary classifications V01-V83	5,068	778	1,028	2,001	1,261
Females with deliveries V27	4,136	589	802	1,679	1,066

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 18. Rate of discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
All conditions	1,192.3	1,355.2	1,170.0	1,258.2	978.2
Infectious and parasitic diseases 001-139	32.4	36.0	31.0	33.7	29.0
Septicemia 038	14.0	14.7	15.5	13.1	13.4
Neoplasms 140-239	56.1	68.4	52.3	58.9	45.2
Malignant neoplasms 140-208,230-234	41.2	52.3	39.4	42.2	32.4
Malignant neoplasm of large intestine and rectum 153-154,197.5	5.5	7.3	6.1	5.7	3.2
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	5.6	6.2	6.4	6.2	3.4
Benign neoplasms 210-229	13.0	14.1	11.0	14.8	11.2
Benign neoplasm of uterus 218-219	7.4	6.9	6.2	9.2	6.1
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	59.9	73.2	58.2	63.8	44.8
Diabetes mellitus 250	20.5	26.5	16.5	22.5	16.3
Volume depletion 276.5	17.7	18.6	20.9	19.3	11.4
Diseases of the blood and blood-forming organs 280-289	15.3	20.4	14.3	16.9	9.9
Anemias 280-285	10.8	14.9	10.3	12.0	6.1
Mental disorders 290-319	79.0	129.6	72.6	81.8	39.8
Psychoses 290-299	54.8	83.7	54.2	59.6	24.2
Schizophrenic disorders 295	11.3	*	7.9	13.8	4.6
Major depressive disorder 296.2-296.3	15.8	21.4	17.5	19.2	4.4
Diseases of the nervous system and sense organs 320-389	18.9	23.9	18.0	19.3	15.2
Diseases of the circulatory system 390-459	217.6	264.2	224.0	227.8	157.4
Essential hypertension 401	11.7	13.7	10.5	13.8	7.9
Heart disease 391-392.0,393-398,402,404,410-416,420-429	149.6	185.4	156.7	154.0	106.8
Acute myocardial infarction 410	25.0	31.0	26.2	24.6	19.8
Coronary atherosclerosis 414.0	36.4	42.4	36.1	40.0	26.2
Other ischemic heart disease 411-413,414.1-414.9	6.3	8.1	5.9	6.8	4.4
Cardiac dysrhythmias 427	26.1	35.6	27.5	25.7	17.5
Congestive heart failure 428.0,428.2-428.4	37.3	45.3	42.7	38.9	23.1
Cerebrovascular disease 430-438	31.0	35.2	32.1	32.9	23.4
Diseases of the respiratory system 460-519	121.4	141.0	131.0	124.1	91.8
Acute bronchitis and bronchiolitis 466	9.1	7.6	8.1	9.8	10.0
Pneumonia 480-486	45.5	49.8	57.9	43.2	33.3
Chronic bronchitis 491	16.6	21.5	16.7	19.2	8.5
Asthma 493	17.0	23.8	16.7	15.7	13.8
Diseases of the digestive system 520-579	122.4	138.6	123.2	124.0	106.0
Appendicitis 540-543	10.7	12.4	9.8	8.9	13.1
Noninfectious enteritis and colitis 555-558	10.8	13.4	11.7	10.9	7.7
Intestinal obstruction 560	10.4	12.0	11.1	10.0	9.2
Diverticula of intestine 562	10.7	14.7	11.4	10.3	7.2
Cholelithiasis 574	12.1	13.4	10.8	11.7	13.0
Acute pancreatitis 577.0	8.4	8.1	8.4	9.5	6.9
Diseases of the genitourinary system 580-629	66.4	66.3	69.6	73.8	51.7
Calculus of kidney and ureter 592	5.9	6.3	7.1	6.2	3.7
Urinary tract infection 599.0	14.7	15.6	15.5	16.1	10.7
Complications of pregnancy, childbirth, and the puerperium ² 630-677	17.9	16.0	13.7	23.1	15.3
Diseases of the skin and subcutaneous tissue 680-709	26.1	27.5	23.1	33.4	16.5
Cellulitis and abscess 681-682	19.2	22.0	18.8	22.3	12.7
Diseases of the musculoskeletal system and connective tissue 710-739	65.7	74.0	73.0	63.0	56.0
Osteoarthritis and allied disorders 715	23.8	29.2	29.0	21.0	18.8
Intervertebral disc disorders 722	12.6	11.5	13.5	13.3	11.5
Congenital anomalies 740-759	7.0	9.8	*	6.9	*
Certain conditions originating in the perinatal period 760-779	7.0	4.8	*	6.8	*
Symptoms, signs, and ill-defined conditions 780-799	8.5	10.5	5.6	10.3	6.8
Injury and poisoning 800-999	97.3	108.4	93.3	100.3	87.4
Fractures, all sites ³ 800-829	35.5	39.3	35.8	35.6	32.0
Fracture of neck of femur ³ 820	11.3	12.5	10.7	12.0	9.6
Poisonings 960-989	8.8	9.5	8.3	9.0	8.2
Certain complications of surgical and medical care 996-999	29.8	34.3	26.7	31.1	27.1
Supplementary classifications V01-V83	173.3	142.6	156.6	190.2	188.1
Females with deliveries V27	141.4	108.0	122.2	159.6	159.0

* Figure does not meet standard of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."³Excludes fractures coded as 733.1, pathologic fracture.

Table 19. Average length of stay for discharges from short-stay hospitals, by geographic region and first-listed diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Category of first-listed diagnosis and ICD-9-CM code	United States	Northeast	Midwest	South	West
		Average length of stay in days			
All conditions	4.8	5.4	4.4	4.9	4.5
Infectious and parasitic diseases 001-139	6.6	7.8	5.4	6.7	6.4
Septicemia 038	8.4	10.2	6.7	8.9	8.0
Neoplasms 140-239	5.7	6.3	5.3	5.9	5.2
Malignant neoplasms 140-208,230-234	6.6	7.1	5.9	6.9	6.0
Malignant neoplasm of large intestine and rectum 153-154,197.5	8.6	9.3	7.9	8.9	7.7
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	6.9	8.1	6.3	6.9	6.4
Benign neoplasms 210-229	3.2	3.4	3.2	3.3	3.0
Benign neoplasm of uterus 218-219	2.5	2.5	2.6	2.4	2.5
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	4.0	4.5	3.4	4.1	3.8
Diabetes mellitus 250	4.7	5.1	3.7	5.0	4.5
Volume depletion 276.5	3.3	4.0	3.1	3.3	2.5
Diseases of the blood and blood-forming organs 280-289	4.7	4.7	3.6	5.3	4.6
Anemias 280-285	4.9	4.8	3.5	5.7	4.8
Mental disorders 290-319	7.1	8.2	5.8	7.2	6.1
Psychoses 290-299	7.9	9.5	6.4	7.7	7.4
Schizophrenic disorders 295	10.8	13.5	9.3	9.8	8.9
Major depressive disorder 296.2-296.3	6.8	8.8	5.4	6.7	5.2
Diseases of the nervous system and sense organs 320-389	5.3	5.1	5.0	5.5	5.3
Diseases of the circulatory system 390-459	4.7	5.1	4.1	4.8	4.7
Essential hypertension 401	2.3	2.8	2.0	2.3	2.0
Heart disease 391-392.0,393-398,402,404,410-416,420-429	4.6	5.0	4.0	4.8	4.6
Acute myocardial infarction 410	5.8	5.7	5.3	6.2	5.5
Coronary atherosclerosis 414.0	3.5	3.5	3.1	3.7	3.6
Other ischemic heart disease 411-413,414.1-414.9	2.7	2.9	2.4	2.4	*
Cardiac dysrhythmias 427	3.7	3.9	2.9	4.1	3.8
Congestive heart failure 428.0,428.2-428.4	5.4	6.4	4.5	5.5	4.8
Cerebrovascular disease 430-438	5.2	5.3	4.5	5.5	5.5
Diseases of the respiratory system 460-519	5.2	5.6	4.6	5.4	5.1
Acute bronchitis and bronchiolitis 466	3.3	3.5	2.8	3.3	3.4
Pneumonia 480-486	5.5	5.8	5.1	5.9	5.3
Chronic bronchitis 491	5.0	5.9	4.3	4.9	4.4
Asthma 493	3.2	3.6	2.6	3.3	3.1
Diseases of the digestive system 520-579	4.7	5.0	4.3	4.9	4.5
Appendicitis 540-543	3.1	3.0	3.1	3.2	3.2
Noninfectious enteritis and colitis 555-558	4.7	5.5	4.2	4.6	4.4
Intestinal obstruction 560	6.2	6.3	5.5	6.5	6.2
Diverticula of intestine 562	5.0	5.8	4.5	4.9	4.9
Cholelithiasis 574	4.0	4.6	3.9	3.9	3.7
Acute pancreatitis 577.0	5.7	5.8	5.5	5.7	5.7
Diseases of the genitourinary system 580-629	3.8	4.6	3.3	4.0	3.5
Calculus of kidney and ureter 592	2.4	3.0	2.0	2.3	2.3
Urinary tract infection 599.0	4.9	5.4	4.1	5.0	4.8
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	2.7	2.7	3.0	2.6	2.6
Diseases of the skin and subcutaneous tissue 680-709	5.1	5.4	4.4	5.2	5.1
Cellulitis and abscess 681-682	4.8	5.2	4.1	4.9	4.8
Diseases of the musculoskeletal system and connective tissue 710-739	3.9	4.1	3.8	3.9	3.6
Osteoarthritis and allied disorders 715	3.9	3.9	3.9	4.0	3.6
Intervertebral disc disorders 722	2.8	3.3	2.6	2.6	2.9
Congenital anomalies 740-759	6.8	5.9	10.2	5.8	7.1
Certain conditions originating in the perinatal period 760-779	10.7	7.9	10.1	13.1	9.5
Symptoms, signs, and ill-defined conditions 780-799	2.7	3.1	2.0	2.7	2.5
Injury and poisoning 800-999	5.5	5.7	5.2	5.8	4.9
Fractures, all sites ² 800-829	5.6	5.8	5.0	6.0	5.4
Fracture of neck of femur ² 820	6.4	6.5	6.3	6.6	6.0
Poisonings 960-989	3.1	3.0	*	3.1	2.7
Certain complications of surgical and medical care 996-999	6.2	6.9	5.6	6.7	5.1
Supplementary classifications V01-V83	3.7	4.3	4.0	3.5	3.5
Females with deliveries V27	2.6	2.9	2.7	2.6	2.4

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 20. Number of all-listed diagnoses for discharges from short-stay hospitals, by age and diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnoses and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All conditions	165,517	7,277	41,163	41,446	75,631
Infectious and parasitic diseases 001-139	4,143	457	897	1,117	1,671
Septicemia 038	803	23	73	206	501
Neoplasms 140-239	4,532	118	665	1,548	2,201
Malignant neoplasms 140-208,230-234	3,486	99	308	1,169	1,910
Malignant neoplasm of large intestine and rectum 153-154,197.5	273	*	17	91	164
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	495	*	29	167	296
Benign neoplasms 210-229	901	14	337	347	204
Benign neoplasm of uterus 218-219	479	*	249	211	18
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	19,280	617	2,537	5,946	10,179
Diabetes mellitus 250	5,435	43	588	1,879	2,926
Volume depletion 276.5	2,528	338	357	465	1,369
Diseases of the blood and blood-forming organs 280-289	5,051	187	1,050	1,225	2,590
Anemias 280-285	3,864	106	817	878	2,063
Mental disorders 290-319	12,158	379	5,076	3,732	2,971
Psychoses 290-299	3,949	107	1,331	950	1,562
Schizophrenic disorders 295	568	*	245	223	97
Major depressive disorder 296.2-296.3	653	*	344	182	99
Diseases of the nervous system and sense organs 320-389	3,873	368	649	926	1,931
Diseases of the circulatory system 390-459	34,384	158	2,060	8,929	23,236
Essential hypertension 401	8,936	21	711	2,805	5,399
Heart disease 391-392.0,393-398,402,404,410-416,420-429	19,679	90	884	4,590	14,114
Acute myocardial infarction 410	902	*	48	272	580
Coronary atherosclerosis 414.0	4,457	*	133	1,261	3,060
Other ischemic heart disease 411-413,414.1-414.9	2,126	*	93	683	1,345
Cardiac dysrhythmias 427	4,371	26	198	735	3,412
Congestive heart failure 428.0,428.2-428.4	4,016	15	115	777	3,109
Cerebrovascular disease 430-438	1,906	*	91	457	1,346
Diseases of the respiratory system 460-519	11,855	1,235	1,419	2,828	6,373
Acute bronchitis and bronchiolitis 466	439	234	38	56	111
Pneumonia 480-486	2,326	270	200	461	1,395
Chronic bronchitis 491	1,056	*	30	294	729
Asthma 493	1,870	348	501	504	517
Diseases of the digestive system 520-579	11,144	538	2,225	3,453	4,928
Appendicitis 540-543	349	86	175	62	26
Noninfectious enteritis and colitis 555-558	687	122	178	158	229
Intestinal obstruction 560	776	29	111	211	425
Diverticula of intestine 562	646	*	59	160	426
Cholelithiasis 574	627	4	192	179	252
Acute pancreatitis 577.0	337	6	109	122	100
Diseases of the genitourinary system 580-629	8,642	195	2,092	2,141	4,215
Calculus of kidney and ureter 592	293	*	121	106	64
Urinary tract infection 599.0	2,126	51	235	370	1,470
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	10,337	16	10,292	29	...
Diseases of the skin and subcutaneous tissue 680-709	2,511	210	520	719	1,063
Cellulitis and abscess 681-682	1,185	75	287	373	450
Diseases of the musculoskeletal system and connective tissue 710-739	6,868	88	1,012	2,047	3,721
Osteoarthritis and allied disorders 715	1,822	*	74	479	1,268
Intervertebral disc disorders 722	623	*	187	255	180
Congenital anomalies 740-759	730	427	138	97	67
Certain conditions originating in the perinatal period 760-779	677	669	*	*	*
Symptoms, signs, and ill-defined conditions 780-799	8,836	747	1,920	2,536	3,633
Injury and poisoning 800-999	6,914	489	2,144	1,739	2,542
Fractures, all sites ² 800-829	1,661	99	469	337	757
Fracture of neck of femur ² 820	377	*	10	36	329
Poisonings 960-989	458	30	279	103	46
Certain complications of surgical and medical care 996-999	2,221	97	413	761	951
Supplementary classifications V01-V83	13,582	379	6,462	2,434	4,306
Females with deliveries V27	4,136	*5	4,120	10	...

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 21. Number of all-listed diagnoses for discharges from short-stay hospitals, by sex and diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnoses and ICD-9-CM code	Both sexes	Number in thousands	
		Male	Female
All conditions	165,517	67,051	98,465
Infectious and parasitic diseases 001-139	4,143	1,887	2,256
Septicemia 038	803	394	410
Neoplasms 140-239	4,532	1,928	2,603
Malignant neoplasms 140-208,230-234	3,486	1,714	1,772
Malignant neoplasm of large intestine and rectum 153-154,197.5	273	139	134
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	495	253	242
Benign neoplasms 210-229	901	147	755
Benign neoplasm of uterus 218-219	479	...	479
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	19,280	8,122	11,158
Diabetes mellitus 250	5,435	2,541	2,894
Volume depletion 276.5	2,528	1,053	1,475
Diseases of the blood and blood-forming organs 280-289	5,051	1,962	3,089
Anemias 280-285	3,864	1,405	2,459
Mental disorders 290-319	12,158	5,754	6,404
Psychoses 290-299	3,949	1,778	2,171
Schizophrenic disorders 295	568	296	272
Major depressive disorder 296.2-296.3	653	259	395
Diseases of the nervous system and sense organs 320-389	3,873	1,705	2,168
Diseases of the circulatory system 390-459	34,384	16,462	17,922
Essential hypertension 401	8,936	3,769	5,166
Heart disease 391-392.0,393-398,402,404,410-416,420-429	19,679	9,874	9,805
Acute myocardial infarction 410	902	507	395
Coronary atherosclerosis 414.0	4,457	2,558	1,899
Other ischemic heart disease 411-413,414.1-414.9	2,126	1,247	879
Cardiac dysrhythmias 427	4,371	2,129	2,242
Congestive heart failure 428.0,428.2-428.4	4,016	1,792	2,225
Cerebrovascular disease 430-438	1,906	877	1,029
Diseases of the respiratory system 460-519	11,855	5,605	6,250
Acute bronchitis and bronchiolitis 466	439	216	222
Pneumonia 480-486	2,326	1,149	1,178
Chronic bronchitis 491	1,056	507	550
Asthma 493	1,870	641	1,229
Diseases of the digestive system 520-579	11,144	4,758	6,386
Appendicitis 540-543	349	197	152
Noninfectious enteritis and colitis 555-558	687	262	425
Intestinal obstruction 560	776	342	434
Diverticula of intestine 562	646	255	391
Cholelithiasis 574	627	209	418
Acute pancreatitis 577.0	337	159	179
Diseases of the genitourinary system 580-629	8,642	3,040	5,602
Calculus of kidney and ureter 592	293	140	153
Urinary tract infection 599.0	2,126	596	1,530
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	10,337	...	10,337
Diseases of the skin and subcutaneous tissue 680-709	2,511	1,228	1,284
Cellulitis and abscess 681-682	1,185	596	589
Diseases of the musculoskeletal system and connective tissue 710-739	6,868	2,363	4,504
Osteoarthritis and allied disorders 715	1,822	591	1,231
Intervertebral disc disorders 722	623	290	333
Congenital anomalies 740-759	730	374	356
Certain conditions originating in the perinatal period 760-779	677	381	296
Symptoms, signs, and ill-defined conditions 780-799	8,836	3,862	4,974
Injury and poisoning 800-999	6,914	3,599	3,315
Fractures, all sites ² 800-829	1,661	814	847
Fracture of neck of femur ² 820	377	112	265
Poisonings 960-989	458	182	276
Certain complications of surgical and medical care 996-999	2,221	1,064	1,158
Supplementary classifications V01-V83	13,582	4,021	9,561
Females with deliveries V27	4,136	...	4,136

... Category not applicable.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 22. Number of all-listed diagnoses for discharges from short-stay hospitals, by geographic region and diagnosis: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

All-listed diagnoses and ICD-9-CM code	United States	Northeast	Midwest	South	West
	Number in thousands				
All conditions	165,517	35,532	39,478	59,942	30,565
Infectious and parasitic diseases 001-139	4,143	908	875	1,564	795
Septicemia 038	803	169	178	289	168
Neoplasms 140-239	4,532	1,051	1,012	1,666	803
Malignant neoplasms 140-208,230-234	3,486	842	793	1,254	598
Malignant neoplasm of large intestine and rectum 153-154,197.5	273	66	65	103	40
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	495	117	130	180	68
Benign neoplasms 210-229	901	178	185	362	176
Benign neoplasm of uterus 218-219	479	78	96	211	93
Endocrine, nutritional and metabolic diseases, and immunity disorders 240-279	19,280	4,273	4,752	6,956	3,299
Diabetes mellitus 250	5,435	1,246	1,237	2,065	887
Volume depletion 276.5	2,528	500	655	935	438
Diseases of the blood and blood-forming organs 280-289	5,051	1,045	1,216	1,865	925
Anemias 280-285	3,864	802	945	1,424	693
Mental disorders 290-319	12,158	3,139	2,958	4,124	1,937
Psychoses 290-299	3,949	1,058	930	1,408	553
Schizophrenic disorders 295	568	162	98	229	78
Major depressive disorder 296.2-296.3	653	160	161	277	55
Diseases of the nervous system and sense organs 320-389	3,873	879	946	1,312	736
Diseases of the circulatory system 390-459	34,384	7,917	8,420	12,365	5,682
Essential hypertension 401	8,936	2,046	2,131	3,349	1,411
Heart disease 391-392.0,393-398,402,404,410-416,420-429	19,679	4,642	4,935	6,869	3,233
Acute myocardial infarction 410	902	215	212	310	165
Coronary atherosclerosis 414.0	4,457	1,079	1,098	1,602	678
Other ischemic heart disease 411-413,414.1-414.9	2,126	511	530	726	360
Cardiac dysrhythmias 427	4,371	1,047	1,125	1,439	760
Congestive heart failure 428.0,428.2-428.4	4,016	883	1,090	1,457	587
Cerebrovascular disease 430-438	1,906	406	454	679	367
Diseases of the respiratory system 460-519	11,855	2,565	2,893	4,290	2,106
Acute bronchitis and bronchiolitis 466	439	76	89	177	97
Pneumonia 480-486	2,326	475	621	814	416
Chronic bronchitis 491	1,056	240	253	421	142
Asthma 493	1,870	475	434	601	359
Diseases of the digestive system 520-579	11,144	2,381	2,604	4,109	2,051
Appendicitis 540-543	349	73	73	107	96
Noninfectious enteritis and colitis 555-558	687	152	167	259	110
Intestinal obstruction 560	776	148	190	291	147
Diverticula of intestine 562	646	158	156	232	99
Cholelithiasis 574	627	137	126	224	140
Acute pancreatitis 577.0	337	63	71	131	71
Diseases of the genitourinary system 580-629	8,642	1,727	2,072	3,249	1,594
Calculus of kidney and ureter 592	293	59	75	109	50
Urinary tract infection 599.0	2,126	458	510	793	365
Complications of pregnancy, childbirth, and the puerperium ¹ 630-677	10,337	1,556	2,068	4,109	2,604
Diseases of the skin and subcutaneous tissue 680-709	2,511	556	572	962	422
Cellulitis and abscess 681-682	1,185	260	262	460	202
Diseases of the musculoskeletal system and connective tissue 710-739	6,868	1,432	1,804	2,368	1,265
Osteoarthritis and allied disorders 715	1,822	404	524	615	279
Intervertebral disc disorders 722	623	96	152	232	143
Congenital anomalies 740-759	730	174	134	250	*
Certain conditions originating in the perinatal period 760-779	677	80	*	248	*
Symptoms, signs, and ill-defined conditions 780-799	8,836	1,880	2,336	3,041	1,578
Injury and poisoning 800-999	6,914	1,400	1,570	2,471	1,474
Fractures, all sites ² 800-829	1,661	321	380	601	359
Fracture of neck of femur ² 820	377	77	82	147	72
Poisonings 960-989	458	90	104	157	107
Certain complications of surgical and medical care 996-999	2,221	477	474	803	467
Supplementary classifications V01-V83	13,582	2,569	3,119	4,992	2,902
Females with deliveries V27	4,136	589	802	1,679	1,066

* Figure does not meet standard of reliability or precision.

¹First-listed diagnosis for females with deliveries is coded V27, shown under "Supplementary classifications."²Excludes fractures coded as 733.1, pathologic fracture.

Table 23. Number and rate of discharges from short-stay hospitals with a human immunodeficiency virus diagnosis, by selected characteristics: United States, selected years 1990–2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Data are for discharges with at least one of the following *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) codes: 042–044, 279.19, 795.8 (prior to 1995); 042, V08 (beginning in 1995)]

Characteristic	1990	1995	2000	2001	2002	2003	2004
Number in thousands							
All HIV discharges ¹	146	249	173	185	189	207	204
Sex							
Male	114	183	115	115	124	136	132
Female	32	66	58	70	65	71	73
Age							
Under 25 years	14	17	7	10	10	9	10
25–29 years	20	26	11	8	11	13	8
30–34 years	38	48	28	28	23	21	15
35–39 years	36	59	35	41	35	38	35
40–44 years	17	49	34	43	42	38	46
45 years and over	21	51	58	56	68	87	90
Region							
Northeast	66	95	68	61	57	70	67
Midwest	20	27	20	18	20	17	21
South	37	87	64	86	81	94	99
West	24	41	21	21	31	27	17
Rate per 100,000 population ²							
All HIV discharges ¹	58.1	93.9	61.6	65.2	65.8	71.6	69.9
Sex							
Male	93.4	141.6	83.9	82.8	88.5	95.9	91.7
Female	24.7	48.4	40.1	48.2	44.1	48.2	48.9
Age							
Under 25 years	14.8	17.6	7.5	10.1	9.5	8.9	10.2
25–29 years	95.1	132.2	56.6	41.3	57.5	71.3	38.9
30–34 years	172.8	216.2	134.9	137.6	113.0	104.6	71.9
35–39 years	179.0	262.7	154.4	182.9	159.7	180.8	168.3
40–44 years	94.8	241.4	152.6	187.4	185.2	166.6	201.3
45 years and over	27.4	60.2	59.8	55.8	66.6	83.1	84.6
Region							
Northeast	128.8	183.3	126.6	112.3	107.9	129.4	122.1
Midwest	33.6	43.5	30.8	27.9	31.7	25.9	32.6
South	42.1	93.1	64.2	84.5	81.7	90.1	94.3
West	44.1	69.5	33.4	32.6	49.9	40.2	25.7

¹HIV is human immunodeficiency virus.

²Rates are based on the U.S. Census Bureau estimates of the civilian population. Populations for 1990–99 were adjusted for the net underenumeration in the 1990 census. Population for 2000–2004 were 2000-based postcensal estimates. Rates for 2000–2001 differ from rates shown in a previous report because they were calculated using revised population estimates.

Table 24. Annual average number and rate of discharges from short-stay hospitals with a first-listed injury diagnosis, by Barell injury diagnosis matrix categories: United States, 2002–2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. *International Classification of Diseases, 9th Revision, Clinical Modification* code numbers for Barell matrix categories are shown in Table III]

Nature of injury	All body regions	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
Number in thousands ¹									
All injuries	1,903	191	127	12	92	240	240	686	26
Fracture	1,029	52	38	9	80	98	153	598	*
Dislocation	18	...	*	...	3	*	6	8	–
Sprains and strains	51	...	–	...	9	3	16	20	*2
Internal	224	138	...	3	...	82	*
Open wound	94	...	39	13	23	18	*
Amputation	9	7	*1	...
Blood vessels	8	...	*	*2	4	*1	*
Contusions and superficial injuries	63	...	14	19	7	19	4
Crush	3	...	*	*	*2	*	–
Burns	79	...	13	*	18	17	*
Nerves and spinal cord	4	*	*	*	*3	...	*
Unspecified	32	...	22	5	*	*2	*3
System-wide	289
Rate per 10,000 population ²									
All injuries	65.7	6.6	4.4	0.4	3.2	8.3	8.3	23.7	0.9
Fracture	35.5	1.8	1.3	0.3	2.8	3.4	5.3	20.7	*
Dislocation	0.6	...	*	...	0.1	*	0.2	0.3	–
Sprains and strains	1.8	...	–	...	0.3	0.1	0.6	0.7	*0.1
Internal	7.7	4.8	...	0.1	...	2.8	*
Open wound	3.2	...	1.3	0.5	0.8	0.6	*
Amputation	0.3	0.3	*0.1	...
Blood vessels	0.3	...	*	*0.1	0.1	*0.0	*
Contusions and superficial injuries	2.2	...	0.5	0.6	0.2	0.7	0.1
Crush	0.1	...	*	*	*0.1	*	–
Burns	2.7	...	0.5	*	0.6	0.6	*
Nerves and spinal cord	0.1	*	*	*	*0.1	...	*
Unspecified	1.1	...	0.8	0.2	*	*0.1	*0.1
System-wide	10.0

* Figure does not meet standard of reliability or precision.

... Category not applicable.

– Quantity zero.

¹The annual average number of discharges was computed by adding the number of discharges with a first-listed injury diagnosis during the 3-year period and dividing by three.²Rates were calculated by dividing the annual average number of discharges by the 2003 population. The population is the U.S. Census Bureau 2000-based postcensal estimate of the civilian population.**Table 25. Number and rate of deaths for discharges from short-stay hospitals, by age and selected first-listed diagnosis: United States, 2004**[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM)]

First-listed diagnosis and ICD–9–CM code	All ages	Under 65 years	65 years and over	All ages	Under 65 years	65 years and over
	Number of deaths in thousands			Rate per 100 discharges		
All conditions ¹	802	212	590	2.3	1.0	4.5
Heart disease 391–392.0,393–398,402,404,410–416,420–429	146	21	125	3.3	1.3	4.5
Acute myocardial infarction 410	64	7	56	8.7	2.6	12.2
Cardiac dysrhythmias 427	19	*3	16	2.5	*1.5	3.0
Congestive heart failure 428.0,428.2–428.4	40	*5	35	3.7	*2.0	4.3
Malignant neoplasms 140–208,230–234	84	33	50	6.9	5.8	8.0
Malignant neoplasm of trachea, bronchus, and lung 162,176.4,197.0,197.3	21	8	13	12.9	11.9	13.6
Septicemia 038	73	14	58	17.8	11.5	20.5
Pneumonia 480–486	72	13	58	5.4	2.5	7.3
Cerebrovascular disease 430–438	60	17	43	6.7	6.5	6.8
Respiratory failure 518.81,518.83,518.84	50	12	38	21.9	13.6	27.3
Pneumonitis due to solids and liquids 507	26	*4	23	14.5	*10.2	15.5
Nephritis, nephrotic syndrome and nephrosis 580–589	24	*	19	9.1	*	11.6
Fractures, all sites ² 800–829	17	*4	13	1.7	*0.8	2.4

* Figure does not meet standard of reliability or precision.

¹Includes data for deaths not shown in table.²Excludes fractures coded as 733.1, pathologic fracture.

Table 26. Number of discharges from short-stay hospitals with and without procedures and percentage with procedures, by selected characteristics: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants]

Characteristic	All discharges	Discharges without procedures	Discharges with procedures			
			Total	With surgical procedures	Total	With surgical procedures
			Number in thousands		Percent	
All patients	34,864	13,322	21,543	15,960	61.8	45.8
Age						
Under 15 years	2,571	1,334	1,237	689	48.1	26.8
15–44 years	10,800	3,024	7,776	6,136	72.0	56.8
45–64 years	8,323	3,126	5,197	3,947	62.4	47.4
65 years and over	13,171	5,838	7,333	5,188	55.7	39.4
65–74 years	4,786	1,795	2,992	2,270	62.5	47.4
75–84 years	5,538	2,494	3,044	2,125	55.0	38.4
85 years and over	2,846	1,549	1,297	793	45.6	27.9
Sex						
Male	13,844	5,807	8,038	5,723	58.1	41.3
Female	21,020	7,515	13,505	10,237	64.2	48.7
Region						
Northeast	7,388	2,724	4,664	3,082	63.1	41.7
Midwest	7,680	3,333	4,347	3,281	56.6	42.7
South	13,239	5,222	8,017	6,209	60.6	46.9
West	6,557	2,043	4,515	3,388	68.8	51.7

NOTE: All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.

Table 27. Number and rate of all-listed surgical and nonsurgical procedures for discharges from short-stay hospitals by selected procedure categories: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM)]

Procedure category and ICD–9–CM code	Number in thousands	Rate per 10,000 population ¹
Surgical procedures ²	26,632	910.7
Cardiac catheterization 37.21–37.23	1,297	44.3
Repair of current obstetric laceration ³ 75.5–75.6	1,241	42.4
Cesarean section ³ 74.0–74.2,74.4,74.99	1,236	42.3
Artificial rupture of membranes ³ 73.0	941	32.2
Reduction of fracture 76.7,79.0–79.3	667	22.8
Open reduction of fracture with internal fixation 79.3	441	15.1
Episiotomy with or without forceps or vacuum extraction ³ 72.1,72.21,72.31,72.71,73.6	667	22.8
Balloon angioplasty of coronary artery or coronary atherectomy 36.01–36.02,36.05	664	22.7
Endoscopy of small intestine with biopsy 45.14,45.16	658	22.5
Hysterectomy 68.3–68.7,68.9	617	21.1
Insertion of coronary artery stent(s) 36.06–36.07	615	21.0
Total knee replacement 81.54	478	16.3
Oophorectomy and salpingo-oophorectomy 65.3–65.6	451	15.4
Cholecystectomy 51.2	435	14.9
Coronary artery bypass graft ⁴ 36.1	427	14.6
Debridement of wound, infection, or burn 86.22,86.28	368	12.6
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7–37.8	362	12.4
Bilateral destruction or occlusion of fallopian tubes 66.2–66.3	357	12.2
Lysis of peritoneal adhesions 54.5	342	11.7
Appendectomy, excluding incidental 47.0	332	11.3
Excision or destruction of intervertebral disc 80.5	326	11.2
Partial excision of large intestine 45.7	259	8.8
Total hip replacement 81.51	234	8.0
Endoscopy of large intestine with biopsy 45.25	220	7.5
Bronchoscopy with biopsy 33.24,33.27	177	6.1
Partial excision of bone 76.2–76.3,77.6–77.8	172	5.9
Prostatectomy 60.2–60.6	161	5.5
Cystoscopy with biopsy 57.33	13	0.4
Nonsurgical procedures ⁵	18,391	628.9
Arteriography and angiocardiology using contrast material 88.4–88.5	2,110	72.1
Transfusion of blood and blood components 99.0	1,815	62.1
Respiratory therapy 93.9,96.7	1,157	39.6
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	828	28.3
Diagnostic ultrasound 88.7	813	27.8
Hemodialysis 39.95	607	20.8
Medical induction of labor ³ 73.4	573	19.6
Insertion of endotracheal tube 96.04	511	17.5
Endoscopy of small intestine without biopsy 45.11–45.13	457	15.6
Physical therapy procedures 93.1–93.3	415	14.2
Endoscopy of large intestine without biopsy 45.21–45.24	376	12.9
Spinal tap 03.31	357	12.2
Injection or infusion of cancer chemotherapeutic substance 99.25	235	8.0
Cystoscopy without biopsy 57.31–57.32	133	4.5
Bronchoscopy without biopsy 33.21–33.23	92	3.2

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.³See Table 38 for rate per 100 deliveries.⁴See Table 29 for comparison of number of procedures with number of discharges with procedures.⁵Includes data for nonsurgical procedures not shown in table. The ICD–9–CM codes for nonsurgical procedures are listed in the appendix.

Table 28. Number and rate of all-listed surgical and nonsurgical procedures for patients 65 years of age and over discharged from short-stay hospitals, by age and selected procedure categories: United States, 2004[Discharges of inpatients from nonfederal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Procedure category and ICD-9-CM code	65 years and over	65-74 years	75-84 years	85 years and over
	Number in thousands			
Surgical procedures ¹	8,769	3,982	3,576	1,212
Cardiac catheterization 37.21-37.23	638	323	266	49
Endoscopy of small intestine with biopsy 45.14,45.16	368	135	158	76
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	342	175	137	30
Insertion of coronary artery stent(s) 36.06-36.07	316	161	127	28
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	306	73	156	77
Total knee replacement 81.54	292	162	116	13
Reduction of fracture 76.7,79.0-79.3	267	78	100	89
Open reduction of fracture with internal fixation 79.3	194	58	71	65
Coronary artery bypass graft ² 36.1	215	116	92	7
Cholecystectomy 51.2	148	70	62	16
Partial excision of large intestine 45.7	138	65	53	20
Total hip replacement 81.51	135	66	59	11
Debridement of wound, infection, or burn 86.22,86.28	126	52	47	27
Prostatectomy 60.2-60.6	106	57	39	10
Nonsurgical procedures ³	7,136	2,757	3,044	1,335
Transfusion of blood and blood components 99.0	1,053	363	441	250
Arteriography and angiocardiology using contrast material 88.4-88.5	1,053	521	437	95
Respiratory therapy 93.9,96.7	497	183	209	105
Diagnostic ultrasound 88.7	405	141	184	80
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	395	127	174	94
Hemodialysis 39.95	296	144	121	31
Physical therapy procedures 93.1-93.3	274	87	125	62
Insertion of endotracheal tube 96.04	261	93	119	49
Endoscopy of small intestine without biopsy 45.11-45.13	250	80	112	59
Endoscopy of large intestine without biopsy 45.21-45.24	233	70	110	53
	Rate per 10,000 population ⁴			
Surgical procedures ¹	2,416.0	2,156.5	2,756.6	2,493.1
Cardiac catheterization 37.21-37.23	175.7	174.8	204.8	101.5
Endoscopy of small intestine with biopsy 45.14,45.16	101.5	73.1	121.8	155.5
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	94.3	94.7	105.8	62.0
Insertion of coronary artery stent(s) 36.06-36.07	87.2	87.3	97.6	58.6
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	84.3	39.7	120.0	158.1
Total knee replacement 81.54	80.3	87.9	89.5	27.0
Reduction of fracture 76.7,79.0-79.3	73.6	42.4	77.0	182.8
Open reduction of fracture with internal fixation 79.3	53.4	31.4	54.7	133.9
Coronary artery bypass graft ² 36.1	59.3	62.8	71.0	15.1
Cholecystectomy 51.2	40.9	38.0	47.6	33.9
Partial excision of large intestine 45.7	38.1	35.5	40.7	41.4
Total hip replacement 81.51	37.3	35.8	45.1	22.1
Debridement of wound, infection, or burn 86.22,86.28	34.7	28.2	35.9	56.3
Prostatectomy 60.2-60.6	29.1	30.9	29.7	20.9
Nonsurgical procedures ³	1,966.3	1,493.5	2,346.9	2,746.7
Transfusion of blood and blood components 99.0	290.2	196.7	339.7	513.5
Arteriography and angiocardiology using contrast material 88.4-88.5	290.1	282.1	337.0	195.3
Respiratory therapy 93.9,96.7	137.0	99.2	161.5	215.3
Diagnostic ultrasound 88.7	111.7	76.4	141.6	165.5
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	109.0	68.9	134.2	194.0
Hemodialysis 39.95	81.5	77.8	93.3	64.4
Physical therapy procedures 93.1-93.3	75.5	47.2	96.3	127.4
Insertion of endotracheal tube 96.04	71.9	50.5	91.5	101.3
Endoscopy of small intestine without biopsy 45.11-45.13	69.0	43.1	86.6	120.5
Endoscopy of large intestine without biopsy 45.21-45.24	64.3	37.9	85.2	108.9

¹Includes data for surgical procedures not shown in table. All procedures except those listed under "nonsurgical procedures" in the appendix are surgical procedures.²See Table 29 for a comparison of number of procedures with number of discharges with procedures.³Includes data for nonsurgical procedures not shown in table. The ICD-9-CM codes for nonsurgical procedures are listed in the appendix.⁴Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.

Table 29. Number and rate of discharges from short-stay hospitals with a coronary artery bypass graft and number and rate of procedures, by sex, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Sex, age, and region	Discharges with procedures		Procedures	
	Number in thousands	Rate per 10,000 population ¹	Number in thousands	Rate per 10,000 population ¹
Total	249	8.5	427	14.6
Sex				
Male	174	12.1	303	21.1
Female	76	5.1	124	8.3
Age				
Under 15 years	*	*	*	*
15-44 years	9	0.8	17	1.4
45-64 years	111	15.7	195	27.6
65 years and over	129	35.6	215	59.3
65-74 years	71	38.3	116	62.8
75-84 years	54	41.3	92	71.0
85 years and over	5	10.3	7	15.1
Region				
Northeast	47	8.6	81	14.8
Midwest	52	8.0	89	13.6
South	100	9.5	170	16.2
West	50	7.5	87	13.0

* Figure does not meet standard of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.

NOTE: The number of procedures exceeds the number of discharges because these procedures are often assigned more than one code for a single operation. See "Methods" section.

Table 30. Number of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All procedures	45,023	2,310	15,110	11,697	15,905
Operations on the nervous system	1,291	236	398	318	339
Spinal tap	357	153	85	65	54
Operations on the endocrine system	103	*	32	41	28
Operations on the eye	71	9	19	20	24
Operations on the ear	45	22	12	*6	*6
Operations on the nose, mouth, and pharynx	309	63	115	62	68
Operations on the respiratory system	1,045	60	169	317	498
Bronchoscopy with or without biopsy	270	18	40	87	124
Operations on the cardiovascular system	6,993	226	675	2,511	3,581
Balloon angioplasty of coronary artery or coronary atherectomy	664	*	37	284	342
Insertion of coronary artery stent(s)	615	*	35	264	316
Coronary artery bypass graft ¹	427	*	17	195	215
Cardiac catheterization	1,297	11	101	547	638
Insertion, replacement, removal, and revision of pacemaker leads or device	362	*	8	46	306
Hemodialysis	607	*	79	230	296
Operations on the hemic and lymphatic system	341	24	61	117	138
Operations on the digestive system	5,824	261	1,372	1,716	2,475
Endoscopy of small intestine with or without biopsy	1,115	18	160	319	619
Endoscopy of large intestine with or without biopsy	596	5	89	155	347
Partial excision of large intestine	259	*	34	81	138
Appendectomy, excluding incidental	332	81	172	57	22
Cholecystectomy	435	4	146	137	148
Lysis of peritoneal adhesions	342	6	141	104	91
Operations on the urinary system	963	32	217	305	409
Cystoscopy with or without biopsy	145	*	27	40	74
Operations on the male genital organs	242	24	15	72	130
Prostatectomy	161	*	*	54	106
Operations on the female genital organs	2,050	10	1,188	621	231
Oophorectomy and salpingo-oophorectomy	451	*	175	213	60
Bilateral destruction or occlusion of fallopian tubes	357	*	354	*	*
Hysterectomy	617	*	310	248	59
Obstetrical procedures	6,971	*	6,947	16	...
Episiotomy with or without forceps or vacuum extraction	667	*	665	*	...
Artificial rupture of membranes	941	*	937	*	...
Medical induction of labor	573	*	572	*	...
Cesarean section	1,236	*	1,230	*5	...
Repair of current obstetric laceration	1,241	*	1,239	*	...
Operations on the musculoskeletal system	4,040	168	945	1,349	1,578
Partial excision of bone	172	11	52	65	44
Reduction of fracture	667	50	198	152	267
Open reduction of fracture with internal fixation	441	20	120	107	194
Excision or destruction of intervertebral disc	326	*	122	142	61
Total hip replacement	234	*	15	83	135
Total knee replacement	478	*	9	177	292
Operations on the integumentary system	1,498	*	440	468	436
Debridement of wound, infection, or burn	368	*	97	120	126
Miscellaneous diagnostic and therapeutic procedures and new technologies ²	13,239	1,012	2,505	3,758	5,964
Computerized axial tomography	828	54	167	212	395
Arteriography and angiocardiology using contrast material	2,110	19	204	834	1,053
Diagnostic ultrasound	813	*	133	223	405
Physical therapy procedures	415	6	40	95	274
Respiratory therapy	1,157	228	156	276	497
Insertion of endotracheal tube	511	45	67	138	261
Transfusion of blood and blood components	1,815	69	229	464	1,053
Injection or infusion of cancer chemotherapeutic substance	235	50	40	89	55

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 31. Rate of all-listed procedures for discharges from short-stay hospitals, by age and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	All ages	Under 15 years	15-44 years	45-64 years	65 years and over
All procedures	1,539.7	379.9	1,212.3	1,655.5	4,382.3
Operations on the nervous system 01-05	44.1	38.8	31.9	45.0	93.3
Spinal tap 03.31	12.2	25.2	6.8	9.2	14.9
Operations on the endocrine system 06-07	3.5	*	2.6	5.8	7.6
Operations on the eye 08-16	2.4	1.5	1.5	2.8	6.6
Operations on the ear 18-20	1.5	3.6	1.0	*0.8	*1.5
Operations on the nose, mouth, and pharynx 21-29	10.6	10.3	9.2	8.8	18.9
Operations on the respiratory system 30-34	35.7	9.9	13.6	44.9	137.2
Bronchoscopy with or without biopsy 33.21-33.24,33.27	9.2	3.0	3.2	12.3	34.1
Operations on the cardiovascular system 35-39	239.2	37.1	54.2	355.4	986.7
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	22.7	*	3.0	40.2	94.3
Insertion of coronary artery stent(s) 36.06-36.07	21.0	*	2.8	37.3	87.2
Coronary artery bypass graft ² 36.1	14.6	*	1.4	27.6	59.3
Cardiac catheterization 37.21-37.23	44.3	1.8	8.1	77.4	175.7
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	12.4	*	0.6	6.6	84.3
Hemodialysis 39.95	20.8	*	6.3	32.5	81.5
Operations on the hemic and lymphatic system 40-41	11.7	4.0	4.9	16.6	38.1
Operations on the digestive system 42-54	199.2	42.8	110.0	242.9	682.0
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	38.1	2.9	12.8	45.1	170.5
Endoscopy of large intestine with or without biopsy 45.21-45.25	20.4	0.9	7.1	22.0	95.6
Partial excision of large intestine 45.7	8.8	*	2.8	11.4	38.1
Appendectomy, excluding incidental 47.0	11.3	13.3	13.8	8.0	5.9
Cholecystectomy 51.2	14.9	0.6	11.7	19.3	40.9
Lysis of peritoneal adhesions 54.5	11.7	1.1	11.3	14.7	25.1
Operations on the urinary system 55-59	32.9	5.3	17.4	43.2	112.7
Cystoscopy with or without biopsy 57.31-57.33	5.0	*	2.2	5.7	20.5
Operations on the male genital organs 60-64	8.3	3.9	1.2	10.2	35.8
Prostatectomy 60.2-60.6	5.5	*	*	7.7	29.1
Operations on the female genital organs 65-71	70.1	1.6	95.3	87.9	63.8
Oophorectomy and salpingo-oophorectomy 65.3-65.6	15.4	*	14.1	30.2	16.7
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	12.2	*	28.4	*	*
Hysterectomy 68.3-68.7,68.9	21.1	*	24.9	35.1	16.2
Obstetrical procedures 72-75	238.4	*	557.3	2.3	...
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	22.8	*	53.3	*	...
Artificial rupture of membranes 73.0	32.2	*	75.2	*	...
Medical induction of labor 73.4	19.6	*	45.9	*	...
Cesarean section 74.0-74.2,74.4,74.99	42.3	*	98.7	*0.7	...
Repair of current obstetric laceration 75.5-75.6	42.4	*	99.4	*	...
Operations on the musculoskeletal system 76-84	138.2	27.7	75.8	190.9	434.8
Partial excision of bone 76.2-76.3,77.6-77.8	5.9	1.9	4.2	9.2	12.2
Reduction of fracture 76.7,79.0-79.3	22.8	8.3	15.9	21.5	73.6
Open reduction of fracture with internal fixation 79.3	15.1	3.3	9.7	15.2	53.4
Excision or destruction of intervertebral disc 80.5	11.2	*	9.8	20.1	16.8
Total hip replacement 81.51	8.0	*	1.2	11.7	37.3
Total knee replacement 81.54	16.3	*	0.7	25.1	80.3
Operations on the integumentary system 85-86	51.2	*	35.3	66.3	120.0
Debridement of wound, infection, or burn 86.22,86.28	12.6	*	7.8	17.0	34.7
Miscellaneous diagnostic and therapeutic procedures and new technologies ³ 87-99.00	452.7	166.4	201.0	531.8	1,643.3
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	28.3	8.9	13.4	29.9	109.0
Arteriography and angiocardiology using contrast material 88.4-88.5	72.1	3.1	16.4	118.0	290.1
Diagnostic ultrasound 88.7	27.8	*	10.7	31.6	111.7
Physical therapy procedures 93.1-93.3	14.2	1.0	3.2	13.4	75.5
Respiratory therapy 93.9,96.7	39.6	37.5	12.5	39.1	137.0
Insertion of endotracheal tube 96.04	17.5	7.4	5.3	19.6	71.9
Transfusion of blood and blood components 99.0	62.1	11.3	18.4	65.7	290.2
Injection or infusion of cancer chemotherapeutic substance 99.25	8.0	8.3	3.2	12.6	15.3

* Figure does not meet standard of reliability or precision.

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²See Table 29 for comparison of rate of procedures with rate of discharges with procedures.³Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 32. Number of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
All procedures	45,023	17,671	27,352
Operations on the nervous system 01-05	1,291	562	729
Spinal tap03.31	357	181	176
Operations on the endocrine system 06-07	103	26	76
Operations on the eye 08-16	71	45	26
Operations on the ear 18-20	45	31	13
Operations on the nose, mouth, and pharynx 21-29	309	198	110
Operations on the respiratory system 30-34	1,045	572	473
Bronchoscopy with or without biopsy 33.21-33.24,33.27	270	145	125
Operations on the cardiovascular system 35-39	6,993	3,985	3,009
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	664	438	226
Insertion of coronary artery stent(s) 36.06-36.07	615	412	203
Coronary artery bypass graft ¹ 36.1	427	303	124
Cardiac catheterization 37.21-37.23	1,297	771	526
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	362	179	183
Hemodialysis 39.95	607	297	311
Operations on the hemic and lymphatic system 40-41	341	160	180
Operations on the digestive system 42-54	5,824	2,447	3,377
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	1,115	495	620
Endoscopy of large intestine with or without biopsy 45.21-45.25	596	240	356
Partial excision of large intestine 45.7	259	114	145
Appendectomy, excluding incidental 47.0	332	182	150
Cholecystectomy 51.2	435	136	299
Lysis of peritoneal adhesions 54.5	342	74	268
Operations on the urinary system 55-59	963	443	519
Cystoscopy with or without biopsy 57.31-57.33	145	68	77
Operations on the male genital organs 60-64	242	242	...
Prostatectomy 60.2-60.6	161	161	...
Operations on the female genital organs 65-71	2,050	...	2,050
Oophorectomy and salpingo-oophorectomy 65.3-65.6	451	...	451
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	357	...	357
Hysterectomy 68.3-68.7,68.9	617	...	617
Obstetrical procedures 72-75	6,971	...	6,971
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	667	...	667
Artificial rupture of membranes 73.0	941	...	941
Medical induction of labor 73.4	573	...	573
Cesarean section 74.0-74.2,74.4,74.99	1,236	...	1,236
Repair of current obstetric laceration 75.5-75.6	1,241	...	1,241
Operations on the musculoskeletal system 76-84	4,040	1,902	2,138
Partial excision of bone 76.2-76.3,77.6-77.8	172	89	84
Reduction of fracture 76.7,79.0-79.3	667	321	347
Open reduction of fracture with internal fixation 79.3	441	197	244
Excision or destruction of intervertebral disc 80.5	326	162	164
Total hip replacement 81.51	234	99	135
Total knee replacement 81.54	478	167	311
Operations on the integumentary system 85-86	1,498	701	797
Debridement of wound, infection, or burn 86.22,86.28	368	207	161
Miscellaneous diagnostic and therapeutic procedures and new technologies ² 87-99.00	13,239	6,357	6,882
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	828	389	439
Arteriography and angiocardiology using contrast material 88.4-88.5	2,110	1,171	938
Diagnostic ultrasound 88.7	813	368	444
Physical therapy procedures 93.1-93.3	415	168	247
Respiratory therapy 93.9,96.7	1,157	588	569
Insertion of endotracheal tube 96.04	511	258	253
Transfusion of blood and blood components 99.0	1,815	759	1,055
Injection or infusion of cancer chemotherapeutic substance 99.25	235	131	104

... Category not applicable.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 33. Rate of all-listed procedures for discharges from short-stay hospitals, by sex and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	Both sexes	Male	Female
All procedures	1,539.7	1,231.6	1,836.5
Operations on the nervous system 01-05	44.1	39.1	49.0
Spinal tap03.31	12.2	12.6	11.8
Operations on the endocrine system 06-07	3.5	1.8	5.1
Operations on the eye 08-16	2.4	3.2	1.7
Operations on the ear 18-20	1.5	2.2	0.9
Operations on the nose, mouth, and pharynx 21-29	10.6	13.8	7.4
Operations on the respiratory system 30-34	35.7	39.9	31.7
Bronchoscopy with or without biopsy 33.21-33.24,33.27	9.2	10.1	8.4
Operations on the cardiovascular system 35-39	239.2	277.7	202.0
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	22.7	30.5	15.1
Insertion of coronary artery stent(s) 36.06-36.07	21.0	28.7	13.6
Coronary artery bypass graft ² 36.1	14.6	21.1	8.3
Cardiac catheterization 37.21-37.23	44.3	53.7	35.3
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	12.4	12.5	12.3
Hemodialysis 39.95	20.8	20.7	20.9
Operations on the hemic and lymphatic system 40-41	11.7	11.2	12.1
Operations on the digestive system 42-54	199.2	170.5	226.7
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	38.1	34.5	41.6
Endoscopy of large intestine with or without biopsy 45.21-45.25	20.4	16.7	23.9
Partial excision of large intestine 45.7	8.8	7.9	9.7
Appendectomy, excluding incidental 47.0	11.3	12.7	10.0
Cholecystectomy 51.2	14.9	9.5	20.0
Lysis of peritoneal adhesions 54.5	11.7	5.2	18.0
Operations on the urinary system 55-59	32.9	30.9	34.9
Cystoscopy with or without biopsy 57.31-57.33	5.0	4.7	5.2
Operations on the male genital organs 60-64	8.3	16.8	...
Prostatectomy 60.2-60.6	5.5	11.2	...
Operations on the female genital organs 65-71	70.1	...	137.6
Oophorectomy and salpingo-oophorectomy 65.3-65.6	15.4	...	30.3
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	12.2	...	24.0
Hysterectomy 68.3-68.7,68.9	21.1	...	41.4
Obstetrical procedures 72-75	238.4	...	468.1
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	22.8	...	44.8
Artificial rupture of membranes 73.0	32.2	...	63.2
Medical induction of labor 73.4	19.6	...	38.5
Cesarean section 74.0-74.2,74.4,74.99	42.3	...	83.0
Repair of current obstetric laceration 75.5-75.6	42.4	...	83.3
Operations on the musculoskeletal system 76-84	138.2	132.5	143.6
Partial excision of bone 76.2-76.3,77.6-77.8	5.9	6.2	5.6
Reduction of fracture 76.7,79.0-79.3	22.8	22.4	23.3
Open reduction of fracture with internal fixation 79.3	15.1	13.7	16.4
Excision or destruction of intervertebral disc 80.5	11.2	11.3	11.0
Total hip replacement 81.51	8.0	6.9	9.0
Total knee replacement 81.54	16.3	11.6	20.9
Operations on the integumentary system 85-86	51.2	48.8	53.5
Debridement of wound, infection, or burn 86.22,86.28	12.6	14.4	10.8
Miscellaneous diagnostic and therapeutic procedures and new technologies ³ 87-99.00	452.7	443.0	462.1
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	28.3	27.1	29.5
Arteriography and angiocardiology using contrast material 88.4-88.5	72.1	81.6	63.0
Diagnostic ultrasound 88.7	27.8	25.7	29.8
Physical therapy procedures 93.1-93.3	14.2	11.7	16.6
Respiratory therapy 93.9,96.7	39.6	41.0	38.2
Insertion of endotracheal tube 96.04	17.5	18.0	17.0
Transfusion of blood and blood components 99.0	62.1	52.9	70.9
Injection or infusion of cancer chemotherapeutic substance 99.25	8.0	9.2	7.0

... Category not applicable.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²See Table 29 for comparison of rate of procedures with rate of discharges with procedures.³Chapter 00, "Procedures and interventions not else where classified," is included in this category; see "Use of Tables."

Table 34. Number of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures	45,023	9,935	9,295	16,280	9,513
Operations on the nervous system 01-05	1,291	275	298	427	291
Spinal tap 03.31	357	61	75	137	83
Operations on the endocrine system 06-07	103	25	18	33	27
Operations on the eye 08-16	71	14	10	31	15
Operations on the ear 18-20	45	10	*	16	*
Operations on the nose, mouth, and pharynx 21-29	309	75	49	116	68
Operations on the respiratory system 30-34	1,045	229	211	387	218
Bronchoscopy with or without biopsy 33.21-33.24,33.27	270	64	55	101	50
Operations on the cardiovascular system 35-39	6,993	1,388	1,440	2,731	1,434
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	664	119	170	242	133
Insertion of coronary artery stent(s) 36.06-36.07	615	105	163	218	129
Coronary artery bypass graft ¹ 36.1	427	81	89	170	87
Cardiac catheterization 37.21-37.23	1,297	235	275	533	254
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	362	105	61	125	71
Hemodialysis 39.95	607	125	101	246	135
Operations on the hemic and lymphatic system 40-41	341	74	67	126	74
Operations on the digestive system 42-54	5,824	1,230	1,210	2,194	1,190
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	1,115	239	242	424	209
Endoscopy of large intestine with or without biopsy 45.21-45.25	596	130	127	223	116
Partial excision of large intestine 45.7	259	64	62	88	44
Appendectomy, excluding incidental 47.0	332	70	69	103	90
Cholecystectomy 51.2	435	89	87	164	94
Lysis of peritoneal adhesions 54.5	342	64	71	145	62
Operations on the urinary system 55-59	963	225	204	342	192
Cystoscopy with or without biopsy 57.31-57.33	145	37	33	53	22
Operations on the male genital organs 60-64	242	50	54	94	43
Prostatectomy 60.2-60.6	161	35	37	54	34
Operations on the female genital organs 65-71	2,050	296	424	916	413
Oophorectomy and salpingo-oophorectomy 65.3-65.6	451	68	111	188	84
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	357	40	58	187	72
Hysterectomy 68.3-68.7,68.9	617	86	139	269	122
Obstetrical procedures 72-75	6,971	1,056	1,490	2,633	1,793
Episiotomy with or without forceps or vacuum extraction 72.1,72.2,72.31,72.71,73.6	667	88	137	288	154
Artificial rupture of membranes 73.0	941	125	255	333	229
Medical induction of labor 73.4	573	79	118	215	160
Cesarean section 74.0-74.2,74.4,74.99	1,236	182	221	533	299
Repair of current obstetric laceration 75.5-75.6	1,241	200	252	425	364
Operations on the musculoskeletal system 76-84	4,040	786	909	1,460	885
Partial excision of bone 76.2-76.3,77.6-77.8	172	35	36	61	40
Reduction of fracture 76.7,79.0-79.3	667	137	141	243	147
Open reduction of fracture with internal fixation 79.3	441	87	97	164	94
Excision or destruction of intervertebral disc 80.5	326	48	74	127	77
Total hip replacement 81.51	234	61	58	69	45
Total knee replacement 81.54	478	100	142	158	77
Operations on the integumentary system 85-86	1,498	298	283	682	235
Debridement of wound, infection, or burn 86.22,86.28	368	73	74	171	50
Miscellaneous diagnostic and therapeutic procedures and new technologies ² 87-99,00	13,239	3,902	2,619	4,092	2,626
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	828	400	*	*	150
Arteriography and angiocardiology using contrast material 88.4-88.5	2,110	409	470	831	399
Diagnostic ultrasound 88.7	813	325	133	205	150
Physical therapy procedures 93.1-93.3	415	123	*	153	*
Respiratory therapy 93.9,96.7	1,157	299	233	344	282
Insertion of endotracheal tube 96.04	511	107	103	190	110
Transfusion of blood and blood components 99.0	1,815	354	376	718	367
Injection or infusion of cancer chemotherapeutic substance 99.25	235	60	27	99	50

* Figure does not meet standard of reliability or precision.

¹See Table 29 for comparison of number of procedures with number of discharges with procedures.²Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 35. Rate of all-listed procedures for discharges from short-stay hospitals, by geographic region and procedure category: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*]

Procedure category and ICD-9-CM code	United States	North-east	Midwest	South	West
All procedures	1,539.7	1,822.4	1,416.0	1,547.2	1,419.1
Operations on the nervous system 01-05	44.1	50.5	45.3	40.6	43.4
Spinal tap03.31	12.2	11.2	11.4	13.1	12.4
Operations on the endocrine system 06-07	3.5	4.7	2.8	3.1	4.0
Operations on the eye08-16	2.4	2.6	1.6	3.0	2.3
Operations on the ear18-20	1.5	1.9	*	1.5	*
Operations on the nose, mouth, and pharynx 21-29	10.6	13.7	7.5	11.1	10.2
Operations on the respiratory system 30-34	35.7	42.1	32.2	36.8	32.5
Bronchoscopy with or without biopsy 33.21-33.24,33.27	9.2	11.7	8.4	9.6	7.4
Operations on the cardiovascular system 35-39	239.2	254.6	219.4	259.6	213.9
Balloon angioplasty of coronary artery or coronary atherectomy 36.01-36.02,36.05	22.7	21.9	25.8	23.0	19.8
Insertion of coronary artery stent(s) 36.06-36.07	21.0	19.3	24.8	20.7	19.2
Coronary artery bypass graft ² 36.1	14.6	14.8	13.6	16.2	13.0
Cardiac catheterization 37.21-37.23	44.3	43.1	41.8	50.6	37.9
Insertion, replacement, removal, and revision of pacemaker leads or device 37.7-37.8	12.4	19.3	9.2	11.9	10.6
Hemodialysis39.95	20.8	23.0	15.4	23.4	20.2
Operations on the hemic and lymphatic system 40-41	11.7	13.5	10.2	12.0	11.0
Operations on the digestive system 42-54	199.2	225.7	184.3	208.5	177.5
Endoscopy of small intestine with or without biopsy 45.11-45.14,45.16	38.1	43.8	36.9	40.3	31.2
Endoscopy of large intestine with or without biopsy 45.21-45.25	20.4	23.9	19.4	21.2	17.3
Partial excision of large intestine 45.7	8.8	11.8	9.4	8.4	6.6
Appendectomy, excluding incidental 47.0	11.3	12.8	10.6	9.7	13.5
Cholecystectomy 51.2	14.9	16.4	13.2	15.6	14.0
Lysis of peritoneal adhesions 54.5	11.7	11.7	10.9	13.8	9.2
Operations on the urinary system 55-59	32.9	41.2	31.2	32.5	28.6
Cystoscopy with or without biopsy 57.31-57.33	5.0	6.7	5.0	5.1	3.4
Operations on the male genital organs 60-64	8.3	9.2	8.3	9.0	6.4
Prostatectomy 60.2-60.6	5.5	6.5	5.7	5.1	5.1
Operations on the female genital organs 65-71	70.1	54.4	64.6	87.0	61.7
Oophorectomy and salpingo-oophorectomy 65.3-65.6	15.4	12.4	16.9	17.9	12.6
Bilateral destruction or occlusion of fallopian tubes 66.2-66.3	12.2	7.3	8.9	17.8	10.8
Hysterectomy 68.3-68.7,68.9	21.1	15.9	21.2	25.6	18.2
Obstetrical procedures 72-75	238.4	193.7	227.0	250.2	267.5
Episiotomy with or without forceps or vacuum extraction 72.1,72.21,72.31,72.71,73.6	22.8	16.1	20.9	27.4	23.0
Artificial rupture of membranes 73.0	32.2	22.9	38.8	31.6	34.1
Medical induction of labor 73.4	19.6	14.5	17.9	20.5	23.9
Cesarean section 74.0-74.2,74.4,74.99	42.3	33.5	33.7	50.7	44.6
Repair of current obstetric laceration 75.5-75.6	42.4	36.7	38.4	40.4	54.3
Operations on the musculoskeletal system 76-84	138.2	144.2	138.5	138.7	132.0
Partial excision of bone 76.2-76.3,77.6-77.8	5.9	6.4	5.5	5.8	6.0
Reduction of fracture 76.7,79.0-79.3	22.8	25.1	21.4	23.1	22.0
Open reduction of fracture with internal fixation 79.3	15.1	15.9	14.8	15.6	14.0
Excision or destruction of intervertebral disc 80.5	11.2	8.7	11.3	12.1	11.6
Total hip replacement81.51	8.0	11.2	8.9	6.6	6.7
Total knee replacement81.54	16.3	18.4	21.6	15.1	11.5
Operations on the integumentary system 85-86	51.2	54.7	43.1	64.8	35.0
Debridement of wound, infection, or burn 86.22,86.28	12.6	13.3	11.3	16.3	7.4
Miscellaneous diagnostic and therapeutic procedures and new technologies ³ 87-99,00	452.7	715.7	398.9	388.9	391.8
Computerized axial tomography 87.03,87.41,87.71,88.01,88.38	28.3	73.3	*	*	22.4
Arteriography and angiocardiology using contrast material 88.4-88.5	72.1	75.1	71.6	79.0	59.5
Diagnostic ultrasound 88.7	27.8	59.5	20.3	19.5	22.4
Physical therapy procedures 93.1-93.3	14.2	22.6	*	14.5	*
Respiratory therapy 93.9,96.7	39.6	54.9	35.5	32.7	42.0
Insertion of endotracheal tube96.04	17.5	19.7	15.7	18.1	16.5
Transfusion of blood and blood components 99.0	62.1	64.9	57.2	68.2	54.8
Injection or infusion of cancer chemotherapeutic substance99.25	8.0	11.0	4.1	9.4	7.4

* Figure does not meet standard of reliability or precision.

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²See Table 29 for comparison of number of procedures with number of discharges with procedures.³Chapter 00, "Procedures and interventions not elsewhere classified," is included in this category; see "Use of Tables."

Table 36. Number, percent distribution, and rate of discharges and days of care, with average length of stay for females with deliveries discharged from short-stay hospitals, by type of delivery, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Discharges			Days of care			Average length of stay in days
	Number in thousands	Percent distribution	Rate per 1,000 population ¹	Number in thousands	Percent distribution	Rate per 1,000 population ¹	
All deliveries							
10–54 years	4,136	100.0	44.3	10,791	100.0	115.6	2.6
10–19 years	438	10.6	21.5	1,127	10.4	55.3	2.6
20–24 years	1,034	25.0	102.4	2,585	24.0	255.9	2.5
25–29 years	1,109	26.8	116.4	2,799	25.9	293.8	2.5
30–34 years	968	23.4	95.8	2,610	24.2	258.2	2.7
35–39 years	470	11.4	44.9	1,309	12.1	125.1	2.8
40–54 years	117	2.8	3.6	361	3.3	11.0	3.1
All regions	4,136	100.0	44.3	10,791	100.0	115.6	2.6
Northeast	589	14.2	33.9	1,695	15.7	97.6	2.9
Midwest	802	19.4	38.4	2,131	19.8	102.0	2.7
South	1,679	40.6	50.0	4,370	40.5	130.1	2.6
West	1,066	25.8	49.6	2,595	24.0	120.8	2.4
Vaginal deliveries ²							
10–54 years	2,900	100.0	31.1	6,208	100.0	66.5	2.1
10–19 years	350	12.1	17.2	789	12.7	38.7	2.3
20–24 years	765	26.4	75.8	1,632	26.3	161.6	2.1
25–29 years	779	26.9	81.8	1,643	26.5	172.5	2.1
30–34 years	652	22.5	64.5	1,382	22.3	136.7	2.1
35–39 years	289	10.0	27.6	607	9.8	58.0	2.1
40–54 years	66	2.3	2.0	155	2.5	4.7	2.4
All regions	2,900	100.0	31.1	6,208	100.0	66.5	2.1
Northeast	407	14.0	23.4	955	15.4	55.0	2.3
Midwest	580	20.0	27.8	1,247	20.1	59.7	2.1
South	1,146	39.5	34.1	2,488	40.1	74.1	2.2
West	767	26.5	35.7	1,519	24.5	70.7	2.0
Cesarean deliveries ³							
10–54 years	1,236	100.0	13.2	4,583	100.0	49.1	3.7
10–19 years	88	7.2	4.3	339	7.4	16.6	3.8
20–24 years	269	21.8	26.6	952	20.8	94.3	3.5
25–29 years	330	26.7	34.6	1,156	25.2	121.4	3.5
30–34 years	316	25.6	31.3	1,228	26.8	121.5	3.9
35–39 years	181	14.7	17.3	702	15.3	67.1	3.9
40–54 years	51	4.2	1.6	205	4.5	6.3	4.0
All regions	1,236	100.0	13.2	4,583	100.0	49.1	3.7
Northeast	182	14.8	10.5	740	16.2	42.6	4.1
Midwest	221	17.9	10.6	884	19.3	42.3	4.0
South	533	43.1	15.9	1,882	41.1	56.0	3.5
West	299	24.2	13.9	1,076	23.5	50.1	3.6

¹Rates were calculated using U.S. Census Bureau 2000-based postcensal estimates of the civilian population as of July 1, 2004.²Discharges of females with deliveries except those with cesarean deliveries.³Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 37. Number and percent distribution of females with deliveries discharged from short-stay hospitals, by length of stay and type of delivery: United States, selected years 1980–2004[Discharges of inpatients from nonfederal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Length of stay	1980	1985	1990	1995	2000	2004
Number in thousands						
All deliveries	3,762	3,854	4,025	3,766	3,738	4,136
Less than 1 day	23	34	37	36	16	*11
1 day	216	373	623	1,368	551	517
2 days	747	1,037	1,649	1,418	1,863	1,962
3 days	1,205	1,140	900	606	863	1,085
4 days	691	586	468	193	292	372
5–7 days	726	598	282	100	112	137
8 days or more	154	86	67	46	41	51
Vaginal deliveries ¹	3,143	2,977	3,080	2,981	2,882	2,900
Less than 1 day	23	33	35	34	14	*10
1 day	215	371	620	1,362	546	512
2 days	743	1,030	1,624	1,271	1,726	1,729
3 days	1,185	1,054	618	228	500	554
4 days	618	338	106	39	53	58
5–7 days	317	125	55	27	25	20
8 days or more	42	26	23	21	17	16
Cesarean deliveries ²	619	877	945	785	855	1,236
Less than 1 day	*	*	*	*	*	*
1 day	*	*	*	*6	*	*
2 days	*	*	25	147	137	233
3 days	20	86	283	379	363	531
4 days	73	249	361	154	239	314
5–7 days	409	473	227	73	87	117
8 days or more	112	60	44	25	24	35
Percent distribution						
All deliveries	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.6	0.9	0.9	1.0	0.4	*0.3
1 day	5.7	9.7	15.5	36.3	14.7	12.5
2 days	19.9	26.9	41.0	37.7	49.9	47.5
3 days	32.0	29.6	22.4	16.1	23.1	26.2
4 days	18.4	15.2	11.6	5.1	7.8	9.0
5–7 days	19.3	15.5	7.0	2.7	3.0	3.3
8 days or more	4.1	2.2	1.7	1.2	1.1	1.2
Vaginal deliveries ¹	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	0.7	1.1	1.1	1.1	0.5	*0.4
1 day	6.8	12.5	20.1	45.7	19.0	17.6
2 days	23.6	34.6	52.7	42.6	59.9	59.6
3 days	37.7	35.4	20.1	7.6	17.3	19.1
4 days	19.7	11.4	3.4	1.3	1.8	2.0
5–7 days	10.1	4.2	1.8	0.9	0.9	0.7
8 days or more	1.3	0.9	0.7	0.7	0.6	0.6
Cesarean deliveries ²	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	*	*	*	*	*	*
1 day	*	*	*	*0.8	*	*
2 days	*	*	2.6	18.7	16.0	18.9
3 days	3.2	9.8	29.9	48.3	42.4	42.9
4 days	11.8	28.4	38.2	19.6	27.9	25.4
5–7 days	66.1	53.9	24.0	9.3	10.2	9.5
8 days or more	18.1	6.8	4.7	3.2	2.8	2.8
Average length of stay in days						
All deliveries	3.8	3.3	2.8	2.1	2.5	2.6
Vaginal deliveries ¹	3.2	2.7	2.3	1.7	2.1	2.1
Cesarean deliveries ²	6.5	5.2	4.5	3.6	3.7	3.7

* Figure does not meet standard of reliability or precision.

¹Discharges of females with deliveries except those with cesarean deliveries.²Discharges of females with deliveries with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 38. Number and rate of selected obstetrical procedures for females with deliveries discharged from short-stay hospitals: United States, selected years 1980–2004

[Discharges of inpatients from nonfederal hospitals. Procedure groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Data are for selected procedures performed on discharges with code V27, females with deliveries]

Procedure category and ICD–9–CM code	1980	1985	1990	1995	2000	2004
Number of procedures in thousands						
All obstetrical procedures 72–75	3,526	4,239	6,660	6,282	6,145	6,908
Repair of current obstetric laceration 75.5–75.6	350	543	789	961	1,134	1,240
Cesarean section 74.0–74.2,74.4,74.99	619	877	945	785	855	1,236
Manually assisted delivery 73.5	70	193	750	866	898	1,000
Artificial rupture of membranes 73.0	120	315	691	752	833	941
Episiotomy 72.1,72.21,72.31,72.71,73.6	2,012	1,820	1,714	1,410	944	667
Medical induction of labor 73.4	42	66	225	366	435	567
Vacuum extraction ¹ 72.7	22	66	190	279	245	249
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	555	376	269	176	117	68
Rate per 100 deliveries ²						
All obstetrical procedures 72–75	93.7	110.0	165.5	166.8	164.4	167.0
Cesarean section 74.0–74.2,74.4,74.99	16.5	22.7	23.5	20.8	22.9	29.9
Artificial rupture of membranes 73.0	3.2	8.2	17.2	20.0	22.3	22.8
Medical induction of labor 73.4	1.1	1.7	5.6	9.7	11.7	13.7
Rate per 100 vaginal deliveries ³						
Repair of current obstetric laceration 75.5–75.6	11.1	18.1	25.5	32.0	39.2	42.6
Manually assisted delivery 73.5	2.2	6.5	24.3	29.0	31.1	34.5
Episiotomy 72.1,72.21,72.31,72.71,73.6	64.0	61.1	55.6	47.2	32.7	23.0
Vacuum extraction ¹ 72.7	0.7	2.2	6.1	9.2	8.4	8.2
Forceps delivery ¹ 72.0–72.4,72.51,72.53,72.6,73.3	17.6	12.5	8.6	5.8	4.0	2.3

¹Forceps delivery and vacuum extraction with episiotomy are also included in the episiotomy category.

²Procedures per 100 discharges of females with deliveries.

³Procedures for females with vaginal deliveries per 100 discharges with vaginal deliveries. Vaginal deliveries are all deliveries except those with a cesarean section.

Table 39. Number and rate of females with deliveries discharged from short-stay hospitals, by type of delivery: United States, selected years 1980–2004

[Discharges of inpatients from nonfederal hospitals. Data are for discharges with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Type of delivery	1980	1985	1990	1995	2000	2004
Number in thousands						
Females with deliveries	3,762	3,854	4,025	3,766	3,738	4,136
Previous cesarean delivery	192	324	426	413	447	559
No previous cesarean delivery	3,571	3,530	3,600	3,353	3,290	3,577
Vaginal deliveries	3,143	2,977	3,080	2,981	2,882	2,900
Vaginal birth after cesarean delivery (VBAC)	*6	21	87	147	123	61
Other vaginal deliveries	3,137	2,956	2,993	2,835	2,759	2,839
Cesarean deliveries	619	877	945	785	855	1,236
No previous cesarean delivery (primary)	434	574	606	519	531	738
Previous cesarean delivery (repeat)	185	303	339	266	324	498
Rate per 100 deliveries						
Cesarean deliveries per 100 deliveries	16.5	22.7	23.5	20.8	22.9	29.9
Primary cesarean deliveries per 100 deliveries without previous cesarean delivery	12.1	16.3	16.8	15.5	16.1	20.6
Vaginal birth after cesarean deliveries (VBAC) per 100 deliveries with previous cesarean delivery	*3.4	6.6	20.4	35.5	27.6	10.9

* Figure does not meet standard of reliability or precision.

NOTE: Previous cesarean deliveries are discharges with ICD–9–CM diagnostic code 654.2. Cesarean deliveries are discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99.

Table 40. Number and rate of females with deliveries discharged from short-stay hospitals by type of delivery, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Data are for discharges 10–54 years of age with *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM) code V27, females with deliveries]

Age and region	Cesarean deliveries		
	Total ¹	Primary ²	VBAC ^{3,4}
		Number in thousands	
10–54 years	1,236	738	61
10–19 years	88	77	*
20–24 years	269	174	12
25–29 years	330	190	17
30–34 years	316	173	15
35–39 years	181	95	12
40–54 years	51	30	*3
All regions	1,236	738	61
Northeast	182	115	13
Midwest	221	133	11
South	533	312	17
West	299	178	20
		Rate per 100 deliveries	
10–54 years	29.9	20.6	10.9
10–19 years	20.2	18.1	*
20–24 years	26.0	18.8	11.2
25–29 years	29.7	20.0	10.6
30–34 years	32.7	21.4	9.6
35–39 years	38.6	25.5	12.3
40–54 years	43.9	32.1	*12.1
All regions	29.9	20.6	10.9
Northeast	31.0	22.6	16.1
Midwest	27.6	19.0	11.0
South	31.8	21.7	7.3
West	28.0	19.2	14.1

* Figure does not meet standard of reliability or precision.

¹Discharges with ICD–9–CM procedure codes 74.0–74.2, 74.4, or 74.99. Rate per 100 total deliveries.²Cesarean deliveries without ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries without previous cesarean delivery.³VBAC is vaginal birth after cesarean delivery.⁴Vaginal deliveries with ICD–9–CM diagnostic code 654.2. Rate per 100 deliveries with previous cesarean delivery.

Table 41. Number, percent distribution, and average length of stay for newborn infants discharged from short-stay hospitals by sex, geographic region, and health status: United States, 2004

[Infants born in nonfederal hospitals]

Sex, region, and health status	Discharges		Average length of stay in days
	Number in thousands	Percent distribution	
All newborn infants	3,908	100.0	3.3
Sex			
Male	2,020	51.7	3.5
Female	1,888	48.3	3.2
Region			
Northeast	702	18.0	3.4
Midwest	803	20.5	2.9
South	1,403	35.9	3.6
West	1,000	25.6	3.2
Health status			
Well ¹	2,241	57.4	2.1
Male	1,116	28.6	2.1
Female	1,125	28.8	2.1
Sick ²	1,667	42.6	5.0
Male	903	23.1	5.2
Female	763	19.5	4.8

¹Without any illness or risk-related diagnoses.²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.**Table 42. Number and percent distribution of newborn infants discharged from short-stay hospitals, by length of stay, and average length of stay, by health status: United States, selected years 1980-2004**

[Infants born in nonfederal hospitals]

Length of stay and health status	1980	1985	1990	1995	2000	2004
Length of stay						
	Number in thousands					
All newborn infants	3,824	3,794	3,869	3,631	3,643	3,908
Less than 1 day	69	84	79	87	45	47
1 day	270	428	693	1,523	722	700
2 days	842	1,057	1,624	1,145	1,877	1,933
3 days	1,175	1,034	636	486	546	663
4 days	618	528	447	134	205	273
5-7 days	633	497	232	106	94	118
8 days or more	218	167	157	149	153	175
Percent distribution						
All newborn infants	100.0	100.0	100.0	100.0	100.0	100.0
Less than 1 day	1.8	2.2	2.0	2.4	1.2	1.2
1 day	7.1	11.3	17.9	41.9	19.8	17.9
2 days	22.0	27.9	42.0	31.5	51.5	49.5
3 days	30.7	27.3	16.4	13.4	15.0	17.0
4 days	16.2	13.9	11.6	3.7	5.6	7.0
5-7 days	16.5	13.1	6.0	2.9	2.6	3.0
8 days or more	5.7	4.4	4.1	4.1	4.2	4.5
Health status						
	Average length of stay in days					
All newborn infants	4.3	3.8	3.3	2.8	3.2	3.3
Well ¹	3.2	2.8	2.3	1.7	2.1	2.1
Sick ²	7.1	5.7	4.8	4.4	5.1	5.0

¹Without any illness or risk-related diagnoses.²With at least one illness or risk-related diagnosis, ICD-9-CM codes 001-999.

Table 43. Number of all-listed diagnoses for sick newborn infants discharged from short-stay hospitals, by sex and selected diagnostic categories: United States, 2004[Infants born in nonfederal hospitals. Diagnostic groupings and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM)]

Diagnostic category and ICD-9-CM code	Both sexes	Male	Female
	Number in thousands		
Sick newborn diagnoses 001-999	3,404	1,905	1,499
Hemolytic disease of fetus or newborn, due to isoimmunization, and other perinatal jaundice 773-774	714	379	335
Hemolytic disease due to ABO isoimmunization 773.1	51	24	27
Neonatal jaundice associated with preterm delivery 774.2	145	79	66
Unspecified fetal and neonatal jaundice 774.6	497	266	231
Respiratory conditions of fetus and newborn 768-770	391	226	165
Respiratory distress syndrome 769	83	46	38
Transitory tachypnea of newborn 770.6	117	68	49
Disorders relating to short gestation and low birthweight (prematurity) 765.0-765.1	301	159	143
Congenital anomalies 740-759	275	152	122
Congenital anomalies of circulatory system 745-747	79	43	37
Congenital anomalies of musculoskeletal system 754-756	47	22	26
Congenital anomalies of the integument (skin) 757	58	25	33
Disorders relating to long gestation and high birthweight 766	237	151	86
Endocrine and metabolic disturbances specific to the fetus and newborn 775	134	81	53
Neonatal hypoglycemia 775.6	60	39	21
Maternal causes of perinatal morbidity and mortality 760-763	125	61	64
Birth trauma 767	95	56	39
Injuries to scalp 767.1	70	43	27
Conditions involving the integument (skin) and temperature regulation of fetus and newborn 778	94	62	32
Infections specific to the perinatal period 771	86	55	31
Septicemia (sepsis) of newborn 771.81	70	46	24
Feeding problems in newborn 779.3	85	48	38
Slow fetal growth and fetal malnutrition 764	50	18	32
Fetal and neonatal hemorrhage 772	51	30	21

Table 44. Number and percentage of male newborn infants circumcised during hospitalization, by geographic region: United States, selected years 1980-2004[Infants born in nonfederal hospitals with *International Classification of Diseases, 9th Revision, Clinical Modification*, procedure code 64.0]

Region	1980	1985	1990	1995	2000	2004
	Number circumcised in thousands					
United States	1,261	1,162	1,169	1,200	1,166	1,160
Northeast	253	221	238	249	256	241
Midwest	403	331	341	317	330	333
South	379	393	396	454	438	423
West	226	218	195	180	143	163
	Percentage circumcised					
United States	64.7	59.5	59.0	64.1	62.4	57.4
Northeast	67.4	65.2	62.6	68.3	64.6	66.4
Midwest	75.9	70.5	76.0	79.8	81.4	79.5
South	56.0	56.0	57.1	66.1	63.9	58.5
West	61.8	49.0	42.4	42.6	37.3	31.7

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
All codes	34,864	13,844	21,020	2,571	10,800	8,323	13,171	7,388	7,680	13,239	6,557
001-139	949	441	507	155	172	212	409	196	204	354	195
003	*6	*	*	*	*	*	*	*	*	*	*
008	160	63	97	51	22	26	62	34	39	62	26
008.4	68	24	44	*	*9	12	44	18	18	22	10
008.45	64	22	42	*	*7	11	43	16	17	21	9
008.6	30	16	14	29	*	*	*	*	*7	15	*6
008.61	28	16	13	28	*	*	*	*	*7	14	*5
008.8	60	23	37	18	14	13	16	12	14	24	10
009	18	*7	11	*	*8	*	*	*	*	*7	*
009.0	10	*	*6	*	*	*	*	*	*	*	*
009.1	*6	*	*	*	*	*	*	*	*	*	*
011	*7	*	*	*	*	*	*	*	*	*	*
034	14	*7	*7	*8	*	*	*	*	*	*	*
034.0	13	*7	*6	*7	*	*	*	*	*	*	*
038	410	188	222	*7	28	90	285	80	102	138	90
038.0	22	12	9	*	*	*	18	*6	*5	*6	*5
038.1	56	29	27	*	*	15	38	11	12	18	15
038.11	35	21	14	*	*	*9	24	*7	*7	11	10
038.19	16	*7	*9	*	*	*	10	*	*	*5	*
038.2	*5	*	*	—	*	*	*	*	*	*	*
038.4	97	40	58	*	*6	24	66	22	26	29	20
038.40	*6	*	*	*	*	*	*	*	*	*	*
038.42	56	17	39	*	*	14	40	13	15	15	13
038.43	*7	*	*	*	*	*	*	*	*	*	*
038.49	24	13	12	*	*	*6	16	*5	*6	10	*
038.8	17	9	*7	*	*	*	11	*	*	*8	*
038.9	209	94	115	*	14	43	148	37	53	75	45
041	10	*	*6	*	*	*	*	*	*	*	*
042	65	41	24	*	38	25	*	16	*7	37	*
047	44	22	22	22	16	*5	*	*6	*7	20	12
047.8	*6	*	*	*	*	*	—	*	*	*	*
047.9	38	20	18	18	14	*	*	*5	*6	17	10
053	17	*	13	*	*	*6	9	*	*	*6	*
053.1	*5	*	*	*	*	*	*	*	*	*	*
053.9	*8	*	*6	*	*	*	*	*	*	*	*
054	*7	*	*	*	*	*	*	*	*	*	*
070	25	14	11	*	*6	15	*	*5	*	11	*
070.5	12	*7	*5	—	*	*9	*	*	*	*	*
070.51	*7	*	*	—	*	*5	*	*	*	*	*
070.54	*5	*	*	—	*	*	*	*	*	*	*
075	10	*6	*	*	*6	*	—	*	*	*	*
079	59	29	30	31	13	*6	*9	16	9	23	12
079.9	53	27	26	28	12	*6	*9	14	*8	21	10
079.99	53	27	26	28	12	*6	*9	14	*8	21	10
112	19	*7	12	*	*	*5	10	*	*	*7	*5
112.8	*6	*	*	*	*	*	*	*	*	*	*
135	10	*	*7	*	*	*6	*	*	*	*	*
140-239	1,640	655	985	38	286	606	709	373	343	620	303
150	14	12	*	—	*	*5	*7	*	*	*	*
150.5	*6	*5	*	—	*	*	*	*	*	*	*
150.9	*5	*	*	—	*	*	*	*	*	*	*
151	23	15	*8	—	*	*6	15	*6	*	*5	*7
151.0	9	*9	*	—	*	*	*6	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
153	120	57	63	—	*6	32	82	30	28	46	16	
153.0	*5	*	*	—	*	*	*	*	*	*	*	
153.1	10	*	*5	—	*	*	*8	*	*	*	*	
153.2	*5	*	*	—	*	*	*	*	*	*	*	
153.3	28	16	12	—	*	*8	19	*7	*8	9	*	
153.4	24	10	15	—	*	*5	17	*7	*5	*9	*	
153.6	22	10	12	—	*	*7	14	*	*	10	*5	
153.9	16	*8	*8	—	*	*	11	*	*	*7	*	
154	38	23	16	*	*	15	19	10	11	13	*5	
154.0	11	*6	*5	—	*	*	*6	*	*	*	*	
154.1	23	14	*8	*	*	10	11	*	*8	*8	*	
155	19	13	*5	*	*	*7	11	*	*	10	*	
155.0	14	10	*	*	*	*6	*7	*	*	*7	*	
156	*8	*	*	—	*	*	*	*	*	*	*	
157	32	16	17	*	*	10	22	9	*7	10	*7	
157.0	10	*	*6	—	*	*	*7	*	*	*	*	
157.8	*6	*	*	—	*	*	*	*	*	*	*	
157.9	12	*6	*7	—	*	*	*8	*	*	*	*	
161	*7	*6	*	—	*	*	*	*	*	*	*	
162	143	78	65	*	10	46	86	29	38	58	18	
162.2	*8	*	*	—	*	*	*	*	*	*	*	
162.3	39	19	20	*	*	12	25	*9	*9	17	*	
162.5	24	11	13	—	*	*5	18	*	*7	9	*	
162.8	18	12	*5	—	*	*6	*9	*5	*	*	*	
162.9	49	28	21	—	*	18	28	*8	16	21	*	
170	*6	*	*	*	*	*	*	*	*	*	*	
171	*8	*	*	*	*	*	*	*	*	*	*	
173	*5	*	*	*	*	*	*	*	*	*	*	
174	81	—	81	*	9	30	41	16	17	31	17	
174.4	10	—	10	—	*	*	*	*	*	*	*	
174.8	26	—	26	—	*	10	13	*9	*	*6	*8	
174.9	32	—	32	—	*	11	17	*	*8	17	*	
180	12	—	12	—	*5	*	*	*	*	*	*	
180.9	*7	—	*7	—	*	*	*	*	*	*	*	
182	34	—	34	—	*	16	16	*8	10	9	*7	
182.0	34	—	34	—	*	16	16	*8	10	9	*7	
183	26	—	26	*	*	12	11	*7	*	*9	*6	
183.0	25	—	25	*	*	11	11	*7	*	*7	*6	
184	*5	—	*5	—	*	*	*	*	*	*	*	
185	80	80	—	—	*	39	40	14	19	27	19	
188	36	29	*7	*	*	*7	27	12	*9	*7	*8	
188.8	13	11	*	—	*	*	10	*5	*	*	*	
188.9	11	*9	*	*	*	*	*8	*	*	*	*	
189	41	25	16	*	*	17	20	*8	*7	18	*8	
189.0	36	22	14	*	*	15	17	*7	*6	16	*7	
191	34	18	16	*	*6	16	*8	*8	*	18	*6	
191.3	*6	*	*	*	*	*	*	*	*	*	*	
191.9	*8	*	*	*	*	*	*	*	*	*	*	
193	25	*5	19	*	12	9	*	*7	*	*9	*	
196	12	*6	*5	*	*	*	*	*	*	*	*	
197	107	45	61	*	*7	39	60	27	19	42	18	
197.0	20	11	9	*	*	*7	11	*	*	*8	*	
197.2	24	9	14	*	*	*8	15	*7	*	*9	*	
197.6	23	9	13	—	*	*8	12	*6	*	9	*	
197.7	31	11	20	—	*	12	18	*8	*6	14	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
198	84	39	45	*	*7	33	43	27	17	26	15
198.3	36	16	20	*	*	18	15	12	*5	12	*7
198.5	29	13	15	*	*	*9	17	*9	*6	*8	*6
198.8	12	*6	*6	*	*	*	*7	*	*	*	*
198.89	11	*6	*	*	*	*	*6	*	*	*	*
199	20	10	11	—	*	*8	*8	*	*5	*8	*
199.0	11	*	*7	—	*	*6	*	*	*	*	*
199.1	9	*5	*	—	*	*	*	*	*	*	*
200	9	*	*	*	*	*	*5	*	*	*	*
200.0	*6	*	*	*	*	*	*	*	*	*	*
201	*7	*	*	*	*	*	*	*	*	*	*
202	32	16	17	*	*7	*7	17	*7	*8	12	*
202.8	27	12	15	*	*5	*6	15	*7	*6	10	*
202.80	17	*8	*9	*	*	*	9	*	*	*6	*
203	20	10	10	—	*	11	*8	*	*	*9	*
203.0	19	9	10	—	*	11	*8	*	*	*9	*
203.00	18	9	9	—	*	10	*8	*	*	*9	*
204	25	12	13	*8	*	*	11	*	*	*9	*8
204.0	12	*8	*	*7	*	*	*	*	*	*	*
204.00	10	*7	*	*5	*	*	*	*	*	*	*
204.1	12	*	9	—	*	*	10	*	*	*	*
204.10	11	*	*9	—	—	*	10	*	*	*	*
205	20	10	10	*	*	*7	*9	*	*5	*9	*
205.0	16	*8	*8	*	*	*	*8	*	*	*6	*
205.00	15	*8	*8	*	*	*	*8	*	*	*6	*
210	*7	*	*	*	*	*	*	*	*	*	*
210.2	*5	*	*	—	*	*	*	*	*	*	*
211	42	19	23	*	*	13	24	11	10	14	*6
211.3	30	13	16	*	*	10	17	*8	*8	10	*
218	215	—	215	—	108	104	*	37	41	97	41
218.0	28	—	28	—	14	14	*	*7	*5	10	*6
218.1	64	—	64	—	29	33	*	12	11	32	*9
218.2	22	—	22	—	11	11	*	*5	*	9	*
218.9	100	—	100	—	54	45	*	14	20	45	22
220	36	—	36	*	13	12	10	*7	*8	16	*5
225	23	*9	15	*	*6	10	*7	*5	*	*7	*8
225.2	17	*6	11	*	*	*7	*	*	*	*5	*5
226	*9	*	*8	—	*	*	*	*	*	*	*
227	17	*6	11	*	*	*8	*5	*	*	*6	*
227.1	*8	*	*6	*	*	*	*	*	*	*	*
227.3	*7	*	*	—	*	*	*	*	*	*	*
228	*7	*	*	*	*	*	*	*	*	*	*
228.0	*5	*	*	*	*	*	*	*	*	*	*
233	25	*	24	—	9	9	*6	*6	*	*9	*5
233.0	12	*	12	—	*	*5	*	*	*	*	*
233.1	10	—	10	—	*7	*	*	*	*	*	*
235	13	*8	*5	*	*	*	*8	*	*	*	*
235.2	*7	*	*	*	*	*	*	*	*	*	*
236	*6	*	*	—	*	*	*	*	*	*	*
237	*7	*	*	*	*	*	*	*	*	*	*
238	14	*8	*7	*	*	*	*8	*	*	*7	*
238.7	10	*7	*	*	*	*	*7	*	*	*5	*
239	14	*6	*8	*	*	*	*7	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
240-279	1,753	723	1,030	189	386	484	693	399	382	671	300
241	17	*	14	*	*7	*6	*	*	*	*8	*
241.0	*6	*	*6	*	*	*	*	*	*	*	*
241.1	10	*	*8	—	*	*	*	*	*	*	*
242	13	*	10	*	*	*	*	*	*	*5	*
242.0	*6	*	*	*	*	*	*	*	*	*	*
242.9	*6	*	*	*	*	*	*	*	*	*	*
242.90	*6	*	*	*	*	*	*	*	*	*	*
244	24	*	19	—	*	*7	12	*6	*7	*6	*
244.9	21	*	16	—	*	*6	11	*	*5	*6	*
250	599	299	300	31	148	208	211	144	108	237	109
250.0	157	76	81	10	30	65	52	39	33	59	26
250.00	86	44	42	*	16	41	27	22	17	33	13
250.01	18	*9	10	*6	*	*	*	*	*5	*6	*
250.02	43	19	24	*	*6	19	18	10	*7	16	9
250.03	10	*	*6	*	*	*	*	*	*	*	*
250.1	119	63	56	18	68	24	*8	24	21	50	25
250.10	20	11	*8	*	11	*6	*	*	*	*9	*
250.11	59	31	28	14	33	9	*	11	13	21	14
250.12	11	*6	*5	*	*	*	*	*	*	*	*
250.13	30	15	15	*	21	*	*	*6	*	16	*
250.2	15	*8	*7	—	*	*6	*6	*	*	*6	*
250.20	*7	*	*	—	*	*	*	*	*	*	*
250.22	*5	*	*	—	*	*	*	*	*	*	*
250.4	34	21	14	—	*	13	17	*7	*	11	12
250.40	23	15	*7	—	*	*8	13	*	*	*7	10
250.42	*8	*	*	—	*	*	*	*	*	*	*
250.6	60	25	35	*	14	27	19	15	11	26	*8
250.60	30	13	17	—	*	15	13	*8	*	14	*
250.61	11	*	*8	*	*	*	*	*	*	*	*
250.62	12	*6	*7	—	*	*6	*	*	*	*	*
250.63	*6	*	*	—	*	*	*	*	*	*	*
250.7	49	30	19	*	*	17	28	14	*8	19	*9
250.70	33	19	14	—	*	9	21	9	*6	12	*6
250.71	*8	*5	*	*	*	*	*	*	*	*	*
250.72	*7	*5	*	—	*	*	*	*	*	*	*
250.8	147	67	80	*	21	49	77	37	28	57	25
250.80	96	44	52	*	*7	29	60	22	20	34	19
250.81	24	11	13	*	*8	9	*7	*7	*	11	*
250.82	21	*9	12	—	*	*9	10	*6	*	11	*
250.83	*7	*	*	*	*	*	*	*	*	*	*
250.9	9	*6	*	*	*	*	*	*	*	*5	*
251	14	*7	*8	*	*	*	*	*	*	*	*
251.2	10	*	*6	*	*	*	*	*	*	*	*
252	9	*	*6	*	*	*	*	*	*	*	*
252.0	*8	*	*	—	*	*	*	*	*	*	*
253	19	*7	13	*	*	*	13	*	*	*9	*
253.6	16	*5	10	—	*	*	13	*	*	*8	*
255	*8	*	*	*	*	*	*	*	*	*	*
255.4	*5	*	*	*	*	*	*	*	*	*	*
263	13	*6	*7	*	*	*	*7	*	*	*5	*
263.9	11	*6	*6	*	*	*	*7	*	*	*	*
272	50	27	23	*	14	26	10	13	15	13	9
272.0	21	10	10	*	*5	11	*	*6	*6	*	*
272.4	25	16	10	*	*7	14	*	*5	*8	*8	*
274	17	12	*	—	*	*5	10	*6	*	*6	*
274.0	12	*8	*	—	*	*	*8	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
275	18	*7	11	*	*	*5	10	*6	*	*6	*	
275.4	14	*5	9	*	*	*	*9	*5	*	*	*	
275.42	*9	*	*6	—	*	*	*	*	*	*	*	
276	762	295	466	137	93	142	390	149	198	294	121	
276.0	14	*	9	*	*	*	10	*	*	*5	*	
276.1	125	39	86	*	*8	29	85	24	33	47	22	
276.2	12	*	*8	*	*	*	*	*	*	*5	*	
276.5	518	216	302	127	64	82	245	101	137	203	76	
276.7	34	15	19	*	*	*8	21	*	*8	13	*8	
276.8	46	11	35	*	12	16	16	11	9	16	*8	
276.9	*8	*	*5	*	*	*	*5	*	*	*	*	
277	23	12	11	*9	11	*	*	*6	*6	*7	*	
277.0	16	*7	*8	*8	*7	*	—	*5	*	*	*	
277.02	13	*7	*7	*6	*6	*	—	*5	*	*	*	
278	141	25	116	*	85	52	*	36	21	56	27	
278.0	132	24	108	*	82	48	*	34	19	53	27	
278.00	13	*	*8	*	*7	*	*	*	*	*5	*	
278.01	120	20	100	*	75	44	*	30	17	48	25	
278.1	*7	*	*7	—	*	*	*	*	*	*	*	
280-289	449	175	274	58	109	98	184	111	94	178	66	
280	87	25	62	*	14	23	48	22	22	34	9	
280.0	41	13	28	—	*8	10	23	*8	12	18	*	
280.9	44	12	33	*	*6	12	24	14	*9	16	*6	
282	85	36	49	18	58	*8	*	29	17	35	*	
282.6	77	34	43	15	55	*7	*	27	16	33	*	
282.62	66	27	39	11	48	*6	—	24	13	27	*	
283	14	*	9	*	*	*	*9	*	*	*7	*	
283.0	*6	*	*	*	*	*	*	*	*	*	*	
283.9	*7	*	*	*	*	*	*	*	*	*5	*	
284	29	13	15	*	*	10	10	*7	*	13	*5	
284.8	27	12	15	*	*	10	9	*7	*	11	*5	
285	97	35	62	*	11	19	65	20	21	35	20	
285.1	12	*	*8	—	*	*	10	*	*	*	*	
285.2	16	*6	10	*	*	*	12	*	*	*6	*	
285.22	*6	*	*	—	*	*	*	*	*	*	*	
285.29	*8	*	*	*	*	*	*6	*	*	*	*	
285.8	*8	*	*	*	*	*	*5	*	*	*	*	
285.9	61	22	39	*	*8	13	39	13	13	22	12	
286	18	*8	11	*	*	*	11	*	*5	*8	*	
286.7	*5	*	*	—	*	*	*	*	*	*	*	
287	39	15	24	10	*6	11	12	*9	*9	15	*7	
287.3	16	*5	11	*5	*	*	*	*	*	*6	*	
287.5	15	*6	*9	*	*	*6	*6	*	*	*5	*	
288	60	30	31	14	*9	16	22	16	10	24	11	
288.0	43	21	22	*8	*6	12	16	13	*7	15	*7	
288.8	13	*6	*7	*	*	*	*	*	*	*7	*	
289	15	*8	*8	*6	*	*	*	*	*	*	*5	
289.3	*6	*	*	*	*	*	*	*	*	*	*	
290-319	2,311	1,176	1,135	127	1,253	662	268	707	477	861	267	
290	36	16	20	—	*	*	33	13	*7	13	*	
290.3	*6	*	*	—	—	—	*6	*	*	*	*	
290.4	22	11	11	—	—	*	18	*8	*	9	*	
290.40	*8	*	*	—	—	*	*8	*	*	*	*	
290.41	*5	*	*	—	—	*	*	*	*	*	—	
290.42	*5	*	*	—	—	*	*	*	*	*	*	
291	136	96	41	—	59	66	11	31	49	29	27	
291.0	22	17	*6	—	9	10	*	*	*	*9	*5	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
291.8	105	72	33	—	47	51	*7	26	44	17	19	
291.81	102	69	33	—	45	50	*7	25	43	16	18	
292	90	51	39	*	54	23	13	39	18	18	15	
292.0	54	33	21	*	38	14	*	26	13	*7	*9	
292.8	28	13	15	*	12	*7	9	12	*	9	*	
292.81	14	*5	*9	—	*	*	*8	*	*	*5	*	
292.84	13	*7	*6	*	11	*	*	*7	*	*	*	
293	21	10	11	*	*	*	13	*6	*6	*7	*	
293.0	15	*7	*8	*	*	*	11	*	*5	*5	*	
293.8	*5	*	*	—	*	*	*	*	*	*	*	
294	37	17	19	—	*	*	33	11	*7	13	*5	
294.8	33	15	18	—	*	*	32	10	*7	12	*	
295	331	179	151	*	177	123	28	102	52	145	31	
295.3	114	70	44	*	59	42	12	32	19	50	14	
295.30	48	31	18	—	24	19	*6	11	11	21	*6	
295.32	22	13	*9	—	11	*8	*	*5	*	11	*	
295.34	42	26	16	*	24	15	*	14	*5	18	*	
295.6	11	*7	*	—	*	*5	*	*	*	*	*	
295.64	*5	*	*	—	*	*	*	*	*	*	*	
295.7	166	80	86	*	91	64	9	54	27	73	12	
295.70	127	60	66	*	68	51	*5	37	22	57	11	
295.74	34	17	17	*	19	11	*	15	*	14	*	
295.9	31	17	14	*	17	10	*	11	*	14	*	
295.90	21	12	9	*	10	*7	*	*6	*	12	*	
295.94	*5	*	*	—	*	*	*	*	*	*	*	
296	847	361	486	65	496	217	70	222	200	366	59	
296.2	193	86	107	16	115	41	22	34	50	98	11	
296.20	96	46	49	*6	60	20	10	18	32	40	*5	
296.22	17	*6	11	*	11	*	*	*	*	14	*	
296.23	49	19	30	*5	28	10	*5	9	10	27	*	
296.24	30	13	17	*	15	*7	*5	*6	*5	17	*	
296.3	269	104	166	*7	153	83	26	83	65	103	19	
296.30	60	24	36	*	34	22	*	19	20	18	*	
296.32	18	*6	12	*	11	*6	*	*	*	*8	*	
296.33	137	54	83	*	82	39	12	42	29	59	*8	
296.34	49	19	31	*	25	14	10	18	11	18	*	
296.4	59	28	31	*	33	18	*6	18	12	21	*8	
296.40	26	12	14	*	14	*8	*	*8	*6	*9	*	
296.43	*8	*	*	*	*	*	*	*	*	*	*	
296.44	23	11	13	*	14	*8	*	*7	*	*8	*	
296.5	84	33	51	*	51	26	*5	27	22	29	*6	
296.50	44	18	26	*	28	13	*	11	14	16	*	
296.53	19	*8	11	*	12	*5	*	10	*	*	*	
296.54	17	*6	11	*	*9	*6	*	*6	*	*7	*	
296.6	89	42	47	16	55	14	*	20	13	52	*	
296.60	40	19	21	*	27	*7	*	*6	*8	24	*	
296.63	18	*9	*9	*	11	*	*	*5	*	10	*	
296.64	30	14	16	*8	15	*	*	*8	*	18	*	
296.7	72	33	39	10	40	18	*	13	21	34	*	
296.8	46	18	27	*6	28	10	*	*7	13	21	*	
296.80	26	12	14	*	13	*6	*	*	*7	15	*	
296.89	18	*6	12	*	13	*	*	*	*6	*	*	
296.9	32	16	16	*6	20	*5	*	18	*	*8	*	
296.90	31	15	15	*6	19	*	*	17	*	*7	*	
297	*7	*	*	—	*	*	*	*	*	*	*	
298	90	43	47	*	50	21	17	26	13	32	20	
298.9	89	43	46	*	49	21	16	26	13	31	20	
299	*6	*	*	*	*	*	—	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
300	72	22	50	*	34	23	11	23	15	22	13	
300.0	32	9	22	*	12	12	*7	11	*6	*9	*7	
300.00	22	*6	16	*	*7	*8	*5	*7	*	*7	*	
300.01	*7	*	*	*	*	*	*	*	*	*	*	
300.1	*8	*	*7	*	*5	*	*	*	*	*	*	
300.11	*6	*	*	*	*	*	*	*	*	*	*	
300.4	19	*6	13	*	*9	*7	*	*6	*	*6	*	
300.9	*7	*	*	*	*	*	*	*	*	*	*	
303	136	90	45	*	64	58	13	39	23	50	23	
303.0	39	23	16	—	18	18	*	13	*7	*8	10	
303.00	12	*6	*6	—	*6	*6	*	*	*	*	*	
303.01	24	16	*8	—	11	12	*	12	*	*	*	
303.9	97	67	30	*	45	40	11	26	16	43	13	
303.90	43	30	13	—	23	16	*	*	*	29	*	
303.91	53	36	16	*	22	23	*7	20	11	13	*8	
304	98	62	36	*	70	27	*	30	*8	40	20	
304.0	48	31	18	—	34	14	*	19	*	14	12	
304.00	9	*	*	—	*6	*	*	*	*	*5	*	
304.01	38	25	13	—	27	11	*	18	*	*9	9	
304.1	*7	*	*	—	*	*	*	*	*	*	*	
304.2	13	10	*	*	11	*	*	*	*	*8	*	
304.20	*6	*	*	—	*	*	—	*	*	*5	*	
304.21	*7	*6	*	*	*6	*	*	*	*	*	*	
304.7	*7	*5	*	—	*	*	*	*	*	*	*	
304.71	*6	*	*	—	*	*	*	*	*	*	*	
304.8	*7	*	*	—	*6	*	—	*	*	*5	*	
304.80	*6	*	*	—	*	*	—	*	*	*5	*	
304.9	*8	*	*	—	*	*	*	*	*	*6	*	
304.90	*7	*	*	—	*	*	*	*	*	*6	*	
305	92	62	30	*	53	33	*	16	23	35	18	
305.0	38	29	9	*	20	14	*	*5	13	14	*5	
305.00	30	23	*8	*	17	10	*	*	10	13	*	
305.01	*6	*	*	—	*	*	*	*	*	*	*	
305.1	24	14	10	—	13	10	*	*	*7	10	*	
305.5	*6	*	*	—	*	*	—	*	*	*	*	
305.6	10	*7	*	—	*6	*	*	*	*	*	*	
305.60	*8	*6	*	—	*	*	*	*	*	*	*	
305.9	*9	*5	*	*	*7	*	*	*	*	*	*	
305.90	*7	*	*	*	*5	*	*	*	*	*	*	
307	10	*	*6	*	*	*	*	*	*	*	*	
309	71	33	37	*6	51	12	*	32	13	15	11	
309.0	22	13	9	*	16	*	*	13	*	*	*	
309.2	12	*5	*7	*	*8	*	*	*	*	*	*	
309.28	11	*	*6	*	*7	*	*	*	*	*	*	
309.4	16	*9	*7	*	13	*	*	*9	*	*	*	
309.8	15	*	12	*	9	*	*	*	*6	*	*	
309.81	15	*	11	*	9	*	—	*6	*	*	*	
310	*7	*	*	*	*	*	*	*	*	*	*	
311	159	74	86	10	103	36	10	83	24	41	11	
312	22	16	*7	*7	13	*	*	11	*	*7	*	
312.3	17	13	*	*	12	*	*	10	*	*6	—	
312.30	13	*9	*	*	*9	*	*	*8	*	*	—	
313	9	*6	*	*6	*	—	—	*	*	*	*	
313.8	*8	*5	*	*6	*	—	—	*	*	*	*	
313.81	*8	*5	*	*6	*	—	—	*	*	*	*	
314	16	11	*	12	*	*	—	*6	*	*7	*	
314.0	16	11	*	12	*	*	—	*6	*	*7	*	
314.01	14	11	*	11	*	—	—	*5	*	*7	—	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
320-389	553	238	316	80	133	132	208	130	118	203	102
320	*6	*	*	*	*	*	*	*	*	*	*
322	*7	*	*	*	*	*	*	*	*	*	*
322.9	*7	*	*	*	*	*	*	*	*	*	*
324	*8	*6	*	*	*	*	*	*	*	*	*
324.1	*6	*5	*	—	*	*	*	*	*	*	*
331	85	39	47	*	*	*7	70	18	22	31	13
331.0	63	28	35	*	*	*	60	12	18	25	*8
331.3	*5	*	*	*	*	*	*	*	*	*	*
331.4	11	*5	*6	*	*	*	*	*	*	*	*
332	21	12	*9	—	*	*	18	*5	*	*8	*
332.0	20	12	*8	—	*	*	18	*5	*	*8	*
333	11	*	*6	*	*	*	*	*	*	*5	*
336	*6	*	*	*	*	*	*	*	*	*	*
337	*6	*	*	*	*	*	*	*	*	*	*
340	28	*6	21	*	10	14	*	*8	*5	10	*
343	*6	*	*	*	*	*	*	*	*	*	*
344	*7	*	*	*	*	*	*	*	*	*	*
345	60	30	31	25	18	11	*6	20	*9	18	14
345.1	*7	*	*	*	*	*	*	*	*	*	*
345.10	*5	*	*	*	*	*	*	*	*	*	*
345.3	19	10	9	*8	*5	*	*	*	*	*6	*6
345.4	*7	*	*	*	*	*	*	*	*	*	*
345.9	14	*7	*7	*5	*	*	*	*	*	*	*
345.90	10	*5	*	*	*	*	*	*	*	*	*
346	59	13	46	*	34	16	*6	12	15	25	*7
346.2	*5	*	*	*	*	*	*	*	*	*	*
346.8	11	*	9	*	*6	*	*	*	*	*5	*
346.80	*8	*	*6	*	*	*	*	*	*	*	*
346.9	38	*7	31	*	23	11	*	*7	11	16	*
346.90	25	*	20	*	15	*6	*	*7	*6	9	*
346.91	14	*	12	*	*8	*5	*	*	*	*7	*
348	44	18	27	*	14	*9	17	12	*9	16	*8
348.2	*8	*	*8	*	*7	*	—	*	*	*	*
348.3	20	*8	12	*	*	*	14	*	*	9	*
348.30	9	*	*6	*	*	*	*	*	*	*6	*
348.31	*8	*	*5	—	*	*	*7	*	*	*	*
349	12	*	*8	*	*	*	*	*	*	*6	*
349.8	*6	*	*	*	*	*	*	*	*	*	*
349.82	*5	*	*	—	*	*	*	*	*	*	*
350	*7	*	*	*	*	*	*	*	*	*	*
350.1	*7	*	*	—	*	*	*	*	*	*	*
351	*6	*	*	*	*	*	*	*	*	*	*
351.0	*5	*	*	*	*	*	*	*	*	*	*
355	*9	*	*7	*	*	*	*	*	*	*	*
357	10	*7	*	*	*	*	*	*	*	*	*
357.0	*5	*	*	*	*	*	*	*	*	*	*
358	*8	*	*	*	*	*	*	*	*	*	*
358.0	*7	*	*	*	*	*	*	*	*	*	*
362	*6	*	*	*	*	*	*	*	*	*	*
368	*9	*	*	*	*	*	*	*	*	*5	*
376	*8	*	*	*	*	*	*	*	*	*	*
376.0	*7	*	*	*	*	*	*	*	*	*	*
376.01	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
380	*6	*	*	*	*	*	*	*	*	*	*
380.1	*6	*	*	*	*	*	*	*	*	*	*
382	10	*7	*	9	*	*	*	*	*	*	*
382.9	9	*6	*	*9	*	*	*	*	*	*	*
386	37	14	23	*	*	11	24	13	12	*8	*
386.1	20	*8	12	*	*	*7	12	*8	*7	*	*
386.11	13	*5	*8	—	*	*	*8	*	*5	*	*
386.3	13	*5	*8	—	*	*	*9	*	*	*	*
386.30	12	*	*7	—	*	*	*9	*	*	*	*
390-459	6,363	3,227	3,136	31	448	1,881	4,003	1,441	1,471	2,397	1,055
394	*6	*	*5	—	*	*	*	*	*	*	*
396	18	*9	9	*	*	*	14	*	*5	*5	*
396.3	*5	*	*	—	—	*	*	*	*	*	*
396.8	*5	*	*	—	—	*	*	*	*	*	*
398	33	*8	25	—	*	*	29	10	*6	10	*8
398.9	33	*8	25	—	*	*	29	10	*6	10	*8
398.91	33	*8	25	—	*	*	29	10	*6	9	*8
401	341	136	206	*	49	140	152	74	69	145	53
401.0	31	12	19	*	*5	10	16	*	*6	14	*7
401.9	307	123	184	*	43	128	136	67	63	131	46
402	65	26	39	—	*	17	43	11	15	32	*7
402.9	60	24	36	—	*	15	41	11	14	29	*6
402.90	10	*	*6	—	*	*	*	*	*	*	*
402.91	50	19	30	—	*	11	36	*9	12	24	*
403	120	57	63	*	24	45	48	25	21	49	25
403.0	9	*	*6	*	*	*	*	*	*	*	*
403.01	*9	*	*6	*	*	*	*	*	*	*	*
403.9	109	53	56	*	20	41	47	22	19	46	22
403.91	105	51	54	*	20	39	45	22	19	44	20
404	25	11	14	*	*	*7	16	*	*7	10	*
404.9	21	9	12	*	*	*6	14	*	*7	*8	*
404.93	17	*7	10	*	*	*	12	*	*6	*6	*
410	732	423	309	*	42	230	460	169	172	259	132
410.0	17	9	*8	—	*	*5	10	*	*	*7	*
410.01	17	9	*8	—	*	*5	10	*	*	*7	*
410.1	77	45	32	—	*7	33	37	17	18	25	18
410.11	75	44	31	—	*6	33	36	16	17	24	18
410.2	13	*8	*	—	*	*	*8	*	*	*5	*
410.21	13	*8	*	—	*	*	*8	*	*	*	*
410.3	*9	*6	*	—	*	*	*	*	*	*	*
410.31	*9	*6	*	—	*	*	*	*	*	*	*
410.4	93	62	31	—	*7	38	48	18	20	35	19
410.41	90	61	30	—	*7	37	47	17	20	34	19
410.5	10	*5	*	—	*	*	*5	*	*	*	*
410.51	10	*5	*	—	*	*	*5	*	*	*	*
410.7	404	228	177	—	19	117	268	109	85	143	67
410.70	*8	*	*	—	*	*	*6	*	*	*	*
410.71	391	220	171	—	18	115	259	105	83	136	66
410.72	*5	*	*	—	*	*	*	*	*	*	—
410.8	17	10	*7	*	*	*6	11	*	*	*7	*
410.81	16	10	*7	*	*	*6	10	*	*	*6	*
410.9	91	48	43	*	*	19	68	12	33	31	14
410.91	85	45	40	*	*	16	65	11	33	29	12
411	108	53	55	*	9	47	51	26	22	42	18
411.1	98	49	50	*	*8	44	46	22	21	39	16
411.8	*8	*	*	*	*	*	*	*	*	*	*
411.89	*8	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
412	*6	*	*	—	*	*	*	*	*	*	*	*
413	47	23	24	*	*	19	23	9	11	20	*7	
413.9	46	22	24	*	*	19	23	*9	11	19	*7	
414	1,088	677	411	*	49	447	590	239	241	429	180	
414.0	1,065	662	403	*	48	440	576	231	237	421	175	
414.00	148	85	63	—	*7	60	82	35	26	75	13	
414.01	875	547	328	*	41	367	466	187	202	334	152	
414.02	33	22	10	—	*	9	23	*7	*8	*9	9	
414.8	17	12	*5	*	*	*	12	*7	*	*	*	
415	122	54	68	—	21	38	63	24	33	38	26	
415.1	121	53	68	—	21	38	62	24	33	38	25	
415.11	*7	*	*	—	*	*	*	*	*	*	*	
415.19	114	49	65	—	20	35	59	23	31	36	24	
416	21	10	11	*	*	9	*8	*	*6	*7	*	
416.8	*9	*	*6	*	*	*	*	*	*	*	*	
416.9	*8	*6	*	—	*	*	*	*	*	*	*	
420	12	*6	*6	*	*	*	*	*	*	*5	*	
420.9	12	*6	*6	*	*	*	*	*	*	*5	*	
420.90	*7	*	*	*	*	*	*	*	*	*	*	
421	12	*7	*	*	*	*	*	*	*	*	*	
421.0	12	*7	*	*	*	*	*	*	*	*	*	
423	17	10	*8	*	*	*6	*9	*6	*	*	*	
423.9	12	*6	*7	*	*	*	*	*	*	*	*	
424	94	40	54	*	10	25	58	24	19	31	19	
424.0	38	11	27	*	*5	14	19	10	*6	15	*7	
424.1	49	26	23	*	*	10	36	12	13	14	10	
425	33	19	14	*	*6	12	13	*7	*	15	*7	
425.4	28	17	12	*	*5	10	12	*6	*	13	*5	
426	55	29	26	*	*	*9	41	16	12	16	11	
426.0	26	13	13	—	*	*	21	*7	*6	*8	*5	
426.1	17	9	*8	*	*	*	13	*6	*	*	*	
426.13	10	*	*	*	*	*	*7	*	*	*	*	
427	762	364	398	*	56	175	526	194	180	270	118	
427.0	24	11	14	*	*	*8	14	*6	*5	*8	*5	
427.1	56	38	18	*	*	17	34	13	11	23	*9	
427.3	444	204	240	*	28	108	307	112	116	154	63	
427.31	403	180	223	*	26	94	283	103	105	138	57	
427.32	41	23	17	*	*	14	24	9	10	15	*6	
427.4	*6	*	*	*	*	*	*	*	*	*	*	
427.41	*6	*	*	*	*	*	*	*	*	*	*	
427.5	12	*7	*	*	*	*	*8	*	*	*	*	
427.6	11	*	*7	*	*	*	*5	*	*	*	*	
427.69	10	*	*6	*	*	*	*	*	*	*	*	
427.8	204	94	110	*	17	34	151	57	41	73	33	
427.81	77	35	43	*	*	*6	70	23	14	29	12	
427.89	127	60	67	*	15	28	81	34	27	44	22	
428	1,099	524	575	*	38	235	822	248	281	412	158	
428.0	1,021	490	531	*	35	219	764	238	261	378	143	
428.2	24	14	11	—	*	*6	17	*	9	10	*	
428.20	*9	*5	*	—	*	*	*6	*	*	*	*	
428.21	*5	*	*	—	*	*	*	*	*	*	*	
428.23	*8	*	*	—	*	*	*5	*	*	*	*	
428.3	41	14	28	—	*	*8	32	*7	*9	18	*7	
428.30	22	*7	14	—	*	*	17	*	*6	10	*	
428.31	*9	*	*6	—	*	*	*6	*	*	*	*	
428.33	10	*	*7	—	*	*	*8	*	*	*	*	
429	14	*7	*7	*	*	*6	*6	*	*	*7	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
430	25	*7	17	*	*6	11	*7	*7	*5	*8	*	
431	65	29	36	*	*	14	46	14	11	26	15	
432	26	13	13	*	*	*	19	*5	*	10	*6	
432.1	21	12	*8	*	*	*	16	*	*	*7	*5	
432.9	*5	*	*	—	*	*	*	*	*	*	*	
433	144	85	59	—	*	35	107	26	36	58	24	
433.1	111	66	45	—	*	25	84	21	29	44	17	
433.10	95	58	38	—	*	19	75	18	24	39	14	
433.11	15	*8	*7	—	*	*6	*9	*	*	*	*	
433.3	26	15	10	—	—	*6	19	*5	*	11	*5	
433.30	21	13	*9	—	—	*5	16	*	*	9	*	
434	318	142	176	*	15	78	224	71	69	121	57	
434.0	12	*6	*7	—	*	*	*9	*	*	*	*	
434.01	10	*	*5	—	*	*	*7	*	*	*	*	
434.1	36	14	23	—	*	*7	28	11	*7	13	*5	
434.11	33	12	21	—	*	*6	25	10	*6	12	*	
434.9	269	122	147	*	13	68	187	57	60	104	48	
434.90	*7	*	*	—	*	*	*6	*	*	*	*	
434.91	262	120	142	*	13	67	182	56	59	100	47	
435	180	77	103	*	*7	42	131	45	46	65	24	
435.3	*6	*	*	—	*	*	*5	*	*	*	*	
435.8	10	*	*7	—	*	*	*7	*	*	*	*	
435.9	162	71	91	*	*6	38	118	43	39	58	23	
436	82	34	48	*	*6	15	61	9	18	38	17	
437	35	14	22	*	*	10	22	*6	15	11	*	
437.1	11	*	*7	—	—	*	10	*	*8	*	*	
437.2	*5	*	*	*	*	*	*	*	*	*	*	
437.3	*9	*	*5	*	*	*6	*	*	*	*	*	
438	31	15	16	—	*	*7	20	*9	*7	9	*6	
438.2	12	*	*7	—	*	*	*6	*	*	*5	*	
438.20	10	*	*5	—	*	*	*6	*	*	*5	*	
438.8	14	*8	*6	—	*	*	10	*	*5	*	*	
438.89	11	*7	*	—	*	*	*8	*	*5	*	*	
440	123	63	60	*	*	33	88	25	31	48	19	
440.1	9	*	*5	*	*	*	*8	*	*	*	*	
440.2	103	52	51	—	*	28	73	22	27	40	14	
440.20	11	*6	*	—	*	*	*8	*	*	*5	*	
440.21	28	15	13	—	*	13	15	*	*8	10	*	
440.22	*9	*	*6	—	*	*	*6	*	*	*	*	
440.23	20	*9	11	—	*	*	17	*6	*	*7	*	
440.24	33	18	15	—	*	*7	26	*8	9	12	*	
440.3	*7	*	*	—	*	*	*	*	*	*	*	
441	61	47	14	*	*	14	46	15	13	22	11	
441.0	9	*6	*	—	*	*	*5	*	*	*	*	
441.01	*5	*	*	—	*	*	*	*	*	*	*	
441.3	*5	*	*	—	—	*	*	*	*	*	*	
441.4	39	32	*7	—	*	*7	32	10	*9	13	*8	
442	9	*7	*	*	*	*	*5	*	*	*	*	
443	24	14	10	*	*	*8	14	*5	*	13	*	
443.9	21	12	*8	—	*	*7	13	*	*	12	*	
444	37	18	18	—	*	12	20	*5	*8	13	11	
444.2	26	13	13	—	*	*8	16	*	*5	10	*7	
444.21	*7	*	*	—	*	*	*	*	*	*	*	
444.22	20	9	10	—	*	*6	11	*	*	*8	*5	
446	12	*7	*5	*	*	*	*	*	*	*	*	
447	12	*7	*5	*	*	*	*6	*	*	*5	*	
447.1	*6	*	*	—	*	*	*	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
451	15	*	10	*	*	*5	*8	*	*	*	*
451.1	*7	*	*	—	*	*	*	*	*	*	*
453	145	60	85	*	23	41	81	31	39	58	18
453.8	140	58	82	*	22	40	79	30	39	54	17
454	9	*	*6	—	*	*	*6	*	*	*	*
455	31	14	17	—	*8	*9	15	*8	*6	12	*5
455.2	21	10	11	—	*5	*6	10	*6	*	*8	*
456	*6	*	*	*	*	*	*	*	*	*	*
458	99	48	51	*	*8	25	66	24	21	35	19
458.0	43	18	24	*	*	*7	32	11	*8	16	*7
458.2	15	*8	*8	*	*	*	11	*	*	*	*
458.29	14	*7	*7	—	*	*	10	*	*	*	*
458.8	*5	*	*	—	*	*	*	*	*	*	*
458.9	36	19	17	*	*	12	21	9	*8	12	*6
459	32	14	18	*	*	12	16	*6	*8	12	*6
459.0	*9	*	*6	*	*	*	*	*	*	*	*
459.8	18	9	9	—	*	*7	10	*	*	*7	*
459.81	18	*9	*9	—	*	*7	10	*	*	*7	*
460-519	3,550	1,712	1,838	687	346	736	1,780	769	860	1,306	615
461	10	*5	*	*	*	*	*	*	*	*	*
462	*9	*5	*	*	*	*	*	*	*	*	*
463	*8	*	*	*	*	*	—	*	*	*	*
464	20	10	10	16	*	*	*	*	*	*8	*5
464.4	13	*7	*6	13	—	*	—	*	*	*6	*
465	31	17	15	20	*	*	*6	*7	*6	12	*6
465.9	30	16	14	19	*	*	*6	*7	*6	11	*6
466	265	142	123	200	16	15	34	42	53	104	67
466.0	65	25	40	*5	14	14	32	16	12	28	9
466.1	200	117	83	194	*	*	*	26	41	75	58
466.11	121	75	46	119	*	*	*	14	26	43	38
466.19	80	42	37	76	*	*	*	12	16	32	20
473	13	*	*8	*	*	*	*	*	*	*6	*
474	19	12	*8	14	*	*	—	*6	*	*7	*
474.0	*7	*	*	*	*	*	—	*	*	*	*
474.00	*5	*	*	*	*	*	—	*	*	*	*
474.1	12	*9	*	9	*	—	—	*	*	*	*
474.10	*7	*	*	*7	*	—	—	*	*	*	*
475	13	*8	*6	*	*8	*	*	*	*	*	*
478	24	14	10	*7	*7	*6	*	*	*	11	*
478.1	*6	*	*	*	*	*	*	*	*	*	*
478.2	*8	*6	*	*	*	*	*	*	*	*	*
478.7	*6	*	*	*	*	*	*	*	*	*	*
480	32	17	15	26	*	*	*	*	11	10	*7
480.1	17	10	*7	17	*	—	*	*	*6	*	*
480.9	14	*6	*7	*8	*	*	*	*	*	*5	*
481	28	15	13	*	*	*8	16	*7	*6	10	*6
482	125	64	61	*7	9	23	87	21	35	45	25
482.1	25	13	12	*	*	*	16	*	*7	10	*
482.2	*7	*	*	*	*	*	*5	*	*	*	*
482.3	*7	*	*	*	*	*	*	*	*	*	*
482.4	42	22	20	*	*	*6	33	*8	10	16	*8
482.41	35	19	17	*	*	*6	27	*7	*7	13	*7
482.8	27	14	13	*	*	*	21	*	10	11	*
482.82	*5	*	*	*	—	*	*	*	*	*	*
482.83	13	*8	*	*	*	*	10	*	*	*7	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
482.9	14	*6	*7	*	*	*	*8	*	*	*	*
483	*8	*	*	*	*	*	*	*	*	*	*
483.0	*7	*	*	*	*	*	*	*	*	*	*
485	22	10	12	*6	*	*	12	*	*	11	*
486	1,114	528	586	140	88	206	680	234	322	377	181
487	20	9	11	*6	*	*	11	*7	*7	*	*
487.0	*9	*	*	*	—	*	*6	*	*	*	*
487.1	11	*5	*6	*	*	*	*5	*	*	*	*
490	25	12	14	*6	*	*	13	*	*	15	*
491	486	225	261	*	11	150	323	117	109	202	57
491.2	482	223	259	*	10	149	321	116	109	201	56
491.20	18	*7	11	—	*	*	14	*	*	11	*
491.21	464	216	248	*	10	144	308	115	106	190	53
492	17	*7	10	—	*	*6	11	*	*	*	*
492.8	15	*6	*9	—	*	*6	9	*	*	*	*
493	497	207	290	190	91	112	104	130	110	166	92
493.0	47	30	17	37	*5	*	*	*7	*	10	26
493.01	10	*6	*	*8	*	*	*	*	*	*	*
493.02	33	21	12	26	*	*	*	*	*	*6	20
493.2	124	35	89	*	10	48	66	37	29	38	20
493.20	24	*5	18	—	*	*7	15	*6	*6	*8	*
493.21	*5	*	*	*	*	*	*	*	*	*	*
493.22	96	29	67	*	*8	38	49	31	22	28	15
493.9	324	142	183	152	75	62	36	85	77	117	46
493.90	56	29	27	23	12	13	*8	12	11	24	9
493.91	64	34	30	45	13	*	*	17	13	26	*7
493.92	205	79	126	84	51	45	26	56	52	67	30
494	*8	*	*	*	*	*	*5	*	*	*	*
496	100	47	53	*	*	32	64	22	22	41	14
507	182	106	76	*7	*6	22	147	45	38	66	32
507.0	180	105	76	*6	*6	22	146	45	38	65	32
510	12	*8	*	*	*	*5	*	*	*	*	*
510.9	11	*7	*	*	*	*	*	*	*	*	*
511	57	32	25	*	*	15	37	13	11	23	10
511.0	*5	*	*	—	*	*	*	*	*	*	*
511.8	10	*6	*	*	*	*	*7	*	*	*	*
511.9	41	23	19	*	*	10	28	11	*7	17	*6
512	38	22	16	*	17	*9	11	*6	*7	18	*7
512.1	9	*	*6	*	*	*	*5	*	*	*	*
512.8	27	18	10	*	15	*6	*6	*	*5	14	*
513	*7	*	*	*	*	*	*	*	*	*	*
513.0	*7	*	*	*	*	*	*	*	*	*	*
514	9	*	*5	*	*	*	*	*	*	*	*
515	24	*7	17	*	*	*8	14	*	*5	10	*6
516	11	*6	*5	*	*	*	*	*	*	*	*
516.8	*6	*	*	*	*	*	*	*	*	*	*
518	278	130	148	14	29	73	162	50	59	116	53
518.0	*8	*	*	*	*	*	*	*	*	*	*
518.4	*6	*	*	*	*	*	*	*	*	*	*
518.8	254	117	136	11	25	68	149	46	53	105	50
518.81	190	88	102	*7	19	49	115	37	38	75	40
518.82	15	*8	*7	*	*	*	*6	*	*	*8	*
518.84	34	14	20	*	*	11	20	*	10	14	*5
518.89	11	*5	*6	*	*	*	*6	*	*	*6	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
519	21	13	*8	*7	*	*5	*6	*	*6	*7	*
519.0	*6	*	*	*	*	*	*	*	*	*	*
519.1	*6	*	*	*	*	*	*	*	*	*	*
519.8	*7	*5	*	*	*	*	*	*	*	*	*
520-579	3,580	1,580	2,001	248	859	1,070	1,404	756	809	1,305	710
522	*7	*	*	*	*	*	*	*	*	*	*
522.5	*6	*	*	*	*	*	*	*	*	*	*
524	19	*	14	*	14	*	*	*	*	*9	*5
524.0	9	*	*6	*	*7	*	*	*	*	*	*
524.03	*5	*	*	*	*	*	*	*	*	*	*
527	*7	*	*	*	*	*	*	*	*	*	*
527.2	*6	*	*	*	*	*	*	*	*	*	*
528	*8	*5	*	*	*	*	*	*	*	*	*
530	243	113	130	28	45	81	88	55	55	89	43
530.1	39	20	19	*	9	14	14	*6	*8	16	*9
530.10	*7	*	*	*	*	*	*	*	*	*	*
530.11	13	*6	*7	*	*	*5	*	*	*	*	*
530.19	15	*8	*8	*	*	*7	*5	*	*	9	*
530.2	17	*8	*9	*	*	*	12	*	*	*7	*
530.20	*5	*	*	*	*	*	*	*	*	*	*
530.21	12	*6	*6	*	*	*	9	*	*	*	*
530.3	11	*5	*6	*	*	*	*5	*	*	*	*
530.5	*8	*	*	—	*	*	*	*	*	*	*
530.7	13	*8	*6	*	*	*	*	*	*	*	*
530.8	147	65	83	23	28	51	46	36	36	52	23
530.81	139	59	79	23	27	49	40	34	34	50	21
531	108	50	58	*	*7	34	66	20	24	42	22
531.0	14	*5	*8	—	*	*	*8	*	*	*	*
531.00	13	*5	*8	—	*	*	*8	*	*	*	*
531.4	68	32	36	*	*	20	45	14	13	26	15
531.40	65	30	35	*	*	18	43	13	13	25	13
531.9	14	*6	*7	*	*	*	*8	*	*	*6	*
531.90	13	*6	*6	*	*	*	*7	*	*	*5	*
532	71	39	31	*	*9	18	44	16	20	23	12
532.0	10	*6	*	*	*	*	*6	*	*	*	*
532.00	10	*6	*	*	*	*	*6	*	*	*	*
532.4	42	26	16	—	*5	12	25	10	11	14	*7
532.40	41	26	16	—	*	12	25	10	10	14	*7
532.5	*7	*	*	*	*	*	*	*	*	*	*
532.50	*7	*	*	*	*	*	*	*	*	*	*
532.9	*6	*	*	*	*	*	*	*	*	*	*
532.90	*6	*	*	*	*	*	*	*	*	*	*
533	14	*5	*8	—	*	*	*8	*	*	*	*
533.4	*6	*	*	—	*	*	*	*	*	*	*
533.40	*6	*	*	—	*	*	*	*	*	*	*
534	11	*	*8	—	*	*6	*	*	*	*	*
534.4	*7	*	*5	—	*	*	*	*	*	*	*
534.40	*7	*	*5	—	*	*	*	*	*	*	*
535	157	64	94	*6	35	44	72	33	39	63	22
535.0	25	9	15	*	*6	*6	10	*6	*6	12	*
535.00	19	*7	11	*	*	*	*7	*	*	10	*
535.01	*6	*	*	*	*	*	*	*	*	*	*
535.1	19	*7	12	*	*	*	12	*6	*	*8	*
535.10	11	*	*8	*	*	*	*6	*	*	*7	*
535.11	*8	*	*	*	*	*	*6	*	*	*	*
535.3	*6	*	*	—	*	*	*	*	*	*	*
535.4	48	20	28	*	10	13	24	11	12	15	9
535.40	23	*8	15	*	*7	*6	*8	*6	*	*8	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
535.41	25	12	14	*	*	*7	16	*6	*7	*7	*	
535.5	52	19	32	*	11	15	23	*8	13	24	*6	
535.50	31	11	19	*	*8	10	10	*6	*8	14	*	
535.51	21	*8	13	*	*	*	13	*	*5	11	*	
535.6	*8	*	*	*	*	*	*	*	*	*	*	
536	59	21	39	*6	15	13	25	*8	15	25	11	
536.2	16	*	11	*	*	*	*	*	*	*6	*	
536.3	11	*	*8	*	*	*	*	*	*	*5	*	
536.4	19	9	10	*	*	*	12	*	*	*9	*	
536.41	*6	*	*	*	*	*	*	*	*	*	*	
536.42	*7	*	*	*	*	*	*5	*	*	*	*	
536.49	*5	*	*	*	*	*	*	*	*	*	*	
536.8	13	*	9	*	*	*	*6	*	*6	*	*	
537	30	12	18	*	*	*7	19	*	*6	11	*9	
537.0	9	*	*7	*	*	*	*	*	*	*	*	
537.8	17	*8	9	*	*	*	14	*	*	*7	*	
537.83	13	*7	*6	—	*	*	11	*	*	*	*	
540	300	175	124	79	149	52	20	65	61	88	85	
540.0	57	33	23	16	19	13	9	10	15	14	16	
540.1	35	21	14	11	11	*9	*	*8	*5	10	12	
540.9	208	121	87	52	119	31	*7	47	41	64	57	
541	*8	*	*	*	*	*	*	*	*	*	*	
550	38	32	*6	*7	*	*7	20	9	*6	14	*9	
550.1	19	16	*	*	*	*	10	*5	*	*7	*	
550.10	12	9	*	*	*	*	*6	*	*	*	*	
550.9	18	16	*	*	*	*	10	*	*	*7	*6	
550.90	12	10	*	*	*	*	*6	*	*	*	*	
552	71	26	44	*	16	22	32	15	18	26	12	
552.1	*8	*	*	*	*	*	*	*	*	*	*	
552.2	52	19	33	*	12	18	22	11	14	18	9	
552.20	12	*	*8	*	*	*	*6	*	*	*5	*	
552.21	36	14	21	—	*9	12	14	*8	11	10	*7	
553	93	30	62	*	22	39	29	16	16	42	19	
553.1	*8	*	*	*	*	*	*	*	*	*	*	
553.2	62	21	42	*	15	30	17	11	11	27	14	
553.20	13	*	9	—	*	*7	*	*	*	*7	*	
553.21	47	15	31	*	11	21	14	*8	*8	19	11	
553.3	21	*5	16	*	*	*8	10	*	*	11	*	
555	47	20	27	*	28	11	*6	11	12	18	*6	
555.0	11	*	*6	*	*6	*	*	*	*	*	*	
555.1	12	*6	*6	*	*7	*	*	*	*	*5	*	
555.2	*6	*	*	*	*	*	*	*	*	*	*	
555.9	19	*7	12	*	12	*	*	*5	*	*7	*	
556	37	21	15	*	18	10	*6	*9	11	10	*7	
556.6	*7	*	*	*	*	*	*	*	*	*	*	
556.9	19	12	*8	*	10	*	*	*	*7	*	*	
557	53	16	37	*	*6	12	34	14	11	16	12	
557.0	25	*8	16	*	*	*	15	*6	*5	*7	*7	
557.9	24	*6	19	*	*	*6	16	*6	*5	*8	*5	
558	179	66	113	51	41	37	50	39	42	71	27	
558.9	174	63	110	50	41	35	48	38	40	68	27	
560	305	128	177	14	32	85	174	65	73	105	62	
560.1	50	23	27	*	*	13	31	9	15	17	9	
560.2	14	*8	*6	*	*	*	*8	*	*	*	*	
560.3	31	10	21	*	*	*	22	*6	*7	12	*7	
560.39	27	9	18	*	*	*	19	*5	*6	10	*6	
560.8	90	36	54	*	11	28	48	18	21	31	20	
560.81	61	23	38	*	*8	18	33	12	13	22	14	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
560.89	29	12	16	*	*	10	15	*7	*8	*9	*6	
560.9	117	50	67	*	12	35	65	28	28	39	22	
562	312	132	180	*	42	93	177	80	75	109	48	
562.1	308	131	177	*	41	93	174	80	74	106	47	
562.10	29	10	19	—	*	*8	18	*7	*	12	*5	
562.11	188	80	108	*	35	70	82	51	50	60	27	
562.12	79	35	43	*	*	12	65	19	15	31	13	
562.13	12	*6	*7	—	*	*	*9	*	*	*	*	
564	76	26	50	*8	18	22	28	16	22	21	16	
564.0	48	19	29	*7	*8	11	22	10	14	14	*9	
564.00	36	14	22	*	*7	*8	16	*7	11	11	*7	
564.09	11	*	*6	*	*	*	*	*	*	*	*	
564.1	17	*	14	*	*8	*6	*	*	*	*	*	
566	28	19	9	*	12	10	*	*	*7	13	*	
567	24	12	11	*	*7	*8	*6	*	*	*9	*6	
567.2	15	*8	*7	*	*	*	*	*	*	*	*	
568	17	*	12	*	*7	*6	*	*	*	*8	*	
568.0	11	*	*9	*	*6	*	*	*	*	*5	*	
568.8	*6	*	*	*	*	*	*	*	*	*	*	
569	104	38	66	*	14	28	58	20	22	43	19	
569.1	*7	*	*6	*	*	*	*	*	*	*	*	
569.3	28	*9	19	*	*	*8	18	*	*	15	*	
569.4	*8	*	*	—	*	*	*	*	*	*	*	
569.6	21	*8	13	*	*5	*5	*9	*	*6	*6	*	
569.69	12	*	*8	*	*	*	*7	*	*	*	*	
569.8	37	15	22	*	*	10	21	*5	*9	16	*7	
569.81	*5	*	*	—	*	*	*	*	*	*	—	
569.83	13	*6	*7	*	*	*	*7	*	*	*6	*	
569.85	12	*	*7	*	*	*	10	*	*	*	*	
570	*8	*	*	*	*	*	*	*	*	*	*	
571	114	68	46	*	18	68	27	20	21	48	25	
571.1	12	*8	*	—	*	*7	*	*	*	*	*6	
571.2	49	37	12	—	9	33	*7	10	*8	19	12	
571.5	40	19	22	*	*	21	17	*6	*8	22	*	
572	54	32	22	*	*6	33	14	13	*6	22	13	
572.2	44	27	17	—	*	29	11	11	*	18	11	
573	13	*	*8	*	*	*	*5	*	*	*	*	
573.3	*8	*	*6	*	*	*	*	*	*	*	*	
574	354	112	242	*	122	103	127	73	71	123	87	
574.0	130	51	79	*	40	45	45	25	29	50	26	
574.00	121	47	74	*	38	43	39	23	28	46	23	
574.01	10	*	*5	—	*	*	*6	*	*	*	*	
574.1	115	26	89	*	43	35	37	27	20	38	30	
574.10	111	25	86	*	42	33	36	26	20	37	29	
574.2	42	12	30	*	17	10	15	*7	*9	15	12	
574.20	40	11	30	*	16	9	14	*7	*8	14	11	
574.3	*6	*	*	—	*	*	*	*	*	*	*	
574.5	20	*9	11	—	*6	*	11	*	*	*6	*5	
574.50	13	*	*8	—	*	*	*6	*	*	*	*	
574.51	*7	*	*	—	*	*	*	*	*	*	*	
574.6	*8	*	*	—	*	*	*	*	*	*	*	
574.7	15	*	11	*	*6	*	*	*	*	*	*	
574.70	9	*	*7	*	*	*	*	*	*	*	*	
574.71	*6	*	*	*	*	*	*	*	*	*	*	
574.8	*9	*	*5	—	*	*	*	*	*	*	*	
574.81	*6	*	*	—	*	*	*	*	*	*	*	
574.9	*6	*	*5	*	*	*	*	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
575	81	31	50	*	27	25	28	13	24	28	15
575.0	25	10	15	*	*	*9	11	*5	*8	*7	*
575.1	46	18	27	*	17	14	15	*6	13	18	*9
575.10	12	*	*7	—	*	*	*5	*	*	*5	*
575.11	23	*9	14	*	12	*7	*	*	*7	*8	*
575.12	11	*5	*6	—	*	*	*6	*	*	*	*
575.8	*8	*	*6	*	*	*	*	*	*	*	*
576	24	11	13	*	*6	*6	12	*8	*	*7	*
576.1	*7	*	*	*	*	*	*	*	*	*	*
576.2	10	*5	*5	*	*	*	*	*	*	*	*
576.8	*6	*	*	*	*	*	*	*	*	*	*
577	288	136	151	*6	88	114	79	52	63	117	55
577.0	246	117	129	*	76	94	72	44	55	100	46
577.1	26	12	14	*	10	12	*	*	*6	11	*6
577.2	*7	*	*	*	*	*	*	*	*	*	*
577.9	*7	*	*	*	*	*	*	*	*	*	*
578	196	97	99	*	18	52	121	44	55	66	31
578.0	23	10	12	*	*6	*6	10	*	*6	*9	*
578.1	42	17	24	*	*	9	28	11	9	14	*7
578.9	131	69	62	*	9	36	84	29	40	43	19
580-629	1,941	583	1,358	84	580	496	782	361	457	776	347
581	*8	*	*	*	*	*	*	*	*	*	*
581.9	*6	*	*	*	*	*	*	*	*	*	*
584	221	113	108	*	15	58	146	54	48	81	38
584.5	13	*7	*6	*	*	*	*7	*	*	*	*
584.9	204	104	100	*	14	52	137	49	45	76	35
585	19	9	10	*	*	*6	*9	*	*	*9	*
590	150	27	123	23	57	30	39	20	42	56	31
590.1	68	13	55	*8	33	11	17	12	21	23	12
590.10	68	13	55	*8	33	11	17	12	21	23	12
590.8	77	13	64	14	24	19	21	*8	21	31	18
590.80	77	13	64	14	24	19	21	*8	21	31	18
591	13	*5	*7	*	*	*	*	*	*	*5	*
592	171	88	83	*	73	68	29	34	46	65	25
592.0	56	26	29	*	23	23	9	13	16	17	10
592.1	113	61	52	*	47	45	19	22	30	46	15
593	39	18	21	10	*7	11	12	10	10	13	*6
593.4	*7	*	*	*	*	*	*	*	*	*	*
593.7	*7	*	*5	*7	*	*	—	*	*	*	*
593.70	*6	*	*	*6	*	—	—	*	*	*	*
593.9	14	*7	*7	*	*	*6	*7	*	*	*5	*
594	*7	*5	*	—	*	*	*	*	*	*	*
594.1	*6	*	*	—	*	*	*	*	*	*	*
595	24	12	13	*	*	*	18	*9	*	9	*
595.0	*7	*	*	*	*	*	*6	*	*	*	*
595.9	*7	*	*	*	*	*	*6	*	*	*	*
596	13	*9	*	*	*	*	*8	*	*	*	*
596.0	*6	*5	*	—	*	*	*5	*	*	*	*
599	470	147	323	27	37	78	327	98	109	181	81
599.0	429	125	304	26	35	66	302	85	102	170	72
599.7	31	20	11	*	*	*8	20	12	*6	*8	*5
600	94	94	—	—	*	20	74	22	26	30	17
600.0	85	85	—	—	*	18	66	20	24	26	15
600.00	23	23	—	—	*	*6	17	*7	*5	*5	*5
600.01	62	62	—	—	*	12	49	12	18	21	10

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
601	*8	*8	—	—	*	*	*	*	*	*	*
604	11	11	—	*	*	*	*6	*	*	*	*
604.9	*9	*9	—	*	*	*	*	*	*	*	*
604.90	*6	*6	—	*	*	*	*	*	*	*	*
607	*6	*6	—	*	*	*	*	*	*	*	*
608	13	13	—	*	*	*6	*	*	*	*	*
608.4	*6	*6	—	*	*	*	*	*	*	*	*
611	29	*	28	*	14	12	*	*6	*6	13	*
611.0	12	*	11	*	*6	*	*	*	*	*6	*
611.1	12	*	12	*	*7	*	*	*	*	*	*
614	41	—	41	*	34	*	*	*8	9	17	*7
614.2	*7	—	*7	*	*6	*	*	*	*	*	*
614.6	*9	—	*9	—	*7	*	*	*	*	*	*
614.9	13	—	13	*	12	*	*	*	*	*6	*
616	9	—	9	*	*	*	*	*	*	*6	*
617	78	—	78	—	61	18	*	10	19	33	17
617.0	46	—	46	—	33	13	*	*6	11	20	10
617.1	14	—	14	—	12	*	—	*	*	*6	*
617.3	12	—	12	—	10	*	*	*	*	*	*
618	147	—	147	—	27	58	61	19	32	67	30
618.0	30	—	30	—	*	12	13	*6	*6	14	*
618.1	15	—	15	—	*8	*	*	*	*	*	*
618.2	20	—	20	—	*	11	*	*	*	*9	*
618.3	15	—	15	—	*	*	*7	*	*	*	*
618.4	27	—	27	—	*5	12	10	*	*8	14	*
618.5	27	—	27	—	*	*9	18	*	*5	14	*6
618.8	*8	—	*8	—	*	*	*	*	*	*	*
619	*7	—	*7	*	*	*	*	*	*	*	*
620	83	—	83	*	59	16	*	13	20	37	13
620.0	*6	—	*6	*	*	*	—	*	*	*	*
620.1	12	—	12	*	9	*	—	*	*	*6	*
620.2	54	—	54	*	40	10	*	*8	13	24	*9
620.5	*5	—	*5	*	*	*	*	*	*	*	*
620.8	*6	—	*6	*	*	*	*	*	*	*	*
621	21	—	21	—	*8	10	*	*	*	11	*
621.3	11	—	11	—	*	*6	*	*	*	*	*
622	*7	—	*7	—	*	*	*	*	*	*	*
622.1	*6	—	*6	—	*	*	*	*	*	*	*
625	80	—	80	*	54	20	*6	9	20	33	18
625.3	25	—	25	—	22	*	—	*	*	12	*5
625.6	27	—	27	—	*7	15	*5	*	*8	*8	*7
625.9	25	—	25	*	22	*	*	*	*6	10	*
626	124	—	124	*	84	39	*	13	27	58	26
626.2	94	—	94	*	64	29	*	*8	23	43	19
626.8	26	—	26	*	16	9	*	*	*	11	*7
627	15	—	15	—	*	*9	*	*	*	*5	*
627.1	11	—	11	—	*	*7	*	*	*	*	*
630-677	523	—	523	*	520	*	—	87	90	243	103
632	21	—	21	—	21	—	—	*	*	10	*
633	32	—	32	*	31	*	—	*6	*	15	*8
633.1	28	—	28	*	27	*	—	*6	*	12	*7
633.10	26	—	26	*	26	*	—	*	*	12	*7
634	17	—	17	*	17	*	—	*6	*	*8	*
634.9	12	—	12	*	12	*	—	*	*	*6	*
634.91	10	—	10	*	10	*	—	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
640	*7	—	*7	—	*7	—	—	*	*	*	*	
641	19	—	19	—	19	—	—	*	*6	*7	*	
641.1	*6	—	*6	—	*6	—	—	*	*	*	*	
641.13	*6	—	*6	—	*6	—	—	*	*	*	*	
641.9	*5	—	*5	—	*5	—	—	*	*	*	*	
641.93	*5	—	*5	—	*5	—	—	*	*	*	*	
642	30	—	30	—	30	*	—	*5	*7	14	*	
642.3	*8	—	*8	—	*8	*	—	*	*	*6	*	
642.33	*8	—	*8	—	*8	*	—	*	*	*5	*	
642.4	11	—	11	—	11	—	—	*	*	*	*	
642.43	*8	—	*8	—	*8	—	—	*	*	*	*	
643	35	—	35	—	35	*	—	*6	*8	18	*	
643.0	16	—	16	—	16	—	—	*	*	*8	*	
643.03	16	—	16	—	16	—	—	*	*	*8	*	
643.1	13	—	13	—	13	—	—	*	*	*6	*	
643.13	13	—	13	—	13	—	—	*	*	*6	*	
644	104	—	104	*	104	*	—	19	18	49	19	
644.0	93	—	93	*	93	*	—	18	16	44	15	
644.03	92	—	92	*	92	*	—	18	16	43	15	
644.1	11	—	11	—	11	*	—	*	*	*	*	
644.13	11	—	11	—	11	*	—	*	*	*	*	
646	66	—	66	*	66	*	—	*6	10	38	13	
646.2	*5	—	*5	—	*5	—	—	*	*	*	*	
646.23	*5	—	*5	—	*5	—	—	*	*	*	*	
646.6	40	—	40	*	40	*	—	*	*7	21	*8	
646.63	36	—	36	*	36	—	—	*	*7	18	*7	
646.8	18	—	18	—	18	—	—	*	*	12	*	
646.83	16	—	16	—	16	—	—	*	*	12	*	
647	*5	—	*5	—	*5	—	—	*	*	*	*	
648	95	—	95	*	94	*	—	14	17	42	22	
648.0	10	—	10	—	10	—	—	*	*	*7	*	
648.03	10	—	10	—	10	—	—	*	*	*7	*	
648.4	9	—	9	—	9	—	—	*	*	*	*	
648.43	*7	—	*7	—	*7	—	—	*	*	*	*	
648.8	*8	—	*8	—	*8	—	—	*	*	*	*	
648.83	*8	—	*8	—	*8	—	—	*	*	*	*	
648.9	59	—	59	*	59	*	—	10	11	23	14	
648.93	56	—	56	*	56	*	—	9	11	22	14	
654	13	—	13	*	12	—	—	*	*	*	*	
654.5	*5	—	*5	—	*5	—	—	*	*	*	*	
654.53	*5	—	*5	—	*5	—	—	*	*	*	*	
658	10	—	10	—	10	—	—	*	*	*	*	
658.1	*5	—	*5	—	*5	—	—	*	*	*	*	
658.13	*5	—	*5	—	*5	—	—	*	*	*	*	
659	*5	—	*5	—	*5	*	—	*	*	*	*	
666	*6	—	*6	—	*6	—	—	*	*	*	*	
666.2	*6	—	*6	—	*6	—	—	*	*	*	*	
666.24	*6	—	*6	—	*6	—	—	*	*	*	*	
670	10	—	10	—	10	—	—	*	*	*	*	
670.0	10	—	10	—	10	—	—	*	*	*	*	
670.04	10	—	10	—	10	—	—	*	*	*	*	
674	14	—	14	—	14	—	—	*	*	*	*	
674.8	*7	—	*7	—	*7	—	—	*	*	*	*	
674.84	*7	—	*7	—	*7	—	—	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
680-709	764	395	368	105	214	210	234	150	151	351	111
681	33	19	14	*6	11	9	*7	*7	*7	13	*6
681.0	18	10	*8	*	*8	*	*	*	*	*6	*
681.00	16	*8	*7	*	*6	*	*	*	*	*5	*
681.1	15	9	*6	*	*	*6	*	*	*	*7	*
681.10	15	*9	*6	*	*	*6	*	*	*	*7	*
682	529	276	253	48	154	161	166	112	116	221	79
682.0	44	20	24	*9	18	11	*6	*7	*8	22	*6
682.1	11	*6	*	*	*5	*	*	*	*	*6	*
682.2	45	21	24	*	18	13	10	*7	10	22	*6
682.3	59	35	24	*	23	18	14	14	10	25	9
682.4	31	19	12	*	11	13	*5	*5	*6	11	*7
682.5	23	13	10	*8	*9	*6	*	*	*	16	*
682.6	254	129	126	12	57	78	108	63	63	94	35
682.7	54	31	23	*6	10	20	18	10	15	21	*8
692	*5	*	*	*	*	*	*	*	*	*	*
693	*6	*	*	*	*	*	*	*	*	*	*
693.0	*5	*	*	*	*	*	*	*	*	*	*
701	13	*7	*6	*7	*	*	*	*	*	11	*
701.4	*8	*	*	*7	*	—	—	*	*	*8	—
707	82	42	41	*	13	24	45	17	16	36	14
707.0	48	25	22	*	*8	14	25	*8	10	21	9
707.1	30	15	15	*	*	*8	19	*8	*5	12	*
707.15	13	*7	*6	—	*	*5	*7	*	*	*7	*
708	*7	*	*6	*	*	*	*	*	*	*	*
709	51	28	23	31	16	*	*	*	*	48	*
709.2	47	26	21	30	15	*	*	*	*	46	*
710-739	1,920	805	1,115	38	311	681	889	403	479	663	375
710	23	*	20	*	11	*6	*	*6	*	9	*
710.0	16	*	14	*	9	*	*	*	*	*7	*
711	23	16	*7	*	*	*9	*7	*	*	*9	*6
711.0	22	15	*6	*	*	*9	*6	*	*	*9	*6
711.06	11	*8	*	*	*	*6	*	*	*	*	*
714	27	*6	21	*	*	*9	14	*7	*6	*9	*
714.0	21	*	16	*	*	*7	11	*6	*	*6	*
715	696	259	436	—	23	246	427	159	190	221	126
715.1	53	22	31	—	*	17	34	15	*8	22	*8
715.15	13	*	*9	—	*	*	*9	*5	*	*	*
715.16	34	13	21	—	*	12	22	*8	*	16	*6
715.2	*6	*	*	—	*	*	*	*	*	*	*
715.3	382	147	236	—	11	139	233	93	108	110	72
715.31	10	*	*6	—	—	*	*7	*	*	*	*
715.35	105	48	58	—	*	37	65	27	29	28	21
715.36	262	91	171	—	*6	97	159	61	75	78	47
715.8	*7	*	*	—	—	*	*6	*	*	*	*
715.89	*7	*	*	—	—	*	*6	*	*	*	*
715.9	246	85	161	—	*9	86	151	48	70	83	44
715.90	*8	*	*	—	*	*	*6	*	*	*	*
715.91	*7	*	*	—	*	*	*5	*	*	*	*
715.95	75	30	45	—	*5	27	43	20	19	22	14
715.96	153	48	105	—	*	54	96	25	49	53	25
716	18	*9	9	*	*	*5	9	*	*6	*6	*
716.9	10	*	*6	*	*	*	*7	*	*	*	*
717	*7	*	*	*	*	*	*	*	*	*	*
718	11	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
719	38	14	24	*	*6	12	19	*8	12	12	*7	
719.4	27	*7	19	*	*	*8	14	*6	*8	*8	*5	
719.41	*8	*	*5	*	*	*	*	*	*	*	*	
719.45	11	*	*9	*	*	*	*8	*	*	*	*	
721	98	41	57	*	13	40	46	16	26	36	20	
721.0	25	12	12	—	*5	14	*5	*	*6	10	*5	
721.1	22	13	*9	*	*	10	*9	*	*	*8	*6	
721.3	44	14	31	—	*	14	26	*7	13	15	*9	
722	369	178	191	*	130	153	85	62	89	140	77	
722.0	64	34	30	—	28	32	*	10	13	30	11	
722.1	183	91	91	*	74	66	43	34	44	64	39	
722.10	179	91	88	*	74	64	41	34	43	62	39	
722.4	12	*	*8	—	*	*8	*	*	*	*	*	
722.5	63	29	35	—	16	24	24	*8	18	22	16	
722.52	62	27	34	—	15	24	23	*8	17	22	15	
722.7	20	10	10	—	*	11	*	*	*6	*7	*	
722.71	14	*6	*7	—	*	*8	*	*	*	*6	*	
722.73	*6	*	*	—	*	*	*	*	*	*	*	
722.8	16	*6	10	—	*	*8	*	*	*	*7	*	
722.83	14	*5	*8	—	*	*7	*	*	*	*6	*	
722.9	10	*	*6	*	*	*	*	*	*	*	*	
722.93	*7	*	*	*	*	*	*	*	*	*	*	
723	27	12	15	*	*6	12	*9	*	*5	14	*5	
723.0	10	*	*5	*	*	*5	*	*	*	*5	*	
723.1	11	*	*6	*	*	*	*	*	*	*6	*	
724	155	66	88	*	19	52	83	37	36	53	29	
724.0	86	37	50	*	*	30	53	20	18	31	18	
724.02	84	36	48	*	*	29	52	20	17	30	17	
724.2	34	16	19	*	10	11	14	*7	12	11	*	
724.4	*8	*	*	—	*	*	*	*	*	*	*	
724.5	17	*5	12	—	*	*6	*8	*5	*	*	*	
726	42	21	20	*	*7	16	18	11	11	*9	11	
726.1	20	10	10	—	*	*7	12	*	*	*	*8	
726.10	19	9	10	—	*	*6	12	*	*	*	*8	
726.3	10	*7	*	—	*	*5	*	*	*	*	*	
726.33	10	*7	*	—	*	*5	*	*	*	*	*	
727	22	12	10	*	*	*5	10	*	*	*9	*5	
727.0	*7	*	*	*	*	*	*	*	*	*	*	
727.4	*6	*	*	—	*	*	*	*	*	*	*	
727.40	*6	*	*	—	*	*	*	*	*	*	*	
727.6	*5	*	*	—	*	*	*	*	*	*	*	
728	36	19	18	*	10	12	13	*9	*8	15	*	
728.8	33	17	16	*	9	10	12	*8	*7	14	*	
728.88	18	10	*8	*	*5	*	*8	*5	*	*8	*	
729	52	23	29	*	12	18	19	9	13	21	*8	
729.1	10	*	*7	*	*	*	*	*	*	*	*	
729.5	19	10	10	*	*	*7	*8	*	*	*7	*	
729.8	17	*7	9	*	*	*5	*8	*	*	9	*	
729.89	11	*	*7	*	*	*	*6	*	*	*7	*	
730	48	31	17	*	10	18	15	10	13	16	10	
730.0	12	9	*	*	*	*	*	*	*	*	*	
730.07	*6	*	*	*	*	*	*	*	*	*	*	
730.1	*7	*	*	*	*	*	*	*	*	*	*	
730.2	29	17	12	*	*6	12	10	*5	*8	11	*	
730.27	15	*7	*8	*	*	*5	*7	*	*	*6	*	
730.28	*6	*	*	*	*	*	*	*	*	*	*	
732	*5	*	*	*	*	*	*	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
733	160	55	105	*	26	42	88	35	36	55	33
733.1	73	20	53	*	*	10	57	16	20	22	15
733.13	51	12	40	—	*	*6	45	12	12	15	13
733.14	9	*	*8	*	*	*	*	*	*	*	*
733.15	*6	*	*	*	*	*	*	*	*	*	*
733.4	26	13	13	*	*6	12	*8	*	*	13	*6
733.42	21	11	10	*	*5	10	*5	*	*	10	*5
733.6	22	*6	16	*	*	9	10	*6	*6	*7	*
733.8	26	13	13	*	9	*9	*7	*	*5	9	*7
733.81	*5	*	*	*	*	*	*	*	*	*	*
733.82	20	9	11	*	*6	*7	*7	*	*	*7	*6
733.9	*8	*	*6	*	*	*	*	*	*	*	*
736	9	*5	*	*	*	*	*	*	*	*	*
737	12	*	*9	*	*	*	*	*	*	*	*
737.3	11	*	*8	*	*	*	*	*	*	*	*
737.30	*8	*	*7	*	*	*	*	*	*	*	*
738	33	12	21	*	*7	13	12	*6	*6	*9	12
738.4	27	10	17	*	*	11	12	*	*5	*7	10
740-759	206	112	94	143	32	22	*9	53	31	72	49
742	*7	*	*	*7	*	*	*	*	*	*	*
744	*6	*	*	*	*	*	—	*	*	*	*
745	28	15	13	20	*5	*	*	10	*	10	*6
745.2	*7	*	*	*6	*	—	—	*	*	*	*
745.4	*7	*	*	*6	*	—	—	*	*	*	*
745.5	*8	*	*	*	*	*	*	*	*	*	*
746	19	11	*8	11	*	*	*	*6	*	*5	*6
746.4	*5	*	*	*	*	*	*	*	*	*	*
747	25	9	16	12	*	*7	*	*8	*6	*7	*
747.8	10	*	*8	*	*	*6	*	*	*	*	*
747.81	*9	*	*7	*	*	*6	*	*	*	*	*
749	14	*7	*6	13	*	—	—	*	*	*	*6
749.0	*6	*	*	*5	*	—	—	*	*	*	*
749.2	*7	*	*	*6	*	—	—	*	*	*	*
750	16	13	*	15	—	*	*	*	*	*6	*
750.5	14	12	*	14	—	*	—	*	*	*5	*
751	13	*8	*	9	*	*	*	*	*	*5	*
752	*5	*	*	*	*	*	*	*	*	*	*
753	11	*6	*	*	*	*	*	*	*	*5	*
754	13	9	*	9	*	*	*	*5	*	*	*
755	*6	*	*	*	*	*	—	*	*	*	*
756	16	*7	*8	*9	*	*	*	*	*	*5	*
756.1	*7	*	*	*	*	*	*	*	*	*	*
756.12	*5	*	*	*	*	*	*	*	*	*	*
757	9	*	*5	*8	*	—	—	*	*	*8	*
757.3	*9	*	*5	*8	*	—	—	*	*	*8	*
757.32	*9	*	*5	*8	*	—	—	—	*	*8	*
759	*6	*	*	*	*	*	*	*	*	*	*
760-779	203	112	91	203	*	—	*	26	39	72	67
765	34	18	16	34	—	—	—	*	*	19	9
765.1	29	16	13	29	—	—	—	*	*	17	*7
765.17	*7	*	*	*7	—	—	—	*	*	*	*
765.18	*6	*	*	*6	—	—	—	*	*	*	*
769	19	12	*7	19	*	—	—	*	*	*	*9

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
770	40	24	16	40	—	—	—	*	9	11	16
770.6	*7	*6	*	*7	—	—	—	*	*	*	*
770.8	23	13	10	23	—	—	—	*	*6	*6	*9
770.81	*9	*	*	*9	—	—	—	*	*	*	*
770.89	11	*7	*	11	—	—	—	*	*	*	*
771	15	*8	*7	15	—	—	—	*	*	*6	*
771.8	12	*7	*	12	—	—	—	*	*	*	*
771.81	*8	*	*	*8	—	—	—	*	*	*	*
774	52	27	25	52	—	—	—	*9	*8	17	17
774.2	*6	*	*	*6	—	—	—	*	*	*	*
774.6	41	21	20	41	—	—	—	*7	*7	14	13
777	*6	*	*	*6	—	—	—	*	*	*	*
778	10	*5	*	10	—	—	—	*	*	*	*
778.4	*8	*	*	*8	—	—	—	*	*	*	*
779	12	*6	*5	12	*	—	—	*	*	*	*
779.3	*5	*	*	*5	—	—	—	*	*	*	*
780-799	248	115	133	59	79	64	47	57	37	109	46
780	81	43	38	36	18	14	13	20	14	32	15
780.2	19	11	*8	*	*	*7	*8	*6	*	*7	*
780.3	30	14	17	15	10	*	*	10	*6	10	*6
780.39	27	11	15	12	10	*	*	9	*5	*8	*
780.6	19	12	*7	16	*	*	*	*	*	9	*
784	*7	*	*	*	*	*	*	*	*	*	*
786	64	32	32	*	17	28	14	16	*7	31	10
786.0	*7	*	*	*	*	*	*	*	*	*	*
786.5	56	28	27	*	15	27	13	14	*6	26	9
786.50	17	*9	*9	—	*5	*7	*	*	*	*8	*
786.59	36	19	18	*	9	19	*8	*9	*	17	*7
787	*7	*	*	*	*	*	*	*	*	*5	*
789	35	10	25	*6	21	*6	*	9	*6	14	*6
789.0	34	10	24	*6	20	*5	*	9	*6	13	*6
789.00	*7	*	*	*	*	*	*	*	*	*	*
789.03	13	*	10	*	*9	*	*	*	*	*6	*
799	38	16	23	*	13	10	12	*5	*6	18	10
799.9	38	16	22	*	13	10	11	*5	*5	17	10
800-999	2,845	1,392	1,452	240	828	709	1,069	591	612	1,056	586
800	16	10	*6	*7	*6	*	*	*	*	*	*
801	29	21	*8	*6	15	*	*	*5	*7	*7	11
801.0	9	*7	*	*	*	*	*	*	*	*	*
801.2	12	*8	*	*	*6	*	*	*	*	*	*6
802	40	33	*6	*	25	*7	*	*7	*7	17	*8
802.2	14	13	*	*	10	*	*	*	*	*7	*
802.3	*5	*	*	*	*	*	—	*	*	*	*
802.4	*8	*7	*	*	*	*	*	*	*	*	*
805	83	36	47	*	21	16	44	16	21	28	18
805.0	21	13	*8	*	*9	*	*8	*	*	*8	*5
805.02	*8	*	*	*	*	*	*	*	*	*	*
805.2	22	*7	16	*	*	*	13	*	*7	*6	*
805.4	34	15	19	*	*8	*7	18	*6	*9	13	*7
806	*9	*6	*	*	*	*	*	*	*	*	*
807	39	22	17	*	*8	10	21	12	*7	11	*8
807.0	33	20	13	*	*6	9	18	11	*6	10	*6
807.01	*6	*	*	*	*	*	*	*	*	*	*
807.02	*6	*	*	*	*	*	*	*	*	*	*
807.03	*6	*	*	—	*	*	*	*	*	*	*
807.09	*6	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
808	50	15	35	*	*8	*8	33	9	17	15	*8
808.0	13	*9	*	*	*	*	*6	*	*	*	*
808.2	31	*5	26	*	*	*	23	*5	11	10	*5
812	78	27	51	17	11	13	37	17	15	28	17
812.0	39	9	30	*	*	*9	28	*9	*8	17	*6
812.00	15	*	12	*	*	*	10	*	*	10	*
812.01	12	*	10	—	*	*	10	*	*	*	*
812.09	10	*	*7	*	*	*	*6	*	*	*	*
812.2	10	*5	*5	*	*	*	*	*	*	*	*
812.21	*8	*	*	*	*	*	*	*	*	*	*
812.4	24	10	14	15	*	*	*	*6	*	*7	*8
812.41	15	*6	*9	12	*	*	*	*	*	*	*5
813	60	31	29	*8	20	11	22	13	12	20	15
813.0	13	*5	*8	*	*	*	*5	*	*	*	*
813.01	*6	*	*	*	*	*	*	*	*	*	*
813.2	*5	*	*	*	*	*	*	*	*	*	*
813.4	26	10	16	*	*6	*	14	*7	*7	*7	*5
813.41	*7	*	*5	*	*	*	*	*	*	*	*
813.42	10	*5	*	*	*	*	*5	*	*	*	*
813.44	*9	*	*5	*	*	*	*	*	*	*	*
816	*7	*5	*	*	*	*	*	*	*	*	*
820	329	93	236	*	*7	31	289	68	71	126	64
820.0	80	25	55	*	*	*5	72	20	20	27	11
820.00	*5	*	*	—	*	*	*	*	*	*	*
820.02	*7	*	*	*	*	*	*7	*	*	*	*
820.03	*9	*	*5	—	*	*	*8	*	*	*	*
820.09	*9	16	41	*	*	*	53	15	16	20	*6
820.2	172	47	124	*	*	18	149	33	36	67	36
820.20	*8	*	*5	—	*	*	*7	*	*	*	*
820.21	149	38	110	—	*	14	133	27	31	58	32
820.22	15	*6	*9	*	*	*	9	*	*	*5	*
820.8	74	20	54	*	*	*7	66	15	14	30	15
821	68	34	34	13	19	12	24	10	15	29	13
821.0	40	24	17	10	13	*7	11	*6	*8	19	*8
821.00	*6	*	*	*	*	*	*	*	*	*	*
821.01	34	21	13	9	12	*5	*8	*	*7	16	*7
821.1	*5	*	*	—	*	*	*	*	*	*	*
821.2	21	*5	16	*	*	*	12	*	*6	*7	*
821.23	10	*	*8	*	*	*	*6	*	*	*	*
822	18	10	*8	*	*5	*6	*7	*	*	*8	*
822.0	16	*8	*8	*	*	*5	*7	*	*	*7	*
823	67	38	29	*6	25	21	15	15	16	22	14
823.0	28	12	16	*	*6	12	9	*6	*8	*9	*6
823.00	16	*7	10	*	*	*6	*5	*	*	*5	*
823.02	10	*	*6	*	*	*5	*	*	*	*	*
823.2	18	11	*7	*	*9	*	*	*	*	*7	*5
823.20	*6	*	*	*	*	*	*	*	*	*	*
823.22	11	*6	*	*	*5	*	*	*	*	*	*
823.8	10	*6	*	*	*	*	*	*	*	*	*
823.82	*7	*	*	*	*	*	*	*	*	*	*
824	108	48	59	*	43	36	24	23	30	37	18
824.0	*7	*5	*	*	*	*	*	*	*	*	*
824.2	*6	*	*	*	*	*	*	*	*	*	*
824.4	31	14	18	*	13	10	*8	*7	*7	12	*5
824.6	30	9	21	*	12	10	*8	*7	*8	9	*6
824.8	21	11	10	*	*9	*6	*	*5	*7	*7	*
825	18	13	*6	*	11	*	*	*5	*	*6	*
825.0	*6	*	*	*	*	*	*	*	*	*	*
825.2	*9	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
840	14	*7	*7	*	*	*	*9	*	*	*6	*	
840.4	11	*	*6	—	*	*	*8	*	*	*	*	
844	14	*8	*6	*	*6	*	*	*	*	*	*	
844.8	*6	*5	*	—	*	*	*	*	*	*	*	
847	*8	*	*	*	*	*	*	*	*	*	*	
850	57	39	19	*8	26	9	13	15	16	10	17	
850.0	10	*6	*	*	*	*	*	*	*	*	*	
850.1	18	13	*	*	*8	*	*	*	*5	*	*6	
850.11	18	13	*	*	*8	*	*	*	*5	*	*6	
850.5	21	15	*6	*	11	*	*	*	*5	*	*6	
850.9	*6	*	*	*	*	*	*	*	*	*	*	
851	24	15	*8	*	13	*	*5	*	*7	*8	*6	
851.8	19	13	*6	*	9	*	*	*	*	*7	*6	
852	50	26	24	*	*9	9	29	12	11	16	12	
852.0	17	*9	*8	*	*	*	*7	*	*	*6	*	
852.02	*5	*	*	*	*	*	*	*	*	*	*	
852.2	30	15	15	*	*	*	22	*8	*6	*8	*8	
852.20	*8	*	*	*	*	*	*6	*	*	*	*	
852.21	13	*6	*8	*	*	*	11	*	*	*	*	
853	11	*6	*	*	*	*	*6	*	*	*	*	
853.0	10	*6	*	*	*	*	*6	*	*	*	*	
854	*5	*	*	*	*	*	*	*	*	*	*	
860	25	17	*8	*	12	*8	*	*5	*5	*9	*6	
860.0	15	10	*5	*	*7	*6	*	*	*	*6	*	
861	*8	*6	*	*	*	*	*	*	*	*	*	
861.2	*6	*	*	*	*	*	*	*	*	*	*	
861.21	*5	*	*	*	*	*	*	*	*	*	*	
863	12	10	*	*	*8	*	*	*	*	*6	*	
864	11	*7	*	*	*5	*	*	*	*	*	*	
864.0	*8	*5	*	*	*	*	*	*	*	*	*	
865	16	12	*	*	10	*	*	*	*	*6	*	
865.0	15	11	*	*	10	*	*	*	*	*6	*	
873	28	19	*9	*	9	*6	9	*7	*5	*8	*8	
873.0	11	*7	*	*	*	*	*5	*	*	*	*	
873.4	11	*8	*	*	*	*	*	*	*	*	*	
873.42	*5	*	*	*	*	*	*	*	*	*	*	
874	*5	*	*	*	*	*	*	*	*	*	*	
879	*5	*	*	*	*	*	*	*	*	*	*	
881	*7	*6	*	*	*	*	*	*	*	*	*	
882	*5	*	*	*	*	*	*	*	*	*	*	
883	*5	*	*	*	*	*	*	*	*	*	*	
891	9	*6	*	*	*	*	*	*	*	*	*	
920	*9	*	*	*	*	*	*	*	*	*	*	
922	15	*7	*8	*	*	*	*8	*	*	*6	*	
922.1	*8	*	*	*	*	*	*	*	*	*	*	
924	20	*8	12	*	*	*	14	*	*8	*	*	
924.0	10	*	*7	*	*	*	*8	*	*	*	*	
924.01	*8	*	*6	—	*	*	*7	*	*	*	*	
924.1	*7	*	*	—	*	*	*	*	*	*	*	
933	*6	*	*	*	*	*	*	*	*	*	*	
933.1	*6	*	*	*	*	*	*	*	*	*	*	
935	*9	*5	*	*	*	*	*	*	*	*	*	
935.1	*6	*	*	*	*	*	*	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
941	16	10	*6	9	*5	*	*	*	*	13	*
941.2	10	*6	*	*	*	*	*	*	*	*7	*
941.3	*5	*	*	*	*	*	*	*	*	*	—
942	18	10	*8	*8	*	*	*	*	*	14	*
942.2	9	*5	*	*6	*	*	*	*	*	*8	—
942.3	9	*	*	*	*	*	*	*	*	*6	*
943	15	*9	*6	*6	*7	*	*	*	*	11	*
943.2	*6	*	*	*	*	*	—	*	*	*	*
943.3	*8	*	*	*	*	*	*	*	*	*7	*
944	*8	*7	*	*	*	*	—	*	*	*5	*
945	16	11	*	*	*7	*	*	*	*	10	*
945.2	10	*7	*	*	*	*	*	*	*	*6	*
945.3	*6	*	*	*	*	*	*	*	*	*	*
949	*6	*	*	*5	*	*	—	—	*	*6	—
949.0	*6	*	*	*5	*	*	—	—	*	*6	—
958	*7	*	*	*	*	*	*	*	*	*	*
959	32	19	12	*	12	*7	10	*7	*8	*8	*8
959.0	19	11	*8	*	*8	*	*6	*	*5	*	*
959.01	17	11	*7	*	*7	*	*6	*	*	*	*
959.1	*7	*	*	*	*	*	*	*	*	*	*
962	*6	*	*	*	*	*	*	*	*	*	*
962.3	*6	*	*	*	*	*	*	*	*	*	*
963	*6	*	*	*	*	*	*	*	*	*	*
963.0	*6	*	*	*	*	*	—	*	*	*	*
965	67	25	42	*	46	14	*	14	15	25	14
965.0	26	10	16	*	14	9	*	*	*	11	*5
965.02	*7	*	*	*	*	*	*	*	*	*	*
965.09	12	*	*8	*	*6	*	*	*	*	*	*
965.1	*6	*	*	*	*	*	*	*	*	*	*
965.4	26	9	17	*	23	*	*	*6	*6	10	*
966	12	*	*8	*	*6	*	*	*	*	*	*5
966.3	*8	*	*6	*	*	*	*	*	*	*	*
967	12	*	*9	*	*6	*	*	*	*	*6	*
967.8	*6	*	*	*	*	*	*	*	*	*	*
968	*7	*	*	*	*6	*	*	*	*	*	*
969	75	27	48	*	49	17	*5	16	17	26	15
969.0	16	*	13	*	11	*	*	*	*	*6	*
969.3	*6	*	*	*	*5	*	*	*	*	*	*
969.4	36	14	22	*	22	11	*	*9	*9	12	*6
969.8	10	*	*7	*	*6	*	*	*	*	*	*
970	12	*8	*	*	*7	*	*	*	*	*	*
970.8	11	*7	*	*	*7	*	*	*	*	*	*
972	*8	*	*	*	*	*	*	*	*	*	*
977	15	*6	*9	*	*7	*6	*	*	*	*8	*
977.8	*9	*	*6	*	*	*	*	*	*	*	*
977.9	*6	*	*	*	*	*	*	*	*	*	*
980	*7	*5	*	*	*	*	*	*	*	*	*
980.0	*6	*	*	*	*	*	*	*	*	*	*
989	*8	*5	*	*	*	*	*	*	*	*	*
989.5	*6	*	*	*	*	*	*	*	*	*	*
991	*6	*5	*	*	*	*	*	*	*	*	*
994	*8	*6	*	*	*	*	*	*	*	*	*
995	37	15	22	*	*8	11	15	*8	*8	16	*6
995.1	15	*	10	*	*	*	*6	*	*	*7	*

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of first-listed diagnoses in thousands												
995.2	*6	*	*	—	*	*	*	*	*	*	*	*
995.9	*7	*	*	*	*	*	*	*	*	*	*	*
996	501	240	262	31	90	169	210	107	103	188	102	
996.0	18	13	*5	*	*	*	12	*	*	*7	*	
996.01	9	*6	*	—	*	*	*7	*	*	*	*	
996.04	*7	*6	*	*	*	*	*	*	*	*	*	
996.1	11	*6	*	*	*	*	*	*	*	*	*	
996.2	24	10	14	10	*7	*6	*	*	*5	9	*	
996.3	*6	*	*	*	*	*	*	*	*	*	*	
996.4	104	36	69	*	12	27	65	24	27	33	21	
996.5	12	*5	*7	*	*	*	*	*	*	*5	*	
996.59	*6	*	*	*	*	*	*	*	*	*	*	
996.6	173	89	84	12	32	65	64	36	38	67	32	
996.61	*8	*	*	—	*	*	*5	*	*	*	*	
996.62	84	42	42	*8	15	35	26	21	15	35	13	
996.63	10	*7	*	*	*	*	*	*	*	*	*	
996.64	12	*7	*	*	*	*	*8	*	*	*	*	
996.66	24	10	14	*	*	*8	13	*	*7	*7	*6	
996.67	13	*7	*6	*	*	*5	*	*	*	*	*	
996.68	*5	*	*	*	*	*	*	*	*	*	*	
996.69	11	*	*7	*	*	*	*	*	*	*6	*	
996.7	124	62	62	*	20	46	57	25	22	46	31	
996.72	27	16	11	*	*	*8	16	*6	*6	10	*	
996.73	41	21	20	—	*7	16	18	*7	*	13	16	
996.74	20	10	10	*	*	*8	10	*	*	9	*	
996.77	11	*	*6	—	*	*	*6	*	*	*	*	
996.78	12	*5	*6	*	*5	*	*	*	*	*	*	
996.8	28	14	14	*	11	13	*	*6	*	16	*	
996.81	18	*8	10	*	*8	*8	*	*	*	11	*	
997	109	53	56	*	27	44	34	22	19	40	29	
997.0	*5	*	*	*	*	*	*	*	*	*	*	
997.1	10	*6	*	*	*	*	*	*	*	*	*	
997.2	*9	*	*5	—	*	*	*	*	*	*	*	
997.3	10	*5	*	*	*	*5	*	*	*	*	*	
997.4	42	15	28	*	16	14	12	*8	*8	14	12	
997.5	10	*7	*	*	*	*6	*	*	*	*	*	
997.6	19	12	*7	*	*	*8	*8	*	*	11	*	
997.62	13	*8	*	*	*	*6	*	*	*	*8	*	
997.69	*6	*	*	—	*	*	*	*	*	*	*	
998	251	110	142	10	61	100	80	56	52	95	49	
998.1	60	28	31	*	14	20	23	13	11	22	14	
998.11	31	19	13	*	*7	*8	14	*7	*5	11	*7	
998.12	23	*9	14	*	*6	10	*7	*5	*	*8	*6	
998.13	*5	*	*	*	*	*	*	*	*	*	*	
998.2	11	*	*8	*	*	*	*6	*	*	*	*	
998.3	16	*8	*8	*	*	*5	*7	*	*5	*	*	
998.32	14	*7	*7	*	*	*	*6	*	*	*	*	
998.5	139	61	78	*	34	63	38	30	30	54	24	
998.51	*7	*	*5	*	*	*	*	*	*	*	*	
998.59	132	60	72	*	33	59	36	29	29	52	22	
998.6	*6	*	*	*	*	*	*	*	*	*	*	
998.8	19	*9	10	*	*	*7	*6	*	*	*7	*	
998.83	*7	*	*	*	*	*	*	*	*	*	*	
998.89	12	*	*7	*	*	*	*	*	*	*	*	
999	10	*5	*	*	*	*	*	*	*	*	*	
V01-V83	5,068	403	4,665	82	4,244	259	483	778	1,028	2,001	1,261	
V10	*5	*	*	*	*	*	*	*	*	*	*	
V12	12	*	*8	*	*	*	*	*	*	*	*	
V12.5	*7	*	*6	*	*	*	*	*	*	*	*	

See footnotes at end of table.

Table 45. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of first-listed diagnoses in thousands											
V27	4,136	—	4,136	*5	4,120	10	—	589	802	1,679	1,066
V27.0	3,909	—	3,909	*	3,896	*8	—	536	744	1,609	1,019
V27.1	23	—	23	*	23	*	—	*	*	11	*
V27.2	50	—	50	—	49	*	—	*8	*8	20	13
V27.9	150	—	150	*	148	*	—	41	45	37	27
V30	9	*	*5	9	—	—	—	*	*	*	*
V30.1	9	*	*5	9	—	—	—	*	*	*	*
V42	*8	*	*	*	*	*	*	*	*	*	*
V45	20	*6	13	*	*5	*5	*7	*	*	*6	*
V45.8	9	*	*6	*	*	*	*	*	*	*	*
V45.89	*6	*	*	*	*	*	*	*	*	*	*
V50	*6	*	*5	*	*	*	*	*	*	*	*
V53	12	*6	*5	*	*	*	*7	*	*	*	*
V53.3	10	*6	*	*	*	*	*6	*	*	*	*
V53.31	*6	*	*	*	*	*	*	*	*	*	*
V54	21	*8	13	*	*	*	15	*	*8	*5	*
V54.1	*8	*	*	*	*	*	*6	*	*	*	*
V54.8	*9	*	*6	*	*	*	*7	*	*	*	*
V54.81	*6	*	*	—	*	*	*5	*	*	*	*
V55	40	22	19	*	9	15	13	11	9	13	*7
V55.2	*9	*	*	*	*	*	*	*	*	*	*
V55.3	25	15	11	*	*6	10	*8	*7	*5	*9	*
V56	*7	*	*	*	*	*	*	*	*	*	*
V56.0	*6	*	*	*	*	*	*	*	*	*	*
V57	509	205	305	*	34	115	357	91	153	156	110
V57.1	87	36	50	*	*7	19	61	23	9	36	18
V57.8	421	168	253	*	27	95	296	68	144	118	91
V57.89	421	168	253	*	27	95	296	68	144	118	91
V58	198	105	92	40	27	73	57	47	27	81	43
V58.0	10	*6	*	*	*	*	*6	*	*	*6	*
V58.1	154	85	69	38	22	62	32	38	15	66	34
V58.4	10	*	*6	*	*	*	*6	*	*	*	*
V58.49	*8	*	*5	*	*	*	*	*	*	*	*
V58.8	15	*5	9	*	*	*	10	*	*8	*	*
V58.89	12	*	*8	*	*	*	*9	*	*8	*	*
V59	*7	*	*	*	*	*	*	*	*	*	*
V59.4	*6	*	*	*	*	*	*	*	*	*	*
V66	*5	*	*	—	*	*	*	*	*	*	*
V70	12	*	*7	*	*6	*	*	*	*	*	*
V70.7	10	*	*6	*	*6	*	*	*	*	*	*
V71	10	*7	*	*	*	*	*	*	*	*	*
V71.4	*6	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
All codes	167,881	72,449	95,432	11,652	40,079	41,798	74,352	39,876	33,668	64,896	29,441
001-139	6,250	3,025	3,225	458	1,032	1,658	3,101	1,528	1,096	2,385	1,240
003	*25	*	*	*	*	*	*	*	*	*	*
008	698	243	455	123	81	107	387	184	142	263	109
008.4	440	157	283	*	*46	70	310	145	97	142	56
008.45	417	148	269	*	*43	66	298	137	95	137	48
008.6	73	34	39	71	*	*	*	*	*14	33	*20
008.61	63	33	30	62	*	*	*	*	*14	31	*13
008.8	176	50	126	37	34	36	70	31	31	86	28
009	67	*32	35	*	*32	*	*	*	*	*16	*
009.0	28	*	*16	*	*	*	*	*	*	*	*
009.1	*34	*	*	*	*	*	*	*	*	*	*
011	*50	*	*	*	*	*	*	*	*	*	*
034	30	*18	*13	*13	*	*	*	*	*	*	*
034.0	29	*17	*11	*12	*	*	*	*	*	*	*
038	3,438	1,652	1,786	*37	264	816	2,321	817	681	1,223	717
038.0	201	100	101	*	*	*	163	*74	*46	*48	*33
038.1	760	406	354	*	*	197	525	183	121	282	174
038.11	470	295	175	*	*	*115	329	*114	*68	170	118
038.19	225	*86	*139	*	*	*	142	*	*	*72	*
038.2	*45	*	*	-	*	*	*	*	*	*	*
038.4	771	327	444	*	*51	194	510	197	177	246	151
038.40	*39	*	*	*	*	*	*	*	*	*	*
038.42	425	130	295	*	*	105	297	94	95	136	100
038.43	*65	*	*	*	*	*	*	*	*	*	*
038.49	219	114	105	*	*	*54	137	*70	*46	73	*
038.8	132	71	*60	*	*	*	89	*	*	*73	*
038.9	1,487	697	789	*	127	351	999	327	299	555	305
041	69	*	*40	*	*	*	*	*	*	*	*
042	609	372	237	*	309	270	*	203	*60	316	*
047	124	63	61	52	47	*17	*	*17	*17	61	30
047.8	*15	*	*	*	*	*	-	*	*	*	*
047.9	109	58	52	42	43	*	*	*16	*15	53	25
053	95	*	75	*	*	*31	57	*	*	*34	*
053.1	*30	*	*	*	*	*	*	*	*	*	*
053.9	*41	*	*32	*	*	*	*	*	*	*	*
054	*35	*	*	*	*	*	*	*	*	*	*
070	146	86	60	*	*31	94	*	*17	*	48	*
070.5	53	*22	*31	-	*	*40	*	*	*	*	*
070.51	*37	*	*	-	*	*31	*	*	*	*	*
070.54	*17	*	*	-	*	*	*	*	*	*	*
075	24	*12	*	*	*14	*	-	*	*	*	*
079	160	86	75	67	53	*18	*22	38	24	57	41
079.9	144	80	64	58	49	*15	*22	34	*21	51	38
079.99	144	80	64	58	49	*15	*22	34	*21	51	38
112	157	*63	94	*	*	*51	81	*	*	*58	*41
112.8	*29	*	*	*	*	*	*	*	*	*	*
135	43	*	*29	*	*	*24	*	*	*	*	*
140-239	9,420	4,334	5,086	275	1,167	3,262	4,716	2,360	1,815	3,668	1,576
150	136	104	*	-	*	*53	*80	*	*	*	*
150.5	*63	*57	*	-	*	*	*	*	*	*	*
150.9	*37	*	*	-	*	*	*	*	*	*	*
151	248	162	*86	-	*	*63	172	*62	*	*72	*73
151.0	109	*100	*	-	*	*	*75	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
153	1,057	490	568	—	*35	234	789	293	230	407	127
153.0	*37	*	*	—	*	*	*	*	*	*	*
153.1	96	*	*57	—	*	*	*85	*	*	*	*
153.2	*69	*	*	—	*	*	*	*	*	*	*
153.3	258	141	117	—	*	*61	185	*66	*66	95	*
153.4	237	99	138	—	*	*46	185	*72	*54	*88	*
153.6	179	75	103	—	*	*47	129	*	*	72	*43
153.9	99	*55	*44	—	*	*	74	*	*	*44	*
154	294	162	132	*	*	112	156	73	72	115	*35
154.0	93	*45	*49	—	*	*	*56	*	*	*	*
154.1	179	105	*74	*	*	76	86	*	*55	*69	*
155	121	87	*33	*	*	*46	68	*	*	74	*
155.0	95	74	*	*	*	*36	*53	*	*	*61	*
156	*69	*	*	—	*	*	*	*	*	*	*
157	228	103	125	*	*	69	154	68	*43	74	*41
157.0	90	*	*49	—	*	*	*65	*	*	*	*
157.8	*35	*	*	—	*	*	*	*	*	*	*
157.9	69	*19	*49	—	*	*	*42	*	*	*	*
161	*44	*34	*	—	*	*	*	*	*	*	*
162	1,002	520	482	*	85	308	609	238	243	402	119
162.2	*54	*	*	—	*	*	*	*	*	*	*
162.3	299	150	149	*	*	80	203	*77	*54	127	*
162.5	204	99	105	—	*	*45	145	*	*60	91	*
162.8	136	80	*55	—	*	*43	*78	*51	*	*	*
162.9	278	143	135	—	*	99	144	*56	84	115	*
170	*38	*	*	*	*	*	*	*	*	*	*
171	*61	*	*	*	*	*	*	*	*	*	*
173	*23	*	*	*	*	*	*	*	*	*	*
174	225	—	225	*	28	102	93	42	47	84	51
174.4	23	—	23	—	*	*	*	*	*	*	*
174.8	88	—	88	—	*	44	33	*29	*	*14	*31
174.9	91	—	91	—	*	38	40	*	*23	52	*
180	55	—	55	—	*23	*	*	*	*	*	*
180.9	*29	—	*29	—	*	*	*	*	*	*	*
182	164	—	164	—	*	76	80	*37	43	45	*39
182.0	162	—	162	—	*	75	79	*37	43	44	*39
183	196	—	196	*	*	82	102	*52	*	*64	*44
183.0	188	—	188	*	*	77	99	*52	*	*57	*44
184	*33	—	*33	—	*	*	*	*	*	*	*
185	252	252	—	—	*	110	142	39	59	96	57
188	213	173	*40	*	*	*41	168	102	*40	*50	*21
188.8	102	85	*	—	*	*	85	*56	*	*	*
188.9	61	*47	*	*	*	*	*38	*	*	*	*
189	237	146	91	*	*	92	127	*49	*40	100	*48
189.0	209	129	80	*	*	84	107	*40	*39	88	*43
191	211	108	104	*	*41	93	*51	*58	*	108	*25
191.3	*32	*	*	*	*	*	*	*	*	*	*
191.9	*59	*	*	*	*	*	*	*	*	*	*
193	57	*16	42	*	27	18	*	*11	*	*28	*
196	58	*37	*21	*	*	*	*	*	*	*	*
197	757	307	450	*	*44	287	424	201	135	290	132
197.0	130	80	50	*	*	*60	58	*	*	*50	*
197.2	185	66	119	*	*	*69	112	*46	*	*73	*
197.6	173	64	109	—	*	*60	98	*56	*	61	*
197.7	185	60	125	—	*	68	107	*49	*39	77	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
198	528	238	289	*	*43	209	271	168	91	173	95
198.3	195	85	110	*	*	98	82	71	*27	61	*37
198.5	196	90	106	*	*	*72	106	*51	*31	*67	*47
198.8	75	*37	*38	*	*	*	*44	*	*	*	*
198.89	68	*35	*	*	*	*	*41	*	*	*	*
199	157	75	82	—	*	*71	*54	*	*26	*74	*
199.0	93	*	*55	—	*	*54	*	*	*	*	*
199.1	64	*37	*	—	*	*	*	*	*	*	*
200	93	*	*	*	*	*	*47	*	*	*	*
200.0	*64	*	*	*	*	*	*	*	*	*	*
201	*65	*	*	*	*	*	*	*	*	*	*
202	242	141	100	*	*48	*70	111	*62	*46	107	*
202.8	198	112	86	*	*32	*62	100	*58	*36	87	*
202.80	104	*64	*40	*	*	*	57	*	*	*43	*
203	201	129	72	—	*	133	*57	*	*	*101	*
203.0	184	113	71	—	*	120	*56	*	*	*101	*
203.00	179	113	66	—	*	115	*56	*	*	*100	*
204	205	115	90	*82	*	*	76	*	*	*72	*56
204.0	115	*68	*	*56	*	*	*	*	*	*	*
204.00	105	*64	*	*49	*	*	*	*	*	*	*
204.1	64	*	43	—	*	*	53	*	*	*	*
204.10	62	*	*41	—	—	*	52	*	*	*	*
205	303	148	155	*	*	*117	*137	*	*51	*140	*
205.0	256	*120	*136	*	*	*	*131	*	*	*114	*
205.00	251	*116	*135	*	*	*	*131	*	*	*114	*
210	*19	*	*	*	*	*	*	*	*	*	*
210.2	*9	*	*	—	*	*	*	*	*	*	*
211	234	113	121	*	*	55	153	71	53	83	*27
211.3	151	69	82	*	*	46	95	*39	*40	57	*
218	535	—	535	—	271	255	*	93	104	236	101
218.0	72	—	72	—	34	38	*	*17	*14	27	*14
218.1	161	—	161	—	75	83	*	33	33	74	*21
218.2	55	—	55	—	26	28	*	*13	*	24	*
218.9	247	—	247	—	135	106	*	31	48	111	57
220	107	—	107	*	30	34	41	*21	*26	45	*16
225	150	*76	73	*	*40	66	*42	*24	*	*71	*36
225.2	112	*59	53	*	*	*50	*	*	*	*59	*22
226	*13	*	*11	—	*	*	*	*	*	*	*
227	65	*25	40	*	*	*33	*20	*	*	*18	*
227.1	*25	*	*21	*	*	*	*	*	*	*	*
227.3	*28	*	*	—	*	*	*	*	*	*	*
228	*28	*	*	*	*	*	*	*	*	*	*
228.0	*24	*	*	*	*	*	*	*	*	*	*
233	58	*	50	—	19	25	*14	*12	*	*19	*17
233.0	25	*	25	—	*	*12	*	*	*	*	*
233.1	20	—	20	—	*13	*	*	*	*	*	*
235	58	*33	*25	*	*	*	*33	*	*	*	*
235.2	*33	*	*	*	*	*	*	*	*	*	*
236	*23	*	*	—	*	*	*	*	*	*	*
237	*32	*	*	*	*	*	*	*	*	*	*
238	77	*40	*37	*	*	*	*43	*	*	*39	*
238.7	57	*38	*	*	*	*	*37	*	*	*34	*
239	75	*31	*44	*	*	*	*51	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
240-279	6,957	2,963	3,994	570	1,331	1,944	3,112	1,799	1,301	2,723	1,134
241	33	*	28	*	*15	*11	*	*	*	*16	*
241.0	*8	*	*7	*	*	*	*	*	*	*	*
241.1	22	*	*19	—	*	*	*	*	*	*	*
242	37	*	27	*	*	*	*	*	*	*10	*
242.0	*16	*	*	*	*	*	*	*	*	*	*
242.9	*17	*	*	*	*	*	*	*	*	*	*
242.90	*16	*	*	*	*	*	*	*	*	*	*
244	67	*	49	—	*	*14	46	*25	*17	*15	*
244.9	54	*	42	—	*	*11	36	*	*14	*14	*
250	2,825	1,435	1,390	81	597	1,020	1,127	737	403	1,193	492
250.0	422	199	223	23	69	160	169	128	74	152	68
250.00	203	104	98	*	35	92	71	61	31	78	33
250.01	45	*16	29	*15	*	*	*	*	*14	*17	*
250.02	138	67	71	*	*17	56	65	38	*22	46	31
250.03	37	*	*25	*	*	*	*	*	*	*	*
250.1	420	208	213	49	230	98	*43	82	66	196	77
250.10	86	54	*33	*	45	*28	*	*	*	*43	*
250.11	190	93	97	37	108	33	*	35	39	76	40
250.12	54	*20	*34	*	*	*	*	*	*	*	*
250.13	90	41	49	*	63	*	*	*18	*	48	*
250.2	77	*31	*46	—	*	*31	*37	*	*	*34	*
250.20	*43	*	*	—	*	*	*	*	*	*	*
250.22	*28	*	*	—	*	*	*	*	*	*	*
250.4	181	104	77	—	*	60	96	*30	*	67	62
250.40	118	75	*43	—	*	*34	76	*	*	*38	49
250.42	*42	*	*	—	*	*	*	*	*	*	*
250.6	356	162	194	*	78	150	127	105	47	165	*39
250.60	183	92	91	—	*	77	98	*65	*	71	*
250.61	59	*	*37	*	*	*	*	*	*	*	*
250.62	77	*35	*41	—	*	*37	*	*	*	*	*
250.63	*38	*	*	—	*	*	*	*	*	*	*
250.7	482	290	193	*	*	172	279	135	*58	206	*85
250.70	293	161	133	—	*	74	199	83	*40	103	*68
250.71	*83	*50	*	*	*	*	*	*	*	*	*
250.72	*92	*66	*	—	*	*	*	*	*	*	*
250.8	803	409	395	*	136	310	357	212	119	334	138
250.80	497	250	248	*	*30	196	272	113	86	211	87
250.81	146	71	75	*	*74	45	*26	*46	*	51	*
250.82	128	*73	55	—	*	*59	56	*44	*	67	*
250.83	*33	*	*	*	*	*	*	*	*	*	*
250.9	36	*16	*	*	*	*	*	*	*	*21	*
251	62	*35	*27	*	*	*	*	*	*	*	*
251.2	47	*	*21	*	*	*	*	*	*	*	*
252	31	*	*18	*	*	*	*	*	*	*	*
252.0	*25	*	*	—	*	*	*	*	*	*	*
253	104	*31	73	*	*	*	72	*	*	*43	*
253.6	84	*26	57	—	*	*	70	*	*	*36	*
255	*49	*	*	*	*	*	*	*	*	*	*
255.4	*41	*	*	*	*	*	*	*	*	*	*
263	73	*29	*44	*	*	*	*43	*	*	*33	*
263.9	61	*25	*35	*	*	*	*41	*	*	*	*
272	84	43	41	*	22	46	16	23	24	23	14
272.0	36	17	19	*	*8	20	*	*10	*10	*	*
272.4	41	23	17	*	*10	23	*	*9	*12	*10	*
274	77	47	*	—	*	*28	44	*30	*	*33	*
274.0	56	*35	*	—	*	*	*34	*	*	*	*
275	87	*41	45	*	*	*29	47	*45	*	*22	*
275.4	76	*37	39	*	*	*	*44	*41	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
275.42	*57	*	*27	—	*	*	*	*	*	*	*
276	2,704	1,030	1,674	302	293	518	1,591	614	617	1,072	402
276.0	64	*	40	*	*	*	49	*	*	*22	*
276.1	568	204	364	*	*39	164	355	113	103	219	133
276.2	45	*	*24	*	*	*	*	*	*	*22	*
276.5	1,699	664	1,035	267	187	260	984	409	423	679	188
276.7	127	59	68	*	*	*21	86	*	*19	51	*30
276.8	142	33	108	*	37	42	62	34	28	57	*22
276.9	*32	*	*20	*	*	*	*22	*	*	*	*
277	190	90	100	*99	75	*	*	*64	*55	*31	*
277.0	161	*69	*91	*91	*63	*	—	*55	*	*	*
277.02	147	*65	*82	*80	*60	*	—	*52	*	*	*
278	422	75	347	*	255	159	*	110	63	163	86
278.0	403	72	331	*	249	149	*	105	59	154	85
278.00	34	*	*21	*	*25	*	*	*	*	*14	*
278.01	369	60	309	*	224	141	*	97	55	140	77
278.1	*15	*	*14	—	*	*	*	*	*	*	*
280-289	2,099	797	1,301	234	649	470	746	517	341	935	305
280	307	87	220	*	33	78	192	84	73	127	23
280.0	139	48	91	—	*14	36	90	*23	42	66	*
280.9	161	38	122	*	*19	41	97	60	*29	58	*14
282	469	196	273	67	356	*44	*	168	70	214	*
282.6	439	188	251	58	339	*42	*	159	64	201	*
282.62	374	144	230	43	292	*39	—	132	58	173	*
283	64	*	41	*	*	*	*29	*	*	*34	*
283.0	*33	*	*	*	*	*	*	*	*	*	*
283.9	*21	*	*	*	*	*	*	*	*	*16	*
284	218	116	102	*	*	81	58	*39	*	94	*68
284.8	203	110	92	*	*	81	53	*37	*	81	*68
285	470	118	352	*	140	81	244	89	66	245	70
285.1	45	*	*29	—	*	*	37	*	*	*	*
285.2	77	*22	55	*	*	*	55	*	*	*35	*
285.22	*19	*	*	—	*	*	*	*	*	*	*
285.29	*49	*	*	*	*	*	*37	*	*	*	*
285.8	*21	*	*	*	*	*	*11	*	*	*	*
285.9	327	74	253	*	*128	52	141	58	40	184	44
286	74	*28	46	*	*	*	48	*	*18	*34	*
286.7	*22	*	*	—	*	*	*	*	*	*	*
287	151	60	91	25	*21	57	48	*30	*29	61	*31
287.3	61	*21	40	*12	*	*	*	*	*	*25	*
287.5	58	*22	*35	*	*	*32	*19	*	*	*20	*
288	270	132	138	64	*32	69	105	73	45	99	54
288.0	205	104	101	*41	*27	59	78	61	*39	74	*30
288.8	47	*20	*27	*	*	*	*	*	*	*19	*
289	62	*37	*25	*23	*	*	*	*	*	*	*13
289.3	*20	*	*	*	*	*	*	*	*	*	*
290-319	16,464	8,271	8,193	1,319	8,010	4,793	2,343	5,824	2,769	6,234	1,637
290	361	193	168	—	*	*	280	149	*54	134	*
290.3	*38	*	*	—	—	—	*38	*	*	*	*
290.4	232	136	96	—	—	*	154	*94	*	112	*
290.40	*53	*	*	—	—	*	*51	*	*	*	*
290.41	*84	*	*	—	—	*	*	*	*	*	—
290.42	*65	*	*	—	—	*	*	*	*	*	*
291	734	509	225	—	310	341	83	174	270	147	143
291.0	165	135	*30	—	78	64	*	*	*	*77	*27
291.8	532	345	187	—	222	258	*52	136	236	55	104
291.81	517	336	181	—	211	254	*52	130	235	53	100

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
292	389	216	174	*	226	103	58	173	76	74	68
292.0	233	135	98	*	161	62	*	103	54	*28	*48
292.8	129	60	69	*	50	*34	44	63	*	36	*
292.81	61	*18	*44	—	*	*	*38	*	*	*24	*
292.84	55	*33	*23	*	43	*	*	*33	*	*	*
293	114	41	73	*	*	*	68	*32	*29	*35	*
293.0	68	*29	*40	*	*	*	51	*	*23	*24	*
293.8	*44	*	*	—	*	*	*	*	*	*	*
294	347	229	118	—	*	*	320	89	*36	102	*119
294.8	322	214	108	—	*	*	311	86	*35	90	*
295	3,557	1,848	1,710	*	1,809	1,389	335	1,373	484	1,427	274
295.3	1,272	757	515	*	649	482	140	451	197	497	127
295.30	495	304	191	—	218	207	*70	127	115	211	*42
295.32	232	133	*98	—	132	*73	*	*80	*	108	*
295.34	503	301	203	*	279	191	*	227	*58	168	*
295.6	112	*56	*	—	*	*58	*	*	*	*	*
295.64	*66	*	*	—	*	*	*	*	*	*	*
295.7	1,701	785	916	*	819	727	134	677	222	714	89
295.70	1,210	540	670	*	571	547	*72	446	170	526	68
295.74	417	198	219	*	200	157	*	205	*	156	*
295.9	366	177	188	*	237	95	*	168	*	125	*
295.90	166	95	71	*	92	*52	*	*41	*	102	*
295.94	*95	*	*	—	*	*	*	*	*	*	*
296	6,316	2,726	3,590	545	3,247	1,762	762	2,091	1,196	2,626	402
296.2	1,167	513	654	96	613	257	201	241	257	614	54
296.20	540	254	286	*31	300	124	86	130	168	223	*19
296.22	118	*47	72	*	67	*	*	*	*	97	*
296.23	254	95	159	*33	130	59	*33	50	37	153	*
296.24	246	111	135	*	109	*51	*61	*53	*41	140	*
296.3	1,981	781	1,199	*37	937	672	335	782	358	741	99
296.30	390	169	220	*	206	141	*	150	112	114	*
296.32	92	*33	59	*	47	*38	*	*	*	*47	*
296.33	1,017	424	593	*	490	380	129	402	153	404	*58
296.34	436	153	283	*	182	107	139	200	75	147	*
296.4	561	223	338	*	310	178	*68	218	100	178	*66
296.40	250	88	161	*	136	*83	*	*101	*46	*72	*
296.43	*78	*	*	*	*	*	*	*	*	*	*
296.44	211	86	126	*	123	*72	*	*81	*	*64	*
296.5	633	270	363	*	314	237	*69	246	134	225	*28
296.50	295	132	163	*	162	111	*	88	81	114	*
296.53	148	*64	84	*	82	*47	*	76	*	*	*
296.54	162	*70	92	*	*64	*64	*	*75	*	*56	*
296.6	684	342	342	162	367	110	*	165	99	406	*
296.60	325	172	152	*	176	*55	*	*60	*59	202	*
296.63	110	*52	*58	*	67	*	*	*36	*	61	*
296.64	240	116	124	*58	118	*	*	*64	*	141	*
296.7	649	287	362	74	370	182	*	149	134	263	*
296.8	300	118	183	*41	167	74	*	*48	87	148	*
296.80	182	80	101	*	86	*52	*	*	*42	114	*
296.89	113	*36	77	*	77	*	*	*	*44	*	*
296.9	320	187	133	*117	161	*38	*	229	*	*47	*
296.90	312	181	130	*117	156	*	*	224	*	*45	*
297	*69	*	*	—	*	*	*	*	*	*	*
298	707	313	394	*	371	158	148	221	102	236	148
298.9	696	310	386	*	367	156	144	218	98	234	146
299	*49	*	*	*	*	*	—	*	*	*	*
300	393	91	302	*	142	96	41	123	43	190	37
300.0	108	43	65	*	29	42	*26	43	*15	*36	*14
300.00	68	*32	36	*	*14	*34	*11	*22	*	*33	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
300.01	*18	*	*	*	*	*	*	*	*	*	*
300.1	*37	*	*31	*	*18	*	*	*	*	*	*
300.11	*22	*	*	*	*	*	*	*	*	*	*
300.4	190	*22	168	*	*61	*21	*	*42	*	*126	*
300.9	*14	*	*	*	*	*	*	*	*	*	*
303	640	428	212	*	300	269	70	200	126	213	101
303.0	134	72	62	—	53	72	*	44	*21	*27	42
303.00	40	*15	*25	—	*15	*25	*	*	*	*	*
303.01	87	55	*32	—	33	45	*	41	*	*	*
303.9	506	356	150	*	246	197	61	157	105	185	59
303.90	173	119	54	—	95	60	*	*	*	120	*
303.91	328	233	95	*	149	136	*43	134	92	65	*37
304	572	355	217	*	419	145	*	138	*61	249	123
304.0	245	150	95	—	174	69	*	77	*	67	80
304.00	50	*	*	—	*35	*	*	*	*	*24	*
304.01	189	123	67	—	132	55	*	69	*	*42	59
304.1	*37	*	*	—	*	*	*	*	*	*	*
304.2	89	66	*	*	72	*	*	*	*	*57	*
304.20	*39	*	*	—	*	*	—	*	*	*36	*
304.21	*49	*31	*	*	*39	*	*	*	*	*	*
304.7	*39	*26	*	—	*	*	*	*	*	*	*
304.71	*34	*	*	—	*	*	*	*	*	*	*
304.8	*53	*	*	—	*41	*	—	*	*	*42	*
304.80	*43	*	*	—	*	*	—	*	*	*40	*
304.9	*34	*	*	—	*	*	*	*	*	*27	*
304.90	*31	*	*	—	*	*	*	*	*	*27	*
305	250	170	80	*	135	100	*	48	47	111	45
305.0	110	79	30	*	44	56	*	*22	27	51	*10
305.00	85	64	*21	*	35	44	*	*	22	46	*
305.01	*22	*	*	—	*	*	*	*	*	*	*
305.1	39	22	17	—	23	15	*	*	*9	20	*
305.5	*22	*	*	—	*	*	—	*	*	*	*
305.6	25	*18	*	—	*15	*	*	*	*	*	*
305.60	*20	*15	*	—	*	*	*	*	*	*	*
305.9	*39	*24	*	*	*33	*	*	*	*	*	*
305.90	*28	*	*	*	*22	*	*	*	*	*	*
307	68	*	*52	*	*	*	*	*	*	*	*
309	336	143	193	*48	214	64	*	191	49	60	35
309.0	94	56	37	*	60	*	*	64	*	*	*
309.2	47	*20	*27	*	*28	*	*	*	*	*	*
309.28	46	*	*27	*	*27	*	*	*	*	*	*
309.4	61	*32	*29	*	46	*	*	*47	*	*	*
309.8	119	*	91	*	67	*	*	*	*33	*	*
309.81	118	*	90	*	67	*	—	*	*33	*	*
310	*42	*	*	*	*	*	*	*	*	*	*
311	837	391	446	56	520	190	71	496	87	204	50
312	220	163	*58	*112	97	*	*	118	*	*73	*
312.3	155	113	*	*	87	*	*	86	*	*58	—
312.30	97	*67	*	*	*53	*	*	*50	*	*	—
313	146	*118	*	*111	*	—	—	*	*	*	*
313.8	*128	*99	*	*110	*	—	—	*	*	*	*
313.81	*126	*98	*	*109	*	—	—	*	*	*	*
314	197	160	*	153	*	*	—	*75	*	*101	*
314.0	197	160	*	153	*	*	—	*75	*	*101	*
314.01	195	159	*	152	*	—	—	*73	*	*101	—
320-389	2,918	1,354	1,564	372	562	630	1,354	663	596	1,118	542
320	*69	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
322	*36	*	*	*	*	*	*	*	*	*	*
322.9	*35	*	*	*	*	*	*	*	*	*	*
324	*100	*80	*	*	*	*	*	*	*	*	*
324.1	*88	*74	*	—	*	*	*	*	*	*	*
331	670	312	359	*	*	*63	562	137	159	243	130
331.0	535	234	302	*	*	*	506	103	129	216	*87
331.3	*37	*	*	*	*	*	*	*	*	*	*
331.4	64	*41	*23	*	*	*	*	*	*	*	*
332	99	47	*52	—	*	*	90	*29	*	*39	*
332.0	96	47	*49	—	*	*	88	*28	*	*39	*
333	49	*	*27	*	*	*	*	*	*	*22	*
336	*31	*	*	*	*	*	*	*	*	*	*
337	*25	*	*	*	*	*	*	*	*	*	*
340	125	*24	101	*	37	65	*	*34	*26	42	*
343	*16	*	*	*	*	*	*	*	*	*	*
344	*33	*	*	*	*	*	*	*	*	*	*
345	273	152	122	98	82	61	*32	101	*35	82	55
345.1	*24	*	*	*	*	*	*	*	*	*	*
345.10	*19	*	*	*	*	*	*	*	*	*	*
345.3	101	70	30	*22	*35	*	*	*	*	*48	*17
345.4	*27	*	*	*	*	*	*	*	*	*	*
345.9	61	*37	*24	*31	*	*	*	*	*	*	*
345.90	45	*31	*	*	*	*	*	*	*	*	*
346	173	28	145	*	103	49	*16	37	47	74	*14
346.2	*14	*	*	*	*	*	*	*	*	*	*
346.8	38	*	34	*	*24	*	*	*	*	*20	*
346.80	*22	*	*18	*	*	*	*	*	*	*	*
346.9	108	*12	95	*	65	33	*	*23	34	43	*
346.90	62	*	54	*	36	*18	*	*18	*14	25	*
346.91	46	*	41	*	*30	*16	*	*	*	*18	*
348	342	135	207	*	56	*48	171	77	*42	142	*80
348.2	*29	*	*27	*	*24	*	—	*	*	*	*
348.3	184	*86	98	*	*	*	151	*	*	102	*
348.30	90	*	*39	*	*	*	*	*	*	*70	*
348.31	*82	*	*49	—	*	*	*77	*	*	*	*
349	62	*	*43	*	*	*	*	*	*	*29	*
349.8	*37	*	*	*	*	*	*	*	*	*	*
349.82	*31	*	*	—	*	*	*	*	*	*	*
350	*23	*	*	*	*	*	*	*	*	*	*
350.1	*23	*	*	—	*	*	*	*	*	*	*
351	*15	*	*	*	*	*	*	*	*	*	*
351.0	*12	*	*	*	*	*	*	*	*	*	*
355	*48	*	*41	*	*	*	*	*	*	*	*
357	91	*74	*	*	*	*	*	*	*	*	*
357.0	*49	*	*	*	*	*	*	*	*	*	*
358	*115	*	*	*	*	*	*	*	*	*	*
358.0	*114	*	*	*	*	*	*	*	*	*	*
362	*27	*	*	*	*	*	*	*	*	*	*
368	*24	*	*	*	*	*	*	*	*	*16	*
376	*33	*	*	*	*	*	*	*	*	*	*
376.0	*32	*	*	*	*	*	*	*	*	*	*
376.01	*32	*	*	*	*	*	*	*	*	*	*
380	*25	*	*	*	*	*	*	*	*	*	*
380.1	*24	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
382	23	*16	*	21	*	*	*	*	*	*	*
382.9	22	*15	*	*20	*	*	*	*	*	*	*
386	96	36	60	*	*	26	63	35	26	*24	*
386.1	48	*16	32	*	*	*16	29	*19	*15	*	*
386.11	30	*9	*21	—	*	*	*19	*	*11	*	*
386.3	39	*19	*20	—	*	*	*27	*	*	*	*
386.30	35	*	*18	—	*	*	*26	*	*	*	*
390-459	29,823	14,607	15,216	215	1,837	8,099	19,672	7,289	6,024	11,598	4,912
394	*45	*	*40	—	*	*	*	*	*	*	*
396	111	*53	57	*	*	*	81	*	*21	*28	*
396.3	*27	*	*	—	*	*	*	*	*	*	*
396.8	*35	*	*	—	—	*	*	*	*	*	*
398	202	*55	147	—	*	*	174	62	*32	61	*47
398.9	202	*55	147	—	*	*	174	62	*32	61	*47
398.91	199	*55	144	—	*	*	174	62	*32	58	*47
401	787	296	491	*	114	309	361	210	135	337	105
401.0	97	34	63	*	*16	39	42	*	*15	42	*18
401.9	684	260	424	*	96	268	316	182	121	294	87
402	297	109	188	—	*	61	221	58	60	153	*26
402.9	275	102	173	—	*	49	214	55	57	139	*24
402.90	24	*	*15	—	*	*	*	*	*	*	*
402.91	251	92	158	—	*	43	198	*49	54	128	*
403	598	297	301	*	102	236	248	145	103	226	125
403.0	47	*	*32	*	*	*	*	*	*	*	*
403.01	*45	*	*31	*	*	*	*	*	*	*	*
403.9	546	280	266	*	85	217	239	136	96	206	109
403.91	530	274	256	*	83	208	235	134	93	200	103
404	212	94	119	*	*	*51	152	*	*51	81	*
404.9	175	73	102	*	*	*46	123	*	*47	*62	*
404.93	143	*52	90	*	*	*	106	*	*44	*49	*
410	4,218	2,414	1,805	*	144	1,238	2,833	957	917	1,615	730
410.0	95	57	*38	—	*	*34	54	*	*	*39	*
410.01	94	56	*38	—	*	*34	54	*	*	*39	*
410.1	441	272	169	—	*34	178	229	88	122	127	103
410.11	434	266	169	—	*32	176	226	88	119	124	103
410.2	199	*179	*	—	*	*	*139	*	*	*130	*
410.21	199	*179	*	—	*	*	*139	*	*	*	*
410.3	*37	*25	*	—	*	*	*	*	*	*	*
410.31	*37	*25	*	—	*	*	*	*	*	*	*
410.4	497	283	213	—	*23	212	262	103	99	211	83
410.41	491	282	208	—	*23	210	258	101	99	207	83
410.5	41	*18	*	—	*	*	*29	*	*	*	*
410.51	41	*18	*	—	*	*	*29	*	*	*	*
410.7	2,359	1,284	1,075	—	59	586	1,713	650	470	859	379
410.70	*41	*	*	—	*	*	*32	*	*	*	*
410.71	2,273	1,243	1,030	—	58	572	1,643	617	465	819	373
410.72	*44	*	*	—	*	*	*	*	*	*	—
410.8	116	51	*65	*	*	*55	59	*	*	*66	*
410.81	114	50	*65	*	*	*55	57	*	*	*64	*
410.9	421	232	189	*	*	82	329	51	145	156	70
410.91	390	223	167	*	*	69	312	48	143	133	66
411	244	112	132	*	22	96	125	57	49	100	38
411.1	216	101	115	*	*18	89	107	48	48	89	31
411.8	*21	*	*	*	*	*	*	*	*	*	*
411.89	*20	*	*	*	*	*	*	*	*	*	*
412	*10	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
413	112	46	66	*	*	37	67	27	30	41	*13
413.9	110	45	65	*	*	36	66	*26	30	40	*13
414	3,869	2,313	1,556	*	135	1,422	2,269	858	750	1,573	687
414.0	3,743	2,253	1,489	*	134	1,389	2,214	817	741	1,549	636
414.00	411	254	157	—	*13	148	249	119	58	190	43
414.01	3,169	1,900	1,269	*	120	1,201	1,845	665	650	1,302	553
414.02	129	70	59	—	*	25	103	*22	*28	*42	36
414.8	71	47	*24	*	*	*	45	*37	*	*	*
415	752	347	406	—	140	211	402	154	191	252	155
415.1	746	341	405	—	139	210	396	153	191	247	154
415.11	*43	*	*	—	*	*	*	*	*	*	*
415.19	703	317	385	—	129	196	377	147	176	232	147
416	132	58	74	*	*	58	*53	*	*42	*36	*
416.8	*54	*	*38	*	*	*	*	*	*	*	*
416.9	*47	*40	*	—	*	*	*	*	*	*	*
420	46	*23	*23	*	*	*	*	*	*	*14	*
420.9	46	*23	*23	*	*	*	*	*	*	*14	*
420.90	*28	*	*	*	*	*	*	*	*	*	*
421	214	*136	*	*	*	*	*	*	*	*	*
421.0	208	*135	*	*	*	*	*	*	*	*	*
423	96	38	*58	*	*	*31	*56	*19	*	*	*
423.9	72	*22	*49	*	*	*	*	*	*	*	*
424	550	271	280	*	45	127	373	150	98	182	121
424.0	187	69	118	*	*18	60	107	60	*31	60	*36
424.1	326	188	137	*	*	59	247	82	64	104	75
425	123	70	53	*	*24	46	40	*32	*	52	*22
425.4	100	60	40	*	*19	40	36	*30	*	43	*15
426	184	85	99	*	*	*25	143	54	44	57	29
426.0	96	43	52	—	*	*	80	*27	*20	*31	*18
426.1	55	27	*28	*	*	*	46	*19	*	*	*
426.13	25	*	*	*	*	*	*22	*	*	*	*
427	2,822	1,202	1,620	*	211	543	2,045	761	514	1,098	449
427.0	72	37	35	*	*	*16	50	*17	*18	*23	*15
427.1	289	162	126	*	*	81	191	56	30	163	*40
427.3	1,554	645	910	*	68	339	1,146	444	332	586	193
427.31	1,406	561	844	*	61	286	1,057	408	298	524	175
427.32	149	83	65	*	*	53	88	35	34	61	*18
427.4	*36	*	*	*	*	*	*	*	*	*	*
427.41	*36	*	*	*	*	*	*	*	*	*	*
427.5	112	*19	*	*	*	*	*25	*	*	*	*
427.6	23	*	*16	*	*	*	*13	*	*	*	*
427.69	20	*	*15	*	*	*	*	*	*	*	*
427.8	721	304	417	*	32	86	590	221	109	288	102
427.81	312	119	193	*	*	*20	289	113	44	112	43
427.89	409	185	223	*	28	66	301	109	65	176	60
428	5,876	2,724	3,152	*	173	1,256	4,409	1,574	1,276	2,271	754
428.0	5,472	2,544	2,928	*	157	1,178	4,101	1,519	1,176	2,091	686
428.2	116	73	44	—	*	*26	82	*	42	49	*
428.20	*37	*23	*	—	*	*	*25	*	*	*	*
428.21	*23	*	*	—	*	*	*	*	*	*	*
428.23	*40	*	*	—	*	*	*28	*	*	*	*
428.3	223	68	155	—	*	*35	181	*38	*49	100	*37
428.30	105	*37	68	—	*	*	87	*	*28	50	*
428.31	*63	*	*48	—	*	*	*50	*	*	*	*
428.33	49	*	*34	—	*	*	*38	*	*	*	*
429	78	*22	*56	*	*	*15	*56	*	*	*53	*
430	251	*57	194	*	*53	118	*75	*59	*52	*78	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
431	500	235	266	*	*	143	315	114	71	197	118
432	178	101	77	*	*	*	133	*45	*	69	*38
432.1	153	93	*60	*	*	*	119	*	*	*57	*34
432.9	*22	*	*	—	*	*	*	*	*	*	*
433	487	275	212	—	*	154	322	86	115	208	79
433.1	344	189	154	—	*	86	247	68	89	128	59
433.10	229	136	93	—	*	42	184	48	56	95	30
433.11	115	*53	*62	—	*	*44	*64	*	*	*	*
433.3	95	65	30	—	—	*39	56	*16	*	56	*11
433.30	77	55	*22	—	—	*35	42	*	*	49	*
434	1,919	847	1,072	*	74	469	1,362	462	375	738	344
434.0	79	*35	*44	—	*	*	*57	*	*	*	*
434.01	67	*	*39	—	*	*	*48	*	*	*	*
434.1	260	89	171	—	*	*63	185	77	*46	107	*30
434.11	245	81	164	—	*	*62	171	73	*42	104	*
434.9	1,580	722	858	*	62	385	1,120	362	317	603	299
434.90	*47	*	*	—	*	*	*42	*	*	*	*
434.91	1,534	713	821	*	62	380	1,078	360	313	567	292
435	548	216	332	*	*18	104	426	141	139	200	68
435.3	*17	*	*	—	*	*	*16	*	*	*	*
435.8	32	*	*24	—	*	*	*24	*	*	*	*
435.9	489	200	289	*	*15	95	379	132	118	175	64
436	413	162	251	*	*45	78	288	48	65	199	101
437	146	44	102	*	*	40	89	*21	60	45	*
437.1	41	*	*28	—	—	*	39	*	*30	*	*
437.2	*19	*	*	*	*	*	*	*	*	*	*
437.3	*51	*	*41	*	*	*28	*	*	*	*	*
438	277	135	142	—	*	*49	223	*36	*42	170	*29
438.2	172	*	*96	—	*	*	*136	*	*	*142	*
438.20	170	*	*94	—	*	*	*135	*	*	*142	*
438.8	78	*53	*25	—	*	*	63	*	*37	*	*
438.89	62	*42	*	—	*	*	*51	*	*36	*	*
440	813	444	370	*	*	202	592	180	194	321	118
440.1	24	*	*16	*	*	*	*20	*	*	*	*
440.2	713	386	327	—	*	169	532	168	172	284	89
440.20	65	*39	*	—	*	*	*54	*	*	*48	*
440.21	74	45	29	—	*	28	46	*	*17	30	*
440.22	*47	*	*32	—	*	*	*29	*	*	*	*
440.23	143	*66	77	—	*	*	120	*60	*	*38	*
440.24	376	217	159	—	*	*88	278	*84	101	140	*
440.3	*51	*	*	—	*	*	*	*	*	*	*
441	451	309	142	*	*	123	317	110	97	181	63
441.0	88	*58	*	—	*	*	*34	*	*	*	*
441.01	*69	*	*	—	*	*	*	*	*	*	*
441.3	*43	*	*	—	—	*	*	*	*	*	*
441.4	237	186	*51	—	*	*34	201	57	*55	81	*44
442	66	*55	*	*	*	*	*32	*	*	*	*
443	154	82	72	*	*	*56	94	*22	*	97	*
443.9	135	74	*62	—	*	*51	83	*	*	88	*
444	241	94	147	—	*	84	131	*26	*42	89	83
444.2	177	61	115	—	*	*62	99	*	*26	68	*62
444.21	*35	*	*	—	*	*	*	*	*	*	*
444.22	142	49	93	—	*	*50	78	*	*	*51	*55
446	104	*60	*44	*	*	*	*	*	*	*	*
447	55	*30	*25	*	*	*	*27	*	*	*23	*
447.1	*21	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
451	62	*	40	*	*	*21	*35	*	*	*	*
451.1	*27	*	*	—	*	*	*	*	*	*	*
453	694	301	393	*	103	193	396	174	160	288	72
453.8	651	281	370	*	95	173	382	157	159	263	71
454	90	*	*74	—	*	*	*76	*	*	*	*
455	102	44	58	—	*20	*20	62	*35	*15	40	*13
455.2	63	26	37	—	*14	*12	37	*25	*	*22	*
456	*24	*	*	*	*	*	*	*	*	*	*
458	414	172	242	*	*19	126	266	134	67	139	74
458.0	182	84	98	*	*	*27	145	48	*38	70	*25
458.2	50	*24	*26	*	*	*	34	*	*	*	*
458.29	46	*22	*24	—	*	*	31	*	*	*	*
458.8	*14	*	*	—	*	*	*	*	*	*	*
458.9	168	55	112	*	*	81	83	71	*19	45	*32
459	191	92	99	*	*	77	89	*50	*36	78	*26
459.0	*53	*	*38	*	*	*	*	*	*	*	*
459.8	105	56	49	—	*	*44	56	*	*	*35	*
459.81	103	*55	*48	—	*	*42	56	*	*	*33	*
460-519	18,406	8,821	9,585	2,028	1,542	3,966	10,871	4,339	3,950	6,998	3,119
461	30	*17	*	*	*	*	*	*	*	*	*
462	*21	*10	*	*	*	*	*	*	*	*	*
463	*13	*	*	*	*	*	—	*	*	*	*
464	50	24	26	38	*	*	*	*	*	*20	*12
464.4	21	*10	*12	21	—	*	—	*	*	*11	*
465	68	31	38	43	*	*	*15	*18	*14	23	*14
465.9	65	29	36	42	*	*	*15	*17	*12	22	*14
466	868	448	420	618	49	71	131	146	150	342	230
466.0	246	89	157	*15	43	63	126	64	37	110	36
466.1	621	359	263	602	*	*	*	83	112	232	194
466.11	393	241	153	384	*	*	*	48	75	150	120
466.19	228	118	110	219	*	*	*	35	37	82	74
473	35	*	*18	*	*	*	*	*	*	*13	*
474	35	21	*14	24	*	*	—	*12	*	*10	*
474.0	*13	*	*	*	*	*	—	*	*	*	*
474.00	*10	*	*	*	*	*	—	*	*	*	*
474.1	22	*17	*	18	*	—	—	*	*	*	*
474.10	*14	*	*	*14	*	—	—	*	*	*	*
475	27	*16	*11	*	*16	*	*	*	*	*	*
478	89	45	44	*24	*24	*20	*	*	*	52	*
478.1	*14	*	*	*	*	*	*	*	*	*	*
478.2	*34	*22	*	*	*	*	*	*	*	*	*
478.7	*24	*	*	*	*	*	*	*	*	*	*
480	110	55	54	77	*	*	*	*	30	38	*23
480.1	52	29	*22	51	*	—	*	*	*14	*	*
480.9	49	*23	*27	*22	*	*	*	*	*	*21	*
481	206	103	103	*	*	*87	101	*35	*58	76	*36
482	1,120	566	554	*58	78	191	794	189	222	443	266
482.1	268	138	130	*	*	*	179	*	*61	119	*
482.2	*57	*	*	*	*	*	*39	*	*	*	*
482.3	*35	*	*	*	*	*	*	*	*	*	*
482.4	423	224	199	*	*	*81	314	*85	69	172	*96
482.41	383	202	181	*	*	*80	278	*80	*59	152	*92
482.8	234	114	120	*	*	*	180	*	52	109	*
482.820	*21	*	*	*	—	*	*	*	*	*	*
482.83	117	*80	*	*	*	*	88	*	*	*73	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
482.9	68	*29	*39	*	*	*	*49	*	*	*	*
483	*61	*	*	*	*	*	*	*	*	*	*
483.0	*56	*	*	*	*	*	*	*	*	*	*
485	102	47	55	*17	*	*	54	*	*	36	*
486	5,741	2,619	3,122	418	378	1,016	3,930	1,276	1,583	2,059	823
487	99	42	57	*17	*	*	67	*29	*36	*	*
487.0	*55	*	*	*	—	*	*40	*	*	*	*
487.1	41	*17	*24	*	*	*	*27	*	*	*	*
490	78	29	49	*16	*	*	40	*	*	48	*
491	2,413	1,097	1,316	*	60	684	1,655	691	475	997	251
491.2	2,395	1,085	1,310	*	57	682	1,649	686	473	991	245
491.20	80	*34	46	—	*	*	66	*	*	45	*
491.21	2,315	1,051	1,264	*	57	668	1,583	674	463	945	232
492	84	*32	51	—	*	*39	42	*	*	*	*
492.8	61	*19	*42	—	*	*29	31	*	*	*	*
493	1,599	582	1,017	415	264	418	502	472	290	550	287
493.0	109	68	41	79	*18	*	*	*16	*	24	57
493.01	23	*15	*	*18	*	*	*	*	*	*	*
493.02	77	48	29	54	*	*	*	*	*	*16	44
493.2	589	167	422	*	34	212	343	181	112	205	92
493.20	90	*16	75	—	*	*20	67	*23	*20	*27	*
493.21	*32	*	*	*	*	*	*	*	*	*	*
493.22	466	138	328	*	*29	170	267	146	87	167	67
493.9	898	346	552	335	211	200	153	274	167	321	137
493.90	147	68	79	50	23	40	*35	31	25	67	23
493.91	182	94	88	114	45	*	*	53	28	73	*28
493.92	570	185	385	171	144	149	106	190	114	180	86
494	*54	*	*	*	*	*	*38	*	*	*	*
496	380	201	179	*	*	105	266	82	63	183	53
507	1,492	934	557	*44	*43	229	1,176	425	238	520	310
507.0	1,476	920	556	*42	*42	229	1,162	424	236	517	299
510	162	*102	*	*	*	*66	*	*	*	*	*
510.9	134	*95	*	*	*	*	*	*	*	*	*
511	353	213	140	*	*	71	258	72	67	131	83
511.0	*14	*	*	—	*	*	*	*	*	*	*
511.8	81	*53	*	*	*	*	*62	*	*	*	*
511.9	253	146	107	*	*	47	191	66	*39	94	*53
512	225	138	86	*	80	*64	75	*34	*34	118	*38
512.1	45	*	*33	*	*	*	*22	*	*	*	*
512.8	163	111	53	*	66	*45	*51	*	*26	89	*
513	*83	*	*	*	*	*	*	*	*	*	*
513.0	*79	*	*	*	*	*	*	*	*	*	*
514	33	*	*22	*	*	*	*	*	*	*	*
515	169	*42	127	*	*	*51	83	*	*27	95	*25
516	101	*43	*58	*	*	*	*	*	*	*	*
516.8	*59	*	*	*	*	*	*	*	*	*	*
518	2,366	1,159	1,208	99	292	616	1,358	509	423	983	451
518.0	*34	*	*	*	*	*	*	*	*	*	*
518.4	*17	*	*	*	*	*	*	*	*	*	*
518.8	2,249	1,099	1,150	88	279	589	1,293	478	399	928	444
518.81	1,718	807	910	*68	201	462	987	424	272	669	352
518.82	104	*62	*42	*	*	*	*41	*	*	*72	*
518.84	311	145	166	*	*	91	185	*	102	137	*41
518.89	60	*40	*20	*	*	*	*44	*	*	*43	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
519	87	47	*40	*15	*	*27	*33	*	*18	*23	*
519.0	*35	*	*	*	*	*	*	*	*	*	*
519.1	*20	*	*	*	*	*	*	*	*	*	*
519.8	*27	*21	*	*	*	*	*	*	*	*	*
520-579	16,820	7,484	9,336	903	3,228	5,171	7,518	3,806	3,450	6,378	3,187
522	*23	*	*	*	*	*	*	*	*	*	*
522.5	*21	*	*	*	*	*	*	*	*	*	*
524	35	*	28	*	26	*	*	*	*	*13	*11
524.0	17	*	*13	*	*13	*	*	*	*	*	*
524.03	*9	*	*	*	*	*	*	*	*	*	*
527	*25	*	*	*	*	*	*	*	*	*	*
527.2	*20	*	*	*	*	*	*	*	*	*	*
528	*41	*22	*	*	*	*	*	*	*	*	*
530	790	366	424	126	145	216	304	187	151	325	128
530.1	128	63	65	*	26	46	50	*19	*24	57	*29
530.10	*19	*	*	*	*	*	*	*	*	*	*
530.11	37	*17	*20	*	*	*13	*	*	*	*	*
530.19	62	*31	*30	*	*	*26	*27	*	*	38	*
530.2	71	*33	*38	*	*	*	51	*	*	*29	*
530.20	*26	*	*	*	*	*	*	*	*	*	*
530.21	44	*23	*21	*	*	*	34	*	*	*	*
530.3	44	*20	*24	*	*	*	*22	*	*	*	*
530.5	*19	*	*	—	*	*	*	*	*	*	*
530.7	83	*59	*24	*	*	*	*	*	*	*	*
530.8	398	171	227	91	62	107	137	110	90	137	60
530.81	350	143	208	91	61	98	101	101	81	119	49
531	614	332	281	*	*46	226	340	92	113	231	177
531.0	79	*26	*53	—	*	*	*42	*	*	*	*
531.00	77	*26	*51	—	*	*	*42	*	*	*	*
531.4	367	228	139	*	*	153	201	65	54	110	137
531.40	262	130	132	*	*	61	188	61	54	102	45
531.9	66	*29	*38	*	*	*	*45	*	*	*34	*
531.90	51	*23	*28	*	*	*	*30	*	*	*21	*
532	416	233	183	*	*44	86	286	97	118	142	60
532.0	55	*25	*	*	*	*	*27	*	*	*	*
532.00	55	*25	*	*	*	*	*27	*	*	*	*
532.4	217	141	76	—	*17	43	156	58	51	72	*36
532.40	215	140	75	—	*	43	155	57	51	70	*36
532.5	*80	*	*	*	*	*	*	*	*	*	*
532.50	*80	*	*	*	*	*	*	*	*	*	*
532.9	*24	*	*	*	*	*	*	*	*	*	*
532.90	*21	*	*	*	*	*	*	*	*	*	*
533	55	*21	*35	—	*	*	*39	*	*	*	*
533.4	*29	*	*	—	*	*	*	*	*	*	*
533.40	*28	*	*	—	*	*	*	*	*	*	*
534	59	*	*45	—	*	*40	*	*	*	*	*
534.4	*30	*	*22	—	*	*	*	*	*	*	*
534.40	*29	*	*21	—	*	*	*	*	*	*	*
535	576	217	359	*13	90	150	324	142	119	230	85
535.0	79	28	51	*	*16	*19	41	*18	*15	40	*
535.00	60	*22	38	*	*	*	*29	*	*	33	*
535.01	*19	*	*	*	*	*	*	*	*	*	*
535.1	103	*30	72	*	*	*	84	*41	*	*37	*
535.10	51	*	*40	*	*	*	*39	*	*	*31	*
535.11	*51	*	*	*	*	*	*46	*	*	*	*
535.3	*18	*	*	—	*	*	*	*	*	*	*
535.4	174	66	107	*	27	44	100	44	39	53	38
535.40	79	*23	55	*	*20	*23	*34	*21	*	*27	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
535.41	95	43	52	*	*	*21	66	*23	*28	*26	*
535.5	176	70	106	*	27	53	91	*31	36	85	*24
535.50	91	36	55	*	*17	35	34	*16	*17	41	*
535.51	85	*34	50	*	*	*	57	*	*19	44	*
535.6	*27	*	*	*	*	*	*	*	*	*	*
536	255	89	166	*22	62	54	117	*39	54	113	49
536.2	63	*	47	*	*	*	*	*	*	*30	*
536.3	66	*	*52	*	*	*	*	*	*	*30	*
536.4	97	53	44	*	*	*	60	*	*	*38	*
536.41	*45	*	*	*	*	*	*	*	*	*	*
536.42	*32	*	*	*	*	*	*21	*	*	*	*
536.49	*20	*	*	*	*	*	*	*	*	*	*
536.8	28	*	22	*	*	*	*11	*	*9	*	*
537	173	61	112	*	*	*62	94	*	*25	76	*50
537.0	66	*	*58	*	*	*	*	*	*	*	*
537.8	92	*43	49	*	*	*	67	*	*	*40	*
537.83	66	*36	*30	—	*	*	56	*	*	*	*
540	950	565	385	299	360	195	96	195	195	289	271
540.0	276	171	105	89	73	64	49	41	77	83	75
540.1	216	137	79	90	56	*41	*	*55	*32	63	65
540.9	459	257	202	119	230	91	*18	98	86	143	131
541	*20	*	*	*	*	*	*	*	*	*	*
550	160	139	*21	*16	*	*40	94	34	*42	58	*26
550.1	99	84	*	*	*	*	69	*25	*	*43	*
550.10	68	53	*	*	*	*	*43	*	*	*	*
550.9	41	36	*	*	*	*	24	*	*	*15	*11
550.90	25	23	*	*	*	*	*13	*	*	*	*
552	376	134	243	*	56	108	211	89	76	156	55
552.1	*33	*	*	*	*	*	*	*	*	*	*
552.2	281	111	170	*	41	96	145	69	61	105	47
552.20	72	*	*46	*	*	*	*43	*	*	*44	*
552.21	181	75	106	—	*31	64	85	*47	45	51	*38
553	334	122	212	*	66	134	129	56	47	146	85
553.1	*18	*	*	*	*	*	*	*	*	*	*
553.2	245	102	143	*	50	108	85	36	36	104	69
553.20	46	*	28	—	*	*26	*	*	*	*27	*
553.21	193	80	112	*	39	78	72	*28	*30	77	58
553.3	63	*14	49	*	*	*23	31	*	*	25	*
555	248	95	154	*	139	57	*36	54	53	113	*28
555.0	61	*	*34	*	*28	*	*	*	*	*	*
555.1	79	*26	*52	*	*47	*	*	*	*	*42	*
555.2	*33	*	*	*	*	*	*	*	*	*	*
555.9	75	*30	46	*	44	*	*	*22	*	*30	*
556	294	193	100	*	85	151	*49	*100	79	67	*48
556.6	*97	*	*	*	*	*	*	*	*	*	*
556.9	97	57	*40	*	36	*	*	*	*25	*	*
557	439	128	310	*	*32	108	290	124	80	159	76
557.0	276	*88	188	*	*	*	176	*59	*48	*115	*54
557.9	143	*31	112	*	*	*32	101	*52	*27	*42	*22
558	500	169	332	103	105	102	190	123	113	192	72
558.9	461	151	310	98	103	88	172	119	92	177	72
560	1,881	733	1,148	79	165	555	1,082	413	399	686	384
560.1	285	119	166	*	*	95	174	49	76	109	51
560.2	146	*91	*55	*	*	*	*71	*	*	*	*
560.3	132	34	98	*	*	*	94	*25	*32	52	*23
560.39	110	32	78	*	*	*	74	*25	*19	48	*17
560.8	773	274	498	*	80	236	438	178	140	294	161
560.81	608	219	389	*	*62	169	360	144	106	239	120

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
560.89	165	55	110	*	*	67	78	*34	*35	*55	*41
560.9	528	206	322	*	45	156	303	123	127	175	102
562	1,567	643	924	*	197	419	950	461	334	537	236
562.1	1,546	640	906	*	194	419	931	459	329	531	227
562.10	140	48	92	—	*	*29	98	*31	*	51	*31
562.11	995	422	573	*	166	331	498	312	213	324	147
562.12	344	133	211	*	*	49	290	105	63	137	39
562.13	67	*37	*29	—	*	*	*45	*	*	*	*
564	272	94	178	*24	55	82	110	59	86	75	51
564.0	161	62	99	*18	*27	32	84	37	53	39	*32
564.00	110	45	65	*	*21	*20	57	*21	35	29	*25
564.09	48	*	*31	*	*	*	*	*	*	*	*
564.1	54	*	47	*	*19	*22	*	*	*	*	*
566	116	78	38	*	38	45	*	*	*26	61	*
567	156	80	76	*	*38	*58	*46	*	*	*58	*35
567.2	115	*53	*62	*	*	*	*	*	*	*	*
568	95	*	70	*	*33	*28	*	*	*	*51	*
568.0	51	*	*41	*	*31	*	*	*	*	*21	*
568.8	*44	*	*	*	*	*	*	*	*	*	*
569	660	215	444	*	123	197	323	110	119	324	108
569.1	*28	*	*25	*	*	*	*	*	*	*	*
569.3	95	*27	68	*	*	*19	68	*	*	58	*
569.4	*28	*	*	—	*	*	*	*	*	*	*
569.6	165	*57	107	*	*69	*41	*46	*	*32	*83	*
569.69	79	*	*46	*	*	*	*36	*	*	*	*
569.8	315	105	210	*	*	118	158	*41	*61	156	*57
569.81	*86	*	*	—	*	*	*	*	*	*	—
569.83	140	*53	*87	*	*	*	*79	*	*	*60	*
569.85	52	*	*26	*	*	*	42	*	*	*	*
570	*44	*	*	*	*	*	*	*	*	*	*
571	708	430	278	*	116	426	165	140	105	332	131
571.1	72	*45	*	—	*	*43	*	*	*	*	*28
571.2	318	242	76	—	68	223	*27	81	*43	122	72
571.5	268	118	150	*	*	127	128	*43	*31	173	*
572	306	187	119	*	*34	176	86	83	*39	120	64
572.2	247	148	100	—	*	152	72	62	*	104	55
573	55	*	*28	*	*	*	*31	*	*	*	*
573.3	*29	*	*19	*	*	*	*	*	*	*	*
574	1,426	517	909	*	390	385	644	339	280	486	321
574.0	521	236	285	*	123	160	237	112	104	200	105
574.00	458	200	259	*	118	154	185	98	100	177	84
574.01	63	*	*26	—	*	*	*52	*	*	*	*
574.1	392	98	295	*	110	119	161	109	66	133	83
574.10	376	93	283	*	109	111	154	105	65	128	78
574.2	143	43	100	*	68	25	48	*20	*50	40	33
574.20	134	36	99	*	66	24	41	*20	*49	37	28
574.3	*41	*	*	—	*	*	*	*	*	*	*
574.5	92	*41	51	—	*17	*	61	*	*	*26	*22
574.50	48	*	*32	—	*	*	*28	*	*	*	*
574.51	*43	*	*	—	*	*	*	*	*	*	*
574.6	*51	*	*	—	*	*	*	*	*	*	*
574.7	80	*	58	*	*29	*	*	*	*	*	*
574.70	47	*	*36	*	*	*	*	*	*	*	*
574.71	*33	*	*	*	*	*	*	*	*	*	*
574.8	*54	*	*28	—	*	*	*	*	*	*	*
574.81	*35	*	*	—	*	*	*	*	*	*	*
574.9	*38	*	*27	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
575	349	124	224	*	70	97	177	58	85	125	81
575.0	125	54	71	*	*	*41	70	*32	*35	*36	*
575.1	186	61	125	*	42	48	96	*23	40	72	*51
575.10	37	*	*22	—	*	*	*21	*	*	*21	*
575.11	69	*25	44	*	31	*21	*	*	*21	*34	*
575.12	80	*22	*59	—	*	*	*59	*	*	*	*
575.8	*26	*	*18	*	*	*	*	*	*	*	*
576	136	71	65	*	*32	*27	71	*53	*	*43	*
576.1	*40	*	*	*	*	*	*	*	*	*	*
576.2	58	*35	*23	*	*	*	*	*	*	*	*
576.8	*35	*	*	*	*	*	*	*	*	*	*
577	1,644	794	850	*33	509	620	482	304	355	662	323
577.0	1,399	677	722	*	430	494	450	257	302	577	263
577.1	153	73	80	*	59	73	*	*	*39	60	*30
577.2	*55	*	*	*	*	*	*	*	*	*	*
577.9	*26	*	*	*	*	*	*	*	*	*	*
578	920	467	454	*	65	248	592	248	224	335	114
578.0	90	43	46	*	*28	*26	33	*	*17	*37	*
578.1	172	83	89	*	*	32	129	56	30	58	*28
578.9	658	340	318	*	28	190	430	169	177	239	73
580-629	7,462	2,521	4,941	258	1,637	1,754	3,812	1,657	1,506	3,071	1,228
581	*39	*	*	*	*	*	*	*	*	*	*
581.9	*30	*	*	*	*	*	*	*	*	*	*
584	1,569	795	774	*	78	424	1,049	471	274	591	234
584.5	114	*71	*43	*	*	*	*59	*	*	*	*
584.9	1,421	714	707	*	68	372	966	404	248	551	218
585	99	49	50	*	*	*23	*58	*	*	*41	*
590	523	122	402	62	192	115	155	76	138	204	104
590.1	254	70	184	*24	113	42	75	49	70	86	49
590.10	254	70	184	*24	113	42	75	49	70	86	49
590.8	250	49	200	35	74	67	74	*25	65	110	50
590.80	250	49	200	35	74	67	74	*25	65	110	50
591	46	*20	*26	*	*	*	*	*	*	*16	*
592	403	189	215	*	160	152	88	105	91	149	58
592.0	158	64	93	*	63	64	29	44	42	44	28
592.1	237	120	118	*	93	88	54	61	48	98	30
593	139	67	72	30	*19	44	46	36	33	52	*17
593.4	*26	*	*	*	*	*	*	*	*	*	*
593.7	*24	*	*19	*23	*	*	—	*	*	*	*
593.70	*17	*	*	*16	*	—	—	*	*	*	*
593.9	53	*30	*23	*	*	*26	*23	*	*	*23	*
594	*23	*18	*	—	*	*	*	*	*	*	*
594.1	*22	*	*	—	*	*	*	*	*	*	*
595	140	65	74	*	*	*	113	*67	*	45	*
595.0	*40	*	*	*	*	*	*35	*	*	*	*
595.9	*28	*	*	*	*	*	*22	*	*	*	*
596	79	*49	*	*	*	*	*43	*	*	*	*
596.0	*18	*14	*	—	*	*	*18	*	*	*	*
599	2,203	679	1,524	84	153	322	1,644	502	438	889	375
599.0	2,081	610	1,471	82	146	291	1,562	461	419	855	346
599.7	95	61	34	*	*	*19	69	38	*15	*26	*16
600	262	262	—	—	*	46	214	65	74	82	40
600.0	235	235	—	—	*	40	194	59	67	70	38
600.00	66	66	—	—	*	*11	54	*23	*11	*21	*11
600.01	169	169	—	—	*	28	139	36	56	49	28

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
601	*21	*21	—	—	*	*	*	*	*	*	*
604	42	42	—	*	*	*	*20	*	*	*	*
604.9	*29	*29	—	*	*	*	*	*	*	*	*
604.90	*20	*20	—	*	*	*	*	*	*	*	*
607	*14	*14	—	*	*	*	*	*	*	*	*
608	65	65	—	*	*	*36	*	*	*	*	*
608.4	*33	*33	—	*	*	*	*	*	*	*	*
611	128	*	126	*	82	38	*	*14	*12	91	*
611.0	103	*	102	*	*71	*	*	*	*	*80	*
611.1	15	*	14	*	*9	*	*	*	*	*	*
614	149	—	149	*	114	*	*	*32	26	70	*21
614.2	*34	—	*34	*	*25	*	*	*	*	*	*
614.6	*23	—	*23	—	*18	*	*	*	*	*	*
614.9	45	—	45	*	38	*	*	*	*	*25	*
616	29	—	29	*	*	*	*	*	*	*19	*
617	196	—	196	—	148	47	*	27	47	81	41
617.0	110	—	110	—	77	33	*	*17	25	45	23
617.1	42	—	42	—	36	*	—	*	*	*19	*
617.3	28	—	28	—	22	*	*	*	*	*	*
618	343	—	343	—	58	135	151	37	74	158	74
618.0	54	—	54	—	*	22	23	*9	*10	27	*
618.1	32	—	32	—	*16	*	*	*	*	*	*
618.2	43	—	43	—	*	23	*	*	*	*18	*
618.3	35	—	35	—	*	*	*17	*	*	*	*
618.4	65	—	65	—	*12	28	26	*	*20	33	*
618.5	72	—	72	—	*	*16	56	*	*16	38	*15
618.8	*31	—	*31	—	*	*	*	*	*	*	*
619	*35	—	*35	*	*	*	*	*	*	*	*
620	218	—	218	*	133	52	*	29	56	92	43
620.0	*14	—	*14	*	*	*	—	*	*	*	*
620.1	28	—	28	*	22	*	—	*	*	*14	*
620.2	134	—	134	*	85	31	*	*16	34	61	*23
620.5	*13	—	*13	*	*	*	*	*	*	*	*
620.8	*28	—	*28	*	*	*	*	*	*	*	*
621	48	—	48	—	*17	23	*	*	*	23	*
621.3	27	—	27	—	*	*14	*	*	*	*	*
622	*15	—	*15	—	*	*	*	*	*	*	*
622.1	*13	—	*13	—	*	*	*	*	*	*	*
625	175	—	175	*	121	41	*13	21	41	76	37
625.3	58	—	58	—	53	*	—	*	*	29	*13
625.6	48	—	48	—	*10	28	*10	*	*14	*15	*12
625.9	58	—	58	*	51	*	*	*	*16	25	*
626	271	—	271	*	189	80	*	30	61	120	60
626.2	205	—	205	*	144	61	*	*18	51	89	47
626.8	56	—	56	*	37	18	*	*	*	24	*12
627	59	—	59	—	*	*24	*	*	*	*33	*
627.1	51	—	51	—	*	*20	*	*	*	*	*
630-677	1,410	—	1,410	*	1,404	*	—	239	267	640	265
632	33	—	33	—	33	—	—	*	*	16	*
633	62	—	62	*	61	*	—	*11	*	30	*15
633.1	53	—	53	*	52	*	—	*10	*	25	*13
633.10	51	—	51	*	50	*	—	*	*	25	*13
634	31	—	31	*	30	*	—	*12	*	*13	*
634.9	20	—	20	*	20	*	—	*	*	*10	*
634.91	17	—	17	*	17	*	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
640	*20	—	*20	—	*20	—	—	*	*	*	*
641	43	—	43	—	43	—	—	*	*10	*19	*
641.1	*15	—	*15	—	*15	—	—	*	*	*	*
641.13	*15	—	*15	—	*15	—	—	*	*	*	*
641.9	*10	—	*10	—	*10	—	—	*	*	*	*
641.93	*10	—	*10	—	*10	—	—	*	*	*	*
642	64	—	64	—	64	*	—	*14	*11	29	*
642.3	*14	—	*14	—	*14	*	—	*	*	*11	*
642.33	*13	—	*13	—	*13	*	—	*	*	*10	*
642.4	21	—	21	—	21	—	—	*	*	*	*
642.43	*14	—	*14	—	*14	—	—	*	*	*	*
643	87	—	87	—	87	*	—	*14	*18	45	*
643.0	40	—	40	—	40	—	—	*	*	*20	*
643.03	40	—	40	—	40	—	—	*	*	*20	*
643.1	33	—	33	—	33	—	—	*	*	*16	*
643.13	33	—	33	—	33	—	—	*	*	*16	*
644	337	—	337	*	336	*	—	64	67	140	66
644.0	320	—	320	*	319	*	—	61	64	135	60
644.03	318	—	318	*	317	*	—	61	64	134	60
644.1	17	—	17	—	17	*	—	*	*	*	*
644.13	16	—	16	—	16	*	—	*	*	*	*
646	164	—	164	*	163	*	—	*14	21	100	29
646.2	*12	—	*12	—	*12	—	—	*	*	*	*
646.23	*12	—	*12	—	*12	—	—	*	*	*	*
646.6	97	—	97	*	97	*	—	*	*14	57	*17
646.63	87	—	87	*	87	—	—	*	*12	50	*16
646.8	43	—	43	—	43	—	—	*	*	26	*
646.83	39	—	39	—	39	—	—	*	*	25	*
647	*19	—	*19	—	*19	—	—	*	*	*	*
648	267	—	267	*	267	*	—	48	58	109	52
648.0	24	—	24	—	24	—	—	*	*	*16	*
648.03	22	—	22	—	22	—	—	*	*	*13	*
648.4	35	—	35	—	35	—	—	*	*	*	*
648.43	*28	—	*28	—	*28	—	—	*	*	*	*
648.8	*23	—	*23	—	*23	—	—	*	*	*	*
648.83	*23	—	*23	—	*23	—	—	*	*	*	*
648.9	155	—	155	*	154	*	—	32	36	56	31
648.93	146	—	146	*	146	*	—	27	34	54	30
654	42	—	42	*	42	—	—	*	*	*	*
654.5	*21	—	*21	—	*21	—	—	*	*	*	*
654.53	*21	—	*21	—	*21	—	—	*	*	*	*
658	36	—	36	—	36	—	—	*	*	*	*
658.1	*16	—	*16	—	*16	—	—	*	*	*	*
658.13	*16	—	*16	—	*16	—	—	*	*	*	*
659	*10	—	*10	—	*10	*	—	*	*	*	*
666	*13	—	*13	—	*13	—	—	*	*	*	*
666.2	*13	—	*13	—	*13	—	—	*	*	*	*
666.24	*13	—	*13	—	*13	—	—	*	*	*	*
670	32	—	32	—	32	—	—	*	*	*	*
670.0	32	—	32	—	32	—	—	*	*	*	*
670.04	32	—	32	—	32	—	—	*	*	*	*
674	69	—	69	—	69	—	—	*	*	*	*
674.8	*21	—	*21	—	*21	—	—	*	*	*	*
674.84	*21	—	*21	—	*21	—	—	*	*	*	*
680-709	3,867	1,959	1,908	276	915	1,146	1,530	808	671	1,825	563
681	131	68	63	*18	37	37	*38	*21	*24	61	*24
681.0	60	33	*27	*	*25	*	*	*	*	*20	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
681.00	53	*27	*27	*	*22	*	*	*	*	*18	*
681.1	66	35	*31	*	*	*26	*	*	*	*37	*
681.10	66	*35	*31	*	*	*26	*	*	*	*36	*
682	2,554	1,275	1,279	139	623	804	988	598	475	1,093	388
682.0	156	63	93	*21	74	37	*24	*25	*25	88	*17
682.1	42	*27	*	*	*19	*	*	*	*	*23	*
682.2	232	97	135	*	84	73	64	*43	40	108	*41
682.3	282	172	111	*	104	94	76	60	42	129	51
682.4	107	65	42	*	33	50	*19	*25	*20	42	*20
682.5	85	47	39	*27	*28	*21	*	*	*	55	*
682.6	1,345	637	707	36	228	409	672	373	278	512	182
682.7	273	156	117	*17	45	104	107	46	57	119	*51
692	*13	*	*	*	*	*	*	*	*	*	*
693	*22	*	*	*	*	*	*	*	*	*	*
693.0	*21	*	*	*	*	*	*	*	*	*	*
701	29	*17	*12	*17	*	*	*	*	*	25	*
701.4	*21	*	*	*17	*	—	—	*	*	*20	—
707	835	443	391	*	154	233	446	145	131	444	115
707.0	593	328	265	*	*124	175	291	*81	99	335	78
707.1	213	104	109	*	*	*48	143	*59	*30	91	*
707.15	95	*42	*53	—	*	*30	*53	*	*	*45	*
708	*22	*	*19	*	*	*	*	*	*	*	*
709	109	63	46	60	33	*	*	*	*	103	*
709.2	97	58	38	58	33	*	*	*	*	94	*
710-739	7,416	2,962	4,454	163	1,088	2,407	3,758	1,651	1,814	2,592	1,359
710	148	*	129	*	79	*37	*	*47	*	55	*
710.0	95	*	85	*	67	*	*	*	*	*36	*
711	150	91	*59	*	*	*61	*58	*	*	*57	*35
711.0	146	87	*59	*	*	*61	*55	*	*	*56	*35
711.06	75	*49	*	*	*	*37	*	*	*	*	*
714	125	*31	93	*	*	*34	72	*24	*25	*61	*
714.0	87	*	64	*	*	*27	48	*21	*	*34	*
715	2,697	978	1,719	—	81	897	1,719	625	735	878	459
715.1	205	82	122	—	*	67	131	58	*30	92	*26
715.15	49	*	*34	—	*	*	*33	*20	*	*	*
715.16	138	52	86	—	*	47	89	*34	*	70	*19
715.2	*24	*	*	—	*	*	*	*	*	*	*
715.3	1,462	546	916	—	38	499	925	356	413	417	275
715.31	22	*	*12	—	—	*	*14	*	*	*	*
715.35	391	172	219	—	*	134	243	100	110	109	72
715.36	1,028	352	676	—	*18	353	657	238	295	301	194
715.8	*41	*	*	—	—	*	*39	*	*	*	*
715.89	*41	*	*	—	—	*	*39	*	*	*	*
715.9	956	319	637	—	*33	318	605	192	280	336	148
715.90	*25	*	*	—	*	*	*18	*	*	*	*
715.91	*18	*	*	—	*	*	*14	*	*	*	*
715.95	298	115	183	—	*22	102	175	82	76	90	51
715.96	607	188	419	—	*	204	395	101	196	223	87
716	76	*36	40	*	*	*17	39	*	*22	*34	*
716.9	35	*	*22	*	*	*	*27	*	*	*	*
717	*13	*	*	*	*	*	*	*	*	*	*
718	30	*20	*	*	*	*	*	*	*	*	*
719	119	47	72	*	*15	43	57	*30	29	43	*17
719.4	81	*21	60	*	*	*23	43	*20	*19	*29	*12
719.41	*16	*	*11	*	*	*	*	*	*	*	*
719.45	36	*	*27	*	*	*	*26	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
721	323	121	202	*	40	106	177	65	82	116	59
721.0	46	22	24	—	*12	22	*12	*	*11	18	*12
721.1	79	46	*33	*	*	33	*38	*	*	*32	*17
721.3	173	45	128	—	*	47	111	*35	53	56	*29
722	1,022	439	583	*	311	396	314	204	230	365	223
722.0	105	55	50	—	46	48	*	14	22	46	23
722.1	510	209	300	*	175	173	160	117	112	170	110
722.10	498	208	290	*	173	167	156	116	109	164	109
722.4	27	*	*17	—	*	*17	*	*	*	*	*
722.5	228	94	133	—	55	86	87	*37	59	78	54
722.52	221	91	131	—	55	82	84	*36	56	76	53
722.7	55	28	27	—	*	27	*	*	*16	*16	*
722.71	34	*18	*16	—	*	*21	*	*	*	*12	*
722.73	*17	*	*	—	*	*	*	*	*	*	*
722.8	53	*26	27	—	*	*27	*	*	*	*29	*
722.83	46	*24	*22	—	*	*21	*	*	*	*25	*
722.9	37	*	*21	*	*	*	*	*	*	*	*
722.93	*30	*	*	*	*	*	*	*	*	*	*
723	74	33	41	*	*15	34	*24	*	*11	39	*18
723.0	30	*	*18	*	*	*14	*	*	*	*20	*
723.1	22	*	*16	*	*	*	*	*	*	*15	*
724	542	207	336	*	49	156	337	147	128	170	97
724.0	304	113	191	*	*	89	205	71	71	102	60
724.02	293	111	182	*	*	85	199	70	70	97	56
724.2	117	44	73	*	24	34	58	*31	38	35	*
724.4	*25	*	*	—	*	*	*	*	*	*	*
724.5	69	*20	49	—	*	*18	*44	*32	*	*	*
726	107	54	54	*	*21	38	44	34	27	*18	28
726.1	35	18	17	—	*	*11	23	*	*	*	*15
726.10	34	18	16	—	*	*10	23	*	*	*	*15
726.3	34	*19	*	—	*	*15	*	*	*	*	*
726.33	34	*19	*	—	*	*15	*	*	*	*	*
727	65	40	24	*	*	*12	29	*	*	*29	*15
727.0	*27	*	*	*	*	*	*	*	*	*	*
727.4	*11	*	*	—	*	*	*	*	*	*	*
727.40	*10	*	*	—	*	*	*	*	*	*	*
727.6	*16	*	*	—	*	*	*	*	*	*	*
728	241	96	145	*	68	95	71	*46	*46	121	*
728.8	228	87	141	*	64	90	69	*43	*42	115	*
728.88	82	39	*44	*	*18	*	*50	*23	*	*29	*
729	172	71	101	*	40	58	68	43	31	79	*19
729.1	28	*	*19	*	*	*	*	*	*	*	*
729.5	67	29	38	*	*	*18	*36	*	*	*26	*
729.8	53	*27	26	*	*	*18	*21	*	*	33	*
729.89	35	*	*18	*	*	*	*19	*	*	*24	*
730	476	301	175	*	114	153	166	93	142	140	101
730.0	153	92	*	*	*	*	*	*	*	*	*
730.07	*48	*	*	*	*	*	*	*	*	*	*
730.1	*62	*	*	*	*	*	*	*	*	*	*
730.2	260	167	94	*	*61	95	91	*54	*74	83	*
730.27	121	*59	*62	*	*	*46	*56	*	*	*44	*
730.28	*70	*	*	*	*	*	*	*	*	*	*
732	*13	*	*	*	*	*	*	*	*	*	*
733	731	256	475	*	93	181	445	159	177	255	140
733.1	438	141	296	*	*	86	312	100	112	136	89
733.13	289	88	202	—	*	*50	237	78	51	87	74
733.14	87	*	*64	*	*	*	*	*	*	*	*
733.15	*28	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
733.4	102	54	48	*	*19	46	*36	*	*	55	*17
733.42	85	47	39	*	*16	40	*29	*	*	50	*14
733.6	44	*13	31	*	*	17	18	*14	*9	*13	*
733.8	98	34	64	*	24	*23	*50	*	*39	26	*20
733.81	*13	*	*	*	*	*	*	*	*	*	*
733.82	85	26	59	*	*16	*20	*49	*	*	*22	*19
733.9	*38	*	*26	*	*	*	*	*	*	*	*
736	39	*25	*	*	*	*	*	*	*	*	*
737	79	*	*59	*	*	*	*	*	*	*	*
737.3	73	*	*56	*	*	*	*	*	*	*	*
737.30	*50	*	*42	*	*	*	*	*	*	*	*
738	127	42	85	*	*21	54	50	*29	*22	*37	40
738.4	111	34	77	*	*	50	49	*	*20	*30	35
740-759	1,400	751	649	1,024	114	214	*49	314	311	423	352
742	*33	*	*	*30	*	*	*	*	*	*	*
744	*9	*	*	*	*	*	—	*	*	*	*
745	287	207	81	262	*16	*	*	69	*	76	*130
745.2	*87	*	*	*85	*	—	—	*	*	*	*
745.4	*97	*	*	*95	*	—	—	*	*	*	*
745.5	*25	*	*	*	*	*	*	*	*	*	*
746	176	91	*85	125	*	*	*	*39	*	*64	*54
746.4	*52	*	*	*	*	*	*	*	*	*	*
747	284	92	193	118	*	*134	*	*57	*142	*56	*
747.8	160	*	*144	*	*	*129	*	*	*	*	*
747.81	*140	*	*136	*	*	*129	*	*	*	*	*
749	32	*12	*20	31	*	—	—	*	*	*	*8
749.0	*13	*	*	*13	*	—	—	*	*	*	*
749.2	*9	*	*	*9	*	—	—	*	*	*	*
750	58	43	*	54	—	*	*	*	*	*21	*
750.5	39	33	*	39	—	*	—	*	*	*15	*
751	109	*70	*	92	*	*	*	*	*	*57	*
752	*19	*	*	*	*	*	*	*	*	*	*
753	43	*26	*	*	*	*	*	*	*	*17	*
754	45	30	*	32	*	*	*	*15	*	*	*
755	*21	*	*	*	*	*	—	*	*	*	*
756	156	*67	*88	*121	*	*	*	*	*	*50	*
756.1	*37	*	*	*	*	*	*	*	*	*	*
756.12	*25	*	*	*	*	*	*	*	*	*	*
757	10	*	*5	*8	*	—	—	*	*	*9	*
757.3	*9	*	*5	*8	*	—	—	*	*	*9	*
757.32	*9	*	*5	*8	*	—	—	—	*	*9	*
759	*47	*	*	*	*	*	*	*	*	*	*
760-779	2,173	1,226	947	2,170	*	—	*	205	392	937	639
765	926	476	450	926	—	—	—	*	*	522	208
765.1	598	327	271	598	—	—	—	*	*	362	*102
765.17	*115	*	*	*115	—	—	—	*	*	*	*
765.18	*75	*	*	*75	—	—	—	*	*	*	*
769	362	222	*140	361	*	—	—	*	*	*	*183
770	320	213	108	320	—	—	—	*	44	121	104
770.6	*51	*38	*	*51	—	—	—	*	*	*	*
770.8	156	97	60	156	—	—	—	*	*24	*62	*54
770.81	*36	*	*	*36	—	—	—	*	*	*	*
770.89	102	*63	*	102	—	—	—	*	*	*	*
771	88	*47	*42	88	—	—	—	*	*	*27	*
771.8	71	*40	*	71	—	—	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
771.81	*52	*	*	*52	—	—	—	*	*	*	*
774	122	73	49	122	—	—	—	*15	*17	54	36
774.2	*12	*	*	*12	—	—	—	*	*	*	*
774.6	100	60	40	100	—	—	—	*13	*13	47	27
777	*69	*	*	*69	—	—	—	*	*	*	*
778	28	*15	*	28	—	—	—	*	*	*	*
778.4	*18	*	*	*18	—	—	—	*	*	*	*
779	128	*50	*78	127	*	—	—	*	*	*	*
779.3	*62	*	*	*62	—	—	—	*	*	*	*
780-799	659	314	345	117	180	159	203	176	74	296	113
780	179	87	92	71	40	38	31	41	22	80	36
780.2	36	16	*20	*	*	*14	*14	*9	*	*15	*
780.3	71	31	40	30	24	*	*	22	*10	27	*12
780.39	65	28	37	24	24	*	*	22	*9	*24	*
780.6	37	25	*12	31	*	*	*	*	*	20	*
784	*13	*	*	*	*	*	*	*	*	*	*
786	117	54	63	*	26	46	37	27	*8	67	14
786.0	*26	*	*	*	*	*	*	*	*	*	*
786.5	88	46	43	*	19	44	25	21	*7	46	14
786.50	24	*11	*13	—	*7	*9	*	*	*	*11	*
786.59	59	31	28	*	12	33	*13	*14	*	32	*10
787	*15	*	*	*	*	*	*	*	*	*11	*
789	71	17	55	*9	46	*12	*	23	*10	30	*8
789.0	69	16	53	*9	45	*11	*	22	*10	29	*8
789.00	*10	*	*	*	*	*	*	*	*	*	*
789.03	31	*	27	*	*25	*	*	*	*	*16	*
799	227	125	102	*	49	50	114	*68	*25	89	45
799.9	223	125	98	*	48	50	113	*65	*25	88	45
800-999	15,530	7,689	7,841	865	3,829	4,210	6,625	3,373	3,187	6,106	2,864
800	59	36	*22	*14	*32	*	*	*	*	*	*
801	247	182	*65	*17	148	*	*	*40	*54	*42	110
801.0	35	*26	*	*	*	*	*	*	*	*	*
801.2	136	*88	*	*	*65	*	*	*	*	*	*66
802	139	120	*19	*	82	*34	*	*25	*25	61	*27
802.2	49	46	*	*	31	*	*	*	*	*22	*
802.3	*18	*	*	*	*	*	—	*	*	*	*
802.4	*37	*35	*	*	*	*	*	*	*	*	*
805	559	251	308	*	96	133	319	104	122	241	91
805.0	182	123	*59	*	*47	*	*84	*	*	*93	*40
805.02	*66	*	*	*	*	*	*	*	*	*	*
805.2	120	*35	85	*	*	*	63	*	*40	*51	*
805.4	228	85	143	*	*25	*52	148	*60	*40	94	*35
806	*179	*117	*	*	*	*	*	*	*	*	*
807	232	135	97	*	*32	39	160	93	*38	40	*60
807.0	162	117	44	*	*23	35	103	51	*23	34	*53
807.01	*16	*	*	*	*	*	*	*	*	*	*
807.02	*26	*	*	*	*	*	*	*	*	*	*
807.03	*25	*	*	—	*	*	*	*	*	*	*
807.09	*25	*	*	—	*	*	*	*	*	*	*
808	330	136	195	*	*53	*113	157	52	96	106	*77
808.0	101	*70	*	*	*	*	*38	*	*	*	*
808.2	198	*61	137	*	*	*	104	*23	49	64	*61
812	302	95	207	26	37	47	192	71	49	116	66
812.0	190	49	141	*	*	*33	150	*53	*30	79	*28
812.00	73	*	51	*	*	*	51	*	*	41	*
812.01	77	*	65	—	*	*	71	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
812.09	33	*	*21	*	*	*	*23	*	*	*	*
812.2	40	*18	*22	*	*	*	*	*	*	*	*
812.21	*32	*	*	*	*	*	*	*	*	*	*
812.4	57	20	37	21	*	*	*	*10	*	*16	*22
812.41	28	*10	*18	17	*	*	*	*	*	*	*13
813	178	98	80	*15	61	34	68	39	33	67	38
813.0	28	*11	*17	*	*	*	*15	*	*	*	*
813.01	*16	*	*	*	*	*	*	*	*	*	*
813.2	*13	*	*	*	*	*	*	*	*	*	*
813.4	75	25	50	*	*15	*	47	*19	*19	*18	*19
813.41	*21	*	*15	*	*	*	*	*	*	*	*
813.42	25	*12	*	*	*	*	*15	*	*	*	*
813.44	*27	*	*20	*	*	*	*	*	*	*	*
816	*14	*11	*	*	*	*	*	*	*	*	*
820	2,107	612	1,495	*	*40	199	1,864	444	441	835	388
820.0	503	176	327	*	*	*24	466	129	136	176	62
820.00	*32	*	*	—	*	*	*	*	*	*	*
820.02	*49	*	*	*	*	*	*44	*	*	*	*
820.03	*56	*	*24	—	*	*	*52	*	*	*	*
820.09	364	110	254	*	*	*	346	99	111	123	*32
820.2	1,093	300	792	*	*	118	952	216	228	421	227
820.20	*54	*	*34	—	*	*	*46	*	*	*	*
820.21	940	240	700	—	*	92	840	180	192	364	204
820.22	99	*40	*58	*	*	*	67	*	*	*33	*
820.8	485	131	354	*	*	*49	431	93	76	231	84
821	464	220	244	44	145	110	165	72	76	251	65
821.0	276	156	120	35	100	*64	77	*42	*39	159	*36
821.00	*54	*	*	*	*	*	*	*	*	*	*
821.01	222	140	82	33	92	*52	*45	*	*34	125	*27
821.1	*34	*	*	—	*	*	*	*	*	*	*
821.2	144	*31	113	*	*	*	84	*	*31	*68	*
821.23	69	*	*57	*	*	*	*44	*	*	*	*
822	101	74	*27	*	*22	*48	*31	*	*	*34	*
822.0	60	*35	*25	*	*	*14	*31	*	*	*31	*
823	368	227	141	*12	128	147	81	91	73	148	57
823.0	149	70	79	*	*22	72	52	*29	*37	*60	*22
823.00	87	*37	50	*	*	*36	*29	*	*	*41	*
823.02	54	*	*28	*	*	*34	*	*	*	*	*
823.2	72	44	*27	*	*37	*	*	*	*	*33	*15
823.20	*21	*	*	*	*	*	*	*	*	*	*
823.22	46	*27	*	*	*25	*	*	*	*	*	*
823.8	43	*25	*	*	*	*	*	*	*	*	*
823.82	*31	*	*	*	*	*	*	*	*	*	*
824	372	179	193	*	152	115	96	104	92	126	51
824.0	*21	*15	*	*	*	*	*	*	*	*	*
824.2	*13	*	*	*	*	*	*	*	*	*	*
824.4	122	65	57	*	64	24	*33	*56	*20	31	*16
824.6	91	26	65	*	34	25	*31	*19	*24	33	*15
824.8	57	25	32	*	*20	*16	*	*15	*18	*19	*
825	66	45	*22	*	38	*	*	*16	*	*30	*
825.0	*17	*	*	*	*	*	*	*	*	*	*
825.2	*27	*19	*	*	*	*	*	*	*	*	*
840	28	*13	*16	*	*	*	*15	*	*	*10	*
840.4	22	*	*12	—	*	*	*14	*	*	*	*
844	34	*17	*17	*	*10	*	*	*	*	*	*
844.8	*17	*11	*	—	*	*	*	*	*	*	*
847	*29	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
850	147	81	65	*11	61	23	51	40	36	37	33
850.0	30	*9	*	*	*	*	*	*	*	*	*
850.1	37	24	*	*	*16	*	*	*	*14	*	*11
850.11	36	23	*	*	*16	*	*	*	*14	*	*11
850.5	57	35	*22	*	29	*	*	*	*12	*	*15
850.9	*17	*	*	*	*	*	*	*	*	*	*
851	314	236	*78	*	169	*	*40	*	*98	*59	*137
851.8	289	221	*68	*	153	*	*	*	*	*56	*131
852	416	228	189	*	*46	93	261	100	91	143	82
852.0	113	*67	*46	*	*	*	*48	*	*	*54	*
852.02	*40	*	*	*	*	*	*	*	*	*	*
852.2	282	145	137	*	*	*	209	*76	*60	*84	*63
852.20	*74	*	*	*	*	*	*57	*	*	*	*
852.21	121	*42	*79	*	*	*	104	*	*	*	*
853	64	*34	*	*	*	*	*37	*	*	*	*
853.0	61	*31	*	*	*	*	*37	*	*	*	*
854	*22	*	*	*	*	*	*	*	*	*	*
860	159	96	*64	*	86	*37	*	*37	*37	*57	*29
860.0	66	34	*32	*	*28	*16	*	*	*	*15	*
861	*44	*29	*	*	*	*	*	*	*	*	*
861.2	*25	*	*	*	*	*	*	*	*	*	*
861.21	*23	*	*	*	*	*	*	*	*	*	*
863	109	96	*	*	*72	*	*	*	*	*58	*
864	67	*50	*	*	*45	*	*	*	*	*	*
864.0	*47	*34	*	*	*	*	*	*	*	*	*
865	113	92	*	*	78	*	*	*	*	*51	*
865.0	106	88	*	*	75	*	*	*	*	*50	*
873	99	55	*43	*	35	*22	36	*17	*13	*35	*33
873.0	44	*20	*	*	*	*	*17	*	*	*	*
873.4	36	*19	*	*	*	*	*	*	*	*	*
873.42	*21	*	*	*	*	*	*	*	*	*	*
874	*12	*	*	*	*	*	*	*	*	*	*
879	*18	*	*	*	*	*	*	*	*	*	*
881	*21	*19	*	*	*	*	*	*	*	*	*
882	*13	*	*	*	*	*	*	*	*	*	*
883	*13	*	*	*	*	*	*	*	*	*	*
891	40	*22	*	*	*	*	*	*	*	*	*
920	*19	*	*	*	*	*	*	*	*	*	*
922	40	*14	*26	*	*	*	*25	*	*	*17	*
922.1	*19	*	*	*	*	*	*	*	*	*	*
924	73	*27	46	*	*	*	55	*	*26	*	*
924.0	31	*	*23	*	*	*	*25	*	*	*	*
924.01	*26	*	*20	—	*	*	*21	*	*	*	*
924.1	*26	*	*	—	*	*	*	*	*	*	*
933	*26	*	*	*	*	*	*	*	*	*	*
933.1	*25	*	*	*	*	*	*	*	*	*	*
935	*29	*14	*	*	*	*	*	*	*	*	*
935.1	*17	*	*	*	*	*	*	*	*	*	*
941	99	67	*32	54	*41	*	*	*	*	86	*
941.2	24	*16	*	*	*	*	*	*	*	*14	*
941.3	*74	*	*	*	*	*	*	*	*	*	—
942	272	139	*134	*41	*	*	*	*	*	142	*
942.2	22	*13	*	*11	*	*	*	*	*	*17	—
942.3	251	*	*	*	*	*	*	*	*	*125	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
943	122	*63	*59	*19	*75	*	*	*	*	84	*
943.2	*18	*	*	*	*	*	—	*	*	*	*
943.3	*75	*	*	*	*	*	*	*	*	*69	*
944	*32	*27	*	*	*	*	—	*	*	*21	*
945	83	50	*	*	*29	*	*	*	*	39	*
945.2	37	*20	*	*	*	*	*	*	*	*23	*
945.3	*31	*	*	*	*	*	*	*	*	*	*
949	*54	*	*	*29	*	*	—	—	*	*29	—
949.0	*54	*	*	*29	*	*	—	—	*	*29	—
958	*60	*	*	*	*	*	*	*	*	*	*
959	92	60	32	*	34	*21	33	*24	*18	*25	*24
959.0	50	32	*18	*	*13	*	*21	*	*10	*	*
959.01	48	31	*17	*	*12	*	*20	*	*	*	*
959.1	*19	*	*	*	*	*	*	*	*	*	*
962	*12	*	*	*	*	*	*	*	*	*	*
962.3	*10	*	*	*	*	*	*	*	*	*	*
963	*9	*	*	*	*	*	*	*	*	*	*
963.0	*9	*	*	*	*	*	—	*	*	*	*
965	184	73	112	*	112	57	*	41	29	67	47
965.0	83	31	52	*	34	41	*	*	*	24	*29
965.02	*16	*	*	*	*	*	*	*	*	*	*
965.09	36	*	*23	*	*14	*	*	*	*	*	*
965.1	*14	*	*	*	*	*	*	*	*	*	*
965.4	70	27	43	*	56	*	*	*15	*12	32	*
966	33	*	*23	*	*17	*	*	*	*	*	*14
966.3	*24	*	*17	*	*	*	*	*	*	*	*
967	28	*	*20	*	*13	*	*	*	*	*16	*
967.8	*13	*	*	*	*	*	*	*	*	*	*
968	*14	*	*	*	*12	*	*	*	*	*	*
969	240	118	121	*	165	52	*17	42	38	117	42
969.0	44	*	33	*	27	*	*	*	*	*15	*
969.3	*19	*	*	*	*14	*	*	*	*	*	*
969.4	142	89	52	*	103	33	*	*27	*14	84	*17
969.8	21	*	*16	*	*13	*	*	*	*	*	*
970	42	*27	*	*	*23	*	*	*	*	*	*
970.8	40	*25	*	*	*21	*	*	*	*	*	*
972	*25	*	*	*	*	*	*	*	*	*	*
977	37	*11	*27	*	*15	*19	*	*	*	*16	*
977.8	*24	*	*18	*	*	*	*	*	*	*	*
977.9	*12	*	*	*	*	*	*	*	*	*	*
980	*16	*12	*	*	*	*	*	*	*	*	*
980.0	*13	*	*	*	*	*	*	*	*	*	*
989	*90	*10	*	*	*	*	*	*	*	*	*
989.5	*85	*	*	*	*	*	*	*	*	*	*
991	*33	*28	*	*	*	*	*	*	*	*	*
994	*32	*24	*	*	*	*	*	*	*	*	*
995	138	53	85	*	*21	32	73	*30	*29	64	*14
995.1	28	*	20	*	*	*	*13	*	*	*11	*
995.2	*22	*	*	—	*	*	*	*	*	*	*
995.9	*63	*	*	*	*	*	*	*	*	*	*
996	3,078	1,551	1,527	213	483	1,094	1,288	770	559	1,219	530
996.0	89	67	*22	*	*	*	62	*	*	*32	*
996.01	30	*17	*	—	*	*	*23	*	*	*	*
996.04	*25	*22	*	*	*	*	*	*	*	*	*
996.1	73	*49	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
996.2	98	48	50	42	*35	*17	*	*	*19	37	*
996.3	*17	*	*	*	*	*	*	*	*	*	*
996.4	447	155	292	*	37	112	295	100	101	151	95
996.5	63	*36	*27	*	*	*	*	*	*	*32	*
996.59	*29	*	*	*	*	*	*	*	*	*	*
996.6	1,485	802	684	114	230	592	549	332	294	621	238
996.61	*74	*	*	—	*	*	*38	*	*	*	*
996.62	817	427	390	*76	106	385	251	190	127	386	114
996.63	95	*61	*	*	*	*	*	*	*	*	*
996.64	76	*54	*	*	*	*	*40	*	*	*	*
996.66	195	95	100	*	*	*61	116	*	*55	*61	*30
996.67	84	*44	*40	*	*	*27	*	*	*	*	*
996.68	*35	*	*	*	*	*	*	*	*	*	*
996.69	73	*	*37	*	*	*	*	*	*	*35	*
996.7	560	254	306	*	74	188	283	150	98	199	114
996.72	84	52	32	*	*	*19	49	*17	*16	33	*
996.73	169	87	82	—	*25	69	76	*31	*	51	68
996.74	114	59	55	*	*	*48	57	*	*	56	*
996.77	39	*	*26	—	*	*	*21	*	*	*	*
996.78	36	*14	*21	*	*12	*	*	*	*	*	*
996.8	246	133	112	*	70	131	*	*74	*	117	*
996.81	126	*46	80	*	*47	*62	*	*	*	73	*
997	629	293	336	*	134	254	226	125	115	260	129
997.0	*31	*	*	*	*	*	*	*	*	*	*
997.1	53	*29	*	*	*	*	*	*	*	*	*
997.2	*40	*	*23	—	*	*	*	*	*	*	*
997.3	38	*18	*	*	*	*22	*	*	*	*	*
997.4	236	81	155	*	68	92	72	*51	*54	78	52
997.5	42	*24	*	*	*	*15	*	*	*	*	*
997.6	179	104	*74	*	*	*76	*80	*	*	110	*
997.62	145	*82	*	*	*	*63	*	*	*	*96	*
997.69	*33	*	*	—	*	*	*	*	*	*	*
998	1,631	713	918	33	278	698	621	383	308	684	256
998.1	233	91	143	*	40	73	112	65	38	83	47
998.11	111	57	54	*	*17	*25	63	*27	*19	43	*23
998.12	103	*31	72	*	*18	43	*41	*36	*	*32	*18
998.13	*19	*	*	*	*	*	*	*	*	*	*
998.2	92	*	*73	*	*	*	*53	*	*	*	*
998.3	138	*43	*96	*	*	*52	*67	*	*27	*	*
998.32	102	*32	*70	*	*	*	*64	*	*	*	*
998.5	1,003	492	511	*	179	480	327	202	205	443	153
998.51	*33	*	*25	*	*	*	*	*	*	*	*
998.59	971	485	486	*	176	461	316	196	203	430	142
998.6	*51	*	*	*	*	*	*	*	*	*	*
998.8	108	*56	52	*	*	*40	*51	*	*	*43	*
998.83	*55	*	*	*	*	*	*	*	*	*	*
998.89	51	*	*33	*	*	*	*	*	*	*	*
999	43	*12	*	*	*	*	*	*	*	*	*
V01-V83	18,806	3,368	15,438	401	11,551	1,913	4,941	3,327	4,104	6,969	4,406
V10	*10	*	*	*	*	*	*	*	*	*	*
V12	28	*	*22	*	*	*	*	*	*	*	*
V12.5	*18	*	*14	*	*	*	*	*	*	*	*
V27	10,791	—	10,791	*14	10,742	35	—	1,695	2,131	4,370	2,595
V27.0	9,851	—	9,851	*	9,813	*26	—	1,501	1,848	4,080	2,421
V27.1	46	—	46	*	45	*	—	*	*	23	*
V27.2	239	—	239	—	235	*	—	*39	*36	112	53
V27.9	643	—	643	*	636	*	—	148	237	148	109
V30	78	*	*46	78	—	—	—	*	*	*	*
V30.1	78	*	*46	78	—	—	—	*	*	*	*

See footnotes at end of table.

Table 46. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of days of care in thousands											
V42	*25	*	*	*	*	*	*	*	*	*	*
V45	53	*19	34	*	*11	*13	*26	*	*	*18	*
V45.8	26	*	*13	*	*	*	*	*	*	*	*
V45.89	*13	*	*	*	*	*	*	*	*	*	*
V50	*13	*	*12	*	*	*	*	*	*	*	*
V53	19	*12	*7	*	*	*	*13	*	*	*	*
V53.3	17	*11	*	*	*	*	*12	*	*	*	*
V53.31	*11	*	*	*	*	*	*	*	*	*	*
V54	139	*41	98	*	*	*	119	*	*60	*43	*
V54.1	*64	*	*	*	*	*	*60	*	*	*	*
V54.8	*63	*	*42	*	*	*	*51	*	*	*	*
V54.81	*42	*	*	—	*	*	*35	*	*	*	*
V55	243	136	107	*	53	90	83	71	49	82	*41
V55.2	*52	*	*	*	*	*	*	*	*	*	*
V55.3	167	97	70	*	*39	66	*58	*50	*33	*53	*
V56	*52	*	*	*	*	*	*	*	*	*	*
V56.0	*45	*	*	*	*	*	*	*	*	*	*
V57	6,102	2,489	3,613	*	520	1,330	4,213	1,185	1,656	1,871	1,390
V57.1	1,038	426	612	*	*102	230	700	348	101	428	162
V57.8	5,048	2,061	2,987	*	418	1,099	3,501	827	1,554	1,441	1,226
V57.89	5,048	2,061	2,987	*	418	1,099	3,501	827	1,554	1,441	1,226
V58	969	514	455	183	115	313	358	210	139	389	230
V58.0	65	*32	*	*	*	*	*54	*	*	*17	*
V58.1	702	412	290	175	107	267	154	155	66	309	172
V58.4	52	*	*34	*	*	*	*41	*	*	*	*
V58.49	*38	*	*25	*	*	*	*	*	*	*	*
V58.8	89	*26	62	*	*	*	66	*	*49	*	*
V58.89	83	*	*60	*	*	*	*61	*	*49	*	*
V59	*20	*	*	*	*	*	*	*	*	*	*
V59.4	*18	*	*	*	*	*	*	*	*	*	*
V66	*23	*	*	—	*	*	*	*	*	*	*
V70	48	*	*35	*	*16	*	*	*	*	*	*
V70.7	24	*	*10	*	*15	*	*	*	*	*	*
V71	38	*12	*	*	*	*	*	*	*	*	*
V71.4	*9	*	*	*	*	*	*	*	*	*	*

* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate is preceded by asterisk, use according to size of corresponding estimate in Table 45; see "Methods" section.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes	165,517	67,051	98,465	7,277	41,163	41,446	75,631	35,532	39,478	59,942	30,565
001-139	4,143	1,887	2,256	457	897	1,117	1,671	908	875	1,564	795
003	*8	*	*	*	*	*	*	*	*	*	*
003.0	*5	*	*	*	*	*	*	*	*	*	*
005	*9	*	*6	*	*	*	*	*	*	*	*
005.9	*8	*	*5	*	*	*	*	*	*	*	*
008	393	170	223	96	50	63	184	93	98	142	59
008.4	230	98	132	*8	23	43	155	65	61	71	32
008.45	222	95	127	*7	21	42	152	63	60	70	30
008.6	56	30	26	52	*	*	*	*	14	27	10
008.61	52	29	23	50	*	*	*	*	13	25	9
008.8	103	40	63	34	25	19	26	22	22	43	16
009	24	10	14	*	10	*6	*7	*5	*	10	*5
009.0	14	*6	*8	*	*6	*	*	*	*	*5	*
009.1	*7	*	*	*	*	*	*	*	*	*	*
011	13	*8	*	*	*	*	*	*	*	*6	*
011.9	9	*7	*	*	*	*	*	*	*	*	*
031	*8	*	*	*	*	*	*	*	*	*	*
033	*6	*	*	*5	*	*	*	*	*	*	*
034	31	15	16	17	*8	*	*	*5	9	10	*6
034.0	29	14	15	16	*8	*	*	*5	9	9	*5
036	*5	*	*	*	*	*	-	*	*	*	*
038	803	394	410	23	73	206	501	169	178	289	168
038.0	42	23	19	*	*	*9	29	13	10	11	*8
038.1	140	77	63	*	14	42	80	30	28	49	33
038.10	13	*6	*7	*	*	*	*9	*	*	*	*
038.11	93	54	39	*	10	31	50	19	17	35	22
038.19	34	18	16	*	*	*9	20	*8	*8	11	*7
038.2	*8	*	*5	*	*	*	*	*	*	*	*
038.3	*6	*	*	*	*	*	*	*	*	*	*
038.4	151	67	83	*5	13	39	93	34	37	46	34
038.40	11	*7	*	*	*	*	*7	*	*	*	*
038.42	75	25	50	*	*	18	51	17	18	21	19
038.43	15	10	*	*	*	*	9	*	*	*	*
038.49	43	23	20	*	*	13	24	9	*9	17	*9
038.8	35	18	17	*	*	11	18	*8	*6	15	*5
038.9	421	202	219	11	37	100	274	81	93	163	84
041	1,293	517	776	124	270	286	613	264	300	484	245
041.0	205	82	123	13	54	39	99	45	49	68	43
041.00	20	9	10	*	*	*6	*8	*	*	*8	*
041.01	11	*5	*6	*	*	*	*	*	*	*	*
041.02	45	12	32	*	26	*7	9	*9	12	18	*6
041.04	97	40	57	*5	12	16	64	22	20	32	22
041.09	30	13	16	*	*6	*7	13	*7	*9	*7	*7
041.1	345	190	156	41	90	102	112	64	72	142	67
041.10	21	12	9	*	*5	*7	*7	*	*6	*	*5
041.11	276	153	124	33	76	80	87	51	54	117	54
041.19	48	26	23	*6	*9	15	18	*8	12	20	*8
041.3	79	19	60	*	*8	17	51	21	24	23	11
041.4	336	73	263	35	60	58	183	66	79	116	74
041.6	59	24	35	*	*	10	43	11	11	31	*6
041.7	90	52	38	11	15	22	42	19	23	36	12
041.8	155	68	87	14	31	34	77	35	36	55	29
041.84	11	*6	*6	*	*	*	*5	*	*	*	*
041.85	60	27	33	*8	10	11	31	16	14	20	10
041.86	55	24	31	*	10	15	30	13	12	20	10
041.89	28	11	17	*	*7	*6	10	*	*6	10	*6
041.9	16	*7	10	*	*6	*	*	*	*	10	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
042	138	91	47	*	76	57	*	42	15	66	14
047	48	24	25	23	17	*6	*	*6	*8	21	13
047.8	*6	*	*	*	*	*	—	*	*	*	*
047.9	42	22	21	19	16	*5	*	*6	*7	18	11
053	71	22	49	*	*	19	46	19	16	24	13
053.1	26	*8	18	*	*	*6	19	*8	*7	*6	*
053.19	22	*6	15	*	*	*5	16	*6	*7	*5	*
053.9	36	11	25	*	*	*8	23	*8	*7	15	*5
054	58	14	44	*8	33	10	*8	10	12	27	10
054.1	17	*	15	*	13	*	*	*	*	*7	*
054.10	14	*	11	*	10	*	*	*	*	*5	*
054.2	*6	*	*	*	*	*	*	*	*	*	*
054.9	28	*6	22	*	18	*	*	*	*6	15	*
057	*7	*	*	*	*	*	*	*	*	*	*
070	431	256	175	*	140	249	41	101	56	171	103
070.3	51	30	22	*	21	24	*6	13	*7	22	9
070.30	29	17	13	*	13	13	*	*7	*6	12	*6
070.32	21	12	9	—	*8	11	*	*6	*	10	*
070.4	11	*6	*	—	*	*7	*	*	*	*	*
070.44	*7	*	*	—	*	*	*	*	*	*	*
070.5	355	214	141	*	114	213	28	82	47	137	89
070.51	173	103	70	*	58	102	12	37	30	75	31
070.54	179	110	69	*	55	109	15	44	17	61	56
070.9	*9	*	*	*	*	*	*	*	*	*5	*
075	15	*8	*7	*6	*8	*	*	*	*5	*	*
078	18	*5	13	*	12	*	*	*	*	9	*
078.1	11	*	9	*	*9	*	*	*	*	*5	*
078.10	*6	*	*5	*	*5	*	*	*	*	*	*
079	145	66	79	74	33	15	22	32	29	58	27
079.6	9	*	*6	*8	*	*	*	*	*	*	*
079.8	*9	*	*5	*	*	*	*	*	*	*	*
079.89	*7	*	*	*	*	*	*	*	*	*	*
079.9	120	58	63	60	25	14	21	27	24	47	22
079.99	118	58	60	60	22	14	21	27	23	46	22
088	*8	*	*	*	*	*	*	*	*	*	*
088.8	*7	*	*	*	*	*	*	*	*	*	*
088.81	*6	*	*	*	*	*	*	*	*	*	—
094	*9	*6	*	—	*	*	*	*	*	*	*
094.0	*5	*	*	—	*	*	*	*	*	*	*
097	*6	*	*	—	*	*	*	*	*	*	*
098	*5	*	*	*	*	*	*	*	*	*	*
099	*6	*	*5	*	*5	*	*	*	*	*	*
099.5	*5	—	*5	*	*	*	—	*	*	*	*
110	64	39	25	*	16	13	32	17	15	24	*8
110.1	31	18	13	*	*	*6	20	*9	*9	11	*
110.3	9	*	*	*	*	*	*6	*	*	*	*
110.4	15	12	*	*	*7	*5	*	*	*	*	*
112	299	118	181	33	57	79	129	68	71	107	53
112.0	167	71	95	20	30	48	69	39	41	59	28
112.1	24	—	24	*	12	*5	*	*5	*	12	*
112.2	25	10	15	*	*	*	17	*6	*7	*8	*
112.3	17	*5	12	*7	*	*	*	*	*	*6	*
112.4	10	*6	*	*	*	*	*6	*	*	*	*
112.5	9	*5	*	*	*	*	*	*	*	*	*
112.8	42	20	22	*	*	13	22	9	*8	13	11
112.84	33	16	17	*	*	*9	20	*8	*7	10	*9
112.89	*8	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
117	24	16	*9	*	*	*7	10	*	*	11	*5
117.3	*8	*	*5	*	*	*	*	*	*	*	*
117.9	11	*8	*	*	*	*	*6	*	*	*6	*
131	11	*	10	*	*9	*	*	*	*	*7	*
131.0	*7	*	*7	*	*6	*	*	*	*	*	*
135	56	18	38	*	9	33	13	17	12	19	*8
136	18	*9	9	*	9	*5	*	*	*	*7	*
136.3	12	*7	*	*	*7	*	*	*	*	*5	*
138	29	11	18	*	*	*8	20	*5	*7	*8	*9
139	*5	*	*	*	*	*	*	*	*	*	*
140-239	4,532	1,928	2,603	118	665	1,548	2,201	1,051	1,012	1,666	803
141	*8	*6	*	—	*	*5	*	*	*	*	*
145	*6	*	*	—	*	*	*	*	*	*	*
146	*8	*7	*	—	*	*	*	*	*	*	*
147	*7	*	*	*	*	*	*	*	*	*	*
150	32	27	*	—	*	9	22	*6	*7	14	*5
150.5	*8	*7	*	—	*	*	*5	*	*	*	*
150.9	18	15	*	—	*	*	13	*	*	9	*
151	40	27	14	*	*	12	26	10	*8	12	11
151.0	12	11	*	—	*	*	*7	*	*	*	*
151.8	*5	*	*	—	*	*	*	*	*	*	*
151.9	15	9	*6	*	*	*	10	*	*	*	*
152	*7	*	*	—	*	*	*	*	*	*	*
153	197	95	101	*	11	62	124	46	43	78	29
153.0	*8	*	*	—	*	*	*	*	*	*	*
153.1	12	*6	*6	—	*	*	10	*	*	*	*
153.2	*6	*	*	—	*	*	*	*	*	*	*
153.3	36	21	15	—	*	11	24	*8	10	13	*5
153.4	30	12	18	—	*	*7	21	*9	*7	11	*
153.5	*5	*	*	—	*	*	*	*	*	*	*
153.6	26	11	14	—	*	*8	16	*	*	12	*5
153.8	10	*	*	—	*	*	*	*	*	*	*
153.9	62	30	32	*	*	21	37	11	16	27	*8
154	65	37	27	*	*5	24	35	16	18	22	*8
154.0	18	10	*9	—	*	*8	10	*7	*	*6	*
154.1	37	23	14	*	*	14	18	*8	11	13	*5
154.8	*7	*	*	—	*	*	*6	*	*	*	*
155	41	25	16	*	*	17	20	9	*7	19	*5
155.0	29	19	10	*	*	13	13	*6	*	14	*
155.1	*6	*	*	—	*	*	*	*	*	*	*
155.2	*5	*	*	*	*	*	*	*	*	*	*
156	15	*8	*7	—	*	*	10	*	*	*6	*
156.0	*5	*	*	—	*	*	*	*	*	*	*
157	63	36	27	*	*	20	41	18	13	19	13
157.0	15	*8	*7	—	*	*5	10	*	*	*5	*
157.8	11	*6	*	—	*	*	*8	*	*	*	*
157.9	30	17	13	—	*	10	19	*7	*	9	*8
158	*6	*	*	*	*	*	*	*	*	*	*
161	15	11	*	—	*	10	*	*	*	*8	*
161.9	*6	*	*	—	*	*	*	*	*	*	*
162	362	186	176	*	18	113	231	83	100	133	46
162.2	16	9	*7	—	*	*9	*7	*	*	*8	*
162.3	68	35	32	*	*	18	46	14	18	27	*8
162.4	*8	*	*	—	*	*	*5	*	*	*	*
162.5	43	18	25	*	*	10	30	*6	14	15	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
162.8	36	22	14	–	*	12	20	18	*6	*8	*
162.9	192	97	95	*	*8	62	122	40	57	72	23
170	24	*8	16	9	10	*	*	*7	*	*7	*7
170.7	12	*	*8	*7	*6	*	*	*	*	*	*
171	22	12	10	*	*	*8	*6	*	*	*9	*
171.3	*7	*	*	*	*	*	*	*	*	*	*
172	12	*6	*6	–	*	*5	*6	*	*	*	*
173	20	12	*8	*	*	*	15	*	*6	*7	*
173.3	*6	*	*	–	–	*	*	*	*	*	*
174	154	–	154	*	18	60	75	36	35	57	26
174.4	11	–	11	–	*	*	*	*	*	*	*
174.8	34	–	34	–	*	13	17	13	*	*8	*9
174.9	94	–	94	–	11	37	46	18	25	40	11
180	20	–	20	–	*8	*8	*	*	*	*8	*5
180.8	*5	–	*5	–	*	*	*	*	*	*	*
180.9	12	–	12	–	*	*5	*	*	*	*6	*
182	48	–	48	–	*	22	23	10	12	16	10
182.0	47	–	47	–	*	22	23	10	12	16	10
183	62	–	62	*	*6	28	28	17	11	25	9
183.0	60	–	60	*	*6	26	27	16	10	24	9
184	*8	–	*8	*	*	*	*	*	*	*	*
185	193	193	–	*	*	51	142	44	48	65	37
186	*5	*5	–	*	*	*	*	*	*	*	*
186.9	*5	*5	–	*	*	*	*	*	*	*	*
188	67	51	16	*	*	14	51	22	15	17	13
188.8	20	16	*	–	*	*6	14	*7	*5	*	*
188.9	31	23	*8	*	*	*6	24	9	*7	9	*5
189	66	39	27	*	*	21	39	16	11	29	10
189.0	57	34	24	*	*	19	33	13	10	25	9
191	52	30	22	11	*9	21	10	13	*7	23	9
191.1	*6	*	*	*	*	*	*	*	*	*	*
191.3	*8	*5	*	*	*	*6	*	*	*	*5	*
191.8	*8	*	*	*	*	*	*	*	*	*	*
191.9	14	*8	*6	*	*	*6	*	*	*	*7	*
193	29	*7	23	*	15	10	*	*7	*	10	*7
194	*8	*	*	*	*	*	*	*	*	*	*
194.0	*6	*	*	*	*	*	*	*	*	*	*
195	12	*5	*7	*	*	*	*6	*	*	*7	*
195.0	*6	*	*	*	*	*	*	*	*	*	*
196	158	67	90	*	20	58	78	37	43	48	29
196.0	21	14	*7	*	*	*8	*9	*	*6	*8	*
196.1	23	11	12	*	*	*9	12	*7	*6	*7	*
196.2	56	27	30	*	*	21	31	13	15	19	*9
196.3	29	*	26	*	*5	10	13	*7	*9	*7	*6
196.5	*6	*	*	*	*	*	*	*	*	*	*
196.6	9	*	*6	–	*	*	*	*	*	*	*
196.9	11	*6	*	–	*	*	*7	*	*	*	*
197	532	245	287	*	37	206	285	139	120	185	89
197.0	130	66	64	*	11	52	64	33	29	45	22
197.1	10	*5	*	*	*	*	*	*	*	*	*
197.2	54	26	28	*	*	20	31	13	11	20	10
197.4	9	*	*7	–	*	*	*	*	*	*	*
197.5	12	*6	*5	–	*	*	*6	*	*	*	*
197.6	84	29	55	*	*6	34	43	24	15	30	15
197.7	213	100	112	*	12	81	119	57	51	71	33
197.8	18	*9	9	*	*	*	12	*	*	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
198	434	214	220	*	41	155	233	109	108	146	71
198.1	10	*6	*	*	*	*	*6	*	*	*	*
198.2	9	*	*6	*	*	*	*6	*	*	*	*
198.3	106	49	57	*	*8	48	50	29	22	38	16
198.4	*6	*	*	*	*	*	*	*	*	*	*
198.5	186	102	85	*	19	56	109	43	57	57	29
198.6	*7	—	*7	—	*	*	*	*	*	*	*
198.7	21	13	*8	*	*	*8	13	*6	*	*7	*
198.8	83	36	47	*	*8	31	44	22	17	31	14
198.82	15	*5	10	—	*	*7	*7	*	*	*6	*
198.89	66	30	36	*	*7	22	36	17	15	24	10
199	148	65	82	*	24	53	70	31	33	50	34
199.0	46	17	29	—	10	23	13	*7	*8	19	11
199.1	102	48	54	*	14	30	57	24	25	31	23
200	28	15	13	*	*	*6	16	10	*	*8	*6
200.0	17	10	*7	*	*	*	10	*8	*	*	*
200.00	*8	*	*	*	*	*	*	*	*	*	*
200.1	*6	*	*	*	*	*	*	*	*	*	*
201	25	14	11	*6	*7	*7	*	*7	*5	*6	*8
201.5	*5	*	*	*	*	*	*	*	*	*	*
201.9	19	10	*8	*	*	*7	*	*6	*	*	*
201.90	14	*7	*6	*	*	*6	*	*	*	*	*
202	137	72	65	*	12	34	87	30	29	56	22
202.1	*7	*	*	*	*	*	*	*	*	*	*
202.8	121	61	60	*	10	29	80	28	23	51	19
202.80	85	44	41	*	*8	20	56	21	16	34	15
202.81	*9	*5	*	*	*	*	*	*	*	*	*
202.82	*8	*	*	*	*	*	*7	*	*	*6	*
202.83	*9	*	*	*	*	*	*6	*	*	*	*
202.88	*5	*	*	—	*	*	*	*	*	*	*
203	76	37	39	—	*	31	43	19	14	33	11
203.0	74	35	39	—	*	30	42	18	13	32	10
203.00	72	34	38	—	*	29	41	17	13	31	10
204	123	63	59	31	*6	17	69	24	29	43	27
204.0	44	29	15	29	*5	*5	*	*7	*7	14	15
204.00	30	20	10	17	*	*	*	*5	*5	12	*7
204.01	14	*9	*5	12	*	*	*	*	*	*	*8
204.1	76	32	44	—	*	11	64	16	21	28	10
204.10	72	30	42	—	*	11	62	15	20	27	10
205	58	31	27	*	*7	16	31	11	15	24	*8
205.0	41	23	18	*	*5	11	20	*8	11	16	*6
205.00	36	20	16	*	*	9	19	*7	10	14	*5
205.1	16	*8	*8	*	*	*	10	*	*	*8	*
205.10	15	*7	*8	*	*	*	9	*	*	*7	*
208	12	*7	*	*	*	*	9	*	*	*7	*
208.0	*6	*	*	*	*	*	*	*	*	*	*
208.00	*6	*	*	*	*	*	*	*	*	*	*
210	*8	*	*	*	*	*	*	*	*	*	*
210.2	*5	*	*	*	*	*	*	*	*	*	*
211	185	84	101	*	12	53	119	45	42	62	36
211.1	23	*7	16	*	*	*6	16	*5	*	*8	*5
211.2	*6	*	*	*	*	*	*	*	*	*	*
211.3	133	62	72	*	*7	38	88	33	31	43	27
211.4	13	9	*	*	*	*	*8	*	*	*5	*
212	*7	*	*	*	*	*	*	*	*	*	*
213	*6	*	*	*	*	*	*	*	*	*	*
214	17	*7	10	*	*	*6	*7	*	*	*6	*
214.1	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
215	*6	*	*	*	*	*	*	*	*	*	*
216	13	*	*8	*	*6	*	*	*	*	*6	*
216.5	*6	*	*5	*	*	*	*	*	*	*	*
218	475	–	475	–	247	209	18	78	96	209	92
218.0	51	–	51	–	24	26	*	11	9	21	9
218.1	122	–	122	–	57	61	*	22	25	57	19
218.2	76	–	76	–	39	34	*	14	15	34	13
218.9	226	–	226	–	127	88	11	31	46	98	52
220	64	–	64	*	25	21	16	11	13	29	10
221	*9	–	*9	–	*7	*	*	*	*	*	*
225	38	14	24	*	*7	13	16	10	*7	11	9
225.2	28	10	18	*	*5	9	14	*8	*6	*8	*7
226	10	*	*9	–	*5	*	*	*	*	*	*
227	29	10	19	*	*6	11	12	*7	*	10	*8
227.0	*8	*	*5	*	*	*	*	*	*	*	*
227.1	11	*	*8	*	*	*6	*	*	*	*	*
227.3	11	*	*6	*	*	*	*	*	*	*	*
228	22	10	12	*	*6	*8	*	*	*5	*8	*
228.0	20	10	11	*	*6	*8	*	*	*	*8	*
228.04	*7	*	*	*	*	*	*	*	*	*	*
228.09	*7	*	*	*	*	*	*	*	*	*	*
233	37	*	33	–	12	14	11	10	*7	12	*8
233.0	17	*	17	–	*	*8	*7	*	*	*5	*
233.1	14	–	14	–	*9	*	*	*	*	*	*
235	20	11	*9	*	*	*6	11	*	*6	*5	*
235.2	12	*	*7	*	*	*	*6	*	*	*	*
235.7	*5	*	*	*	*	*	*	*	*	*	*
236	*8	*	*6	–	*	*	*	*	*	*	*
237	15	*7	*8	*	*6	*	*	*	*	*5	*
237.7	*8	*	*6	*	*	*	*	*	*	*	*
237.70	*6	*	*	*	*	*	*	*	*	*	*
238	73	36	36	*	*6	10	55	16	16	27	14
238.4	15	*8	*7	*	*	*	9	*	*	*	*
238.7	50	25	24	*	*	*	44	11	10	20	*8
239	28	11	17	*	*	*9	14	*6	*6	9	*7
239.1	*5	*	*	–	*	*	*	*	*	*	*
239.6	*7	*	*	*	*	*	*	*	*	*	*
240–279	19,280	8,122	11,158	617	2,537	5,946	10,179	4,273	4,752	6,956	3,299
240	16	*	13	*	*	*	*7	*	*	*7	*
240.9	15	*	13	*	*	*	*7	*	*	*7	*
241	38	*6	32	*	12	12	14	10	*5	15	*8
241.0	19	*	17	*	*6	*7	*6	*6	*	*7	*
241.1	16	*	13	–	*	*5	*7	*	*	*7	*
242	92	21	72	*	23	25	42	29	21	27	16
242.0	24	*	21	*	*9	*8	*7	*7	*5	*7	*
242.00	22	*	20	*	*8	*7	*7	*7	*	*6	*
242.9	59	15	44	*	14	16	29	19	15	17	*9
242.90	56	14	42	*	13	14	28	18	15	15	*9
244	1,684	319	1,365	*8	209	416	1,051	411	444	533	297
244.0	45	*	42	*	*5	12	27	11	11	14	10
244.1	13	*	10	*	*	*	*	*	*	*7	*
244.8	10	*	*8	–	*	*	*5	*	*	*	*
244.9	1,614	309	1,305	*8	198	396	1,012	395	426	509	284
245	16	*	15	*	*7	*7	*	*5	*	*	*
245.2	12	*	11	*	*5	*5	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
246	22	*	19	*	*5	*7	10	*5	*	*9	*
246.8	*5	*	*	—	*	*	*	*	*	*	*
246.9	15	*	13	*	*	*	*6	*	*	*7	*
250	5,435	2,541	2,894	43	588	1,879	2,926	1,246	1,237	2,065	887
250.0	4,038	1,863	2,175	19	353	1,356	2,310	930	939	1,548	621
250.00	3,403	1,583	1,821	*6	264	1,131	2,002	781	819	1,280	523
250.01	303	139	164	11	54	97	142	80	61	116	46
250.02	295	126	169	*	26	115	154	61	52	135	47
250.03	36	15	21	*	9	13	12	*9	*6	17	*
250.1	151	81	70	20	82	33	16	28	28	65	30
250.10	28	17	11	*	13	10	*	*	*	13	*6
250.11	70	36	33	14	39	11	*5	14	15	26	15
250.12	18	*9	*9	*	*	*6	*6	*	*	*8	*
250.13	36	19	17	*	25	*5	*	*7	*	17	*6
250.2	21	11	10	*	*5	*8	*8	*7	*	*8	*
250.20	10	*5	*	—	*	*	*	*	*	*	*
250.22	*7	*	*	—	*	*	*	*	*	*	*
250.3	*6	*	*	*	*	*	*	*	*	*	*
250.4	285	138	148	*	30	120	134	58	57	94	76
250.40	194	96	97	*	10	81	102	35	37	63	59
250.41	42	20	22	—	13	19	10	9	11	13	9
250.42	39	17	22	—	*	16	18	11	*8	14	*6
250.43	11	*	*6	—	*	*	*	*	*	*	*
250.5	113	50	62	—	15	55	44	28	31	33	21
250.50	61	28	34	—	*	30	28	14	16	17	14
250.51	30	14	16	—	*8	14	*8	*8	9	*8	*
250.52	13	*6	*7	—	*	*7	*	*	*	*	*
250.53	*9	*	*5	—	*	*	*	*	*	*	*
250.6	373	176	197	*	48	154	170	90	90	133	61
250.60	230	109	121	—	15	93	122	55	56	77	43
250.61	68	32	36	*	19	31	18	18	16	23	10
250.62	55	26	29	—	*7	22	26	11	14	26	*5
250.63	20	9	11	—	*8	*8	*	*6	*5	*7	*
250.7	94	58	36	*	*6	34	53	24	17	34	19
250.70	61	36	25	—	*	19	40	15	12	22	12
250.71	18	11	*7	*	*	*8	*8	*7	*	*5	*
250.72	11	*8	*	—	*	*6	*5	*	*	*6	*
250.8	295	135	160	*	34	95	164	75	61	111	47
250.80	194	86	107	*	12	58	123	46	44	68	35
250.81	45	22	23	*	12	15	16	14	*7	18	*7
250.82	42	20	22	—	*	17	21	11	*6	21	*
250.83	14	*7	*7	*	*	*5	*	*	*	*	*
250.9	58	26	32	*	13	21	23	*6	9	36	*8
250.90	14	*7	*7	*	*5	*	*	*	*	11	*
250.92	31	14	17	*	*	12	15	*	*6	17	*5
250.93	*8	*	*	*	*	*	*	*	*	*5	*
251	91	35	56	13	18	21	39	26	19	31	15
251.1	*8	*	*	*	*	*	*	*	*	*	*
251.2	35	14	20	9	9	*6	11	*9	*7	12	*7
251.8	46	16	29	*	*7	13	25	15	10	15	*6
252	41	10	31	*	*5	15	20	9	*8	15	9
252.0	26	*6	20	*	*	11	13	*6	*	*9	*6
252.1	14	*	10	*	*	*	*6	*	*	*6	*
253	83	37	47	*6	14	23	40	19	14	32	18
253.2	12	*6	*6	*	*	*	*	*	*	*	*
253.5	12	*7	*6	*	*	*	*	*	*	*	*
253.6	48	20	28	*	*	11	33	12	*8	20	*8
255	85	30	56	*	11	25	46	18	24	28	15
255.0	*9	*	*8	*	*	*	*	*	*	*	*
255.1	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
255.4	55	21	34	*	*7	15	32	12	15	18	9
255.9	*8	*	*6	—	*	*	*5	*	*	*	*
256	12	—	12	*	10	*	*	*	*	*	*
256.4	10	—	10	*	9	*	*	*	*	*	*
257	*5	*5	—	*	*	*	*	*	*	*	*
257.2	*5	*5	—	*	*	*	*	*	*	*	*
259	*6	*	*	*	*	*	*	*	*	*	*
260	27	12	15	*	*	*8	18	*	*	20	*
261	26	12	15	*	*	*7	17	*	*9	11	*
262	10	*	*6	*	*	*	*5	*	*	*	*
263	408	189	219	*6	36	94	271	73	107	165	63
263.1	*8	*	*	*	*	*	*	*	*	*	*
263.8	26	14	12	*	*	*6	16	*	*6	10	*7
263.9	370	169	201	*5	31	85	248	64	99	152	55
266	42	13	30	—	*	*6	32	13	10	14	*6
266.2	41	12	29	—	*	*6	32	13	10	13	*6
269	11	*	*7	*	*	*	*	*	*	*	*
270	23	11	12	*5	*	*8	*6	*	*	10	*6
270.4	9	*	*	*	*	*	*	*	*	*	*
270.6	*6	*	*	*	*	*	*	*	*	*	*
271	24	12	13	*	*5	9	*	*8	*6	*5	*
271.3	19	*8	11	*	*	*8	*	*	*6	*	*
272	3,451	1,735	1,715	*	259	1,280	1,907	900	846	1,128	577
272.0	1,275	616	660	*	91	492	690	426	303	357	189
272.1	47	29	18	*	14	19	13	*8	14	15	10
272.2	19	9	10	—	*	*	12	*	*	*8	*7
272.4	2,019	1,036	983	*	141	732	1,145	441	504	719	355
272.6	*5	*	*	—	*	*	*	*	*	*	*
272.8	13	*7	*7	*	*	*5	*7	*5	*	*	*
272.9	65	33	32	*	*6	22	37	15	18	18	13
273	66	31	35	*	12	18	34	15	17	27	*7
273.1	*8	*	*5	—	*	*	*6	*	*	*	*
273.3	*5	*	*	*	*	*	*	*	*	*	*
273.8	49	24	25	*	11	13	23	10	15	19	*6
274	256	177	79	*	14	78	164	68	69	77	41
274.0	59	41	18	—	*	16	37	17	13	21	*8
274.9	196	135	60	*	*9	61	125	51	56	56	33
275	272	109	164	10	54	95	113	54	77	102	38
275.0	15	*9	*6	*	*	*5	*	*	*	*6	*
275.2	113	43	69	*	22	41	48	18	38	48	*9
275.3	34	17	17	*	*9	11	10	*6	*9	12	*7
275.4	108	37	71	*	19	37	49	27	27	34	20
275.41	54	20	34	*	14	20	18	11	15	18	10
275.42	47	14	32	*	*	16	25	14	10	15	*8
275.49	*7	*	*	*	*	*	*6	*	*	*	*
276	5,705	2,346	3,359	455	791	1,337	3,121	1,067	1,502	2,124	1,012
276.0	148	69	79	*6	12	27	103	34	38	54	22
276.1	962	385	578	26	89	244	603	164	265	351	183
276.2	307	151	155	43	45	86	134	45	67	122	72
276.3	56	23	33	*	10	18	25	*6	17	24	9
276.4	23	10	13	*	*	*7	11	*	*8	*5	*6
276.5	2,528	1,053	1,475	338	357	465	1,369	500	655	935	438
276.6	91	40	51	*	12	34	42	13	22	38	18
276.7	457	225	232	10	42	118	286	91	123	169	74
276.8	1,029	345	684	20	209	310	490	200	272	391	166
276.9	104	46	58	*	13	29	58	12	35	34	23

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
277	65	31	33	19	25	12	*8	15	15	21	13
277.0	31	14	18	16	14	*	*	9	9	*8	*5
277.00	*6	*	*	*	*	*	*	*	*	*	*
277.02	23	11	12	12	10	*	—	*7	*7	*	*
277.4	*6	*	*	*	*	*	*	*	*	*	*
277.8	*6	*	*	*	*	*	—	*	*	*	*
277.89	*6	*	*	*	*	*	—	*	*	*	*
277.9	*5	*	*	*	*	*	*	*	*	*	*
278	1,229	409	820	20	407	539	264	249	291	451	238
278.0	1,207	404	803	20	400	528	260	242	285	445	235
278.00	734	270	464	16	216	315	187	144	180	260	150
278.01	473	134	338	*	184	213	73	99	105	185	85
278.1	14	*	12	—	*	*7	*	*	*	*5	*
278.8	*7	*	*	—	*	*	*	*	*	*	*
279	25	10	14	*8	*5	*6	*6	*7	*	10	*
279.0	10	*	*6	*	*	*	*	*	*	*5	*
280-289	5,051	1,962	3,089	187	1,050	1,225	2,590	1,045	1,216	1,865	925
280	792	277	514	19	147	182	444	160	200	298	132
280.0	439	164	275	*	78	106	251	84	111	175	69
280.8	*7	*	*5	*	*	*	*	*	*	*	*
280.9	342	110	232	14	66	75	188	74	87	119	62
281	92	33	59	*	*8	15	68	15	31	28	18
281.0	28	*8	21	*	*	*	23	*6	10	*8	*5
281.1	*9	*	*6	—	*	*	*7	*	*	*	*
281.9	53	22	31	*	*6	9	37	*7	18	18	11
282	155	55	101	28	97	22	*7	52	25	64	14
282.4	24	*8	17	*	12	*5	*	10	*	*6	*5
282.49	19	*6	13	*	9	*	*	*8	*	*	*
282.5	27	*	23	*	20	*	*	*7	*	14	*
282.6	97	41	56	21	65	11	*	32	19	42	*
282.60	14	*	10	*	*7	*	*	*	*	*7	*
282.62	72	31	42	13	52	*7	*	25	14	31	*
283	41	18	24	*	*8	9	21	*5	9	21	*6
283.0	15	*7	*9	*	*	*	10	*	*6	*	*
283.1	*5	*	*	*	*	*	*	*	*	*	*
283.9	19	*7	12	*	*	*	10	*	*	12	*
284	205	104	100	14	30	67	93	52	43	75	34
284.8	198	102	96	12	29	66	90	50	42	72	34
284.9	*6	*	*	*	*	*	*	*	*	*	*
285	2,579	918	1,661	40	526	582	1,431	518	635	938	488
285.1	696	249	447	*	107	169	416	127	226	202	141
285.2	289	127	162	*	37	74	176	68	60	108	53
285.21	77	36	41	*	12	24	40	16	15	31	14
285.22	51	21	31	*	*	17	29	11	10	24	*7
285.29	160	70	91	*	20	32	107	40	35	53	32
285.8	54	23	31	*	12	14	27	*	*7	28	15
285.9	1,537	518	1,019	34	370	324	808	319	341	599	278
286	202	95	107	10	29	60	103	44	42	82	34
286.0	*6	*	*	*	*	*	*	*	*	*	*
286.3	17	*7	11	*	*	*7	*6	*	*	*6	*
286.4	*8	*	*6	*	*	*	*	*	*	*	*
286.5	12	*	*8	—	*	*	11	*	*	*7	*
286.6	23	9	14	*	*	*6	12	*5	*5	*7	*6
286.7	36	17	19	*	*	12	20	11	*7	11	*8
286.9	93	47	46	*	9	30	51	16	19	43	14
287	494	253	240	22	91	146	234	103	114	178	99
287.3	45	17	27	*8	11	*8	18	13	*8	14	*9
287.4	76	42	34	*	18	29	28	17	19	20	19

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
287.5	361	188	173	9	59	107	185	70	86	138	67
288	352	157	195	37	71	100	145	64	84	135	70
288.0	149	66	83	18	25	47	60	36	34	52	28
288.3	*9	*	*	*	*	*	*6	*	*	*	*
288.8	182	80	103	15	42	50	76	24	47	75	36
288.9	10	*6	*	*	*	*	*	*	*	*5	*
289	139	51	88	12	41	40	45	33	30	47	29
289.0	9	*6	*	—	*	*	*	*	*	*	*
289.3	14	*5	*9	*	*	*	*	*	*	*	*
289.4	*6	*	*	*	*	*	*	*	*	*	*
289.5	*7	*	*	*	*	*	*	*	*	*	*
289.59	*5	*	*	*	*	*	*	*	*	*	*
289.8	73	22	51	*	28	22	23	18	16	25	14
289.81	41	11	30	*	19	15	*7	10	9	17	*6
289.89	28	10	19	*	*8	*5	14	*8	*7	*7	*7
289.9	24	10	14	*	*5	*	12	*6	*6	*7	*
290-319	12,158	5,754	6,404	379	5,076	3,732	2,971	3,139	2,958	4,124	1,937
290	211	78	133	*	*	10	200	68	45	69	30
290.0	87	26	61	*	*	*	86	34	18	25	10
290.2	16	*5	10	—	—	*	15	*	*6	*	*
290.20	*9	*	*	—	—	*	*9	*	*	*	*
290.21	*7	*	*5	—	—	*	*7	*	*	*	*
290.3	17	*6	10	—	—	*	17	*6	*	*	*
290.4	86	38	48	—	*	*8	78	23	16	35	13
290.40	60	27	34	—	*	*	55	13	11	25	11
290.41	12	*6	*6	—	—	*	10	*6	*	*	*
290.42	*8	*	*	—	—	*	*6	*	*	*	*
290.43	*6	*	*	—	—	*	*6	*	*	*	*
291	316	229	88	*	129	148	39	79	101	74	63
291.0	51	39	12	—	15	26	10	11	9	21	10
291.2	16	14	*	—	*	9	*6	*	*6	*	*
291.3	*7	*6	*	—	*	*	*	*	*	*	*
291.8	235	165	70	*	108	106	21	63	83	44	46
291.81	216	151	64	*	98	97	20	60	70	42	43
291.89	20	13	*6	—	10	*8	*	*	13	*	*
292	238	122	115	*	121	59	54	79	59	56	44
292.0	109	61	48	*	74	29	*	47	25	18	18
292.1	16	9	*6	*	*	*	*7	*	*	*5	*
292.12	14	*8	*6	*	*	*	*6	*	*	*	*
292.8	101	44	57	*	38	22	41	27	29	29	16
292.81	61	23	38	*	*7	15	39	11	22	18	10
292.84	35	18	17	*	29	*5	*	14	*5	11	*
292.9	12	*8	*	*	*6	*	*	*	*	*	*6
293	113	57	56	*	15	21	77	28	34	38	14
293.0	71	36	34	*	*	12	54	16	21	25	*9
293.8	29	16	13	*	11	*8	10	10	*9	*8	*
293.83	21	13	*8	*	*9	*7	*5	*8	*7	*	*
293.9	12	*	*7	*	*	*	11	*	*	*	*
294	888	326	562	*	10	41	835	231	210	315	132
294.1	344	131	214	*	*	13	329	97	85	110	52
294.10	278	98	180	*	*	*7	270	84	64	86	44
294.11	66	32	34	*	*	*7	58	13	21	23	*8
294.8	513	180	333	*	*	17	493	127	119	195	72
294.9	26	14	12	*	*5	10	11	*6	*5	*9	*6
295	568	296	272	*	245	223	97	162	98	229	78
295.3	163	97	66	*	74	63	26	43	29	65	26
295.30	76	48	28	—	31	30	15	18	18	29	10
295.32	38	19	18	—	16	14	*8	9	*	16	*9
295.34	47	28	19	*	27	16	*	15	*6	20	*6
295.6	30	18	12	*	*7	15	*9	9	*5	*9	*7

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
295.60	*8	*	*	—	*	*	*	*	*	*	*
295.62	16	10	*6	*	*	*7	*6	*6	*	*	*5
295.64	*6	*	*	—	*	*	*	*	*	*	*
295.7	224	105	119	*	114	86	22	66	42	93	22
295.70	176	82	94	*	88	72	14	47	36	74	20
295.72	10	*	*5	—	*	*	*	*	*	*	*
295.74	36	17	18	*	20	11	*	16	*	14	*
295.9	138	70	68	*	42	57	38	39	21	58	19
295.90	123	62	61	*	33	54	36	32	20	54	16
295.92	*9	*	*	—	*5	*	*	*	*	*	*
295.94	*6	*	*	*	*	*	*	*	*	*	*
296	1,352	543	809	82	719	391	161	344	325	540	143
296.2	299	126	173	21	153	74	52	55	72	147	25
296.20	173	80	94	*8	90	44	30	34	51	72	15
296.22	21	*7	15	*	12	*	*	*	*	16	*
296.23	64	24	40	*5	33	15	11	11	11	38	*
296.24	37	15	22	*	17	*8	*7	*8	*6	20	*
296.3	354	132	222	*8	190	109	48	105	89	130	30
296.30	97	35	62	*	50	31	14	31	33	29	*
296.32	27	11	16	*	14	10	*	*5	*7	10	*
296.33	159	60	99	*	93	46	15	47	34	67	11
296.34	63	24	39	*	30	18	14	20	12	23	*7
296.4	68	31	37	*	36	23	*8	20	14	23	11
296.40	32	14	18	*	16	11	*	10	*7	10	*
296.43	*8	*	*	*	*	*	*	*	*	*	*
296.44	27	12	15	*	15	10	*	*7	*6	*9	*6
296.5	112	41	72	*	68	36	*6	36	29	37	10
296.50	65	25	41	*	41	20	*	17	21	21	*7
296.53	21	*9	13	*	13	*6	*	10	*	*6	*
296.54	21	*6	15	*	11	*8	*	*8	*	*9	*
296.6	99	48	52	17	59	20	*	23	15	56	*7
296.60	45	22	23	*	30	10	*	*7	9	26	*
296.63	19	9	10	*	12	*	*	*6	*	10	*
296.64	34	16	17	*8	16	*8	*	9	*	19	*
296.7	246	92	155	16	117	85	28	55	63	92	36
296.8	104	39	65	*8	56	30	10	20	30	40	14
296.80	73	28	45	*6	35	24	*8	13	22	28	10
296.89	29	10	19	*	19	*6	*	*7	*8	10	*
296.9	64	33	32	11	38	12	*	29	12	14	10
296.90	59	29	30	10	34	11	*	27	10	13	9
296.99	*6	*	*	*	*	*	*	*	*	*	*
297	31	13	19	*	*6	*6	19	12	*6	10	*
297.1	*7	*	*	*	*	*	*	*	*	*	*
297.9	22	9	13	—	*	*	15	11	*	*6	*
298	205	93	112	*	74	47	80	45	46	71	42
298.9	198	91	107	*	71	46	76	44	44	69	41
299	26	20	*6	11	12	*	*	10	*5	*7	*
299.0	18	13	*	*8	*8	*	*	*5	*	*	*
299.00	18	13	*	*8	*8	*	*	*5	*	*	*
299.8	*6	*6	*	*	*	*	*	*	*	*	—
299.80	*6	*6	*	*	*	*	*	*	*	*	—
300	1,207	357	850	24	396	403	384	287	334	418	168
300.0	659	196	463	10	183	226	240	161	171	233	94
300.00	539	161	378	*8	131	183	218	133	138	191	77
300.01	76	19	57	*	34	29	12	18	19	26	13
300.02	43	15	28	*	17	15	10	10	13	16	*
300.1	30	*5	25	*	18	*9	*	*6	*8	11	*
300.11	13	*	9	*	*7	*	*	*	*	*	*
300.14	10	*	9	*	*6	*	—	*	*	*5	*
300.2	20	*6	14	—	13	*	*	*7	*	*6	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
300.21	12	*	*8	–	*8	*	*	*	*	*	*
300.3	32	15	18	*	21	*6	*	15	*7	*	*
300.4	370	95	276	*8	98	134	131	87	105	129	49
300.8	*6	*	*	*	*	*	*	*	*	*	*
300.9	90	40	50	*	60	20	*6	9	37	32	11
301	330	138	192	*8	222	87	12	130	77	85	38
301.2	*7	*	*	*	*	*	*	*	*	*	*
301.20	*5	*	*	*	*	*	*	*	*	*	*
301.5	13	*	*8	*	*7	*	*	*	*	10	*
301.50	12	*	*7	*	*7	*	*	*	*	*9	*
301.6	33	15	18	*	18	9	*	*7	*5	16	*
301.7	30	24	*6	*	23	*7	*	*7	14	*6	*
301.8	126	29	98	*	94	29	*	40	43	28	14
301.81	*7	*	*	–	*	*	*	*	*	*	*
301.83	97	15	82	*	75	20	*	30	34	22	10
301.89	18	*7	11	*	13	*	*	*9	*	*	*
301.9	108	55	53	*	70	32	*	70	*8	18	12
302	*8	*7	*	*	*	*	*	*	*	*	*
303	752	520	232	*	320	336	95	216	194	204	138
303.0	145	96	49	*	68	66	11	43	45	26	31
303.00	44	28	15	*	19	21	*	*	14	10	15
303.01	96	64	32	–	46	43	*6	37	29	15	15
303.02	*6	*	*	–	*	*	*	*	*	*	*
303.9	607	424	183	*	252	270	83	172	149	179	107
303.90	266	190	75	*	104	120	40	52	60	106	47
303.91	275	194	81	*	119	121	34	101	66	59	48
303.92	16	10	*6	*	9	*5	*	*6	*5	*	*
303.93	51	30	21	*	19	24	*8	14	17	11	*9
304	584	336	248	*	403	164	13	226	104	153	100
304.0	210	122	89	*	125	78	*6	89	34	45	43
304.00	61	31	30	*	37	21	*	10	14	21	16
304.01	139	84	54	*	82	54	*	74	19	23	23
304.03	*8	*	*	–	*	*	*	*	*	*	*
304.1	35	17	18	*	20	12	*	13	*	*8	10
304.10	12	*6	*6	*	*6	*	*	*	*	*	*
304.11	21	9	12	–	13	*7	*	*8	*	*	*7
304.2	122	76	46	*	92	30	*	61	17	36	*9
304.20	46	29	18	*	35	11	*	10	10	21	*
304.21	69	43	25	*	51	17	*	46	*	14	*
304.23	*6	*	*	–	*	*	–	*	*	*	*
304.3	55	35	20	*	49	*5	*	13	26	10	*6
304.30	16	11	*6	*	14	*	*	*	*7	*	*
304.31	33	20	12	*	30	*	–	10	15	*	*
304.4	22	13	9	–	19	*	–	*	*	*	14
304.40	12	*7	*	–	10	*	–	*	*	*	*8
304.41	*8	*	*	–	*7	*	–	*	*	*	*
304.6	11	*	*7	–	*6	*	*	*	*	*	*
304.60	*5	*	*	–	*	*	*	*	*	*	*
304.7	40	20	19	–	29	10	*	19	*6	9	*5
304.70	14	*7	*7	–	11	*	*	*	*	*5	*
304.71	24	13	11	–	17	*7	*	13	*	*	*
304.8	38	22	17	*	31	*6	*	13	*	19	*
304.80	22	12	10	*	18	*	*	*5	*	15	*
304.81	9	*6	*	–	*8	*	*	*	*	*	*
304.9	49	26	23	*	31	14	*	14	*5	19	10
304.90	35	16	18	*	21	11	*	*6	*	17	*8
304.91	*8	*	*	*	*6	*	*	*	*	*	*
305	3,205	1,768	1,437	20	1,582	1,209	395	625	770	1,199	610
305.0	535	373	161	*	259	200	72	134	126	182	93
305.00	370	263	107	*	186	132	48	68	101	137	65

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
305.01	99	69	29	*	42	42	14	35	19	28	17
305.02	28	17	11	*	16	10	*	18	*	*	*
305.03	37	23	14	—	15	15	*7	13	*	14	*8
305.1	1,963	994	969	*	823	823	313	327	511	769	355
305.2	159	101	57	*7	125	25	*	41	35	55	28
305.20	111	68	43	*6	87	18	*	18	30	43	19
305.21	32	21	10	*	26	*	*	15	*	*8	*
305.22	10	*8	*	*	*8	*	—	*6	*	*	*
305.23	*6	*	*	—	*	*	*	*	*	*	*
305.4	16	*8	*9	*	10	*6	*	*6	*	*6	*
305.40	12	*6	*6	*	*7	*	*	*	*	*	*
305.5	84	41	43	*	51	32	*	19	17	19	29
305.50	56	26	30	*	35	19	*	9	15	13	18
305.51	22	12	10	—	14	*8	*	*8	*	*	*8
305.53	*6	*	*	—	*	*	*	*	*	*	*
305.6	236	137	99	*	158	73	*	57	48	89	42
305.60	175	98	77	*	116	55	*	29	43	75	29
305.61	39	25	14	*	28	11	*	18	*	*9	9
305.62	10	*7	*	*	*8	*	—	*7	*	*	*
305.63	11	*7	*	*	*6	*	*	*	*	*	*
305.7	52	25	27	*	43	*8	—	*	*	11	35
305.70	41	19	22	*	35	*5	—	*	*	10	26
305.71	*6	*	*	*	*	*	—	*	*	*	*5
305.9	155	86	70	*	109	41	*	39	25	65	25
305.90	115	62	53	*	82	29	*	19	22	52	22
305.91	16	9	*6	*	11	*	*	*8	*	*	*
305.93	21	12	*9	*	13	*8	*	10	*	*8	*
306	*7	*	*	*	*	*	*	*	*	*	*
307	71	23	48	*5	37	13	16	20	21	17	13
307.1	*8	*	*7	*	*6	*	*	*	*	*	*
307.2	*6	*5	*	*	*	*	*	*	*	*	*
307.5	20	*	18	*	16	*	*	*6	*6	*	*
307.50	11	*	10	*	*9	*	*	*6	*	*	*
307.51	*6	*	*5	*	*5	*	—	*	*	*	*
307.8	15	*	11	*	*7	*5	*	*	*	*	*
307.81	13	*	10	*	*6	*	*	*	*	*	*
307.9	17	*9	*8	*	*	*	11	*	*6	*5	*
308	17	*6	10	*	*8	*5	*	*	*	*7	*5
308.3	*7	*	*	*	*	*	*	*	*	*	*
308.9	*6	*	*	*	*	*	*	*	*	*	*
309	288	112	176	22	171	75	19	101	81	65	41
309.0	55	27	29	*	34	12	*8	19	14	14	*8
309.2	37	15	21	*	21	*8	*	9	16	*6	*5
309.28	32	13	18	*	19	*7	*	*8	15	*	*5
309.4	22	12	10	*	16	*	*	11	*	*	*5
309.8	151	47	105	11	86	51	*	53	43	35	19
309.81	149	45	104	11	84	50	*	53	43	34	18
309.9	18	9	*9	*	10	*	*	*6	*	*6	*
310	41	21	20	*	*8	10	22	11	*6	13	11
310.1	9	*7	*	*	*	*	*	*	*	*	*
310.2	10	*6	*	*	*	*	*	*	*	*	*
310.9	19	*7	12	—	*	*	14	*6	*	*9	*
311	1,261	426	834	19	413	407	422	333	334	394	200
312	67	44	23	20	35	*5	*8	27	14	21	*
312.3	42	27	15	9	27	*	*	18	10	12	*
312.30	28	17	11	*6	18	*	*	13	*5	*7	*
312.34	10	*8	*	*	*6	*	*	*	*	*	*
312.8	*5	*	*	*	*	*	*	*	*	*	*
312.9	19	12	*6	*7	*6	*	*	*6	*	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
313	40	24	16	27	14	*	—	*9	*8	18	*5
313.8	38	23	15	26	12	—	-	*7	*8	18	*5
313.81	34	21	13	23	11	—	-	*7	*7	16	*
314	131	84	47	65	56	11	*	33	32	54	11
314.0	131	84	46	64	55	11	*	33	32	54	11
314.00	26	13	13	*5	15	*6	*	*8	*6	9	*
314.01	104	71	33	59	40	*	*	26	27	45	*7
315	50	31	19	33	12	*	*	11	14	10	15
315.2	10	*6	*	*	*5	*	—	*	*6	*	*
315.3	*8	*	*	*7	*	*	—	*	*	*	*
315.39	*6	*	*	*6	*	—	-	*	*	*	*
315.8	14	10	*	12	*	*	—	*	*	*	*
315.9	15	*9	*7	*7	*6	*	*	*	*	*	*
317	38	17	22	*	20	12	*	11	*6	18	*
318	29	16	13	*	12	10	*	12	*8	*6	*
318.0	*8	*	*	*	*	*	*	*	*	*	*
318.1	12	*7	*	*	*	*	*	*	*	*	*
318.2	*9	*	*	*	*	*	*	*5	*	*	*
319	81	42	39	12	29	28	12	23	18	26	14
320-389	3,873	1,705	2,168	368	649	926	1,931	879	946	1,312	736
320	14	*9	*6	*5	*	*	*	*	*	*7	*
322	12	*5	*6	*	*	*	*	*	*	*	*
322.9	10	*	*5	*	*	*	*	*	*	*	*
323	12	*	*8	*	*	*	*	*	*	*	*
323.9	*6	*	*	*	*	*	*	*	*	*	*
324	16	11	*	*	*	*6	*	*	*	*6	*5
324.1	12	9	*	*	*	*5	*	*	*	*	*
331	549	214	334	20	17	32	479	140	134	193	83
331.0	418	150	268	*	*	*9	408	109	105	149	55
331.3	16	*8	*8	*	*	*	10	*	*	*5	*
331.4	67	33	34	17	13	16	20	16	13	23	15
331.8	33	16	17	*	*	*	30	*8	*8	12	*6
331.82	31	15	16	*	—	*	29	*7	*7	11	*6
331.9	11	*6	*	*	*	*	*9	*	*	*	*
332	273	157	116	*	*	27	245	69	74	86	44
332.0	271	156	115	*	*	26	244	68	73	86	44
333	99	33	66	*	12	25	59	22	27	31	19
333.0	*6	*	*	*	*	*	*	*	*	*	*
333.1	27	*8	19	*	*	*5	20	*6	*7	*9	*
333.2	*6	*	*	*	*	*	*	*	*	*	*
333.8	*7	*	*	—	*	*	*	*	*	*	*
333.82	*7	*	*	—	*	*	*	*	*	*	*
333.9	46	13	33	*	*7	12	25	*9	13	14	10
333.99	41	10	31	*	*5	11	23	*8	12	12	*9
334	10	*	*	*	*	*	*	*	*	*	*
335	12	*6	*6	*	*	*	*5	*	*	*	*
335.2	10	*6	*	—	*	*	*	*	*	*	*
335.20	*8	*	*	—	*	*	*	*	*	*	*
336	24	16	*9	*	*	*8	10	*5	*	11	*
336.0	*7	*	*	*	*	*	*	*	*	*	*
336.3	*6	*	*	*	*	*	*	*	*	*	*
336.9	*6	*	*	*	*	*	*	*	*	*	*
337	75	33	42	*	21	33	21	19	12	32	12
337.1	48	24	24	*	13	21	14	11	*7	23	*7
337.2	16	*	14	*	*	9	*	*	*	*	*
337.20	*6	*	*5	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
337.9	*6	*	*	*	*	*	*	*	*	*	*
340	118	25	93	*	25	60	32	34	31	29	24
341	*6	*	*	*	*	*	*	*	*	*	*
342	169	72	97	*	15	44	106	31	37	61	40
342.0	*7	*	*	*	*	*	*6	*	*	*	*
342.00	*5	*	*	*	—	*	*	*	*	*	*
342.8	14	*6	*8	*	*	*	*7	*	*	*7	*
342.80	*9	*	*5	*	*	*	*	*	*	*	*
342.9	144	62	82	*	12	37	93	25	32	52	35
342.90	123	53	70	*	10	29	81	22	27	44	30
342.91	11	*6	*	*	*	*	*7	*	*	*	*
342.92	10	*	*7	—	*	*	*5	*	*	*	*
343	90	47	43	33	32	16	*9	21	28	25	16
343.2	12	*7	*5	*	*	*	*	*	*	*	*
343.9	68	33	34	24	24	12	*8	15	22	20	11
344	146	99	47	*	50	57	36	22	32	62	30
344.0	58	45	13	*	20	22	14	*8	12	27	11
344.00	47	36	11	*	15	17	13	*6	9	25	*7
344.1	52	36	17	*	22	20	10	10	10	19	13
344.6	10	*	*5	*	*	*	*	*	*	*	*
344.60	*8	*	*	*	*	*	*	*	*	*	*
344.8	*6	*	*	*	*	*	*	*	*	*	*
344.89	*6	*	*	*	*	*	*	*	*	*	*
344.9	11	*6	*6	*	*	*6	*	*	*	*	*
345	107	50	57	32	35	26	14	30	21	33	24
345.1	15	*7	*8	*	*	*	*	*6	*	*	*
345.10	13	*6	*7	*	*	*	*	*5	*	*	*
345.3	24	12	12	*9	*6	*7	*	*	*5	*8	*7
345.4	*8	*	*	*	*	*	*	*	*	*	*
345.5	*9	*6	*	*	*	*	*	*	*	*	*
345.50	*6	*	*	*	*	*	*	*	*	*	*
345.9	38	16	22	*9	14	10	*5	9	10	13	*6
345.90	34	14	19	*7	12	9	*5	*8	10	11	*6
346	245	45	201	*7	124	87	27	48	67	89	42
346.2	10	*6	*	*	*	*	*	*	*	*	*
346.20	*8	*	*	*	*	*	*	*	*	*	*
346.8	17	*	14	*	10	*	*	*	*	*7	*
346.80	13	*	10	*	*7	*	*	*	*	*	*
346.9	212	34	178	*5	108	75	24	40	59	76	36
346.90	196	32	164	*	98	69	23	39	54	68	35
346.91	16	*	14	*	9	*6	*	*	*5	*8	*
347	*7	*	*	*	*	*	*	*	*	*	*
348	215	102	113	23	42	56	94	45	45	75	49
348.0	*8	*6	*	*	*	*	*	*	*	*	*
348.1	59	31	29	*5	*8	20	26	9	15	18	16
348.2	13	*	11	*	11	*	*	*	*	*	*
348.3	83	37	46	*	*8	18	53	18	15	36	14
348.30	40	17	23	*	*5	*9	24	*8	*7	18	*7
348.31	25	12	13	*	*	*	20	*9	*	*8	*
348.39	19	*8	11	*	*	*6	9	*	*	10	*
348.4	13	*6	*7	*	*	*	*	*	*	*	*
348.5	12	*8	*	*	*	*	*	*	*	*	*
348.8	19	10	10	*	*6	*	*	*	*	*6	*
348.9	*7	*	*	*	*	*	*	*	*	*	*
349	63	19	43	*	30	12	18	12	16	23	12
349.0	28	*	24	*	24	*	*	*	*8	10	*6
349.2	*6	*	*	*	*	*	*	*	*	*	*
349.8	27	12	15	*	*5	*5	15	*6	*7	9	*
349.82	21	*9	13	*	*	*	14	*6	*6	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
350	22	*6	16	*	*	9	11	*5	*8	*5	*
350.1	21	*6	15	—	*	*8	11	*5	*8	*5	*
351	28	14	14	*	*6	10	11	*7	*6	11	*5
351.0	26	14	13	*	*6	*9	11	*6	*5	10	*
353	11	*5	*6	*	*5	*	*	*	*	*	*
353.0	*6	*	*	*	*	*	*	*	*	*	*
354	36	17	20	*	11	14	11	*8	*8	13	*7
354.0	22	9	13	*	*8	*7	*7	*	*5	*8	*
354.2	*5	*	*	*	*	*	*	*	*	*	*
355	62	26	36	*	*9	21	31	13	15	18	16
355.3	*6	*	*	*	*	*	*	*	*	*	*
355.8	26	11	15	*	*	*7	16	*	*7	*8	*7
355.9	21	11	10	*	*	*7	11	*	*	*6	*6
356	94	34	61	*	*7	23	65	12	24	44	14
356.9	88	31	57	*	*6	21	61	11	23	41	13
357	242	124	119	*	26	97	118	58	61	81	43
357.0	12	*8	*	*	*	*	*	*	*	*5	*
357.2	214	105	109	—	20	86	108	52	53	72	37
357.5	*6	*	*	—	*	*	*	*	*	*	*
357.8	*6	*	*	*	*	*	*	*	*	*	*
357.81	*5	*	*	*	*	*	*	*	*	*	*
358	30	13	17	*	*	*8	18	*6	*7	*9	*8
358.0	25	11	15	*	*	*7	16	*5	*6	*7	*7
358.00	21	9	12	*	*	*	14	*	*5	*7	*
359	28	15	13	*	*8	*9	9	*9	*	10	*5
359.1	12	*7	*	*	*	*	*	*	*	*	*
359.4	*7	*	*	—	*	*	*	*	*	*	*
362	156	64	92	*6	14	47	90	38	47	43	27
362.0	81	36	45	*	11	38	32	20	22	25	14
362.01	78	34	43	*	11	36	31	19	22	24	14
362.2	*5	*	*	*	—	—	*	*	*	*	*
362.3	*8	*	*	—	*	*	*6	*	*	*	*
362.5	51	15	36	—	*	*	49	12	19	12	*8
362.50	50	14	36	—	*	*	48	12	19	12	*7
364	*7	*	*	*	*	*	*	*	*	*	*
365	169	70	99	*	*6	31	132	51	48	46	24
365.9	163	67	96	*	*6	30	127	49	46	44	24
366	30	11	19	*	*	*6	22	*8	*6	12	*
366.9	24	9	14	—	*	*	19	*7	*6	*7	*
368	69	31	38	*6	15	24	26	11	18	25	16
368.1	10	*	*6	*	*	*	*	*	*	*	*
368.16	*6	*	*	*	*	*	*	*	*	*	*
368.2	22	9	13	*	*	*8	*7	*	*6	*7	*6
368.4	22	11	10	*	*	*7	9	*	*	*8	*7
368.46	14	*9	*	*	*	*	*8	*	*	*6	*
368.8	11	*	*6	*	*	*	*	*	*	*	*
369	85	43	42	*	14	23	44	16	20	29	19
369.0	27	14	14	*	*	*6	16	*6	*6	10	*
369.00	26	12	13	*	*	*6	15	*6	*5	9	*
369.4	23	9	13	*	*	*6	13	*6	*6	*7	*
369.6	22	13	*8	*	*	*8	*8	*	*	*9	*6
369.60	21	13	*8	*	*	*8	*8	*	*	*9	*6
369.9	*8	*	*	*	*	*	*	*	*	*	*
372	47	23	25	15	10	9	13	11	10	14	12
372.0	*9	*	*6	*	*	*	*	*	*	*	*
372.00	*8	*	*5	*	*	*	*	*	*	*	*
372.3	28	13	15	12	*	*	9	*5	*6	*8	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
372.30	26	12	14	11	*	*	*9	*5	*6	*7	*7
372.7	*8	*	*	*	*	*	*	*	*	*	*
372.72	*7	*	*	*	*	*	*	*	*	*	*
373	*9	*	*	*	*	*	*	*	*	*	*
373.1	*5	*	*	*	*	*	*	*	*	*	*
374	11	*	*7	*	*	*6	*	*	*	*	*
374.3	*8	*	*5	*	*	*	*	*	*	*	*
374.30	*7	*	*	*	*	*	*	*	*	*	*
376	18	9	*8	*6	*6	*	*	*	*	*6	*
376.0	13	*7	*6	*5	*	*	*	*	*	*	*
376.01	12	*7	*6	*5	*	*	*	*	*	*	*
377	13	*5	*7	*	*5	*	*	*	*	*6	*
377.3	*5	*	*	*	*	*	*	*	*	*	*
378	21	12	9	*	*6	*6	*	*	*6	*8	*
378.5	13	*8	*	*	*	*	*	*	*	*	*
378.51	*6	*	*	*	*	*	*	*	*	*	*
378.54	*5	*	*	*	*	*	*	*	*	*	*
379	17	10	*7	*	*5	*	*	*	*	*6	*
379.5	*6	*	*	*	*	*	*	*	*	*	*
380	32	17	15	*7	*8	*6	12	*8	*8	11	*
380.1	18	11	*7	*	*6	*	*5	*	*5	*6	*
380.10	15	*8	*7	*	*	*	*	*	*	*5	*
380.4	12	*5	*6	*	*	*	*6	*	*	*	*
381	19	10	*8	14	*	*	*	*6	*	*	*
381.4	*7	*	*	*5	*	*	*	*	*	*	*
382	136	81	56	117	*9	*6	*5	20	30	53	34
382.9	131	78	53	112	*8	*5	*5	19	27	52	34
383	*7	*	*	*	*	*	*	*	*	*	*
386	75	25	51	*	*6	21	48	24	22	18	12
386.0	17	*	12	—	*	*5	9	*	*	*	*
386.00	16	*	12	—	*	*5	9	*	*	*	*
386.1	32	11	20	*	*	10	20	11	10	*7	*
386.11	22	*8	14	*	*	*7	14	*6	*8	*	*
386.12	*5	*	*	*	*	*	*	*	*	*	*
386.3	23	*8	15	—	*	*	16	*8	*5	*5	*
386.30	20	*8	13	—	*	*	15	*7	*	*	*
388	17	10	*7	*	*6	*	*5	*	*5	*	*
388.3	*7	*	*	*	*	*	*	*	*	*	*
388.30	*6	*	*	*	*	*	*	*	*	*	*
389	80	39	41	*7	12	*9	53	18	22	25	15
389.1	*6	*	*	*	*	*	*	*	*	*	*
389.9	64	31	33	*	*8	*8	45	14	17	20	13
390-459	34,384	16,462	17,922	158	2,060	8,929	23,236	7,917	8,420	12,365	5,682
394	27	*6	21	*	*	*8	16	*6	*8	*8	*6
394.0	12	*	10	*	*	*	*7	*	*	*	*
394.1	*7	*	*5	*	*	*	*5	*	*	*	*
394.2	*5	*	*	*	*	*	*	*	*	*	*
396	178	65	113	*	*	19	155	48	44	58	27
396.0	*7	*	*6	*	*	*	*5	*	*	*	*
396.2	49	18	31	—	*	*	45	15	11	16	*8
396.3	89	35	54	*	*	10	77	21	22	32	14
396.8	20	*6	14	*	*	*	18	*6	*7	*	*
396.9	10	*	*7	—	—	*	*9	*	*	*	*
397	180	61	119	*	9	26	142	46	41	58	36
397.0	177	59	117	*	*9	26	139	45	40	57	35

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
398	86	25	61	—	*	11	71	25	17	27	17
398.9	86	25	61	—	*	11	71	25	17	27	17
398.90	*7	*	*5	—	*	*	*	*	*	*	*
398.91	77	23	55	—	*	*9	66	24	15	25	15
401	8,936	3,769	5,166	21	711	2,805	5,399	2,046	2,131	3,349	1,411
401.0	83	32	50	*	10	25	47	14	13	38	17
401.1	115	48	66	*	9	37	69	84	*	25	*
401.9	8,739	3,689	5,050	20	692	2,743	5,283	1,947	2,115	3,285	1,391
402	328	128	200	*	12	74	240	74	78	129	47
402.0	15	*7	*9	*	*	*5	*7	*	*	11	*
402.01	12	*6	*7	*	*	*	*6	*	*	10	*
402.9	308	120	188	*	10	68	230	70	76	117	46
402.90	162	62	101	*	*6	36	120	40	39	60	23
402.91	146	59	87	—	*	32	110	30	37	57	23
403	1,112	562	550	*7	108	327	670	220	236	444	212
403.0	26	*9	17	*	*8	10	*7	*	*	14	*5
403.01	24	*8	17	*	*7	9	*7	*	*	13	*5
403.1	*9	*	*	—	*	*	*5	*6	*	*	*
403.11	*9	*	*	—	*	*	*	*6	*	*	*
403.9	1,078	549	528	*6	99	314	658	211	232	428	206
403.90	58	31	27	*	*	16	39	10	14	24	10
403.91	1,020	518	502	*	97	299	619	201	219	404	196
404	90	41	49	*	*	27	59	17	18	39	15
404.0	*8	*	*	—	*	*	*	*	*	*	*
404.9	81	37	44	*	*	24	54	15	17	34	14
404.91	*6	*	*	—	*	*	*	*	*	*	*
404.92	23	10	13	—	*	10	11	*	*	13	*
404.93	48	22	25	*	*	11	35	11	13	16	*7
405	*7	*	*	*	*	*	*	*	*	*	*
405.9	*7	*	*	*	*	*	*	*	*	*	*
410	902	507	395	*	48	272	580	215	212	310	165
410.0	19	10	*8	—	*	*5	11	*	*	*7	*
410.01	18	10	*8	—	*	*5	11	*	*	*7	*
410.1	91	52	39	—	*8	38	46	19	22	32	19
410.11	82	46	35	—	*7	34	41	17	18	27	18
410.12	*8	*	*	—	*	*	*	*	*	*	*
410.2	16	10	*6	—	*	*	10	*	*	*6	*
410.21	13	*8	*5	—	*	*	*9	*	*	*5	*
410.3	9	*7	*	—	*	*	*	*	*	*	*
410.31	*9	*6	*	—	*	*	*	*	*	*	*
410.4	103	69	35	—	*7	43	53	21	23	38	22
410.41	95	63	32	—	*7	39	49	19	22	35	20
410.42	*7	*6	*	—	*	*	*	*	*	*	*
410.5	11	*6	*5	—	*	*	*6	*	*	*	*
410.51	11	*6	*	—	*	*	*6	*	*	*	*
410.7	503	275	228	*	22	141	340	141	107	170	85
410.70	9	*	*	—	*	*	*7	*	*	*	*
410.71	466	255	211	*	20	131	315	132	99	156	80
410.72	28	15	13	—	*	*8	18	*6	*8	9	*
410.8	20	11	*8	*	*	*6	13	*	*5	*8	*
410.81	19	11	*8	*	*	*6	13	*	*5	*7	*
410.9	127	65	61	*	*5	26	95	20	41	45	21
410.90	*6	*	*	*	*	*	*	*	*	*	*
410.91	107	56	51	*	*	20	83	16	37	37	17
410.92	14	*8	*6	—	*	*	10	*	*	*	*
411	670	385	285	*	33	255	381	168	145	244	114
411.0	*7	*	*	—	*	*	*	*	*	*	*
411.1	637	365	271	*	30	244	361	156	138	235	108
411.8	27	16	11	*	*	*7	17	10	*	*7	*5
411.89	23	13	10	*	*	*7	15	10	*	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
412	711	434	278	*	34	220	456	170	189	234	119
413	429	230	199	*	16	135	278	93	117	148	70
413.9	425	229	196	*	15	133	277	93	117	146	69
414	4,772	2,756	2,015	*	143	1,333	3,290	1,159	1,177	1,701	735
414.0	4,457	2,558	1,899	*	133	1,261	3,060	1,079	1,098	1,602	678
414.00	1,244	741	504	*	24	290	930	324	354	442	125
414.01	3,079	1,721	1,358	*	106	932	2,039	729	712	1,110	528
414.02	100	75	25	*	*	29	70	20	27	31	22
414.04	12	*9	*	—	*	*	*8	*	*	*	*
414.05	20	12	*8	—	*	*6	13	*	*	13	*
414.1	17	*8	*8	*	*	*	*8	*	*6	*	*6
414.10	10	*	*7	*	*	*	*6	*	*	*	*
414.8	257	165	91	*	*5	58	193	73	59	78	47
414.9	42	24	17	*	*	10	30	*	14	19	*
415	194	80	114	*	28	57	108	40	51	67	36
415.1	191	78	113	*	28	56	107	39	51	66	35
415.11	16	*7	*9	—	*	*6	*7	*	*	*6	*
415.19	175	70	105	*	25	50	100	36	47	60	32
416	246	98	148	*6	23	69	148	50	60	87	49
416.0	25	*8	18	*	*	*6	14	*6	*	11	*
416.8	155	55	99	*	15	40	96	31	37	53	34
416.9	66	35	31	*	*	23	39	13	19	24	10
417	*5	*	*	*	*	*	*	*	*	*	*
420	27	14	13	*	*	10	13	*6	*7	*8	*6
420.9	25	13	12	*	*	*9	12	*6	*6	*7	*6
420.90	16	*9	*8	*	*	*6	*8	*	*	*5	*
420.91	*7	*	*	*	*	*	*	*	*	*	*
421	27	15	12	*	*8	*7	11	10	*	*6	*7
421.0	26	15	11	*	*8	*6	11	9	*	*5	*7
423	53	27	26	*	10	15	25	19	*9	16	10
423.2	*6	*	*	—	*	*	*	*	*	*	*
423.8	*6	*	*	*	*	*	*	*	*	*	*
423.9	37	15	22	*	*9	12	14	11	*7	12	*8
424	1,069	393	677	*8	95	197	770	280	248	351	191
424.0	594	197	397	*	77	138	375	144	140	212	98
424.1	397	167	230	*	11	44	339	119	88	112	78
424.2	32	11	20	*	*	*6	23	*7	*6	11	*8
424.3	*6	*	*	*	*	*	*	*	*	*	*
424.9	41	16	25	*	*	*7	29	*9	13	14	*
424.90	38	15	23	*	*	*7	28	*8	13	13	*
425	641	346	295	*6	52	197	386	133	135	265	108
425.1	22	*6	15	*	*	*	14	*	*6	*7	*6
425.4	572	309	263	*	42	170	356	119	120	239	94
425.5	18	15	*	—	*	12	*	*5	*	*	*
425.8	18	*9	10	*	*	*7	*9	*	*	11	*
425.9	*8	*	*	*	*	*	*	*	*	*	*
426	427	236	192	*	23	82	317	99	111	123	94
426.0	67	34	32	*	*	12	51	16	15	19	16
426.1	115	63	51	*	*6	19	89	28	34	28	25
426.10	*7	*	*	—	*	*	*6	*	*	*	*
426.11	65	35	29	*	*	11	51	15	23	13	14
426.12	10	*6	*	*	*	*	*8	*	*	*	*
426.13	34	19	15	*	*	*5	25	*8	*8	9	*8
426.2	9	*6	*	—	*	*	*7	*	*	*	*
426.3	93	44	49	*	*	18	73	23	23	27	21
426.4	79	47	32	*	*6	20	53	17	21	25	16
426.5	33	21	12	—	*	*	27	*5	10	10	*7
426.52	*5	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
426.53	18	13	*5	—	*	*	15	*	*7	*	*
426.6	*6	*	*	*	*	*	*	*	*	*	*
426.7	13	*8	*	*	*	*	*	*	*	*5	*
426.9	*8	*5	*	*	*	*	*5	*	*	*	*
427	4,371	2,129	2,242	26	198	735	3,412	1,047	1,125	1,439	760
427.0	93	37	55	*	10	20	62	19	19	36	19
427.1	269	177	92	*	18	77	172	59	68	99	43
427.3	2,936	1,396	1,539	*	63	399	2,472	747	760	934	495
427.31	2,739	1,280	1,459	*	55	350	2,332	701	715	863	460
427.32	197	117	80	*	*7	49	140	45	45	72	34
427.4	50	30	20	*	*	17	30	*7	10	19	14
427.41	49	29	20	*	*	16	29	*7	10	19	14
427.5	98	53	44	*	10	27	58	14	21	36	26
427.6	95	44	51	*	13	21	59	21	24	30	19
427.61	20	*8	12	*	*	*	14	*	*	*6	*6
427.69	73	34	39	*	11	18	44	18	19	24	13
427.8	771	366	405	16	77	161	517	171	207	258	136
427.81	180	77	103	*	*	17	160	47	40	61	31
427.89	591	289	302	16	75	144	356	124	166	197	104
427.9	58	25	33	*	*	12	41	*8	16	26	*8
428	4,047	1,805	2,242	16	116	782	3,133	887	1,096	1,468	596
428.0	3,773	1,685	2,088	14	106	725	2,928	845	1,027	1,351	550
428.1	*8	*	*	—	*	*	*6	*	*	*	*
428.2	64	37	27	*	*	15	45	*5	25	26	*8
428.20	30	18	13	*	*	*7	22	*	10	13	*
428.21	10	*6	*	—	*	*	*6	*	*	*5	*
428.22	12	*7	*5	—	*	*	11	*	*7	*	*
428.23	12	*6	*5	—	*	*	*7	*	*	*	*
428.3	166	63	103	*	*	33	129	31	36	72	27
428.30	109	42	67	*	*	21	85	20	24	50	15
428.31	21	*9	12	—	*	*	15	*	*	*9	*
428.32	18	*6	12	—	*	*	15	*	*	*7	*
428.33	18	*6	12	—	*	*	14	*	*	*7	*
428.4	13	*7	*7	*	*	*	*8	*	*	*8	*
428.43	*6	*	*	*	*	*	*	*	*	*	*
428.9	22	10	13	*	*	*	17	*	*	10	*6
429	196	91	106	*5	16	54	121	49	43	78	26
429.2	64	27	37	—	*	13	51	24	*9	27	*
429.3	74	34	39	*	*9	22	40	13	22	28	11
429.4	*9	*5	*	*	*	*	*6	*	*	*	*
429.7	*6	*	*	*	*	*	*	*	*	*	*
429.8	*9	*6	*	*	*	*	*	*	*	*	*
429.89	*6	*	*	*	*	*	*	*	*	*	*
429.9	32	13	18	*	*	12	16	*5	9	12	*5
430	29	*8	20	*	*7	12	*9	*7	*6	9	*7
431	86	38	49	*	*8	20	58	17	16	34	20
432	38	20	18	*	*	*6	27	*8	*7	14	10
432.1	31	18	13	*	*	*	23	*6	*6	10	*9
432.9	*6	*	*	—	*	*	*	*	*	*	*
433	291	157	134	*	*	61	225	67	71	109	45
433.0	*5	*	*	—	*	*	*	*	*	*	*
433.1	219	117	101	*	*	43	173	51	56	79	33
433.10	198	106	92	—	*	36	159	47	51	72	28
433.11	21	11	10	*	*	*7	13	*	*6	*7	*5
433.2	*6	*	*	—	*	*	*	*	*	*	*
433.3	57	31	26	—	*	12	45	15	11	22	9
433.30	51	28	23	—	*	11	40	13	10	21	*8
433.31	*6	*	*	—	—	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
434	394	179	215	*	19	96	276	84	84	154	73
434.0	14	*7	*7	*	*	*	9	*	*	*	*
434.01	11	*6	*6	*	*	*	*7	*	*	*	*
434.1	48	20	28	*	*	*9	37	12	10	19	*7
434.11	43	18	26	*	*	*8	33	11	*8	17	*7
434.9	332	152	180	*	16	83	231	69	72	131	61
434.90	12	*	*7	*	*	*	*8	*	*	*6	*
434.91	321	148	173	*	16	79	223	67	70	125	58
435	240	101	140	*	*9	56	175	63	59	89	29
435.3	9	*	*6	—	*	*	*8	*	*	*	*
435.8	12	*	*8	—	*	*	*8	*	*	*6	*
435.9	213	92	121	*	*8	50	155	58	51	78	27
436	121	50	71	*	*7	23	90	16	27	53	25
437	157	64	93	*	10	29	116	27	46	59	25
437.0	62	28	34	*	*	*8	53	13	16	26	*8
437.1	21	*9	13	*	*	*	17	*	11	*5	*
437.2	12	*	*8	*	*	*	*5	*	*	*	*
437.3	16	*6	10	*	*	*9	*	*	*	*6	*
437.7	*5	*	*	—	*	*	*	*	*	*	*
437.9	30	12	18	—	*	*	29	*	9	12	*
438	550	261	289	*	23	155	369	118	138	159	135
438.0	11	*5	*6	—	*	*	*7	*	*	*	*
438.1	99	46	53	*	*	26	69	18	26	28	27
438.10	*7	*	*	*	*	*	*6	*	*	*	*
438.11	71	30	40	*	*	21	47	13	21	20	17
438.12	*7	*	*	—	*	*	*7	*	*	*	*
438.19	13	*8	*	*	*	*	*9	*	*	*	*
438.2	228	106	123	*	10	65	151	65	55	64	45
438.20	193	86	107	*	*7	54	131	55	45	58	35
438.21	19	10	9	—	*	*6	11	*6	*7	*	*
438.22	16	10	*6	—	*	*6	*9	*	*	*	*7
438.5	*8	*	*	*	*	*	*6	*	*	*	*
438.50	*6	*	*	—	*	*	*	*	*	*	*
438.7	10	*	*6	—	*	*	*5	*	*	*	*5
438.8	175	88	87	*	*7	49	118	26	47	52	49
438.82	45	24	22	—	*	17	27	*	10	14	16
438.83	*7	*	*	—	*	*	*	*	*	*	*
438.84	11	*7	*	—	*	*	*8	*	*	*	*
438.89	107	52	55	*	*5	25	77	17	31	33	27
438.9	14	*	10	*	*	*	10	*	*	*6	*
440	357	189	169	*	9	88	260	75	92	138	53
440.0	22	9	13	—	*	*5	16	*7	*7	*	*
440.1	48	23	26	*	*	9	36	*8	14	20	*6
440.2	228	129	99	—	*5	62	160	40	59	91	38
440.20	60	36	24	—	*	15	44	*6	15	29	10
440.21	51	30	22	—	*	21	29	*9	13	18	12
440.22	13	*6	*7	—	*	*	9	*	*	*8	*
440.23	32	17	16	—	*	*6	26	*9	*8	11	*
440.24	63	36	27	—	*	15	46	15	16	23	*9
440.29	*8	*	*	—	*	*	*6	*	*	*	*
440.3	11	*6	*	—	*	*	*7	*	*	*6	*
440.31	*6	*5	*	—	*	*	*5	*	*	*	*
440.8	*6	*	*	—	*	*	*	*	*	*	*
440.9	43	18	25	—	*	*6	36	17	*8	14	*
441	184	124	60	*	*	32	149	40	52	60	31
441.0	15	*8	*6	—	*	*5	*9	*5	*	*6	*
441.01	*7	*	*	—	*	*	*	*	*	*	*
441.2	19	12	*7	*	*	*7	12	*6	*6	*	*
441.3	*7	*5	*	—	*	*	*5	*	*	*	*
441.4	131	93	39	—	*	17	114	27	38	43	24

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
441.7	*6	*	*	—	*	*	*	*	*	*	*
441.9	*5	*	*	*	*	*	*	*	*	*	*
442	33	24	*8	*	*	10	19	*8	*8	11	*6
442.2	9	*8	*	—	*	*	*7	*	*	*	*
442.3	13	*9	*	—	*	*	*8	*	*	*	*
442.8	*6	*	*	—	*	*	*	*	*	*	*
443	502	275	227	*	18	126	357	118	117	199	68
443.0	16	*	11	*	*	*	*8	*	*6	*	*
443.2	*6	*	*	*	*	*	*	*	*	*	*
443.8	35	20	15	*	*	13	20	9	*7	14	*6
443.81	21	13	*8	—	*	*8	12	*6	*	*7	*
443.89	15	*7	*7	*	*	*6	*9	*	*	*6	*
443.9	443	245	198	—	10	105	328	105	102	179	57
444	91	49	42	*	*9	29	52	15	18	33	25
444.0	12	*8	*	—	*	*	*8	*	*	*	*
444.2	58	30	28	*	*5	20	33	10	12	21	16
444.21	12	*6	*6	—	*	*	*6	*	*	*	*
444.22	46	24	22	*	*	15	26	*7	*9	17	12
444.8	17	*9	*8	*	*	*	10	*	*	*7	*
444.81	11	*7	*	—	*	*	*8	*	*	*	*
444.89	*5	*	*	*	*	*	*	*	*	*	*
446	47	22	25	*6	*6	*7	28	14	12	11	11
446.1	*6	*	*	*5	*	*	*	*	*	*	*
446.2	*5	*	*	*	*	*	*	*	*	*	*
446.4	*8	*	*	*	*	*	*	*	*	*	*
446.5	19	*7	12	—	-	*	18	*6	*5	*	*
446.6	*6	*	*	*	*	*	*	*	*	*	*
447	57	28	29	*	*8	18	30	10	11	24	12
447.1	26	14	13	*	*	*8	17	*	*6	11	*6
447.6	12	*5	*7	*	*	*	*	*	*	*	*
447.8	*7	*	*	*	*	*	*	*	*	*	*
447.9	*5	*	*	—	*	*	*	*	*	*	*
448	*5	*	*	*	*	*	*	*	*	*	*
451	51	20	31	*	*9	15	26	13	11	20	*6
451.0	*8	*	*	—	*	*	*5	*	*	*	*
451.1	13	*	*9	—	*	*	*8	*5	*	*	*
451.19	9	*	*5	—	*	*	*5	*	*	*	*
451.2	*7	*	*5	*	*	*	*	*	*	*	*
451.8	20	*8	12	*	*5	*7	*7	*	*	9	*
451.82	*9	*	*	*	*	*	*	*	*	*	*
451.84	*7	*	*	*	*	*	*	*	*	*	*
452	*7	*	*	*	*	*	*	*	*	*	*
453	385	167	218	*	53	111	218	87	95	146	57
453.8	351	153	198	*	50	101	199	78	90	133	51
453.9	26	10	16	*	*	*7	17	*7	*	10	*5
454	33	13	20	—	*	10	19	*7	9	*9	*8
454.0	9	*	*6	—	*	*	*6	*	*	*	*
454.9	14	*5	9	—	*	*	*7	*	*	*	*
455	200	76	124	*	43	56	100	39	34	89	38
455.0	72	28	44	—	11	23	39	14	12	30	16
455.2	36	16	20	—	*9	10	17	9	*5	16	*5
455.3	22	*7	15	*	*	*8	10	*	*	13	*
455.5	11	*	*6	—	*	*	*6	*	*	*6	*
455.6	38	13	25	*	12	*8	18	*8	*8	14	*8
455.8	12	*	*7	—	*	*	*7	*	*	*7	*
456	72	45	27	*	15	38	18	16	12	26	17
456.0	*5	*	*	*	*	*	*	*	*	*	*
456.1	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
456.2	46	31	15	*	*8	26	11	10	*8	17	11
456.20	29	20	9	*	*7	16	*6	*6	*	12	*9
456.21	16	10	*6	*	*	10	*	*	*	*	*
456.8	10	*6	*	*	*	*	*	*	*	*	*
457	32	12	20	*	*	11	16	*5	10	11	*7
457.1	23	*8	15	*	*	*8	13	*	*7	*7	*
458	501	233	267	10	61	130	300	105	128	176	92
458.0	119	56	63	*	13	27	78	27	28	46	18
458.2	67	30	37	*	*6	18	42	16	18	20	13
458.29	65	29	36	*	*6	17	41	16	18	18	12
458.8	35	14	21	*	*	*9	21	*	10	13	*7
458.9	277	133	145	*7	37	75	158	57	71	96	54
459	177	88	89	*	15	58	101	44	48	62	23
459.0	26	*9	17	*	*	*6	17	*	*6	10	*
459.2	14	*8	*6	*	*	*9	*	*	*	*	*
459.8	124	66	58	*	*9	41	73	32	37	43	13
459.81	117	64	54	*	*8	39	70	30	36	39	12
459.89	*7	*	*	*	*	*	*	*	*	*	*
459.9	*8	*	*	*	*	*	*	*	*	*	*
460-519	11,855	5,605	6,250	1,235	1,419	2,828	6,373	2,565	2,893	4,290	2,106
461	56	22	34	*7	13	12	25	*8	20	20	*8
461.0	14	*	10	*	*	*	*7	*	*6	*	*
461.2	*7	*	*	*	*	*	*	*	*	*	*
461.9	26	10	16	*	*8	*6	*9	*	*8	9	*
462	55	26	29	14	24	*7	9	12	13	23	*7
463	19	10	9	*8	9	*	*	*5	*	*8	*
464	33	14	20	23	*	*	*	*5	*7	12	*9
464.1	*7	*	*	*6	*	*	*	*	*	*	*
464.10	*7	*	*	*6	*	*	*	*	*	*	*
464.4	16	*7	*9	16	*	*	-	*	*	*7	*
465	154	66	87	68	35	20	31	34	35	56	28
465.9	150	64	86	67	33	19	31	33	34	55	28
466	439	216	222	234	38	56	111	76	89	177	97
466.0	207	82	125	9	35	54	108	45	43	87	31
466.1	232	135	98	224	*	*	*	31	46	89	66
466.11	133	81	52	131	*	*	*	15	28	48	42
466.19	100	54	46	94	*	*	*	15	18	42	24
470	13	*6	*7	*	*6	*	*	*	*	*7	*
471	*7	*	*	*	*	*	*	*	*	*	*
472	12	*5	*7	*	*5	*	*	*	*	*	*
472.0	11	*5	*6	*	*	*	*	*	*	*	*
473	148	66	82	22	41	45	41	26	32	60	31
473.0	20	*7	13	*	*	*5	*6	*	*	*8	*
473.2	17	*9	*8	*	*	*7	*	*	*	*6	*
473.3	*6	*	*	*	*	*	*	*	*	*	*
473.8	10	*	*	*	*	*	*	*	*	*	*
473.9	92	40	52	11	28	27	26	16	20	40	16
474	36	20	16	22	12	*	*	10	*6	11	10
474.0	13	*6	*7	*6	*6	*	*	*	*	*6	*
474.00	10	*	*6	*	*6	*	*	*	*	*	*
474.1	22	14	*8	16	*6	*	*	*8	*	*5	*6
474.10	11	*6	*	10	*	-	-	*	*	*	*
474.11	10	*7	*	*	*5	*	*	*	*	*	*
475	15	*8	*6	*	9	*	*	*	*	*	*
477	68	28	40	*	17	25	23	12	16	30	10
477.9	65	26	39	*	15	24	22	11	16	29	9

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
478	113	61	52	22	30	36	24	18	22	45	28
478.0	*5	*	*	—	*	*	*	*	*	*	*
478.1	25	14	11	*7	10	*	*	*	*	14	*
478.2	43	26	17	*7	11	18	*8	*6	9	17	10
478.24	*5	*	*	*	*	*	*	*	*	*	*
478.29	30	17	13	*	*8	14	*5	*5	*6	10	*9
478.3	15	*8	*7	*	*	*	*7	*	*	*	*
478.30	*5	*	*	*	*	*	*	*	*	*	*
478.31	*6	*	*	*	*	*	*	*	*	*	*
478.6	*5	*	*	*	*	*	*	*	*	*	*
478.7	13	*5	*8	*	*	*	*	*	*	*5	*
478.74	*6	*	*	*	*	*	*	*	*	*	*
480	45	24	21	35	*	*	*5	*	13	14	14
480.1	22	13	9	22	*	*	*	*	*7	*6	*7
480.9	20	9	11	12	*	*	*	*	*6	*7	*6
481	51	29	23	*	*6	14	27	11	11	19	10
482	244	133	112	12	25	50	157	42	63	87	52
482.0	15	*8	*8	*	*	*	11	*	*	*5	*
482.1	47	28	19	*	*8	*7	29	*8	12	17	*9
482.2	12	*6	*6	*	*	*	*7	*	*	*	*
482.3	13	*	*8	*	*	*	*7	*	*	*	*
482.30	*6	*	*	*	*	*	*	*	*	*	*
482.4	87	48	39	*	*8	18	58	15	20	33	20
482.40	*5	*	*	*	*	*	*	*	*	*	*
482.41	76	43	33	*	*8	16	50	13	16	30	17
482.49	*5	*	*	*	*	*	*	*	*	*	*
482.8	50	28	22	*	*	12	33	*7	16	19	*9
482.82	12	*8	*	*	*	*	*9	*	*	*	*
482.83	27	16	11	*	*	*6	18	*	*6	14	*
482.9	20	10	10	*	*	*	12	*6	*5	*	*
483	13	*6	*7	*	*	*	*	*	*	*	*
483.0	10	*	*5	*	*	*	*	*	*	*	*
484	*5	*	*	*	*	*	*	*	*	*	*
485	31	15	16	*8	*	*6	15	*5	*5	15	*
486	1,937	939	998	206	163	385	1,183	408	524	672	332
487	37	16	21	*8	*	*6	21	11	13	9	*
487.0	13	*6	*7	*	*	*	10	*	*6	*	*
487.1	22	10	13	*6	*	*	11	*8	*6	*6	*
490	103	45	58	12	19	25	47	15	22	50	17
491	1,056	507	550	*	30	294	729	240	253	421	142
491.2	1,018	490	528	*	25	281	710	234	242	404	137
491.20	100	53	47	*	*	28	67	18	24	39	20
491.21	917	437	481	*	21	253	643	216	218	366	118
491.8	*5	*	*	*	*	*	*	*	*	*	*
491.9	29	12	16	*	*	11	13	*	*7	14	*
492	228	117	111	*	9	63	156	70	59	71	28
492.0	14	10	*	—	*	*	*6	*	*	*6	*
492.8	214	107	107	*	*7	58	150	67	56	65	26
493	1,870	641	1,229	348	501	504	517	475	434	601	359
493.0	81	42	39	54	14	*8	*	17	*8	17	38
493.00	32	13	19	16	*8	*6	*	10	*	*6	11
493.01	10	*7	*	*8	*	*	*	*	*	*	*
493.02	39	23	16	30	*5	*	*	*5	*	*8	22
493.2	432	139	294	*	35	144	253	109	110	132	80
493.20	269	88	181	*	23	83	163	64	68	86	51
493.21	*8	*	*6	*	*	*	*	*	*	*	*
493.22	155	49	107	*	11	57	87	44	40	44	28
493.8	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
493.9	1,350	458	892	292	447	350	260	346	314	449	240
493.90	975	306	669	131	355	281	207	256	226	308	185
493.91	78	39	38	51	17	*5	*	19	15	35	*8
493.92	297	113	185	110	75	64	48	71	74	106	46
494	31	15	16	*	*	*	21	10	*	13	*
494.0	20	*9	11	*	*	*	13	*6	*	*7	*
494.1	11	*6	*	*	*	*	*8	*	*	*6	*
495	*5	*	*	*	*	*	*	*	*	*	*
496	2,142	1,101	1,041	*	70	527	1,545	468	581	781	312
500	*5	*	*	—	*	*	*	*	*	*	*
501	21	20	*	—	*	*	18	*7	*	*8	*
507	373	212	161	12	30	58	273	87	76	126	84
507.0	370	210	160	11	30	58	271	87	76	125	83
510	27	19	*8	*	*	11	*8	*5	*	10	*8
510.9	23	16	*7	*	*	9	*7	*	*	*8	*7
511	434	217	217	16	46	103	269	102	99	155	78
511.0	21	10	11	*	*	*8	*9	*	*5	*8	*
511.8	32	17	16	*	*	9	18	*5	*8	12	*8
511.9	379	189	190	15	37	85	241	91	86	135	66
512	98	57	41	*	25	27	42	20	20	38	20
512.1	46	25	21	*	*6	15	24	9	*9	15	13
512.8	48	30	18	*	18	11	17	10	10	22	*7
513	14	*8	*5	*	*	*	*	*	*	*	*
513.0	13	*8	*5	*	*	*	*	*	*	*	*
514	34	16	19	*	*5	*9	18	*7	*7	17	*
515	186	82	104	*	11	36	137	38	47	66	35
516	33	16	17	*	*	*9	20	*7	*9	10	*8
516.3	17	*8	9	—	*	*	13	*	*	*	*
516.8	14	*7	*7	*	*	*	*7	*	*	*	*
517	31	10	22	*	*7	16	*7	*9	*6	12	*
517.3	*6	*	*	*	*	*	*	*	*	*	*
517.8	22	*6	16	*	*	14	*6	*6	*	*7	*
518	1,533	745	787	98	191	421	823	285	345	587	315
518.0	384	171	213	29	64	110	181	69	89	142	85
518.1	*7	*	*	*	*	*	*	*	*	*	*
518.3	23	13	10	*	*	*6	12	*	*5	11	*
518.4	19	10	*9	*	*	*	11	*	*5	*7	*
518.5	114	65	49	*6	22	36	51	24	25	43	23
518.8	984	482	502	60	95	262	566	184	220	382	199
518.81	669	331	338	24	69	180	397	131	142	261	135
518.82	96	49	46	25	*9	17	44	13	20	39	25
518.83	50	23	27	*	*	14	31	*6	12	23	9
518.84	62	28	34	*	*	20	37	10	18	25	9
518.89	108	50	58	*9	11	31	57	24	28	35	21
519	84	48	36	17	11	22	34	12	27	31	13
519.0	15	*8	*6	*	*	*	*6	*	*	*6	*
519.09	*7	*	*	*	*	*	*	*	*	*	*
519.1	32	17	14	*8	*	*7	12	*	10	11	*6
519.8	28	16	12	*	*	*8	13	*	12	10	*
520-579	11,144	4,758	6,386	538	2,225	3,453	4,928	2,381	2,604	4,109	2,051
521	22	11	11	*	11	*6	*	*5	*6	*9	*
521.0	21	11	10	*	11	*	*	*5	*6	*8	*
521.00	15	*7	*8	*	*6	*	*	*	*	*6	*
521.09	*5	*	*	*	*	*	*	*	*	*	*
522	27	11	15	*	10	*8	*5	*6	*8	9	*
522.5	23	11	12	*	*8	*7	*	*6	*6	*8	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
523	13	*	*9	*	*	*5	*	*	*	*5	*
524	39	12	27	*	27	*	*	*8	*	20	*8
524.0	17	*6	11	*	14	*	*	*5	*	*7	*
524.03	*6	*	*	*	*	*	*	*	*	*	*
524.6	10	*	*9	*	*	*	*	*	*	*	*
524.60	*5	*	*	—	*	*	*	*	*	*	*
525	12	*8	*	*	*7	*	*	*	*	*	*
525.9	*5	*	*	*	*	*	*	*	*	*	*
526	*8	*	*	*	*	*	*	*	*	*5	*
527	16	*6	10	*	*	*6	*7	*	*	*5	*
527.2	11	*	*7	*	*	*	*5	*	*	*	*
528	44	23	21	12	*9	13	10	*8	10	16	10
528.0	19	9	10	*7	*	*6	*	*	*	*7	*
528.3	*6	*	*	*	*	*	*	*	*	*	*
528.5	*5	*	*	*	*	*	*	*	*	*	*
528.9	11	*5	*6	*	*	*	*	*	*	*	*
529	*7	*	*	*	*	*	*	*	*	*	*
530	2,391	971	1,420	105	370	807	1,109	543	585	870	394
530.0	*6	*	*	*	*	*	*	*	*	*	*
530.1	208	102	106	*	41	65	97	49	45	76	38
530.10	74	36	38	*	17	21	35	21	17	26	11
530.11	70	29	41	*	13	23	32	17	14	24	16
530.12	*9	*6	*	*	*	*	*	*	*	*	*
530.19	55	31	24	*	*9	19	25	10	10	24	10
530.2	38	18	20	*	*6	10	22	9	*7	15	*7
530.20	21	9	12	*	*	*7	10	*5	*	*7	*
530.21	17	9	*8	*	*	*	13	*	*	*8	*
530.3	54	22	32	*	*	11	36	10	12	20	13
530.5	23	10	12	*	*	*8	12	*	*6	*8	*
530.6	*8	*	*	—	*	*	*6	*	*	*	*
530.7	27	16	10	*	*7	10	10	*5	*7	*9	*6
530.8	2,023	793	1,230	96	308	698	921	462	503	735	322
530.81	1,947	753	1,194	95	298	676	878	443	486	713	304
530.82	15	*8	*7	*	*	*	*8	*	*	*7	*
530.85	43	26	18	*	*	14	25	12	11	11	10
530.89	17	*6	11	*	*	*	10	*5	*	*	*
531	189	81	108	*	16	57	114	37	45	70	38
531.0	19	*7	11	—	*	*6	11	*	*5	*6	*
531.00	19	*7	11	—	*	*6	11	*	*5	*6	*
531.3	*7	*	*6	—	*	*	*	*	*	*	*
531.30	*7	*	*6	—	*	*	*	*	*	*	*
531.4	94	43	51	*	*6	26	62	18	19	35	21
531.40	90	40	50	*	*5	25	60	18	19	35	19
531.5	*6	*	*	*	*	*	*	*	*	*	*
531.50	*5	*	*	*	*	*	*	*	*	*	*
531.7	*6	*	*	—	*	*	*	*	*	*	*
531.70	*6	*	*	—	*	*	*	*	*	*	*
531.9	55	22	33	*	*6	19	29	12	15	21	*8
531.90	53	22	32	*	*6	19	27	11	15	20	*7
532	116	67	49	*	12	31	71	25	27	42	23
532.0	11	*6	*	*	*	*	*7	*	*	*	*
532.00	11	*6	*	*	*	*	*7	*	*	*	*
532.4	57	36	21	*	*6	18	34	13	13	21	11
532.40	56	35	21	*	*5	18	33	13	13	20	11
532.5	10	*5	*	*	*	*	*8	*	*	*	*
532.50	10	*5	*	*	*	*	*8	*	*	*	*
532.9	30	15	15	*	*	*7	18	*6	*6	12	*6
532.90	29	14	14	*	*	*7	18	*6	*6	12	*6
533	121	61	60	*	15	37	70	28	33	44	17

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
533.4	10	*	*6	—	*	*	*7	*	*	*	*
533.40	9	*	*6	—	*	*	*7	*	*	*	*
533.7	*6	*	*	—	*	*	*	*	*	*	*
533.70	*6	*	*	—	*	*	*	*	*	*	*
533.9	101	53	48	*	13	33	55	23	29	36	14
533.90	101	53	48	*	13	33	55	23	29	36	13
534	16	*	11	*	*	*9	*	*	*	*5	*
534.4	*8	*	*6	—	*	*	*	*	*	*	*
534.40	*8	*	*5	—	*	*	*	*	*	*	*
535	628	277	351	15	117	180	316	131	144	254	98
535.0	59	23	35	*	12	15	28	15	12	28	*
535.00	46	18	28	*	10	13	21	11	*9	23	*
535.01	13	*6	*7	*	*	*	*8	*	*	*5	*
535.1	83	35	47	*	10	22	49	20	13	33	16
535.10	64	27	37	*	*8	18	37	14	10	29	12
535.11	18	*8	10	*	*	*	13	*6	*	*	*
535.3	21	15	*6	—	*8	12	*	*	*	*7	*6
535.30	15	12	*	—	*6	*8	*	*	*	*	*
535.31	*7	*	*	—	*	*	*	*	*	*	*
535.4	147	63	84	*	25	39	81	29	34	55	30
535.40	103	43	61	*	21	29	52	20	25	37	21
535.41	44	21	23	*	*	10	29	*8	*9	18	*9
535.5	250	100	150	*7	48	65	130	49	61	108	32
535.50	202	78	124	*7	43	54	99	41	50	87	24
535.51	48	22	26	*	*5	12	31	*8	11	22	*8
535.6	67	39	28	*	14	26	25	14	19	24	10
535.60	54	31	23	*	14	21	18	12	17	19	*7
535.61	12	*8	*	*	*	*	*7	*	*	*	*
536	268	102	166	15	62	83	107	43	76	107	41
536.2	33	*9	24	*7	10	*6	10	*5	*7	14	*7
536.3	95	31	64	*	30	39	26	17	17	45	16
536.4	47	23	24	*	*	11	28	11	*8	20	*9
536.41	12	*5	*7	*	*	*	*8	*	*	*5	*
536.42	20	11	9	*	*	*	12	*	*	*8	*
536.49	14	*7	*7	*	*	*	*9	*	*	*7	*
536.8	88	36	52	*	19	26	41	9	43	25	10
537	79	33	46	*	*7	18	53	17	15	29	18
537.0	25	*9	16	*	*	*7	15	*	*	10	*6
537.8	44	18	26	*	*	*9	32	10	*9	15	10
537.82	*6	*	*	—	*	*	*5	*	*	*	*
537.83	23	10	13	—	*	*	19	*5	*	*8	*6
537.89	13	*	*8	*	*	*	*6	*	*	*	*
540	311	180	131	80	155	53	23	67	64	93	87
540.0	59	35	24	16	20	13	10	11	16	16	17
540.1	37	22	15	11	12	9	*	*8	*6	11	12
540.9	215	123	92	53	124	31	*8	48	43	66	58
541	12	*7	*5	*	*6	*	*	*	*	*5	*
542	*6	*	*	*	*	*	*	*	*	*	*
543	20	*8	13	*	11	*	*	*	*6	*7	*
543.9	19	*7	12	*	10	*	*	*	*5	*7	*
550	71	58	14	12	*8	13	38	17	14	23	17
550.1	23	19	*	*	*	*	13	*6	*	*8	*
550.10	16	12	*	*	*	*	*9	*	*	*5	*
550.9	47	37	9	*9	*5	*8	24	11	*9	15	12
550.90	34	26	*8	*6	*	*6	18	9	*6	11	*7
550.92	*9	*8	*	*	*	*	*	*	*	*	*
552	95	36	59	*	21	30	43	20	24	35	17
552.1	17	*7	10	*	*6	*6	*5	*	*5	*6	*
552.2	61	22	39	*	14	21	27	14	16	21	11

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
552.20	14	*	10	*	*	*	*6	*	*	*6	*
552.21	42	16	25	—	10	14	18	10	12	13	*8
552.29	*6	*	*	—	*	*	*	*	*	*	*
552.8	*8	*	*	*	*	*	*	*	*	*	*
553	532	191	341	11	78	176	268	103	128	211	90
553.1	55	23	32	*5	15	17	18	13	12	17	14
553.2	108	37	71	*	25	46	36	21	21	43	23
553.20	32	11	21	—	*6	12	14	*7	*7	13	*5
553.21	72	24	48	*	18	32	21	13	14	28	16
553.3	361	130	231	*	36	112	209	68	92	149	51
555	106	39	67	*	49	34	20	29	27	38	12
555.0	14	*7	*7	*	*7	*	*	*	*	*5	*
555.1	16	*7	9	*	*9	*5	*	*	*	*7	*
555.2	*7	*	*	*	*	*	*	*	*	*	*
555.9	69	22	47	*	29	23	15	21	17	23	*8
556	79	41	38	*	32	24	21	20	23	23	13
556.6	10	*	*6	*	*	*	*	*	*	*	*
556.8	*8	*	*	*	*5	*	*	*	*	*	*
556.9	51	26	25	*	20	14	16	12	16	14	*9
557	84	25	59	*	10	18	54	22	18	26	18
557.0	43	15	28	*	*7	*9	25	10	10	12	10
557.1	*6	*	*	*	*	*	*	*	*	*	*
557.9	35	*8	28	*	*	*8	25	9	*7	12	*7
558	417	157	261	114	86	82	135	81	99	172	66
558.1	*6	*	*	—	*	*	*	*	*	*	*
558.9	407	152	255	112	86	79	130	78	95	168	65
560	776	342	434	29	111	211	425	148	190	291	147
560.0	*8	*	*	*	*	*	*	*	*	*	*
560.1	341	166	175	9	57	97	177	55	92	132	63
560.2	20	10	10	*	*	*	10	*	*	*7	*6
560.3	84	30	54	*6	*5	*9	64	17	16	37	14
560.39	79	29	50	*6	*	*8	60	16	15	35	13
560.8	155	61	94	*	22	48	82	31	36	55	34
560.81	86	31	54	*	13	24	46	15	17	32	21
560.89	70	30	40	*	9	24	36	16	19	23	12
560.9	167	71	97	*5	20	51	91	39	41	57	30
562	646	255	391	*	59	160	426	158	156	232	99
562.0	14	*8	*5	—	*	*	*9	*	*	*6	*
562.00	10	*7	*	—	*	*	*6	*	*	*	*
562.1	632	246	386	*	57	158	417	156	154	225	96
562.10	272	100	172	—	15	56	201	64	67	99	42
562.11	246	96	150	*	38	82	125	66	63	80	36
562.12	99	44	55	*	*	17	80	23	18	41	17
562.13	15	*6	9	—	*	*	11	*	*5	*	*
564	664	227	437	39	133	174	319	137	191	214	122
564.0	462	186	276	34	71	110	246	93	137	153	79
564.00	410	166	244	28	63	97	221	79	124	138	69
564.09	52	20	32	*6	*8	13	24	14	13	15	10
564.1	163	23	140	*	51	53	58	36	46	50	32
564.8	19	11	*8	*	*6	*	*6	*5	*	*5	*
564.81	*6	*	*	*	*	*	*	*	*	*	*
564.89	13	*6	*7	*	*	*	*5	*	*	*	*
564.9	*8	*	*	*	*	*	*	*	*	*	*
565	13	*6	*6	*	*5	*	*	*	*	*7	*
565.1	*8	*	*	*	*	*	*	*	*	*5	*
566	39	26	13	*	16	14	*6	*7	9	17	*5
567	112	57	55	*	24	42	41	21	26	37	28
567.2	75	39	36	*	17	28	27	14	18	24	18

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
567.8	16	*8	*8	*	*	*6	*5	*	*	*	*
567.9	21	10	11	*	*	*8	*9	*	*6	*8	*
568	171	46	125	*5	63	62	41	35	36	68	33
568.0	134	37	97	*	46	53	31	29	25	54	26
568.8	35	*7	28	*	16	*8	10	*6	11	13	*5
568.81	20	*	17	*	11	*	*6	*	*7	*6	*
568.89	15	*	10	*	*	*	*	*	*	*6	*
569	289	123	166	*6	44	89	149	63	60	114	52
569.0	11	*	*6	*	*	*	*6	*	*	*5	*
569.1	10	*	*8	*	*	*	*5	*	*	*	*
569.3	72	29	43	*	10	23	39	13	14	36	9
569.4	31	16	15	*	*5	*8	18	*6	*7	13	*6
569.4	*9	*	*	*	*	*	*5	*	*	*	*
569.49	18	10	*8	*	*	*	11	*	*	*7	*
569.5	26	12	15	—	*5	12	9	*7	*5	*8	*7
569.6	34	15	19	*	*8	9	15	*7	*8	9	*9
569.61	*5	*	*	*	*	*	*	*	*	*	*
569.62	*9	*	*5	*	*	*	*	*	*	*	*
569.69	19	*8	11	*	*	*	10	*	*	*6	*
569.8	97	41	56	*	14	28	54	23	21	37	15
569.81	14	*	11	*	*	*6	*	*	*	*8	*
569.82	12	*5	*6	*	*	*	*6	*	*	*	*
569.83	27	13	14	*	*	10	13	*	*6	13	*5
569.84	*6	*	*	—	*	*	*	*	*	*	*
569.85	22	9	13	*	*	*	19	*	*5	*8	*
569.89	16	*8	*8	*	*	*	*7	*	*	*	*
569.9	*6	*	*	*	*	*	*	*	*	*	*
570	39	18	22	*	*9	13	16	10	*7	13	10
571	513	300	214	*	103	269	139	109	92	198	115
571.0	*7	*	*	—	*	*	*	*	*	*	*
571.1	53	38	15	—	20	29	*	*9	12	18	14
571.2	163	121	42	*	33	96	34	39	30	51	44
571.3	25	18	*7	—	*9	12	*	*	*7	*	10
571.4	15	*	11	*	*	*6	*6	*	*	*	*
571.49	11	*	9	*	*	*	*5	*	*	*	*
571.5	189	95	94	*	18	94	76	43	29	88	28
571.6	12	*	11	*	*	*8	*	*	*	*5	*
571.8	44	14	30	*	17	17	9	*8	*7	20	*8
571.9	*6	*	*	*	*	*	*	*	*	*	*
572	163	100	63	*	22	96	43	38	26	60	39
572.0	*6	*	*	*	*	*	*	*	*	*	*
572.2	91	57	34	*	*9	57	25	21	13	35	22
572.3	37	24	13	*	*8	19	*9	*7	*8	14	*8
572.4	11	*7	*	*	*	*7	*	*	*	*	*
572.8	18	10	*9	*	*	11	*	*5	*	*6	*
573	93	41	52	*	21	32	37	22	16	37	19
573.3	38	16	21	*	12	14	10	10	*6	13	*8
573.8	33	14	19	*	*	11	17	*7	*6	14	*6
573.9	16	*7	*8	*	*	*6	*7	*	*	*7	*
574	627	209	418	*	192	179	252	137	126	224	140
574.0	159	63	97	*	47	52	60	31	36	60	33
574.00	147	57	90	*	45	50	52	29	35	55	29
574.01	12	*5	*7	—	*	*	*8	*	*	*5	*
574.1	218	58	159	*	74	67	76	55	36	76	51
574.10	209	55	154	*	71	65	72	52	36	72	49
574.11	*9	*	*6	—	*	*	*	*	*	*	*
574.2	140	50	89	*	40	32	65	31	29	51	29
574.20	136	48	88	*	39	31	63	31	29	49	27
574.3	11	*	*8	—	*	*	*	*	*	*5	*
574.31	*7	*	*5	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
574.4	10	*	*7	*	*	*	*	*	*	*	*
574.41	*6	*	*	*	*	*	*	*	*	*	*
574.5	40	18	22	*	9	*8	22	*9	9	13	*9
574.50	23	9	14	*	*7	*	12	*6	*5	*6	*6
574.51	16	*8	*8	—	*	*	10	*	*	*7	*
574.6	9	*	*6	—	*	*	*	*	*	*	*
574.7	22	*5	17	*	*8	*5	9	*	*6	*7	*6
574.70	13	*	11	*	*6	*	*	*	*	*	*
574.71	*9	*	*6	*	*	*	*5	*	*	*	*
574.8	10	*	*6	—	*	*	*	*	*	*	*
574.81	*6	*	*	—	*	*	*	*	*	*	*
574.9	9	*	*7	*	*	*	*	*	*	*	*
574.90	*6	*	*	*	*	*	*	*	*	*	*
575	151	58	93	*	48	46	54	26	40	52	33
575.0	33	15	18	*	*6	10	17	*7	10	10	*6
575.1	67	26	42	*	24	20	22	10	17	27	13
575.10	18	*7	10	*	*	*5	*7	*	*	*7	*
575.11	38	13	25	*	17	12	*9	*5	10	15	*6
575.12	12	*6	*6	—	*	*	*6	*	*	*	*
575.3	*7	*	*	*	*	*	*	*	*	*	*
575.6	*7	*	*	—	*	*	*	*	*	*	*
575.8	30	9	21	*	13	9	*8	*	*9	10	*7
576	106	43	63	*	26	32	47	27	19	39	22
576.1	34	17	17	*	*5	11	17	9	*	12	*8
576.2	30	14	16	*	*7	*7	16	*8	*6	9	*6
576.8	40	12	29	*	13	12	13	*8	*8	17	*7
577	479	232	247	12	152	186	129	95	100	184	100
577.0	337	159	179	*6	109	122	100	63	71	131	71
577.1	84	43	41	*	30	40	11	16	17	32	19
577.2	26	14	12	*	*6	13	*6	*7	*5	*9	*
577.8	16	*6	10	*	*5	*	*5	*	*	*6	*
577.9	16	10	*6	*	*	*6	*7	*	*	*6	*
578	478	229	249	9	47	123	299	103	130	168	77
578.0	67	34	33	*	15	20	30	14	18	24	12
578.1	130	60	71	*	16	37	74	26	34	47	24
578.9	280	136	145	*	16	66	194	63	78	98	41
579	48	21	27	*9	11	13	15	11	12	17	*9
579.0	*7	*	*5	*	*	*	*	*	*	*	*
579.3	22	10	12	*	*	*7	*6	*	*	9	*
579.8	14	*6	*8	*	*	*	*	*	*	*	*
580-629	8,642	3,040	5,602	195	2,092	2,141	4,215	1,727	2,072	3,249	1,594
580	*8	*	*5	*	*	*	*	*	*	*	*
581	42	23	19	*8	*9	15	11	9	*6	17	10
581.8	19	9	10	*	*	*9	*5	*	*	*7	*
581.81	18	*9	9	*	*	*9	*	*	*	*6	*
581.9	18	10	*7	*6	*	*	*	*	*	*8	*
582	18	10	*8	*	*8	*	*5	*	*	9	*
582.1	*7	*	*	*	*	*	*	*	*	*	*
582.8	*6	*	*	*	*	*	*	*	*	*	*
582.9	*5	*	*	*	*	*	*	*	*	*	*
583	144	69	76	*	25	56	61	32	34	48	30
583.8	127	58	69	*	19	50	57	28	30	43	26
583.81	120	54	65	*	17	48	54	25	29	41	25
583.89	*7	*	*	*	*	*	*	*	*	*	*
583.9	14	*8	*6	*	*5	*	*	*	*	*	*
584	947	488	459	*8	79	252	609	204	211	358	174
584.5	83	42	41	*	*8	27	47	20	17	30	16
584.8	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
584.9	856	442	414	*6	70	222	558	183	192	324	157
585	247	134	113	*5	22	64	156	60	65	79	44
586	32	16	16	*	*	*8	20	*7	*7	12	*
588	23	11	12	*	*	*9	*9	*	*	11	*
588.8	17	*7	10	*	*	*6	*6	*	*	10	*
590	246	42	204	26	107	47	67	31	68	95	53
590.0	*7	*	*5	*	*	*	*	*	*	*	*
590.00	*7	*	*5	*	*	*	*	*	*	*	*
590.1	100	16	83	*9	45	16	29	16	30	34	20
590.10	99	16	83	*9	45	16	29	16	30	34	20
590.8	136	23	113	16	59	29	33	13	35	57	31
590.80	136	23	113	16	59	29	33	13	35	57	31
591	226	97	129	10	65	62	88	49	55	79	42
592	293	140	153	*	121	106	64	59	75	109	50
592.0	138	59	79	*	58	48	32	33	35	48	23
592.1	150	80	71	*	60	58	31	26	39	59	27
593	806	421	385	24	51	173	557	169	209	288	141
593.2	15	*9	*7	*	*	*7	*7	*7	*	*	*
593.3	10	*	*6	*	*	*	*	*	*	*	*
593.4	24	10	15	*	*	*8	11	*	*	*8	*7
593.5	15	*6	*9	*	*5	*	*	*	*5	*	*
593.7	17	*6	11	15	*	*	*	*	*	*7	*
593.70	13	*	*9	12	*	*	*	*	*	*	*
593.8	17	*8	*8	*	*	*6	*5	*	*	*6	*
593.89	12	*5	*6	*	*	*	*	*	*	*	*
593.9	706	378	329	*	33	144	525	144	186	255	121
594	20	16	*	*	*	*	14	*	*	*7	*
594.1	18	14	*	*	*	*	13	*	*	*7	*
595	81	30	51	*	10	18	52	21	18	29	12
595.0	19	*7	11	*	*	*	12	*	*	*7	*
595.1	11	*	9	—	*	*	*	*	*	*	*
595.2	*8	*	*	*	*	*	*6	*	*	*	*
595.8	11	*7	*	*	—	*	*8	*	*	*	*
595.82	*6	*	*	—	—	*	*	*	*	*	*
595.9	30	10	20	*	*	*	21	*6	*6	13	*
596	174	102	72	*	27	36	107	41	52	48	33
596.0	35	31	*	*	*	*	29	*8	*8	10	*7
596.1	*7	*	*	*	*	*	*	*	*	*	*
596.4	*7	*	*	*	*	*	*	*	*	*	*
596.5	81	38	43	*	17	20	42	22	24	21	14
596.51	11	*	9	—	*	*	*8	*	*	*	*
596.54	62	32	30	*	15	17	27	15	19	16	12
596.59	*9	*	*	*	*	*	*6	*	*	*	*
596.8	31	15	16	*	*7	*7	17	*	12	10	*5
598	26	21	*	*	*	*6	16	*7	*6	*7	*6
598.9	19	16	*	*	*	*	12	*6	*	*6	*
599	2,497	819	1,678	62	281	453	1,701	539	601	917	439
599.0	2,126	596	1,530	51	235	370	1,470	458	510	793	365
599.6	27	20	*7	*	*	*7	18	*7	*8	*7	*
599.7	313	195	118	*9	42	66	197	68	80	104	60
599.8	13	*	12	*	*	*7	*	*	*	*7	*
599.81	*5	—	*5	—	*	*	*	*	*	*	*
599.9	13	*	*8	*	*	*	*8	*	*	*	*6
600	450	450	—	*	*	72	375	128	105	125	92
600.0	423	423	—	*	*	68	352	123	98	115	86
600.00	317	317	—	*	*	49	266	97	67	83	69
600.01	106	106	—	—	*	19	86	26	31	32	17
600.1	*5	*5	—	—	—	—	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
600.9	19	19	—	—	—	*	16	*	*	*8	*
600.90	11	11	—	—	—	*	9	*	*	*	*
600.91	*8	*8	—	—	—	*	*7	*	*	*	*
601	39	39	—	—	*	12	24	11	*8	15	*5
601.0	13	13	—	—	*	*	*8	*	*	*6	*
601.1	10	10	—	—	*	*	*6	*	*	*	*
601.9	13	13	—	—	*	*	*9	*	*	*5	*
603	11	11	—	*	*	*	*	*	*	*5	*
603.9	*8	*8	—	*	*	*	*	*	*	*	*
604	17	17	—	*	*	*	*7	*	*	*7	*
604.9	14	14	—	*	*	*	*6	*	*	*	*
604.90	11	11	—	*	*	*	*	*	*	*	*
605	10	10	—	*	*	*	*	*	*	*6	*
607	18	18	—	*	*	*	*7	*	*	*8	*
607.8	10	10	—	*	*	*	*5	*	*	*	*
607.84	*6	*6	—	—	*	*	*	*	*	*	*
608	37	37	—	*	*8	15	11	*	10	17	*5
608.4	16	16	—	*	*	*7	*	*	*	*9	*
608.8	15	15	—	*	*	*6	*5	*	*	*7	*
608.86	*8	*8	—	*	*	*	*	*	*	*	*
610	13	*	13	—	*	*8	*	*	*	*5	*
610.1	*9	*	*9	—	*	*	*	*	*	*	*
611	64	*	60	*	26	26	11	16	12	25	11
611.0	21	*	21	*	10	*8	*	*	*	9	*
611.1	18	*	16	*	*8	*7	*	*6	*	*6	*
611.7	17	*	16	—	*	*8	*5	*	*	*7	*
611.72	14	*	13	—	*	*7	*	*	*	*5	*
611.8	*6	*	*6	*	*	*	*	*	*	*	*
614	240	—	240	*	170	54	14	36	52	106	47
614.0	*6	—	*6	*	*	*	*	*	*	*	*
614.1	20	—	20	*	11	*8	*	*	*	10	*
614.2	12	—	12	*	*9	*	*	*	*	*	*
614.3	*7	—	*7	*	*	*	*	*	*	*	*
614.4	*6	—	*6	*	*	*	*	*	*	*	*
614.6	167	—	167	*	116	39	11	24	36	75	32
614.9	21	—	21	*	19	*	*	*	*6	9	*
615	29	—	29	*	28	*	*	*	*	15	*5
615.9	23	—	23	*	22	*	*	*	*	13	*
616	113	—	113	*	61	41	*9	19	27	45	22
616.0	78	—	78	*	39	33	*	16	19	27	16
616.1	22	—	22	*	13	*6	*	*	*6	10	*
616.10	21	—	21	*	13	*6	*	*	*5	10	*
616.4	*6	—	*6	*	*	*	*	*	*	*	*
617	281	—	281	*	186	88	*6	42	71	111	57
617.0	149	—	149	*	85	59	*	22	36	62	28
617.1	42	—	42	*	31	11	*	*7	11	14	11
617.2	11	—	11	—	*8	*	*	*	*	*	*
617.3	51	—	51	*	39	11	*	*	15	20	10
617.8	*6	—	*6	—	*	*	—	*	*	*	*
617.9	17	—	17	*	15	*	*	*	*	*7	*
618	253	—	253	—	56	103	94	31	57	109	56
618.0	64	—	64	—	14	29	21	11	16	26	11
618.1	30	—	30	—	15	*7	*8	*	*7	12	*7
618.2	25	—	25	—	*7	13	*6	*	*6	10	*
618.3	17	—	17	—	*	*6	*8	*	*	*	*5
618.4	34	—	34	—	*8	15	11	*	10	16	*6
618.5	36	—	36	—	*	12	23	*	*9	17	*8
618.6	27	—	27	—	*	10	12	*	*	13	*7

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
618.8	16	—	16	—	*	*8	*	*	*	*8	*6
619	14	—	14	*	*	*	*6	*	*	*5	*
619.1	9	—	9	*	*	*	*	*	*	*	*
620	326	—	326	*9	202	98	17	53	81	135	58
620.0	47	—	47	*	29	17	*	*6	13	21	*6
620.1	55	—	55	*	38	17	—	10	14	25	*6
620.2	161	—	161	*5	106	40	9	25	40	66	30
620.5	*8	—	*8	*	*	*	*	*	*	*	*
620.8	44	—	44	*	22	17	*	*8	10	16	10
620.9	*6	—	*6	*	*	*	*	*	*	*	*
621	83	—	83	—	33	39	11	17	20	32	14
621.0	35	—	35	—	11	18	*5	*9	9	12	*5
621.2	15	—	15	—	10	*5	*	*	*5	*6	*
621.3	20	—	20	—	*	12	*	*	*	*8	*
621.8	*8	—	*8	—	*	*	*	*	*	*	*
622	30	—	30	—	18	9	*	*6	*8	11	*
622.1	21	—	21	—	16	*	*	*	*	10	*
623	21	—	21	*	11	*	*	*6	*	*8	*
623.8	16	—	16	*	*9	*	*	*	*	*6	*
624	*6	—	*6	*	*	*	*	*	*	*	*
625	371	—	371	*	229	100	42	34	89	172	75
625.0	29	—	29	*	26	*	—	*	*8	14	*6
625.3	99	—	99	—	78	21	—	*7	22	50	20
625.6	141	—	141	*	43	59	38	14	33	60	34
625.8	*8	—	*8	—	*	*	*	*	*	*	*
625.9	84	—	84	*	69	13	*	10	21	40	14
626	316	—	316	*	205	106	*	42	72	143	59
626.2	239	—	239	*	154	83	*	29	57	108	44
626.4	*6	—	*6	*	*	*	—	*	*	*	*
626.6	*5	—	*5	—	*	*	—	*	*	*	*
626.8	53	—	53	*	34	17	*	*8	10	22	13
626.9	*7	—	*7	—	*6	*	*	*	*	*	—
627	48	—	48	—	*6	29	13	*8	12	19	9
627.0	*8	—	*8	—	*	*	*	*	*	*	*
627.1	26	—	26	—	*	19	*7	*	*8	10	*
627.9	*6	—	*6	—	*	*	*	*	*	*	*
628	*5	—	*5	*	*	*	—	*	*	*	*
630-677	10,337	—	10,337	16	10,292	29	—	1,556	2,068	4,109	2,604
632	22	—	22	—	22	—	—	*	*	10	*
633	34	—	34	*	32	*	—	*7	*	15	*9
633.1	29	—	29	*	28	*	—	*6	*	12	*8
633.10	28	—	28	*	27	*	—	*5	*	12	*7
634	19	—	19	*	19	*	—	*6	*	*9	*
634.9	13	—	13	*	13	*	—	*	*	*6	*
634.91	11	—	11	*	11	*	—	*	*	*5	*
639	11	—	11	—	11	—	—	*	*	*	*
640	12	—	12	—	12	—	—	*	*	*7	*
640.0	*7	—	*7	—	*7	—	—	*	*	*	*
640.03	*7	—	*7	—	*7	—	—	*	*	*	*
641	101	—	101	—	100	*	—	13	24	42	23
641.0	12	—	12	—	12	*	—	*	*	*7	*
641.01	11	—	11	—	11	*	—	*	*	*7	*
641.1	19	—	19	—	19	—	—	*	*	*7	*6
641.11	12	—	12	—	12	—	—	*	*	*	*
641.13	*6	—	*6	—	*6	—	—	*	*	*	*
641.2	43	—	43	—	43	—	—	*6	10	17	10

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
641.21	40	—	40	—	40	—	—	*5	9	15	10
641.8	*7	—	*7	—	*7	—	—	*	*	*	*
641.9	15	—	15	—	15	*	—	*	*	*5	*
641.91	*7	—	*7	—	*7	*	—	*	*	*	*
641.93	*8	—	*8	—	*8	*	—	*	*	*	*
642	384	—	384	*	381	*	—	59	79	165	80
642.0	46	—	46	—	46	*	—	*7	10	21	*8
642.01	37	—	37	—	37	*	—	*6	9	16	*6
642.03	*8	—	*8	—	*8	—	—	*	*	*	*
642.3	134	—	134	*	133	*	—	18	28	56	32
642.31	120	—	120	*	119	*	—	17	25	47	30
642.33	11	—	11	—	11	*	—	*	*	*8	*
642.4	101	—	101	*	101	*	—	18	24	37	23
642.41	84	—	84	*	84	*	—	14	19	31	20
642.43	*8	—	*8	—	*8	*	—	*	*	*	*
642.5	48	—	48	*	48	*	—	*7	*7	24	10
642.51	43	—	43	*	43	*	—	*6	*7	22	9
642.6	*5	—	*5	—	*5	—	—	*	*	*	*
642.7	16	—	16	—	16	*	—	*	*	10	*
642.71	14	—	14	—	14	*	—	*	*	*9	*
642.9	30	—	30	*	29	*	—	*	*9	14	*
642.91	22	—	22	*	21	*	—	*	*6	11	*
643	46	—	46	—	46	*	—	*6	11	24	*5
643.0	19	—	19	—	19	*	—	*	*	10	*
643.03	18	—	18	—	18	*	—	*	*	9	*
643.1	16	—	16	—	16	—	—	*	*	*8	*
643.13	16	—	16	—	16	—	—	*	*	*8	*
643.2	*7	—	*7	—	*7	*	—	*	*	*	*
643.23	*5	—	*5	—	*5	*	—	*	*	*	*
644	437	—	437	*	436	*	—	65	78	201	94
644.0	107	—	107	*	107	*	—	19	20	49	18
644.03	106	—	106	*	106	*	—	19	20	49	18
644.1	13	—	13	—	13	*	—	*	*	*6	*
644.13	13	—	13	—	13	*	—	*	*	*6	*
644.2	317	—	317	*	316	*	—	44	56	145	73
644.21	317	—	317	*	316	*	—	44	56	145	73
645	376	—	376	*	375	*	—	68	74	124	109
645.1	370	—	370	*	369	*	—	67	74	121	107
645.11	365	—	365	*	364	*	—	67	73	121	104
645.2	*6	—	*6	—	*6	—	—	*	*	*	*
645.21	*6	—	*6	—	*6	—	—	*	*	*	*
646	236	—	236	*	235	*	—	25	44	114	54
646.1	25	—	25	—	25	*	—	*	*6	11	*6
646.11	22	—	22	—	22	*	—	*	*6	10	*5
646.2	12	—	12	—	12	—	—	*	*	*7	*
646.23	*7	—	*7	—	*7	—	—	*	*	*6	*
646.3	*7	—	*7	—	*7	—	—	*	*	*	*
646.31	*6	—	*6	—	*6	—	—	*	*	*	*
646.6	120	—	120	*	119	*	—	14	23	60	23
646.61	44	—	44	—	44	*	—	*5	*9	21	10
646.62	*6	—	*6	—	*6	—	—	*	*	*	*
646.63	63	—	63	*	62	*	—	*7	13	34	9
646.64	*7	—	*7	—	*7	*	—	*	*	*	*
646.7	*7	—	*7	*	*7	—	—	*	*	*	*
646.8	63	—	63	*	63	—	—	*7	11	28	17
646.81	36	—	36	*	36	—	—	*	*7	13	12
646.83	23	—	23	—	23	—	—	*	*	15	*
647	91	—	91	*	91	*	—	11	17	52	11
647.6	49	—	49	*	49	*	—	*6	*8	28	*7
647.61	44	—	44	*	43	*	—	*5	*7	24	*7
647.63	*5	—	*5	—	*5	*	—	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
647.8	30	—	30	—	30	—	—	*	*7	17	*
647.81	20	—	20	—	20	—	—	*	*6	11	*
647.83	*8	—	*8	—	*8	—	—	*	*	*	*
648	1,826	—	1,826	*	1,818	*6	—	273	383	745	426
648.0	50	—	50	—	50	*	—	*9	9	24	*9
648.01	33	—	33	—	33	*	—	*6	*6	13	*7
648.03	16	—	16	—	16	—	—	*	*	10	*
648.1	81	—	81	—	80	*	—	15	17	31	19
648.11	71	—	71	—	71	*	—	14	14	27	16
648.13	*8	—	*8	—	*8	*	—	*	*	*	*
648.2	308	—	308	*	305	*	—	47	59	125	76
648.21	185	—	185	*	184	*	—	24	31	82	48
648.22	90	—	90	*	90	*	—	17	22	28	23
648.23	25	—	25	*	24	—	—	*	*	12	*
648.24	*7	—	*7	—	*7	*	—	*	*	*	*
648.3	*9	—	*9	—	*9	—	—	*	*	*	*
648.4	227	—	227	*	226	*	—	28	60	84	55
648.41	181	—	181	*	180	*	—	20	47	65	48
648.43	34	—	34	*	34	*	—	*6	9	15	*
648.44	*6	—	*6	—	*6	—	—	*	*	*	*
648.6	41	—	41	—	40	*	—	*8	*8	19	*6
648.61	32	—	32	—	31	*	—	*7	*6	14	*
648.63	*5	—	*5	—	*5	—	—	*	*	*	*
648.7	12	—	12	—	12	—	—	*	*	*6	*
648.71	9	—	9	—	9	—	—	*	*	*	*
648.8	198	—	198	—	197	*	—	29	45	68	55
648.81	181	—	181	—	180	*	—	27	42	63	49
648.83	15	—	15	—	15	*	—	*	*	*	*6
648.9	897	—	897	*	895	*	—	132	181	384	200
648.91	769	—	769	*	768	*	—	112	157	326	174
648.92	12	—	12	—	12	—	—	*	*	*	*
648.93	109	—	109	*	109	*	—	16	19	51	22
648.94	*6	—	*6	—	*6	—	—	*	*	*	*
650	375	—	375	—	375	*	—	37	66	179	93
651	65	—	65	*	64	*	—	14	10	23	18
651.0	60	—	60	*	59	*	—	12	10	21	17
651.01	41	—	41	*	41	*	—	*7	*7	15	12
651.03	19	—	19	—	18	*	—	*5	*	*6	*
652	297	—	297	*	296	*	—	49	57	118	73
652.1	*8	—	*8	—	*8	*	—	*	*	*6	*
652.11	*8	—	*8	—	*8	*	—	*	*	*6	*
652.2	125	—	125	—	124	*	—	18	24	49	34
652.21	121	—	121	—	120	*	—	18	24	47	32
652.3	21	—	21	—	21	*	—	*	*	*6	*6
652.31	21	—	21	—	21	*	—	*	*	*6	*6
652.5	60	—	60	*	60	*	—	*8	12	27	12
652.51	60	—	60	*	60	*	—	*8	12	27	12
652.6	22	—	22	—	21	*	—	*	*	*9	*5
652.61	20	—	20	—	20	*	—	*	*	*8	*
652.8	49	—	49	—	49	*	—	12	9	17	10
652.81	48	—	48	—	48	*	—	12	9	17	10
653	104	—	104	*	104	*	—	13	22	40	29
653.4	85	—	85	*	85	—	—	*8	19	34	23
653.41	85	—	85	*	85	—	—	*8	19	34	23
653.5	13	—	13	*	13	*	—	*	*	*	*
653.51	13	—	13	*	13	*	—	*	*	*	*
654	717	—	717	*	712	*	—	111	127	311	168
654.0	12	—	12	—	12	—	—	*	*	*5	*
654.01	10	—	10	—	10	—	—	*	*	*5	*
654.1	40	—	40	—	40	*	—	*6	*6	19	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
654.11	33	—	33	—	33	*	—	*	*5	16	*8
654.2	574	—	574	*	570	*	—	83	101	246	143
654.21	559	—	559	*	555	*	—	81	99	238	141
654.23	15	—	15	—	15	—	—	*	*	*8	*
654.4	32	—	32	*	32	*	—	*7	*7	13	*5
654.41	26	—	26	—	26	*	—	*6	*6	10	*
654.43	*6	—	*6	*	*6	—	—	*	*	*	*
654.5	29	—	29	—	29	*	—	*9	*5	11	*
654.51	16	—	16	—	16	*	—	*	*	*6	*
654.53	13	—	13	—	13	—	—	*	*	*5	*
654.6	21	—	21	—	21	—	—	*	*	11	*
654.61	15	—	15	—	15	—	—	*	*	*8	*
654.63	*6	—	*6	—	*6	—	—	*	*	*	*
654.9	*5	—	*5	—	*	*	—	*	*	*	*
655	56	—	56	*	56	*	—	*9	10	26	11
655.7	21	—	21	*	21	*	—	*	*5	10	*
655.71	20	—	20	*	20	—	—	*	*	*8	*
655.8	24	—	24	—	24	*	—	*5	*	11	*5
655.81	21	—	21	—	21	*	—	*	*	9	*5
656	384	—	384	*	384	*	—	54	86	159	85
656.1	80	—	80	*	79	—	—	10	13	41	15
656.11	77	—	77	*	77	—	—	10	13	40	15
656.3	24	—	24	—	24	*	—	*	*5	13	*
656.31	24	—	24	—	24	*	—	*	*5	13	*
656.4	20	—	20	*	20	*	—	*	*	9	*
656.41	20	—	20	*	19	*	—	*	*	*9	*
656.5	68	—	68	—	67	*	—	9	13	26	18
656.51	64	—	64	—	64	*	—	9	12	25	18
656.6	94	—	94	*	94	—	—	11	27	35	21
656.61	92	—	92	*	92	—	—	11	27	34	20
656.7	21	—	21	—	21	—	—	*	9	*6	*
656.71	20	—	20	—	20	—	—	*	9	*	*
656.8	74	—	74	—	74	*	—	14	13	27	19
656.81	73	—	73	—	73	*	—	14	13	27	19
657	31	—	31	—	31	*	—	*7	*	11	9
657.0	31	—	31	—	31	*	—	*7	*	11	9
657.01	27	—	27	—	27	*	—	*6	*	9	*9
658	380	—	380	*	379	*	—	69	65	150	96
658.0	112	—	112	*	111	*	—	23	17	50	22
658.01	105	—	105	*	104	*	—	22	17	46	20
658.03	*7	—	*7	—	*7	—	—	*	*	*	*
658.1	149	—	149	*	149	*	—	26	28	57	38
658.11	142	—	142	*	141	*	—	26	26	54	36
658.13	*7	—	*7	—	*7	—	—	*	*	*	*
658.2	50	—	50	—	50	*	—	10	11	17	12
658.21	47	—	47	—	47	*	—	*9	11	15	12
658.4	66	—	66	—	66	*	—	9	*9	27	22
658.41	65	—	65	—	65	*	—	9	*9	26	21
659	1,020	—	1,020	*	1,008	*8	—	150	191	392	286
659.1	45	—	45	*	44	*	—	*6	*7	20	12
659.11	39	—	39	*	38	*	—	*6	*6	18	*8
659.13	*6	—	*6	—	*6	—	—	*	*	*	*
659.2	42	—	42	*	42	*	—	*7	*7	18	10
659.21	41	—	41	*	41	*	—	*7	*7	17	10
659.4	18	—	18	—	18	*	—	*	*	*8	*
659.41	17	—	17	—	17	*	—	*	*	*8	*
659.5	50	—	50	—	49	*	—	11	9	18	12
659.51	47	—	47	—	46	*	—	10	9	16	11
659.6	295	—	295	—	289	*5	—	49	57	98	91
659.61	281	—	281	—	275	*5	—	47	54	93	87

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
659.63	14	—	14	—	14	*	—	*	*	*	*
659.7	494	—	494	*	493	*	—	68	95	194	137
659.71	488	—	488	*	488	*	—	67	94	191	137
659.73	*6	—	*6	—	*6	*	—	*	*	*	*
659.8	55	—	55	*	52	—	—	*7	*9	28	12
659.81	54	—	54	*	51	—	—	*7	*8	27	11
659.9	17	—	17	*	16	*	—	*	*	*8	*8
659.91	16	—	16	*	15	*	—	*	*	*8	*7
660	222	—	222	*	222	*	—	26	49	77	70
660.0	44	—	44	—	43	*	—	*8	*7	14	14
660.01	44	—	44	—	43	*	—	*8	*7	14	14
660.1	40	—	40	*	40	—	—	*	*8	12	15
660.11	40	—	40	*	40	—	—	*	*8	12	15
660.2	*6	—	*6	—	*6	—	—	*	*	*	*
660.21	*6	—	*6	—	*6	—	—	*	*	*	*
660.3	54	—	54	*	54	—	—	*6	11	21	15
660.31	54	—	54	*	54	—	—	*6	11	21	15
660.4	61	—	61	—	61	—	—	*5	16	20	19
660.41	61	—	61	—	61	—	—	*5	16	20	19
660.6	*6	—	*6	—	*6	—	—	*	*	*	*
660.61	*6	—	*6	—	*6	—	—	*	*	*	*
660.7	9	—	9	—	9	—	—	*	*	*	*
660.71	9	—	9	—	9	—	—	*	*	*	*
661	351	—	351	*	350	*	—	59	66	137	89
661.0	88	—	88	*	88	*	—	13	21	35	20
661.01	88	—	88	*	88	*	—	13	21	34	20
661.1	82	—	82	*	82	*	—	18	11	38	16
661.11	82	—	82	*	82	*	—	18	11	38	16
661.2	82	—	82	*	81	*	—	13	17	33	19
661.21	81	—	81	*	80	*	—	13	17	33	18
661.3	86	—	86	*	86	*	—	13	16	26	30
661.31	86	—	86	*	86	*	—	13	16	26	30
661.4	*6	—	*6	—	*6	—	—	*	*	*	*
661.41	*6	—	*6	—	*6	—	—	*	*	*	*
661.9	*7	—	*7	—	*7	—	—	*	*	*	*
661.91	*7	—	*7	—	*7	—	—	*	*	*	*
662	35	—	35	—	35	—	—	*	11	*7	13
662.0	*5	—	*5	—	*5	—	—	*	*	*	*
662.01	*5	—	*5	—	*5	—	—	*	*	*	*
662.1	9	—	9	—	9	—	—	*	*	*	*
662.11	9	—	9	—	9	—	—	*	*	*	*
662.2	20	—	20	—	20	—	—	*	*	*	*8
662.21	20	—	20	—	20	—	—	*	*	*	*8
663	831	—	831	*	829	*	—	124	185	294	227
663.0	11	—	11	—	11	—	—	*	*	*	*5
663.01	11	—	11	—	11	—	—	*	*	*	*5
663.1	136	—	136	*	136	—	—	19	33	42	42
663.11	136	—	136	*	136	—	—	19	33	42	42
663.2	48	—	48	—	48	—	—	*6	10	17	14
663.21	48	—	48	—	48	—	—	*6	10	17	14
663.3	591	—	591	*	589	*	—	91	132	215	152
663.31	591	—	591	*	588	*	—	91	132	215	152
663.4	15	—	15	*	15	—	—	*	*	*	*
663.41	15	—	15	*	15	—	—	*	*	*	*
663.8	27	—	27	—	27	—	—	*	*	10	9
663.81	27	—	27	—	27	—	—	*	*	10	9
664	1,238	—	1,238	*	1,236	*	—	198	252	422	366
664.0	546	—	546	*	545	*	—	87	103	195	161
664.01	546	—	546	*	545	*	—	87	103	195	161
664.1	550	—	550	*	549	*	—	92	121	171	166

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
664.11	550	—	550	*	549	*	—	91	121	171	166
664.2	89	—	89	*	89	—	—	14	18	30	27
664.21	89	—	89	*	88	—	—	14	18	30	27
664.3	27	—	27	*	27	*	—	*	*5	15	*
664.31	27	—	27	*	27	*	—	*	*5	15	*
664.4	10	—	10	—	10	—	—	*	*	*6	*
664.41	10	—	10	—	10	—	—	*	*	*6	*
664.8	*6	—	*6	—	*6	—	—	*	*	*	*
664.81	*6	—	*6	—	*6	—	—	*	*	*	*
664.9	*8	—	*8	—	*8	—	—	*	—	*	*
664.91	*8	—	*8	—	*8	—	—	*	—	*	*
665	181	—	181	*	180	*	—	27	53	67	35
665.3	*9	—	*9	—	*9	—	—	*	*	*	*
665.31	*9	—	*9	—	*9	—	—	*	*	*	*
665.4	58	—	58	*	58	*	—	*8	15	26	9
665.41	58	—	58	*	58	*	—	*8	15	26	9
665.5	102	—	102	*	102	*	—	17	31	33	21
665.51	102	—	102	*	101	*	—	17	31	33	21
666	113	—	113	*	113	*	—	15	27	40	31
666.0	11	—	11	—	11	—	—	*	*	*	*
666.02	10	—	10	—	10	—	—	*	*	*	*
666.1	77	—	77	*	77	*	—	10	20	26	22
666.12	77	—	77	*	77	*	—	10	19	26	22
666.2	22	—	22	—	22	—	—	*	*	10	*
666.22	16	—	16	—	16	—	—	*	*	*7	*
666.24	*6	—	*6	—	*6	—	—	*	*	*	*
667	24	—	24	—	24	—	—	*	*7	*7	*7
667.0	20	—	20	—	20	—	—	*	*6	*5	*
667.02	19	—	19	—	19	—	—	*	*6	*5	*
668	15	—	15	—	15	—	—	*	*5	*	*
668.8	14	—	14	—	14	—	—	*	*5	*	*
668.81	*6	—	*6	—	*6	—	—	*	*	*	*
668.82	*9	—	*9	—	*9	—	—	*	*	*	*
669	153	—	153	*	153	—	—	17	26	71	40
669.4	*6	—	*6	—	*6	—	—	*	*	*	*
669.5	76	—	76	*	76	—	—	*6	*9	42	19
669.51	76	—	76	*	75	—	—	*6	*9	42	19
669.7	12	—	12	—	12	—	—	*	*	*7	*
669.71	12	—	12	—	12	—	—	*	*	*7	*
669.8	52	—	52	*	52	—	—	*7	10	17	17
669.81	49	—	49	*	49	—	—	*6	9	17	17
670	25	—	25	*	25	—	—	*	*	12	*5
670.0	25	—	25	*	25	—	—	*	*	12	*5
670.02	14	—	14	*	14	—	—	*	*	*7	*
670.04	11	—	11	—	11	—	—	*	*	*6	*
671	27	—	27	*	27	—	—	*	*	11	*7
671.8	15	—	15	*	15	—	—	*	*	*6	*
671.81	*8	—	*8	—	*8	—	—	*	*	*	*
671.82	*6	—	*6	*	*6	—	—	*	*	*	*
672	22	—	22	*	22	—	—	*	*	*8	*7
672.0	22	—	22	*	22	—	—	*	*	*8	*7
672.02	21	—	21	*	21	—	—	*	*	*7	*7
674	51	—	51	*	50	*	—	*8	10	21	11
674.3	17	—	17	—	17	*	—	*	*	*6	*
674.32	13	—	13	—	13	*	—	*	*	*	*
674.8	25	—	25	*	25	—	—	*	*	11	*6
674.82	15	—	15	*	15	—	—	*	*	*7	*
674.84	10	—	10	—	10	—	—	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
676	10	—	10	*	10	—	—	*	*	*7	*
680-709	2,511	1,228	1,284	210	520	719	1,063	556	572	962	422
680	15	*9	*6	*	*8	*	*	*	*	*9	*
681	70	40	31	*8	18	24	21	16	17	26	12
681.0	26	14	12	*5	10	*6	*	*6	*7	*8	*
681.00	21	11	9	*	*8	*5	*	*	*6	*7	*
681.1	44	26	18	*	*8	17	17	10	10	17	*8
681.10	42	24	18	*	*8	16	16	9	9	16	*7
682	1,114	556	558	67	269	349	429	244	245	435	190
682.0	61	27	35	14	23	14	10	11	11	28	11
682.1	19	10	*9	*	*8	*	*	*	*	*9	*5
682.2	142	57	85	*6	50	48	38	25	31	60	26
682.3	125	69	56	*	39	38	45	29	23	49	23
682.4	47	27	19	*	17	18	10	*9	10	18	10
682.5	36	20	16	*9	13	*9	*5	*	*	21	*7
682.6	493	244	249	16	90	148	239	121	117	179	75
682.7	160	87	72	*8	21	62	68	38	40	54	27
682.8	15	*7	*7	*	*	*	*	*	*	*8	*
682.9	16	*7	10	*	*	*	*7	*	*	*8	*
683	*7	*	*	*	*	*	*	*	*	*	*
684	*9	*	*	*	*	*	*	*	*	*	*
685	*6	*	*	*	*	*	*	*	*	*	*
686	17	9	*8	*	*5	*	*5	*	*	*7	*
686.9	10	*	*5	*	*	*	*	*	*	*	*
690	12	*8	*	*	*	*	*	*	*	*	*
690.1	10	*8	*	*	*	*	*	*	*	*	*
690.10	*8	*7	*	*	*	*	*	*	*	*	*
691	34	19	15	30	*	*	*	*	*9	12	*9
691.0	25	13	12	24	*	*	*	*	*7	*8	*7
691.8	*9	*6	*	*5	*	*	*	*	*	*	*
692	66	28	38	18	15	13	20	15	17	22	13
692.9	56	23	32	16	13	10	16	12	14	17	12
693	39	15	24	*5	10	10	13	12	*7	13	*7
693.0	37	14	23	*	10	10	13	11	*7	13	*7
694	*7	*	*	*	*	*	*	*	*	*	*
695	34	10	24	*	*7	14	9	11	*8	10	*
695.1	*7	*	*	*	*	*	*	*	*	*	*
695.3	9	*	*7	—	*	*	*	*	*	*	*
695.4	*7	*	*5	—	*	*	*	*	*	*	*
695.8	*6	*	*	*	*	*	*	*	*	*	*
695.89	*6	*	*	*	*	*	*	*	*	*	*
696	51	25	25	*	10	19	21	13	16	13	10
696.0	11	*6	*	*	*	*6	*	*	*	*	*
696.1	40	19	21	*	*9	14	17	10	13	10	*7
698	35	10	25	*	17	*9	*7	*6	9	13	*6
698.9	27	*9	18	*	13	*7	*	*	*7	11	*
701	34	12	22	*8	11	*9	*6	*7	*	19	*
701.4	12	*5	*7	*7	*	*	*	*	*	9	*
701.9	*7	*	*6	*	*	*	*	*	*	*	*
702	*6	*	*	*	*	*	*	*	*	*	*
703	14	*6	*8	*	*	*	*7	*	*	*6	*
703.0	*8	*	*	*	*	*	*	*	*	*	*
704	16	*9	*8	*	*8	*	*	*	*	*7	*
704.0	*7	*	*	*	*	*	*	*	*	*	*
704.8	*7	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
705	10	*	*6	*	*	*	*	*	*	*6	*
705.8	*6	*	*	*	*	*	*	*	*	*	*
705.83	*6	*	*	*	*	*	*	*	*	*	*
706	24	14	11	*	13	*	*	*7	*5	*9	*
706.1	12	*7	*5	*	*9	*	*	*	*	*	*
706.2	10	*5	*	*	*	*	*	*	*	*	*
707	763	381	382	*	69	210	480	182	183	270	128
707.0	360	170	190	*	30	74	253	72	84	137	67
707.1	372	199	173	*	32	126	213	104	94	119	55
707.10	39	20	19	*	*	16	20	11	9	14	*5
707.12	26	11	15	—	*	*8	16	*8	11	*6	*
707.13	27	14	13	*	*	*6	19	*7	*7	10	*
707.14	51	28	23	*	*	11	37	13	14	15	*9
707.15	177	99	78	*	15	73	89	51	39	59	29
707.19	46	25	21	*	*	11	31	13	13	14	*
707.8	24	*9	15	*	*6	*9	*9	*6	*5	*9	*
707.9	*7	*	*	*	*	*	*	*	*	*	*
708	27	*9	18	*	*8	*7	*7	*	10	*8	*
708.0	12	*	*8	*	*	*	*	*	*	*	*
708.9	11	*	*7	*	*	*	*5	*	*	*	*
709	97	47	50	36	31	19	11	11	13	63	10
709.0	*6	*	*	*	*	*	*	*	*	*	*
709.2	61	30	31	32	21	*5	*	*	*	50	*
709.8	14	*7	*7	*	*	*5	*	*	*	*	*
709.9	10	*5	*	*	*	*	*	*	*	*	*
710-739	6,868	2,363	4,504	88	1,012	2,047	3,721	1,432	1,804	2,368	1,265
710	192	24	168	*	65	63	60	40	38	83	32
710.0	137	16	121	*	54	45	36	25	27	65	19
710.1	23	*	21	*	*	*7	11	*5	*	*7	*6
710.2	13	*	12	*	*	*	*7	*	*	*5	*
710.4	*6	*	*	—	*	*	*	*	*	*	*
710.9	*5	*	*	*	*	*	*	*	*	*	*
711	51	32	19	*5	*9	19	18	11	11	18	11
711.0	46	29	17	*	*8	18	16	10	9	17	10
711.06	23	13	10	*	*	11	*9	*	*	*9	*5
711.07	*7	*	*	*	*	*	*	*	*	*	*
712	*5	*	*	—	*	*	*	*	*	*	*
713	*9	*	*	*	*	*6	*	*	*	*	*
713.5	*7	*	*	—	*	*6	*	*	*	*	*
714	257	62	196	*	27	79	148	57	67	91	42
714.0	237	57	180	*	24	75	139	53	63	83	39
714.3	*8	*	*6	*	*	*	*	*	*	*	*
714.30	*7	*	*6	*	*	*	*	*	*	*	*
714.8	*6	*	*5	—	*	*	*	*	*	*	*
715	1,822	591	1,231	*	74	479	1,268	404	524	615	279
715.0	55	16	39	—	*	*6	48	13	20	16	*6
715.00	*7	*	*	—	*	*	*7	*	*	*	*
715.09	47	13	34	—	*	*6	41	12	19	12	*5
715.1	67	29	38	—	*	20	43	20	11	25	10
715.15	16	*6	10	—	*	*	10	*7	*	*	*
715.16	43	17	26	—	*	14	28	11	*6	19	*8
715.2	10	*	*5	—	*	*5	*	*	*	*	*
715.3	502	187	314	—	19	171	312	121	140	145	96
715.30	*5	*	*	—	*	*	*	*	*	*	*
715.31	20	*9	11	—	*	*5	15	*	*5	*6	*6
715.35	132	57	75	—	*5	42	84	34	36	36	26
715.36	327	111	216	—	10	116	201	78	92	98	59
715.37	*7	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
715.38	*5	*	*	—	*	*	*	*	*	*	*
715.8	71	18	53	—	*	12	56	*7	26	28	10
715.89	71	18	53	—	*	12	56	*7	26	28	10
715.9	1,117	336	781	*	46	264	807	238	326	398	154
715.90	742	209	534	*	29	144	569	158	215	277	92
715.91	19	*9	10	—	*	*	14	*	*6	*5	*
715.94	*6	*	*	—	*	*	*	*	*	*	*
715.95	104	41	63	—	*7	34	63	26	29	31	18
715.96	220	69	151	—	*6	74	140	41	71	74	34
715.97	*6	*	*	—	*	*	*	*	*	*	*
715.98	19	*	14	—	*	*	15	*7	*	*7	*
716	318	109	210	*	24	99	193	53	86	133	46
716.1	*8	*	*	*	*	*	*	*	*	*	*
716.5	20	*5	14	*	*	*5	13	*	*	11	*
716.59	12	*	*9	*	*	*	*8	*	*	*8	*
716.9	285	96	188	*	21	88	176	46	80	117	42
716.90	213	70	143	*	13	68	132	36	61	89	28
716.91	*7	*	*	*	*	*	*	*	*	*	*
716.95	9	*	*6	*	*	*	*7	*	*	*	*
716.96	22	*8	14	*	*	*7	11	*	*6	*8	*
716.98	10	*	*6	—	*	*	*6	*	*	*	*
716.99	13	*	9	*	*	*	*8	*	*	*7	*
717	28	18	10	*	15	*9	*	*6	*	11	*6
717.7	*7	*	*	—	*	*	*	*	*	*	*
717.8	*6	*5	*	*	*6	*	*	*	*	*	*
717.83	*5	*	*	*	*5	*	—	*	*	*	*
718	42	19	23	*	*8	14	17	*7	*7	19	*9
718.3	*6	*	*	*	*	*	*	*	*	*	*
718.4	18	*9	10	*	*	*6	10	*	*	10	*
718.5	*5	*	*	*	*	*	*	*	*	*	*
719	255	100	154	*6	50	69	130	57	71	75	52
719.0	24	12	12	*	*	*7	12	*6	*7	*8	*
719.06	18	*9	10	*	*	*6	10	*	*5	*6	*
719.1	*7	*	*	*	*	*	*	*	*	*	*
719.16	*5	*	*	*	*	*	*	*	*	*	*
719.4	165	57	108	*	39	47	75	26	45	57	37
719.41	52	20	32	*	12	20	20	*9	13	21	10
719.45	51	17	35	*	*6	11	33	*6	15	16	14
719.46	30	10	20	*	10	*9	10	*6	*7	10	*7
719.47	10	*	*8	*	*	*	*	*	*	*	*
719.49	9	*	*7	*	*	*	*	*	*	*	*
719.7	47	22	25	*	*	10	33	22	16	*	*6
720	12	*9	*	*	*	*6	*	*	*5	*	*
720.0	9	*8	*	—	*	*6	*	*	*	*	*
721	287	116	171	*	34	95	158	49	86	101	52
721.0	75	35	40	—	11	33	31	11	19	31	14
721.1	30	18	12	*	*	13	13	*6	*5	13	*6
721.3	118	43	75	*	12	31	74	20	41	35	22
721.8	*7	*	*	—	*	*	*	*	*	*	*
721.9	45	13	33	—	*	11	31	*8	15	15	*7
721.90	45	12	33	—	*	11	31	*8	15	15	*7
722	623	290	333	*	187	255	180	96	152	232	143
722.0	77	41	37	—	31	41	*6	11	16	36	14
722.1	230	115	115	*	86	85	59	42	56	78	53
722.10	225	114	111	*	85	82	57	42	55	75	53
722.11	*5	*	*	—	*	*	*	*	*	*	*
722.2	*7	*	*	—	*	*	*	*	*	*	*
722.4	42	15	28	*	*9	22	11	*	10	17	11
722.5	145	62	83	*	29	54	62	20	40	48	36
722.51	*8	*	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West	
Number of all-listed diagnoses in thousands												
722.52	137	58	79	*	28	52	57	19	37	46	35	
722.6	24	10	14	—	*6	*6	11	*	*6	13	*	
722.7	28	14	14	—	*6	15	*8	*	*7	10	*7	
722.71	19	9	9	—	*	11	*	*	*	*8	*	
722.73	*8	*	*	—	*	*	*	*	*	*	*	
722.8	32	12	21	—	*5	15	12	*	*9	13	*5	
722.83	28	10	18	—	*	12	11	*	*8	11	*	
722.9	37	19	18	*	11	15	11	*6	*6	15	11	
722.91	*8	*	*	—	*	*	*	*	*	*	*	
722.93	24	12	12	*	*8	9	*7	*	*	*9	*7	
723	98	40	58	*	26	34	34	16	24	38	20	
723.0	23	*9	14	*	*	10	12	*	*5	11	*	
723.1	47	20	28	*	19	14	13	*	14	18	11	
723.4	18	*8	10	*	*	*7	*6	*6	*	*5	*	
724	641	258	383	*	153	216	269	131	174	214	122	
724.0	173	64	109	*	*8	49	116	43	45	53	32	
724.00	31	9	22	—	*	*	26	10	11	*6	*	
724.02	140	54	86	*	*7	44	88	32	34	46	28	
724.2	208	87	121	*	65	77	66	34	58	71	44	
724.3	22	*6	16	—	*	*6	11	*8	*	*	*5	
724.4	35	13	22	*	9	11	16	*9	*8	15	*	
724.5	179	80	100	*	59	67	52	35	50	63	32	
724.8	11	*	*6	*	*	*	*	*	*	*	*	
725	49	13	36	—	—	*	45	10	16	12	11	
726	107	57	50	*	17	40	49	26	24	28	28	
726.1	40	20	20	—	*	15	22	10	*9	*8	13	
726.10	36	19	18	—	*	13	20	*9	*9	*6	12	
726.2	11	*	*7	*	*	*	*6	*	*	*	*	
726.3	20	16	*	—	*	11	*	*7	*5	*	*	
726.33	19	14	*	—	*	10	*	*6	*	*	*	
726.5	10	*	*8	*	*	*	*7	*	*	*	*	
726.6	11	*7	*	*	*	*	*	*	*	*	*	
726.65	*6	*5	*	*	*	*	*	*	*	*	*	
726.9	*5	*	*	*	*	*	*	*	*	*	*	
727	79	33	45	*	12	29	33	14	15	29	20	
727.0	30	15	14	*	*6	12	*8	*6	*	12	*7	
727.00	15	*7	*7	*	*	*6	*5	*	*	*6	*	
727.05	*7	*	*	*	*	*	*	*	*	*	*	
727.4	20	*6	14	—	*	*8	*9	*	*	*7	*6	
727.40	16	*5	11	—	*	*7	*7	*	*	*7	*5	
727.5	*9	*	*8	—	*	*	*7	*	*5	*	*	
727.51	*9	*	*8	—	*	*	*7	*	*5	*	*	
727.6	*8	*	*	—	*	*	*5	*	*	*	*	
727.61	*6	*	*	—	*	*	*	*	*	*	*	
727.8	*8	*6	*	*	*	*	*	*	*	*	*	
728	185	92	93	9	40	56	80	37	40	69	38	
728.2	*8	*	*	*	*	*	*	*	*	*	*	
728.3	10	*	*6	*	*	*	*	*	*	*6	*	
728.8	152	78	74	*	35	46	68	29	33	57	32	
728.85	24	10	14	*	*7	9	*6	*	*6	9	*	
728.86	13	*8	*5	—	*	*7	*	*	*	*6	*	
728.87	16	*8	*8	*	*	*	11	*	*	*6	*	
728.88	85	47	38	*	19	21	44	18	17	31	19	
728.89	12	*6	*6	*	*	*	*	*	*	*	*	
729	395	119	276	*7	93	153	143	63	122	137	73	
729.1	164	18	146	*	45	84	33	27	51	54	32	
729.2	9	*	*6	*	*	*	*	*	*	*	*	
729.3	*8	*	*6	*	*	*	*	*	*	*	*	
729.39	*7	*	*	*	*	*	*	*	*	*	*	
729.5	87	36	52	*	21	29	35	14	31	28	15	

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
729.8	117	54	63	*	17	29	69	17	33	45	22
729.81	24	12	12	*	*6	*6	11	*	*8	*8	*
729.82	13	*	9	—	*	*	*7	*	*	*6	*
729.89	81	39	42	*	*9	19	52	10	23	31	16
730	154	94	59	*7	31	60	56	35	34	53	32
730.0	35	20	15	*	*	13	15	*7	*8	10	10
730.07	22	14	*8	*	*	10	10	*	*6	*6	*5
730.1	28	19	10	*	9	*8	10	*8	*	10	*6
730.15	*5	*	*	*	*	*	*	*	*	*	*
730.17	14	9	*	*	*	*	*7	*6	*	*	*
730.2	87	52	35	*	16	38	30	18	22	32	14
730.25	*7	*	*	*	*	*	*	*	*	*	*
730.26	*6	*	*	*	*	*	*	*	*	*	*
730.27	46	28	18	*	*7	21	17	11	10	17	*8
730.28	16	11	*	*	*	*9	*5	*	*	*7	*
731	46	26	19	—	*6	23	16	12	*7	16	11
731.0	*7	*	*	—	*	*	*5	*	*	*	*
731.8	38	22	16	—	*6	22	10	10	*6	13	9
732	9	*7	*	*	*	*	*	*	*	*	*
733	949	166	783	10	65	174	700	235	226	312	176
733.0	577	45	531	*	12	77	487	154	142	179	101
733.00	556	43	514	*	11	74	472	148	140	173	96
733.01	*6	*	*6	—	—	*	*6	*	*	*	*
733.09	14	*	11	*	*	*	9	*	*	*	*
733.1	164	45	120	*	10	25	127	35	38	54	37
733.13	125	31	94	*	*	17	103	26	27	41	31
733.14	13	*	9	*	*	*	*7	*	*	*	*
733.15	*7	*	*	*	*	*	*6	*	*	*	*
733.19	11	*	*7	*	*	*	*6	*	*	*	*
733.4	52	23	29	*	13	22	16	10	10	23	10
733.42	39	17	22	*	10	17	11	*7	*6	18	*8
733.6	37	11	27	*	*8	15	12	12	*8	12	*5
733.8	42	21	21	*	13	14	15	*8	*8	16	10
733.81	*9	*5	*	*	*	*	*	*	*	*	*
733.82	33	16	18	*	*9	11	13	*6	*6	13	*8
733.9	71	18	53	*	*9	21	40	15	19	25	12
733.90	56	12	45	*	*6	16	34	13	16	19	*8
733.99	*8	*	*	*	*	*	*	*	*	*	*
735	12	*	*8	*	*	*	*6	*	*	*	*
735.4	*6	*	*	—	*	*	*	*	*	*	*
736	50	25	25	*	*8	14	23	*9	18	14	10
736.4	*8	*	*6	*	*	*	*	*	*	*	*
736.42	*6	*	*	*	*	*	*	*	*	*	*
736.6	*6	*	*	*	*	*	*	*	*	*	*
736.7	21	12	10	*	*5	*6	9	*	*8	*	*
736.79	18	10	*8	*	*	*5	*9	*	*7	*	*
736.8	*7	*	*	*	*	*	*	*	*	*	*
737	91	23	68	10	23	16	42	16	23	33	19
737.1	18	*	14	*	*	*	13	*	*8	*6	*
737.10	15	*	12	*	*	*	11	*	*7	*	*
737.3	67	18	49	9	20	12	25	12	13	26	16
737.30	57	14	43	*8	16	11	21	10	11	22	13
737.39	9	*	*	*	*	*	*	*	*	*	*
738	71	26	45	*	12	24	33	12	16	21	22
738.1	*5	*	*	*	*	*	*	*	*	*	*
738.19	*5	*	*	*	*	*	*	*	*	*	*
738.4	59	20	39	*	*6	21	32	9	13	17	19
739	30	*	26	*	21	*	*6	22	*7	*	—
739.2	*5	*	*	*	*	*	*	*	*	*	—

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
739.3	17	*	15	*	14	*	*	15	*	*	—
740-759	730	374	356	427	138	97	67	174	134	250	173
741	29	13	16	12	12	*	*	*7	*7	10	*
741.0	11	*5	*5	*6	*	*	*	*	*	*	*
741.00	*7	*	*	*	*	*	*	*	*	*	*
741.9	19	*7	11	*6	*9	*	*	*	*	*7	*
741.90	16	*7	10	*5	*8	*	*	*	*	*6	*
742	37	18	18	29	*	*	*	*8	11	*8	*9
742.1	*6	*	*	*	*	*	—	*	*	*	*
742.2	*6	*	*	*5	*	—	*	*	*	*	*
742.3	13	*7	*6	11	*	*	*	*	*	*	*
742.5	*6	*	*	*	*	*	*	*	*	*	*
742.59	*5	*	*	*	*	*	*	*	*	*	*
744	10	*6	*	*9	*	*	—	*	*	*	*
745	107	56	51	77	17	10	*	28	15	36	28
745.1	*7	*	*	*6	*	*	*	*	*	*	*
745.2	12	*8	*	10	*	*	*	*	*	*5	*
745.4	30	14	16	23	*5	*	*	*7	*	10	*8
745.5	51	26	25	32	*8	*8	*	13	*7	17	14
746	63	33	30	36	15	*8	*	16	11	21	16
746.0	10	*	*	*7	*	*	*	*	*	*	*
746.02	*7	*	*	*5	*	*	*	*	*	*	*
746.4	15	*9	*6	*	*	*	*	*	*	*	*5
746.8	19	9	10	10	*	*	*	*	*	9	*
746.89	9	*5	*	*6	*	*	*	*	*	*	*
746.9	*5	*	*	*	*	*	*	*	*	*	*
747	78	39	40	52	*7	9	10	20	15	24	19
747.0	24	14	10	23	*	*	*	*	*	*8	*8
747.1	*6	*	*	*5	*	*	*	*	*	*	*
747.10	*5	*	*	*	*	*	*	*	*	*	*
747.3	13	*7	*7	10	*	*	*	*	*	*	*
747.4	*5	*	*	*5	*	*	*	*	*	*	*
747.6	11	*	*7	*	*	*	*6	*	*	*	*
747.61	*7	*	*	*	*	*	*	*	*	*	*
747.8	14	*	10	*	*	*6	*	*	*6	*	*
747.81	10	*	*7	*	*	*6	*	*	*	*	*
748	19	11	*7	17	*	*	*	*5	*	*6	*
748.3	12	*8	*	11	*	*	*	*	*	*	*
749	19	11	*8	16	*	*	*	*	*	*7	*6
749.0	*9	*	*	*7	*	—	*	*	*	*	*
749.00	*8	*	*	*6	*	—	*	*	*	*	*
749.2	*9	*6	*	*8	*	*	*	*	*	*	*
750	22	17	*	18	*	*	*	*	*	*7	*6
750.5	15	13	*	15	—	*	—	*	*	*5	*
751	34	19	15	17	*6	*5	*	*7	*7	12	*8
751.0	*7	*	*	*	*	*	*	*	*	*	*
751.3	*5	*	*	*	*	*	—	*	*	*	*
751.5	*6	*	*	*	*	*	*	*	*	*	*
751.6	*7	*	*	*	*	*	*	*	*	*	*
752	31	13	17	14	13	*	*	*	*5	12	*8
752.1	*7	—	*7	*	*	*	—	*	*	*	*
752.11	*7	—	*7	*	*	*	—	*	*	*	*
752.3	*7	—	*7	*	*6	*	*	*	*	*	*
752.5	*5	*5	—	*	*	*	*	*	*	*	*
752.6	*8	*8	—	*7	*	*	—	*	*	*	*
752.61	*5	*5	—	*	*	*	—	*	*	*	*
753	68	33	35	20	13	21	14	15	12	25	17
753.0	*9	*	*5	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
753.1	38	19	19	*	*7	18	9	*8	*7	15	*8
753.10	13	*6	*6	*	*	*6	*5	*	*	*6	*
753.12	11	*	*6	*	*	*6	*	*	*	*	*
753.13	*6	*	*	*	*	*	*	*	*	*	*
753.2	*7	*	*	*6	*	*	*	*	*	*	*
753.3	*8	*	*	*	*	*	*	*	*	*	*
754	27	16	11	18	*	*	*	*8	*	10	*6
754.2	*8	*	*	*6	*	*	*	*	*	*5	*
754.8	*7	*	*	*	*	*	*	*	*	*	*
754.81	*5	*	*	*	*	*	*	*	*	*	*
755	18	*8	10	*8	*5	*	*	*6	*	*	*
755.6	12	*	*7	*	*	*	*	*	*	*	*
755.63	*8	*	*6	*	*	*	*	*	*	*	*
756	64	25	39	27	13	13	11	13	15	24	13
756.0	9	*	*	*7	*	*	*	*	*	*	*
756.1	29	10	19	*	*	9	11	*5	*8	11	*
756.12	18	*	15	*	*	*7	*9	*	*	*8	*
756.5	*6	*	*	*	*	*	*	*	*	*	*
756.8	*8	*	*5	*	*	*	*	*	*	*	*
757	18	9	*9	10	*	*	*	*	*	12	*
757.3	16	*9	*8	10	*	*	*	*	*	11	*
757.32	10	*	*5	*8	*	—	—	*	*	*9	*
757.39	*5	*	*	*	*	*	*	*	*	*	*
758	44	20	23	22	*9	10	*	14	*7	15	*8
758.0	31	15	16	15	*7	*8	*	11	*6	10	*
759	36	23	13	21	*9	*	*	*7	9	10	10
759.8	22	14	*9	13	*7	*	*	*	*5	*7	*6
759.89	14	*9	*	10	*	*	*	*	*	*	*
760-779	677	381	296	669	*	*	*	80	127	248	222
760	*9	*	*	*7	*	—	—	*	*	*	*
760.7	*7	*	*	*5	*	—	—	*	*	*	*
764	*5	*	*	*5	—	*	—	*	*	*	*
765	141	79	62	140	*	*	*	13	28	57	44
765.0	*8	*	*	*8	—	—	*	*	*	*	*
765.1	66	37	29	66	*	*	—	*5	13	27	21
765.16	*8	*	*	*8	—	—	—	*	*	*	*
765.17	*9	*	*5	*9	—	—	—	*	*	*	*
765.18	18	10	*8	18	*	—	—	*	*	*7	*
765.19	17	12	*6	17	—	*	—	*	*	*7	*7
765.2	68	38	30	67	*	*	*	*7	14	26	21
765.26	11	*7	*	11	—	—	—	*	*	*	*
765.27	17	*8	*9	17	*	—	—	*	*	*7	*6
765.28	22	12	10	22	*	*	—	*	*5	*9	*6
765.29	*5	*	*	*	*	—	*	*	*	*	*
767	*9	*5	*	*9	*	—	—	*	*	*	*
769	34	19	14	33	*	—	—	*	*7	12	12
770	120	72	48	119	*	*	*	16	24	44	36
770.0	*6	*	*	*6	—	—	—	*	*	*	*
770.1	*6	*	*	*6	—	—	—	*	*	*	*
770.2	*6	*	*	*6	—	—	—	*	*	*	*
770.6	15	10	*5	15	—	—	—	*	*	*6	*5
770.7	17	11	*7	17	*	*	*	*	*	*5	*
770.8	64	36	28	63	*	*	*	*7	14	22	21
770.81	24	13	11	24	*	*	*	*	*7	*7	*8
770.84	*6	*	*	*6	—	—	—	*	*	*	*
770.89	25	15	10	25	—	*	—	*	*	11	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
771	48	27	21	47	*	*	*	*6	*8	18	17
771.7	*6	*	*	*6	—	*	*	*	*	*	*
771.8	36	22	15	36	*	*	*	*	*6	12	14
771.81	27	15	12	27	*	*	*	*	*	*9	11
772	16	10	*5	15	*	—	*	*	*	*7	*5
772.1	9	*6	*	*9	*	—	—	*	*	*	*
773	11	*6	*	11	*	—	—	*	*	*	*
773.2	*6	*	*	*6	*	—	—	*	*	*	*
774	117	66	51	117	*	*	*	14	20	43	41
774.2	46	27	18	45	*	—	*	*	*7	19	16
774.3	*6	*	*	*6	—	—	—	*	*	*	*
774.39	*6	*	*	*6	—	—	—	*	*	*	*
774.6	64	34	30	63	*	*	*	10	11	22	21
775	41	24	17	40	*	—	*	*	*7	17	11
775.4	*9	*5	*	*9	—	—	—	*	*	*	*
775.5	17	10	*7	17	—	—	*	*	*	*6	*5
775.6	*7	*	*	*7	—	—	—	*	*	*	*
776	20	12	*9	20	—	—	*	*	*	10	*5
776.1	*6	*	*	*6	—	—	—	*	*	*	*
776.6	9	*6	*	9	—	—	—	*	*	*5	*
777	13	*6	*7	13	—	—	—	*	*	*	*5
777.8	*7	*	*	*7	—	—	—	*	*	*	*
778	20	11	*9	20	—	—	—	*	*	*	*8
778.4	11	*6	*	11	—	—	—	*	*	*	*
779	63	33	30	62	*	*	*	*9	15	16	22
779.0	*6	*	*	*6	—	—	—	*	*	*	*
779.3	37	19	18	37	*	*	*	*	*6	11	16
779.8	16	*9	*7	16	*	—	—	*	*5	*	*
779.81	*8	*	*	*8	—	—	—	*	*	*	*
779.89	*6	*	*	*5	*	—	—	*	*	*	*
780-799	8,836	3,862	4,974	747	1,920	2,536	3,633	1,880	2,336	3,041	1,578
780	2,664	1,215	1,450	251	556	751	1,107	644	671	905	444
780.0	114	45	68	*	28	27	54	20	24	39	31
780.01	17	*7	10	*	*	*	10	*	*	*7	*
780.02	9	*	*6	*	*	*	*	*	*	*5	*
780.09	84	33	51	*	22	18	41	17	17	25	25
780.1	18	*8	11	*	*7	*	*8	*	*	*8	*
780.2	460	211	249	*7	54	106	292	134	113	155	58
780.3	922	432	490	109	227	286	301	238	223	305	156
780.31	19	11	*7	18	*	*	*	*	*	*5	*6
780.39	904	421	483	91	227	286	300	235	219	300	150
780.4	149	48	100	*	21	39	86	43	42	45	18
780.5	354	188	166	18	96	152	89	92	91	111	60
780.50	*5	*	*	*	*	*	*	*	*	*	*
780.52	43	18	25	*	13	11	18	*8	13	14	*8
780.57	294	163	131	16	77	135	67	78	74	92	50
780.59	*5	*	*	*	*	*	*	*	*	*	*
780.6	322	151	171	98	71	65	87	53	75	131	63
780.7	207	78	129	*6	26	44	131	36	68	68	35
780.79	203	77	126	*6	24	42	131	35	68	66	33
780.8	9	*	*	*	*	*	*	*	*	*	*
780.9	110	49	61	*7	22	24	56	22	28	41	19
780.93	15	*6	*9	*	*7	*	*6	*	*6	*	*
780.99	93	43	50	*	16	23	50	20	21	36	16
781	294	115	179	17	28	66	183	69	81	91	54
781.0	37	18	19	*5	*7	*9	16	*6	14	10	*7
781.2	168	55	113	*	*9	33	123	50	42	52	24
781.3	56	26	30	*6	*9	15	26	*8	15	18	15

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
781.9	27	13	14	*	*	*7	15	*	*7	9	*6
781.94	19	*9	10	*	*	*5	12	*	*6	*6	*
781.99	*7	*	*	*	*	*	*	*	*	*	*
782	222	99	123	27	46	66	83	42	66	68	46
782.0	58	25	32	*	18	24	14	*9	20	17	12
782.1	40	21	19	14	11	9	*7	*8	12	11	10
782.3	86	33	53	*	11	24	49	18	25	26	16
782.4	22	12	9	*	*5	*7	*8	*	*5	*8	*
782.5	*7	*	*	*5	*	*	*	*	*	*	*
783	225	95	130	79	30	25	90	49	64	71	40
783.0	38	13	24	*	*	*6	23	*6	13	15	*
783.2	60	26	34	*7	14	10	29	10	23	18	*9
783.21	59	26	33	*6	14	10	29	10	22	18	*8
783.3	17	*7	10	13	*	*	*	*	*	*	*6
783.4	58	30	28	53	*	*	*	18	14	14	12
783.40	20	11	10	17	*	*	*	*8	*5	*	*
783.41	35	18	17	34	*	*	*	10	*8	*8	*9
783.7	43	16	27	*	*	*6	35	10	9	16	*7
784	402	158	244	24	103	121	154	78	104	151	69
784.0	194	61	134	11	77	64	43	34	56	75	29
784.2	16	*7	*9	*	*	*6	*	*	*	*6	*
784.3	55	24	31	*	*	15	36	12	13	18	11
784.4	*7	*	*	*	*	*	*	*	*	*	*
784.49	*6	*	*	*	*	*	*	*	*	*	*
784.5	67	30	37	*	*7	20	38	10	16	28	13
784.7	51	26	25	*	*7	12	29	13	11	18	*9
784.9	*6	*	*	*	*	*	*	*	*	*	*
785	442	199	242	36	86	118	202	91	123	154	74
785.0	113	45	68	12	37	28	35	16	33	48	15
785.1	50	17	33	*	13	15	22	12	15	16	*7
785.2	40	16	24	10	*8	10	11	*6	14	12	*8
785.4	67	38	30	*	*	20	42	18	13	25	10
785.5	119	59	60	*	10	30	73	26	31	35	26
785.50	10	*	*5	*	*	*	*5	*	*	*	*
785.51	28	14	14	*	*	*7	20	*6	*7	10	*5
785.52	66	34	32	*	*5	17	41	15	17	18	15
785.59	15	*6	*9	*	*	*	*8	*	*	*	*
785.6	48	23	25	*7	12	13	15	11	14	16	*7
785.9	*6	*	*	*	*	*	*	*	*	*	*
786	1,596	743	852	82	303	621	590	332	368	600	296
786.0	216	97	120	62	32	49	73	31	56	78	51
786.03	24	12	11	17	*	*	*	*	*6	*6	*8
786.05	72	25	47	*	15	25	31	9	17	30	16
786.06	*6	*	*	*	*	*	*	*	*	*	*
786.07	12	*7	*5	*7	*	*	*	*	*	*5	*
786.09	98	47	50	32	12	19	35	15	26	33	23
786.1	*7	*	*	*	*	*	*	*	*	*	*
786.2	33	16	18	*8	*8	*8	10	*	10	11	*8
786.3	54	34	20	*	12	17	23	13	15	19	*8
786.5	1,237	563	674	*	244	533	455	272	273	471	221
786.50	491	231	260	*	90	212	188	98	112	196	85
786.51	33	14	19	*	*6	11	16	*6	9	10	*8
786.52	46	19	27	*	13	17	15	*7	14	18	*7
786.59	667	299	367	*	136	293	236	160	138	248	121
786.6	38	22	16	*	*	9	24	*7	11	14	*6
786.8	*8	*7	*	*	*	*	*	*	*	*	*
787	796	297	499	101	202	189	303	138	243	276	138
787.0	407	134	274	64	131	97	115	61	128	143	75
787.01	238	73	165	16	83	63	77	36	75	85	43
787.02	93	28	65	*6	32	27	28	15	33	27	18

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
787.03	76	33	43	42	16	*6	11	11	20	31	14
787.1	*7	*	*5	*	*	*	*	*	*	*	*
787.2	131	59	72	*	15	29	83	28	36	43	25
787.3	19	11	*8	*	*	*7	*6	*	*5	10	*
787.6	13	*	9	*	*	*	*	*	*	*	*
787.9	218	87	131	27	47	50	94	45	67	73	32
787.91	212	84	128	27	47	48	91	43	67	72	31
787.99	*6	*	*	*	*	*	*	*	*	*	*
788	413	203	210	10	59	91	253	93	120	124	77
788.0	*7	*	*	*	*	*	*	*	*	*	*
788.1	11	*	*8	*	*	*	*	*	*	*	*
788.2	223	141	82	*	30	43	148	54	61	64	44
788.20	183	114	69	*	25	34	122	45	52	54	32
788.29	36	24	12	*	*	*8	23	9	*8	*9	10
788.3	117	33	83	*	12	35	66	23	36	36	22
788.30	86	26	60	*	*9	24	51	17	28	27	14
788.31	*8	*	*7	—	*	*	*	*	*	*	*
788.33	10	*	10	—	*	*	*	*	*	*	*
788.39	*9	*	*6	*	*	*	*	*	*	*	*
788.4	17	*7	10	*	*	*	*9	*	*	*6	*
788.41	10	*	*7	*	*	*	*6	*	*	*	*
788.5	*8	*	*	*	*	*	*	*	*	*	*
788.6	*5	*	*	*	*	*	*	*	*	*	*
788.9	25	9	15	*	*	*	18	*6	9	*6	*
789	689	253	436	38	234	210	207	123	183	245	138
789.0	497	166	331	32	200	126	139	82	136	181	99
789.00	165	58	108	14	62	43	46	26	42	64	33
789.01	51	15	36	*	23	12	15	*8	16	17	10
789.02	12	*	*8	*	*7	*	*	*	*	*	*
789.03	68	20	48	*7	37	14	10	10	18	25	15
789.04	28	*7	21	*	*9	*7	11	*	*6	12	*
789.05	*7	*	*	*	*	*	*	*	*	*	*
789.06	65	24	41	*	20	20	23	*9	23	21	12
789.07	20	*7	13	*	*7	*6	*5	*	*	9	*
789.09	80	27	53	*	32	20	25	17	21	26	16
789.1	13	*8	*	*	*5	*	*	*	*	*	*5
789.2	14	*7	*7	*	*	*5	*	*	*	*	*
789.3	17	*	15	*	*	*	*8	*	*	*	*
789.30	*8	*	*7	—	*	*	*	*	*	*	*
789.5	145	70	75	*	18	69	54	32	37	49	26
790	428	207	221	27	89	119	193	96	129	130	74
790.2	45	16	29	*	19	13	12	*8	16	14	*8
790.22	*5	*	*	*	*	*	*	*	*	*	*
790.29	37	13	24	*	18	10	*9	*6	13	12	*7
790.4	17	10	*7	*	*8	*	*	*	*6	*	*
790.5	44	23	21	*	*7	15	20	*7	22	*9	*6
790.6	128	58	70	*7	27	37	58	26	36	48	18
790.7	108	56	52	13	19	32	44	29	23	33	23
790.9	79	40	38	*	*7	16	54	21	25	19	14
790.92	56	26	31	*	*	10	42	15	20	12	10
790.93	10	10	—	—	—	*	*8	*	*	*	*
790.99	10	*	*6	*	*	*	*	*	*	*	*
791	32	15	17	*6	12	*7	*6	*7	*7	11	*7
791.0	15	*8	*8	*	*6	*	*	*	*	*	*
791.9	11	*	*7	*	*	*	*	*	*	*	*
792	60	*6	54	*	43	*	13	15	13	18	14
792.1	20	*6	14	*	*	*	13	*9	*	*	*
792.3	40	—	40	—	40	—	—	*6	*9	13	12
793	25	12	13	*	*5	*7	11	*	*8	*7	*6
793.1	11	*7	*	*	*	*	*6	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
794	170	79	90	*	41	63	61	32	51	59	27
794.3	84	41	43	*	14	33	37	15	30	30	10
794.30	*7	*	*	—	*	*	*	*	*	*	*
794.31	54	24	30	*	10	20	23	12	18	18	*6
794.39	23	14	*9	*	*	10	10	*	10	*8	*
794.8	74	35	39	*	24	27	20	15	17	25	17
795	33	*9	24	*	20	*7	*5	10	*7	*7	9
795.5	16	*7	10	*	*9	*	*	*8	*	*	*
795.7	10	*	*8	*	*7	*	*	*	*	*	*
795.79	10	*	*8	*	*7	*	*	*	*	*	*
796	66	25	41	*	28	14	19	10	18	23	15
796.0	*7	*	*	*	*	*	*	*	*	*	*
796.2	42	17	26	*	22	9	*9	*	14	16	*8
796.4	9	*	*7	*	*	*	*	*	*	*	*
799	274	129	145	34	34	57	149	45	78	101	50
799.0	108	53	56	28	12	20	48	19	34	33	23
799.1	14	*8	*5	*	*	*	*7	*	*	*	*
799.3	68	28	39	*	*	11	54	*8	28	26	*6
799.4	42	21	20	*	*	11	26	*8	*8	19	*7
799.9	38	16	22	*	13	10	11	*5	*5	17	10
800-999	6,914	3,599	3,315	489	2,144	1,739	2,542	1,400	1,570	2,471	1,474
800	20	13	*7	9	*8	*	*	*	*	*6	*6
800.0	*7	*	*	*	*	*	*	*	*	*	*
800.2	*6	*	*	*	*	*	*	*	*	*	*
801	47	35	12	*7	26	*8	*6	9	11	12	15
801.0	19	15	*	*	*9	*	*	*5	*	*6	*
801.01	*8	*6	*	*	*	*	*	*	*	*	*
801.2	17	11	*5	*	*8	*	*	*	*	*	*7
801.26	*5	*	*	*	*	*	*	*	*	*	*
802	117	89	28	*6	68	25	19	23	22	45	28
802.0	25	17	*9	*	13	*5	*5	*	*6	*9	*6
802.2	26	24	*	*	17	*5	*	*5	*	11	*6
802.25	*6	*5	*	*	*	*	*	*	*	*	*
802.26	*5	*	*	*	*	*	*	*	*	*	*
802.3	*6	*5	*	*	*6	*	—	*	*	*	*
802.4	22	16	*6	*	12	*	*5	*5	*	*8	*
802.6	15	11	*	*	*8	*	*	*	*	*	*5
802.8	21	14	*7	*	11	*5	*	*	*	*7	*
803	*6	*	*	*	*	*	*	*	*	*	*
805	185	93	93	*	58	35	87	30	51	64	41
805.0	42	27	16	*	18	11	13	*6	*8	17	11
805.02	14	*8	*6	*	*	*	*7	*	*	*7	*
805.06	*6	*	*	—	*	*	*	*	*	*	*
805.07	*8	*	*	—	*	*	*	*	*	*	*
805.2	51	23	28	*	13	*9	27	10	16	15	11
805.4	67	33	35	*	20	12	34	10	20	24	14
805.6	18	*7	11	*	*6	*	*8	*	*5	*6	*
805.8	*5	*	*	—	*	*	*	*	*	*	*
806	12	*8	*	*	*6	*	*	*	*	*	*
806.0	*7	*5	*	*	*	*	*	*	*	*	*
807	119	73	46	*	33	34	49	24	27	38	30
807.0	103	64	39	*	27	31	43	21	24	33	25
807.00	*9	*5	*	*	*	*	*	*	*	*	*
807.01	26	17	10	*	*6	*8	12	*	*7	*8	*6
807.02	15	9	*6	*	*	*	*7	*	*	*6	*
807.03	11	*6	*5	*	*	*	*6	*	*	*	*
807.04	*9	*	*	—	*	*	*	*	*	*	*
807.09	21	14	*7	*	*5	*7	*8	*	*	*7	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
807.2	10	*	*5	—	*	*	*	*	*	*	*
808	102	39	63	*	28	17	56	17	29	37	19
808.0	27	18	9	*	10	*6	10	*	*7	11	*
808.2	58	15	44	*	12	*8	37	*9	18	21	11
808.4	9	*	*5	*	*	*	*	*	*	*	*
808.8	*7	*	*	*	*	*	*	*	*	*	*
810	16	12	*	*	*7	*	*	*	*	*	*
810.0	16	12	*	*	*7	*	*	*	*	*	*
810.00	*9	*7	*	*	*	*	*	*	*	*	*
810.02	*5	*	*	*	*	*	*	*	*	*	*
811	12	10	*	*	*	*5	*	*	*	*	*
811.0	12	10	*	*	*	*5	*	*	*	*	*
811.00	*6	*	*	—	*	*	*	*	*	*	*
812	108	39	69	19	15	21	52	22	21	40	25
812.0	60	18	43	*	*	15	40	12	12	26	10
812.00	20	*6	14	*	*	*6	12	*	*	12	*
812.01	16	*	13	—	*	*	14	*	*	*5	*
812.03	*7	*	*5	—	*	*	*	*	*	*	*
812.09	16	*6	9	*	*	*	*9	*	*	*	*
812.2	16	*7	*9	*	*	*	*7	*	*	*5	*
812.21	11	*	*7	*	*	*	*5	*	*	*	*
812.4	27	11	16	15	*	*	*	*7	*	*7	*9
812.40	*5	*	*	*	*	*	*	*	*	*	*
812.41	16	*6	9	13	*	*	*	*	*	*	*6
813	99	48	51	11	33	19	36	20	25	32	22
813.0	20	*8	12	*	*6	*	*8	*	*	*7	*
813.01	*8	*	*	*	*	*	*	*	*	*	*
813.2	*9	*	*	*	*5	*	*	*	*	*	*
813.4	51	20	31	*	13	10	25	11	15	14	11
813.41	9	*	*7	*	*	*	*5	*	*	*	*
813.42	23	10	13	*	*6	*	12	*6	*6	*7	*
813.44	16	*7	9	*	*	*	*7	*	*5	*	*
813.5	*6	*	*	*	*	*	*	*	*	*	*
814	*7	*	*	*	*	*	*	*	*	*	*
814.0	*7	*	*	*	*	*	*	*	*	*	*
815	15	11	*	*	*7	*	*	*	*	*	*
815.0	12	*8	*	*	*5	*	*	*	*	*	*
815.00	*5	*	*	*	*	*	*	*	*	*	*
816	19	13	*6	*	*7	*7	*	*	*	*7	*5
816.0	11	*7	*	*	*	*	*	*	*	*	*
816.1	*8	*6	*	*	*	*	*	*	*	*	*
820	377	112	265	*	10	36	329	77	82	147	72
820.0	89	27	62	*	*	*6	80	22	23	30	13
820.00	*5	*	*	—	*	*	*	*	*	*	*
820.02	*8	*	*5	*	*	*	*7	*	*	*	*
820.03	9	*	*5	—	*	*	*8	*	*	*	*
820.09	65	18	47	*	*	*	60	17	18	22	*7
820.2	193	54	139	*	*5	20	166	36	42	74	41
820.20	10	*	*7	*	*	*	*8	*	*	*	*
820.21	162	43	119	*	*	16	144	29	34	63	36
820.22	21	*8	13	*	*	*	14	*	*5	*8	*
820.8	90	29	61	*	*	*7	80	17	17	40	17
821	85	42	43	15	23	17	30	13	18	38	17
821.0	50	28	22	11	15	10	14	*7	9	24	10
821.00	10	*5	*5	*	*	*	*	*	*	*5	*
821.01	39	23	16	9	13	*7	10	*6	*7	18	*8
821.1	*6	*5	*	*	*5	*	*	*	*	*	*
821.2	28	*8	20	*	*	*6	16	*	*7	11	*5
821.20	*8	*	*7	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
821.23	12	*	*9	*	*	*	*7	*	*	*	*
822	27	15	11	*	9	*8	10	*	*6	12	*
822.0	24	13	11	*	*7	*7	10	*	*5	11	*
823	91	52	39	*7	33	30	21	20	20	31	20
823.0	40	19	21	*	9	16	13	*8	10	13	9
823.00	24	10	14	*	*6	*9	*9	*	*	*8	*6
823.01	*5	*	*	—	*	*	*	*	*	*	*
823.02	11	*5	*6	*	*	*6	*	*	*	*	*
823.2	22	14	*8	*	10	*6	*	*	*	*8	*5
823.20	*7	*	*	*	*	*	*	*	*	*	*
823.22	13	*7	*5	*	*6	*	*	*	*	*	*
823.3	*6	*	*	*	*	*	*	*	*	*	*
823.8	15	9	*6	*	*6	*	*	*5	*	*5	*
823.82	*8	*6	*	*	*	*	*	*	*	*	*
824	134	61	73	*6	52	43	32	29	34	47	24
824.0	11	*7	*	*	*	*	*	*	*	*	*
824.2	11	*	*7	*	*	*	*	*	*	*	*
824.4	35	16	19	*	14	11	9	*7	*8	13	*6
824.6	31	9	21	*	12	10	*8	*8	*8	9	*6
824.8	29	15	14	*	10	*9	*7	*7	*9	*8	*5
824.9	*7	*	*	*	*	*	*	*	*	*	*
825	47	29	18	*	27	13	*7	12	10	16	9
825.0	12	*7	*	*	*7	*	*	*	*	*	*
825.2	26	15	11	*	14	*7	*5	*6	*5	*8	*6
825.21	*6	*	*	—	*	*	*	*	*	*	*
825.25	14	*7	*7	*	*	*	*5	*	*	*	*
825.3	*6	*	*	*	*	*	*	*	*	*	*
826	*8	*	*	*	*	*	*	*	*	*	*
831	*9	*5	*	*	*	*	*	*	*	*	*
831.0	*9	*5	*	*	*	*	*	*	*	*	*
835	10	*8	*	*	*	*	*	*	*	*	*
835.0	10	*8	*	*	*	*	*	*	*	*	*
835.00	*7	*	*	*	*	*	*	*	*	*	*
836	16	10	*6	*	*7	*6	*	*	*	*6	*
836.0	*6	*	*	*	*	*	*	*	*	*	*
839	15	10	*	*	*8	*	*	*	*	*7	*
839.0	*7	*	*	*	*	*	*	*	*	*	*
840	34	17	17	*	*5	9	19	*8	*8	12	*5
840.4	20	*8	12	—	*	*6	13	*5	*	*8	*
840.9	*6	*	*	—	*	*	*	*	*	*	*
843	*5	*	*	*	*	*	*	*	*	*	*
844	26	16	10	*	12	*8	*5	*7	*6	9	*
844.2	*8	*	*	*	*6	*	*	*	*	*	*
844.8	*7	*6	*	*	*	*	*	*	*	*	*
844.9	*7	*	*	*	*	*	*	*	*	*	*
845	20	11	*8	*	9	*6	*	*6	*6	*	*
845.0	19	11	*8	*	9	*6	*	*6	*6	*	*
845.00	9	*	*	*	*	*	*	*	*	*	*
847	38	22	16	*	16	13	*8	*	12	9	11
847.0	22	14	*8	*	11	*5	*	*	*6	*	*9
847.2	11	*6	*5	—	*	*	*	*	*	*	*
848	*7	*	*	*	*	*	*	*	*	*	*
850	93	61	32	11	47	15	21	26	22	24	21
850.0	17	*8	*8	*	*7	*	*5	*6	*5	*	*
850.1	29	21	*8	*	16	*	*	*8	*7	*5	*8
850.11	29	21	*8	*	16	*	*	*8	*7	*5	*8
850.5	33	21	11	*	19	*6	*	*6	*7	12	*8

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
850.9	12	*8	*	*	*	*	*5	*	*	*	*
851	28	18	11	*	17	*	*6	*	*8	10	*7
851.4	*5	*	*	—	*	*	*	*	*	*	*
851.8	22	14	*8	*	12	*	*	*	*	*8	*6
852	69	37	31	*	16	13	36	15	16	21	17
852.0	23	13	10	*	*8	*6	*8	*	*6	*8	*5
852.01	*7	*	*	*	*	*	*	*	*	*	*
852.02	*6	*	*	*	*	*	*	*	*	*	*
852.2	41	22	19	*	*6	*6	27	10	9	12	10
852.20	12	*7	*5	*	*	*	*7	*	*	*	*
852.21	16	*7	*9	*	*	*	12	*	*	*	*
852.26	*6	*	*	*	*	*	*	*	*	*	*
853	13	*8	*	*	*	*	*7	*	*	*5	*
853.0	13	*8	*	*	*	*	*7	*	*	*5	*
854	12	*7	*	*	*8	*	*	*	*	*	*
854.0	12	*7	*	*	*8	*	*	*	*	*	*
860	59	44	15	*	27	19	11	13	11	17	17
860.0	35	25	9	*	15	13	*6	*7	*7	11	10
860.2	*9	*6	*	*	*	*	*	*	*	*	*
860.4	11	*8	*	*	*	*	*	*	*	*	*
861	35	23	12	*	19	*6	*6	*8	*7	9	10
861.2	28	18	10	*	15	*	*	*7	*6	*8	*8
861.21	26	17	9	*	13	*	*	*7	*6	*7	*7
862	11	*6	*	*	*5	*	*	*	*	*	*
862.2	*7	*	*	*	*	*	*	*	*	*	*
862.29	*7	*	*	*	*	*	*	*	*	*	*
863	22	16	*6	*	15	*	*	*	*	12	*
864	26	18	*7	*	16	*	*	*	*	13	*6
864.0	21	15	*7	*	13	*	*	*	*	9	*5
864.05	*7	*	*	*	*5	*	*	*	*	*	*
865	29	22	*7	*	18	*	*	*6	*6	11	*7
865.0	27	20	*7	*	16	*	*	*5	*6	9	*7
865.00	*8	*6	*	*	*5	*	*	*	*	*	*
865.03	*8	*5	*	*	*5	*	*	*	*	*	*
866	10	*9	*	*	*6	*	*	*	*	*	*
866.0	*8	*8	*	*	*5	*	*	*	*	*	*
867	19	16	*	*	*6	*	10	*	*	*8	*
867.0	16	14	*	*	*	*	9	*	*	*6	*
868	15	11	*	*	10	*	*	*	*	*7	*
868.0	11	*8	*	*	*7	*	*	*	*	*5	*
868.03	*5	*	*	*	*	*	*	*	*	*	*
870	15	13	*	*	*8	*	*	*	*	*7	*
870.8	*6	*5	*	*	*	*	*	*	*	*	*
871	*6	*	*	*	*	*	*	*	*	*	*
872	10	*9	*	*	*7	*	*	*	*	*	*
873	188	126	63	12	85	37	54	44	42	61	41
873.0	51	30	22	*	15	15	21	14	12	16	9
873.2	9	*7	*	*	*5	*	*	*	*	*	*
873.20	*8	*5	*	*	*	*	*	*	*	*	*
873.4	92	62	30	*7	42	17	26	21	20	28	23
873.40	*8	*5	*	*	*	*	*	*	*	*	*
873.41	*8	*6	*	*	*	*	*	*	*	*	*
873.42	44	29	15	*	15	10	17	*8	10	14	12
873.43	20	14	*6	*	12	*	*	*	*	*7	*
873.44	*8	*	*	*	*	*	*	*	*	*	*
873.49	*5	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
873.5	*6	*	*	*	*	*	*	*	*	*	*
873.6	19	15	*	*	14	*	*	*	*	*6	*
873.63	12	10	*	*	9	*	*	*	*	*	*
874	*8	*6	*	*	*5	*	*	*	*	*	*
874.8	*6	*	*	*	*	*	*	*	*	*	*
875	9	*9	*	*	*6	*	*	*	*	*5	*
875.0	*7	*7	*	*	*	*	*	*	*	*	*
879	19	11	*8	*	10	*	*	*	*	*7	*5
879.2	*7	*	*	*	*	*	*	*	*	*	*
880	*8	*6	*	*	*	*	*	*	*	*	*
880.0	*6	*	*	*	*	*	*	*	*	*	*
881	47	27	20	*	29	*8	*8	*8	11	14	13
881.0	32	14	17	*	19	*5	*7	*6	*8	*8	10
881.00	12	*6	*6	*	*7	*	*	*	*	*	*
881.01	*9	*5	*	*	*	*	*	*	*	*	*
881.02	11	*	*8	*	*8	*	*	*	*	*	*
881.1	*8	*6	*	*	*5	*	*	*	*	*	*
881.10	*5	*	*	*	*	*	*	*	*	*	*
881.2	*8	*7	*	*	*6	*	*	*	*	*	*
882	19	10	*9	*	10	*	*	*	*	*8	*
882.0	*8	*	*	*	*	*	*	*	*	*	*
882.1	*9	*	*	*	*	*	*	*	*	*	*
883	23	19	*	*	13	*7	*	*	*6	*9	*6
883.0	9	*7	*	*	*6	*	*	*	*	*	*
883.1	*5	*	*	*	*	*	*	*	*	*	*
883.2	*9	*8	*	*	*	*	*	*	*	*	*
884	*7	*	*	*	*	*	*	*	*	*	*
884.0	*5	*	*	*	*	*	*	*	*	*	*
886	*6	*	*	*	*	*	*	*	*	*	*
886.0	*5	*	*	*	*	*	*	*	*	*	*
890	13	10	*	*	9	*	*	*	*	*	*
890.0	*8	*6	*	*	*	*	*	*	*	*	*
891	33	23	11	*	18	*6	*6	*5	*6	14	*8
891.0	21	13	*8	*	12	*	*	*	*	10	*
891.1	11	*9	*	*	*	*	*	*	*	*	*
892	13	*6	*7	*	*	*	*	*	*	*5	*
892.1	*8	*	*	*	*	*	*	*	*	*	*
903	*8	*7	*	*	*6	*	*	*	*	*	*
905	62	34	28	*	18	20	21	12	18	15	17
905.0	*5	*	*	*	*	*	*	*	*	*	*
905.1	*9	*	*	*	*	*	*	*	*	*	*
905.2	11	*7	*	*	*	*6	*	*	*	*	*
905.3	*8	*	*6	—	*	*	*6	*	*	*	*
905.4	22	13	*9	*	*8	*6	*6	*	*	*5	*8
906	18	11	*7	*	*6	*	*	*	*	9	*
907	58	46	12	*	33	17	*7	*8	15	21	15
907.0	24	16	*8	*	13	*8	*	*	*8	*	*6
907.2	31	27	*	*	19	*7	*	*	*	15	*8
908	19	12	*6	*	*8	*6	*	*	*6	*	*5
908.9	13	*8	*5	*	*	*5	*	*	*5	*	*
909	15	*8	*7	*	*	*	*6	*	*	*	*
909.2	*7	*	*	*	*	*	*	*	*	*	*
910	34	21	13	*5	13	*7	*9	*6	11	10	*7
910.0	31	20	11	*	12	*6	*8	*6	10	9	*6

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
911	19	10	*8	*	12	*	*	*	*	*7	*
911.0	14	9	*	*	9	*	*	*	*	*	*
912	*6	*	*	*	*	*	*	*	*	*	*
912.0	*5	*	*	*	*	*	*	*	*	*	*
913	30	18	13	*	12	*6	10	*	11	11	*
913.0	22	13	*9	*	*9	*	*7	*	*8	*7	*
914	11	*6	*	*	*5	*	*	*	*	*	*
914.0	*7	*	*	*	*	*	*	*	*	*	*
916	34	21	13	*	14	*7	9	*	13	11	*6
916.0	24	15	*9	*	11	*	*7	*	10	*6	*
916.5	*6	*	*	*	*	*	*	*	*	*	*
917	*7	*	*	*	*	*	*	*	*	*	*
918	*8	*	*	*	*	*	*	*	*	*	*
918.1	*7	*	*	*	*	*	*	*	*	*	*
919	16	*9	*7	*	*8	*	*	*	*5	*	*
919.0	13	*7	*6	*	*7	*	*	*	*	*	*
920	81	39	43	10	23	13	35	21	22	22	16
921	16	*8	*8	*	*6	*	*6	*	*	*6	*
922	64	30	34	*5	21	13	25	12	16	20	15
922.1	30	15	16	*	*9	*9	12	*6	*7	9	*8
922.2	16	*9	*7	*	*7	*	*	*	*	*	*5
922.3	14	*5	*9	*	*	*	*6	*	*	*	*
922.31	*8	*	*	*	*	*	*	*	*	*	*
922.32	*6	*	*	*	*	*	*	*	*	*	*
923	41	20	21	*	13	*8	19	9	13	*8	11
923.0	15	*7	*8	*	*	*	*8	*	*	*	*
923.00	9	*	*5	*	*	*	*6	*	*	*	*
923.1	14	*8	*7	*	*7	*	*6	*	*	*	*
923.10	*6	*	*	*	*	*	*	*	*	*	*
923.11	*9	*	*	*	*	*	*	*	*	*	*
923.2	*8	*	*	*	*	*	*	*	*	*	*
923.20	*6	*	*	*	*	*	*	*	*	*	*
924	85	37	48	*	24	11	46	17	29	19	19
924.0	27	9	18	*	*6	*	18	*6	*9	*6	*6
924.00	*8	*	*	*	*	*	*	*	*	*	*
924.01	19	*5	14	*	*	*	14	*	*6	*	*
924.1	31	15	15	*	*8	*5	16	*5	10	*8	*8
924.10	15	*8	*7	*	*	*	*6	*	*	*	*
924.11	16	*8	*8	*	*	*	10	*	*	*	*
924.8	18	*8	9	*	*7	*	*8	*	*8	*	*
933	15	10	*6	*	*	*	11	*	*	*5	*
933.1	14	*9	*5	*	*	*	10	*	*	*5	*
934	12	*5	*7	*	*	*	*6	*	*	*	*
935	13	*8	*	*	*	*	*	*	*	*	*
935.1	*7	*	*	*	*	*	*	*	*	*	*
935.2	*5	*	*	*	*	*	*	*	*	*	*
941	31	20	11	15	13	*	*	*5	*6	19	*
941.2	20	13	*7	9	*9	*	*	*	*	11	*
941.29	*6	*	*	*	*	*	*	*	*	*	*
941.3	*7	*	*	*	*	*	*	*	*	*6	—
942	31	18	14	13	10	*	*	*	*6	21	*
942.2	16	10	*7	9	*	*	*	*	*	11	*
942.3	13	*6	*7	*	*	*	*	*	*	*9	*
943	30	20	10	10	16	*	*	*	*6	18	*
943.2	17	12	*5	*6	10	*	*	*	*	*9	*
943.21	*6	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
943.3	9	*5	*	*	*	*	*	*	*	*8	*
944	26	19	*7	*7	13	*5	*	*	*	17	*
944.2	14	11	*	*5	*6	*	*	*	*	*8	*
944.28	*6	*5	*	*	*	*	*	*	*	*5	—
944.3	*8	*5	*	*	*	*	*	*	*	*7	*
944.38	*5	*	*	*	*	*	*	*	—	*	*
945	29	19	11	*7	13	*6	*	*6	*5	15	*
945.2	19	12	*7	*5	*8	*	*	*	*	11	*
945.22	*5	*	*	*	*	*	*	*	*	*	*
945.3	*7	*5	*	*	*	*	*	*	*	*	*
948	41	25	16	12	17	9	*	*7	*8	25	*
948.0	30	19	11	*8	13	*7	*	*6	*5	18	*
948.00	30	19	11	*8	13	*7	*	*6	*5	18	*
948.1	*7	*	*	*	*	*	*	*	*	*	*
949	*6	*	*	*6	*	*	—	*	*	*6	*
949.0	*6	*	*	*5	*	*	—	—	*	*6	*
952	9	*7	*	*	*	*	*	*	*	*	*
952.0	*6	*	*	*	*	*	*	*	*	*	*
955	10	9	*	*	*6	*	*	*	*	*	*
958	39	25	15	*	16	11	*9	*6	11	14	*8
958.3	12	*	*7	*	*	*	*	*	*	*	*
958.4	*6	*	*	*	*	*	*	*	*	*	*
958.7	9	*7	*	*	*	*	*	*	*	*	*
958.8	*9	*7	*	*	*	*	*	*	*	*	*
959	90	51	39	*6	38	18	27	19	21	27	23
959.0	48	26	22	*	18	*8	17	*9	13	14	12
959.01	43	24	19	*	16	*8	16	*8	11	14	11
959.1	18	10	*8	*	10	*	*	*	*5	*	*5
959.12	*8	*	*	*	*	*	*	*	*	*	*
959.7	*9	*6	*	*	*	*	*	*	*	*	*
959.8	*6	*	*	*	*	*	*	*	*	*	*
962	10	*	*5	*	*	*	*	*	*	*	*
962.3	*9	*	*	*	*	*	*	*	*	*	*
963	16	*6	9	*	10	*	*	*	*5	*	*
963.0	13	*	*8	*	*9	*	*	*	*	*	*
965	110	40	70	*6	74	22	*8	21	27	38	25
965.0	45	19	26	*	23	15	*5	*8	*8	18	10
965.00	*9	*	*	*	*	*	*	*	*	*5	*
965.01	*5	*	*	—	*	*	—	*	*	*	*
965.02	*8	*	*	*	*	*	*	*	*	*5	*
965.09	22	*7	15	*	13	*7	*	*	*	*7	*7
965.1	12	*	*9	*	*8	*	*	*	*6	*	*
965.4	36	12	24	*	31	*	*	*8	9	12	*8
965.6	*8	*	*7	*	*7	*	*	*	*	*	*
965.61	*8	*	*6	*	*6	*	*	*	*	*	*
966	19	*6	13	*	11	*	*	*	*	*	*7
966.3	13	*	9	*	9	*	*	*	*	*	*6
967	21	*5	16	*	10	*6	*	*	*	*8	*
967.0	*6	*	*	*	*	*	*	*	*	*	*
967.8	*9	*	*7	*	*	*	*	*	*	*	*
967.9	*5	*	*	—	*	*	*	*	*	*	*
968	11	*6	*5	*	*9	*	*	*	*	*	*
968.0	*7	*	*	—	*6	*	*	*	*	*	*
969	128	44	85	*	86	28	9	26	30	44	29
969.0	31	*8	23	*	22	*6	*	*7	*8	10	*6
969.3	13	*	9	*	11	*	*	*	*	*	*
969.4	57	21	36	*	35	16	*	12	13	20	12

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
969.6	*5	*	*	*	*	*	*	*	*	*	*
969.8	10	*	*8	*	*7	*	*	*	*	*	*
970	19	12	*7	*	14	*	*	*6	*	*	*
970.8	18	11	*7	*	13	*	*	*6	*	*	*
972	14	*	10	*	*	*	*	*	*	*	*
972.6	*7	*	*	*	*	*	*	*	*	*	*
975	*5	*	*	*	*	*	—	*	*	*	*
977	22	*9	13	*	12	*8	*	*	*	11	*
977.8	13	*	*8	*	*6	*5	*	*	*	*5	*
977.9	*8	*	*	*	*	*	*	*	*	*5	*
980	25	13	12	*	17	*6	*	*	*6	9	*6
980.0	21	11	11	*	15	*	*	*	*	*8	*
987	*8	*	*	*	*	*	*	*	*	*	*
989	16	10	*5	*	*6	*	*	*	*	*8	*
989.5	11	*8	*	*	*	*	*	*	*	*6	*
991	14	10	*	*	*	*	*	*	*	*5	*
991.6	11	*7	*	*	*	*	*	*	*	*5	*
994	9	*7	*	*	*	*	*	*	*	*	*
995	397	187	210	22	47	97	231	90	93	141	73
995.0	*7	*	*	*	*	*	*	*	*	*	*
995.1	21	*7	14	*	*6	*6	*8	*	*	*9	*
995.2	25	*9	16	*	*	*6	16	*5	*8	9	*
995.5	12	*7	*	11	*	—	*	*	*	*8	*
995.9	321	158	162	*8	32	79	201	74	76	109	61
995.90	*9	*	*	*	*	*	*	*	*	*	*
995.91	195	92	102	*5	21	48	121	41	49	69	35
995.92	112	59	52	*	*9	28	72	30	24	36	22
996	832	410	422	49	149	282	352	176	170	316	171
996.0	28	19	9	*	*	*5	19	*8	*	10	*5
996.01	15	*8	*8	*	*	*	12	*	*	*7	*
996.04	*9	*8	*	*	*	*	*5	*	*	*	*
996.1	29	14	15	*	*7	*7	12	*7	*	10	*8
996.2	28	13	15	12	*8	*8	*	*5	*6	11	*6
996.3	12	*8	*	*	*	*	*5	*	*	*6	*
996.39	*8	*6	*	*	*	*	*	*	*	*	*
996.4	132	45	87	*	19	37	75	30	33	44	25
996.5	25	10	15	*	*	9	*9	*	*	11	*6
996.59	16	*7	*8	*	*	*6	*6	*	*	*7	*
996.6	277	142	134	20	52	100	104	57	60	106	53
996.61	11	*5	*5	—	*	*	*7	*	*	*	*
996.62	144	73	71	13	29	56	46	35	26	59	23
996.63	12	*7	*	*	*	*	*	*	*	*	*
996.64	22	13	9	*	*	*6	15	*	*6	*8	*6
996.65	*5	*	*	—	*	*	*	*	*	*	*
996.66	33	15	18	*	*	11	18	*6	*9	9	*8
996.67	19	11	*8	*	*5	*9	*	*	*6	*7	*
996.68	*7	*5	*	*	*	*	*	*	*	*	*
996.69	22	*8	13	*	*	*8	*9	*	*	11	*
996.7	242	127	114	*	33	84	120	50	49	89	53
996.72	56	35	21	*	*	18	33	13	13	22	9
996.73	70	34	36	*	12	26	32	12	11	26	22
996.74	46	23	22	*	*6	15	23	12	9	18	*7
996.76	16	12	*	*	*	*	11	*	*5	*	*
996.77	15	*6	*8	*	*	*6	*8	*	*	*	*
996.78	17	*8	9	*	*6	*6	*	*	*	*7	*
996.79	12	*	*7	*	*	*6	*	*	*	*	*
996.8	59	31	28	*	19	29	*7	11	*6	30	13
996.81	39	20	19	*	14	20	*	*6	*	20	*9

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
996.82	*6	*	*	*	*	*	*	*	*	*	*
996.85	*6	*	*	*	*	*	*	*	*	*	*
997	589	299	290	19	97	190	283	129	127	197	136
997.0	27	14	13	*	*	*9	12	*5	*8	*7	*7
997.02	13	*7	*6	*	*	*	*9	*	*	*	*
997.09	11	*6	*6	*	*	*	*	*	*	*	*
997.1	114	62	53	*	9	29	73	31	22	34	27
997.2	23	11	12	—	*	11	9	*6	*	*8	*5
997.3	113	56	56	*5	17	34	56	24	28	38	23
997.4	201	88	112	*7	48	69	77	37	44	71	49
997.5	65	40	25	*	9	17	37	16	13	19	17
997.6	29	19	10	*	*	12	13	*6	*	15	*
997.62	20	14	*6	*	*	*8	*8	*	*	11	*
997.69	9	*	*	*	*	*	*	*	*	*	*
997.9	15	*7	*8	*	*	*7	*	*	*	*	*
997.91	*5	*	*	*	*	*	*	*	*	*	*
997.99	9	*	*5	*	*	*	*	*	*	*	*
998	751	328	422	24	160	270	297	162	167	271	151
998.1	241	111	130	*8	49	73	111	50	55	81	54
998.11	122	63	59	*6	26	32	59	26	28	42	27
998.12	105	44	61	*	19	37	47	22	24	35	24
998.13	13	*	9	*	*	*	*	*	*	*	*
998.2	79	27	51	*	15	32	29	17	20	27	15
998.3	60	29	31	*	14	21	22	11	14	21	13
998.31	*8	*5	*	*	*	*	*	*	*	*	*
998.32	52	24	28	*	13	17	20	10	13	19	10
998.5	230	102	128	*7	49	96	78	52	50	87	41
998.51	10	*	*8	*	*	*5	*	*	*	*	*
998.59	220	100	121	*7	48	91	74	50	48	83	38
998.6	19	*6	13	*	*	10	*	*	*	*5	*5
998.8	109	46	63	*	27	35	44	25	22	41	21
998.83	19	*9	11	*	*	11	*6	*	*	9	*
998.89	87	36	50	*	24	23	37	21	19	30	17
998.9	9	*6	*	—	*	*	*7	*	*	*8	*
999	49	26	23	*	*8	19	18	11	10	19	10
999.2	14	*9	*6	*	*	*5	*6	*	*	*5	*
999.3	10	*	*	*	*	*	*	*	*	*	*
999.8	13	*7	*6	*	*	*6	*	*	*	*5	*
999.9	10	*	*5	*	*	*	*	*	*	*	*
V01-V83	13,582	4,021	9,561	379	6,462	2,434	4,306	2,569	3,119	4,992	2,902
V01	*6	*	*	*	*	*	*	*	*	*	*
V02	572	28	544	*	543	20	*7	99	102	241	130
V02.5	539	10	529	*	528	*	*5	81	99	234	125
V02.51	516	*	516	*	515	*	—	76	96	225	120
V02.59	19	9	10	*	10	*	*5	*	*	*9	*
V02.6	30	18	13	—	13	16	*	18	*	*6	*
V02.61	*8	*	*6	—	*	*	*	*	*	*	*
V02.62	22	15	*7	—	*8	12	*	15	*	*	*
V03	*8	*	*	*	*	*	*	*	*	*	*
V03.8	*7	*	*	—	*	*	*	*	*	*	*
V03.82	*7	*	*	—	*	*	*	*	*	*	*
V04	20	*	19	*	15	*	*	*6	*	*7	*
V04.3	13	—	13	—	13	—	—	*	*	*6	*
V04.8	*6	*	*	*	*	*	*	*	*	*	*
V04.81	*6	*	*	*	*	*	*	*	*	*	*
V05	*9	*	*	*6	*	*	*	*	*	*	*
V05.3	*7	*	*	*5	*	*	—	*	*	*	*
V06	*7	*	*5	*	*5	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V07	45	*	43	*	38	*	*	13	*9	10	12
V07.2	33	*	33	*	33	*	—	10	*7	*	11
V08	67	41	26	*	35	29	*	25	*6	33	*
V09	126	70	56	14	35	36	42	25	28	53	20
V09.0	121	68	53	14	34	35	39	23	27	51	19
V10	949	402	546	*	59	248	637	256	228	287	177
V10.0	209	113	96	*	*7	56	145	58	49	64	38
V10.01	*6	*	*	—	*	*	*	*	*	*	*
V10.02	*8	*5	*	*	*	*	*	*	*	*	*
V10.03	10	*8	*	*	*	*	*5	*	*	*	*
V10.04	*7	*	*	*	*	*	*6	*	*	*	*
V10.05	143	74	69	*	*	35	104	42	33	43	24
V10.06	23	12	11	—	*	*8	15	*6	*6	*6	*
V10.09	*9	*	*	*	*	*	*6	*	*	*	*
V10.1	60	24	36	*	*	17	41	18	15	19	*8
V10.11	60	24	36	*	*	17	41	18	15	19	*8
V10.2	16	11	*6	—	*	*7	*8	*	*	*	*
V10.21	14	10	*	—	*	*6	*8	*	*	*	*
V10.3	245	*	242	*	13	65	166	62	59	75	48
V10.4	224	144	80	*	15	49	160	59	59	67	41
V10.41	24	—	24	—	*7	12	*6	*6	*7	*7	*
V10.42	27	—	27	*	*	*8	18	*7	*7	*7	*6
V10.43	21	—	21	—	*	*9	11	*7	*	*7	*
V10.44	*7	—	*7	—	*	*	*	*	*	*	*
V10.46	132	132	—	*	*	15	117	34	32	42	25
V10.47	*8	*8	—	—	*	*	*	*	*	*	*
V10.5	67	42	25	*	*	15	51	23	14	19	11
V10.51	41	30	11	—	*	*6	34	13	9	11	*7
V10.52	24	12	12	*	*	*8	15	9	*	*7	*
V10.7	18	12	*6	—	*	*6	*9	*	*	*8	*
V10.79	15	10	*	—	*	*	*8	*	*	*7	*
V10.8	104	52	52	*	16	31	55	29	22	28	25
V10.82	27	14	13	—	*	10	13	*6	*5	*8	*7
V10.83	39	24	15	—	*	12	25	11	*8	12	*8
V10.85	*7	*	*	*	*	*	*	*	*	*	*
V10.87	15	*	12	—	*	*	*7	*5	*	*	*
V10.89	12	*6	*6	*	*	*	*8	*	*	*	*5
V11	31	19	11	*	*9	12	10	*	10	*8	*8
V11.3	18	14	*	—	*	*7	*9	*	*5	*	*7
V11.8	*6	*	*	*	*	*	*	*	*	*	*
V12	809	353	456	10	93	209	497	175	225	270	140
V12.0	43	18	25	*	14	13	15	*8	*8	16	11
V12.01	15	*6	9	*	*	*	*8	*	*	*5	*
V12.02	*6	*	*	—	*	*	*	*	*	*	*
V12.09	19	*8	12	*	10	*6	*	*	*	*8	*
V12.2	*8	*	*	*	*	*	*	*	*	*	*
V12.4	11	*5	*6	*	*	*	*5	*	*	*	*
V12.49	*7	*	*	*	*	*	*	*	*	*	*
V12.5	604	262	342	*	55	144	401	140	169	199	95
V12.50	10	*	*	*	*	*	*7	*	*	*	*
V12.51	172	64	108	*	33	57	81	43	51	46	32
V12.52	*9	*	*7	—	*	*	*5	*	*	*	*
V12.59	413	192	221	*	21	82	308	94	115	146	58
V12.6	14	*6	*8	*	*	*	*6	*	*6	*	*
V12.7	126	57	69	*	17	41	66	17	34	45	30
V12.71	69	31	38	*	11	23	35	10	15	27	18
V12.72	31	18	13	*	*	11	19	*	10	10	*8
V12.79	22	*7	15	*	*	*6	11	*	*8	*6	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V13	81	36	45	*7	27	23	25	14	26	23	19
V13.0	54	28	27	*	16	19	18	*8	17	18	12
V13.01	46	26	20	—	14	17	15	*7	14	15	10
V13.09	*8	*	*6	*	*	*	*	*	*	*	*
V13.5	*5	*	*	*	*	*	*	*	*	*	*
V14	139	39	99	*	46	46	44	28	35	56	19
V14.0	70	21	49	*	24	23	22	16	19	25	10
V14.1	*8	*	*	*	*	*	*	*	*	*	*
V14.2	19	*	16	*	*5	*6	*8	*	*	*9	*
V14.5	11	*	*8	*	*	*	*	*	*	*6	*
V14.6	11	*	*8	*	*	*	*	*	*	*	*
V14.8	13	*	10	*	*	*	*	*	*	*6	*
V15	899	494	405	36	219	317	327	164	225	321	189
V15.0	27	11	17	*8	*8	*7	*	*7	*7	*8	*
V15.09	*7	*	*	*	*	*	*	*	*	*	*
V15.1	26	16	10	14	*5	*	*	*8	*	*9	*
V15.2	*6	*	*	*	*	*	*	*	*	*	*
V15.3	31	15	16	*	*	14	11	*6	*7	10	*8
V15.4	15	*	12	*	9	*	*	*	*	*9	*
V15.41	13	*	10	*	*8	*	*	*	*	*7	*
V15.5	27	16	10	*	12	*8	*	*	9	9	*
V15.8	767	431	336	*6	178	278	305	137	196	272	163
V15.81	226	129	96	*	105	82	34	47	62	85	32
V15.82	523	292	231	*	70	191	262	83	130	185	126
V15.89	14	*5	*8	*	*	*	*5	*6	*	*	*
V16	57	16	42	*	15	29	12	*7	15	19	15
V16.0	14	*	9	*	*	*7	*	*	*	*6	*
V16.3	18	*	16	*	*6	*8	*	*	*5	*	*6
V16.4	11	*	*9	*	*	*7	*	*	*	*	*
V16.41	*7	—	*7	*	*	*	*	*	*	*	*
V17	194	106	88	16	48	90	40	33	51	68	43
V17.1	*5	*	*	*	*	*	*	*	*	*	*
V17.3	124	72	52	*	31	66	26	23	28	43	30
V17.4	48	22	26	*	13	20	11	*8	13	17	9
V17.5	13	*8	*	11	*	*	*	*	*6	*	*
V18	32	14	18	*7	11	11	*	*	*8	14	*6
V18.0	25	11	14	*	9	*9	*	*	*6	12	*
V19	*5	*	*	*	*	*	*	*	*	*	*
V21	11	*6	*5	10	*	*	—	*	*	*	*6
V21.3	10	*6	*	10	*	*	—	*	*	*	*6
V22	*9	—	*9	*	*9	*	—	*	*	*	*
V22.2	*6	—	*6	*	*5	*	—	*	*	*	*
V23	128	—	128	*	127	*	—	10	24	69	25
V23.2	13	—	13	—	13	—	—	*	*	10	*
V23.4	13	—	13	—	13	—	—	*	*	*6	*
V23.41	*7	—	*7	—	*7	—	—	*	*	*	*
V23.49	*6	—	*6	—	*6	—	—	*	*	*	*
V23.5	*8	—	*8	—	*8	*	—	*	*	*	*
V23.7	82	—	82	*	81	*	—	*6	15	43	18
V23.8	*7	—	*7	*	*7	—	—	*	*	*	*
V25	332	*	332	—	329	*	—	34	55	180	64
V25.0	*8	—	*8	—	*8	—	—	—	*	*8	—
V25.02	*7	—	*7	—	*7	—	—	—	*	*7	—
V25.2	322	*	322	—	319	*	—	34	54	171	63
V26	15	*	15	—	10	*	*	*	*	10	*
V26.5	13	*	13	—	*8	*	*	*	*	*8	*
V26.51	13	—	13	—	*8	*	*	*	*	*8	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V27	4,136	—	4,136	*5	4,120	10	—	589	802	1,679	1,066
V27.0	3,909	—	3,909	*	3,896	*8	—	536	744	1,609	1,019
V27.1	23	—	23	*	23	*	—	*	*	11	*
V27.2	50	—	50	—	49	*	—	*8	*8	20	13
V27.9	150	—	150	*	148	*	—	41	45	37	27
V29	19	10	10	19	—	—	—	*	*	13	*
V29.0	19	9	9	19	—	—	—	*	*	13	*
V30	9	*	*6	9	—	—	—	*	*	*	*
V30.1	9	*	*6	9	—	—	—	*	*	*	*
V42	127	65	62	14	28	44	40	37	21	40	29
V42.0	51	25	25	*	15	25	*9	12	*8	17	13
V42.1	10	*7	*	*	*	*	*	*	*	*	*
V42.2	26	12	13	*	*	*	22	*9	*5	*6	*6
V42.7	16	*9	*7	*	*	*6	*	*	*	*6	*
V42.8	21	10	11	*7	*6	*7	*	*8	*	*6	*
V42.81	*7	*	*	*	*	*	*	*	*	*	*
V42.83	*5	*	*	*	*	*	—	*	*	*	*
V43	484	186	297	*	23	113	346	99	147	149	89
V43.3	128	68	60	*	12	35	81	29	29	45	25
V43.6	350	115	235	*	11	77	263	68	117	103	63
V43.64	149	53	96	*	*7	29	112	26	49	43	30
V43.65	194	60	134	*	*	45	145	39	66	58	30
V44	200	108	92	43	36	53	68	57	51	59	33
V44.0	38	24	14	12	*9	*8	9	10	*8	17	*
V44.1	80	44	36	27	*8	18	28	23	24	19	14
V44.2	19	*9	11	*	*6	*7	*6	*7	*	*	*5
V44.3	46	24	22	*	10	14	21	11	13	16	*7
V44.6	*9	*	*5	*	*	*	*	*	*	*	*
V45	1,809	998	810	39	172	532	1,066	420	470	607	311
V45.0	303	171	132	*	*8	45	249	80	83	102	38
V45.01	252	132	120	*	*6	25	220	67	73	80	31
V45.02	46	36	10	*	*	17	26	12	*8	20	*6
V45.1	106	51	54	*	17	40	47	21	24	35	26
V45.2	31	15	16	13	*8	*	*7	*8	*7	9	*7
V45.3	46	14	32	*	22	17	*6	*8	12	15	10
V45.4	54	26	28	*	12	25	17	*	12	17	20
V45.6	*6	*	*	*	—	*	*	*	*	*	*
V45.7	168	42	126	*	35	58	72	32	47	55	34
V45.71	30	*	30	—	*	10	15	*	*6	12	*7
V45.72	19	*8	11	*	*	*	10	*	*8	*5	*
V45.73	30	13	17	*	*6	*9	15	*9	*7	*9	*6
V45.76	*8	*	*	*	*	*	*	*	*	*	*
V45.77	45	*	42	*	11	19	16	*6	13	16	11
V45.79	30	10	20	*	9	10	10	*8	*9	*8	*5
V45.8	1,094	676	417	19	69	343	662	266	284	370	174
V45.81	557	380	177	*	*8	141	407	157	133	191	76
V45.82	351	222	129	*	17	140	193	78	92	118	62
V45.89	183	73	110	18	43	60	61	30	59	59	35
V46	59	30	29	*6	*6	12	36	12	17	19	10
V46.1	22	13	*9	*	*	*	10	*	*7	9	*
V46.2	36	17	19	*	*	*8	25	*8	10	10	*8
V49	85	53	32	*	9	29	45	20	22	31	12
V49.6	*5	*	*	*	*	*	*	*	*	*	*
V49.7	73	49	25	*	*8	23	42	18	19	26	11
V49.72	*8	*	*	—	*	*	*	*	*	*	*
V49.75	35	24	11	*	*	11	20	*8	*8	13	*6
V49.76	23	14	*9	*	*	*	17	*6	*6	*9	*
V49.8	*6	*	*6	*	*	*	*	*	*	*	*
V49.81	*6	—	*6	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V50	12	*	*8	*	*	*6	*	*	*	*	*
V53	18	11	*6	*	*	*	10	*	*	*6	*
V53.3	13	*8	*5	*	*	*	*9	*	*	*	*
V53.31	*8	*	*	*	*	*	*6	*	*	*	*
V53.32	*5	*	*	–	*	*	*	*	*	*	*
V54	157	55	102	*	17	30	108	21	55	36	45
V54.0	*8	*	*	*	*	*	*	*	*	*	*
V54.01	*6	*	*	*	*	*	*	*	*	*	*
V54.1	74	30	44	*	10	13	51	11	27	15	21
V54.11	*7	*	*5	*	*	*	*5	*	*	*	*
V54.13	24	*8	16	–	*	*	22	*	*6	*9	*7
V54.16	9	*	*5	–	*	*	*5	*	*	*	*
V54.19	21	10	11	*	*	*	12	*	9	*	*6
V54.8	70	20	50	*	*	15	51	*8	25	16	20
V54.81	54	14	40	–	*	11	41	*6	20	11	17
V54.89	16	*6	10	*	*	*	10	*	*6	*	*
V55	54	30	24	*7	12	18	17	14	12	17	11
V55.1	*7	*	*	*	*	*	*	*	*	*	*
V55.2	11	*	*6	*	*	*	*	*	*	*	*
V55.3	27	16	11	*	*6	11	*9	*7	*6	10	*5
V56	130	56	74	*	17	55	56	28	*6	53	43
V56.0	126	55	71	*	16	53	55	28	*6	51	41
V57	594	239	355	*	51	134	406	111	178	181	124
V57.1	127	53	74	*	11	29	87	32	23	45	28
V57.2	27	12	15	*	*8	*6	13	*	9	11	*
V57.21	27	12	15	*	*8	*6	13	*	9	11	*
V57.8	436	173	263	*	30	99	304	74	145	124	93
V57.89	435	173	262	*	30	99	304	74	145	124	93
V58	667	315	352	50	82	191	344	141	164	221	140
V58.0	21	13	*8	*	*	*7	10	*	*	9	*5
V58.1	182	102	80	43	28	73	39	45	18	77	43
V58.4	17	*7	10	*	*	*	9	*5	*	*	*7
V58.49	11	*	*7	*	*	*	*6	*	*	*	*
V58.6	402	176	226	*	42	97	259	81	128	121	72
V58.61	243	114	129	*	20	47	176	55	64	70	54
V58.62	22	*8	14	*	*	*	19	*	20	*	*
V58.63	*6	*	*	–	*	*	*	*	*	*	*
V58.64	*5	*	*	–	*	*	*	*	*	*	*
V58.65	35	13	22	*	*7	10	16	*7	18	*6	*
V58.69	91	35	56	*	14	34	41	16	26	38	11
V58.7	17	*7	10	*	*	*	11	*	*	*	*6
V58.78	*8	*	*5	*	*	*	*6	*	*	*	*
V58.8	22	*7	15	*	*	*	13	*	*9	*5	*5
V58.83	*6	*	*	–	*	*	*	*	*	*	*
V58.89	13	*	9	*	*	*	*9	*	*8	*	*
V59	*9	*	*	*	*	*	*	*	*	*	*
V59.4	*7	*	*	*	*	*	*	*	*	*	*
V60	55	34	21	*	29	14	12	13	13	15	14
V60.0	32	24	*8	*	19	11	*	*6	*6	10	10
V60.2	9	*5	*	*	*7	*	*	*	*	*	*
V60.3	*5	*	*	–	*	*	*	*	*	*	*
V61	60	20	39	11	40	*8	*	*8	17	21	14
V61.0	10	*	*5	*	*7	*	*	*	*	*	*
V61.1	11	*	*7	–	*8	*	*	*	*	*	*
V61.10	11	*	*7	–	*8	*	*	*	*	*	*
V61.2	15	*6	*9	*8	*7	*	–	*	*	*8	*
V61.20	13	*6	*7	*7	*6	*	–	*	*	*7	*
V61.5	13	–	13	–	12	*	–	*	*	*7	*

See footnotes at end of table.

Table 47. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of all-listed diagnoses in thousands											
V62	51	27	24	*6	32	9	*	11	16	15	9
V62.0	*9	*	*	-	*7	*	-	*	*	*	*
V62.8	29	15	14	*	20	*5	*	*9	*9	*7	*
V62.81	*9	*	*5	*	*7	*	*	*	*	*	*
V62.89	17	11	*6	*	12	*	*	*6	*	*	*
V64	123	48	75	*5	44	41	33	25	24	47	26
V64.1	23	12	11	*	*6	*6	10	*	*	*9	*5
V64.2	16	*7	*9	*	*5	*6	*5	*	*	9	*
V64.3	20	*8	12	*	*5	*8	*5	*	*	10	*
V64.4	64	21	44	*	28	22	11	14	15	20	16
V64.41	64	20	43	*	28	21	11	14	15	20	15
V65	34	17	17	11	13	*5	*	*5	*	21	*
V65.2	*5	*	*	*	*	*	*	*	*	*	*
V65.3	23	9	13	11	*6	*	*	*	*	16	*
V65.4	*6	*	*	*	*	*	*	*	*	*	*
V66	47	25	22	*	*	10	34	*8	16	18	*
V66.7	41	22	19	*	*	*8	32	*6	16	16	*
V70	21	11	10	*	9	*5	*	*7	*	10	*
V70.7	19	10	*9	*	*8	*	*	*6	*	*8	*
V71	18	12	*6	*	*8	*	*	*	*	*8	*
V71.0	*6	*	*	*	*	*	*	*	*	*	*
V71.09	*5	*	*	*	*	*	*	-	*	*	*
V71.4	*7	*	*	*	*	*	*	*	*	*	*
V72	15	*6	*9	*	*	*	*9	*	*	9	*
V72.8	14	*6	*8	*	*	*	*9	*	*	*9	-
V72.83	*7	*	*	*	*	*	*	*	*	*	-

* Figure does not meet standard of reliability or precision.
 - Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
All codes	45,023	17,671	27,352	2,310	15,110	11,697	15,905	9,935	9,295	16,280	9,513
00	92	52	39	*	*	21	67	20	27	38	*7
00.1	48	21	26	*	*	13	32	10	13	22	*
00.11	*8	*	*	—	*	*	*	*	*	*	*
00.13	30	13	17	—	*	*	25	*	*9	17	*
00.14	*6	*	*	—	*	*	*	*	*	*	*
00.5	43	30	13	*	*	*7	34	9	14	15	*
00.50	*9	*	*	—	*	*	*7	*	*	*	*
00.51	31	23	*7	*	*	*6	24	*8	11	9	*
01-05	1,291	562	729	236	398	318	339	275	298	427	291
01	151	77	74	19	45	48	39	29	32	54	36
01.0	11	*6	*6	*	*	*	*	*	*	*	*
01.02	9	*	*	*	*	*	*	*	*	*	*
01.1	22	12	11	*	*7	*8	*	*	*7	*7	*
01.18	13	*7	*6	*	*5	*	*	*	*6	*	*
01.2	21	*9	12	*	9	*	*	*5	*	*5	*7
01.24	13	*6	*7	*	*5	*	*	*	*	*	*
01.3	36	21	15	*	*8	*7	19	*7	9	11	9
01.31	27	17	10	*	*	*5	16	*5	*6	*8	*7
01.39	*9	*	*	*	*	*	*	*	*	*	*
01.5	57	29	29	*6	18	24	10	11	*9	25	13
01.51	12	*	*7	*	*	*	*	*	*	*	*
01.59	44	24	20	*5	13	19	*7	10	*6	20	*8
02	119	61	58	33	41	29	17	25	25	41	28
02.0	23	12	11	*5	10	*7	*	*6	*	9	*
02.06	11	*5	*6	*	*	*	*	*	*	*5	*
02.1	13	*	*8	*	*6	*	*	*	*	*	*
02.12	12	*	*8	*	*6	*	*	*	*	*	*
02.2	27	13	13	*	9	9	*	*	*	10	*8
02.3	18	10	*8	*7	*	*	*5	*	*	*6	*
02.34	15	*8	*7	*6	*	*	*	*	*	*5	*
02.4	26	14	12	13	*8	*	*	*	*7	*8	*7
02.42	20	11	*9	11	*6	*	*	*	*5	*6	*
02.43	*6	*	*	*	*	*	*	*	*	*	*
02.9	12	*7	*5	*	*	*	*	*	*	*	*
02.94	*7	*	*	*	*	*	*	*	*	*	*
03	939	387	552	178	295	213	253	197	226	300	216
03.0	224	107	117	*	23	77	121	48	49	75	52
03.09	219	105	115	*	22	75	119	46	48	74	52
03.3	358	182	176	154	85	65	54	61	75	138	84
03.31	357	181	176	153	85	65	54	61	75	137	83
03.4	10	*	*6	*	*	*	*	*	*	*	*
03.5	30	15	15	*	*7	10	*9	*	*7	11	*8
03.53	12	*7	*6	—	*5	*	*	*	*	*	*
03.59	16	*8	*8	*	*	*6	*5	*	*	*5	*
03.6	*7	*	*	*	*	*	*	*	*	*5	*
03.9	302	74	228	15	173	53	61	77	92	66	67
03.90	123	27	96	*	80	20	20	40	25	26	32
03.91	86	10	76	*	60	*9	17	16	44	11	15
03.92	60	28	31	11	11	14	23	13	12	20	14
03.93	*8	*	*	—	*	*6	*	*	*	*	*
03.95	18	*	16	*	16	*	*	*	*7	*7	*
04	70	32	38	*5	14	26	25	22	13	26	9
04.0	*7	*	*	*	*	*	*	*	*	*	*
04.4	18	*8	10	*	*	*6	*6	*	*	10	*
04.43	10	*	*7	*	*	*	*	*	*	*6	*
04.49	*5	*	*	*	*	*	*	*	*	*	*
04.8	31	13	18	—	*	12	15	15	*7	*7	*
04.81	30	12	17	—	*	12	15	15	*7	*6	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
05	11	*5	*6	*	*	*	*	*	*	*6	*
05.3	*6	*	*	—	*	*	*	*	*	*	*
05.31	*6	*	*	—	*	*	*	*	*	*	*
06-07	103	26	76	*	32	41	28	25	18	33	27
06	84	18	66	*	28	34	21	22	15	28	19
06.1	*6	*	*5	—	*	*	*	*	*	*	*
06.2	19	*	15	*	*8	*7	*	*5	*	*6	*
06.3	10	*	*9	*	*	*	*	*	*	*	*
06.39	*9	*	*8	*	*	*	*	*	*	*	*
06.4	21	*	17	*	10	*8	*	*6	*	*7	*
06.8	16	*	12	*	*	*9	*6	*	*	*	*5
06.89	14	*	11	*	*	*8	*5	*	*	*	*
07	19	*9	10	*	*	*7	*6	*	*	*	*7
07.2	*6	*	*	*	*	*	*	*	*	*	*
07.22	*5	*	*	*	*	*	*	*	*	*	*
07.6	*7	*	*	*	*	*	*	*	*	*	*
07.62	*5	*	*	—	*	*	*	*	*	*	*
08-16	71	45	26	*9	19	20	24	14	10	31	15
08	29	18	10	*	10	*7	*8	*7	*6	10	*
08.8	17	11	*6	*	*7	*	*6	*	*	*6	*
08.81	15	10	*	*	*7	*	*	*	*	*	*
13	12	*	*7	*	*	*6	*5	*	*	9	*
14	17	13	*	*	*	*	*8	*	*	*7	*
14.7	*8	*6	*	*	*	*	*	*	*	*	*
16	*6	*	*	*	*	*	*	*	*	*	*
18-20	45	31	13	22	12	*6	*6	10	*8	16	11
18	20	15	*5	*7	*8	*	*	*	*	9	*5
18.2	*5	*	*	*	*	*	*	*	*	*	*
18.29	*5	*	*	*	*	*	*	*	*	*	*
18.4	*5	*	*	*	*	*	*	*	*	*	*
18.7	*6	*	*	*	*	*	*	*	*	*	*
20	20	13	*7	15	*	*	*	*7	*	*	*
20.0	15	*9	*6	13	*	*	*	*	*	*	*
20.01	14	*8	*5	13	*	*	*	*	*	*	*
21-29	309	198	110	63	115	62	68	75	49	116	68
21	86	58	28	*	28	19	34	17	17	33	19
21.0	39	27	12	*	*	*9	26	9	9	12	*8
21.01	17	11	*6	*	*	*	11	*	*	*6	*
21.02	*9	*6	*	—	*	*	*6	*	*	*	*
21.03	*8	*5	*	*	*	*	*6	*	*	*	*
21.3	*6	*	*	*	*	*	*	*	*	*	*
21.6	*7	*5	*	*	*5	*	*	*	*	*	*
21.69	*6	*	*	—	*	*	*	*	*	*	*
21.7	*6	*	*	*	*	*	*	*	*	*	*
21.8	20	13	*7	*	12	*	*	*	*	10	*
21.81	*8	*7	*	*	*	*	*	*	*	*	*
21.88	*7	*	*	*	*	*	*	*	*	*	*
22	29	16	13	*7	11	*	*7	*	*	13	*7
22.1	*5	*	*	*	*	*	*	*	*	*	*
22.6	13	*8	*6	*	*	*	*	*	*	*6	*
22.63	*7	*	*	*	*	*	*	*	*	*	*
23	27	17	10	*	14	*	*	*7	*	13	*
23.0	10	*6	*	*	*	*	*	*	*	*	*
23.09	10	*5	*	*	*	*	*	*	*	*	*
23.1	14	10	*	*	*8	*	*	*	*	*8	*
23.19	13	10	*	*	*8	*	*	*	*	*8	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
24	*7	*	*	*	*	*	*	*	*	*	*
25	*9	*5	*	*	*	*	*	*	*	*	*
26	13	*8	*	*	*	*5	*	*	*	*6	*
26.3	11	*7	*	*	*	*	*	*	*	*5	*
26.31	*5	*	*	—	*	*	*	*	*	*	*
27	66	42	24	19	26	12	*8	17	*7	24	18
27.0	10	*7	*	*	*7	*	*	*	*	*7	*
27.4	*8	*	*	*	*	*	*	*	*	*	*
27.5	22	15	*7	*7	*9	*	*	*	*	*7	*8
27.51	10	*6	*	*	*	*	*	*	*	*	*
27.54	*5	*	*	*	*	—	—	*	*	*	*
27.6	19	11	*8	10	*7	*	*	*6	*	*6	*5
27.62	*8	*	*	*8	*	—	—	*	*	*	*
27.69	*9	*6	*	*	*6	*	*	*	*	*	*
28	48	31	17	24	19	*	*	14	*9	14	11
28.0	*8	*5	*	*	*6	*	*	*	*	*	*
28.2	16	10	*5	*	*8	*	*	*6	*	*	*
28.3	18	11	*7	16	*	*	—	*	*	*6	*6
29	24	17	*7	*	*8	*8	*6	*8	*	*7	*6
29.1	*7	*	*	*	*	*	*	*	*	*	*
29.11	*6	*	*	*	*	*	*	*	*	*	*
29.3	*6	*	*	*	*	*	*	*	*	*	*
29.4	*9	*6	*	*	*6	*	*	*	*	*	*
30-34	1,045	572	473	60	169	317	498	229	211	387	218
30	*9	*	*	*	*	*	*	*	*	*	*
31	160	88	72	14	28	54	64	35	31	59	35
31.1	87	50	37	*	18	29	38	16	16	34	21
31.2	13	*7	*6	*	*	*	*6	*	*	*	*
31.29	13	*7	*6	*	*	*	*6	*	*	*	*
31.4	48	25	23	*8	*7	17	16	13	10	18	*7
31.42	40	21	19	*8	*7	12	13	12	9	13	*6
31.43	*6	*	*	—	*	*	*	*	*	*	*
31.7	*6	*	*	*	*	*	*	*	*	*	*
32	72	35	37	*	*9	24	38	16	14	29	13
32.2	34	17	17	*	*7	11	15	*8	*8	13	*5
32.29	33	16	16	*	*6	11	15	*8	*7	13	*
32.3	*6	*	*	*	*	*	*	*	*	*	*
32.4	28	13	15	*	*	*9	17	*6	*	11	*6
33	328	172	156	21	48	105	155	75	68	121	64
33.2	318	167	151	20	46	103	150	74	64	118	62
33.22	45	27	18	*	*7	15	19	15	*7	14	*8
33.23	43	25	18	9	*7	14	13	10	9	14	9
33.24	131	67	63	*	19	38	69	28	28	48	26
33.26	26	12	13	*	*	*7	17	*	*6	10	*5
33.27	47	23	23	*	*7	18	21	*8	10	23	*6
33.28	22	10	12	*	*	*8	9	*	*	*7	*7
34	476	272	203	22	84	131	239	102	98	174	102
34.0	174	107	67	11	46	50	67	31	37	68	39
34.03	*8	*6	*	*	*	*	*	*	*	*	*
34.04	128	77	52	*9	35	37	49	26	28	48	27
34.09	32	20	13	*	*8	9	13	*	*7	13	9
34.2	52	26	26	*	*8	16	26	11	10	19	13
34.21	21	12	9	*	*	*6	9	*	*	*7	*6
34.22	11	*6	*	—	*	*	*6	*	*	*	*
34.24	*8	*	*	*	*	*	*	*	*	*	*
34.25	*6	*	*	*	*	*	*	*	*	*	*
34.5	24	18	*6	*	*	*7	10	*	*	*9	*6
34.51	17	13	*	*	*	*	*7	*	*	*6	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
34.59	*7	*5	*	*	*	*	*	*	*	*	*
34.6	*7	*5	*	*	*	*	*	*	*	*	*
34.7	*9	*7	*	*	*	*	*	*	*	*	*
34.9	198	103	96	*	18	50	128	49	43	67	39
34.91	176	89	87	*	15	43	116	43	38	60	36
34.92	21	13	*8	*	*	*6	13	*5	*	*7	*
35-39	6,993	3,985	3,009	226	675	2,511	3,581	1,388	1,440	2,731	1,434
35	147	79	68	33	18	33	64	42	25	46	34
35.1	18	*8	9	*	*	*5	*7	*6	*	*7	*
35.12	12	*6	*6	*	*	*	*	*	*	*5	*
35.2	81	44	38	*	10	20	51	19	17	26	20
35.21	24	15	9	*	*	*	20	*8	*	*	*7
35.22	28	16	12	—	*	*9	15	*5	*7	12	*
35.23	11	*	*6	—	*	*	*9	*	*	*	*
35.24	16	*7	*9	*	*	*6	*7	*	*	*7	*
35.3	*8	*	*	*	*	*	*	*	*	*	*
35.33	*7	*	*	*	*	*	*	*	*	*	*
35.5	*7	*	*	*	*	*	*	*	*	*	*
35.6	*8	*	*	*6	*	*	*	*	*	*	*
35.7	12	*7	*6	*7	*	*	*	*	*	*	*
35.71	*9	*5	*	*	*	*	*	*	*	*	*
35.8	*6	*	*	*6	*	*	*	*	*	*	*
36	1,718	1,161	558	*	89	748	880	310	424	634	350
36.0	1,285	853	432	*	72	550	663	227	333	463	262
36.01	537	353	183	*	30	234	272	99	134	201	103
36.02	16	11	*6	—	*	*8	*7	*	*	*7	*
36.05	111	74	37	*	*5	42	63	19	33	34	25
36.06	157	113	44	—	12	66	78	19	41	59	37
36.07	459	300	159	*	23	197	238	86	122	160	92
36.1	427	303	124	—	17	195	215	81	89	170	87
36.11	45	29	16	—	*	18	25	11	*8	19	*8
36.12	81	56	26	—	*	38	41	19	18	31	14
36.13	70	52	18	—	*	33	34	10	15	31	14
36.14	39	30	9	—	*	18	20	*	*8	16	11
36.15	177	126	51	—	*8	81	89	32	38	71	37
36.16	*8	*7	*	—	*	*	*	*	*	*	*
37	1,977	1,149	828	21	143	708	1,107	417	406	767	387
37.0	11	*6	*	*	*	*	*	*	*	*	*
37.1	11	*7	*	*	*	*6	*	*	*	*	*
37.12	10	*6	*	*	*	*	*	*	*	*	*
37.2	1,421	843	578	14	114	588	706	268	305	573	276
37.21	26	16	10	*	*	11	12	*7	*	11	*
37.22	1,136	684	452	*	87	495	553	192	245	478	221
37.23	134	71	63	9	11	42	73	36	26	43	29
37.25	*5	*	*	*	*	*	*	*	*	*	*
37.26	77	47	30	*	*6	25	44	20	18	27	12
37.27	27	13	13	*	*	*8	15	*	*8	*9	*6
37.29	12	*6	*6	*	*	*	*6	*6	*	*	*
37.3	57	32	25	*	*6	19	28	13	13	19	13
37.33	11	*6	*5	*	*	*	*	*	*	*	*
37.34	41	23	18	*	*	16	20	9	10	15	*7
37.6	39	25	13	*	*	17	18	10	*5	16	*7
37.61	37	24	13	—	*	17	17	*9	*5	15	*7
37.7	192	95	98	*	*	24	162	54	32	67	40
37.71	34	16	18	—	*	*	31	13	*	11	*
37.72	113	57	57	*	*	15	96	31	20	40	23
37.75	*5	*	*	—	*	*	*5	*	*	*	*
37.76	*7	*	*	—	*	*	*5	*	*	*	*
37.78	19	9	9	*	*	*	14	*	*	*6	*5

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
37.8	170	85	85	*	*	22	144	51	29	59	31
37.81	10	*	*6	*	*	*	*8	*	*	*	*
37.82	21	11	10	—	*	*	20	9	*	*6	*
37.83	115	58	56	*	*	16	96	31	20	40	23
37.87	13	*6	*7	*	*	*	11	*	*	*6	*
37.89	*6	*	*	*	*	*	*	*	*	*	*
37.9	72	54	18	*	*6	27	39	16	16	25	16
37.94	57	42	15	*	*	22	31	12	13	19	13
37.98	*5	*	*	—	*	*	*	*	*	*	*
37.99	*6	*	*	—	*	*	*	*	*	*	*
38	1,786	858	928	129	277	537	843	344	329	758	355
38.0	28	15	13	*	*	*9	16	*	*6	10	*8
38.03	*9	*	*	—	*	*	*	*	*	*	*
38.08	12	*8	*	—	*	*	*8	*	*	*	*
38.1	131	83	48	*	*	27	103	25	27	56	24
38.12	98	60	38	—	*	19	78	19	21	43	15
38.16	*7	*	*	—	*	*	*6	*	*	*	*
38.18	21	14	*6	—	*	*6	15	*	*	*8	*
38.2	9	*	*6	—	*	*	*7	*	*	*	*
38.21	*8	*	*6	—	*	*	*6	*	*	*	*
38.4	41	30	11	*	*	10	28	10	10	12	11
38.44	21	15	*6	*	*	*	18	*	*6	*6	*
38.45	10	*6	*	*	*	*	*	*	*	*	*
38.7	82	33	48	*	11	20	51	24	16	32	9
38.8	26	14	12	9	*9	*	*	*7	*	10	*
38.85	9	*5	*	9	*	*	*	*	*	*	*
38.9	1,457	673	784	117	246	462	632	271	262	632	291
38.91	95	45	50	18	13	27	36	14	21	41	19
38.92	13	*8	*5	13	—	*	*	*	*	*6	*
38.93	1,132	513	619	79	202	363	488	217	207	485	222
38.94	*8	*5	*	*	*	*	*	*	*	*	*
38.95	171	82	89	*	24	59	86	27	29	76	39
38.98	29	16	13	*	*	*6	17	*8	*	17	*
38.99	10	*	*5	*	*	*5	*	*	*	*	*
39	1,364	738	627	42	149	486	687	274	257	526	308
39.2	125	75	50	*	*7	41	74	25	21	50	29
39.25	18	14	*	—	*	*7	11	*	*5	*6	*5
39.27	33	17	16	—	*	10	18	*7	*	13	10
39.29	66	39	27	—	*	22	43	15	11	28	12
39.3	21	12	*9	*	9	*5	*7	*	*	*8	*5
39.31	13	*8	*5	*	*	*	*	*	*	*	*
39.4	68	39	29	*	9	25	33	9	11	28	20
39.42	19	10	*9	—	*	*8	*9	*	*	*6	*7
39.43	*5	*	*	—	*	*	*	*	*	*	*
39.49	44	27	17	*	*5	15	22	*5	*7	19	12
39.5	170	86	84	*7	16	54	93	35	37	65	34
39.50	120	61	59	*	*8	37	73	27	27	45	21
39.51	11	*	*8	*	*	*7	*	*	*	*	*
39.52	10	*7	*	*	*	*	*7	*	*	*	*
39.56	*7	*	*	*	*	*	*	*	*	*	*
39.57	*6	*	*	*	*	*	*	*	*	*	*
39.59	10	*	*6	*	*	*	*6	*	*	*	*
39.6	256	164	92	24	18	96	118	49	55	87	65
39.61	227	145	82	22	16	86	104	46	51	80	50
39.62	*9	*7	*	*	*	*	*6	*	*	*	*7
39.63	*8	*6	*	*	*	*	*5	*	*	*	*6
39.64	10	*5	*	*	*	*5	*	*	*	*	*
39.7	32	23	9	*	*	9	19	10	*6	9	*6
39.71	14	12	*	—	—	*	12	*	*	*	*
39.72	*9	*	*5	*	*	*	*	*	*	*	*
39.79	*9	*7	*	*	*	*	*5	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
39.9	686	335	350	*	85	253	343	140	123	275	148
39.90	62	33	29	*	*	18	41	11	18	23	11
39.95	607	297	311	*	79	230	296	125	101	246	135
39.98	*8	*	*5	*	*	*	*	*	*	*	*
40-41	341	160	180	24	61	117	138	74	67	126	74
40	212	94	117	*6	32	88	86	44	45	76	48
40.1	58	23	35	*	13	19	23	11	12	22	13
40.11	55	23	32	*	12	18	22	10	12	22	12
40.2	53	15	39	*	*8	21	23	9	13	16	15
40.23	16	*	15	-	*	*6	*7	*	*	*6	*
40.24	*6	*	*	*	*	*	*	*	*	*	*
40.29	27	10	17	*	*	10	11	*	*7	*7	*9
40.3	69	35	34	*	*6	35	28	17	15	22	14
40.4	14	10	*	*	*	*5	*5	*	*	*6	*
40.41	11	*8	*	-	*	*	*	*	*	*	*
40.5	16	11	*	-	*	*8	*6	*	*	*8	*
40.59	*7	*5	*	-	*	*	*	*	*	*	*
41	129	66	63	18	30	29	52	30	22	50	26
41.0	*7	*	*	*	*	*	*	*	*	*	*
41.3	92	46	46	13	16	20	43	21	18	35	18
41.31	91	45	46	13	15	20	43	21	18	35	18
41.5	25	12	13	*	10	*6	*7	*	*	12	*
42-54	5,824	2,447	3,377	261	1,372	1,716	2,475	1,230	1,210	2,194	1,190
42	132	74	59	10	15	48	60	30	22	49	33
42.2	43	25	19	*6	*6	16	15	12	*6	17	*7
42.23	39	23	17	*5	*6	15	13	10	*6	16	*7
42.3	31	19	13	*	*	18	10	*7	*	12	9
42.33	30	18	12	*	*	17	*9	*7	*	12	*8
42.4	*6	*5	*	-	*	*	*	*	*	*	*
42.9	41	18	24	*	*	*7	28	*7	11	13	11
42.92	41	17	24	*	*	*7	28	*7	11	13	11
43	243	124	119	27	23	49	145	51	43	95	54
43.1	180	87	93	11	17	35	117	38	34	70	37
43.11	145	68	77	*	10	30	102	33	27	57	28
43.19	36	19	17	*8	*7	*5	15	*6	*7	14	*9
43.3	16	14	*	15	*	*	*	*	*	*5	*
43.4	19	*6	13	*	*	*	12	*	*	*8	*
43.41	16	*	12	*	*	*	11	*	*	*6	*
43.7	11	*7	*	-	*	*	*7	*	*	*6	*
43.9	*5	*	*	-	*	*	*	*	*	*	*
44	319	115	204	12	116	102	89	62	56	133	68
44.0	*5	*	*	-	*	*	*	*	*	*	*
44.1	26	10	16	*	9	*6	10	*	*6	14	*
44.13	14	*	10	*	*6	*	*	*	*	*8	*
44.14	10	*	*5	*	*	*	*5	*	*	*6	*
44.2	13	*	*8	*	*	*	*	*	*	*5	*
44.22	*6	*	*5	*	*	*	*	*	*	*	*
44.29	*6	*	*	*	*	*	*	*	*	*	*
44.3	114	20	93	*	67	40	*6	28	16	46	24
44.31	22	*	19	-	15	*7	*	*	*	*8	*5
44.39	88	15	73	*	52	32	*	22	11	36	18
44.4	92	51	41	*	11	25	56	15	19	39	18
44.41	*6	*	*	-	*	*	*	*	*	*	*
44.42	11	*5	*6	*	*	*	*7	*	*	*	*
44.43	73	41	32	*	*7	20	46	12	14	32	15
44.6	65	24	41	10	23	21	11	13	10	24	18
44.66	38	14	23	*8	*9	14	*7	*	*7	15	11
44.69	20	*	15	*	10	*7	*	*6	*	*5	*6

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
45	2,287	990	1,296	36	326	643	1,280	509	501	847	430
45.0	*7	*	*	*	*	*	*	*	*	*	*
45.1	1,120	498	622	18	160	322	620	241	243	426	211
45.13	453	201	251	*	66	133	249	95	104	166	87
45.14	*6	*	*	*	*	*	*	*	*	*	*
45.16	653	291	361	12	90	183	367	141	137	254	121
45.2	602	242	360	*6	89	157	350	131	127	226	118
45.23	332	132	200	*	40	85	207	71	71	132	59
45.24	41	17	23	*	*7	*8	25	*6	9	18	*7
45.25	220	88	132	*	41	61	114	53	46	71	50
45.3	15	*8	*8	*	*	*	*8	*	*	*	*
45.30	*5	*	*	—	*	*	*	*	*	*	*
45.33	*7	*	*	*	*	*	*	*	*	*	*
45.4	141	64	77	*	*8	35	98	38	32	45	26
45.42	117	54	63	*	*6	30	82	31	26	38	22
45.43	20	*8	12	—	*	*	14	*6	*	*6	*
45.6	73	30	43	*	16	22	31	15	16	28	14
45.62	69	29	40	*	15	20	30	14	16	27	13
45.7	259	114	145	*5	34	81	138	64	62	88	44
45.72	11	*6	*6	*	*	*	*	*	*	*	*
45.73	95	37	58	*	*8	28	59	23	21	36	15
45.74	10	*	*6	*	*	*	*7	*	*	*	*
45.75	23	11	12	*	*	*9	12	*6	*	9	*
45.76	95	45	50	*	18	31	45	25	25	26	19
45.79	22	10	12	*	*	*8	10	*5	*	10	*
45.8	*9	*6	*	*	*	*	*	*	*	*	*
45.9	58	26	32	*	12	17	27	12	13	24	10
45.91	16	*7	9	*	*	*	*7	*	*	*7	*
45.93	22	*9	14	*	*	*6	12	*6	*5	*8	*
45.94	16	*8	*8	*	*	*6	*7	*	*	*7	*
46	219	107	112	13	49	67	90	52	46	77	45
46.0	15	*7	*8	*	*	*	*6	*	*	*	*
46.01	*6	*	*	*	*	*	*	*	*	*	*
46.03	*8	*	*	*	*	*	*	*	*	*	*
46.1	42	19	23	*	*5	12	22	11	*8	16	*7
46.10	23	10	14	*	*	*7	14	*7	*	10	*
46.11	13	*7	*6	*	*	*	*	*	*	*	*
46.13	*6	*	*	*	*	*	*	*	*	*	*
46.2	20	11	9	*	*6	*6	*6	*5	*	*7	*
46.20	10	*7	*	*	*	*	*	*	*	*	*
46.3	28	17	11	*	*	9	14	*6	*6	*9	*8
46.32	*5	*	*	*	*	*	*	*	*	*	*
46.39	23	14	9	*	*	*7	11	*5	*6	*7	*5
46.4	12	*	*8	*	*	*	*7	*	*	*	*
46.5	42	23	19	*	11	17	12	11	9	14	*9
46.51	12	*6	*6	*	*	*	*	*	*	*	*
46.52	30	17	13	*	*7	12	*9	*8	*6	10	*6
46.7	40	15	25	*	15	10	14	9	*7	17	*7
46.73	14	*7	*7	*	*	*	*	*	*	*6	*
46.75	12	*	*8	*	*5	*	*5	*	*	*6	*
46.79	*8	*	*7	*	*	*	*	*	*	*	*
46.8	16	9	*7	*	*	*	*7	*	*	*6	*
46.85	11	*6	*	*	*	*	*7	*	*	*	*
47	391	202	188	91	195	71	34	77	82	125	106
47.0	332	182	150	81	172	57	22	70	69	103	90
47.01	151	83	68	34	84	26	*8	29	33	46	43
47.09	181	99	81	47	88	31	14	41	36	56	47
47.1	55	17	37	*8	22	13	12	*7	13	20	15
47.11	*7	*	*	*	*	*	*	*	*	*	*
47.19	48	15	33	*7	18	12	11	*6	11	18	12

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
48	168	89	79	10	32	53	73	37	44	60	28
48.2	63	31	32	*6	12	16	28	15	16	21	11
48.23	25	10	15	*	*5	*7	11	*	*5	10	*5
48.24	35	19	17	*	*6	*9	17	9	11	10	*
48.3	30	17	14	*	*	10	19	*6	*7	12	*7
48.36	25	13	12	*	*	*9	15	*	*6	10	*
48.5	*8	*5	*	—	*	*	*	*	*	*	*
48.6	33	17	17	*	*6	12	14	*8	9	11	*
48.63	18	*9	9	*	*	*7	*9	*	*6	*6	*
48.69	11	*5	*5	*	*	*	*	*	*	*	*
48.7	*6	*	*	*	*	*	*	*	*	*	*
48.8	22	16	*7	*	9	*8	*	*	*7	10	*
48.81	21	14	*6	*	9	*8	*	*	*7	*9	*
49	42	21	21	*	15	12	12	*8	*7	18	*9
49.0	9	*	*	*	*	*	*	*	*	*	*
49.01	*8	*	*	*	*	*	*	*	*	*	*
49.2	*9	*6	*	*	*	*	*	*	*	*	*
49.21	*6	*	*	*	*	*	*	*	*	*	*
49.4	10	*	*6	—	*	*	*	*	*	*	*
49.46	*6	*	*	—	*	*	*	*	*	*	*
49.7	*6	*	*	*	*	*	*	*	*	*	*
50	119	58	60	*	27	46	42	25	22	50	22
50.1	90	44	47	*	20	35	33	18	18	35	19
50.11	65	34	31	*	12	27	24	11	15	26	13
50.12	22	9	13	*	*6	*7	*8	*6	*	*8	*5
50.2	11	*	*6	*	*	*	*5	*	*	*6	*
50.29	*6	*	*	*	*	*	*	*	*	*	*
50.9	*5	*	*	*	*	*	*	*	*	*	*
51	696	238	458	*6	214	205	271	141	136	261	159
51.0	*7	*	*	*	*	*	*5	*	*	*	*
51.1	53	25	28	*	11	15	27	11	9	21	12
51.10	39	17	22	*	9	11	18	*9	*6	16	*8
51.14	*8	*	*	—	*	*	*6	*	*	*	*
51.2	435	136	299	*	146	137	148	89	87	164	94
51.22	99	38	61	*	20	33	46	24	22	34	19
51.23	335	98	237	*	126	104	103	66	65	130	75
51.3	*7	*	*	*	*	*	*	*	*	*	*
51.4	13	*	9	—	*	*6	*6	*	*	*6	*
51.41	*8	*	*6	—	*	*	*	*	*	*	*
51.8	163	56	107	*	51	41	71	31	33	55	44
51.84	*6	*	*	—	*	*	*	*	*	*	*
51.85	80	24	55	*	25	22	32	15	17	26	23
51.87	34	11	23	*	*8	*8	18	*7	*6	12	*9
51.88	44	18	26	*	16	9	18	*9	*9	15	12
51.9	10	*5	*5	*	*	*	*7	*	*	*5	*
51.98	9	*	*5	*	*	*	*6	*	*	*	*
52	34	15	19	*	*8	13	12	10	*7	13	*
52.1	9	*	*7	*	*	*	*5	*	*	*	*
52.7	*5	*	*	—	*	*	*	*	*	*	*
52.9	10	*	*6	*	*	*	*	*	*	*	*
52.93	*8	*	*	—	*	*	*	*	*	*	*
53	257	114	143	12	54	89	102	54	55	95	53
53.0	38	31	*6	*	*	*8	21	10	*7	13	*7
53.03	9	*8	*	—	*	*	*6	*	*	*	*
53.04	9	*7	*	*	*	*	*6	*	*	*	*
53.05	*7	*6	*	*	*	*	*	*	*	*	*
53.1	9	*8	*	*	*	*	*	*	*	*	*
53.2	*6	*	*	—	*	*	*	*	*	*	*
53.4	53	20	32	*	13	16	22	11	12	19	11
53.41	9	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
53.49	44	15	28	*	10	12	20	*9	10	16	9
53.5	48	15	32	*	14	18	15	*8	10	21	*8
53.51	31	11	21	*	10	11	9	*	*7	13	*7
53.59	17	*	12	*	*	*7	*6	*	*	*8	*
53.6	91	32	59	—	20	39	31	18	20	32	20
53.61	67	23	44	—	14	30	24	14	16	23	14
53.69	23	*8	15	—	*7	9	*7	*	*	9	*6
53.7	*6	*	*	*	*	*	*	*	*	*	*
53.9	*6	*	*	*	*	*	*	*	*	*	*
54	916	298	618	37	297	317	265	177	189	372	178
54.0	19	*6	13	*	*	*8	*6	*	*	*9	*
54.1	60	24	37	*	22	18	17	14	13	21	13
54.11	26	*9	17	*	11	*7	*6	*7	*	*9	*6
54.12	17	*8	9	*	*5	*	*6	*	*5	*6	*
54.19	17	*7	10	*	*5	*6	*	*6	*	*6	*
54.2	145	43	102	*5	57	44	39	27	29	61	28
54.21	93	28	65	*	41	30	18	15	21	39	18
54.23	24	*6	17	*	*7	*5	11	*	*	11	*
54.24	*9	*	*6	*	*	*	*	*	*	*	*
54.25	16	*6	11	*	*5	*6	*	*5	*	*6	*
54.3	18	*6	13	*	*6	*6	*6	*	*	*9	*
54.4	66	11	55	*	17	28	18	13	14	28	11
54.5	342	74	268	*6	141	104	91	64	71	145	62
54.51	57	16	42	*	25	17	13	13	13	22	10
54.59	285	59	226	*	116	87	78	51	59	123	52
54.6	11	*7	*	*	*	*	*	*	*	*	*
54.7	10	*	*6	*	*	*	*	*	*	*	*
54.72	*5	*	*	*	*	*	*	*	*	*	*
54.9	245	123	122	16	43	102	83	47	50	93	55
54.91	184	91	93	*5	30	82	67	40	38	69	37
54.93	9	*5	*	*	*	*	*	*	*	*	*
54.95	16	*8	*8	*6	*	*	*	*	*	*7	*
54.98	29	16	13	*	*	11	9	*	*	11	9
55-59	963	443	519	32	217	305	409	225	204	342	192
55	155	84	70	*9	38	66	42	36	28	65	26
55.0	38	20	18	*	*8	16	14	9	10	12	*6
55.03	24	13	11	*	*	*8	12	*7	*	*8	*
55.04	*8	*	*	—	*	*	*	*	*	*	*
55.2	31	17	14	*	12	11	*6	*8	*	13	*7
55.23	28	14	13	*	11	10	*	*7	*	12	*6
55.4	*8	*	*	*	*	*	*	*	*	*	*
55.5	45	25	20	*	*9	20	14	*8	*7	20	*9
55.51	42	23	19	*	*8	19	14	*7	*7	19	*9
55.6	16	9	*7	*	*6	9	*	*	*	9	*
55.69	16	9	*7	*	*6	*9	*	*	*	9	*
55.8	*6	*	*	*	*	*	*	*	*	*	*
55.87	*6	*	*	*	*	*	*	*	*	*	*
55.9	*7	*	*	*	*	*	*	*	*	*	*
55.93	*5	*	*	*	*	*	*	*	*	*	*
56	100	52	49	*9	33	33	26	24	19	36	21
56.0	50	27	23	*	21	20	*9	9	10	21	9
56.3	18	*9	9	—	*7	*	*6	*6	*	*7	*
56.31	17	*8	*8	—	*7	*	*5	*5	*	*6	*
56.5	10	*6	*	—	*	*	*6	*	*	*	*
56.51	9	*6	*	—	*	*	*6	*	*	*	*
56.7	12	*5	*7	*7	*	*	*	*	*	*	*
56.74	11	*	*7	*7	*	*	*	*	*	*	*
57	383	204	179	10	54	91	228	100	83	116	83
57.0	27	22	*	*	*	*	23	*8	*8	*8	*
57.1	30	15	15	*	*	*8	17	*	*8	*8	*8

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
57.17	11	*6	*6	*	*	*	*8	*	*	*	*
57.18	12	*6	*6	*	*	*	*6	*	*	*	*
57.19	*5	*	*	-	*	*	*	*	*	*	*
57.3	146	68	78	*	27	40	75	37	33	54	22
57.32	132	61	72	*	26	37	65	33	31	50	19
57.33	13	*7	*6	-	*	*	9	*	*	*	*
57.4	46	36	10	*	*	*8	36	15	*9	13	9
57.49	46	36	10	*	*	*8	36	15	*9	13	9
57.7	10	*6	*	-	*	*	*6	*	*	*	*
57.71	*9	*6	*	-	-	*	*5	*	*	*	*
57.8	23	*7	16	*	*6	9	*6	*	*7	*7	*5
57.81	11	*	*8	*	*	*5	*	*	*	*	*
57.9	95	45	50	*	11	18	64	28	15	22	30
57.94	86	38	48	*	10	17	57	25	13	20	28
58	42	31	11	*	*7	*8	25	11	11	14	*6
58.5	*5	*	*	*	*	*	*	*	*	*	*
58.6	22	16	*6	*	*	*5	14	*6	*7	*8	*
59	283	73	210	*	85	107	89	54	63	110	56
59.0	*8	*	*7	*	*	*	*	*	*	*	*
59.02	*6	*	*	*	*	*	*	*	*	*	*
59.4	*7	-	*7	-	*	*	*	*	*	*	*
59.5	30	*	30	-	13	11	*6	*	*6	16	*6
59.7	80	*	79	*	17	35	28	*8	18	34	20
59.79	78	*	77	-	17	34	27	*8	18	32	19
59.8	148	68	80	*	47	50	48	37	33	53	25
60-64	242	242	-	24	15	72	130	50	54	94	43
60	171	171	-	-	*	57	113	40	39	57	35
60.2	99	99	-	-	*	18	81	25	23	32	19
60.29	96	96	-	-	*	17	79	24	23	32	18
60.5	55	55	-	-	*	35	20	9	13	20	13
61	18	18	-	*	*	*8	*5	*	*	11	*
61.0	14	14	-	*	*	*6	*	*	*	*9	*
62	11	11	-	*	*	*	*	*	*	*	*
62.3	*5	*5	-	*	*	*	*	*	*	*	*
63	*7	*7	-	*	*	*	*	*	*	*	*
64	35	35	-	19	*6	*	*6	*	*7	19	*
64.0	21	21	-	18	*	*	*	*	*	13	*
64.9	10	10	-	*	*	*	*	*	*	*5	*
65-71	2,050	-	2,050	10	1,188	621	231	296	424	916	413
65	551	-	551	*	252	231	64	83	131	229	108
65.0	*7	-	*7	*	*	*	*	*	*	*	*
65.1	*5	-	*5	*	*	*	*	*	*	*	*
65.2	52	-	52	*	41	*8	*	*8	*8	23	12
65.25	*9	-	*9	*	*8	*	-	*	*	*	*
65.29	41	-	41	*	32	*7	*	*6	*6	19	10
65.3	21	-	21	*	14	*	*	*5	*	9	*
65.39	20	-	20	*	13	*	*	*	*	9	*
65.4	85	-	85	*	49	28	*7	12	23	32	18
65.41	13	-	13	*	9	*	*	*	*	*7	*
65.49	73	-	73	*	40	25	*7	11	21	25	16
65.5	*7	-	*7	-	*	*	*	*	*	*	*
65.6	338	-	338	-	110	178	50	49	83	144	62
65.61	292	-	292	-	93	154	44	44	71	121	56
65.62	9	-	9	-	*6	*	*	*	*	*	*
65.63	35	-	35	-	10	20	*	*	*9	18	*5
65.8	26	-	26	*	17	*7	*	*	*6	10	*7
65.81	*6	-	*6	-	*	*	*	*	*	*	*
65.89	20	-	20	*	12	*6	*	*	*6	*7	*5

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
65.9	*9	—	*9	*	*7	*	—	*	*	*	*
65.91	*7	—	*7	*	*6	*	—	*	*	*	*
66	422	—	422	*	413	*8	*	51	68	217	86
66.0	*5	—	*5	—	*5	*	—	*	*	*	*
66.2	*6	—	*6	—	*5	*	—	*	*	*	*
66.3	352	—	352	—	349	*	—	39	58	184	71
66.32	248	—	248	—	246	*	—	26	34	136	53
66.39	103	—	103	—	101	*	—	13	24	48	18
66.4	*9	—	*9	*	*7	*	*	*	*	*	*
66.5	9	—	9	—	*6	*	*	*	*	*	*
66.51	*8	—	*8	—	*5	*	*	*	*	*	*
66.6	33	—	33	*	33	*	*	*5	*	15	*7
66.61	*7	—	*7	*	*6	*	—	*	*	*	*
66.62	22	—	22	—	22	—	—	*	*	11	*
67	18	—	18	—	12	*	*	*	*	*7	*
67.5	*6	—	*6	—	*6	—	—	*	*	*	*
67.59	*6	—	*6	—	*6	—	—	*	*	*	*
68	680	—	680	*	354	262	64	99	147	298	136
68.1	14	—	14	*	*7	*	*	*	*	*6	*
68.12	*6	—	*6	—	*	*	*	*	*	*	*
68.16	*5	—	*5	—	*	*	*	*	*	*	*
68.2	47	—	47	*	37	*9	*	10	*	21	11
68.29	42	—	42	*	34	*7	*	*9	*	19	10
68.3	40	—	40	*	22	15	*	11	*6	14	*9
68.31	*6	—	*6	—	*	*	—	*	*	*	*
68.39	34	—	34	*	18	13	*	10	*	12	*8
68.4	374	—	374	—	188	155	31	53	91	167	63
68.5	197	—	197	—	98	75	24	22	41	86	49
68.51	60	—	60	—	33	23	*	*6	17	28	10
68.59	136	—	136	—	65	52	20	16	25	57	39
68.6	*6	—	*6	—	*	*	*	*	*	*	*
69	98	—	98	*	82	10	*	23	15	41	19
69.0	53	—	53	*	43	*7	*	12	*7	23	11
69.02	31	—	31	—	31	*	—	*6	*	13	*8
69.09	22	—	22	—	12	*7	*	*5	*	10	*
69.1	*7	—	*7	*	*5	*	*	*	*	*	*
69.19	*7	—	*7	*	*5	*	*	*	*	*	*
69.5	22	—	22	*	21	*	*	*7	*	*8	*
69.52	17	—	17	*	17	—	—	*5	*	*5	*
69.9	9	—	9	—	9	*	*	*	*	*	*
69.96	*7	—	*7	—	*7	*	—	*	*	*	*
70	246	—	246	*	57	96	91	31	56	106	53
70.1	*6	—	*6	*	*	*	*	*	*	*	*
70.3	*9	—	*9	*	*6	*	*	*	*	*	*
70.32	*5	—	*5	—	*	*	*	*	*	*	*
70.5	133	—	133	—	26	60	47	18	32	55	28
70.50	65	—	65	—	10	30	25	*7	16	29	14
70.51	40	—	40	—	*9	17	14	*7	*7	16	10
70.52	27	—	27	—	*7	13	*7	*	*9	11	*
70.7	44	—	44	*	10	14	20	*	*9	20	11
70.77	28	—	28	—	*	10	16	*	*7	14	*6
70.79	*8	—	*8	*	*	*	*	*	*	*	*
70.9	39	—	39	*	*8	16	14	*5	*7	17	9
70.92	38	—	38	—	*8	16	14	*5	*7	17	9
71	35	—	35	*	18	*9	*6	*	*6	18	*8
71.0	*6	—	*6	*	*	*	*	*	*	*	*
71.09	*6	—	*6	*	*	*	*	*	*	*	*
71.3	11	—	11	*	9	*	*	*	*	*6	*
71.7	*9	—	*9	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
72-75	6,971	—	6,971	*9	6,947	16	—	1,056	1,490	2,633	1,793
72	322	—	322	*	322	*	—	35	57	142	88
72.0	21	—	21	—	21	—	—	*	*	13	*
72.1	34	—	34	—	34	—	—	*	*	22	*
72.2	*6	—	*6	—	*6	*	—	*	*	*5	*
72.4	*5	—	*5	—	*5	—	—	*	*	*	—
72.7	249	—	249	*	249	*	—	28	47	94	80
72.71	113	—	113	*	112	—	—	15	20	44	34
72.79	137	—	137	*	137	*	—	13	27	51	47
73	3,047	—	3,047	*	3,037	*6	—	424	686	1,172	766
73.0	941	—	941	*	937	*	—	125	255	333	229
73.01	147	—	147	—	147	*	—	15	39	56	38
73.09	793	—	793	*	790	*	—	110	216	277	191
73.1	11	—	11	—	11	—	—	*	*	*	*
73.4	573	—	573	*	572	*	—	79	118	215	160
73.5	1,000	—	1,000	*	997	*	—	145	199	399	257
73.51	*7	—	*7	*	*7	—	—	*	*	*	*
73.59	993	—	993	*	990	*	—	144	197	397	255
73.6	516	—	516	*	514	*	—	70	112	218	116
74	1,238	—	1,238	*	1,231	*5	—	183	221	534	300
74.0	14	—	14	—	14	—	—	*	*	*6	*
74.1	1,216	—	1,216	*	1,210	*5	—	179	216	525	296
75	2,364	—	2,364	*	2,357	*	—	414	525	785	640
75.1	12	—	12	—	12	*	—	*	*	*6	*
75.3	1,026	—	1,026	*	1,022	*	—	198	252	323	254
75.32	179	—	179	*	178	*	—	21	49	60	49
75.34	676	—	676	*	673	*	—	154	172	219	131
75.35	117	—	117	*	117	*	—	16	25	27	50
75.37	52	—	52	—	52	*	—	*5	*7	16	24
75.4	45	—	45	—	45	—	—	*7	12	10	16
75.5	*8	—	*8	—	*8	—	—	*	*	*	*
75.51	*7	—	*7	—	*7	—	—	*	*	*	*
75.6	1,233	—	1,233	*	1,231	*	—	199	249	422	362
75.61	34	—	34	*	33	—	—	*5	*6	13	9
75.62	99	—	99	*	98	*	—	15	19	39	26
75.69	1,100	—	1,100	*	1,099	*	—	179	224	370	327
75.7	34	—	34	—	34	—	—	*7	*	20	*
76-84	4,040	1,902	2,138	168	945	1,349	1,578	786	909	1,460	885
76	97	59	38	*8	65	14	10	18	12	45	23
76.4	*5	*	*	*	*	*	*	*	*	*	*
76.6	25	*9	16	*	21	*	*	*5	*	12	*5
76.65	*7	*	*	*	*6	*	*	*	*	*	*
76.7	42	37	*	*	29	*8	*	*6	*7	18	11
76.75	*6	*6	*	*	*	*	*	*	*	*	*
76.76	19	17	*	*	15	*	*	*	*	*9	*
76.79	*6	*5	*	*	*	*	*	*	*	*	*
76.9	15	*7	*8	*	*8	*	*	*	*	*7	*
76.92	*9	*5	*	*	*5	*	*	*	*	*	*
77	240	125	115	19	65	81	74	52	53	80	55
77.1	*5	*	*	*	*	*	*	*	*	*	*
77.3	14	*8	*6	*	*	*	*	*	*	*	*
77.4	37	21	17	*	*6	*8	22	*7	*9	12	11
77.49	28	15	14	*	*	*	18	*	*7	*8	9
77.5	10	*	*7	*	*	*	*6	*	*	*	*
77.6	50	32	19	*6	13	17	14	10	11	17	13
77.61	*7	*5	*	*	*	*	*	*	*	*	*
77.65	*8	*	*	*	*	*	*	*	*	*	*
77.67	*7	*	*	*	*	*	*	*	*	*	*
77.68	*9	*6	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
77.69	15	*8	*7	*	*	*5	*	*	*	*5	*
77.7	78	37	41	*	26	34	14	16	17	29	17
77.79	69	31	38	*	24	30	12	14	16	25	14
77.8	38	18	20	*	11	13	14	*8	*8	12	*9
77.81	13	*	*9	*	*	*	*7	*	*	*	*
77.85	*5	*	*	*	*	*	*	*	*	*	*
77.88	*7	*	*	*	*	*	*	*	*	*	*
77.89	*7	*	*	*	*	*	*	*	*	*	*
78	254	123	131	27	66	69	93	51	48	97	59
78.0	31	18	14	*	9	12	*6	*	*8	10	*9
78.05	*9	*	*	*	*	*	*	*	*	*	*
78.07	10	*6	*	*	*	*	*	*	*	*	*
78.1	33	20	13	*	13	10	*7	*9	*7	12	*5
78.13	10	*6	*	*	*	*	*	*	*	*	*
78.17	13	10	*	*	*6	*	*	*	*	*6	*
78.4	40	13	28	*6	*	*	28	*6	*6	18	11
78.41	*5	*	*	*5	*	*	*	*	*	*	—
78.49	32	*8	24	*	*	*	28	*	*5	12	10
78.5	59	28	31	*	14	12	28	16	10	19	14
78.55	37	15	22	*	*	*	24	11	*7	11	*7
78.57	9	*	*	*	*6	*	*	*	*	*	*
78.59	*5	*	*	*	*	*	*	*	*	*	*
78.6	82	40	42	*6	25	30	22	15	15	34	18
78.65	23	11	12	*	*5	*7	10	*5	*5	*8	*
78.67	18	11	*7	*	*5	*9	*	*	*	*8	*
78.69	29	12	18	*	11	12	*6	*	*6	13	*6
79	728	348	380	60	212	170	286	151	157	260	161
79.0	71	35	37	12	20	17	22	17	15	20	19
79.01	*7	*	*	*	*	*	*	*	*	*	*
79.02	17	*5	12	*	*	*	*8	*	*	*6	*
79.05	11	*6	*	*	*	*	*	*	*	*	*
79.06	27	14	13	*	*8	9	*5	*7	*9	*6	*6
79.1	108	50	58	16	26	19	47	25	20	39	23
79.11	15	*6	*8	11	*	*	*	*	*	*	*
79.12	11	*5	*6	*	*	*	*	*	*	*	*
79.15	59	23	36	*	10	10	37	14	12	22	11
79.16	14	10	*	*	*7	*	*	*	*	*6	*
79.3	441	197	244	20	120	107	194	87	97	164	94
79.31	34	12	22	*	*6	10	13	*8	*7	10	*9
79.32	46	26	20	*	18	11	13	*8	11	18	10
79.33	*6	*5	*	*	*	*	*	*	*	*	*
79.35	178	60	118	*	15	25	134	33	34	73	38
79.36	143	70	73	*5	56	51	31	31	36	47	30
79.37	12	*8	*	*	*8	*	*	*	*	*	*
79.39	14	10	*	*	*9	*	*	*	*	*6	*
79.6	52	38	14	*5	27	15	*5	*9	9	22	12
79.62	10	*8	*	*	*5	*	*	*	*	*	*
79.64	*7	*	*	*	*	*	*	*	*	*	*
79.66	19	13	*6	*	9	*6	*	*	*	*7	*
79.7	36	17	19	*	11	*9	14	*8	11	10	*7
79.71	*5	*	*	—	*	*	*	*	*	*	*
79.75	23	10	14	*	*6	*7	9	*6	*8	*5	*
79.8	*9	*5	*	*	*	*	*	*	*	*	*
80	472	246	226	*6	160	196	111	78	102	180	112
80.0	12	*	*8	*	*	*	*6	*	*	*	*
80.06	*7	*	*5	—	*	*	*	*	*	*	*
80.1	31	18	13	*	*9	10	*9	*	*8	11	*8
80.16	14	*8	*6	*	*	*	*6	*	*	*6	*
80.2	24	14	10	*	*7	*8	10	*	*	13	*
80.21	10	*6	*	—	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
80.26	14	*8	*6	*	*5	*	*6	*	*	10	*
80.4	10	*	*5	*	*	*	*	*	*	*	*
80.46	*6	*	*	*	*	*	*	*	*	*	*
80.5	326	162	164	*	122	142	61	48	74	127	77
80.51	324	161	163	*	121	142	60	47	74	125	77
80.6	13	*8	*5	*	*6	*	*	*	*	*5	*
80.7	13	*8	*6	*	*	*5	*	*	*	*	*
80.76	10	*5	*	*	*	*	*	*	*	*	*
80.8	35	22	13	*	*7	15	12	*9	*8	11	*7
80.81	*7	*	*	*	*	*	*	*	*	*	*
80.85	*5	*	*	*	*	*	*	*	*	*	*
80.86	16	9	*6	*	*	*8	*	*	*	*6	*
81	1,721	706	1,014	19	246	632	824	340	428	596	357
81.0	317	146	171	*7	91	145	74	47	70	125	76
81.02	115	59	56	—	37	63	15	17	26	49	23
81.03	15	*8	*7	*	*	*7	*	*	*	*	*5
81.04	*6	*	*	*	*	*	*	*	*	*	*
81.05	17	*8	10	*	*6	*5	*	*	*	*7	*
81.06	26	12	14	*	12	10	*	*	*8	*9	*5
81.07	13	*6	*6	—	*	*	*6	*	*	*	*
81.08	122	49	73	*	26	53	42	17	25	46	33
81.1	18	10	*7	*	*5	*7	*	*	*	*	*
81.11	*8	*6	*	—	*	*	*	*	*	*	*
81.3	17	*6	10	*	*	*8	*	*	*	*8	*
81.38	11	*	*6	*	*	*	*	*	*	*	*
81.4	30	16	14	*	17	*8	*5	*6	*6	11	*7
81.45	9	*6	*	*	*7	*	*	*	*	*	*
81.47	13	*7	*6	*	*6	*	*	*	*	*	*
81.5	1,039	387	653	*	70	350	616	232	287	331	189
81.51	234	99	135	—	15	83	135	61	58	69	45
81.52	240	88	152	*	41	66	130	49	68	77	46
81.53	46	20	27	—	*	11	32	13	9	14	10
81.54	478	167	311	*	*9	177	292	100	142	158	77
81.55	40	12	28	—	*	12	26	*9	9	12	10
81.6	144	67	77	*	37	67	37	11	22	65	46
81.62	117	56	61	*	33	55	28	*7	18	53	39
81.63	19	*7	12	*	*	*9	*8	*	*	10	*
81.8	47	21	26	*	*	15	27	*9	11	15	12
81.80	13	*6	*7	—	*	*	*9	*	*	*	*
81.81	16	*8	*9	*	*	*6	10	*	*	*7	*
81.83	12	*6	*6	*	*	*	*7	*	*	*	*
81.9	103	50	53	*	14	29	55	27	25	34	17
81.91	57	34	23	*	*9	15	30	18	12	19	*9
81.92	38	12	26	*	*	11	24	*8	12	11	*6
82	22	15	*7	*	12	*7	*	*	*	11	*
82.0	*8	*5	*	*	*	*	*	*	*	*	*
82.01	*6	*	*	*	*	*	*	*	*	*	*
82.4	*8	*7	*	*	*5	*	*	*	*	*	*
83	246	136	110	24	65	79	78	53	50	83	59
83.0	28	21	*7	*	*7	11	*6	*	*6	12	*6
83.03	*6	*5	*	*	*	*	*	*	*	*	*
83.09	19	14	*5	*	*5	*7	*	*	*	9	*
83.1	27	15	12	*5	9	*9	*	*5	*5	9	*8
83.13	*7	*	*	*	*	*	*	*	*	*	*
83.14	14	*9	*6	*	*6	*5	*	*	*	*5	*
83.2	16	*8	*9	*	*	*6	*5	*	*	*6	*
83.21	16	*8	*9	*	*	*6	*5	*	*	*6	*
83.3	24	*8	15	*	*	*9	*8	*	*6	9	*
83.39	19	*7	12	*	*	*7	*8	*	*	*8	*
83.4	33	17	16	*	13	*9	9	10	*5	10	*9

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
83.45	24	13	10	*	10	*5	*7	*8	*	*7	*6
83.5	11	*5	*6	*	*	*	*	*	*	*	*
83.6	58	34	24	*	13	16	27	12	10	20	15
83.63	33	16	18	—	*	*9	24	*8	*6	10	10
83.64	13	11	*	*	*8	*	*	*	*	*6	*
83.65	10	*6	*	*	*	*	*	*	*	*	*
83.7	*8	*	*5	*	*	*	*	*	*	*	*
83.8	28	18	10	*6	*5	9	*8	*8	*6	*8	*5
83.82	*6	*	*	*	*	*	*	*	*	*	*
83.85	*9	*6	*	*	*	*	*	*	*	*	*
83.88	9	*6	*	*	*	*	*	*	*	*	*
83.9	12	*6	*7	*	*	*5	*	*	*	*	*
84	261	145	116	*	53	103	101	40	57	107	56
84.0	*8	*6	*	*	*	*	*	*	*	*	*
84.01	*6	*	*	*	*	*	*	*	*	*	*
84.1	129	80	49	*	13	48	68	22	24	62	21
84.11	54	35	19	*	*6	25	23	10	10	25	10
84.12	13	*8	*5	—	*	*5	*7	*	*	*6	*
84.15	28	19	*9	*	*	10	14	*	*6	12	*
84.17	30	14	15	*	*	*6	22	*5	*	17	*
84.3	19	12	*7	*	*6	*8	*5	*	*	10	*
84.5	101	44	57	*	30	44	27	12	27	31	31
84.51	73	30	42	*	22	31	19	10	19	21	22
84.52	28	14	14	*	*8	13	*8	*	*8	10	*8
85-86	1,498	701	797	154	440	468	436	298	283	682	235
85	187	*	186	*	43	83	59	40	35	73	39
85.0	10	*	10	*	*	*	*	*	*	*7	*
85.1	12	*	12	—	*	*	*6	*	*	*	*
85.12	*6	*	*6	—	*	*	*	*	*	*	*
85.2	15	—	15	*	*	*6	*7	*	*	*5	*
85.21	*9	—	*9	*	*	*	*	*	*	*	*
85.23	*5	—	*5	—	*	*	*	*	*	*	*
85.3	16	*	15	*	*8	*6	*	*5	*	*5	*
85.32	13	*	13	*	*8	*	*	*	*	*	*
85.4	80	*	80	*	10	31	38	15	16	30	18
85.41	25	*	24	—	*	11	*9	*	*	10	*
85.43	48	*	47	*	*	16	28	*9	*9	18	12
85.5	*6	*	*6	*	*	*	*	*	*	*	*
85.7	13	*	13	—	*	9	*	*	*	*	*
85.8	11	*	11	*	*	*6	*	*	*	*	*
85.89	*5	*	*	—	*	*	*	*	*	*	*
85.9	20	*	20	*	*	14	*	*	*	*8	*
85.95	13	—	13	—	*	*8	*	*	*	*	*
86	1,311	699	611	152	397	385	376	258	248	609	196
86.0	379	191	188	40	116	127	97	82	73	163	62
86.01	10	*	*5	*	*	*	*	*	*	*	*
86.04	192	105	87	19	72	61	41	36	39	84	33
86.05	48	21	28	*6	13	18	11	10	10	20	*9
86.06	10	*	*6	*	*	*	*	*	*	*	*
86.07	81	38	43	*8	14	28	31	22	13	37	9
86.09	35	18	16	*	*8	12	10	*7	*8	15	*
86.1	29	18	11	*	*	13	10	*7	*5	12	*5
86.11	29	18	11	*	*	13	10	*7	*5	12	*5
86.2	424	235	189	25	103	131	166	91	91	187	54
86.22	302	170	132	19	84	104	95	61	60	144	37
86.23	*7	*	*	*	*	*	*	*	*	*	*
86.27	47	24	22	—	*	*7	36	17	16	12	*
86.28	66	37	29	*5	13	16	31	11	14	27	13
86.3	85	33	52	12	33	21	19	13	18	42	12
86.4	*9	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
86.5	151	92	58	9	63	34	45	27	32	50	41
86.59	151	92	58	9	63	34	45	27	32	50	41
86.6	115	70	45	33	37	29	16	15	12	81	*6
86.62	*7	*	*	*	*	*	*	*	*	*6	*
86.66	15	11	*	*	*	*	*	*	*	14	—
86.67	20	12	*8	*7	*8	*	*	*	*	17	*
86.69	62	36	27	19	18	14	11	11	10	36	*5
86.7	43	27	15	*8	13	11	11	*7	*7	22	*6
86.74	28	17	10	*6	*8	*6	*9	*	*	15	*
86.8	72	26	45	21	27	18	*7	13	*7	45	*7
86.83	24	*	22	*	10	11	*	*8	*	10	*
86.84	32	17	14	19	11	*	—	*	*	30	*
86.89	12	*6	*6	*	*	*	*	*	*	*	*5
87-99	13,147	6,305	6,842	1,012	2,501	3,737	5,898	3,882	2,591	4,054	2,620
87	887	409	478	56	191	240	399	286	177	214	210
87.0	330	161	169	25	53	75	177	153	54	50	73
87.03	324	158	166	24	51	73	176	151	53	49	72
87.2	18	*9	10	*	*	*8	*6	*	*	9	*
87.21	11	*5	*5	—	*	*	*	*	*	*6	*
87.24	*5	*	*	*	*	*	*	*	*	*	*
87.3	*8	*	*5	*	*	*	*	*	*	*	*6
87.39	*7	*	*	*	*	*	*	*	—	*	*6
87.4	223	113	110	12	39	67	105	67	59	41	55
87.41	133	62	71	*	18	40	72	58	30	28	17
87.44	82	46	36	*8	18	25	31	*9	26	12	36
87.49	*8	*	*	*	*	*	*	*	*	*	*
87.5	146	44	102	*	47	48	50	23	25	60	38
87.51	*6	*	*	*	*	*	*	*	*	*	*
87.53	133	39	94	*	45	43	44	19	24	56	34
87.54	*6	*	*	*	*	*	*	*	*	*	*
87.6	30	14	16	*6	*	*7	12	*8	*	*6	14
87.61	*7	*	*	*	*	*	*	*	*	*	*
87.62	10	*	*7	*	*	*	*	*	*	*	*
87.69	*5	*	*	*	*	*	*	*	*	*	*
87.7	128	62	66	10	41	32	44	29	32	47	19
87.73	11	*	*7	*	*5	*	*	*	*	*	*
87.74	83	40	43	*	26	22	32	19	21	31	12
87.75	*6	*	*	*	*	*	*	*	*	*	*
87.76	*9	*5	*	*7	*	*	*	*	*	*6	*
87.79	*7	*	*	*	*	*	*	*	*	*	*
88	3,641	1,873	1,768	133	510	1,254	1,744	1,077	739	1,151	674
88.0	267	119	148	17	69	75	106	133	46	37	52
88.01	263	115	147	17	69	72	105	132	45	34	51
88.1	18	*9	9	*	*8	*	*	*	*	*6	*6
88.19	12	*6	*7	*	*	*	*	*	*	*	*
88.2	16	11	*	*	*6	*	*	*	*7	*	*
88.3	112	55	56	12	29	27	43	57	27	15	13
88.38	103	52	51	9	28	25	41	56	26	12	10
88.4	496	258	238	12	61	158	265	104	119	187	86
88.41	130	61	68	*	22	45	59	32	27	51	20
88.42	130	69	61	*	11	40	76	21	36	53	21
88.43	23	10	12	*	*	*6	*9	*	*	*7	*8
88.44	13	*6	*7	*	*	*5	*6	*5	*	*	*
88.45	30	13	17	*	*	10	18	*	*7	15	*
88.47	31	18	13	*	*	10	18	10	9	*8	*
88.48	106	64	42	*	12	32	62	23	25	38	20
88.49	33	16	16	*	*6	9	17	*	9	11	*8
88.5	1,614	913	701	*7	144	676	787	305	351	645	312
88.50	*5	*	*	*	*	*	*	*	*	*	—
88.51	19	*8	11	*	*	*5	12	*5	*5	*7	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
88.52	*5	*	*	*	*	*	*	*	*	*	*
88.53	592	324	268	*	57	254	279	117	128	243	104
88.54	25	14	11	*	*	11	11	*7	*	12	*
88.55	15	9	*6	*	*	*6	*6	*	*	*	*
88.56	908	520	388	*	75	372	460	156	202	356	194
88.57	44	30	15	*	*	22	17	*9	10	19	*6
88.6	23	13	10	*	*6	*8	*9	*7	*5	*7	*
88.67	11	*6	*	*	*	*	*	*	*	*	*
88.7	813	368	444	51	133	223	405	325	133	205	150
88.71	52	24	27	*	*	12	33	22	*8	11	11
88.72	502	238	264	25	62	153	263	187	75	159	81
88.73	*7	*	*	*	*	*	*	*	*	*	*
88.74	14	*8	*6	*	*	*5	*	*	*	*	*
88.75	37	18	19	*6	*8	*6	17	20	*	*	10
88.76	95	38	57	11	22	24	38	41	17	*9	28
88.77	77	32	45	*	14	18	44	37	21	9	10
88.78	12	—	12	—	12	—	*	*5	*	*	*
88.79	17	*6	10	*	*7	*	*	*9	*	*	*
88.9	283	127	156	31	54	80	117	141	47	47	46
88.91	169	75	94	21	30	46	72	85	28	25	31
88.92	*5	*	*	*	*	*	*	*	*	*	*
88.93	39	19	20	*	11	*9	16	20	*8	*7	*
88.94	20	9	10	*	*	*6	*8	12	*	*	*
88.97	46	19	27	*	*8	16	18	20	*9	11	*6
89	684	331	353	62	114	219	290	240	111	157	176
89.0	44	26	18	*	*8	19	15	*	*	*5	36
89.06	18	10	*8	—	*	*7	*8	—	—	*	18
89.07	*7	*	*	—	*	*	*	—	—	—	*7
89.08	*7	*	*	*	*	*	*	*	*	*	*
89.09	12	*8	*	—	*	*6	*	*	—	*	11
89.1	97	43	54	34	21	18	23	41	*5	30	21
89.14	59	25	34	19	10	11	19	24	*	15	17
89.15	*8	*	*	*	*	*	*	*	*	*	*
89.19	27	12	15	12	10	*	*	15	*	10	*
89.2	17	*7	11	*	*	*	10	*	*6	*5	*
89.22	9	*	*6	—	*	*	*7	*	*5	*	*
89.3	21	11	9	*9	*	*	*	*5	*	*9	*
89.38	*5	*	*	*	*	*	*	*	*	*	*
89.39	10	*6	*	*6	*	*	*	*	*	*	*
89.4	146	63	83	*	16	59	70	48	15	33	49
89.41	45	20	25	*	*6	22	17	19	*	11	13
89.44	92	40	52	*	9	37	45	27	11	19	35
89.45	*7	*	*	*	*	*	*5	*	*	*	*
89.5	251	123	128	10	42	79	119	121	55	40	36
89.50	*6	*	*	*	*	*	*	*	—	*	*
89.52	103	55	49	*9	24	36	35	29	29	21	25
89.54	117	55	62	*	15	34	67	81	14	13	9
89.59	25	11	13	*	*	*8	15	*6	11	*6	*
89.6	108	57	51	*5	18	36	49	19	24	35	30
89.61	11	*7	*	*	*	*	*	*	*	*	*
89.62	12	*	*7	*	*	*	*	*	*	*6	*
89.64	23	13	10	*	*	*8	11	*	*6	*7	*6
89.65	43	22	22	*	*7	14	20	*7	*6	15	15
89.68	15	*8	*7	*	*	*	*6	*	*6	*5	*
90	21	9	12	*	*5	*	*7	*	*5	*8	*7
90.5	12	*6	*7	*	*	*	*	*	*	*	*7
90.59	*9	*	*5	*	*	*	*	*	*	*	*6
91	*9	*	*5	*	*	*	*	*	*	*	*
92	234	107	128	*6	31	83	114	90	26	57	62
92.0	99	43	57	*	11	35	51	36	*8	21	34

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see “Methods” section]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
92.02	*9	*	*	*	*	*	*	*	*	*	*
92.04	10	*	*6	*	*	*	*	*	*	*	*
92.05	75	32	43	*	*	28	42	26	*6	16	27
92.1	52	22	31	*	*6	18	26	28	*	10	11
92.14	19	*7	11	*	*	*6	9	12	*	*	*
92.15	25	10	15	*	*	9	13	13	*	*	*6
92.2	77	41	36	*	13	27	37	23	14	25	16
92.24	13	*7	*6	*	*	*	*8	*6	*	*	*
92.27	13	*7	*7	–	*	*	*7	*	*	*	*
92.29	45	25	20	*	*8	18	19	12	9	16	*9
92.3	*5	*	*	*	*	*	*	*	*	*	*
93	1,474	664	811	186	183	331	774	437	251	467	319
93.0	35	18	17	*	*5	9	19	23	*	*	*7
93.01	30	14	15	*	*	*8	17	22	*	*	*
93.1	24	10	14	*	*	*6	15	20	*	*	*
93.11	*5	*	*	*	*	*	*	*5	–	*	*
93.12	*8	*	*	*	*	*	*6	*6	*	*	–
93.17	*6	*	*	*	*	*	*	*6	–	*	–
93.2	33	15	19	*	*	10	18	24	*	*6	*
93.22	25	9	16	*	*	*6	15	20	*	*	*
93.3	357	143	214	*	32	80	241	79	59	144	75
93.38	43	18	25	–	*	10	30	*	–	21	18
93.39	309	122	187	*	28	67	209	73	57	121	58
93.4	14	*5	*8	*	*	*	*8	*	*	*	*
93.46	*7	*	*6	*	*	*	*6	*	*	*	*
93.5	102	54	48	26	24	23	29	29	20	41	12
93.53	17	*8	*9	*	*	*	*6	*7	*	*	*
93.54	24	12	12	*	*6	*6	*9	*6	*8	*6	*
93.57	27	15	12	16	*6	*	*	*6	*	17	*
93.59	20	10	11	*	*	*9	*7	*6	*	*8	*
93.7	36	17	19	*	*	10	21	*7	*8	10	11
93.75	35	17	18	*	*	9	21	*6	*8	10	11
93.8	397	170	227	*9	51	92	245	92	60	160	85
93.81	48	25	23	*	16	12	18	*6	*5	20	17
93.83	216	91	124	*5	24	50	137	49	41	85	40
93.89	128	52	76	*	*8	29	89	37	13	51	28
93.9	473	232	241	141	54	101	177	158	93	97	126
93.90	112	56	56	11	11	31	59	41	27	28	17
93.94	175	86	89	73	17	29	56	66	38	26	45
93.96	167	79	88	51	22	37	56	44	25	38	60
93.99	9	*	*	*	*	*	*	*	*	*	*
94	712	424	288	37	412	219	43	355	133	149	75
94.0	9	*	*	*	*	*	*	*	–	*7	*
94.1	*7	*	*	*	*	*	*	*5	–	*	–
94.2	184	97	86	12	103	53	16	142	16	22	*
94.22	*7	*	*	*	*	*	*	*6	*	*	–
94.23	48	26	21	*	31	10	*	46	–	*	–
94.25	102	56	46	*6	61	30	*	81	*9	10	*
94.27	26	10	16	–	*7	10	9	*8	*7	9	*
94.3	64	32	32	*9	39	12	*	28	25	10	–
94.38	23	11	12	*	16	*5	*	23	–	–	–
94.39	38	20	19	*7	22	*7	*	*	25	10	–
94.4	93	49	44	12	54	22	*	31	29	33	–
94.42	11	*	*6	*	*6	*	*	*	*	*	–
94.44	70	36	34	*8	40	19	*	16	26	29	–
94.6	355	237	118	*	207	129	18	145	63	74	72
94.61	*6	*	*	*	*	*	*	*	*	*	*
94.62	179	123	56	–	81	83	16	70	39	32	38
94.63	10	*7	*	–	*	*5	*	–	*	*	*
94.64	*7	*	*	*	*5	*	*	*	*	*	*
94.65	84	49	35	*	63	20	*	36	10	19	19

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
94.66	*7	*	*	—	*6	*	*	*	*	*	*
94.67	*7	*	*	*	*5	*	—	*	*	*	*
94.68	49	38	12	—	34	15	*	32	*	*7	*8
94.69	*5	*	*	—	*	*	—	*	*	*	*
95	10	*	*	*9	*	*	*	*	*	*6	*
95.4	*9	*	*	*8	*	*	*	*	*	*6	*
95.47	*7	*	*	*7	—	*	—	*	*	*5	*
96	1,621	774	847	173	324	384	739	370	340	547	365
96.0	616	311	306	51	83	167	315	132	124	220	141
96.04	511	258	253	45	67	138	261	107	103	190	110
96.05	10	*6	*	*	*	*	*6	*	*	*	*
96.07	86	41	45	*	14	23	44	21	17	22	27
96.08	*6	*	*	*	*	*	*	*	*	*	*
96.2	*8	*	*	*	*	*	*	*	*	*	*
96.3	22	10	11	*	*7	*6	*6	*6	*	*7	*
96.33	11	*	*6	*	*6	*	*	*	*	*	*
96.38	*5	*	*	*	*	*	*	*	*	*	*
96.4	112	*	109	*	107	*	*	12	29	34	37
96.49	110	*	108	*	107	*	*	11	28	34	37
96.5	42	26	16	*	11	10	17	*9	10	15	*8
96.52	*5	*	*	*	*	*	*	*	*	*	*
96.56	14	*9	*5	*	*	*	*9	*	*	*	*
96.59	21	13	*8	*	*8	*6	*	*	*	*9	*
96.6	136	64	72	26	11	25	74	68	32	21	15
96.7	684	356	328	87	103	175	320	141	140	247	156
96.70	*8	*	*	*	*	*	*	*	*	*	*
96.71	432	224	208	54	73	100	205	90	94	150	98
96.72	245	129	116	32	28	73	112	50	45	93	57
97	123	62	61	14	24	30	55	26	26	50	21
97.0	46	24	22	*6	*	10	27	9	*8	18	10
97.02	31	15	16	*	*	*6	20	*6	*5	13	*7
97.03	*5	*	*	*	*	*	*	*	*	*	—
97.05	*8	*	*	*	*	*	*	*	*	*	*
97.2	11	*6	*	*	*	*	*	*	*	*	*
97.23	10	*6	*	*	*	*	*	*	*	*	*
97.3	*5	*	*	*	*	*	*	*	*	*	*
97.4	11	*5	*6	*	*	*	*	*	*	*	*
97.49	*7	*	*	*	*	*	*	*	*	*	*
97.5	12	*5	*7	*	*	*	*6	*	*	*7	*
97.6	16	*7	*8	*	*	*	*7	*	*	*6	*
97.62	14	*7	*7	*	*	*	*6	*	*	*6	*
97.8	15	9	*5	*	*	*	*	*	*	*8	*
98	28	13	15	*	*8	*8	*7	*7	*	10	*7
98.0	10	*5	*	*	*	*	*	*	*	*	*
98.02	*7	*	*	*	*	*	*	*	*	*	*
98.5	10	*	*	*	*	*	*	*	*	*	*
98.51	10	*	*	*	*	*	*	*	*	*	*
99	3,703	1,632	2,072	325	695	960	1,723	990	779	1,237	698
99.0	1,815	759	1,055	69	229	464	1,053	354	376	718	367
99.00	26	12	14	*	*	*9	14	*6	10	*6	*
99.02	54	19	35	*	*	23	26	11	11	16	16
99.03	11	*	*7	*	*	*	*8	*	*	*	*
99.04	1,473	599	873	49	186	357	881	285	296	600	292
99.05	86	43	43	12	16	28	30	15	17	31	23
99.07	150	74	76	*	15	38	91	33	31	53	32
99.09	*5	*	*	*	*	*	*	*	*	*	*
99.1	392	144	248	46	136	87	123	135	71	96	90
99.10	42	21	22	*	*	20	17	9	*6	14	13
99.11	78	*	78	*	77	*	*	18	12	26	23
99.14	14	*6	*8	*5	*	*	*	*6	*	*	*

See footnotes at end of table.

Table 48. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex, age, and geographic region: United States, 2004—Con.

[Discharges of inpatients from nonfederal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see "Methods" section]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	North-east	Midwest	South	West
Number of procedures in thousands											
99.15	142	63	79	32	23	32	55	42	32	39	28
99.17	31	14	17	*	*7	12	10	17	*8	*	*
99.18	38	17	21	*	9	*6	18	22	*	*	11
99.19	48	24	24	*	11	15	22	20	11	*7	10
99.2	1,157	555	601	134	267	339	416	429	268	304	155
99.20	137	93	45	*	10	64	63	23	33	55	26
99.21	310	135	175	40	97	64	109	137	76	61	35
99.23	96	34	62	*5	14	27	50	30	27	28	10
99.25	235	131	104	50	40	89	55	60	27	99	50
99.29	367	156	211	38	104	89	136	173	102	58	34
99.3	9	*6	*	*	*	*	*	*	*	*	*
99.4	21	*	20	*	20	*	-	*	*	9	*6
99.47	16	*	16	*	16	-	-	*	*	*8	*
99.5	32	15	17	*7	*6	*8	11	12	10	*7	*
99.52	*9	*	*5	*	*	*	*	*	*	*	-
99.55	21	11	10	*6	*	*	*7	*8	*6	*5	*
99.6	162	96	66	*	15	47	97	33	37	54	37
99.60	70	42	28	*	*8	20	41	12	17	24	17
99.61	23	15	*9	*	*	*7	15	*6	*6	*7	*
99.62	59	33	26	*	*	17	37	13	13	21	13
99.69	*7	*	*	*	*	*	*	*	*	*	*
99.7	35	15	21	*	14	*8	11	12	*	14	*5
99.71	12	*	*8	*	*	*	*	*	*	*5	*
99.74	*8	*5	*	*	*	*	*	*	*	*	*
99.77	11	*	*8	*	*	*	*	*	*	*	*
99.8	69	35	34	64	*	*	*	*7	*9	25	29
99.82	*6	*	*	*6	-	-	-	-	*	*	*
99.83	59	30	29	58	*	*	*	*7	*7	20	25
99.9	12	*6	*6	*	*	*	*6	*	*	*6	*
99.99	10	*	*6	*	*	*	*5	*	*	*5	*

* Figure does not meet standard of reliability or precision.

- Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see "Methods" section.

Appendix

Technical Notes

Abstract Form

Figure I shows the medical abstract form used for the 2004 NHDS. The hospital number, date of birth, and patient ZIP code are confidential information and are not released to the public, although they can be used with privacy safeguards through the NCHS Research Data Center.

Population Estimates

Population estimates used to compute rates are shown in Table I. These estimates are for the civilian population of the United States, including institutionalized persons, as of July 1, 2004. The data are postcensal estimates based on the 2000 census and are from unpublished tabulations provided by the U.S. Census Bureau.

Reliability of Estimates

The statistics presented in this report are based on a sample survey, and therefore may differ from the figures that would be obtained if a complete census had been taken. The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than an entire universe is surveyed. Standard errors for the estimates from the NHDS were calculated with SUDAAN software, which takes into account the complex sample design. A description of the software and the approach it uses has been published (22).

The chances are about 32 in 100 that an estimate from the sample would differ from a complete census by more than the standard error. The chances are 5 in 100 that the difference would be more than twice the standard error, and about 1 in 100 that the difference would be more than 2.5 times as large as the standard error. The relative standard error (RSE) expresses the standard error as a percent of the estimate and can be multiplied by the estimate to obtain the standard error.

Table I. Civilian population by age, geographic region, and sex: United States, 2004

Age and geographic region	Both sexes	Male	Female
All ages			
Total	292,414	143,480	148,934
Region:			
Northeast	54,515	26,423	28,091
Midwest	65,641	32,231	33,409
South	105,223	51,445	53,778
West	67,036	33,381	33,655
Under 15 years			
Total	60,822	31,123	29,699
Under 1 year	4,077	2,085	1,992
1–4 years	15,994	8,178	7,817
5–14 years	40,751	20,860	19,890
Region:			
Northeast	10,584	5,414	5,170
Midwest	13,384	6,851	6,533
South	22,098	11,300	10,798
West	14,755	7,558	7,197
15–44 years			
Total	124,644	62,789	61,855
15–24 years	41,184	21,007	20,177
25–34 years	39,599	19,967	19,632
35–44 years	43,860	21,814	22,046
Region:			
Northeast	22,782	11,378	11,404
Midwest	27,803	14,034	13,768
South	44,793	22,425	22,368
West	29,266	14,952	14,315
45–64 years			
Total	70,655	34,415	36,240
45–54 years	41,578	20,417	21,161
55–64 years	29,077	13,998	15,079
Region:			
Northeast	13,728	6,623	7,104
Midwest	16,071	7,887	8,184
South	25,270	12,235	13,035
West	15,586	7,670	7,916
65 years and over			
Total	36,294	15,154	21,140
65–74 years	18,463	8,428	10,036
75–84 years	12,971	5,218	7,753
85 years and over	4,860	1,508	3,352
Region:			
Northeast	7,421	3,008	4,413
Midwest	8,383	3,459	4,923
South	13,061	5,484	7,577
West	7,428	3,202	4,227

NOTE: Unpublished 2000-based postcensal estimates from the U.S. Census Bureau.

To derive approximate standard errors for the estimates in this report, first calculate the approximate RSE of the estimate using the parameters shown in Table II. Then multiply the approximate RSE by the estimate. The approximate RSE of an estimate X , may be estimated from the formula:

$$\text{RSE}(X) = \sqrt{a + b/X}$$

where a and b are defined in Table II. When multiplied by 100, $\text{RSE}(X)$ is expressed as a percent of X .

The approximate RSE for a percent, 100p, in which the numerator is a

Notice – All information which would permit identification of an individual or an establishment will be held confidential, will be used only by persons engaged in and for the purposes of the survey, and will not be disclosed or released to other persons or used for any other purpose. Public reporting burden of this collection of information is estimated to average 4 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Reports Clearance Officer, 1600 Clifton Road, MS D-24, Atlanta, GA 30333, ATTN: PRA (0920-0212)

FORM **HDS-1**
(3-27-2003)

U.S. DEPARTMENT OF COMMERCE
Economics and Statistics Administration
U.S. CENSUS BUREAU
ACTING AS COLLECTING AGENT FOR
DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION
NATIONAL CENTER FOR HEALTH STATISTICS

MEDICAL ABSTRACT – NATIONAL HOSPITAL DISCHARGE SURVEY

A. PATIENT IDENTIFICATION

1. Hospital number	<input type="text"/>	4. Date of admission	Month <input type="text"/> <input type="text"/> – Day <input type="text"/> <input type="text"/> – Year <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
2. HDS number	<input type="text"/>	5. Date of discharge	Month <input type="text"/> <input type="text"/> – Day <input type="text"/> <input type="text"/> – Year <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
3. (Item deleted)		6. Residence ZIP Code	<input type="text"/>

B. PATIENT CHARACTERISTICS

7. Date of birth	Month <input type="text"/> <input type="text"/> – Day <input type="text"/> <input type="text"/> – Year <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>	11. Race – Mark all that apply 1 <input type="checkbox"/> White 2 <input type="checkbox"/> Black or African American 3 <input type="checkbox"/> American Indian or Alaska Native 4 <input type="checkbox"/> Asian 5 <input type="checkbox"/> Native Hawaiian or Other Pacific Islander 6 <input type="checkbox"/> Other – Specify <u> </u> 7 <input type="checkbox"/> Not stated
8. Age – Complete only if date of birth not given	Units <input type="text"/> <input type="text"/> <input type="text"/> { 1 <input type="checkbox"/> Years 2 <input type="checkbox"/> Months 3 <input type="checkbox"/> Days	
9. Sex – Mark (X) one 1 <input type="checkbox"/> Male 2 <input type="checkbox"/> Female 3 <input type="checkbox"/> Not stated		
10. Ethnicity – Mark (X) one 1 <input type="checkbox"/> Hispanic or Latino 2 <input type="checkbox"/> Not Hispanic or Latino 3 <input type="checkbox"/> Not stated	12. Marital status – Mark (X) one 1 <input type="checkbox"/> Married 3 <input type="checkbox"/> Widowed 5 <input type="checkbox"/> Separated 2 <input type="checkbox"/> Single 4 <input type="checkbox"/> Divorced 6 <input type="checkbox"/> Not stated	

C. ADMINISTRATIVE INFORMATION

13. Type of Admission – Mark (X) one 1 <input type="checkbox"/> Emergency 3 <input type="checkbox"/> Elective 5 <input type="checkbox"/> Items not available/ 2 <input type="checkbox"/> Urgent 4 <input type="checkbox"/> Newborn unknown	16. Expected source(s) of payment	Principal Mark one only	Other additional sources Mark all that apply
14. Source of Admission – Mark (X) one 1 <input type="checkbox"/> Physician referral 7 <input type="checkbox"/> Emergency room 2 <input type="checkbox"/> Clinical referral 8 <input type="checkbox"/> Court/Law enforcement 3 <input type="checkbox"/> HMO referral 9 <input type="checkbox"/> Other – Specify <u> </u> 4 <input type="checkbox"/> Transfer from a hospital 5 <input type="checkbox"/> Transfer from SNF 6 <input type="checkbox"/> Transfer from other health facility 10 <input type="checkbox"/> Item not available			
15. Status/Disposition of patient – Mark (X) appropriate box(es)			
Status	Disposition		
1 <input type="checkbox"/> Alive	a. <input type="checkbox"/> Routine discharge/discharged home b. <input type="checkbox"/> Left against medical advice c. <input type="checkbox"/> Discharged, transferred to another short-term hospital d. <input type="checkbox"/> Discharged, transferred to long-term care institution e. <input type="checkbox"/> Other disposition/not stated		
2 <input type="checkbox"/> Died			
3 <input type="checkbox"/> Status not stated			
	1. Worker's compensation 2. Medicare 3. Medicaid 4. Other government payments 5. Blue Cross/Blue Shield 6. HMO/PPO 7. Other private or commercial insurance 8. Self pay 9. No charge 10. Other – Specify <u> </u> _____ _____ <input type="checkbox"/> No source of payment indicated	<input type="checkbox"/>	<input type="checkbox"/>

(Over)

Figure I. Medical abstract form for the National Hospital Discharge Survey, 2004

Table II. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics, by selected characteristics: 2004

Characteristic	Discharges or first-listed diagnosis		Days of care		All-listed diagnosis		All-listed procedures	
	a	b	a	b	a	b	a	b
Total	0.00193	348.537	0.00298	1,102.846	0.00244	368.046	0.00327	385.864
Sex								
Male	0.00211	355.129	0.00346	1,420.176	0.00294	303.325	0.00375	352.677
Female	0.00202	305.682	0.00297	979.291	0.00403	379.773	0.00663	319.316
Age								
Under 15 years	0.02165	252.708	0.02950	652.859	0.02385	269.440	0.03286	211.058
15–44 years	0.00238	261.974	0.00417	886.487	0.00306	287.451	0.00305	299.209
45–64 years	0.00180	334.793	0.00330	1,198.332	0.00254	329.647	0.00353	322.197
65 years and over	0.00278	342.258	0.00342	1,679.155	0.00366	361.861	0.00322	276.671
Region								
Northeast	0.00900	217.924	0.01201	655.356	0.01786	268.205	0.01325	136.233
Midwest	0.01269	244.190	0.01558	730.556	0.01453	248.977	0.01850	183.671
South	0.00258	319.022	0.00377	1,229.085	0.00341	398.944	0.00382	344.749
West	0.00682	244.576	0.01016	1,061.770	0.00879	369.545	0.00759	300.855
Principal expected source of payment								
Medicare	0.00247	347.898	0.00381	1,468.303	0.00431	382.152	0.00358	251.689
Medicaid	0.00685	275.268	0.00944	1,011.549	0.00516	341.450	0.00660	312.522
Worker's compensation and other government payment	0.00672	327.736	0.01296	1,331.432	0.00940	322.407	0.00861	324.239
HMO and PPO ¹	0.00511	281.902	0.00694	881.873	0.00515	253.947	0.00698	231.569
Blue Cross Blue Shield and other private insurance	0.00352	299.179	0.00577	921.851	0.00411	341.241	0.00546	324.355
Self pay	0.00507	238.592	0.01181	445.933	0.00495	287.494	0.00575	298.687
No charge and other	0.02980	216.556	0.04234	452.271	0.02688	225.121	0.03263	204.223

¹Health maintenance organization and preferred provider organization.

NOTE: The relative standard error (RSE) can be determined from the equation RSE(x)=SQRT(a+b/X).

subclass of the denominator, may be calculated directly using the formula:

$$RSE(p) = 100 \sqrt{b(1-p)/(pX)}$$

where 100p is the percent of interest, X is the denominator of the percent, and b is the parameter b in the formula for approximating the RSE(X). The values for b are given in Table II. When multiplied by 100, RSE(p) is expressed as a percent of p. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (23,24).

The approximate RSE of a population-based rate is equivalent to the approximate RSE of the numerator (obtained by the method described above using figures from Table II), if the denominator is an estimate produced by the U.S. Bureau of the Census for the total population or one or more of the age-sex groups of the total population.

The approximate RSE of a ratio (X/Y) in which the numerator (X) and the denominator (Y) are both estimated from the same survey, but the numerator

is not a subclass of the denominator, is calculated using the formula:

$$RSE(X/Y) = \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

When multiplied by 100, the RSE(X/Y) is expressed as a percent of the ratio X/Y. This approximation is valid if the RSE of the denominator is less than 5 percent or the RSEs of the numerator and denominator are both less than 10 percent (23,24).

Definitions of Selected Terms

Admission source—The source of admission variable was added to the NHDS beginning in 2001. It was modeled after the variable in the National Uniform Bill (UB-92), the standard billing form used by Medicare for reimbursement of hospital inpatient care. The categories for admission source shown in this report are:

Physician referral—The patient was admitted to the hospital upon the recommendation of his/her personal physician, the hospital's clinic

physician, or a health maintenance organization physician.

Emergency department—The patient was admitted to the hospital upon the recommendation of the hospital's emergency department physician. Emergency department and emergency room are used interchangeably in this report.

Transfer from a hospital—The patient was admitted to the hospital as a transfer from an acute care hospital where he/she was an inpatient.

Transfer from other health facility—The patient was admitted to the hospital as a transfer from a skilled nursing facility or other health facility where he/she was an inpatient.

Other sources—The patient was admitted to the hospital upon the direction of a court of law, on the request of a law enforcement agency representative, or for other reasons not categorized above.

Item not available—Information about the source of admission to the hospital was not available for the patient.

Admission type—Like admission source, the admission type variable was added to the NHDS in 2001 and is modeled after the variable in the UB-92. Four admission types are shown in this report:

Emergency—The patient required immediate medical intervention as a result of severe life threatening or potentially disabling conditions. Patients in this group are generally admitted through the emergency room.

Urgent—The patient required immediate care and treatment of a physical or mental disorder. Patients in this group are usually admitted to the first available and suitable accommodation.

Elective—The patient's condition permitted adequate time to schedule the availability of suitable accommodation.

Item not available—Information about the admission type for the patient was not available.

Age—Patient's age at the birthday prior to admission to the hospital.

Average length of stay—The mean length of stay for discharges. It is calculated by dividing the number of days of care by the number of discharges.

Barell injury diagnosis matrix—A two-dimensional array that includes all the *International Classification of Diseases, 9th revision, Clinical Modification (ICD-9-CM)* codes describing injury (18). The code numbers from the matrix are shown in [Table III](#). This matrix was developed to provide a standard format for describing the site and type of injuries for use in epidemiological and clinical analyses of injury data.

Bed size of hospital—The number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for patients, not including bassinets for newborn infants. In this report, the classification of hospitals by

bed size is based on the number of beds listed in the Verispan files (15).

Days of care—The total number of days a patient spent in the hospital. A stay of less than 1 day (patient admitted as an inpatient and discharged on the same day) is counted as one day. For patients admitted and discharged on different days, the number of days of care is computed by counting all the days from (and including) the day of admission to (but not including) the day of discharge.

Death rate—Calculated by dividing the number of deaths for a category by the total number of discharges in that category and multiplying by 100. Discharges whose status is unknown (1.1 percent) were assumed to be discharged alive.

Diagnosis—A disease or injury (or other factor that influences a patient's health status or causes contact with health services) that is listed on the medical record of the patient at discharge. The diagnosis established after study to be chiefly responsible for the admission of the patient to the hospital is the principal diagnosis. The *first-listed diagnosis* is the one specified as the principal diagnosis on the face sheet or discharge summary of the medical record, or if the principal diagnosis is not specified, the first one listed on the face sheet or discharge summary of the medical record. It is usually the main cause of the hospitalization. The number of first-listed diagnoses is the same as the number of discharges. *All-listed* diagnoses include first-listed diagnoses and secondary diagnoses and are included for users who want to know the total number of times conditions have been reported for hospitalized patients. A minimum of one and a maximum of seven diagnoses are coded per discharge in the NHDS. The HIV data ([Table 23](#)) are for discharges with an HIV diagnosis listed in any position among the coded diagnoses. The leading diagnostic categories shown in [Figure 3](#) were identified by ranking the ICD-9-CM Chapters 1-17 and childbirth (code V27) in order of frequency. Discharge rates do not differ significantly for all of the rankings, but the discharge rate was significantly

higher for the top-ranked diagnostic category than for any other category for both males and females, and the rate for second-ranked category was significantly higher than the rate for any of the remaining categories for females.

Discharge—A completed inpatient hospitalization. A hospitalization may be completed by death or by releasing the patient to the customary place of residence, a nursing home, another hospital, or other locations.

Disposition—The placement or status of the patient when discharged from the hospital. Six disposition categories are shown in this report:

Routine or discharged home—Patients who were discharged to return to their homes, including those referred to home care services, discharged to a relative or foster home, or transferred to a corrections institute.

Transferred to other short-term hospital—Patients discharged and transferred to another short-term hospital for additional care.

Transferred to long-term care institution—Patients discharged and transferred or returned to a nursing home, skilled nursing facility (SNF), an extended care facility, an intermediate care facility (ICF), or a custodial care facility.

Other discharged alive—Includes patients discharged alive whose disposition is unknown, patients who left the hospital against medical advice, patients transferred to another facility that could not be classified as short term or long term, and patients transferred to a state facility, psychiatric facility, terminal care/hospice facility, rehabilitation facility, or specialty hospital.

Died—Patients who died after being admitted to the hospital as inpatients.

Status not stated—No information was available on whether the patient was discharged alive or died in the hospital.

Hospital—Hospitals with an average length of stay for all inpatients

Table III. Code numbers for Barell injury diagnosis matrix categories in Table 24

[Code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification*]

Type of injury	All body sites	Traumatic brain injury	Other head, face, and neck	Spinal cord	Vertebral column	Torso	Upper extremities	Lower extremities	Other and unspecified
All types of injury . . .	800–908, 909.0–909.2, 909.4, 909.9, 910–994, 995.50–995.59, 995.80–995.85	800–801, 803, 804, 850–854, 950.1–950.3, 995.55	802, 807.5–807.6, 830, 848.0–848.2, 870–874, 900, 910, 918, 920–921, 925, 940–941, 947.0, 950.0, 950.9, 951, 953.0, 954.0, 957.0, 959.0	806, 952	805, 839.0–839.5, 847.0–847.4	807.0–807.4, 808–809, 839.6–839.7, 846, 847.9, 848.3–848.5, 860–868, 875–878, 879.0–879.7, 901, 902.0–902.5, 902.81–902.82, 911, 922, 926, 942, 947.3–947.4, 953.1–953.3, 953.5, 954.1–954.9, 959.1	810–818, 831–834, 840–842, 880–887, 903, 912–915, 923, 927, 943–944, 953.4, 955, 959.2–959.5	820–827, 835–838, 843–845, 890–897, 904.0–904.8, 916, 917, 924.0–924.5, 928, 945, 959.6–959.7	819, 828–829, 839.8–839.9, 848.8–848.9, 869, 879.8–879.9, 902.87, 902.89, 902.9, 904.9, 919, 924.8–924.9, 929, 946, 947.1–947.2, 947.8–947.9, 948, 949, 953.8–953.9, 956, 957.1–957.9, 959.8–959.9
Fracture	800–829	800–801, 803, 804	802, 807.5–807.6	806	805	807.0–807.4, 808, 809	810–818	820–827	819, 828–829
Dislocation	830–839	None	830	None	839.0–839.5	839.6–839.7	831–834	835–838	839.8–839.9
Sprains and strains . .	840–848	None	848.0–848.2	None	847.0–847.4	846, 847.9, 848.3–848.5	840–842	843–845	848.8–848.9
Internal	850–854, 860–869, 952, 995.55	850–854, 995.55	None	952	None	860–868	None	None	869
Open wound	870–884, 890–894	None	870–874	None	None	875–878, 879.0–879.7	880–884	890–894	879.8–879.9
Amputation	885–887, 895–897	None	None	None	None	None	885–887	895–897	None
Blood vessels	900–904	None	900	None	None	901, 902.0–902.5, 902.81–902.82	903	904.0–904.8	902.87, 902.89, 902.9, 904.9
Contusions and superficial injuries . .	910–924	None	910, 918, 920–921	None	None	911, 922	912–915, 923	916–917, 924.0–924.5	919, 924.8–924.9
Crush	925–929	None	925	None	None	926	927	928	929
Burns	940–949	None	940–941, 947.0	None	None	942, 947.3–947.4	943–944	945	946, 947.1–947.2, 947.8–947.9, 948–949
Nerves and spinal cord	950–951, 953–957	950.1–950.3	950.0, 950.9, 951, 953.0, 954.0, 957.0	None	None	953.1–953.3, 953.5, 954.1–954.9	953.4, 955	None	953.8–953.9, 956, 957.1–957.9
Unspecified	959	None	959.0	None	None	959.1	959.2–959.5	959.6–959.7	959.8–959.9
System-wide	905–908, 909.0–909.2, 909.4, 909.9, 930–939, 958, 960–994, 995.50–995.54, 995.59, 995.80–995.85								

SOURCE: <http://www.cdc.gov/nchs/about/otheract/ice/barellmatrix.htm>.

of less than 30 days (short stay) or those whose specialty is general (medical or surgical) or children’s general. Federal hospitals, hospital units of institutions, and hospitals with fewer than six beds staffed for patient use are excluded.

Newborn infant—Inpatients admitted to a hospital by birth. Sick newborn infants are those that had at least one of the ICD–9–CM (1) diagnostic codes 001–999.

Ownership of hospital—The type of ownership is determined by the organization that controls and operates the hospital. Hospitals are grouped as follows:

Proprietary—Hospitals operated by individuals, partnerships, or corporations for profit.

Government—Hospitals operated by State or local governments.

Nonprofit—Hospitals operated by a nonprofit organization or church.

Principal expected source of payment—The primary expected method of payment for the hospitalization as reported on the patient’s medical record. Types of payment are:

Medicare—The health insurance program for the aged and disabled administered by the Centers for Medicare and Medicaid Services (formerly the Health Care Financing Administration). A person 65 years

of age and over is included in this category only if Medicare is reported as the principal expected source of payment.

Medicaid—A jointly funded federal-state health insurance program providing medical care to those unable to afford it. If Medicaid and HMO or PPO are reported, the discharge is included in the category listed first.

Worker’s compensation—A state or municipal disability insurance or industrial accident insurance.

Other government—This includes payments under the Title V Program, the state-funded Maternal and Child Health Program that provides coverage for children up to 21 years of age with specific chronic medical conditions (e.g., Crippled Children’s Programs), CHAMPUS (Civilian Health and Medical Program of the Uniformed Services) or TRICARE (the Defense Department’s health care program for service families), casualty insurance paid by the state, vocational rehabilitation, federal or state research grant (medical research), or legal hold (prisoner in medical detention).

HMO or PPO—Any health maintenance organization (HMO) or preferred provider organization

(PPO) sponsored by consumers, insurers, organizations, physicians, or hospitals. Medicare and Medicaid discharges are included in this category when HMO or PPO is reported as their principal expected source of payment.

Blue Cross Blue Shield and other private—A private insurance plan not specified as an HMO or PPO. This includes Blue Cross Blue Shield plans, medical coverage provided by life insurance companies, private casualty insurance companies, health insurance companies, and independent plans such as employer- and union-sponsored plans and self-funded plans (partial or total).

Self-pay—The majority of the costs for the hospitalization were expected to be paid by the patient, spouse, family, or next of kin.

No charge—Patients admitted with the understanding that the medical services would be free. This includes hospital-sponsored special research or teaching patients.

Other—Includes any nonprofit source of payment, such as church welfare, United Way (United Appeal), or Shriner’s Hospitals for Children.

Table IV. Code numbers for procedures considered nonsurgical in the National Hospital Discharge Survey, by ICD–9–CM category, 2004

[Procedure categories and code numbers are based on the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD–9–CM)*]

Procedure category and ICD–9–CM code	Nonsurgical codes
Operations on the nervous system 01–05	01.18–01.19,03.31,03.39,04.19,05.19
Operations on the endocrine system 06–07	06.19,07.19
Operations on the eye 08–16	08.19,09.19,09.41–09.44,09.49,10.29,11.29,12.29,14.19,15.09,16.21,16.29
Operations on the ear 18–20	18.01,18.11,18.19,20.31,20.39
Operations on the nose, mouth, and pharynx 21–29	21.00–21.02,21.21,21.29,22.19,24.19,25.09,26.19,27.29,28.19,29.11,29.19
Operations on the respiratory system 30–34	31.41–31.42,31.48–31.49,33.21–33.23,33.29,34.21–34.22,34.28–34.29
Operations on the cardiovascular system 35–39	37.26–37.28,37.29,38.29,39.95
Operations on the hemic and lymphatic system 40–41	40.19,41.38–41.39
Operations on the digestive system 42–54	42.22–42.23,42.29,44.11–44.13,44.19,44.32,45.11–45.13,45.19,45.21–45.24,45.28–45.29,48.21–48.23,48.29,49.21,49.29,50.19,51.10–51.11,51.19,52.19,54.21,54.29
Operations on the urinary system 55–59	55.21–55.22,55.29,56.31,56.35,56.39,57.31–57.32,57.39,57.94–57.95,58.21–58.22,58.29,59.29
Operations on the male genital organs 60–64	60.18–60.19,61.19,62.19,63.09,64.19,64.94
Operations on the female genital organs 65–71	65.19,66.19,67.19,68.11,68.19,69.92,70.21–70.22,70.29,71.19
Obstetrical procedures 72–75	73.4–73.59,73.91–73.92,75.31–75.32,75.34–75.35,75.38,75.94
Operations on the musculoskeletal system 76–84	76.19,78.80–78.89,80.20–80.29,81.98,83.29,84.41–84.43,84.45–84.47
Operations on the integumentary system 85–86	85.19,86.19,86.92
Miscellaneous diagnostic and therapeutic procedures and new technologies 87–99,00	87.01–99.99,00.01–00.15

Procedure—Surgical procedures (such as appendectomy), diagnostic procedures (such as spinal tap), and therapeutic treatments (such as injection or infusion of cancer chemotherapeutic substance) reported on a patient's medical record. Data are for *all-listed* procedures, which include all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded per discharge in the NHDS. The procedures classified as nonsurgical in this report are listed in [Table IV](#) by ICD-9-CM code numbers (1). All other procedures were considered surgical. This classification was begun in 1979 for the presentation of NHDS data, and still remains an important distinction. However, the development of minimally invasive and noninvasive procedures has resulted in less difference between surgical and nonsurgical procedures with regard to operative or anesthetic risk and requirements for highly trained personnel or special equipment.

Region—Hospitals are classified by location in one of the four geographic regions of the United States that correspond to those used by the U.S. Census Bureau.

<i>Region</i>	<i>States included</i>
Northeast	Maine, Vermont, New Hampshire, Massachusetts, Connecticut, Rhode Island, New York, New Jersey, and Pennsylvania;
Midwest	Ohio, Illinois, Indiana, Michigan, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Kansas, and Nebraska;
South	Delaware, Maryland, District of Columbia, West Virginia, Virginia, Kentucky, Tennessee, North Carolina, South Carolina, Georgia, Florida, Alabama, Mississippi, Louisiana, Oklahoma, Arkansas, and Texas;

West Washington, Oregon, California, Nevada, New Mexico, Arizona, Idaho, Utah, Colorado, Montana, Wyoming, Alaska, and Hawaii.

Changes in ICD-9-CM Codes

Although the ICD-9-CM has been used for coding NHDS data since 1979, the classification system undergoes annual updating. Assignment of new diagnostic and procedure codes, fourth- and fifth-digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Centers for Medicare and Medicaid Services (formerly Health Care Financing Administration). Addenda to the ICD-9-CM have become effective October 1 of each year since 1986, except in 1999. No addendum was released in 1999 because of concerns about instituting coding changes just before the millennium crossover.

The data in this report were coded using the addenda for October 1, 1986-2003. When hospitals reported data using code changes introduced in the October 1, 2004 addendum, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

The new ICD-9-CM codes from the 1986-2003 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned before development of the new code are shown in [Table V](#) (diagnoses codes) and [Table VI](#) (procedure codes). Specific titles and more detailed information about the coding system can be found in appropriate volumes of the ICD-9-CM (1).

Table V. Changes in ICD-9-CM diagnosis codes, 1986-2003

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
005.81	1995	005.8	204.21	1991	V10.61
005.89	1995	005.8	204.80	1991	204.8
007.4	1997	007.8	204.81	1991	V10.61
007.5	2000	007.8	204.90	1991	204.9
008.00-008.09	1992	008.0	204.91	1991	V10.61
008.43-008.47	1992	008.49	205.00	1991	205.0
008.61-008.69	1992	008.6	205.01	1991	V10.62
031.2	1997	031.8	205.10	1991	205.1
038.10	1997	038.1	205.11	1991	V10.62
038.11	1997	038.1	205.20	1991	205.2
038.19	1997	038.1	205.21	1991	V10.62
040.82	2002	040.89	205.30	1991	205.3
041.00-041.03	1992	041.0	205.31	1991	V10.62
041.04	1992	041.0	205.80	1991	205.8
041.04 (code title restated)	1997	041.04	205.81	1991	V10.62
041.05	1992	041.0	205.90	1991	205.9
041.09	1992	041.0	205.91	1991	V10.62
041.10-041.19	1992	041.1	206.00	1991	206.0
041.81-041.85	1992	041.8	206.01	1991	V10.63
041.86	1995	041.84	206.10	1991	206.1
041.89	1992	041.8	206.11	1991	V10.63
042	1994	042.0-042.2, 042.9, 043.0-043.3, 043.9, 044.0, 044.9 (Codes deleted)	206.20	1991	206.2
042.0-042.9	1986	279.19	206.21	1991	V10.63
043.0-043.9	1986	279.19	206.80	1991	206.8
044.0-044.9	1986	279.19	206.81	1991	V10.63
066.4	2002	066.3	206.90	1991	206.9
070.20-070.21	1991	070.2	206.91	1991	V10.63
070.22	1994	070.20	207.00	1991	207.0
070.23	1994	070.21	207.01	1991	V10.69
070.30-070.31	1991	070.3	207.10	1991	207.1
070.32	1994	070.30	207.11	1991	V10.69
070.33	1994	070.31	207.20	1991	207.2
070.41-070.43	1991	070.4	207.21	1991	V10.69
070.44	1994	070.41	207.80	1991	207.8
070.49	1991	070.4	207.81	1991	V10.69
070.51-070.53	1991	070.5	208.00	1991	208.0
070.54	1994	070.51	208.01	1991	V10.60
070.59	1991	070.5	208.10	1991	208.1
077.98-077.99	1993	077.9	208.11	1991	V10.60
078.10-078.11,078.19	1993	078.1	208.20	1991	208.2
078.88	1993	078.89	208.21	1991	V10.60
079.4	1993	079.8	208.80	1991	208.8
079.50-079.53,079.59	1993	079.8	208.81	1991	V10.60
079.6	1996	079.89	208.90	1991	208.9
079.81	1995	079.89	208.91	1991	V10.60
079.82	2003	079.89	237.70-237.72	1990	237.7
079.88-079.89	1993	079.8	250.02	1993	250.90
079.98-079.99	1993	079.9	250.03	1993	250.91
082.40-082.41, 082.49	2000	082.8	250.12	1993	250.10
088.81,088.89	1989	088.8	250.13	1993	250.11
088.82	1993	088.89	250.22	1993	250.20
099.40-099.49	1992	099.4	250.23	1993	250.21
099.50-099.59	1992	078.89	250.32	1993	250.30
112.84-112.85	1992	112.89	250.33	1993	250.31
114.4-114.5	1993	114.3	250.42	1993	250.40
176.0-176.9	1991	173.0-173.9	250.43	1993	250.41
203.00	1991	203.0	250.52	1993	250.50
203.01	1991	V10.79	250.53	1993	250.51
203.10	1991	203.1	250.62	1993	250.60
203.11	1991	V10.79	250.63	1993	250.61
203.80	1991	203.8	250.72	1993	250.70
203.81	1991	V10.79	250.73	1993	250.71
204.00	1991	204.0	250.82	1993	250.80
204.01	1991	V10.61	250.83	1993	250.81
204.10	1991	204.1	250.92	1993	250.90
204.11	1991	V10.61	250.93	1993	250.91
204.20	1991	204.2	255.10-255.14	2003	255.1
			256.31-256.39	2001	256.3
			275.40-275.42	1997	275.4
			275.49	1997	275.4

Table V. Changes in ICD-9-CM diagnosis codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
277.02-277.03,277.09	2002	277.00	410.30-410.32	1989	410.3
277.7	2001	277.8	410.40-410.42	1989	410.4
277.81-277.84,277.89	2003	277.8	410.50-410.52	1989	410.5
278.00-278.01	1995	278.0	410.60-410.62	1989	410.6
282.41-282.42,282.49	2003	282.4	410.70-410.72	1989	410.7
282.64	2003	282.63	410.80-410.82	1989	410.8
282.68	2003	282.69	410.90-410.92	1989	410.9
283.10-283.11,283.19	1993	283.1	411.81	1989	410.9
285.21-285.22,285.29	2000	285.8	411.89	1989	411.8
289.52	2003	289.59	414.00-414.01	1994	414.0
289.81-289.82,289.89	2003	289.8	414.02-414.03	1994	996.03
291.81	1996	291.8	414.04-414.05	1996	414.00
291.89	1996	291.8	414.06	2002	414.00
293.84	1996	293.89	414.07	2003	414.06
294.10-294.11	2000	294.1	414.12	2002	414.11
300.82	1996	300.81	415.11	1995	997.3 & 415.1
305.1	1994	305.10, 305.11, 305.12, 305.13 (Codes deleted)	415.19	1995	415.1
312.81-312.82,312.89	1994	312.8	428.20-428.23	2002	428.0
315.32	1996	315.39	428.30-428.33	2002	428.0
320.81-320.89	1992	320.8	428.40-428.43	2002	428.0
331.11	2003	331.1	429.71	1989	410.0-410.9
331.19	2003	331.1	429.79	1989	410.0-410.9
331.82	2003	331.89	433.00-433.01	1993	433.0
333.92-333.93	1994	333.99	433.10-433.11	1993	433.1
337.20-337.22,337.29	1993	337.9	433.20-433.21	1993	433.2
337.3	1998	337.9	433.30-433.31	1993	433.3
342.00-342.02	1994	342.0	433.80-433.81	1993	433.8
342.10-342.12	1994	342.1	433.90-433.91	1993	433.9
342.80-342.82	1994	342.9	434.00-434.01	1993	434.0
342.90-342.92	1994	342.9	434.10-434.11	1993	434.1
344.00-344.04,344.09	1994	344.0	434.90-434.91	1993	434.9
344.30-344.32	1994	344.3	435.3	1995	435.0 & 435.1
344.40-344.42	1994	344.4	437.7	1992	780.9
344.81,344.89	1993	344.8	438.0	1997	294.9 & 438
345.00-345.01	1989	345.0	438.10	1997	784.5 & 438
345.10-345.11	1989	345.1	438.11	1997	784.3 & 438
345.40-345.41	1989	345.4	438.12	1997	784.4 & 438
345.50-345.51	1989	345.5	438.19	1997	784.5 & 438
345.60-345.61	1989	345.6	438.20	1997	342.90 & 438
345.70-345.71	1989	345.7	438.21	1997	342.91 & 438
345.80-345.81	1989	345.8	438.22	1997	342.92 & 438
345.90-345.91	1989	345.9	438.30	1997	344.40 & 438
346.00-346.01	1992	346.0	438.31	1997	344.41 & 438
346.10-346.11	1992	346.1	438.32	1997	344.42 & 438
346.20-346.21	1992	346.2	438.40	1997	344.30 & 438
346.80-346.81	1992	346.8	438.41	1997	344.31 & 438
346.90-346.91	1992	346.9	438.42	1997	344.32 & 438
348.30-348.31,348.39	2003	348.3	438.50-438.52	1997	344.89 & 438
355.71	1993	354.4	438.53	1998	438.50
355.79	1993	355.7	438.6-438.7	2002	438.89
357.81-357.82,357.89	2002	357.8	438.81	1997	784.69 & 438
358.00-358.01	2003	358.0	438.82	1997	787.2 & 438
359.81,359.89	2002	359.8	438.83-438.85	2002	438.89
365.83	2002	365.89	438.89	1997	438
371.82	1992	371.89	438.9	1997	438
372.81,372.89	2000	372.8	440.20-440.22	1992	440.2
374.87	1990	374.89	440.23	1993	440.20 & 707.1 or 707.8 or 707.9
403.00-403.01	1989	403.0	440.24	1993	440.20 & 785.4
403.10-403.11	1989	403.1	440.29	1993	440.2
403.90-403.91	1989	403.9	440.30-440.32	1994	996.1
404.00-404.03	1989	404.0	441.00-441.03	1994	441.0
404.10-404.13	1989	404.1	441.6	1993	441.1 & 441.3
404.90-404.93	1989	404.9	441.7	1993	441.2 & 441.4
410.00-410.02	1989	410.0	443.21	2002	442.81
410.10-410.12	1989	410.1	443.22	2002	442.2
410.20-410.22	1989	410.2	443.23	2002	442.1
			443.24	2002	442.89

Table V. Changes in ICD-9-CM diagnosis codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
443.29	2002	442.89	536.40-536.42,536.49	1998	997.4
445.01-445.02	2002	440.29	537.82	1990	537.89
445.81	2002	440.1	537.83	1991	537.82
445.89	2002	440.8	537.84	2002	531.00
446.20-446.21,446.29	1990	446.2	556.0-556.6	1994	556
451.82-451.84	1993	451.89	556.8-556.9	1994	556
454.8	2002	454.9	558.3	2000	558.9
458.2	1995	997.9 & 458.9	562.02	1991	562.00
458.21,458.29	2003	458.2	562.03	1991	562.01
458.8	1997	458.9	562.12	1991	562.10
459.10-459.13,459.19	2002	459.1	562.13	1991	562.11
459.30-459.33,459.39	2002	459.89	564.00-564.09	2001	564.0
464.00-464.01	2001	464.0	564.81	1998	564.8
464.50-464.51	2001	464.0	564.89	1998	564.8
466.11	1996	466.1	569.60-569.61	1995	569.6
466.19	1996	466.1	569.62	1998	569.69
474.0 (Code title restated)	1997	474.0	569.69	1995	569.6
474.00-474.02	1997	474.0	569.84	1990	557.1
477.1	2000	477.8	569.85	1991	569.84
480.3	2003	480.8	569.86	2002	569.82
482.30-482.39	1992	482.3	574.60	1996	574.00 & 574.30
482.40	1998	482.4	574.61	1996	574.01 & 574.31
482.41	1998	482.4	574.70	1996	574.10 & 574.40
482.49	1998	482.4	574.71	1996	574.11 & 574.41
482.81-482.83,482.89	1992	482.8	574.80	1996	574.00 & 574.10 574.30 & 574.40
482.84	1997	482.83	574.81	1996	574.01 & 574.11 574.31 & 574.41
483.0	1992	483	574.90	1996	574.20 & 574.50
483.1	1996	078.88 & 484.8	574.91	1996	574.21 & 574.51
483.8	1992	483	575.10-575.11	1996	575.1
491.20-491.21	1991	491.2	575.12	1996	575.0 & 575.1
493.20	1989	493.90	593.70-593.73	1994	593.7
493.21	1989	493.91	596.51-596.53	1992	596.5
493.02	2000	493.00	596.54	1992	344.61
493.12	2000	493.10	596.55-596.59	1992	596.5
493.22	2000	493.20	599.81-599.89	1992	599.8
493.81	2003	519.1	600.0-600.3, 600.9	2000	600
493.82	2003	493.90-493.91	600.00-600.01	2003	600.0
493.92	2000	493.90	600.10-600.11	2003	600.1
494.0-494.1	2000	494	600.20-600.21	2003	600.2
512.1	1994	997.3	600.90-600.91	2003	600.9
517.3	2003	282.62	602.3	2001	602.8
518.6	1997	518.89	607.85	2003	607.89
518.81	1987	799.1	608.82	2001	608.83
518.82-518.89	1987	518.8	608.87	2001	608.89
518.83	1998	518.81	633.00-633.01	2002	633.0
518.84	1998	518.81	633.10-633.11	2002	633.1
519.00-519.02, 519.09	1998	519.0	633.20-633.21	2002	633.2
521.00-521.09	2001	521.0	633.80-633.81	2002	633.8
524.00-524.09	1992	524.0	633.90-633.91	2002	633.9
524.10-524.19	1992	524.1	645.0	1991	645
524.60-524.69	1991	524.6	645.10-645.11, 645.13	2000	645.00-645.01, 645.03
524.70-524.79	1992	524.8	645.20-645.21, 645.23	2000	645.00-645.01, 645.03
525.10-525.19	2001	525.1	651.30-651.31,651.33	1989	651.00-651.01, 651.03
530.10-530.11, 530.19	1993	530.1	651.40-651.41,651.43	1989	651.10-651.11, 651.13
530.12	2001	530.10	651.50-651.51,651.53	1989	651.20-651.21, 651.23
530.20-530.21	2003	530.2	651.60-651.61,651.63	1989	651.80-651.81, 651.83
530.81	1993	530.1	654.20-654.21,654.23	1990	654.2,654.9
530.82-530.84	1993	530.8	654.90-654.94	1990	654.2,654.9
530.85	2003	530.2	655.70 & 655.71	1997	655.8
530.89	1993	530.8	655.73	1997	655.8
535.00-535.01	1991	535.0	657.0	1991	657
535.10-535.11	1991	535.1			
535.20-535.21	1991	535.2			
535.30-535.31	1991	535.3			
535.40-535.41	1991	535.4			
535.50-535.51	1991	535.5			
535.60-535.61	1991	535.6			
536.3	1994	536.8			

Table V. Changes in ICD-9-CM diagnosis codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
659.60,659.61,659.63	1992	659.80-659.81, 659.83	771.81-771.83,771.89	2002	771.8
659.70,659.71,659.73	1998	656.30, 656.31, 656.33	772.10-772.14	2001	772.1
665.10,665.11	1992	665.10,665.11,665.12, 665.14 ¹	779.7	2001	772.1
670.0	1991	670	779.81-779.82,779.89	2002	779.8
672.0	1991	672	779.83	2003	779.89
674.50-674.54	2003	674.80,674.82,674.84	780.01-780.02	1992	780.0
677	1994	None	780.03	1993	780.01
686.00 & 686.01	1997	686.0	780.09	1992	780.0
686.09	1997	686.0	780.31	1997	780.3
690.10	1995	690	780.39	1997	780.3
690.11	1995	691.8 & 704.8	780.57	1992	780.51,780.53
690.12	1995	691.8	780.71	1998	780.7
690.18	1995	690	780.79	1998	780.7
690.8	1995	690	780.91-780.92	2002	780.9
692.72-692.74	1992	692.79	780.93	2003	780.99
692.75	2000	692.79	780.94	2003	780.99
692.76-692.77	2001	692.71	780.99	2002	780.9
692.82-692.83	1992	692.89	781.8	1994	781.9
702.0-702.8	1991	702	781.91-781.92	2000	781.9
702.11,702.19	1994	702.1	781.93	2002	723.5
704.02	1993	704.09	781.94	2003	781.99
707.10-707.15,707.19	2000	707.1	781.99	2000	781.9
709.00-709.01,709.09	1994	709.0	783.21	2000	783.2
710.5	1992	288.3,729.1	783.22	2000	783.4
718.70-718.79	2001	718.80-718.89	783.40-783.43	2000	783.4
719.7	2003	719.70,719.75-719.79 (Codes deleted)	783.7	2000	783.4
727.83	2000	727.89	785.52	2003	785.59
728.86	1995	729.4	786.03	1998	786.09
728.87	2003	728.9	786.04	1998	786.09
728.88	2003	728.89	786.05	1998	786.09
733.10-733.16,733.19	1993	733.1	786.06	1998	786.09
733.93	2001	733.16	786.07	1998	786.09
733.94	2001	733.19	787.01-787.03	1994	787.0
733.95	2001	733.19	787.91	1995	558.9
738.10-738.19	1992	738.1	787.99	1995	787.9
747.60-747.64,747.69	1993	747.6	788.20-788.21, 788.29	1993	788.2
747.82	1993	747.89	788.30-788.39	1992	788.3
747.83	2002	747.89,747.9	788.41-788.43	1993	788.4
752.51-752.52	1996	752.5	788.61-788.62	1993	788.6
752.61-752.63	1996	752.6	788.63	2003	788.69
752.64-752.65	1996	752.8	788.69	1993	788.6
752.69	1996	752.8	789.00-789.07, 789.09	1994	789.0
752.81,752.89	2003	752.8	789.30-789.37, 789.39	1994	789.3
753.10-753.17,753.19	1990	753.1	789.40-789.47, 789.49	1994	789.4
753.20-753.23	1996	753.2	789.60-789.67, 789.69	1994	789.6
753.29	1996	753.2	790.01, 790.09	2000	790.0
756.70-756.71	1997	756.7	790.21-790.22,790.29	2003	790.2
756.79	1997	756.7	790.91	1993	790.9
758.81	1996	758.8	790.92	1993	286.9
758.89	1996	758.9	790.93, 790.99	1993	790.9
759.81-759.82	1989	759.8	790.94	1997	790.99
759.83	1994	759.89	792.5	2000	792.9
759.89	1989	759.8	793.80-793.81, 793.89	2001	793.8
760.75	1991	760.79	795.00-795.02,795.09	2002	795.0
760.76	1994	760.79	795.31-795.39	2002	795.3
763.81-763.83,763.89	1998	763.8	795.71	1994	795.8 (Code deleted)
764.00-764.09	1988	764.0	795.79	1994	795.7
764.10-764.19	1988	764.1	795.8	1986	795.7
764.20-764.29	1988	764.2	796.5	1997	796.9
764.90-764.99	1988	764.9	799.81	2003	799.8
765.00-765.09	1988	765.0	799.89	2003	799.8
765.10-765.19	1988	765.1	813.45	2002	813.42
765.20-765.24	2002	765.00-765.09	823.40-823.42	2002	823.80-823.82
765.25-765.29	2002	765.10-765.19	840.7	2001	840.8
766.21-766.22	2003	766.2	850.11-850.12	2003	850.1
767.11,767.19	2003	767.1	864.05	1992	864.09
770.81-770.84,770.89	2002	770.8	864.15	1992	864.19

See footnote at end of table.

Table V. Changes ICD-9-CM in diagnosis codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
909.5	1994	909.9	V10.48	1998	V10.49
922.31-922.33	1996	922.3	V10.53	2001	V10.59
925.1-925.2	1993	925	V12.00-V12.03,V12.09	1994	V12.0
959.0 (Code title restated)	1997	959.0	V12.40-V12.41	1997	V12.4
959.01	1997	854.00	V12.49	1997	V12.4
959.09	1997	959.0	V12.50-V12.52	1995	V12.5
959.11-959.14,959.19	2003	959.1	V12.59	1995	V12.5
965.61	1998	965.6	V12.70-V12.72,V12.79	1994	V12.7
965.69	1998	965.6	V13.00-V13.01,V13.09	1994	V13.0
989.81-989.84	1995	989.8	V13.21,V13.29	2002	V13.2
989.89	1995	989.8	V13.61	1998	V13.6
995.50-995.55	1996	995.5	V13.69	1998	V13.6
995.59	1996	995.5	V15.01-V15.09	2000	V15.0
995.60-995.69	1993	995.0	V15.41-V15.42	1996	V15.4
995.7	2000	None	V15.49	1996	V15.4
995.80	1996	995.81	V15.82	1994	305.13 (Codes deleted)
995.81 (Code title restated)	1996	995.81	V15.84-V15.86	1995	V15.89
995.82-995.85	1996	995.81	V15.87	2003	V15.89
995.86	1998	995.89	V16.40-V16.43	1997	V16.4
995.90	2002	038.0-038.9	V16.49	1997	V16.4
995.91	202	038.0-038.9	V16.51	1998	V16.5
995.92	2002	038.0-038.9	V16.59	1998	V16.5
995.93	2002	038.0-038.9	V18.61	1998	V18.6
995.94	2002	038.0-038.9	V18.69	1998	V18.6
996.04	1994	996.09	V21.30-V21.35	2000	None
996.51-996.59	1987	996.5	V23.41,V23.49	2002	V23.4
996.55	1998	996.52	V23.7	1989	V23.8
996.56	1998	996.59	V23.81	1998	V23.8
996.57	2003	996.59	V23.82	1998	V23.8
996.60-996.69	1989	996.6	V23.83	1998	V23.8
996.68	1998	996.69	V23.84	1998	V23.8
996.70-996.79	1989	996.7	V23.89	1998	V23.8
996.80-996.84	1987	996.8	V25.03	2003	V25.01
996.85	1990	996.8	V25.43	1992	V25.49
996.86	1987	996.8	V25.5	1992	V25.8
996.87	2000	996.89	V26.21-V26.22,V26.29	2000	V26.2
996.89	1987	996.8	V26.51	1998	None
997.00-997.01	1995	997.0	V26.52	1998	None
997.02	1995	997.9 & 430-434, 436	V28.6	1997	V28.8
997.09	1995	997.0	V29.0-V29.1,V29.8	1992	V71.8
997.71	2001	997.4	V29.2	1994	V29.8
997.72	2001	997.5	V29.3	1998	V29.8
997.79	2001	997.2	V29.8	1992	V71.8
997.91	1995	997.9	V29.9	1992	V71.9
997.99	1995	997.9	V30.00-V30.01	1989	V30.0
998.11-998.12	1996	998.1	V31.00-V31.01	1989	V31.0
998.13	1996	998.89	V32.00-V32.01	1989	V32.0
998.31-998.32	2002	998.3	V33.00-V33.01	1989	V33.0
998.51	1996	998.5	V34.00-V34.01	1989	V34.0
998.59	1996	998.5	V35.00-V35.01	1989	V35.0
998.81-998.82,998.89	1994	998.8	V36.00-V36.01	1989	V36.0
998.83	1996	998.89	V37.00-V37.01	1989	V37.0
V01.81,V01.89	2002	V01.8	V39.00-V39.01	1989	V39.0
V01.82	2003	V01.89	V42.81-V42.83	1997	V42.8
V02.51	1998	V02.5	V42.84	2000	V42.89
V02.52	1998	V02.5	V42.89	1997	V42.8
V02.59	1998	V02.5	V43.21-V43.22	2003	V43.2
V02.60-V02.62	1997	V02.6	V43.60-V43.66,V43.69	1994	V43.6
V02.69	1997	V02.6	V43.81-V43.82	1995	V43.8
V03.81-V03.82,V03.89	1994	V03.8	V43.83	1998	V43.89
V04.81-V04.82,V04.89	2003	V04.8	V43.89	1995	V43.8
V05.3-V05.4	1993	V05.8	V44.50	1998	V44.5
V06.5-V06.6	1994	V06.8	V44.51	1998	V44.5
V07.31,V07.39	1994	V07.3	V44.52	1998	V44.5
V07.4	1992	V07.8	V44.59	1998	V44.5
V08	1994	044.9,795.8 (Codes deleted)	V45.00	1994	V45.89
V09.0-V09.91	1993	None	V45.01	1994	V45.0
			V45.02,V45.09	1994	V45.89

Table V. Changes in ICD-9-CM diagnosis codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
V45.51	1994	V45.5	V66.7	1996	None
V45.52,V45.59	1994	V45.89	V67.00-V67.01,V67.09	2000	V67.0
V45.61	1997	V45.6	V69.0-V69.3	1994	None
V45.69	1997	V45.6	V69.8-V69.9	1994	None
V45.71	1997	611.8	V71.81	2000	V71.8
V45.72	1997	569.89	V71.82-V71.83	2002	V71.89
V45.73	1997	593.89	V71.89	2000	V71.8
V45.74	2000	593.89,596.8	V72.81-V72.85	1993	V72.8
V45.75	2000	V45.89	V73.88-V73.89	1993	V73.8
V45.76	2000	518.89	V73.98-V73.99	1993	V73.9
V45.77	2000	602.8,607.89, 608.89,620.8, 621.8,622.8	V76.10-V76.12,V76.19	1997	V76.1
V45.78	2000	360.89	V76.44	1998	V76.49
V45.79	2000	255.8,289.59, 388.8,569.49, 577.8,V45.89	V76.45	1998	V76.49
V45.82	1994	V45.89	V76.46-V76.47	2000	V76.49
V45.83	1995	V45.89	V76.50-V76.52	2000	V76.49
V45.84	2001	None	V76.81, V76.89	2000	V76.8
V45.85	2003	V45.89	V77.91, V77.99	2000	V77.9
V46.2	2002	V46.8	V82.81, V82.89	2000	V82.8
V49.60-V49.67	1994	V49.5	V83.01-V83.02	2001	None
V49.70-V49.77	1994	V49.5	V83.81	2002	None
V49.81	2000	None	V83.89	2002	V19.8
V49.82	2001	None	E854.8	1995	E858.8
V49.89	2000	V49.8	E869.4	1994	E869.8
V50.41-V50.42,V50.49	1994	V50.8	E880.1	1995	E884.9
V53.01-V53.02	1997	V53.0	E884.3-E884.4	1995	E884.2
V53.09	1997	V53.0	E884.5-E884.6	1995	E884.9
V53.31	1994	V53.3	E885.0	2002	E885.9
V53.32,V53.39	1994	V53.9	E885.1-E885.4,E885.9	2000	E885
V53.90-V53.91,V53.99	2003	V53.9	E888.0	2001	E920
V54.01-V54.02,V54.09	2003	V54.0	E888.1	2001	E888
V54.10-V54.17,V54.19	2002	V54.8	E888.8-E888.9	2001	E888
V54.20-V54.27,V54.29	2002	V54.8	E906.5	1995	E906.3
V54.81,V54.89	2002	V54.8	E908.0-E908.4	1995	E908
V56.1	1995	V58.89	E908.8-E908.9	1995	E908
V56.1 (Code title restated)	1998	V56.1	E909.0-E909.4	1995	E909
V56.2	1998	V56.1	E909.8-E909.9	1995	E909
V56.31-V56.32	2000	V56.8	E917.3	2001	E917.9
V57.21-V57.22	1994	V57.2	E917.4	2001	E917.9
V58.41	1994	V58.4	E917.5	2001	E886.0
V58.42	2002	V58.49	E917.6	2001	E917.1
V58.43	2002	V58.49	E917.7	2001	E888
V58.49	1994	V58.4	E917.8	2001	E888
V58.61	1995	V67.51	E920.5	1995	E920.4
V58.62	1998	V58.69	E922.4	1997	E917.9
V58.63-V58.65	2003	V58.69	E922.5	2002	E922.8
V58.69	1995	V67.51	E924.2	1995	E924.0
V58.71-V58.78	2002	V58.49	E928.3	2000	E928.8
V58.81,V58.89	1994	V58.8	E928.4	2003	E928.8
V58.82	1995	V58.89	E928.5	2003	E928.8
V58.83	2000	V58.89	E955.6	1997	E955.9
V59.01-V59.02	1995	V59.0	E955.7	2002	E955.4
V59.09	1995	V59.0	E967.2	1996	E967.0
V59.6	1995	V59.8	E967.3	1996	None
V61.10-V61.12	1996	V61.1	E967.4-E967.8	1996	E967.1
V61.22	1996	V61.21	E968.5	1995	E968.8
V62.83	1996	V65.49	E968.6	1997	E968.8
V64.4	1997	None	E968.7	2000	E968.8
V64.41-V64.43	2003	V64.4	E985.6	1997	E985.4
V65.11,V65.19	2003	V65.1	E985.7	2002	E985.4
V65.40-V65.45	1994	V65.4	E979.0-E979.9	2002	E960.0- E966,E968.0- E968.9
V65.46	2003	V65.49	E999.0	2002	E999
V65.49	1994	V65.4	E999.1	2002	E969

¹The title for the subcategory 665.1 has been changed, making the fifth-digit subclassifications 665.12 and 665.14 invalid.

Table VI. Changes ICD-9-CM in procedure codes, 1986-2003

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
00.01	2002	99.99	37.70 (Leads only)	1987	37.70 (Leads/Device)
00.02	2002	99.99	37.71-37.72 (Leads only)	1987	37.74 (Leads/Device)
00.03	2002	99.99	37.73 (Leads only)	1987	37.73 (Leads/Device)
00.09	2002	99.99	37.74 (Leads only)	1987	37.76 (Leads/Device)
00.10	2002	99.25	37.75 (Leads only)	1987	37.89 (Leads/Device)
00.11	2002	99.19	37.76 (Leads only)	1987	37.81 (Leads/Device)
00.12	2002	93.98	37.77 (Leads only)	1987	37.83-37.84 (Leads/Device)
00.13	2002	99.99	37.78	1987	37.71-37.72
00.14	2002	99.21	37.79	1987	86.09
00.15	2003	99.28	37.80-37.87	1992	89.49 (Code deleted, this procedure is included in the code for pacemaker insertion/replacement)
00.50	2002	37.70-37.74,37.76,37.80-37.87	37.80 (Device only)	1987	37.73-37.77 (Leads/Device)
00.51	2002	37.94	37.81 (Device only)	1987	37.73-37.77 (Leads/Device)
00.52	2002	None	37.82 (Device only)	1987	37.73-37.77 (Leads/Device)
00.53	2002	37.80-37.87	37.83 (Device only)	1987	37.73-37.77 (Leads/Device)
00.54	2002	37.96	37.85-37.87	1987	37.85
00.55	2002	39.90	37.89	1987	37.86+37.89
02.96	1992	89.19	37.94-37.98	1986	37.99
03.90	1987	03.99 (Insertion of catheter)	38.22	1986	38.29
05.25	1995	39.7	38.44 (Abdominal aorta only)	1986	38.44 (entire aorta)
11.75	1989	11.79	38.45 (Thoracic aorta added)	1986	38.44-38.45
11.76	1989	11.62	38.95	1989	38.93
20.96-20.98	1986	20.95	39.28	1991	39.29
22.12	1988	22.11	39.50	1995	39.59
26.12	1988	26.11	39.65	1988	39.61
29.31	1991	83.02	39.66	1990	39.65
29.32	1991	29.3	39.71	2000	39.52
29.33	1991	29.3	39.72	2002	39.79
29.39	1991	29.3	39.79	2000	39.52
31.45	1988	31.43-31.44	39.90	1996	39.50
31.95	1989	31.75	41.00-41.03	1988	41.0
32.01	1989	32.0	41.04	1994	99.79
32.09	1989	32.0	41.05	1997	None
32.22	1995	32.29,32.9	41.06	1997	None
32.28	1989	32.29	41.07	2000	41.04
33.27	1987	33.22+33.27	41.08	2000	41.05
33.28	1987	33.27	41.09	2000	41.01
33.29	1987	33.28-33.29	42.25	1988	42.24
33.50	1995	33.5	42.33	1989	42.32,42.39
33.51	1995	33.5	42.33	1990	42.91
33.52	1995	33.5	43.11	1989	43.1
33.6	1990	33.5+37.5	43.19	1989	43.1,43.2
34.05	1994	34.99	43.41	1989	43.41,43.49
35.84	1988	35.82	44.21	1986	44.2
35.96	1986	35.03	44.22	1986	44.99
36.00-36.03	1986	36.0	44.29	1986	44.2
36.04	1986	39.97	44.32	2001	44.39
36.05	1986	36.01 ¹ ,36.02	44.43	1989	43.49,45.32
36.05	1987	36.01	44.44	1989	38.86
36.06	1995	36.01,36.02,36.03,36.05	44.49	1989	43.0
36.07	2002	36.06	44.93-44.94	1986	44.99
36.09	1986	36.0	45.16	1988	45.14 (45.15 before 1987)
36.09	1991	36.00 (Code deleted)	45.30	1989	45.31,45.32
36.17	1996	36.19	45.42	1988	45.41
36.31	1998	36.3	45.43	1989	45.49
36.32	1998	36.3	45.75 (Hartmann resection added)	1988	48.66 (Code deleted)
36.39	1998	36.3	45.95	1987	45.93
37.26-37.27	1988	37.29	46.13	1992	46.12 (Code deleted)
37.28	2001	88.72	46.32	1989	46.39
37.34	1988	37.33	46.85	1989	46.99
37.35	1997	37.33	46.97	2000	46.99
37.51	2003	37.5			
37.52	2003	37.62			
37.53	2003	37.63			
37.54	2003	37.63			
37.65	1995	37.62			
37.66	1995	37.62			
37.67	1998	37.4			

See footnotes at end of table.

Table VI. Changes in ICD-9-CM procedure codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
47.01	1996	47.0	65.54	1996	65.52
47.09	1996	47.0	65.63	1996	65.61
47.11	1996	47.1	65.64	1996	65.62
47.19	1996	47.1	65.74	1996	65.71
48.36	1995	45.42	65.75	1996	65.72
49.31	1989	49.3	65.76	1996	65.73
49.39	1989	49.3	65.81	1996	65.8
49.75	2002	49.79	65.89	1996	65.8
49.76	2002	49.79	66.01	1992	66.0
51.10	1989	51.97	66.02	1992	66.73
51.11	1989	51.11,51.97	67.51	2001	67.5
51.14	1989	51.12	67.59	2001	67.5
51.15	1989	51.97	68.15	1987	68.14
51.21	1996	51.22,51.23	68.16	1987	68.13
51.22	1991	51.21 (Code deleted), 51.22	68.23	1996	68.29
51.23	1991	51.22	68.31	2003	68.3
51.24	1996	51.22,51.23	68.39	2003	68.3
51.64	1989	51.69	68.51	1996	68.5
51.84-51.88	1989	51.97	68.59	1996	68.5
51.97	1986	52.91,51.99, or 51.82	68.9	1992	68.4
51.98	1986	51.99	74.3	1992	69.11 (Code deleted)
52.13	1989	51.97,52.91	75.37	1998	99.29
52.14	1989	52.11	75.38	2001	75.34
52.21	1989	52.2	77.56	1989	77.89,78.49,81.18
52.22	1989	52.2	77.57	1989	77.89,80.48,81.18,83.85
52.84	1996	99.29	77.58	1989	77.59,81.18
52.85	1996	99.29	78.10	1991	78.40
52.86	1996	99.29	78.11	1991	78.41
52.93	1989	52.93+52.91	78.12	1991	78.42
52.94	1989	52.09	78.13	1991	78.43
52.97	1989	52.91	78.14	1991	78.44
52.98	1989	52.91	78.15	1991	78.45
52.99	1989	52.93,52.94,52.99	78.16	1991	78.46
54.24	1987	54.23	78.17	1991	78.47
54.25	1993	54.98	78.18	1991	78.48
54.51	1996	54.5	78.19	1991	78.49
54.59	1996	54.5	78.20	1991	78.10,78.20,78.30
55.03-55.04	1986	55.02	78.21	1991	78.11,78.31
56.33-56.34	1987	56.33	78.22	1991	78.12,78.22,78.32
56.35	1987	45.12	78.23	1991	78.13,78.23,78.33
57.17-57.18	1989	57.21	78.24	1991	78.14,78.34
57.22	1989	57.22,57.82	78.25	1991	78.15,78.25,78.35
58.31	1990	58.3	78.27	1991	78.17,78.27,78.37
58.39	1990	58.3	78.28	1991	78.18,78.38
58.93	1986	57.99	78.29	1991	78.11,78.16,78.19,78.29, 78.39
59.03	1996	59.02	78.39	1991	78.31
59.12	1996	59.11	78.90 ²	1987	78.40
59.72	1995	59.79	78.91 ²	1987	78.41
59.96	1986	59.95	78.92 ²	1987	78.42
60.21	1995	60.2	78.93 ²	1987	78.43
60.29	1995	60.2	78.94 ²	1987	78.44
60.95	1991	60.99	78.95 ²	1987	78.45
60.96	2000	60.29	78.96 ²	1987	78.46
60.97	2000	60.29	78.97 ²	1987	78.47
64.97	1986	64.95	78.98 ²	1987	78.48
65.01	1996	65.0	78.99 ²	1987	78.49
65.09	1996	65.0	80.50-80.59	1986	80.5
65.13	1996	65.12	81.03	1989	81.02
65.14	1996	65.19	81.04-81.05	1989	81.03,81.04,81.05
65.23	1996	65.21	81.06-81.07	1989	81.06,81.07
65.24	1996	65.22	81.08	1989	81.06,81.07,81.08
65.25	1996	65.29	81.09	1989	81.08
65.31	1996	65.3	81.30	2001	81.09
65.39	1996	65.3	81.31	2001	81.09
65.41	1996	65.4	81.32	2001	81.09
65.49	1996	65.4	81.33	2001	81.09
65.53	1996	65.51	81.34	2001	81.09

See footnotes at end of table.

Table VI. Changes in ICD-9-CM procedure codes, 1986-2003—Con.

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
81.35	2001	81.09	88.95	1986	89.29
81.36	2001	81.09	88.96	2002	88.91-88.97
81.37	2001	81.09	88.97	1989	88.99
81.38	2001	81.09	88.98	1989	88.90
81.39	2001	81.09	88.99	1986	89.39
81.40	1989	81.69	89.10	1989	89.15
81.51	1989	81.51,81.59	89.17-89.18	1988	89.15
81.52	1989	81.61,81.62,81.63,81.64	89.19	1989	89.15
81.53	1989	81.51,81.59,81.61,81.62, 81.63,81.64	89.50	1991	89.54
81.54-81.55	1989	81.41	89.60	2002	89.65
81.56	1989	81.48	92.3	1995	01.59,04.07,07.63,07.68
81.57	1989	81.31,81.39	92.30	1998	92.3
81.59	1989	81.39	92.31	1998	92.3
81.61	2002	81.00-81.08,81.30-81.39	92.32	1998	92.3
81.62	2003	None	92.33	1998	92.3
81.63	2003	None	92.39	1998	92.3
81.64	2003	None	93.90	1988	93.92
81.72	1989	81.79	94.61-94.69	1989	94.25
81.73-81.74	1989	81.86	96.29	1998	96.39
81.75	1989	81.87	96.6	1986	96.35
81.79	1989	81.79,81.87	96.70	1991	93.92 (Code deleted)
81.80	1989	81.81	96.71	1991	93.92 (Code deleted)
81.97	1992	81.59	96.72	1991	93.92 (Code deleted)
84.51	2002	None	97.05	1989	51.97
84.52	2002	None	97.44	2001	37.64
85.95	1987	85.99	98.51-98.52	1989	59.96 (Code deleted)
85.96	1987	85.99	98.59	1989	59.96 (Code deleted)
86.06	1987	86.09	99.00	1995	99.02
86.07	1990	86.09	99.10	1998	99.29
86.27	1986	86.22-86.23	99.15	1986	99.29
86.28	1988	86.22	99.20	1998	99.29
86.67	1998	86.65	99.28	1994	99.25
86.93	1987	86.89	99.71-99.79 ³	1988	99.07
88.90	1986	88.39	99.75	2000	99.29
88.91	1986	89.15	99.76	2002	99.79
88.92	1986	89.39	99.77	2002	None
88.93	1986	89.15	99.85	1987	93.35
88.94	1986	89.39	99.86	1987	93.39
			99.88	1988	99.83

¹Before October 1986 contents of current code 36.05 would have been assigned to 36.0.²Codes 78.90-78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90-78.99 were reassigned to codes 78.40-78.49.³Codes 99.71-99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71-99.79 were reclassified as "Therapeutic apheresis."

Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *National Vital Statistics Report* (NVSR). These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the NVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Information Dissemination Staff
National Center for Health Statistics
Centers for Disease Control and Prevention
3311 Toledo Road, Room 5412
Hyattsville, MD 20782
1-866-441-NCHS (6247)
E-mail: nchsquery@cdc.gov
Internet: www.cdc.gov/nchs

**U.S. DEPARTMENT OF
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention
National Center for Health Statistics
3311 Toledo Road
Hyattsville, MD 20782

MEDIA MAIL
POSTAGE & FEES PAID
CDC/NCHS
PERMIT NO. G-284

06-0136 (10/06)
CS105422
T26540
DHHS Publication No. (PHS) 2006-1733, Series 13, No. 162