

NATIONAL HOSPITAL AMBULATORY MEDICAL CARE SURVEY

2009 CERVICAL CANCER SCREENING SUPPLEMENT

NOTICE – Public reporting burden of this collection of information is estimated to average 15 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. An agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a currently valid OMB control number. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to CDC/ATSDR Information Collection Review Office; 1600 Clifton Road, MS D-74, Atlanta, GA 30333, ATTN: PRA (0920-0234).

Assurance of Confidentiality – All information which would permit identification of an individual, a practice, or an establishment will be held confidential, will be used for statistical purposes only by NCHS staff, contractors, and agents only when required and with necessary controls, and will not be disclosed or released to other persons without the consent of the individual or establishment in accordance with section 308(d) of the Public Health Service Act (42 USC 242m) and the Confidential Information Protection and Statistical Efficiency Act (PL-107-347).

BACKGROUND INFORMATION

A. Hospital number	B. Clinic type – <i>Mark (X) only ONE.</i> 1 <input type="checkbox"/> OB/GYN 2 <input type="checkbox"/> GM	C. AU number
D. Census contact name	E. Census contact telephone	Area code Number

INTRODUCTION

The Centers for Disease Control and Prevention is conducting a special survey on cervical cancer screening performed in hospital outpatient clinics. Please answer the following questions. We appreciate your time on this important public health concern.

	<i>Mark (X) one interval for routine screening.</i>				
	Annually	Every 2 years	Every 3 years	More than 3 years	No routine interval recommended
1. Does this clinic use any of the following methods to screen for cervical cancer? <i>Mark (X) all that apply.</i>					
a. Conventional Pap test (<i>Definition – Smear spread on glass slide and fixed</i>) 1 <input type="checkbox"/> Yes – How often does this clinic routinely screen women using this method? → 2 <input type="checkbox"/> No 3 <input type="checkbox"/> Unknown } <i>Continue with item 1b</i>	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
b. Liquid-based cytology (<i>Definition – Specimen suspended in liquid solution</i>) 1 <input type="checkbox"/> Yes – How often does this clinic routinely screen women using this method? → 2 <input type="checkbox"/> No 3 <input type="checkbox"/> Unknown } <i>Continue with item 1c</i>	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
c. Other – <i>Specify</i> ↴ <div style="border: 1px solid black; height: 20px; width: 100%; margin: 5px 0;"></div> 1 <input type="checkbox"/> Yes – How often does this clinic routinely screen women using this method? → 2 <input type="checkbox"/> No 3 <input type="checkbox"/> Unknown	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

2. Does this clinic perform colposcopy?

1 Yes
2 No
3 Unknown

9. As it relates to the HPV vaccine, how often does your clinic – <i>Mark (X) only ONE for each row.</i>	Rarely or never	Sometimes	Usually	Always or almost always	Unknown/Not applicable/ Do not ask
a. Use the number of sexual partners to determine who should get the HPV vaccine?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
b. Perform a Pap test to determine who should get the HPV vaccine?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
c. Recommend the HPV vaccine to females with a history of an abnormal Pap test result (ASC-US or higher)?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>
d. Recommend the HPV vaccine to females with a positive HPV test?	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>

10. Will this clinic's cervical cancer screening and management procedures change for females who have been fully vaccinated with the HPV vaccine?

1 Yes
2 No – *SKIP to item 14*

11. How will this clinic determine when to start routine cervical cancer screening for fully HPV vaccinated females?

Mark (X) all that apply.

1 By age

 1 At same age as non-HPV vaccinated females –
 Specify age → _____

 2 At a later age –
 Specify age → _____

2 By onset of sexual activity –
 How many year(s) since onset of sexual activity? → _____

3 Will not be screening fully HPV vaccinated females

4 Unknown

CLOSING STATEMENT

Thank you for completing this special survey. We appreciate your time and cooperation.

3a. Does this clinic ever order or collect the Human Papillomavirus (HPV) DNA test?

- 1 Yes – Go to item 3b
 - 2 No – SKIP to item 3c
 - 3 Not aware of HPV DNA test
 - 4 Unknown
- } SKIP to item 9 on page 4

b. Which of the following HPV DNA tests are ordered or collected in this clinic? Mark (X) all that apply.

- 1 High risk (HR) HPV DNA test
 - 2 Low risk (LR) HPV DNA test
 - 3 Not aware there was a high risk or low risk HPV DNA test
 - 4 Type-specific HPV DNA test
 - 5 Unknown
- } SKIP to item 4a

c. Why is the HPV DNA test not ordered or collected in this clinic? – Mark (X) all that apply.

- 1 This clinic does not see the types of patients for whom the HPV DNA test is indicated.
- 2 This clinic uses other tests, procedures, or examination methods to manage patients for whom the HPV DNA test is indicated.
- 3 The patients in this clinic have timely access to colposcopy.
- 4 Assessing patients' HPV infection status is not a priority at this clinic.
- 5 The labs affiliated with this clinic do not offer the HPV DNA test.
- 6 The health plans or health systems affiliated with this clinic do not recommend the HPV DNA test.
- 7 The HPV DNA test is not a reimbursed or covered service for most patients in this clinic.
- 8 Discussing cervical cancer screening in the context of an STD is avoided in this clinic.
- 9 Notifying or counseling patients about positive HPV DNA test results would take too much time.
- 10 Notifying or counseling patients about positive HPV DNA test results might make clinicians in this clinic feel uncomfortable.
- 11 Notifying or counseling patients about positive HPV DNA test results might make patients in this clinic feel uncomfortable, angry, or upset.

SKIP to item 7 on page 3.

4a. If a patient's Pap test result is borderline or abnormal, does this clinic routinely order an HPV DNA test to be performed on that sample (commonly called reflex HPV DNA testing)? (An HPV DNA test may be run on the same liquid-based medium as the Pap test or an HPV DNA test specimen may be collected at the same time as the conventional Pap test.)

- 1 Yes – Go to item 4b
 - 2 No
 - 3 Unknown
- } SKIP to item 5a

b. For which borderline or abnormal Pap test result would this clinic order or collect a reflex HPV DNA test? Mark (X) all that apply.

- 1 ASC-US (atypical squamous cells of undetermined significance)
- 2 ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion)
- 3 LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1)
- 4 HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ)
- 5 AGC (atypical glandular cells)

c. For which patients does this clinic usually order reflex HPV DNA testing? – Mark (X) all that apply.

- 1 Women under 21 years old
- 2 Women 21 years old to 29 years old
- 3 Women 30 years old and over
- 4 Other – Specify

5a. Does this clinic routinely recall patients to come back for a second sample collection for an HPV DNA test if their Pap test is abnormal or borderline (recall testing)?

- 1 Yes – Go to item 5b
 - 2 No
 - 3 Unknown
- } SKIP to item 6a on page 3

b. For which abnormal or borderline Pap test result would this clinic recall a patient for an HPV DNA test? Mark (X) all that apply.

- 1 ASC-US (atypical squamous cells of undetermined significance)
- 2 ASC-H (atypical squamous cells of undetermined significance – cannot exclude high-grade intraepithelial lesion)
- 3 LSIL (low-grade squamous intraepithelial lesion, encompassing mild dysplasia/CIN1)
- 4 HSIL (high-grade squamous intraepithelial lesion, moderate dysplasia/CIN2, severe dysplasia/CIN3, and carcinoma in situ)
- 5 AGC (atypical glandular cells)

6a. Does this clinic routinely order or collect an HPV DNA test at the same time as the Pap test as part of routine cervical cancer screening (commonly called adjunct HPV testing or cotesting)?

- 1 Yes – Go to item 6b
 2 No
 3 Unknown } SKIP to item 7

b. For which patients does this clinic routinely order or collect an HPV DNA test along with the Pap test (commonly called adjunct HPV testing or cotesting)? Mark (X) all that apply.

- 1 Women under 21 years old
 2 Women 21 years old to 29 years old
 3 Women 30 years old and over
 4 Women who request the test for cervical cancer screening
 5 Women who request the test to check their HPV infection status
 6 Other – Specify

7. Given the following screening histories, when would this clinic recommend that a woman between 30 and 60 years of age return for her next Pap test?

Prior Pap test results in past 5 years (excluding current normal results)	Current HPV DNA test results	Current Pap test result	For each of the following scenarios, mark (X) only ONE for each row.							Have no experience with this type of patient or test
			No follow-up needed	Less than 6 months	6 months to less than 1 year	1 year	2 years	3 years or more		
(a) Two consecutive normal Pap tests	Has not had test	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(b) Two consecutive normal Pap tests	Negative	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(c) Two consecutive normal Pap tests	Positive	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(d) Has not had a Pap test	Negative	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(e) Has not had a Pap test	Positive	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(f) Abnormal Pap test	Negative	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	
(g) Abnormal Pap test	Positive	Normal	1 <input type="checkbox"/>	2 <input type="checkbox"/>	3 <input type="checkbox"/>	4 <input type="checkbox"/>	5 <input type="checkbox"/>	6 <input type="checkbox"/>	7 <input type="checkbox"/>	

QUESTIONS 8-14 ASK ABOUT THE HPV VACCINE

8. How often does this clinic use an HPV test to determine who should get the HPV vaccine? Mark (X) only one.

- 1 Rarely or never
 2 Sometimes
 3 Usually
 4 Always or almost always
 5 Do not recommend the HPV vaccine –SKIP to item 10.