

ICD 10 Diagnosis Codes for Bronchiolitis Obliterans Syndrome (BOS)

Dr. Howard J. Huang, MD

Lung transplantation section chief, Houston Methodist J.C. Walter Jr. Transplant Center

Dr. Ajay Sheshadri, MD, MSCI


Assistant Professor, Department of Pulmonary Medicine, Division of Internal Medicine,
The University of Texas MD Anderson Cancer Center

September, 2021




Bronchiolitis Obliterans Syndrome



- Bronchiolitis obliterans syndrome (BOS) is a clinical syndrome characterized by airflow limitation not reversible with inhaled bronchodilators which may be associated with progressive dyspnea.
- It was first clearly described in early 1980s in the context of lung transplant as a rare fibrotic disorder involving terminal and respiratory bronchioles.^{1,2}
- The histologic hallmark of BOS is obliterative bronchiolitis (OB), which consists of a fibrotic luminal obliteration of the respiratory and terminal bronchioles.
- BOS is classified as a rare disease.³



Normal bronchiole



Bronchiole at late-stage BO




Bronchioles with BO, a patchy infestation

Reference(s):

[1]. Seminars in Respiratory and Critical Care Medicine. 33 (5): 509–32; [2]. Pathology of Solid Organ Transplantation. doi.org/10.1007/978-3-540-79343-4_7; [3]. Orphanet ORPHA: 1303

Pathology of Bronchiolitis Obliterans

Compared to normal lung bronchiole, the airway lumen is narrowed in BOS



Lung biopsy specimens of a normal bronchiole (left) and a bronchiole from a patient with obliterative bronchiolitis (right)

Thickened bronchiolar wall by inflammatory fibrosis located between epithelium and smooth muscle

Cross section of a normal bronchiole (left) and a bronchiole affected by obliterative bronchiolitis (right)

BOS as a manifestation of CLAD after lung transplant

- BOS is associated with injury of small airways due to infections, systemic and autoimmune diseases, and certain inhaled agents, but appears most frequently after:
 - Lung transplantation as a manifestation of chronic lung allograft dysfunction (CLAD), or
 - Allogenic hematopoietic stem cell transplantation (alloHSCT) as the pulmonary manifestation of chronic graft-versus-host-disease (cGVHD).
- BOS is the most common manifestation (or phenotype) of CLAD, accounting for nearly 50% to 70% of the cases. Up to 30% of patients with CLAD develop a restrictive defect called restrictive allograft syndrome (RAS).⁴
- The pulmonary council of the International Society for Heart and Lung Transplantation (ISHLT) classifies CLAD into four clinical sub-types—BOS, RAS, mixed and unknown.⁴

Classification of CLAD after lung transplant


| | |
|---|---|
| Bronchiolitis Obliterans Syndrome (BOS) | Most common manifestation (or phenotype) of CLAD, accounting for nearly 50% to 70% of the cases |
| Restrictive Allograft Syndrome (RAS) | ~30% of patients with CLAD develop a restrictive defect |
| Mixed | BOS with other measures of chronic lung allograft dysfunction |
| Unknown | BOS due to environmental, chemical exposure and other reasons |

Reference(s):

[4]. J Heart Lung Transplant. 2019 May;38(5):493-503


Incidence of BOS after lung transplant

Within first 4.5 years after lung transplant, half of the patients develop BOS



BOS as a manifestation of cGVHD after allogeneic stem cell transplant

- In alloHSCT patients, BOS occurs as a pulmonary manifestation of cGVHD.
- In alloHSCT patients afflicted by cGVHD, BOS is an important contributor to morbidity and mortality. It is associated with poor prognosis with a 5-year survival rate of 60%.⁷
- It is the most common non-infectious pulmonary complication of alloHSCT, typically presenting after the first 100 days following transplantation, with most cases presenting between 12 to 18 months after transplantation.⁸
- The exact incidence and prevalence of BOS after alloHSCT are not known but individual studies have reported that the prevalence of BOS in alloHSCT patients ranges from 3.4% to 10%.⁹



Reference(s):

[7]. Bone Marrow Transplant. 2019 Mar;54(3):383-392; [8]. Bone Marrow Transplant. 2013 Sep;48(9):1224-9 [9]. Blood 2019; 134 (Supplement_1): 5678.

Challenges due to absence of an ICD code specific to BOS

- Currently, there is no specific code in the ICD-10 system for CLAD and its phenotypes (BOS, RAS, mixed and undefined) following lung transplant.
- Similarly, there are no codes available for BOS and other pulmonary manifestations of cGVHD after alloHSCT.
- Because BOS after these procedures requires frequent healthcare encounters, therapeutic interventions and regular monitoring, a coding strategy that helps to accurately identify BOS after lung transplant and alloHSCT could help in
 - accurate measurement of the quality, safety and efficacy of care in these two separate patient populations
 - conducting more effective research, epidemiological studies, and clinical trials
 - effective monitoring of resource utilization in these patients

Support for ICD 10 code application from an international panel of clinical experts

| KOL | Specialty | Affiliation | |
|------------------|----------------------------------|---|-----------|
| Adriana Balduzzi | Allo-HSCT (pediatric hematology) | University of Milan-Bicocca, San Gerardo Hospital | Italy |
| Ajay Sheshadri | Pulmonary medicine | MD Anderson Cancer Center | USA |
| Alan Glanville | Lung transplant | University of New South Wales | Australia |
| Aldo Iacono | Pulmonary medicine, Lung Tx | University of Maryland | USA |
| Anne Bergeron | Pulmonary medicine | Hôpital Saint-Louis, Service de Pneumologie | France |
| Claus Neurohr | Pulmonary medicine, Lung Tx | Department for Pneumology and Respiratory Medicine, Robert Bosch Hospital | Germany |
| Daniel Wolff | Hematology, HSCT | University of Regensburg Faculty of Medicine | Germany |
| Fabio Ciceri | Hematology, HSCT | Bone Marrow Transplantation, IRCCS Ospedale San Raffaele | Italy |

| KOL | Specialty | Affiliation | |
|---------------------|--------------------------------|---|---------|
| Federica Melloni | Pulmonary medicine, Lung Tx | Department of Hematological, Pneumological and Cardiovascular Sciences, University of Pavia | Italy |
| Geert Verleden | Pulmonary medicine, Lung Tx | Dept. of Respiratory disease, University Hospital Gasthuisberg KU Leuven | Belgium |
| Howard Huang | Pulmonary medicine, Lung Tx | Houston Methodist J.C. Walter Jr. Transplant Center | USA |
| Jens Gottlieb | Pulmonary medicine, Lung Tx | Hannover Medical School, Center for Internal Medicine | Germany |
| Kirsten M Williams | Pediatric Hematology/ Oncology | Aflac Cancer and Blood Disorders Center, Children's Healthcare of Atlanta | USA |
| Michael Perch | Pulmonary medicine, Lung Tx | Dept. of Cardiology, section for Lung transplantation, Copenhagen Uni Hospital | Denmark |
| Nikolaus Kneidinger | Pulmonary medicine, Lung Tx | Department of Internal Medicine V, Comprehensive Pneumology Center; University of Munich | Germany |



Thank you!