

Table 22. Number of deaths, death rates, and age-adjusted death rates for drug-induced causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. A listing of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition codes included in this table can be found in the note at the bottom of the table]

Year	All origins ¹			Hispanic			Non-Hispanic ²			Non-Hispanic white ³			Non-Hispanic black ³		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Number															
2007	38,371	23,883	14,488	2,723	2,045	678	35,532	21,771	13,761	30,755	18,760	11,995	4,126	2,662	1,464
2006	38,396	24,507	13,889	2,871	2,135	736	35,379	22,262	13,117	29,970	18,634	11,336	4,722	3,212	1,510
2005	33,541	21,208	12,333	2,596	1,969	627	30,809	19,140	11,669	26,186	16,170	10,016	4,019	2,612	1,407
2004	30,711	19,362	11,349	2,257	1,671	586	28,339	17,605	10,734	24,201	14,952	9,249	3,577	2,309	1,268
2003	28,723	18,426	10,297	2,358	1,800	558	26,199	16,497	9,702	22,245	13,959	8,286	3,466	2,256	1,210
2002	26,040	16,734	9,306	2,137	1,647	490	23,756	14,978	8,778	19,949	12,478	7,471	3,404	2,264	1,140
2001	21,705	14,253	7,452	1,731	1,335	396	19,799	12,778	7,021	16,367	10,465	5,902	3,099	2,113	986
2000	19,720	13,137	6,583	1,700	1,348	352	17,835	11,656	6,179	14,585	9,439	5,146	2,977	2,050	927
1999	19,128	12,885	6,243	1,965	1,605	360	16,966	11,136	5,830	13,644	8,831	4,813	3,030	2,134	896
Rate															
2007	12.7	16.1	9.5	6.0	8.7	3.1	13.9	17.4	10.5	15.3	19.0	11.7	10.8	14.6	7.3
2006	12.8	16.6	9.1	6.5	9.3	3.4	13.9	17.9	10.1	14.9	18.9	11.1	12.5	17.8	7.6
2005	11.3	14.5	8.2	6.1	8.9	3.0	12.1	15.4	9.0	13.1	16.4	9.8	10.8	14.7	7.2
2004	10.5	13.4	7.6	5.5	7.8	2.9	11.2	14.3	8.3	12.1	15.3	9.1	9.7	13.1	6.6
2003	9.9	12.9	7.0	5.9	8.7	2.9	10.4	13.5	7.6	11.2	14.3	8.2	9.5	13.0	6.3
2002	9.0	11.8	6.3	5.5	8.2	2.6	9.5	12.3	6.9	10.0	12.8	7.4	9.4	13.2	6.0
2001	7.6	10.2	5.1	4.7	7.0	2.2	8.0	10.6	5.5	8.3	10.8	5.8	8.7	12.5	5.3
2000	7.0	9.5	4.6	4.8	7.4	2.1	7.2	9.7	4.9	7.4	9.8	5.1	8.5	12.3	5.0
1999	6.9	9.4	4.4	5.8	9.2	2.2	6.9	9.3	4.6	6.9	9.2	4.8	8.7	12.9	4.9
Age-adjusted rate ⁴															
2007	12.6	15.8	9.3	6.5	9.5	3.4	13.6	17.0	10.2	15.1	18.7	11.4	11.4	15.8	7.5
2006	12.7	16.4	9.0	7.2	10.3	3.9	13.6	17.5	9.8	14.7	18.6	10.8	13.2	19.4	7.9
2005	11.3	14.4	8.1	6.8	10.0	3.5	11.9	15.2	8.8	12.8	16.2	9.6	11.4	16.1	7.4
2004	10.4	13.3	7.6	6.2	8.9	3.4	11.1	14.1	8.2	12.0	15.0	8.9	10.4	14.6	6.8
2003	9.9	12.8	7.0	6.7	9.9	3.3	10.3	13.3	7.4	11.0	14.1	8.0	10.1	14.4	6.6
2002	9.0	11.7	6.3	6.2	9.3	3.0	9.4	12.1	6.8	9.9	12.6	7.2	10.1	14.5	6.3
2001	7.6	10.1	5.1	5.3	8.0	2.5	7.9	10.4	5.4	8.1	10.6	5.7	9.3	13.8	5.5
2000	7.0	9.5	4.6	5.4	8.3	2.4	7.1	9.5	4.8	7.2	9.6	4.9	9.1	13.6	5.3
1999	6.8	9.4	4.4	6.4	10.3	2.5	6.8	9.2	4.6	6.8	8.9	4.6	9.4	14.4	5.2

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

²Includes races other than white and black.

³Multiple-race data were reported by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴For method of computation, see "Technical Notes."

NOTE: Causes of death attributable to drug-induced mortality include ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0-F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. In 2006, the list of drug-induced codes was modified to include two new ICD-10 codes, Drug-induced acute pancreatitis (ICD-10 code K85.3) and Drug-induced fever (ICD-10 code R50.2).