

Series 13

No. 130



---

# Vital and Health Statistics

---

From the CENTERS FOR DISEASE CONTROL AND PREVENTION / National Center for Health Statistics

## Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1995

November 1997



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics



**Copyright Information**

All material appearing in this report is in the public domain and may be reproduced or copied without permission; citation as to source, however, is appreciated.

---

**Suggested citation**

Graves EJ, Gillum BS. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1995. National Center for Health Statistics. Vital Health Stat 13(130). 1997.

---

**Library of Congress Catalog Card Number 90-13573**

---

For sale by the U.S. Government Printing Office  
Superintendent of Documents  
Mail Stop: SSOP  
Washington, DC 20402-9328

---

# Vital and Health Statistics

---

## Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1995

Series 13:  
Data From the National Health  
Survey  
No. 130

---

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Centers for Disease Control and Prevention  
National Center for Health Statistics

Hyattsville, Maryland  
November 1997  
DHHS Publication No. (PHS) 98-1791

## **National Center for Health Statistics**

Edward J. Sondik, Ph.D., *Director*

Jack R. Anderson, *Deputy Director*

Jack R. Anderson, *Acting Associate Director for International Statistics*

Lester R. Curtin, Ph.D., *Acting Associate Director for Research and Methodology*

Jacob J. Feldman, Ph.D., *Associate Director for Analysis, Epidemiology, and Health Promotion*

Gail F. Fisher, Ph.D., *Associate Director for Data Standards, Program Development, and Extramural Programs*

Edward L. Hunter, *Associate Director for Planning, Budget, and Legislation*

Jennifer H. Madans, Ph.D., *Acting Associate Director for Vital and Health Statistics Systems*

Stephen E. Nieberding, *Associate Director for Management*

Charles J. Rothwell, *Associate Director for Data Processing and Services*

## **Division of Health Care Statistics**

W. Edward Bacon, Ph.D., *Director*

Thomas McLemore, *Deputy Director*

Robert Pokras, *Chief, Hospital Care Statistics Branch*

---

### *Cooperation of the U.S. Bureau of the Census*

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the U.S. Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

---

# Contents

---

Abstract .....	1
Introduction.....	1
Methods.....	1
Data Source .....	2
Estimation .....	2
Use of Tables .....	2
References.....	3
Appendix I .....	140
Technical Notes on Methods .....	140
Reliability of Estimates .....	140
Population Estimates.....	140
Appendix II.....	141
Conversion Table of Changes in ICD-9-CM Codes .....	141

## List of detailed tables

1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995 .....	4
2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995 .....	32
3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995 .....	60
4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995 .....	119

## Appendix tables

I. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics: United States, 1995.....	140
II. Civilian population by sex, age, and geographic region: United States, 1995.....	141
III. Diagnosis codes.....	141
IV. Procedure codes .....	145

## Abstract

### Objectives

This report presents national estimates of the use of non-Federal short-stay hospitals in the United States during 1995. Estimates of first-listed diagnoses, all-listed diagnoses, days of care for first-listed diagnoses, and all-listed procedures are shown by sex and age of patient and geographic region of hospital.

### Methods

The estimates are based on data collected through the National Hospital Discharge Survey for 1995. The survey has been conducted annually by the National Center for Health Statistics since 1965. In 1995, data were collected for approximately 263,000 discharges from 466 non-Federal short-stay hospitals. Diagnoses and procedures are presented according to their code number in the *International Classification of Diseases, 9th Revision, Clinical Modification (ICD-9-CM)*.

**Keywords:** *inpatients • diagnoses • procedures • ICD-9-CM*

# Detailed Diagnoses and Procedures, National Hospital Discharge Survey, 1995

by Edmund J. Graves and Brenda S. Gillum, Division of Health Care Statistics

## Introduction

This report presents data from the 1995 National Hospital Discharge Survey (NHDS). The survey has been conducted continuously by the National Center for Health Statistics (NCHS) since 1965. National estimates of hospital use derived from the NHDS are published for each calendar year by NCHS. This report is intended as a source for researchers and others who need data for detailed diagnostic and procedure categories. Estimates of specific diagnoses and procedures and days of care for first-listed diagnoses are shown by age and sex of patients and geographic region of the hospitals. Similar reports have been published with data for 1977-79 (1-3) and 1983-94 (4-15). Other data from the NHDS are also published in Series 13 reports of *Vital and Health Statistics*.

The NHDS is the principal source for national data on the characteristics of discharges from non-Federal short-stay hospitals. Data from the NHDS are used to examine important topics in public health (16-20) and for a variety of planning, administrative, and evaluation activities by governmental, scientific, academic, and commercial institutions.

Estimates of the number of procedures shown in this report are for inpatients only. Data on ambulatory surgery are collected through the

National Survey of Ambulatory Surgery (NSAS), another survey conducted by NCHS. The NSAS was implemented in 1994 and covers ambulatory surgery provided in hospitals and freestanding ambulatory surgery centers (21). Data from NSAS are not included in the tables of this report.

## Methods

### Data Source

Through NHDS, data are collected from a sample of inpatient records acquired from a national sample of hospitals. Data from NHDS represent a sample of discharges, not individuals. Persons with multiple discharges during the year may be sampled more than once. Only hospitals with an average length of stay of fewer than 30 days for all patients, general hospitals, or children's general hospitals are included in the survey. Federal, military, and Department of Veterans Affairs hospitals, as well as hospital units of institutions (such as prison hospitals) and hospitals with fewer than six beds staffed for patient use, are excluded. Beginning in 1988, the sampling frame for hospitals has been the *SMG Hospital Market Database* (22-24).

Prior to 1988, NHDS had a two-stage design, but in 1988 the survey

This report was prepared in the Division of Health Care Statistics. David Lei, of the OAO Corporation, produced estimated parameters for relative standard error equations. Rong Cai, OAO Corporation, did the computer programming for the report. This report was edited by Klaudia Cox and typeset by Zung T. N. Le of the Publications Branch, Division of Data Services.

was redesigned (25). Beginning in 1988, all hospitals with 1,000 beds or more or 40,000 discharges or more annually were included in the sample with certainty, while the remaining sample of hospitals was based on a stratified three-stage design. The first stage consisted of the selection of 112 primary sampling units (PSU's) that comprise a probability sample of PSU's used in the 1985–94 National Health Interview Survey (26). The second stage consisted of the selection of noncertainty hospitals from the sample PSU's. At the third stage, a systematic random sample of discharges within hospitals was selected.

For 1995, the sample consisted of 525 hospitals, 17 of which were out of scope (ineligible) because they had gone out of business or otherwise failed to meet the criteria for the NHDS universe. Of the 508 in-scope (eligible) hospitals, 466 (92 percent) responded to the survey. Data were collected for approximately 263,000 patient discharges from the 466 responding hospitals.

Two data collection procedures were used for the survey. One was a manual system in which sample selection and transcription of information from the hospital records to abstract forms were performed by the hospital's staff or by staff of the U.S. Bureau of the Census on behalf of NCHS. Completed forms were sent to NCHS for coding, editing, and postsurvey weighting adjustments.

The other data collection procedure was an automated system in which NCHS purchased tapes containing machine-readable medical record data from commercial organizations, State data systems, hospitals, or hospital associations. Records from these tapes were systematically sampled by NCHS. In 1995, approximately 36 percent of respondent hospitals provided data through the automated system.

The medical abstract form and the automated data tapes contain items that relate to the personal characteristics of the patient. These items include birth date (or age), sex, race, ethnicity, marital status, ZIP Code, and expected sources of payment. Administrative items such as admission and discharge dates, discharge status, and medical

record number are also included. The medical information about patients includes diagnoses, surgical and nonsurgical procedures, and dates of surgery. (The medical record number, date of birth, and ZIP Code are confidential information not available to the public.)

For the manual data collection, an ongoing quality control program is undertaken on the coding and keying of data from abstracts to machine-readable form. Approximately 5 percent of the abstracts are independently recoded by an NHDS coder, with discrepancies resolved by the chief coder. The overall error rate for records manually coded by NCHS for the 1995 data year was 4.3 percent for medical (ICD–9–CM) coding and keying and 1.4 percent for demographic coding and keying.

## Estimation

Because of the nature of sample data, the survey data must be inflated or weighted to produce national estimates. The estimation procedure produces essentially unbiased national estimates and has three basic components: (a) inflation by reciprocals of the probabilities of sample selection, (b) adjustment for nonresponse, and (c) population weighting ratio adjustments. These three components of the final weight are described in more detail in an earlier report (14). Information about the standard errors of statistics for the 1995 NHDS is in [appendix I](#) of this report.

## Use of Tables

The tables in this report present diagnoses and procedures according to the detailed code numbers of the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM)(27), which is updated periodically (see [appendix II](#)). Estimates shown are the number of first-listed diagnoses ([table 1](#)), number of days of care ([table 2](#)), number of all-listed diagnoses ([table 3](#)), and number of all-listed procedures ([table 4](#)) for discharges from short-stay hospitals.

The first-listed diagnosis ([table 1](#)) is the one specified as the principal

diagnosis on the face sheet or discharge summary of the medical record.

However, if the principal diagnosis is not specified, it is the first one listed on the face sheet or discharge summary of the medical record. Data for all-listed diagnoses ([table 3](#)) include the first-listed diagnosis and other diagnoses appearing in the medical record. A minimum of one and a maximum of seven diagnoses are coded.

Estimates of procedures, which consist of surgical and nonsurgical procedures, are also published. Estimates of all-listed procedures ([table 4](#)) contain all occurrences of the procedure coded, regardless of their order in the medical record. Up to four procedures are coded for each discharge.

[Tables 1–3](#) provide information by 3-digit, 4-digit, and 5-digit ICD–9–CM diagnostic codes. [Table 4](#) shows 2-, 3-, and 4-digit procedure codes. Only ICD–9–CM code numbers appear in [tables 1–4](#). Specific titles and more detailed information about the coding scheme can be found in appropriate volumes of the ICD–9–CM (27).

A specific code appears in [tables 1, 3, and 4](#) if the estimate of the total for the code is 5,000 or more. These estimates generally have a relative standard error (a measure of the reliability of the data) no higher than 30 percent and a sample size greater than 30. The codes that appear in [table 2](#) for days of care are identical to those in [table 1](#) for first-listed diagnoses. Only an asterisk (\*) is shown in [tables 1, 3, and 4](#) if the estimate of diagnoses or procedures for a sex or age group or region is less than 5,000. In [table 2](#) only an asterisk is shown for estimates of days of care derived from fewer than 5,000 estimated discharges.

In addition, all estimates of diagnoses and procedures less than 9,000 are preceded by an asterisk to indicate that they are based on a relatively small sample of discharge records. These estimates are generally based on fewer than 60 records. Days-of-care estimates derived from fewer than 9,000 estimated discharges are also preceded by asterisks.

The total for all codes presented at the beginning of each table includes estimates for codes not shown in the

table. Likewise, the total for each diagnostic chapter or procedure category includes estimates for all the codes in the chapter or category, whether or not they are shown individually. Estimates for 5-digit codes may not add to the 4-digit total, 4-digit codes to the 3-digit total, or, for procedures, 3-digit codes to the 2-digit total because the totals include estimates for codes not shown or because the data were rounded to the nearest thousand for presentation.

Although 4- and 5-digit diagnostic codes are presented together under the 3-digit codes of which they are a part, the estimates for 4- and 5-digit codes cannot be added to obtain the estimate for the 3-digit code. For example, in [table 1](#), the estimate for first-listed diagnosis code 198 is 113,000. If the estimates for the 4- and 5-digit codes included in 198 are added, the total would be 122,000. This occurs because the estimate of 18,000 for code 198.89 is part of the estimate of 22,000 for code 198.8 and thus cannot be added to it. Similarly, estimates for 3- and 4-digit codes cannot be added together to obtain the estimate for a 2-digit code.

## References

- Haupt BJ. Detailed diagnoses and surgical procedures for patients discharged from short-stay hospitals: United States, 1977. National Center for Health Statistics. 1979.
- Haupt BJ. Detailed diagnoses and surgical procedures for patients discharged from short-stay hospitals: United States, 1978. National Center for Health Statistics. 1980.
- Haupt BJ, Graves EJ. Detailed diagnoses and surgical procedures for patients discharged from short-stay hospitals: United States, 1979. National Center for Health Statistics. 1982.
- Kozak LJ, Moien M. Detailed diagnoses and surgical procedures for patients discharged from short-stay hospitals: United States, 1983. National Center for Health Statistics. *Vital Health Stat* 13(82). 1985.
- Lawrence L. Detailed diagnoses and procedures for patients discharged from short-stay hospitals: United States, 1984. National Center for Health Statistics. *Vital Health Stat* 13(86). 1986.
- Pokras R. Detailed diagnoses and procedures for patients discharged from short-stay hospitals: United States, 1985. National Center for Health Statistics. *Vital Health Stat* 13(90). 1987.
- Graham D. Detailed diagnoses and procedures for patients discharged from short-stay hospitals: United States, 1986. National Center for Health Statistics. *Vital Health Stat* 13(95). 1988.
- Graves EJ. Detailed diagnoses and procedures for patients discharged from short-stay hospitals: United States, 1987. National Center for Health Statistics. *Vital Health Stat* 13(100). 1989.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1988. National Center for Health Statistics. *Vital Health Stat* 13(107). 1991.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1989. National Center for Health Statistics. *Vital Health Stat* 13(108). 1991.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1990. National Center for Health Statistics. *Vital Health Stat* 13(113). 1992.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1991. National Center for Health Statistics. *Vital Health Stat* 13(115). 1994.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1992. National Center for Health Statistics. *Vital Health Stat* 13(118). 1994.
- Graves EJ. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1993. National Center for Health Statistics. *Vital Health Stat* 13(122). 1995.
- Graves EJ, Gillum BS. Detailed diagnoses and procedures, National Hospital Discharge Survey, 1994. National Center for Health Statistics. *Vital Health Stat* 13(127). 1997.
- Velebil P, Wingo PA, Xia Z, Wilcox LS, Peterson HB. Rate of hospitalization for gynecologic disorders among reproductive-age women in the United States. *Obstet and Gynecol* 86(5): 764–9. 1995.
- Gillum RF. Epidemiology of carotid endarterectomy and cerebral arteriography in the United States. *Stroke* 26(9): 1724–8. 1995.
- Rosenblum LS, Castro KG, Dooley S, Morgan M. Effect of HIV infection and tuberculosis on hospitalizations and cost of care for young adults in the United States, 1985 to 1990. *Ann Intern Med* 121(10):786–92. 1994.
- Fingerhut LA, Gillum BS. Injury among persons 1–24 years of age in the United States: data from the National Center for Health Statistics. In: *Proceedings of the international collaborative effort on injury statistics*. Hyattsville, Maryland: Public Health Service. 1995.
- Fox K, Merrill JC, Chang HH, Califano JA Jr. Estimating the costs of substance abuse to the Medicaid hospital care program. *Am J Public Health* 85(1):48–54. 1995.
- Kozak LJ, Hall MJ, Pokras R, Lawrence L. Ambulatory surgery in the United States, 1994. Advance data from vital and health statistics; no 283. Hyattsville, Maryland: National Center for Health Statistics. 1997.
- SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. 1987.
- SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. April 1991.
- SMG Marketing Group, Inc. Hospital Market Database. Chicago: Healthcare Information Specialists. April 1994.
- Haupt BJ, Kozak LJ. Estimates from two survey designs: National Hospital Discharge Survey. National Center for Health Statistics. *Vital Health Stat* 13(111). 1992.
- Massey JT, Moore TF, Parsons VL, Tadros W. Design and estimation for the National Health Interview Survey, 1985–94. National Center for Health Statistics. *Vital Health Stat* 2(110). 1989.
- Public Health Service and Health Care Financing Administration. *International Classification of Diseases, 9th Revision, Clinical Modification*. Washington: Public Health Service. 4th ed. 1991.
- Shah BV, Barnwell BG, Bieler GS. SUDAAN User's Manual: Software for Analysis of Correlated Data, Release 6.40. Research Triangle Park, NC: Research Triangle Institute. 1995.
- Hansen MH, Hurwitz WN, Madow WG. *Sample survey methods and theory, vol 1*. New York: John Wiley & Sons. 1953.
- Cochran WG. *Sampling techniques*, 3rd ed. New York: John Wiley & Sons. 1977.



**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region				
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West	
Number of first-listed diagnoses in thousands												
All codes . . . . .	30,722	12,198	18,525	2,405	10,593	6,168	11,556	7,051	6,994	11,373	5,303	
001-139 . . . . .	885	445	440	193	242	141	309	212	169	348	156	
003 . . . . .	12	*7	*5	*5	*	*	*	*	*	*	*	
003.0 . . . . .	11	*6	*	*	*	*	*	*	*	*	*	
008 . . . . .	119	45	74	46	21	12	40	25	27	44	23	
008.4 . . . . .	31	*7	24	*	*	*	22	*8	*7	10	*5	
008.45 . . . . .	27	*6	21	*	*	*	20	*7	*7	9	*	
008.6 . . . . .	23	12	11	22	*	*	*	*	*6	9	*5	
008.61 . . . . .	21	11	10	21	*	*	-	*	*6	*9	*5	
008.8 . . . . .	61	23	38	20	16	*8	17	15	13	22	11	
009 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*	
009.0 . . . . .	*5	*	*	-	*	*	*	*	*	*	*	
011 . . . . .	11	*8	*	*	*	*	*	*	*	*6	*	
011.9 . . . . .	*8	*6	*	*	*	*	*	*	*	*5	*	
034 . . . . .	12	*5	*7	*7	*	*	*	*	*	*5	*	
034.0 . . . . .	11	*5	*6	*6	*	*	*	*	*	*	*	
038 . . . . .	308	133	175	19	27	52	210	61	65	132	50	
038.0 . . . . .	26	13	13	*	*	*	16	*6	*7	11	*	
038.1 . . . . .	46	22	24	*	*	*8	32	12	10	18	*7	
038.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*	
038.4 . . . . .	88	32	56	*	*7	16	64	18	19	33	18	
038.40 . . . . .	*8	*	*	*	*	*	*	*	*	*	*	
038.42 . . . . .	50	15	36	*	*	10	36	9	12	18	11	
038.43 . . . . .	*6	*	*	*	*	*	*	*	*	*	*	
038.49 . . . . .	23	10	14	*	*	*	19	*	*	11	*	
038.8 . . . . .	13	*	*9	*	*	*	10	*	*	*6	*	
038.9 . . . . .	124	56	68	12	10	20	82	20	24	61	19	
041 . . . . .	15	9	*6	*	*	*	*7	*	*	*7	*	
042 . . . . .	114	83	31	*5	87	20	*	45	11	41	18	
047 . . . . .	38	19	19	21	14	*	*	*8	*	17	*7	
047.9 . . . . .	36	18	18	20	14	*	*	*7	*	17	*7	
052 . . . . .	13	9	*	*9	*	*	*	*	*	*	*	
052.9 . . . . .	*7	*6	*	*5	*	*	*	*	*	*	*	
053 . . . . .	19	*8	11	*	*	*5	12	*6	*	*7	*	
053.9 . . . . .	12	*	*8	*	*	*	*8	*	*	*	*	
054 . . . . .	*8	*	*	*	*	*	*	*	*	*	*	
070 . . . . .	15	*9	*6	*	10	*	*	*	*	*5	*	
070.5 . . . . .	*5	*	*	-	*	*	*	*	*	*	*	
075 . . . . .	*8	*	*	*	*5	*	*	*	*	*	*	
078 . . . . .	9	*6	*	*	*	*	*	*	*	*	*	
079 . . . . .	74	34	40	50	12	*8	*	20	15	27	12	
079.9 . . . . .	69	31	38	45	12	*7	*	19	13	26	11	
079.99 . . . . .	69	31	38	44	12	*7	*	18	13	26	11	
112 . . . . .	17	*8	*9	*	*	*	*6	*	*	*7	*	
112.4 . . . . .	*5	*	*	-	*	*	*	*	*	*	*	
117 . . . . .	*6	*	*	*	*	*	*	*	*	*	*	
135 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*	
136 . . . . .	13	10	*	*	*6	*	*	*	*	*	*	
136.3 . . . . .	10	*9	*	*	*6	*	*	*	*	*	*	
140-239 . . . . .	1,802	691	1,112	36	327	589	851	454	343	670	336	
150 . . . . .	15	10	*	-	*	*5	*9	*	*	*5	*	
150.9 . . . . .	*5	*	*	-	*	*	*	*	*	*	*	

\*Figure does not meet standard of reliability or precision.

-Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
151 . . . . .	28	15	13	—	*	*8	18	*6	*6	10	*6
151.0 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
151.8 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
153 . . . . .	92	42	50	—	*	15	74	24	28	27	12
153.1 . . . . .	*8	*	*	—	*	*	*7	*	*	*	*
153.2 . . . . .	*7	*	*	—	*	*	*6	*	*	*	*
153.3 . . . . .	25	10	15	—	*	*	19	*7	*7	*8	*
153.4 . . . . .	16	*8	*8	—	*	*	13	*	*	*	*
153.6 . . . . .	13	*	*8	—	*	*	11	*	*5	*	*
153.9 . . . . .	10	*	*6	—	*	*	*8	*	*	*	*
154 . . . . .	45	24	22	—	*	12	30	15	10	12	*9
154.0 . . . . .	16	*8	*7	—	*	*	11	*	*	*6	*
154.1 . . . . .	25	14	11	—	*	*7	16	10	*6	*	*
155 . . . . .	12	*9	*	*	*	*	*7	*	*	*5	*
155.0 . . . . .	*9	*6	*	*	*	*	*5	*	*	*	*
156 . . . . .	*8	*	*	—	*	*	*5	*	*	*	*
157 . . . . .	29	14	14	—	*	9	19	*7	*	11	*6
157.0 . . . . .	13	*7	*5	—	*	*	9	*	*	*5	*
157.8 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
157.9 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
158 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
161 . . . . .	11	*7	*	—	*	*6	*	*	*	*	*
162 . . . . .	168	92	76	*	*8	64	95	44	32	68	24
162.2 . . . . .	10	*5	*5	—	*	*	*6	*	*	*	*
162.3 . . . . .	49	24	25	—	*	19	28	13	12	18	*7
162.5 . . . . .	22	13	*8	—	*	*8	14	*6	*	*8	*
162.8 . . . . .	18	12	*6	*	*	*6	12	10	*	*	*
162.9 . . . . .	65	36	29	—	*5	27	33	11	10	33	10
170 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
171 . . . . .	10	*5	*	*	*	*	*	*	*	*	*
172 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
173 . . . . .	14	*7	*6	*	—	*	9	*	*	*5	*
174 . . . . .	139	—	139	—	22	56	62	37	27	47	28
174.1 . . . . .	*5	—	*5	—	*	*	*	*	*	*	*
174.2 . . . . .	*7	—	*7	—	*	*	*	*	*	*	*
174.4 . . . . .	25	—	25	—	*	10	10	*7	*7	*7	*
174.5 . . . . .	*6	—	*6	—	*	*	*	*	*	*	*
174.8 . . . . .	40	—	40	—	*	15	20	13	*7	*8	12
174.9 . . . . .	50	—	50	—	*9	23	18	*9	10	25	*6
180 . . . . .	19	—	19	—	*8	*8	*	*5	*	11	*
180.8 . . . . .	*6	—	*6	—	*	*	*	*	*	*	*
180.9 . . . . .	10	—	10	—	*7	*	*	*	*	*8	*
182 . . . . .	42	—	42	—	*	18	21	*8	*7	18	10
182.0 . . . . .	42	—	42	—	*	18	20	*8	*7	18	9
183 . . . . .	39	—	39	*	*	15	20	10	*6	17	*6
183.0 . . . . .	38	—	38	*	*	15	20	10	*6	17	*6
185 . . . . .	97	97	—	*	*	32	64	19	22	33	21
188 . . . . .	51	34	18	—	*	11	40	19	11	16	*6
188.8 . . . . .	15	10	*6	—	*	*	13	*9	*	*	*
188.9 . . . . .	17	12	*5	—	*	*5	12	*	*	*7	*
189 . . . . .	38	20	18	*	*	11	20	*7	*7	17	*7
189.0 . . . . .	32	17	16	*	*	9	16	*6	*6	15	*6

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
191 . . . . .	28	18	10	*	*8	10	*8	*9	*	9	*5
191.1 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
191.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
193 . . . . .	10	*	*7	*	*	*	*	*	*	*	*
194 . . . . .	*6	*	*	*	*	*	–	*	*	*	*
196 . . . . .	10	*6	*	–	*	*	*5	*	*	*	*
197 . . . . .	124	46	77	*	11	42	70	32	22	49	21
197.0 . . . . .	29	13	16	*	*	11	13	*7	*6	12	*
197.2 . . . . .	27	11	15	–	*	*7	19	*8	*	*8	*6
197.6 . . . . .	17	*	14	–	*	*	11	*	*	*6	*
197.7 . . . . .	36	13	23	*	*	12	20	10	*6	15	*5
198 . . . . .	113	47	66	*	*9	42	60	30	23	39	21
198.3 . . . . .	36	16	20	*	*	15	18	*8	*8	13	*8
198.5 . . . . .	46	19	27	*	*	17	25	15	*7	19	*
198.8 . . . . .	22	*9	13	*	*	*6	13	*	*6	*	*7
198.89 . . . . .	18	*8	10	*	*	*5	10	*	*5	*	*5
199 . . . . .	19	*7	12	–	*	*6	11	*	*	*9	*6
199.0 . . . . .	9	*	*5	–	*	*	*7	*	*	*	*
199.1 . . . . .	9	*	*7	–	*	*	*	*	*	*	*
200 . . . . .	21	12	*9	*	*5	*	11	*5	*	11	*
200.0 . . . . .	13	*8	*	*	*	*	*9	*	*	*8	*
200.00 . . . . .	*7	*6	*	*	*	*	*6	*	*	*6	*
201 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
201.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
202 . . . . .	44	24	20	*	*8	13	23	*9	*5	23	*7
202.8 . . . . .	40	23	17	*	*8	11	21	*8	*5	21	*7
202.80 . . . . .	14	*8	*6	*	*	*	*5	*	*	*	*
202.81 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
202.83 . . . . .	*8	*	*	–	*	*	*	*	*	*	*
202.88 . . . . .	*8	*6	*	–	*	*	*6	*	*	*5	*
203 . . . . .	23	11	12	–	*	*7	15	*	*	12	*
203.0 . . . . .	23	11	12	–	*	*7	15	*	*	12	*
203.00 . . . . .	23	11	12	–	*	*7	15	*	*	12	*
204 . . . . .	24	12	12	9	*	*	*7	*6	*	*7	*7
204.0 . . . . .	15	*8	*8	9	*	*	*	*	*	*	*6
204.00 . . . . .	11	*7	*	*6	*	*	*	*	*	*	*
204.1 . . . . .	*8	*	*	–	*	*	*6	*	*	*	*
204.10 . . . . .	*8	*	*	–	*	*	*6	*	*	*	*
205 . . . . .	16	*9	*8	*	*	*	9	*6	*	*	*
205.0 . . . . .	12	*7	*5	*	*	*	*8	*5	*	*	*
205.00 . . . . .	11	*7	*	*	*	*	*8	*5	*	*	*
208 . . . . .	*6	*	*	–	*	*	*	*	*	*	*
208.0 . . . . .	*5	*	*	–	*	*	*	*	*	*	*
210 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
210.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
211 . . . . .	41	20	21	*	*6	10	24	19	*7	10	*5
211.3 . . . . .	32	16	17	*	*	10	19	16	*	*8	*
214 . . . . .	*8	*	*	–	*	*	*	*	*	*	*
218 . . . . .	192	–	192	–	115	74	*	36	38	76	41
218.0 . . . . .	27	–	27	–	16	11	*	*8	*5	*9	*5
218.1 . . . . .	49	–	49	–	29	20	*	14	11	17	*8
218.2 . . . . .	21	–	21	–	13	*8	*	*	*	10	*
218.9 . . . . .	94	–	94	–	57	35	*	10	19	41	23
220 . . . . .	30	–	30	*	17	*6	*7	*8	*8	9	*5

\*Figure does not meet standard of reliability or precision.  
–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
225 . . . . .	15	*	11	*	*	*6	*5	*	*	*6	*
225.2 . . . . .	9	*	*7	*	*	*	*	*	*	*	*
226 . . . . .	9	*	*7	—	*	*	*	*	*	*	*
227 . . . . .	15	*	10	*	*5	*6	*	*	*	*5	*
227.1 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
227.3 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
228 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
233 . . . . .	30	*	29	—	15	11	*	*7	*7	11	*5
233.0 . . . . .	12	*	11	—	*	*6	*	*	*	*	*
233.1 . . . . .	16	—	16	—	12	*	*	*	*	*7	*
235 . . . . .	*9	*	*	*	*	*	*6	*	*	*	*
235.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
236 . . . . .	*7	*	*6	*	*	*	*	*	*	*	*
237 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
237.5 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
238 . . . . .	13	*7	*6	*	*	*	*9	*	*	*	*
238.7 . . . . .	9	*6	*	*	*	*	*7	*	*	*	*
239 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
240-279 . . . . .	1,278	523	755	143	248	322	565	283	293	495	207
241 . . . . .	12	*	11	*	*	*	*	*	*	*	*
241.1 . . . . .	*6	*	*6	—	*	*	*	*	*	*	*
242 . . . . .	15	*	13	*	*7	*	*5	*	*	*	*
242.0 . . . . .	*7	*	*6	*	*5	*	*	*	*	*	*
242.00 . . . . .	*6	*	*6	—	*	*	*	*	*	*	*
242.9 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
242.90 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
244 . . . . .	13	*	10	—	*	*	*7	*6	*	*	*
244.9 . . . . .	11	*	9	—	*	*	*6	*5	*	*	*
250 . . . . .	493	232	261	21	112	172	188	107	104	201	80
250.0 . . . . .	163	69	94	*7	29	65	62	35	34	73	21
250.00 . . . . .	50	23	27	*	*8	22	20	*9	*9	22	11
250.01 . . . . .	33	14	19	*	*	13	12	10	*7	13	*
250.02 . . . . .	39	15	24	*	*6	16	17	10	9	18	*
250.03 . . . . .	41	16	24	*	11	14	13	*6	*9	21	*
250.1 . . . . .	89	42	47	10	51	19	*9	19	22	30	18
250.10 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
250.11 . . . . .	31	14	17	*	19	*	*	*6	*8	12	*
250.13 . . . . .	45	20	26	*	27	9	*	11	11	13	10
250.2 . . . . .	13	*6	*7	—	*	*	*8	*	*	*	*
250.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
250.4 . . . . .	26	12	14	—	*	12	9	*	*	12	*6
250.40 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
250.41 . . . . .	*8	*	*	—	*	*5	*	*	*	*	*
250.42 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
250.5 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
250.6 . . . . .	32	15	17	*	*6	15	12	*7	*7	13	*6
250.60 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
250.61 . . . . .	12	*	*7	—	*	*6	*	*	*	*5	*
250.63 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
250.7 . . . . .	59	34	25	—	*	19	35	13	11	21	14
250.70 . . . . .	20	10	10	—	*	*	14	*	*	*6	*5
250.71 . . . . .	23	14	9	—	*	*9	13	*6	*	*8	*5
250.72 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
250.73 . . . . .	10	*7	*	—	*	*	*	*	*	*6	*
250.8 . . . . .	82	44	38	*	9	29	42	20	17	35	10

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
250.80	20	13	*7	—	*	*6	12	*6	*	*7	*
250.81	34	17	17	*	*	12	17	*6	*7	16	*6
250.82	11	*	*6	—	*	*	*6	*	*	*6	*
250.83	17	*9	*8	*	*	*7	*7	*5	*	*6	*
250.9	18	*7	11	*	*	*7	*6	*	*	9	*
250.92	*5	*	*	*	*	*	*	*	*	*	*
250.93	*8	*	*	*	*	*	*	*	*	*6	—
251	12	*7	*5	*	*	*	*5	*	*	*	*
251.2	*9	*5	*	*	*	*	*	*	*	*	*
253	12	*5	*6	*	*	*	*8	*	*	*	*
253.6	10	*	*5	*	*	*	*8	*	*	*	*
255	*8	*	*6	*	*	*	*	*	*	*	*
263	15	*7	*8	*	*	*	11	*	*	*6	*
263.9	12	*6	*6	*	*	*	9	*	*	*6	*
272	26	12	14	*	*8	12	*5	*7	*	11	*
272.0	15	9	*5	—	*	*8	*	*	*	*5	*
272.4	*7	*	*6	—	*	*	*	*	*	*	*
274	16	11	*5	—	*	*	10	*6	*	*	*
274.0	11	*7	*	—	*	*	*8	*	*	*	*
275	15	*7	*8	*	*	*6	*6	*	*	*6	*
275.4	12	*5	*7	*	*	*	*5	*	*	*5	*
276	573	212	361	108	70	90	306	114	148	223	88
276.0	*8	*	*5	*	*	*	*	*	*	*	*
276.1	60	17	43	*	*	*7	47	*8	17	24	11
276.2	*8	*	*	*	*	*	*	*	*	*	*
276.5	416	163	253	100	48	58	210	90	104	163	59
276.6	*6	*	*	*	*	*	*	*	*	*	*
276.7	17	*8	*8	*	*	*5	10	*	*	*6	*
276.8	51	13	38	*	10	14	26	*9	13	19	10
276.9	*7	*	*5	*	*	*	*	*	*	*	*
277	14	*6	*8	*	*6	*	*	*6	*	*	*
277.0	*8	*	*	*	*	*	—	*	*	*	—
277.00	*8	*	*	*	*	*	—	*	*	*	—
278	24	*7	17	*	16	*8	*	*	*	*9	*8
278.0	22	*6	16	*	14	*7	*	*	*	*8	*8
280–289	339	142	197	59	101	58	121	80	80	131	49
280	38	10	28	*	*8	*6	23	*7	11	17	*
280.0	18	*	14	*	*5	*	11	*	*	*8	*
280.9	20	*6	14	*	*	*	12	*	*8	*9	*
281	*7	*	*6	*	*	*	*	*	*	*	*
282	69	33	36	22	43	*	*	14	15	29	12
282.6	64	30	34	20	42	*	*	13	13	28	10
282.60	*6	*	*	*	*	*	—	*	*	*	*
282.62	55	25	30	15	38	*	*	11	11	25	*8
284	23	11	12	*	*5	*	11	*6	*6	*7	*
284.8	22	10	11	*	*	*	11	*6	*6	*7	*
285	69	27	41	*	13	13	41	12	15	34	*8
285.1	21	*6	15	*	*	*	12	*	*7	*8	*
285.9	44	19	25	*	9	*7	27	*7	*7	25	*5
286	17	10	*6	*	*	*	*8	*6	*	*	*
286.7	*6	*	*	—	*	*	*	*	*	*	*
287	36	16	20	*9	11	*7	10	11	*9	10	*6
287.3	13	*	*9	*	*6	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
287.4	*6	*	*	*	*	*	*	*	*	*	*
287.5	14	*5	*8	*	*	*	*5	*	*	*	*
288	51	19	32	*8	9	18	15	13	13	17	*8
288.0	43	16	27	*7	*7	17	12	11	11	15	*6
288.8	*7	*	*	*	*	*	*	*	*	*	*
289	24	13	11	9	*7	*	*	*6	*6	*6	*6
289.2	10	*5	*	*	*	*	—	*	*	*	*
290-319	2,009	1,023	986	90	1,216	387	316	659	516	652	182
290	67	26	41	*	*	*	64	16	16	27	*7
290.0	23	10	13	—	*	*	22	*	*	11	*
290.2	10	*	*7	—	—	*	9	*	*	*	*
290.20	*6	*	*	—	—	*	*6	*	*	*	*
290.3	13	*6	*7	*	—	*	13	*	*	*	*
290.4	16	*6	10	—	—	*	16	*	*	*5	*
290.40	*8	*	*5	—	—	*	*7	*	*	*	*
291	84	64	20	*	45	32	*6	18	38	18	9
291.0	20	16	*	*	10	*8	*	*	*6	*7	*
291.8	60	46	14	*	33	23	*	13	31	10	*6
292	34	17	18	*	21	*	10	13	*7	10	*
292.0	13	*7	*6	*	10	*	*	*7	*	*	*
292.1	*6	*	*	*	*	—	*	*	*	*	*
292.8	15	*6	*9	*	*7	*	*6	*	*	*	*
292.81	*7	*	*5	—	*	*	*5	*	*	*	*
292.84	*5	*	*	*	*	*	*	*	*	*	*
293	38	15	23	—	10	11	18	17	*	13	*
293.0	14	*	9	—	*	*	11	*	*	*	*
293.8	23	10	13	—	*8	*9	*6	12	*	*8	*
293.81	*8	*	*6	—	*	*	*	*	*	*	*
293.83	13	*7	*6	—	*6	*	*	9	*	*	*
294	23	10	13	*	*	*	19	*6	*6	*7	*
294.8	17	*7	10	—	*	*	15	*	*5	*	*
295	274	155	119	*	184	67	23	125	65	63	21
295.3	78	44	34	—	49	20	9	39	10	22	*7
295.30	12	*6	*6	—	*7	*	*	*	*	*	*
295.32	11	*7	*	—	*7	*	*	*6	*	*	*
295.34	52	30	22	—	34	15	*	30	*6	13	*
295.4	*6	*	*	*	*6	*	*	*	*	*	*
295.6	27	16	10	*	14	9	*	10	*8	*8	*
295.62	*8	*6	*	—	*	*	*	*	*	*	*
295.64	14	*8	*6	—	*7	*6	*	*5	*	*	*
295.7	115	59	56	*	84	25	*6	46	36	25	*9
295.70	61	32	28	*	44	14	*	13	24	18	*5
295.74	53	26	27	*	39	11	*	31	11	*7	*
295.9	41	28	14	*	27	11	*	26	*8	*	*
295.90	13	*6	*7	*	*8	*	*	*	*7	*	*
295.94	26	20	*6	—	17	*7	*	24	*	*	*
296	598	211	387	31	360	112	95	163	160	223	51
296.2	206	76	129	19	121	36	30	42	54	93	17
296.20	101	40	62	*9	60	19	14	24	36	34	*7
296.22	21	*8	14	*	13	*	*	*	*	17	*
296.23	54	18	36	*	33	*8	*9	10	9	30	*
296.24	26	11	15	*	14	*5	*	*6	*7	10	*
296.3	219	70	149	*6	135	35	43	70	61	75	13
296.30	65	19	46	*	37	10	16	20	24	18	*
296.32	13	*	10	*	9	*	*	*	*	*7	*
296.33	98	31	67	*	67	15	13	35	22	35	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
296.34	37	13	24	*	20	*7	9	11	11	12	*
296.4	44	19	26	*	25	12	*7	18	9	10	*7
296.40	24	10	14	*	13	*6	*	*6	*7	*8	*
296.44	15	*6	9	–	*8	*	*	*9	*	*	*
296.5	35	11	24	*	20	*9	*5	*8	*8	14	*
296.50	18	*	13	*	11	*	*	*	*6	*6	*
296.53	9	*	*6	*	*	*	*	*	*	*	*
296.54	*6	*	*	*	*	*	*	*	*	*	*
296.6	42	14	28	*	26	11	*	11	11	15	*
296.60	22	*8	14	*	15	*	*	*	*8	9	*
296.63	*7	*	*	*	*	*	*	*	*	*	*
296.64	10	*	*7	–	*6	*	*	*	*	*	*
296.7	29	12	17	*	19	*5	*	*7	11	*7	*
296.8	13	*6	*7	*	*8	*	*	*6	*	*	*
296.80	*5	*	*	*	*	*	*	*	*	*	*
296.82	*6	*	*	*	*	*	*	*	*	*	*
297	13	*	*8	*	*	*	*5	*	*	*	*
297.1	*5	*	*	–	*	*	*	*	*	*	*
297.9	*7	*	*	*	*	*	*	*	*	*	*
298	72	35	37	*	44	11	16	29	11	25	*8
298.9	66	31	35	*	41	*8	16	27	9	22	*7
300	87	33	54	*6	45	19	18	22	18	34	13
300.0	34	13	20	*	14	*9	10	*8	*	16	*
300.00	22	*9	13	*	*8	*6	*8	*5	*	10	*
300.01	*9	*	*6	*	*5	*	*	*	*	*	*
300.1	11	*	*8	*	*7	*	*	*	*	*	*
300.11	*8	*	*6	*	*	*	*	*	*	*	*
300.4	25	10	16	*5	14	*	*	9	*7	*6	*
300.9	11	*	*7	*	*6	*	*	*	*	*5	*
301	18	9	*9	*	15	*	*	10	*	*	*
301.8	11	*6	*5	*	11	*	*	*8	*	*	*
301.83	*6	*	*	*	*6	–	*	*	*	*	*
303	247	183	64	*	163	71	12	68	74	87	18
303.0	43	32	11	–	24	16	*	14	13	11	*6
303.00	11	*7	*	–	*7	*	*	*	*	*	*
303.01	31	24	*7	–	17	13	*	13	*9	*6	*
303.9	204	150	54	*	139	56	*9	54	62	76	12
303.90	60	42	18	–	40	16	*	14	12	29	*5
303.91	141	106	35	*	97	39	*	40	49	46	*6
304	131	91	40	*	115	14	*	57	21	50	*
304.0	32	20	12	–	27	*	*	21	*	*7	*
304.00	*6	*	*	–	*	*	*	*	*	*	*
304.01	26	15	10	–	22	*	–	20	*	*	–
304.2	48	35	13	–	44	*	*	23	*6	18	*
304.20	13	*9	*	–	12	*	*	*	*	11	*
304.21	34	25	9	–	31	*	–	22	*	*7	*
304.3	*8	*6	*	*	*7	–	–	*	*	*	*
304.31	*7	*5	*	*	*6	–	–	*	*	*	–
304.7	15	10	*	–	13	*	–	10	*	*	*
304.71	12	*9	*	–	10	*	–	*9	*	*	*
304.8	13	11	*	–	11	*	–	*	*	*9	*
304.81	*8	*7	*	–	*7	*	–	*	*	*6	*
304.9	*8	*	*	*	*7	*	*	*	*	*5	*
304.90	*6	*	*	*	*	*	*	*	*	*	–
305	51	36	15	*	35	11	*	10	14	19	*8
305.0	21	16	*6	*	14	*	*	*	*6	*8	*
305.00	16	11	*5	*	10	*	*	*	*5	*7	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
305.1	*7	*5	*	—	*	*	*	*	*	*	*
305.6	*9	*6	*	—	*8	*	—	*	*	*	*
305.60	*6	*	*	—	*6	*	—	*	*	*	*
305.9	*8	*	*	*	*6	*	*	*	*	*	*
305.90	*7	*	*	*	*5	*	*	*	*	*	*
306	*6	*	*	*	*	*	*	*	*	*	*
307	11	*	*8	*	*6	*	*	*	*	*	*
308	*7	*	*6	*	*	*	*	*	*	*	*
309	110	55	55	10	85	11	*	52	22	20	15
309.0	45	22	23	*	36	*	*	20	*7	10	*8
309.2	12	*	*8	*	9	*	*	*	*	*	*
309.28	11	*	*7	*	*9	*	*	*	*	*	*
309.4	27	17	11	*	21	*	*	18	*5	*	*
309.8	12	*	*7	*	*8	*	*	*5	*	*	*
309.81	11	*	*7	*	*7	*	—	*	*	*	*
309.9	*8	*	*	*	*7	*	*	*	*	*	*
310	14	*7	*7	*	*	*	*7	*	*	*	*
310.9	*6	*	*	—	*	*	*	*	*	*	*
311	71	31	40	*8	49	*7	*7	21	22	20	*7
312	23	16	*7	*	17	*	*	12	*5	*	*
312.3	15	10	*5	*	13	*	*	*8	*	*	*
312.30	*7	*5	*	*	*6	*	—	*	*	*	—
312.34	*6	*	*	*	*	*	*	*	*	*	*
313	*8	*5	*	*5	*	—	—	*	*	*	*
313.8	*8	*5	*	*5	*	—	—	*	*	*	*
313.81	*8	*5	*	*5	*	—	—	*	*	*	*
314	12	10	*	10	*	—	—	*	*	*6	*
314.0	11	10	*	10	*	—	—	*	*	*5	*
314.01	11	9	*	10	*	—	—	*	*	*5	*
320-389	561	247	314	102	136	113	210	161	116	200	84
320	14	*7	*7	10	*	*	*	*	*	9	*
320.9	*5	*	*	*	*	—	*	*	*	*	*
322	*6	*	*	*	*	*	*	*	*	*	*
322.9	*5	*	*	*	*	*	*	*	*	*	*
323	*7	*	*5	*	*5	*	*	*	*	*5	*
323.9	*7	*	*5	*	*5	*	*	*	*	*5	*
331	25	13	12	*	*	*	17	*6	*	9	*
331.0	9	*	*	—	—	*	9	*	*	*	*
331.3	*5	*	*	*	*	*	*	*	*	*	*
331.4	*7	*	*	*	*	*	*	*	*	*	*
332	16	*9	*7	—	*	*	15	*	*7	*	*
332.0	16	*9	*7	—	*	*	15	*	*6	*	*
333	9	*	*6	*	*	*	*	*	*	*	*
337	*7	*	*6	*	*	*	*	*	*	*	*
337.2	*5	*	*	*	*	*	*	*	*	*	*
340	22	*	18	*	11	9	*	*	*7	*8	*
342	26	15	12	*	*	*7	16	*	*	13	*
342.9	26	14	12	*	*	*7	16	*	*	13	*
342.90	20	11	*9	*	*	*5	13	*	*	11	*
344	12	*5	*7	*	*5	*	*	*	*	*	*
345	59	30	28	20	22	*7	10	14	16	19	9
345.1	13	*7	*6	*	*	*	*	*	*6	*	*
345.10	11	*6	*	*	*	*	*	*	*	*	*
345.3	13	*7	*6	*5	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).



**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
345.5	*6	*	*	*	*	*	*	*	*	*	*
345.9	17	*9	*8	*5	*8	*	*	*	*	*8	*
345.90	13	*7	*6	*	*5	*	*	*	*	*7	*
346	36	*7	29	*	20	10	*	*7	10	15	*
346.9	19	*	15	*	10	*5	*	*	*6	*9	*
346.90	13	*	10	*	*8	*	*	*	*	*5	*
346.91	*5	*	*5	*	*	*	*	*	*	*	*
348	28	12	15	*6	*5	*5	11	*7	*7	*7	*7
348.1	*8	*	*	*	*	*	*	*	*	*	*
348.3	*8	*	*	*	*	*	*	*	*	*	*
349	13	*	*9	*	*7	*	*	*	*	*	*
349.8	*6	*	*	*	*	*	*	*	*	*	*
351	*5	*	*	*	*	*	*	*	*	*	*
353	*5	*	*	*	*	*	*	*	*	*	*
354	11	*	*6	*	*6	*	*	*7	*	*	*
354.0	*5	*	*	–	*	*	*	*	–	*	*
355	*8	*	*5	*	*	*	*	*	*	*	*
356	*6	*	*	*	*	*	*	*	*	*	*
357	*7	*	*	*	*	*	*	*	*	*	*
358	*7	*	*	*	*	*	*	*	*	*	*
358.0	*7	*	*	*	*	*	*	*	*	*	*
361	16	10	*6	*	*	*7	*7	*	*	10	*
361.0	*9	*6	*	*	*	*	*	*	*	*6	*
362	*7	*	*	*	*	*	*	*	*	*	*
365	*6	*	*	*	*	*	*	*	*	*	*
366	54	21	33	*	*	*7	44	38	*	11	*
366.1	12	*	9	–	–	*	11	9	*	*	*
366.10	*7	*	*5	–	–	*	*7	*7	–	–	*
366.8	*8	*5	*	–	–	*	*6	*	*	*	*
366.9	34	13	20	*	*	*	27	24	*	*9	–
376	*7	*	*	*	*	*	*	*	*	*	*
376.0	*7	*	*	*	*	*	*	*	*	*	*
376.01	*7	*	*	*	*	*	*	*	*	*	*
380	*6	*	*	*	*	*	*	*	*	*	*
380.1	*5	*	*	*	*	*	*	*	*	*	*
381	*8	*	*	*7	*	*	*	*	*	*	*
382	26	15	12	23	*	*	*	*6	*7	*9	*
382.9	24	13	11	21	*	*	*	*	*6	*9	*
386	28	*8	20	*	*	9	15	*7	*9	*9	*
386.1	*6	*	*	*	*	*	*	*	*	*	*
386.3	18	*6	12	*	*	*5	10	*5	*5	*5	*
386.30	16	*6	10	*	*	*5	*8	*5	*	*	*
390–459	5,830	2,973	2,857	21	421	1,589	3,799	1,384	1,380	2,156	911
394	*6	*	*5	*	*	*	*	*	*	*	*
396	12	*5	*7	*	*	*	*8	*	*	*	*
401	195	82	113	*	31	82	81	45	39	84	27
401.0	27	9	18	*	*	10	12	*7	*5	12	*
401.9	164	70	94	*	25	70	68	36	33	70	24
402	90	39	51	–	*	26	61	20	29	29	11
402.9	84	36	48	–	*	23	58	18	28	27	10
402.90	15	*7	*8	–	*	*8	*7	*	*	*	*
402.91	69	29	39	–	*	15	52	13	23	24	*9

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
403	67	38	29	*	18	20	28	11	11	37	*8
403.0	*6	*	*	—	*	*	*	*	*	*	*
403.01	*5	*	*	—	*	*	*	*	*	*	*
403.9	61	34	26	*	14	19	27	9	10	34	*8
403.90	*5	*	*	*	*	*	*	*	*	*	*
403.91	55	32	24	*	13	18	24	9	10	29	*7
404	32	15	17	*	*	*	23	*8	*6	13	*
404.9	27	13	14	*	*	*	22	*7	*6	10	*
404.93	21	11	10	—	*	*	16	*5	*	*8	*
410	771	468	303	—	43	256	471	188	175	268	140
410.0	37	23	14	—	*	12	22	*5	10	15	*7
410.01	32	19	13	—	*	11	18	*	9	12	*6
410.1	140	86	54	—	11	48	80	34	32	47	27
410.11	122	77	46	—	10	42	71	28	29	43	23
410.12	13	*7	*6	—	*	*6	*7	*	*	*	*
410.2	26	16	10	—	*	11	14	*5	*6	12	*
410.21	26	15	10	—	*	11	13	*5	*6	11	*
410.3	20	13	*6	—	*	*9	10	*7	*	*7	*
410.31	17	12	*5	—	*	*8	*8	*6	*	*5	*
410.4	170	110	60	—	12	72	86	44	43	54	29
410.41	159	101	58	—	12	68	80	41	41	50	27
410.42	9	*7	*	—	*	*	*6	*	*	*	*
410.5	23	15	*8	—	*	*9	12	*	*5	9	*
410.51	21	14	*8	—	*	*8	11	*	*	*9	*
410.6	*5	*	*	—	*	*	*	*	*	*	*
410.61	*5	*	*	—	*	*	*	*	*	*	*
410.7	251	145	105	—	*8	68	174	70	53	82	46
410.70	*6	*	*	—	*	*	*	*	*	*	*
410.71	227	134	93	—	*7	62	157	65	46	76	40
410.72	18	9	*9	—	*	*	12	*	*	*5	*
410.8	11	*6	*	—	*	*	*9	*	*	*	*
410.81	9	*5	*	—	*	*	*8	*	*	*	*
410.9	89	52	37	—	*	24	62	14	18	38	18
410.90	22	14	*8	—	*	*5	16	*	*	10	*6
410.91	66	38	28	—	*	19	45	11	14	28	13
411	312	157	155	*	26	104	182	83	83	104	43
411.1	285	141	144	—	23	95	167	76	78	96	36
411.8	25	15	10	*	*	*8	14	*7	*	*6	*7
411.81	11	*7	*	*	*	*	*6	*	*	*	*
411.89	14	*8	*6	—	*	*	*8	*5	*	*	*
412	10	*7	*	—	*	*	*	*	*	*	*
413	115	52	62	—	12	39	63	24	37	42	12
413.9	111	50	62	—	10	38	63	23	36	40	12
414	922	585	338	—	46	359	517	209	200	345	168
414.0	895	567	328	—	44	350	501	203	195	335	163
414.00	202	116	86	—	*9	74	119	47	35	96	24
414.01	647	414	232	—	35	260	352	150	149	222	126
414.02	47	37	10	—	*	16	30	*7	10	17	13
414.8	*8	*6	*	—	*	*	*	*	*	*	*
414.9	18	11	*7	—	*	*5	11	*	*	*8	*
415	65	26	38	*	*8	19	37	12	19	22	12
415.1	63	25	38	*	*8	19	36	11	19	22	11
416	11	*5	*6	*	*	*	*7	*	*	*	*
416.0	*6	*	*	*	*	*	*	*	*	*	*
420	14	*6	*7	*	*	*5	*6	*	*	*	*
420.9	14	*6	*7	*	*	*5	*6	*	*	*	*
420.90	10	*	*5	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
421 . . . . .	17	12	*5	*	*8	*	*5	*	*	*8	*
421.0 . . . . .	17	12	*5	*	*7	*	*5	*	*	*8	*
423 . . . . .	10	*	*7	*	*	*	*	*	*	*	*
423.9 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
424 . . . . .	82	39	43	*	15	19	47	20	13	37	11
424.0 . . . . .	32	14	18	*	10	12	11	*6	*6	16	*
424.1 . . . . .	46	23	23	*	*6	*6	34	14	*7	19	*6
425 . . . . .	34	21	13	*	*5	13	15	*9	*8	11	*7
425.4 . . . . .	30	19	12	*	*	12	14	*7	*8	9	*6
426 . . . . .	53	24	29	*	*6	*6	40	16	11	21	*5
426.0 . . . . .	26	13	13	*	*	*	22	*7	*6	10	*
426.1 . . . . .	11	*5	*6	*	*	*	*9	*	*	*	*
426.13 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
427 . . . . .	585	277	307	*5	48	131	400	148	131	205	101
427.0 . . . . .	35	15	20	*	*6	12	17	*8	*9	13	*
427.1 . . . . .	40	23	17	*	*	11	26	11	11	13	*
427.3 . . . . .	302	143	158	*	14	64	222	75	66	103	58
427.31 . . . . .	270	122	148	*	12	55	203	67	60	96	47
427.32 . . . . .	31	21	10	*	*	9	19	*7	*6	*7	11
427.4 . . . . .	11	*6	*	*	*	*	*6	*	*	*	*
427.41 . . . . .	10	*6	*	*	*	*	*5	*	*	*	*
427.5 . . . . .	13	*9	*	*	*	*	9	*	*	*6	*
427.6 . . . . .	21	12	9	*	*	*	12	*5	*	9	*
427.69 . . . . .	18	11	*7	*	*	*	10	*	*	*7	*
427.8 . . . . .	153	63	90	*	17	32	101	42	34	52	26
427.81 . . . . .	56	26	29	*	*	*8	46	16	*9	21	10
427.89 . . . . .	98	37	61	*	16	23	55	26	25	31	15
427.9 . . . . .	9	*6	*	*	*	*	*6	*	*	*	*
428 . . . . .	880	384	497	*	32	143	704	214	226	331	109
428.0 . . . . .	872	378	494	*	31	142	698	212	224	329	107
428.1 . . . . .	*7	*	*	*	*	*	*6	*	*	*	*
429 . . . . .	18	*9	10	*	*	*	12	*	*	10	*
429.2 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
429.4 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
430 . . . . .	28	11	17	*	*7	10	10	*6	*8	9	*
431 . . . . .	57	25	32	*	*5	13	38	10	12	24	12
432 . . . . .	16	*8	*8	*	*	*	11	*	*5	*6	*
432.1 . . . . .	13	*7	*6	*	*	*	*9	*	*	*	*
433 . . . . .	175	101	74	*	*	37	136	38	42	73	23
433.1 . . . . .	127	72	55	*	*	29	97	28	33	51	16
433.10 . . . . .	107	61	46	*	*	24	82	24	27	43	14
433.11 . . . . .	20	11	*9	—	*	*5	14	*	*6	*8	*
433.3 . . . . .	42	24	18	*	*	*7	35	*8	*9	20	*6
433.30 . . . . .	32	19	14	*	*	*6	26	*6	*6	16	*
433.31 . . . . .	10	*5	*	—	—	*	*8	*	*	*	*
434 . . . . .	275	114	161	*	*8	46	220	75	63	93	44
434.0 . . . . .	18	10	*8	—	*	*	15	*5	*	*6	*
434.01 . . . . .	16	*8	*7	—	*	*	13	*	*	*5	*
434.1 . . . . .	33	15	18	—	*	*6	26	14	*6	*8	*6
434.11 . . . . .	29	13	16	—	*	*6	22	13	*5	*6	*6
434.9 . . . . .	224	89	135	*	*7	38	179	56	54	79	35
434.90 . . . . .	*8	*	*	*	*	*	*5	*	*	*	*
434.91 . . . . .	217	86	131	*	*6	35	174	55	51	77	34
435 . . . . .	175	83	92	*	*	33	139	48	47	63	18
435.0 . . . . .	*6	*	*	—	—	*	*6	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
435.8	*5	*	*	—	*	*	*	*	*	*	*
435.9	160	78	82	*	*	31	126	44	42	58	16
436	149	61	88	*	*	28	116	18	41	67	23
437	36	14	22	*	*	*8	24	*6	10	18	*
437.1	*9	*	*6	*	—	*	*8	*	*	*	*
437.2	*8	*	*	—	*	*	*	*	*	*	*
437.3	*6	*	*	*	*	*	*	*	*	*	*
438	15	*7	*7	*	*	*	9	*	*	*6	*
440	108	59	49	*	*	25	81	26	25	38	18
440.2	92	50	41	*	*	20	70	22	21	34	15
440.20	10	*6	*	—	*	*	*8	*	*	*	*
440.21	20	13	*7	—	*	*7	13	*	*	9	*
440.22	*9	*	*	—	*	*	*6	*	*	*	*
440.23	13	*7	*6	—	*	*	10	*	*	*	*
440.24	38	20	18	—	*	*5	32	9	9	13	*6
441	61	47	14	*	*	11	49	14	13	22	12
441.0	*8	*	*	—	*	*	*6	*	*	*	*
441.01	*5	*	*	—	*	*	*	*	*	*	*
441.3	*7	*	*	—	—	*	*6	*	*	*	*
441.4	40	32	*7	—	*	*7	33	*8	10	15	*7
442	10	*5	*	*	*	*	*6	*	*	*	*
442.3	*6	*	*	—	*	*	*5	*	*	*	*
443	20	9	10	—	*	*	14	*6	*	*8	*
443.9	17	*8	*8	—	*	*	12	*5	*	*6	*
444	53	25	28	*	*	15	36	13	13	19	*8
444.0	*6	*	*	*	*	*	*	*	*	*	*
444.2	36	17	20	*	*	*9	26	10	9	11	*6
444.22	32	15	17	—	*	*8	23	*9	*8	9	*6
444.8	*9	*	*	—	*	*	*	*	*	*	*
444.81	*7	*	*	—	*	*	*	*	*	*	*
446	*8	*	*	*	*	*	*	*	*	*	*
447	16	*8	*8	*	*	*5	*8	*	*	*6	*
447.1	11	*5	*6	—	*	*	*6	*	*	*	*
451	46	16	30	*	11	15	21	9	12	18	*8
451.0	*8	*	*6	—	*	*	*	*	*	*	*
451.1	24	9	15	*	*	*7	13	*5	*6	*8	*
451.11	*6	*	*	*	*	*	*	*	*	*	*
451.19	18	*7	11	—	*	*6	*8	*	*	*6	*
451.2	*9	*	*7	—	*	*	*	*	*	*	*
453	128	55	73	*	19	35	74	29	34	46	18
453.8	121	52	69	*	18	32	70	28	33	42	17
453.9	*6	*	*	—	*	*	*	*	*	*	*
454	21	*9	13	—	*6	*	11	*5	*6	*6	*
454.2	12	*5	*6	—	*	*	*8	*	*	*	*
455	38	17	20	*	13	10	14	15	*	13	*5
455.2	21	*8	13	*	*7	*	9	*7	*	*8	*
456	*5	*	*	*	*	*	*	*	*	*	*
458	51	16	35	*	*	12	35	10	14	20	*7
458.0	24	*	19	—	*	*	17	*5	*9	*6	*
458.9	27	11	16	*	*	*7	17	*	*6	14	*
459	18	*8	10	*	*	*6	11	*	*7	*	*
459.8	*7	*	*	*	*	*	*5	*	*	*	*
459.81	*7	*	*	—	*	*	*	*	*	*	*
459.9	*5	*	*	—	—	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
460-519 . . . . .	3,329	1,590	1,738	766	462	572	1,528	715	816	1,315	483
461 . . . . .	17	*9	*8	*5	*	*	*	*	*6	*7	*
461.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
462 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
463 . . . . .	14	*7	*8	*8	*6	*	—	*	*	*7	*
464 . . . . .	40	27	13	34	*	*	*	10	12	11	*7
464.4 . . . . .	31	21	10	31	—	*	*	*7	10	*9	*
465 . . . . .	33	17	17	21	*	*	*	9	*6	13	*
465.9 . . . . .	32	16	16	20	*	*	*	*8	*6	13	*
466 . . . . .	270	148	122	168	25	27	51	52	69	104	45
466.0 . . . . .	130	57	72	29	24	26	50	26	40	57	*7
466.1 . . . . .	140	91	49	138	*	*	*	26	30	48	37
470 . . . . .	11	*	*6	—	*6	*	*	*6	*	*	*
473 . . . . .	21	*9	12	*	*8	*	*	*6	*	*9	*
473.9 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
474 . . . . .	32	17	15	25	*6	*	*	21	*	*	*
474.0 . . . . .	13	*6	*7	*9	*	—	—	*7	*	*	*
474.1 . . . . .	18	11	*7	16	*	*	*	13	*	*	*
474.10 . . . . .	14	*9	*6	13	*	—	—	11	*	*	*
475 . . . . .	15	*9	*6	*	9	*	*	*	*	*6	*
478 . . . . .	25	17	*8	*6	*8	*6	*6	*8	*6	*8	*
478.2 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
478.7 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
480 . . . . .	38	21	16	29	*	*	*5	*6	10	14	*8
480.1 . . . . .	19	11	*8	19	*	*	—	*	*7	*6	*
480.9 . . . . .	16	*9	*8	*8	*	*	*	*	*	*7	*
481 . . . . .	60	33	27	*	14	11	32	10	15	20	15
482 . . . . .	241	121	120	11	29	30	171	32	53	124	32
482.0 . . . . .	20	10	10	*	*	*	18	*	*	11	*
482.1 . . . . .	43	23	20	*	*8	*	28	*	*9	22	*7
482.2 . . . . .	28	14	14	*	*	*	20	*	*7	12	*
482.3 . . . . .	15	10	*6	*	*	*	9	*	*	*6	*
482.30 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
482.39 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
482.4 . . . . .	32	14	19	*	*	*	26	*5	*8	15	*
482.8 . . . . .	83	41	43	*	*6	11	63	*7	16	51	*9
482.82 . . . . .	*7	*	*	—	*	*	*6	*	*	*	*
482.83 . . . . .	44	21	23	*	*	*8	32	*	11	25	*5
482.89 . . . . .	30	15	15	*	*	*	23	*	*	22	*
482.9 . . . . .	19	10	9	*	*5	*	*8	*5	*	*7	*
483 . . . . .	14	*7	*6	*	*	*	*	*	*	*6	*
483.0 . . . . .	12	*6	*6	*	*	*	*	*	*	*6	*
485 . . . . .	45	23	22	22	*	*	17	*8	12	22	*
486 . . . . .	849	405	444	176	102	113	458	179	220	332	117
487 . . . . .	19	*7	12	*	*	*	*7	*	*	*8	*
487.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
487.1 . . . . .	10	*	*6	*	*	*	*	*	*	*6	*
490 . . . . .	20	12	*8	*	*	*	11	*	*6	9	*
491 . . . . .	343	141	202	*	17	92	232	64	87	153	40
491.2 . . . . .	339	139	200	*	17	90	230	63	85	152	39
491.20 . . . . .	25	12	14	*	*	*5	18	*6	*	12	*
491.21 . . . . .	313	127	186	*	15	85	212	57	81	139	35
492 . . . . .	33	19	14	*	*	13	19	*6	9	14	*
492.8 . . . . .	30	17	13	*	*	12	18	*5	*8	13	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
493 . . . . .	511	210	301	212	135	87	77	141	135	157	79
493.0 . . . . .	34	18	16	22	*9	*	*	11	*6	*6	11
493.00 . . . . .	13	*7	*6	*8	*	*	*	*	*	*	*5
493.01 . . . . .	21	10	10	14	*6	*	—	9	*	*	*5
493.2 . . . . .	69	22	48	*	*7	25	35	21	15	24	10
493.20 . . . . .	52	17	35	*	*6	18	27	16	11	17	*8
493.21 . . . . .	18	*	13	*	*	*8	*8	*5	*	*7	*
493.9 . . . . .	406	170	236	188	118	59	41	109	114	126	58
493.90 . . . . .	216	84	132	80	66	38	31	49	56	78	33
493.91 . . . . .	190	86	104	107	51	22	10	59	58	48	25
496 . . . . .	157	76	81	*	*	44	109	31	30	77	18
507 . . . . .	124	63	61	*7	*6	12	99	29	30	46	19
507.0 . . . . .	123	63	60	*6	*6	12	98	29	29	46	19
510 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
510.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
511 . . . . .	59	25	34	*	9	17	31	11	13	24	11
511.0 . . . . .	15	*7	*8	—	*5	*	*5	*	*6	*5	*
511.8 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
511.9 . . . . .	34	13	21	*	*	10	22	*6	*6	16	*6
512 . . . . .	30	18	12	*	14	*7	9	*6	10	11	*
512.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
512.8 . . . . .	21	13	*9	*	11	*	*	*5	*7	*8	*
513 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
513.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
514 . . . . .	11	*6	*5	*	*	*	*	*	*	*5	*
515 . . . . .	16	*7	*9	—	*	*	11	*	*5	*	*
516 . . . . .	15	9	*6	*	*	*5	*7	*	*	*6	*
516.3 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
516.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
518 . . . . .	202	92	110	*7	16	51	128	36	38	85	43
518.0 . . . . .	*9	*	*5	*	*	*	*	*	*	*	*
518.4 . . . . .	10	*	*	*	*	*	*5	*	*	*	*
518.5 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
518.8 . . . . .	171	79	93	*6	10	42	114	29	31	74	37
518.81 . . . . .	149	66	82	*	*8	36	103	24	26	66	33
518.82 . . . . .	10	*	*6	*	*	*	*	*	*	*	*
518.89 . . . . .	13	*8	*	*	*	*	*8	*	*	*	*
519 . . . . .	25	11	14	*7	*	*6	10	*7	*5	12	*
519.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
519.8 . . . . .	12	*	*7	*	*	*	*6	*	*	*9	*
520-579 . . . . .	3,031	1,323	1,707	219	807	803	1,202	677	678	1,166	509
522 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
524 . . . . .	19	*6	13	*	16	*	*	*	*	*8	*
524.0 . . . . .	*7	*	*	*	*6	*	—	*	*	*	*
527 . . . . .	12	*	*8	*	*	*	*	*	*	*6	*
527.2 . . . . .	10	*	*6	*	*	*	*	*	*	*5	*
528 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
530 . . . . .	186	78	108	22	32	62	69	49	34	71	32
530.1 . . . . .	48	17	31	*	11	17	19	16	*6	18	*8
530.10 . . . . .	22	*9	13	*	*	*6	10	9	*	9	*
530.11 . . . . .	20	*5	15	*	*5	*8	*6	*6	*	*7	*
530.19 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
530.2 . . . . .	19	9	10	—	*	*7	9	*	*	*9	*
530.3 . . . . .	15	*7	*8	*	*	*	10	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
530.5	*9	*	*	*	*	*	*	*	*	*	*
530.7	*8	*	*	*	*	*	*	*	*	*	*
530.8	79	31	48	19	14	26	20	20	18	31	11
530.81	74	30	44	19	14	25	17	18	18	29	10
531	104	46	58	*	14	25	64	20	25	38	21
531.0	27	11	15	*	*	*	18	*	*7	10	*
531.00	26	11	15	*	*	*	18	*	*7	10	*
531.3	*7	*	*	—	*	*	*	*	*	*	*
531.30	*6	*	*	—	*	*	*	*	*	*	*
531.4	47	24	23	*	*6	12	29	10	10	17	*9
531.40	45	23	22	*	*6	11	28	10	10	17	*9
531.5	*6	*	*	*	*	*	*	*	*	*	*
531.9	12	*	*8	*	*	*	*7	*	*	*	*
531.90	12	*	*8	*	*	*	*6	*	*	*	*
532	80	51	29	*	15	19	46	17	21	26	16
532.0	21	14	*6	—	*	*	13	*	*	*7	*6
532.00	20	14	*5	—	*	*	12	*	*	*6	*6
532.3	*6	*	*	—	*	*	*	*	*	*	*
532.30	*5	*	*	—	*	*	*	*	*	*	*
532.4	31	23	*7	—	*6	*7	18	*8	*7	*9	*7
532.40	29	22	*7	—	*6	*7	17	*7	*7	*8	*6
532.9	12	*	*7	*	*	*	*5	*	*5	*	*
532.90	11	*	*7	*	*	*	*	*	*5	*	*
533	26	*8	18	*	*5	*8	12	*5	*6	12	*
533.4	*8	*	*6	—	*	*	*6	*	*	*5	*
533.40	*8	*	*6	—	*	*	*6	*	*	*5	*
533.9	10	*	*7	*	*	*	*	*	*	*	*
533.90	10	*	*7	*	*	*	*	*	*	*	*
535	169	70	99	*9	46	43	71	35	43	66	24
535.0	40	15	25	*	10	10	18	*7	*9	18	*6
535.00	28	10	18	*	*8	*7	10	*	*7	14	*
535.01	13	*5	*7	*	*	*	*8	*	*	*	*
535.1	13	*	9	*	*	*	*7	*	*	*	*
535.10	*8	*	*6	*	*	*	*	*	*	*	*
535.3	12	*6	*6	—	*6	*	*	*	*	*	*
535.30	*7	*	*	—	*	*	*	*	*	*	*
535.31	*5	*	*	—	*	*	*	*	*	*	*
535.4	38	15	23	*	10	*7	21	*8	12	10	*7
535.40	20	*6	15	*	*7	*	*9	*6	*5	*	*
535.41	18	10	*8	*	*	*	12	*	*7	*6	*
535.5	55	23	32	*	17	14	19	11	10	27	*7
535.50	41	18	24	*	14	11	12	*9	*8	20	*
535.51	13	*5	*8	*	*	*	*7	*	*	*7	*
535.6	10	*6	*	*	*	*	*	*	*	*	*
535.60	*8	*5	*	*	*	*	*	*	*	*	*
536	33	14	18	*	10	*9	10	*8	*	16	*
536.2	11	*	*6	*	*	*	*	*	*	*6	*
536.3	*5	*	*	*	*	*	*	*	*	*	*
536.8	14	*7	*7	*	*5	*	*6	*	*	*8	*
537	21	*8	12	*	*	*	12	*6	*	*6	*5
537.0	12	*5	*7	*	*	*	*6	*	*	*	*
537.8	*8	*	*	*	*	*	*5	*	*	*	*
537.83	*5	*	*	—	*	*	*	*	*	*	*
540	190	116	74	46	109	23	12	37	42	65	46
540.0	44	28	16	13	19	*9	*	*9	10	16	9
540.1	25	18	*6	*6	11	*	*	*	*5	*9	*6
540.9	121	69	51	27	79	12	*	23	26	40	31

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
541 . . . . .	12	*6	*6	*	*8	*	*	*	*	*6	*
550 . . . . .	66	60	*6	*9	13	14	30	28	12	19	*8
550.1 . . . . .	22	19	*	*	*	*	13	*7	*	*7	*
550.10 . . . . .	17	15	*	*	*	*	10	*5	*	*6	*
550.9 . . . . .	43	40	*	*6	*9	12	17	20	*8	11	*
550.90 . . . . .	28	26	*	*	*7	*8	10	14	*	*8	*
550.91 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
550.92 . . . . .	*8	*7	*	*	*	*	*	*	*	*	*
552 . . . . .	40	14	26	*	*8	14	17	13	9	12	*6
552.1 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
552.2 . . . . .	22	*6	16	*	*	*9	9	*7	*	*7	*
552.20 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
552.21 . . . . .	13	*	10	–	*	*5	*6	*	*	*	*
553 . . . . .	91	34	57	*	19	35	36	20	16	43	12
553.1 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
553.2 . . . . .	49	20	29	*	11	18	20	11	11	22	*5
553.20 . . . . .	10	*	*7	–	*	*	*	*	*	*	*
553.21 . . . . .	36	16	21	*	*8	14	15	*7	*8	18	*
553.3 . . . . .	32	11	20	*	*	13	13	*6	*	17	*
555 . . . . .	35	16	20	*	21	*8	*6	*9	9	13	*
555.0 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
555.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
555.9 . . . . .	15	*6	9	*	10	*	*	*	*	*8	*
556 . . . . .	24	11	13	*	10	*6	*7	*5	*6	*9	*
556.9 . . . . .	14	*6	*9	*	*6	*	*	*	*	*6	*
557 . . . . .	30	*7	23	*	*	*6	21	*6	*7	11	*6
557.0 . . . . .	16	*	12	*	*	*	11	*	*	*6	*
557.1 . . . . .	*6	*	*5	*	*	*	*	*	*	*	*
557.9 . . . . .	*7	*	*6	–	*	*	*6	*	*	*	*
558 . . . . .	244	101	143	76	67	40	62	49	52	109	34
558.9 . . . . .	237	98	139	76	66	37	59	45	51	108	33
560 . . . . .	252	115	137	12	33	64	143	46	59	98	48
560.1 . . . . .	43	22	21	*	*5	10	27	*8	11	22	*
560.2 . . . . .	12	*7	*	*	*	*	*8	*	*	*6	*
560.3 . . . . .	26	14	13	*	*	*	19	*5	*	12	*
560.39 . . . . .	24	13	11	*	*	*	17	*5	*	12	*
560.8 . . . . .	78	33	46	*	13	23	41	17	17	23	21
560.81 . . . . .	60	24	36	*	10	18	31	13	13	19	14
560.89 . . . . .	19	9	10	*	*	*5	10	*	*	*	*7
560.9 . . . . .	87	36	51	*	10	26	49	13	24	34	16
562 . . . . .	219	88	131	*	18	65	136	55	52	82	30
562.1 . . . . .	217	87	130	*	18	65	135	55	52	82	29
562.10 . . . . .	31	9	22	*	*	*9	19	*9	*	12	*
562.11 . . . . .	117	50	68	–	14	44	59	31	31	40	16
562.12 . . . . .	53	22	31	–	*	*8	44	13	11	23	*6
562.13 . . . . .	16	*6	10	–	*	*	12	*	*	*7	*
564 . . . . .	44	14	30	*6	13	*8	17	9	12	14	9
564.0 . . . . .	17	*7	11	*	*	*	*7	*	*	*6	*
564.1 . . . . .	17	*	15	*	*8	*	*	*	*6	*	*
564.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
565 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
566 . . . . .	20	14	*7	*	11	*6	*	*5	*	*6	*
567 . . . . .	18	*8	10	*	*	*7	*5	*	*	*7	*
567.2 . . . . .	14	*7	*8	*	*	*6	*	*	*	*6	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
568	11	*	*8	*	*6	*	*	*	*	*	*
568.0	*9	*	*7	*	*5	*	*	*	*	*	*
569	72	32	40	*	10	20	41	20	13	28	11
569.3	12	*5	*6	*	*	*	*6	*	*	*5	*
569.4	*7	*	*	*	*	*	*	*	*	*	*
569.6	11	*	*6	*	*	*	*6	*	*	*	*
569.8	31	14	16	*	*	9	18	*8	*6	12	*5
569.81	*5	*	*	—	*	*	*	*	*	*	*
569.83	*7	*	*	*	*	*	*	*	*	*	*
569.85	12	*6	*6	—	*	*	9	*	*	*	*
570	*6	*	*	*	*	*	*	*	*	*	*
571	77	44	32	*	19	36	21	18	16	31	12
571.1	13	10	*	—	*6	*7	*	*	*	*6	*
571.2	28	20	*8	—	*6	15	*7	*8	*7	10	*
571.5	23	11	12	*	*	*8	13	*5	*	10	*
572	27	15	12	*	*7	9	11	*	*	14	*
572.2	18	9	*8	—	*	*7	*7	*	*	*8	*
573	14	*	10	*	*	*6	*	*	*	*8	*
573.3	*9	*	*6	*	*	*	*	*	*	*	*
574	430	125	306	*	145	129	155	101	95	157	77
574.0	140	54	86	*	41	47	53	26	36	55	23
574.00	127	50	76	—	38	41	48	24	33	50	20
574.01	13	*	9	*	*	*6	*	*	*	*	*
574.1	203	46	156	*	77	63	61	58	37	69	38
574.10	198	45	153	*	75	61	60	58	36	67	37
574.2	46	10	36	*	17	13	15	*8	13	18	*7
574.20	45	10	36	*	17	13	15	*8	12	18	*7
574.3	13	*6	*7	*	*	*	9	*	*	*7	*
574.30	*5	*	*	*	*	*	*	*	*	*	*
574.31	*8	*	*	—	*	*	*6	*	*	*	*
574.4	*9	*	*6	*	*	*	*	*	*	*	*
574.5	19	*5	14	*	*5	*	12	*	*	*	*6
574.50	10	*	*7	*	*	*	*6	*	*	*	*
574.51	9	*	*7	—	*	*	*6	*	*	*	*
575	55	20	34	*	23	17	15	14	14	20	*6
575.0	23	11	12	*	*9	*7	*7	*6	*8	*7	*
575.1	24	*6	18	*	12	*7	*	*6	*5	10	*
575.8	*6	*	*	*	*	*	*	*	*	*	—
576	29	11	18	*	*	*6	19	*7	*	10	*8
576.1	12	*5	*6	*	*	*	*7	*	*	*	*
576.2	*9	*	*7	*	*	*	*5	*	*	*	*
576.8	*6	*	*	*	*	*	*5	*	*	*	*
577	181	86	95	*	62	62	52	31	53	69	27
577.0	148	73	75	*	51	51	44	25	46	58	19
577.1	24	10	14	*	9	*8	*7	*	*6	9	*6
578	143	69	75	*	20	32	89	28	34	63	17
578.0	13	*7	*6	*	*	*	9	*	*	*6	*
578.1	24	11	13	*	*	*	16	*	*7	9	*
578.9	106	51	56	*	15	26	64	20	26	48	12
580-629	1,768	587	1,181	72	654	395	647	373	388	727	281
584	63	37	26	*	*	13	45	16	16	21	10
584.9	59	35	25	*	*	11	44	15	16	19	9
585	37	18	19	*	*8	9	19	11	*7	12	*7
586	10	*	*6	*	*	*	*7	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
590 . . . . .	113	22	91	17	53	21	23	19	29	42	22
590.1 . . . . .	74	16	59	11	37	11	15	13	22	25	14
590.10 . . . . .	74	15	59	11	37	11	15	12	22	25	14
590.8 . . . . .	34	*	29	*6	15	*7	*6	*6	*6	15	*7
590.80 . . . . .	34	*	29	*6	15	*7	*6	*6	*6	15	*7
591 . . . . .	14	*6	*8	*	*	*	*7	*6	*	*	*
592 . . . . .	195	127	68	*	90	76	27	43	50	74	28
592.0 . . . . .	55	32	24	*	27	20	*8	12	14	19	10
592.1 . . . . .	137	93	43	*	61	55	19	30	35	54	17
593 . . . . .	44	20	24	*7	11	*9	17	10	*8	15	11
593.4 . . . . .	17	*9	*8	*	*6	*	*7	*	*	*6	*5
593.7 . . . . .	*6	*	*	*5	*	*	—	*	*	*	*
593.70 . . . . .	*5	*	*	*	*	—	—	*	*	*	*
593.9 . . . . .	11	*	*6	*	*	*	*6	*	*	*	*
594 . . . . .	*9	*7	*	*	*	*	*	*	*	*	*
594.1 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
595 . . . . .	15	*	11	*	*	*	*9	*	*	*	*
595.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
596 . . . . .	26	20	*6	*	*	*6	18	*6	*	10	*
596.0 . . . . .	12	12	*	—	*	*	10	*	*	*	*
598 . . . . .	*7	*6	*	*	*	*	*	*	*	*	*
599 . . . . .	350	123	228	28	39	46	237	78	73	157	43
599.0 . . . . .	319	106	213	27	30	39	223	70	65	146	38
599.7 . . . . .	23	13	10	*	*7	*6	10	*6	*5	*8	*
600 . . . . .	136	136	—	—	*	22	112	31	33	48	23
601 . . . . .	14	14	—	—	*	*	*9	*	*	*7	*
601.0 . . . . .	11	11	—	—	*	*	*7	*	*	*	*
604 . . . . .	9	9	—	*	*	*	*	*	*	*	*
604.9 . . . . .	*9	*9	—	*	*	*	*	*	*	*	*
604.90 . . . . .	*5	*5	—	*	*	*	*	*	*	*	*
607 . . . . .	*8	*8	—	*	*	*	*	*	*	*	*
607.8 . . . . .	*6	*6	—	—	*	*	*	*	*	*	*
608 . . . . .	*8	*8	—	*	*	*	*	*	*	*	*
610 . . . . .	*8	*	*8	*	*	*	*	*	*	*	*
610.1 . . . . .	*7	—	*7	—	*	*	*	*	*	*	*
611 . . . . .	34	*	31	*	20	*8	*	10	*5	13	*
611.1 . . . . .	24	*	21	*	17	*	*	*8	*	*9	*
614 . . . . .	74	—	74	*	64	*7	*	17	16	31	9
614.2 . . . . .	*7	—	*7	*	*6	*	—	*	*	*	*
614.3 . . . . .	11	—	11	*	10	*	*	*	*	*	*
614.6 . . . . .	24	—	24	—	20	*	*	*	*	12	*
614.9 . . . . .	18	—	18	*	17	*	—	*	*	*8	*
616 . . . . .	10	—	10	*	*8	*	*	*	*	*6	*
616.0 . . . . .	*6	—	*6	—	*	*	*	*	*	*	*
617 . . . . .	92	—	92	—	74	16	*	14	18	47	13
617.0 . . . . .	50	—	50	—	36	12	*	*6	*8	28	*8
617.1 . . . . .	14	—	14	—	12	*	*	*	*	*5	*
617.3 . . . . .	19	—	19	—	17	*	—	*	*	*7	*
618 . . . . .	163	—	163	*	39	60	64	30	37	66	30
618.0 . . . . .	42	—	42	—	*	18	20	*6	14	14	*8
618.1 . . . . .	17	—	17	—	11	*	*	*	*	*8	*
618.2 . . . . .	28	—	28	—	*8	9	10	*8	*5	10	*
618.3 . . . . .	19	—	19	—	*	*6	11	*5	*	*5	*
618.4 . . . . .	23	—	23	—	*8	10	*5	*	*	13	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
618.5	19	–	19	–	*	*7	11	*	*	*8	*
618.8	10	–	10	–	*	*	*	*	*	*	*
619	*7	–	*7	–	*5	*	*	*	*	*	*
619.1	*6	–	*6	–	*	*	*	*	*	*	*
620	67	–	67	*	53	10	*	15	16	25	11
620.0	*7	–	*7	*	*5	*	*	*	*	*	*
620.1	13	–	13	*	11	*	–	*	*	*	*
620.2	40	–	40	*	31	*6	*	*9	10	14	*7
621	19	–	19	*	*8	*6	*	*6	*	*6	*
621.3	10	–	10	–	*	*	*	*	*	*	*
622	12	–	12	–	10	*	*	*	*	*8	*
622.1	10	–	10	–	9	*	*	*	*	*7	*
625	97	–	97	*	57	28	12	10	19	47	21
625.3	*8	–	*8	*	*8	*	–	*	*	*6	*
625.6	49	–	49	–	14	25	10	*7	9	21	12
625.9	33	–	33	*	29	*	*	*	*7	18	*6
626	80	–	80	*	61	17	*	*9	17	41	13
626.2	56	–	56	*	44	11	*	*6	10	31	10
626.8	18	–	18	*	14	*	*	*	*6	*9	*
627	14	–	14	–	*	9	*	*	*	*5	*
627.1	12	–	12	–	*	*8	*	*	*	*	*
630–676	542	–	542	*	540	*	–	115	104	215	107
632	16	–	16	–	16	–	–	*	*	9	*
633	41	–	41	–	41	*	–	11	*7	14	*7
633.1	38	–	38	–	37	*	–	10	*7	14	*6
634	38	–	38	*	37	*	–	18	*	12	*
634.9	32	–	32	*	31	*	–	15	*	11	*
634.91	25	–	25	*	24	*	–	13	*	*8	*
635	*6	–	*6	*	*6	*	–	*	*	*	*
637	*6	–	*6	–	*6	–	–	*	*	*	*
637.9	*6	–	*6	–	*6	–	–	*	*	*	*
640	*7	–	*7	–	*7	–	–	*	*	*	*
640.0	*5	–	*5	–	*5	–	–	*	*	*	*
640.03	*5	–	*5	–	*5	–	–	*	*	*	*
641	17	–	17	–	17	–	–	*	*	*6	*
641.1	*6	–	*6	–	*6	–	–	*	*	*	*
641.13	*6	–	*6	–	*6	–	–	*	*	*	*
642	25	–	25	–	25	–	–	*	*5	11	*
642.3	*8	–	*8	–	*8	–	–	*	*	*	*
642.33	*7	–	*7	–	*7	–	–	*	*	*	*
642.4	10	–	10	–	10	–	–	*	*	*	*
642.43	*9	–	*9	–	*9	–	–	*	*	*	*
643	41	–	41	–	41	–	–	*8	10	19	*
643.0	15	–	15	–	15	–	–	*	*5	*7	*
643.03	15	–	15	–	15	–	–	*	*	*7	*
643.1	22	–	22	–	22	–	–	*5	*	10	*
643.13	22	–	22	–	22	–	–	*5	*	10	*
644	142	–	142	*	142	–	–	23	31	53	35
644.0	107	–	107	*	107	–	–	17	24	41	25
644.03	107	–	107	*	106	–	–	17	24	41	24
644.1	35	–	35	–	35	–	–	*5	*7	12	10
644.13	34	–	34	–	34	–	–	*5	*7	12	10
646	45	–	45	*	44	–	–	*8	*8	18	11
646.6	34	–	34	*	34	–	–	*6	*	15	*8

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
646.63	30	–	30	*	30	–	–	*6	*	13	*7
646.8	*7	–	*7	*	*7	–	–	*	*	*	*
646.83	*7	–	*7	*	*6	–	–	*	*	*	*
647	11	–	11	*	11	–	–	*	*	*	*
647.6	*5	–	*5	*	*5	–	–	*	*	*	*
648	69	–	69	*	69	*	–	17	15	26	12
648.0	*6	–	*6	–	*6	–	–	*	*	*	*
648.03	*6	–	*6	–	*6	–	–	*	*	*	*
648.4	*6	–	*6	*	*5	–	–	*	*	*	*
648.43	*5	–	*5	*	*	–	–	*	*	*	*
648.8	15	–	15	–	15	–	–	*	*	*6	*
648.83	15	–	15	–	15	–	–	*	*	*6	*
648.9	37	–	37	*	37	*	–	9	*7	14	*7
648.93	35	–	35	*	35	*	–	*9	*7	12	*7
654	10	–	10	–	10	–	–	*	*	*5	*
654.5	*6	–	*6	–	*6	–	–	*	*	*	*
654.53	*6	–	*6	–	*6	–	–	*	*	*	*
656	*5	–	*5	–	*5	–	–	*	*	*	*
658	10	–	10	–	10	–	–	*	*	*5	*
658.1	*6	–	*6	–	*6	–	–	*	*	*	*
658.13	*5	–	*5	–	*5	–	–	*	*	*	*
659	11	–	11	–	11	–	–	*	*	*	*
659.1	*5	–	*5	–	*5	–	–	*	*	*	*
659.13	*5	–	*5	–	*5	–	–	*	*	*	*
666	*8	–	*8	*	*8	–	–	*	*	*6	*
666.2	*7	–	*7	*	*7	–	–	*	*	*6	*
666.24	*7	–	*7	*	*7	–	–	*	*	*6	*
670	*7	–	*7	–	*7	–	–	*	*	*	*
670.0	*7	–	*7	–	*7	–	–	*	*	*	*
670.04	*7	–	*7	–	*7	–	–	*	*	*	*
680–709	482	235	246	41	126	118	197	118	119	170	74
681	24	13	11	*	*8	*6	*7	*6	*7	*8	*
681.0	12	*8	*	*	*5	*	*	*	*	*	*
681.00	11	*7	*	*	*	*	*	*	*	*	*
681.1	12	*6	*6	*	*	*	*5	*	*	*5	*
681.10	11	*	*6	*	*	*	*	*	*	*5	*
682	325	168	157	23	86	89	127	81	81	112	52
682.0	26	13	13	*	13	*	*	*	*6	*8	*7
682.1	11	*8	*	*	*7	*	*	*	*	*5	*
682.2	26	12	14	*	9	*6	*9	*8	*	*9	*
682.3	33	17	16	*	12	*9	9	*9	10	10	*
682.4	20	12	*8	*	*8	*	*7	*6	*	*8	*
682.6	151	73	78	*6	26	45	73	43	36	52	21
682.7	46	25	22	*	*8	13	22	*8	16	16	*6
686	*7	*	*	*	*	*	*	*	*	*	*
692	*5	*	*	*	*	*	*	*	*	*	*
695	*8	*	*6	*	*	*	*	*	*	*	*
695.1	*5	*	*	*	*	*	*	*	*	*	*
706	*6	*	*	*	*	*	*	*	*	*	*
706.2	*6	*	*	*	*	*	*	*	*	*	*
707	69	29	40	*	10	10	49	12	20	29	*8
707.0	36	15	22	*	*7	*	27	*	11	16	*
707.1	31	14	17	*	*	*6	21	*7	*7	13	*
709	*7	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
710-739 . . . . .	1,479	648	831	37	391	417	634	333	322	547	277
710 . . . . .	25	*	21	*	13	*8	*	*	*	12	*5
710.0 . . . . .	19	*	15	*	11	*5	*	*	*	9	*
711 . . . . .	19	10	*9	*	*	*	*8	*	*6	*	*
711.0 . . . . .	17	9	*8	*	*	*	*7	*	*6	*	*
711.06 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
714 . . . . .	32	*9	23	*	*	10	17	*9	*8	9	*6
714.0 . . . . .	28	*6	21	*	*	*9	16	*7	*7	*8	*5
715 . . . . .	355	126	230	*	14	86	255	78	90	117	70
715.1 . . . . .	32	15	18	—	*	*7	24	*	*8	14	*7
715.15 . . . . .	11	*	*7	—	*	*	*7	*	*	*	*
715.16 . . . . .	20	11	*9	—	*	*	15	*	*5	9	*
715.3 . . . . .	232	79	153	—	*9	56	167	47	44	91	50
715.35 . . . . .	75	32	43	—	*	15	55	14	10	32	19
715.36 . . . . .	149	45	104	—	*	37	108	32	32	57	29
715.9 . . . . .	83	30	54	*	*	21	59	26	37	10	11
715.90 . . . . .	*6	*	*	—	—	*	*	*	*	*	—
715.95 . . . . .	25	10	16	—	*	*8	16	*9	10	*	*
715.96 . . . . .	47	15	32	*	*	11	34	13	24	*	*
716 . . . . .	25	11	14	*	*	*8	11	*7	*	*9	*
716.9 . . . . .	13	*	9	—	*	*	*7	*	*	*5	*
717 . . . . .	37	22	14	*	29	*6	*	17	*8	*7	*5
717.8 . . . . .	22	14	*9	*	19	*	*	*8	*7	*	*
717.83 . . . . .	21	13	*9	*	18	*	—	*8	*7	*	*
718 . . . . .	29	17	12	*	19	*	*	*8	*	10	*7
718.3 . . . . .	11	*7	*	*	*8	*	*	*	*	*	*
718.31 . . . . .	*7	*6	*	—	*6	*	*	*	*	*	*
718.8 . . . . .	*8	*	*	—	*6	*	*	*	*	*	*
719 . . . . .	39	17	22	*	*8	*5	23	*8	*7	15	*9
719.4 . . . . .	17	*7	10	*	*5	*	*8	*	*	*	*5
719.45 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
719.7 . . . . .	11	*	*8	—	*	*	10	*	*	*6	*
720 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
721 . . . . .	66	28	39	—	*8	24	34	11	15	27	14
721.0 . . . . .	22	*9	13	—	*	11	*8	*	*5	11	*
721.1 . . . . .	13	*8	*	—	*	*6	*5	*	*	*6	*
721.3 . . . . .	27	10	17	—	*	*7	17	*	*6	*8	*7
722 . . . . .	327	187	140	—	158	115	54	62	67	140	58
722.0 . . . . .	59	36	23	—	29	24	*6	10	11	30	*7
722.1 . . . . .	208	120	89	—	108	71	30	41	44	87	37
722.10 . . . . .	207	119	87	—	107	70	29	41	43	86	36
722.5 . . . . .	22	10	12	—	*6	*6	10	*	*7	*7	*
722.52 . . . . .	22	10	12	—	*6	*6	9	*	*6	*7	*
722.7 . . . . .	15	10	*	—	*7	*	*	*	*	*	*5
722.71 . . . . .	*7	*5	*	—	*	*	*	*	*	*	*
722.73 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
722.8 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
722.83 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
722.9 . . . . .	*9	*	*5	—	*	*	*	*	*	*	*
722.93 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
723 . . . . .	17	11	*6	*	*6	*7	*	*6	*	*7	*
723.4 . . . . .	*8	*6	*	—	*	*	*	*	*	*	*
724 . . . . .	104	41	63	*	25	30	48	20	26	39	19
724.0 . . . . .	45	19	26	—	*	13	29	10	9	17	9

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
724.02	43	18	25	—	*	13	28	10	9	16	*9
724.2	26	*9	17	*	10	*8	*7	*	*6	12	*
724.4	11	*	*6	—	*	*	*	*	*	*	*
724.5	12	*	*8	*	*	*	*	*	*	*	*
726	49	26	23	*	12	26	11	16	9	15	9
726.1	14	*8	*6	—	*	*9	*	*	*	*	*
726.10	13	*8	*5	—	*	*8	*	*	*	*	*
726.2	18	10	*8	—	*	12	*	*	*	*7	*
726.6	*5	*	*	*	*	*	*	*	*	*	*
727	28	13	15	*	11	*9	*6	11	*5	*8	*
727.0	10	*	*5	*	*5	*	*	*	*	*	*
727.1	*6	*	*	—	*	*	*	*	*	*	*
728	18	*8	10	*	*6	*	*6	*5	*6	*5	*
728.8	9	*	*	*	*	*	*	*	*	*	*
728.89	*7	*	*	*	*	*	*	*	*	*	*
729	30	11	19	*	*8	11	11	*	*5	15	*5
729.1	10	*	*7	*	*	*	*	*	*	*6	*
729.5	10	*	*6	*	*	*	*	*	*	*	*
729.8	*6	*	*	*	*	*	*	*	*	*	*
730	52	29	22	*5	11	14	22	13	10	20	*9
730.0	16	9	*7	*	*	*	*7	*	*	*5	*
730.07	9	*	*	*	*	*	*	*	*	*	*
730.1	10	*6	*	*	*	*	*	*	*	*	*
730.2	25	14	11	*	*6	*7	10	*8	*	10	*
730.27	11	*7	*	*	*	*	*	*	*	*	*
730.28	*5	*	*	*	*	*	*	*	*	*	—
732	*6	*	*	*	*	*	—	*	*	*	*
733	168	55	113	*	31	33	101	32	33	70	33
733.1	83	17	66	*	*	*8	72	15	19	32	17
733.13	51	9	41	—	*	*	46	*8	12	20	10
733.14	16	*	12	*	*	*	13	*	*	*	*
733.15	*5	*	*	*	*	*	*	*	*	*	*
733.19	*6	*	*	—	*	*	*	*	*	*	*
733.4	19	*9	10	—	*8	*	*6	*	*	*7	*5
733.42	14	*7	*7	—	*6	*	*	*	*	*5	*
733.6	23	*7	16	—	*7	*8	*9	*	*	13	*
733.8	33	19	14	*	11	9	11	*7	*	15	*6
733.81	*5	*	*	*	*	*	*	*	*	*	*
733.82	28	15	12	*	*9	*8	9	*6	*	13	*
735	*8	*	*6	*	*	*	*	*	*	*	*
735.0	*5	*	*	—	*	*	*	*	*	*	*
736	*9	*5	*	*	*	*	*	*	*	*	*
737	12	*	9	*	*	*	*	*	*	*	*
737.3	11	*	9	*	*	*	*	*	*	*	*
737.30	*8	*	*7	*	*	*	*	*	*	*	*
738	17	*9	*8	*	*6	*6	*	*	*	*6	*
738.4	12	*6	*6	—	*	*6	*	*	*	*	*
740-759	159	87	73	120	24	10	*5	56	28	37	38
742	*6	*	*	*	*	—	*	*	*	*	*
745	26	14	12	21	*	*	*	9	*	*5	*8
745.2	*6	*	*	*5	*	*	—	*	*	*	*
745.4	*5	*	*	*	*	*	—	*	*	*	*
745.5	*7	*	*	*	*	*	*	*	*	*	*
746	13	*6	*7	11	*	*	*	*5	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
747	16	*9	*8	11	*	*	*	*5	*	*	*
747.0	*5	*	*	*	*	*	–	*	*	*	*
749	*9	*5	*	*8	*	–	–	*	*	*	*
750	10	*7	*	9	*	*	–	*	*	*	*
750.5	*7	*5	*	*7	–	–	–	*	*	*	*
751	11	*7	*	*8	*	*	*	*	*	*	*
752	*8	*	*	*	*	*	*	*	*	*	*
753	*9	*	*	*7	*	*	*	*	*	*	*
754	10	*6	*	*8	*	*	–	*	*	*	*
756	11	*6	*5	*5	*	*	*	*	*	*	*
759	*6	*	*	*	*	*	*	*	*	*	*
760–779	142	86	56	141	*	–	–	23	41	41	37
765	18	10	*8	18	*	–	–	*	*	*6	*
765.1	16	*9	*7	16	*	–	–	*	*	*6	*
769	14	*8	*6	14	*	–	–	*	*	*	*7
770	29	18	11	29	*	–	–	*5	*8	*9	*6
770.8	17	*8	*8	16	*	–	–	*	*	*7	*
771	12	*7	*5	12	–	–	–	*	*	*	*
771.8	11	*6	*	11	–	–	–	*	*	*	*
774	35	22	13	35	–	–	–	*	12	*7	10
774.6	30	19	11	30	–	–	–	*	9	*7	10
775	*7	*6	*	*7	–	–	–	*	*	*5	*
775.5	*5	*	*	*5	–	–	–	*	*	*	*
778	*7	*	*	*7	–	–	–	*	*	*	*
778.4	*6	*	*	*6	–	–	–	*	*	*	*
779	*7	*	*	*7	*	–	–	*	*	*	*
780–799	312	143	170	50	121	77	64	66	61	144	41
780	64	30	34	22	18	10	14	17	14	23	10
780.2	15	*9	*7	*	*	*	*8	*	*	*5	*
780.3	28	14	13	12	11	*	*	*8	*6	10	*
780.6	11	*	*7	*7	*	*	*	*	*	*	*
784	*8	*	*	*	*	*	*	*	*	*	*
785	*6	*	*	*	*	*	*	*	*	*	*
786	94	49	45	*8	36	36	14	16	20	44	13
786.0	*8	*	*5	*6	*	*	*	*	*	*	*
786.09	*8	*	*	*6	*	*	*	*	*	*	*
786.5	81	44	37	*	35	36	11	15	18	37	11
786.50	70	38	32	*	31	29	10	13	16	32	10
786.52	*8	*5	*	–	*	*	*	*	*	*	*
787	*6	*	*	*	*	*	*	*	*	*	*
788	*8	*	*	–	*5	*	*	*	*	*	*
788.0	*6	*	*	–	*	*	*	*	*	*	–
789	49	14	35	*6	31	*7	*5	9	10	21	*8
789.0	47	14	33	*6	30	*6	*	*9	10	20	*8
789.00	18	*5	12	*	12	*	*	*	*	*9	*
789.01	*6	*	*	*	*	*	*	*	*	*	*
789.03	13	*	*8	*	*8	*	*	*	*	*	*
799	69	31	37	*8	21	18	23	17	11	38	*
799.9	68	31	37	*8	21	18	22	17	11	37	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
800-999 . . . . .	2,591	1,289	1,302	241	925	482	943	593	577	911	510
800 . . . . .	14	10	*	*9	*	*	*	*	*	*	*
800.0 . . . . .	*7	*	*	*6	*	*	*	*	*	*	*
801 . . . . .	27	21	*6	*	14	*	*	*7	*	*7	*9
801.0 . . . . .	12	10	*	*	*8	*	*	*	*	*	*
802 . . . . .	44	34	10	*	34	*	*	*9	*8	17	10
802.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
802.2 . . . . .	12	*9	*	*	10	*	*	*	*	*	*
802.4 . . . . .	11	*7	*	*	*9	*	*	*	*	*5	*
805 . . . . .	63	30	33	*	21	10	31	11	21	21	10
805.0 . . . . .	13	*8	*	*	*7	*	*	*	*	*	*
805.2 . . . . .	17	*7	10	*	*	*	9	*	*6	*	*
805.4 . . . . .	31	15	16	*	*8	*	17	*	10	12	*
806 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
807 . . . . .	44	22	22	*	16	*9	19	9	11	14	10
807.0 . . . . .	37	20	18	*	14	*7	16	*7	*9	13	*9
807.01 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
807.04 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
807.09 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
807.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
808 . . . . .	57	24	34	*	13	*7	36	9	15	19	14
808.0 . . . . .	11	*8	*	*	*5	*	*	*	*	*	*
808.2 . . . . .	32	9	22	*	*	*	22	*5	*8	*9	9
808.4 . . . . .	*7	*	*	—	*	*	*5	*	*	*	*
808.8 . . . . .	*7	*	*	*	*	*	*5	*	*	*	*
810 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
810.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
812 . . . . .	67	20	47	11	11	11	34	16	13	21	17
812.0 . . . . .	34	9	25	*	*	*6	25	9	*6	11	*7
812.00 . . . . .	11	*	*9	*	*	*	*8	*	*	*	*
812.01 . . . . .	15	*	10	*	*	*	10	*	*	*6	*
812.09 . . . . .	*6	*	*	*	*	*	*5	*	*	*	*
812.2 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
812.21 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
812.4 . . . . .	23	*8	14	10	*5	*	*6	*6	*	*7	*5
812.40 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
812.41 . . . . .	11	*	*7	*8	*	*	*	*	*	*	*
813 . . . . .	51	26	25	*9	20	10	13	14	13	16	*7
813.0 . . . . .	11	*	*6	*	*	*	*	*	*	*	*
813.01 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
813.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
813.4 . . . . .	20	11	10	*	*5	*	*7	*6	*5	*7	*
813.42 . . . . .	12	*7	*	*	*	*	*	*	*	*5	*
815 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
816 . . . . .	*8	*7	*	*	*	*	*	*	*	*	*
816.1 . . . . .	*6	*6	*	*	*	*	*	*	*	*	*
820 . . . . .	301	79	222	*	*6	18	274	63	77	106	55
820.0 . . . . .	97	24	73	*	*	*	90	23	26	29	18
820.02 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
820.03 . . . . .	*8	*	*7	—	*	*	*7	*	*	*	*
820.09 . . . . .	78	19	59	*	*	*	74	19	23	25	11
820.2 . . . . .	137	32	104	*	*	12	122	30	35	48	23
820.20 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
820.21 . . . . .	121	27	94	*	*	10	110	27	30	43	21

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).



**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
820.22	*8	*	*5	*	*	*	*7	*	*	*	*
820.8	66	22	44	*	*	*	61	10	14	28	14
821	61	30	31	12	18	*7	24	16	12	25	*9
821.0	32	21	11	10	12	*	*8	9	*6	12	*5
821.00	*7	*	*	*	*	*	*	*	*	*	*
821.01	25	17	*8	*8	11	*	*	*7	*5	*8	*
821.2	24	*7	17	*	*	*	16	*6	*6	11	*
821.21	*6	*	*	–	*	*	*	*	*	*	*
821.23	11	*	*8	*	*	*	*7	*	*	*	*
822	18	*8	10	*	*6	*	*9	*	*	*5	*7
822.0	17	*8	9	*	*5	*	*9	*	*	*	*6
823	68	44	25	*8	30	17	14	14	14	27	13
823.0	24	11	13	*	*7	*7	*9	*5	*6	*9	*
823.00	20	*9	11	*	*5	*6	*8	*	*	*7	*
823.2	12	*8	*	*	*7	*	*	*	*	*	*
823.22	*9	*6	*	*	*6	*	*	*	*	*	*
823.3	*8	*6	*	*	*5	*	*	*	*	*	*
823.32	*6	*	*	*	*	*	*	*	*	*	*
823.8	16	12	*	*	*7	*5	*	*	*	*6	*
823.80	*6	*5	*	*	*	*	*	*	*	*	*
823.82	*9	*6	*	*	*	*	*	*	*	*	*
824	104	45	59	*	52	30	17	23	25	38	18
824.0	*5	*	*	*	*	*	*	*	*	*	*
824.2	11	*	*7	*	*6	*	*	*	*	*5	*
824.4	24	10	14	*	12	*6	*	*5	*6	*8	*
824.6	26	*7	18	–	10	11	*	*7	*8	*6	*5
824.8	30	16	14	*	16	*7	*	*6	*6	14	*
825	19	12	*7	*	10	*6	*	*	*6	*6	*
825.0	*9	*7	*	*	*	*	*	*	*	*	*
825.2	9	*	*5	*	*	*	*	*	*	*	*
825.25	*6	*	*	*	*	*	*	*	*	*	*
831	*6	*	*	*	*	*	*	*	*	*	*
831.0	*6	*	*	*	*	*	*	*	*	*	*
836	22	13	*8	*	12	*6	*	11	*	*6	*
836.0	12	*9	*	*	*7	*	*	*7	*	*	*
836.1	*7	*	*	*	*	*	*	*	*	*	*
840	41	22	18	*	*7	17	17	11	11	13	*5
840.4	38	21	17	–	*5	16	17	10	10	13	*
844	24	14	9	*	16	*	*	*5	*6	*8	*
844.2	13	*8	*	*	12	*	–	*	*	*	*
845	*7	*	*	*	*	*	*	*	*	*	*
845.0	*6	*	*	–	*	*	*	*	*	*	*
846	*8	*	*	–	*	*	*	*	*	*	*
846.0	*7	*	*	–	*	*	*	*	*	*	*
847	18	*8	10	*	*9	*5	*	*	*6	*	*
847.0	*8	*	*	*	*	*	*	*	*	*	*
847.2	*8	*	*	–	*	*	*	*	*	*	*
850	45	29	16	*6	26	*	9	17	11	12	*
850.0	11	*6	*	*	*	*	*	*	*	*	*
850.1	14	*9	*5	*	*8	*	*	*7	*	*	*
850.5	*8	*6	*	*	*7	*	*	*	*	*	*
850.9	11	*7	*	*	*6	*	*	*	*	*	*
851	14	*9	*5	*	*6	*	*	*	*	*	*
851.8	9	*6	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
852 . . . . .	33	21	11	*	*8	*6	14	*8	*6	*8	10
852.0 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*
852.2 . . . . .	23	15	*9	*	*5	*	12	*5	*	*	9
852.20 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
852.21 . . . . .	12	*8	*	*	*	*	*6	*	*	*	*5
853 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
853.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
854 . . . . .	58	33	24	14	28	*8	*9	9	17	18	13
854.0 . . . . .	56	32	24	13	27	*7	*8	*9	17	18	12
854.00 . . . . .	18	10	*8	*	*7	*	*	*	*	*8	*
854.01 . . . . .	13	*6	*7	*	*6	*	*	*	*	*	*
854.02 . . . . .	14	10	*	*	*8	*	*	*	*	*	*
854.06 . . . . .	*7	*	*	*	*5	*	*	*	*	*	*
860 . . . . .	27	17	10	*	16	*	*6	*	*7	10	*
860.0 . . . . .	13	*8	*5	*	*6	*	*	*	*	*	*
860.4 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
861 . . . . .	12	9	*	*	*6	*	*	*	*	*6	*
861.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
861.21 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
863 . . . . .	11	*9	*	*	*7	*	*	*	*	*	*
864 . . . . .	10	*6	*	*	*6	*	*	*	*	*	*
864.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
865 . . . . .	16	12	*	*	12	*	*	*	*	*6	*
865.0 . . . . .	13	9	*	*	10	*	*	*	*	*	*
866 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
871 . . . . .	*9	*	*	*	*5	*	*	*	*	*	*
873 . . . . .	24	15	10	*	14	*	*	*6	*6	*8	*
873.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
873.4 . . . . .	12	*6	*6	*	*8	*	*	*	*	*	*
873.42 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
874 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
875 . . . . .	10	9	*	*	*7	*	*	*	*	*7	*
875.0 . . . . .	*8	*7	*	*	*6	*	*	*	*	*5	*
879 . . . . .	9	*7	*	*	*8	*	*	*	*	*	*
881 . . . . .	*9	*7	*	*	*7	*	*	*	*	*	*
882 . . . . .	*7	*6	*	*	*5	*	*	*	*	*	*
883 . . . . .	12	*8	*	*	9	*	*	*	*	*	*
886 . . . . .	*6	*6	*	*	*	*	*	*	*	*	*
891 . . . . .	13	10	*	*	*7	*	*	*	*	*	*
891.0 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*
892 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*
892.1 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
920 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
922 . . . . .	19	*6	13	*	*6	*	9	*	*6	*	*
922.1 . . . . .	9	*	*7	*	*	*	*	*	*	*	*
924 . . . . .	21	*7	13	*	*	*	14	*	*	*8	*5
924.0 . . . . .	11	*	*8	*	*	*	10	*	*	*	*
924.01 . . . . .	10	*	*7	—	*	*	10	*	*	*	*
924.8 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
935 . . . . .	10	*7	*	*	*	*	*	*	*	*	*
935.1 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
941 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
942 . . . . .	*6	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
945	11	*9	*	*	*6	*	*	*	*	*	*
945.2	*6	*	*	*	*	*	*	*	*	*	*
946	13	*8	*5	*	*7	*	*	*	*	*7	*
946.2	*7	*	*	*	*	*	—	*	*	*	*
952	*5	*	*	*	*	*	*	*	*	*	*
958	*8	*6	*	*	*	*	*	*	*	*	*
959	19	9	10	*	13	*	*	*	*	*8	*
959.1	9	*	*6	*	*6	*	*	*	*	*	*
963	*5	*	*	*	*	—	*	*	*	*	*
965	47	18	28	*6	34	*	*	11	12	15	*9
965.0	*8	*6	*	*	*6	*	*	*	*	*	*
965.1	*8	*	*7	*	*	*	*	*	*	*	*
965.4	21	*8	13	*	17	*	*	*	*5	*7	*
965.6	*6	*	*	*	*	*	*	*	*	*	*
966	11	*	*6	*	*	*	*	*	*	*	*
966.3	*6	*	*	*	*	*	*	*	*	*	*
967	*7	*	*	*	*	*	*	*	*	*	*
968	*8	*6	*	*	*7	*	—	*	*	*	*
968.5	*5	*	*	—	*	*	—	*	*	*	*
969	59	22	37	*7	37	12	*	11	13	25	11
969.0	29	10	19	*	19	*	*	*5	*6	14	*
969.4	19	*8	11	*	11	*	*	*	*	*8	*
970	*5	*	*	—	*5	—	—	*	*	*	—
970.8	*5	*	*	—	*5	—	—	*	*	*	—
972	*9	*	*5	*	*	*	*	*	*	*	*
977	15	*	10	*	11	*	*	*	*	*9	*
977.8	*5	*	*	*	*	*	*	*	*	*	*
977.9	*8	*	*	*	*7	—	*	*	*	*6	*
989	*8	*	*	*	*	*	*	*	*	*	*
989.5	*6	*	*	*	*	*	*	*	*	*	*
992	*6	*	*	*	*	*	*	*	*	*	*
995	23	*8	15	*5	*6	*	*7	*	*8	*7	*
995.1	*6	*	*	*	*	*	*	*	*	*	*
995.2	*9	*	*6	*	*	*	*	*	*	*	*
996	365	171	194	22	76	105	162	92	67	132	75
996.0	25	12	13	*	*	*	18	*8	*	*6	*6
996.01	18	*8	11	*	*	*	14	*	*	*	*5
996.1	13	*6	*6	*	*	*	*7	*	*	*	*
996.2	16	10	*6	*8	*6	*	*	*	*	*5	*
996.4	74	30	44	*	12	19	42	19	13	26	16
996.5	13	*5	*7	*	*	*	*	*	*	*5	*
996.59	9	*	*6	*	*	*	*	*	*	*	*
996.6	87	40	48	*8	21	24	34	23	21	30	13
996.62	39	17	23	*	12	11	12	12	10	12	*7
996.64	*6	*	*	*	*	*	*	*	*	*	*
996.66	13	*6	*7	*	*	*	*8	*	*	*5	*
996.67	*7	*	*	*	*	*	*	*	*	*	*
996.69	14	*6	*7	*	*	*5	*	*	*	*	*
996.7	113	53	60	*	21	38	52	26	17	43	27
996.72	*6	*	*	*	*	*	*	*	*	*	*
996.73	50	25	25	*	10	19	21	11	*6	20	14
996.74	30	15	15	*	*	10	17	*7	*	13	*7
996.77	*7	*	*	—	*	*	*	*	*	*	*
996.78	9	*	*5	*	*5	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 1. Number of first-listed diagnoses for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of first-listed diagnoses in thousands											
996.8	20	12	*8	*	10	*8	*	*	*	11	*
996.81	14	*8	*6	*	*8	*5	*	*	*	9	*
997	94	42	52	*	19	26	45	23	22	24	25
997.1	10	*	*6	*	*	*	*	*	*	*	*
997.2	*8	*	*6	–	*	*	*	*	*	*	*
997.3	11	*7	*	*	*	*	*	*	*	*	*
997.4	36	12	24	*	*8	11	15	*8	10	*8	10
997.5	11	*8	*	*	*	*	*6	*	*	*	*
997.6	12	*7	*5	*	*	*	*7	*	*	*	*
997.62	*8	*	*	*	*	*	*5	*	*	*	*
998	172	74	98	10	49	48	65	42	34	59	37
998.1	46	22	24	*	14	14	15	14	*8	14	11
998.2	*7	*	*5	–	*	*	*	*	*	*	*
998.3	*9	*	*6	*	*	*	*	*	*	*	*
998.5	89	38	51	*	26	22	38	21	18	32	18
998.8	16	*7	*9	*	*	*	*	*	*	*	*
998.89	14	*7	*7	*	*	*	*	*	*	*	*
999	*7	*	*	*	*	*	*	*	*	*	*
V01–V82	4,183	165	4,017	74	3,852	93	164	748	964	1,449	1,022
V10	*5	*	*	*	*	*	*	*	*	*	*
V12	*5	*	*	*	–	*	*	*	*	*	*
V22	22	–	22	–	22	–	–	*	*7	*8	*6
V22.2	17	–	17	–	17	–	–	*	*6	*7	*
V24	*5	–	*5	*	*5	–	–	*	*	*	*
V24.0	*5	–	*5	*	*5	–	–	*	*	*	*
V25	*6	*	*6	–	*6	–	–	*	*	*	*
V25.2	*6	*	*6	–	*6	–	–	*	*	*	*
V27	3,766	–	3,766	12	3,752	*	–	633	821	1,359	953
V27.0	2,844	–	2,844	10	2,832	*	–	473	596	1,059	716
V27.1	22	–	22	*	22	–	–	*	*	12	*
V27.2	37	–	37	–	37	*	–	*	*8	15	10
V27.9	861	–	861	*	858	*	–	150	214	272	225
V30	16	*8	*8	16	–	–	*	*7	*	*5	*
V30.1	16	*8	*8	16	–	–	–	*7	*	*5	*
V45	*8	*	*	*	*	*	*	*	*	*	*
V45.8	*6	*	*	*	*	*	*	*	*	*	*
V54	*7	*	*	*	*	*	*	*	*	*	*
V54.0	*6	*	*	*	*	*	*	*	*	*	*
V55	14	*7	*7	*	*	*	*	*6	*	*	*
V55.3	11	*	*6	*	*	*	*	*	*	*	*
V57	132	51	80	*	*8	22	101	20	72	19	22
V57.1	34	14	20	*	*	*5	26	12	9	*9	*
V57.8	97	37	59	*	*6	17	74	*7	62	10	17
V57.89	97	37	59	*	*6	17	74	*7	62	10	17
V58	117	51	66	20	20	41	36	49	37	21	10
V58.0	*5	*	*	*	*	*	*	*	*	*	*
V58.1	106	46	60	19	17	38	32	45	34	20	*8
V70	*7	*	*	*	*	*	*	*	*	*	*
V70.7	*7	*	*	*	*	*	*	*	*	*	*
V71	20	9	10	*6	*9	*	*	*7	*6	*	*
V71.8	11	*	*7	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.

– Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
All codes . . . . .	164,627	71,340	93,287	10,715	40,825	34,207	78,880	44,481	36,382	59,798	23,965
001-139 . . . . .	6,089	3,128	2,961	706	1,661	1,189	2,532	1,777	1,037	2,404	871
003 . . . . .	57	*27	*29	*22	*	*	*	*	*	*	*
003.0 . . . . .	44	*23	*	*	*	*	*	*	*	*	*
008 . . . . .	453	154	299	123	53	45	232	123	98	138	94
008.4 . . . . .	216	*56	159	*	*	*	161	*80	*46	56	*34
008.45 . . . . .	197	*49	148	*	*	*	150	*72	*42	53	*
008.6 . . . . .	67	35	31	64	*	*	*	*	*19	26	*18
008.61 . . . . .	61	34	28	61	*	*	-	*	*18	*25	*17
008.8 . . . . .	160	54	106	44	31	*19	67	39	32	53	36
009 . . . . .	34	*17	*17	*	*	*	*	*	*	*	*
009.0 . . . . .	*15	*	*	-	*	*	*	*	*	*	*
011 . . . . .	122	*82	*	*	*	*	*	*	*	*59	*
011.9 . . . . .	*83	*53	*	*	*	*	*	*	*	*51	*
034 . . . . .	29	*14	*15	*16	*	*	*	*	*	*13	*
034.0 . . . . .	28	*14	*14	*15	*	*	*	*	*	*	*
038 . . . . .	2,678	1,146	1,533	103	203	514	1,858	642	537	1,199	300
038.0 . . . . .	236	100	136	*	*	*	156	*65	*48	98	*
038.1 . . . . .	523	248	274	*	*	*125	345	147	119	209	*49
038.2 . . . . .	*75	*	*	*	*	*	*	*	*	*	*
038.4 . . . . .	769	284	485	*	*54	137	556	179	170	303	117
038.40 . . . . .	*68	*	*	*	*	*	*	*	*	*	*
038.42 . . . . .	388	107	280	*	*	79	279	91	105	133	59
038.43 . . . . .	*79	*	*	*	*	*	*	*	*	*	*
038.49 . . . . .	227	84	143	*	*	*	186	*	*	118	*
038.8 . . . . .	114	*	*80	*	*	*	83	*	*	*51	*
038.9 . . . . .	941	433	508	46	73	162	660	180	165	510	86
041 . . . . .	109	66	*43	*	*	*	*43	*	*	*55	*
042 . . . . .	1,233	892	341	*43	916	238	*	569	86	401	177
047 . . . . .	132	65	67	60	49	*	*	*26	*	64	*28
047.9 . . . . .	120	63	57	57	48	*	*	*23	*	55	*27
052 . . . . .	44	26	*	*27	*	*	*	*	*	*	*
052.9 . . . . .	*21	*13	*	*14	*	*	*	*	*	*	*
053 . . . . .	109	*38	71	*	*	*25	76	*48	*	*37	*
053.9 . . . . .	70	*	*50	*	*	*	*47	*	*	*	*
054 . . . . .	*35	*	*	*	*	*	*	*	*	*	*
070 . . . . .	60	*33	*27	*	32	*	*	*	*	*18	*
070.5 . . . . .	*18	*	*	-	*	*	*	*	*	*	*
075 . . . . .	*24	*	*	*	*17	*	*	*	*	*	*
078 . . . . .	49	*25	*	*	*	*	*	*	*	*	*
079 . . . . .	221	101	120	145	41	*20	*	68	43	75	34
079.9 . . . . .	196	88	108	122	39	*20	*	60	36	72	28
079.99 . . . . .	195	87	108	121	39	*20	*	59	36	72	28
112 . . . . .	140	*63	*77	*	*	*	*54	*	*	*67	*
112.4 . . . . .	*49	*	*	-	*	*	*	*	*	*	*
117 . . . . .	*82	*	*	*	*	*	*	*	*	*	*
135 . . . . .	*48	*	*32	*	*	*	*	*	*	*	*
136 . . . . .	138	96	*	*	*50	*	*	*	*	*	*
136.3 . . . . .	115	*85	*	*	*40	*	*	*	*	*	*
140-239 . . . . .	11,833	5,163	6,670	246	1,502	3,493	6,593	3,440	2,269	4,166	1,959
150 . . . . .	106	72	*	-	*	*37	*62	*	*	*35	*
150.9 . . . . .	*24	*	*	-	*	*	*	*	*	*	*
151 . . . . .	349	195	155	-	*	*95	217	*109	*88	102	*51

\*Figure does not meet standard of reliability or precision.  
-Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
151.0	*97	*	*	—	*	*	*	*	*	*	*
151.8	*111	*	*	—	*	*	*	*	*	*	*
153	978	442	536	—	*	122	837	319	273	289	97
153.1	*97	*	*	—	*	*	*81	*	*	*	*
153.2	*73	*	*	—	*	*	*62	*	*	*	*
153.3	260	110	150	—	*	*	210	*70	*75	*85	*
153.4	226	*86	*140	—	*	*	205	*	*	*	*
153.6	138	*	*81	—	*	*	120	*	*45	*	*
153.9	73	*	*42	—	*	*	*61	*	*	*	*
154	434	214	220	—	*	102	306	165	83	110	*76
154.0	155	*74	*81	—	*	*	121	*	*	*63	*
154.1	240	124	115	—	*	*67	160	103	*55	*	*
155	84	*57	*	*	*	*	*44	*	*	*42	*
155.0	*65	*46	*	*	*	*	*31	*	*	*	*
156	*92	*	*	—	*	*	*57	*	*	*	*
157	293	126	167	—	*	67	220	*112	*	91	*55
157.0	140	*66	*74	—	*	*	113	*	*	*51	*
157.8	*81	*	*	—	*	*	*	*	*	*	*
157.9	*41	*	*	—	*	*	*	*	*	*	*
158	*70	*	*53	*	*	*	*	*	*	*	*
161	110	*72	*	—	*	*68	*	*	*	*	*
162	1,254	698	556	*	*51	407	796	405	229	481	139
162.2	86	*31	*54	—	*	*	*67	*	*	*	*
162.3	424	204	221	—	*	149	263	124	98	160	*42
162.5	202	128	*74	—	*	*71	129	*77	*	*61	*
162.8	161	113	*48	*	*	*54	104	107	*	*	*
162.9	352	202	151	—	*30	114	209	49	56	190	58
170	*51	*	*	*	*	*	*	*	*	*	*
171	51	*28	*	*	*	*	*	*	*	*	*
172	*23	*	*	—	*	*	*	*	*	*	*
173	81	*32	*49	*	—	*	51	*	*	*43	*
174	486	—	486	—	98	201	187	132	108	174	73
174.1	*12	—	*12	—	*	*	*	*	*	*	*
174.2	*14	—	*14	—	*	*	*	*	*	*	*
174.4	75	—	75	—	*	26	28	*19	*24	*22	*
174.5	*13	—	*13	—	*	*	*	*	*	*	*
174.8	140	—	140	—	*	56	72	63	*18	*26	33
174.9	212	—	212	—	*58	101	54	*29	54	108	*21
180	80	—	80	—	*28	*41	*	*34	*	35	*
180.8	*34	—	*34	—	*	*	*	*	*	*	*
180.9	35	—	35	—	*22	*	*	*	*	*27	*
182	181	—	181	—	*	74	94	*34	*32	80	35
182.0	178	—	178	—	*	74	91	*32	*31	80	34
183	173	—	173	*	*	70	87	39	*45	58	*32
183.0	173	—	173	*	*	70	87	38	*45	58	*32
185	480	480	—	*	*	163	315	128	109	159	84
188	316	222	94	—	*	61	249	121	74	100	*22
188.8	118	89	*29	—	*	*	96	*77	*	*	*
188.9	100	79	*20	—	*	*30	66	*	*	*39	*
189	267	129	138	*	*	60	160	*50	*51	119	*46
189.0	224	106	118	*	*	52	126	*42	*39	100	*43

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
191	242	160	82	*	*50	65	*99	*71	*	94	*34
191.1	*50	*	*	*	*	*	*	*	*	*	*
191.3	*57	*	*	*	*	*	*	*	*	*	*
193	25	*	*15	*	*	*	*	*	*	*	*
194	*33	*	*	*	*	*	—	*	*	*	*
196	58	*40	*	—	*	*	*34	*	*	*	*
197	996	328	668	*	70	297	624	293	152	397	154
197.0	195	97	98	*	*	64	108	*61	*41	61	*
197.2	226	84	143	—	*	*59	163	*82	*	*68	*42
197.6	155	*	128	—	*	*	108	*	*	*67	*
197.7	280	84	196	*	*	89	171	67	*34	147	*33
198	941	405	536	*	*56	336	534	291	174	283	193
198.3	300	146	155	*	*	122	155	*79	*44	93	*85
198.5	436	181	255	*	*	151	264	157	*58	150	*
198.8	147	*56	91	*	*	*38	86	*	*54	*	*29
198.89	129	*55	74	*	*	*34	72	*	*47	*	*26
199	147	*61	86	—	*	*51	90	*	*	*71	*54
199.0	82	*	*38	—	*	*	*66	*	*	*	*
199.1	65	*	*48	—	*	*	*	*	*	*	*
200	157	86	*71	*	*39	*	76	*49	*	63	*
200.0	89	*54	*	*	*	*	*52	*	*	*46	*
200.00	*49	*38	*	*	*	*	*33	*	*	*31	*
201	*42	*	*	*	*	*	*	*	*	*	*
201.9	*32	*	*	*	*	*	*	*	*	*	*
202	344	176	168	*	*64	115	163	*82	*47	142	*73
202.8	305	162	143	*	*60	92	151	*68	*43	125	*69
202.80	125	*68	*57	*	*	*	*49	*	*	*	*
202.81	*37	*	*	*	*	*	*	*	*	*	*
202.83	*60	*	*	—	*	*	*	*	*	*	*
202.88	*53	*43	*	—	*	*	*36	*	*	*31	*
203	261	109	152	—	*	*60	198	*	*	119	*
203.0	254	108	146	—	*	*59	193	*	*	118	*
203.00	254	108	146	—	*	*59	193	*	*	118	*
204	201	112	90	62	*	*	*52	*46	*	*55	*80
204.0	149	*87	*62	62	*	*	*	*	*	*	*72
204.00	107	*68	*	*34	*	*	*	*	*	*	*
204.1	*51	*	*	—	*	*	*45	*	*	*	*
204.10	*51	*	*	—	*	*	*44	*	*	*	*
205	279	*152	*127	*	*	*	159	*137	*	*	*
205.0	227	*135	*92	*	*	*	*151	*122	*	*	*
205.00	209	*128	*	*	*	*	*150	*112	*	*	*
208	*63	*	*	—	*	*	*	*	*	*	*
208.0	*54	*	*	—	*	*	*	*	*	*	*
210	*9	*	*	*	*	*	*	*	*	*	*
210.2	*9	*	*	*	*	*	*	*	*	*	*
211	188	87	101	*	*24	41	122	55	*33	64	*35
211.3	136	65	71	*	*	28	92	44	*	*55	*
214	*23	*	*	—	*	*	*	*	*	*	*
218	600	—	600	—	351	231	*	132	121	230	117
218.0	93	—	93	—	53	36	*	*35	*17	*24	*17
218.1	158	—	158	—	87	67	*	52	31	53	*22
218.2	73	—	73	—	46	*26	*	*	*	36	*
218.9	276	—	276	—	165	101	*	31	63	118	64
220	117	—	117	*	57	*23	*36	*37	*27	36	*17

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
225 . . . . .	118	*	83	*	*	*52	*41	*	*	*53	*
225.2 . . . . .	77	*	*56	*	*	*	*	*	*	*	*
226 . . . . .	15	*	*10	—	*	*	*	*	*	*	*
227 . . . . .	70	*	50	*	*17	*23	*	*	*	*18	*
227.1 . . . . .	*15	*	*	—	*	*	*	*	*	*	*
227.3 . . . . .	*49	*	*	*	*	*	*	*	*	*	*
228 . . . . .	*26	*	*	*	*	*	*	*	*	*	*
233 . . . . .	76	*	72	—	35	30	*	*17	*20	28	*12
233.0 . . . . .	26	*	25	—	*	*13	*	*	*	*	*
233.1 . . . . .	43	—	43	—	31	*	*	*	*	*19	*
235 . . . . .	*76	*	*	*	*	*	*59	*	*	*	*
235.2 . . . . .	*58	*	*	*	*	*	*	*	*	*	*
236 . . . . .	*28	*	*25	*	*	*	*	*	*	*	*
237 . . . . .	*38	*	*	*	*	*	*	*	*	*	*
237.5 . . . . .	*33	*	*	*	*	*	*	*	*	*	*
238 . . . . .	84	*54	*31	*	*	*	*47	*	*	*	*
238.7 . . . . .	66	*48	*	*	*	*	*40	*	*	*	*
239 . . . . .	55	*	*31	*	*	*	*	*	*	*	*
240-279 . . . . .	6,823	2,759	4,064	430	1,076	1,699	3,619	1,845	1,372	2,623	983
241 . . . . .	22	*	19	*	*	*	*	*	*	*	*
241.1 . . . . .	*12	*	*11	—	*	*	*	*	*	*	*
242 . . . . .	55	*	44	*	*22	*	*23	*	*	*	*
242.0 . . . . .	*20	*	*19	*	*14	*	*	*	*	*	*
242.00 . . . . .	*17	*	*16	—	*	*	*	*	*	*	*
242.9 . . . . .	*27	*	*	*	*	*	*	*	*	*	*
242.90 . . . . .	*24	*	*	*	*	*	*	*	*	*	*
244 . . . . .	63	*	55	—	*	*	*40	*26	*	*	*
244.9 . . . . .	59	*	52	—	*	*	*38	*25	*	*	*
250 . . . . .	3,072	1,460	1,613	76	518	1,082	1,397	873	521	1,208	471
250.0 . . . . .	731	295	437	*31	98	257	345	215	124	316	76
250.00 . . . . .	177	79	98	*	*17	67	93	*36	*21	81	38
250.01 . . . . .	174	95	79	*	*	43	99	84	*28	54	*
250.02 . . . . .	184	58	127	*	*20	81	82	56	36	82	*
250.03 . . . . .	196	63	133	*	43	66	71	*39	*38	99	*
250.1 . . . . .	403	192	212	34	209	94	*66	112	92	133	66
250.10 . . . . .	*43	*30	*	*	*	*	*	*	*	*	*
250.11 . . . . .	140	67	73	*	84	*	*	*44	*27	54	*
250.13 . . . . .	196	84	112	*	105	42	*	55	44	57	40
250.2 . . . . .	94	*36	*57	—	*	*	*56	*	*	*	*
250.3 . . . . .	*30	*	*	*	*	*	*	*	*	*	*
250.4 . . . . .	193	86	107	—	*	67	77	*	*	96	*38
250.40 . . . . .	*62	*	*	—	*	*	*	*	*	*	*
250.41 . . . . .	*47	*	*	—	*	*25	*	*	*	*	*
250.42 . . . . .	*48	*	*	—	*	*	*	*	*	*	*
250.5 . . . . .	*26	*	*	—	*	*	*	*	*	*	*
250.6 . . . . .	188	70	118	*	*29	89	69	*44	*46	73	*25
250.60 . . . . .	*52	*	*	—	*	*	*	*	*	*	*
250.61 . . . . .	68	*	*53	—	*	*37	*	*	*	*36	*
250.63 . . . . .	*43	*	*	*	*	*	*	*	*	*	*
250.7 . . . . .	712	424	288	—	*	259	395	204	102	238	168
250.70 . . . . .	247	138	109	—	*	*	143	*	*	*77	*66
250.71 . . . . .	294	167	127	—	*	*107	178	*106	*	*90	*53
250.72 . . . . .	*48	*	*	—	*	*	*	*	*	*	*
250.73 . . . . .	123	*90	*	—	*	*	*	*	*	*61	*
250.8 . . . . .	611	315	297	*	50	239	320	188	92	269	63

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).



**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
250.80	131	85	*46	—	*	*38	86	*41	*	*49	*
250.81	239	132	107	*	*	104	109	*54	*35	118	*33
250.82	110	*	*79	—	*	*	*79	*	*	*51	*
250.83	131	*67	*64	*	*	*70	*45	*54	*	*50	*
250.9	84	*28	56	*	*	*32	*36	*	*	38	*
250.92	*30	*	*	*	*	*	*	*	*	*	*
250.93	*37	*	*	*	*	*	*	*	*	*24	—
251	51	*26	*24	*	*	*	*26	*	*	*	*
251.2	*30	*16	*	*	*	*	*	*	*	*	*
253	75	*36	*38	*	*	*	*59	*	*	*	*
253.6	64	*	*32	*	*	*	*59	*	*	*	*
255	*32	*	*22	*	*	*	*	*	*	*	*
263	98	*40	*58	*	*	*	75	*	*	*38	*
263.9	82	*32	*50	*	*	*	65	*	*	*36	*
272	63	28	35	*	*15	29	*19	*12	*	32	*
272.0	38	21	*16	—	*	*20	*	*	*	*17	*
272.4	*19	*	*15	—	*	*	*	*	*	*	*
274	67	43	*23	—	*	*	41	*37	*	*	*
274.0	46	*26	*	—	*	*	*35	*	*	*	*
275	101	*37	*63	*	*	*28	*43	*	*	*44	*
275.4	85	*25	*61	*	*	*	*42	*	*	*43	*
276	2,715	924	1,791	255	223	424	1,812	651	658	1,041	364
276.0	*54	*	*40	*	*	*	*	*	*	*	*
276.1	292	73	219	*	*	*38	230	*40	89	117	47
276.2	*31	*	*	*	*	*	*	*	*	*	*
276.5	1,996	711	1,285	226	151	278	1,340	515	460	766	255
276.6	*37	*	*	*	*	*	*	*	*	*	*
276.7	65	*35	*31	*	*	*18	44	*	*	*19	*
276.8	196	52	144	*	29	54	113	*45	43	79	29
276.9	*36	*	*25	*	*	*	*	*	*	*	*
277	159	*51	*107	*	*82	*	*	*59	*	*	*
277.0	*96	*	*	*	*	*	—	*	*	*	—
277.00	*96	*	*	*	*	*	—	*	*	*	—
278	83	*24	59	*	59	*22	*	*	*	*28	*30
278.0	79	*24	55	*	55	*22	*	*	*	*27	*30
280-289	1,674	658	1,016	201	530	308	635	472	356	639	208
280	171	39	133	*	*27	*22	118	*50	48	67	*
280.0	78	*	62	*	*19	*	52	*	*	*28	*
280.9	90	*22	68	*	*	*	64	*	*33	*37	*
281	*38	*	*34	*	*	*	*	*	*	*	*
282	380	176	204	76	290	*	*	89	63	166	62
282.6	363	165	197	66	286	*	*	85	60	162	55
282.60	*20	*	*	*	*	*	—	*	*	*	*
282.62	332	150	182	53	269	*	*	77	54	153	*48
284	136	54	82	*	*41	*	64	*51	*31	*40	*
284.8	120	45	74	*	*	*	64	*42	*29	*36	*
285	294	102	192	*	41	59	188	54	59	159	*22
285.1	102	*28	75	*	*	*	66	*	*29	*48	*
285.9	180	69	111	*	30	*34	113	*28	*26	110	*15
286	66	44	*22	*	*	*	*30	*29	*	*	*
286.7	*22	*	*	—	*	*	*	*	*	*	*
287	159	68	91	*21	46	*37	56	48	*39	49	*23
287.3	53	*	*35	*	*24	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.  
 — Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
287.4	*33	*	*	*	*	*	*	*	*	*	*
287.5	61	*22	*39	*	*	*	*32	*	*	*	*
288	288	107	181	*38	42	115	93	85	61	98	*44
288.0	257	94	163	*35	*35	105	82	78	56	91	*32
288.8	*27	*	*	*	*	*	*	*	*	*	*
289	97	53	45	27	*21	*	*	*23	*32	*20	*23
289.2	23	*13	*	*	*	*	—	*	*	*	*
290-319	17,856	8,731	9,125	1,088	9,803	3,483	3,482	6,588	4,423	5,608	1,237
290	652	231	421	*	*	*	629	189	134	286	*43
290.0	180	89	91	—	*	*	177	*	*	93	*
290.2	102	*	*74	—	—	*	99	*	*	*	*
290.20	*73	*	*	—	—	*	*72	*	*	*	*
290.3	93	*38	*55	*	—	*	93	*	*	*	*
290.4	194	*64	130	—	—	*	188	*	*	*60	*
290.40	*85	*	*60	—	—	*	*81	*	*	*	*
291	447	338	109	*	239	168	*39	91	200	105	51
291.0	123	100	*	*	46	*65	*	*	*34	*44	*
291.8	300	225	75	*	180	97	*	59	162	57	*23
292	247	131	116	*	148	*	71	100	*64	67	*
292.0	90	*62	*27	*	71	*	*	*33	*	*	*
292.1	*43	*	*	*	*	—	*	*	*	*	*
292.8	107	*45	*62	*	*39	*	*53	*	*	*	*
292.81	*51	*	*36	—	*	*	*38	*	*	*	*
292.84	*42	*	*	*	*	*	*	*	*	*	*
293	314	132	182	—	90	73	151	171	*	95	*
293.0	73	*	48	—	*	*	62	*	*	*	*
293.8	222	97	125	—	*78	*64	*81	141	*	*56	*
293.81	*86	*	*62	—	*	*	*	*	*	*	*
293.83	128	*70	*58	—	*69	*	*	94	*	*	*
294	222	94	128	*	*	*	175	*55	*79	*60	*
294.8	156	*61	96	—	*	*	135	*	*66	*	*
295	3,640	2,032	1,608	*	2,253	1,028	349	1,804	805	815	217
295.3	1,097	578	519	—	627	311	159	599	116	323	*59
295.30	131	*53	*78	—	*64	*	*	*	*	*	*
295.32	210	*108	*	—	*125	*	*	*108	*	*	*
295.34	722	401	321	—	423	231	*	438	*65	183	*
295.4	*93	*	*	*	*87	*	*	*	*	*	*
295.6	369	259	110	*	168	166	*	170	*97	*86	*
295.62	*139	*114	*	—	*	*	*	*	*	*	*
295.64	165	*94	*71	—	*89	*65	*	*69	*	*	*
295.7	1,521	781	741	*	1,033	369	*113	643	451	306	*121
295.70	788	446	342	*	548	198	*	219	272	204	*93
295.74	716	325	390	*	471	171	*	419	171	*98	*
295.9	484	318	165	*	288	161	*	309	*110	*	*
295.90	171	*70	*101	*	*101	*	*	*	*99	*	*
295.94	288	228	*60	—	175	*90	*	262	*	*	*
296	5,957	2,110	3,847	391	3,029	1,115	1,422	1,939	1,453	2,125	441
296.2	1,873	732	1,140	212	937	306	417	455	458	846	114
296.20	857	335	523	*85	451	147	174	243	292	286	*36
296.22	161	*67	94	*	101	*	*	*	*	134	*
296.23	483	166	317	*	234	*82	*111	85	84	294	*
296.24	329	162	167	*	142	*65	*	*112	*73	99	*
296.3	2,195	701	1,494	*91	1,080	346	679	796	556	710	133
296.30	639	183	456	*	265	112	238	254	198	175	*
296.32	109	*	82	*	67	*	*	*	*	*53	*
296.33	922	306	615	*	544	136	193	347	202	317	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
296.34	452	146	307	*	191	*82	168	158	128	125	*
296.4	570	247	323	*	307	134	*122	311	100	106	*53
296.40	244	94	150	*	121	*61	*	*91	*67	*65	*
296.44	267	*122	144	—	*148	*	*	*187	*	*	*
296.5	361	103	258	*	150	*111	*89	*82	*86	143	*
296.50	176	*	137	*	75	*	*	*	*64	*67	*
296.53	111	*	*77	*	*	*	*	*	*	*	*
296.54	*61	*	*	*	*	*	*	*	*	*	*
296.6	482	157	325	*	278	131	*	139	103	191	*
296.60	262	*103	158	*	162	*	*	*	*81	127	*
296.63	*70	*	*	*	*	*	*	*	*	*	*
296.64	115	*	*93	—	*60	*	*	*	*	*	*
296.7	279	96	183	*	179	*49	*	*84	101	*69	*
296.8	113	*52	*61	*	*54	*	*	*51	*	*	*
296.80	*42	*	*	*	*	*	*	*	*	*	*
296.82	*47	*	*	*	*	*	*	*	*	*	*
297	113	*	*79	*	*	*	*51	*	*	*	*
297.1	*46	*	*	—	*	*	*	*	*	*	*
297.9	*63	*	*	*	*	*	*	*	*	*	*
298	814	330	484	*	542	91	173	417	86	260	*52
298.9	756	289	467	*	514	*66	168	392	76	237	*51
300	450	176	274	*54	207	94	95	112	115	170	54
300.0	101	40	62	*	31	*31	37	*25	*	51	*
300.00	54	*20	34	*	*16	*14	*22	*15	*	23	*
300.01	*25	*	*18	*	*13	*	*	*	*	*	*
300.1	58	*	*47	*	*36	*	*	*	*	*	*
300.11	*31	*	*22	*	*	*	*	*	*	*	*
300.4	158	63	94	*45	70	*	*	55	*50	*35	*
300.9	80	*	*47	*	*28	*	*	*	*	*45	*
301	127	67	*60	*	104	*	*	84	*	*	*
301.8	80	*39	*41	*	78	*	*	*70	*	*	*
301.83	*40	*	*	*	*39	—	*	*	*	*	*
303	1,792	1,284	508	*	1,186	506	92	442	633	608	109
303.0	195	146	49	—	112	68	*	51	63	51	*29
303.00	37	*25	*	—	*22	*	*	*	*	*	*
303.01	154	118	*37	—	87	60	*	48	*55	*30	*
303.9	1,597	1,138	459	*	1,074	438	*76	391	570	557	80
303.90	430	303	127	—	288	106	*	88	98	214	*30
303.91	1,150	825	325	*	780	325	*	302	465	338	*45
304	934	637	297	*	791	101	*	329	206	373	*
304.0	179	129	50	—	143	*	*	105	*	*48	*
304.00	*59	*	*	—	*	*	*	*	*	*	*
304.01	119	79	40	—	99	*	—	99	*	*	—
304.2	322	226	96	—	295	*	*	133	*52	134	*
304.20	98	*75	*	—	94	*	*	*	*	83	*
304.21	220	148	71	—	197	*	—	124	*	*49	*
304.3	*109	*71	*	*	*76	—	—	*	*	*	*
304.31	*76	*57	*	*	*62	—	—	*	*	*	—
304.7	115	68	*	—	100	*	—	62	*	*	*
304.71	68	*46	*	—	53	*	—	*60	*	*	*
304.8	96	83	*	—	75	*	—	*	*	*64	*
304.81	*64	*58	*	—	*44	*	—	*	*	*47	*
304.9	*59	*	*	*	*49	*	*	*	*	*42	*
304.90	*47	*	*	*	*	*	*	*	*	*	—
305	173	130	43	*	112	47	*	33	53	71	*16
305.0	60	49	*12	*	36	*	*	*	*18	*28	*
305.00	45	34	*11	*	25	*	*	*	*11	*25	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
305.1	*15	*9	*	—	*	*	*	*	*	*	*
305.6	*34	*26	*	—	*28	*	—	*	*	*	*
305.60	*27	*	*	—	*21	*	—	*	*	*	*
305.9	*32	*	*	*	*24	*	*	*	*	*	*
305.90	*26	*	*	*	*21	*	*	*	*	*	*
306	*24	*	*	*	*	*	*	*	*	*	*
307	120	*	*109	*	*85	*	*	*	*	*	*
308	*49	*	*44	*	*	*	*	*	*	*	*
309	633	311	321	119	413	59	*	325	128	130	50
309.0	232	113	119	*	168	*	*	112	*28	75	*18
309.2	59	*	*42	*	43	*	*	*	*	*	*
309.28	52	*	*38	*	*37	*	*	*	*	*	*
309.4	143	88	55	*	99	*	*	85	*30	*	*
309.8	140	*	*74	*	*63	*	*	*81	*	*	*
309.81	126	*	*73	*	*61	*	—	*	*	*	*
309.9	*23	*	*	*	*17	*	*	*	*	*	*
310	128	*75	*53	*	*	*	*64	*	*	*	*
310.9	*44	*	*	—	*	*	*	*	*	*	*
311	498	206	293	*93	300	*36	*70	164	162	142	*30
312	191	146	*44	*	125	*	*	101	*44	*	*
312.3	119	86	*33	*	96	*	*	*56	*	*	*
312.30	*49	*37	*	*	*44	*	—	*	*	*	—
312.34	*45	*	*	*	*	*	*	*	*	*	*
313	*72	*45	*	*46	*	—	—	*	*	*	*
313.8	*72	*45	*	*46	*	—	—	*	*	*	*
313.81	*72	*45	*	*46	*	—	—	*	*	*	*
314	155	128	*	141	*	—	—	*	*	*81	*
314.0	153	127	*	141	*	—	—	*	*	*79	*
314.01	144	117	*	132	*	—	—	*	*	*76	*
320-389	2,869	1,311	1,559	384	656	620	1,210	722	602	1,082	463
320	120	*55	*64	67	*	*	*	*	*	63	*
320.9	*37	*	*	*	*	—	*	*	*	*	*
322	*63	*	*	*	*	*	*	*	*	*	*
322.9	*60	*	*	*	*	*	*	*	*	*	*
323	*50	*	*31	*	*35	*	*	*	*	*36	*
323.9	*48	*	*30	*	*35	*	*	*	*	*35	*
331	144	74	70	*	*	*	113	*43	*	46	*
331.0	55	*	*	—	—	*	53	*	*	*	*
331.3	*35	*	*	*	*	*	*	*	*	*	*
331.4	*44	*	*	*	*	*	*	*	*	*	*
332	136	*62	*74	—	*	*	128	*	*64	*	*
332.0	133	*62	*71	—	*	*	126	*	*62	*	*
333	66	*	*29	*	*	*	*	*	*	*	*
337	*37	*	*28	*	*	*	*	*	*	*	*
337.2	*26	*	*	*	*	*	*	*	*	*	*
340	127	*	101	*	62	57	*	*	*37	*52	*
342	411	230	181	*	*	*112	257	*	*	210	*
342.9	403	222	181	*	*	*112	253	*	*	209	*
342.90	296	164	*132	*	*	*70	190	*	*	162	*
344	122	*63	*59	*	*46	*	*	*	*	*	*
345	281	119	162	83	87	*31	80	66	69	113	34
345.1	47	*23	*24	*	*	*	*	*	*17	*	*
345.10	36	*19	*	*	*	*	*	*	*	*	*
345.3	73	*31	*43	*31	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
345.5	*28	*	*	*	*	*	*	*	*	*	*
345.9	71	*32	*39	*15	*29	*	*	*	*	*30	*
345.90	50	*23	*26	*	*16	*	*	*	*	*23	*
346	125	*17	108	*	70	37	*	*22	39	50	*
346.9	50	*	44	*	28	*18	*	*	*16	*22	*
346.90	32	*	25	*	*21	*	*	*	*	*12	*
346.91	*19	*	*18	*	*	*	*	*	*	*	*
348	172	95	77	*31	*25	*54	61	*36	*33	*55	*48
348.1	*61	*	*	*	*	*	*	*	*	*	*
348.3	*54	*	*	*	*	*	*	*	*	*	*
349	48	*	*30	*	*19	*	*	*	*	*	*
349.8	*25	*	*	*	*	*	*	*	*	*	*
351	*27	*	*	*	*	*	*	*	*	*	*
353	*24	*	*	*	*	*	*	*	*	*	*
354	29	*	*20	*	*18	*	*	*18	*	*	*
354.0	*8	*	*	—	*	*	*	*	—	*	*
355	*51	*	*39	*	*	*	*	*	*	*	*
356	*39	*	*	*	*	*	*	*	*	*	*
357	*76	*	*	*	*	*	*	*	*	*	*
358	*37	*	*	*	*	*	*	*	*	*	*
358.0	*33	*	*	*	*	*	*	*	*	*	*
361	27	16	*11	*	*	*10	*11	*	*	20	*
361.0	*15	*10	*	*	*	*	*	*	*	*11	*
362	*27	*	*	*	*	*	*	*	*	*	*
365	*16	*	*	*	*	*	*	*	*	*	*
366	58	22	36	*	*	*7	48	40	*	12	*
366.1	12	*	9	—	—	*	11	9	*	*	*
366.10	*7	*	*6	—	—	*	*7	*7	—	—	*
366.8	*8	*6	*	—	—	*	*6	*	*	*	*
366.9	38	14	23	*	*	*	31	27	*	*10	—
376	*24	*	*	*	*	*	*	*	*	*	*
376.0	*22	*	*	*	*	*	*	*	*	*	*
376.01	*22	*	*	*	*	*	*	*	*	*	*
380	*28	*	*	*	*	*	*	*	*	*	*
380.1	*26	*	*	*	*	*	*	*	*	*	*
381	*15	*	*	*12	*	*	*	*	*	*	*
382	55	27	28	45	*	*	*	*19	*13	*15	*
382.9	45	24	22	41	*	*	*	*	*13	*15	*
386	83	*24	59	*	*	20	56	*27	*24	*25	*
386.1	*17	*	*	*	*	*	*	*	*	*	*
386.3	52	*18	34	*	*	*10	38	*20	*15	*13	*
386.30	47	*18	29	*	*	*10	*34	*20	*	*	*
390-459	33,638	16,637	17,001	101	1,976	8,007	23,554	9,152	7,575	12,248	4,663
394	*50	*	*46	*	*	*	*	*	*	*	*
396	103	*51	*52	*	*	*	*78	*	*	*	*
401	601	228	373	*	79	206	313	165	108	261	68
401.0	93	37	56	*	*	29	48	*27	*19	42	*
401.9	498	186	312	*	59	174	263	134	86	215	63
402	538	216	322	—	*	125	393	142	184	158	54
402.9	508	205	303	—	*	115	377	129	177	151	52
402.90	54	*29	*25	—	*	*29	*24	*	*	*	*
402.91	454	176	278	—	*	86	353	114	154	141	*45

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
403 . . . . .	411	227	184	*	95	114	198	90	88	198	*35
403.0 . . . . .	*47	*	*	—	*	*	*	*	*	*	*
403.01 . . . . .	*44	*	*	—	*	*	*	*	*	*	*
403.9 . . . . .	363	197	166	*	68	109	181	72	86	171	*34
403.90 . . . . .	*23	*	*	*	*	*	*	*	*	*	*
403.91 . . . . .	340	185	155	*	63	104	169	71	85	150	*34
404 . . . . .	258	125	132	*	*	*	191	*77	*49	99	*
404.9 . . . . .	230	114	116	*	*	*	180	*70	*45	83	*
404.93 . . . . .	191	98	93	—	*	*	145	*57	*	*69	*
410 . . . . .	5,121	2,953	2,168	—	210	1,477	3,434	1,405	1,095	1,788	833
410.0 . . . . .	254	142	112	—	*	76	165	*48	62	112	*32
410.01 . . . . .	224	116	107	—	*	73	139	*	61	88	*28
410.1 . . . . .	957	572	385	—	58	303	596	253	213	322	169
410.11 . . . . .	858	518	340	—	49	274	535	214	193	299	152
410.12 . . . . .	78	*44	*35	—	*	*27	*43	*	*	*	*
410.2 . . . . .	157	83	74	—	*	61	92	*28	*39	67	*
410.21 . . . . .	154	80	74	—	*	61	89	*28	*37	66	*
410.3 . . . . .	129	81	*49	—	*	*42	82	*39	*	*47	*
410.31 . . . . .	107	72	*34	—	*	*40	*63	*37	*	*32	*
410.4 . . . . .	1,070	676	394	—	73	376	621	304	282	312	173
410.41 . . . . .	1,019	636	382	—	71	362	585	292	269	294	164
410.42 . . . . .	46	*38	*	—	*	*	*32	*	*	*	*
410.5 . . . . .	135	72	*63	—	*	*48	82	*	*34	54	*
410.51 . . . . .	127	69	*58	—	*	*47	75	*	*	*50	*
410.6 . . . . .	*34	*	*	—	*	*	*	*	*	*	*
410.61 . . . . .	*34	*	*	—	*	*	*	*	*	*	*
410.7 . . . . .	1,793	995	798	—	*41	432	1,320	584	322	617	269
410.70 . . . . .	*42	*	*	—	*	*	*	*	*	*	*
410.71 . . . . .	1,651	943	708	—	*37	397	1,217	550	299	581	221
410.72 . . . . .	100	43	*57	—	*	*	74	*	*	*25	*
410.8 . . . . .	73	*41	*	—	*	*	*62	*	*	*	*
410.81 . . . . .	66	*36	*	—	*	*	*54	*	*	*	*
410.9 . . . . .	518	275	243	—	*	118	392	88	89	230	112
410.90 . . . . .	147	60	*88	—	*	*30	115	*	*	46	*58
410.91 . . . . .	367	214	153	—	*	86	274	63	67	182	55
411 . . . . .	1,286	605	681	*	77	374	834	422	265	454	145
411.1 . . . . .	1,158	539	619	—	66	339	753	392	240	420	106
411.8 . . . . .	121	65	56	*	*	*31	78	*30	*	*28	*38
411.81 . . . . .	75	*40	*	*	*	*	*47	*	*	*	*
411.89 . . . . .	46	*25	*21	—	*	*	*31	*17	*	*	*
412 . . . . .	32	*22	*	—	*	*	*	*	*	*	*
413 . . . . .	325	134	191	—	27	102	195	82	81	131	30
413.9 . . . . .	317	128	190	—	24	98	195	81	80	129	27
414 . . . . .	4,464	2,850	1,613	—	187	1,547	2,730	1,086	1,053	1,646	679
414.0 . . . . .	4,337	2,768	1,569	—	180	1,505	2,652	1,049	1,027	1,601	660
414.00 . . . . .	847	478	368	—	*53	277	517	217	142	407	81
414.01 . . . . .	3,242	2,095	1,146	—	126	1,140	1,976	795	816	1,104	527
414.02 . . . . .	246	191	54	—	*	85	159	*37	68	88	52
414.8 . . . . .	*41	*28	*	—	*	*	*	*	*	*	*
414.9 . . . . .	78	49	*29	—	*	*21	53	*	*	*26	*
415 . . . . .	518	215	304	*	*50	143	325	114	158	171	75
415.1 . . . . .	507	206	301	*	*50	142	315	109	157	171	69
416 . . . . .	62	*29	*34	*	*	*	*34	*	*	*	*
416.0 . . . . .	*25	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
420 . . . . .	72	*28	*43	*	*	*21	*44	*	*	*	*
420.9 . . . . .	72	*28	*43	*	*	*21	*44	*	*	*	*
420.90 . . . . .	45	*	*28	*	*	*	*	*	*	*	*
421 . . . . .	294	204	*90	*	*125	*	*63	*	*	*118	*
421.0 . . . . .	283	195	*89	*	*120	*	*63	*	*	*117	*
423 . . . . .	62	*	*40	*	*	*	*	*	*	*	*
423.9 . . . . .	*48	*	*34	*	*	*	*	*	*	*	*
424 . . . . .	502	267	234	*	62	83	353	137	72	219	74
424.0 . . . . .	141	70	70	*	30	42	67	*35	*19	60	*
424.1 . . . . .	336	179	157	*	*31	*40	264	102	*53	134	*47
425 . . . . .	185	95	89	*	*29	68	81	*53	*29	76	*26
425.4 . . . . .	162	86	76	*	*	55	74	*38	*28	71	*24
426 . . . . .	223	99	124	*	*16	*19	184	70	39	95	*18
426.0 . . . . .	139	66	72	*	*	*	123	*41	*24	60	*
426.1 . . . . .	35	*16	*19	*	*	*	*27	*	*	*	*
426.13 . . . . .	*19	*	*	—	*	*	*	*	*	*	*
427 . . . . .	2,455	1,125	1,330	*18	130	485	1,821	739	542	822	352
427.0 . . . . .	102	45	56	*	*9	30	62	*27	*21	36	*
427.1 . . . . .	249	148	101	*	*	63	163	81	65	78	*
427.3 . . . . .	1,291	557	734	*	33	251	1,006	374	275	448	194
427.31 . . . . .	1,162	465	697	*	26	208	927	339	252	409	161
427.32 . . . . .	129	92	37	*	*	42	78	*34	*23	*38	33
427.4 . . . . .	73	*38	*	*	*	*	*38	*	*	*	*
427.41 . . . . .	69	*34	*	*	*	*	*36	*	*	*	*
427.5 . . . . .	67	*56	*	*	*	*	51	*	*	*22	*
427.6 . . . . .	68	39	29	*	*	*	44	*24	*	23	*
427.69 . . . . .	58	37	*21	*	*	*	39	*	*	*19	*
427.8 . . . . .	569	222	347	*	35	95	428	195	114	171	89
427.81 . . . . .	242	104	139	*	*	*32	207	88	*37	83	34
427.89 . . . . .	327	118	208	*	31	62	222	107	77	88	54
427.9 . . . . .	36	*19	*	*	*	*	*29	*	*	*	*
428 . . . . .	5,580	2,278	3,301	*	197	804	4,568	1,617	1,306	2,089	569
428.0 . . . . .	5,543	2,255	3,288	*	194	799	4,540	1,609	1,295	2,079	560
428.1 . . . . .	*31	*	*	*	*	*	*25	*	*	*	*
429 . . . . .	91	*40	51	*	*	*	68	*	*	58	*
429.2 . . . . .	*21	*	*	—	*	*	*	*	*	*	*
429.4 . . . . .	*39	*	*	*	*	*	*	*	*	*	*
430 . . . . .	352	162	190	*	*62	172	111	*86	*104	69	*
431 . . . . .	664	319	345	*	*75	160	426	129	101	287	146
432 . . . . .	156	*90	*66	*	*	*	95	*	*33	*58	*
432.1 . . . . .	113	*76	*37	*	*	*	*78	*	*	*	*
433 . . . . .	845	483	362	*	*	175	661	203	184	369	90
433.1 . . . . .	542	311	232	*	*	130	408	124	141	209	68
433.10 . . . . .	398	224	174	*	*	88	306	93	102	151	52
433.11 . . . . .	144	86	*58	—	*	*42	102	*	*39	*58	*
433.3 . . . . .	255	146	109	*	*	*29	225	*63	*35	140	*17
433.30 . . . . .	155	86	69	*	*	*16	138	*39	*22	87	*
433.31 . . . . .	99	*59	*	—	—	*	*87	*	*	*	*
434 . . . . .	2,216	914	1,302	*	*57	309	1,846	716	467	713	320
434.0 . . . . .	148	87	*61	—	*	*	128	*44	*	*51	*
434.01 . . . . .	133	*79	*55	—	*	*	116	*	*	*46	*
434.1 . . . . .	268	109	159	—	*	*49	213	125	*51	*56	*36
434.11 . . . . .	245	100	145	—	*	*47	192	116	*49	*44	*35
434.9 . . . . .	1,800	718	1,082	*	*47	245	1,505	547	393	606	253

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
434.90	*66	*	*	*	*	*	*51	*	*	*	*
434.91	1,734	693	1,041	*	*44	233	1,454	536	369	579	250
435	723	343	380	*	*	118	594	244	179	235	66
435.0	*30	*	*	—	—	*	*24	*	*	*	*
435.8	*24	*	*	—	*	*	*	*	*	*	*
435.9	637	302	335	*	*	106	520	205	155	217	60
436	1,024	408	616	*	*	174	829	162	251	465	146
437	191	75	116	*	*	*42	124	*36	54	89	*
437.1	*64	*	*44	*	—	*	*57	*	*	*	*
437.2	*28	*	*	—	*	*	*	*	*	*	*
437.3	*42	*	*	*	*	*	*	*	*	*	*
438	123	*55	*68	*	*	*	70	*	*	*44	*
440	955	464	491	*	*	214	734	272	244	322	118
440.2	837	404	433	*	*	174	658	239	223	276	99
440.20	66	*37	*	—	*	*	*58	*	*	*	*
440.21	95	53	*42	—	*	*25	68	*	*	46	*
440.22	*72	*	*	—	*	*	*58	*	*	*	*
440.23	146	*72	*74	—	*	*	107	*	*	*	*
440.24	442	210	232	—	*	*85	355	137	112	141	*52
441	498	382	116	*	*	86	399	128	112	183	75
441.0	*54	*	*	—	*	*	*31	*	*	*	*
441.01	*39	*	*	—	*	*	*	*	*	*	*
441.3	*71	*	*	—	—	*	*64	*	*	*	*
441.4	330	261	*69	—	*	*53	277	*76	89	120	*45
442	57	*36	*	*	*	*	*32	*	*	*	*
442.3	*33	*	*	—	*	*	*26	*	*	*	*
443	125	48	77	—	*	*	99	*45	*	*50	*
443.9	103	*39	*64	—	*	*	86	*39	*	*35	*
444	468	198	270	*	*	106	329	116	108	159	*86
444.0	*83	*	*	*	*	*	*	*	*	*	*
444.2	315	126	189	*	*	*68	232	80	72	100	*63
444.22	291	119	172	—	*	*64	217	*75	*68	86	*61
444.8	*54	*	*	—	*	*	*	*	*	*	*
444.81	*42	*	*	—	*	*	*	*	*	*	*
446	*57	*	*	*	*	*	*	*	*	*	*
447	75	*35	*40	*	*	*25	*39	*	*	*22	*
447.1	48	*20	*28	—	*	*	*32	*	*	*	*
451	275	97	178	*	46	98	131	59	69	101	*46
451.0	*39	*	*29	—	*	*	*	*	*	*	*
451.1	160	56	105	*	*	*54	81	*37	*43	*53	*
451.11	*52	*	*	*	*	*	*	*	*	*	*
451.19	109	*41	68	—	*	*38	*46	*	*	*32	*
451.2	*45	*	*33	—	*	*	*	*	*	*	*
453	868	369	499	*	110	216	541	239	234	297	98
453.8	823	348	475	*	106	200	516	232	228	269	94
453.9	*32	*	*	—	*	*	*	*	*	*	*
454	136	*51	85	—	*14	*	108	*33	*29	*53	*
454.2	106	*40	*67	—	*	*	*92	*	*	*	*
455	94	39	55	*	30	24	41	30	*	36	*13
455.2	58	*18	39	*	*17	*	28	*16	*	*25	*
456	*29	*	*	*	*	*	*	*	*	*	*
458	232	85	147	*	*	50	170	53	62	73	*42
458.0	96	*	74	—	*	*	73	*24	*38	*22	*
458.9	136	63	73	*	*	*34	97	*	*25	51	*

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).



**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
459 . . . . .	90	*46	44	*	*	*25	56	*	*34	*	*
459.8 . . . . .	*38	*	*	*	*	*	*25	*	*	*	*
459.81 . . . . .	*32	*	*	—	*	*	*	*	*	*	*
459.9 . . . . .	*29	*	*	—	—	*	*	*	*	*	*
460-519 . . . . .	19,888	9,147	10,741	2,490	2,272	3,571	11,556	4,972	4,405	7,736	2,774
461 . . . . .	63	*30	*33	*16	*	*	*	*	*17	*30	*
461.0 . . . . .	*22	*	*	*	*	*	*	*	*	*	*
462 . . . . .	*25	*	*	*	*	*	*	*	*	*	*
463 . . . . .	42	*20	*22	*24	*16	*	—	*	*	*21	*
464 . . . . .	81	51	30	64	*	*	*	25	23	20	*12
464.4 . . . . .	51	34	17	50	—	*	*	*15	15	*14	*
465 . . . . .	107	54	53	51	*	*	*	29	*13	51	*
465.9 . . . . .	102	50	52	47	*	*	*	*25	*13	51	*
466 . . . . .	1,031	539	491	554	92	128	257	234	251	388	158
466.0 . . . . .	548	235	312	77	89	126	256	133	166	223	*25
466.1 . . . . .	483	304	179	477	*	*	*	101	84	165	133
470 . . . . .	13	*	*6	—	*7	*	*	*7	*	*	*
473 . . . . .	56	*23	33	*	*21	*	*	*13	*	*26	*
473.9 . . . . .	*23	*	*	*	*	*	*	*	*	*	*
474 . . . . .	40	23	17	30	*9	*	*	26	*	*	*
474.0 . . . . .	15	*7	*7	*10	*	—	—	*7	*	*	*
474.1 . . . . .	25	16	*9	20	*	*	*	19	*	*	*
474.10 . . . . .	20	*13	*7	17	*	—	—	16	*	*	*
475 . . . . .	29	*16	*13	*	19	*	*	*	*	*10	*
478 . . . . .	114	63	*51	*27	*29	*26	*32	*51	*17	*35	*
478.2 . . . . .	*37	*26	*	*	*	*	*	*	*	*	*
478.7 . . . . .	*33	*	*	*	*	*	*	*	*	*	*
480 . . . . .	192	122	70	136	*	*	*33	*22	43	81	*47
480.1 . . . . .	84	48	*36	83	*	*	—	*	*25	*27	*
480.9 . . . . .	72	*42	*30	*26	*	*	*	*	*	*38	*
481 . . . . .	409	231	178	*	98	71	225	58	111	140	100
482 . . . . .	2,128	1,083	1,045	76	231	262	1,559	351	420	1,037	319
482.0 . . . . .	206	122	83	*	*	*	189	*	*	80	*
482.1 . . . . .	462	244	218	*	*97	*	291	*	*98	225	*84
482.2 . . . . .	189	90	98	*	*	*	140	*	*42	79	*
482.3 . . . . .	116	59	*57	*	*	*	81	*	*	*58	*
482.30 . . . . .	*48	*	*	*	*	*	*	*	*	*	*
482.39 . . . . .	*50	*	*	*	*	*	*	*	*	*	*
482.4 . . . . .	372	173	199	*	*	*	296	*67	*85	167	*
482.8 . . . . .	659	321	339	*	*47	93	493	*81	97	384	*97
482.82 . . . . .	*69	*	*	—	*	*	*64	*	*	*	*
482.83 . . . . .	354	159	194	*	*	*67	253	*	66	199	*55
482.89 . . . . .	218	105	113	*	*	*	165	*	*	151	*
482.9 . . . . .	124	73	52	*	*28	*	*69	*40	*	*45	*
483 . . . . .	98	*53	*45	*	*	*	*	*	*	*48	*
483.0 . . . . .	82	*46	*37	*	*	*	*	*	*	*40	*
485 . . . . .	247	121	126	72	*	*	125	*72	56	102	*
486 . . . . .	5,262	2,382	2,880	599	552	709	3,402	1,473	1,305	1,882	602
487 . . . . .	78	*35	44	*	*	*	*42	*	*	*27	*
487.0 . . . . .	*35	*	*	*	*	*	*	*	*	*	*
487.1 . . . . .	40	*	*23	*	*	*	*	*	*	*21	*
490 . . . . .	170	65	*105	*	*	*	70	*	*23	49	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
491 . . . . .	2,029	790	1,239	*	73	483	1,463	455	471	844	259
491.2 . . . . .	2,004	782	1,222	*	72	470	1,454	451	464	832	257
491.20 . . . . .	124	52	72	*	*	*26	92	*39	*	52	*
491.21 . . . . .	1,880	730	1,150	*	69	444	1,361	412	450	781	237
492 . . . . .	339	235	104	*	*	150	185	*64	59	164	*
492.8 . . . . .	306	210	95	*	*	142	163	*56	*52	150	*
493 . . . . .	1,903	661	1,243	576	453	420	454	586	428	651	237
493.0 . . . . .	87	38	49	54	*25	*	*	31	*16	*17	23
493.00 . . . . .	33	*16	*16	*20	*	*	*	*	*	*	*11
493.01 . . . . .	54	22	32	34	*16	*	—	24	*	*	*12
493.2 . . . . .	385	97	288	*	*38	145	199	144	70	129	42
493.20 . . . . .	280	73	208	*	*32	103	145	104	53	90	*33
493.21 . . . . .	105	*	81	*	*	*42	*54	*40	*	*38	*
493.9 . . . . .	1,425	523	902	518	388	267	252	408	341	504	172
493.90 . . . . .	809	267	542	231	213	167	197	181	186	350	91
493.91 . . . . .	617	256	361	287	175	100	55	228	155	154	80
496 . . . . .	946	420	526	*	*	207	723	249	150	448	100
507 . . . . .	1,237	643	593	*44	*62	165	966	301	277	450	208
507.0 . . . . .	1,225	643	582	*42	*62	165	957	301	276	450	198
510 . . . . .	*118	*96	*	*	*	*	*	*	*	*	*
510.9 . . . . .	*103	*	*	*	*	*	*	*	*	*	*
511 . . . . .	368	144	223	*	36	93	231	85	57	153	72
511.0 . . . . .	44	*22	*22	—	*10	*	*23	*	*18	*11	*
511.8 . . . . .	*69	*	*	*	*	*	*	*	*	*	*
511.9 . . . . .	227	81	145	*	*	53	156	*57	*27	109	*33
512 . . . . .	235	127	109	*	85	*65	80	*42	69	93	*
512.1 . . . . .	*48	*	*	*	*	*	*	*	*	*	*
512.8 . . . . .	162	91	*71	*	63	*	*	*34	*39	*65	*
513 . . . . .	*82	*	*	*	*	*	*	*	*	*	*
513.0 . . . . .	*82	*	*	*	*	*	*	*	*	*	*
514 . . . . .	44	*16	*28	*	*	*	*	*	*	*18	*
515 . . . . .	153	*46	*107	—	*	*	119	*	*28	*	*
516 . . . . .	126	65	*61	*	*	*40	*63	*	*	*44	*
516.3 . . . . .	*60	*	*	—	*	*	*	*	*	*	*
516.8 . . . . .	*63	*	*	*	*	*	*	*	*	*	*
518 . . . . .	1,836	808	1,029	*84	141	428	1,184	382	385	678	391
518.0 . . . . .	*64	*	*45	*	*	*	*	*	*	*	*
518.4 . . . . .	68	*	*	*	*	*	*43	*	*	*	*
518.5 . . . . .	*59	*	*	*	*	*	*	*	*	*	*
518.8 . . . . .	1,600	693	907	*79	113	353	1,055	313	342	602	343
518.81 . . . . .	1,444	610	834	*	*79	334	990	277	303	558	306
518.82 . . . . .	88	*	*48	*	*	*	*	*	*	*	*
518.89 . . . . .	68	*42	*	*	*	*	*40	*	*	*	*
519 . . . . .	210	103	107	*22	*	*79	92	*100	*25	83	*
519.1 . . . . .	*28	*	*	*	*	*	*	*	*	*	*
519.8 . . . . .	88	*	*58	*	*	*	*63	*	*	*66	*
520-579 . . . . .	15,790	6,823	8,967	833	3,108	4,286	7,563	3,899	3,372	6,236	2,283
522 . . . . .	*14	*	*	*	*	*	*	*	*	*	*
524 . . . . .	38	*12	26	*	31	*	*	*	*	*16	*
524.0 . . . . .	*13	*	*	*	*9	*	—	*	*	*	*
527 . . . . .	42	*	*24	*	*	*	*	*	*	*21	*
527.2 . . . . .	36	*	*20	*	*	*	*	*	*	*19	*
528 . . . . .	*44	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
530	728	319	409	98	89	198	343	191	128	297	111
530.1	163	69	93	*	26	55	67	46	*22	70	*25
530.10	77	*38	40	*	*	*23	32	22	*	41	*
530.11	66	*20	45	*	*10	*24	*29	*19	*	*23	*
530.19	*20	*	*	*	*	*	*	*	*	*	*
530.2	106	39	67	—	*	*26	70	*	*	*54	*
530.3	77	*46	*31	*	*	*	61	*	*	*	*
530.5	*35	*	*	*	*	*	*	*	*	*	*
530.7	*31	*	*	*	*	*	*	*	*	*	*
530.8	255	100	155	77	41	65	72	77	64	88	26
530.81	220	94	126	77	36	56	51	58	60	77	25
531	605	255	350	*	69	136	396	139	131	211	125
531.0	144	63	82	*	*	*	102	*	*38	52	*
531.00	138	59	79	*	*	*	99	*	*36	52	*
531.3	*28	*	*	—	*	*	*	*	*	*	*
531.30	*19	*	*	—	*	*	*	*	*	*	*
531.4	247	122	125	*	*22	63	161	59	50	95	*43
531.40	235	113	123	*	*22	52	160	58	50	85	*43
531.5	*59	*	*	*	*	*	*	*	*	*	*
531.9	52	*	*33	*	*	*	*29	*	*	*	*
531.90	46	*	*28	*	*	*	*24	*	*	*	*
532	556	332	225	*	66	96	394	154	125	193	84
532.0	159	108	*51	—	*	*	118	*	*	*57	*28
532.00	148	107	*41	—	*	*	108	*	*	*53	*27
532.3	*25	*	*	—	*	*	*	*	*	*	*
532.30	*24	*	*	—	*	*	*	*	*	*	*
532.4	196	136	*60	—	*22	*31	144	*62	*38	*61	*35
532.40	182	122	*60	—	*22	*26	134	*59	*32	*59	*31
532.9	44	*	*29	*	*	*	*26	*	*19	*	*
532.90	42	*	*27	*	*	*	*	*	*19	*	*
533	105	*29	76	*	*19	*27	58	*20	*26	51	*
533.4	*42	*	*33	—	*	*	*33	*	*	*29	*
533.40	*42	*	*33	—	*	*	*33	*	*	*29	*
533.9	32	*	*24	*	*	*	*	*	*	*	*
533.90	31	*	*23	*	*	*	*	*	*	*	*
535	615	242	373	*31	143	134	307	136	155	247	77
535.0	142	58	84	*	23	37	75	*39	*28	56	*18
535.00	83	31	52	*	*19	*20	39	*	*22	40	*
535.01	59	*27	*32	*	*	*	*36	*	*	*	*
535.1	51	*	39	*	*	*	*40	*	*	*	*
535.10	*31	*	*26	*	*	*	*	*	*	*	*
535.3	45	*19	*26	—	*22	*	*	*	*	*	*
535.30	*18	*	*	—	*	*	*	*	*	*	*
535.31	*27	*	*	—	*	*	*	*	*	*	*
535.4	155	62	92	*	39	*19	90	*29	55	49	*21
535.40	70	*18	53	*	*21	*	*36	*15	*23	*	*
535.41	84	45	*39	*	*	*	54	*	*33	*29	*
535.5	180	71	109	*	44	48	76	37	34	93	*17
535.50	124	51	73	*	35	31	46	*29	*22	62	*
535.51	57	*20	*37	*	*	*	*30	*	*	*31	*
535.6	39	*18	*	*	*	*	*	*	*	*	*
535.60	*29	*15	*	*	*	*	*	*	*	*	*
536	173	57	116	*	41	*27	98	*57	*	69	*
536.2	51	*	*27	*	*	*	*	*	*	*24	*
536.3	*27	*	*	*	*	*	*	*	*	*	*
536.8	83	*23	*60	*	*11	*	*63	*	*	*32	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
537 . . . . .	162	*54	107	*	*	*	93	*46	*	*66	*37
537.0 . . . . .	105	*36	*68	*	*	*	*52	*	*	*	*
537.8 . . . . .	*47	*	*	*	*	*	*34	*	*	*	*
537.83 . . . . .	*30	*	*	—	*	*	*	*	*	*	*
540 . . . . .	749	454	295	177	364	103	105	196	161	235	157
540.0 . . . . .	258	156	102	72	86	*55	*	*76	57	81	44
540.1 . . . . .	181	133	*48	*41	77	*	*	*	*43	*61	*41
540.9 . . . . .	310	165	145	64	200	32	*	84	61	93	72
541 . . . . .	35	*18	*17	*	*26	*	*	*	*	*17	*
550 . . . . .	161	143	*18	*14	26	25	96	60	39	44	*19
550.1 . . . . .	77	67	*	*	*	*	53	*25	*	*24	*
550.10 . . . . .	61	53	*	*	*	*	42	*19	*	*19	*
550.9 . . . . .	81	75	*	*10	*14	17	41	33	*21	20	*
550.90 . . . . .	48	45	*	*	*10	*12	22	21	*	*14	*
550.91 . . . . .	*16	*16	*	*	*	*	*	*	*	*	*
550.92 . . . . .	*16	*13	*	*	*	*	*	*	*	*	*
552 . . . . .	238	74	164	*	*26	87	119	76	56	73	*33
552.1 . . . . .	*30	*	*	*	*	*	*	*	*	*	*
552.2 . . . . .	119	*29	89	*	*	*49	58	*37	*	*35	*
552.20 . . . . .	*32	*	*28	*	*	*	*	*	*	*	*
552.21 . . . . .	81	*	58	—	*	*35	*36	*	*	*	*
553 . . . . .	315	118	197	*	51	138	123	71	57	144	43
553.1 . . . . .	*36	*	*30	*	*	*	*	*	*	*	*
553.2 . . . . .	176	77	99	*	35	73	67	37	42	83	*15
553.20 . . . . .	30	*	*23	—	*	*	*	*	*	*	*
553.21 . . . . .	139	66	73	*	*25	63	51	*28	*32	69	*
553.3 . . . . .	94	34	60	*	*	38	45	*24	*	48	*
555 . . . . .	251	119	133	*	127	*67	*50	*65	75	86	*
555.0 . . . . .	*61	*	*	*	*	*	*	*	*	*	*
555.1 . . . . .	*59	*	*	*	*	*	*	*	*	*	*
555.9 . . . . .	83	*28	54	*	52	*	*	*	*	*38	*
556 . . . . .	201	112	89	*	64	*70	*57	*44	*43	*63	*
556.9 . . . . .	84	*31	*53	*	*28	*	*	*	*	*29	*
557 . . . . .	264	*66	198	*	*	*69	163	*57	*72	91	*44
557.0 . . . . .	162	*	123	*	*	*	100	*	*	*51	*
557.1 . . . . .	*51	*	*37	*	*	*	*	*	*	*	*
557.9 . . . . .	*51	*	*39	—	*	*	*40	*	*	*	*
558 . . . . .	751	280	471	191	138	139	284	165	171	325	90
558.9 . . . . .	716	270	446	191	137	124	264	145	165	320	85
560 . . . . .	1,796	795	1,002	57	187	464	1,088	380	394	711	312
560.1 . . . . .	265	141	124	*	*24	56	178	*49	54	150	*
560.2 . . . . .	125	*72	*	*	*	*	*92	*	*	*65	*
560.3 . . . . .	146	73	73	*	*	*	118	*39	*	66	*
560.39 . . . . .	136	71	65	*	*	*	108	*39	*	64	*
560.8 . . . . .	732	274	458	*	93	205	413	168	176	223	165
560.81 . . . . .	602	217	386	*	74	174	336	145	134	194	131
560.89 . . . . .	129	58	72	*	*	*31	77	*	*	*	*35
560.9 . . . . .	515	227	287	*	52	166	287	88	120	204	103
562 . . . . .	1,388	499	889	*	101	387	900	390	290	548	159
562.1 . . . . .	1,378	495	883	*	101	386	891	387	288	547	156
562.10 . . . . .	173	52	121	*	*	*40	119	*42	*	65	*
562.11 . . . . .	757	291	466	—	82	263	413	233	167	272	86
562.12 . . . . .	346	124	222	—	*	*72	274	99	60	161	*27
562.13 . . . . .	102	*27	75	—	*	*	85	*	*	*49	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
564 . . . . .	182	60	122	*11	53	*24	93	43	41	62	36
564.0 . . . . .	55	*19	37	*	*	*	*32	*	*	*14	*
564.1 . . . . .	63	*	55	*	*28	*	*	*	*21	*	*
564.8 . . . . .	*49	*	*	*	*	*	*	*	*	*	*
565 . . . . .	*20	*	*	*	*	*	*	*	*	*	*
566 . . . . .	85	57	*27	*	28	*29	*	*26	*	*33	*
567 . . . . .	186	*100	87	*	*	*53	*96	*	*	*74	*
567.2 . . . . .	157	*88	*69	*	*	*44	*	*	*	*67	*
568 . . . . .	59	*	*43	*	*29	*	*	*	*	*	*
568.0 . . . . .	*43	*	*33	*	*22	*	*	*	*	*	*
569 . . . . .	558	202	357	*	100	143	310	157	101	228	73
569.3 . . . . .	62	*23	*39	*	*	*	*49	*	*	*40	*
569.4 . . . . .	*31	*	*	*	*	*	*	*	*	*	*
569.6 . . . . .	86	*	*53	*	*	*	*49	*	*	*	*
569.8 . . . . .	310	120	189	*	*	94	147	*90	*58	118	*43
569.81 . . . . .	*81	*	*	—	*	*	*	*	*	*	*
569.83 . . . . .	*100	*	*	*	*	*	*	*	*	*	*
569.85 . . . . .	78	*36	*42	—	*	*	59	*	*	*	*
570 . . . . .	*91	*	*	*	*	*	*	*	*	*	*
571 . . . . .	603	324	279	*	115	328	152	139	108	293	63
571.1 . . . . .	101	82	*	—	*39	*59	*	*	*	*45	*
571.2 . . . . .	222	156	*67	—	*46	118	*58	*70	*53	75	*
571.5 . . . . .	140	60	80	*	*	*45	81	*34	*	69	*
572 . . . . .	245	114	132	*	*51	85	91	*	*	139	*
572.2 . . . . .	136	53	*83	—	*	*60	*50	*	*	*77	*
573 . . . . .	77	*	55	*	*	*22	*	*	*	*49	*
573.3 . . . . .	*43	*	*27	*	*	*	*	*	*	*	*
574 . . . . .	1,804	624	1,180	*	399	510	886	487	390	670	257
574.0 . . . . .	707	289	418	*	142	213	352	177	182	260	88
574.00 . . . . .	620	265	355	—	134	188	297	144	163	235	77
574.01 . . . . .	88	*	63	*	*	*25	*	*	*	*	*
574.1 . . . . .	692	184	508	*	177	209	301	219	114	250	109
574.10 . . . . .	668	173	495	*	170	199	293	215	109	241	103
574.2 . . . . .	155	57	98	*	38	55	61	*25	41	73	*16
574.20 . . . . .	152	56	96	*	38	53	61	*25	40	72	*15
574.3 . . . . .	102	*46	*56	*	*	*	76	*	*	*58	*
574.30 . . . . .	*35	*	*	*	*	*	*	*	*	*	*
574.31 . . . . .	*68	*	*	—	*	*	*52	*	*	*	*
574.4 . . . . .	*51	*	*37	*	*	*	*	*	*	*	*
574.5 . . . . .	97	*34	63	*	*19	*	71	*	*	*	*22
574.50 . . . . .	56	*	*33	*	*	*	*38	*	*	*	*
574.51 . . . . .	41	*	*30	—	*	*	*33	*	*	*	*
575 . . . . .	256	108	148	*	70	78	107	57	64	115	*20
575.0 . . . . .	117	61	56	*	*25	*34	*57	*30	*34	*45	*
575.1 . . . . .	102	*36	66	*	40	*26	*	*19	*13	62	*
575.8 . . . . .	*24	*	*	*	*	*	*	*	*	*	—
576 . . . . .	174	64	110	*	*	*29	123	*44	*	68	*36
576.1 . . . . .	74	*40	*35	*	*	*	*48	*	*	*	*
576.2 . . . . .	*55	*	*45	*	*	*	*41	*	*	*	*
576.8 . . . . .	*33	*	*	*	*	*	*26	*	*	*	*
577 . . . . .	1,281	608	673	*	413	458	380	264	345	491	181
577.0 . . . . .	1,024	513	511	*	320	381	300	217	273	410	123
577.1 . . . . .	180	64	116	*	67	*50	*61	*	*57	61	*29
578 . . . . .	725	311	414	*	80	149	483	163	154	324	84
578.0 . . . . .	92	*61	*32	*	*	*	68	*	*	*41	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
578.1	107	47	60	*	*	*	70	*	*34	36	*
578.9	527	204	323	*	54	121	346	124	114	248	41
580-629	7,183	2,498	4,685	274	1,963	1,432	3,514	1,769	1,492	2,941	981
584	576	322	254	*	*	122	420	171	137	173	95
584.9	514	300	214	*	*	81	404	134	132	157	92
585	283	162	121	*	*59	62	156	89	*49	93	*52
586	62	*	*44	*	*	*	*34	*	*	*	*
590	491	105	386	68	182	105	136	106	121	172	91
590.1	314	68	246	45	129	56	84	77	82	103	52
590.10	312	66	246	45	129	56	82	75	82	103	52
590.8	146	*	117	*23	48	*33	*41	*23	*33	60	*31
590.80	146	*	117	*23	48	*33	*41	*23	*33	60	*31
591	92	*26	*65	*	*	*	*59	*22	*	*	*
592	438	252	185	*	184	169	80	114	114	152	58
592.0	141	64	77	*	64	48	*26	41	30	43	27
592.1	294	186	107	*	118	120	54	72	82	108	31
593	206	92	114	*31	43	*55	77	54	*47	73	32
593.4	67	*34	*33	*	*20	*	*26	*	*	*24	*15
593.7	*27	*	*	*20	*	*	—	*	*	*	*
593.70	*21	*	*	*	*	—	—	*	*	*	*
593.9	64	*	*30	*	*	*	*31	*	*	*	*
594	*26	*22	*	*	*	*	*	*	*	*	*
594.1	*21	*	*	*	*	*	*	*	*	*	*
595	76	*	50	*	*	*	*55	*	*	*	*
595.0	*34	*	*	*	*	*	*	*	*	*	*
596	143	100	*43	*	*	*23	106	*30	*	69	*
596.0	52	49	*	—	*	*	44	*	*	*	*
598	*25	*24	*	*	*	*	*	*	*	*	*
599	2,007	684	1,324	112	140	228	1,527	557	364	886	201
599.0	1,888	616	1,273	111	120	207	1,450	523	342	836	188
599.7	95	57	38	*	*15	*17	61	*26	*14	*45	*
600	419	419	—	—	*	56	361	109	93	161	56
601	57	57	—	—	*	*	*39	*	*	*27	*
601.0	42	42	—	—	*	*	*31	*	*	*	*
604	33	33	—	*	*	*	*	*	*	*	*
604.9	*31	*31	—	*	*	*	*	*	*	*	*
604.90	*19	*19	—	*	*	*	*	*	*	*	*
607	*21	*21	—	*	*	*	*	*	*	*	*
607.8	*10	*10	—	—	*	*	*	*	*	*	*
608	*44	*44	—	*	*	*	*	*	*	*	*
610	*12	*	*12	*	*	*	*	*	*	*	*
610.1	*10	—	*10	—	*	*	*	*	*	*	*
611	66	*	62	*	36	*17	*	16	*9	32	*
611.1	34	*	30	*	25	*	*	*12	*	*12	*
614	283	—	283	*	229	*33	*	84	50	120	29
614.2	*43	—	*43	*	*30	*	—	*	*	*	*
614.3	44	—	44	*	42	*	*	*	*	*	*
614.6	75	—	75	—	53	*	*	*	*	42	*
614.9	62	—	62	*	59	*	—	*	*	*28	*
616	33	—	33	*	*22	*	*	*	*	*16	*
616.0	*15	—	*15	—	*	*	*	*	*	*	*
617	312	—	312	—	254	53	*	41	50	179	42
617.0	183	—	183	—	138	40	*	*18	*23	120	*22

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
617.1	45	—	45	—	39	*	*	*	*	*15	*
617.3	57	—	57	—	52	*	—	*	*	*25	*
618	508	—	508	*	120	173	214	98	118	213	79
618.0	114	—	114	—	*	46	58	*18	36	41	*19
618.1	62	—	62	—	38	*	*	*	*	*26	*
618.2	92	—	92	—	*26	30	36	*26	*16	33	*
618.3	56	—	56	—	*	*15	34	*17	*	*13	*
618.4	67	—	67	—	*24	28	*15	*	*	41	*
618.5	59	—	59	—	*	*18	37	*	*	*25	*
618.8	47	—	47	—	*	*	*	*	*	*	*
619	*44	—	*44	—	*10	*	*	*	*	*	*
619.1	*43	—	*43	—	*	*	*	*	*	*	*
620	182	—	182	*	135	29	*	40	41	74	27
620.0	*22	—	*22	*	*19	*	*	*	*	*	*
620.1	36	—	36	*	30	—	—	*	*	*	*
620.2	104	—	104	*	72	*18	*	*21	26	39	*18
621	56	—	56	*	*24	*16	*	*16	*	*21	*
621.3	29	—	29	—	*	*	*	*	*	*	*
622	28	—	28	—	24	*	*	*	*	*19	*
622.1	26	—	26	—	23	*	*	*	*	*19	*
625	259	—	259	*	141	78	38	29	56	121	52
625.3	*23	—	*23	*	*22	*	—	*	*	*15	*
625.6	141	—	141	—	38	70	33	*22	30	59	29
625.9	81	—	81	*	69	*	*	*	*18	42	*16
626	210	—	210	*	159	47	*	*17	50	110	33
626.2	150	—	150	*	117	31	*	*9	31	86	24
626.8	46	—	46	*	36	*	*	*	*17	*19	*
627	36	—	36	—	*	23	*	*	*	*15	*
627.1	29	—	29	—	*	*20	*	*	*	*	*
630-676	1,341	—	1,341	*	1,332	*	—	306	271	517	247
632	23	—	23	—	23	—	—	*	*	15	*
633	94	—	94	—	93	*	—	26	*19	32	*16
633.1	86	—	86	—	86	*	—	24	*18	30	*14
634	55	—	55	*	54	*	—	26	*	16	*
634.9	42	—	42	*	42	*	—	20	*	13	*
634.91	30	—	30	*	29	*	—	17	*	*10	*
635	*12	—	*12	*	*12	*	—	*	*	*	*
637	*7	—	*7	—	*7	—	—	*	*	*	*
637.9	*6	—	*6	—	*6	—	—	*	*	*	*
640	*12	—	*12	—	*12	—	—	*	*	*	*
640.0	*8	—	*8	—	*8	—	—	*	*	*	*
640.03	*8	—	*8	—	*8	—	—	*	*	*	*
641	40	—	40	—	40	—	—	*	*	*12	*
641.1	*18	—	*18	—	*18	—	—	*	*	*	*
641.13	*18	—	*18	—	*18	—	—	*	*	*	*
642	60	—	60	—	60	—	—	*	*11	31	*
642.3	*21	—	*21	—	*21	—	—	*	*	*	*
642.33	*20	—	*20	—	*20	—	—	*	*	*	*
642.4	19	—	19	—	19	—	—	*	*	*	*
642.43	*18	—	*18	—	*18	—	—	*	*	*	*
643	121	—	121	—	121	—	—	*26	29	47	*
643.0	41	—	41	—	41	—	—	*	*14	*14	*
643.03	40	—	40	—	40	—	—	*	*	*13	*

\* Figure does not meet standard of reliability or precision.  
 — Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
643.1	69	—	69	—	69	—	—	*18	*	27	*
643.13	68	—	68	—	68	—	—	*18	*	26	*
644	324	—	324	*	323	—	—	48	71	131	74
644.0	278	—	278	*	278	—	—	41	62	113	63
644.03	278	—	278	*	277	—	—	41	62	112	62
644.1	42	—	42	—	42	—	—	*7	*8	15	11
644.13	41	—	41	—	41	—	—	*7	*8	15	11
646	117	—	117	*	113	—	—	*26	*18	47	26
646.6	88	—	88	*	87	—	—	*17	*	39	*20
646.63	78	—	78	*	78	—	—	*16	*	33	*17
646.8	*21	—	*21	*	*17	—	—	*	*	*	*
646.83	*19	—	*19	*	*16	—	—	*	*	*	*
647	29	—	29	*	29	—	—	*	*	*	*
647.6	*15	—	*15	*	*14	—	—	*	*	*	*
648	221	—	221	*	218	*	—	83	41	67	31
648.0	*24	—	*24	—	*24	—	—	*	*	*	*
648.03	*24	—	*24	—	*24	—	—	*	*	*	*
648.4	*57	—	*57	*	*55	—	—	*	*	*	*
648.43	*56	—	*56	*	*	—	—	*	*	*	*
648.8	38	—	38	—	38	—	—	*	*	*15	*
648.83	38	—	38	—	38	—	—	*	*	*15	*
648.9	79	—	79	*	78	*	—	21	*16	28	*14
648.93	75	—	75	*	75	*	—	*21	*16	25	*14
654	27	—	27	—	27	—	—	*	*	*17	*
654.5	*14	—	*14	—	*14	—	—	*	*	*	*
654.53	*14	—	*14	—	*14	—	—	*	*	*	*
656	*11	—	*11	—	*11	—	—	*	*	*	*
658	53	—	53	—	53	—	—	*	*	*21	*
658.1	*38	—	*38	—	*38	—	—	*	*	*	*
658.13	*37	—	*37	—	*37	—	—	*	*	*	*
659	15	—	15	—	15	—	—	*	*	*	*
659.1	*6	—	*6	—	*6	—	—	*	*	*	*
659.13	*6	—	*6	—	*6	—	—	*	*	*	*
666	*18	—	*18	*	*18	—	—	*	*	*15	*
666.2	*16	—	*16	*	*16	—	—	*	*	*15	*
666.24	*16	—	*16	*	*16	—	—	*	*	*15	*
670	*23	—	*23	—	*23	—	—	*	*	*	*
670.0	*23	—	*23	—	*23	—	—	*	*	*	*
670.04	*23	—	*23	—	*23	—	—	*	*	*	*
680-709	3,206	1,482	1,724	143	677	755	1,631	881	803	1,109	412
681	118	57	62	*	*41	*23	*46	*39	*31	*33	*
681.0	59	*32	*	*	*29	*	*	*	*	*	*
681.00	54	*31	*	*	*	*	*	*	*	*	*
681.1	59	*25	*34	*	*	*	*34	*	*	*21	*
681.10	56	*	*34	*	*	*	*	*	*	*21	*
682	2,010	986	1,024	82	425	533	970	618	496	627	270
682.0	119	65	54	*	61	*	*	*	*24	*52	*29
682.1	46	*31	*	*	*25	*	*	*	*	*18	*
682.2	185	86	99	*	62	*44	*72	*56	*	*69	*
682.3	227	134	93	*	59	*54	105	*111	62	34	*
682.4	96	59	*37	*	*32	*	*33	*27	*	*43	*
682.6	965	417	549	*23	137	257	548	323	250	283	110
682.7	303	156	147	*	*35	92	168	*59	94	111	*39
686	*39	*	*	*	*	*	*	*	*	*	*
692	*23	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).



**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
695 . . . . .	*50	*	*38	*	*	*	*	*	*	*	*
695.1 . . . . .	*37	*	*	*	*	*	*	*	*	*	*
706 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
706.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
707 . . . . .	808	347	461	*	136	129	537	133	227	362	*86
707.0 . . . . .	560	244	316	*	*112	*	375	*	168	253	*
707.1 . . . . .	240	102	138	*	*	*57	156	*57	*54	107	*
709 . . . . .	*17	*	*	*	*	*	*	*	*	*	*
710-739 . . . . .	7,299	2,964	4,336	168	1,285	1,813	4,034	1,794	1,453	2,729	1,324
710 . . . . .	209	*	176	*	90	*75	*	*	*	78	*65
710.0 . . . . .	158	*	135	*	74	*65	*	*	*	56	*
711 . . . . .	130	69	*61	*	*	*	*58	*	*42	*	*
711.0 . . . . .	121	64	*57	*	*	*	*54	*	*41	*	*
711.06 . . . . .	*46	*	*	*	*	*	*	*	*	*	*
714 . . . . .	176	*51	125	*	*	54	96	*47	*35	47	*48
714.0 . . . . .	140	*32	108	*	*	*41	89	*33	*32	*37	*39
715 . . . . .	2,130	763	1,367	*	68	444	1,618	513	473	717	428
715.1 . . . . .	191	90	101	—	*	*30	155	*	*48	90	*35
715.15 . . . . .	67	*	*42	—	*	*	*50	*	*	*	*
715.16 . . . . .	117	64	*52	—	*	*	99	*	*30	65	*
715.3 . . . . .	1,397	504	892	—	*45	289	1,063	292	229	539	337
715.35 . . . . .	504	247	257	—	*	87	396	85	54	186	179
715.36 . . . . .	876	254	622	—	*	194	659	205	168	347	156
715.9 . . . . .	501	159	343	*	*	116	373	186	190	78	47
715.90 . . . . .	*77	*	*	—	—	*	*	*	*	*	—
715.95 . . . . .	146	52	94	—	*	*45	94	*59	54	*	*
715.96 . . . . .	259	79	180	*	*	64	190	83	129	*	*
716 . . . . .	114	49	65	*	*	*37	57	*30	*	*52	*
716.9 . . . . .	64	*	46	—	*	*	*38	*	*	*32	*
717 . . . . .	59	35	24	*	48	*9	*	25	*16	*9	*8
717.8 . . . . .	40	25	*16	*	34	*	*	*14	*14	*	*
717.83 . . . . .	37	22	*15	*	32	*	—	*14	*14	*	*
718 . . . . .	62	33	29	*	33	*	*	*16	*	20	*17
718.3 . . . . .	19	*10	*	*	*12	*	*	*	*	*	*
718.31 . . . . .	*9	*8	*	—	*8	*	*	*	*	*	*
718.8 . . . . .	*14	*	*	—	*9	*	*	*	*	*	*
719 . . . . .	257	97	160	*	*34	*21	189	*47	*30	143	*38
719.4 . . . . .	86	*43	43	*	*27	*	*52	*	*	*	*27
719.45 . . . . .	*29	*	*	*	*	*	*	*	*	*	*
719.7 . . . . .	134	*	*104	—	*	*	123	*	*	*85	*
720 . . . . .	*44	*	*	—	*	*	*	*	*	*	*
721 . . . . .	327	151	176	—	*45	92	190	88	61	104	74
721.0 . . . . .	63	*25	38	—	*	26	*30	*	*16	27	*
721.1 . . . . .	88	*70	*	—	*	*29	*37	*	*	*33	*
721.3 . . . . .	142	51	92	—	*	*35	95	*	*29	*41	*40
722 . . . . .	1,082	562	520	—	409	392	282	247	209	431	196
722.0 . . . . .	126	73	53	—	54	52	*20	24	25	61	*16
722.1 . . . . .	655	354	302	—	277	243	134	153	132	255	115
722.10 . . . . .	642	353	290	—	276	234	133	153	131	252	107
722.5 . . . . .	119	41	77	—	*21	*29	68	*	*26	*44	*
722.52 . . . . .	115	40	75	—	*21	*28	66	*	*25	*44	*
722.7 . . . . .	68	49	*	—	*21	*	*	*	*	*	*27
722.71 . . . . .	*40	*29	*	—	*	*	*	*	*	*	*
722.73 . . . . .	*22	*	*	—	*	*	*	*	*	*	*
722.8 . . . . .	*24	*	*	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
722.83	*23	*	*	—	*	*	*	*	*	*	*
722.9	*66	*	*44	—	*	*	*	*	*	*	*
722.93	*35	*	*	—	*	*	*	*	*	*	*
723	68	48	*20	*	*18	*36	*	*30	*	*24	*
723.4	*33	*27	*	—	*	*	*	*	*	*	*
724	470	173	297	*	97	107	265	105	124	174	67
724.0	211	83	128	—	*	47	157	52	50	73	37
724.02	192	74	118	—	*	43	143	49	47	65	*31
724.2	128	*44	84	*	46	*33	*49	*	*33	63	*
724.4	49	*	*24	—	*	*	*	*	*	*	*
724.5	38	*	*31	*	*	*	*	*	*	*	*
726	90	48	42	*	25	42	23	30	17	25	18
726.1	25	*14	*11	—	*	*15	*	*	*	*	*
726.10	22	*13	*9	—	*	*14	*	*	*	*	*
726.2	23	11	*12	—	*	15	*	*	*	*9	*
726.6	*17	*	*	*	*	*	*	*	*	*	*
727	68	37	31	*	27	*17	*19	17	*14	*29	*
727.0	31	*	*15	*	*17	*	*	*	*	*	*
727.1	*7	*	*	—	*	*	*	*	*	*	*
728	126	*56	70	*	*16	*	*75	*25	*50	*37	*
728.8	41	*	*	*	*	*	*	*	*	*	*
728.89	*37	*	*	*	*	*	*	*	*	*	*
729	132	46	86	*	*23	53	52	*	*23	63	*21
729.1	47	*	*31	*	*	*	*	*	*	*34	*
729.5	30	*	*19	*	*	*	*	*	*	*	*
729.8	*16	*	*	*	*	*	*	*	*	*	*
730	584	322	262	*32	117	128	307	186	86	254	*58
730.0	200	126	*74	*	*	*	*118	*	*	*86	*
730.07	108	*	*	*	*	*	*	*	*	*	*
730.1	93	*54	*	*	*	*	*	*	*	*	*
730.2	281	132	149	*	*49	*72	146	*106	*	125	*
730.27	95	*58	*	*	*	*	*	*	*	*	*
730.28	*99	*	*	*	*	*	*	*	*	*	—
732	*15	*	*	*	*	*	—	*	*	*	*
733	939	280	659	*	96	176	659	213	163	383	180
733.1	598	120	478	*	*	*79	509	142	112	221	123
733.13	342	56	286	—	*	*	316	*67	71	144	59
733.14	147	*	105	*	*	*	117	*	*	*	*
733.15	*42	*	*	*	*	*	*	*	*	*	*
733.19	*27	*	*	—	*	*	*	*	*	*	*
733.4	106	*47	58	—	*36	*	*44	*	*	*44	*26
733.42	82	*38	*44	—	*30	*	*	*	*	*31	*
733.6	58	*20	39	—	*18	*20	*20	*	*	32	*
733.8	141	88	52	*	28	46	63	*24	*	78	*18
733.81	*21	*	*	*	*	*	*	*	*	*	*
733.82	120	73	46	*	*20	*41	56	*22	*	69	*
735	*10	*	*8	*	*	*	*	*	*	*	*
735.0	*7	*	*	—	*	*	*	*	*	*	*
736	*28	*18	*	*	*	*	*	*	*	*	*
737	84	*	57	*	*	*	*	*	*	*	*
737.3	67	*	54	*	*	*	*	*	*	*	*
737.30	*44	*	*38	*	*	*	*	*	*	*	*
738	75	*34	*41	*	*21	*29	*	*	*	*30	*
738.4	60	*28	*32	—	*	*27	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
740-759 . . . . .	999	552	447	830	101	36	*32	359	186	201	253
742 . . . . .	*40	*	*	*	*	—	*	*	*	*	*
745 . . . . .	225	139	86	201	*	*	*	79	*	*29	*90
745.2 . . . . .	*46	*	*	*44	*	*	—	*	*	*	*
745.4 . . . . .	*37	*	*	*	*	*	—	*	*	*	*
745.5 . . . . .	*59	*	*	*	*	*	*	*	*	*	*
746 . . . . .	122	*63	*60	112	*	*	*	*62	*	*	*
747 . . . . .	122	*68	*53	93	*	*	*	*35	*	*	*
747.0 . . . . .	*20	*	*	*	*	*	—	*	*	*	*
749 . . . . .	*18	*12	*	*17	*	—	—	*	*	*	*
750 . . . . .	48	*33	*	47	*	*	—	*	*	*	*
750.5 . . . . .	*21	*15	*	*21	—	—	—	*	*	*	*
751 . . . . .	107	*46	*	*93	*	*	*	*	*	*	*
752 . . . . .	*19	*	*	*	*	*	*	*	*	*	*
753 . . . . .	*38	*	*	*28	*	*	*	*	*	*	*
754 . . . . .	28	*16	*	*22	*	*	—	*	*	*	*
756 . . . . .	81	*39	*43	*55	*	*	*	*	*	*	*
759 . . . . .	*21	*	*	*	*	*	*	*	*	*	*
760-779 . . . . .	1,360	824	535	1,355	*	—	—	202	401	371	386
765 . . . . .	462	279	*182	461	*	—	—	*	*	*165	*
765.1 . . . . .	349	*221	*127	348	*	—	—	*	*	*137	*
769 . . . . .	334	*199	*135	332	*	—	—	*	*	*	*156
770 . . . . .	224	135	89	223	*	—	—	*37	*101	*54	*32
770.8 . . . . .	66	*35	*31	65	*	—	—	*	*	*28	*
771 . . . . .	78	*46	*32	78	—	—	—	*	*	*	*
771.8 . . . . .	70	*42	*	70	—	—	—	*	*	*	*
774 . . . . .	85	59	26	85	—	—	—	*	24	*13	36
774.6 . . . . .	74	52	22	74	—	—	—	*	19	*10	36
775 . . . . .	*23	*20	*	*23	—	—	—	*	*	*16	*
775.5 . . . . .	*15	*	*	*15	—	—	—	*	*	*	*
778 . . . . .	*23	*	*	*23	—	—	—	*	*	*	*
778.4 . . . . .	*18	*	*	*18	—	—	—	*	*	*	*
779 . . . . .	*32	*	*	*31	*	—	—	*	*	*	*
780-799 . . . . .	922	370	551	125	259	223	315	228	151	453	90
780 . . . . .	159	65	94	58	37	19	45	49	27	62	20
780.2 . . . . .	33	*18	*14	*	*	*	*19	*	*	*12	*
780.3 . . . . .	63	27	36	29	21	*	*	*25	*10	20	*
780.6 . . . . .	28	*	*17	*20	*	*	*	*	*	*	*
784 . . . . .	*19	*	*	*	*	*	*	*	*	*	*
785 . . . . .	*16	*	*	*	*	*	*	*	*	*	*
786 . . . . .	184	79	105	*20	58	61	45	28	32	106	17
786.0 . . . . .	*24	*	*14	*18	*	*	*	*	*	*	*
786.09 . . . . .	*23	*	*	*18	*	*	*	*	*	*	*
786.5 . . . . .	130	62	68	*	56	55	19	26	27	65	13
786.50 . . . . .	113	53	60	*	52	46	16	22	23	57	11
786.52 . . . . .	*12	*7	*	—	*	*	*	*	*	*	*
787 . . . . .	*11	*	*	*	*	*	*	*	*	*	*
788 . . . . .	*16	*	*	—	*9	*	*	*	*	*	*
788.0 . . . . .	*9	*	*	—	*	*	*	*	*	*	—
789 . . . . .	110	28	82	*9	70	*17	*14	18	23	52	*17
789.0 . . . . .	102	25	77	*9	67	*14	*	*17	22	47	*16
789.00 . . . . .	38	*9	29	*	24	*	*	*	*	*20	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
789.01	*19	*	*	*	*	*	*	*	*	*	*
789.03	23	*	*14	*	*16	*	*	*	*	*	*
799	361	154	207	*14	63	112	172	98	56	187	*
799.9	357	154	204	*14	63	112	168	98	56	183	*
800-999	14,576	6,902	7,673	974	4,242	2,635	6,724	3,646	3,066	5,233	2,630
800	54	35	*	*23	*	*	*	*	*	*	*
800.0	*14	*	*	*12	*	*	*	*	*	*	*
801	159	107	*52	*	75	*	*	*42	*	*55	*38
801.0	38	22	*	*	*23	*	*	*	*	*	*
802	160	131	29	*	124	*	*	*26	*43	54	36
802.0	*15	*	*	*	*	*	*	*	*	*	*
802.2	46	*35	*	*	41	*	*	*	*	*	*
802.4	37	*28	*	*	*29	*	*	*	*	*12	*
805	375	195	180	*	118	58	190	76	110	133	55
805.0	77	*55	*	*	*37	*	*	*	*	*	*
805.2	102	*48	53	*	*	*	58	*	*32	*	*
805.4	180	91	90	*	*50	*	99	*	53	78	*
806	*118	*	*	*	*	*	*	*	*	*	*
807	249	110	140	*	68	*55	126	60	49	73	66
807.0	213	98	115	*	61	*36	114	*51	*42	58	*61
807.01	*32	*	*	—	*	*	*	*	*	*	*
807.04	*25	*	*	*	*	*	*	*	*	*	*
807.09	*52	*	*	—	*	*	*	*	*	*	*
807.2	*26	*	*	*	*	*	*	*	*	*	*
808	445	236	209	*	141	*82	217	75	89	186	95
808.0	148	*115	*	*	*93	*	*	*	*	*	*
808.2	196	69	127	*	*	*	125	*35	*41	*51	69
808.4	*41	*	*	—	*	*	*25	*	*	*	*
808.8	*45	*	*	*	*	*	*34	*	*	*	*
810	*25	*	*	*	*	*	*	*	*	*	*
810.0	*23	*	*	*	*	*	*	*	*	*	*
812	277	78	200	21	37	35	184	67	52	89	69
812.0	169	49	120	*	*	*18	134	43	*36	48	*42
812.00	56	*	*46	*	*	*	*47	*	*	*	*
812.01	76	*	46	*	*	*	58	*	*	*26	*
812.09	*31	*	*	*	*	*	*26	*	*	*	*
812.2	*26	*	*20	*	*	*	*	*	*	*	*
812.21	*20	*	*16	*	*	*	*	*	*	*	*
812.4	64	*13	51	15	*13	*	*31	*12	*	*30	*11
812.40	*15	*	*14	*	*	*	*	*	*	*	*
812.41	39	*	*33	*12	*	*	*	*	*	*	*
813	161	81	79	*15	72	30	43	50	37	58	*16
813.0	27	*	*16	*	*	*	*	*	*	*	*
813.01	*14	*	*	*	*	*	*	*	*	*	*
813.2	*21	*	*	*	*	*	*	*	*	*	*
813.4	51	21	30	*	*13	*	*23	*19	*12	*17	*
813.42	27	*14	*	*	*	*	*	*	*	*10	*
815	*14	*	*	*	*	*	*	*	*	*	*
816	*31	*16	*	*	*	*	*	*	*	*	*
816.1	*28	*13	*	*	*	*	*	*	*	*	*
820	2,519	607	1,911	*	*48	149	2,308	594	559	939	427
820.0	787	182	605	*	*	*	742	203	197	238	148
820.02	*43	*	*	—	*	*	*	*	*	*	*
820.03	*83	*	*79	—	*	*	*81	*	*	*	*
820.09	616	148	469	*	*	*	584	173	175	198	70

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
820.2	1,145	257	888	*	*	109	1,007	309	239	430	168
820.20	*44	*	*	*	*	*	*	*	*	*	*
820.21	1,005	224	780	*	*	84	909	275	204	366	161
820.22	*97	*	*72	*	*	*	*66	*	*	*	*
820.8	575	164	411	*	*	*	549	79	117	268	111
821	570	280	290	126	134	*75	235	138	146	230	*56
821.0	339	222	117	110	95	*	*89	80	*84	145	*31
821.00	*90	*	*	*	*	*	*	*	*	*	*
821.01	249	158	*91	*88	70	*	*	*64	*76	*81	*
821.2	193	*40	153	*	*	*	142	*48	*56	75	*
821.21	*39	*	*	—	*	*	*	*	*	*	*
821.23	101	*	*86	*	*	*	*74	*	*	*	*
822	85	*31	54	*	*18	*	*41	*	*	*35	*23
822.0	66	*28	38	*	*16	*	*41	*	*	*	*22
823	390	239	151	*26	155	100	109	106	65	155	64
823.0	130	46	84	*	*33	*31	*62	*37	*27	*46	*
823.00	106	*38	68	*	*27	*27	*51	*	*	*40	*
823.2	73	*53	*	*	*30	*	*	*	*	*	*
823.22	*55	*44	*	*	*28	*	*	*	*	*	*
823.3	*66	*54	*	*	*46	*	*	*	*	*	*
823.32	*58	*	*	*	*	*	*	*	*	*	*
823.8	63	43	*	*	*26	*24	*	*	*	*22	*
823.80	*27	*17	*	*	*	*	*	*	*	*	*
823.82	*35	*25	*	*	*	*	*	*	*	*	*
824	368	157	211	*	164	101	86	87	92	133	55
824.0	*13	*	*	*	*	*	*	*	*	*	*
824.2	34	*	*18	*	*15	*	*	*	*	*15	*
824.4	80	36	45	*	41	*19	*	*21	*19	*31	*
824.6	82	*21	61	—	27	35	*	*22	*27	*18	*15
824.8	116	52	64	*	48	*25	*	*28	*30	48	*
825	85	56	*29	*	47	*20	*	*	*19	*28	*
825.0	*43	*36	*	*	*	*	*	*	*	*	*
825.2	37	*	*21	*	*	*	*	*	*	*	*
825.25	*20	*	*	*	*	*	*	*	*	*	*
831	*10	*	*	*	*	*	*	*	*	*	*
831.0	*10	*	*	*	*	*	*	*	*	*	*
836	48	34	*14	*	33	*8	*	15	*	*17	*
836.0	25	*20	*	*	*18	*	*	*9	*	*	*
836.1	*8	*	*	*	*	*	*	*	*	*	*
840	79	46	34	*	*14	34	31	25	19	24	*11
840.4	71	39	32	—	*7	34	30	24	17	23	*
844	67	35	32	*	35	*	*	*12	*24	*20	*
844.2	26	*17	*	*	23	*	—	*	*	*	*
845	*19	*	*	*	*	*	*	*	*	*	*
845.0	*18	*	*	—	*	*	*	*	*	*	*
846	*40	*	*	—	*	*	*	*	*	*	*
846.0	*38	*	*	—	*	*	*	*	*	*	*
847	58	*26	32	*	*25	*17	*	*	*15	*	*
847.0	*26	*	*	*	*	*	*	*	*	*	*
847.2	*27	*	*	—	*	*	*	*	*	*	*
850	144	104	40	*8	62	*	36	76	26	34	*
850.0	26	*14	*	*	*	*	*	*	*	*	*
850.1	56	*46	*9	*	*15	*	*	*40	*	*	*
850.5	*21	*17	*	*	*18	*	*	*	*	*	*
850.9	32	*17	*	*	*13	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
851 . . . . .	289	*194	*95	*	*149	*	*	*	*	*	*
851.8 . . . . .	208	*174	*	*	*	*	*	*	*	*	*
852 . . . . .	274	168	106	*	*76	*49	113	*61	*45	*55	114
852.0 . . . . .	*61	*45	*	*	*	*	*	*	*	*	*
852.2 . . . . .	208	120	*88	*	*61	*	90	*36	*	*	103
852.20 . . . . .	*51	*	*	*	*	*	*	*	*	*	*
852.21 . . . . .	107	*64	*	*	*	*	*46	*	*	*	*52
853 . . . . .	*73	*	*	*	*	*	*	*	*	*	*
853.0 . . . . .	*73	*	*	*	*	*	*	*	*	*	*
854 . . . . .	210	101	109	33	109	*29	*40	31	48	62	69
854.0 . . . . .	203	95	108	32	102	*29	*40	*30	46	62	65
854.00 . . . . .	59	37	*22	*	*18	*	*	*	*	*24	*
854.01 . . . . .	30	*11	*19	*	*9	*	*	*	*	*	*
854.02 . . . . .	35	23	*	*	*20	*	*	*	*	*	*
854.06 . . . . .	*64	*	*	*	*53	*	*	*	*	*	*
860 . . . . .	199	112	88	*	87	*	*76	*	*48	68	*
860.0 . . . . .	107	*55	*52	*	*34	*	*	*	*	*	*
860.4 . . . . .	*42	*	*	—	*	*	*	*	*	*	*
861 . . . . .	88	71	*	*	*48	*	*	*	*	*42	*
861.2 . . . . .	*39	*	*	*	*	*	*	*	*	*	*
861.21 . . . . .	*37	*	*	*	*	*	*	*	*	*	*
863 . . . . .	118	*70	*	*	*80	*	*	*	*	*	*
864 . . . . .	63	*40	*	*	*37	*	*	*	*	*	*
864.0 . . . . .	*49	*	*	*	*	*	*	*	*	*	*
865 . . . . .	119	87	*	*	94	*	*	*	*	*45	*
865.0 . . . . .	89	59	*	*	69	*	*	*	*	*	*
866 . . . . .	*40	*	*	*	*	*	*	*	*	*	*
871 . . . . .	*36	*	*	*	*21	*	*	*	*	*	*
873 . . . . .	72	32	40	*	30	*	*	*15	*16	*18	*
873.0 . . . . .	*14	*	*	*	*	*	*	*	*	*	*
873.4 . . . . .	39	*13	*26	*	*15	*	*	*	*	*	*
873.42 . . . . .	*24	*	*	*	*	*	*	*	*	*	*
874 . . . . .	*14	*	*	*	*	*	*	*	*	*	*
875 . . . . .	33	31	*	*	*28	*	*	*	*	*29	*
875.0 . . . . .	*21	*20	*	*	*18	*	*	*	*	*18	*
879 . . . . .	31	*20	*	*	*27	*	*	*	*	*	*
881 . . . . .	*27	*17	*	*	*23	*	*	*	*	*	*
882 . . . . .	*28	*22	*	*	*21	*	*	*	*	*	*
883 . . . . .	26	*20	*	*	20	*	*	*	*	*	*
886 . . . . .	*10	*10	*	*	*	*	*	*	*	*	*
891 . . . . .	58	46	*	*	*37	*	*	*	*	*	*
891.0 . . . . .	*26	*21	*	*	*	*	*	*	*	*	*
892 . . . . .	*41	*35	*	*	*	*	*	*	*	*	*
892.1 . . . . .	*37	*	*	*	*	*	*	*	*	*	*
920 . . . . .	*20	*	*	*	*	*	*	*	*	*	*
922 . . . . .	68	*22	46	*	*13	*	47	*	*22	*	*
922.1 . . . . .	31	*	*22	*	*	*	*	*	*	*	*
924 . . . . .	93	*42	51	*	*	*	69	*	*	*30	*29
924.0 . . . . .	67	*	*37	*	*	*	59	*	*	*	*
924.01 . . . . .	54	*	*30	—	*	*	53	*	*	*	*
924.8 . . . . .	*13	*	*	—	*	*	*	*	*	*	*
935 . . . . .	24	*16	*	*	*	*	*	*	*	*	*
935.1 . . . . .	*19	*15	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
941	*29	*	*	*	*	*	*	*	*	*	*
942	*81	*	*	*	*	*	*	*	*	*	*
945	90	*71	*	*	*40	*	*	*	*	*	*
945.2	*30	*	*	*	*	*	*	*	*	*	*
946	214	*121	*92	*	*112	*	*	*	*	*119	*
946.2	*41	*	*	*	*	—	*	*	*	*	*
952	*63	*	*	*	*	*	*	*	*	*	*
958	*44	*23	*	*	*	*	*	*	*	*	*
959	58	27	31	*	34	*	*	*	*	*29	*
959.1	20	*	*14	*	*9	*	*	*	*	*	*
963	*8	*	*	*	*	—	*	*	*	*	*
965	123	51	72	*11	91	*	*	43	25	33	*22
965.0	*27	*21	*	*	*18	*	*	*	*	*	*
965.1	*17	*	*16	*	*	*	*	*	*	*	*
965.4	56	*23	33	*	45	*	*	*	*13	*18	*
965.6	*10	*	*	*	*	*	*	*	*	*	*
966	36	*	*19	*	*	*	*	*	*	*	*
966.3	*12	*	*	*	*	*	*	*	*	*	*
967	*14	*	*	*	*	*	*	*	*	*	*
968	*17	*13	*	*	*14	*	—	*	*	*	*
968.5	*9	*	*	—	*	*	—	*	*	*	*
969	144	44	100	*20	80	34	*	29	26	61	29
969.0	69	22	47	*	39	*	*	*13	*13	35	*
969.4	40	*15	24	*	21	*	*	*	*	*15	*
970	*11	*	*	—	*11	—	—	*	*	*	—
970.8	*11	*	*	—	*11	—	—	*	*	*	—
972	*31	*	*17	*	*	*	*	*	*	*	*
977	32	*	21	*	23	*	*	*	*	*13	*
977.8	*13	*	*	*	*	*	*	*	*	*	*
977.9	*16	*	*	*	*14	—	*	*	*	*8	*
989	*19	*	*	*	*	*	*	*	*	*	*
989.5	*14	*	*	*	*	*	*	*	*	*	*
992	*17	*	*	*	*	*	*	*	*	*	*
995	77	*24	53	*14	*14	*	*35	*	*26	*24	*
995.1	*19	*	*	*	*	*	*	*	*	*	*
995.2	*32	*	*23	*	*	*	*	*	*	*	*
996	2,319	1,030	1,289	135	472	599	1,113	656	427	879	357
996.0	91	49	42	*	*	*	64	*42	*	*21	*17
996.01	54	*25	28	*	*	*	42	*	*	*	*12
996.1	53	*29	*24	*	*	*	*28	*	*	*	*
996.2	68	38	*30	*26	*33	*	*	*	*	*25	*
996.4	409	152	258	*	57	91	258	112	76	150	71
996.5	74	*48	*26	*	*	*	*	*	*	*35	*
996.59	35	*	*21	*	*	*	*	*	*	*	*
996.6	865	369	496	*68	160	216	421	275	190	296	105
996.62	424	158	266	*	103	89	192	145	102	131	*45
996.64	*44	*	*	*	*	*	*	*	*	*	*
996.66	152	*63	*89	*	*	*	*107	*	*	*55	*
996.67	*59	*	*	*	*	*	*	*	*	*	*
996.69	107	*46	*62	*	*	*38	*	*	*	*	*
996.7	544	234	310	*	60	163	307	136	90	204	114
996.72	*37	*	*	*	*	*	*	*	*	*	*
996.73	200	93	106	*	26	68	106	39	*32	86	42
996.74	190	88	102	*	*	52	123	*61	*	58	*46

\* Figure does not meet standard of reliability or precision.  
 — Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).

**Table 2. Number of days of care for discharges from short-stay hospitals, by ICD-9-CM code of first-listed diagnosis, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of days of care in thousands											
996.77	*36	*	*	—	*	*	*	*	*	*	*
996.78	30	*	*20	*	*11	*	*	*	*	*	*
996.8	172	106	*67	*	93	*65	*	*	*	107	*
996.81	110	*61	*49	*	*63	*42	*	*	*	71	*
997	514	226	288	*	87	140	272	140	115	142	118
997.1	31	*	*23	*	*	*	*	*	*	*	*
997.2	*52	*	*38	—	*	*	*	*	*	*	*
997.3	64	*32	*	*	*	*	*	*	*	*	*
997.4	189	75	114	*	*29	66	86	*43	50	*54	42
997.5	47	*35	*	*	*	*	*31	*	*	*	*
997.6	106	*55	*51	*	*	*	*62	*	*	*	*
997.62	*77	*	*	*	*	*	*48	*	*	*	*
998	1,112	458	654	26	248	313	525	292	191	395	234
998.1	147	69	79	*	34	50	60	42	*28	45	32
998.2	*63	*	*31	—	*	*	*	*	*	*	*
998.3	*49	*	*34	*	*	*	*	*	*	*	*
998.5	751	307	445	*	174	193	373	216	126	255	155
998.8	65	*24	*41	*	*	*	*	*	*	*	*
998.89	51	*22	*29	*	*	*	*	*	*	*	*
999	*27	*	*	*	*	*	*	*	*	*	*
V01-V82	11,280	1,389	9,891	358	8,379	657	1,887	2,428	3,150	3,502	2,200
V10	*11	*	*	*	*	*	*	*	*	*	*
V12	*14	*	*	*	—	*	*	*	*	*	*
V22	42	—	42	—	42	—	—	*	*12	*20	*9
V22.2	36	—	36	—	36	—	—	*	*11	*18	*
V24	*11	—	*11	*	*11	—	—	*	*	*	*
V24.0	*11	—	*11	*	*11	—	—	*	*	*	*
V25	*8	*	*7	—	*8	—	—	*	*	*	*
V25.2	*7	*	*7	—	*7	—	—	*	*	*	*
V27	8,033	—	8,033	27	8,000	*	—	1,543	1,726	2,960	1,804
V27.0	5,855	—	5,855	22	5,829	*	—	1,103	1,192	2,267	1,293
V27.1	58	—	58	*	58	—	—	*	*	27	*
V27.2	162	—	162	—	162	*	—	*	*54	56	28
V27.9	1,948	—	1,948	*	1,942	*	—	394	472	604	477
V30	155	*90	*65	148	—	—	*	*65	*	*70	*
V30.1	148	*90	*58	148	—	—	—	*65	*	*70	*
V45	*22	*	*	*	*	*	*	*	*	*	*
V45.8	*14	*	*	*	*	*	*	*	*	*	*
V54	*16	*	*	*	*	*	*	*	*	*	*
V54.0	*11	*	*	*	*	*	*	*	*	*	*
V55	110	*54	*56	*	*	*	*	*40	*	*	*
V55.3	85	*	*46	*	*	*	*	*	*	*	*
V57	2,135	879	1,256	*	*143	383	1,573	462	1,159	270	244
V57.1	590	258	332	*	*	*94	430	278	161	*108	*
V57.8	1,533	616	916	*	*103	289	1,132	*180	993	160	200
V57.89	1,533	616	916	*	*103	289	1,132	*180	993	160	200
V58	468	225	243	64	74	168	162	203	147	76	43
V58.0	*31	*	*	*	*	*	*	*	*	*	*
V58.1	404	189	215	63	63	155	123	184	111	70	*38
V70	*33	*	*	*	*	*	*	*	*	*	*
V70.7	*32	*	*	*	*	*	*	*	*	*	*
V71	38	20	18	*12	*13	*	*	*13	*12	*	*
V71.8	20	*	*11	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: If estimate preceded by asterisk, use according to size of corresponding estimate in table 1; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
All codes . . . . .	118,255	47,703	70,552	6,100	32,357	24,198	55,600	27,146	28,479	42,352	20,278
001-139 . . . . .	3,473	1,585	1,887	537	985	603	1,348	818	760	1,286	609
003 . . . . .	18	10	*8	*6	*8	*	*	*5	*	*6	*
003.0 . . . . .	13	*7	*6	*5	*6	*	*	*	*	*	*
008 . . . . .	239	96	142	80	38	28	92	64	53	80	41
008.4 . . . . .	94	33	61	*5	11	16	61	32	24	24	14
008.43 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
008.45 . . . . .	77	26	51	*	9	11	52	28	20	20	9
008.47 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
008.6 . . . . .	37	21	16	36	*	*	*	*5	10	14	*9
008.61 . . . . .	34	20	15	34	*	*	—	*	9	13	*8
008.8 . . . . .	103	39	64	36	25	11	30	26	20	40	16
009 . . . . .	17	*8	*9	*	*6	*	*	*	*	*6	*
009.0 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
011 . . . . .	30	23	*7	*	*8	11	*9	*	*	18	*
011.9 . . . . .	25	19	*6	*	*7	9	*7	*	*	16	*
011.90 . . . . .	17	13	*	*	*	*7	*	*	*	13	*
031 . . . . .	21	14	*7	*	13	*	*	*6	*	*9	*
031.0 . . . . .	13	*9	*	*	*8	*	*	*	*	*7	*
034 . . . . .	34	15	19	19	12	*	*	11	*7	11	*5
034.0 . . . . .	33	15	18	18	12	*	*	10	*6	11	*5
038 . . . . .	588	275	312	40	72	116	360	132	120	235	100
038.0 . . . . .	50	25	25	*5	*7	11	27	10	13	22	*5
038.1 . . . . .	121	59	62	*6	21	27	67	32	26	45	18
038.2 . . . . .	16	*8	*8	*	*	*	*9	*	*	*6	*
038.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
038.4 . . . . .	141	58	83	*	17	27	93	34	31	49	27
038.40 . . . . .	14	*6	*7	*	*	*	*7	*	*	*	*
038.42 . . . . .	74	25	49	*	*9	15	49	16	17	25	15
038.43 . . . . .	14	*9	*5	*	*	*	*8	*	*	*	*
038.49 . . . . .	37	17	20	*	*	*6	28	*9	*6	16	*6
038.8 . . . . .	32	15	17	*	*	*9	17	12	*	9	*7
038.9 . . . . .	222	108	114	20	19	40	143	40	41	103	38
041 . . . . .	1,190	442	748	92	273	209	615	236	313	435	205
041.0 . . . . .	197	66	131	12	74	31	80	43	47	70	37
041.00 . . . . .	51	19	32	*	13	*8	28	14	12	16	*9
041.01 . . . . .	16	10	*6	*	*7	*	*	*	*	*5	*5
041.02 . . . . .	54	*7	47	*	38	*6	*9	12	12	19	11
041.04 . . . . .	40	17	24	*	*6	*6	28	*	12	18	*6
041.09 . . . . .	34	13	21	*	10	*8	13	10	*9	10	*6
041.1 . . . . .	218	117	101	20	58	51	89	42	57	78	41
041.10 . . . . .	32	14	18	*	9	*6	14	*8	*8	13	*
041.11 . . . . .	144	82	62	14	37	33	60	26	34	52	31
041.19 . . . . .	42	21	22	*	12	11	16	*8	14	13	*7
041.3 . . . . .	85	24	61	*	11	14	56	18	22	32	13
041.4 . . . . .	344	85	259	31	66	49	198	69	91	120	64
041.5 . . . . .	19	*9	10	*	*	*	10	*	*6	*6	*
041.6 . . . . .	49	16	33	*	*5	*6	37	11	13	21	*
041.7 . . . . .	96	45	51	*7	15	17	58	14	26	45	11
041.8 . . . . .	158	69	89	11	37	33	77	29	49	52	28
041.84 . . . . .	17	9	*8	*	*6	*	*6	*	*7	*	*
041.85 . . . . .	88	38	50	*	19	18	47	17	27	30	14
041.89 . . . . .	50	20	30	*	11	9	25	10	13	17	10
041.9 . . . . .	20	*8	11	*	*	*	10	*	*	11	*
042 . . . . .	216	160	55	*8	163	42	*	77	22	78	38

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
047 . . . . .	40	20	20	22	15	*	*	*8	*6	18	*8
047.9 . . . . .	39	20	19	21	14	*	*	*8	*6	17	*8
052 . . . . .	20	12	*7	13	*	*	*	*7	*	*	*6
052.7 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
052.9 . . . . .	11	*8	*	*9	*	*	*	*	*	*	*
053 . . . . .	54	23	30	*	*	12	37	14	11	19	10
053.1 . . . . .	16	*6	9	—	*	*	13	*	*	*6	*
053.19 . . . . .	14	*5	*9	—	*	*	11	*	*	*6	*
053.9 . . . . .	32	14	18	*	*	*9	20	*8	*6	11	*7
054 . . . . .	56	22	34	*7	28	*8	12	14	10	20	12
054.1 . . . . .	14	*	10	*	11	*	*	*	*	*	*
054.10 . . . . .	*8	*	*6	*	*7	*	*	*	*	*	*
054.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
054.7 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
054.9 . . . . .	25	9	16	*	12	*	*6	*6	*	9	*5
057 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
070 . . . . .	91	51	40	*	49	25	16	22	11	38	20
070.3 . . . . .	24	15	*9	*	13	*6	*	*7	*	9	*
070.30 . . . . .	13	*8	*5	—	*8	*	*	*	*	*	*
070.32 . . . . .	10	*7	*	*	*5	*	*	*	*	*	*
070.5 . . . . .	53	30	23	*	30	14	9	12	*7	22	12
070.51 . . . . .	24	14	10	*	14	*6	*	*7	*	*8	*
070.54 . . . . .	29	16	13	—	15	*8	*6	*5	*	14	*8
070.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
075 . . . . .	13	*7	*6	*	*9	*	*	*	*	*6	*
078 . . . . .	46	24	22	*6	31	*7	*	14	10	15	*7
078.1 . . . . .	14	*	11	*	12	*	*	*	*	*	*
078.10 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
078.11 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
078.5 . . . . .	24	17	*7	*	17	*	*	*8	*	*8	*
078.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
078.89 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
079 . . . . .	223	112	111	160	39	12	12	48	49	86	40
079.8 . . . . .	98	61	36	94	*	*	*	17	22	38	20
079.89 . . . . .	94	60	34	93	*	*	*	15	22	37	20
079.9 . . . . .	120	49	71	64	33	12	12	30	25	46	19
079.99 . . . . .	115	49	67	63	29	12	12	29	24	44	19
088 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
088.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
088.81 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
094 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
097 . . . . .	12	*	*8	—	*8	*	*	*	*	*6	*
097.1 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
098 . . . . .	9	*	*8	*	*8	*	—	*	*	*5	*
110 . . . . .	43	24	19	*	11	*7	23	15	14	9	*5
110.1 . . . . .	20	11	9	*	*	*	13	*8	*8	*	*
110.4 . . . . .	13	*7	*6	*	*	*	*6	*5	*	*	*
111 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
112 . . . . .	257	108	149	40	73	47	97	68	57	91	42
112.0 . . . . .	128	63	65	25	42	26	35	40	29	40	18
112.1 . . . . .	19	—	19	*	9	*	*7	*	*	*8	*
112.2 . . . . .	30	10	19	*	*	*	21	*7	*7	10	*6
112.3 . . . . .	16	*7	9	*8	*	*	*7	*	*	*6	*
112.4 . . . . .	10	*	*5	—	*	*	*6	*	*	*5	*
112.8 . . . . .	41	19	23	*	14	*9	16	11	*7	17	*6

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
112.84	27	12	16	*	9	*7	11	*8	*	11	*
112.89	12	*6	*6	*	*	*	*	*	*	*5	*
112.9	*9	*	*6	*	*	*	*	*	*	*	*
117	19	11	*8	*	11	*	*	*	*	*7	*
117.3	*7	*	*	*	*	*	*	*	*	*	*
117.5	*7	*6	*	*	*6	*	*	*	*	*	*
130	*7	*6	*	*	*6	*	—	*	*	*	*
131	20	*	18	*	15	*	*	*	*	11	*
131.0	18	*	17	*	14	*	*	*	*	10	*
131.01	11	—	11	*	*9	*	*	*	*	*6	*
135	35	*9	26	*	13	13	9	*7	11	12	*5
136	42	30	12	*	26	10	*	12	*7	13	10
136.3	34	28	*6	*	24	*8	*	12	*	11	*8
136.9	*6	*	*	*	*	*	*	*	*	*	*
138	21	14	*6	*	*	*5	15	*6	*5	*	*5
140-239	4,356	1,829	2,527	96	682	1,331	2,248	1,088	960	1,563	744
141	*7	*	*	—	*	*	*	*	*	*	*
150	31	23	*8	—	*	10	19	9	*	10	*6
150.5	*7	*6	*	—	*	*	*5	*	*	*	*
150.9	14	12	*	—	*	*6	*8	*	*	*5	*
151	46	25	21	—	*	14	29	10	*8	20	*8
151.0	10	*8	*	—	*	*	*6	*	*	*	*
151.8	9	*	*	—	*	*	*5	*	*	*	*
151.9	14	*5	*9	—	*	*	*9	*	*	*9	*
152	*6	*	*	—	*	*	*	*	*	*	*
153	135	67	69	*	*	24	107	32	43	42	18
153.0	*5	*	*	—	—	*	*	*	*	*	*
153.1	10	*6	*	—	*	*	*8	*	*	*	*
153.2	*8	*	*	—	*	*	*6	*	*	*	*
153.3	30	13	17	—	*	*6	23	*9	*8	9	*
153.4	18	10	*9	—	*	*	16	*	*5	*6	*
153.6	16	*6	10	—	*	*	13	*	*6	*	*
153.8	10	*	*7	—	*	*	*7	*	*	*	*
153.9	32	18	14	*	*	*7	24	*7	11	10	*
154	64	33	30	—	*	18	41	20	14	17	12
154.0	22	12	10	—	*	*	15	*6	*	*8	*
154.1	35	19	16	—	*	12	22	13	*9	*7	*6
155	23	16	*7	*	*	*6	15	*	*	*8	*6
155.0	15	11	*	*	*	*	9	*	*	*	*
155.2	*5	*	*	—	—	*	*	*	*	*	*
156	15	*8	*7	—	*	*5	9	*	*5	*	*
156.1	*7	*5	*	—	—	*	*	*	*	*	*
157	50	25	25	—	*	15	33	12	*8	19	10
157.0	17	*9	*8	—	*	*	13	*	*	*7	*
157.8	*9	*	*5	—	*	*	*6	*	*	*	*
157.9	20	11	*9	—	*	*6	13	*	*	*8	*
158	10	*	*8	*	*	*	*	*	*	*	*
158.8	*5	*	*5	—	*	*	*	*	*	*	*
161	20	13	*7	—	*	9	10	*	*	10	*
161.9	*9	*5	*	—	*	*	*5	*	*	*6	*
162	359	205	154	*	15	123	221	90	81	136	52
162.2	20	10	9	—	*	*5	14	*	*	*8	*
162.3	84	45	39	—	*	27	53	22	21	28	12
162.4	*6	*	*	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
162.5	35	20	15	—	*	13	21	10	*7	13	*5
162.8	38	23	15	*	*	13	24	20	*	*9	*
162.9	175	102	73	*	10	63	103	32	42	76	25
164	*6	*	*	*	*	*	*	*	*	*	*
170	16	*9	*7	*	*6	*	*	*	*5	*	*
170.7	*5	*	*	*	*	*	*	*	*	*	*
171	19	*9	11	*	*	*8	*6	*	*	*6	*5
171.3	*5	*	*	*	*	*	*	*	*	*	*
171.5	*5	*	*	*	*	*	*	—	*	*	*
172	11	*8	*	—	*	*	*5	*	*	*	*
172.9	*5	*	*	—	*	*	*	*	*	*	*
173	30	19	12	*	*	*9	20	*7	*6	13	*
173.3	*7	*	*	*	*	*	*	*	*	*	*
173.4	*7	*6	*	*	*	*	*	*	*	*	*
174	215	—	215	—	32	82	101	57	44	79	36
174.1	*5	—	*5	—	*	*	*	*	*	*	*
174.2	*8	—	*8	—	*	*	*	*	*	*	*
174.4	28	—	28	—	*6	11	12	*7	*8	*9	*
174.5	*6	—	*6	—	*	*	*	*	*	*	*
174.8	52	—	52	—	*7	21	25	17	*9	14	13
174.9	107	—	107	—	16	43	48	23	22	49	13
176	17	16	*	—	15	*	*	*	*	*	10
176.0	*7	*7	*	—	*6	*	*	*	*	*	*
179	*5	—	*5	—	*	*	*	*	*	*	*
180	33	—	33	—	11	14	*8	11	*	17	*
180.8	10	—	10	—	*	*6	*	*6	*	*	*
180.9	20	—	20	—	*8	*6	*	*	*	12	*
182	49	—	49	—	*	21	24	11	*8	19	12
182.0	48	—	48	—	*	21	24	10	*8	19	12
183	70	—	70	*	*	28	38	20	13	27	10
183.0	70	—	70	*	*	28	37	19	13	27	10
184	*7	—	*7	—	*	*	*5	*	*	*	*
184.4	*5	—	*5	—	*	*	*	*	*	*	*
185	225	225	—	*	*	47	177	51	53	83	38
186	*6	*6	—	*	*	*	*	*	*	*	*
186.9	*6	*6	—	*	*	*	*	*	*	*	*
188	79	54	25	*	*	14	63	29	16	25	*9
188.2	*7	*	*	*	*	*	*5	*	*	*	*
188.4	*5	*	*	—	—	*	*	*	*	*	*
188.8	21	14	*7	—	*	*	17	10	*	*	*
188.9	33	25	*8	—	*	*7	25	*9	*8	13	*
189	64	35	29	*7	*	17	36	11	13	26	14
189.0	56	30	26	*7	*	14	31	10	11	23	12
191	43	26	18	*9	10	14	11	13	10	13	*8
191.1	*6	*	*	*	*	*	*	*	*	*	*
191.2	*5	*	*	*	*	*	*	*	*	*	*
191.3	*6	*	*	*	*	*	*	*	*	*	*
191.6	*6	*	*	*	*	*	—	*	*	*	*
191.9	9	*5	*	*	*	*	*	*	*	*	*
193	14	*	10	*	*6	*	*	*	*	*5	*
194	11	*	*7	*	*	*	*	*	*	*	*
194.0	*6	*	*	*	*	*	*	*	*	*	*
195	13	*7	*6	*	*	*	*9	*	*	*	*
195.2	*6	*	*	*	*	*	*5	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
196 . . . . .	192	74	119	*	22	67	102	56	47	58	31
196.0 . . . . .	26	14	12	*	*	10	11	*7	*	9	*5
196.1 . . . . .	26	13	14	*	*	9	15	9	*5	*8	*
196.2 . . . . .	64	30	34	*	*	18	41	18	14	21	10
196.3 . . . . .	44	*	40	—	*7	20	17	15	12	12	*
196.5 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
196.6 . . . . .	12	*	*8	*	*	*	*9	*	*	*	*
196.9 . . . . .	10	*	*6	—	*	*	*5	*	*	*	*
197 . . . . .	509	217	292	*6	43	164	295	128	112	180	89
197.0 . . . . .	124	53	71	*	16	41	63	28	27	51	19
197.1 . . . . .	17	11	*6	*	*	*6	10	*	*6	*	*
197.2 . . . . .	59	24	35	*	*	17	39	16	10	20	13
197.4 . . . . .	12	*	*8	—	*	*	*8	*	*	*	*
197.5 . . . . .	14	*	9	—	*	*5	*	*	*	*	*
197.6 . . . . .	65	20	45	*	*	24	38	16	13	21	16
197.7 . . . . .	198	89	109	*	15	63	118	55	46	67	31
197.8 . . . . .	16	10	*7	*	*	*	11	*	*	*6	*
198 . . . . .	482	221	260	*9	42	158	274	122	115	169	76
198.1 . . . . .	13	*6	*7	—	*	*	*9	*	*	*5	*
198.2 . . . . .	13	*	*8	—	*	*	*8	*	*	*	*
198.3 . . . . .	100	50	50	*	*9	38	49	23	25	33	18
198.4 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
198.5 . . . . .	216	109	107	*	15	63	135	56	47	84	30
198.6 . . . . .	*7	—	*7	—	*	*	*	*	*	*	*
198.7 . . . . .	17	*8	*9	*	*	*6	*8	*	*6	*	*
198.8 . . . . .	105	38	67	*	10	35	57	26	26	33	20
198.82 . . . . .	18	*	13	—	*	*7	9	*	*	*5	*6
198.89 . . . . .	85	33	52	*	*9	27	47	23	23	26	14
199 . . . . .	106	46	61	*	*8	34	64	17	19	46	24
199.0 . . . . .	23	*9	14	*	*	*7	15	*	*5	10	*
199.1 . . . . .	83	37	46	*	*6	27	49	15	14	36	19
200 . . . . .	40	23	17	*	*7	10	21	10	*8	16	*5
200.0 . . . . .	22	14	*8	*	*	*	15	*5	*	12	*
200.00 . . . . .	11	*8	*	*	*	*	*8	*	*	*7	*
200.1 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
200.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
200.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
201 . . . . .	17	*8	*9	*	10	*	*	*	*5	*	*
201.9 . . . . .	13	*6	*7	*	*7	*	*	*	*	*	*
201.90 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
202 . . . . .	114	62	52	*	18	36	59	31	22	44	17
202.0 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
202.8 . . . . .	102	57	46	*	17	31	53	26	20	41	15
202.80 . . . . .	55	29	26	*	*8	17	29	17	12	17	*8
202.81 . . . . .	10	*6	*	*	*	*	*6	*	*	*	*
202.82 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
202.83 . . . . .	13	*7	*7	—	*	*	*6	*	*	*6	*
202.88 . . . . .	12	*8	*	*	*	*	*7	*	*	*7	*
203 . . . . .	63	33	30	—	*	21	39	13	12	28	10
203.0 . . . . .	62	32	29	—	*	21	39	13	12	28	9
203.00 . . . . .	59	32	27	—	*	19	38	13	11	27	9
204 . . . . .	96	47	49	21	9	9	57	26	21	33	16
204.0 . . . . .	32	19	14	21	*8	*	*	9	*7	*9	*7
204.00 . . . . .	22	13	*9	13	*5	*	*	*6	*5	*8	*
204.01 . . . . .	10	*5	*	*7	*	*	—	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
204.1	63	28	35	—	*	*7	55	17	14	24	*9
204.10	60	28	33	—	*	*7	52	16	13	23	*9
205	43	24	19	*	10	*9	20	15	*8	15	*5
205.0	29	19	11	*	*7	*6	13	10	*6	11	*
205.00	25	17	*9	*	*6	*	13	*8	*5	10	*
205.1	13	*	*8	*	*	*	*6	*	*	*	*
205.10	12	*	*8	*	*	*	*6	*	*	*	*
208	11	*5	*6	*	*	*	*6	*	*	*6	*
208.0	*6	*	*	*	*	*	*	*	*	*	*
208.00	*5	*	*	*	*	*	*	*	*	*	*
210	*8	*	*	*	*	*	*5	*	*	*	*
210.2	*7	*	*	*	*	*	*	*	*	*	*
211	136	62	74	*	14	28	94	41	37	41	18
211.1	12	*	9	*	*	*	9	*	*	*	*
211.3	106	51	55	*	*8	24	74	31	29	34	11
211.4	*8	*	*	—	*	*	*7	*	*	*	*
212	*8	*	*	*	*	*	*	*	*	*	*
214	25	12	13	*	*6	*9	10	*7	*	11	*
214.1	13	*	*9	—	*	*	*	*	*	*7	*
214.8	*5	*	*	*	*	*	*	*	*	*	*
215	*6	*	*	*	*	*	*	*	*	*	*
216	18	*6	12	*	9	*	*	*6	*	*5	*
216.5	*9	*	*6	*	*5	*	*	*	*	*	*
217	*9	*	*8	—	*	*	*	*	*	*	*
218	402	—	402	—	227	151	24	75	87	159	80
218.0	43	—	43	—	23	18	*	12	*8	14	*8
218.1	102	—	102	—	52	41	*8	27	24	36	15
218.2	59	—	59	—	36	20	*	14	14	22	10
218.9	198	—	198	—	116	71	11	22	43	87	46
219	*7	—	*7	—	*	*	*	*	*	*	*
220	49	—	49	*	25	14	10	13	11	17	*8
225	25	*6	19	*	*	*7	13	*6	*6	10	*
225.2	17	*	13	*	*	*5	10	*	*	*7	*
226	11	*	*9	—	*5	*	*	*	*	*	*
227	18	*5	13	*	*6	*7	*	*5	*	*5	*
227.1	*7	*	*5	—	*	*	*	*	*	*	*
227.3	10	*	*6	*	*	*	*	*	*	*	*
228	15	*7	*8	*	*	*	*	*	*	*	*
228.0	13	*6	*8	*	*	*	*	*	*	*	*
230	*5	*	*	—	*	*	*	*	*	*	*
233	43	*	41	—	19	15	*9	10	10	16	*7
233.0	19	*	18	—	*	*8	*7	*6	*	*5	*
233.1	20	—	20	—	14	*	*	*	*	9	*
235	18	9	*8	*	*	*	13	*	*5	*6	*
235.2	*9	*	*	*	*	*	*6	*	*	*	*
236	10	*	*8	*	*	*	*	*	*	*	*
236.2	*5	—	*5	—	*	*	*	*	*	*	*
237	15	*7	*9	*	*	*6	*	*5	*	*6	*
237.5	*6	*	*	*	*	*	*	*	*	*	*
237.7	*7	*	*	*	*	*	*	*	*	*	*
238	62	35	27	*	*	12	45	18	14	17	13
238.4	17	10	*8	*	*	*	11	*6	*	*	*
238.7	38	24	14	*	*	*	32	11	*8	9	10

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
239 . . . . .	25	12	13	*	*6	*	13	*	*7	9	*
239.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
239.6 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
240-279 . . . . .	11,960	4,834	7,126	487	1,680	3,098	6,695	2,666	3,024	4,382	1,888
240 . . . . .	12	*	11	—	*	*	*7	*5	*	*	*
240.9 . . . . .	12	*	11	—	*	*	*7	*5	*	*	*
241 . . . . .	28	*	24	*	*7	9	12	*8	*6	9	*5
241.0 . . . . .	10	*	10	*	*	*	*	*	*	*	*
241.1 . . . . .	14	*	12	*	*	*5	*7	*	*	*5	*
242 . . . . .	62	10	52	*	15	13	34	14	15	21	12
242.0 . . . . .	17	*	15	*	*8	*	*	*	*	*7	*
242.00 . . . . .	16	*	14	*	*7	*	*	*	*	*6	*
242.9 . . . . .	40	*8	32	*	*5	*7	27	*9	10	12	*9
242.90 . . . . .	38	*8	30	*	*	*7	26	*8	10	12	*8
244 . . . . .	822	158	664	*	79	184	555	215	213	246	148
244.0 . . . . .	34	*	30	*	*	11	20	*7	*7	9	11
244.1 . . . . .	*6	*	*5	—	*	*	*	*	*	*	*
244.9 . . . . .	775	151	624	*	74	167	530	204	203	233	134
245 . . . . .	14	*	12	*	*7	*	*	*6	*	*	*
245.2 . . . . .	11	*	10	—	*7	*	*	*5	*	*	*
246 . . . . .	*6	*	*6	*	*	*	*	*	*	*	*
250 . . . . .	3,952	1,742	2,210	26	417	1,227	2,282	915	924	1,482	631
250.0 . . . . .	2,691	1,155	1,536	11	214	799	1,667	642	649	1,012	388
250.00 . . . . .	1,565	710	855	*	95	442	1,027	387	364	562	252
250.01 . . . . .	693	266	428	*7	70	213	404	171	178	256	89
250.02 . . . . .	258	111	146	*	18	86	153	53	59	118	28
250.03 . . . . .	175	68	107	*	31	58	83	32	49	76	19
250.1 . . . . .	118	54	64	10	62	28	18	26	26	40	26
250.10 . . . . .	12	*7	*6	*	*	*	*	*	*	*	*5
250.11 . . . . .	41	17	23	*	23	*6	*6	11	9	15	*6
250.12 . . . . .	11	*6	*	*	*	*	*	*	*	*5	*
250.13 . . . . .	54	24	30	*	31	13	*5	12	13	17	13
250.2 . . . . .	19	9	10	—	*	*	13	*	*	*6	*
250.20 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
250.22 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
250.23 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
250.3 . . . . .	*9	*	*7	*	*	*	*	*	*	*	*
250.4 . . . . .	297	135	162	*	39	109	149	64	53	108	73
250.40 . . . . .	109	49	60	—	*	39	66	22	20	37	29
250.41 . . . . .	128	55	73	*	22	47	58	30	20	47	31
250.42 . . . . .	21	*9	12	—	*	*8	11	*	*	12	*
250.43 . . . . .	40	23	17	*	11	15	14	*8	9	11	11
250.5 . . . . .	132	59	73	—	25	52	56	30	33	48	21
250.50 . . . . .	29	13	16	—	*	13	15	*8	*7	10	*
250.51 . . . . .	70	29	41	—	15	26	29	17	16	24	12
250.52 . . . . .	11	*	*6	—	*	*	*6	*	*	*7	*
250.53 . . . . .	22	12	10	—	*8	*9	*6	*	*8	*6	*
250.6 . . . . .	247	113	134	*	37	93	117	49	64	87	47
250.60 . . . . .	72	38	34	—	*	22	47	17	18	21	16
250.61 . . . . .	109	44	65	—	17	43	49	20	30	39	20
250.62 . . . . .	24	13	11	—	*	9	12	*	*8	10	*
250.63 . . . . .	43	18	25	*	14	20	*9	*7	9	17	9
250.7 . . . . .	159	91	68	—	9	46	103	36	32	55	35
250.70 . . . . .	61	33	28	—	*	15	44	12	14	19	16
250.71 . . . . .	63	35	29	—	*	20	40	18	11	22	12
250.72 . . . . .	13	10	*	—	*	*	9	*	*	*	*
250.73 . . . . .	21	13	*8	—	*	*7	10	*	*	10	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
250.8	200	95	105	*	19	67	112	47	45	79	28
250.80	70	35	35	*	*5	18	47	17	17	25	12
250.81	70	35	35	*	*8	25	35	19	14	27	11
250.82	25	9	16	—	*	*8	16	*	*6	14	*
250.83	34	16	19	*	*5	15	13	*8	*8	14	*
250.9	80	29	51	*	10	26	42	15	17	41	*7
250.90	*9	*	*6	*	*	*	*	*	*	*	*
250.91	*9	*	*6	*	*	*	*	*	*	*	*
250.92	26	*8	18	*	*	9	14	*7	*7	*9	*
250.93	37	15	21	*	*	12	20	*	*7	25	*
251	51	24	27	*	13	11	23	12	10	21	*8
251.2	30	14	15	*	*7	*6	13	*6	*6	12	*5
251.8	15	*7	*8	*	*	*	*7	*	*	*6	*
252	23	*	18	*	*	*7	10	*6	*6	*7	*
252.0	19	*	15	*	*	*7	*9	*5	*	*6	*
253	61	27	34	*	*9	14	33	15	15	17	13
253.2	*8	*	*	*	*	*	*	*	*	*	*
253.5	*5	*	*	*	*	*	*	*	*	*	*
253.6	39	15	24	*	*	*8	28	10	9	11	9
255	49	15	34	*	11	12	22	13	10	16	10
255.0	11	*	10	*	*	*	*5	*	*	*6	*
255.4	25	*9	16	*	*6	*5	13	*7	*6	*7	*
256	10	—	10	—	*9	*	*	*	*	*	*
256.4	*6	—	*6	—	*6	*	—	*	*	*	*
260	10	*5	*5	*	*	*	*7	*	*	*	*
261	24	12	12	*	*	*	16	*6	*9	*6	*
262	13	*5	*7	*	*	*	*7	*	*	*	*
263	345	153	192	*8	39	57	241	75	94	134	42
263.0	16	*7	9	*	*	*	11	*8	*	*	*
263.1	19	*6	13	*	*	*	14	9	*	*5	*
263.8	22	10	12	*	*	*	15	*	*7	*9	*
263.9	289	130	159	*	34	49	201	55	83	115	36
266	27	10	18	—	*	*	23	11	*6	*8	*
266.2	27	10	17	—	*	*	23	10	*6	*8	*
269	10	*	*6	*	*	*	*	*	*	*6	*
269.9	*6	*	*	*	*	*	*	*	*	*	*
270	*9	*	*5	*	*	*	*	*	*	*	*
271	31	13	18	*	*5	10	13	13	*	11	*
271.3	28	12	17	*	*	*9	12	12	*	10	*
272	900	440	460	*	103	368	427	229	239	287	145
272.0	518	249	268	*	55	213	248	153	139	147	78
272.1	33	20	12	*	*7	14	12	*	11	13	*
272.2	16	*6	10	—	*	10	*5	*	*5	*7	*
272.4	317	159	158	*	35	126	155	65	80	116	57
272.9	*8	*	*7	*	*	*	*	*	*	*	*
273	63	28	35	*	11	15	35	16	13	23	11
273.3	*5	*	*	—	*	*	*	*	*	*	*
273.8	52	22	30	*	10	12	28	12	12	21	*8
274	175	112	64	*	11	37	127	48	43	59	25
274.0	61	37	24	*	*	12	45	14	17	20	9
274.9	112	73	39	—	*8	25	79	33	25	38	16
275	218	91	127	*5	47	67	99	42	52	88	36
275.0	10	*6	*	*	*	*	*	*	*	*	*
275.2	71	28	42	*	15	23	31	17	20	24	10

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
275.3	38	18	20	*	11	11	15	*	*7	15	12
275.4	99	39	60	*	19	30	49	17	23	47	12
276	4,421	1,743	2,679	389	667	809	2,556	872	1,175	1,697	677
276.0	96	42	54	9	*7	13	67	19	21	40	15
276.1	852	348	504	26	92	175	559	152	227	334	139
276.2	160	71	90	20	31	34	75	30	43	57	30
276.3	35	15	20	*	*7	*7	18	*6	*8	16	*
276.4	16	9	*7	*	*	*	10	*	*	10	*
276.5	1,721	711	1,010	291	285	236	909	350	439	660	271
276.6	60	26	34	*	10	17	32	*7	14	26	14
276.7	249	116	133	*8	31	48	161	49	65	97	37
276.8	1,124	362	762	25	185	253	661	241	313	419	151
276.9	108	44	64	*5	17	22	63	14	43	37	14
277	48	22	25	14	17	*9	*8	16	13	13	*
277.0	23	10	13	12	11	*	—	*8	*7	*9	*
277.00	23	10	13	12	10	*	—	*8	*7	*8	*
277.3	*6	*	*	—	*	*	*	*	*	*	*
277.6	*7	*	*5	*	*	*	*	*	*	*	*
278	541	187	354	*	177	219	140	108	148	196	89
278.0	530	183	347	*	172	216	137	105	145	192	87
278.8	*6	*	*	*	*	*	*	*	*	*	*
279	20	10	10	*6	*	*5	*	*	*8	*6	*
279.0	9	*	*5	*	*	*	*	*	*	*	*
280-289	3,874	1,535	2,338	194	951	807	1,922	870	892	1,440	671
280	614	211	403	25	128	122	338	133	152	239	90
280.0	368	129	239	*	69	78	217	86	81	144	56
280.9	238	80	158	22	58	42	117	45	68	92	32
281	114	45	69	*	12	18	82	22	33	43	16
281.0	32	10	22	*	*	*5	25	*7	*9	11	*
281.1	10	*	*6	—	*	*	*9	*	*	*	*
281.9	65	29	36	*	*8	10	44	11	20	24	10
282	130	59	71	40	70	10	*9	33	26	49	22
282.4	18	*7	11	*	*6	*	*	*8	*	*	*
282.5	13	*	10	*	*8	*	*	*	*	*6	*
282.6	87	42	45	29	53	*	*	19	19	35	13
282.60	18	9	*9	*7	*9	*	*	*5	*	*6	*
282.61	*7	*	*	*6	*	*	*	*	*	*	*
282.62	60	27	33	16	42	*	*	12	12	28	*8
282.9	*5	*	*	*	*	*	*	*	*	*	*
283	16	*7	10	*	*	*	*9	*6	*	*5	*
283.0	*8	*	*	*	*	*	*	*	*	*	*
284	153	73	81	*8	35	40	70	43	31	48	32
284.8	142	68	74	*7	32	38	65	40	29	43	30
284.9	9	*	*5	*	*	*	*	*	*	*	*
285	2,074	776	1,298	48	502	429	1,096	433	471	793	377
285.1	805	305	500	12	183	165	446	158	196	282	169
285.8	37	21	16	*	*6	10	20	*7	10	12	*8
285.9	1,229	448	781	35	312	254	628	267	265	498	199
286	128	68	60	*7	32	31	57	33	27	43	24
286.0	*7	*7	*	*	*5	*	*	*	*	*	*
286.4	*9	*	*6	*	*	*	*	*	*	*	*
286.5	14	*	10	*	*	*	*8	*	*	*6	*
286.6	20	*8	12	*	*	*	10	*5	*	*7	*
286.7	26	15	11	*	*	*8	14	*7	*6	11	*
286.9	43	24	19	*	13	10	20	11	9	14	9

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
287 . . . . .	274	127	147	19	71	64	119	74	60	92	48
287.3 . . . . .	35	9	26	*	11	*	15	11	*6	13	*6
287.4 . . . . .	51	26	25	*	14	18	16	16	11	11	13
287.5 . . . . .	179	89	90	9	44	42	84	46	41	64	28
288 . . . . .	267	120	147	26	68	67	106	68	66	94	38
288.0 . . . . .	157	73	84	17	38	46	56	45	36	52	24
288.3 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
288.8 . . . . .	99	42	57	*8	24	20	47	22	24	39	14
289 . . . . .	104	51	54	16	27	26	35	25	25	33	22
289.0 . . . . .	*8	*	*	—	*	*	*	*	*	*	*
289.2 . . . . .	14	*6	*8	*7	*7	*	*	*	*	*6	*
289.3 . . . . .	14	*7	*7	*5	*	*	*	*	*	*	*
289.4 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
289.5 . . . . .	12	*8	*	*	*	*	*	*	*	*	*
289.59 . . . . .	10	*7	*	*	*	*	*	*	*	*	*
289.8 . . . . .	25	13	12	*	*	*6	14	*6	*7	*8	*
289.9 . . . . .	20	*7	13	*	*5	*	10	*6	*	*6	*
290-319 . . . . .	7,279	3,623	3,656	269	3,548	1,520	1,942	2,052	1,914	2,367	946
290 . . . . .	509	178	331	*	*	14	492	115	122	193	80
290.0 . . . . .	328	103	225	*	*	*	322	72	76	129	51
290.1 . . . . .	16	*	12	—	*	*	13	*	*	*7	*
290.10 . . . . .	10	*	*7	—	*	*	*7	*	*	*	*
290.2 . . . . .	22	*	17	—	—	*	21	*	*6	*8	*
290.20 . . . . .	11	*	*9	—	—	*	11	*	*	*	*
290.21 . . . . .	10	*	*8	—	—	*	10	*	*	*	*
290.3 . . . . .	41	18	24	*	—	*	41	11	11	11	9
290.4 . . . . .	97	45	52	—	*	*6	90	20	25	36	16
290.40 . . . . .	79	37	42	—	*	*5	74	16	18	30	15
290.41 . . . . .	10	*5	*	—	—	*	10	*	*	*	*
290.43 . . . . .	*6	*	*	—	—	*	*5	*	*	*	*
291 . . . . .	186	142	43	*	91	70	24	37	67	50	33
291.0 . . . . .	49	37	12	*	23	18	*8	*7	15	17	10
291.2 . . . . .	13	11	*	—	*	*	*6	*	*	*	*
291.8 . . . . .	113	87	26	*	61	43	*9	24	44	26	18
292 . . . . .	99	54	45	*	56	14	28	34	21	29	15
292.0 . . . . .	38	23	16	*	27	*7	*	19	*8	*7	*
292.1 . . . . .	15	*9	*7	*	10	*	*	*	*	*9	*
292.12 . . . . .	11	*7	*	*	*6	*	*	*	*	*7	*
292.8 . . . . .	41	21	20	*	16	*	20	11	10	12	*8
292.81 . . . . .	26	12	14	*	*	*	18	*7	*5	*9	*5
292.84 . . . . .	*8	*	*	*	*6	*	*	*	*	*	*
292.89 . . . . .	*6	*	*	—	*5	*	*	*	*	*	*
293 . . . . .	110	46	65	*	18	23	69	31	27	35	18
293.0 . . . . .	60	25	34	*	*	*6	49	12	19	17	11
293.8 . . . . .	38	16	22	*	12	14	12	16	*	14	*
293.81 . . . . .	10	*	*7	—	*	*	*	*	*	*	*
293.82 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
293.83 . . . . .	21	11	10	*	*8	*	*8	11	*	*5	*
293.9 . . . . .	12	*	*7	—	*	*	*8	*	*	*	*
294 . . . . .	209	82	126	*	9	11	188	47	57	76	28
294.1 . . . . .	11	*7	*	*	*	*	*7	*	*	*	*
294.8 . . . . .	180	68	112	—	*	*8	170	40	51	65	25
294.9 . . . . .	15	*7	*9	*	*	*	9	*	*	*6	*
295 . . . . .	395	210	185	*	225	107	63	165	90	102	38
295.3 . . . . .	105	60	45	—	59	29	17	51	13	30	11

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
295.30	25	14	11	—	12	*6	*7	*6	*	*9	*
295.32	21	13	*8	—	*9	*7	*5	12	*	*5	*
295.34	55	32	24	—	36	15	*	31	*6	14	*
295.4	*7	*	*	*	*6	*	*	*	*	*	*
295.40	*5	*	*	*	*	*	*	*	*	*	*
295.6	44	25	20	*	19	15	11	14	12	14	*
295.60	11	*6	*	*	*	*	*	*	*	*	*
295.62	18	10	*8	—	*7	*6	*	*6	*	*6	*
295.64	15	*8	*7	—	*8	*6	*	*6	*	*	*
295.7	138	68	70	*	94	34	10	54	41	32	12
295.70	77	39	38	*	51	21	*5	20	28	22	*7
295.74	57	27	30	*	41	12	*	32	12	*8	*
295.8	*6	*	*	—	*	*	*	*	*	*	*
295.9	88	47	42	*	41	24	23	39	19	21	9
295.90	54	22	32	*	20	15	20	13	17	17	*7
295.92	*6	*	*	—	*	*	*	*	*	*	*
295.94	27	21	*6	—	18	*7	*	24	*	*	*
296	853	317	536	41	482	176	154	223	225	316	88
296.0	*6	*	*	—	*	*	*	*	*	*	*
296.2	305	122	182	24	172	55	53	64	77	133	30
296.20	165	66	100	11	94	31	29	40	55	55	15
296.22	25	*9	16	*	15	*	*	*	*	18	*
296.23	76	31	45	*6	45	11	14	13	11	45	*7
296.24	34	15	19	*	17	*9	*6	*9	*8	13	*5
296.3	277	89	188	*9	162	50	55	84	76	97	20
296.30	93	28	65	*	50	18	22	28	33	27	*
296.31	*6	*	*	*	*	*	*	*	*	*	*
296.32	16	*	12	*	11	*	*	*	*	*8	*
296.33	113	36	76	*	74	19	16	40	25	42	*7
296.34	46	16	30	*	23	9	11	13	12	16	*5
296.4	52	23	29	*	29	14	*8	19	12	12	9
296.40	28	12	16	*	15	*8	*	*7	*8	*8	*
296.43	*6	*	*	*	*	*	*	*	*	*	*
296.44	16	*6	10	—	*9	*	*	*9	*	*	*
296.5	44	14	29	*	25	11	*7	10	11	16	*6
296.50	24	*7	17	*	14	*	*	*6	*9	*7	*
296.53	10	*	*7	*	*	*	*	*	*	*	*
296.54	*7	*	*	*	*	*	*	*	*	*	*
296.6	48	15	32	*	28	13	*	12	12	18	*5
296.60	25	9	16	*	16	*5	*	*5	*8	11	*
296.63	*7	*	*	*	*	*	*	*	*	*	*
296.64	11	*	*8	—	*6	*	*	*	*	*	*
296.7	79	33	46	*	40	21	16	20	25	23	11
296.8	33	14	19	*	16	*8	*8	12	*6	11	*
296.80	20	*8	12	*	*7	*6	*5	*	*	*9	*
296.82	*8	*	*	*	*6	*	*	*6	*	*	*
296.9	9	*	*6	*	*	*	*	*	*	*	*
296.90	*7	*	*	—	*	*	*	*	*	*	*
297	25	*9	16	*	*7	*5	13	*6	*7	*8	*
297.1	*8	*	*	*	*	*	*	*	*	*	*
297.9	16	*5	11	*	*	*	9	*	*	*	*
298	152	70	82	*	60	22	68	46	28	62	17
298.8	*5	*	*	*	*	*	*	*	*	*	*
298.9	142	65	78	*	56	19	65	42	26	58	16
299	*7	*	*	*5	*	*	—	*	*	*	*
300	578	186	391	16	218	149	195	145	160	203	69
300.0	286	95	191	*	80	79	125	69	82	106	29

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
300.00	234	77	157	*	55	64	113	57	70	84	23
300.01	37	13	24	*	21	10	*6	*9	9	15	*
300.02	14	*5	*9	*	*	*	*6	*	*	*7	*
300.1	34	*6	28	*	23	*6	*	12	*8	10	*5
300.11	15	*	10	*	*8	*	*	*	*	*	*
300.14	*9	*	*8	—	*8	*	—	*	*	*	*
300.15	*6	*	*5	*	*	*	*	*	*	*	*
300.2	12	*	*9	*	*8	*	*	*	*	*5	*
300.21	*9	*	*7	—	*6	*	*	*	*	*	*
300.3	17	*7	10	*	10	*	*	*	*	*6	*
300.4	158	45	112	*7	54	45	51	45	44	52	17
300.8	*7	*	*5	—	*	*	*	*	*	*	*
300.81	*6	*	*	—	*	*	*	*	*	*	*
300.9	62	27	35	*	41	10	*8	10	16	21	14
301	354	161	193	10	268	55	21	139	91	87	36
301.0	*6	*	*	*	*	*	*	*	*	*	*
301.2	11	*7	*	*	*8	*	*	*5	*	*	*
301.20	*8	*	*	*	*	*	*	*	*	*	*
301.3	*6	*	*	*	*	*	*	*	*	*	*
301.4	*7	*	*	—	*	*	*	*	*	*	*
301.5	11	*	*7	*	*8	*	*	*	*	*	*
301.50	10	*	*6	*	*7	*	*	*	*	*	*
301.6	36	14	21	*	21	*8	*6	10	*9	14	*
301.7	31	26	*	*	27	*	*	*7	15	*	*
301.8	175	65	110	*6	140	26	*	83	36	41	16
301.81	*7	*	*	*	*	*	*	*	*	*	*
301.82	*6	*	*	*	*	*	*	*	*	*	*
301.83	89	21	68	*	72	12	*	29	26	26	*8
301.84	10	*5	*	*	*7	*	*	*	*	*	*
301.89	64	33	31	*	53	*8	*	51	*	*5	*
301.9	66	30	36	*	51	10	*	26	18	14	9
303	846	615	232	*	478	272	92	221	243	277	106
303.0	119	87	32	*	67	42	10	27	46	25	21
303.00	36	25	11	*	21	13	*	*	11	11	11
303.01	77	58	20	—	43	28	*6	23	33	13	*7
303.9	727	528	199	*	412	229	83	194	197	252	85
303.90	294	204	89	*	152	98	43	57	79	120	38
303.91	380	287	93	*	232	113	33	123	103	117	38
303.92	16	12	*	*	11	*	*	*	*5	*	*
303.93	37	24	13	—	17	14	*7	10	10	11	*7
304	410	258	152	*6	335	61	*8	162	92	117	39
304.0	90	55	35	*	69	20	*	48	17	15	10
304.00	26	17	*9	*	18	*7	*	*5	*8	*7	*6
304.01	61	36	25	*	48	11	*	41	*8	*7	*
304.1	20	*8	11	—	10	*6	*	*7	*	*6	*
304.10	*8	*	*	—	*	*	*	*	*	*	*
304.11	11	*	*6	—	*5	*	*	*	*	*	*
304.2	124	79	44	*	111	12	*	58	24	35	*6
304.20	36	21	16	*	33	*	*	*6	*8	18	*
304.21	82	56	26	*	74	*8	—	49	14	17	*
304.3	46	34	12	*	41	*	—	10	25	9	*
304.30	10	*6	*	*	*9	*	—	*	*	*	*
304.31	35	27	*8	*	32	*	—	*7	21	*5	*
304.4	9	*6	*	*	*9	*	—	*	*	*	*
304.41	*6	*	*	*	*6	*	—	*	*	*	*
304.6	14	*9	*5	*	*9	*	*	*	*	*	*
304.60	*7	*	*	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
304.61	*7	*	*	*	*	*	—	*	*	*	*
304.7	32	20	12	—	26	*5	*	18	*	*8	*
304.70	*8	*	*	—	*6	*	*	*	*	*	*
304.71	24	15	*9	—	19	*	*	15	*	*5	*
304.8	47	34	12	*	38	*7	—	11	*8	24	*
304.80	19	14	*	*	17	*	—	*	*	*9	*
304.81	25	19	*6	*	19	*6	—	*6	*	15	*
304.9	27	11	16	*	21	*	*	*6	*5	12	*
304.90	14	*6	*8	*	11	*	*	*	*	*7	*
304.91	9	*	*	—	*7	*	*	*	*	*	*
305	1,248	732	517	13	736	337	163	301	329	408	210
305.0	248	174	74	*	164	56	26	82	62	71	33
305.00	143	95	48	*	96	31	15	31	47	45	20
305.01	66	51	14	*	44	15	*5	28	11	19	*8
305.02	24	17	*7	*	18	*	*	17	*	*	*
305.03	15	10	*5	*	*6	*5	*	*6	*	*	*
305.1	591	312	279	*	218	238	131	86	167	221	116
305.2	86	54	31	*	77	*	*	19	23	29	14
305.20	56	35	21	*	51	*	*	*9	18	20	10
305.21	21	15	*7	*	19	*	—	*8	*	*8	*
305.22	*6	*	*	*	*5	*	—	*	*	*	*
305.4	*6	*	*	—	*5	*	*	*	*	*	*
305.5	30	19	10	*	21	*8	*	12	*7	*6	*
305.50	19	13	*7	*	14	*5	*	*7	*	*	*
305.51	*7	*	*	—	*5	*	—	*	*	*	*
305.6	141	80	61	*	124	16	*	46	36	45	14
305.60	93	53	41	*	82	10	*	19	29	33	12
305.61	33	19	14	*	29	*	*	18	*	*9	*
305.62	*9	*5	*	—	*8	*	—	*6	*	*	*
305.63	*6	*	*	—	*	*	—	*	*	*	*
305.7	13	*8	*5	*	12	*	—	*	*	*	*7
305.70	*8	*	*	—	*8	*	—	*	*	*	*
305.9	127	77	51	*	108	12	*	51	27	31	18
305.90	78	47	32	*	66	*7	*	23	21	23	11
305.91	23	14	*9	*	19	*	*	10	*	*6	*
305.92	13	9	*	*	12	*	—	12	*	*	*
305.93	14	*7	*7	*	11	*	*	*6	*	*	*
306	15	*	11	*	*6	*	*7	*	*	*8	*
306.1	*6	*	*	*	*	*	*	*	*	*	*
307	63	15	49	*6	33	*9	15	13	16	26	*9
307.1	*8	*	*8	*	*6	*	*	*	*	*	*
307.5	19	*	18	*	14	*	*	*	*5	*6	*
307.50	*7	*	*7	*	*5	*	—	*	*	*	*
307.51	9	*	9	*	*8	*	—	*	*	*	*
307.8	18	*	13	*	*9	*	*5	*	*5	*7	*
307.81	15	*	11	—	*7	*	*	*	*5	*5	*
307.9	12	*6	*6	*	*	*	*8	*	*	*5	*
308	32	*6	26	*	21	*6	*	12	10	*7	*
308.0	*6	*	*	—	*	*	*	*	*	*	*
308.3	19	*	16	*	13	*	*	*8	*6	*	*
308.9	*7	*	*6	—	*	*	*	*	*	*	*
309	257	114	143	25	175	33	24	96	67	58	36
309.0	90	38	52	*6	60	11	14	32	16	26	17
309.2	31	13	18	*	18	*5	*	*7	11	*8	*
309.24	*5	*	*	*	*	*	*	*	*	*	*
309.28	22	9	13	*	15	*	*	*5	*8	*5	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
309.3	*6	*	*	*	*	*	*	*	*	*	*
309.4	38	23	14	*	27	*7	*	22	*8	*	*
309.8	61	22	40	*9	43	*7	*	22	21	12	*6
309.81	59	21	38	*8	42	*7	*	21	21	12	*6
309.9	26	11	15	*	20	*	*	*8	*8	*6	*
310	126	52	74	*	11	*8	103	29	28	48	21
310.1	10	*6	*	*	*	*	*	*	*	*	*
310.2	*8	*	*	*	*	*	*	*	*	*	*
310.9	104	41	63	*	*	*	95	26	22	44	13
311	483	172	311	14	176	102	190	119	136	164	64
312	66	42	24	19	42	*	*	29	16	16	*
312.3	34	23	11	*5	26	*	*	16	*8	*8	*
312.30	13	9	*	*	11	*	*	*9	*	*	*
312.34	13	9	*	*	9	*	*	*	*	*	*
312.8	13	*6	*7	*7	*6	—	*	*7	*	*	*
312.89	*5	*	*	*	*	—	*	*	*	*	*
312.9	13	*8	*	*	*8	*	*	*	*6	*	*
313	30	16	14	19	11	—	*	*8	11	*8	*
313.8	28	15	13	18	10	—	—	*8	11	*7	*
313.81	26	14	12	16	10	—	—	*8	*8	*7	*
314	51	40	11	32	18	*	*	13	20	15	*
314.0	49	39	10	31	17	*	*	13	19	14	*
314.00	*9	*6	*	*	*5	*	—	*	*	*	*
314.01	40	32	*7	28	12	—	*	11	15	12	*
315	32	19	13	24	*7	*	*	*9	13	*6	*
315.8	*5	*	*	*	*	*	—	*	*	*	*
315.9	12	*7	*	9	*	*	—	*	*7	*	*
317	39	24	15	*	21	13	*	18	*7	10	*
318	33	21	12	*6	15	11	*	12	9	*8	*
318.0	11	*5	*6	*	*7	*	*	*	*	*	*
318.1	11	*8	*	*	*	*	*	*	*	*	*
318.2	11	*8	*	*	*	*5	—	*8	*	*	*
319	62	30	33	12	22	14	14	17	14	23	*8
320-389	3,391	1,571	1,820	463	547	631	1,750	847	850	1,121	573
320	20	10	9	12	*	*	*	*	*	11	*
320.9	*6	*	*	*	*	*	*	*	*	*	*
322	13	*9	*	*	*	*	*	*	*	*5	*
322.9	10	*6	*	*	*	*	*	*	*	*	*
323	12	*	*7	*	*8	*	*	*	*	*7	*
323.9	10	*	*6	*	*7	*	*	*	*	*6	*
324	*5	*	*	*	*	*	*	*	*	*	*
331	309	122	188	15	11	17	267	72	76	117	45
331.0	224	75	148	*	*	*	220	51	52	89	32
331.3	13	*6	*6	*	*	*	*7	*	*	*	*
331.4	44	25	19	12	*6	*9	17	11	13	14	*
331.9	23	12	10	*	*	*	18	*	*7	*7	*
332	202	111	92	—	*	12	190	52	55	65	31
332.0	199	110	89	—	*	11	188	51	54	64	31
333	60	25	36	*	10	11	37	12	19	17	11
333.0	*5	*	*	*	*	*	*	*	*	*	*
333.1	15	*6	9	—	*	*	12	*	*7	*	*
333.2	*5	*	*	*	*	*	*	*	*	*	*
333.8	11	*	*9	*	*	*	*7	*	*	*	*
333.82	10	*	*7	—	*	*	*6	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
333.9 . . . . .	13	*5	*7	*	*	*	*6	*	*	*	*
333.90 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
333.99 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
334 . . . . .	11	*6	*	*	*	*	*6	*	*	*	*
335 . . . . .	10	*5	*5	*	*	*	*6	*	*	*	*
335.2 . . . . .	*8	*	*	—	*	*	*6	*	*	*	*
336 . . . . .	19	11	*8	*	*	*7	*7	*6	*	*	*
336.9 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
337 . . . . .	64	24	40	*	19	18	27	13	15	18	19
337.1 . . . . .	45	17	27	—	12	13	20	10	12	12	10
337.2 . . . . .	11	*	9	*	*6	*	*	*	*	*	*
337.22 . . . . .	*6	*	*5	*	*	*	*	*	*	*	*
340 . . . . .	64	16	48	*	23	26	14	13	23	19	10
342 . . . . .	435	196	240	*5	25	88	317	91	120	138	86
342.0 . . . . .	15	*5	9	*	*	*	11	*	*5	*	*
342.00 . . . . .	12	*	*7	*	*	*	*9	*	*	*	*
342.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
342.8 . . . . .	9	*	*	*	*	*	*6	*	*	*	*
342.80 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
342.9 . . . . .	406	183	223	*	22	83	297	87	109	131	79
342.90 . . . . .	345	157	188	*	17	68	255	76	93	109	67
342.91 . . . . .	37	16	21	*	*	*8	25	*7	9	13	*8
342.92 . . . . .	24	9	15	*	*	*6	16	*	*6	*9	*
343 . . . . .	63	36	27	28	18	13	*	26	11	15	11
343.2 . . . . .	12	*6	*5	*	*	*	*	*6	*	*	*
343.9 . . . . .	45	25	20	21	13	9	*	17	*7	12	*9
344 . . . . .	134	81	53	*	43	34	51	28	39	40	27
344.0 . . . . .	49	37	12	*	20	12	15	11	14	14	10
344.00 . . . . .	36	26	9	*	12	10	13	10	10	11	*5
344.03 . . . . .	*6	*	*	—	*5	*	*	*	*	*	*
344.1 . . . . .	43	25	19	*	14	11	17	*8	11	15	*8
344.6 . . . . .	14	*8	*6	*	*	*	*5	*	*5	*	*
344.61 . . . . .	9	*	*	*	*	*	*	*	*	*	*
344.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
344.89 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
344.9 . . . . .	16	*6	10	*	*	*5	*9	*	*7	*6	*
345 . . . . .	125	59	66	31	43	24	27	34	39	36	17
345.1 . . . . .	25	13	12	*	9	*5	*5	*6	11	*5	*
345.10 . . . . .	22	11	11	*	*8	*5	*5	*5	10	*	*
345.3 . . . . .	19	10	*9	*5	*	*	*5	*	*	*6	*
345.4 . . . . .	10	*	*6	*	*6	*	*	*	*	*	*
345.40 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
345.5 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
345.50 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
345.9 . . . . .	51	22	28	13	16	11	11	14	15	16	*6
345.90 . . . . .	45	20	24	9	14	11	11	11	14	14	*6
345.91 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
346 . . . . .	107	20	87	*	59	34	12	24	29	39	15
346.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
346.1 . . . . .	*7	*	*7	—	*	*	—	*	*	*	*
346.10 . . . . .	*6	*	*6	—	*	*	—	*	*	*	*
346.2 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
346.8 . . . . .	*9	*	*7	*	*	*	*	*	*	*	*
346.80 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
346.9 . . . . .	79	15	64	*	43	26	*8	18	23	29	10

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
346.90	73	15	58	*	40	24	*7	17	22	25	9
346.91	*6	*	*6	*	*	*	*	*	*	*	*
348	127	65	62	19	27	27	54	33	29	42	25
348.0	*7	*	*	*	*	*	*	*	*	*	*
348.1	43	23	20	*	*5	10	23	10	10	14	9
348.3	41	20	21	*5	9	*6	21	11	*6	16	*8
348.4	*6	*	*	*	*	*	*	*	*	*	*
348.5	*8	*	*	*	*	*	*	*	*	*	*
348.8	16	*9	*7	*	*6	*	*	*	*	*5	*
349	41	13	28	*	22	*7	11	10	*8	14	*9
349.0	18	*	14	—	14	*	*	*	*	*7	*
349.8	15	*7	*8	*	*	*	*6	*	*	*	*
349.81	*6	*	*	*	*	*	*	*	*	*	*
349.82	*8	*	*	*	*	*	*5	*	*	*	*
350	12	*	*8	—	*	*	*6	*	*	*	*
350.1	10	*	*7	—	*	*	*6	*	*	*	*
351	22	*8	14	*	*	*6	12	*6	*	*8	*5
351.0	19	*8	12	*	*	*5	10	*	*	*7	*
353	13	*7	*6	*	*5	*	*	*	*	*	*
353.0	*7	*	*	*	*	*	*	*	*	*	*
354	46	24	22	*	13	17	14	16	*7	15	*8
354.0	24	10	14	*	*7	9	*7	*7	*	10	*
354.2	*8	*6	*	*	*	*	*	*	*	*	*
354.9	*6	*5	*	—	*	*	*	*	*	*	*
355	30	12	17	*	10	*6	13	*8	*5	9	*7
355.8	11	*6	*6	*	*	*	*6	*	*	*	*
355.9	*6	*	*	—	*	*	*	*	*	*	*
356	51	24	27	*	9	12	29	15	12	17	*7
356.9	42	22	20	*	*7	10	25	12	9	14	*7
357	137	72	65	*	17	52	67	26	39	48	23
357.0	9	*	*	*	*	*	*	*	*	*	*
357.2	118	60	58	—	14	46	58	22	34	43	20
358	19	9	10	*	*	*	13	*6	*5	*6	*
358.0	15	*7	*8	*	*	*	10	*	*	*	*
359	23	14	*9	*	*6	*	9	*9	*	*	*6
359.1	10	*8	*	*	*	*	*	*	*	*	*
360	*8	*	*	*	*	*	*	*	*	*5	*
360.0	*5	*	*	*	*	*	*	*	*	*	*
361	25	14	11	*	*	11	9	*	*	16	*
361.0	12	*7	*	*	*	*	*	*	*	*7	*
361.8	*6	*	*	—	*	*	*	*	*	*	*
361.9	*5	*	*	*	*	*	*	*	*	*	*
362	141	64	77	*6	20	42	73	31	37	50	23
362.0	79	34	45	—	14	32	33	20	23	26	11
362.01	71	33	38	—	13	27	31	18	22	21	10
362.02	*8	*	*6	—	*	*	*	*	*	*5	*
362.1	*6	*	*	*	*	*	*	*	*	*	*
362.2	*8	*5	*	*	—	*	*	*	*	*	*
362.3	13	*7	*7	—	*	*	*9	*	*	*	*
362.34	9	*	*	—	*	*	*6	*	*	*	*
362.5	28	9	19	*	*	*	27	*	*7	10	*6
362.50	23	*8	15	*	*	*	22	*	*7	*6	*6
363	17	13	*	*	10	*	*	*7	*	*6	*
363.2	13	10	*	*	10	*	*	*6	*	*	*
363.20	13	10	*	*	10	*	*	*6	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
364	*6	*	*	*	*	*	*	*	*	*	*
365	156	60	95	*	*	20	132	43	36	57	20
365.9	141	54	87	*	*	17	121	40	32	52	17
366	84	33	51	*	*	11	69	47	*7	23	*7
366.1	17	*	13	—	—	*	15	11	*	*	*
366.10	10	*	*7	—	—	*	9	*8	*	*	*
366.8	*8	*5	*	—	—	*	*6	*	*	*	*
366.9	56	21	35	*	*	*7	46	31	*6	16	*
368	42	19	24	*	10	12	18	11	9	14	*8
368.1	*8	*	*6	*	*	*	*	*	*	*	*
368.2	*9	*	*6	*	*	*	*	*	*	*	*
368.4	15	*8	*7	*	*	*7	*7	*	*	*	*
368.46	13	*7	*5	*	*	*6	*5	*	*	*	*
368.8	*7	*	*	*	*	*	*	*	*	*	*
369	84	40	44	*	11	14	57	16	18	32	18
369.0	35	14	20	*	*	*	26	*6	*5	15	*9
369.00	31	13	18	*	*	*	24	*6	*	13	*8
369.4	16	*7	*9	*	*	*	11	*	*	*6	*
369.6	21	12	*9	*	*	*7	11	*	*6	*8	*
369.60	19	10	*9	*	*	*6	10	*	*6	*7	*
370	11	*	*6	*	*	*	*7	*6	*	*	*
372	48	27	21	15	9	*5	18	10	12	16	10
372.0	11	*6	*6	*	*	*	*6	*	*	*	*
372.00	10	*	*5	*	*	*	*6	*	*	*	*
372.3	24	12	12	*9	*	*	*8	*	*7	*8	*
372.30	23	11	12	*8	*	*	*8	*	*6	*7	*
372.7	10	*8	*	*	*	*	*	*	*	*5	*
372.72	*8	*6	*	*	*	*	*	*	*	*	*
373	10	*	*6	*	*	*	*	*5	*	*	*
373.1	*5	*	*	*	*	*	*	*	*	*	*
374	10	*	*7	*	*	*	*	*	*	*	*
374.3	*7	*	*5	*	*	*	*	*	*	*	*
376	14	9	*5	*7	*	*	*	*	*	*	*
376.0	*9	*6	*	*6	*	*	*	*	*	*	*
376.01	*9	*6	*	*6	*	*	*	*	*	*	*
377	14	*6	*7	*	*	*	*	*	*	*	*
378	20	10	10	*	*	*	*7	*7	*	*5	*
378.5	11	*6	*5	*	*	*	*5	*	*	*	*
379	22	10	12	*	*5	*6	*7	*5	*	*8	*
379.2	9	*	*5	*	*	*	*	*	*	*	*
379.23	*7	*	*	*	*	*	*	*	*	*	*
380	33	16	17	10	*7	*	12	9	*9	12	*
380.1	20	*8	12	*	*6	*	*7	*	*6	*8	*
380.10	19	*7	11	*	*6	*	*7	*	*6	*8	*
380.4	11	*7	*	*5	*	*	*	*	*	*	*
381	36	20	16	29	*	*	*	12	9	10	*
381.0	10	*	*5	*8	*	*	*	*	*	*	*
381.01	*6	*	*	*	*	*	*	*	*	*	*
381.1	9	*5	*	*8	*	*	*	*	*	*	*
381.10	9	*5	*	*8	*	*	*	*	*	*	*
381.4	10	*6	*	*8	*	*	*	*	*	*	*
382	239	136	102	213	13	*7	*6	48	59	83	49
382.9	231	133	98	207	12	*6	*5	44	57	81	48
383	*8	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
384 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
384.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
385 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
386 . . . . .	55	16	39	*	*5	16	34	15	20	13	*7
386.0 . . . . .	*9	*	*7	—	*	*	*5	*	*	*	*
386.00 . . . . .	*8	*	*6	—	*	*	*5	*	*	*	*
386.1 . . . . .	12	*	*9	*	*	*	*7	*	*	*	*
386.11 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
386.3 . . . . .	30	9	21	*	*	*7	19	*8	11	*8	*
386.30 . . . . .	28	*9	19	*	*	*7	18	*8	10	*7	*
388 . . . . .	13	*6	*7	*	*	*	*6	*	*	*	*
389 . . . . .	74	35	39	*6	11	*9	48	18	21	23	11
389.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
389.00 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
389.1 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
389.10 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
389.9 . . . . .	54	25	29	*	*7	*	41	12	16	18	*8
390-459 . . . . .	25,390	12,238	13,152	97	1,503	5,901	17,889	6,174	6,100	9,189	3,928
394 . . . . .	38	*7	31	*	*5	*8	25	14	*	12	*7
394.0 . . . . .	13	*	12	*	*	*	*7	*6	*	*	*
394.1 . . . . .	10	*	*8	*	*	*	*7	*	*	*	*
394.2 . . . . .	*7	*	*6	—	*	*	*	*	*	*	*
394.9 . . . . .	*9	*	*6	—	*	*	*6	*	*	*	*
395 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
396 . . . . .	159	69	90	*	*	14	141	44	41	56	19
396.0 . . . . .	*7	*	*5	*	*	*	*7	*	*	*	*
396.2 . . . . .	35	18	17	*	*	*	33	10	12	*9	*
396.3 . . . . .	94	40	54	*	*	*8	84	23	21	37	13
396.8 . . . . .	14	*6	*8	—	*	*	12	*6	*	*	*
396.9 . . . . .	*5	*	*	—	*	*	*	*	*	*	—
397 . . . . .	104	39	65	*	*5	18	80	33	21	35	14
397.0 . . . . .	102	38	63	*	*	18	79	33	21	34	14
398 . . . . .	24	*7	17	—	*	*6	15	11	*	*8	*
398.9 . . . . .	24	*7	17	—	*	*6	15	11	*	*8	*
398.90 . . . . .	14	*	9	—	*	*	*7	*	*	*6	*
398.91 . . . . .	9	*	*7	—	*	*	*8	*5	*	*	*
401 . . . . .	4,876	2,027	2,849	*8	384	1,381	3,104	1,155	1,181	1,798	743
401.0 . . . . .	66	27	39	*	9	22	34	16	13	26	10
401.1 . . . . .	155	63	93	*	12	41	103	92	25	38	*
401.9 . . . . .	4,655	1,938	2,717	*7	363	1,318	2,966	1,046	1,143	1,733	732
402 . . . . .	405	150	254	*	11	80	313	106	93	155	50
402.0 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
402.1 . . . . .	12	*	*7	—	*	*	*9	*6	*	*	*
402.10 . . . . .	*6	*	*	—	*	*	*	*	*	*	—
402.11 . . . . .	*6	*	*	—	—	*	*6	*	*	*	*
402.9 . . . . .	385	141	244	*	10	73	301	99	90	147	49
402.90 . . . . .	215	80	135	*	*7	46	161	61	46	85	23
402.91 . . . . .	170	61	109	—	*	27	140	38	44	62	26
403 . . . . .	436	227	209	*	66	130	239	88	80	186	81
403.0 . . . . .	11	*	*6	*	*	*	*	*	*	*	*
403.01 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
403.1 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
403.11 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
403.9 . . . . .	419	220	199	*	61	127	230	84	76	180	79

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
403.90	38	20	18	*	*6	12	20	*7	*8	19	*
403.91	381	200	181	*	55	115	211	77	68	161	75
404	79	37	42	*	*6	15	59	18	16	33	12
404.0	*5	*	*	—	*	*	*	*	*	*	*
404.9	71	33	38	*	*	14	54	17	14	28	12
404.90	*6	*	*	*	*	*	*	*	*	*	*
404.91	*6	*	*	—	*	*	*	*	*	*	*
404.92	13	*6	*7	*	*	*	*8	*	*	*	*
404.93	46	23	23	—	*	*8	36	10	10	19	*8
405	11	*	*6	*	*	*	*6	*5	*	*	*
405.9	10	*	*6	*	*	*	*6	*	*	*	*
405.91	*8	*	*	*	*	*	*5	*	*	*	*
410	858	517	341	*	46	274	537	215	196	290	157
410.0	39	23	16	—	*	12	24	*6	10	16	*8
410.01	34	19	14	—	*	11	20	*	10	13	*7
410.1	151	91	60	—	12	50	89	37	36	50	29
410.11	131	80	51	—	10	43	78	30	31	45	24
410.12	15	*9	*6	—	*	*6	*8	*6	*	*	*
410.2	28	17	11	—	*	12	15	*6	*7	12	*
410.21	27	17	11	—	*	11	15	*6	*6	11	*
410.3	21	14	*7	—	*	9	10	*7	*	*7	*
410.31	18	12	*5	—	*	*8	*9	*7	*	*5	*
410.4	178	114	64	—	13	75	91	46	45	56	31
410.41	165	105	60	—	12	70	83	43	42	52	28
410.42	10	*8	*	—	*	*	*7	*	*	*	*
410.5	24	15	9	—	*	9	13	*	*5	9	*
410.51	22	14	*8	—	*	*9	12	*	*5	*9	*
410.6	*6	*	*	—	*	*	*	*	*	*	*
410.61	*6	*	*	—	*	*	*	*	*	*	*
410.7	273	158	116	*	*8	72	193	79	58	87	49
410.70	*7	*	*	—	*	*	*5	*	*	*	*
410.71	244	143	101	*	*7	65	172	71	51	80	42
410.72	23	12	10	—	*	*6	16	*6	*	*6	*6
410.8	12	*7	*6	—	*	*	10	*6	*	*	*
410.81	11	*6	*	—	*	*	*9	*	*	*	*
410.9	125	74	51	*	*	32	89	24	26	49	26
410.90	29	19	11	—	*	*8	20	*	*	13	*8
410.91	89	51	38	*	*	22	65	18	20	34	17
410.92	*7	*	*	*	*	*	*	*	*	*	*
411	1,034	580	455	*	59	351	624	247	229	392	166
411.0	14	*8	*6	—	*	*5	*8	*	*	*	*
411.1	952	529	422	*	53	325	573	224	209	369	149
411.8	69	42	27	*	*5	20	43	18	17	19	14
411.81	28	19	9	*	*	10	16	*5	*8	*7	*8
411.89	41	23	18	—	*	10	27	13	9	12	*7
412	575	349	226	*	28	168	379	149	134	190	101
413	585	284	301	*	25	145	414	153	147	212	73
413.1	*6	*	*	—	*	*	*	*	*	*	*
413.9	579	279	299	*	23	143	412	150	147	210	72
414	3,741	2,103	1,637	*	118	969	2,653	960	875	1,317	589
414.0	3,450	1,931	1,519	*	108	906	2,435	865	805	1,224	555
414.00	1,340	691	649	*	30	285	1,025	342	336	531	131
414.01	1,993	1,154	839	*	77	582	1,333	503	443	652	395
414.02	116	86	30	—	*	39	76	20	27	41	29
414.1	14	10	*	—	*	*	11	*	*	*	*6
414.10	11	*8	*	—	*	*	*8	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
414.8	137	86	51	*	*6	29	103	49	27	44	18
414.9	139	75	64	*	*	31	105	44	40	45	10
415	113	49	63	*	16	32	64	24	29	39	21
415.0	*7	*	*	*	*	*	*	*	*	*	*
415.1	106	46	59	*	16	30	60	23	28	37	19
416	184	83	101	*6	15	41	122	43	40	59	42
416.0	105	43	62	*	9	22	69	23	25	34	23
416.8	16	*7	9	*	*	*	*8	*7	*	*	*
416.9	63	33	30	*	*	14	45	13	13	21	17
420	30	18	12	*	*6	10	13	*7	*5	10	*8
420.9	27	15	12	*	*	10	12	*7	*5	*8	*7
420.90	21	13	*9	*	*	*7	10	*5	*	*7	*5
421	25	17	*8	*	10	*7	*8	*7	*	10	*
421.0	23	16	*7	*	9	*6	*8	*7	*	10	*
423	35	18	17	*	*7	12	16	*9	*7	12	*8
423.9	29	15	15	*	*7	11	12	*8	*5	10	*7
424	752	289	463	*	95	136	517	196	175	260	121
424.0	435	147	288	*	79	100	255	111	100	157	67
424.1	269	122	147	*	12	29	226	73	64	87	46
424.2	26	10	16	*	*	*	20	*7	*6	*9	*
424.3	*6	*	*	*	*	*	*	*	*	*	*
424.9	15	*7	*8	*	*	*	12	*	*	*	*
424.90	14	*7	*7	*	*	*	12	*	*	*	*
425	383	209	175	*	34	117	230	95	89	144	56
425.1	18	*	14	—	*	*5	11	*8	*	*6	*
425.4	331	185	146	*	28	102	200	79	77	124	51
425.5	11	9	*	—	*	*6	*	*	*	*	*
425.8	14	*7	*7	*	*	*	10	*	*	*7	*
426	531	281	250	*	26	88	414	123	132	193	83
426.0	69	34	35	*	*	12	53	20	17	23	10
426.1	148	79	68	*	*7	21	120	33	38	56	21
426.10	9	*	*	—	*	*	*7	*	*	*	*
426.11	93	48	45	*	*	12	77	19	23	37	14
426.12	*7	*	*	—	*	*	*6	*	*	*	*
426.13	39	24	15	*	*	*6	30	10	10	14	*6
426.2	32	14	18	*	*	*	27	*8	*5	14	*
426.3	100	44	56	*	*	18	80	20	26	39	14
426.4	74	43	31	*	*	15	54	21	16	21	16
426.5	53	35	18	—	*	*6	46	*7	19	20	*7
426.52	14	10	*	—	—	*	13	*	*6	*5	*
426.53	30	19	11	—	*	*	28	*	12	12	*
426.6	13	*	*8	*	*	*	10	*	*	*	*
426.7	14	*9	*	*	*	*	*	*7	*	*	*
426.8	9	*7	*	*	*	*	*7	*	*	*	*
426.89	9	*7	*	*	*	*	*7	*	*	*	*
426.9	19	11	*8	*	*	*	13	*	*	11	*
427	3,145	1,546	1,599	24	170	511	2,440	762	744	1,108	532
427.0	111	45	65	*	10	25	75	22	22	44	23
427.1	210	126	84	*	12	49	148	55	45	82	27
427.3	1,779	856	922	*	37	218	1,521	433	446	599	302
427.31	1,618	765	853	*	29	185	1,402	395	407	548	269
427.32	160	91	69	*	*7	33	119	38	39	51	33
427.4	55	32	23	*	9	15	30	14	12	17	12
427.41	54	31	23	*	*9	15	29	14	12	17	11
427.5	116	64	52	*	12	23	78	25	25	42	25
427.6	213	114	99	*	15	44	152	50	45	78	39

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
427.60	*5	*	*	—	*	*	*	*	*	*	*
427.61	63	31	32	*	*	10	50	14	*8	26	14
427.69	145	79	66	*	13	31	100	35	36	49	25
427.8	573	259	314	12	67	120	373	146	126	209	91
427.81	131	57	74	*	*	15	113	32	21	55	23
427.89	442	201	240	12	64	106	260	115	105	154	68
427.9	87	48	38	*	*8	15	63	18	22	34	13
428	2,672	1,133	1,539	14	81	409	2,168	623	705	987	357
428.0	2,639	1,114	1,525	13	79	402	2,145	616	698	973	352
428.1	18	11	*7	*	*	*	13	*	*5	*5	*
428.9	14	*8	*6	*	*	*	10	*	*	*8	*
429	384	183	200	*	14	68	299	93	92	153	46
429.2	204	95	109	*	*	33	169	51	50	84	20
429.3	127	59	68	*	*9	26	91	29	29	49	20
429.4	13	*7	*6	*	*	*	10	*	*	*7	*
429.7	*7	*	*	—	*	*	*6	*	*	*	*
429.9	24	14	11	*	*	*	18	*6	*9	*7	*
430	33	14	19	*	*7	12	14	*6	*9	12	*6
431	73	32	40	*	*7	18	47	17	15	26	15
432	24	15	10	*	*	*	16	*	*8	9	*
432.1	19	12	*7	*	*	*	13	*	*6	*7	*
433	317	169	148	*	*	61	252	77	82	121	38
433.0	*7	*5	*	—	*	*	*	*	*	*	*
433.00	*5	*	*	—	*	*	*	*	*	*	—
433.1	221	118	103	*	*	44	174	54	57	83	26
433.10	193	102	91	*	*	37	154	47	49	73	24
433.11	28	16	12	—	*	*7	21	*7	*8	10	*
433.3	82	43	39	*	*	14	67	19	20	33	10
433.30	68	35	33	*	*	12	55	15	17	28	*8
433.31	14	*7	*7	—	*	*	12	*	*	*	*
434	358	151	206	*	12	62	283	96	85	118	58
434.0	24	12	11	—	*	*	20	*6	*6	*7	*
434.01	19	10	9	—	*	*	17	*6	*	*6	*
434.1	47	20	28	*	*	10	36	18	*8	12	9
434.10	*8	*	*	—	—	*	*6	*	*	*	*
434.11	39	17	23	*	*	*8	30	16	*7	*8	*8
434.9	287	119	167	*	10	49	226	72	71	98	45
434.90	11	*	*6	*	*	*	*6	*	*	*	*
434.91	276	115	161	*	9	46	220	70	68	95	43
435	259	123	136	*	*6	44	209	63	70	101	25
435.0	18	*6	11	—	—	*	17	*	*5	*7	*
435.1	19	10	*8	—	*	*	17	*	9	*	*
435.8	*9	*	*	—	*	*	*7	*	*	*	*
435.9	212	102	111	*	*	40	168	53	53	85	21
436	207	86	121	*	*6	38	163	24	60	89	34
437	161	68	93	*	*7	27	126	35	40	69	17
437.0	70	27	43	—	*	10	61	16	14	32	*9
437.1	16	*7	*9	*	*	*	15	*	*6	*7	*
437.2	13	*8	*5	*	*	*	*7	*	*	*5	*
437.3	14	*	10	*	*	*	*7	*	*	*5	*
437.7	*5	*	*	*	*	*	*	*	*	*	*
437.9	38	19	19	*	*	*	33	*8	11	16	*
438	453	226	227	*	14	89	349	97	109	169	79
440	473	233	239	*	*6	89	378	110	122	173	67
440.0	24	*8	16	—	*	*5	18	*7	*	*9	*
440.1	24	*8	16	—	*	*7	17	*8	*	*7	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
440.2	225	126	99	*	*	52	170	44	57	82	42
440.20	64	34	30	—	*	14	50	13	18	24	*9
440.21	46	30	17	—	*	13	32	*6	10	20	10
440.22	13	*7	*6	—	*	*	*8	*	*	*	*
440.23	28	16	12	—	*	*6	22	*6	*8	*6	*7
440.24	67	37	30	—	*	14	54	14	15	25	13
440.29	*7	*	*	*	*	*	*6	*	*	*	*
440.3	*7	*5	*	—	*	*	*5	*	*	*	*
440.9	188	83	104	*	*	22	164	49	54	68	17
441	164	106	58	*	*	28	135	40	45	54	26
441.0	15	*7	*7	—	*	*	11	*	*	*	*
441.01	*8	*	*	—	*	*	*6	*	*	*	*
441.2	12	*8	*	*	*	*	9	*	*7	*	*
441.3	*8	*	*	—	—	*	*6	*	*	*	*
441.4	120	79	42	—	*	18	102	28	32	40	20
442	33	22	11	*	*	*8	24	*7	*8	11	*7
442.2	10	*9	*	—	*	*	*9	*	*	*	*
442.3	14	10	*	—	*	*	12	*	*	*	*
443	359	192	167	*	12	82	265	95	91	128	45
443.0	11	*	*7	—	*	*	*	*	*	*	*
443.8	64	34	30	*	*	18	43	17	11	24	12
443.81	52	29	24	—	*	16	34	14	*8	20	10
443.89	11	*6	*6	*	*	*	*9	*	*	*	*
443.9	282	153	129	*	*5	58	218	76	76	100	30
444	117	58	60	*	*	34	78	27	29	42	20
444.0	13	*7	*6	*	*	*	*9	*	*	*	*
444.2	78	39	39	*	*	21	53	18	20	26	14
444.21	12	*	*7	*	*	*5	*5	*	*	*7	*
444.22	66	34	32	*	*	16	48	16	17	20	13
444.8	21	9	12	*	*	*8	12	*6	*	*7	*
444.81	14	*7	*7	*	*	*6	*7	*	*	*	*
444.89	*7	*	*5	*	*	*	*5	*	*	*	*
446	39	13	26	*	*7	*	25	10	*8	11	9
446.4	*7	*	*	—	*	*	*	*	*	*	*
446.5	18	*	14	—	*	*	17	*	*6	*	*
447	53	25	28	*	*5	14	33	17	9	19	*7
447.1	29	12	17	*	*	*8	19	10	*7	10	*
447.6	10	*	*5	*	*	*	*	*	*	*	*
447.8	*6	*	*	*	*	*	*	*	*	*	*
448	*8	*	*	—	*	*	*	*	*	*	*
451	101	39	62	*	18	28	54	21	27	36	17
451.0	16	*6	10	*	*	*	9	*	*	*7	*
451.1	45	18	28	*	*6	13	26	10	13	13	10
451.11	15	*6	9	*	*	*	10	*	*	*	*
451.19	31	12	19	*	*6	*9	16	*7	9	*8	*6
451.2	17	*	12	*	*	*	9	*	*	*7	*
451.8	15	*8	*7	*	*	*	*6	*	*	*5	*
451.84	*6	*	*	*	*	*	*	*	*	*	*
451.9	*7	*	*	—	*	*	*	*	*	*	*
453	245	103	142	*	35	70	140	59	66	88	32
453.8	221	93	128	*	32	62	126	56	60	76	29
453.9	17	*7	9	—	*	*	11	*	*	9	*
454	57	20	36	—	*8	12	37	13	19	15	9
454.0	15	*5	10	—	*	*	11	*	*5	*	*
454.2	18	*8	10	—	*	*	13	*	*6	*6	*
454.9	19	*6	14	—	*	*	11	*	*7	*6	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
455 . . . . .	155	66	89	*	39	45	71	53	27	54	22
455.0 . . . . .	45	17	28	—	9	14	22	18	9	12	*6
455.2 . . . . .	35	17	18	*	*9	9	17	12	*	13	*5
455.3 . . . . .	19	*7	12	—	*	*5	*9	*7	*	*5	*
455.5 . . . . .	13	*6	*6	—	*	*	*6	*	*	*5	*
455.6 . . . . .	23	9	14	—	*6	*7	9	*5	*	11	*
455.8 . . . . .	10	*	*6	—	*	*	*6	*	*	*	*
456 . . . . .	55	33	23	*	12	24	19	12	11	21	11
456.0 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
456.1 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
456.2 . . . . .	32	19	12	*	*	16	10	*8	*5	12	*6
456.20 . . . . .	20	12	*7	*	*	10	*6	*6	*	*7	*
456.21 . . . . .	12	*7	*5	*	*	*6	*	*	*	*	*
456.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
457 . . . . .	23	9	14	*	*6	*7	10	9	*5	*6	*
457.1 . . . . .	16	*5	11	*	*	*	*8	*7	*	*	*
458 . . . . .	319	143	176	*	40	78	197	71	72	117	59
458.0 . . . . .	93	37	56	*	12	23	58	23	25	31	15
458.9 . . . . .	225	106	119	*	27	55	139	48	47	85	44
459 . . . . .	107	47	60	*	*7	25	74	25	34	33	15
459.0 . . . . .	10	*	*5	*	*	*	*5	*	*	*	*
459.2 . . . . .	15	*8	*6	*	*	*	9	*	*	*6	*
459.8 . . . . .	67	28	38	*	*	16	48	16	24	20	*8
459.81 . . . . .	59	26	33	*	*	14	42	13	22	18	*7
459.89 . . . . .	*8	*	*5	*	*	*	*6	*	*	*	*
459.9 . . . . .	13	*6	*7	*	*	*	*9	*	*	*	*
460-519 . . . . .	9,338	4,501	4,837	1,248	1,290	1,843	4,958	2,073	2,315	3,503	1,448
460 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
461 . . . . .	84	35	49	18	22	16	28	11	25	40	*8
461.0 . . . . .	28	12	16	*7	*7	*	10	*	10	10	*
461.2 . . . . .	12	*7	*	*	*	*	*	*	*	*	*
461.8 . . . . .	10	*6	*	*	*	*	*	*	*	*	*
461.9 . . . . .	30	9	21	*6	10	*6	*7	*	*7	18	*
462 . . . . .	55	27	28	19	21	*7	*8	9	17	20	*8
463 . . . . .	27	12	15	15	*9	*	*	*	*8	14	*
464 . . . . .	54	33	21	42	*	*	*	13	14	17	*9
464.0 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
464.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
464.4 . . . . .	36	24	12	36	—	*	*	*8	11	11	*5
465 . . . . .	144	64	79	61	37	16	30	38	38	47	20
465.9 . . . . .	140	62	77	58	35	16	30	36	38	46	20
466 . . . . .	450	218	232	210	65	59	116	84	118	178	70
466.0 . . . . .	278	108	169	41	63	57	115	52	81	118	27
466.1 . . . . .	173	110	63	169	*	*	*	32	37	60	43
470 . . . . .	22	10	12	*	12	*8	*	10	*	*6	*
471 . . . . .	10	*6	*	*	*5	*	*	*	*	*	*
472 . . . . .	14	*8	*5	*	*	*	*	*	*5	*5	*
472.0 . . . . .	13	*7	*5	*	*	*	*	*	*5	*	*
473 . . . . .	162	66	95	23	52	42	45	33	43	65	21
473.0 . . . . .	29	11	17	*	*8	*6	10	*7	*7	12	*
473.2 . . . . .	16	*9	*7	*	*	*	*6	*	*	*	*
473.8 . . . . .	13	*7	*6	*	*5	*	*	*	*	*5	*
473.9 . . . . .	97	37	60	13	31	30	23	18	27	40	12
474 . . . . .	47	27	20	35	10	*	*	27	*8	*8	*
474.0 . . . . .	18	*9	9	11	*6	*	—	*8	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
474.1	29	19	10	23	*	*	*	18	*	*	*
474.10	18	11	*7	17	*	*	—	12	*	*	*
474.11	*5	*	*	*	*	*	*	*	*	*	*
475	19	10	*9	*	12	*	*	*	*5	*7	*
477	26	13	12	*	*9	*8	*7	*	*7	11	*
477.9	23	12	10	*	*7	*7	*6	*	*7	10	*
478	81	46	35	18	21	19	23	28	17	24	13
478.0	*7	*	*	*	*	*	*	*	*	*	*
478.1	16	*8	*7	*	*5	*	*	*5	*	*	*
478.2	16	11	*5	*	*6	*5	*	*	*	*	*
478.29	11	*7	*	*	*	*	*	*	*	*	*
478.3	16	*8	*8	*	*	*	*9	*	*	*7	*
478.30	*5	*	*	*	*	*	*	*	*	*	*
478.7	17	9	*8	*8	*	*	*5	*7	*	*	*
478.74	*6	*	*	*	*	*	*	*	*	*	*
478.79	*6	*	*	*	*	*	*	*	*	*	*
480	45	25	20	33	*	*	*6	*7	11	16	11
480.1	21	12	9	21	*	*	—	*	*8	*7	*5
480.9	21	11	10	11	*	*	*6	*	*	*8	*5
481	87	49	39	*	22	18	43	14	23	30	21
482	385	203	182	18	50	61	256	68	84	177	55
482.0	36	19	17	*	*	*	29	*7	*8	16	*
482.1	66	38	28	*	12	11	39	*9	14	31	12
482.2	47	24	23	*	*7	*8	30	*8	11	19	10
482.3	30	17	12	*	*	*	18	*	10	11	*
482.30	15	*8	*7	*	*	*	*7	*	*5	*	*
482.39	10	*7	*	*	*	*	*6	*	*	*	*
482.4	59	29	29	*	*6	10	41	12	13	26	*7
482.8	114	58	55	*	10	18	82	14	22	64	13
482.82	11	*7	*	*	*	*	*8	*	*	*	*
482.83	59	29	30	*	*	10	42	*7	14	31	*7
482.89	41	20	20	*	*	*5	30	*5	*	27	*
482.9	34	17	17	*	*9	*	17	13	*6	11	*
483	19	*8	11	*	*	*	*8	*	*5	*8	*
483.0	17	*7	10	*	*	*	*7	*	*	*8	*
484	*7	*	*	*	*	*	*	*	*	*	*
485	68	36	32	29	*6	*6	27	11	21	32	*
486	1,311	632	679	227	167	197	721	279	330	502	200
487	34	12	21	*6	10	*	13	*6	*8	16	*
487.0	*8	*	*5	*	*	*	*	*	*	*	*
487.1	22	*9	14	*	*6	*	9	*	*	12	*
490	80	34	46	*8	16	18	38	16	19	35	10
491	694	318	376	*	41	181	469	137	166	300	92
491.2	643	290	354	*	30	165	446	125	154	281	83
491.20	126	62	63	*	*9	29	88	29	29	51	17
491.21	518	227	291	*	21	137	358	96	125	231	66
491.8	*7	*	*	*	*	*	*	*	*	*	*
491.9	36	20	17	*	*7	12	17	9	*8	14	*5
492	245	141	104	*	*7	56	182	52	69	94	31
492.0	17	13	*	—	*	*	9	*	*	*6	*
492.8	228	128	100	*	*	52	172	47	66	87	28
493	1,176	444	732	324	342	239	271	321	304	348	203
493.0	53	28	25	34	14	*	*	17	*9	9	18
493.00	28	15	13	17	*7	*	*	*6	*5	*6	11
493.01	25	13	12	18	*7	*	—	11	*	*	*7

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
493.2	210	75	134	*	19	65	124	51	52	74	33
493.20	183	67	116	*	17	53	111	44	45	65	29
493.21	27	*9	18	*	*	11	12	*7	*7	*9	*
493.9	908	339	569	287	307	169	145	252	242	264	151
493.90	683	239	445	163	248	142	132	184	176	204	119
493.91	225	100	125	125	60	28	13	68	66	60	32
494	35	13	22	*	*5	*5	23	*9	*	15	*6
496	1,682	898	785	*	44	363	1,274	366	413	649	254
500	16	14	*	—	*	*	12	11	*	*	*
501	9	*9	*	—	*	*	*7	*	*	*	*
506	*5	*	*	*	*	*	*	*	*	*	*
507	232	126	107	14	18	29	171	56	50	80	47
507.0	230	125	105	13	18	29	170	55	49	80	47
508	*6	*	*	*	*	*	*	*	*	*	*
510	18	11	*7	*	*5	*6	*6	*	*	*7	*
510.9	15	*9	*6	*	*	*	*5	*	*	*5	*
511	426	182	244	14	48	96	268	96	118	147	65
511.0	33	14	18	*	12	*8	13	*	12	12	*5
511.1	*6	*	*	*	*	*	*	*	*	*	*
511.8	32	18	15	*	*	*8	18	*7	*6	*9	11
511.9	354	148	207	11	31	78	234	85	98	124	47
512	106	58	48	*	27	29	46	24	24	42	16
512.0	*6	*	*	*	*	*	*	*	*	*	*
512.1	51	27	24	*	*6	15	28	12	10	19	9
512.8	49	27	22	*	18	13	15	9	11	22	*6
513	14	*8	*6	*	*	*	*6	*	*	*7	*
513.0	14	*8	*6	*	*	*	*6	*	*	*7	*
514	35	16	19	*	*9	*6	20	*8	*8	13	*6
515	179	78	101	*	*8	29	141	26	61	67	25
516	32	17	15	*	*	*9	17	*7	*8	13	*
516.3	19	10	10	*	*	*	13	*	*	9	*
516.8	11	*6	*	*	*	*	*	*	*	*	*
517	20	*	16	—	*	*8	*8	*	*6	*	*6
517.8	16	*	14	—	*	*	*6	*	*6	*	*
518	1,069	522	546	65	139	255	610	243	230	404	191
518.0	349	167	183	30	54	87	179	75	84	132	59
518.1	*7	*	*	*	*	*	*	*	*	*	*
518.3	17	*7	10	*	*	*	12	*	*	*6	*
518.4	27	15	13	*	*	*8	14	*5	*8	10	*
518.5	63	35	28	*	14	19	28	13	10	27	12
518.8	605	295	309	29	62	138	375	145	121	228	111
518.81	437	218	219	9	39	95	294	104	81	169	83
518.82	92	42	50	15	14	22	42	19	24	35	15
518.89	75	35	40	*5	10	20	39	22	16	24	13
519	94	47	46	30	14	14	36	24	25	35	10
519.0	*8	*	*	*	*	*	*	*	*	*	*
519.1	43	23	20	17	*6	*	15	10	17	11	*
519.8	32	14	18	*8	*5	*	14	*7	*	18	*
519.9	*6	*	*	*	*	*	*	*	*	*	*
520-579	8,228	3,520	4,708	445	1,787	2,080	3,916	1,868	1,953	3,049	1,358
520	*7	*	*	*	*6	*	*	*	*	*	*
521	20	13	*7	*	10	*	*	*6	*	*7	*
521.0	20	13	*7	*	10	*	*	*5	*	*7	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
522 . . . . .	18	*8	9	*	10	*	*	*	*	*7	*
522.5 . . . . .	14	*6	*8	*	*8	*	*	*	*	*6	*
523 . . . . .	14	*8	*6	*	*6	*	*	*	*	*6	*
524 . . . . .	39	12	27	*	30	*	*	12	*7	15	*
524.0 . . . . .	13	*5	*8	*	10	*	*	*	*	*6	*
524.1 . . . . .	*8	*	*7	*	*7	*	—	*	*	*	*
524.10 . . . . .	*7	*	*5	*	*6	*	—	*	*	*	*
524.6 . . . . .	9	*	*7	*	*7	*	*	*	*	*	*
525 . . . . .	*9	*	*5	*	*	*	*	*	*	*	*
526 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
527 . . . . .	19	*8	11	*	*6	*5	*8	*5	*	*8	*
527.2 . . . . .	14	*6	*9	*	*	*	*7	*	*	*7	*
528 . . . . .	39	18	20	*6	16	*7	9	10	*8	14	*7
528.0 . . . . .	20	*8	12	*	*6	*	*6	*7	*	*5	*
528.3 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
528.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
529 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
530 . . . . .	861	350	511	63	137	221	440	193	197	323	148
530.0 . . . . .	*9	*	*	*	*	*	*6	*	*	*	*
530.1 . . . . .	258	113	145	9	54	62	133	58	58	96	48
530.10 . . . . .	116	56	60	*6	25	27	58	27	26	45	17
530.11 . . . . .	111	40	71	*	23	27	59	25	25	38	23
530.19 . . . . .	31	17	14	*	*6	*8	15	*5	*6	13	*8
530.2 . . . . .	62	31	31	*	9	19	34	16	11	23	12
530.3 . . . . .	67	30	37	*	*	14	49	17	14	23	14
530.5 . . . . .	26	10	15	*	*	*9	13	*7	*	9	*7
530.6 . . . . .	*9	*	*	—	*	*	*7	*	*	*	*
530.7 . . . . .	20	11	*9	*	*6	*7	*6	*	*	9	*
530.8 . . . . .	408	144	263	52	60	106	190	87	103	158	60
530.81 . . . . .	379	135	244	52	58	98	171	82	97	148	53
530.82 . . . . .	14	*5	*9	*	*	*5	*7	*	*	*	*
530.89 . . . . .	14	*	10	*	*	*	11	*	*	*	*
531 . . . . .	192	85	107	*	25	46	118	38	46	73	35
531.0 . . . . .	34	15	19	*	*	*5	24	*6	*8	14	*6
531.00 . . . . .	34	15	19	*	*	*5	24	*6	*8	14	*6
531.3 . . . . .	18	*8	10	—	*	*6	*8	*	*	*8	*
531.30 . . . . .	17	*8	9	—	*	*6	*7	*	*	*7	*
531.4 . . . . .	68	32	35	*	*7	16	44	14	15	26	13
531.40 . . . . .	65	31	34	*	*7	15	43	14	13	25	13
531.5 . . . . .	*7	*	*5	*	*	*	*	*	*	*	*
531.50 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
531.7 . . . . .	*7	*	*	—	*	*	*5	*	*	*	*
531.70 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
531.9 . . . . .	52	23	29	*	*8	15	28	10	15	19	*7
531.90 . . . . .	51	22	28	*	*7	15	27	10	15	19	*7
532 . . . . .	136	85	51	*	23	32	79	29	34	48	25
532.0 . . . . .	24	17	*7	*	*	*5	16	*6	*	*8	*6
532.00 . . . . .	23	17	*6	*	*	*5	14	*6	*	*7	*6
532.3 . . . . .	12	*8	*	—	*	*	*7	*	*	*6	*
532.30 . . . . .	12	*8	*	—	*	*	*7	*	*	*6	*
532.4 . . . . .	45	32	13	—	*7	11	26	10	11	15	9
532.40 . . . . .	42	30	12	—	*7	10	24	*9	10	14	*9
532.5 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
532.9 . . . . .	38	20	18	*	*8	*9	20	*7	11	13	*6
532.90 . . . . .	37	20	17	*	*8	*9	20	*7	11	13	*6

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
533 . . . . .	219	91	128	*	35	64	119	57	65	73	23
533.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
533.00 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
533.4 . . . . .	18	*8	10	—	*	*	13	*	*	*7	*
533.40 . . . . .	18	*8	10	—	*	*	13	*	*	*7	*
533.7 . . . . .	24	10	14	—	*	*7	13	*6	*8	*6	*
533.70 . . . . .	24	10	14	—	*	*7	13	*6	*8	*6	*
533.9 . . . . .	164	68	96	*	29	49	86	47	46	55	16
533.90 . . . . .	164	68	96	*	29	49	86	47	46	55	16
535 . . . . .	666	281	386	17	144	165	340	147	172	251	96
535.0 . . . . .	107	37	70	*	23	24	58	18	33	43	14
535.00 . . . . .	85	27	58	*	20	19	42	13	28	36	*8
535.01 . . . . .	23	10	13	*	*	*	15	*5	*	*7	*6
535.1 . . . . .	67	22	44	*	*9	13	44	21	13	24	*9
535.10 . . . . .	58	19	39	*	*8	11	39	19	11	22	*7
535.11 . . . . .	*8	*	*5	*	*	*	*5	*	*	*	*
535.3 . . . . .	29	20	*9	—	15	11	*	*8	10	9	*
535.30 . . . . .	22	15	*7	—	11	*9	*	*6	*8	*5	*
535.31 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
535.4 . . . . .	130	50	80	*	22	29	77	29	38	40	23
535.40 . . . . .	93	33	60	*	16	23	51	20	26	30	16
535.41 . . . . .	37	17	20	*	*6	*6	26	9	11	10	*7
535.5 . . . . .	237	101	136	*9	52	61	115	52	53	101	32
535.50 . . . . .	206	85	121	*8	48	53	97	46	48	85	28
535.51 . . . . .	31	15	16	*	*	*8	18	*6	*5	16	*
535.6 . . . . .	94	49	45	*	23	28	42	20	24	34	16
535.60 . . . . .	82	44	38	*	19	26	36	17	22	29	14
535.61 . . . . .	12	*	*7	*	*	*	*6	*	*	*5	*
536 . . . . .	182	71	111	*7	43	53	79	35	38	81	29
536.2 . . . . .	27	10	17	*5	*8	*6	*8	*5	*	14	*
536.3 . . . . .	38	14	24	*	13	10	15	10	*8	12	*8
536.8 . . . . .	89	35	54	*	18	27	43	16	21	38	14
536.9 . . . . .	27	12	15	*	*	9	13	*	*	17	*
537 . . . . .	50	19	31	*	*7	11	30	15	*8	15	12
537.0 . . . . .	23	9	14	*	*	*5	13	*6	*	*7	*7
537.8 . . . . .	21	*8	14	*	*	*	14	*8	*	*6	*
537.83 . . . . .	*8	*	*6	—	*	*	*6	*	*	*	*
537.89 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
540 . . . . .	203	120	83	47	115	26	15	41	44	69	49
540.0 . . . . .	48	31	18	13	20	9	*6	9	11	18	10
540.1 . . . . .	28	20	*8	*7	12	*	*5	*6	*6	9	*7
540.9 . . . . .	127	70	57	27	83	13	*	26	27	42	31
541 . . . . .	19	9	9	*	12	*	*	*	*	*8	*
542 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
543 . . . . .	14	*6	*8	*	*8	*	*	*	*5	*	*
543.9 . . . . .	9	*	*	*	*6	*	*	*	*	*	*
550 . . . . .	104	93	10	12	14	23	55	37	25	30	12
550.1 . . . . .	28	24	*	*	*	*	17	*9	*	9	*
550.10 . . . . .	22	19	*	*	*	*	13	*7	*	*8	*
550.9 . . . . .	76	69	*6	9	10	19	38	28	20	20	*8
550.90 . . . . .	52	47	*	*	*8	13	27	19	13	15	*5
550.91 . . . . .	9	*9	*	*	*	*	*6	*	*	*	*
550.92 . . . . .	12	12	*	*	*	*	*	*6	*	*	*
552 . . . . .	59	19	39	*	10	20	28	17	14	19	*9
552.0 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
552.00 . . . . .	*5	*	*	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
552.1	14	*	10	*	*	*	*6	*	*	*	*
552.2	29	9	20	*	*	12	13	*9	*6	9	*
552.20	*8	*	*7	*	*	*	*	*	*	*	*
552.21	17	*	12	—	*	*7	*7	*6	*	*	*
553	547	209	338	*7	77	148	314	126	140	208	72
553.1	50	23	27	*	10	15	21	12	10	18	10
553.2	84	35	49	*	15	29	39	16	24	34	10
553.20	27	*8	19	*	*	*8	14	*	9	10	*
553.21	52	24	27	*	10	20	22	10	14	22	*6
553.29	*5	*	*	*	*	*	*	*	*	*	*
553.3	406	148	258	*	51	102	250	96	104	154	52
555	80	34	46	*	36	22	20	20	20	31	9
555.0	14	*7	*7	*	*7	*	*	*	*	*	*
555.1	12	*5	*7	*	*	*	*	*	*	*	*
555.2	*6	*	*	*	*	*	*	*	*	*	*
555.9	49	20	29	*	22	13	14	11	12	21	*
556	47	20	27	*	19	11	15	12	15	13	*7
556.9	32	13	19	*	14	*6	11	*7	11	10	*
557	61	17	43	*	*	14	41	15	13	21	12
557.0	35	11	25	*	*	*8	23	*9	*6	13	*7
557.1	10	*	*7	*	*	*	*7	*	*	*	*
557.9	15	*	12	—	*	*	11	*	*	*	*
558	582	242	340	159	134	103	185	122	136	231	94
558.1	*9	*	*6	—	*	*	*6	*	*	*	*
558.2	*7	*	*	*	*	*	*	*	*	*	*
558.9	567	237	330	159	133	99	177	115	133	227	92
560	613	269	345	23	87	140	364	116	148	234	115
560.0	*5	*	*	*	*	*	*	*	*	*	*
560.1	252	116	136	9	46	57	139	44	64	100	45
560.2	17	11	*6	*	*	*	12	*	*	*8	*
560.3	87	36	51	*	*	*8	72	17	20	39	12
560.39	83	35	48	*	*	*7	68	17	19	36	11
560.8	120	50	70	*	18	34	65	26	26	38	30
560.81	78	32	46	*	13	23	40	16	17	28	17
560.89	42	18	24	*	*5	11	25	10	*8	11	12
560.9	132	53	79	*	17	37	75	24	35	47	25
562	495	194	302	*	27	109	359	131	110	185	69
562.0	*8	*	*5	—	*	*	*7	*	*	*	*
562.00	*7	*	*	—	*	*	*5	*	*	*	*
562.1	487	191	297	*	26	108	353	130	108	183	67
562.10	233	88	146	*	10	44	180	65	47	87	34
562.11	162	64	98	—	15	51	96	42	42	57	22
562.12	70	30	40	—	*	10	60	19	13	30	*9
562.13	22	*9	13	—	*	*	17	*	*6	9	*
564	264	91	172	17	53	53	141	57	69	95	43
564.0	167	67	100	13	24	32	97	37	43	60	26
564.1	70	12	58	*	22	16	30	17	21	23	*8
564.8	14	*7	*7	*	*	*	*7	*	*	*6	*
565	20	11	*9	*	*8	*6	*	*	*	10	*
565.0	10	*5	*5	*	*	*	*	*	*	*5	*
565.1	10	*6	*	*	*	*	*	*	*	*5	*
566	26	18	*9	*	12	*8	*	*6	*	10	*5
567	74	35	39	*	18	25	30	18	14	26	17
567.2	52	26	27	*	13	18	20	12	10	18	12
567.8	*6	*	*	*	*	*	*	*	*	*	*
567.9	16	*6	10	*	*	*6	*6	*	*	*6	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
568 . . . . .	141	36	105	*	55	34	49	35	34	49	23
568.0 . . . . .	123	33	90	*	45	32	44	31	30	42	19
568.8 . . . . .	18	*	15	*	11	*	*	*	*	*6	*
568.81 . . . . .	11	*	10	*	*8	*	*	*	*	*	*
568.89 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
569 . . . . .	212	96	117	*	34	59	115	62	47	69	34
569.0 . . . . .	13	*7	*6	*	*	*	*7	*	*	*	*
569.1 . . . . .	*9	*	*6	*	*	*	*5	*	*	*	*
569.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
569.3 . . . . .	37	15	21	*	*6	*8	22	11	*6	15	*6
569.4 . . . . .	28	14	14	*	*7	*7	14	11	*6	*7	*
569.41 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
569.49 . . . . .	22	11	11	*	*6	*6	10	*9	*	*5	*
569.5 . . . . .	15	*6	*9	—	*	*8	*	*	*	*	*
569.6 . . . . .	18	*8	9	*	*	*5	9	*6	*	*7	*
569.8 . . . . .	83	38	45	*	9	24	48	21	22	27	14
569.81 . . . . .	12	*	*7	*	*	*	*	*	*	*	*
569.82 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
569.83 . . . . .	20	10	10	*	*	*	10	*	*6	*5	*
569.84 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
569.85 . . . . .	22	11	11	—	*	*6	16	*	*5	*8	*
569.89 . . . . .	15	*6	9	*	*	*	*8	*6	*	*	*
570 . . . . .	15	*8	*7	*	*	*	*5	*	*	*	*
571 . . . . .	334	194	141	*	89	138	104	76	79	119	60
571.0 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
571.1 . . . . .	52	37	15	—	28	21	*	9	19	14	10
571.2 . . . . .	98	67	31	—	25	46	28	24	25	33	15
571.3 . . . . .	31	22	10	—	10	16	*6	*6	*7	9	9
571.4 . . . . .	19	*8	11	*	*8	*	*6	*	*	*6	*
571.40 . . . . .	10	*	*6	*	*	*	*	*	*	*	*
571.49 . . . . .	9	*	*	*	*	*	*	*	*	*	*
571.5 . . . . .	101	47	54	*	*9	39	51	25	19	42	16
571.6 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
571.8 . . . . .	15	*6	9	*	*6	*	*	*	*	*7	*
572 . . . . .	105	59	46	*	21	40	41	22	21	42	19
572.0 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
572.2 . . . . .	44	26	17	*	*8	18	17	*8	9	19	*7
572.3 . . . . .	33	17	16	*	*6	15	11	*8	*7	12	*6
572.4 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
572.8 . . . . .	13	*5	*7	*	*	*	*6	*	*	*	*
573 . . . . .	77	33	44	*	22	23	31	14	18	30	15
573.0 . . . . .	10	*6	*	—	*	*	*7	*	*	*5	*
573.3 . . . . .	28	10	18	*	12	10	*5	*6	*6	9	*6
573.8 . . . . .	23	10	13	*	*	*7	13	*	*5	*8	*5
573.9 . . . . .	13	*5	*8	*	*	*	*5	*	*	*7	*
574 . . . . .	721	227	494	*	197	198	322	160	165	263	132
574.0 . . . . .	160	62	98	*	43	50	67	29	43	62	27
574.00 . . . . .	144	57	87	*	40	44	60	26	39	56	23
574.01 . . . . .	16	*5	11	*	*	*6	*7	*	*	*6	*
574.1 . . . . .	298	81	217	*	99	90	108	76	58	108	56
574.10 . . . . .	287	77	210	*	96	86	104	74	56	104	53
574.11 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
574.2 . . . . .	164	52	112	*	31	42	90	35	42	60	27
574.20 . . . . .	158	49	109	*	30	40	86	35	41	58	24
574.21 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
574.3 . . . . .	27	12	16	*	*6	*	19	*5	*6	14	*
574.30 . . . . .	13	*5	*7	*	*	*	*7	*	*	*6	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
574.31	15	*6	*8	—	*	*	11	*	*	*8	*
574.4	27	9	18	*	*9	*5	12	*6	*7	*8	*6
574.40	16	*	11	*	*6	*	*7	*	*	*5	*
574.41	11	*	*6	—	*	*	*6	*	*	*	*
574.5	44	11	33	*	10	*8	26	*8	10	12	13
574.50	25	*7	17	*	*6	*	14	*	*8	*5	*7
574.51	20	*	16	*	*	*	12	*	*	*7	*6
575	122	44	78	*	47	34	41	31	28	43	19
575.0	29	14	15	*	9	*8	11	*8	*9	*8	*
575.1	50	15	36	*	21	16	14	11	12	22	*6
575.3	*6	*	*	*	*	*	*	*	*	*	*
575.6	13	*	10	—	*7	*	*	*	*	*	*5
575.8	17	*6	11	*	*6	*5	*6	*5	*	*7	*
576	87	36	51	*	12	18	55	22	14	29	22
576.1	33	16	17	*	*5	*5	22	9	*6	9	*9
576.2	25	*8	17	*	*	*7	15	*6	*	*9	*8
576.8	25	11	14	*	*	*6	15	*7	*	9	*
577	325	148	176	*7	113	104	102	58	86	126	54
577.0	226	103	123	*	78	70	73	40	63	88	36
577.1	63	28	35	*	24	21	18	11	15	25	12
577.2	19	11	*8	*	*7	*7	11	*	*	*8	*
577.8	13	*	*9	*	*	*	*	*	*	*	*
578	351	172	180	11	43	79	218	83	81	137	50
578.0	41	22	18	*	*7	*9	24	10	*7	16	*7
578.1	90	44	46	*	10	22	54	21	22	33	14
578.9	220	106	115	*5	26	48	141	51	52	88	29
579	31	11	20	*9	*	*7	12	*8	10	*9	*
579.3	13	*	10	*	*	*	*5	*	*	*	*
579.8	9	*	*	*	*	*	*	*	*	*	*
580-629	6,715	2,247	4,468	148	2,030	1,453	3,083	1,427	1,586	2,636	1,065
580	16	*7	*9	*	*	*	9	*	*5	*6	*
580.9	13	*6	*7	*	*	*	*9	*	*	*	*
581	34	13	20	*	9	12	9	*6	*6	16	*
581.8	17	*6	11	*	*5	*7	*	*	*	*8	*
581.81	17	*6	11	*	*5	*7	*	*	*	*8	*
581.9	14	*7	*7	*	*	*	*	*	*	*7	*
582	64	33	32	*	*7	12	44	19	17	21	*7
582.8	*7	*	*	*	*	*	*	*	*	*	*
582.9	53	28	25	*	*	*9	40	16	15	17	*
583	139	65	74	*	27	45	65	37	29	49	25
583.8	127	58	68	*	23	41	62	35	26	45	21
583.81	120	55	64	*	20	38	60	33	25	41	20
583.89	*7	*	*	*	*	*	*	*	*	*	*
583.9	11	*6	*	*	*	*	*	*	*	*	*
584	237	130	107	*	26	48	160	62	53	85	38
584.5	27	16	11	*	*5	*7	14	*8	*7	10	*
584.9	206	112	94	*	19	41	144	53	45	73	35
585	305	156	148	*	41	75	186	83	70	93	58
586	73	33	40	*	*6	12	55	12	13	30	18
588	14	*6	*7	*	*	*	*	*	*	*6	*
588.8	10	*	*5	*	*	*	*	*	*	*	*
590	186	36	149	21	86	32	47	32	49	69	35
590.0	*8	*	*	*	*	*	*	*	*	*	*
590.00	*8	*	*	*	*	*	*	*	*	*	*
590.1	108	23	85	12	53	16	28	18	33	35	22

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
590.10	107	22	85	12	53	16	27	18	33	35	22
590.2	*6	*	*	*	*	*	*	*	*	*	*
590.8	61	*9	52	*8	27	12	14	12	11	27	11
590.80	61	*9	52	*8	27	12	14	12	11	27	11
591	120	57	63	*5	30	25	60	34	28	37	21
592	270	167	103	*	112	97	58	61	67	104	39
592.0	108	61	47	*	41	35	31	26	25	40	18
592.1	156	103	54	*	69	61	25	34	41	62	20
592.9	*5	*	*	—	*	*	*	*	*	*	*
593	345	185	160	16	40	59	231	81	88	116	61
593.2	26	14	12	*	*	*5	19	*8	*5	9	*
593.3	10	*	*7	*	*	*	*	*	*	*	*
593.4	32	15	17	*	*8	*7	15	*8	*6	10	*8
593.7	13	*	*8	9	*	*	*	*	*	*	*
593.70	10	*	*7	*9	*	*	*	*	*	*	*
593.8	29	17	12	*	*7	*	15	*7	*8	10	*
593.89	25	14	10	*	*6	*	13	*6	*7	*8	*
593.9	230	129	102	*	15	38	175	49	62	79	40
594	23	20	*	*	*	*7	13	*8	*5	*	*6
594.1	18	16	*	*	*	*5	12	*7	*	*	*
595	65	17	48	*	10	9	44	18	19	19	9
595.0	16	*	12	*	*	*	13	*	*	*	*
595.1	*7	*	*	—	*	*	*	*	*	*	*
595.2	9	*	*6	*	*	*	*7	*	*	*	*
595.8	*8	*	*	*	*	*	*6	*	*	*	*
595.9	23	*	19	*	*5	*	14	*	*6	*9	*
596	202	143	59	*	19	32	148	46	57	69	30
596.0	71	70	*	—	*	*9	61	12	20	25	13
596.1	*5	*	*	—	*	*	*	*	*	*	*
596.3	14	13	*	*	*	*	11	*7	*	*	*
596.4	*9	*5	*	*	*	*	*7	*	*	*	*
596.5	71	32	38	*	12	14	43	16	21	23	11
596.54	64	31	34	*	*9	12	41	14	17	21	11
596.8	26	16	10	*	*5	*	16	*5	10	10	*
597	*8	*	*5	—	*	*	*	*	*	*	*
597.8	*8	*	*5	—	*	*	*	*	*	*	*
598	50	42	*9	*	*5	11	33	10	12	21	*7
598.8	*8	*7	*	*	*	*	*5	*	*	*	*
598.9	40	33	*7	*	*	*9	26	*8	10	17	*6
599	1,841	603	1,237	56	246	271	1,267	401	463	725	252
599.0	1,557	426	1,131	51	190	216	1,101	333	392	619	212
599.6	25	20	*5	*	*	*	18	*5	*8	*5	*6
599.7	245	152	92	*	50	49	141	59	60	94	31
599.8	*6	*	*	*	*	*	*	*	*	*	*
600	350	350	—	*	*	62	286	83	88	125	55
601	67	67	—	—	*8	12	47	14	12	31	10
601.0	26	26	—	—	*	*	17	*	*	13	*
601.1	23	23	—	—	*	*	19	*5	*5	*9	*
601.9	16	16	—	—	*	*	10	*	*	*9	*
602	*8	*8	—	—	*	*	*7	*	*	*	*
603	15	15	—	*	*	*	*8	*	*6	*5	*
603.9	14	14	—	*	*	*	*8	*	*5	*	*
604	21	21	—	*	*5	*	12	*	*6	10	*
604.9	21	21	—	*	*5	*	11	*	*5	10	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
604.90	13	13	—	*	*	*	*6	*	*	*6	*
604.99	*8	*8	—	*	*	*	*	*	*	*	*
605	13	13	—	*	*	*	*	*	*	*5	*
607	25	25	—	*	*	9	10	*	*6	11	*
607.8	16	16	—	*	*	*7	*6	*	*	*8	*
607.84	11	11	—	—	*	*	*	*	*	*5	*
608	21	21	—	*	*7	*6	*5	*7	*	*7	*
608.8	*9	*9	—	*	*	*	*	*	*	*	*
610	42	*	41	*	14	20	*8	15	*6	15	*6
610.1	34	*	33	—	12	16	*6	11	*5	12	*6
611	66	*	61	*	30	18	15	18	13	24	10
611.0	*9	*	*8	*	*	*	*	*	*	*	*
611.1	28	*	24	*	19	*6	*	*9	*	11	*
611.7	18	*	18	—	*	*6	*8	*5	*	*6	*
611.72	16	*	16	—	*	*6	*8	*	*	*6	*
611.8	*9	—	*9	—	*	*	*	*	*	*	*
614	292	—	292	*	215	58	17	55	61	134	42
614.0	*7	—	*7	—	*6	*	*	*	*	*	*
614.1	30	—	30	—	21	*7	*	*5	*6	11	*7
614.2	16	—	16	*	11	*	*	*	*	*7	*
614.3	16	—	16	*	14	*	*	*6	*	*6	*
614.4	10	—	10	—	*7	*	*	*	*	*	*
614.6	179	—	179	*	127	42	10	31	37	85	27
614.9	30	—	30	*	27	*	—	*6	*7	16	*
615	19	—	19	—	16	*	*	*	*	*7	*
615.0	*5	—	*5	—	*	*	*	*	*	*	*
615.9	10	—	10	—	10	*	*	*	*	*	*
616	142	—	142	*	87	42	12	29	32	61	21
616.0	109	—	109	—	67	34	*7	22	23	47	17
616.1	27	—	27	*	16	*6	*	*5	*6	12	*
616.10	27	—	27	*	16	*6	*	*5	*6	12	*
617	287	—	287	—	202	77	*8	46	61	129	51
617.0	165	—	165	—	100	58	*7	26	35	73	31
617.1	43	—	43	—	33	9	*	9	*8	17	*8
617.2	11	—	11	—	*8	*	*	*	*	*	*
617.3	44	—	44	—	38	*6	*	*7	12	19	*7
617.9	19	—	19	—	18	*	*	*	*	14	*
618	278	—	278	*	70	104	104	46	61	116	54
618.0	81	—	81	—	14	36	31	12	23	30	17
618.1	33	—	33	—	18	*8	*6	*6	*6	15	*5
618.2	32	—	32	—	10	11	11	*9	*5	14	*
618.3	22	—	22	—	*	*7	11	*6	*5	*6	*
618.4	30	—	30	—	11	12	*7	*	*5	17	*
618.5	25	—	25	—	*	*8	15	*	*6	10	*6
618.6	31	—	31	—	*	12	15	*	*8	14	*
618.8	22	—	22	—	*7	9	*6	*	*	*9	*8
619	13	—	13	*	*7	*	*	*	*	*	*
619.1	11	—	11	—	*6	*	*	*	*	*	*
620	276	—	276	*	193	65	14	56	66	109	45
620.0	42	—	42	*	32	*9	*	*8	10	18	*5
620.1	63	—	63	*	48	15	—	11	19	22	11
620.2	124	—	124	*	88	25	*9	28	28	50	18
620.5	*6	—	*6	*	*	*	*	*	*	*	*
620.8	38	—	38	*	19	14	*	*7	*6	16	*9
621	89	—	89	*	42	32	15	18	21	34	15
621.0	25	—	25	—	*6	14	*	*5	*6	*8	*6

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
621.2	11	—	11	—	*8	*	*	*	*	*5	*
621.3	27	—	27	—	10	11	*6	*7	*6	9	*
621.5	9	—	9	—	*7	*	*	*	*	*	*
621.8	12	—	12	—	*7	*	*	*	*	*	*
622	39	—	39	—	28	*7	*	*6	*8	20	*5
622.1	25	—	25	—	22	*	*	*	*	15	*
622.7	*7	—	*7	—	*	*	*	*	*	*	*
623	17	—	17	*	*8	*	*	*	*	*	*
623.8	12	—	12	*	*5	*	*	*	*	*	*
624	*7	—	*7	*	*	*	*	*	*	*	*
625	331	—	331	*	214	78	38	35	65	166	64
625.0	23	—	23	—	23	*	*	*	*	11	*7
625.3	70	—	70	*	63	*7	—	*	11	44	11
625.6	141	—	141	—	47	61	33	18	27	63	33
625.8	*6	—	*6	—	*	*	*	*	*	*	*
625.9	82	—	82	*	72	*8	*	*8	19	43	12
626	236	—	236	*	172	59	*	33	50	115	37
626.0	*6	—	*6	*	*6	*	*	*	*	*	*
626.2	172	—	172	*	128	42	*	23	34	86	29
626.8	40	—	40	*	28	10	*	*7	11	18	*
626.9	*6	—	*6	—	*	*	*	*	*	*	*
627	45	—	45	—	*5	25	15	*9	12	17	*9
627.1	22	—	22	—	*	13	9	*5	*5	*8	*
627.2	11	—	11	—	*	*7	*	*	*	*	*
628	12	—	12	—	12	—	—	*	*	*	*
628.2	*6	—	*6	—	*6	—	—	*	*	*	*
630-676	7,858	—	7,858	21	7,831	*6	—	1,384	1,728	2,777	1,969
632	17	—	17	—	17	—	—	*	*	9	*
633	42	—	42	—	42	*	—	12	*7	15	*8
633.1	38	—	38	—	38	*	—	10	*7	14	*7
634	40	—	40	*	39	*	—	19	*	13	*
634.9	33	—	33	*	33	*	—	16	*	11	*
634.91	25	—	25	*	25	*	—	14	*	*8	*
635	*7	—	*7	*	*7	*	—	*	*	*	*
635.9	*5	—	*5	*	*	*	—	*	*	*	*
637	*7	—	*7	—	*7	—	—	*	*	*	*
637.9	*6	—	*6	—	*6	—	—	*	*	*	*
639	*8	—	*8	—	*8	*	—	*	*	*	*
640	9	—	9	—	9	—	—	*	*	*	*
640.0	*6	—	*6	—	*6	—	—	*	*	*	*
640.03	*6	—	*6	—	*6	—	—	*	*	*	*
641	95	—	95	*	95	—	—	16	18	37	25
641.0	*5	—	*5	—	*5	—	—	*	*	*	*
641.1	21	—	21	*	21	—	—	*	*	*6	*6
641.11	15	—	15	*	14	—	—	*	*	*5	*
641.13	*6	—	*6	—	*6	—	—	*	*	*	*
641.2	49	—	49	*	48	—	—	*7	*9	22	12
641.21	42	—	42	*	42	—	—	*6	*8	19	9
641.23	*6	—	*6	—	*6	—	—	*	*	*	*
641.9	14	—	14	—	14	—	—	*	*	*	*
641.91	*6	—	*6	—	*6	—	—	*	*	*	*
641.93	*8	—	*8	—	*8	—	—	*	*	*	*
642	272	—	272	*	270	*	—	40	62	112	58
642.0	25	—	25	—	25	—	—	*	*6	11	*5
642.01	21	—	21	—	21	—	—	*	*5	9	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
642.3	78	—	78	*	77	*	—	11	15	31	21
642.31	67	—	67	*	66	*	—	*9	14	25	19
642.33	9	—	9	—	9	—	—	*	*	*5	*
642.4	96	—	96	*	95	*	—	14	24	40	18
642.41	82	—	82	*	81	*	—	12	20	34	17
642.43	11	—	11	—	11	—	—	*	*	*	*
642.5	27	—	27	—	27	—	—	*5	*	9	*7
642.51	25	—	25	—	25	—	—	*	*	*9	*7
642.6	*6	—	*6	*	*6	—	—	*	*	*	*
642.7	*6	—	*6	—	*6	—	—	*	*	*	*
642.71	*5	—	*5	—	*5	—	—	*	*	*	*
642.9	30	—	30	*	30	*	—	*	9	13	*
642.91	25	—	25	—	24	*	—	*	*8	10	*
643	52	—	52	*	52	—	—	10	11	23	*8
643.0	18	—	18	—	18	—	—	*	*6	*8	*
643.03	16	—	16	—	16	—	—	*	*5	*7	*
643.1	26	—	26	*	26	—	—	*6	*	12	*
643.13	25	—	25	*	25	—	—	*6	*	11	*
644	394	—	394	*	393	*	—	65	84	157	89
644.0	123	—	123	*	123	—	—	20	28	49	26
644.03	123	—	123	*	122	—	—	20	28	49	26
644.1	36	—	36	—	36	—	—	*6	*7	12	11
644.13	36	—	36	—	36	—	—	*5	*7	12	11
644.2	235	—	235	*	234	*	—	39	49	95	52
644.21	233	—	233	*	233	*	—	39	49	94	51
645	151	—	151	*	151	*	—	21	27	52	51
645.0	151	—	151	*	151	*	—	21	27	52	51
645.01	147	—	147	*	146	*	—	21	27	50	49
646	223	—	223	*	221	—	—	36	39	97	51
646.1	13	—	13	—	13	—	—	*	*	*6	*
646.11	12	—	12	—	12	—	—	*	*	*5	*
646.2	9	—	9	*	9	—	—	*	*	*	*
646.21	*6	—	*6	*	*6	—	—	*	*	*	*
646.6	159	—	159	*	158	—	—	26	25	73	35
646.61	92	—	92	*	91	—	—	14	13	45	19
646.62	*8	—	*8	—	*8	—	—	*	*	*	*
646.63	54	—	54	*	53	—	—	10	10	22	11
646.64	*5	—	*5	*	*5	—	—	*	*	*	*
646.8	33	—	33	*	32	—	—	*6	*9	14	*
646.81	17	—	17	—	17	—	—	*	*	*7	*
646.83	14	—	14	*	14	—	—	*	*	*6	*
647	87	—	87	*	87	—	—	14	19	34	19
647.0	*6	—	*6	—	*6	—	—	*	*	*	*
647.6	38	—	38	*	38	—	—	*8	*8	13	10
647.61	30	—	30	*	29	—	—	*7	*6	*9	*8
647.63	*8	—	*8	*	*7	—	—	*	*	*	*
647.8	33	—	33	*	33	—	—	*5	*8	13	*7
647.81	22	—	22	*	22	—	—	*	*	9	*6
647.83	10	—	10	—	10	—	—	*	*	*	*
648	784	—	784	*	782	*	—	157	170	280	177
648.0	31	—	31	—	31	—	—	*	*7	13	*6
648.01	20	—	20	—	20	—	—	*	*	10	*
648.03	10	—	10	—	10	—	—	*	*	*	*
648.1	22	—	22	—	22	—	—	*	*	*8	*
648.11	19	—	19	—	19	—	—	*	*	*8	*
648.2	338	—	338	*	336	—	—	58	70	128	81
648.21	190	—	190	*	189	—	—	26	40	86	37

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
648.22	113	—	113	*	113	—	—	27	22	28	36
648.23	26	—	26	*	26	—	—	*	*6	9	*6
648.24	*7	—	*7	*	*7	—	—	*	*	*	*
648.3	19	—	19	—	19	—	—	*	*	*7	*
648.31	14	—	14	—	14	—	—	*	*	*	*
648.4	61	—	61	*	60	—	—	13	12	20	16
648.41	43	—	43	—	43	—	—	10	*8	11	13
648.43	14	—	14	*	14	—	—	*	*	*6	*
648.6	34	—	34	—	34	—	—	*9	*8	12	*
648.61	28	—	28	—	28	—	—	*7	*6	10	*
648.7	*5	—	*5	—	*5	—	—	*	*	*	*
648.8	121	—	121	—	121	*	—	29	23	42	27
648.81	100	—	100	—	100	*	—	23	18	34	25
648.83	20	—	20	—	20	—	—	*6	*	*8	*
648.9	153	—	153	*	152	*	—	34	38	49	32
648.91	81	—	81	*	81	*	—	16	22	24	18
648.92	*9	—	*9	—	*9	—	—	*	*	*	*
648.93	59	—	59	*	59	*	—	13	14	20	11
650	686	—	686	*	682	*	—	109	134	269	175
651	64	—	64	*	64	—	—	10	13	24	17
651.0	60	—	60	*	59	—	—	9	12	23	16
651.01	42	—	42	—	42	—	—	*6	*8	16	11
651.03	18	—	18	*	18	—	—	*	*	*6	*
652	252	—	252	*	251	*	—	46	60	85	61
652.1	*7	—	*7	—	*7	—	—	*	*	*	*
652.11	*7	—	*7	—	*7	—	—	*	*	*	*
652.2	139	—	139	*	138	*	—	23	36	49	32
652.21	134	—	134	*	133	*	—	22	35	48	29
652.3	21	—	21	*	21	—	—	*	*5	*7	*5
652.31	21	—	21	*	21	—	—	*	*	*7	*5
652.4	*7	—	*7	—	*7	—	—	*	*	*	*
652.41	*7	—	*7	—	*7	—	—	*	*	*	*
652.5	27	—	27	—	27	—	—	*6	*6	12	*
652.51	27	—	27	—	27	—	—	*6	*6	12	*
652.6	10	—	10	—	10	—	—	*	*	*	*
652.61	10	—	10	—	10	—	—	*	*	*	*
652.8	36	—	36	—	36	*	—	*8	*6	9	12
652.81	36	—	36	—	36	*	—	*8	*6	9	12
653	143	—	143	—	143	*	—	20	24	67	33
653.4	114	—	114	—	114	*	—	15	19	53	27
653.41	114	—	114	—	114	*	—	15	19	53	27
653.5	20	—	20	—	20	—	—	*	*	*8	*
653.51	20	—	20	—	20	—	—	*	*	*8	*
654	507	—	507	*	507	*	—	90	113	194	111
654.0	10	—	10	—	10	—	—	*	*	*	*
654.01	9	—	9	—	9	—	—	*	*	*	*
654.1	25	—	25	—	25	*	—	*	*6	12	*
654.11	21	—	21	—	21	*	—	*	*5	10	*
654.2	422	—	422	—	422	*	—	69	97	161	94
654.21	412	—	412	—	412	*	—	66	95	159	92
654.23	9	—	9	—	9	—	—	*	*	*	*
654.4	14	—	14	—	14	—	—	*	*	*	*
654.41	10	—	10	—	10	—	—	*	*	*	*
654.5	22	—	22	*	22	—	—	*7	*	*9	*
654.51	12	—	12	*	12	—	—	*	*	*	*
654.53	9	—	9	—	9	—	—	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
654.6	*6	—	*6	—	*6	—	—	*	*	*	*
654.61	*6	—	*6	—	*6	—	—	*	*	*	*
655	17	—	17	—	17	*	—	*	*	*6	*5
655.8	*7	—	*7	—	*7	—	—	*	*	*	*
655.81	*6	—	*6	—	*6	—	—	*	*	*	*
656	583	—	583	*	581	*	—	104	132	196	150
656.1	63	—	63	*	63	—	—	10	17	22	14
656.11	60	—	60	*	60	—	—	9	15	22	14
656.3	341	—	341	*	340	*	—	63	71	118	89
656.31	337	—	337	*	335	*	—	62	70	117	88
656.4	19	—	19	—	19	—	—	*	*	10	*
656.41	19	—	19	—	19	—	—	*	*	10	*
656.5	58	—	58	*	58	—	—	12	11	19	15
656.51	54	—	54	*	54	—	—	11	11	18	14
656.6	68	—	68	—	68	—	—	11	19	20	18
656.61	66	—	66	—	66	—	—	11	18	19	18
656.7	13	—	13	—	13	—	—	*5	*	*	*
656.71	13	—	13	—	13	—	—	*	*	*	*
656.8	15	—	15	—	15	—	—	*	*7	*	*7
656.81	14	—	14	—	14	—	—	*	*7	*	*6
657	21	—	21	—	21	—	—	*6	*	*6	*6
657.0	21	—	21	—	21	—	—	*6	*	*6	*6
657.01	19	—	19	—	19	—	—	*5	*	*6	*
658	385	—	385	*	384	*	—	69	86	137	93
658.0	82	—	82	—	82	*	—	16	18	27	21
658.01	77	—	77	—	77	*	—	15	17	26	19
658.1	157	—	157	—	157	*	—	26	41	55	35
658.11	147	—	147	—	147	*	—	24	39	50	33
658.13	10	—	10	—	10	—	—	*	*	*	*
658.2	62	—	62	—	62	*	—	15	13	20	14
658.21	59	—	59	—	59	*	—	14	12	18	14
658.4	77	—	77	*	77	*	—	12	12	34	19
658.41	75	—	75	*	75	*	—	11	12	33	18
659	185	—	185	*	184	*	—	28	48	59	50
659.1	45	—	45	*	45	*	—	*6	10	18	12
659.11	37	—	37	*	37	*	—	*6	*8	16	*8
659.13	*8	—	*8	—	*8	—	—	*	*	*	*
659.2	46	—	46	*	46	—	—	*7	10	19	10
659.21	46	—	46	*	46	—	—	*7	10	19	10
659.4	12	—	12	—	12	—	—	*	*	*	*
659.41	12	—	12	—	12	—	—	*	*	*	*
659.5	*6	—	*6	—	*6	—	—	*	*	*	*
659.51	*5	—	*5	—	*5	—	—	*	*	*	*
659.6	32	—	32	—	31	*	—	*	*9	*9	10
659.61	30	—	30	—	30	*	—	*	*8	*8	9
659.8	37	—	37	—	37	*	—	*6	14	*7	10
659.81	33	—	33	—	33	*	—	*6	13	*	9
660	229	—	229	*	229	*	—	34	58	70	68
660.0	36	—	36	—	35	*	—	*6	*8	12	10
660.01	35	—	35	—	35	*	—	*6	*8	12	10
660.1	45	—	45	—	45	—	—	*6	*8	15	16
660.11	45	—	45	—	45	—	—	*6	*8	15	16
660.3	60	—	60	—	60	—	—	10	17	15	18
660.31	60	—	60	—	60	—	—	10	17	15	18
660.4	52	—	52	—	52	—	—	*6	15	17	14
660.41	52	—	52	—	52	—	—	*6	15	17	14
660.6	18	—	18	—	18	—	—	*	*5	*6	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
660.61	18	—	18	—	18	—	—	*	*5	*6	*
660.7	11	—	11	—	11	—	—	*	*	*	*
660.71	11	—	11	—	11	—	—	*	*	*	*
661	347	—	347	*	347	—	—	63	77	107	101
661.0	77	—	77	—	77	—	—	11	15	28	23
661.01	76	—	76	—	76	—	—	11	15	28	23
661.1	60	—	60	—	60	—	—	16	10	17	17
661.11	59	—	59	—	59	—	—	16	10	17	16
661.2	114	—	114	*	114	—	—	21	30	38	24
661.21	112	—	112	*	112	—	—	21	30	37	24
661.3	74	—	74	*	74	—	—	11	16	17	29
661.31	74	—	74	*	73	—	—	11	16	17	29
661.4	*9	—	*9	—	*9	—	—	*	*	*	*
661.41	*8	—	*8	—	*8	—	—	*	*	*	*
661.9	14	—	14	—	14	—	—	*	*	*5	*7
661.91	14	—	14	—	14	—	—	*	*	*5	*6
662	60	—	60	—	60	*	—	14	11	12	22
662.0	11	—	11	—	11	*	—	*	*	*	*5
662.01	11	—	11	—	11	*	—	*	*	*	*5
662.1	12	—	12	—	12	—	—	*	*	*	*
662.11	12	—	12	—	12	—	—	*	*	*	*
662.2	37	—	37	—	37	*	—	9	*8	*8	13
662.21	37	—	37	—	37	*	—	9	*8	*8	13
663	620	—	620	*	619	*	—	122	133	202	162
663.0	11	—	11	—	11	—	—	*	*	*	*
663.01	11	—	11	—	11	—	—	*	*	*	*
663.1	78	—	78	*	78	—	—	15	16	27	20
663.11	78	—	78	*	78	—	—	15	16	27	20
663.2	31	—	31	*	31	—	—	*5	*5	14	*6
663.21	30	—	30	*	30	—	—	*5	*5	14	*6
663.3	471	—	471	*	470	*	—	93	102	150	126
663.31	471	—	471	*	470	*	—	93	102	150	126
663.4	*8	—	*8	*	*8	—	—	*	*	*	*
663.41	*8	—	*8	*	*8	—	—	*	*	*	*
663.8	15	—	15	*	15	—	—	*	*	*5	*
663.81	15	—	15	*	15	—	—	*	*	*5	*
664	967	—	967	*	963	*	—	178	226	288	275
664.0	399	—	399	*	398	*	—	78	85	119	118
664.01	398	—	398	*	397	*	—	78	84	119	117
664.1	345	—	345	*	343	*	—	60	84	101	100
664.11	344	—	344	*	342	*	—	60	84	101	100
664.2	134	—	134	*	133	*	—	26	37	39	32
664.21	132	—	132	*	132	*	—	26	37	38	32
664.3	73	—	73	*	72	—	—	12	16	25	19
664.31	73	—	73	*	72	—	—	12	16	25	19
664.4	*7	—	*7	—	*7	—	—	*	*	*	*
664.41	*7	—	*7	—	*7	—	—	*	*	*	*
665	154	—	154	*	154	—	—	22	43	57	32
665.3	9	—	9	—	9	—	—	*	*	*	*
665.31	*9	—	*9	—	*9	—	—	*	*	*	*
665.4	46	—	46	*	46	—	—	*7	*8	22	*9
665.41	46	—	46	*	45	—	—	*7	*8	22	*9
665.5	86	—	86	*	86	—	—	13	28	28	18
665.51	86	—	86	*	86	—	—	13	28	28	18
666	84	—	84	*	83	—	—	14	16	24	29
666.0	17	—	17	*	16	—	—	*	*	*5	*5
666.02	14	—	14	*	14	—	—	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
666.1 . . . . .	48	—	48	*	47	—	—	*9	*8	10	21
666.12 . . . . .	45	—	45	*	45	—	—	*8	*8	9	20
666.2 . . . . .	16	—	16	*	16	—	—	*	*	*8	*
666.22 . . . . .	*8	—	*8	—	*8	—	—	*	*	*	*
666.24 . . . . .	*8	—	*8	*	*8	—	—	*	*	*6	*
667 . . . . .	28	—	28	—	28	—	—	*	*8	9	*6
667.0 . . . . .	22	—	22	—	22	—	—	*	*7	*7	*
667.02 . . . . .	21	—	21	—	21	—	—	*	*7	*6	*
667.1 . . . . .	*6	—	*6	—	*6	—	—	*	*	*	*
667.12 . . . . .	*6	—	*6	—	*6	—	—	*	*	*	*
669 . . . . .	207	—	207	*	206	*	—	21	61	75	50
669.4 . . . . .	*6	—	*6	—	*6	—	—	*	*	*	*
669.5 . . . . .	122	—	122	*	122	*	—	12	44	46	21
669.51 . . . . .	122	—	122	*	122	*	—	12	44	46	21
669.7 . . . . .	10	—	10	—	10	*	—	*	*	*6	*
669.71 . . . . .	10	—	10	—	10	*	—	*	*	*6	*
669.8 . . . . .	43	—	43	—	43	*	—	*5	*8	11	18
669.81 . . . . .	40	—	40	—	40	*	—	*5	*6	10	18
669.9 . . . . .	16	—	16	—	16	—	—	*	*	*6	*6
669.91 . . . . .	16	—	16	—	16	—	—	*	*	*6	*6
670 . . . . .	35	—	35	*	35	—	—	*5	*5	16	*9
670.0 . . . . .	35	—	35	*	35	—	—	*5	*5	16	*9
670.02 . . . . .	26	—	26	*	26	—	—	*	*	12	*5
670.04 . . . . .	*9	—	*9	—	*9	—	—	*	*	*	*
671 . . . . .	32	—	32	—	32	—	—	*8	*7	11	*6
671.0 . . . . .	*6	—	*6	—	*6	—	—	*	*	*	*
671.8 . . . . .	17	—	17	—	17	—	—	*	*	*7	*
671.81 . . . . .	*9	—	*9	—	*9	—	—	*	*	*5	*
671.82 . . . . .	*8	—	*8	—	*8	—	—	*	*	*	*
672 . . . . .	14	—	14	*	13	—	—	*	*	*5	*
672.0 . . . . .	14	—	14	*	13	—	—	*	*	*5	*
672.02 . . . . .	13	—	13	*	13	—	—	*	*	*	*
674 . . . . .	29	—	29	—	29	*	—	*5	*7	11	*6
674.3 . . . . .	9	—	9	—	*9	*	—	*	*	*	*
674.32 . . . . .	*7	—	*7	—	*7	*	—	*	*	*	*
674.8 . . . . .	16	—	16	—	16	—	—	*	*5	*5	*
674.82 . . . . .	14	—	14	—	14	—	—	*	*	*	*
676 . . . . .	*8	—	*8	*	*8	—	—	*	*	*	*
676.2 . . . . .	*7	—	*7	*	*7	—	—	*	*	*	*
680-709 . . . . .	1,503	702	801	118	317	361	707	373	369	525	235
681 . . . . .	59	29	30	*	14	16	26	13	14	23	9
681.0 . . . . .	21	12	9	*	*8	*	*7	*5	*6	*6	*
681.00 . . . . .	15	10	*	*	*6	*	*	*	*	*	*
681.1 . . . . .	38	18	21	*	*7	11	19	*7	*9	17	*
681.10 . . . . .	33	16	17	*	*6	10	16	*7	*7	16	*
681.11 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
682 . . . . .	613	305	308	37	148	162	266	149	153	213	99
682.0 . . . . .	41	20	21	*7	20	*7	*6	*7	9	15	9
682.1 . . . . .	19	13	*6	*	9	*6	*	*	*	*6	*
682.2 . . . . .	62	25	36	*5	19	15	23	14	13	23	11
682.3 . . . . .	58	29	28	*	19	15	19	14	18	19	*6
682.4 . . . . .	30	17	13	*	11	*7	9	*7	*5	12	*6
682.5 . . . . .	9	*	*	*	*	*	*	*	*	*	*
682.6 . . . . .	269	128	142	9	45	70	145	72	68	91	39
682.7 . . . . .	106	58	48	*	18	32	52	23	30	38	15

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
682.8	*8	*5	*	*	*	*	*	*	*	*	*
682.9	11	*6	*5	*	*	*	*5	*	*	*	*
683	*7	*	*	*	*	*	*	*	*	*	*
684	*7	*	*	*	*	*	*	*	*	*	*
685	*6	*	*	*	*	*	*	*	*	*	*
686	23	10	13	*	*7	*7	*7	*6	*	*9	*
686.9	17	*7	11	*	*5	*	*6	*5	*	*5	*
690	*7	*	*	*	*	*	*	*	*	*	*
691	32	15	17	27	*	*	*	*6	*7	13	*6
691.0	24	11	14	24	*	*	*	*	*6	9	*5
691.8	*8	*	*	*	*	*	*	*	*	*	*
692	48	24	24	9	10	10	19	13	12	16	*7
692.9	37	19	19	*8	*6	*8	15	10	11	12	*
693	40	17	23	*	11	*9	16	11	13	12	*
693.0	39	16	23	*	10	*9	16	11	12	12	*
694	*7	*	*5	*	*	*	*6	*	*	*	*
695	31	10	21	*	*7	9	11	*5	*8	10	*8
695.1	*8	*	*6	*	*	*	*	*	*	*	*
695.3	*6	*	*	—	*	*	*	*	*	*	*
695.4	*6	*	*	—	*	*	*	*	*	*	*
695.8	*6	*	*	*	*	*	*	*	*	*	*
695.89	*6	*	*	*	*	*	*	*	*	*	*
696	36	16	20	*	9	15	11	11	*7	14	*
696.1	31	14	16	*	*8	14	*9	10	*6	11	*
698	19	*6	13	*	*7	*	*7	*	*6	*5	*
698.9	14	*	11	*	*6	*	*5	*	*	*	*
700	*5	*	*	—	*	*	*	*	*	*	*
701	21	*6	16	*	*8	*6	*5	*7	*5	*6	*
701.4	*7	*	*	*	*	*	*	*	*	*	*
702	9	*	*	*	*	*	*5	*	*	*	*
702.1	*6	*	*	*	*	*	*	*	*	*	*
702.19	*6	*	*	*	*	*	*	*	*	*	*
703	10	*5	*5	*	*	*	*8	*	*	*	*
703.0	*6	*	*	*	*	*	*	*	*	*	*
704	*9	*	*	*	*	*	*	*	*	*	*
704.8	*5	*	*	*	*	*	*	*	*	*	*
706	28	14	14	*	15	*	*5	9	*6	9	*
706.1	11	*6	*5	*	*9	*	—	*	*	*	*
706.2	15	*7	*8	*	*6	*	*	*5	*	*	*
707	420	190	231	*	39	90	288	107	101	153	59
707.0	240	97	143	*	22	33	182	59	55	94	31
707.1	169	87	82	*	15	53	100	45	43	54	27
707.8	*8	*	*5	*	*	*	*	*	*	*	*
708	22	*9	13	*	*7	*	*6	*6	*6	*7	*
708.0	14	*6	*7	*	*	*	*	*	*	*	*
708.9	*7	*	*5	*	*	*	*	*	*	*	*
709	33	15	18	*6	13	*7	*7	*6	*8	12	*8
709.2	13	*7	*6	*	*6	*	*	*	*	*	*
709.8	*6	*	*	*	*	*	*	*	*	*	*
709.9	*9	*	*	*	*	*	*	*	*	*	*
710-739	4,374	1,617	2,758	65	797	1,050	2,462	996	1,045	1,554	779
710	121	15	106	*	48	41	31	26	21	50	24
710.0	79	*8	71	*	37	26	15	17	13	33	16

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
710.1	19	*	15	—	*7	*7	*5	*	*	10	*
710.2	*9	*	*8	*	*	*	*6	*	*	*	*
710.4	*7	*	*5	—	*	*	*	*	*	*	*
711	34	19	15	*	9	*6	15	*8	10	*9	*8
711.0	29	17	13	*	*7	*6	14	*7	10	*6	*7
711.06	13	*8	*	*	*	*	*5	*	*	*	*
712	*6	*	*	—	*	*	*5	*	*	*	*
713	*7	*	*	—	*	*6	*	*	*	*	*
714	221	54	167	*	17	63	136	46	58	74	43
714.0	203	47	157	*	14	58	131	40	56	70	37
714.3	*8	*	*5	*	*	*	*	*	*	*	*
714.30	*7	*	*5	*	*	*	*	*	*	*	*
714.8	*7	*	*	—	*	*	*	*	*	*	*
714.81	*5	*	*	—	*	*	*	*	*	*	*
715	1,075	339	736	*	34	190	850	253	280	370	171
715.0	30	*8	22	—	*	*	26	*9	11	*8	*
715.09	26	*7	18	—	*	*	22	*8	*9	*7	*
715.1	40	17	23	—	*	*8	30	*	10	16	11
715.15	14	*	*9	—	*	*	9	*	*	*	*5
715.16	22	12	11	—	*	*	18	*	*6	10	*
715.3	352	119	233	—	15	79	258	81	74	127	70
715.31	14	*5	9	—	*	*	11	*6	*	*	*
715.35	105	43	62	—	*6	21	78	22	18	41	25
715.36	208	67	141	—	*7	49	153	46	48	74	40
715.37	*6	*	*	—	*	*	*	*	*	*	*
715.38	10	*	*9	—	*	*	*8	*	*	*5	*
715.8	57	16	41	*	*	*8	48	17	16	15	9
715.89	55	15	40	*	*	*8	46	16	16	14	9
715.9	591	176	415	*	12	91	487	141	170	204	77
715.90	450	126	324	*	*7	58	384	101	107	185	57
715.91	9	*	*5	*	*	*	*6	*	*5	*	*
715.95	38	13	25	—	*	9	27	14	15	*	*5
715.96	72	25	47	*	*	15	55	18	37	*9	*9
715.98	16	*6	10	—	*	*	12	*	*	*	*
716	188	69	119	*	20	50	117	35	52	73	27
716.1	9	*6	*	—	*	*	*	*	*	*	*
716.16	*5	*	*	—	*	*	*	*	*	*	*
716.5	*6	*	*	*	*	*	*	*	*	*	*
716.6	*6	*	*	*	*	*	*	*	*	*	*
716.9	161	55	106	*	13	43	105	29	48	63	21
716.90	111	38	73	*	*7	28	76	17	31	47	16
716.91	*9	*	*5	—	*	*	*	*	*	*	*
716.96	16	*5	11	*	*	*	9	*	*5	*5	*
716.98	*6	*	*	—	*	*	*	*	*	*	*
716.99	*8	*	*6	*	*	*	*	*	*	*	*
717	84	49	34	*	58	19	*6	39	18	13	14
717.2	*5	*	*	*	*	*	*	*	*	*	*
717.3	9	*6	*	*	*6	*	*	*5	*	*	*
717.4	10	*7	*	—	*6	*	*	*5	*	*	*
717.7	19	10	9	*	12	*6	*	12	*	*	*
717.8	29	18	11	*	25	*	*	9	*8	*	*7
717.83	27	16	10	*	23	*	*	9	*8	*	*7
718	67	31	36	*	31	17	17	16	14	23	15
718.3	15	*8	*7	*	11	*	*	*	*	*6	*
718.31	9	*7	*	—	*8	*	*	*	*	*	*
718.4	19	*6	13	*	*	*6	10	*5	*	*7	*
718.46	*5	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
718.49	*5	*	*	*	*	*	*	*	*	*	*
718.5	*6	*	*	*	*	*	*	*	*	*	*
718.8	13	*8	*5		9	*	*	*	*	*	*
719	137	53	84	*6	31	24	77	29	30	48	30
719.0	18	*8	10	*	*	*	10	*	*	*	*
719.06	15	*7	*8	*	*	*	*9	*	*	*	*
719.4	69	25	45	*	22	15	30	*9	15	29	17
719.41	23	10	13	*	*6	*5	11	*	*	*8	*7
719.45	20	*7	13	*	*6	*	12	*	*	10	*
719.46	12	*	*8	*	*	*	*	*	*	*	*
719.7	32	9	22	*	*	*	28	11	*8	*8	*
719.70	*9	*	*	*	*	*	*7	*5	*	*	*
719.75	*8	*	*7	—	*	*	*7	*	*	*	*
719.76	9	*	*7	—	*	*	*9	*	*	*	*
720	22	15	*7	—	*	12	*7	10	*	*8	*
720.0	14	12	*	—	*	10	*	*9	*	*	*
721	253	105	148	—	23	60	169	51	70	93	38
721.0	67	34	33	—	9	24	34	12	17	29	10
721.1	21	13	*8	—	*	10	*9	*7	*	*8	*
721.2	11	*	*7	—	*	*	10	*	*	*	*
721.3	92	32	61	—	*6	18	68	17	29	28	19
721.4	*5	*	*	—	*	*	*	*	*	*	*
721.8	*6	*5	*	—	*	*	*	*	*	*	*
721.9	49	13	36	—	*	*5	41	10	14	21	*
721.90	49	13	36	—	*	*5	41	10	14	21	*
722	448	244	205	*	183	158	107	89	98	182	79
722.0	68	40	28	—	32	28	*8	12	13	35	*8
722.1	229	129	100	—	114	78	37	46	48	96	40
722.10	226	129	98	—	113	77	36	46	47	95	39
722.4	17	9	*7	—	*	*7	*8	*	*	*6	*
722.5	63	29	34	*	14	17	31	13	20	17	14
722.52	60	27	33	*	14	17	29	12	19	16	13
722.6	*7	*	*	—	*	*	*	*	*	*	*
722.7	22	15	*7	*	*7	*8	*6	*7	*	*5	*5
722.71	12	*9	*	*	*	*5	*	*5	*	*	*
722.73	*9	*	*	—	*	*	*	*	*	*	*
722.8	16	*6	9	—	*	*8	*	*	*	*8	*
722.83	15	*6	*9	—	*	*7	*	*	*	*8	*
722.9	22	10	12	*	*6	*8	*7	*	*5	10	*
722.91	*6	*	*	*	*	*	*	*	*	*	*
722.93	14	*7	*7	—	*5	*	*	*	*	*6	*
723	45	24	22	*	15	17	11	14	*9	15	*7
723.0	*8	*	*	*	*	*	*	*	*	*	*
723.1	16	*7	9	*	*9	*	*	*	*	*6	*
723.4	16	9	*7	*	*5	*8	*	*6	*	*5	*
724	294	121	172	*	80	86	126	62	73	106	52
724.0	87	36	51	—	*7	20	60	22	18	30	17
724.00	14	*7	*7	—	*	*	12	*	*	*	*
724.02	72	28	44	—	*6	18	48	18	16	25	13
724.2	87	37	51	*	32	27	28	15	24	35	13
724.3	13	*	*9	*	*	*	*8	*	*	*	*
724.4	37	16	21	—	13	12	12	*8	10	13	*5
724.5	49	20	29	*	17	17	13	10	13	16	10
724.6	*6	*	*	—	*	*	*	*	*	*	*
724.8	10	*	*6	*	*	*	*	*	*	*	*
725	43	12	31	—	—	*	41	*7	11	16	9

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
726 . . . . .	117	58	59	*	27	51	38	32	28	36	21
726.0 . . . . .	*9	*	*5	—	*	*	*	*	*	*	*
726.1 . . . . .	34	19	15	—	*7	15	12	10	*7	11	*6
726.10 . . . . .	27	15	12	—	*	13	9	*9	*5	*8	*5
726.2 . . . . .	36	18	18	—	*7	20	*9	*8	*9	11	*7
726.3 . . . . .	9	*6	*	—	*	*	*	*	*	*	*
726.33 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
726.5 . . . . .	*8	*	*5	*	*	*	*5	*	*	*	*
726.6 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
726.9 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
726.91 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
727 . . . . .	67	28	39	*	20	23	21	24	12	21	10
727.0 . . . . .	30	13	18	*	11	10	*8	*9	*	12	*
727.00 . . . . .	14	*6	*8	*	*	*	*	*	*	*6	*
727.05 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
727.1 . . . . .	*8	*	*7	*	*	*	*	*	*	*	*
727.4 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
727.6 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
727.8 . . . . .	*9	*5	*	*	*	*	*	*	*	*	*
728 . . . . .	81	41	40	*	20	18	39	21	20	25	14
728.2 . . . . .	10	*	*7	*	*	*	*	*	*	*	*
728.3 . . . . .	10	*	*	*	*	*	*6	*	*	*	*
728.8 . . . . .	42	23	19	*	13	10	17	13	*8	11	10
728.85 . . . . .	9	*6	*	*	*	*	*	*	*	*	*
728.89 . . . . .	26	14	12	*	*7	*	13	*9	*	*7	*6
728.9 . . . . .	12	*6	*6	*	*	*	*9	*	*	*6	*
729 . . . . .	144	53	92	*	39	47	56	30	34	49	31
729.1 . . . . .	55	16	40	*	19	21	14	12	14	18	11
729.2 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
729.5 . . . . .	40	15	25	*	12	12	15	*8	*8	14	10
729.8 . . . . .	36	18	19	*	*	11	20	*7	9	14	*7
729.82 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
729.89 . . . . .	27	13	14	*	*	*8	17	*5	*7	10	*6
730 . . . . .	124	67	57	*6	23	36	59	31	24	50	19
730.0 . . . . .	36	19	17	*	*7	*8	18	*8	*7	16	*6
730.06 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
730.07 . . . . .	21	12	10	*	*	*6	11	*5	*	10	*
730.08 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
730.1 . . . . .	27	15	12	*	*	10	13	*6	*6	*9	*7
730.15 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
730.16 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
730.17 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
730.2 . . . . .	50	26	24	*	11	14	23	15	9	21	*5
730.25 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
730.26 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
730.27 . . . . .	23	14	9	*	*	9	11	*8	*	*8	*
730.28 . . . . .	*9	*	*5	*	*	*	*	*	*	*	*
730.8 . . . . .	10	*7	*	*	*	*	*	*	*	*	*
730.87 . . . . .	*8	*5	*	—	*	*	*	*	*	*	*
731 . . . . .	22	12	10	—	*	*	17	*7	*6	*8	*
731.0 . . . . .	18	*8	10	—	*	*	17	*6	*5	*6	*
732 . . . . .	*8	*	*	*5	*	*	*	*	*	*	*
733 . . . . .	595	136	459	*7	65	84	439	114	127	222	132
733.0 . . . . .	256	26	230	*	*6	19	231	49	57	94	56
733.00 . . . . .	225	22	202	*	*	16	205	43	53	86	43
733.01 . . . . .	16	*	15	—	*	*	14	*	*	*	*5

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
733.09	15	*	12	*	*	*	12	*	*	*	*7
733.1	175	41	134	*	*6	24	145	34	39	65	38
733.11	*7	*	*6	—	*	*	*	*	*	*	*
733.13	117	28	89	*	*	11	102	22	26	42	26
733.14	20	*	15	*	*	*	17	*	*	*6	*
733.15	*8	*	*7	*	*	*	*	*	*	*	*
733.19	16	*	12	—	*	*	12	*	*	*5	*
733.2	10	*	*6	*	*	*	*	*	*	*	*6
733.20	*7	*	*	*	*	*	*	*	*	*	*5
733.4	34	16	18	*	12	*8	14	*7	*	13	10
733.42	25	13	12	*	*7	*6	11	*6	*	9	*6
733.49	*6	*	*	*	*	*	*	*	*	*	*
733.6	37	10	27	*	13	11	12	*6	*8	17	*5
733.8	44	24	20	*	14	12	16	*8	*7	19	10
733.81	*6	*	*	*	*	*	*	*	*	*	*
733.82	37	20	17	*	11	11	14	*7	*	17	*8
733.9	37	14	23	*	10	*8	18	*8	10	11	*8
733.90	22	*7	15	*	*	*	14	*	*	*7	*6
733.92	*6	*	*	—	*	*	*	*	*	*	*
733.99	*9	*	*5	*	*	*	*	*	*	*	*
735	17	*	13	*	*5	*6	*6	*9	*	*	*
735.0	*7	*	*6	—	*	*	*	*	*	*	*
735.4	*7	*	*	—	*	*	*	*	*	*	*
736	28	14	15	*	*6	*5	14	*7	*7	*7	*7
736.4	*7	*	*	*	*	*	*	*	*	*	*
736.7	10	*	*6	*	*	*	*	*	*	*	*
736.79	*8	*	*	*	*	*	*	*	*	*	*
736.8	*6	*	*	*	*	*	*	*	*	*	*
737	70	18	52	*8	15	9	38	16	14	30	10
737.1	16	*8	*8	*	*	*	12	*	*	*9	*
737.10	13	*6	*7	*	*	*	11	*	*	*7	*
737.3	48	*9	39	*7	13	*7	21	12	11	20	*6
737.30	40	*7	33	*	10	*6	18	9	*8	17	*5
737.39	*7	*	*5	*	*	*	*	*	*	*	*
737.4	*5	*	*	*	*	*	*	*	*	*	*
738	42	22	20	*	16	12	14	*8	11	12	10
738.4	29	16	13	—	*7	10	11	*	*9	*8	*8
739	12	*	9	*	*	*	*5	*	*5	*	—
740-759	577	289	288	340	115	63	60	210	123	144	100
741	18	*8	10	*9	*8	*	*	*6	*6	*	*
741.0	*8	*	*	*6	*	*	*	*	*	*	*
741.9	10	*5	*	*	*6	*	*	*	*	*	*
741.90	*6	*	*	*	*	*	*	*	*	*	*
742	31	16	16	21	*	*	*	12	*8	*8	*
742.1	*5	*	*	*	*	*	—	*	*	*	*
742.3	10	*5	*	*7	*	*	*	*	*	*	*
742.5	*7	*	*	*	*	—	*	*	*	*	*
742.59	*7	*	*	*	*	—	*	*	*	*	*
743	*6	*	*	*	*	—	*	*	*	*	*
744	*6	*	*	*	*	*	*	*	*	*	*
745	78	36	41	64	*7	*	*	32	14	16	16
745.1	*8	*5	*	*7	*	—	—	*	*	*	*
745.2	*9	*	*	*7	*	*	*	*	*	*	*
745.4	26	12	14	23	*	*	*	11	*6	*	*
745.5	26	11	15	19	*	*	*	11	*	*6	*5

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
746 . . . . .	52	23	28	35	*9	*	*	21	*9	12	10
746.0 . . . . .	10	*	*7	*9	*	—	—	*	*	*	*
746.02 . . . . .	*8	*	*5	*7	*	—	—	*	*	*	*
746.8 . . . . .	18	10	*8	11	*	*	*	*8	*	*	*
746.89 . . . . .	9	*5	*	*6	*	*	*	*	*	*	*
746.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
747 . . . . .	63	33	31	42	11	*	*8	19	14	19	11
747.0 . . . . .	22	11	11	21	*	*	*	*6	*	*6	*7
747.1 . . . . .	*7	*	*	*6	*	—	*	*	*	*	*
747.10 . . . . .	*7	*	*	*6	*	—	*	*	*	*	*
747.2 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
747.3 . . . . .	12	*	*7	*7	*	*	—	*6	*	*	*
747.6 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
747.8 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
747.81 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
748 . . . . .	16	10	*6	14	*	*	*	*7	*	*	*
748.3 . . . . .	*7	*	*	*7	*	—	*	*	*	*	*
749 . . . . .	14	*8	*5	13	*	—	*	*5	*	*	*
749.0 . . . . .	*6	*	*	*5	*	—	*	*	*	*	*
749.2 . . . . .	*5	*	*	*5	*	—	—	*	*	*	*
750 . . . . .	17	11	*6	11	*	*	*	*	*	*	*
750.5 . . . . .	*8	*6	*	*8	—	—	—	*	*	*	*
751 . . . . .	33	19	14	15	*6	*6	*6	10	*9	*8	*7
751.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
751.5 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
752 . . . . .	38	19	19	11	16	*9	*	13	*8	13	*
752.1 . . . . .	9	—	9	*	*5	*	*	*	*	*	*
752.11 . . . . .	*8	—	*8	*	*	*	*	*	*	*	*
752.5 . . . . .	*8	*8	—	*	*	*	*	*	*	*	*
752.6 . . . . .	*8	*8	—	*	*	*	*	*	*	*	*
753 . . . . .	54	26	28	19	10	*9	17	18	11	17	*8
753.1 . . . . .	30	15	15	*	*7	*7	14	*8	*5	12	*
753.10 . . . . .	*8	*	*	*	*	*	*6	*	*	*	*
753.12 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
753.13 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
753.2 . . . . .	11	*	*6	9	*	*	*	*	*	*	*
754 . . . . .	23	11	12	16	*	*	*	*8	*5	*6	*
754.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
754.7 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
755 . . . . .	14	*5	9	*8	*	*	*	*8	*	*	*
755.6 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
756 . . . . .	41	22	19	20	9	*6	*6	13	*9	12	*7
756.0 . . . . .	10	*6	*	9	*	*	*	*	*	*	*
756.1 . . . . .	18	9	9	*	*6	*	*5	*6	*	*5	*
756.12 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
756.5 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
757 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
758 . . . . .	31	18	13	17	*6	*7	*	11	10	*5	*5
758.0 . . . . .	24	16	*9	12	*5	*6	*	*8	*8	*	*
759 . . . . .	34	13	21	14	14	*	*	15	*6	*8	*
759.8 . . . . .	19	*8	11	*9	*8	*	*	10	*	*	*
759.89 . . . . .	16	*6	9	*8	*7	*	*	9	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
760-779 . . . . .	389	239	150	382	*	*	*	76	112	109	93
765 . . . . .	46	27	19	45	*	-	-	10	12	12	12
765.0 . . . . .	*6	*	*	*6	-	-	-	*	*	*	*
765.1 . . . . .	39	24	16	39	*	-	-	*9	11	10	10
765.16 . . . . .	*5	*	*	*5	-	-	-	*	*	*	*
765.18 . . . . .	*8	*6	*	*8	-	-	-	*	*	*	*
765.19 . . . . .	*9	*5	*	*9	*	-	-	*	*	*	*
767 . . . . .	*6	*	*	*6	-	-	-	*	*	*	*
769 . . . . .	24	14	9	23	*	*	*	*	*5	*6	*9
770 . . . . .	88	55	32	87	*	*	-	19	26	24	18
770.2 . . . . .	*7	*	*	*7	*	*	-	*	*	*	*
770.6 . . . . .	11	9	*	11	-	-	-	*	*	*	*
770.7 . . . . .	17	11	*6	17	-	*	-	*	*6	*	*
770.8 . . . . .	42	24	18	42	*	*	-	10	11	12	10
771 . . . . .	37	23	14	37	*	*	-	*7	11	11	*8
771.8 . . . . .	30	20	11	30	-	*	-	*6	*9	9	*7
772 . . . . .	*5	*	*	*	-	*	-	*	*	*	*
773 . . . . .	*6	*	*	*6	*	-	*	*	*	*	*
774 . . . . .	71	45	27	70	*	*	-	12	22	18	18
774.2 . . . . .	23	15	*8	23	*	-	-	*	*8	*7	*6
774.6 . . . . .	45	27	17	44	*	*	-	9	13	10	12
775 . . . . .	25	17	*8	25	*	-	-	*	*5	12	*
775.5 . . . . .	14	10	*	14	-	-	-	*	*	*8	*
776 . . . . .	11	*7	*	11	*	-	*	*	*	*	*
776.6 . . . . .	*6	*	*	*6	-	-	-	*	*	*	*
777 . . . . .	*7	*	*	*7	*	-	-	*	*	*	*
778 . . . . .	13	*8	*6	13	*	*	-	*	*6	*	*
778.4 . . . . .	9	*5	*	*9	*	-	-	*	*	*	*
779 . . . . .	30	17	13	29	*	*	*	*	*8	12	*7
779.0 . . . . .	*6	*	*	*6	-	-	-	*	*	*	*
779.3 . . . . .	19	12	*7	18	*	*	*	*	*	*9	*
780-799 . . . . .	5,673	2,578	3,095	492	1,317	1,373	2,490	1,292	1,411	2,019	951
780 . . . . .	1,658	789	869	186	358	361	753	411	410	572	266
780.0 . . . . .	64	32	32	*	12	10	39	15	15	16	19
780.01 . . . . .	21	11	10	*	*	*	15	*5	*	*5	*6
780.09 . . . . .	38	19	19	*	*8	*6	22	*8	9	9	10
780.1 . . . . .	11	*6	*6	*	*	*	*6	*	*	*6	*
780.2 . . . . .	312	132	180	*	36	64	207	83	74	106	48
780.3 . . . . .	656	332	324	84	174	151	248	173	174	222	87
780.4 . . . . .	91	32	60	*	11	20	59	19	27	29	16
780.5 . . . . .	89	58	32	10	24	33	23	23	20	30	16
780.52 . . . . .	12	*	*8	*	*	*	*7	*	*	*6	*
780.57 . . . . .	68	47	20	*9	20	26	13	18	15	23	11
780.6 . . . . .	275	134	141	79	71	51	73	69	61	100	44
780.7 . . . . .	109	39	70	*	14	19	75	15	27	41	26
780.9 . . . . .	46	21	26	*	13	9	23	9	10	20	*7
781 . . . . .	122	57	66	*8	15	20	80	29	38	39	16
781.0 . . . . .	15	*7	*8	*	*	*	*6	*	*	*	*
781.2 . . . . .	58	22	36	*	*	*7	47	12	21	19	*7
781.3 . . . . .	39	22	17	*	*	*8	24	*9	11	14	*
781.9 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
782 . . . . .	133	61	72	17	29	27	60	36	35	38	25
782.0 . . . . .	23	11	12	*	*8	*7	*7	*5	*6	*6	*6
782.1 . . . . .	28	12	16	*6	*9	*	*8	*6	9	9	*

\*Figure does not meet standard of reliability or precision.  
-Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
782.3	48	19	29	*	*6	10	31	14	13	14	*8
782.4	21	13	*8	*	*	*	9	*	*6	*6	*
782.7	*6	*	*	*	*	*	*	*	*	*	*
783	127	59	67	51	20	10	46	31	30	39	25
783.0	26	10	16	*	*6	*	15	*7	*7	*9	*
783.2	29	12	17	*	*7	*	17	*	*8	12	*
783.3	11	*5	*6	*5	—	*	*5	*	*	*	*
783.4	55	30	25	41	*5	*	*8	17	11	13	15
784	321	133	188	14	81	73	154	59	80	116	66
784.0	116	38	78	*8	53	30	25	17	30	49	18
784.2	*9	*	*5	*	*	*	*	*	*	*	*
784.3	83	41	42	*	*	14	63	19	23	23	18
784.5	54	22	32	*	*	11	38	10	14	15	15
784.7	50	25	25	*	16	13	19	9	10	21	10
785	277	132	145	20	53	64	140	71	68	90	47
785.0	43	19	24	*	11	*9	18	*6	14	16	*7
785.1	22	*7	15	*	*7	*6	*9	*7	*5	*8	*
785.2	19	*7	11	*7	*6	*	*	*	*	*7	*
785.4	86	45	41	*	*9	23	55	25	19	23	19
785.5	67	34	33	*	*	13	47	18	17	20	12
785.50	*9	*	*	*	*	*	*6	*	*	*5	*
785.51	30	15	15	*	*	*7	23	*9	*8	*8	*6
785.59	28	14	14	*	*	*5	18	*9	*7	*7	*5
785.6	34	18	15	*6	16	*8	*	10	*6	14	*
785.9	*6	*	*	—	*	*	*	*	*	*	*
786	996	472	525	63	201	371	360	219	227	396	155
786.0	150	74	76	54	15	24	57	28	37	61	23
786.01	*5	*	*	*	*	*	*	*	*	*	*
786.09	142	71	70	53	13	22	54	27	35	59	20
786.1	*7	*	*	*	*	*	*	*	*	*	*
786.2	13	*5	*7	*	*	*	*	*	*	*6	*
786.3	41	24	18	*	*8	15	17	11	11	15	*
786.5	751	346	405	*	173	319	257	168	166	298	119
786.50	610	288	321	*	140	263	206	135	139	244	91
786.51	19	*7	12	—	*	9	*6	*6	*	9	*
786.52	86	36	50	*	21	33	31	18	20	32	16
786.59	36	15	21	*	*8	14	14	9	*6	12	*9
786.6	28	15	13	*	*	*6	20	*7	*8	10	*
786.8	*6	*	*	—	*	*	*	*	*	*	*
787	403	156	248	50	91	81	182	73	96	146	89
787.0	272	95	177	43	75	60	94	50	63	100	59
787.01	167	54	114	14	53	40	61	30	37	65	35
787.02	51	15	36	*	12	13	23	9	12	18	12
787.03	54	27	27	27	10	*8	10	11	14	17	11
787.1	*5	*	*	—	*	*	*	*	*	*	*
787.2	95	46	49	*	*9	13	70	17	27	30	22
787.3	*7	*	*	*	*	*	*	*	*	*	*
787.6	11	*	*7	*	*	*	*	*	*	*	*
787.9	13	*	10	*	*	*	*7	*	*	*9	*
788	399	235	164	*6	79	79	234	99	100	137	63
788.0	32	20	12	*	18	9	*5	9	*6	13	*
788.1	*6	*	*	*	*	*	*	*	*	*	*
788.2	245	162	83	*	45	45	153	63	58	85	39
788.20	203	132	71	*	37	39	126	52	46	76	29
788.21	*8	*	*	—	*	*	*6	*	*	*	*
788.29	34	25	*9	*	*7	*5	22	9	*8	*8	*8
788.3	78	30	48	*	10	14	52	16	23	25	14

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
788.30	54	20	34	*	*5	*9	39	12	15	17	10
788.33	*7	*	*6	—	*	*	*	*	*	*	*
788.39	*7	*	*	*	*	*	*	*	*	*	*
788.4	*9	*5	*	*	*	*	*5	*	*	*	*
788.41	*5	*	*	*	*	*	*	*	*	*	*
788.9	20	*9	11	*	*	*	14	*	*7	*5	*
789	460	162	298	29	165	121	145	92	116	173	79
789.0	336	107	229	24	139	76	96	63	88	129	57
789.00	153	50	104	11	61	32	50	26	36	67	24
789.01	31	10	21	*	13	*9	*9	*6	*8	13	*
789.02	*9	*	*5	*	*	*	*	*	*	*	*
789.03	51	15	36	*7	30	9	*6	13	12	17	10
789.04	22	*	17	*	9	*6	*6	*	*6	*8	*
789.06	25	11	15	*	*8	*7	*9	*5	*8	*7	*
789.07	13	*	*9	*	*	*	*	*	*	*	*
789.09	29	*8	21	*	10	*8	*9	*	9	*8	*6
789.1	13	*6	*7	*	*	*	*5	*	*	*5	*
789.2	12	*6	*6	*	*	*	*	*	*	*	*
789.3	13	*	12	*	*	*	*	*	*	*	*
789.30	*5	*	*	*	*	*	*	*	*	*	*
789.5	84	42	42	*	16	32	35	21	19	32	12
790	291	122	169	17	85	58	132	68	82	93	48
790.1	*7	*	*6	—	*	*	*5	*	*	*	*
790.2	62	*8	54	—	47	*	10	11	15	21	15
790.4	10	*	*5	*	*	*	*	*	*	*	*
790.5	24	15	*9	*	*7	*	12	*7	*	10	*
790.6	128	58	70	*5	15	35	73	28	41	41	18
790.7	29	14	15	*8	*7	*	10	*8	*7	*9	*5
790.9	24	17	*7	*	*	*	15	*6	*8	*7	*
790.92	14	*8	*6	*	*	*	*8	*	*	*	*
790.93	*6	*6	—	—	—	*	*6	*	*	*	*
791	22	*9	13	*	*8	*	*7	*6	*6	*8	*
791.0	12	*6	*6	*	*	*	*	*	*	*	*
792	62	*8	54	*	42	*6	12	12	16	19	15
792.1	22	*8	13	*	*	*6	12	*7	*5	*5	*
792.3	40	—	40	*	40	—	—	*5	10	14	11
793	13	*5	*8	*	*	*	*6	*	*	*6	*
793.1	*9	*	*	*	*	*	*	*	*	*5	*
794	83	40	43	*	21	27	32	18	23	30	12
794.3	33	14	19	*	*8	11	13	*6	11	12	*
794.31	24	10	14	*	*7	*8	*8	*	*8	*8	*
794.39	*6	*	*	*	*	*	*	*	*	*	*
794.8	44	22	22	*	12	14	18	11	*9	17	*7
795	31	13	18	*	15	*8	*7	11	*6	9	*5
795.5	23	10	13	*	11	*7	*	9	*	*7	*
796	28	14	14	*	*7	*8	11	*6	*9	*9	*
796.2	16	*7	*8	*	*	*5	*5	*	*5	*	*
796.3	*6	*	*	*	*	*	*	*	*	*	*
797	9	*	*7	—	—	—	9	*	*	*	*
799	235	109	126	19	45	52	119	46	63	95	31
799.0	99	44	55	10	14	22	54	15	30	36	18
799.1	14	*8	*7	*	*	*	*8	*	*	*	*
799.3	15	*5	10	—	*	*	13	*	*8	*	*
799.4	34	19	15	*	*7	*7	20	9	*9	12	*
799.9	68	31	37	*8	21	18	22	17	11	37	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
800-999 . . . . .	5,965	3,081	2,884	454	2,169	1,174	2,168	1,325	1,384	2,013	1,244
800 . . . . .	17	12	*	10	*	*	*	*	*	*	*
800.0 . . . . .	*8	*6	*	*6	*	*	*	*	*	*	*
801 . . . . .	39	30	10	*8	21	*6	*	10	*9	11	10
801.0 . . . . .	20	15	*	*6	11	*	*	*6	*	*	*6
801.01 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
801.2 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
802 . . . . .	115	85	30	*8	82	14	11	20	26	37	33
802.0 . . . . .	21	14	*7	*	13	*	*	*6	*	*7	*
802.2 . . . . .	24	18	*6	*	20	*	*	*	*	*7	*8
802.3 . . . . .	*6	*6	*	*	*	*	—	*	*	*	*
802.4 . . . . .	26	17	*9	*	18	*	*	*	*6	9	*7
802.6 . . . . .	14	12	*	*	11	*	*	*	*	*	*5
802.8 . . . . .	17	13	*	*	11	*	*	*	*	*	*
803 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
805 . . . . .	118	56	62	*	38	21	56	21	38	37	23
805.0 . . . . .	21	14	*8	*	12	*5	*	*	*6	*7	*5
805.02 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
805.2 . . . . .	33	13	19	*	*8	*7	17	*8	11	*8	*6
805.4 . . . . .	50	24	27	*	15	*5	28	*8	17	18	*8
805.6 . . . . .	11	*	*7	*	*	*	*6	*	*	*	*
806 . . . . .	*8	*6	*	*	*6	*	*	*	*	*	*
807 . . . . .	135	74	60	*	46	28	58	29	32	43	30
807.0 . . . . .	117	66	51	*	41	23	51	23	28	40	26
807.00 . . . . .	13	*7	*6	*	*	*	*5	*	*	*6	*
807.01 . . . . .	28	15	14	*	10	*	13	*8	*8	*6	*6
807.02 . . . . .	17	10	*7	*	*5	*	*9	*	*	*7	*
807.03 . . . . .	13	*8	*	*	*	*	*6	*	*	*	*
807.04 . . . . .	13	*8	*	*	*	*	*	*	*	*	*
807.05 . . . . .	9	*	*5	*	*	*	*	*	*	*	*
807.09 . . . . .	18	10	*8	*	*7	*	*7	*	*	*7	*
807.2 . . . . .	12	*	*8	*	*	*	*6	*	*	*	*
808 . . . . .	98	42	56	*	27	13	56	16	28	32	23
808.0 . . . . .	17	12	*	*	*9	*	*	*	*	*7	*
808.2 . . . . .	53	17	36	*	11	*8	34	10	14	15	15
808.4 . . . . .	15	*8	*7	*	*	*	*8	*	*6	*	*
808.42 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
808.8 . . . . .	11	*	*8	*	*	*	*8	*	*	*5	*
810 . . . . .	26	16	10	*	13	*	*7	*5	*	*8	*8
810.0 . . . . .	25	15	10	*	13	*	*7	*5	*	*8	*7
810.00 . . . . .	13	*8	*5	*	*7	*	*	*	*	*5	*
810.02 . . . . .	*9	*6	*	*	*5	*	*	*	*	*	*5
811 . . . . .	12	*7	*5	*	*7	*	*	*	*	*	*5
811.0 . . . . .	12	*7	*5	*	*7	*	*	*	*	*	*5
812 . . . . .	105	36	70	13	21	17	55	23	21	34	27
812.0 . . . . .	53	15	38	*	*	*9	39	13	11	17	12
812.00 . . . . .	16	*	14	*	*	*	12	*	*5	*	*
812.01 . . . . .	21	*6	15	*	*	*	15	*6	*	*8	*
812.09 . . . . .	11	*5	*6	*	*	*	10	*	*	*	*
812.2 . . . . .	17	*6	11	*	*6	*	*6	*	*	*5	*7
812.20 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
812.21 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
812.4 . . . . .	29	9	20	11	*7	*	*9	*7	*6	10	*6
812.40 . . . . .	*9	*	*7	*	*	*	*	*	*	*	*
812.41 . . . . .	13	*	*8	*8	*	*	*	*	*	*5	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
813 . . . . .	94	45	49	12	35	15	31	23	24	31	15
813.0 . . . . .	23	10	13	*	9	*	*9	*8	*6	*6	*
813.01 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
813.05 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
813.2 . . . . .	*9	*6	*	*	*5	*	*	*	*	*	*
813.23 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
813.4 . . . . .	43	18	24	*6	12	*7	18	11	11	14	*7
813.41 . . . . .	10	*	*7	*	*	*	*6	*	*	*	*
813.42 . . . . .	22	11	11	*	*7	*	*8	*6	*6	*7	*
813.44 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
813.8 . . . . .	*8	*	*	*	*	*	*	*	*	*5	*
814 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
814.0 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
815 . . . . .	16	10	*5	*	10	*	*	*	*	*6	*
815.0 . . . . .	12	*6	*5	*	*8	*	*	*	*	*	*
815.00 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
816 . . . . .	23	16	*7	*	*9	*6	*7	*	*	*9	*
816.0 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*
816.01 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
816.1 . . . . .	11	10	*	*	*	*	*	*	*	*	*
820 . . . . .	337	91	247	*	9	22	304	70	86	117	64
820.0 . . . . .	104	27	77	*	*	*	96	25	28	31	20
820.00 . . . . .	*5	*	*	—	*	*	*5	*	*	*	*
820.02 . . . . .	*6	*	*	—	*	*	*5	*	*	*	*
820.03 . . . . .	*8	*	*7	—	*	*	*7	*	*	*	*
820.09 . . . . .	84	21	63	*	*	*	79	20	25	26	14
820.2 . . . . .	153	38	115	*	*	13	136	34	40	53	26
820.20 . . . . .	*9	*	*6	*	*	*	*7	*	*	*	*
820.21 . . . . .	132	32	100	*	*	11	119	29	33	46	24
820.22 . . . . .	13	*	9	*	*	*	10	*	*	*	*
820.8 . . . . .	78	25	53	*	*	*	71	11	17	33	17
821 . . . . .	78	38	40	14	26	*8	30	19	17	29	13
821.0 . . . . .	39	25	15	11	15	*	10	10	*8	14	*7
821.00 . . . . .	9	*5	*	*	*	*	*	*	*	*	*
821.01 . . . . .	30	20	10	9	13	*	*5	*8	*7	9	*6
821.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
821.11 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
821.2 . . . . .	30	*8	22	*	*	*	20	*7	*7	13	*
821.20 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
821.21 . . . . .	9	*	*8	—	*	*	*7	*	*	*	*
821.23 . . . . .	12	*	9	*	*	*	*8	*	*	*5	*
822 . . . . .	24	11	13	*	9	*	10	*	*	*7	*9
822.0 . . . . .	20	9	11	*	*7	*	10	*	*	*5	*8
823 . . . . .	104	63	42	11	46	23	24	24	19	41	21
823.0 . . . . .	43	19	24	*	15	11	16	10	*9	16	*8
823.00 . . . . .	29	13	16	*	10	*7	12	*7	*6	10	*6
823.01 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
823.02 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
823.1 . . . . .	*6	*6	*	*	*	*	*	*	*	*	*
823.2 . . . . .	17	11	*6	*	*9	*	*	*	*	*7	*
823.22 . . . . .	11	*7	*	*	*6	*	*	*	*	*	*
823.3 . . . . .	10	*9	*	*	*7	*	*	*	*	*	*
823.32 . . . . .	*8	*7	*	*	*6	*	*	*	*	*	*
823.8 . . . . .	22	15	*7	*	*9	*7	*	*5	*	*9	*
823.80 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
823.82 . . . . .	12	*9	*	*	*	*	*	*	*	*	*
823.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
824 . . . . .	124	54	70	*6	61	35	22	26	29	47	22
824.0 . . . . .	9	*	*	*	*5	*	*	*	*	*	*
824.2 . . . . .	14	*	*9	*	*6	*	*	*	*	*7	*
824.4 . . . . .	26	11	15	*	13	*7	*6	*6	*7	*8	*
824.6 . . . . .	27	*7	20	—	10	12	*	*7	*8	*6	*5
824.8 . . . . .	39	20	19	*	20	*9	*7	*8	*8	18	*5
825 . . . . .	43	26	18	*	21	13	*7	*8	10	15	10
825.0 . . . . .	12	9	*	*	*6	*	*	*	*	*	*
825.2 . . . . .	27	13	13	*	13	*7	*6	*5	*6	*8	*7
825.21 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
825.25 . . . . .	16	*8	*8	*	*7	*	*	*	*	*5	*
826 . . . . .	11	*6	*6	*	*	*	*	*	*	*	*
826.0 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
831 . . . . .	16	11	*	*	9	*	*	*	*	*9	*
831.0 . . . . .	16	11	*	*	9	*	*	*	*	*9	*
831.01 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
831.04 . . . . .	*7	*6	*	*	*6	*	*	*	*	*	*
835 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
835.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
836 . . . . .	38	25	13	*	24	*7	*	14	*5	13	*5
836.0 . . . . .	19	15	*	*	12	*	*	*8	*	*6	*
836.1 . . . . .	14	*8	*6	*	9	*	*	*6	*	*6	*
839 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
840 . . . . .	60	33	28	*	15	24	21	13	15	20	12
840.4 . . . . .	50	27	24	—	*7	22	21	12	12	17	9
840.9 . . . . .	*7	*	*	*	*5	*	*	*	*	*	*
842 . . . . .	*7	*	*	*	*5	*	*	*	*	*	*
844 . . . . .	48	29	19	*	28	*7	11	10	12	15	11
844.1 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
844.2 . . . . .	17	11	*6	*	14	*	*	*	*	*6	*
844.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
844.9 . . . . .	15	*7	*8	*	*6	*	*5	*	*6	*	*
845 . . . . .	33	13	21	*	18	*8	*6	*6	*	12	12
845.0 . . . . .	31	12	19	*	16	*8	*6	*6	*	12	10
845.00 . . . . .	15	*	12	*	*7	*	*	*	*	*	*6
845.01 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
845.09 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
846 . . . . .	12	*	*8	—	*	*	*5	*	*	*	*
846.0 . . . . .	10	*	*6	—	*	*	*	*	*	*	*
847 . . . . .	47	22	25	*	27	11	*8	*8	15	11	12
847.0 . . . . .	25	12	13	*	15	*	*	*	*8	*	*8
847.1 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
847.2 . . . . .	13	*6	*7	*	*6	*	*	*	*	*	*
848 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
850 . . . . .	70	40	30	*7	39	*8	15	24	18	20	*9
850.0 . . . . .	17	9	*7	*	*7	*	*6	*6	*	*	*
850.1 . . . . .	23	13	10	*	12	*	*	10	*	*6	*
850.5 . . . . .	10	*7	*	*	*9	*	*	*	*	*	*
850.9 . . . . .	19	10	9	*	10	*	*5	*	*7	*6	*
851 . . . . .	18	12	*6	*	*8	*	*	*5	*	*	*5
851.8 . . . . .	12	*8	*	*	*5	*	*	*	*	*	*
852 . . . . .	40	26	14	*5	12	*7	16	11	*8	9	13
852.0 . . . . .	10	*6	*	*	*	*	*	*	*	*	*
852.2 . . . . .	27	17	10	*	*6	*	14	*7	*5	*5	10

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
852.20	*9	*	*	*	*	*	*6	*	*	*	*
852.21	13	*9	*	*	*	*	*6	*	*	*	*5
853	*7	*	*	*	*	*	*	*	*	*	*
853.0	*7	*	*	*	*	*	*	*	*	*	*
854	86	51	35	18	40	11	16	14	25	26	20
854.0	84	50	34	17	39	11	16	13	25	26	19
854.00	26	15	12	*	11	*	*8	*	*8	9	*6
854.01	21	11	9	*7	*7	*	*	*6	*6	*6	*
854.02	20	14	*6	*	12	*	*	*	*6	*6	*6
854.06	11	*6	*	*	*7	*	*	*	*	*	*
860	60	42	18	*	36	11	11	9	15	23	13
860.0	30	20	10	*	17	*5	*7	*6	*8	11	*6
860.2	*7	*	*	*	*	*	*	*	*	*	*
860.4	12	*8	*	*	*8	*	*	*	*	*	*
861	36	25	11	*	21	*6	*7	*6	*8	16	*7
861.0	*6	*	*	*	*	*	*	*	*	*	*
861.01	*5	*	*	*	*	*	*	*	*	*	*
861.2	24	16	*8	*	16	*	*	*	*5	11	*
861.21	23	15	*8	*	15	*	*	*	*5	11	*
862	*8	*7	*	*	*5	*	*	*	*	*	*
863	23	17	*6	*	15	*	*	*	*6	*8	*7
863.5	*5	*	*	*	*	*	—	*	*	*	*
864	22	16	*7	*	14	*	*	*5	*	9	*
864.0	17	10	*6	*	*9	*	*	*	*	*7	*
864.01	*6	*	*	*	*	*	*	*	*	*	*
864.1	*6	*5	*	*	*	*	*	*	*	*	*
865	25	18	*7	*	18	*	*	*	*6	12	*
865.0	22	15	*6	*	15	*	*	*	*5	10	*
866	16	12	*	*	10	*	*	*	*	*5	*
866.0	13	9	*	*	*6	*	*	*	*	*	*
866.01	*7	*5	*	*	*	*	*	*	*	*	*
867	10	*8	*	*	*	*	*5	*	*	*	*
867.0	10	*8	*	*	*	*	*5	*	*	*	*
868	16	10	*6	*	9	*	*	*	*	*7	*5
868.0	12	*7	*5	*	*6	*	*	*	*	*	*
870	15	11	*	*	*8	*	*	*	*	*	*
870.0	*6	*	*	*	*	*	*	*	*	*	*
870.8	*6	*	*	*	*	*	*	*	*	*	*
871	12	*6	*5	*	*7	*	*	*	*	*5	*
871.1	*6	*	*	*	*	*	*	*	*	*	*
872	12	10	*	*	*9	*	*	*	*	*	*
872.0	*8	*6	*	*	*6	*	*	*	*	*	*
873	189	117	72	19	102	23	45	47	50	55	37
873.0	52	30	22	*	23	*8	16	14	15	13	10
873.2	12	9	*	*	*7	*	*	*	*	*	*
873.20	10	*7	*	*	*5	*	*	*	*	*	*
873.4	101	62	39	10	58	13	21	24	25	32	20
873.40	*6	*	*	*	*	*	*	*	*	*	*
873.41	9	*5	*	*	*6	*	*	*	*	*	*
873.42	56	31	25	*7	27	*7	15	13	14	16	13
873.43	14	10	*	*	11	*	*	*	*	*5	*
873.44	9	*6	*	*	*5	*	*	*	*	*	*
873.49	*7	*5	*	*	*	*	*	*	*	*	*
873.6	13	9	*	*	*8	*	*	*	*	*	*
873.63	*7	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
874 . . . . .	*9	*7	*	*	*6	*	*	*	*	*	*
874.8 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
875 . . . . .	14	12	*	*	10	*	*	*	*	10	*
875.0 . . . . .	11	10	*	*	*8	*	*	*	*	*8	*
876 . . . . .	*6	*5	*	*	*6	—	—	*	*	*	*
877 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
878 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
879 . . . . .	17	13	*	*	13	*	*	*	*	*7	*
880 . . . . .	10	*7	*	*	*8	*	*	*	*	*	*
880.0 . . . . .	*8	*6	*	*	*6	*	*	*	*	*	*
881 . . . . .	43	27	16	*	28	*	11	*9	11	14	10
881.0 . . . . .	33	19	14	*	20	*	10	*7	9	*9	*8
881.00 . . . . .	9	*6	*	*	*5	*	*	*	*	*	*
881.01 . . . . .	14	*8	*6	*	*6	*	*6	*	*	*	*
881.02 . . . . .	10	*6	*	*	*8	*	*	*	*	*	*
881.2 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
882 . . . . .	25	19	*5	*	17	*	*	*6	*7	*6	*6
882.0 . . . . .	14	10	*	*	9	*	*	*	*	*	*
882.1 . . . . .	*8	*7	*	*	*6	*	*	*	*	*	*
883 . . . . .	33	25	*8	*	23	*	*	*7	*7	12	*7
883.0 . . . . .	12	9	*	*	*9	*	*	*	*	*	*
883.1 . . . . .	9	*8	*	*	*6	*	*	*	*	*	*
883.2 . . . . .	11	*8	*	*	*8	*	*	*	*	*6	*
884 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
886 . . . . .	*7	*7	*	*	*	*	*	*	*	*	*
886.0 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
890 . . . . .	12	10	*	*	*8	*	*	*	*	*6	*
890.0 . . . . .	*8	*7	*	*	*5	*	*	*	*	*	*
891 . . . . .	30	19	11	*	17	*5	*	*6	*7	10	*8
891.0 . . . . .	19	12	*7	*	12	*	*	*	*	*5	*
891.1 . . . . .	*9	*5	*	*	*	*	*	*	*	*	*
892 . . . . .	13	9	*	*	*6	*	*	*	*	*	*
892.1 . . . . .	9	*7	*	*	*	*	*	*	*	*	*
893 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
902 . . . . .	*6	*5	*	*	*6	*	*	*	*	*	*
903 . . . . .	*8	*6	*	*	*6	*	*	*	*	*	*
904 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
905 . . . . .	46	22	24	*	18	12	14	10	11	15	*9
905.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
905.3 . . . . .	*6	*	*	*	*	*	*5	*	*	*	*
905.4 . . . . .	16	*8	*8	*	*6	*5	*	*	*	*5	*
906 . . . . .	15	12	*	*	*7	*	*	*	*	*	*
907 . . . . .	26	20	*7	*	16	*5	*	*6	*6	9	*6
907.0 . . . . .	10	*8	*	*	*5	*	*	*	*	*	*
907.2 . . . . .	13	11	*	*	10	*	*	*	*	*5	*
908 . . . . .	10	*6	*	*	*	*	*	*	*	*	*
908.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
909 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
910 . . . . .	35	27	*9	*8	18	*	*6	12	10	*8	*5
910.0 . . . . .	33	26	*7	*7	17	*	*	12	*9	*7	*5
911 . . . . .	*9	*6	*	*	*	*	*	*	*	*	*
911.0 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
912 . . . . .	*7	*5	*	*	*6	*	*	*	*	*	*
912.0 . . . . .	*7	*5	*	*	*6	*	*	*	*	*	*
913 . . . . .	18	11	*7	*	*8	*	*	*	*	*	*5
913.0 . . . . .	13	*7	*6	*	*7	*	*	*	*	*	*
914 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
916 . . . . .	21	14	*7	*	10	*	*6	*5	*5	*5	*
916.0 . . . . .	17	12	*6	*	*8	*	*5	*	*	*	*
917 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
917.0 . . . . .	*6	*5	*	*	*	*	*	*	*	*	*
919 . . . . .	21	13	*8	*	14	*	*	*	*6	*8	*
919.0 . . . . .	19	13	*7	*	13	*	*	*	*5	*8	*
920 . . . . .	47	24	23	*7	17	*6	16	11	14	14	*9
921 . . . . .	10	*6	*	*	*	*	*	*	*	*	*
922 . . . . .	62	23	39	*	25	12	23	14	23	13	12
922.1 . . . . .	33	13	19	*	14	*7	12	*7	11	*8	*7
922.2 . . . . .	13	*6	*7	*	*	*	*5	*	*	*	*
922.3 . . . . .	11	*	9	*	*	*	*6	*	*	*	*
923 . . . . .	27	12	15	*	10	*	11	*	*7	9	*6
923.0 . . . . .	13	*6	*7	*	*5	*	*6	*	*	*	*
923.00 . . . . .	10	*6	*	*	*	*	*	*	*	*	*
923.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
924 . . . . .	88	34	54	*	32	14	39	15	24	28	20
924.0 . . . . .	27	9	18	*	*	*	18	*	*8	9	*6
924.00 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
924.01 . . . . .	22	*7	15	*	*	*	15	*	*6	*9	*
924.1 . . . . .	21	*6	15	*	*8	*	9	*6	*	*	*7
924.10 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
924.11 . . . . .	13	*	10	*	*6	*	*	*	*	*	*
924.8 . . . . .	31	15	15	*	16	*	*8	*	10	11	*6
933 . . . . .	14	*7	*7	*	*	*	*8	*	*	*	*
933.1 . . . . .	11	*5	*6	*	*	*	*6	*	*	*	*
934 . . . . .	12	*8	*	*	*	*	*6	*	*	*	*
934.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
935 . . . . .	13	*9	*	*	*	*	*	*	*	*6	*
935.1 . . . . .	10	*7	*	*	*	*	*	*	*	*	*
941 . . . . .	12	*7	*5	*5	*	*	*	*	*	*	*
941.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
942 . . . . .	15	9	*5	*	*6	*	*	*	*5	*5	*
942.2 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
943 . . . . .	15	10	*6	*	*8	*	*	*	*5	*5	*
943.2 . . . . .	10	*6	*	*	*5	*	*	*	*	*	*
944 . . . . .	15	12	*	*	*7	*	*	*	*	*	*
944.2 . . . . .	*9	*7	*	*	*	*	*	*	*	*	*
945 . . . . .	26	18	*8	*6	13	*	*	*6	*7	*8	*
945.2 . . . . .	15	10	*	*	*7	*	*	*	*5	*	*
945.3 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*
946 . . . . .	16	10	*6	*	*9	*	*	*	*	*9	*
946.2 . . . . .	*9	*6	*	*	*5	*	*	*	*	*	*
946.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	—
948 . . . . .	37	26	11	10	16	*5	*6	10	*8	16	*
948.0 . . . . .	18	14	*	*7	*8	*	*	*6	*	*6	*
948.00 . . . . .	18	14	*	*7	*8	*	*	*6	*	*6	*
948.1 . . . . .	10	*7	*	*	*	*	*	*	*	*6	*
948.10 . . . . .	*7	*5	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
 —Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
952	*8	*7	*	*	*	*	*	*	*	*	*
955	15	9	*6	*	10	*	*	*	*	*7	*
958	32	23	9	*	16	*7	*7	*6	*7	11	*8
958.3	9	*6	*	*	*	*	*	*	*	*	*
958.7	*7	*	*	*	*	*	*	*	*	*	*
958.8	*8	*7	*	*	*	*	*	*	*	*	*
959	46	22	24	*	30	*	10	*8	13	16	10
959.1	21	*9	12	*	14	*	*	*	*6	*	*6
959.8	*7	*	*	*	*	*	*	*	*	*	*
962	*7	*	*	*	*	*	*	*	*	*	*
962.3	*6	*	*	*	*	*	*	*	*	*	*
963	10	*	*7	*	*8	*	*	*	*	*	*
963.0	*8	*	*6	*	*7	*	—	*	*	*	*
964	*5	*	*	*	*	*	*	*	*	*	*
965	74	29	45	*8	54	*8	*	16	20	22	15
965.0	15	9	*6	*	11	*	*	*	*	*	*5
965.09	*6	*	*	*	*	*	*	*	*	*	*
965.1	12	*	10	*	*8	*	*	*	*	*	*
965.4	29	10	19	*	23	*	*	*6	*8	9	*5
965.6	*9	*	*6	*	*7	*	*	*	*	*	*
965.8	*6	*	*	*	*	*	*	*	*	*	*
966	17	*8	*9	*	*6	*	*	*	*	*7	*
966.1	*7	*	*	*	*	*	*	*	*	*	*
966.3	9	*	*	*	*	*	*	*	*	*	*
967	13	*6	*7	*	9	*	*	*	*	*6	*
967.0	*6	*	*	*	*	*	*	*	*	*	*
968	13	*8	*5	*	11	*	—	*	*	*	*
968.5	*8	*6	*	—	*7	*	—	*	*	*	*
969	91	34	57	*9	59	19	*	16	22	34	18
969.0	39	14	26	*5	26	*7	*	*7	10	17	*6
969.4	33	13	20	*	21	*8	*	*6	*7	12	*8
969.7	*5	*	*	*	*	*	—	*	*	*	*
970	*6	*	*	—	*6	—	—	*	*	*	*
970.8	*6	*	*	—	*6	—	—	*	*	*	*
971	*5	*	*	*	*	*	—	*	*	*	*
972	21	*8	13	*	*	*	13	*	*7	*8	*
972.1	14	*5	9	*	*	*	12	*	*6	*6	*
975	*6	*	*	*	*	*	*	*	*	*	*
977	18	*6	13	*	13	*	*	*	*	11	*
977.8	*6	*	*5	*	*	*	*	*	*	*	*
977.9	11	*	*7	*	*8	*	*	*	*	*8	*
980	19	*7	12	*	15	*	*	*6	*	*5	*
980.0	14	*	*9	*	12	*	*	*	*	*	*
983	*6	*	*	*	*	*	*	*	*	*	*
986	*5	*	*	*	*	*	*	*	*	*	*
987	*6	*	*	*	*	*	*	*	*	*	*
989	12	*7	*	*	*6	*	*	*	*	*6	*
989.5	*9	*5	*	*	*	*	*	*	*	*5	*
991	11	*8	*	*	*	*	*5	*	*	*	*
991.6	10	*6	*	*	*	*	*	*	*	*	*
992	9	*5	*	*	*	*	*	*	*	*5	*
994	*6	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
995 . . . . .	84	30	54	10	16	18	39	23	26	27	*8
995.0 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
995.1 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
995.2 . . . . .	50	17	34	*	*5	12	31	15	13	18	*5
995.5 . . . . .	*7	*	*	*6	*	*	—	*	*	*	*
996 . . . . .	608	287	322	35	121	168	284	146	111	224	127
996.0 . . . . .	40	20	20	*	*	*6	31	12	*8	10	10
996.01 . . . . .	26	12	14	*	*	*	22	*7	*5	*7	*7
996.1 . . . . .	24	12	12	*	*	*5	12	*6	*6	*7	*
996.2 . . . . .	20	12	*8	9	*7	*	*	*5	*	*7	*
996.3 . . . . .	10	*5	*5	—	*	*	*6	*	*	*	*
996.39 . . . . .	*7	*	*	—	*	*	*	*	*	*	*
996.4 . . . . .	92	38	54	*	16	21	54	23	17	29	22
996.5 . . . . .	33	13	20	*	*5	12	14	*8	*	12	*8
996.59 . . . . .	25	10	16	*	*	9	12	*6	*	*9	*7
996.6 . . . . .	174	77	97	14	36	49	75	38	39	65	32
996.61 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
996.62 . . . . .	79	34	45	*7	21	24	26	18	17	26	17
996.64 . . . . .	25	10	15	*	*	*	19	*	*8	10	*
996.66 . . . . .	16	*7	9	*	*	*	10	*	*	*7	*
996.67 . . . . .	10	*5	*	*	*	*	*	*	*	*	*
996.69 . . . . .	30	13	17	*	*6	10	12	*8	*5	11	*7
996.7 . . . . .	183	89	94	*	31	57	92	43	27	73	40
996.72 . . . . .	14	*6	*8	*	*	*	9	*	*	*	*
996.73 . . . . .	71	35	36	*	14	26	31	16	*8	28	19
996.74 . . . . .	49	24	25	*	*	15	27	12	*7	21	9
996.76 . . . . .	11	9	*	*	*	*	*8	*	*	*	*
996.77 . . . . .	9	*	*6	—	*	*	*5	*	*	*	*
996.78 . . . . .	13	*5	*8	*	*7	*	*	*	*	*	*
996.79 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
996.8 . . . . .	32	21	12	*	17	12	*	*7	*	18	*
996.81 . . . . .	20	13	*7	*	12	*7	*	*	*	13	*
997 . . . . .	639	319	320	23	98	171	346	169	139	193	138
997.0 . . . . .	19	10	9	*	*	*	12	*	*	*7	*
997.1 . . . . .	146	89	57	*	*9	38	95	47	30	40	29
997.2 . . . . .	23	*8	15	*	*	*6	12	*	*5	*7	*6
997.3 . . . . .	152	74	78	*7	23	46	75	39	35	46	32
997.4 . . . . .	178	70	108	11	39	47	81	37	46	55	41
997.5 . . . . .	89	50	38	*	17	20	51	30	14	28	17
997.6 . . . . .	20	11	*9	*	*	*6	12	*	*	*6	*5
997.62 . . . . .	14	*8	*6	*	*	*	*9	*	*	*	*
997.9 . . . . .	12	*6	*7	*	*	*	*8	*	*	*	*
998 . . . . .	608	265	343	21	146	170	271	140	128	209	131
998.1 . . . . .	198	91	107	*5	43	50	100	54	40	60	44
998.2 . . . . .	62	22	40	*	16	18	26	13	13	22	14
998.3 . . . . .	40	19	21	*	*7	14	17	*7	9	14	9
998.5 . . . . .	170	71	99	*6	41	51	71	40	35	58	37
998.6 . . . . .	9	*	*8	*	*	*	*	*	*	*5	*
998.8 . . . . .	117	54	63	*7	32	30	47	23	26	46	22
998.81 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
998.89 . . . . .	111	51	60	*7	31	28	46	23	25	44	19
999 . . . . .	55	24	31	*	14	14	23	15	11	19	11
999.2 . . . . .	20	*8	12	*	*5	*	*9	*7	*	*7	*
999.3 . . . . .	14	*7	*7	*	*5	*	*	*	*	*	*
999.8 . . . . .	*9	*	*	*	*	*	*5	*	*	*	*
999.9 . . . . .	9	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V01-V82 . . . . .	7,913	1,714	6,198	245	4,804	902	1,961	1,608	1,953	2,674	1,677
V01 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V02 . . . . .	17	*	15	*	15	*	*	*	*6	*	*
V02.5 . . . . .	13	*	12	*	12	*	*	*	*	*	*
V04 . . . . .	10	*	10	—	10	*	*	*	*	*	*
V04.3 . . . . .	10	—	10	—	10	—	—	*	*	*	*
V05 . . . . .	*6	*	*	*5	*	*	—	*	*	*	*
V05.3 . . . . .	*6	*	*	*5	*	—	—	*	*	*	*
V07 . . . . .	27	*	26	*	24	*	*	10	*	*5	*8
V07.2 . . . . .	23	*	22	—	22	*	*	10	*	*	*6
V08 . . . . .	34	23	10	*	26	*6	*	18	*5	*8	*
V09 . . . . .	18	*9	9	*	*	*	10	*	*6	*6	*
V09.0 . . . . .	16	*8	*8	*	*	*	10	*	*6	*6	*
V10 . . . . .	799	317	482	*	48	202	544	218	210	226	145
V10.0 . . . . .	184	85	99	*	*6	39	140	52	45	56	32
V10.02 . . . . .	*8	*	*	—	*	*	*5	*	*	*	*
V10.04 . . . . .	12	*7	*5	—	*	*	11	*	*	*	*
V10.05 . . . . .	126	55	70	—	*	23	99	35	33	38	19
V10.06 . . . . .	19	*9	10	*	*	*	14	*	*5	*	*
V10.09 . . . . .	*9	*	*	—	*	*	*6	*	*	*	*
V10.1 . . . . .	52	33	19	*	*	18	34	14	14	15	*9
V10.11 . . . . .	52	32	19	*	*	18	34	14	14	14	*9
V10.2 . . . . .	21	17	*	*	*	11	10	*	*6	*8	*
V10.21 . . . . .	18	16	*	—	*	9	*9	*	*5	*7	*
V10.3 . . . . .	208	*	205	*	12	58	137	56	56	58	37
V10.4 . . . . .	177	95	82	*	13	36	128	46	49	44	38
V10.41 . . . . .	20	—	20	—	*	*7	*7	*	*	*7	*
V10.42 . . . . .	29	—	29	—	*	*5	23	*7	11	*	*7
V10.43 . . . . .	26	—	26	*	*	12	12	*7	*7	*8	*
V10.44 . . . . .	*6	—	*6	—	*	*	*	*	*	*	*
V10.46 . . . . .	90	90	—	—	*	*8	81	25	24	22	19
V10.5 . . . . .	60	40	21	*	*	15	43	17	16	19	9
V10.51 . . . . .	33	24	10	*	*	*7	26	9	11	*8	*6
V10.52 . . . . .	25	15	11	*	*	*8	16	*6	*5	10	*
V10.7 . . . . .	13	*5	*8	*	*	*	*6	*	*	*	*
V10.72 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V10.79 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V10.8 . . . . .	76	36	40	*	11	21	41	22	20	22	13
V10.82 . . . . .	19	9	10	—	*	*6	10	*	*	*6	*
V10.83 . . . . .	30	15	15	*	*	*8	21	9	11	*7	*
V10.87 . . . . .	9	*	*6	*	*	*	*	*	*	*	*
V10.89 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V11 . . . . .	28	17	11	*	*7	11	9	*	*7	12	*
V11.3 . . . . .	13	11	*	—	*	*6	*	*	*	*5	*
V11.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V12 . . . . .	480	229	251	14	64	105	296	99	128	165	89
V12.0 . . . . .	38	18	20	*	13	*7	16	*8	*6	16	*9
V12.00 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
V12.01 . . . . .	14	*8	*6	—	*	*	*9	*	*	*5	*
V12.09 . . . . .	12	*	*8	*	*8	*	*	*	*	*6	*
V12.2 . . . . .	16	*8	*8	*	*	*	10	*	*	*7	*
V12.3 . . . . .	10	*	*8	*	*	*	*6	*	*	*	*
V12.4 . . . . .	19	11	*8	*	*5	*	*7	*6	*	*	*
V12.5 . . . . .	245	120	125	*	16	50	175	54	69	83	38
V12.6 . . . . .	35	19	16	*	10	10	13	*7	13	9	*7

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V12.7	118	51	67	*	16	31	70	21	31	40	25
V12.70	*9	*	*6	*	*	*	*6	*	*	*	*
V12.71	65	28	37	—	*6	17	42	10	17	24	14
V12.72	10	*6	*	—	*	*	*6	*	*	*	*
V12.79	34	15	19	*	*7	10	17	*6	*9	12	*8
V13	63	32	32	13	17	13	21	13	22	19	9
V13.0	21	12	10	*	*6	*7	*8	*6	*	*8	*
V13.01	13	10	*	—	*	*	*5	*	*	*	*
V13.09	*7	*	*5	*	*	*	*	*	*	*	*
V13.2	*9	*	*6	*	*6	*	*	*	*	*	*
V13.5	11	*	*7	*	*	*	*5	*	*	*	*
V13.7	11	*7	*	11	*	—	—	*	*8	*	*
V14	14	*6	*8	*	*	*5	*6	*	*5	*	*
V14.0	*6	*	*	*	*	*	*	*	*	*	*
V15	285	159	126	23	71	90	102	55	71	89	70
V15.0	*5	*	*	*	*	*	*	*	*	*	*
V15.1	16	*7	*9	10	*	*	*	*7	*	*	*
V15.3	15	*7	*8	*	*	*	*6	*	*	*5	*
V15.5	18	10	*8	*	*7	*	*8	*	*	9	*
V15.8	224	128	96	9	54	79	81	37	58	69	59
V15.81	82	42	39	*9	35	25	17	14	22	30	16
V15.82	121	77	44	—	14	50	57	19	33	31	37
V15.89	21	*9	12	*	*6	*	*7	*	*	*8	*6
V16	11	*	9	—	*	*	*	*	*	*	*
V17	17	9	*8	*	*	*7	*6	*	*6	*	*
V17.3	12	*6	*6	*	*	*	*5	*	*	*	*
V22	56	—	56	*	55	*	—	*7	16	22	11
V22.1	*7	—	*7	—	*7	—	—	*	*	*	*
V22.2	48	—	48	*	47	*	—	*6	14	20	*8
V23	64	—	64	*	63	—	—	*9	12	30	13
V23.7	43	—	43	*	43	—	—	*5	*7	22	*9
V23.8	*8	—	*8	*	*8	—	—	*	*	*	*
V24	*7	—	*7	*	*7	—	—	*	*	*	*
V24.0	*7	—	*7	*	*7	—	—	*	*	*	*
V25	312	*	311	*	310	*	*	41	59	157	54
V25.2	305	*	304	—	304	*	*	41	59	151	54
V27	3,766	—	3,766	12	3,752	*	—	633	821	1,359	953
V27.0	2,844	—	2,844	10	2,833	*	—	474	596	1,059	716
V27.1	22	—	22	*	22	—	—	*	*	12	*
V27.2	37	—	37	—	37	*	—	*	*8	15	10
V27.9	861	—	861	*	858	*	—	150	214	272	225
V29	17	*9	*9	16	*	—	—	*	9	*	*
V29.0	16	*8	*8	15	*	—	—	*	*9	*	*
V30	16	*8	*8	16	—	—	*	*7	*	*5	*
V30.1	16	*8	*8	16	—	—	—	*7	*	*5	*
V42	34	18	15	*6	11	12	*	13	*7	*8	*5
V42.0	15	*9	*6	*	*6	*7	*	*	*	*	*
V42.8	*5	*	*	*	*	*	*	*	*	*	*
V43	111	49	61	*	*7	21	82	27	47	21	15
V43.3	42	22	19	*	*	*9	29	13	16	*8	*
V43.6	65	25	39	*	*	12	50	13	29	12	11
V43.64	36	15	21	—	*	*6	28	*8	15	*6	*6
V43.65	28	10	18	—	*	*5	22	*	14	*5	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V44 . . . . .	63	29	34	15	*9	10	29	24	25	9	*5
V44.0 . . . . .	11	*6	*	*	*	*	*	*	*	*	*
V44.1 . . . . .	17	*8	9	*8	*	*	*6	*7	*6	*	*
V44.2 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V44.3 . . . . .	23	9	14	*	*	*	15	*8	10	*	*
V45 . . . . .	497	285	212	19	36	122	319	143	199	94	61
V45.0 . . . . .	85	42	43	*	*	9	72	27	32	17	*8
V45.01 . . . . .	75	37	38	*	*	*7	65	23	29	16	*7
V45.02 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
V45.1 . . . . .	26	10	16	*	*7	*8	10	*5	*	*9	*9
V45.2 . . . . .	16	9	*6	10	*	*	*	*6	*6	*	*
V45.8 . . . . .	362	221	142	*8	21	101	232	102	156	64	41
V45.81 . . . . .	230	152	77	*	*	60	166	72	93	42	23
V45.82 . . . . .	62	40	22	—	*	27	30	17	23	13	*9
V45.89 . . . . .	71	28	42	*8	13	14	36	12	40	10	*9
V46 . . . . .	11	*7	*	*	*	*	*	*	*	*	*
V46.1 . . . . .	11	*7	*	*	*	*	*	*	*	*	*
V49 . . . . .	42	27	16	*	*	14	23	9	14	13	*6
V49.7 . . . . .	40	25	15	*	*	14	22	9	14	12	*6
V49.75 . . . . .	19	12	*7	*	*	*6	11	*	*6	*6	*
V49.76 . . . . .	12	*7	*5	*	*	*	*8	*	*5	*	*
V53 . . . . .	*7	*	*	*	*	*	*5	*	*	*	*
V53.3 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V54 . . . . .	26	10	16	*	*8	*5	10	*	*6	9	*6
V54.0 . . . . .	10	*6	*	*	*6	*	*	*	*	*	*
V54.8 . . . . .	9	*	*7	*	*	*	*6	*	*	*	*
V54.9 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V55 . . . . .	34	18	16	*7	*8	*9	10	12	10	*8	*
V55.3 . . . . .	21	10	10	*	*5	*6	*8	*6	*7	*6	*
V57 . . . . .	360	145	215	*	34	58	265	41	109	142	68
V57.1 . . . . .	128	48	79	*	*7	16	103	19	24	67	18
V57.2 . . . . .	33	13	20	*	*	*5	25	*	11	12	*7
V57.21 . . . . .	32	13	20	*	*	*5	24	*	11	12	*7
V57.8 . . . . .	191	79	111	*	23	36	130	20	73	58	40
V57.89 . . . . .	190	79	111	*	23	36	130	20	72	58	40
V58 . . . . .	369	156	213	32	62	135	140	108	67	136	58
V58.0 . . . . .	59	28	31	*	*7	22	29	14	*8	26	11
V58.1 . . . . .	284	119	166	30	50	108	96	88	50	106	40
V58.4 . . . . .	12	*	*7	*	*	*	*	*	*	*	*
V58.49 . . . . .	12	*	*7	*	*	*	*	*	*	*	*
V58.8 . . . . .	11	*	*8	*	*	*	9	*	*5	*	*
V58.89 . . . . .	11	*	*7	*	*	*	9	*	*5	*	*
V59 . . . . .	*5	*	*	*	*	*	—	*	*	*	*
V60 . . . . .	15	*8	*7	*	*8	*	*	*	*6	*	*
V60.0 . . . . .	*7	*	*	*	*5	*	*	*	*	*	*
V61 . . . . .	58	22	36	14	38	*	*	15	17	22	*
V61.1 . . . . .	16	*8	*8	*	13	*	*	*	*	*7	*
V61.2 . . . . .	18	*7	10	10	*7	*	—	*7	*8	*	*
V61.20 . . . . .	13	*5	*8	*6	*7	*	—	*6	*	*	*
V61.5 . . . . .	*9	*	*8	*	*8	—	—	*	*	*6	*
V61.8 . . . . .	*9	*	*	*	*	*	*	*	*	*5	*
V62 . . . . .	37	20	17	*	25	*6	*	13	*6	17	*
V62.5 . . . . .	9	*8	*	*	*7	*	*	*	*	*8	—
V62.8 . . . . .	23	9	13	*	14	*	*	10	*	*7	*
V62.81 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
V62.89 . . . . .	13	*6	*8	*	*9	*	*	*6	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 3. Number of all-listed diagnoses for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table; see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of all-listed diagnoses in thousands											
V64 . . . . .	50	22	27	*	14	15	19	12	11	17	9
V64.1 . . . . .	26	12	14	*	*6	*8	10	*7	*6	*9	*
V64.2 . . . . .	12	*6	*7	*	*	*	*	*	*	*	*
V64.3 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
V65 . . . . .	21	10	11	*	*8	*	*7	*	*	*	11
V65.3 . . . . .	13	*6	*7	*	*	*	*	*	*	*	*8
V66 . . . . .	20	9	11	*	*	*	10	*	*	*5	9
V66.1 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V67 . . . . .	11	*	*6	*	*	*	*7	*	*	*	*
V67.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
V70 . . . . .	13	*7	*7	*	*6	*	*	*6	*	*	*
V70.7 . . . . .	12	*6	*6	*	*6	*	*	*6	*	*	*
V71 . . . . .	32	15	17	11	11	*	*6	*8	12	*7	*
V71.4 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
V71.8 . . . . .	18	*7	11	10	*5	*	*	*	9	*	*
V72 . . . . .	10	*5	*	*	*	*	*	*	*	*	*8
V72.5 . . . . .	*8	*	*	—	*	*	*	—	—	—	*8

\* Figure does not meet standard of reliability or precision.

— Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
All codes . . . . .	39,807	15,649	24,158	2,023	14,651	8,915	14,218	10,253	8,511	13,074	7,969
01–05 . . . . .	954	442	512	223	324	187	220	230	216	321	187
01 . . . . .	117	64	53	17	37	27	36	28	27	36	26
01.0 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
01.1 . . . . .	22	14	*7	*	*8	*5	*6	*5	*7	*	*
01.14 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
01.18 . . . . .	11	*6	*	*	*	*	*	*	*	*	*
01.2 . . . . .	22	12	9	*6	*8	*	*	*5	*	*6	*7
01.24 . . . . .	18	10	*7	*	*7	*	*	*	*	*	*6
01.3 . . . . .	30	15	15	*	*8	*6	14	*7	*7	10	*6
01.31 . . . . .	19	10	*8	*	*	*	11	*	*5	*6	*
01.39 . . . . .	11	*	*6	*	*	*	*	*	*	*	*
01.5 . . . . .	37	19	19	*	12	12	9	10	*7	13	*7
01.51 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
01.59 . . . . .	30	15	15	*	10	10	*7	*8	*5	10	*6
02 . . . . .	88	52	36	31	27	16	15	24	22	30	13
02.0 . . . . .	15	*9	*6	*	*7	*	*	*	*	*6	*
02.06 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
02.1 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
02.12 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
02.2 . . . . .	15	*9	*6	*	*	*	*	*	*	*	*
02.3 . . . . .	15	*8	*8	*6	*	*	*	*	*	*5	*
02.34 . . . . .	12	*6	*6	*	*	*	*	*	*	*	*
02.4 . . . . .	22	15	*7	12	*5	*	*	*6	*6	*6	*
02.42 . . . . .	19	13	*5	11	*	*	*	*5	*5	*5	*
02.9 . . . . .	13	*7	*5	*	*6	*	*	*	*	*	*
02.94 . . . . .	*8	*5	*	*	*	*	*	*	*	*	*
03 . . . . .	673	292	381	169	227	125	152	160	155	228	130
03.0 . . . . .	102	47	56	*	24	38	39	23	21	34	25
03.09 . . . . .	97	44	53	*	22	36	37	22	20	32	23
03.3 . . . . .	334	176	158	158	85	41	50	79	71	119	66
03.31 . . . . .	332	175	157	157	85	41	49	79	70	118	66
03.4 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
03.5 . . . . .	17	*8	*8	*	*9	*	*	*	*	*	*
03.53 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
03.59 . . . . .	11	*	*7	*	*5	*	*	*	*	*	*
03.6 . . . . .	9	*	*5	*	*	*	*	*	*	*5	*
03.8 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
03.9 . . . . .	195	49	146	*	101	37	54	50	55	57	32
03.90 . . . . .	78	22	56	*	37	16	24	23	23	22	10
03.91 . . . . .	49	9	40	*	30	*7	11	13	17	9	10
03.92 . . . . .	51	14	37	*	23	10	17	11	11	19	10
03.95 . . . . .	11	*	10	*	*8	*	*	*	*	*5	*
04 . . . . .	64	29	35	*	29	16	16	17	9	24	14
04.0 . . . . .	13	*6	*8	*	*	*	*	*	*	*	*
04.07 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
04.3 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
04.4 . . . . .	23	10	13	*	11	*6	*	*8	*	*8	*
04.43 . . . . .	11	*5	*6	*	*	*	*	*5	*	*	*
04.49 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
04.7 . . . . .	*5	*	*	*	*	*	–	*	*	*	*
04.8 . . . . .	11	*	*9	–	*	*	*	*	*	*5	*
04.81 . . . . .	11	*	*8	–	*	*	*	*	*	*5	*
05 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
05.3 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
05.31 . . . . .	*8	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
06–07	85	24	60	*	38	28	16	26	15	30	14
06	67	15	52	*	32	23	12	20	11	25	12
06.2	20	*	16	*	10	*6	*	*7	*	*6	*
06.3	13	*	11	*	*7	*	*	*	*	*5	*
06.39	12	*	10	*	*6	*	*	*	*	*5	*
06.4	11	*	*9	*	*6	*	*	*	*	*	*
06.8	13	*	*8	*	*	*7	*	*	*	*6	*
06.89	11	*	*7	*	*	*6	*	*	*	*6	*
07	17	9	*8	*	*6	*5	*	*6	*	*	*
07.6	*7	*	*	*	*	*	*	*	*	*	*
08–16	269	126	143	15	45	55	154	113	26	103	27
08	33	19	14	*	13	*6	10	11	*8	*7	*7
08.8	22	15	*7	*	11	*	*	*6	*7	*6	*
08.81	20	15	*6	*	11	*	*	*5	*6	*5	*
11	*8	*	*	*	*	*	*	*	*	*	*
12	16	9	*7	*	*	*	*8	*	*	*8	*
12.9	*6	*	*	*	*	*	*	*	*	*	*
13	117	47	70	*	*	17	95	78	*	27	*8
13.4	45	20	26	–	*	*8	36	32	*	10	*
13.41	45	19	26	–	*	*8	36	32	*	9	*
13.5	10	*	*7	*	*	*	*9	*5	*	*	*
13.59	10	*	*7	*	*	*	*9	*5	*	*	*
13.7	55	21	34	*	*	*8	45	38	*	11	*
13.71	54	21	33	*	*	*8	45	38	*	11	*
14	67	32	35	*	9	23	32	9	*8	47	*
14.4	12	*7	*5	*	*	*	*7	*	*	*7	*
14.49	*9	*	*	*	*	*	*	*	*	*	*
14.5	11	*5	*6	*	*	*	*	*	*	*8	*
14.7	29	13	16	*	*	9	13	*	*	21	*
14.74	16	*6	10	*	*	*6	*8	*	*	12	*
14.75	*5	*	*	*	*	*	*	*	*	*	*
14.9	*6	*	*	*	*	*	*	*	*	*	*
16	19	10	*8	*	9	*	*5	*	*	*6	*6
18–20	68	41	27	37	12	9	10	29	11	18	11
18	20	14	*6	*5	*7	*	*	*	*	*7	*
18.4	*7	*	*	*	*	*	*	*	*	*	*
19	*8	*	*	*	*	*	*	*	*	*	*
20	41	23	18	29	*	*	*5	19	*5	10	*6
20.0	31	17	15	25	*	*	*	14	*	*8	*
20.01	26	13	13	22	*	*	*	11	*	*8	*
21–29	353	202	151	75	135	78	64	121	57	112	62
21	102	60	42	*7	43	31	22	33	16	37	17
21.0	27	16	11	*	*	11	12	*6	*6	10	*6
21.01	*8	*	*	–	*	*	*	*	*	*	*
21.02	11	*7	*	–	*	*	*5	*	*	*5	*
21.03	*5	*	*	*	*	*	*	*	*	*	*
21.2	*6	*	*	*	*	*	*	*	*	*	*
21.3	*8	*6	*	*	*	*	*	*	*	*	*
21.31	*6	*	*	*	*	*	*	*	*	*	*
21.6	14	9	*	*	*7	*5	*	*7	*	*	*
21.69	10	*6	*	*	*6	*	*	*6	*	*	*
21.7	10	*7	*	*	*7	*	*	*	*	*	*
21.71	*6	*	*	*	*	*	*	*	*	*	*
21.8	31	17	15	*	16	*9	*	12	*	11	*

\*Figure does not meet standard of reliability or precision.  
–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
21.81	*7	*	*	*	*	*	*	*	*	*	*
21.88	15	*7	*8	*	*7	*7	*	*7	*	*5	*
22	55	27	28	*9	21	19	*8	18	*7	19	12
22.1	*7	*	*	*	*	*	*	*	*	*	*
22.19	*6	*	*	*	*	*	*	*	*	*	*
22.2	11	*5	*6	*	*	*	*	*	*	*	*
22.6	24	12	12	*	9	*8	*	*8	*	*8	*
22.63	18	9	*8	*	*8	*5	*	*7	*	*	*
23	35	21	14	*6	18	*	*7	12	*6	10	*7
23.0	10	*6	*	*	*6	*	*	*	*	*	*
23.09	*8	*	*	*	*6	*	*	*	*	*	*
23.1	20	11	*9	*	11	*	*	*6	*	*6	*
23.19	19	11	*8	*	11	*	*	*6	*	*6	*
24	*7	*	*	*	*	*	*	*	*	*	*
25	*7	*	*	*	*	*	*	*	*	*	*
26	18	9	*9	*	*	*	11	*6	*	*6	*
26.3	14	*7	*6	*	*	*	*8	*	*	*	*
26.31	*6	*	*	*	*	*	*	*	*	*	*
26.32	*6	*	*	*	*	*	*	*	*	*	*
27	47	29	18	13	20	*6	*8	13	9	15	10
27.0	*9	*5	*	*	*	*	*	*	*	*	*
27.4	*7	*	*	*	*	*	*	*	*	*	*
27.5	17	13	*	*	10	*	*	*	*	*	*
27.51	9	*7	*	*	*7	*	*	*	*	*	*
27.6	*8	*	*	*6	*	*	*	*	*	*	*
27.62	*6	*	*	*5	*	—	*	*	*	*	*
28	61	35	27	37	18	*6	*	29	11	15	*6
28.0	10	*6	*	*	*	*	*	*	*	*	*
28.2	16	9	*6	*	10	*	*	*6	*	*	*
28.3	27	14	13	25	*	—	—	16	*	*	*
29	19	13	*6	*	*6	*7	*5	*	*	*8	*
29.1	*6	*	*	*	*	*	*	*	*	*	*
29.4	*9	*7	*	*	*	*5	*	*	*	*	*
30-34	1,041	588	454	56	201	285	498	286	224	348	183
30	15	*7	*7	*	*	*5	*6	*6	*	*	*
30.0	*6	*	*	*	*	*	*	*	*	*	*
31	135	78	56	19	24	32	61	42	28	46	19
31.1	50	30	20	*	12	13	24	14	11	18	*7
31.2	18	10	*7	*	*	*	10	*	*	*6	*
31.29	17	10	*7	*	*	*	9	*	*	*5	*
31.4	57	33	25	13	*8	12	24	19	11	21	*7
31.42	48	27	21	12	*8	*9	19	16	10	16	*7
31.43	*8	*	*	*	*	*	*	*	*	*	*
32	69	34	34	*	9	26	33	21	12	27	*9
32.2	28	13	14	*	*7	*7	13	*7	*	11	*
32.29	24	11	13	*	*6	*6	12	*6	*	10	*
32.3	*8	*	*	*	*	*	*	*	*	*	*
32.4	26	12	14	—	*	12	13	*8	*6	11	*
33	357	207	150	19	63	101	173	100	80	115	61
33.2	346	200	145	18	59	98	170	97	77	112	59
33.22	61	36	26	*5	11	16	30	14	11	20	17
33.23	52	29	23	10	12	13	18	18	10	18	*6
33.24	127	74	54	*	21	41	64	37	32	39	19
33.26	21	12	*9	*	*	*6	14	*7	*6	*6	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
33.27	58	33	25	*	11	16	32	14	12	23	*9
33.28	22	14	*8	*	*	*6	11	*8	*5	*5	*
34	466	260	206	16	102	122	225	117	102	156	92
34.0	180	109	71	10	54	47	69	45	42	58	35
34.03	*6	*	*	*	*	*	*	*	*	*	*
34.04	147	88	59	*7	47	39	53	38	34	49	26
34.09	20	13	*7	*	*6	*6	*7	*	*	*6	*6
34.2	58	34	24	*	12	19	26	18	13	15	11
34.21	19	12	*6	*	*5	*7	*6	*7	*	*	*
34.22	13	*7	*6	—	*	*6	*5	*	*	*	*
34.24	14	*7	*7	—	*	*	9	*5	*	*	*
34.5	11	*9	*	*	*	*	*	*	*	*	*
34.51	*8	*6	*	*	*	*	*	*	*	*	*
34.6	*9	*5	*	*	*	*	*	*	*	*	*
34.7	9	*8	*	*	*	*	*	*	*	*	*
34.79	*7	*5	*	*	*	*	*	*	*	*	*
34.9	183	85	98	*	18	45	118	43	39	65	37
34.91	164	74	90	*	15	41	105	38	35	60	31
34.92	18	10	*8	—	*	*	11	*	*	*	*6
35-39	4,840	2,873	1,967	160	475	1,646	2,558	1,070	1,076	1,743	951
35	110	64	45	30	12	22	46	36	16	33	25
35.1	10	*7	*	*	*	*	*	*	*	*	*
35.12	*5	*	*	*	*	*	*	*	*	*	*
35.2	61	35	26	*	*8	14	37	17	10	23	11
35.21	*6	*	*	*	*	*	*	*	*	*	*
35.22	36	23	13	*	*	*8	23	10	*7	13	*
35.24	16	*7	*8	*	*	*6	*8	*	*	*6	*
35.3	*7	*	*	*	*	*	*	*	*	*	*
35.33	*5	*	*	*	*	*	*	*	*	*	*
35.5	*5	*	*	*	*	*	—	*	*	*	*
35.6	*6	*	*	*	*	*	*	*	*	*	*
35.7	*7	*	*	*	*	*	*	*	*	*	*
35.71	*5	*	*	*	*	*	*	*	*	*	*
35.9	*6	*	*	*	*	*	*	*	*	*	*
36	1,009	710	299	*	50	430	529	202	244	350	212
36.0	434	285	149	*	32	193	209	72	109	153	100
36.01	361	233	128	*	27	160	173	65	90	124	82
36.02	11	*7	*	—	*	*6	*	*	*	*	*
36.05	47	35	13	—	*	20	25	*	12	19	12
36.09	9	*7	*	—	*	*	*	*	*	*	*
36.1	573	423	150	—	17	236	320	130	134	197	112
36.11	49	34	14	—	*	20	25	11	12	15	11
36.12	103	72	31	—	*	41	61	29	24	31	19
36.13	117	86	31	—	*	48	68	22	31	37	27
36.14	75	58	17	—	*	24	48	13	17	32	12
36.15	213	161	53	—	*6	96	112	53	48	74	39
36.16	*9	*7	*	—	*	*5	*	*	*	*	*
37	1,579	943	636	22	124	552	881	362	341	599	278
37.0	11	*	*6	*	*	*	*	*	*	*	*
37.1	*8	*	*	*	*	*	*	*	*	*	*
37.12	*7	*	*	*	*	*	*	*	*	*	*
37.2	1,137	701	436	16	100	460	561	240	251	449	197
37.21	47	28	18	*	*	18	21	12	*7	16	12
37.22	802	505	297	*	67	347	387	151	187	323	141
37.23	219	127	93	*9	17	73	121	54	41	91	34
37.26	52	32	21	*	*8	18	25	18	11	15	*9
37.27	*7	*	*	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
37.3	29	15	14	*	*5	11	10	12	*	10	*
37.34	19	10	9	*	*	*7	*7	*8	*	*7	*
37.6	51	38	13	*	*	17	31	11	13	14	13
37.61	48	36	12	*	*	15	30	10	12	13	13
37.7	178	88	90	*	*	29	143	47	38	62	31
37.71	43	22	21	*	*	*	37	12	*7	18	*6
37.72	66	33	32	—	*	10	53	17	15	24	10
37.75	*7	*	*	*	—	*	*7	*	*	*	*
37.76	12	*5	*7	*	*	*	11	*	*	*	*
37.78	40	20	20	*	*	11	27	11	10	12	*7
37.8	136	69	67	*	*	18	113	36	26	51	23
37.80	9	*	*	*	*	*	*8	*	*	*7	*
37.81	22	11	12	*	*	*	20	*6	*	11	*
37.82	17	10	*7	*	*	*	15	*	*	*	*
37.83	66	35	31	*	*	10	53	18	15	23	10
37.86	*5	*	*	*	*	*	*	*	*	*	*
37.87	10	*	*5	*	*	*	*7	*	*	*	*
37.9	27	20	*6	*	*	*8	14	*9	*	*7	*7
37.94	17	14	*	*	*	*6	9	*	*	*6	*
38	1,021	525	496	76	155	262	528	219	253	361	187
38.0	42	21	21	*	*	14	24	10	*7	14	10
38.03	9	*	*6	*	*	*	*	*	*	*	*
38.08	21	11	10	*	*	*6	15	*6	*	*6	*5
38.1	157	90	67	*	*	38	118	31	41	66	19
38.12	132	78	54	—	*	30	101	26	34	56	16
38.18	19	10	*9	*	*	*5	13	*	*	*7	*
38.2	*7	*	*	—	*	*	*6	*	*	*	*
38.21	*6	*	*	—	*	*	*6	*	*	*	*
38.4	49	39	10	*	*	9	37	*9	11	18	10
38.44	32	26	*6	*	—	*6	26	*8	*8	11	*5
38.45	*8	*5	*	*	*	*	*	*	*	*	*
38.48	*6	*5	*	—	*	*	*	*	*	*	*
38.5	*9	*	*	—	*5	*	*	*	*	*	*
38.59	*8	*	*	—	*5	*	*	*	*	*	*
38.6	12	*7	*5	*	*	*	*8	*	*	*	*
38.7	33	16	18	*	*	9	20	12	*9	10	*
38.8	33	16	17	10	*8	*9	*6	*9	*5	10	*8
38.85	10	*6	*	*8	*	*	*	*	*	*	*
38.86	*7	*	*	*	*	*	*	*	*	*	*
38.9	674	326	349	62	129	176	307	139	172	233	129
38.91	115	67	49	20	15	28	53	29	38	32	17
38.92	*8	*	*	*8	*	*	—	*	*	*	*
38.93	436	197	238	30	95	112	199	87	102	161	86
38.94	*9	*	*	*	*	*	*	*	*	*	*
38.95	87	44	43	*	16	32	38	20	18	30	19
38.98	12	*6	*7	—	*	*	9	*	10	*	—
38.99	*7	*	*	*	*	*	*	*	*	*	*
39	1,121	630	491	32	133	380	575	250	222	400	249
39.2	164	87	77	*	10	49	103	39	33	60	32
39.25	24	12	12	—	*	*7	16	*	*7	*9	*
39.27	47	22	25	*	*8	16	24	11	*6	19	11
39.29	83	49	34	*	*	24	57	21	18	28	16
39.3	23	14	10	*	10	*	*8	*6	*6	*7	*
39.31	14	*8	*6	*	*6	*	*5	*	*	*	*
39.32	*7	*	*	*	*	*	*	*	*	*	*
39.4	94	45	49	*	14	31	48	19	15	33	27
39.42	28	15	13	—	*6	*9	13	*5	*	*9	9

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).



**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
39.43	*8	*	*6	*	*	*	*5	*	*	*	*
39.49	57	27	30	*	*8	20	29	13	10	21	13
39.5	112	54	58	*	13	39	57	29	25	37	21
39.51	13	*	*8	—	*	*7	*	*	*	*	*
39.52	11	*	*6	—	*	*	*8	*	*	*	*
39.56	*5	*	*	*	*	*	*	*	*	*	*
39.57	11	*7	*	*	*	*	*7	*	*	*	*
39.59	64	30	33	*	*6	23	32	16	12	24	11
39.6	362	254	108	21	19	132	190	79	85	120	79
39.61	314	220	95	19	16	116	164	71	73	101	69
39.62	16	13	*	*	*	*7	*8	*	*	*6	*
39.63	*9	*6	*	*	*	*	*	*	*	*	*
39.64	21	14	*7	*	*	*6	12	*	*	10	*
39.9	359	172	187	*	67	124	167	75	58	141	85
39.93	11	*	*6	*	*	*	*7	*	*	*	*
39.95	339	163	176	*	62	119	157	72	53	133	81
40-41	363	188	175	19	70	117	156	110	73	118	61
40	198	102	96	*	27	80	86	60	42	63	33
40.1	48	24	24	*	11	14	21	15	10	17	*7
40.11	46	23	23	*	10	14	20	14	9	16	*7
40.2	56	24	32	*	*8	22	24	16	13	18	*8
40.23	16	*	15	—	*	*7	*6	*5	*	*	*
40.24	*5	*	*	*	*	*	*	*	*	*	*
40.29	31	18	12	*	*	11	16	*8	*8	11	*
40.3	56	29	26	*	*5	23	28	19	9	17	11
40.4	12	*8	*	*	*	*6	*	*	*	*	*
40.41	*9	*6	*	*	*	*	*	*	*	*	*
40.5	25	16	*9	—	*	15	*8	*7	*8	*6	*
40.53	11	*8	*	—	*	*8	*	*	*	*	*
40.59	*6	*	*	—	*	*	*	*	*	*	*
41	165	86	79	14	43	37	70	49	32	56	28
41.0	10	*	*6	*	*	*	*	*	*	*	*
41.04	*5	*	*	*	*	*	*	*	*	*	*
41.3	122	62	60	11	22	25	64	40	25	38	18
41.31	121	62	60	11	22	25	63	40	25	38	18
41.5	26	17	10	*	13	*7	*	*	*	10	*7
41.9	*6	*	*	*	*	*	*	*	*	*	*
42-54	5,074	2,113	2,962	193	1,288	1,270	2,324	1,221	1,159	1,774	920
42	164	82	82	11	23	53	78	50	31	54	29
42.2	82	44	38	*8	11	27	36	26	13	28	15
42.23	72	39	33	*	*9	25	33	22	12	25	13
42.24	*7	*	*	*	*	*	*	*	*	*	*
42.3	27	17	10	*	*	10	13	*7	*6	10	*6
42.33	25	16	9	*	*	9	11	*6	*	9	*
42.9	40	14	27	*	*	10	26	11	9	13	*7
42.92	40	14	26	*	*	9	25	11	9	13	*7
43	225	99	126	16	20	40	149	49	50	90	36
43.1	169	74	95	*8	12	27	121	36	36	73	24
43.11	128	58	70	*	*8	18	100	28	26	55	19
43.19	40	16	25	*6	*	*9	21	*8	10	18	*
43.3	*8	*6	*	*7	*	*	*	*	*	*	*
43.4	16	*	12	*	*	*	10	*	*5	*	*
43.41	10	*	*8	—	*	*	*7	*	*	*	*
43.42	*6	*	*	*	*	*	*	*	*	*	*
43.6	*6	*	*	—	*	*	*	*	*	*	*
43.7	14	*6	*8	—	*	*	*8	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
43.9 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
43.99 . . . . .	*5	*	*	—	*	*	*	*	*	*	*
44 . . . . .	192	94	99	*9	49	45	89	34	40	72	46
44.0 . . . . .	14	9	*	*	*5	*	*6	*	*	*	*
44.00 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
44.01 . . . . .	*6	*	*	—	*	*	*	*	*	*	*
44.1 . . . . .	37	14	22	*	*6	*7	22	*5	10	20	*
44.13 . . . . .	18	*7	11	*	*	*	11	*	*7	*6	*
44.14 . . . . .	17	*6	11	*	*	*	10	*	*	13	*
44.2 . . . . .	11	*7	*	*	*	*	*6	*	*	*	*
44.29 . . . . .	*8	*6	*	*	*	*	*	*	*	*	*
44.3 . . . . .	21	*6	15	*	10	*	*6	*	*	*8	*7
44.39 . . . . .	18	*6	12	*	*7	*	*6	*	*	*5	*7
44.4 . . . . .	64	36	27	*	10	13	40	11	12	21	20
44.41 . . . . .	*9	*	*6	—	*	*	*5	*	*	*	*
44.42 . . . . .	9	*6	*	*	*	*	*6	*	*	*	*
44.43 . . . . .	44	27	17	*	*6	9	28	*8	*8	13	14
44.6 . . . . .	44	20	24	*7	16	15	*7	*8	11	15	11
44.66 . . . . .	30	13	17	*6	10	10	*	*	*6	11	*8
44.69 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
45 . . . . .	1,846	816	1,030	32	264	445	1,105	466	439	637	304
45.0 . . . . .	10	*	*5	*	*	*	*	*	*	*	*
45.1 . . . . .	896	401	495	18	145	221	511	212	212	321	150
45.13 . . . . .	447	212	235	9	69	117	252	94	108	164	81
45.14 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
45.16 . . . . .	435	182	253	*8	73	98	256	117	98	153	67
45.2 . . . . .	515	211	305	*5	64	112	333	144	117	176	78
45.23 . . . . .	239	105	134	*	26	53	159	71	49	82	36
45.24 . . . . .	91	32	59	*	*9	17	64	18	23	34	16
45.25 . . . . .	177	69	108	*	28	41	105	54	40	57	26
45.3 . . . . .	17	*9	*8	*	*6	*	*6	*	*6	*	*
45.33 . . . . .	*8	*	*5	*	*	*	*	*	*	*	*
45.4 . . . . .	94	46	48	*	*	22	66	30	26	28	9
45.42 . . . . .	76	37	38	*	*	18	54	24	23	22	*7
45.43 . . . . .	13	*7	*6	*	*	*	9	*	*	*	*
45.6 . . . . .	50	21	29	*	10	12	25	11	13	16	11
45.62 . . . . .	47	19	28	*	10	11	24	10	12	15	10
45.7 . . . . .	206	96	110	*	23	54	127	49	51	70	36
45.72 . . . . .	*9	*	*	*	*	*	*	*	*	*	*
45.73 . . . . .	59	30	28	*	*	16	38	14	15	19	11
45.74 . . . . .	*7	*	*	*	*	*	*6	*	*	*	*
45.75 . . . . .	28	14	13	*	*	*	20	*6	*8	9	*
45.76 . . . . .	81	35	45	*	10	25	46	21	16	27	17
45.79 . . . . .	23	*8	15	*	*	*5	15	*6	*6	9	*
45.8 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
45.9 . . . . .	50	23	27	*	*7	13	29	12	11	17	10
45.91 . . . . .	12	*	*7	*	*	*	*	*	*	*	*
45.93 . . . . .	17	*8	*8	*	*	*	10	*	*	*	*
45.94 . . . . .	16	*7	*9	*	*	*	11	*	*	*7	*
46 . . . . .	195	93	101	13	38	50	93	47	57	60	31
46.0 . . . . .	13	*7	*7	*	*	*	*8	*	*	*	*
46.03 . . . . .	11	*6	*	*	*	*	*7	*	*	*	*
46.1 . . . . .	50	23	27	*	*8	13	28	10	15	17	*8
46.10 . . . . .	23	12	11	*	*	*6	12	*	*6	*8	*
46.11 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
46.13 . . . . .	17	*7	10	*	*	*	11	*	*6	*5	*
46.2 . . . . .	15	*8	*7	*	*	*	*6	*	*	*	*
46.20 . . . . .	*7	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
46.23	*6	*	*	—	*	*	*	*	*	*	*
46.3	29	15	14	*	*	10	13	10	*8	*7	*
46.32	*6	*	*	*	*	*	*	*	*	*	*
46.39	23	13	9	*	*	*9	9	*8	*7	*5	*
46.4	10	*	*5	*	*	*	*6	*	*	*	*
46.5	32	17	15	*	*8	*7	13	*9	10	9	*
46.51	*6	*	*	*	*	*	*	*	*	*	*
46.52	26	13	13	*	*7	*6	11	*7	*8	*7	*
46.7	27	10	17	*	*8	*7	11	*	*6	10	*6
46.73	11	*	*7	*	*	*	*	*	*	*	*
46.75	*8	*	*6	*	*	*	*	*	*	*	*
46.8	15	*7	*8	*	*	*	*7	*	*6	*	*
46.85	*7	*	*	*	*	*	*	*	*	*	*
47	313	155	158	59	177	44	32	62	76	109	65
47.0	237	135	102	52	139	27	19	48	54	81	55
47.1	73	18	55	*	39	16	13	14	21	27	11
48	125	62	63	*6	23	29	68	41	28	40	16
48.2	46	23	23	*	9	10	22	14	13	12	*6
48.23	25	12	13	*	*6	*6	12	*7	*6	*7	*
48.24	19	10	*8	*	*	*	10	*6	*6	*	*
48.3	19	*9	10	*	*	*	14	*7	*	*7	*
48.35	16	*7	*9	*	*	*	12	*6	*	*6	*
48.5	11	*5	*5	—	*	*	*6	*	*	*	*
48.6	28	13	15	*	*	*7	20	11	*	*9	*
48.63	17	*7	9	—	*	*	12	*8	*	*	*
48.69	*8	*	*	—	*	*	*5	*	*	*	*
48.7	*8	*	*6	*	*	*	*	*	*	*	*
48.8	9	*7	*	*	*5	*	*	*	*	*	*
48.81	*8	*6	*	*	*	*	*	*	*	*	*
49	64	32	32	*	26	19	18	19	11	24	10
49.0	10	*6	*	*	*6	*	*	*	*	*	*
49.01	*9	*6	*	*	*	*	*	*	*	*	*
49.4	26	13	13	—	10	*7	*9	9	*	*9	*
49.46	23	11	12	—	*9	*7	*7	*8	*	*8	*
49.5	*8	*	*	—	*	*	*	*	*	*	*
49.7	*6	*	*	*	*	*	*	*	*	*	*
49.79	*6	*	*	*	*	*	*	*	*	*	*
50	95	47	48	*	25	30	35	23	18	35	20
50.1	69	31	38	*	15	22	29	16	14	26	13
50.11	50	22	28	*	12	15	20	11	10	20	*8
50.12	19	9	10	*	*	*7	*9	*	*	*6	*
50.2	*7	*	*	*	*	*	*	*	*	*	*
50.6	*6	*	*	*	*	*	*	*	*	*	*
50.61	*5	*	*	—	*	*	*	*	*	*	*
50.9	*6	*5	*	*	*	*	*	*	*	*	*
51	724	236	487	*	218	195	306	164	147	258	154
51.0	*6	*	*	*	*	*	*	*	*	*	*
51.1	104	37	66	*	28	26	49	21	23	31	29
51.10	90	32	58	*	26	23	42	19	18	26	27
51.11	*5	*	*	—	*	*	*	*	*	*	*
51.2	470	142	328	*	161	141	165	110	97	175	88
51.22	155	67	87	*	34	48	71	33	34	58	30
51.23	315	74	241	*	127	93	94	77	62	117	58
51.3	*8	*	*6	*	*	*	*	*	*	*	*
51.4	21	9	11	*	*	*	15	*	*	10	*
51.41	14	*7	*7	*	*	*	10	*	*	*7	*
51.5	9	*	*	*	*	*	*5	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
51.51 . . . . .	*9	*	*	*	*	*	*5	*	*	*	*
51.8 . . . . .	92	31	61	*	20	18	53	19	18	31	24
51.85 . . . . .	42	13	29	*	11	*8	22	*8	9	13	12
51.87 . . . . .	18	*8	10	*	*	*	11	*	*	*8	*
51.88 . . . . .	25	*8	17	–	*5	*	16	*	*6	*8	*7
51.9 . . . . .	10	*	*6	*	*	*	*7	*	*	*	*
51.98 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
52 . . . . .	38	15	23	*	*8	15	15	*8	*8	12	10
52.1 . . . . .	14	*7	*7	–	*	*	*8	*	*	*	*5
52.11 . . . . .	*6	*	*	–	*	*	*	*	*	*	*
52.9 . . . . .	10	*	*7	*	*	*	*	*	*	*	*
53 . . . . .	242	132	111	13	51	77	101	68	52	89	33
53.0 . . . . .	68	61	*7	*	12	15	36	25	13	23	*7
53.00 . . . . .	11	*8	*	*	*	*	*6	*	*	*5	*
53.01 . . . . .	10	*9	*	*	*	*	*	*5	*	*	*
53.02 . . . . .	13	11	*	*	*	*	*6	*	*	*	*
53.03 . . . . .	15	14	*	–	*	*	*8	*6	*	*	*
53.04 . . . . .	9	*9	*	–	*	*	*6	*	*	*	*
53.05 . . . . .	10	9	*	–	*	*	*6	*	*	*	*
53.1 . . . . .	11	10	*	*	*	*	*	*	*	*	*
53.2 . . . . .	*7	*	*	–	*	*	*	*	*	*	*
53.4 . . . . .	47	20	27	*	12	17	16	12	10	17	*7
53.41 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
53.49 . . . . .	41	17	24	*	11	14	14	9	10	16	*6
53.5 . . . . .	42	14	28	*	*8	16	17	*7	11	18	*6
53.51 . . . . .	22	*9	14	*	*	*7	11	*	*7	*9	*
53.59 . . . . .	20	*5	14	*	*	*8	*7	*	*	9	*
53.6 . . . . .	49	16	33	*	9	21	19	13	11	20	*
53.61 . . . . .	36	12	24	*	*8	16	13	11	*8	15	*
53.69 . . . . .	13	*	9	–	*	*5	*6	*	*	*6	*
53.7 . . . . .	10	*	*6	*	*	*	*	*	*	*	*
53.9 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
54 . . . . .	851	248	603	23	364	229	236	190	201	296	164
54.0 . . . . .	19	10	*9	*	*5	*6	*6	*	*	*5	*5
54.1 . . . . .	68	30	38	*	27	16	22	18	21	16	13
54.11 . . . . .	46	20	26	*	19	*9	16	11	15	11	*9
54.12 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
54.19 . . . . .	16	*8	*8	*	*5	*	*6	*5	*	*	*
54.2 . . . . .	206	48	158	*6	125	42	34	45	43	69	50
54.21 . . . . .	153	26	127	*5	106	26	15	31	27	55	39
54.23 . . . . .	21	*	16	*	*	*7	*8	*5	*6	*6	*
54.25 . . . . .	24	13	10	*	13	*6	*5	*5	*6	*7	*6
54.3 . . . . .	11	*	*7	*	*	*	*	*	*	*	*
54.4 . . . . .	48	10	38	*	14	14	18	13	11	15	10
54.5 . . . . .	326	59	267	*	148	80	94	69	75	130	52
54.6 . . . . .	11	*6	*	*	*6	*	*	*	*	*	*
54.7 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
54.9 . . . . .	155	79	77	*	33	63	55	35	38	55	28
54.91 . . . . .	82	42	40	*	16	32	33	18	18	32	14
54.93 . . . . .	16	*9	*7	*	*	*5	*6	*	*	*	*
54.95 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
54.98 . . . . .	41	20	21	*	10	19	11	*9	12	13	*8
55–59 . . . . .	1,102	564	538	46	255	285	516	326	240	365	171
55 . . . . .	131	66	65	12	39	38	42	34	26	48	23
55.0 . . . . .	30	14	15	*	*9	*8	12	*8	*7	*7	*8
55.03 . . . . .	18	9	*8	*	*	*	*9	*	*	*5	*
55.2 . . . . .	27	15	13	*	12	*9	*	*7	*	13	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
55.23	24	14	11	*	10	*7	*	*6	*	11	*
55.5	43	20	23	*	9	13	19	10	9	17	*7
55.51	41	19	21	*	*8	13	18	10	9	16	*6
55.6	10	*7	*	*	*	*	*	*	*	*	*
55.69	10	*7	*	*	*	*	*	*	*	*	*
55.8	*8	*	*	*	*	*	*	*	*	*	*
55.87	*7	*	*	*	*	*	*	*	*	*	*
55.9	*7	*	*	*	*	*	*	*	*	*	*
56	130	71	59	10	45	42	33	35	27	50	18
56.0	58	37	21	*	25	22	10	12	13	23	9
56.3	35	21	14	*	13	13	9	10	*5	15	*
56.31	31	19	12	*	12	13	*6	*8	*	14	*
56.5	11	*5	*6	*	*	*	*7	*	*	*6	*
56.51	10	*5	*5	*	*	*	*6	*	*	*5	*
56.7	13	*	*9	*7	*	*	*	*	*	*	*
56.74	12	*	*8	*7	*	*	*	*	*	*	*
57	556	319	236	20	87	114	335	196	122	152	87
57.0	22	21	*	*	*	*6	15	*7	*6	*6	*
57.1	50	19	31	*8	*6	14	22	15	*8	20	*7
57.11	*8	*	*	*6	*	*	*	*	*	*	–
57.17	18	*6	13	*	*	*5	10	*7	*	*	*
57.18	19	*7	12	*	*	*6	*9	*	*	10	*
57.3	235	152	83	*7	38	50	141	73	59	80	23
57.32	213	139	74	*7	36	46	125	62	57	73	22
57.33	22	13	*9	*	*	*	15	11	*	*7	*
57.4	58	44	14	*	*	9	47	22	12	16	*8
57.49	58	44	14	*	*	9	47	22	12	16	*8
57.5	*6	*	*	*	*	*	*	*	*	*	*
57.59	*6	*	*	*	*	*	*	*	*	*	*
57.7	*8	*6	*	–	*	*	*6	*	*	*	*
57.71	*6	*	*	–	*	*	*	*	*	*	*
57.8	20	*	16	*	*7	*5	*6	*	*	*5	*6
57.81	*9	*	*7	*	*	*	*	*	*	*	*
57.89	*6	*	*6	*	*	*	*	*	*	*	*
57.9	151	68	83	*	32	26	91	68	28	16	38
57.91	*7	*7	–	–	*	*	*	*	*	*	*
57.94	138	56	82	*	31	24	82	65	25	13	35
58	77	57	20	*	12	15	47	18	17	34	*8
58.4	9	*	*5	*	*	*	*	*	*	*	*
58.5	15	13	*	*	*	*	11	*	*	*8	*
58.6	40	31	10	*	*6	*6	28	10	10	18	*
59	209	50	158	*	72	77	58	44	50	81	34
59.0	*6	*	*	*	*	*	*	*	*	*	*
59.5	36	–	36	–	13	17	*5	*	11	16	*6
59.7	52	*	51	–	14	22	15	*7	10	22	13
59.79	50	*	49	–	13	22	15	*7	9	21	13
59.8	98	46	52	*	39	27	30	26	25	35	12
60–64	351	351	–	28	17	78	228	88	84	121	58
60	268	268	–	–	*	63	204	60	65	93	50
60.0	*6	*6	–	–	*	*	*	*	*	*	*
60.1	16	16	–	–	*	*	13	*	*	*6	*
60.11	16	16	–	–	*	*	13	*	*	*6	*
60.2	188	188	–	–	*	27	160	42	45	67	35
60.5	43	43	–	–	*	26	17	*8	11	14	11
60.9	*5	*5	–	–	–	*	*	*	*	*	*
61	*8	*8	–	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.  
– Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
62	25	25	—	*8	*6	*	*9	13	*	*5	*
62.3	*7	*7	—	*	*	*	*	*	*	*	*
62.4	*7	*7	—	—	*	*	*6	*	*	*	*
62.41	*6	*6	—	—	*	*	*6	*	*	*	*
62.5	*7	*7	—	*	*	*	—	*	*	*	*
63	14	14	—	*	*5	*	*	*	*	*6	*
63.1	*5	*5	—	*	*	*	*	*	*	*	*
64	37	37	—	18	*	*7	10	9	9	14	*
64.0	21	21	—	17	*	*	*	*6	*7	*7	*
64.9	13	13	—	*	*	*	*6	*	*	*6	*
64.97	*7	*7	—	—	*	*	*	*	*	*	*
65-71	2,093	—	2,093	*6	1,323	513	250	386	426	898	384
65	555	—	555	*	308	185	60	102	125	228	101
65.1	*6	—	*6	*	*	*	*	*	*	*	*
65.12	*6	—	*6	*	*	*	*	*	*	*	*
65.2	57	—	57	*	49	*	*	16	13	18	11
65.29	54	—	54	*	47	*	*	14	12	17	11
65.3	21	—	21	*	14	*	*	*	*6	*9	*
65.4	69	—	69	*	51	12	*	13	18	25	13
65.5	*7	—	*7	*	*	*	*	*	*	*	*
65.51	*6	—	*6	*	*	*	*	*	*	*	*
65.6	349	—	349	*	147	157	45	53	75	156	65
65.61	334	—	334	*	135	153	45	50	71	151	62
65.62	15	—	15	—	11	*	*	*	*	*	*
65.8	24	—	24	—	20	*	*	*7	*	*9	*
65.9	15	—	15	*	13	*	*	*	*	*	*
65.91	15	—	15	*	13	*	—	*	*	*	*
66	416	—	416	*	408	*5	*	65	78	197	76
66.0	9	—	9	—	*9	*	—	*	*	*	*
66.02	*7	—	*7	—	*7	—	—	*	*	*	*
66.2	25	—	25	—	25	—	—	*	*	15	*
66.22	9	—	9	—	9	—	—	*	*	*5	*
66.29	15	—	15	—	15	—	—	*	*	9	*
66.3	302	—	302	—	301	*	*	37	57	148	59
66.32	175	—	175	—	175	*	—	23	32	80	40
66.39	126	—	126	—	126	*	*	14	25	68	20
66.4	*7	—	*7	*	*5	*	*	*	*	*	*
66.5	10	—	10	—	*8	*	*	*	*	*5	*
66.51	*8	—	*8	—	*6	*	*	*	*	*	*
66.6	42	—	42	*	40	*	*	12	*8	15	*7
66.61	*8	—	*8	*	*7	*	*	*	*	*	*
66.62	27	—	27	—	27	—	—	*7	*	11	*5
66.7	10	—	10	—	10	—	—	*	*	*	*
66.79	*6	—	*6	—	*6	—	—	*	*	*	*
66.8	*6	—	*6	—	*6	—	—	*	*	*	*
67	28	—	28	—	19	*7	*	*6	*	14	*
67.3	9	—	9	—	*7	*	*	*	*	*	*
67.39	*7	—	*7	—	*5	*	*	*	*	*	*
67.5	*6	—	*6	—	*6	*	—	*	*	*	*
68	643	—	643	*	369	200	73	108	130	285	120
68.1	20	—	20	*	11	*	*	*7	*	*8	*
68.12	11	—	11	*	*7	*	*	*5	*	*	*
68.16	*7	—	*7	—	*	*	*	*	*	*	*
68.2	38	—	38	—	33	*	*	9	*	16	*8
68.29	38	—	38	—	33	*	*	9	*	16	*7
68.3	*8	—	*8	—	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
68.4 . . . . .	377	–	377	–	217	124	36	57	85	167	67
68.5 . . . . .	189	–	189	–	100	60	29	28	34	88	38
69 . . . . .	156	–	156	*	133	14	*9	51	22	60	23
69.0 . . . . .	107	–	107	*	89	12	*6	36	16	42	12
69.02 . . . . .	55	–	55	*	55	*	–	19	*6	22	*9
69.09 . . . . .	49	–	49	–	31	11	*6	16	10	19	*
69.1 . . . . .	*6	–	*6	–	*5	*	*	*	*	*	*
69.19 . . . . .	*6	–	*6	–	*5	*	*	*	*	*	*
69.2 . . . . .	*5	–	*5	–	*	*	*	*	*	*	*
69.5 . . . . .	22	–	22	*	21	*	*	11	*	*5	*
69.52 . . . . .	17	–	17	*	16	–	–	*8	*	*	*
69.9 . . . . .	13	–	13	–	11	*	*	*	*	*7	*
69.96 . . . . .	*5	–	*5	–	*5	–	–	*	*	*	*
70 . . . . .	264	–	264	*	69	95	98	46	59	106	53
70.1 . . . . .	9	–	9	*	*5	*	*	*	*	*	*
70.2 . . . . .	*7	–	*7	*	*	*	*	*	*	*	*
70.3 . . . . .	*8	–	*8	*	*6	*	*	*	*	*	*
70.5 . . . . .	158	–	158	–	32	64	63	28	36	62	33
70.50 . . . . .	85	–	85	–	15	34	35	15	13	37	19
70.51 . . . . .	40	–	40	–	*7	16	16	*8	11	14	*6
70.52 . . . . .	34	–	34	–	10	13	11	*	11	12	*7
70.7 . . . . .	31	–	31	*	11	9	10	*	*	15	*8
70.73 . . . . .	*5	–	*5	*	*	*	–	*	*	*	*
70.77 . . . . .	18	–	18	*	*	*6	9	*	*	*9	*
70.9 . . . . .	43	–	43	*	*8	17	18	*6	12	16	*9
70.92 . . . . .	41	–	41	–	*8	17	16	*6	12	16	*7
71 . . . . .	30	–	30	*	17	*6	*6	*8	*8	*8	*6
71.3 . . . . .	10	–	10	–	*8	*	*	*	*	*	*
71.7 . . . . .	9	–	9	*	*	*	*	*	*	*	*
72–75 . . . . .	6,365	–	6,365	16	6,345	*	–	1,184	1,505	2,001	1,675
72 . . . . .	463	–	463	*	463	*	–	61	103	161	139
72.0 . . . . .	18	–	18	–	18	–	–	*	*	*8	*6
72.1 . . . . .	133	–	133	*	133	*	–	14	35	62	22
72.2 . . . . .	14	–	14	*	14	–	–	*	*	*8	*
72.21 . . . . .	11	–	11	*	11	–	–	*	*	*7	*
72.4 . . . . .	*8	–	*8	*	*8	–	–	*	*	*	*
72.5 . . . . .	*7	–	*7	–	*7	–	–	*	*	*	*
72.7 . . . . .	279	–	279	*	279	*	–	40	59	77	103
72.71 . . . . .	204	–	204	*	204	*	–	35	43	54	72
72.79 . . . . .	75	–	75	*	75	*	–	*5	16	23	31
73 . . . . .	3,074	–	3,074	9	3,063	*	–	562	708	1,041	763
73.0 . . . . .	752	–	752	*	751	*	–	115	210	195	232
73.01 . . . . .	101	–	101	*	101	–	–	12	34	23	33
73.09 . . . . .	651	–	651	*	650	*	–	103	177	173	199
73.1 . . . . .	*6	–	*6	*	*6	–	–	*	*	*	*
73.4 . . . . .	374	–	374	*	373	*	–	58	89	106	122
73.5 . . . . .	869	–	869	*	866	*	–	171	158	349	190
73.51 . . . . .	10	–	10	–	10	–	–	*	*	*	*
73.59 . . . . .	859	–	859	*	856	*	–	171	155	347	186
73.6 . . . . .	1,062	–	1,062	*	1,057	*	–	216	247	387	213
74 . . . . .	786	–	786	*	784	*	–	120	168	321	176
74.0 . . . . .	10	–	10	*	10	–	–	*	*	*5	*
74.1 . . . . .	756	–	756	*	755	*	–	119	162	307	169
74.4 . . . . .	*5	–	*5	*	*5	–	–	*	*	*	*
74.9 . . . . .	12	–	12	–	12	–	–	–	*	*	*6
74.99 . . . . .	12	–	12	–	12	–	–	–	*	*	*6

\*Figure does not meet standard of reliability or precision.  
–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
75	2,043	—	2,043	*6	2,036	*	—	441	527	478	598
75.1	24	—	24	—	24	*	—	*	*7	*5	*9
75.3	992	—	992	*	990	*	—	250	277	159	306
75.32	314	—	314	*	313	—	—	63	101	57	92
75.34	621	—	621	*	619	*	—	174	156	91	200
75.35	57	—	57	*	56	*	—	11	20	12	14
75.4	37	—	37	*	36	—	—	*8	11	12	*6
75.5	*9	—	*9	—	*9	—	—	*	*	*	*
75.51	*8	—	*8	—	*8	—	—	*	*	*	*
75.6	955	—	955	*	952	*	—	177	220	287	272
75.61	34	—	34	*	34	—	—	*6	9	11	*8
75.62	128	—	128	*	128	*	—	25	27	39	38
75.69	793	—	793	*	790	*	—	146	184	237	226
75.7	20	—	20	*	20	—	—	*	*6	10	*
76-84	3,077	1,543	1,534	146	1,009	791	1,131	763	664	1,082	567
76	98	57	41	*6	70	13	*9	21	21	33	23
76.6	26	*9	17	*	22	*	*	*7	*	10	*
76.64	*7	*	*5	*	*6	*	—	*	*	*	*
76.65	*8	*	*	*	*6	*	—	*	*	*	*
76.7	47	36	11	*	36	*6	*	*8	12	14	14
76.72	11	*8	*	*	*8	*	*	*	*	*	*
76.76	18	13	*	*	14	*	*	*	*	*6	*5
76.79	*8	*6	*	—	*6	*	*	*	*	*	*
76.9	12	*7	*5	*	*8	*	*	*	*	*	*
76.91	*5	*	*	*	*	*	*	*	*	*	*
77	289	150	139	20	102	92	75	78	56	99	56
77.1	*5	*	*	*	*	*	*	*	*	*	*
77.2	*5	*	*	*	*	*	*	*	*	*	*
77.3	16	*8	*8	*7	*5	*	*	*6	*	*	*
77.4	18	*8	11	*	*	*	11	*5	*	*6	*
77.49	*7	*	*	*	*	*	*	*	*	*	*
77.5	22	*6	16	*	*7	*9	*5	12	*	*6	*
77.51	*6	*	*	—	*	*	*	*	*	*	*
77.56	*7	*	*5	—	*	*	*	*5	*	*	*
77.6	44	23	21	*	14	11	15	14	*8	13	*9
77.65	*6	*	*	*	*	*	*	*	*	*	*
77.67	*7	*	*	*	*	*	*	*	*	*	*
77.68	*9	*	*6	*	*	*	*	*5	*	*	*
77.69	12	*5	*7	*	*	*	*	*	*	*	*
77.7	108	60	48	*6	49	36	17	21	21	44	21
77.79	98	55	44	*5	45	33	16	19	20	40	20
77.8	58	35	24	*	14	24	19	14	13	20	12
77.81	22	14	*8	*	*	10	*8	*	*6	*7	*
77.88	*6	*	*	*	*	*	*	*	*	*	*
77.89	18	11	*7	*	*6	*7	*	*	*	*6	*
77.9	10	*	*7	*	*	*	*	*	*	*	*
78	185	92	94	19	73	45	48	49	34	69	34
78.0	39	19	21	*	17	10	*9	*7	*8	16	*8
78.05	*9	*	*5	*	*	*	*	*	*	*	*
78.07	14	*7	*8	*	*5	*	*	*	*	*6	*
78.09	*5	*	*	*	*	*	*	*	*	*	*
78.1	17	11	*5	*	*8	*	*	*	*	*8	*
78.17	*6	*	*	*	*	*	*	*	*	*	*
78.4	*8	*	*	*	*	*	*	*	*	*	*
78.5	52	23	29	*6	16	9	20	14	*9	18	11
78.55	25	9	16	*	*	*	16	*9	*5	*5	*6
78.57	12	*6	*7	*	*8	*	*	*	*	*6	*

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).



**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
78.59	*8	*	*	*	*	*	*	*	*	*	*
78.6	62	30	32	*	26	17	15	19	11	22	10
78.65	22	11	12	*	*7	*	11	*7	*	*8	*
78.67	14	*8	*6	*	*6	*6	*	*	*	*6	*
78.69	14	*5	*9	*	*8	*	*	*	*	*	*
79	657	310	347	52	220	129	256	150	144	242	121
79.0	84	41	43	21	23	15	26	24	19	32	10
79.01	12	*	*8	*	*	*	*6	*	*	*	*
79.02	24	9	14	*7	*	*	9	*8	*6	*7	*
79.05	12	*7	*	*6	*	*	*	*	*	*	*
79.06	32	19	14	*6	11	*8	*6	*8	*6	14	*
79.1	66	32	34	*9	20	11	27	16	18	22	10
79.11	*8	*	*5	*6	*	*	*	*	*	*	*
79.12	*7	*	*	*	*	*	*	*	*	*	*
79.15	36	16	19	*	*9	*	21	*7	12	12	*
79.16	10	*	*	*	*6	*	*	*	*	*	*
79.2	*8	*6	*	*	*	*	*	*	*	*	*
79.3	414	177	238	14	138	84	178	94	88	153	79
79.31	32	11	20	*	10	*6	11	*9	*	10	*9
79.32	31	18	14	*	17	*6	*6	*7	*7	10	*7
79.33	*6	*	*	*	*	—	*	*	*	*	*
79.34	*9	*6	*	*	*	*	*	*	*	*	*
79.35	172	51	121	*	16	21	133	41	38	64	28
79.36	140	69	71	*	72	42	22	29	32	52	27
79.37	11	*7	*	*	*6	*	*	*	*	*	*
79.39	13	9	*	*	9	*	*	*	*	*	*
79.6	41	31	10	*	25	*7	*	*7	*8	18	9
79.62	*6	*	*	*	*	*	*	*	*	*	*
79.64	*6	*5	*	*	*	*	*	*	*	*	*
79.66	20	16	*	*	12	*	*	*	*	9	*
79.7	31	18	13	*	*9	*7	16	*7	*7	10	*8
79.71	*7	*5	*	—	*	*	*	*	*	*	*
79.75	18	9	*8	*	*	*	11	*	*	*	*5
79.8	*8	*	*	*	*	*	*	*	*	*	*
80	502	289	213	9	237	162	94	123	97	190	92
80.0	12	*	*7	*	*	*	*7	*	*	*6	*
80.06	*5	*	*	—	*	*	*	*	*	*	*
80.1	26	16	11	*	*9	*8	*8	*	*8	*7	*7
80.16	10	*5	*5	*	*	*	*	*	*	*	*
80.2	65	37	28	*	38	16	9	22	14	18	10
80.21	19	9	10	*	*8	*7	*	*	*6	*6	*
80.26	44	27	17	*	29	*8	*	18	*8	11	*7
80.4	16	*8	*9	*	*6	*5	*	*7	*	*5	*
80.46	9	*	*5	*	*	*	*	*5	*	*	*
80.5	273	162	111	*	132	98	43	47	54	119	53
80.50	*6	*	*	*	*	*	*	*	*	*	*
80.51	266	160	107	*	129	97	40	47	51	115	53
80.6	45	28	16	*	27	12	*5	22	*7	10	*5
80.7	20	*8	12	*	*7	*	*8	*6	*	*9	*
80.76	14	*6	*8	*	*5	*	*6	*5	*	*6	*
80.8	35	20	14	*	15	11	*9	*9	*6	13	*8
80.81	*7	*	*	*	*	*	*	*	*	*	*
80.86	17	11	*6	*	*8	*6	*	*5	*	*6	*
80.9	*7	*	*	*	*	*	*	*	*	*	*
81	944	415	529	15	203	239	486	238	228	315	162
81.0	160	90	70	*7	67	58	28	26	36	67	31
81.02	72	42	30	—	32	29	11	11	16	34	12
81.03	*7	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
81.05	10	*	*6	*	*	*	*	*	*	*	*
81.06	*8	*	*	*	*	*	*	*	*	*	*
81.07	16	10	*6	–	*5	*6	*	*	*5	*	*
81.08	37	20	17	*	14	13	*8	*6	*7	15	9
81.09	*7	*	*	*	*	*	*	*	*	*	*
81.1	15	*	11	*	*	*7	*	*	*	*6	*
81.11	*5	*	*	–	*	*	*	*	*	*	*
81.2	12	*	*7	–	*5	*	*5	*	*	*	*
81.4	86	49	37	*	60	15	*8	30	16	25	15
81.45	42	28	14	*	37	*	*	14	10	10	*9
81.47	29	14	15	*	15	*9	*5	13	*	*9	*
81.5	495	175	320	*	25	102	369	113	129	164	89
81.51	134	54	80	–	13	31	90	31	26	48	29
81.52	97	30	68	*	*	*7	89	23	28	31	16
81.53	29	9	20	–	*	*7	18	*7	*7	10	*5
81.54	216	75	141	–	*6	53	157	48	65	69	35
81.55	18	*7	11	*	*	*	14	*	*	*6	*
81.7	*5	*	*	*	*	*	*	*	*	*	*
81.8	82	46	36	*	22	37	23	25	20	25	11
81.81	10	*	*6	–	*	*	*	*	*	*	*
81.82	10	*8	*	*	*7	*	*	*	*	*	*
81.83	56	31	25	–	12	29	15	17	15	16	*8
81.9	88	45	43	*5	18	18	47	34	21	24	9
81.91	52	28	24	*	*8	*9	31	21	13	13	*
81.92	24	11	13	–	*	*6	16	10	*6	*7	*
81.96	*5	*	*	*	*	*	*	*	*	*	*
82	32	19	13	*	15	*7	*7	11	*	11	*
82.0	*7	*	*	*	*	*	*	*	*	*	*
82.01	*5	*	*	*	*	*	*	*	*	*	*
82.4	12	9	*	*	*8	*	*	*	*	*5	*
82.45	*6	*	*	*	*	*	*	*	*	*	*
83	213	121	92	22	71	65	56	62	46	67	39
83.0	17	12	*	*	*6	*	*	*	*5	*	*
83.09	*7	*5	*	*	*	*	*	*	*	*	*
83.1	24	13	11	*7	*7	*	*7	*6	*	11	*
83.12	*6	*	*	*	*	*	*	*	*	*	*
83.14	11	*6	*5	*	*5	*	*	*	*	*5	*
83.2	16	*8	*8	*	*	*	*6	*7	*	*5	*
83.21	16	*8	*8	*	*	*	*6	*7	*	*5	*
83.3	16	*8	*8	*	*	*6	*5	*5	*	*	*
83.39	12	*6	*6	*	*	*	*	*	*	*	*
83.4	16	*9	*7	*	*8	*	*	*5	*	*	*
83.45	*7	*	*	*	*	*	*	*	*	*	*
83.5	*9	*6	*	*	*	*	*	*	*	*	*
83.6	72	42	30	*	19	32	20	18	19	21	14
83.63	52	26	25	–	*7	28	17	15	14	14	10
83.64	12	11	*	*	*7	*	*	*	*	*5	*
83.65	*7	*	*	*	*	*	*	*	*	*	*
83.7	*6	*	*	*	*	*	*	*	*	*	*
83.8	29	19	10	*7	12	*7	*	*8	*	10	*7
83.85	*7	*	*	*	*	*	*	*	*	*	*
83.88	10	*6	*	*	*6	*	*	*	*	*	*
83.9	*8	*	*6	*	*	*	*	*	*	*	*
83.95	*5	*	*	*	*	*	*	*	*	*	*
84	156	90	66	*	18	38	99	29	35	56	35
84.0	11	9	*	*	*5	*	*	*	*	*	*
84.01	*8	*8	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.

–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
84.1 . . . . .	132	72	60	*	11	29	92	25	30	51	27
84.11 . . . . .	49	31	18	*	*7	14	28	*9	12	19	10
84.12 . . . . .	15	*8	*7	*	*	*	9	*	*	*	*
84.15 . . . . .	30	18	11	*	*	*6	21	*6	*6	11	*7
84.17 . . . . .	36	14	22	—	*	*	31	*6	10	16	*
84.3 . . . . .	11	*7	*	*	*	*	*6	*	*	*	*
85-86 . . . . .	1,342	588	753	92	423	384	442	359	297	468	218
85 . . . . .	249	*	244	*	64	97	86	72	45	88	43
85.1 . . . . .	31	*	30	—	*6	9	15	10	*	14	*
85.11 . . . . .	13	—	13	—	*	*	*	*	*	*8	*
85.12 . . . . .	15	—	15	—	*	*	*8	*7	*	*	*
85.2 . . . . .	43	*	42	—	*8	15	19	17	*8	10	*7
85.21 . . . . .	29	*	28	—	*6	11	12	12	*6	*7	*
85.23 . . . . .	10	—	10	—	*	*	*	*	*	*	*
85.3 . . . . .	26	*	25	*	20	*	*	9	*	9	*
85.32 . . . . .	23	*	22	*	18	*	*	*9	*	*8	*
85.4 . . . . .	103	*	102	—	14	42	48	26	21	36	21
85.41 . . . . .	11	*	11	—	*	*5	*	*	*	*6	*
85.43 . . . . .	85	*	84	—	10	34	40	22	19	28	16
85.7 . . . . .	11	—	11	—	*	*8	*	*	*	*5	*
85.8 . . . . .	*8	—	*8	—	*	*	*	*	*	*	*
85.9 . . . . .	18	*	17	*	*6	10	*	*	*	*6	*
85.95 . . . . .	*7	—	*7	—	*	*	*	*	*	*	*
86 . . . . .	1,093	584	509	91	359	287	356	287	251	380	175
86.0 . . . . .	296	142	154	25	86	91	95	80	64	107	45
86.01 . . . . .	9	*	*5	*	*	*	*	*	*	*	*
86.04 . . . . .	108	60	48	*7	38	33	29	26	24	40	19
86.05 . . . . .	15	*8	*7	*	*6	*	*	*	*	*6	*
86.06 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
86.07 . . . . .	128	56	72	11	26	42	49	36	28	48	16
86.09 . . . . .	29	12	16	*	10	9	*6	10	*7	*8	*
86.1 . . . . .	31	15	16	*	9	11	*8	11	*6	11	*
86.11 . . . . .	31	15	16	*	9	11	*8	11	*6	11	*
86.2 . . . . .	385	217	168	25	105	100	154	96	95	131	62
86.22 . . . . .	270	159	112	14	76	75	105	63	66	97	45
86.23 . . . . .	9	*6	*	*	*	*	*	*	*	*6	*
86.27 . . . . .	23	*9	15	*	*	*	19	12	*	*	*
86.28 . . . . .	79	43	36	*9	24	19	28	19	23	25	12
86.3 . . . . .	71	28	44	*7	27	21	17	23	13	25	10
86.4 . . . . .	14	*6	*8	*	*	*	*8	*6	*	*	*
86.5 . . . . .	160	98	62	16	78	27	39	37	40	53	29
86.59 . . . . .	159	98	62	16	78	27	39	37	40	53	29
86.6 . . . . .	82	53	30	12	30	18	23	20	18	31	14
86.63 . . . . .	*5	*	*	*	*	*	*	*	*	*	*
86.65 . . . . .	*7	*	*	*	*	*	*	*	*	*5	*
86.69 . . . . .	59	38	21	*7	20	14	17	16	13	20	10
86.7 . . . . .	26	16	10	*	*8	9	*8	*6	*6	*8	*6
86.72 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
86.74 . . . . .	9	*	*	*	*	*	*	*	*	*	*
86.8 . . . . .	26	*9	17	*	12	*8	*	*8	*6	*7	*
86.83 . . . . .	10	*	*8	*	*	*	*	*	*	*	*
86.89 . . . . .	10	*	*	*	*	*	*	*	*	*	*
87-99 . . . . .	12,431	6,007	6,424	906	2,691	3,184	5,651	3,942	2,439	3,571	2,479
87 . . . . .	1,247	574	673	68	296	299	584	348	218	358	323
87.0 . . . . .	526	267	258	36	108	103	278	170	91	137	129
87.03 . . . . .	519	263	256	35	106	102	276	169	89	135	127

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000-9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
87.2 . . . . .	56	26	30	*	20	16	18	9	9	31	*7
87.21 . . . . .	48	23	26	*	17	16	15	*7	*7	30	*
87.3 . . . . .	*8	*	*8	*	*	*	*	*	*	*	*
87.37 . . . . .	*6	*	*6	—	*	*	*	*	*	*	*
87.4 . . . . .	172	81	92	*7	29	46	91	33	17	42	81
87.41 . . . . .	91	45	46	*	15	26	48	29	14	28	19
87.44 . . . . .	73	34	40	*6	13	17	38	*	*	11	58
87.49 . . . . .	*8	*	*6	*	*	*	*	*	*	*	*
87.5 . . . . .	163	51	112	*	48	46	68	34	36	61	33
87.53 . . . . .	148	46	102	*	45	42	60	28	33	56	31
87.54 . . . . .	*9	*	*6	*	*	*	*	*	*	*	*
87.6 . . . . .	106	37	69	10	16	26	54	40	19	19	29
87.61 . . . . .	25	12	12	*	*	*	14	*8	*6	*5	*
87.62 . . . . .	39	12	27	*	*7	10	18	16	*6	*8	9
87.63 . . . . .	12	*	*8	*	*	*	*	*	*	*	*
87.64 . . . . .	23	*6	17	*	*	*6	14	*9	*	*	*9
87.7 . . . . .	212	110	103	11	74	58	70	59	44	67	43
87.73 . . . . .	74	40	34	*	33	20	18	21	12	20	21
87.74 . . . . .	101	52	49	*	32	31	35	28	26	38	*8
87.76 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
87.77 . . . . .	*8	*	*	*	*	*	*	*	*	*	*
87.79 . . . . .	13	*	*8	*	*	*	*6	*	*	*	*6
88 . . . . .	3,676	1,877	1,799	106	614	1,168	1,787	1,011	752	1,180	733
88.0 . . . . .	274	127	147	11	73	75	116	86	52	67	69
88.01 . . . . .	268	125	143	11	71	73	113	86	51	64	66
88.1 . . . . .	22	9	12	*	*6	*5	*8	*	*	12	*6
88.19 . . . . .	21	*9	12	*	*5	*	*8	*	*	11	*6
88.2 . . . . .	11	*6	*5	*	*	*	*	*	*	*	*6
88.3 . . . . .	95	44	51	*5	28	22	39	34	13	29	19
88.38 . . . . .	87	38	48	*	27	21	35	32	12	26	16
88.4 . . . . .	365	195	170	*6	46	110	203	87	84	150	44
88.41 . . . . .	124	63	60	*	21	36	65	26	28	57	12
88.42 . . . . .	93	54	40	*	*7	29	56	21	21	38	13
88.43 . . . . .	14	*7	*8	*	*	*	*	*	*	*	*
88.45 . . . . .	14	*5	*9	*	*	*	*7	*	*	*	*
88.47 . . . . .	19	*9	11	*	*	*7	9	*	*5	9	*
88.48 . . . . .	79	46	33	*	*5	22	52	23	18	27	10
88.49 . . . . .	15	*8	*7	*	*	*6	*7	*	*	*7	*
88.5 . . . . .	1,468	880	588	11	137	615	706	295	343	590	241
88.50 . . . . .	*7	*	*	*	*	*	*5	*	*	*	*
88.51 . . . . .	*7	*	*	*	*	*	*	*	*	*	*
88.53 . . . . .	620	369	251	*	59	265	296	124	141	254	101
88.54 . . . . .	29	18	11	*	*	*7	15	*6	*7	12	*
88.55 . . . . .	16	9	*7	—	*	*6	*9	*	*	*7	*
88.56 . . . . .	680	411	269	*	62	292	326	140	172	255	113
88.57 . . . . .	103	64	39	*	10	41	51	16	16	52	18
88.6 . . . . .	22	10	11	*	*	*7	11	*	*9	*	*
88.66 . . . . .	14	*6	*8	—	*	*	*9	*	*6	*	*
88.67 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
88.7 . . . . .	1,181	499	682	51	258	269	603	423	204	269	285
88.71 . . . . .	107	49	59	*5	*	23	76	33	22	25	27
88.72 . . . . .	554	262	291	18	58	141	337	203	95	141	115
88.73 . . . . .	10	*6	*	*	*	*	*6	*	*	*	*
88.74 . . . . .	43	16	27	*	12	12	18	10	*5	20	*8
88.75 . . . . .	77	42	35	10	21	15	31	31	*9	15	22
88.76 . . . . .	187	71	116	13	65	44	66	66	31	31	59
88.77 . . . . .	108	43	65	*	20	28	60	46	25	13	24
88.78 . . . . .	61	—	61	*	60	*	—	19	12	13	17

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
88.79	34	11	23	*	17	*5	9	13	*	9	*9
88.9	236	105	131	19	60	61	96	77	45	56	58
88.91	136	58	78	12	27	37	61	43	29	35	30
88.93	57	26	31	*	21	13	20	15	13	11	18
88.94	12	*6	*6	*	*	*	*	*	*	*	*
88.97	25	13	12	*	*5	*7	*9	13	*	*	*6
89	1,245	600	646	81	195	323	647	394	227	291	333
89.0	*7	*	*	*	*	*	*	*	*	*	*
89.1	141	69	73	31	29	24	57	63	15	37	28
89.14	115	53	61	21	21	20	54	50	11	30	24
89.17	*7	*	*	*	*	*	*	*	—	*	*
89.19	13	*8	*	*	*5	*	*	*6	*	*	*
89.2	35	14	20	*	*8	*7	19	10	12	12	*
89.22	17	9	*8	*	*	*	13	*6	*6	*	*
89.26	*8	—	*8	*	*	*	*	*	*	*	*
89.3	61	28	33	14	13	14	20	20	15	10	16
89.37	28	12	16	*	*7	9	9	*8	*9	*	*9
89.38	16	*8	*9	*5	*	*	*6	*8	*	*	*
89.39	13	*7	*6	*6	*	*	*	*	*	*	*
89.4	139	67	72	*	18	60	61	50	26	36	27
89.41	57	30	27	*	9	28	20	21	11	*9	16
89.44	81	36	45	*	*9	32	40	29	15	27	10
89.5	486	227	259	13	64	127	282	134	74	108	171
89.50	33	13	20	*	*	*8	22	21	*	*7	*
89.51	10	*	*6	*	*	*	*7	*	—	—	10
89.52	124	55	69	*6	23	25	71	15	*	33	74
89.54	307	148	159	*7	36	88	176	95	64	64	84
89.59	10	*6	*	*	*	*	*5	*	*	*	*
89.6	375	192	183	20	59	89	206	115	84	87	89
89.61	16	10	*6	*	*	*	*5	*	*	*	*
89.62	22	13	9	*	*	*	13	*6	*6	*7	*
89.64	97	56	41	*	*8	24	64	32	24	22	18
89.65	218	101	117	14	40	52	112	69	42	49	59
89.68	15	*8	*6	*	*	*	*8	*	*	*	*6
90	40	18	22	*6	*7	*9	18	*	9	21	*
90.5	20	9	11	*	*	*	10	*	*	14	*
90.52	*5	*	*	*	*	*	*	*	*	*	*
90.53	9	*	*6	*	*	*	*5	*	—	*7	*
90.59	*6	*	*	—	*	*	*	*	—	*5	—
90.8	11	*	*6	—	*	*	*6	*	*6	*	*
90.81	10	*	*6	—	*	*	*6	*	*6	*	—
91	17	*5	12	*	*6	*	9	*	*	10	*
91.3	13	*	*8	*	*	*	*8	*	*	9	*
91.33	10	*	*7	*	*	*	*7	*	—	*8	*
92	416	192	224	*9	57	122	228	141	66	108	101
92.0	144	71	72	*	21	42	78	49	20	35	40
92.02	20	9	10	*	*	*	10	*7	*	*	*6
92.03	11	*7	*	*	*	*	*	*	*	*	*
92.04	25	11	15	*	*	*7	13	*5	*	*8	9
92.05	84	44	40	*	*9	27	48	31	13	19	21
92.1	162	69	93	*	25	41	92	59	24	37	42
92.14	61	26	36	*	*7	12	40	22	10	15	14
92.15	79	33	46	*	12	25	41	26	12	17	24
92.19	11	*5	*5	*	*	*	*6	*5	*	*	*
92.2	110	52	59	*	11	39	58	34	22	35	19
92.23	10	*6	*	—	*	*	*5	*6	*	*	*
92.24	28	14	14	*	*	11	15	10	*7	*	*8

\*Figure does not meet standard of reliability or precision.  
—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD-9-CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD-9-CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD-9-CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15-44 years	45-64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
92.27	21	*	16	*	*	*7	12	*6	*	*9	*
92.29	46	24	22	*	*	17	23	11	11	19	*
93	1,507	687	820	214	237	271	785	515	326	372	293
93.0	17	*8	*8	*	*	*	9	*	*9	*	*
93.01	*7	*	*	*	*	*	*	*	*6	—	*
93.08	*8	*	*	*	*	*	*	*	*	*	*
93.1	22	*8	14	*	*	*	17	11	*8	*	*
93.12	*6	*	*	*	*	*	*5	*	*	*	—
93.2	75	26	49	*	*8	18	48	37	23	*	12
93.22	56	19	37	*	*	14	36	28	16	*	11
93.24	11	*	*7	*	*	*	*9	*5	*5	*	*
93.3	316	126	190	*	40	56	216	103	66	108	40
93.32	11	*6	*	*	*	*	*	*	*	*	*
93.36	*6	*	*	—	—	*	*	*	—	*5	*
93.38	*5	*	*	*	*	*	*	*	*	*	*
93.39	291	113	177	*	36	48	203	98	61	96	36
93.4	26	18	*8	*	11	*	*8	10	*6	*8	*
93.44	11	10	*	*	*6	*	*	*	*	*	*
93.46	11	*7	*	*	*	*	*7	*6	*	*	*
93.5	107	56	51	18	40	16	32	36	26	24	21
93.53	30	16	14	*8	12	*	*7	10	*8	*8	*
93.54	20	10	10	*	11	*	*6	*	*	*5	*6
93.57	17	12	*	*	*7	*	*	*7	*	*	*
93.59	28	11	17	*	*5	*7	14	12	*6	*	*5
93.6	*7	*	*	—	*	*	*6	*	*	*	—
93.7	33	16	18	*	*	*6	24	12	10	*7	*
93.75	32	15	18	*	*	*5	24	12	10	*7	*
93.8	220	93	127	*	44	42	131	88	56	60	16
93.81	29	13	16	*	11	*	14	*8	*7	14	*
93.83	184	76	108	*	30	36	115	78	48	44	14
93.89	*5	*	*	*	*	*	*	*	*	*	*
93.9	683	334	349	182	84	124	293	213	121	153	196
93.90	19	11	*8	*5	*	*	*7	*7	*6	*	*
93.94	287	141	146	101	34	46	106	95	30	73	89
93.96	288	137	151	59	33	57	137	64	66	63	94
93.99	77	39	38	16	12	13	35	45	16	11	*
94	761	454	306	40	480	179	62	343	203	181	33
94.0	17	9	*8	*	11	*	*	*	*	14	*
94.01	13	*7	*6	*	*9	*	*	*	*	12	—
94.1	*7	*	*	*	*	*	*	*	*	*	*
94.2	125	51	75	*6	69	28	22	81	21	13	10
94.22	*9	*	*5	*	*6	*	*	*	*	*	*
94.23	17	10	*7	*	10	*	*	15	—	*	*
94.25	71	32	40	*	46	17	*	52	11	*	*
94.27	28	*5	23	*	*7	*5	16	9	*9	*7	*
94.3	96	43	53	11	56	17	12	50	30	16	*
94.38	32	14	18	*	22	*7	*	32	—	*	—
94.39	57	25	32	*8	31	9	9	11	30	16	*
94.4	125	70	55	16	60	36	13	53	32	38	*
94.42	11	*7	*	*7	*	*	*	*	*7	*	—
94.44	91	47	45	9	50	21	11	43	26	22	*
94.45	*8	*	*	—	*	*6	—	*	—	*5	—
94.46	13	11	*	—	*	*8	*	*	—	10	—
94.6	388	275	112	*	277	94	13	154	119	94	21
94.61	17	11	*6	*	14	*	*	*9	*7	*	*
94.62	123	93	29	*	67	48	*7	43	43	25	12
94.63	54	41	13	—	32	18	*	12	23	15	*
94.64	*6	*	*	*	*6	*	—	*	*	*	—

\*Figure does not meet standard of reliability or precision.

—Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
94.65	56	35	21	*	49	*6	*	38	*6	11	*
94.66	16	9	*7	*	15	*	*	*	*	*7	–
94.67	19	13	*6	*	16	*	–	*	14	*	*
94.68	63	46	16	*	54	*8	*	36	9	16	*
94.69	34	22	12	–	25	*9	*	*5	10	18	*
95	45	17	27	*7	16	11	11	*	*	39	*
95.0	32	12	20	*	13	*9	*8	*	*	30	–
95.04	*5	*	*	*	*	*	*	*	*	*	–
95.1	*8	*	*	*	*	*	*	*	–	*7	–
96	1,064	531	533	109	204	234	518	322	231	309	202
96.0	432	224	209	35	64	107	227	137	93	118	85
96.04	340	179	160	30	48	81	181	97	72	104	65
96.05	*8	*	*	*	*	*	*	*	*	*	*
96.07	75	37	38	*	14	21	35	32	17	9	16
96.08	*5	*	*	–	*	*	*	*	*	*	*
96.2	*6	*	*	*	*	*	*	*	*	*	*
96.3	27	14	13	*	14	*5	*	*7	*6	*8	*6
96.33	21	11	10	*	13	*	*	*5	*5	*6	*
96.4	42	*	39	*	38	*	*	*	13	10	15
96.49	39	*	37	*	37	*	*	*	12	*8	15
96.5	42	24	18	*8	14	*8	12	*9	13	14	*7
96.56	11	*5	*6	*	*	*	*6	*	*	*5	*
96.59	24	14	10	*	11	*	*	*	10	*7	*
96.6	70	27	43	*6	*7	12	44	42	*6	15	*7
96.7	444	238	206	55	65	98	225	122	98	142	82
96.70	14	*6	*9	*	*	*	*9	*	*	*7	*
96.71	289	155	134	33	45	60	150	81	63	91	55
96.72	141	77	63	21	18	35	66	40	32	45	24
97	83	39	44	13	16	18	35	24	20	26	13
97.0	23	9	14	*	*	*	16	*7	*	*7	*
97.02	15	*	11	*	*	*	11	*6	*	*	*
97.05	*6	*	*	–	*	*	*	*	*	*	*
97.2	*8	*	*5	*	*	*	*	*	*	*	*
97.23	*7	*	*	*	*	*	*	*	*	*	*
97.3	*5	*	*	*	*	*	*	*	*	*	*
97.5	*8	*	*	*	*	*	*	*	*	*	*
97.6	11	*7	*	*	*	*	*5	*	*	*	*
97.62	*8	*	*	*	*	*	*	*	*	*	*
97.8	19	11	*9	*7	*	*	*	*6	*6	*	*
97.88	*5	*	*	*	*	–	*	*	*	*	*
97.89	9	*	*	*	*	*	*	*	*	*	*
98	26	16	9	*6	*8	*	*6	*8	*5	*9	*
98.0	9	*7	*	*	*	*	*	*	*	*	*
98.02	*8	*6	*	*	*	*	*	*	*	*	*
98.5	*8	*5	*	–	*	*	*	*	*	*	*
98.51	*8	*5	*	–	*	*	*	*	*	*	*
99	2,305	997	1,308	245	555	544	961	827	376	666	435
99.0	716	301	415	36	109	170	402	212	101	261	142
99.02	38	16	22	*	*5	11	20	12	*8	10	*7
99.03	27	12	14	*	*	*7	16	*	*	16	*6
99.04	568	224	344	22	84	131	332	170	80	206	112
99.05	35	19	15	*	*8	*9	13	12	*5	11	*7
99.06	*7	*6	*	*	*	*	*	*	*	*	*
99.07	37	21	16	*	*6	10	19	*9	*	16	*9
99.1	310	113	197	30	116	59	106	142	52	58	59
99.11	64	*	64	*	64	*	*	14	12	21	17
99.14	*5	*	*	*	*	*	*	*	*	*	*

\*Figure does not meet standard of reliability or precision.  
–Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).

**Table 4. Number of all-listed procedures for discharges from short-stay hospitals, by ICD–9–CM code, sex and age of patient, and geographic region of hospital: United States, 1995—Con.**

[Discharges of inpatients from non-Federal hospitals. Excludes newborn infants. Code numbers are from the *International Classification of Diseases, 9th Revision, Clinical Modification* (ICD–9–CM). Totals include data for categories not listed in table: see [Methods section](#)]

ICD–9–CM code	Total	Sex		Age				Region			
		Male	Female	Under 15 years	15–44 years	45–64 years	65 years and over	Northeast	Midwest	South	West
Number of procedures in thousands											
99.15 . . . . .	89	39	49	18	18	14	38	33	19	14	23
99.17 . . . . .	47	21	26	*	11	14	21	23	*5	11	*7
99.18 . . . . .	37	16	20	*7	9	*6	14	28	*6	*	*
99.19 . . . . .	66	33	33	*	10	24	31	41	10	*6	10
99.2 . . . . .	1,056	459	597	133	300	278	345	415	177	283	181
99.21 . . . . .	361	153	208	60	132	61	109	162	46	97	56
99.23 . . . . .	65	28	37	10	15	15	25	34	13	12	*6
99.24 . . . . .	*5	*	*	–	*	*	*	*	–	*	*
99.25 . . . . .	330	141	189	32	57	125	116	109	54	120	47
99.29 . . . . .	291	134	157	31	91	75	94	107	65	48	71
99.4 . . . . .	11	*	11	*	11	*	–	*	*	*	*
99.47 . . . . .	9	–	9	–	9	–	–	*	*	*	*
99.5 . . . . .	9	*	*	*6	*	*	*	*	*	*	*
99.55 . . . . .	*6	*	*	*5	*	*	*	*	*	*	*
99.6 . . . . .	140	80	60	*	*9	33	97	42	30	36	31
99.60 . . . . .	42	24	18	*	*	*8	29	11	11	10	10
99.61 . . . . .	23	14	9	*	*	*6	16	*6	*	*5	*7
99.62 . . . . .	66	37	29	*	*	17	45	23	13	18	12
99.69 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
99.7 . . . . .	11	*6	*	*	*	*	*	*	*	*	*
99.71 . . . . .	*6	*	*	*	*	*	*	*	*	*	*
99.8 . . . . .	41	26	15	37	*	*	*	*6	*7	14	14
99.83 . . . . .	35	23	12	35	*	*	–	*	*6	12	13
99.9 . . . . .	*7	*	*	*	*	*	*	*	*	*	*

\* Figure does not meet standard of reliability or precision.

– Quantity zero.

NOTE: Estimates of 5,000–9,000 are to be used with caution; see [Methods section](#).



# Appendix I

## Technical Notes on Methods

### Reliability of Estimates

The standard error is primarily a measure of sampling variability that occurs by chance because only a sample rather than the entire universe is surveyed. The standard error, as calculated for the National Hospital Discharge Survey (NHDS), also reflects part of the variation that arises in the measurement process, but does not include estimates of any systematic bias.

The relative standard error (RSE) of an estimate is defined as the standard error divided by the estimate. The resulting value can be multiplied by 100 to express the relative standard error as a percent of the estimate. To obtain the RSE of an estimate, use one of the appropriate formulas described below.

Estimates of sampling variability were calculated with SUDAAN software. A description of the software and the approach it uses has been published (28).

*RSE for aggregate estimates*—The parameters for relative standard error curves for the 1995 NHDS statistics are presented in table I. The relative standard error or RSE of an estimate  $X$ , expressed as a percent of  $X$ , may be estimated from the formula:

$$RSE(X) = 100 \sqrt{a+b/X}$$

where  $a$  and  $b$  are as defined in table I.

*RSE for estimates of percents*—The approximate RSE for a percent  $p$  expressed as a percent of  $p$  may be calculated directly using the formula:

$$RSE(p) = 100 \sqrt{b(1-p)/(pX)}$$

where  $100p$  is the percent of interest,  $X$  is the denominator of the percent, and  $b$  is the parameter  $b$  in the formula for approximating the RSE( $X$ ). The values for  $b$  are given in table I.

This approximation is valid if the relative standard error of the denominator is less than 5 percent or the relative standard errors of the numerator and denominator are both less than 10 percent (29,30).

*RSE where the numerator is not a subclass of the denominator*—If the denominator of the rate is an estimate produced by the U.S. Bureau of the

Census for the total U.S. population or one or more of the age-sex-race groups of the total population, then the approximate relative standard error of the rate is equivalent to the approximate relative standard error of the numerator that can be obtained from table I.

If the numerator  $X$  and the denominator  $Y$  are both estimated from the NHDS, then the approximate relative standard error of the ratio  $X/Y$ , expressed as a percent of  $X/Y$ , is calculated using the formula:

$$RSE(X/Y) = 100 \sqrt{[RSE(X)]^2 + [RSE(Y)]^2}$$

This approximation is valid if the relative standard error of the denominator is less than 5 percent or the relative standard errors of the numerator and denominator are both less than 10 percent (29,30).

### Population Estimates

The population estimates (table II) used in computing rates are for the U.S. civilian population, including institutionalized persons, on July 1, 1995. These estimates are provided by the U.S. Bureau of the Census and are consistent with the population estimates published in *Current Population Reports*, Series P-25.

**Table I. Estimated parameters for approximate relative standard error equations for National Hospital Discharge Survey statistics by selected characteristics: United States, 1995**

Characteristic	Number of discharges or first-listed diagnoses		Number of all-listed diagnoses		Number of days of care		Number of procedures	
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>
Total	0.001470	383.006	0.001830	416.120	0.004678	810.910	0.002292	338.540
Sex								
Male	0.001760	337.999	0.002143	394.996	0.003051	1216.430	0.002633	350.456
Female	0.001318	375.570	0.001880	352.920	0.005842	741.020	0.002195	342.400
Age								
Under 15 years	0.017012	204.502	0.013717	274.540	0.019964	576.270	0.020369	261.921
15–44 years	0.001562	344.452	0.002528	330.180	0.003015	1185.440	0.003580	355.115
45–64 years	0.001346	346.329	0.001584	370.660	0.002950	1225.550	0.002882	319.558
65 years and over	0.001742	362.499	0.001988	404.887	0.002789	1852.770	0.002971	315.118
Region								
Northeast	0.005335	242.463	0.005105	255.647	0.009606	529.150	0.006875	262.270
Midwest	0.007528	206.001	0.007693	192.250	0.009445	652.660	0.009340	178.460
South	0.005192	391.875	0.006951	428.100	0.005597	1333.620	0.004027	330.110
West	0.004464	356.516	0.005627	384.339	0.008591	1042.910	0.007656	381.790

NOTE: The relative standard error (RSE) for an estimate ( $X$ ) can be determined from the equation  $RSE(X) = 100 (\sqrt{a+b/X})$ .

**Table II. Civilian population by sex, age, and geographic region: United States, 1995**

[Population estimates consistent with Series P-25, *Current Population Reports*, U.S. Bureau of the Census]

Sex, age, and geographic region	Population in thousands
Total	261,407
Sex	
Male	127,134
Female	134,273
Age	
Under 15 years	57,726
15-44 years	117,976
45-64 years	52,173
65 years and over	33,532
Geographic region	
Northeast	51,395
Midwest	61,698
South	91,143
West	57,170

## Appendix II

### Conversion Table of Changes in ICD-9-CM Codes

Although the *International Classification of Diseases, 9th Revision, Clinical Modification* has been used for coding NHDS data since 1979, the classification undergoes annual updating. Assignment of new diagnostic and procedure codes, fourth and fifth digit expansion of codes, as well as code deletions, are contained in addenda developed by the ICD-9-CM Coordination and Maintenance Committee and approved by the Director of NCHS and the Administrator of the Health Care Financing Administration. Addenda to the

ICD-9-CM become effective October 1 of the affected year and have been released annually since 1986.

The data in this report were coded using the addenda for October 1, 1986-94. When data were reported using code changes introduced in the October 1, 1995, addenda, the new codes were converted to their previous code assignments so that the annual estimates would be based on consistent coding.

Tables III and IV show the new ICD-9-CM codes from the 1986-94 addenda, the date they were introduced, and the code to which the diagnosis or procedure had been assigned prior to the development of the new code. Specific titles and more detailed information about the coding scheme can be found in appropriate volumes of the ICD-9-CM (27).

**Table III. Diagnosis codes**

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
008.00-008.09	1992	008.0	088.81, 088.89	1989	088.8
008.43-008.47	1992	008.49	088.82	1993	088.89
008.61-008.69	1992	008.6	099.40-099.49	1992	099.4
041.00-041.09	1992	041.0	099.50-099.59	1992	078.89
041.10-041.19	1992	041.1	112.84-112.85	1992	112.89
041.81-041.89	1992	041.8	114.4-114.5	1993	114.3
042	1994	042.0-042.2, 042.9, 043.0-043.3, 043.9, 044.0, 044.9	176.0-176.9	1991	173.0-173.9
042.0-042.9	1986	279.19	203.00	1991	203.0
043.0-043.9	1986	279.19	203.01	1991	V10.79
044.0-044.9	1986	279.19	203.10	1991	203.1
070.20-070.21	1991	070.2	203.11	1991	V10.79
070.22	1994	070.20	203.80	1991	203.8
070.23	1994	070.21	203.81	1991	V10.79
070.30-070.31	1991	070.3	204.00	1991	204.0
070.32	1994	070.30	204.01	1991	V10.61
070.33	1994	070.31	204.10	1991	204.1
070.41-070.49	1991	070.4	204.11	1991	V10.61
070.44	1994	070.41	204.20	1991	204.2
070.51-070.59	1991	070.5	204.21	1991	V10.61
070.54	1994	070.51	204.80	1991	204.8
077.98-077.99	1993	077.9	204.81	1991	V10.61
078.10-078.11, 078.19	1993	078.1	204.90	1991	204.9
078.88	1993	078.89	204.91	1991	V10.61
079.4	1993	079.8	205.00	1991	205.0
079.50-079.53, 079.59	1993	079.8	205.01	1991	V10.62
079.88-079.89	1993	079.8	205.10	1991	205.1
079.98-079.99	1993	079.9	205.11	1991	V10.62
			205.20	1991	205.2
			205.21	1991	V10.62

Table III. Diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
205.30	1991	205.3	342.00–342.02	1994	342.0
205.31	1991	V10.62	342.10–342.12	1994	342.1
205.80	1991	205.8	342.80–342.82	1994	342.9
205.81	1991	V10.62	342.90–342.92	1994	342.9
205.90	1991	205.9	344.00–344.04, 344.09	1994	344.0
205.91	1991	V10.62	344.30–344.32	1994	344.3
206.00	1991	206.0	344.40–344.42	1994	344.4
206.01	1991	V10.63	344.81–344.89	1993	344.8
206.10	1991	206.1	345.00–345.01	1989	345.0
206.11	1991	V10.63	345.10–345.11	1989	345.1
206.20	1991	206.2	345.40–345.41	1989	345.4
206.21	1991	V10.63	345.50–345.51	1989	345.5
206.80	1991	206.8	345.60–345.61	1989	345.6
206.81	1991	V10.63	345.70–345.71	1989	345.7
206.90	1991	206.9	345.80–345.81	1989	345.8
206.91	1991	V10.63	345.90–345.91	1989	345.9
207.00	1991	207.0	346.00–346.01	1992	346.0
207.01	1991	V10.69	346.10–346.11	1992	346.1
207.10	1991	207.1	346.20–346.21	1992	346.2
207.11	1991	V10.69	346.80–346.81	1992	346.8
207.20	1991	207.2	346.90–346.91	1992	346.9
207.21	1991	V10.69	355.71	1993	354.4
207.80	1991	207.8	355.79	1993	355.7
207.81	1991	V10.69	371.82	1992	371.89
208.00	1991	208.0	374.87	1990	374.89
208.01	1991	V10.60	403.00–403.01	1989	403.0
208.10	1991	208.1	403.10–403.11	1989	403.1
208.11	1991	V10.60	403.90–403.91	1989	403.9
208.20	1991	208.2	404.00–404.03	1989	404.0
208.21	1991	V10.60	404.10–404.13	1989	404.1
208.80	1991	208.8	404.90–404.93	1989	404.9
208.81	1991	V10.60	410.00–410.02	1989	410.0
208.90	1991	208.9	410.10–410.12	1989	410.1
208.91	1991	V10.60	410.20–410.22	1989	410.2
237.70–237.72	1990	237.7	410.30–410.32	1989	410.3
250.02	1993	250.90	410.40–410.42	1989	410.4
250.03	1993	250.91	410.50–410.52	1989	410.5
250.12	1993	250.10	410.60–410.62	1989	410.6
250.13	1993	250.11	410.70–410.72	1989	410.7
250.22	1993	250.20	410.80–410.82	1989	410.8
250.23	1993	250.21	410.90–410.92	1989	410.9
250.32	1993	250.30	411.81	1989	410.9
250.33	1993	250.31	411.89	1989	411.8
250.42	1993	250.40	414.00–414.01	1994	414.0
250.43	1993	250.41	414.02–414.03	1994	996.03
250.52	1993	250.50	429.71	1989	410.0–410.9
250.53	1993	250.51	429.79	1989	410.0–410.9
250.62	1993	250.60	433.00–433.01	1993	433.0
250.63	1993	250.61	433.10–433.11	1993	433.1
250.72	1993	250.70	433.20–433.21	1993	433.2
250.73	1993	250.71	433.30–433.31	1993	433.3
250.82	1993	250.80	433.80–433.81	1993	433.8
250.83	1993	250.81	433.90–433.91	1993	433.9
250.92	1993	250.90	434.00–434.01	1993	434.0
250.93	1993	250.91	434.10–434.11	1993	434.1
283.10–283.11, 283.19	1993	283.1	434.90–434.91	1993	434.9
305.1	1994	305.10, 305.11, 305.12, 305.13 (delete code)	437.7	1992	780.9
312.81–312.82, 381.89	1994	312.8	440.20–440.22	1992	440.2
320.81–320.89	1992	320.8	440.23	1993	440.20 & 707.1 or 707.8 or 707.9
333.92–333.93	1994	333.99			
337.20–337.22, 337.29	1993	337.9			

Table III. Diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
440.24 . . . . .	1993	440.20 & 785.4	657.0 . . . . .	1991	657
440.29 . . . . .	1993	440.20	659.60, 659.61, 659.63 . . .	1992	659.80, 659.81, 659.83
440.30–440.32 . . . . .	1994	996.1	665.10, 665.11 . . . . .	1992	665.10, 665.11, 665.12, 665.14 <sup>1</sup>
441.00–441.03 . . . . .	1994	441.0	670.0 . . . . .	1991	670
441.6 . . . . .	1993	441.1 & 441.3	672.0 . . . . .	1991	672
441.7 . . . . .	1993	441.2 & 441.4	677 . . . . .	1994	None <sup>2</sup>
446.20–446.21, 446.29 . . . . .	1990	446.2	692.72–692.74 . . . . .	1992	692.79
451.82–451.84 . . . . .	1993	451.89	692.82–692.83 . . . . .	1992	692.89
482.30–482.39 . . . . .	1992	482.3	702.0–702.8 . . . . .	1991	702
482.81–482.89 . . . . .	1992	482.8	702.11, 702.19 . . . . .	1994	702.1
483.0 . . . . .	1992	483	704.02 . . . . .	1993	704.09
483.8 . . . . .	1992	483	709.00–709.01, 709.09 . . . . .	1994	709.0
491.20–491.21 . . . . .	1991	491.2	710.5 . . . . .	1992	288.3, 729.1
493.20 . . . . .	1989	493.90	733.10–733.16, 733.19 . . . . .	1993	733.1
493.21 . . . . .	1989	493.91	738.10–738.19 . . . . .	1992	738.1
512.1 . . . . .	1994	997.3	747.60–747.64, 747.69 . . . . .	1993	747.6
518.81 . . . . .	1987	799.1	747.82 . . . . .	1993	747.89
518.82–518.89 . . . . .	1987	518.8	753.10–753.17, 753.19 . . . . .	1990	753.1
524.00–524.09 . . . . .	1992	524.0	759.81–759.89 . . . . .	1989	759.8
524.10–524.19 . . . . .	1992	524.1	759.83 . . . . .	1994	759.89
524.60–524.69 . . . . .	1991	524.6	760.75 . . . . .	1991	760.79
524.70–524.79 . . . . .	1992	524.8	760.76 . . . . .	1994	760.79
530.10–530.11, 530.19 . . . . .	1993	530.1	764.00–764.09 . . . . .	1988	764.0
530.81 . . . . .	1993	530.1	764.10–764.19 . . . . .	1988	764.1
530.82–530.84, 530.89 . . . . .	1993	530.8	764.20–764.29 . . . . .	1988	764.2
535.00–535.01 . . . . .	1991	535.0	764.90–764.99 . . . . .	1988	764.9
535.10–535.11 . . . . .	1991	535.1	765.00–765.09 . . . . .	1988	765.0
535.20–535.21 . . . . .	1991	535.2	765.10–765.19 . . . . .	1988	765.1
535.30–535.31 . . . . .	1991	535.3	780.01–780.09 . . . . .	1992	780.0
535.40–535.41 . . . . .	1991	535.4	780.03 . . . . .	1993	780.01
535.50–535.51 . . . . .	1991	535.5	780.57 . . . . .	1992	780.51, 780.53
535.60–535.61 . . . . .	1991	535.6	781.8 . . . . .	1994	781.9
536.3 . . . . .	1994	536.8	787.01–787.03 . . . . .	1994	787.0
537.82 . . . . .	1990	537.89	788.20–788.21, 788.29 . . . . .	1993	788.2
537.83 . . . . .	1991	537.82	788.30–788.39 . . . . .	1992	788.3
556.0–556.6 . . . . .	1994	556	788.41–788.43 . . . . .	1993	788.4
556.8–556.9 . . . . .	1994	556	788.61–788.62, 788.69 . . . . .	1993	788.6
562.02 . . . . .	1991	562.00	789.00–789.07, 789.09 . . . . .	1994	789.0
562.03 . . . . .	1991	562.01	789.30–789.37, 789.39 . . . . .	1994	789.3
562.12 . . . . .	1991	562.10	789.40–789.47, 789.49 . . . . .	1994	789.4
562.13 . . . . .	1991	562.11	789.60–789.67, 789.69 . . . . .	1994	789.6
569.84 . . . . .	1990	557.1	790.91 . . . . .	1993	790.9
569.85 . . . . .	1991	569.84	790.92 . . . . .	1993	286.9
593.70–593.73 . . . . .	1994	593.7	790.93, 790.99 . . . . .	1993	790.9
596.51–596.53 . . . . .	1992	596.5	795.71 . . . . .	1994	795.8 (delete code)
596.54 . . . . .	1992	344.61	795.79 . . . . .	1994	795.7
596.55–596.59 . . . . .	1992	596.5	795.8 . . . . .	1986	795.7
599.81–599.89 . . . . .	1992	599.8	864.05 . . . . .	1992	864.09
645.0 . . . . .	1991	645	864.15 . . . . .	1992	864.19
651.30–651.31, 651.33 . . . . .	1989	651.00–651.01, 651.03	909.5 . . . . .	1994	909.9
651.40–651.41, 651.43 . . . . .	1989	651.10–651.11, 651.13	925.1–925.2 . . . . .	1993	925
651.50–651.51, 651.53 . . . . .	1989	651.20–651.21, 651.23	995.60–995.69 . . . . .	1993	995.0
651.60–651.61, 651.63 . . . . .	1989	651.80–651.81, 651.83			
654.20–654.21, 654.23 . . . . .	1990	654.2, 654.9			
654.90–654.94 . . . . .	1990	654.2, 654.9			

Table III. Diagnosis codes—Con

Current code(s) assignment	Effective October 1	Previous code(s) assignment	Current code(s) assignment	Effective October 1	Previous code(s) assignment
996.04 . . . . .	1994	996.09	V35.00–V35.01 . . . . .	1989	V35.0
996.51–996.59 . . . . .	1987	996.5	V36.00–V36.01 . . . . .	1989	V36.0
996.60–996.69 . . . . .	1989	996.6	V37.00–V37.01 . . . . .	1989	V37.0
996.70–996.79 . . . . .	1989	996.7	V39.00–V39.01 . . . . .	1989	V39.0
996.80–996.89 . . . . .	1987	996.8	V43.60–V43.66, V43.69 . . .	1994	V43.6
996.85 . . . . .	1990	999.8	V45.00 . . . . .	1994	V45.89
998.81–998.82, 998.89 . . . .	1994	998.8	V45.01 . . . . .	1994	V45.0
V03.81–V03.82, V03.89 . . .	1994	V03.8	V45.02, V45.09 . . . . .	1994	V45.89
V05.3–V05.4 . . . . .	1993	V05.8	V45.51 . . . . .	1994	V45.5
V06.5–V06.6 . . . . .	1994	V06.8	V45.52, V45.59 . . . . .	1994	V45.89
V07.31, V07.39 . . . . .	1994	V07.3	V45.82 . . . . .	1994	V45.89
V07.4 . . . . .	1992	V07.8	V49.60–V49.67 . . . . .	1994	V49.5
V08 . . . . .	1994	044.9, 795.8 (delete code)	V49.70–V49.77 . . . . .	1994	V49.5
V09.0–V09.91 . . . . .	1993	None <sup>2</sup>	V50.41–V50.42, V50.49 . . .	1994	V50.8
V12.00–V12.03, V12.09 . . .	1994	V12.0	V53.31 . . . . .	1994	V53.3
V12.70–V12.72, V12.79 . . .	1994	V12.7	V53.32, V53.39 . . . . .	1994	V53.9
V13.00–V13.01, V13.09 . . .	1994	V13.0	V57.21–V57.22 . . . . .	1994	V57.2
V15.82 . . . . .	1994	305.13 (delete code)	V58.41, V58.49 . . . . .	1994	V58.4
V25.43 . . . . .	1992	V25.49	V58.81, V58.89 . . . . .	1994	V58.8
V25.5 . . . . .	1992	V25.8	V65.40–V65.45, V65.49 . . .	1994	V65.4
V29.0–V29.8 . . . . .	1992	V71.8	V69.0–V69.3 . . . . .	1994	None <sup>2</sup>
V29.9 . . . . .	1992	V71.9	V69.8–V69.9 . . . . .	1994	None <sup>2</sup>
V30.00–V30.01 . . . . .	1989	V30.0	V72.81–V72.85 . . . . .	1993	V72.8
V31.00–V31.01 . . . . .	1989	V31.0	V73.88–V73.89 . . . . .	1993	V73.8
V32.00–V32.01 . . . . .	1989	V32.0	V73.98–V73.99 . . . . .	1993	V73.9
V33.00–V33.01 . . . . .	1989	V33.0	E869.4 . . . . .	1994	E869.8
V34.00–V34.01 . . . . .	1989	V34.0			

<sup>1</sup>The title for the subcategory 665.1 has been changed, making the fifth-digit subclassification, 665.12 and 665.14 invalid.<sup>2</sup>There were no previous code assignments for this code.

Table IV. Procedures codes

Current codes(s) assignment	Effective October 1	Previous code(s) assignment	Current codes(s) assignment	Effective October 1	Previous code(s) assignment
02.96	1992	89.19	42.25	1988	42.24
03.90	1987	03.99 (insertion of catheter)	42.33	1989	42.32, 42.39
			42.33	1990	42.91
11.75	1989	11.79	43.11	1989	43.1
11.76	1989	11.62	43.19	1989	43.1, 43.2
20.96–20.98	1986	20.95	43.41	1989	43.41, 43.49
22.12	1988	22.11	44.21	1986	44.2
26.12	1988	26.11	44.22	1986	44.99
29.31	1991	83.02	44.29	1986	44.2
29.32	1991	29.3	44.43	1989	43.49, 45.32
29.33	1991	29.3	44.44	1989	38.86
29.39	1991	29.3	44.49	1989	43.0
31.45	1988	31.43–31.44	44.93–44.94	1986	44.99
31.95	1989	31.75	45.16	1988	45.14 (45.15 before 1987)
32.01	1989	32.0	45.30	1989	45.31, 45.32
32.09	1989	32.0	45.42	1988	45.41
32.28	1989	32.29	45.43	1989	45.49
33.27	1987	33.22+33.27	45.75 (Hartmann resection added)	1988	48.66 (code deleted)
33.28	1987	33.27	45.95	1987	45.93
33.29	1987	33.28–33.29	46.13	1992	46.12 (code deleted)
33.6	1990	33.5+37.5	46.32	1989	46.39
34.05	1994	34.99	46.85	1989	46.99
35.84	1988	35.82	49.31	1989	49.3
35.96	1986	35.03	49.39	1989	49.3
36.00–36.03	1986	36.0	51.10	1989	51.97
36.04	1986	39.97	51.11	1989	51.11, 51.97
36.05	1986	36.01 <sup>1</sup> , 36.02	51.14	1989	51.12
36.05	1987	36.01	51.15	1989	51.97
36.09	1986	36.0	51.22	1991	51.21 (code deleted)
36.09	1991	36.00 (code deleted)	51.23	1991	51.22
37.26–37.27	1988	37.29	51.64	1989	51.69
37.34	1988	37.33	51.84–51.88	1989	51.97
37.70 (leads only)	1987 (leads/device)	37.7	51.97	1986	52.91, 51.99, or 51.82
37.71–37.72 (leads only)	1987 (leads/device)	37.74	51.98	1986	51.99
37.73 (leads only)	1987 (leads/device)	37.73	52.13	1989	51.97, 52.91
37.74 (leads only)	1987 (leads/device)	37.76	52.14	1989	52.11
37.75 (leads only)	1987 (leads/device)	37.89	52.21	1989	52.2
37.76 (leads only)	1987 (leads/device)	37.81	52.22	1989	52.2
37.77 (leads only)	1987 (leads/device)	37.83–37.84	52.93	1989	52.93 + 52.91
37.78	1987	37.71–37.72	52.94	1989	52.09
37.79	1987	86.09	52.97	1989	52.91
37.80–37.87	1992	89.49 <sup>2</sup>	52.98	1989	52.91
37.80 (device only)	1987 (leads/device)	37.73–37.77	52.99	1989	52.93, 52.94, 52.99
37.81 (device only)	1987 (leads/device)	37.73–37.77	54.24	1987	54.23
37.82 (device only)	1987 (leads/device)	37.73–37.77	54.25	1993	54.98
37.83 (device only)	1987 (leads/device)	37.73–37.77	55.03–55.04	1986	55.02
37.85–37.87	1987	37.85	56.33–56.34	1987	56.33
37.89	1987	37.86 + 37.89	56.35	1987	45.12
37.94–37.98	1986	37.99	57.17–57.18	1989	57.21
38.22	1986	38.29	57.22	1989	57.22, 57.82
38.44 (abdominal aorta only)	1986	38.44 (entire aorta)	58.31	1990	58.3
38.45 (thoracic aorta only)	1986	38.44–38.45	58.39	1990	58.3
38.95	1989	38.93	58.93	1986	57.99
39.28	1991	39.29	59.96	1986	59.95
39.65	1988	39.61	60.95	1991	60.99
39.66	1990	39.65	64.97	1986	64.95
41.00–41.03	1988	41.0			
41.04	1994	99.79			

**Table IV. Procedures codes—Con**

Current codes(s) assignment	Effective October 1	Previous code(s) assignment	Current codes(s) assignment	Effective October 1	Previous code(s) assignment
66.01 . . . . .	1992	66.0	81.53 . . . . .	1989	81.51, 81.59, 81.61, 81.62, 81.63, 81.64
66.02 . . . . .	1992	66.73	81.54–81.55 . . . . .	1989	81.41
68.15 . . . . .	1987	68.14	81.56 . . . . .	1989	81.48
68.16 . . . . .	1987	68.13	81.57 . . . . .	1989	81.31, 81.39
68.9 . . . . .	1992	68.4	81.59 . . . . .	1989	81.39
74.3 . . . . .	1992	69.11 (code deleted)	81.72 . . . . .	1989	81.79
77.56 . . . . .	1989	77.89, 78.49, 81.18	81.73–81.74 . . . . .	1989	81.86
77.57 . . . . .	1989	77.89, 80.48, 81.18, 83.85	81.75 . . . . .	1989	81.87
77.58 . . . . .	1989	77.59, 81.18	81.79 . . . . .	1989	81.79, 81.87
78.10 . . . . .	1991	78.40	81.80 . . . . .	1989	81.81
78.11 . . . . .	1991	78.41	81.97 . . . . .	1992	81.59
78.12 . . . . .	1991	78.42	85.95 . . . . .	1987	85.99
78.13 . . . . .	1991	78.43	85.96 . . . . .	1987	85.99
78.14 . . . . .	1991	78.44	86.06 . . . . .	1987	86.09
78.15 . . . . .	1991	78.45	86.07 . . . . .	1990	86.09
78.16 . . . . .	1991	78.46	86.27 . . . . .	1986	86.22–86.23
78.17 . . . . .	1991	78.47	86.28 . . . . .	1988	86.22
78.18 . . . . .	1991	78.48	86.93 . . . . .	1987	86.89
78.19 . . . . .	1991	78.49	88.90 . . . . .	1986	88.39
78.20 . . . . .	1991	78.10, 78.20, 78.30	88.91 . . . . .	1986	89.15
78.21 . . . . .	1991	78.11, 78.31	88.92 . . . . .	1986	89.39
78.22 . . . . .	1991	78.12, 78.22, 78.32	88.93 . . . . .	1986	89.15
78.23 . . . . .	1991	78.13, 78.23, 78.33	88.94 . . . . .	1986	89.39
78.24 . . . . .	1991	78.14, 78.34	88.95 . . . . .	1986	89.29
78.25 . . . . .	1991	78.15, 78.25, 78.35	88.97 . . . . .	1989	88.99
78.27 . . . . .	1991	78.17, 78.27, 78.37	88.98 . . . . .	1989	88.90
78.28 . . . . .	1991	78.18, 78.38	88.99 . . . . .	1986	89.39
78.29 . . . . .	1991	78.11, 78.16, 78.19, 78.29, 78.3	89.10 . . . . .	1989	89.15
78.39 . . . . .	1991	78.31	89.17–89.18 . . . . .	1988	89.15
78.90 <sup>3</sup> . . . . .	1987	78.40	89.19 . . . . .	1989	89.15
78.91 <sup>3</sup> . . . . .	1987	78.41	89.50 . . . . .	1991	89.54
78.92 <sup>3</sup> . . . . .	1987	78.42	93.90 . . . . .	1988	93.92
78.93 <sup>3</sup> . . . . .	1987	78.43	94.61–94.69 . . . . .	1989	94.25
78.94 <sup>3</sup> . . . . .	1987	78.44	96.6 . . . . .	1986	96.35
78.95 <sup>3</sup> . . . . .	1987	78.45	96.70 . . . . .	1991	93.92 (code deleted)
78.96 <sup>3</sup> . . . . .	1987	78.46	96.71 . . . . .	1991	93.92 (code deleted)
78.97 <sup>3</sup> . . . . .	1987	78.47	96.72 . . . . .	1991	93.92 (code deleted)
78.98 <sup>3</sup> . . . . .	1987	78.48	97.05 . . . . .	1989	51.97
78.99 <sup>3</sup> . . . . .	1987	78.49	98.51–98.52 . . . . .	1989	59.96 (code deleted)
80.50–80.59 . . . . .	1986	80.5	98.59 . . . . .	1989	59.96 (code deleted)
81.03 . . . . .	1989	81.02	99.15 . . . . .	1986	99.29
81.04–81.05 . . . . .	1989	81.03, 81.04, 81.05	99.28 . . . . .	1994	99.25
81.06–81.07 . . . . .	1989	81.06, 81.07	99.71–99.79 <sup>4</sup> . . . . .	1988	99.07
81.08 . . . . .	1989	81.06, 81.07, 81.08	99.85 . . . . .	1987	93.35
81.09 . . . . .	1989	81.08	99.86 . . . . .	1987	93.39
81.40 . . . . .	1989	81.69	99.88 . . . . .	1988	99.83
81.51 . . . . .	1989	81.51, 81.59			
81.52 . . . . .	1989	81.61, 81.62, 81.63, 81.64			

<sup>1</sup>Before October 1986, contents of current code 36.05 would have been assigned to 36.0.

<sup>2</sup>Code 89.49 deleted; this procedure is included in the code for pacemaker insertion/replacement.

<sup>3</sup>Codes 78.90–78.99 were retitled as "Insertion of bone growth stimulator" in October 1987; the previous contents of codes 78.90–78.99 were reassigned to codes 78.40–78.49.

<sup>4</sup>Codes 99.71–99.79 were deleted in October 1987; their contents were not transferred elsewhere. In the October 1988 revision, codes 99.71–99.79 were reclassified as "Therapeutic apheresis."

# Vital and Health Statistics series descriptions

- SERIES 1. **Programs and Collection Procedures**—These reports describe the data collection programs of the National Center for Health Statistics. They include descriptions of the methods used to collect and process the data, definitions, and other material necessary for understanding the data.
- SERIES 2. **Data Evaluation and Methods Research**—These reports are studies of new statistical methods and include analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory. These studies also include experimental tests of new survey methods and comparisons of U.S. methodology with those of other countries.
- SERIES 3. **Analytical and Epidemiological Studies**—These reports present analytical or interpretive studies based on vital and health statistics. These reports carry the analyses further than the expository types of reports in the other series.
- SERIES 4. **Documents and Committee Reports**—These are final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- SERIES 5. **International Vital and Health Statistics Reports**—These reports are analytical or descriptive reports that compare U.S. vital and health statistics with those of other countries or present other international data of relevance to the health statistics system of the United States.
- SERIES 6. **Cognition and Survey Measurement**—These reports are from the National Laboratory for Collaborative Research in Cognition and Survey Measurement. They use methods of cognitive science to design, evaluate, and test survey instruments.
- SERIES 10. **Data From the National Health Interview Survey**—These reports contain statistics on illness; unintentional injuries; disability; use of hospital, medical, and other health services; and a wide range of special current health topics covering many aspects of health behaviors, health status, and health care utilization. They are based on data collected in a continuing national household interview survey.
- SERIES 11. **Data From the National Health Examination Survey, the National Health and Nutrition Examination Surveys, and the Hispanic Health and Nutrition Examination Survey**—Data from direct examination, testing, and measurement on representative samples of the civilian noninstitutionalized population provide the basis for (1) medically defined total prevalence of specific diseases or conditions in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics, and (2) analyses of trends and relationships among various measurements and between survey periods.
- SERIES 12. **Data From the Institutionalized Population Surveys**—Discontinued in 1975. Reports from these surveys are included in Series 13.
- SERIES 13. **Data From the National Health Care Survey**—These reports contain statistics on health resources and the public's use of health care resources including ambulatory, hospital, and long-term care services based on data collected directly from health care providers and provider records.
- SERIES 14. **Data on Health Resources: Manpower and Facilities**—Discontinued in 1990. Reports on the numbers, geographic distribution, and characteristics of health resources are now included in Series 13.
- SERIES 15. **Data From Special Surveys**—These reports contain statistics on health and health-related topics collected in special surveys that are not part of the continuing data systems of the National Center for Health Statistics.
- SERIES 16. **Compilations of Advance Data From Vital and Health Statistics**—Advance Data Reports provide early release of information from the National Center for Health Statistics' health and demographic surveys. They are compiled in the order in which they are published. Some of these releases may be followed by detailed reports in Series 10–13.
- SERIES 20. **Data on Mortality**—These reports contain statistics on mortality that are not included in regular, annual, or monthly reports. Special analyses by cause of death, age, other demographic variables, and geographic and trend analyses are included.
- SERIES 21. **Data on Natality, Marriage, and Divorce**—These reports contain statistics on natality, marriage, and divorce that are not included in regular, annual, or monthly reports. Special analyses by health and demographic variables and geographic and trend analyses are included.
- SERIES 22. **Data From the National Mortality and Natality Surveys**—Discontinued in 1975. Reports from these sample surveys, based on vital records, are now published in Series 20 or 21.
- SERIES 23. **Data From the National Survey of Family Growth**—These reports contain statistics on factors that affect birth rates, including contraception, infertility, cohabitation, marriage, divorce, and remarriage; adoption; use of medical care for family planning and infertility; and related maternal and infant health topics. These statistics are based on national surveys of women of childbearing age.
- SERIES 24. **Compilations of Data on Natality, Mortality, Marriage, Divorce, and Induced Terminations of Pregnancy**—These include advance reports of births, deaths, marriages, and divorces based on final data from the National Vital Statistics System that were published as supplements to the *Monthly Vital Statistics Report (MVSR)*. These reports provide highlights and summaries of detailed data subsequently published in *Vital Statistics of the United States*. Other supplements to the MVSR published here provide selected findings based on final data from the National Vital Statistics System and may be followed by detailed reports in Series 20 or 21.

For answers to questions about this report or for a list of reports published in these series, contact:

Data Dissemination Branch  
National Center for Health Statistics  
Centers for Disease Control and Prevention  
6525 Belcrest Road, Room 1064  
Hyattsville, MD 20782-2003  
(301) 436-8500  
E-mail: [nchsquery@cdc.gov](mailto:nchsquery@cdc.gov)  
Internet: <http://www.cdc.gov/nchswwww/nchshome.htm>



**DEPARTMENT OF  
HEALTH & HUMAN SERVICES**

Centers for Disease Control and Prevention  
National Center for Health Statistics  
6525 Belcrest Road  
Hyattsville, Maryland 20782-2003

---

OFFICIAL BUSINESS  
PENALTY FOR PRIVATE USE, \$300

STANDARD MAIL (B)  
POSTAGE & FEES PAID  
PHS/NCHS  
PERMIT NO. G-281