	Baseline	)									Targ
Objective	year	Baseline	1990	1991	1992	1993	1994	1995	1996	1997	200
Adolescent pregnancy											
Pregnancies (per 1,000)/1											
Female 10-14 years	1990	a/3.5		3.4	3.4	3.3	3.3	3.0			
Female 15-17 years	1990	a/80.3		79.8	77.3	76.8	75.5	71.7			
Live births (per 1,000)											
Female 10-14 years	1990	a/1.4		1.4	1.4	1.4	1.4	1.3	1.2	p/1.2	
Female 15-17 years	1990	a/37.5		38.7	37.8	37.8	37.6	36.0	33.8	p/32.6	
Abortions (per 1,000)											
Female 10-14 years	1990	a/1.5		1.4	1.5	1.4	1.3	1.2			
Female 15-17 years	1990	a/26.5		24.3	23.1	22.5	21.4	19.9			
Fetal losses (per 1,000)											
Female 10-14 years	1990	a/0.5		0.5	0.5	0.5	0.6	0.5			
Female 15-17 years	1990	a/16.2		16.8	16.4	16.5	16.5	15.9			
a. Pregnancies, black adolescents (per 1,000)/1											
Female 15-19 years	1990	a/215		215	211	205	195	178			
Female 15-17 years	1990	a/161		160	156	153	146	133			
Live births (per 1,000)											
Female 15-19 years	1990	a/113		116	112	109	105	96	91	p/90	
Female 15-17 years	1990	a/82		84	81	80	76	70	65	p/62	
Abortions (per 1,000)	1000	0,02		01	01	00	10	10	00	p/ 02	
Female 15-19 years/2	1990	a/80		77	76	75	70	63			
Female 15-17 years	1990	a/55		53	52	51	48	44			
Fetal losses (per 1,000)	1000	4/00		00	52	51	40				
Female 15-19 years/2	1990	a/22		22	22	21	21	19			
Female 15-17 years	1990	a/22		23	23	22	21	19			
b. Pregnancies, Hispanic adolescents (per 1,000)/1	1330	a/25		25	25	22	21	15			
Female 15-19 years	1990	a/156		165	168	166	167	163			
Live births (per 1,000)	1330	a/150		105	100	100	107	105			
Female 15-19 years/3	1990	a/100		107	107	107	108	107	102	p/99	
Abortions (per 1,000)	1990	a/100		107	107	107	100	107	102	p/99	
Female 15-19 years	1990	a/39		40	43	42	42	39			
Fetal losses (per 1,000)	1990	a/39		40	43	42	42	39			
· · · ·	1000	o/17		10	10	10	10	40			
Female 15-19 years	1990	a/17		18	18	18	18	18			
Unintended pregnancy (female 15-44 years)	1988	56%						49%			
a. Black female (15-44 years)	1988	78%						3/72%			
b. Hispanic female (15-44 years)	1988	55%						48%			3
Infertility											
Married couples with wives 15-44 years	1988	7.9%						7.1%			6
a. Black couples with wives 15-44 years	1988	12.1%						3/10.5%			
b. Hispanic couples with wives 15-44 years	1988	12.4%						7.0%			

5.4*	Objective Adolescents who ever engaged in sexual intercourse Adolescents 15 years	year	Baseline	1990	1991	1992	1993	1994	1995	1996	1997	Target 2000
5.4*	Adolescents 15 years											2000
5.4	Adolescents 15 years											
	•											
		1000	070/						22%			15%
	All females	1988	27%									
	In-school females			35%	36%		37%		38% 27%			
	All males	1988	33%		 44%				27% 42%			15%
	In-school males			48%			45%					
	a. All black males	1988	69%						58%			15%
	In-school non-Hispanic black males				79%		82%		77%		75%	
	Adolescents 17 years	4000	500/						<b>E40</b> (			400/
	All females	1988	50%						51%			40%
	In-school females			62%	66%		66%		67%			
	All males	1988	66%									40%
	In-school males			73%	68%		68%		65%			
	b. All black males	1988	90%									40%
	In-school non-Hispanic black males				90%		92%		88%			
	c. All black females	1988	66%						3/48%			40%
	In-school non-Hispanic black females				84%		80%		75%		73%	
5.5*	Adolescent abstinence from sexual intercourse for previous 3 months											
	among adolescents who have ever engaged is sexual intercourse											
	All sexually experienced females 15-17 years	1988	24%						27%			40%
	In-school sexually experienced females 15-17 years			24%	25%		25%		23%		23%	
	All sexually experienced males 15-17 years	1988	33%						37%			40%
	In-school sexually experienced males 15-17 years			30%	36%		33%		34%		44%  42%  75%  62%  60%  73%	
5.6	Contraception use by sexually active adolescents											
	Female											
	First intercourse (15-19 years)	1988	65%						76%			90%
	Recent intercourse (15-19 years)	1988	78%						p/84%			90%
	Recent intercourse (In school, 15-17 years)			78%	81%		83%		83%		85%	
	Oral contraceptive and the condom at most											
	recent intercourse (15-19 years)	1988	2%						8%			90%
	Male											
	Contraception use at most recent intercourse (15-19 years)	1990	78%						79%			90%
	Contraception use at most recent intercourse											
	(In school, 15-17 years)				83%		84%		85%		81%	
	Birth control pills and condoms at most recent				0070		0170		0070		0.70	
	intercourse (In school, 15-17 years)	1990	a/2.0%		3.3%		2.9%		3.6%		4 8%	90%
	Condom and pill use at last intercourse (17-19 years)	1988	15%						16%			90%
	Condom and pill use at last intercourse		10,0						1070			0070
	(In school, 17 years and over)			14%							5%	
5.7	Failure of contraceptive method for females 15-44 years	1988	14%									7%
	a. Black female 15-44 years	1988	18%									8%
	b. Hispanic female 15-44 years	1988	16%									8%
5.8	Persons ages 13-18 years who have discussed sexuality with parents	1986	66%					4/73%	5/80%		44%  42%  75%  62%  85%  73%  32%  32%  85%  81% 4.8%  5%  5% 	85%
	Persons ages 10-17 who have discussed human sexuality											
	with parents, church or school							6/89%	7/98%			

		Baseline										Target
	Objective	year	Baseline	1990	1991	1992	1993	1994	1995	1996	1997	2000
5.9	Family planning counseling	1984	60%									90%
5.10*	Age-appropriate preconception counseling											
	by clinicians											60%
	Percent of clinicians routinely providing service											
	to 81-100% of patients											
	Inquiry about family planning											
	(female, childbearing age)											
	Pediatricians	1992	18%								8,9/	60%
	Nurse practitioners	1992	53%								8/42%	60%
	Obstetricians/gynecologists	1992	48%								8,9/	60%
	Internists	1992	24%								8,9/	60%
	Family physicians	1992	28%								8,9/	60%
	Counseling about family planning											
	Pediatricians	1992	36%								8,9/	60%
	Nurse practitioners	1992	53%								8/40%	60%
	Obstetricians/gynecologists	1992	65%								8,9/	60%
	Internists	1992	26%								8,9/	60%
	Family physicians	1992	36%								8,9/	60%
											-,	
5.11*	Clinic services for HIV and other sexually transmitted diseases	1000	400/									500
	Family planning clinics	1989	40%									50%
	Title X funded family planning clinics											
	STD testing (excluding HIV)							95%				
	STD counseling (excluding HIV)							98%				
	STD treatment (excluding HIV)							93%				
	Gonorrhea											
	Client testing/10			97%								
	Client treatment			82%								
	Partner notification/11			23%								
	Partner testing			60%								
	Partner treatment			62%								
	Syphilis											
	Client testing/10			86%								
	Client treatment			48%								
	Partner notification/11			29%								
	Partner testing			57%								
	Partner treatment			40%								
	Chlamydia											
	Client testing/10			66%								
	Client treatment			73%								
	Partner notification/11			15%								
	Partner testing			29%								
	Partner treatment			50%								
	HIV											
	Client pretest counseling			66%				82%				
	Client testing			60%				74%				
E 40												
5.12	Contraception use	1022	00 00/	12/00 19/					02 50/			050/
	Female 15-44 years	1982	88.2%	12/90.1%					92.5%			95%

	Baseline										Target
Objective	year	Baseline	1990	1991	1992	1993	1994	1995	1996	1997	2000
a. Black female 15-44 years	1982	78.9%	12/84.7%					89.9%			95%
b. Female 15-44 years under 100% poverty	1982	79.6%	12/80.2%					92.1%			95%
c. Female 15-19 years under 200% poverty	1982	67.4%	12/74.9%					84.8%			95%

---/Data not available.

.../Category not applicable.

a/Baseline has been revised.

p/Preliminary data.

1/Pregnancy rates are calculated from the number of births, fetal losses, and abortions.

2/Excludes data from States lacking an Hispanic-origin item on their birth or death certificate.

3/Non-Hispanic black female.

4/Data represent the proportion of people aged 10-17 who had discussed human sexuality with parents. Proportions for school and church were 76% and 32%, respectively.

5/1995 data are for females 18-19 years who have ever discussed birth control methods, how pregnancy occurs, or STDs

with a parent, or have had a sex education class on birth control methods, STDs, safe sex, or abstinence.

6/Data represent the proportion of people 10-17 years who had discussed human sexuality with parents or in church or school.

7/1995 data are for females 18-19 years who have ever discussed birth control methods, how pregnancy occurs, or STDs with a parent.

8/1997-98 data.

9/Response rate for this group was too low to produce reliable estimates.

10/Includes testing at initial visit, at annual visit, or if symptomatic.

11/By family planning clinic staff via telephone or mail.

12/1988 data.

NOTE: Data include revisions and, therefore, may differ from data previously published in these reports and other publications.

#### Data Sources:

Objective number	Data source
5.1, 5.1a, b	Abortion Provider Survey, Alan Guttmacher Institute;
	National Vital Statistics System, CDC, NCHS;
	National Survey of Family Growth, CDC, NCHS.
5.2, 5.2a, b	National Survey of Family Growth, CDC, NCHS.
5.3, 5.3a, b	National Survey of Family Growth, CDC, NCHS.
5.4*	Baseline and Update for all females and all black females: National Survey of Family Growth, CDC, NCHS.
	Baseline and Update for all males and all black males: National Survey of Adolescent Males, NIH, NICHD.
	1990-97 data for in-school females and males: Youth Risk Behavior Survey, CDC, NCCDPHP.
5.5*	Baseline and update for all females: National Survey of Family Growth, CDC, NCHS.
	Baseline for all males: National Survey of Adolescent Males, NIH, NICHD.
	1990-97 data for in-school females and males: Youth Risk Behavior Survey, CDC, NCCDPHP.
5.6	All females: National Survey of Family Growth, CDC, NCHS;
	All males: National Survey of Adolescent Males, NIH, NICHD;
	1990-97 data for in-school females and males: Youth Risk Behavior Survey, CDC, NCCDPHP.
5.7, 5.7a, b	National Survey of Family Growth, CDC, NCHS.
5.8	Baseline: Planned Parenthood Federation of America, Inc., 1986.
	1994 Update: National Health Interview Survey, CDC, NCHS.
	1995 Update: National Survey of Family Growth, CDC, NCHS.
5.9	Baseline: Mech EB. Unpublished. Orientation of Pregnancy Counselors toward Adoption. 1984.

	Baseline										Target
Objective	year	Baseline	1990	1991	1992	1993	1994	1995	1996	1997	2000
5.10*	Baseline: Primary Care Provider Surv	eys, OPHS, ODPH	P.								
	Update: Prevention in Primary Care S	tudy, American Co	llege of Prevent	ive Medicine.							
5.11*	1989 baseline: State Family Planning Directors.										
	1990 data: National Questionnaire on Provision of STD and HIV Services by Family Planning Clinics, PHS, OPA.										
	1994 data: The Urban Institute. Family Planning Clinics: Current Status and Recent Changes in Services,										
	Clients, Staffing and Income Source	s. March, 1994.									
5.12, 5.12a-c	Forrest, JD and Sing, S. The Sexual a	and Reproduction E	Behavior of Ame	erican Women,	1982-88, Fam	ily Planning Pe	erspectives 2	2(5):206-14,	1990.		
	1995 updates: National Survey of Fa	mily Growth, CDC,	NCHS.								

\*Duplicate objective. See full text of objective following this table.