Table 21. Number of deaths, death rates, and age-adjusted death rates for drug-induced causes, by race and sex: United States, 1999-2007

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." A listing of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition codes included in this table can be found in the note at the bottom of the table)

							American Indian or								
Year	All races			White ¹			Black ¹			Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both			Both			Both			Both			Both		
	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
							Nun	nber							
2007	38,371	23,883	14,488	33,480	20,795	12,685	4,194	2,708	1,486	388	205	183	309	175	134
2006	38,396	24,507	13,889	32,866	20,786	12,080	4,790	3,261	1,529	407	253	154	333	207	126
2005	33,541	21,208	12,333	28,804	18,152	10,652	4,098	2,677	1,421	362	205	157	277	174	103
2004	30,711	19,362	11,349	26,474	16,634	9,840	3,633	2,352	1,281	354	212	142	250	164	86
2003	28,723	18,426	10,297	24,683	15,824	8,859	3,527	2,303	1,224	295	168	127	218	131	87
2002	26,040	16,734	9,306	22,146	14,170	7,976	3,463	2,307	1,156	230	136	94	201	121	80
2001	21,705	14,253	7,452	18,195	11,882	6,313	3,165	2,163	1,002	184	100	84	161	108	53
2000	19,720	13,137	6,583	16,388	10,857	5,531	3,034	2,094	940	160	99	61	138	87	51
1999	19,128	12,885	6,243	15,714	10,506	5,208	3,100	2,191	909	164	96	68	150	92	58
							Ra	ate							
2007	12.7	16.1	9.5	13.7	17.2	10.3	10.5	14.2	7.1	12.0	12.7	11.3	2.1	2.4	1.8
2006	12.8	16.6	9.1	13.6	17.3	9.9	12.1	17.3	7.4	12.7	15.8	9.6	2.3	2.9	1.7
2005	11.3	14.5	8.2	12.0	15.3	8.8	10.5	14.3	7.0	11.5	13.0	9.9	2.0	2.5	1.4
2004	10.5	13.4	7.6	11.1	14.1	8.2	9.4	12.8	6.3	11.2	13.5	9.0	1.8	2.5	1.2
2003	9.9	12.9	7.0	10.4	13.5	7.4	9.2	12.7	6.1	9.5	10.8	8.2	1.7	2.0	1.3
2002	9.0	11.8	6.3	9.4	12.2	6.7	9.2	12.8	5.8	7.5	8.9	6.1	1.6	2.0	1.2
2001	7.6	10.2	5.1	7.8	10.4	5.4	8.5	12.2	5.1	6.0	6.6	5.5	1.3	1.8	0.8
2000	7.0	9.5	4.6	7.1	9.6	4.7	8.3	12.0	4.9	5.4	6.7	4.1	1.2	1.5	0.8
1999	6.9	9.4	4.4	6.9	9.3	4.5	8.6	12.7	4.8	5.8	6.8	4.8	1.3	1.7	1.0
							Age-adju	sted rate ⁴							
2007	12.6	15.8	9.3	13.6	16.9	10.2	11.0	15.4	7.3	12.1	12.6	11.5	2.0	2.3	1.7
2006	12.7	16.4	9.0	13.5	17.1	9.8	12.9	18.9	7.7	13.4	16.3	10.5	2.3	3.0	1.7
2005	11.3	14.4	8.1	11.9	15.1	8.7	11.2	15.8	7.2	11.9	13.4	10.4	1.9	2.6	1.4
2004	10.4	13.3	7.6	11.1	13.9	8.1	10.1	14.3	6.6	11.6	13.8	9.4	1.8	2.5	1.2
2003	9.9	12.8	7.0	10.4	13.4	7.4	9.9	14.1	6.4	9.9	11.2	8.6	1.6	1.9	1.3
2002	9.0	11.7	6.3	9.4	12.1	6.7	9.9	14.2	6.1	7.8	9.0	6.4	1.6	2.1	1.2
2001	7.6	10.1	5.1	7.8	10.2	5.3	9.2	13.6	5.4	6.6	7.3	5.8	1.3	1.8	0.8
2000	7.0	9.5	4.6	7.1	9.4	4.7	9.0	13.5	5.2	5.6	6.9	4.3	1.1	1.5	0.8
1999	6.8	9.4	4.4	6.8	9.2	4.4	9.3	14.3	5.1	6.1	7.2	5.0	1.4	1.7	1.0

¹Multiple-race data were reported by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

NOTE: Causes of death attributable to drug-induced mortality include ICD-10 codes D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F1

F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85, and Y10-Y14. In 2006, the list of drug-induced codes was modified to include two new ICD-10 codes, Drug-induced acute pancreatitis (ICD-10 code K85.3) and Drug-induced fever (ICD-10 code R50.2); see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴For method of computation, see "Technical Notes."