

Table 23. Number of deaths, death rates, and age-adjusted death rates for alcohol-induced causes, by race and sex: United States, 1999-2007

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys; see "Technical Notes." A listing of the *International Classification of Diseases, Tenth Revision* (ICD-10), Second Edition codes included in this table can be found in the note at the bottom of the table]

Year	All races			White ¹			Black ¹			American Indian or Alaska Native ^{1,2}			Asian or Pacific Islander ^{1,3}		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Number															
2007	23,199	17,428	5,771	19,921	15,027	4,894	2,248	1,703	545	774	495	279	256	203	53
2006	22,073	16,472	5,601	18,917	14,186	4,731	2,172	1,599	573	717	458	259	267	229	38
2005	21,634	16,238	5,396	18,432	13,917	4,515	2,316	1,698	618	655	432	223	231	191	40
2004	21,081	15,906	5,175	17,875	13,525	4,350	2,351	1,784	567	645	417	228	210	180	30
2003	20,687	15,630	5,057	17,437	13,218	4,219	2,406	1,824	582	664	431	233	180	157	23
2002	20,218	15,272	4,946	16,988	12,926	4,062	2,434	1,798	636	597	387	210	199	161	38
2001	20,114	15,149	4,965	16,640	12,588	4,052	2,723	2,048	675	591	387	204	160	126	34
2000	19,643	14,993	4,650	16,223	12,509	3,714	2,712	1,993	719	537	353	184	171	138	33
1999	19,469	14,894	4,575	15,903	12,277	3,626	2,832	2,100	732	589	397	192	145	120	25
Rate															
2007	7.7	11.7	3.8	8.2	12.4	4.0	5.6	8.9	2.6	23.9	30.6	17.2	1.7	2.8	0.7
2006	7.4	11.2	3.7	7.8	11.8	3.9	5.5	8.5	2.8	22.4	28.6	16.2	1.8	3.2	0.5
2005	7.3	11.1	3.6	7.7	11.7	3.7	5.9	9.1	3.0	20.7	27.4	14.1	1.6	2.8	0.6
2004	7.2	11.0	3.5	7.5	11.5	3.6	6.1	9.7	2.8	20.5	26.5	14.5	1.5	2.7	0.4
2003	7.1	10.9	3.4	7.4	11.3	3.5	6.3	10.0	2.9	21.3	27.8	15.0	1.4	2.4	0.3
2002	7.0	10.8	3.4	7.2	11.1	3.4	6.4	10.0	3.2	19.4	25.2	13.6	1.6	2.6	0.6
2001	7.1	10.8	3.4	7.2	11.0	3.4	7.3	11.6	3.5	19.3	25.4	13.3	1.3	2.1	0.5
2000	7.0	10.9	3.2	7.1	11.0	3.2	7.4	11.4	3.7	18.0	23.7	12.3	1.5	2.4	0.5
1999	7.0	10.9	3.2	7.0	10.9	3.1	7.8	12.2	3.9	20.8	28.1	13.5	1.3	2.2	0.4
Age-adjusted rate ⁴															
2007	7.3	11.3	3.5	7.5	11.6	3.6	6.3	11.0	2.7	26.3	35.1	18.2	1.8	3.0	0.7
2006	7.0	10.9	3.4	7.2	11.2	3.5	6.2	10.4	3.0	25.2	33.3	17.8	1.9	3.5	0.5
2005	7.0	11.0	3.4	7.2	11.1	3.4	6.8	11.4	3.3	23.7	32.5	15.6	1.7	3.1	0.6
2004	7.0	11.0	3.3	7.1	11.0	3.3	7.2	12.3	3.1	23.8	32.4	16.1	1.7	3.2	0.5
2003	7.0	11.0	3.3	7.0	11.0	3.3	7.4	12.8	3.3	25.1	34.0	16.9	1.5	2.8	0.4
2002	6.9	11.0	3.3	6.9	10.9	3.2	7.8	13.1	3.6	23.2	31.5	15.6	1.8	3.2	0.6
2001	7.0	11.2	3.3	6.9	10.9	3.3	8.9	15.1	3.9	23.6	33.0	15.4	1.5	2.6	0.6
2000	7.0	11.4	3.2	6.9	11.1	3.0	9.1	15.3	4.3	22.7	31.4	14.9	1.7	2.9	0.7
1999	7.1	11.5	3.2	6.8	11.0	3.0	9.8	16.7	4.5	26.7	38.8	16.2	1.6	2.8	0.5

¹Multiple-race data were reported by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2005, by 15 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

²Includes Aleuts and Eskimos.

³Includes Chinese, Filipino, Hawaiian, Japanese, and Other Asian or Pacific Islander.

⁴For method of computation, see "Technical Notes."

NOTE: Causes of death attributable to alcohol-induced mortality include ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. In 2006, the list of alcohol-induced codes was modified to include a new ICD-10 code, Alcohol-induced acute pancreatitis (ICD-10 code K85.2); see "Technical Notes."