Table 24. Number of deaths, death rates, and age-adjusted death rates for alcohol-induced causes, by Hispanic origin, race for non-Hispanic population, and sex: United States, 1999-2007

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates per 100,000 U.S. standard population; see "Technical Notes." Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see "Technical Notes." Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see "Technical Notes." Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. A listing of the International Classification of Diseases, Tenth Revision (ICD-10), Second Edition codes included in this table can be found in the note at the bottom of the table]

	All origins ¹			Hispanic			Non-Hispanic ²			Non-Hispanic white ³			Non-Hispanic black ³		
	Both			Both			Both			Both			Both		
Year	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female	sexes	Male	Female
								Number							
2007	23,199	17,428	5,771	2,977	2,539	438	20,130	14,811	5,319	16,935	12,474	4,461	2,210	1,674	536
2006	22,073	16,472	5,601	2,804	2,341	463	19,187	14,055	5,132	16,114	11,837	4,277	2,128	1,561	567
2005	21,634	16,238	5,396	2,658	2,265	393	18,877	13,890	4,987	15,729	11,610	4,119	2,282	1,670	612
2004	21,081	15,906	5,175	2,406	2,056	350	18,567	13,761	4,806	15,418	11,428	3,990	2,318	1,754	564
2003	20,687	15,630	5,057	2,422	2,048	374	18,160	13,490	4,670	14,977	11,133	3,844	2,367	1,787	580
2002	20,218	15,272	4,946	2,408	2,065	343	17,661	13,078	4,583	14,494	10,783	3,711	2,396	1,768	628
2001	20,114	15,149	4,965	2,381	2,026	355	17,593	13,009	4,584	14,186	10,497	3,689	2,677	2,016	661
2000	19,643	14,993	4,650	2,323	2,024	299	17,177	12,843	4,334	13,815	10,408	3,407	2,672	1,959	713
1999	19,469	14,894	4,575	2,184	1,864	320	17,143	12,905	4,238	13,633	10,337	3,296	2,794	2,066	728
								Rate							
2007	7.7	11.7	3.8	6.5	10.8	2.0	7.9	11.8	4.1	8.4	12.6	4.4	5.8	9.2	2.7
2006	7.4	11.2	3.7	6.3	10.2	2.2	7.5	11.3	3.9	8.0	12.0	4.2	5.6	8.7	2.9
2005	7.3	11.1	3.6	6.2	10.3	1.9	7.4	11.2	3.8	7.9	11.8	4.0	6.1	9.4	3.1
2004	7.2	11.0	3.5	5.8	9.6	1.8	7.4	11.2	3.7	7.7	11.7	3.9	6.3	10.0	2.9
2003	7.1	10.9	3.4	6.1	9.9	1.9	7.2	11.0	3.6	7.5	11.4	3.8	6.5	10.3	3.0
2002	7.0	10.8	3.4	6.2	10.3	1.8	7.1	10.7	3.6	7.3	11.1	3.7	6.6	10.3	3.3
2001	7.1	10.8	3.4	6.4	10.7	2.0	7.1	10.8	3.6	7.2	10.8	3.6	7.5	11.9	3.5
2000	7.0	10.9	3.2	6.6	11.1	1.7	7.0	10.7	3.4	7.0	10.8	3.4	7.6	11.7	3.9
1999	7.0	10.9	3.2	6.4	10.7	1.9	7.0	10.8	3.4	6.9	10.7	3.3	8.0	12.5	4.0
							Age	-adjusted ra	ıte ⁴						
2007	7.3	11.3	3.5	9.3	16.4	2.7	7.0	10.8	3.6	7.2	10.9	3.8	6.5	11.2	2.8
2006	7.0	10.9	3.4	9.1	15.7	3.0	6.8	10.4	3.5	6.9	10.5	3.6	6.3	10.5	3.0
2005	7.0	11.0	3.4	9.1	16.2	2.6	6.8	10.4	3.5	6.8	10.4	3.5	7.0	11.6	3.4
2004	7.0	11.0	3.3	8.6	15.1	2.5	6.8	10.6	3.4	6.8	10.5	3.4	7.3	12.5	3.2
2003	7.0	11.0	3.3	9.2	16.2	2.8	6.8	10.5	3.3	6.7	10.4	3.4	7.6	12.9	3.3
2002	6.9	11.0	3.3	9.5	17.0	2.7	6.7	10.4	3.3	6.6	10.2	3.3	7.9	13.3	3.7
2001	7.0	11.2	3.3	10.1	18.1	2.9	6.7	10.5	3.4	6.5	10.1	3.3	9.0	15.4	4.0
2000	7.0	11.4	3.2	10.5	19.4	2.6	6.7	10.6	3.2	6.4	10.1	3.1	9.3	15.5	4.4
1999	7.1	11.5	3.2	10.3	18.6	3.0	6.8	10.8	3.2	6.4	10.2	3.0	10.0	16.9	4.6

¹Figures for origin not stated are included in "All origins" but not distributed among specified origins.

NOTE: Causes of death attributable to alcohol-induced mortality include ICD-10 codes E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65, and Y15. In 2006, the list of alcohol-induced codes was modified to include a new ICD-10 code, Alcohol-induced acute pancreatitis (ICD-10 code K85.2).

²Includes races other than white and black.

³Multiple-race data were reported by 27 states and the District of Columbia in 2007, by 25 states and the District of Columbia in 2006, by 21 states and the District of Columbia in 2004, and by 7 states in 2004, and by 7 states in 2003; see "Technical Notes." The multiple-race data for these reporting areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other reporting areas; see "Technical Notes."

⁴For method of computation, see "Technical Notes."