USC REPRODUCTIVE ENDOCRINOLOGY AND INFERTILITY LOS ANGELES, CALIFORNIA

Comparison of success rates across clinics may not be meaningful. Patient medical characteristics and treatment approaches vary (see pages 11-20).

| 2015 ART CYCLE PR | - |
|-------------------|-------|
| ZUIS ABI UTUIE PR | - |

Data verified by Richard J. Paulson, MD

| Type of ART and Procedural Factors ^a | | | | Patient Diagnosis b | | | | |
|---|------|-----------|-----|----------------------------|-----|----------------|-----|---------------------------|
| IVF | 100% | With ICSI | 85% | Tubal factor | 5% | Uterine factor | 5% | Multiple Factors: |
| Unstimulated | 2% | PGD/PGS | <1% | Ovulatory dysfunction | 12% | Male factor | 56% | Female factors only 9% |
| Used gestational carrier | 2% | | | Diminished ovarian reserve | 60% | Other factor | 15% | Female & male factors 42% |
| | | | | Endometriosis | 5% | Unknown factor | 3% | |

| 2015 | ADT CII | CCECC | DATECC |
|------|---------|-------|--------|

Total number of cycles^d: 705 (includes 18 cycle[s] using frozen eggs)

| 2015 ART SUCCESS RATES | Total number of cycles 705 | (include | es 18 cycle | | | ggs) | |
|---|---|----------------------------|-------------|------------|-----------------------------|--------------|-----|
| Time of Cuelo | | Age of Woman | | | | | |
| Type of Cycle | | <35 | 35-37 | 38-40 | 41-42 | 43-44 | >44 |
| Fresh Embryos from Nondonor Eggs | | | | | | | |
| Number of cycles | | 62 | 41 | 70 | 43 | 23 | 8 |
| Percentage of cancellations before retrieval (%) | | 3.2 | 2.4 | 8.6 | 25.6 | 8.7 | 1/8 |
| Average number of embryos transferred | | 1.8 | 2.4 | 2.9 | 3.5 | 4.0 | 4.1 |
| Percentage of embryos transferred resulting in | implantation (%) | 33.0 | 16.7 | 17.2 | 5.6 | 1.3 | 0.0 |
| Percentage of elective single embryo transfers | | 16.7 | 5.4 | 1.9 | 0.0 | 0/16 | 0/6 |
| Outcomes per Cycle | (/ (/ | | | | | | |
| Percentage of cycles resulting in term, normal v | veight & singleton live births ^e (%) | 24.2 | 19.5 | 4.3 | 7.0 | 4.3 | 0/8 |
| Percentage of cycles resulting in singleton live I | | 25.8 | 22.0 | 5.7 | 7.0 | 4.3 | 0/8 |
| Percentage of cycles resulting in twin live births | | 12.9 | 4.9 | 8.6 | 2.3 | 0.0 | 0/8 |
| Percentage of cycles resulting in live births (%) | | 38.7 | 26.8 | 17.1 | 9.3 | 4.3 | 0/8 |
| Percentage of cycles resulting in pregnancies (| %) | 45.2 | 31.7 | 25.7 | 16.3 | 4.3 | 0/8 |
| Outcomes per Transfer | | | | | | | |
| Number of transfers | | 57 | 39 | 57 | 32 | 20 | 7 |
| Percentage of transfers resulting in term, norma | al weight & singleton live births ^e (%) | 26.3 | 20.5 | 5.3 | 9.4 | 5.0 | 0/7 |
| Percentage of transfers resulting in singleton liv | | 28.1 | 23.1 | 7.0 | 9.4 | 5.0 | 0/7 |
| Percentage of transfers resulting in twin live bird | ths (%) | 14.0 | 5.1 | 10.5 | 3.1 | 0.0 | 0/7 |
| Percentage of transfers resulting in live births (9 | %) | 42.1 | 28.2 | 21.1 | 12.5 | 5.0 | 0/7 |
| Percentage of transfers resulting in pregnancies | s (%) | 49.1 | 33.3 | 31.6 | 21.9 | 5.0 | 0/7 |
| Frozen Embryos from Nondonor Eggs | | | | | | | |
| Number of cycles | | 50 | 47 | 43 | 16 | 10 | 2 |
| Number of transfers | | 50 | 47 | 43 | 16 | 10 | 2 |
| Estimated average number of transfers per retri | eval | 0.6 | 0.6 | 0.6 | 0.4 | 0.5 | 0.3 |
| Average number of embryos transferred | Cvai | 1.7 | 1.6 | 2.0 | 2.8 | 3.8 | 3.0 |
| Percentage of embryos transferred resulting in | implantation (%) | 30.5 | 27.0 | 20.0 | 13.6 | 2.6 | 1/6 |
| Percentage of transfers resulting in term, norma | | 18.0 | 27.7 | 18.6 | 3 / 16 | 0/10 | 0/2 |
| Percentage of transfers resulting in singleton liv | | 22.0 | 27.7 | 18.6 | 3 / 16 | 0/10 | 0/2 |
| Percentage of transfers resulting in twin live bird | | 10.0 | 6.4 | 7.0 | 1 / 16 | 0/10 | 0/2 |
| Percentage of transfers resulting in live births (9 | · · | 34.0 | 34.0 | 25.6 | 4 / 16 | 0/10 | 0/2 |
| | | 40.0 | 38.3 | 37.2 | 5/16 | 1/10 | 1/2 |
| Number of Egg/Embryo Banking Cycle | | 58 | 63 | 61 | 35 | 19 | 6 |
| Donor Fage | | Fresh Embryos ^f | | | Frozen Embryos ^f | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | |
| Percentage of transfers resulting in pregnancies | implantation (%) al weight & singleton live births e (%) the births (%) ths (%) | 40.0 58 | 38.3 63 | 37.2 61 | 5 / 16 35 | 1 / 10 19 | 1/2 |

CURRENT SERVICES & PROFILE

Current Name: USC Reproductive Endocrinology and Infertility

| Donor eggs? | Yes | Gestational carriers? | Yes | Single women? | Yes | Verified lab accreditation? | Yes |
|----------------|-----|--------------------------|-----|---------------|-----|-------------------------------|-----|
| Donor embryos? | Yes | Embryo cryopreservation? | Yes | SART member? | Yes | (See Appendix C for details.) | |

a Reflects features of fresh nondonor cycles. If IVF is less than 100%, the remaining cycles are GIFT, ZIFT, or a combination of these procedures with IVF.

^b Total patient diagnosis percentages may be greater than 100% because more than one diagnosis can be reported for each ART cycle.

^c Fractions are used for rates with denominators less than 20. Multiple-infant births (for example, twins) with at least one live infant are counted as one live birth.

d Total cycle number includes those using frozen eggs. It excludes 0 cycle(s) evaluating new procedures. Both cycle types are excluded from ART success rates.

e In this report, births are defined as term if at least 37 full weeks gestation and normal birth weight if at least 2,500 grams (approximately 5 pounds, 8 ounces).

f All ages are reported together because previous data show that patient age does not materially affect success with donor eggs.