JONES INSTITUTE FOR REPRODUCTIVE MEDICINE **NORFOLK, VIRGINIA**

Comparing success rates across clinics may not be meaningful. Patient medical characteristics and treatment approaches vary (see pages 11-21).

2016 ART CYCLE PROFILE

Data verified by Laurel A. Stadtmauer, MD, PhD

| Type of ART and Procedural Factors ^a | | | | Patient Diagnosis a,b | | | | | |
|---|------|-----------|-----|----------------------------|-----|----------------|-----|-----------------------|----|
| IVF | 100% | With ICSI | 73% | Tubal factor | 20% | Uterine factor | 4% | Multiple Factors: | |
| Unstimulated | 0% | PGD/PGS | 5% | Ovulatory dysfunction | 21% | Male factor | 23% | Female factors only | 9% |
| Used gestational carrier | 1% | | | Diminished ovarian reserve | 27% | Other factor | 8% | Female & male factors | 7% |
| | | | | Endometriosis | 5% | Unknown factor | 7% | | |

| 2016 ART SUCCESS RATES c,d (in | tal number of cycles ^d : 356 cludes 0 cycle[s] using fresh embr | yos from fr | ozen nondor | nor eggs) | | | | |
|--|---|----------------|----------------|-----------|---------------|-----------------|--|--|
| T (0.1 | | - | Age of Patient | | | | | |
| Type of Cycle | | <35 | 35-37 | 38-40 | 41-42 | >42 | | |
| Fresh Embryos from Fresh Nondonor Egg | IS | | | | | | | |
| Number of cycles | | 105 | 33 | 27 | 9 | 7 | | |
| Percentage of cancellations before retrieval (%) | | 8.6 | 15.2 | 33.3 | 3/9 | 3/7 | | |
| Number of transfers | | 69 | 27 | 13 | 2 | 3 | | |
| Average number of embryos transferred | | 1.9 | 1.7 | 1.8 | 2.0 | 2.3 | | |
| Percentage of elective single embryo transfers (eSE | T) (%) | 7.6 | 9.1 | 1 / 10 | 0/1 | 0/3 | | |
| Outcomes per Cycle | | | | | | | | |
| Percentage of cycles resulting in pregnancies (%) | | 38.1 | 33.3 | 22.2 | 2/9 | 1/7 | | |
| Percentage of cycles resulting in live births (%) | | 32.4 | 30.3 | 11.1 | 1/9 | 1/7 | | |
| Percentage of cycles resulting in singleton live birth | s (%) | 21.9 | 15.2 | 7.4 | 1/9 | 1/7 | | |
| Percentage of cycles resulting in twin live births (%) | | 9.5 | 15.2 | 3.7 | 0/9 | 0/7 | | |
| Percentage of cycles resulting in term, normal weight | nt and singleton live births (%) | 17.1 | 12.1 | 3.7 | 0/9 | 1/7 | | |
| Outcomes per Transfer | | | | | | | | |
| Percentage of embryos transferred resulting in imple | | 43.4 | 34.0 | 33.3 | 2/4 | 3/7 | | |
| Percentage of transfers resulting in pregnancies (%) | | 58.0 | 40.7 | 6 / 13 | 2/2 | 1/3 | | |
| Percentage of transfers resulting in live births (%) | | 49.3 | 37.0 | 3 / 13 | 1/2 | 1/3 | | |
| Percentage of transfers resulting in singleton live bir | | 33.3 | 18.5 | 2 / 13 | 1/2 | 1/3 | | |
| Percentage of transfers resulting in twin live births (| | 14.5 | 18.5 | 1 / 13 | 0/2 | 0/3 | | |
| Percentage of transfers resulting in term, normal we | eight and singleton live births (%) | 26.1 | 14.8 | 1 / 13 | 0/2 | 1/3 | | |
| Frozen Embryos from Nondonor Eggs | | | | | | | | |
| Number of cycles | | 49 | 15 | 14 | 4 | 7 | | |
| Number of transfers | | 49 | 15 | 14 | 4 | 6 | | |
| Estimated average number of transfers per retrieval | | 1.3 | 1.5 | 1.2 | 0.6 | 3.0 | | |
| Average number of embryos transferred | | 1.7 | 1.9 | 1.4 | 1.5 | 1.8 | | |
| Percentage of embryos transferred resulting in imple | antation (%) | 32.9 | 17.9 | 10 / 19 | 2/6 | 2/11 | | |
| Percentage of transfers resulting in pregnancies (%) | | 44.9 | 5 / 15 | 7 / 14 | 2/4 | 2/6 | | |
| Percentage of transfers resulting in live births (%) | | 38.8 | 5 / 15 | 7 / 14 | 1/4 | 2/6 | | |
| Percentage of transfers resulting in singleton live bir | ths (%) | 28.6 | 5 / 15 | 4 / 14 | 1/4 | 2/6 | | |
| Percentage of transfers resulting in twin live births (| | 10.2 | 0 / 15 | 3 / 14 | 0/4 | 0/6 | | |
| Percentage of transfers resulting in term, normal we | eight and singleton live births ^e (%) | 20.4 | 3 / 15 | 4 / 14 | 0/4 | 1/6 | | |
| Number of Egg or Embryo Banking Cycle | s | 13 | 6 | 7 | 7 | 2 | | |
| Number of fertility preservation cycles | | 2 | 2 | 1 | 0 | 0 | | |
| rumbo. or rommy process rumon cycles | | Fresh | | - | ozen | Donated | | |
| Donor Eggs ^f | | | Froze | | ozen bryos | Embryos | | |
| Number of cycles | | Eggs 20 | Eggs | | 10 | 20 | | |
| Number of transfers | | 18 | 1 | | 9 | 17 | | |
| Average number of embryos transferred | | 1.9 | 2.0 | | 1.7 | 1.7 | | |
| Percentage of embryos transferred resulting in imple | 38.2 | 2.0 | | / 12 | 42.3 | | | |
| Percentage of embryos transferred resulting in Imple Percentage of transfers resulting in pregnancies (%) | * * | 9 / 18 | 1/1 | | / 12 5/9 | 42.3 11 / 17 | | |
| Percentage of transfers resulting in live births (%) | | 9/18 | 1/1 | | 2/9 | 7/17 | | |
| Percentage of transfers resulting in five births (70) | ths (%) | 6/18 | 0/1 | | 2/9 | 5 / 17 | | |
| Percentage of transfers resulting in twin live births (| | 3 / 18 | 1/1 | |)/9 | 2/17 | | |
| Percentage of transfers resulting in term, normal we | | 5 / 18 | 0/1 | | 2/9 | 4/17 | | |
| . 1.13. Lago of transcript fooditing in torni, normal we | .g (70) | 0, 10 | 0,1 | | | ., ., | | |

CURRENT SERVICES & PROFILE

Current Name: Jones Institute for Reproductive Medicine

| Donor eggs? | Yes | Embryo cryopreservation? | Yes | Single women? | Yes | SART member? | Yes |
|----------------|-----|--------------------------|-----|-----------------------|-----|-----------------------------|-----|
| Donor embryos? | Yes | Egg cryopreservation? | Yes | Gestational carriers? | Yes | Verified lab accreditation? | Yes |

^a Excludes cycles evaluating new procedures and banking cycles; unstimulated percentage includes fresh egg cycles only.

b Total patient diagnosis percentages may be greater than 100% because more than one diagnosis can be reported for each ART cycle.

^c Fractions are used for rates with denominators less than 20. Multiple-infant births (for example, twins) with at least one live infant are counted as one live birth.

d Total cycle number and success rates exclude 0 cycle(s) evaluating new procedures. Success rates exclude cycles using fresh embryos from frozen nondonor eggs.

e In this report, births are defined as term if at least 37 full weeks gestation and normal birth weight if at least 2,500 grams (approximately 5 pounds, 8 ounces).

f All ages are reported together because previous data show that patient age does not materially affect success with donor eggs.